

Polynomial realizations of natural Hopf algebras of nonsymmetric operads

Samuele Giraudo^{*1}

¹ *Université du Québec à Montréal, LACIM, Pavillon Président-Kennedy, 201 Avenue du Président-Kennedy, Montréal, H2X 3Y7, Canada.*

Abstract. The natural Hopf algebra $\mathbf{N} \cdot \mathcal{O}$ of an operad \mathcal{O} is a Hopf algebra whose bases are indexed by some words on \mathcal{O} . We introduce new bases of these Hopf algebras deriving from free operads via new lattice structures on their basis elements. We construct polynomial realizations of $\mathbf{N} \cdot \mathcal{O}$ by using alphabets of variables endowed with unary and binary relations. By specializing our polynomial realizations, we discover links between $\mathbf{N} \cdot \mathcal{O}$ and the Hopf algebra of word quasi-symmetric functions of Hivert, the decorated versions of the Connes-Kreimer Hopf algebra of Foissy, the Faà di Bruno Hopf algebra and its deformations, and the noncommutative multi-symmetric functions Hopf algebras of Novelli and Thibon.

Keywords: Combinatorial Hopf algebra; Operad; Polynomial realization; Word quasi-symmetric function; Connes-Kreimer Hopf algebra; Faà di Bruno Hopf algebra.

Introduction

A polynomial realization (abbreviated as *PR*) of a Hopf algebra consists of interpreting its elements as polynomials, either commutative or not, in such a way that its product translates as polynomial multiplication and the coproduct translates as a simple transformation of the alphabet of variables. A great portion of combinatorial Hopf algebras appearing in combinatorics admit PRs [3, 17, 7, 6]. It is striking to note that Hopf algebras involving a variety of different families of combinatorial objects and operations on them can be translated and understood in a common manner through adequate PRs.

Such realizations are crucial for several reasons. First, they allow us to prove easily that the Hopf algebra axioms (like associativity, coassociativity, and the Hopf compatibility between the product and the coproduct) are satisfied. Indeed, if a Hopf algebra admits a PR, such properties are almost immediate on polynomials [12]. Second, given a PR of a Hopf algebra, it is in most cases fruitful to specialize the associated polynomials (for instance by letting the variables commute) in order to get Hopf algebra morphisms to other Hopf algebras. This leads to the construction of new Hopf algebras

^{*}giraudo.samuele@uqam.ca. This research has been partially supported by the projects CARPLO (ANR-20-CE40-0007) and ALCOHOL (ANR-19-CE40-0006) of the Agence nationale de la recherche.

or establishes links between already existing ones [7]. Finally, such families of polynomials realizing a Hopf algebra lead to the definition of new classes of special functions, analogous to Schur or Macdonald functions [3].

Under right conditions, an operad \mathcal{O} produces a Hopf algebra $\mathbf{N} \cdot \mathcal{O}$, called the *natural Hopf algebra* of \mathcal{O} . The bases of $\mathbf{N} \cdot \mathcal{O}$ are indexed by some words on \mathcal{O} , and its coproduct is inherited from the composition map of \mathcal{O} . This construction is considered in [14, 2, 1], and we focus here on a noncommutative variation for nonsymmetric operads [16]. In contrast to many examples of Hopf algebras having PRs, none are known for $\mathbf{N} \cdot \mathcal{O}$. The main contribution of this work is to provide a PR for this family of Hopf algebras built from nonsymmetric operads. The particularity of our approach is that we consider a PR based on variables belonging to alphabets endowed with several unary and binary relations in order to capture the particularities of the coproduct of $\mathbf{N} \cdot \mathcal{O}$. This approach, using what we call *related alphabets*, generalizes the previous approaches using totally ordered alphabets [3, 17, 12], quasi-ordered alphabets [6], or alphabets endowed with a single binary relation [7, 8].

This work is presented as follows. In Section 1, the main notions about natural Hopf algebras of operads, PRs, and free operads are provided. We introduce in Section 2 two new bases of $\mathbf{N} \cdot \mathcal{O}$ in the case where \mathcal{O} is a free operad. These constructions use new lattices on treelike structures. In Section 3, we describe an injection of $\mathbf{N} \cdot \mathcal{O} / \equiv$ into $\mathbf{N} \cdot \mathcal{O}$ when \mathcal{O} is a free operad and \equiv is an operad congruence subjected to some properties. Section 4 contains a PR of the natural Hopf algebra of a free operad \mathcal{O} and of some of its quotients. In the subsequent Sections 5, 6, 7, and 8, we present some consequences of our PR, leading to links between $\mathbf{N} \cdot \mathcal{O}$ and other Hopf algebras. In particular, we show that a specialization of the PR of $\mathbf{N} \cdot \mathcal{O}$ forms a Hopf subalgebra of **WQSym**, the Hopf algebra word quasi-symmetric functions [11, 17]. We provide, by building a free operad \mathcal{O}_D depending on a set D of decorations, a construction of the Hopf algebra of D -decorated forests [4] as a specialization of the PR of $\mathbf{N} \cdot \mathcal{O}_D$. We show that the noncommutative Faà di Bruno **FdB** Hopf algebra and its one-parameter deformation **FdB_r** [5] can be built from the multi-multiassociative operads $\mathbf{MA}_{\mathcal{S}}$, extending a construction of [9]. Finally, we build from $\mathbf{MA}_{\mathcal{S}}$ a Hopf algebra **FdB_{r,s}** depending on two parameters such that **FdB_{0,s}** is the noncommutative multi-symmetric function Hopf algebra [18] and **FdB_{r,0}** is **FdB_r**. By using our framework, we obtain PRs of these Hopf algebras.

General notations and conventions. If f is an entity parameterized by an input x , we write $f \cdot x$ for $f(x)$. As \cdot associates from right to left, $f \cdot g \cdot x$ denotes $f \cdot (g \cdot x)$. For any statement P , the Iverson bracket $[P]$ takes 1 as value if P is true and 0 otherwise. For any integer i , $[i]$ denotes the set $\{1, \dots, i\}$. For any set A , A^* is the set of words on A . For any $w \in A^*$, $\ell \cdot w$ is the length of w , and for any $i \in [\ell \cdot w]$, $w \cdot i$ is the i -th letter of w . The only word of length 0 is the empty word ϵ . For any subset A' of A , $w|_{A'}$ is the subword of w made of letters of A' . Given two words w and w' , the concatenation of w and w' is denoted by ww' or by $w \cdot w'$.

1 Natural Hopf algebras from operads

1.1 Natural Hopf algebras from nonsymmetric operads

We follow the usual notations about nonsymmetric operads [10] (called simply *operads* here). Let \mathcal{O} be an operad with arity map ar , composition map $-[-, \dots, -]$, and unit $\mathbb{1}$. When each $x \in \mathcal{O}$ admits finitely many factorizations $x = y[z_1, \dots, z_{\text{ar} \cdot y}]$ where $y, z_1, \dots, z_{\text{ar} \cdot y} \in \mathcal{O}$, \mathcal{O} is *finitely factorizable*. When there exists a map $\text{dg} : \mathcal{O} \rightarrow \mathbb{N}$ such that $\text{dg}^{-1} \cdot 0 = \{\mathbb{1}\}$ and, for any $y, z_1, \dots, z_{\text{ar} \cdot y} \in \mathcal{O}$, $\text{dg} \cdot y[z_1, \dots, z_{\text{ar} \cdot y}] = \text{dg} \cdot y + \text{dg} \cdot z_1 + \dots + \text{dg} \cdot z_{\text{ar} \cdot y}$, the map dg is a *grading* of \mathcal{O} .

The *natural Hopf algebra* [14, 2, 16, 1] of a finitely factorizable operad \mathcal{O} admitting a grading dg is the Hopf algebra $\mathbf{N} \cdot \mathcal{O}$ defined as follows. Let rd be the map such that $\text{rd} \cdot w$ is the subword of $w \in \mathcal{O}^*$ consisting of its letters different from $\mathbb{1}$. A word w on \mathcal{O} is *reduced* if $\text{rd} \cdot w = w$. Let $\mathbf{N} \cdot \mathcal{O}$ be the \mathbb{K} -linear span of the set $\text{rd} \cdot \mathcal{O}^*$, where \mathbb{K} is any field of characteristic zero. The bases of $\mathbf{N} \cdot \mathcal{O}$ are thus indexed by $\text{rd} \cdot \mathcal{O}^*$, and the *elementary basis* (or *E-basis* for short) of $\mathbf{N} \cdot \mathcal{O}$ is the set $\{E_w : w \in \text{rd} \cdot \mathcal{O}^*\}$. This vector space is endowed with an associative algebra structure through the product \cdot satisfying $E_w \cdot E_{w'} = E_{w \cdot w'}$. Moreover, $\mathbf{N} \cdot \mathcal{O}$ is endowed with the coproduct Δ defined as the unique associative algebra morphism satisfying, for any $x \in \mathcal{O}$,

$$\Delta \cdot E_x = \sum_{y \in \mathcal{O}} \sum_{w \in \mathcal{O}^{\text{ar} \cdot y}} [x = y[w \cdot \mathbb{1}, \dots, w \cdot \ell \cdot w]] E_{\text{rd} \cdot y} \otimes E_{\text{rd} \cdot w}, \quad (1.1)$$

where the outer $[-]$ denotes the Iverson bracket (see *General notations and conventions*). Due to the fact that \mathcal{O} is finitely factorizable, (1.1) is a finite sum. This coproduct endows $\mathbf{N} \cdot \mathcal{O}$ with the structure of a bialgebra. By extending additively dg on \mathcal{O}^* , the map dg defines a grading of $\mathbf{N} \cdot \mathcal{O}$. Thus, $\mathbf{N} \cdot \mathcal{O}$ admits an antipode and becomes a Hopf algebra.

For instance, let us consider the *associative operad* As , defined by $\text{As} := \{\alpha_n : n \geq 1\}$, $\text{ar} \cdot \alpha_n := n$, $\alpha_n[\alpha_{m_1}, \dots, \alpha_{m_n}] := \alpha_{m_1 + \dots + m_n}$, and $\mathbb{1} := \alpha_1$. The map dg satisfying $\text{dg} \cdot \alpha_n = n - 1$ is a grading of As . The bases of $\mathbf{N} \cdot \text{As}$ are indexed by $\text{rd} \cdot \text{As}^*$ and we have, e.g.,

$$\Delta \cdot E_{\alpha_4} = E_\epsilon \otimes E_{\alpha_4} + 2E_{\alpha_2} \otimes E_{\alpha_3} + E_{\alpha_2} \otimes E_{\alpha_2 \alpha_2} + 3E_{\alpha_3} \otimes E_{\alpha_2} + E_{\alpha_4} \otimes E_\epsilon. \quad (1.2)$$

It is shown in [1] that $\mathbf{N} \cdot \text{As}$ is isomorphic to the *noncommutative Faà di Bruno Hopf algebra FdB* (see [5]).

1.2 Polynomial realizations

For any alphabet A , $\mathbb{K}\langle A \rangle$ is the \mathbb{K} -vector space of *A-polynomials*, which are noncommutative polynomials with variables in A , having a possibly infinite support but finite degree. Given two alphabets A and A' , let $\theta_{A, A'} : \mathbb{K}\langle A \sqcup A' \rangle \rightarrow \mathbb{K}\langle A \rangle \otimes \mathbb{K}\langle A' \rangle$ be the linear map such that $\theta_{A, A'} \cdot w = w|_A \otimes w|_{A'}$ for any $w \in (A \sqcup A')^*$.

A *related alphabet* is a set A endowed with a family of n -ary relations, $n \geq 1$. As usual, an n -ary relation \mathcal{R} is a subset of A^n , and we denote by $\mathcal{R}(a_1, \dots, a_n)$ the fact that $(a_1, \dots, a_n) \in \mathcal{R}$, and when $n = 2$, we write $a_1 \mathcal{R} a_2$ instead of $\mathcal{R}(a_1, a_2)$. A *polynomial realization* of a Hopf algebra \mathcal{H} is a quadruple $(\mathcal{A}, \#, r_A, \mathbb{A})$ such that

- (i) \mathcal{A} is a class of related alphabets;
- (ii) $\#$ is an associative operation on \mathcal{A} , called *disjoint sum*;
- (iii) for any related alphabet A of \mathcal{A} , $r_A : \mathcal{H} \rightarrow \mathbb{K}\langle A \rangle$ is a graded linear map, and for any alphabets A and A' of \mathcal{A} and any $x \in \mathcal{H}$, $\theta_{A,A'} \cdot r_{A+A'} \cdot x = (r_A \otimes r_{A'}) \circ \Delta \cdot x$;
- (iv) \mathbb{A} is an alphabet of \mathcal{A} such that the map $r_{\mathbb{A}}$ is injective.

Property (iii) is known as the *alphabet doubling trick*. This property, enjoyed by polynomial realizations of a large number of Hopf algebras, allows us to rephrase their coproduct by such alphabet transformations [3, 17, 12, 8, 7, 6].

1.3 Terms and free operads

A *signature* is a graded set $\mathcal{S} = \bigsqcup_{n \in \mathbb{N}} \mathcal{S} \cdot n$. A signature \mathcal{S} is *positive* if $\mathcal{S} \cdot 0 = \emptyset$. When all $\mathcal{S} \cdot n$ are finite, the *profile* of \mathcal{S} is the infinite word w such that $w \cdot i$ is the cardinality of $\mathcal{S} \cdot i - 1$. An *\mathcal{S} -term* is either the *leaf* \perp or a pair $(g, (t_1, \dots, t_n))$ where $g \in \mathcal{S} \cdot n$ and t_1, \dots, t_n are \mathcal{S} -terms. Thus, an \mathcal{S} -term is a planar rooted tree where internal nodes are decorated on \mathcal{S} . The set of \mathcal{S} -terms is denoted by $\mathfrak{T} \cdot \mathcal{S}$. The *free operad* on \mathcal{S} is the set $\mathfrak{T} \cdot \mathcal{S}$ such that the arity $\text{ar} \cdot t$ of $t \in \mathfrak{T} \cdot \mathcal{S}$ is the number of leaves of t , for any $t, s_1, \dots, s_{\text{ar} \cdot t} \in \mathfrak{T} \cdot \mathcal{S}$, $t[s_1, \dots, s_{\text{ar} \cdot t}]$ is the \mathcal{S} -term obtained by grafting the root of each s_i onto the i -th leaf of t , and the unit of $\mathfrak{T} \cdot \mathcal{S}$ is \perp . The *degree* $\text{dg} \cdot t$ of $t \in \mathfrak{T} \cdot \mathcal{S}$ is the number of internal nodes of t . This map dg is a grading of $\mathfrak{T} \cdot \mathcal{S}$.

An *\mathcal{S} -forest* is any element of $\mathfrak{F} \cdot \mathcal{S} := (\mathfrak{T} \cdot \mathcal{S})^*$. We identify each internal node of f with its position, starting by 1, according to its left to right preorder traversal. Let $\rightsquigarrow_{f,j}$ be the binary relation on the set of internal nodes of f such that $i \rightsquigarrow_{f,j} i'$ holds if i' is the j -th child of i in f . Figure 1 contains examples of these notions.





Figure 1: A \mathcal{S}_e -forest f and $\text{rd} \cdot f$, where \mathcal{S}_e is the signature $\mathcal{S}_e := \mathcal{S}_e \cdot 1 \sqcup \mathcal{S}_e \cdot 2 \sqcup \mathcal{S}_e \cdot 3$ with $\mathcal{S}_e \cdot 1 = \{a\}$, $\mathcal{S}_e \cdot 2 = \{b\}$, and $\mathcal{S}_e \cdot 3 = \{c\}$. Internal nodes are identified by the integers close to them. The degree of f is 7, and we have for instance $1 \rightsquigarrow_{f,1} 2$, $1 \rightsquigarrow_{f,3} 3$, and $5 \rightsquigarrow_{f,2} 6$.

Note that $\mathbb{N} \cdot \mathfrak{F} \cdot \mathcal{S}$ is a Hopf algebra graded by dg and whose bases are indexed by the set $\text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}$ of reduced \mathcal{S} -forests.

2 Lattices on forests and alternative bases

Let \mathcal{S} be a positive signature and \rightarrow be the binary relation on $\text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}$ such that $f \rightarrow f'$ if f' can be obtained from f by selecting an internal node i , detaching the subterm rooted at i , and grafting it to the rightmost leaf of its former immediate left brother. For instance,



Let \preceq be the reflexive and transitive closure of \rightarrow .

Proposition 2.1. *For any positive signature \mathcal{S} , the relation \preceq is a partial order on $\text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}$. Moreover, \rightarrow is the covering relation of this partial order.*

The poset $(\text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}, \preceq)$ is called the *\mathcal{S} -easterly wind poset*. This name is derived from the observation that the covering relation of this poset involves detaching a subterm from the east and then attaching it to the west, as if an easterly breeze is blowing on the forest. Let $\iota : \mathcal{S} \rightarrow \mathfrak{T} \cdot \mathcal{S}$ be the map such that $\iota \cdot g$ is the \mathcal{S} -term of degree 1 whose root is decorated by g . Given $w \in \mathcal{S}^*$, let $\mathcal{W} \cdot w$ be the interval of the \mathcal{S} -easterly wind poset whose lower bound is the \mathcal{S} -forest f_{\min} of length $\ell \cdot w$ such that $f_{\min} \cdot i := \iota \cdot w \cdot i$ for any $i \in [\ell \cdot w]$, and the upper bound is the \mathcal{S} -forest f_{\max} of length 1 such that $f_{\max} \cdot 1 := (\iota \cdot w \cdot 1) \circ_1 \cdots \circ_1 (\iota \cdot w \cdot \ell \cdot w)$, where \circ_i is the partial composition map of the free operad $\mathfrak{T} \cdot \mathcal{S}$. Figure 2 shows an example of an \mathcal{S} -easterly wind lattice.

Theorem 2.2. *For any positive signature \mathcal{S} and any $w \in \mathcal{S}^*$, the poset $\mathcal{W} \cdot w$ is a lattice.*

Let us now use these lattices to build new bases of $\mathbf{N} \cdot \mathfrak{T} \cdot \mathcal{S}$. For any $f \in \text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}$, let

$$F_f := \sum_{f' \in \text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}} [f \preceq f'] \mu_{\preceq}(f, f') E_{f'} \quad \text{and} \quad H_f := \sum_{f' \in \text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}} [f' \preceq f] F_{f'}, \quad (2.2)$$

where μ_{\preceq} is the Möbius function of the \mathcal{S} -easterly wind poset. For instance, in $\mathbf{N} \cdot \mathfrak{T} \cdot \mathcal{S}_e$,

$$F = E - E - E + E, \quad (2.3)$$

$$H = F + F + F. \quad (2.4)$$

By Möbius inversion and triangularity,

$$E_f = \sum_{f' \in \text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}} [f \preceq f'] F_{f'} \quad \text{and} \quad F_f = \sum_{f' \in \text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}} [f' \preceq f] \mu_{\preceq}(f', f) H_{f'}, \quad (2.5)$$




Figure 2: The Hasse diagram of the poset $\mathcal{W} \cdot \text{cab}$ where a , b , and c belong to the signature \mathcal{S}_e defined in Figure 1.

The reduced \mathcal{S}_e -forest at the top is the minimal element of the poset, and the one at the bottom is the maximal element.

so that the sets $\{F_f : f \in \text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}\}$ and $\{H_f : f \in \text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}\}$ are bases of $\mathbf{N} \cdot \mathfrak{T} \cdot \mathcal{S}$, called respectively the *fundamental basis* (or *F-basis* for short) and the *homogeneous basis* (or *H-basis* for short).

The i -th leaf of $t \in \mathfrak{T} \cdot \mathcal{S}$ is *extremal* if, by setting $\mathcal{S}' := \mathcal{S} \sqcup \{\vartheta\}$ where ϑ is a constant of arity 0, the only internal node of the \mathcal{S}' -term $t \circ_i \vartheta$ which is decorated by ϑ is the last one. Let \setminus be the *under* operation on $\text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}$ such that $f \setminus f'$ is obtained by concatenating f and f' , and then, by successively grafting the first term of f' onto the leftmost extremal leaf of the last term of f , the second term of f' onto the second leftmost leaf of the last term f , and so on, until there is no remaining term in f' or there is no extremal leaf in the last term of f . Let also $/$ be the *over* operation on $\text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}$ defined by $f / f' := f \cdot f'$.

Theorem 2.3. For any positive signature \mathcal{S} and any reduced \mathcal{S} -forests f' and f'' ,

$$E_{f'} \cdot E_{f''} = E_{f' / f''}, \quad F_{f'} \cdot F_{f''} = \sum_{f \in \text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}} [f' / f'' \preceq f \preceq f' \setminus f''] F_f, \quad H_{f'} \cdot H_{f''} = H_{f' \setminus f''}. \quad (2.6)$$

The product of $\mathbf{N} \cdot \mathfrak{T} \cdot \mathcal{S}$ on the F-basis is akin to a shuffle of reduced \mathcal{S} -forests. For instance, in $\mathbf{N} \cdot \mathfrak{T} \cdot \mathcal{S}_e$,

$$F \cdot F = F + F + F + F. \quad (2.7)$$

These three bases, together with the over and under operations, and the partial order on reduced \mathcal{S} -forests mimic classical basis constructions of Hopf algebras [3, 15, 13].

3 Quotient operads and Hopf subalgebras

Let \mathcal{S} be a signature and \equiv be an operad congruence of the free operad $\mathfrak{T} \cdot \mathcal{S}$. The equivalence relation \equiv is extended on $\text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}$ by setting $f \equiv f'$ if $\ell \cdot f = \ell \cdot f'$ and, for any $i \in [\ell \cdot f]$, $f \cdot i \equiv f' \cdot i$. The operad congruence \equiv is *compatible with the degree* if $t \equiv t'$ implies $\text{dg} \cdot t = \text{dg} \cdot t'$ for any $t, t' \in \mathfrak{T} \cdot \mathcal{S}$. The operad congruence \equiv is of *finite type* if for any $t \in \mathfrak{T} \cdot \mathcal{S}$, the \equiv -equivalence class $[t]_{\equiv}$ of t is finite. In this case, for any $f \in \text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}$,

$$E_{[f]_{\equiv}} := \sum_{f' \in [f]_{\equiv}} E_{f'} \quad (3.1)$$

is a well-defined element of $\mathbf{N} \cdot \mathfrak{T} \cdot \mathcal{S}$.

Theorem 3.1. *Let \mathcal{S} be a signature and \equiv be an operad congruence of $\mathfrak{T} \cdot \mathcal{S}$ compatible with the degree and of finite type. By denoting by \mathcal{V} the linear span of the $E_{[f]_{\equiv}}$, $f \in \text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}$, the following assertions hold:*

- (i) *The space \mathcal{V} is a Hopf sub-algebra of $\mathbf{N} \cdot \mathfrak{T} \cdot \mathcal{S}$.*
- (ii) *The Hopf algebra \mathcal{V} is isomorphic to $\mathbf{N} \cdot (\mathfrak{T} \cdot \mathcal{S}) / \equiv$.*

Point (i) of Theorem 3.1 is a particular case of [1, Theorem 3.13]. Point (ii) provides an interpretation of $\mathbf{N} \cdot (\mathfrak{T} \cdot \mathcal{S}) / \equiv$ as a Hopf subalgebra of the natural Hopf algebra of a free operad. The condition of compatibility with the degree for \equiv is important for the fact that $\mathbf{N} \cdot (\mathfrak{T} \cdot \mathcal{S}) / \equiv$ inherits the graduation of $\mathbf{N} \cdot \mathfrak{T} \cdot \mathcal{S}$.

4 Polynomial realization

Given a signature \mathcal{S} , the class of *\mathcal{S} -forest-like alphabets* is the class of related alphabets A endowed with relations R , D_g , and \prec_j such that

- (i) R is a unary relation called *root relation*;
- (ii) for any $g \in \mathcal{S}$, D_g is a unary relation called *g -decoration relation*;
- (iii) for any $j \geq 1$, \prec_j is a binary relation called *j -edge relation*.

Let A' and A'' be two \mathcal{S} -forest-like alphabets, each endowed respectively with the relations R' , D'_g , and \prec'_j , and R'' , D''_g , and \prec''_j . The *disjoint sum* of A' and A'' is the \mathcal{S} -forest-like alphabet $A := A' \# A''$ endowed with the relations R , D_g , and \prec_j such that $A := A' \sqcup A''$, $R := R' \sqcup R''$, $D_g := D'_g \sqcup D''_g$, and $a_1 \prec_j a_2$ holds if $a_1, a_2 \in A'$ and $a_1 \prec'_j a_2$, or $a_1, a_2 \in A''$ and $a_1 \prec''_j a_2$, or $a_1 \in A'$, $a_2 \in A''$, and $a_2 \in R''$.

Let A be an \mathcal{S} -forestlike alphabet. Given $f \in \text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}$, $w \in A^*$ is *f -compatible* if the following four assertions are satisfied: (i) $\ell \cdot w = \text{dg} \cdot f$; (ii) for any $i \in [\text{dg} \cdot f]$, i is a root of f implies $w \cdot i \in R$; (iii) for any $i \in [\text{dg} \cdot f]$ the internal node i of f is decorated by $g \in \mathcal{S}$ implies $w \cdot i \in D_g$; (iv) for any $i, i' \in [\text{dg} \cdot f]$, $i \rightsquigarrow_{f,j} i'$ implies $w \cdot i \prec_j w \cdot i'$. This property is denoted by $w \Vdash^A f$. For instance, by considering the reduced \mathcal{S}_e -forest f of Figure 1,

any f -compatible word $w \in A^*$ satisfies $\ell \cdot w = 7$, $w \cdot 1, w \cdot 5 \in R$, $w \cdot 2, w \cdot 4, w \cdot 7 \in D_a$, $w \cdot 3, w \cdot 5, w \cdot 6 \in D_b$, $w \cdot 1 \in D_c$, $w \cdot 1 \prec_1 w \cdot 2$, $w \cdot 1 \prec_3 w \cdot 3$, $w \cdot 3 \prec_1 w \cdot 4$, $w \cdot 5 \prec_2 w \cdot 6$, and $w \cdot 6 \prec_1 w \cdot 7$.

Let $r_A : \mathbb{N} \cdot \mathfrak{T} \cdot \mathcal{S} \rightarrow \mathbb{K}\langle A \rangle$ be the linear map defined for any $f \in \text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}$ by

$$r_A \cdot E_f := \sum_{w \in A^*} [w \Vdash^A f] w. \quad (4.1)$$

The A -polynomial $r_A \cdot E_f$ is the *A-realization* of f .

We now construct a specific \mathcal{S} -forest-like alphabet $\mathbb{A}_{\mathcal{S}}$ in order to obtain an injective map $r_{\mathbb{A}_{\mathcal{S}}}$. For any signature \mathcal{S} , let the \mathcal{S} -forest-like alphabet $\mathbb{A}_{\mathcal{S}}$ such that

$$\mathbb{A}_{\mathcal{S}} := \{\mathbf{a}_{g,u} : g \in \mathcal{S} \text{ and } u \in \mathbb{N}^*\}, \quad (4.2)$$

the root relation is defined by $R := \{\mathbf{a}_{g,u} \in \mathbb{A}_{\mathcal{S}} : u = 0^\ell, \ell \geq 0\}$, the g -decoration relation D_g is defined by $D_g := \{\mathbf{a}_{g',u} \in \mathbb{A}_{\mathcal{S}} : g' = g\}$, the j -edge relation \prec_j is defined by $\mathbf{a}_{g,u} \prec_j \mathbf{a}_{g',u,j \cdot 0^\ell}$ for any $g, g' \in \mathcal{S}$, $u \in \mathbb{N}^*$, and $\ell \geq 0$. Observe that these binary relations are antisymmetric but are neither transitive nor reflexive. For instance, by considering the reduced \mathcal{S}_e -forest f of Figure 1,

$$r_{\mathbb{A}_{\mathcal{S}}} \cdot E_f = \sum_{\ell_1, \dots, \ell_7 \in \mathbb{N}} \mathbf{a}_{c,0^{\ell_1}} \mathbf{a}_{a,0^{\ell_1}10^{\ell_2}} \mathbf{a}_{b,0^{\ell_1}130^{\ell_3}} \mathbf{a}_{a,0^{\ell_1}130^{\ell_3}10^{\ell_4}} \mathbf{a}_{b,0^{\ell_5}} \mathbf{a}_{b,0^{\ell_5}20^{\ell_6}} \mathbf{a}_{b,0^{\ell_5}20^{\ell_6}10^{\ell_7}}. \quad (4.3)$$

Theorem 4.1. *For any signature \mathcal{S} , the class of \mathcal{S} -forest-like alphabets, together with the alphabet disjoint sum operation \ddagger , the map r_A , and the alphabet $\mathbb{A}_{\mathcal{S}}$, forms a polynomial realization of the Hopf algebra $\mathbb{N} \cdot \mathfrak{T} \cdot \mathcal{S}$.*

By Theorems 3.1 and 4.1, the map $r_{\mathbb{A}_{\mathcal{S}}}$ provides also a polynomial realization of the Hopf subalgebra $\mathbb{N} \cdot (\mathfrak{T} \cdot \mathcal{S}) / \equiv$ of $\mathbb{N} \cdot \mathfrak{T} \cdot \mathcal{S}$ when \equiv is an operad congruence of $\mathfrak{T} \cdot \mathcal{S}$ compatible with the degree and of finite type.

5 Word quasi-symmetric functions Hopf algebra

Let A be a totally ordered alphabet. The *packing* of $w \in A^*$ is the word $\text{pck} \cdot w$ on positive integers obtained by replacing each letter a of w by the number of distinct letters less than or equal to a in w . For instance, for $A = \mathbb{N}$, $\text{pck} \cdot 4234473 = 3123342$. When w is a word of positive integers such that $\text{pck} \cdot w = w$, w is *packed*. The set of packed words is denoted by \mathcal{P} . The Hopf algebra of *word quasi-symmetric functions* \mathbf{WQSym} [11, 17] is defined on the linear span of \mathcal{P} . The *fundamental basis* of \mathbf{WQSym} is the set $\{M_u : u \in \mathcal{P}\}$. We need here only to describe the polynomial realization of \mathbf{WQSym} , which satisfies

$$r_A \cdot M_u = \sum_{w \in A^*} [\text{pck}(w) = u] w. \quad (5.1)$$

By setting \mathbb{A} as the related alphabet $\{\mathbf{a}_i : i \in \mathbb{N}\}$ totally ordered by the order relation \prec satisfying $\mathbf{a}_i \prec \mathbf{a}_{i'}$ if $i \leq i'$, we have for instance

$$r_{\mathbb{A}} \cdot M_{2113} = \sum_{\ell_1, \ell_2, \ell_3, \ell_4 \in \mathbb{N}} [\ell_2 = \ell_3 < \ell_1 < \ell_4] \mathbf{a}_{\ell_1} \mathbf{a}_{\ell_2} \mathbf{a}_{\ell_3} \mathbf{a}_{\ell_4}. \quad (5.2)$$

Let $\mathbb{A}_{\mathbb{N}}$ be the \mathcal{S} -forest-like alphabet such that $\mathbb{A}_{\mathbb{N}} := \mathbb{A}$, $\mathbb{R} := \mathbb{A}_{\mathbb{N}}$, $\mathbb{D}_{\mathbf{g}} := \mathbb{A}_{\mathbb{N}}$, and $\mathbf{a}_i \prec_j \mathbf{a}_{i'}$ if $i < i'$. Given $\mathbf{f} \in \text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}$, $u \in \mathcal{P}$ is \mathbf{f} -compatible if $\mathbf{a}_{u \cdot 1} \dots \mathbf{a}_{u \cdot \ell \cdot u} \Vdash^{\mathbb{A}_{\mathbb{N}}} \mathbf{f}$. This property is denoted by $u \vdash \mathbf{f}$. For instance, by considering the reduced \mathcal{S}_e -forest \mathbf{f} defined in Figure 1, $1223123 \vdash \mathbf{f}$ and $1525346 \vdash \mathbf{f}$.

Theorem 5.1. *For any signature \mathcal{S} , the space $r_{\mathbb{A}_{\mathbb{N}}} \cdot \mathbb{N} \cdot \mathfrak{F} \cdot \mathcal{S}$ is a Hopf subalgebra of $r_{\mathbb{A}} \cdot \mathbf{WQSym}$. More precisely, for any $\mathbf{f} \in \text{rd} \cdot \mathfrak{F} \cdot \mathcal{S}$,*

$$r_{\mathbb{A}_{\mathbb{N}}} \cdot E_{\mathbf{f}} = \sum_{u \in \mathcal{P}} [u \vdash \mathbf{f}] r_{\mathbb{A}} \cdot M_u. \quad (5.3)$$

By Theorems 3.1 and 5.1, $r_{\mathbb{A}_{\mathbb{N}}} \cdot \mathbb{N} \cdot (\mathfrak{F} \cdot \mathcal{S}) / \equiv$ is a Hopf subalgebra of $r_{\mathbb{A}} \cdot \mathbf{WQSym}$ when \equiv is an operad congruence compatible with the degree and of finite type. Observe that the map $r_{\mathbb{A}_{\mathbb{N}}}$ is not injective as, for instance,

$$r_{\mathbb{A}_{\mathbb{N}}} \cdot E \quad \begin{array}{c} \circ \\ | \\ \circ \text{---} \text{b} \text{---} \circ \\ | \quad | \\ \circ \text{---} \text{a} \quad \circ \text{---} \text{b} \\ | \quad | \\ \circ \quad \circ \end{array} = M_{122} + M_{123} + M_{132} = r_{\mathbb{A}_{\mathbb{N}}} \cdot E \quad \begin{array}{c} \circ \\ | \\ \circ \text{---} \text{c} \text{---} \circ \\ | \quad | \\ \circ \text{---} \text{b} \quad \circ \text{---} \text{b} \\ | \quad | \\ \circ \quad \circ \end{array}. \quad (5.4)$$

6 Connes-Kreimer and decorated forests Hopf algebras

For any set D , a D -decorated forest is a word of nonempty planar rooted trees where nodes are decorated on D . In [4], Foissy introduces a Hopf algebra structure \mathbf{NCK}_D on the linear span of D -decorated forests. This is a generalization of the noncommutative Connes-Kreimer Hopf algebra as the original case occurs when D is a singleton.

For any set D , let the signature $\mathcal{S}_D := \{\alpha_{d,n} : d \in D, n \in \mathbb{N}\}$ such that $\text{ar} \cdot \alpha_{d,n} = n$. Let also $\mathbb{A}_{D,\mathbb{N}}$ be the \mathcal{S}_D -forest-like alphabet such that $\mathbb{A}_{D,\mathbb{N}} := \{\mathbf{a}_{d,i} : d \in D, i \in \mathbb{N}\}$, $\mathbb{R} := \mathbb{A}_{D,\mathbb{N}}$, $\mathbb{D}_{\alpha_{d,n}} := \{\mathbf{a}_{d,i} \in \mathbb{A}_{D,\mathbb{N}} : i \in \mathbb{N}\}$, and $\mathbf{a}_{d,i} \prec_j \mathbf{a}_{d',i'}$ if $i < i'$.

Theorem 6.1. *For any set D , the Hopf algebra $r_{\mathbb{A}_{D,\mathbb{N}}} \cdot \mathbb{N} \cdot \mathfrak{F} \cdot \mathcal{S}_D$ is isomorphic to the Hopf algebra \mathbf{NCK}_D of D -decorated forests.*

Theorem 6.1 provides a polynomial realization of \mathbf{NCK}_D . Indeed, given a D -decorated forest F , the $\mathbb{A}_{D,\mathbb{N}}$ -polynomial associated with the basis element E_F of \mathbf{NCK}_D , is $r_{\mathbb{A}_{D,\mathbb{N}}} \cdot E_{\mathbf{f}}$ where \mathbf{f} is the \mathcal{S}_D -forest obtained by replacing each internal node decorated by $d \in D$

and having n children by $\alpha_{d,n}$. For instance, for $D := \{1, 2, 3\}$,

$$\begin{aligned}
 r_{\mathbb{A}_{D,\mathbb{N}}} \cdot E & \quad \text{:=} \quad r_{\mathbb{A}_{D,\mathbb{N}}} \cdot E & (6.1) \\
 \begin{array}{c} \textcircled{3} \\ \diagup \quad \diagdown \\ \textcircled{2} \quad \textcircled{3} \\ \diagup \quad \diagdown \\ \textcircled{2} \quad \textcircled{3} \quad \textcircled{1} \end{array} & \quad \begin{array}{c} \textcircled{1} \\ \textcircled{2} \end{array} & \quad \begin{array}{c} \alpha_{3,2} \\ \diagup \quad \diagdown \\ \alpha_{2,0} \quad \alpha_{3,0} \\ \diagup \quad \diagdown \\ \alpha_{2,0} \quad \alpha_{3,0} \quad \alpha_{1,0} \end{array} & \quad \begin{array}{c} \alpha_{1,1} \\ \textcircled{2,0} \end{array}
 \end{aligned}$$

$$= \sum_{\ell_1, \dots, \ell_8 \in \mathbb{N}} [\ell_1 < \ell_2, \ell_3][\ell_3 < \ell_4, \ell_5, \ell_6][\ell_7 < \ell_8] \mathbf{a}_{3,\ell_1} \mathbf{a}_{2,\ell_2} \mathbf{a}_{3,\ell_3} \mathbf{a}_{2,\ell_4} \mathbf{a}_{3,\ell_5} \mathbf{a}_{1,\ell_6} \mathbf{a}_{1,\ell_7} \mathbf{a}_{2,\ell_8}.$$

7 Faà di Bruno Hopf algebras

Let $\text{MA}_{\mathcal{S}}$ be the quotient of the free operad $\mathfrak{T} \cdot \mathcal{S}$ by the congruence \equiv satisfying, for any $g, g' \in \mathcal{S}$, $i \in [\text{ar} \cdot g]$, and $i' \in [\text{ar} \cdot g']$, $g \circ_i g' \equiv g' \circ_{i'} g$. These operads $\text{MA}_{\mathcal{S}}$ are generalizations of multiassociative operads, introduced in [9], which contain originally only binary generators. For this reason, we call $\text{MA}_{\mathcal{S}}$ the *\mathcal{S} -multi-multiassociative operad*.

Proposition 7.1. *For any signature \mathcal{S} , the operad $\text{MA}_{\mathcal{S}}$ admits the following realization. For any $n \geq 0$, $\text{MA}_{\mathcal{S}} \cdot n$ is the set of multisets $\{g_1, \dots, g_\ell\}$ on \mathcal{S} such that $\text{ar} \cdot g_1 + \dots + \text{ar} \cdot g_\ell - \ell + 1 = n$. Moreover, for any $m, m' \in \text{MA}_{\mathcal{S}}$ and $i \in [\text{ar} \cdot m]$, $m \circ_i m'$ is the union of m and m' .*

Recall that \mathbf{FdB} is the noncommutative Faà di Bruno Hopf algebra whose construction is detailed in Section 1.1. In [5], a deformation \mathbf{FdB}_r of \mathbf{FdB} is introduced. The special case \mathbf{FdB}_0 is the Hopf algebra of noncommutative symmetric functions \mathbf{Sym} , and for any $r \geq 1$, each \mathbf{FdB}_r is isomorphic (in a nontrivial way) to \mathbf{FdB} .

Proposition 7.2. *Let \mathcal{S} be a signature of profile $0^{r+1}10\dots$ where $r \in \mathbb{N}$. The Hopf algebra $\mathbb{N} \cdot \text{MA}_{\mathcal{S}}$ is isomorphic to the noncommutative Faà di Bruno Hopf algebra \mathbf{FdB}_r .*

This result is a consequence of an alternative construction of \mathbf{FdB}_r via pros, whose details can be found in [1].

8 Multi-symmetric functions Hopf algebras

Let us now consider the Hopf algebra $\mathbf{Sym}^{(s)}$ of noncommutative multi-symmetric functions of level $s \in \mathbb{N}$. This Hopf algebra is a noncommutative analog of multi-symmetric functions, introduced in [18].

Proposition 8.1. *Let \mathcal{S} be a signature of profile $0s0\dots$ where $s \in \mathbb{N}$. The Hopf algebra $\mathbb{N} \cdot \text{MA}_{\mathcal{S}}$ is isomorphic to the noncommutative multi-symmetric functions Hopf algebra $\mathbf{Sym}^{(s)}$.*

This framework offered by the multi-multiassociative operads leads naturally to Hopf algebras which are to $\mathbf{Sym}^{(s)}$ what \mathbf{FdB}_r is to \mathbf{FdB}_0 . These Hopf algebras depend on the

two parameters $r, s \in \mathbb{N}$, are denoted by $\mathbf{FdB}_{r,s}$, and are defined as $\mathbf{N} \cdot \mathbf{MA}_S$ where the S are signatures of profiles $0^{r+1}s0 \dots$. By Theorems 3.1 and 4.1, we have, for free, polynomial realizations of $\mathbf{FdB}_{r,s}$ on the alphabet \mathbb{A}_S .

By the following result, these polynomial realizations can be expressed on a simpler alphabet $\mathbb{A}_{S,\mathbb{N}}$ defined as follows. Let $\mathbb{A}_{S,\mathbb{N}}$ be the S -forest-like alphabet such that $\mathbb{A}_{S,\mathbb{N}} := \{\mathbf{a}_{g,i} : g \in S, i \in \mathbb{N}\}$, $\mathbb{R} := \mathbb{A}_{S,\mathbb{N}}$, $\mathbb{D}_g := \{\mathbf{a}_{g,i} \in \mathbb{A}_{S,\mathbb{N}} : i \in \mathbb{N}\}$, and $\mathbf{a}_{g,i} \prec_j \mathbf{a}_{g',i'}$ if $i < i'$.

Theorem 8.2. *Let S be a signature of profile $0^{r+1}s0 \dots$ where $r, s \in \mathbb{N}$. The Hopf algebra $r_{\mathbb{A}_{S,\mathbb{N}}} \cdot \mathbf{N} \cdot \mathbf{MA}_S$ is isomorphic to $\mathbf{FdB}_{r,s}$.*

For instance, let $S := \{a, b\}$ be a signature of profile $0^{r+1}s0 \dots$ where $r = 1$ and $s = 2$. In $\mathbf{FdB}_{r,s}$, we have

$$r_{\mathbb{A}_{S,\mathbb{N}}} \cdot E_{\{aab\} \{a\}} = 4 r_{\mathbb{A}_{S,\mathbb{N}}} \cdot E_{\begin{array}{c} \circ \\ | \\ \text{a} \\ / \quad \backslash \\ \text{b} \quad \text{a} \\ | \quad | \\ \circ \quad \circ \end{array}} + 4 r_{\mathbb{A}_{S,\mathbb{N}}} \cdot E_{\begin{array}{c} \circ \\ | \\ \text{a} \\ / \quad \backslash \\ \text{b} \quad \text{a} \\ | \quad | \\ \circ \quad \circ \end{array}} + 4 r_{\mathbb{A}_{S,\mathbb{N}}} \cdot E_{\begin{array}{c} \circ \\ | \\ \text{b} \\ / \quad \backslash \\ \text{a} \quad \text{a} \\ | \quad | \\ \circ \quad \circ \end{array}} \tag{8.1}$$

$$+ r_{\mathbb{A}_{S,\mathbb{N}}} \cdot E_{\begin{array}{c} \circ \\ | \\ \text{a} \\ / \quad \backslash \\ \text{a} \quad \text{b} \\ | \quad | \\ \circ \quad \circ \end{array}} + r_{\mathbb{A}_{S,\mathbb{N}}} \cdot E_{\begin{array}{c} \circ \\ | \\ \text{a} \\ / \quad \backslash \\ \text{b} \quad \text{a} \\ | \quad | \\ \circ \quad \circ \end{array}} + r_{\mathbb{A}_{S,\mathbb{N}}} \cdot E_{\begin{array}{c} \circ \\ | \\ \text{b} \\ / \quad \backslash \\ \text{a} \quad \text{a} \\ | \quad | \\ \circ \quad \circ \end{array}} \dots$$

Open questions and future work

We have introduced polynomial realizations of $\mathbf{N} \cdot \mathcal{O}$ and used them to establish links between this Hopf algebra and other ones. Here are open questions raised by this work.

- (1) This concerns the further study of easterly wind lattices, specifically to investigate properties like semidistributivity, number of covering relations, and number of intervals.
- (2) Another question is to delineate conditions on the operad \mathcal{O} for the fact that $\mathbf{N} \cdot \mathcal{O}$ is self-dual. The F and H-bases (when \mathcal{O} is free) of $\mathbf{N} \cdot \mathcal{O}$, together with its polynomial realization could intervene to solve this question.
- (3) This axis is about the construction of a Hopf algebra $\mathbf{WQSym}^{(s)}$ that is to \mathbf{WQSym} what $\mathbf{Sym}^{(s)}$ is to \mathbf{Sym} with the aim to obtain a stronger version of Theorem 5.1.
- (4) The question of understanding the structure of $\mathbf{N} \cdot \mathbf{MA}_S$ in the general case, when S is any signature (having possibly elements of arity 0) as this Hopf algebra generalizes the Faà du Bruno and the multi-symmetric function Hopf algebras.

References

[1] J.-P. Bultel and S. Giraud. “Combinatorial Hopf algebras from PROs”. *J. Algebr. Comb.* **44.2** (2016), pp. 455–493.

- [2] F. Chapoton and M. Livernet. “Relating two Hopf algebras built from an operad”. *Int. Math. Res. Notices* **2007.24** (2007), Art. ID rnm131, 27 pp.
- [3] G. Duchamp, F. Hivert, and J.-Y. Thibon. “Noncommutative Symmetric Functions VI: Free Quasi-Symmetric Functions and Related Algebras”. *Int. J. of Algebr. Comput.* **12** (2002), pp. 671–717.
- [4] L. Foissy. “Les algèbres de Hopf des arbres enracinés décorés, I”. *Bull. Sci. Math.* **126.3** (2002), pp. 193–239.
- [5] L. Foissy. “Faà di Bruno subalgebras of the Hopf algebra of planar trees from combinatorial Dyson-Schwinger equations”. *Adv. Math.* **218.1** (2008), pp. 136–162.
- [6] L. Foissy. “Realizations of Hopf algebras of graphs by alphabets”. *Algebraic combinatorics, resurgence, moulds and applications CARMA. Vol. 1.* Vol. 31. IRMA Lect. Math. Theor. Phys. EMS Publ. House, Berlin, 2020, pp. 225–261.
- [7] L. Foissy, J.-C. Novelli, and J.-Y. Thibon. “Polynomial realizations of some combinatorial Hopf algebras”. *J. Noncommut. Geom.* **8.1** (2014), pp. 141–162.
- [8] S. Giraud. “Combinatoire algébrique des arbres”. PhD thesis. Université de Marne la Vallée, 2011.
- [9] S. Giraud. “Pluriassociative algebras II: The polydendriform operad and related operads”. *Adv. Appl. Math.* **77** (2016), pp. 43–85.
- [10] S. Giraud. *Nonsymmetric Operads in Combinatorics.* ix+172. Springer Nature Switzerland AG, 2018.
- [11] F. Hivert. “Combinatoire des fonctions quasi-symétriques”. PhD thesis. Université de Marne la Vallée, 1999.
- [12] F. Hivert. “An introduction to combinatorial Hopf algebras—examples and realizations”. *Physics and theoretical computer science.* Vol. 7. NATO Secur. Sci. Ser. D Inf. Commun. Secur. IOS, Amsterdam, 2007, pp. 253–274.
- [13] F. Hivert, J.-C. Novelli, and J.-Y. Thibon. “The Algebra of Binary Search Trees”. *Theor. Comput. Sc.* **339.1** (2005), pp. 129–165.
- [14] P. van der Laan. “Operads. Hopf algebras and coloured Koszul duality”. PhD thesis. Universiteit Utrecht, 2004.
- [15] J.-L. Loday and M. Ronco. “Order Structure on the Algebra of Permutations and of Planar Binary Trees”. *J. Algebr. Comb.* **15.3** (2002), pp. 253–270.
- [16] M. Méndez and J. Liendo. “An antipode formula for the natural Hopf algebra of a set operad”. *Adv. Appl. Math.* **53** (2014), pp. 112–140.
- [17] J.-C. Novelli and J.-Y. Thibon. “Polynomial realizations of some trialgebras”. *Formal Power Series and Algebraic Combinatorics* (2006), pp. 243–254.
- [18] J.-C. Novelli and J.-Y. Thibon. “Free quasi-symmetric functions and descent algebras for wreath products, and noncommutative multi-symmetric functions”. *Discrete Math.* **310.24** (2010), pp. 3584–3606.