

Joint covariance property under geometric image transformations for spatio-temporal receptive fields according to the generalized Gaussian derivative model for visual receptive fields

Tony Lindeberg

Received: date / Accepted: date

Abstract The influence of natural image transformations on receptive field responses is crucial for modelling visual operations in computer vision and biological vision. In this regard, covariance properties with respect to geometric image transformations in the earliest layers of the visual hierarchy are essential for expressing robust image operations and for formulating invariant visual operations at higher levels. This paper defines and proves a joint covariance property under compositions of spatial scaling transformations, spatial affine transformations, Galilean transformations and temporal scaling transformations, which makes it possible to characterize how different types of image transformations interact with each other. Specifically, the derived relations show how the receptive field parameters need to be transformed, in order to match the output from spatio-temporal receptive fields with the underlying spatio-temporal image transformations.

Keywords Covariance · Receptive field · Scaling · Affine · Galilean · Spatial · Temporal · Spatio-temporal · Image transformations · Vision

1 Introduction

When images, video sequences or video streams are acquired from the real world, they are subject to natural image transformations, as caused by variations in the positions, relative orientations and motions between the objects in the world and the observer:

The support from the Swedish Research Council (contracts 2018-03586 and 2022-02969) is gratefully acknowledged.

Tony Lindeberg
Computational Brain Science Lab, Division of Computational Science and Technology, KTH Royal Institute of Technology, SE-100 44 Stockholm, Sweden. E-mail: tony@kth.se

- Depending on the distance between the objects in the world and the observer, the perspective projections of objects onto the image surfaces may become smaller or larger, which to first-order of approximation can be modelled as *spatial scaling transformations*.
- Depending on the orientations of the surface normals of the objects in relation to the viewing direction, the image patterns may be compressed by different amounts in different directions (perspective foreshortening), which to first-order of approximation can be modelled as *spatial affine transformations*.
- Depending on how the objects in the world move relative to the (possibly time-dependent) viewing direction, the image patterns of objects may move in the image plane, which to first-order of approximation can be modelled as *Galilean transformations*.
- Depending on how fast the perspective projections of objects move in the image plane, or how fast spatio-temporal actions occur, the time-line along the temporal dimension may be compressed or expanded, which can be modelled as *temporal scaling transformations*.

These types of geometric image transformations will have a profound effect on the spatio-temporal receptive fields that register and process the image information at the earliest stages in the visual hierarchy, in that the output from the receptive fields to particular image patterns may be strongly dependent on the imaging conditions. Specifically, if the interaction effects between the geometric image transformations and the receptive fields are not properly taken into account, then the robustness of the visual modules can be strongly affected in a negative manner. On the other hand, if the interaction effects between the natural image transformations are properly taken into account, then the robustness of the visual measurements may be substantially improved.

A particular way of handling the interaction effects between the geometric image transformations and the recep-

tive fields is by requiring that the family of receptive fields is *covariant* under the relevant classes of image transformations. Covariance means that the geometric image transformations essentially commute with the image operations induced by the receptive fields, and do in this way provide a way to propagate well-defined relationships between the geometric image transformations and the receptive fields. Specifically, covariance properties of the receptive fields at lower levels in the visual hierarchy make it possible to define invariant image measurements at higher levels in the visual hierarchy.

The subject of this paper is to describe and derive a joint covariance property of receptive fields according to a specific model for spatio-temporal receptive fields, according to the generalized Gaussian derivative model for receptive fields, under joint combinations of spatial scaling transformations, spatial affine transformations, Galilean transformations and temporal scaling transformations.

2 The generalized Gaussian derivative model for spatio-temporal receptive fields

Given spatio-temporal image data of the form $f(x, t)$ for $x = (x_1, x_2)^T \in \mathbb{R}^2$ and $t \in \mathbb{R}$, in (Lindeberg 2011, 2016) a principled model for spatio-temporal receptive fields is derived and applied of the form (with slightly modified notation)

$$T(x, t; s, \Sigma, \tau, v) = g(x - vt; s, \Sigma) h(t; \tau), \quad (1)$$

where

- $s \in \mathbb{R}_+$ denotes a spatial scale parameter corresponding to the spatial variance of a non-negative spatial smoothing kernel,
- Σ denotes a positive definite 2×2 spatial covariance matrix that describes the spatial shape of the spatial smoothing kernel,
- $\tau \in \mathbb{R}_+$ denotes a temporal scale parameter corresponding to the temporal variance of a non-negative temporal smoothing kernel,
- $v \in \mathbb{R}^2$ denotes an image velocity vector,
- $g(x; s, \Sigma)$ denotes a 2-D affine Gaussian kernel of the form

$$g(x; s, \Sigma) = \frac{1}{2\pi s \det \Sigma} e^{-x^T \Sigma^{-1} x / 2s}, \quad (2)$$

- $h(t; \tau)$ denotes a temporal smoothing kernel, that for any temporal scaling factor S_t obeys the *temporal scale covariance property*

$$h(t'; \tau') = \frac{1}{S_t} h(t; \tau) \quad (3)$$

for $t' = S_t t$ and $\tau' = S_t^2 \tau$.

2.1 Temporal smoothing kernels

Based on the treatments in (Lindeberg 2011, 2016), the choice of the temporal smoothing as a 1-D temporal Gaussian kernel

$$h(t; \tau) = g_{1D}(t; \tau) = \frac{1}{\sqrt{2\pi\tau}} e^{-t^2/2\tau} \quad (4)$$

stands out as a canonical choice over a non-causal temporal domain (where the relative future in relation to any time moment can be accessed), whereas the choice of the temporal kernel as the time-causal limit kernel

$$h(t; \tau) = \Psi(t; \tau, c), \quad (5)$$

defined by having a Fourier transform of the form

$$\hat{\Psi}(\omega; \tau, c) = \prod_{k=1}^{\infty} \frac{1}{1 + i c^{-k} \sqrt{c^2 - 1} \sqrt{\tau} \omega}, \quad (6)$$

and corresponding to an infinite number of truncated exponential kernels with specially chosen time constants to obtain temporal scale covariance, stands out as a canonical choice over a time-causal temporal domain, where the future cannot be accessed. The distribution parameter $c > 1$ of this time-causal limit kernel is for practical purposes often chosen as $c = \sqrt{2}$ or $c = 2$.

2.2 Spatio-temporal derivatives

These purely spatio-temporal smoothing components of receptive fields are then to be combined with spatial and temporal derivative operations. Over the spatial domain, we can compute either partial spatial derivatives

$$\partial_{x^\alpha} = \partial_{x_1}^{\alpha_1} \partial_{x_2}^{\alpha_2} \quad (7)$$

for different orders $\alpha = (\alpha_1, \alpha_2)$ of spatial differentiation, or oriented directional derivatives in the directions φ

$$\partial_\varphi^m = (\cos \varphi \partial_{x_1} + \sin \varphi \partial_{x_2})^m \quad (8)$$

or different orientations ϕ and different orders of spatial differentiation m .

Over the temporal domain, we can compute partial temporal derivatives

$$\partial_t^n \quad (9)$$

for different orders n of temporal differentiation, or velocity-adapted temporal derivatives

$$\partial_{\vec{t}}^n = (v_1 \partial_{x_1} + v_2 \partial_{x_2} + \partial_t)^n \quad (10)$$

for different image velocities $v = (v_1, v_2)^T$ and orders n of temporal differentiation.

Specifically, in relation to the parameters Σ and v of the purely spatio-temporal smoothing component of the spatio-temporal receptive fields in (1), the image orientations φ in the directional derivative operators (8) should be chosen in the directions of the eigendirections of the spatial covariance matrix Σ , whereas the image velocities v in the velocity-adapted derivative operator should be chosen equal to the image velocity v in the spatio-temporal smoothing kernel.

2.3 Relations to biological vision

In (Lindeberg (2021)) it is demonstrated that the receptive fields of neurons in the lateral geniculate nucleus (LGN) as well as the receptive fields of simple cells in the primary visual cortex (V1), as measured by neurophysiological cell recordings by DeAngelis *et al.* (1995, 2004), Conway and Livingstone (2006) and Johnson *et al.* (2008), can be well modelled by idealized receptive fields derived from this generalized Gaussian derivative model for receptive fields.

According to the treatment in (Lindeberg (2021)), the spatio-temporal receptive fields of “non-lagged cells” and “lagged cells”, which have rotationally symmetric response properties over the spatial domain, are modelled by idealized receptive fields of the form

$$h_{LGN}(x_1, x_2, t; s, \tau) = \pm(\partial_{x_1x_1} + \partial_{x_2x_2})g(x_1, x_2; s) \partial_{t^n} h(t; \tau), \quad (11)$$

where $\nabla_x^2 = \partial_{x_1x_1} + \partial_{x_2x_2}$ represents the spatial Laplacian operator, and $h(t; \tau)$ represents a set of first-order truncated exponential kernels coupled in cascade, which in the most idealized situation may correspond to the time-causal limit kernel $\Psi(t; \tau, c)$ according to (6).

The spatio-temporal receptive fields of orientation selective simple cells in V1 are, in turn, modelled by idealized receptive fields of the form

$$T_{\varphi^{m_1} \perp \varphi^{m_2} \bar{t}^n}(x_1, x_2, t; s, \Sigma, \tau, v) = \partial_{\varphi}^{m_1} \partial_{\perp \varphi}^{m_2} \partial_{\bar{t}}^n (g(x_1 - v_1 t, x_2 - v_2 t; s \Sigma) h(t; \tau)), \quad (12)$$

where ∂_{φ} and $\partial_{\perp \varphi}$ denote directional derivative operators in the (mutually orthogonal) eigendirections φ and $\perp \varphi$ of the spatial covariance matrix Σ according to (8), $\partial_{\bar{t}}$ denotes a velocity-adapted temporal derivative operator according to (10), and $h(t; \tau)$ again represents a set of first-order truncated exponential kernels coupled in cascade, which in the most idealized situation may correspond to the time-causal limit kernel $\Psi(t; \tau, c)$ according to (6).

3 Covariance properties of the generalized Gaussian derivative model for spatio-temporal receptive fields

For processing spatio-temporal image data $f(x, t)$, convolution of f with the purely spatio-temporal smoothing kernel

T according to (1) defines spatio-temporal smoothed image data L according to

$$L(\cdot, \cdot; s, \Sigma, \tau, v) = T(\cdot, \cdot; s, \Sigma, \tau, v) * f(\cdot, \cdot), \quad (13)$$

which is referred to as a spatio-temporal scale-space representation of f .

3.1 Transformation properties under geometric image transformations

In Lindeberg (2023), covariance properties of the generalized Gaussian derivative model for receptive fields were studied in detail. It was specifically shown that:

- under *spatial scaling transformations* of the form

$$f'(x', t') = f(x, t) \quad (14)$$

for $t' = t$ and

$$x' = S_x x, \quad (15)$$

where $S_x > 0$ denotes a spatial scaling factor, the spatio-temporal scale-space representations L' and L obtained by convolving the input signals f' and f with spatio-temporal convolution kernels of the form (1) are related according to

$$L'(x', t'; s', \Sigma', \tau', v') = L(x, t; s, \Sigma, \tau, v), \quad (16)$$

provided that the spatial scale parameters and the velocity vectors are related according to

$$s' = S_x^2 s \quad \text{and} \quad v' = S_x v, \quad (17)$$

- under *spatial affine transformations* of the form

$$f_R(x_R, t_R) = f_L(x_L, t_L) \quad (18)$$

for $t_R = t_L$ and

$$x_R = A x_L, \quad (19)$$

where A denotes a 2×2 affine transformation matrix, the spatio-temporal scale-space representations L' and L obtained by convolving the input signals f_R and f_L with spatio-temporal convolution kernels of the form (1) are related according to

$$L_R(x_R, t_R; s_R, \Sigma_R, \tau_R, v_R) = L_L(x_L, t_L; s_L, \Sigma_L, \tau_L, v_L), \quad (20)$$

provided that the spatial covariance matrices and the velocity vectors are related according to

$$\Sigma_R = A \Sigma_L A^T \quad \text{and} \quad v_R = A v_L, \quad (21)$$

and provided that the other receptive field parameters are the same,

– under *temporal scaling transformations* of the form

$$f'(x', t') = f(x, t) \quad (22)$$

for $x' = x$ and

$$t' = S_t^2 t, \quad (23)$$

the spatio-temporal scale-space representations L' and L obtained by convolving the input signals f' and f with spatio-temporal convolution kernels of the form (1) are related according to

$$L'(x', t'; s', \Sigma', \tau', v') = L(x, t; s, \Sigma, \tau, v), \quad (24)$$

provided that the temporal scale parameters and the velocity parameters are related according to

$$\tau' = S_t^2 \tau \quad \text{and} \quad v' = v/S_t, \quad (25)$$

and provided that the other receptive field parameters are the same,

– under *Galilean transformations* of the form

$$f'(x', t') = f(x, t) \quad (26)$$

for $t' = t$ and

$$x' = x + u t, \quad (27)$$

the spatio-temporal scale-space representations L' and L obtained by convolving the input signals f' and f with spatio-temporal convolution kernels of the form (1) are related according to

$$L'(x', t'; s', \Sigma', \tau', v') = L(x, t; s, \Sigma, \tau, v), \quad (28)$$

provided that the velocity parameters for the two domains are related according to

$$v' = v + u \quad (29)$$

and provided that the other receptive field parameters are the same.

A notable characteristics of the transformation properties of the scale-space representations under these natural image transformations, is that the four classes of natural image transformations are not independent. Instead, for example, spatial and temporal scaling transformations affect the image velocities, beyond the spatial and temporal scale parameters, respectively. Furthermore, spatial affine transformations also affect the image velocities. For this reason, it is of interest to model all the four classes of image transformations jointly, which we will do in the next section.

4 Joint covariance property under spatial and temporal scaling transformations, spatial affine transformations and Galilean transformations

4.1 Composed geometric image transformation

Consider the composition of:

- spatial scaling transformation with spatial scaling factor S_x ,
- a spatial affine transformation with affine transformation matrix A ,
- a Galilean transformation with velocity vector u , and
- a temporal scaling transformation with temporal scaling factor S_t

of the form

$$x' = S_x A (x + u t), \quad (30)$$

$$t' = S_t t. \quad (31)$$

4.2 Joint transformation property of purely spatio-temporally smoothed image data

Prerequisites: Let us assume that we have two video sequences $f'(x', t')$ and $f(x, t)$ that are related according to (30) and (31) such that

$$f'(x', t') = f(x, t). \quad (32)$$

What we want to derive, is a relationship for how the scale-space representations L' and L of f' and f , respectively, are related, when they are defined according to

$$\begin{aligned} L(x, t; s, \Sigma, \tau, v) &= \\ &= \int_{\xi \in \mathbb{R}^2} \int_{\eta \in \mathbb{R}} T(\xi, \eta; s, \Sigma, \tau, v) f(x - \xi, t - \eta) d\xi d\eta, \end{aligned} \quad (33)$$

$$\begin{aligned} L(x', t'; s', \Sigma', \tau', v') &= \\ &= \int_{\xi' \in \mathbb{R}^2} \int_{\eta' \in \mathbb{R}} T(\xi', \eta'; s', \Sigma', \tau', v') \times \\ &\quad f(x' - \xi', t' - \eta') d\xi' d\eta'. \end{aligned} \quad (34)$$

Step I: Let us first consider how the velocity-adapted Gaussian kernel $g(x' - v't'; s' \Sigma')$ transforms in the expression for the spatio-temporal receptive field according to (1)

$$T(x', t'; s', \Sigma', \tau', v') = g(x' - v't'; s' \Sigma') h(t'; \tau'). \quad (35)$$

Expanding the expression of the function $g(x' - v't'; s' \Sigma')$ according to the definition (2) of the 2-D affine Gaussian

kernel and making use of the explicit expressions for the transformation in (30) and (31) then gives

$$\begin{aligned} g(x' - v't'; s' \Sigma') &= \\ &= \frac{1}{2\pi s' \sqrt{\det \Sigma'}} e^{-(x' - v't')^T \Sigma'^{-1} (x' - v't') / 2s'} \\ &= \frac{1}{2\pi s' \sqrt{\det \Sigma'}} \times \\ &\quad e^{-(S_x A(x+ut) - v' S_t t)^T \Sigma'^{-1} (S_x A(x+ut) - v' S_t t) / 2s'} \end{aligned} \quad (36)$$

If we, inspired by the transformation property of the spatial scale parameters under a spatial scaling transformation in (17), introduce a similar relationship

$$s' = S_x^2 s, \quad (37)$$

as well as inspired by the transformation property of the spatial covariance matrices under a spatial affine transformation in (21), introduce a similar relationship

$$\Sigma' = A \Sigma A^T, \quad (38)$$

which gives

$$\Sigma'^{-1} = (A \Sigma A^T)^{-1} = A^{-T} \Sigma^{-1} A^{-1}, \quad (39)$$

as well as

$$\det \Sigma' = |\det A|^2 \det \Sigma, \quad (40)$$

then we obtain

$$\begin{aligned} g(x' - v't'; s' \Sigma') &= \\ &= \frac{1}{2\pi S_x^2 s |\det A| \sqrt{\det \Sigma}} \times \\ &\quad e^{-(A^{-1}((S_x A x + ut) - v' S_t t))^T \Sigma^{-1} (A^{-1}((S_x A x + ut) - v' S_t t)) / (2S_x^2 s)} \end{aligned} \quad (41)$$

Notably, this expression can be written as

$$\begin{aligned} g(x' - v't'; s' \Sigma') &= \\ &= \frac{1}{2\pi s S_x^2 |\det A| \sqrt{\det \Sigma}} e^{-(x-vt)^T \Sigma^{-1} (x-vt) / 2s}, \end{aligned} \quad (42)$$

provided that the velocity parameters v and v' are related according to

$$v = -A^{-1}u/S_x + A^{-1}v'S_t/S_x, \quad (43)$$

in other words if

$$S_x A v = -u + v' S_t, \quad (44)$$

that is provided that the velocity parameters v' and v are related according to

$$v' = (S_x A v + u) / S_t. \quad (45)$$

Step II: Consider next the scale-space representation L' of f' according to (34) and perform the change of variables

$$\xi' = S_x A (\xi + u t), \quad (46)$$

$$\eta' = S_t \eta, \quad (47)$$

which for $d\xi' = d\xi'_1 d\xi'_2$ and $d\xi = d\xi_1 d\xi_2$ gives

$$d\xi' = S_x^2 |\det A| d\xi, \quad (48)$$

$$d\eta' = S_t d\eta. \quad (49)$$

Let us additionally in the convolution integral (34) transform the kernel $T(x', t'; s', \Sigma', \tau', v')$ according to (35) with its components $g(x' - v't'; s' \Sigma')$ transforming according to (42) and $h(t'; \tau')$ transforming according to (3), with the receptive field parameters transforming according to

$$s' = S_x^2 s, \quad (50)$$

$$\Sigma' = A \Sigma A^T, \quad (51)$$

$$\tau' = S_t \tau, \quad (52)$$

$$v' = (S_x A v + u) / S_t, \quad (53)$$

which gives

$$T(x', t'; s', \Sigma', \tau', v') = \frac{1}{S_x^2 |\det A| S_t} T(x, t; s, \Sigma, \tau, v). \quad (54)$$

Step III: Then, given that the functions f and f' transform according to (32), combined with the previous result that T transforms according to (54) as well as that $d\xi$, $d\xi'$, $d\eta$ and $d\eta'$ transform according to (48) and (49), we obtain

$$\begin{aligned} L(x', t'; s', \Sigma', \tau', v') &= \\ &= \int_{\xi' \in \mathbb{R}^2} \int_{\eta' \in \mathbb{R}} T(\xi', \eta'; s', \Sigma', \tau', v') \times \\ &\quad f(x' - \xi', t' - \eta') d\xi' d\eta' = \\ &= \int_{\xi \in \mathbb{R}^2} \int_{\eta \in \mathbb{R}} T(\xi, \eta; s, \Sigma, \tau, v) f(x - \xi, t - \eta) d\xi d\eta. \end{aligned} \quad (55)$$

$$= L(x, t; s, \Sigma, \tau, v). \quad (56)$$

Summary of main result: To conclude, given two video sequences f' and f that are related according to

$$f'(x', t') = f(x, t) \quad (57)$$

under a composed image transformation of the form

$$x' = S_x A (x + u t), \quad (58)$$

$$t' = S_t t, \quad (59)$$

we have shown that the corresponding spatio-temporal scale-space representations L' and L of f' and f , respectively, are related according to

$$L(x', t'; s', \Sigma', \tau', v') = L(x, t; s, \Sigma, \tau, v), \quad (60)$$

provided that the receptive field parameters transform according to

$$s' = S_x^2 s, \quad (61)$$

$$\Sigma' = A \Sigma A^T, \quad (62)$$

$$\tau' = S_t^2 \tau, \quad (63)$$

$$v' = (S_x A v + u)/S_t. \quad (64)$$

which proves the joint spatio-temporal covariance property.

This result also serves as an explicit proof of all the individual transformation properties in (Lindeberg 2023), where the explicit proofs were omitted because of lack of space.

4.3 Joint transformation properties of spatio-temporal derivatives

Let us denote spatio-temporal coordinates for the original and the transformed domains by $p = (x_1, x_2, t)^T$ and $p' = (x'_1, x'_2, t')^T$, respectively, and let us denote the components of the affine transformation matrix A by a_{ij} for i and $j \in \{1, 2\}$ and let the velocity vector in the Galilean transformation be $u = (u_1, u_2)^T$. Then, the composed image transformation according to (30) and (31) can be written

$$\begin{aligned} p' &= \begin{pmatrix} x'_1 \\ x'_2 \\ t' \end{pmatrix} = S_x \begin{pmatrix} a_{11} & a_{12} & a_{11} u_1 + a_{12} u_2 \\ a_{21} & a_{22} & a_{21} u_1 + a_{22} u_2 \\ 0 & 0 & S_t/S_x \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ t \end{pmatrix} \\ &= Q p. \end{aligned} \quad (65)$$

According to the general transformation property of derivative operators under a linear change of variables, it then follows that the spatio-temporal derivative operators in the original and the transformed domains are related by

$$\begin{aligned} \nabla_p &= \begin{pmatrix} \partial_{x_1} \\ \partial_{x_2} \\ \partial_t \end{pmatrix} \\ &= S_x \begin{pmatrix} a_{11} & a_{21} & 0 \\ a_{12} & a_{22} & 0 \\ a_{11} u_1 + a_{12} u_2 & a_{21} u_1 + a_{22} u_2 & S_t/S_x \end{pmatrix} \begin{pmatrix} \partial_{x'_1} \\ \partial_{x'_2} \\ \partial_{t'} \end{pmatrix} \\ &= Q^T \nabla_{p'}, \end{aligned} \quad (66)$$

which, in turn, gives

$$\nabla_{p'} = Q^{-T} \nabla_p \quad (67)$$

with

$$Q^{-T} = \frac{1}{S_x \det A} \times \begin{pmatrix} a_{22} & -a_{21} & 0 \\ -a_{12} & a_{11} & 0 \\ -S_x \det A u_1/S_t & -S_x \det A u_2/S_t & S_x \det A/S_t \end{pmatrix}. \quad (68)$$

In terms of vector notation, after introducing the purely spatial gradient operators $\nabla_x = (\partial_{x_1}, \partial_{x_2})$ and $\nabla_{x'} = (\partial_{x'_1}, \partial_{x'_2})$, the transformation property (66) of the spatio-temporal gradient operators can be written as

$$\nabla_x = S_x A^T \nabla_{x'}, \quad (69)$$

$$\partial_t = S_t \partial_{t'} + S_x u^T A^T \nabla_{x'}, \quad (70)$$

whereas the corresponding inverse relationship (67) can be expressed as

$$\nabla_{x'} = \frac{1}{S_x} A^{-T} \nabla_x, \quad (71)$$

$$\partial_{t'} = \frac{1}{S_t} (\partial_t - u^T \nabla_x). \quad (72)$$

Based on these relations, expressions for spatio-temporal derivatives can be transformed between the two domains under the composed image transformation, thus extending the transformation property (60) of the spatio-temporal receptive fields to beyond the effect of purely spatio-temporal smoothing operation in the spatio-temporal receptive fields also cover the spatio-temporal derivative operators in the receptive field models.

For example, given the vector notation for the derivative operators, the velocity-adapted derivative operators corresponding to (10) are with $v = (v_1, v_2)^T$ and $v' = (v'_1, v'_2)^T$ given by

$$\partial_{\bar{t}} = v^T \nabla_x + \partial_t \quad \text{and} \quad \partial_{\bar{t}'} = v'^T \nabla_{x'} + \partial_{t'}, \quad (73)$$

where velocity vectors v and v' are related according to (64).

Similarly, the directional derivative operators ∂_φ and $\partial_{\varphi'}$ corresponding to (8) are with the unit vectors

$$e_\varphi = (\cos \varphi, \sin \varphi)^T \quad \text{and} \quad e_{\varphi'} = (\cos \varphi', \sin \varphi')^T \quad (74)$$

given by

$$\partial_\varphi = e_\varphi^T \nabla_x \quad \text{and} \quad \partial_{\varphi'} = e_{\varphi'}^T \nabla_{x'}. \quad (75)$$

In the special case when the affine transformation matrix A is a rotation matrix $A = R_\theta$, the eigenvectors of covariance matrix Σ do also transform according to a rotation, according to (62), implying that the rotational angles φ and φ' will be related according to

$$\varphi' = \varphi + \theta. \quad (76)$$

For general affine transformations, the relation will, however, be more complex.

By exploiting these transformation properties of spatio-temporal gradient operators, we can in a geometric way transform all the spatio-temporal derivatives of the spatio-temporal receptive fields described in Sections 2.2 and 2.3.

5 Summary and discussion

We have presented a detailed theoretical analysis of covariance properties of the spatio-temporal receptive fields according to the generalized Gaussian model for spatio-temporal receptive fields. Specifically, we have derived a joint covariance property for the composition of a spatial scaling transformation, a spatial affine transformation, a Galilean transformation and a temporal scaling transformation, with explicit expressions for how the receptive field parameters should be transformed under the composed image transformation to make it possible to perfectly match the receptive field responses under convolutions with spatio-temporal receptive fields according to the generalized Gaussian derivative model.

While previous work with the generalized Gaussian derivative model for spatio-temporal receptive fields have primarily focused on using either the non-causal 1-D Gaussian kernel or the time-causal limit kernel for temporal smoothing in the spatio-temporal smoothing process, the derivations in this paper have been made under a weaker assumption of only requiring temporal scale covariance (according to (3)) for the temporal smoothing kernels. Thus, the results presented in this article do immediately generalize to the use of other temporal smoothing kernels, provided that they are covariant under temporal scaling transformations.

The theoretical results in this treatment are intended as a theoretical foundation for computer vision modules that make use of populations of spatio-temporal receptive field responses as the first processing layers in the visual hierarchy, as well as for formulating and interpreting models of biological vision.

Acknowledgements

I would like to thank Jens Pedersen for valuable interactions concerning the joint covariance property of the spatio-temporal smoothing transformation.

References

B. R. Conway and M. S. Livingstone. Spatial and temporal properties of cone signals in alert macaque primary visual cortex. *Journal of Neuroscience*, 26(42):10826–10846, 2006.

- G. C. DeAngelis and A. Anzai. A modern view of the classical receptive field: Linear and non-linear spatio-temporal processing by V1 neurons. In L. M. Chalupa and J. S. Werner, editors, *The Visual Neurosciences*, volume 1, pages 704–719. MIT Press, 2004.
- G. C. DeAngelis, I. Ohzawa, and R. D. Freeman. Receptive field dynamics in the central visual pathways. *Trends in Neuroscience*, 18(10):451–457, 1995.
- E. N. Johnson, M. J. Hawken, and R. Shapley. The orientation selectivity of color-responsive neurons in Macaque V1. *The Journal of Neuroscience*, 28(32):8096–8106, 2008.
- T. Lindeberg. Generalized Gaussian scale-space axiomatics comprising linear scale-space, affine scale-space and spatio-temporal scale-space. *Journal of Mathematical Imaging and Vision*, 40(1):36–81, 2011.
- T. Lindeberg. Time-causal and time-recursive spatio-temporal receptive fields. *Journal of Mathematical Imaging and Vision*, 55(1):50–88, 2016.
- T. Lindeberg. Normative theory of visual receptive fields. *Heliyon*, 7(1):e05897:1–20, 2021. doi: 10.1016/j.heliyon.2021.e05897.
- T. Lindeberg. Covariance properties under natural image transformations for the generalized Gaussian derivative model for visual receptive fields. *Frontiers in Computational Neuroscience*, 17:1189949:1–23, 2023.