### SPARSE REGULAR SUBSETS OF THE REALS

JASON BELL AND ALEXI BLOCK GORMAN

ABSTRACT. This paper concerns the expansion of the real ordered additive group by a predicate for a subset of [0, 1] whose base-*r* representations are recognized by a Büchi automaton. In the case that this predicate is closed, a dichotomy is established for when this expansion is interdefinable with the structure  $(\mathbb{R}, <, +, 0, r^{-\mathbb{N}})$  for some  $r \in \mathbb{N}_{>1}$ . In the case that the closure of the predicate has Hausdorff dimension less than 1, the dichotomy further characterizes these expansions of  $(\mathbb{R}, <, +, 0, 1)$  by when they have NIP and NTP<sub>2</sub>, which is precisely when the closure of the predicate has Hausdorff dimension 0.

#### 1. INTRODUCTION

This paper concerns how geometric and model-theoretic notions of tameness interact for expansions of the real ordered additive group by a subset of [0, 1] that is *r*-regular. Notions from the theory of finite automata are adapted to the real additive group setting, and used to establish a partial tameness dividing line for these expansions. Key tools include examining the interactions of automaton-theoretic properties and topological properties of subsets of the real numbers.

A finite automaton is a machine with finitely many states, some subset of which are called accepting states, with the remaining states being called rejecting states. The machine takes finitelength strings over a fixed finite alphabet  $\Sigma$  as input, and, beginning at a fixed starting state, it moves from state to state based upon simple transition rules as it reads the string from left-to-right. The machine then either accepts or rejects a string depending upon whether or not it arrives in an accepting state after it has finished reading the word.

Büchi automata differ from classical finite automata in that they take infinite-length strings over  $\Sigma$  as input. This is accomplished by taking a finite automaton and imposing the acceptance condition that an infinite string  $w \in \Sigma^{\omega}$  is accepted precisely if a run of the automaton on w enters an accept state infinitely often. For both kinds of automata, we say that an automaton *recognizes* a set X (either a subset of  $\Sigma^*$  or, if it is a Büchi automaton, a subset of  $\Sigma^{\omega}$ ) if every element of X is accepted by the automaton, and no element of  $X^c$  (the complement) is accepted. Büchi automata have long been an important object of study in logic, combinatorics on words, and theoretical computer science. Much of the context for the results of this paper and many of the connections between Büchi automata and topological phenomena on the reals come from [CLR15].

We borrow the notion of "sparse" sets recognized by automata. These sets arise naturally in many contexts, including: Derksen's [Der07] extension of the Skolem-Mahler-Lech theorem on zero sets of linear recurrences to positive characteristic base fields; the isotrivial case of the Mordell-Lang

Date: November 21, 2023.

<sup>2020</sup> Mathematics Subject Classification. Primary 03C64, 03D05 Secondary 28A80.

Key words and phrases. Büchi automata, Finite automata, Hausdorff dimension, Model theory, Tame geometry, Definability, D-minimality.

The first-named author was supported by NSERC grant RGPIN-2016-03632. The second-named author was supported by the National Science Foundation Graduate Research Fellowship Program under Grant No. DGE -1746047.

conjecture [BGM20, BM19]; the characterization of which k-automatic sets  $S \subseteq \mathbb{N}$  form an additive basis for the natural numbers [BHS18]; Kedlaya's description of the algebraic closure of function fields in positive characteristic [Ked06], [Ked17]; as well, as other uses in theoretical computer science [GKRS10, IR86, Tro81].

We introduce a natural analog for sparsity in the setting of Büchi automata, one which we demonstrate is useful in the setting of real *r*-regular sets. Among other things, the notion of sparsity for Büchi automata allows for the development of a definability dividing line simultaneously in terms of fractal dimension and model-theoretic notions.

To state this result formally requires first defining what it means for a subset of  $[0,1] \subseteq \mathbb{R}$  to be *r*-regular. Below, let  $r \in \mathbb{N}$  be greater than one, and set  $[r] := \{0, \ldots, r-1\}$ . We also let  $[r]^{\omega}$  denote the set of all functions from the ordinal  $\omega$  to the set [r].

**Definition 1.1.** Say that  $A \subseteq [0,1]$  is *r*-regular if there is a Büchi automaton  $\mathcal{A}$  with alphabet  $\{0,\ldots,r-1\}$  that recognizes a set  $L \subseteq \{0,\ldots,r-1\}^{\omega}$  such that  $(w_i)_{i<\omega} \in L$  if and only if there is  $x \in A$  such that

$$x = \sum_{i=0}^{\infty} \frac{w_i}{r^{i+1}}$$

Moreover, if this holds say that  $\mathcal{A}$  recognizes A.

**Definition 1.2.** Say that  $A \subseteq [0, 1]$  is *r*-sparse if it is *r*-regular and whenever  $\mathcal{A}$  is an automaton that recognizes A, the set of strings in  $\{0, \ldots, r-1\}^n$  with an infinite prolongation that is accepted by  $\mathcal{A}$  grows at most polynomially in n.

It is not hard to show that in Definition 1.2, if one automaton recognizing the set A has the property that the set of strings that have an infinite prolongation that is accepted by the automaton grows polynomially, then this holds for all automata recognizing A.

Recall that in [vdD85], van den Dries proves that the expansion of both the real additive group and the reals as an ordered field by a predicate for  $2^{\mathbb{Z}}$  has desirable tameness properties, including d-minimality (see [Mi05] for an introduction to the notion) and decidability. The results of [vdD85] and the following interdefinability result imply many desirable properties hold for the expansion  $\mathcal{R}_A$  of ( $\mathbb{R}, <, +, 0, 1$ ) by a predicate A that picks out an r-sparse subset of [0, 1]. Below, set  $\mathcal{R}_A :=$ ( $\mathbb{R}, <, +, 0, 1, A$ ), and set  $\mathcal{R}_{r,\ell} := (\mathbb{R}, <, +, 0, 1, r^{-\ell \mathbb{N}})$ .

**Theorem 1** (Corollary 3.10). Let r > 1 be a natural number, and suppose  $A \subseteq [0,1]$  is r-sparse. Then there exists  $\ell \in \mathbb{N}$  such that A is  $\emptyset$ -definable in  $\mathcal{R}_{r,\ell}$ , and the set  $r^{-\ell \mathbb{N}}$  is  $\emptyset$ -definable in  $\mathcal{R}_A$ .

Recall that  $X \subseteq \mathbb{R}^d$  is called a *Cantor set* if it is compact, has no isolated points, and no interior. From d-minimality, we can conclude that for an *r*-sparse set A the structure  $\mathcal{R}_A$  does not define a Cantor set. Theorem 1, in conjunction with the following result, gives us a characterization of what kind of definable sets show up in expansions of the real ordered additive group by a predicate for an *r*-regular subset of [0, 1]. Below, for  $X \subseteq \mathbb{R}$  let  $d_H(X)$  denote the Hausdorff dimension of X.

**Theorem 2** (Theorem 4.5). If A is a closed r-regular subset of [0,1] such that  $0 < d_H(A) < 1$ , then there is a Cantor set definable in  $\mathcal{R}_A$ .

Theorem 2 demonstrates the connection of fractal dimension, namely Hausdorff dimension, to definability of a more "pathological" set, in this case a Cantor set. What makes a Cantor set "pathological" in this setting is that work of Hieronymi and Walsberg in [HW19] shows that the expansion of  $(\mathbb{R}, <, +, 0)$  by a Cantor set is not model-theoretically tame.

A couple notions of Shelah-style tameness in the context of model theory include NIP (also known as not the independence property) and  $NTP_2$  (also known as not the tree property of the second kind). These notions correspond to combinatorial properties of formulas modulo a specific theory, or of the sets definable in the models of a given theory. The property NIP for formulas corresponds to finiteness of V-C dimension, a notion of interest to some computer scientists. For definitions of NIP and  $NTP_2$ , see [Sim15].

Let  $\pi_i : \mathbb{R}^n \to \mathbb{R}$  denote the projection of  $\vec{x} \in \mathbb{R}^n$  onto the *i*-th coordinate. Connecting the work of Hieronymi and Walsberg with Theorem 2, we obtain a theorem that solidifies the notion that for these structures, tameness in the model-theoretic sense and tameness in the sense of fractal geometry completely coincide.

**Theorem 3** (Theorem 4.6). For  $A \subseteq [0,1]^d$  an *r*-regular set such that  $d_H(\overline{\pi_i(A)}) < 1$  for all  $i \in [d]$ , the following are equivalent:

- (1) A is r-sparse; (2)  $d_H(\overline{\pi_i(A)}) = 0$  for all  $i \in [d]$ ;
- (3)  $\mathcal{R}_A$  is d-minimal;
- (4) The theory of  $\mathcal{R}_A$  has NIP;
- (5) The theory of  $\mathcal{R}_A$  has  $NTP_2$ .

Note that in this final theorem, we have dropped the assumption that A is closed, illustrating how well the topological closures of these r-regular sets reflect or control the behavior of even the non-closed r-regular sets.

#### Acknowledgements

We thank Rahim Moosa and Christa Hawthorne for many valuable contributions to this project and for many helpful comments. Many thanks also to Philipp Hieronymi for his insightful comments and suggestions.

1.1. Background on automata theory. Throughout this paper, all terminology, notation, and definitions relating to model theory are drawn from [M02], unless stated otherwise. In recent years, there have been new applications of automata theory to the model theory of tame structures. Uses of finite automata in recent works of model theorists include the introduction of *F*-sets by Moosa and Scanlon in [MS02] and [MS04] with applications to isotrivial Mordell-Lang in positive chracteristic, which were shown to be recognized by finite automata in a particular sense by Bell and Moosa in [BM19]. Other applications include tameness results, namely stability, of the expansion of  $(\mathbb{Z}, +)$  by a predicate for a subset of  $\mathbb{Z}$  recognized by an automaton with a specific form of alphabet [Haw20].

The connections between model theory and automata theory have inspired the work in this paper centered around expansions of the real ordered additive group by sets recognized by Büchi automata. Recent work in this area includes chracterizing continuous functions in expansion of the real additive group by sets recognized by Büchi automata; in [BG et al.20], the authors show that a continuous real-valued function on a closed interval whose graph is recognized by a Büchi automaton must be locally affine.

As is standard, if X is a set, then  $X^{\omega}$  denotes the set of all sequences of elements of X indexed by  $\omega$ , or equivalently all functions from the ordinal  $\omega$  to the set X. The following facts and definitions about finite automata can all be found in [Sip13]. A *finite automaton* is a 5-tuple  $(Q, \Sigma, \delta, q_0, F)$  such that:

• Q is a finite set of states,

- $\Sigma$  is a finite alphabet,
- $\delta: Q \times \Sigma \to Q$  is a transition function, or rather partial function;  $(q_i, \alpha) \mapsto q_j$  or  $(q_i, \alpha) \mapsto \emptyset$ ,
- $q_0$  is the initial state,
- $F \subseteq Q$  is the set of accept states (also called final states).

If  $\mathcal{A}$  is a finite automaton and  $w = \sigma_1 \dots \sigma_n$  is a finite string generated from  $\Sigma$ , i.e.  $\sigma_i \in \Sigma$  for each  $i \leq n$ , then a *run* of  $\mathcal{A}$  on w is a sequence of states  $q_0 \dots q_n$  where each  $q_i$  is a state in  $\mathcal{A}$ , the state  $q_0$  is the start state, and for each  $1 \leq i \leq n$  we have  $\delta(q_{i-1}, \sigma_i) = q_i$ . Say that a finite automaton  $\mathcal{A}$  accepts a finite string w generated from  $\Sigma$  precisely if a run of  $\mathcal{A}$  on w terminates in an accept state, i.e. a state in F.

The definition above is of a *deterministic* finite automaton. We can generalize this definition to *nondeterministic* finite automata by changing  $\delta$  from a partial function  $\delta : Q \times \Sigma \to Q$  to a full function on  $Q \times \Sigma$  that takes values in  $\mathcal{P}(Q)$ , the powerset of Q. In other words, we allow  $\delta : Q \times \Sigma \to \mathcal{P}(Q)$  for the transition function. The notion of a "run" is adapted accordingly. Conveniently, for finite automata the class of deterministic finite automata and the class of nondeterministic finite automata are equivalent, in the sense that for each non-deterministic finite automaton there is a deterministic one that accepts precisely the same words, and vice versa.

For any finite alphabet  $\Sigma$ , let  $\Sigma^*$  denote all finite length strings  $\Sigma$  generates. Given any set X of finite length strings, let  $X^* = \{x_1 x_2 \dots x_n : x_i \in X, n \in \mathbb{N}\}$ , where  $x_i x_j$  denotes the concatenation of  $x_i$  by  $x_j$  on right. We call  $X^*$  the Kleene star of X. Call any subset  $L \subseteq \Sigma^*$  a language. Say that automaton  $\mathcal{A}$  recognizes L if for all  $w \in \Sigma^*$ ,  $\mathcal{A}$  accepts w if and only if  $w \in L$ . Regular languages, the subsets of  $\Sigma^*$  recognized by some finite automaton, are closed under complementation, taking finite unions and intersections, concatenation, and the Kleene star operation.

We will often conflate regular languages with what are called *regular expressions*. Regular expressions of a finite alphabet  $\Sigma$  are a class of strings generated from  $\Sigma$  via union, concatenation, and Kleene star. In other words, the regular expressions on  $\Sigma$  include the empty set  $\emptyset$ , the empty string  $\varepsilon$ , and each character  $\sigma \in \Sigma$ , as well as all expressions one can generate by starting with those symbols and iteratively applying the operations union, concatenation, and Kleene star. For more details on finite-state automata, regular languages, and regular expressions, we refer the reader to [AS03].

Büchi automata are still given by 5-tuples of the form  $(Q, \Sigma, \delta, q_0, F)$  as defined above, the difference is that their inputs are strings from  $\Sigma^{\omega} = \{(\sigma_i)_{i \in \omega} : \sigma_i \in \Sigma\}$  rather than  $\Sigma^*$ , and the acceptance condition differs. We define a run of a Büchi automaton  $\mathcal{A}$  on an element  $w \in \Sigma^{\omega}$  in an analogous manner; a *run* of  $\mathcal{A}$  on w is a sequence of states  $(q_i)_{i < \alpha}$  where  $\alpha \in \omega \cup \{\omega\}$  and each  $q_i$  is a state in  $\mathcal{A}$ , the state  $q_1$  is the start state  $q_0$ , and for each  $i < \alpha$  we have  $\delta(q_i, \sigma_i) = q_{i+1}$ . We say that a Büchi automaton  $\mathcal{A}$  accepts a string  $w \in \Sigma^{\omega}$  precisely if a run of  $\mathcal{A}$  on w includes an accept state infinitely often. Note that in the definition of a run for Büchi automata, if  $\alpha < \omega$ , then the string w cannot be accepted, since the run terminates despite w being an infinite string.

Since for Büchi automata the alphabet  $\Sigma$  is still a finite set, the languages that Büchi automata recognize can all be viewed as subsets of a Cantor space, namely the space  $\{0, \ldots, n-1\}^{\omega}$  if  $|\Sigma| = n$ . Now we can restate Definition 1.1 more clearly, and for subsets of  $[0, 1]^m$ . Set  $[r] := \{0, \ldots, r-1\}$ , so that  $([r]^m)^{\omega} = \{\{0, \ldots, r-1\}^m\}^{\omega}$ . For  $(x_1, \ldots, x_m) \subseteq [0, 1]^m$ , say that  $(w_{1,i}, \ldots, w_{m,i})_{i < \omega} \in ([r]^m)^{\omega}$  is an *r*-representation of  $(x_1, \ldots, x_m)$  if for each  $j \leq m$  the following holds:

$$x_j = \sum_{i=0}^{\infty} \frac{w_{j,i}}{r^{i+1}}.$$

**Definition 1.3.** For  $m, r \in \mathbb{N}_{>0}$  with r > 1, say that  $X \subseteq [0,1]^m$  is *r*-regular if there is a Büchi automaton  $\mathcal{A}$  such that the following holds:

- (1) for each  $w \in ([r]^m)^{\omega}$ , if w is not an r-representation for any  $x \in X$ , then  $\mathcal{A}$  does not accept w;
- (2) for each  $x \in X$ , there exists a  $w \in ([r]^m)^{\omega}$  such that w is an *r*-representation of x and  $\mathcal{A}$  accepts w.

If such an  $\mathcal{A}$  exists, say that  $\mathcal{A}$  recognizes X.

Note that unlike with finite automata, reversing the accept and non-accept states of a Büchi automaton does not yield an automaton that recognizes the complement of the language that the original automaton recognized. Just as there is a correspondence between finite automata and regular languages, there is an analogous correspondence between Büchi automata and regular  $\omega$ languages. Call any set  $L \subseteq \Sigma^{\omega}$  an  $\omega$ -language. Say that the Büchi automaton  $\mathcal{A}$  recognizes L if for all  $w \in \Sigma^{\omega}$ ,  $\mathcal{A}$  accepts w if and only if  $w \in L$ . Say that  $L \subseteq \Sigma^{\omega}$  is a regular  $\omega$ -language if there is a Büchi automaton that recognizes L. It is again true, though less easy to show, that regular  $\omega$ -languages are closed under the boolean operations union, intersection, and complementation. Suppose that  $V \subseteq \Sigma^*$  is a regular language and  $L \subseteq \Sigma^{\omega}$  is a regular  $\omega$ -language. Then the following are regular  $\omega$ -languages as well:

- $VL = \{vw \in \Sigma^{\omega} : v \in V, w \in L\},\$
- $V^{\omega} = \{v_1 v_2 v_3 \ldots \in \Sigma^{\omega} : v_i \in V\}.$

There is an important theorem of Büchi's that makes working with regular  $\omega$ -languages significantly easier. We will use the following theorem frequently in this paper without direct reference.

**Theorem 1.4** ([Büc62]). For every  $L \subseteq \Sigma^{\omega}$  recognized by a Büchi automaton, there are regular languages  $V_1, \ldots, V_k, W_1, \ldots, W_k \in \Sigma^*$  such that

$$L = \bigcup_{i=1}^{k} V_i W_i^{\omega}.$$

This theorem allows us to understand what regular  $\omega$ -languages look like in terms of regular languages on the same alphabet. We can define regular  $\omega$ -expressions analogously to regular expressions, though by the above theorem every regular  $\omega$ -expression is equivalent to one of the form

$$v_1 w_1^\omega \cup \cdots \cup v_n w_n^\omega$$

where  $v_1, \ldots, v_n, w_1, \ldots, w_n$  are each regular expressions generated from  $\Sigma$ , and  $w_i^{\omega}$  is the infinitary version of Kleene star.

**Definition 1.5.** Suppose that  $\mathcal{A}$  is a Büchi automaton.

- (1) For states p and q in an automaton, say q is *accessible* from p if there exists a run of some non-trivial word from p to q.
- (2) Say that  $\mathcal{A}$  is *weak* if for any states p and q in  $\mathcal{A}$ , whenever p and q are accessible to and from each other, then one is an accept state precisely if the other is an accept state.
- (3) Say that  $\mathcal{A}$  is *trim* if each state is accessible from the start state and some accept state is accessible from each state (via a nontrivial path).
- (4) If  $\mathcal{A}$  is trim, say that  $\mathcal{A}$  is *closed* if every state is an accept state.
- (5) Let  $\overline{\mathcal{A}}$  be the automaton resulting from making every state of  $\mathcal{A}$  accepting.

**Fact 1.6** ([CLR15, Remark 59]). If the set  $X \subseteq [0,1]$  is recognized by a Büchi automaton  $\mathcal{A}$ , then its closure in the order topology  $\overline{X}$  is recognized by  $\overline{\mathcal{A}}$ .

Some of the earliest connections made between Büchi automata and model theory come from the work of Boigelot, Rassart, and Wolper in [BRW98]. In that extended abstract, the authors introduce the following ternary predicate.

**Definition 1.7.** Let  $V_r(x, u, k)$  be a ternary predicate on  $\mathbb{R}$  that holds precisely if  $u = r^{-n}$  for some  $n \in \mathbb{N}_{>0}$  and the *n*-th digit of a base-*r* representation of *x* is *k*.

By examining the expansion of the real ordered additive group by this predicate, they prove the following definability result.

**Theorem 1.8** ([BRW98]). A subset  $X \subseteq [0,1]^n$  is r-regular if and only if X is  $\emptyset$ -definable in  $(\mathbb{R}, <, 0, +, V_r)$ .

This theorem has given rise to many further applications of automata theory to model theory.

1.2. Background on fractal dimensions. Finally, we will recall the definitions of Hausdorff measure and Hausdorff dimension for subsets of  $\mathbb{R}$ . For  $X \subseteq \mathbb{R}$  and  $\varepsilon > 0$ , let  $\mathcal{B}_{\varepsilon}$  be the collection of all countable covers of X by closed intervals of diameter at most  $\varepsilon$ . We let  $\text{Diam}(U) \in \mathbb{R} \cup \{\infty\}$  denote the diameter of  $U \subseteq \mathbb{R}$ . Define Hausdorff s-measure of  $X \subseteq \mathbb{R}$  as follows:

$$\mu_{H}^{s}(X) = \lim_{\varepsilon \to 0} \inf_{B \in \mathcal{B}_{\varepsilon}} \Big\{ \sum_{i=1}^{\infty} (\operatorname{Diam} U_{i})^{s} : U_{i} \in B \Big\}.$$

The Hausdorff dimension of X is written  $d_H(X)$ , and is the unique  $s \in \mathbb{R}_{\geq 0}$  such that if s' > s, then  $\mu_H^{s'}(X) = 0$  and if s' < s, then  $\mu_H^{s'}(X) = \infty$ . Note that it is quite possible that  $\mu_H^s(X) = 0$ , and it is possible that s = 0 (indeed, if X is a single point, then  $d_H(X) = 0$ ). If  $X \subseteq \mathbb{R}^n$ , the Lebesgue measure of X is positive and finite if and only if  $\mu_H^n(X)$  is positive and finite. Thus it is not possible that  $\mu_H^s(X) = \infty$  for all s > 0.

Hausdorff dimension behaves well with respect to subsets and unions, as shown by this fact:

**Fact 1.9** ([Fal03]). Let X, Y, and  $(Y_i)_{i < \omega}$  be subsets of  $\mathbb{R}$ .

- (1) If  $X \subseteq Y$ , then  $d_H(X) \leq d_H(Y)$ .
- (2) If  $X = Y_1 \cup Y_2$ , then  $d_H(X) = \max(d_H(Y_1), d_H(Y_2))$ .
- (3) If  $X = \bigcup_{i < \omega} Y_i$ , then  $d_H(X) = \max_{i < \omega} \{ d_H(Y_i) \}$ .

1.3. Background on S-unit theory. In this brief section we give an overview of the theory of S-unit equations. Specifically, we require a quantitative version of a result due to Evertse, Schlickewei and Schmidt (see [ESS02, Theorem 1.1] and also [EG15, Theorem 6.1.3]).

**Definition 1.10.** Let *L* be a field and let  $a_1, \ldots, a_m \in L$  be nonzero elements of *L*. Then a solution  $x_1, \ldots, x_m \in L$  to the equation

$$a_1x_1 + \dots + a_mx_m = 1$$

is called *nondegenerate* if the left side has no non-trivial vanishing subsums; i.e., if  $\sum_{i \in I} a_i x_i \neq 0$  for each non-empty subset I of  $\{1, \ldots, m\}$ . In general, we will say a sum  $\sum_{i=1}^m x_i$  is *nondegenerate* if it has no non-trivial vanishing subsums.

The following theorem is usually referred as the S-unit theorem.

**Theorem 1.11.** Let L be a field of characteristic zero, let  $a_1, \ldots, a_m$  be nonzero elements of L, and let  $H \subset (L^{\times})^m$  be a finitely generated multiplicative group. Then there are only finitely many nondegenerate solutions  $(y_1, \ldots, y_m) \in H$  to the equation  $a_1y_1 + \cdots + a_my_m = 1$ . (See Theorem 6.1.3 in [EG15]—although this theorem is only stated for  $m \ge 2$ , the case m = 1 is immediate.)

To show that sparse k- and  $\ell$ -regular sets have finite intersection when k and  $\ell$  are multiplicatively independent, we will also use the following theorem of Evertse, Schlickewei, and Schmidt (2002) (see [EG15, Theorem 6.1.3]).

**Theorem 1.12.** Let K be a field of characteristic zero, let  $n \ge 2, r \ge 1$ , let  $\Gamma \le K^*$  be a finitely generated subgroup of rank r, and let  $a_1, \ldots, a_n$  be nonzero elements of K. Then the number of nondegenerate solutions to

$$a_1x_1 + \dots + a_nx_n = 1$$

with  $(x_1,\ldots,x_n) \in \Gamma^n$  is at most  $\exp((6n)^{3n}(r+1))$ .

# 2. Preliminaries on sparsity

Let  $r \in \mathbb{N}_{>1}$ , and set  $[r] := \{0, 1, \dots, r-1\}$  for the remainder of this paper. Note that we will often switch between elements of [0, 1] and their *r*-representations, and may sometimes say that an automaton  $\mathcal{A}$  accepts the *r*-representation of  $x \in [0, 1]$ . For the countable subset of [0, 1] whose elements have multiple (i.e. exactly two) *r*-representations, we mean that  $\mathcal{A}$  accepts at least one of the *r*-representations of *x*. For ease of switching between *x* and its *r*-representation, we will define a valuation for elements of  $[r]^{\omega}$ .

**Definition 2.1.** Define  $\nu_r : [r]^{\omega} \to [0, 1]$  by:

$$\nu_r(w) = \sum_{i=0}^{\infty} \frac{w_i}{r^{i+1}}$$

where  $w = w_0 w_1 w_2 \dots$  with  $w_i \in [r]$  for each  $i \in \mathbb{N}$ .

Note that the equivalence relation  $v \equiv w \iff \nu_r(v) = \nu_r(w)$  is not only a finite equivalence relation, but moreover each equivalence class has size at most two. As noted above, only countably many elements in  $[r]^{\omega}$  are not the unique element of their  $\nu_r$ -equivalence class. For  $L \subseteq [r]^{\omega}$  an  $\omega$ -language, we set  $\nu_r(L) := \{\nu_r(w) : w \in L\}$ . If  $L \subseteq ([r]^m)^{\omega}$ , set

$$\nu_r(L) := \{ (\nu_r(w_1), \dots, \nu_r(w_m)) : w_1, \dots, w_m \in [r]^{\omega}, (w_{1,i}, \dots, w_{m,i})_{i < \omega} \in L \}.$$

We can now expand Definition 1.2 to languages in  $([r]^m)^{\omega}$ .

**Definition 2.2.** Say that  $L \subseteq ([r]^m)^{\omega}$  is *sparse* if it is regular (as an  $\omega$ -language) and the set of length-*n* prefixes of the elements of *L* grows at most polynomially in *n*.

Say that  $X \subseteq [0,1]$  is *r*-sparse if  $X = \nu_r(L)$  for some sparse  $L \subseteq ([r]^m)^{\omega}$ , and say that a Büchi automaton is sparse if the language that it recognizes is sparse.

There is a dichotomy for the set of length-*n* prefixes of an *r*-regular  $\omega$ -language: either the set of length-*n* prefixes grows polynomially in *n*, or it grows exponentially in *n* (see Remark 2.5).

For any word  $w = (s_1, \ldots, s_d) \in ([r]^m)^*$ , where  $s_j \in [r]^*$  for each  $j \leq d$ , say that w is a *uniform* word if  $|s_1| = |s_2| = \cdots = |s_d|$ ; i.e., the length of each coordinate word  $s_j$  is the same for all  $j \leq d$ . Suppose  $L \subseteq ([r]^m)^{\omega}$  is r-regular. We will see that a characterization of sparseness in [BM19] implies that if L is sparse, then L is a finite union of sets of the following form:

(1) 
$$u_1 v_1^* u_2 v_2^* \dots u_{d-1} v_{d-1}^* u_d v_d^{\omega}$$

where  $u_i$  and  $v_i$  are all uniform words in  $([r]^m)^*$  for all  $i \leq d$ , and  $v_1, \ldots, v_d$  are non-empty strings. If L is sparse and a union of n sets of form (1), let  $W_L := \{u_{1,1}, v_{1,1}, \ldots, u_{m,d}, v_{m,d}\}$  be the words in  $([r]^m)^*$  as specified below (1) that witness this. Let  $\ell_L$  be the least common multiple of  $\{|w|: w \in W_L\}$ .

By Theorem 1.4, L is a finite union of sets of the form  $V_i W_i^{\omega}$  where  $V_i, W_i \subseteq ([r]^m)^*$ , so we can reduce to showing that if one such  $VW^{\omega}$  is not of the form (1), then the set of length n prefixes of L grows exponentially in n. We will sometimes call these  $V_i W_i^{\omega}$  subsets of L the V-W components of L.

**Proposition 2.3.** A language  $L \subseteq ([r]^m)^{\omega}$  is sparse if and only if L is a finite union of sets of the form

 $u_1 v_1^* u_2 v_2^* \dots v_{d-1}^* u_d v_d^{\omega}$ 

where the  $u_i$  and  $v_i$  are all uniform words in  $([r]^m)^*$  for  $i \leq d$ , and  $v_1, \ldots, v_d$  are non-empty strings.

*Proof.* For the forward direction, we prove the contrapositive, namely that if L is not a finite union of sets of the form  $u_1v_1^*u_2v_2^*\ldots v_{d-1}^*u_dv_d^{\omega}$ , then the set of length n prefixes of L grows exponentially in n.

First, for any finite  $\Sigma$ , we observe that if  $a, b \in \Sigma^*$  and  $\{a, b\}^*$  contains at most one word of length k for each  $k \in \mathbb{N}$ , then ab = ba. So by Corollary 6.2.5 in [Lot02], we conclude that  $\{a, b\}^* \subseteq c^*$  for some word  $c \in \Sigma^*$ . We consider one of the V-W components of L, say  $VW^{\omega}$ , and fix this component for the remainder of the proof. Note that if V is not of the form  $u_1v_1^*u_2v_2^*\ldots v_{d-1}^*u_dv_d^*$ , then by the characterization of sparseness in [BM19] for subsets of  $([r]^m)^*$ , the length n prefixes of V will grow exponentially in n, and hence so too will the length n prefixes of L. If V is of the above form, but W is not generated by a single word, i.e.  $W^*$  contains more than one word of length k for some  $k \in \mathbb{N}$ , call them a and b, then for some  $v \in V$  the set of words of the form  $v\{a, b\}^{\omega}$  is contained in  $VW^{\omega}$ , and the length n prefixes of these words are the same as those of  $v\{a, b\}^*$ , whose length n prefixes grow exponentially with respect to n, as shown in [BM19].

For the backwards direction, if L is a finite union of sets of the form  $u_1v_1^*u_2v_2^*\ldots v_{d-1}^*u_dv_d^\omega$ , then the length n prefixes are the same as those of  $u_1v_1^*u_2v_2^*\ldots v_{d-1}^*u_dv_d^*$ , which by [BM19] grow polynomially in n, as desired.

Assuming the automata for our regular sets are trim, the closure of the automaton accepts the closure of the regular set. For r-sparse subsets of  $[0, 1]^m$ , we will see that the closures of these sets are finite unions of sets of this form:

$$\nu_r(u_1v_1^*\ldots u_nv_n^{\omega}) \cup \nu_r(u_1v_1^*\ldots u_{n-1}v_{n-1}^{\omega}) \cup \cdots \cup \nu_r(u_1v_1^{\omega}).$$

For the remainder of this section, we restrict our attention to the set of closed *r*-regular subsets of  $[0, 1]^m$ . The characterization of non-sparse regular languages from [BM19] implies that if  $X \subseteq [0, 1]^m$  is closed but not *r*-sparse, then X contains a set of the form  $Y = \nu_r(u\{a, b\}^{\omega})$ , i.e., a translate of a Cantor set. It is not clear *a priori* that Y is definable in the language of ordered additive groups expanded by a predicate for X.

The following lemma will be crucial for a subsequent characterization of expansions of the real ordered additive group by a predicate for an r-regular set, among other things.

**Lemma 2.4.** Suppose that X is a closed r-regular subset of  $[0,1]^m$ . The following are equivalent:

- (1) X is r-sparse;
- (2)  $X \subseteq \mathbb{Q}^m$ ;
- (3) X is countable;
- (4) the set of accumulation points of X is countable.

Proof. Let  $L \subseteq ([r]^m)^{\omega}$  be such that  $\nu_r(L) = X$ . For  $(1) \Longrightarrow (2)$ , we observe that since X is r-sparse, by Proposition 2.3 it is a finite union of sets of the form  $\nu_r(u_1v_1^* \dots u_nv_n^{\omega})$ . Hence each element of L is tail-equivalent to one of finitely many sequences of the form  $v_n^{\omega}$ . Recall that an *r*-representation of a real number is eventually periodic if and only if that number is rational. Since  $v_n^{\omega}$  is by construction a periodic *r*-representation, anything element of  $([r]^m)^{\omega}$  that is tail-equivalent to it is eventually periodic as well, and thus in  $\mathbb{Q}^m$ .

It is obvious that  $(2) \Longrightarrow (3)$ . For  $(3) \Longrightarrow (4)$ , since X is closed, it contains all of its accumulation points. So as a subset of a countable set the accumulation points of X are countable as well. Finally, for  $(4) \Longrightarrow (1)$ , if the set of accumulation points of X is countable, then there is no subset of L of the form  $\nu_r(u\{a,b\}^*wv^{\omega})$  with  $a, b, u, v, w \in ([r]^m)^*$  and |a| = |b| but  $a \neq b$ . If there were, the set  $\nu_r(u\{a,b\}^{\omega})$  would all be accumulation points of X. So we deduce by Proposition 2.3 that X is r-sparse.

**Remark 2.5.** As a corollary of Lemma 2.4, it is immediate that for any *r*-regular  $\omega$ -language  $L \subseteq ([r]^m)^{\omega}$ , either the size of the set of length-*n* prefixes of words in *L* eventually is bounded by a polynomial in *n*, or eventually grows exponentially.

**Fact 2.6** ([CHM02]). If there is a V-W set X contained in [0, 1] such that W contains two distinct words of the same length, then  $d \cdot X := X + X + \cdots + X$  contains a closed interval [a, b] with b > a for some d.

The above fact means that there is another characterization of sparseness: a closed r-regular subset X of [0,1] is sparse if and only if there is some  $d = d(X) \ge 1$  such that

 $d \cdot X := \{x_1 + \dots + x_d \colon x_1, \dots, x_d \in X\}$ 

contains an interval [a, b] with b > a.

Finally, we establish the connection between Hausdorff dimension and closed, r-sparse subsets of  $[0,1]^m$ .

**Lemma 2.7.** If X is a closed regular subset of  $[0, 1]^m$ , then X is r-sparse if and only if  $d_H(\pi_i(X)) = 0$  for all  $i \leq m$ .

*Proof.* Suppose that  $X \subseteq [0,1]^m$  is recognized by the automaton  $\mathcal{A}$ . First suppose that the set of words that  $\mathcal{A}$  accepts includes some of the form  $u\{a,b\}^*wv^{\omega}$  with  $a \neq b$  and |a| = |b|. Then the automaton must also accept all words of the form  $u\{a,b\}^{\omega}$ , due to the fact that in a closed automaton all states with a path to itself are accept states. We can write this set as follows:

$$(u_1,\ldots,u_m)\{(a_1,\ldots,a_m),(b_1,\ldots,b_m)\}^{\omega}$$

where for each  $1 \leq i \leq m$  we have  $u_i, a_i, b_i \in [r]^*$  and for any  $1 \leq j \leq m$ , we have  $|u_i| = |u_j|, |a_i| = |a_j|$ , and  $|b_i| = |b_j|$ . We observe that by the pigeonhole principle, there exists some  $i \in \{1, \ldots, m\}$  such that the words in coordinate i are of the form  $u_i\{a_i, b_i\}^{\omega}$  with  $a_i \neq b_i$ . Otherwise, it would have to be the case that for each  $i \in \{1, \ldots, m\}$  we have  $a_i = b_i$ , which contradicts the fact that we supposed a and b are distinct elements of  $([r]^m)^*$ .

Let  $X_i \subseteq \mathbb{R}$  denote the set recognized by the "projection" of the automaton onto the *i*-th coordinate, which we will call  $\mathcal{A}_i$ . We observe that  $u_i\{a_i, b_i\}^{\omega}$ , the language that  $\mathcal{A}_i$  accepts, corresponds to a shift of a Cantor-like set. Any such Cantor-like set can be expressed as a Moran construction [Mor66], i.e. the limit set of a geometric construction that starts with finitely many basic sets, and such that each basic set is the closure of its interior, at each level the interiors are disjoint, each basic set is geometrically similar to a basic set one level above, and moreover the

diameters of the basic sets each successive level is  $\frac{1}{k}$  that of a basic set from the previous level. For the set  $\nu_r(u\{a,b\}^{\omega})$ , we take  $k = r^{\ell}$  where  $\ell = |a| = |b|$ . Then results of Moran [Mor66] tell us the Hausdorff dimension of the limiting set is the unique root of  $n(1/k)^x = 1$ , where n < kis the number of initial basic sets. Since n is at least 2 (there must be distinct initial basic sets corresponding to the words a and b) this root is always nonzero.

Now suppose X is sparse. Then  $X \subseteq \mathbb{Q}^m$  by Lemma 2.4. Since Hausdorff dimension is an increasing function with respect to the subset relation, and Hausdorff dimension of projections of  $\mathbb{Q}^m$  are all zero, we are done.

# 3. *r*-sparse sets and d-minimality

In this section, we look more directly at the structure and properties of sparse automata themselves, rather than just the sets that they recognize.

**Lemma 3.1.** For every sparse Büchi automaton, there is a weak Büchi automaton that recognizes the same language.

*Proof.* Suppose  $\mathcal{A}$  is sparse. Then  $\mathcal{A}$  recognizes a finite union of sets, each of which is recognized by an automaton  $\mathcal{A}_i$  of the following form:



where the labels of transition arrows  $u_i$  and  $v_i$  are a shorthand for a finite string on the alphabet  $\Sigma$ , and the states  $q_i$  are shorthand for the finite sequence of states that allows through only the strings with the corresponding  $u_i$  as a substring, and which loop on the corresponding  $v_i$  substrings.

We construct the automaton that recognizes this union via the automaton union construction. This involves creating a new automaton in which the states are labeled by elements of  $Q_1 \times \cdots \times Q_m$ where  $Q_i$  is the set of states of  $\mathcal{A}_i$  for each  $i \leq m$ , and there is a transition from  $(q_1, \ldots, q_m)$  to  $(q'_1, \ldots, q'_m)$  on character  $\sigma$  precisely if there is  $q_i$  and  $q'_i$  such that  $\delta_i(q_i, \sigma) = q'_i$  where  $\delta_i$  is the transition function of  $\mathcal{A}_i$ . Similarly,  $(q_1, \ldots, q_m)$  is an accept state if at least one  $q_i$  is in  $\mathcal{A}_i$ .

Consequently, in the resulting automaton of the union construction, the accepting states are only those which include as a coordinate one of the states in  $\mathcal{A}_i$  that were in the cycle of states corresponding to  $q_n$  in the diagram. Since each  $q_n$  in the above diagram denotes a strongly connected component of accept states, any state in the union construction with a coordinate state from  $q_n$  inherits that strongly connected component. If a state in the union is not an accept state, then by the nature of the union construction it cannot be accessed from any of the accept states. It follows that the resulting automaton is weak.

**Lemma 3.2.** If  $X \subseteq [0,1]^m$  is r-sparse, then  $\overline{X}$  is r-sparse as well.

*Proof.* By the above remark, it is evident that each sparse Büchi automaton is equivalent to a weak and trim Büchi automaton, since in the proof of the above remark it is clear all states are accessible from the start state, and co-accessible from a final state. We can now apply Remark 59 in [CLR15], which says if X is recognized by a trim automaton  $\mathcal{A}$ , then  $\overline{X}$  the topological closure is recognized by  $\overline{\mathcal{A}}$ . The closure of the automaton  $\mathcal{A}$  is constructed by making all states final states. Hence, the resulting automaton  $\overline{\mathcal{A}}$  for a sparse automaton  $\mathcal{A}$  is equivalent to the union construction applied to the closed automata of the following form:



as explained in the remark above. Fix one such sub-automaton of this form. The closure of this sub-automaton is equivalent to the union construction applied to automata of this form:



where *i* ranges over  $\{0, \ldots, n\}$ , effectively unioning over truncations of the sub-automaton in question at each state preceding  $q_n$ . Hence, the closure  $\overline{A}$  is yet again an *r*-sparse automaton, and recognizes  $\overline{X}$ .

The first step toward proving Theorem 1 is establishing the  $\emptyset$ -definability of each *r*-sparse subset of  $[0,1]^m$  in  $\mathcal{R}_{r,\ell} = (\mathbb{R}, <, +, 0, 1, r^{-\ell \mathbb{N}})$  for some  $\ell \in \mathbb{N}$ . In fact that, we can take this  $\ell$  to be the natural number  $\ell_L$  described below form (1). Below, if  $v = (v_1, \ldots, v_m) \in ([r]^m)^*$  with |v| = k, then let  $\tilde{\nu}_r(v)$  denote

$$(v_{1,0}r^{k} + v_{1,1}r^{k-1} + \ldots + v_{1,k-1}r + v_{1,k}, \ldots, v_{m,0}r^{k} + v_{m,1}r^{k-1} + \ldots + v_{m,k-1}r + v_{m,k}).$$

Throughout, we say a set  $X \subseteq \mathbb{R}$  is *discrete* if for every  $x \in X$  there is an open interval  $I \subseteq \mathbb{R}$  such that  $\{x\} = X \cap I$ . Note that in any linearly ordered structure (with the ordering symbol < in the language) if X is a definable and discrete set, the graph of the predecessor function on X is defined as follows:

$$P_X(x) = y \iff x \in X \land y \in X \land y < x \land \forall z (y < z < x \to z \notin X).$$

**Remark 3.3.** If k is a divisor of  $\ell$ , then  $r^{-k\mathbb{N}}$  is definable in  $r^{-\ell\mathbb{N}}$ . In particular,  $r^{-\mathbb{N}}$  is definable in  $r^{-\ell\mathbb{N}}$ .

**Proposition 3.4.** Every *r*-sparse set  $A \subseteq [0,1]^m$  is  $\emptyset$ -definable in  $\mathcal{R}_{r,\ell}$  for some  $\ell \in \mathbb{N}$ .

Proof. Below, let  $\vec{0}$  denote  $(0, \ldots, 0) \in [0, 1]^m$ . Let  $L \subseteq ([r]^m)^{\omega}$  be such that  $\nu_r(L) = A$ . Let  $\ell_L$  be the least common multiple of the lengths of words in the set  $W_L$  as described below form (1). Since A is r-sparse, we can express L as a finite union of  $\omega$ -languages of the form

$$u_0v_0^*u_1v_1^*\ldots u_nv_n^{\omega}$$

where  $n \in \mathbb{N}$ , and each  $u_i$  and  $v_i$  is in  $([r]^m)^*$ . Fix one such sub-language of L, and note that by showing that the image of this sub-language under  $\nu_r$  is definable, we can infer that  $\nu_r(L)$  is definable, since L is a finite union of such sub-languages.

Recall that for each  $k \in \mathbb{N}$  we have the following identity:

$$\frac{1}{r^k - 1} = \frac{1}{r^k} + \frac{1}{r^{2k}} + \frac{1}{r^{3k}} + \dots$$

Note that since the set  $\nu_r(\{10^{\omega}, 010^{\omega}, 0010^{\omega}, \ldots\})$  is definable in  $\mathcal{R}_{r,\ell_L}$ , then so too is the following, where  $k \in \mathbb{N}$  is fixed, using that  $\frac{1}{r^k-1} = \nu_r((0^k 1)^{\omega})$ :

$$\nu_r(\{0^*(0^k1)^\omega\}) = \frac{1}{r^k - 1} \cdot r^{-\mathbb{N}}.$$

Let  $|v_n| = m_n$ , and observe  $\nu_r(v_n^{\omega}) = \tilde{\nu}_r(v_n) \cdot \frac{1}{r^{m_n-1}}$ .

For each  $u_i$ , if  $|u_i| = \ell_i$  observe that  $\nu_r(\{\vec{0}^* u_i \vec{0}^\omega\})$  is equal to the following:

(2) 
$$\left\{\frac{\tilde{\nu}_r(u_i)}{r^{\ell_i}}x: x \in r^{-\mathbb{N}}\right\}$$

Similarly, for each  $v_i$  if  $|v_i| = m_i$ , then we observe that the set  $\nu_r(\{\vec{0}^* v_i^* \vec{0}^\omega\})$  is equal to the following:

(3) 
$$\bigcup_{k \in [m_i]} \left\{ \frac{\tilde{\nu}_r(v_i)}{r^k(r^{m_i}-1)} (z-x) : x, z \in r^{-m_i \mathbb{N}} \land x < z \right\}$$

Note that in the above expression for  $\nu_r(\{\vec{0}^* v_i^* \vec{0}^\omega\})$ , the number  $1/(r^{m_i} - 1)$  is a fixed constant that depends only on the length of  $v_i$ . Finally, we observe that  $\nu_r(\{\vec{0}^* v_n^\omega\})$  is defined by

(4) 
$$\left\{\frac{\tilde{\nu}_r(v_n)}{r^{m_n}-1} \cdot z : z \in r^{-\mathbb{N}}\right\}.$$

For brevity, we let  $\phi_i(x, \vec{y})$  denote the  $\mathcal{L}_r$ -formula such that  $\exists x \phi_i(x, \vec{y})$  defines the set  $\nu_r(\{0^* u_i^* 0^\omega\})$ as seen in expression (2). Let  $\psi_i(x, \vec{y}, z)$  denote the  $\mathcal{L}_r$ -formula such that  $\exists x \exists z \psi_i(x, \vec{y}, z)$  defines the set  $\nu_r(\{0^* v_i^* 0^\omega\})$  as seen in expression (3). Let  $\theta(z, \vec{y})$  denote the  $\mathcal{L}_r$ -formula such that  $\exists z \theta(z, \vec{y})$ defines the set  $\nu_r(\{0^* v_n^\omega\})$  as seen in expression (4). To define the set of elements of precisely the form  $\nu_r(u_0 v_0^* u_1 v_1^* \dots u_n v_n^\omega)$ , we consider the following formula:

$$\begin{split} \varphi(v) &:= \exists x_0, x_1, \dots, x_{2n-1}, x_{2n} \exists z_0, \dots, z_n \exists \vec{y_0}, \vec{y_1}, \dots, \vec{y_{2n-1}}, \vec{y_{2n+1}} \left( v = \sum_{i=0}^{2n} \vec{y_i} \land \bigwedge_{i=0}^n \phi_i(x_{2i}, \vec{y_{2i}}) \land \\ \bigwedge_{i=0}^{n-1} \psi_i(x_{2i+1}, \vec{y_{2i+1}}, z_i) \land \theta(z_n, \vec{y_{2n+1}}) \land x_0 > x_2 > \dots > x_{2n} \land z_0 > z_1 > \dots > z_n \land \\ \sum_{i=0}^n \frac{r^{\ell_i} - 1}{r^{\ell_i}(r-1)} x_{2i} + \sum_{i=0}^{n-1} \frac{1}{r-1} (z_i - x_{2i+1}) + \frac{1}{r-1} z_n = \frac{1}{r-1} \Big). \end{split}$$

Observe that the final conjunct, which essentially dictates that the (slightly weighted, to compensate for the lengths of the  $u_i$ 's and  $v_i$ 's) sum over the witnesses that mark the spacing for the  $u_i$ 's and  $v_i$ 's has the repeating *r*-representation  $1^{\omega}$ , since  $1/r + 1/r^2 + 1/r^3 + \ldots = 1/(r-1)$ . In other words, it ensures that the components  $y_i$  that contribute the  $u_i$ 's and  $v_i$ 's do not overlap, nor leave gaps. Additionally, the required ordering of the  $x_{2i}$ 's and  $z_i$ 's ensures that the subwords appear in the correct order. This is sufficient to conclude that any element that satisfies  $\varphi(v)$  must be in the set  $\nu_r(\{u_0v_0^*u_1v_1^*\ldots u_nv_n^{\omega}\})$ . Conversely, we have designed  $\varphi(v)$  so any such element will satisfy it. Note that although  $\varphi(v)$  is defined with parameters, all of these parameters are elements of  $\mathbb{Q}$  which are  $\emptyset$ -definable in the structure  $\mathcal{R}_{r,\ell_L}$ , so there is an equivalent formula without parameters.  $\Box$ 

We note that the above result is optimal in the sense that if A is an r-sparse set definable in  $\mathcal{R}_{r,\ell}$  with  $\ell > 1$ , in general A will not also be definable in  $\mathcal{R}_{r,m}$  for some  $1 \leq m < \ell$ . To see this,

let  $T_r := Th(\mathbb{R}, <, 0, r^{-\mathbb{N}})$ . Consider  $(R, <, 0, +, P) \models T_r$  where  $\mathbb{R} \subseteq R$ , and define  $\lambda : R \to R$  as follows:

$$t \mapsto \begin{cases} 1, & t \ge 1\\ 0, & t \le 0\\ \min([t, 1) \cap P), & t \in (0, 1). \end{cases}$$

Effectively, the function  $\lambda$  maps any element t of (0, 1) to the smallest element of P greater than P, and all other positive t get mapped to 1, while all non-positive t are sent to 0. Note that all models of  $T_r$  are interdefinable with a model of  $T_{\lambda} := Th(\mathbb{R}, <, 0, +, \lambda)$  that has the same underlying set, and vice versa. Observe that  $T_{\lambda}$  is axiomized by the axioms of ordered divisible abelian groups plus the following axioms, which are adapted from [MT06]:

(i)  $s \leq t \rightarrow \lambda(s) \leq \lambda(t)$ , (ii)  $t \geq 1 \rightarrow \lambda(t) = 1$ , (iii)  $t \leq 0 \rightarrow \lambda(t) = 0$ , (iv)  $t \in (0,1) \rightarrow \frac{1}{r}\lambda(t) < t \leq \lambda(t)$ , (v)  $\frac{1}{r}\lambda(t) < s \leq \lambda(t) \rightarrow \lambda(s) = \lambda(t)$ , (vi)  $t \in [0,1] \rightarrow (\lambda(t) = t \leftrightarrow \lambda(\frac{t}{r}) = \frac{t}{r})$ .

Note that these axioms are a bit redundant, but it will be helpful to have them written this way. Below, let  $x \ll 1$  denote the statement  $\forall q \in \mathbb{Q}(0 < x < q)$ , i.e. the property that x is infinitesimal with respect to the real numbers. For a subset X of an expansion of  $\mathbb{R}$ , let  $X^{\ll 1}$  denote the set of infinitesimal elements in X (with respect to  $\mathbb{R}$ ).

**Lemma 3.5.** Suppose that  $S := (S, <, +, 0, \lambda)$  is a  $|\mathbb{R}|$ -saturated elementary extension of the structure  $\mathcal{R}_r = (\mathbb{R}, <, +, 0, \lambda) \models T_{\lambda}$ . Let  $\mathcal{L} = \{<, +, 0\}$ , and let  $\mathcal{L}_{\lambda} = \{<, +, 0, \lambda\}$  where the interpretation of  $\lambda(\mathbb{R})$  in  $\mathcal{R}_r$  is  $r^{-\mathbb{N}} \cup \{0\}$ . Suppose b and b' are in  $\lambda(S)$  and satisfy the same  $\mathcal{L}$ -type over  $\mathbb{R}$ . Then b and b' satisfy the same  $\mathcal{L}_{\lambda}$ -type over  $\mathbb{R}$ .

Proof. To prove the claim, it suffices to show that if  $e: S \to S$  is an  $\mathcal{L}$ -elementary map that fixes  $\mathbb{R}$  pointwise and maps b to b', then e is also an  $\mathcal{L}_{\lambda}$ -elementary map. Note that any  $\mathcal{L}$ -embedding  $e: S \to S$  is also an  $\mathcal{L}$ -elementary map because the reduct of  $\mathcal{S}_r$  to  $\mathcal{L}$  has quantifier elimination. By Blum's criterion (see, for example, [M02]), to show quantifier elimination for  $T_{\lambda}$  it suffices to show the following: if  $\mathcal{A} = (A, <, +, 0, \lambda), \mathcal{B} = (B, <, +, 0, \lambda) \models T_{\lambda}$  and  $\mathcal{S}$  is an  $|\mathcal{A}|^+$ -saturated elementary extension of  $\mathcal{A}$ , then for any  $b \in B$  there is  $b' \in S$  such that there is an  $\mathcal{L}_{\lambda}$  embedding  $e: B \to S$  of  $dcl(\mathcal{A} \cup \{b\})$  into  $\mathcal{S}$  that maps A to A pointwise, maps b to b', and the image of e in  $\mathcal{S}$  is  $dcl(\mathcal{A} \cup \{b'\})$ . Hence, both to prove quantifier elimination for  $T_{\lambda}$  and to prove the claim, it suffices to show that for any  $b \in B \setminus A$ , if  $b' \in S$  satisfies the same cut over A as b and  $\lambda^{\mathcal{B}}(b) = b \iff \lambda^{\mathcal{S}}(b') = b'$ , then any  $\mathcal{L}$ -embedding  $e: dcl(\mathcal{A} \cup \{b\}) \to S$  that fixes A pointwise and maps b to b' is also an  $\mathcal{L}_{\lambda}$ -embedding.

Let  $\mathcal{B} = (B, <, +, 0, \lambda) \models T_{\lambda}$ , let  $\mathcal{A}$  be a substructure of  $\mathcal{B}$ , and let  $(\mathcal{S}, \lambda^*)$  be an  $|\mathcal{A}|^+$ -saturated elementary extension of  $\mathcal{A}$ . Suppose that  $b \in \lambda(B) \setminus \lambda(A)$ , and note that by saturation of  $\mathcal{S}$ , there is an element  $b' \in \lambda^*(S)$  such that b and b' satisfy the same cut over  $\mathcal{A}$ . Let  $e : \operatorname{dcl}(\mathcal{A} \cup \{b\}) \to S$ map  $\mathcal{A}$  to itself pointwise, and let e(b) = b'. We note that e is indeed an  $\mathcal{L}$ -embedding over  $\mathcal{A}$ , since the fact that b and b' satisfy the same cut over  $\mathcal{A}$  implies that they satisfy the same  $\mathcal{L}(\mathcal{A})$ -formulas. To demonstrate that e is also an  $\mathcal{L}_{\lambda}$ -embedding over  $\mathcal{A}$ , it suffices to show that  $\lambda(\operatorname{dcl}_{\mathcal{L}}(\mathcal{A} \cup \{b\})) \subseteq \operatorname{dcl}_{\mathcal{L}}(\mathcal{A} \cup \{b\})$  and  $e(\lambda(x)) = \lambda^*(e(x))$  for all  $x \in \operatorname{dcl}_{\mathcal{L}}(\mathcal{A} \cup \{b\})$ , because from this we conclude e respects all atomic  $\mathcal{L}_{\lambda}(\mathcal{A})$ -formulas, and thus all quantifier-free ones by induction.

Given  $s \in S$ , define  $r^{\mathbb{Z}}s := \{r^z s : z \in \mathbb{Z}\}$ , and for any  $X \subseteq S$  let H(X) denote the convex hull of X in S. Recall the following fact from [MT06]: If  $s \ll 1$  and  $A \cap H(r^{\mathbb{Z}}s) = \emptyset$ , then

(5) 
$$\{x \in \operatorname{dcl}_{\mathcal{L}}(A \cup \{s\}) : x \ll 1\} \subseteq \bigcup_{\{a \in A^{\ll 1}\}} H(r^{\mathbb{Z}}\lambda(a)) \cup H(r^{\mathbb{Z}}s).$$

Suppose that  $c \in C := \operatorname{dcl}_{\mathcal{L}}(A \cup \{b\})$ . If  $c \in \operatorname{dcl}_{\mathcal{L}}(A)$ , then it is immediate that  $\lambda(c) \in A$  since  $\mathcal{A}$  is a substructure of  $\mathcal{B}$ . So suppose that  $c \in C \setminus \operatorname{dcl}_{\mathcal{L}}(A)$ , and suppose also that  $0 < c \ll 1$ , since otherwise it is clear by definition of  $\lambda$  that  $\lambda(c) \in \operatorname{dcl}_{\mathcal{L}}(A \cup \{b\})$ . By axiom (iv), we observe that  $c < \lambda(c) < rc$ , hence  $\lambda(c) \in H(r^{\mathbb{Z}}c)$ . Since we assume  $c \in \{x \in \operatorname{dcl}(A \cup \{s\}) : x \ll 1\}$ , we know that either  $c \in H(r^{\mathbb{Z}}\lambda(a))$  for some  $a \in A$ , in which case there is  $z \in \mathbb{Z}$  for which  $r^z\lambda(a) \ge c > r^{z+1}\lambda(a)$  so by axiom (v) we conclude  $\lambda(c) = \lambda(r^z a)$ . Otherwise, we know  $c \in H(r^{\mathbb{Z}}b)$ , so by a similar argument, and the fact that  $b = \lambda(b)$ , we conclude  $\lambda(c) = r^z b$  for some  $z \in \mathbb{Z}$ . In either case, we conclude that  $\lambda(c) \in C$ , as desired.

Finally, suppose that  $x \in C$ , and for contradiction we suppose that  $e(\lambda(x)) \neq \lambda^*(e(x))$ . Without loss of generality, we may assume that  $e(\lambda(x)) < \lambda^*(e(x))$ . Since e(a) = a for all  $a \in \mathcal{A}$ , we know  $\lambda(a) = \lambda^*(a)$  for all  $a \in \mathcal{A}$ . Note also that e preserves cuts over  $A \cup \{b\}$ , so the cut of e(x) over  $A \cup \{b^*\}$  is the image under e of the cut of x over  $A \cup \{b\}$ . If x > q for some  $q \in \mathbb{Q}^S$ , then it must be the case that  $\lambda^*(e(x)) = e(\lambda(x))$ , since  $e(q^{\mathcal{A}}) = q^S$  for all  $q \in \mathbb{Q}$ , and for each  $q \in \mathbb{Q}$  we know that  $\lambda(q)$  is the least power of  $\frac{1}{r}$  greater than q. So we cuclude that  $x \ll 1$ . By (5), we obtain the following:

$$x \in \bigcup_{1 \gg a \in A} H(r^{\mathbb{Z}}\lambda(a)) \cup H(r^{\mathbb{Z}}b).$$

Since e is  $\mathcal{L}$ -elementary, we conclude the following:

 $e(x) \in \bigcup_{1 \gg a \in A} H(r^{\mathbb{Z}}\lambda(a)) \cup H(r^{\mathbb{Z}}b^*).$ 

If  $x \in H(r^{\mathbb{Z}}\lambda(a))$ , then by definition, there exist  $z, z' \in \mathbb{Z}$  such that  $r^{z'}\lambda(a) \leq x \leq r^z\lambda(a)$  for some  $a \in A$ . Hence there is some  $k \in \mathbb{N}$  with  $z < k \leq z'$  such that  $r^{k-1}\lambda(a) \leq x \leq r^k\lambda(a)$ . Since  $r^{k-1}\lambda(a), r^k\lambda(a) \in \lambda(\mathcal{A})$ , we conclude that  $\frac{1}{r}\lambda(r^k\lambda(a)) \leq x \leq \lambda(r^k\lambda(a))$ . So by axiom (v), we know  $\lambda^*(x) = \lambda(r^k\lambda(a))$ . As e is a  $\mathcal{L}$ -elementary map preserving A, we conclude that  $\frac{1}{r}\lambda(r^k\lambda(a)) \leq e(x) \leq \lambda(r^k\lambda(a))$ . Hence we know  $\lambda^*(e(x)) = \lambda(r^k\lambda(a)) = \lambda^*(x)$ .

Otherwise, we must conclude  $x \in H(r^{\mathbb{Z}}b)$  and  $e(x) \in H(r^{\mathbb{Z}}b^*)$ . In this case we similarly conclude that there is some  $k \in \mathbb{Z}$  such that  $r^{k-1}b < x \leq r^k b$ , and that there is some  $k' \in \mathbb{Z}$  such that  $r^{k'-1}b^* < e(x) \leq r^k b^*$ . Since  $\lambda(b) = b$  and  $\lambda(b^*) = b^*$ , we see that  $r^{k-1}\lambda(b) < x \leq r^k\lambda(b)$  and  $r^{k-1}\lambda(b^*) < x \leq r^k\lambda(b^*)$ . Hence  $\frac{1}{r}\lambda(r^k b) < x \leq \lambda(r^k b)$  and  $\frac{1}{r}\lambda(r^{k'}b^*) < e(x) \leq r^k\lambda(r^{k'}b^*)$ , by repeated application of axiom (vi). So by axioms (v) and (vi), we see  $\lambda(x) = r^k b$  and  $\lambda^*(e(x)) =$  $r^{k'}b^*$ . If  $e(\lambda(x)) < \lambda^*(e(x))$ , then  $e(r^{k-1}b) = r^{k-1}b^* < e(x) \leq e(\lambda(x)) = e(r^k b) = r^k b^* < \lambda^*(e(x))$ . As  $r^{k'-1}b^* < e(x) \leq \lambda^*(e(x)) = r^{k'}b^*$ , we conclude that  $r^k b^* < \lambda^*(e(x)) = r^{k'}b^*$ . Yet this yields both  $e(x) \leq r^k b^* < r^{k'}b^*$  and also  $r^k b^* \leq r^{k'-1}b^* < e(x)$ , a contradiction. So it must be the case that  $e(\lambda(x)) = \lambda^*(e(x))$ , and we conclude that e is in fact an  $\mathcal{L}_{\lambda}$ -embedding over A.

**Fact 3.6.** For m > 1 the set  $r^{-m\mathbb{N}}$  is not definable in the structure  $(\mathbb{R}, <, +, r^{-\mathbb{N}})$ .

Proof. Assume for sake of contradiction that the set  $r^{-m\mathbb{N}}$  is defined by the  $\mathcal{L}_{\lambda}$ -formula  $\phi(x)$ , with parameters in  $\mathbb{R}$ . Let  $(\mathcal{R}^*, \lambda^*)$  be an  $\omega$ -saturated elementary extension of  $(\mathbb{R}, <, +, r^{-\mathbb{N}})$ , and let  $b \in \lambda^*(\mathcal{R}^*)$  be such that  $b \ll 1$  and  $(\mathcal{R}^*, \lambda^*) \models \phi(b)$ . Let  $b' \in \lambda^*(\mathcal{R}^*)$  be the largest element of  $\lambda^*(\mathcal{R}^*)$  less than b. Then  $\neg \phi(b')$  holds, since m > 1. Yet we know by Lemma 3.5 that b and b' have the same  $\mathcal{L}_{\lambda}(\mathbb{R})$ -type, a contradiction. So no such  $\phi$  exists, and hence none exists in the language  $(<, +, r^{-\mathbb{N}})$  either. Say a structure is *d*-minimal if for every (m + 1)-ary definable set A, there is  $N \in \mathbb{N}$  such that for each *m*-tuple x the fiber  $\{t : (x, t) \in A\}$  either has interior or is the union of at most N discrete sets, as defined in [Mi05]. The *Cantor-Bendixson derivative* of a set  $X \subseteq \mathbb{R}^d$  is  $X \setminus X^{iso}$ , where  $X^{iso}$  are the isolated points of X. The Cantor-Bendixson derivative of X is often denoted by X', and for  $\alpha$  an ordinal, let  $X^{(\alpha)}$  denote the result of taking the Cantor-Bendixson derivative  $\alpha$  times. The *Cantor-Bendixson rank* of X is the least ordinal  $\alpha$  such that  $X^{(\alpha)} = X^{(\alpha+1)}$ . If the Cantor-Bendixson rank of X is  $\alpha$  and  $X^{(\alpha)} = \emptyset$ , then we say that X has vanishing Cantor-Bendixson derivative.

**Lemma 3.7.** Suppose that  $A \subseteq [0,1]^d$  is r-regular and closed. Then A has vanishing Cantor-Bendixson derivative and finite Cantor-Bendixson rank if and only if A is r-sparse.

*Proof.* First, we observe that Proposition 3.4 tells us that A is definable in a d-minimal structure, namely  $\mathcal{R}_r$ . Since A is a finite union of discrete sets definable in a d-minimal structure, it has finite Cantor-Bendixson rank; see [Mi05] or [MT18]. Moreover, a closed subset of  $\mathbb{R}^d$  has vanishing Cantor-Bendixson derivative if and only if it is countable [Kec95, §6.B]. The backward direction follows from putting together that A must have vanishing Cantor-Bendixson derivative, and the Cantor-Bendixson rank must be finite.

For the forward direction, we suppose that A is closed but not r-sparse. Then Lemma 2.4 tells us that A is uncountable, and hence does not have vanishing Cantor-Bendixson derivative by the above "if and only if" statement.

**Lemma 3.8.** Suppose that  $A \subseteq [0,1]^d$  is r-sparse and infinite. Let  $A^{acc}$  denote the accumulation points of A, i.e. non-isolated points in  $\overline{A}$ . Then  $A^{acc}$  has an isolated point.

Proof. By Lemma 3.7, we know that  $\bar{A}$  has vanishing Cantor-Bendixson derivative, and finite Cantor-Bendixson rank. This means there is  $n \in \mathbb{N}$  such that the *n*-th Cantor-Bendixson derivative is  $\emptyset$ . Since  $\bar{A}$  is an infinite compact subset of  $\mathbb{R}^d$ , it must have at least one accumulation point (by the Bolzano-Weierstrass theorem). Since taking the Cantor-Bendixson derivative removes the isolated points, the first Cantor-Bendixson derivative of  $\bar{A}$  is  $A^{acc} \neq \emptyset$ . Taking the second Cantor-Bendixson derivative, it cannot be the case that we get  $A^{acc}$  again, because this would contradict the fact that  $\bar{A}$  has vanishing Cantor-Bendixson derivative. So we conclude there is at least one isolated point in  $A^{acc}$ .

We now establish some notation concerning subwords. For any  $u, v \in ([r]^m)^*$ , write  $u \subseteq v$  if u is a prefix of v. If it does not hold that  $u \subseteq v$ , then we write  $u \not\subseteq v$ .

We are now able to prove the last result needed for Theorem 1. For the rest of this chapter, set  $\mathcal{R}_A := (\mathbb{R}, <, +, 0, 1, A)$  with A a unary predicate.

**Proposition 3.9.** The set  $r^{-\mathbb{N}}$  is  $\emptyset$ -definable in  $\mathcal{R}_A$ , whenever  $A \subseteq [0,1]^d$  is infinite and r-sparse.

*Proof.* We will write  $\mathcal{L} := \{<, +, 0, 1\}$  and  $\mathcal{L}_A := \mathcal{L} \cup \{A\}$  for our languages. Since A is r-sparse, we can express A as a finite union of sets of the following form:

(6) 
$$\nu_r(u_0v_0^*u_1v_1^*\dots u_nv_n^{\omega})$$

where  $n \in \mathbb{N}$ , and  $u_i, v_i \in [r]^*$  with  $v_i \neq \varepsilon$  for each  $i \leq n$ . For each subset of A of the form (6), we can take n to be minimal. Note that it must be the case that n > 0 for at least one such subset of A, since otherwise the set A would consist of a finite set of points, each of the form  $\nu_r(u_0v_0^{\omega}) \in \mathbb{Q}$ . Let  $L \subseteq [r]^{\omega}$  be such that  $\nu_r(L) = A$ . Let  $V_1W_1^{\omega}, \ldots, V_tW_t^{\omega}$  be a set of V-W components whose union is L, as given to us by Theorem 1.4. Moreover, we can take t to be minimal. Since we assume that A is infinite, there exists some index  $i \in [d]$  such that the projection of A onto the *i*-th coordinate is again infinite. Moreover, the projection of an *r*-sparse set is again *r*-sparse. Let  $A_i$  denote  $\pi_i(A)$ , the projection of A onto the *i*-th coordinate, and observe that  $A_i$  is  $\emptyset$ -definable in  $\mathcal{R}_A$ . Let  $L_i \subseteq [r]^{\omega}$  be such that  $\nu_r(L_i) = A_i$ .

Let  $V_i W_i^{\omega}$  be a V-W component of  $L_i$  containing the words  $u_0 v_0^* u_1 v_1^* \dots u_n v_n^{\omega}$  such that there is an infinite subset of  $A_i$  that has an isolated accumulation point  $p \in A'_i$ . This p exists by Lemma 3.8, and p is of the form  $\nu_r(wv_{n-1}^{\omega})$  where  $w \in [r]^*$  is such that  $wv_{n-1}^* u_n \subseteq V_i$ . There is an open set  $U \subseteq [0,1]$  that witnesses p being isolated as an element of  $A'_i$ . Let  $e \in \mathbb{N}$  be sufficiently large that the image under  $\nu_r$  of every element in  $L_i$  with prefix  $wv_{n-1}^e$  is in U. Consider the subset of elements of  $A_i$  of the form  $\nu_r(wv_{n-1}^e v_{n-1}^* u_n v_n^{\omega})$ , which is defined by the following formula:

$$\nu_r(wv_{n-1}^e 0^\omega) \le x \le \nu_r(wv_{n-1}^e (r-1)^\omega) \land x \in A_i$$

Let this subset with initial r-representation  $wv_{n-1}^e$  be called  $\tilde{A}_i$ . Assume without loss of generality that p is the infimum of this subset. Otherwise, if it is the supremum we can perform the same proof but with the inequalities reversed, and if it is neither we restrict our definable set so that it becomes either the infimum or supremum. Let d be the least common multiple of the minimal (i.e., shortest choice of) generators  $v_n$  for the tails  $v_n^{\omega}$  for the words whose image under  $\nu_r$  are in  $\tilde{A}_i$ . Observe that via change of base from r to  $r^d$ , the tails of the words  $(r^d - 1) \cdot \nu_r(wv_{n-1}^e v_{n-1}^* u_n v_n^{\omega})$ are all eventually zero, for each  $v_n$ . Hence, the set  $(r^d - 1)\tilde{A}_i$  is of the following form:

$$\bigcup_{i \le m} \nu_r(w'v'^*u'_i 0^\omega)$$

for some minimal  $m \in \mathbb{N}$  and words  $w', v', u'_1, \ldots, u'_m \in [r]^*$ .

We consider the following set:

$$D_m = \left\{ x - P^m(x) : x \in \tilde{A}_i \right\}$$

and observe that each element of  $D_m$  is of the following form:

$$\{\nu_r(0^{k'}(0^k)^*w_j0^\omega): j \in [m]\}$$

where  $k' \in \mathbb{N}$  is the length of w' and  $k \in \mathbb{N}$  is the length of v', and  $\nu_r(w_j 0^{\omega}) = \nu_r(u'_j 0^{\omega}) - \nu_r(v'u'_j 0^{\omega})$ for some  $j \leq m$ .

Note that by our assumptions on the minimality of n for each V-W component of L, it must be the case that each  $w_j$  is not the empty string. At this point, we note that if m = 1, i.e. there is only one V-W component of L whose elements are r-representations for the points in  $\tilde{A}_i$ , then the following set contains all elements of  $r^{-k\mathbb{N}}$ :

$$D_{1,1} := \frac{r^{k'}}{\tilde{\nu}_r(w_1)} D_1 = \left\{ \frac{r^{k'}}{\tilde{\nu}_r(w_1)} \nu_r \left( 0^{k'+bk} w_1 0^{\omega} \right) : b \in \mathbb{N} \right\}$$

where  $w_j = w_1$  since m = 1. By inspection, any element of this set is equal to a point of the form  $\nu_r((0^k)^*10^\omega)$ , and hence  $D_{1,1}$  is equal to the set  $r^{-k\mathbb{N}}$ .

Finally, we see that we can define  $r^{-\mathbb{N}}$  via the following observation:

$$r^{-\mathbb{N}} = \left( r^{-k\mathbb{N}} \bigcup r \cdot r^{-k\mathbb{N}} \bigcup r^2 \cdot r^{-k\mathbb{N}} \bigcup \dots \bigcup r^{k-1} \cdot r^{-k\mathbb{N}} \right) \bigcap [0,1].$$

Returning to the case that m > 1, we instead consider a definable subset of the intersection the sets obtained by multiplying  $D_m$  by  $r^{k'}$  and dividing by  $r \cdot \tilde{\nu}_r(w_j)$  for each  $j \in [m]$ :

(7) 
$$\left\{ x : \exists y_1, \dots, y_m \in D_m \left( \bigwedge_{j \in [m]} x = \frac{r^{k'-1}}{\tilde{\nu}_r(w_j)} y_j \wedge \bigwedge_{i,j \in [m]} \left( y_i < y_j \to y_i \in (r^{-k}y_j, y_j) \right) \right) \right\}$$

which we aim to show only contains elements in  $r^{-k\mathbb{N}}$ . Observe that if x is in (7), then the following holds:

$$x \in \bigcap_{j \in [m]} \frac{r^{k'-1}}{\tilde{\nu}_r(w_j)} \cdot D_m$$

but also, the elements  $y_1, \dots, y_m$  that witness x being in this intersection have the additional property that none is bigger than the others by a factor of  $r^{-k}$  or more.

Suppose that there is an element x such that x is an element of the set (7), but not of the form  $\nu_r((0^{k'+bk})^*10^{\omega})$ . Then for each  $j \leq m$  there is some  $b_j \in \mathbb{N}$  and  $i_j \leq m$  with  $j \neq i_j$  such that the following holds:

$$x = \frac{r^{k'}}{\tilde{\nu}_r(w_{i_j})} \nu_r \Big( 0^{k'+b_j k} w_j 0^\omega \Big).$$

Without loss of generality, suppose  $w_{\ell}$  is such that  $\tilde{\nu}_r(w_{\ell})$  is the maximum among the  $w_j$ 's (this exists because they are distinct), and similarly suppose that  $\tilde{\nu}_r(w_1)$  is minimal among the  $w_j$ 's. Since x is in the set (7), we deduce that all the exponents  $b_j$  must be the same, otherwise the witness  $y_i$ 's from the defining formula for set (7) would be off by too much. So there must be some  $w_{i_{\ell}}$  such that the following holds:

$$1 > \frac{\tilde{\nu}_r(w_1)}{\tilde{\nu}_r(w_{i_1})} = \frac{\tilde{\nu}_r(w_\ell)}{\tilde{\nu}_r(w_{i_\ell})}.$$

Yet this is impossible, since for all  $k \neq \ell$  we can deduce the following:

$$\frac{\tilde{\nu}_r(w_\ell)}{\tilde{\nu}_r(w_k)} \ge 1.$$

So we conclude that x must be in  $r^{-k\mathbb{N}}$ .

**Corollary 3.10** (Theorem 1). If A is infinite and r-sparse and  $L \subseteq ([r]^m)^{\omega}$  is such that  $\nu_r(L) = A$ , then there is a reduct  $\tilde{\mathcal{R}}$  of  $\mathcal{R}_{r,\ell_L}$  that expands  $(\mathbb{R}, <, +, r^{-\mathbb{N}})$  and defines the same sets (without parameters) as  $\mathcal{R}_A$ . Hence  $\mathcal{R}_A$  is d-minimal if A is r-sparse.

*Proof.* By Proposition 3.9, if A is r-sparse then the set  $r^{-\mathbb{N}}$  is  $\emptyset$ -definable in the language

$$(<, +, 0, 1, A),$$

and similarly by Proposition 3.4 we know such an A is  $\emptyset$ -definable in the language  $(<, +, 0, r^{-\ell_L \mathbb{N}})$ . Hence  $\mathcal{R}_A$  defines the same sets as some reduct of  $\mathcal{R}_{r,\ell_L}$ , and that reduct includes all sets definable with  $(<, +, r^{-\mathbb{N}})$ . Moreover, the work of van den Dries [vdD85] establishes that the structure  $\mathcal{R}_{r,\ell_L}$  is d-minimal, so  $\mathcal{R}_A$  is d-minimal as well.

**Corollary 3.11.** The structure  $\mathcal{R}_A$  has NIP whenever A is r-sparse.

*Proof.* This is immediate from Theorem 1 and results from [GH11], in which the authors show that certain d-minimal structures (which include our  $\mathcal{R}_A$ ) have NIP.

#### 4. A FRACTAL GEOMETRY AND TAMENESS DICHOTOMY

In this section, we consider the structure  $\mathcal{R}_A := (\mathbb{R}, <, 0, +, A)$  for which we now mandate that A be an r-regular subset of  $[0, 1]^d$ , but not necessarily r-sparse. We will start by proving some key facts about the isolated points of A, denoted  $A^{iso}$ , and the subset A' of A consisting of points that are not isolated points, but for which there exists an open interval surrounding it that only contains a countable subset of A. Throughout this section, let  $L \subseteq ([r]^d)^\omega$  be such that  $\nu_r(L) = A$ , let  $L^{iso} \subseteq ([r]^d)^\omega$  be such that  $\nu_r(L^{iso}) = A^{iso}$ , and let  $L' \subseteq ([r]^d)^\omega$  be such that  $\nu_r(L') = A'$ . We use throughout that by Theorem 1.4 we can write L as the finite union of sets of the form  $V_1W_1^\omega, \ldots, V_nW_n^\omega$  where each  $V_i$  and  $W_i$  are in  $([r]^d)^*$ .

We now introduce some notation and the notion of "tail-equivalence" in the context of rrepresentations. Given an element  $x \in ([r]^m)^{\omega}$ , let  $x_{[a,b]}$  with  $a \in \mathbb{N}$  and  $b \in \mathbb{N} \cup \{\infty\}$  denote the substring of x starting at index  $a \in \omega$  and ending with index  $b \in \omega$ . When  $b = \infty$ , then  $x_{[a,\infty)}$  denotes the tail sequence of x starting at index a, and it is again an element of  $([r]^m)^{\omega}$ . For  $u, v \in ([r]^m)^{\omega}$ , say that  $u \simeq_k v$  precisely if  $u_{[0,k]} = v_{[0,k]}$ . Given a language  $L \subseteq ([r]^m)^{\omega}$ , let  $[u]_{\simeq_k,L}$ denote  $\{v \in L : u_{[0,k]} = v_{[0,k]}\}$ .

**Definition 4.1.** For  $x, y \in ([r]^m)^{\omega}$ , say that x and y are *tail equivalent*, written  $x \simeq y$ , if there exist k and  $\ell$  in  $\mathbb{N}$  such that  $x_{[k,\infty)} = y_{[\ell,\infty)}$ .

**Lemma 4.2.** Let  $L \subseteq [r]^{\omega}$  be regular. For every element  $x \in L$ , the element  $\nu_r(x)$  is not in the closure of any uncountable V-W component of L if and only if  $x \in L^{iso} \cup L'$ . Moreover, the elements of L' and  $L^{iso}$  are in one of finitely many tail-equivalence classes.

Proof. For the backwards direction, we consider the contrapositive; suppose  $\nu_r(x)$  is in the closure of some uncountable V-W component of L. Fix one such component  $V_i W_i^{\omega}$ . Since x is in the closure of  $V_i W_i^{\omega}$ , it has a r-representation that is accepted by the automaton-theoretic closure of  $V_i W_i^{\omega}$ . Let  $(y_k)_{k \in \mathbb{N}}$  witness this, i.e.  $(y_k)_{k \in \mathbb{N}} \subseteq V_i W_i^{\omega}$  and  $(y_k)_{k \in \mathbb{N}} \to x$ . By discarding some  $y_k$ 's and re-indexing if necessary, we may assume that for each  $k \in \mathbb{N}$  it holds that  $x_{[0,k]} = y_{k[0,k]}$ , for some k-representations of x and  $y_k$ . For each  $k \in \mathbb{N}$ , let  $vw_1 \dots w_{m_k} \in V_i W_i^*$  be a prefix of  $y_k$  that witnesses  $x_{[0,k]} = y_{k[0,k]}$ .

Since we know that  $W_i^{\omega}$  contains uncountably many elements, we know there is an uncountable set of elements in  $W_i^{\omega}$  that are tail-inequivalent; let  $(z_{\lambda})_{\lambda < \aleph_1}$  enumerate one such uncountable set. Then the set of sequences given by  $(vw_1 \dots w_{m_k} z_{\lambda})_{k \in \mathbb{N}}$  each converge to x, and there are uncountably many of them. We conclude that for each open interval  $U \subseteq \mathbb{R}$ , if  $\nu_r(x) \in U$ , then  $U \cap A$  contains infinitely many elements of each sequence of the form  $\nu_r(\{vw_1, \dots, w_{m_k} z_{\lambda} : k \in \mathbb{N}\})$ , of which there are uncountably many. Hence x cannot be in L' nor in  $L^{iso}$  if it is in an uncountable  $V_i W_i^{\omega}$  component, or the closure thereof.

For the forward direction, we again show the contrapositive. We note that if there is no open interval  $U \subseteq \mathbb{R}$  containing  $x \in A$  for which  $U \cap A$  is countable, then there is a sequence of elements in an uncountable V-W component whose images under  $\nu_r$  converge to x. Otherwise, there is  $k \in \mathbb{N}$  such that  $x_{[0,k]}$  is not the prefix of any element in any uncountable V-W component, so  $U = \operatorname{int}(\nu_r(\{z : z_{[0,k]} = x_{[0,k]}\}))$  is an open set that has countable intersection with A. Yet we have seen that the existence of a sequence of elements in an uncountable V-W component whose images under  $\nu_r$  converge to x yields uncountably many such sequences. So every open set  $U \ni x$  will include elements from that uncountable set of convergent sequences.

So  $L^{iso}$  and L' must both be contained entirely in the countable V-W components of L, and each such component contains only elements from one tail-equivalence class, as shown in Proposition 2.3.

We will see next that if  $V_i W_i^{\omega}$  is uncountable, then  $\nu_r(V_i W_i^{\omega})$  has infinite Cantor-Bendixson rank, and that the set A' has finite Cantor-Bendixson rank.

# **Lemma 4.3.** Suppose that A is closed, and $x \in L' \setminus L^{iso}$ . Then there exists some $k \in \mathbb{N}$ and some r-sparse subset $B = V'_1 W'^{\omega}_1 \cup \ldots \cup V'_{\ell} W'^{\omega}_{\ell}$ such that if $y \in L$ and $y_{[0,k]} = x_{[0,k]}$ , then $y \in B$ .

Proof. We suppose for contradiction that  $x \in L' \setminus L^{iso}$  and also that for every  $k \in \mathbb{N}$  there is  $y_k \neq x$ with  $x_{[0,k]} = y_{k[0,k]}$  yet  $\{z \in L : z_{[0,k]} = x_{[0,k]}\}$  is not r-sparse. By the pigeonhole principle, we may assume that  $(y_k)_{k \in \mathbb{N}}$  are all contained in the same countable V-W component of L, call it  $V_i W_i^{\omega}$ . By the proof of Proposition 2.3, the fact that  $V_i W_i^{\omega}$  is countable means that we can take  $W_i^{\omega}$  to be generated by a single word; otherwise,  $W_i$  contains two distinct words of the same length, and thus  $W_i^{\omega}$  is uncountable.

So for  $\{y \in V_i W_i^{\omega} : y_{[0,k]} = x_{[0,k]}\}$  to not be *r*-sparse for each  $k \in \mathbb{N}$ , it must be the case that for each  $y_k$  there exist words  $a, b, u, v \in [r]^*$  such that  $x_{[0,k]} \subseteq v$  and  $v\{a, b\}^* u W_i^{\omega} \subseteq V_i W_i^{\omega}$ and  $y_k \in v\{a, b\}^* u W_i^{\omega}$ . Since *A* is closed, we know that *A* also contains all elements whose *r*representations are words of the form  $v\{a, b\}^{\omega}$ . Yet this implies that for each open interval  $U \subseteq \mathbb{R}$ containing  $\nu_r(x)$ , there is a prefix  $x_{[0,k]}$  such that all elements whose *r*-representations have that prefix are in *U*, including the uncountable set of elements with *r*-representations of the form  $v\{a, b\}^{\omega}$ . This contradicts the assumption that  $x \in L'$ .

So for each V-W component there must be a prefix of x such that any other element of that component with the same prefix is in an r-sparse subset.

There is one last piece to put in place before we can employ these lemmas to prove the primary result of this section. The final step is to show that for A a closed r-regular set, not only is the set of points that accumulate to any given  $x \in A'$  an r-sparse set, i.e. one with finite Cantor-Bendixson rank and vanishing Cantor-Bendixson derivative, but also there is a uniform bound on the Cantor-Bendixson rank of the set of elements in A that are sufficiently close to  $x \in A'$ . Here, "sufficiently close" means there is an open interval containing x, and whose diameter depends on x, in which the intersection with A has a Cantor-Bendixson rank less than the uniform bound, which does not depend on x.

**Lemma 4.4.** Suppose that L is closed. Then there exists  $N \in \mathbb{N}$  such that for every  $x \in L'$  there is some  $k \in \mathbb{N}$  such that the Cantor-Bendixson rank of  $\nu_r([x]_{\simeq_k,L})$  is at most N.

Proof. Recall that  $[x]_{\simeq_k,L} = \{y \in L : y_{[0,k]} = x_{[0,k]}\}$ . We suppose not, i.e. that there exists a sequence  $(x_n)_{n\in\mathbb{N}}$  such that for each  $n\in\mathbb{N}$  and each  $k\in\mathbb{N}$  the set  $[x_n]_{\simeq k,L}$  has Cantor-Bendixson rank greater than n. We can decompose L into  $E = int(\nu_r(L))$ , the closure of the interior of L, and  $L \setminus E$ , the maximal subset of L that is nowhere dense on an open subset of [0,1]. Since the Cantor-Bendixson derivative on an interval is itself, we may disregard E when bounding the Cantor-Bendixson rank of  $\nu_r([x]_{\simeq_k,L}\})$  for  $x \in L$ . So, without loss of generality, we may assume  $L = L \setminus E$ , i. e. that L has no interior.

By the pigeonhole principle, we may take all of the  $x_n$ 's to be in the same V-W component of L, call it  $V_i W_i^{\omega}$ . Since we have assumed that L has no interior, for each  $x_n$  there is  $k_n \in \mathbb{N}$  such that  $[x_n]_{\simeq_{k_n},L}$  is r-sparse. By Lemma 2.4, this means the Cantor-Bendixson rank of  $[x_n]_{\simeq_{k_n},L}$  is some finite  $m_n$  with  $m_n > n$  by hypothesis. Applying sequential compactness of the [0, 1] interval to the sequence  $(x_n)_{n\in\mathbb{N}}$ , we conclude that there is an infinite subsequence of  $(x_n)_{n\in\mathbb{N}}$  such that for n' > n it holds that  $x_{n[0,n]} \subseteq x_{n'[0,n']}$ . This also follows from König's Lemma (that any infinite finitely-branching tree has an infinite branch) because for each  $k \in \mathbb{N}$ , inifinitely many  $x_n$ 's have

the same length-k prefix, hence we can construct an infinite sequence in which  $x_{n+1}$  extends  $x_n$  for each  $n \in \mathbb{N}$ . Without loss of generality, let  $(x_n)_{n \in \mathbb{N}}$  now denote this subsequence.

Note that since  $x_{n+1}$  extends  $x_n$  for each  $n \in \mathbb{N}$ , it must be the case that  $k_{n+1} > k_n$ . Otherwise the sequence of  $k_n$ 's would stabilize, implying that eventually  $[x_n]_{\simeq_{k_n},L} = [x'_n]_{\simeq_{k_n'},L}$  for sufficiently large n', which in turn implies the Cantor-Bendixson rank of these sets stabilizes. We reach a contradiction because by definition of Cantor-Bendixson rank, removing the isolated points of  $[x_n]_{\simeq_{k_n},L}$  iteratively  $m_n$  times should yield the empty set. Yet by definition we know that for sufficiently large  $n' > m_n > n$  we have

$$[x_{n'}]_{\simeq_{k_{n'}},L} = \{y \in L : y_{[0,k_{n'}]} = x_{n'[0,k_{n'}]}\} \subseteq \{y \in L : y_{[0,k_n]} = x_{n[0,k_n]}\} = [x_n]_{\simeq_{k_n},L}$$

since  $k_n \leq k_{n'}$  ensures that any such  $y_{[0,k_{n'}]}$  contains the prefix  $x_{n[0,k_n]}$ . This would mean that removing the isolated points of  $(x_{n'})^{\simeq k_{n'}}$  iteratively  $m_n$  times yields the empty set, contradicting that  $m_{n'} > n'$ , the Cantor-Bendixson rank of  $[x_{n'}]_{\simeq k_{n'},L}$ , is strictly greater than  $m_n$ . So we conclude that there is some  $N \in \mathbb{N}$  such that for each  $x \in L'$  there is eventually a  $k \in \mathbb{N}$  such that the Cantor-Bendixson rank of  $[x]_{\simeq_{k},L}$  stabilizes to a number less than or equal to N.

Recall that within Euclidean space a *Cantor set*, used in the most general sense, is a nonempty subset of  $\mathbb{R}^d$  that is compact, has no isolated points, and has no interior.

**Theorem 4.5** (Theorem 2). If A is a closed r-regular subset of  $[0,1]^d$  and there is some  $i \in [d]$  such that  $0 < d_H(\pi_i(A)) < 1$ , then there is a unary set X definable in  $\mathcal{R}_A = (\mathbb{R}, <, +, 0, 1, A)$  such that X is a Cantor set.

*Proof.* Let  $i \in [d]$  be such that  $A_i := \pi_i(A)$  witness the hypothesis of the claim, i.e.  $0 < d_H(A_i) < 1$ . Note that the projection is compact since A is compact. We will definably remove a countable set of points from  $A_i$  such that the remaining set X is closed and has no isolated points. Then because  $d_H(A_i)$  is between 0 and 1, we conclude the same is true for X.

Recall that we use  $A'_i$  to denote the subset of  $A_i$  such that  $x \in A'_i$  precisely if there is some open interval I containing x such that  $A_i \cap I$  is countable. By Lemma 4.4, we know that there is some  $N \in \mathbb{N}$  such that if we strip away the isolated points of  $A_i^{iso} \cup A'_i$  at least N times, we are left with the empty set. By Lemma 4.2, we know that  $A_i^{iso} \cup A'_i$  is the part of  $A_i$  whose r-representations are contained in the complement of the closure of the uncountable  $V_k W_k^{\omega}$  components of L. Call the image under  $\nu_r$  of the closure of these uncountable components X. Hence we can write  $A_i$  as the disjoint union of  $A_i^{iso} \cup A'_i$  and X. When we take the N-th Cantor-Bendixson derivative of  $A_i$ , i.e. strip away the isolated points iteratively N times, what remains is X, since X being a closed set without isolated points makes it a perfect set, and hence its own Cantor-Bendixson derivative. Notably, taking the Cantor-Bendixson derivative N times is definable in the language (<, +, 0, 1), so we can isolate X definably.

Lastly, to see that X is a Cantor set we determine that it cannot have interior. If so, it would have a subset with Hausdorff dimension 1, since the Hausdorff 1-measure is positive whenever Lebesgue measure is positive.  $\Box$ 

Finally, we observe that even when A is not a closed set, its closure is definable in the structure  $\mathcal{R}_A$ , and r-regular, and hence we get our final corollary.

**Corollary 4.6** (Theorem 3). For  $A \subseteq [0,1]^d$  an r-regular set such that  $d_H(\overline{\pi_i(A)}) < 1$  for all  $i \in [d]$ , the following are equivalent:

- (1) A is r-sparse;
- (2)  $d_H(\overline{\pi_i(A)}) = 0$  for all  $i \in [d]$ ;

(3)  $\mathcal{R}_A$  is d-minimal; (4)  $\mathcal{R}_A$  has NIP; (5)  $\mathcal{R}_A$  has NTP<sub>2</sub>.

Proof. As above, set  $A_i := \pi_i(A)$  for  $i \in [d]$ . The equivalence of (1) and (2) follows from Lemma 2.7 and Lemma 3.2. To see that (2) implies (3), we note that if  $d_H(\overline{A_i}) = 0$  for each  $i \in [d]$ , then by Lemma 2.7 we know that  $\overline{A_i}$  is r-sparse, and hence countable, for each  $i \in [d]$ . We know  $\overline{A} \subseteq \overline{A_1} \times \cdots \times \overline{A_d}$  since coordinate projection is a continuous open map, and the latter set is countable. We conclude that A is countable, thus by Lemma 2.4 the set A is r-sparse. Hence by Theorem 3.10, we conclude that  $\mathcal{R}_A$  is d-minimal.

For (3) implies (4), this follows from [GH11], in which that authors show that certain d-minimal structures, which include  $\mathcal{R}_A$ , have NIP. That (4) implies (5) follows from the definitions of NIP and NTP<sub>2</sub>.

Finally, to see that (5) implies (2), we show the contrapositive. Suppose that  $d_H(\overline{\pi_i(A)}) > 0$  for some  $i \in [d]$ . By hypothesis,  $d_H(\overline{\pi_i(A)}) < 1$  for all  $i \in [d]$ , so by Theorem 2 we conclude that a Cantor set is definable in  $\mathcal{R}_A$ . By [HW19], this also means the structure interprets the monadic second order theory of the natural numbers with the successor function, which has TP<sub>2</sub> (and thus does not have NTP<sub>2</sub>).

### 5. SIMULTANEOUS SPARSITY IN MULTIPLICATIVELY INDEPENDENT BASES

In this section, we consider the intersection of two sparse sets that are regular with respect to two multiplicatively independent bases. We will show that sparse k- and  $\ell$ -regular sets have finite intersection when k and  $\ell$  are multiplicatively independent and give upper bounds on the size of the intersection in terms of the accepting Büchi automata for these sets. We note that this work shares some overlap with the paper [AB23].

**Lemma 5.1.** Let  $k \geq 2$  be a natural number and let S be a non-empty sparse k-regular subset of [0,1]. Then S is a finite union of sets  $S_1 \cup S_2 \cup \cdots \cup S_m$  such that for each  $i \in \{1,2,\ldots,m\}$  there exist  $s = s_i \geq 0, c_0, \ldots, c_s \in \mathbb{Q}$  such that  $(k^{\ell} - 1)c_i \in \mathbb{Z}$  for some  $\ell \geq 0, c_0 + c_1 + \cdots + c_s \in \mathbb{Z}_{\geq 0}$  and positive integers  $\delta_1, \ldots, \delta_s$  such that

(8) 
$$S_i \subseteq \left\{ c_0 k^q + c_1 k^{q+\delta_s n_s} + c_2 k^{q+\delta_s n_s+\delta_{s-1}n_{s-1}} \dots + c_s k^{q+\delta_s n_s+\dots+\delta_1 n_1} : n_1, \dots, n_s \ge 0, q \in -\mathbb{N} \right\}.$$

*Proof.* We know that a sparse k-regular subset of [0, 1] in a finite union of sets  $S_1, \ldots, S_m$  with each  $S_i$  a set of the form

$$\{[\bullet u_0 v_0^{n_1} u_1 v_1^{n_1} \cdots v_s^{n_s} u_s v_{s+1}^{\omega}]_k \colon n_1, \dots, n_s \ge 0\}.$$

Notice this set  $S_i$  is contained in the union of all sets of the form  $k^{-q}T_i$  with  $q \ge 0$ , where  $T_i$  is the set of rational numbers of the form

$$\{[u_0v_0^{n_1}u_1v_1^{n_1}\cdots v_s^{n_s}u_{s\bullet}v_{s+1}^{\omega}\}\$$

By a result of Ginsburg and Spanier [GS66], the set of natural numbers of the form  $[u_0v_0^{n_1}u_1v_1^{n_1}\cdots v_s^{n_s}u_s]_k$  is just the set

$$\{c_0 + c_1 k^{\delta_s n_s} + c_2 k^{\delta_s n_s + \delta_{s-1} n_{s-1}} \dots + c_s k^{\delta_s n_s + \dots + \delta_1 n_1} \colon n_1, \dots, n_s \ge 0\}$$

for some  $c_i, \delta_i$  as above. Since  $[\bullet v_{s+1}^{\omega}]_k$  is a rational number of the form  $c/(k^p - 1)$  for some positive integer c and some  $p \ge 1$ , we may replace  $c_0$  by  $c_0 + [\bullet v_{s+1}^{\omega}]_k$  and we get the desired result.  $\Box$ 

We call a k-regular subset of [0, 1] of the form

$$\{[\bullet u_0 v_0^{n_1} u_1 v_1^{n_1} \cdots v_s^{n_s} u_s v_{s+1}^{\omega}]_k \colon n_1, \dots, n_s \ge 0\}$$

a simple sparse k-regular set of length s.

The following result is primarily a consequence of Theorem 1.12.

**Lemma 5.2.** Let k and  $\ell$  be multiplicatively independent integers, let  $n, m \ge 1$ , and let  $a_0, \ldots, a_{n+m+1}$  be nonzero rational numbers, then there are at most

(9) 
$$\exp(3(6(n+m))^{3n+3m})$$

solutions to the equation

(10) 
$$a_0 X_0 + \dots + a_{n+m+1} X_{n+m+1} = 0$$

in which each  $X_i$  is a power of k for i = 0, ..., n and each  $X_i$  is a power of  $\ell$  for  $a_{n+1}, ..., a_{n+m+1}$ .

*Proof.* Letting  $a'_i = -a_i/a_0$  for i = 0, ..., m, we see that after dividing Equation (10) by  $a_0X_0$  that a nondegenerate solution to Equation (10) of the desired form corresponds to a nondegenerate solution  $(Y_1, \ldots, Y_{n+m+1})$  to the equation

$$\sum_{i=1}^{n+m+1} a'_i Y_i = 1$$

in which  $Y_1, \ldots, Y_n$  are integer powers of k, and  $Y_{n+1}, \ldots, Y_{n+m+1}$  are of the form  $k^c$  times an integer power of  $\ell$  for some fixed c. (Here  $k^c = 1/X_0$ .) By Theorem 1.12 there are at most  $\exp(3(6(n+m+1))^{3n+3m+1})$  nondegenerate solutions to this equation. Since the integer power  $k^c$  is uniquely determined by  $Y_{n+1}$ , we can recover our original solution  $(X_1, \ldots, X_{n+m+1})$  from  $(Y_1, \ldots, Y_{n+m})$ . The result follows.

**Lemma 5.3.** Let k and  $\ell$  be multiplicatively independent integers, let  $n, m \ge 1$ , and let  $a_0, \ldots, a_{n+m+1}$  be nonzero rational numbers, then there are at most

(11) 
$$(n+m+2)^{n+m+2} \exp(3(6(n+m+1))^{3(n+m+1)})$$

solutions to the equation

$$a_0 X_0 + \dots + a_{n+m+1} X_{n+m+1} = 0$$

in which each  $X_i$  is a power of k for i = 0, ..., n and each  $X_i$  is a power of  $\ell$  for  $a_{n+1}, ..., a_{n+m+1}$ and in which no non-trivial subsum of either  $a_0X_0 + \cdots + a_nX_n$  or  $a_{n+1}X_{n+1} + \cdots + a_{n+m+1}X_{n+m+1}$ vanishes.

*Proof.* Let  $\Gamma \leq \mathbb{Q}^*$  be the rank two group generated by k and  $\ell$ . For each solution to

$$a_0 X_0 + \dots + a_{n+m+1} X_{n+m+1} = 0$$

such that no subsum of either  $a_0X_0 + \cdots + a_nX_n$  or  $a_{n+1}X_{n+1} + \cdots + a_{n+m+1}X_{n+m+1}$  vanishes, we can form a set partition of the set of variables  $\{X_0, \ldots, X_{n+m+1}\}$  into disjoint non-empty subsets  $V_1, \ldots, V_s$  such that the subsum corresponding to the variables in each  $V_i$  vanishes and no proper subsum vanishes. By assumption, each  $V_i$  must intersect both  $\{X_0, \ldots, X_n\}$  and  $\{X_{n+1}, \ldots, X_{n+m+1}\}$  non-trivially. Then by Lemma 5.2, there are at most  $\exp(3(6(|V_i| - 1))^{3(|V_i|-1)}))$  nondegenerate solutions to the subsum equation

$$\sum_{X_j \in V_i} a_j X_j = 0$$

with each  $X_j$  a power of k for  $j \leq n$  and a power of  $\ell$  for j > n.

Thus for the set partition as above with parts  $V_1, V_2, \ldots, V_s$  we have at most

$$\prod_{i=1}^{s} \exp(3(6(|V_i|-1))^{3(|V_i|-1)}) \le \exp(3(6(n+m+1))^{3(n+m+1)})$$

solutions, where the last step follows from straightforward estimates for the exponential function.

Since a set partition of  $\{0, \ldots, n+m+1\}$  with s parts naturally gives rise to a surjective map from  $\{0, \ldots, n+m+1\}$  to  $\{1, \ldots, s\}$ , and since  $s \leq n+m+2$ , we see there are at most  $(n+m+2)^{n+m+2}$  possible set partitions. Putting this together, we get the desired bound.

**Proposition 5.4.** Let k and  $\ell$  be multiplicatively independent positive integers and let S be a sparse k-regular subset of [0, 1] of the form

$$\{\bullet[v_0w_1^*v_1w_2^*\cdots v_sw_s^*v_{s+1}w_{s+1}^{\omega}]_k\}$$

and let T be sparse  $\ell$ -automatic set of the form

$$\{[\bullet u_0 y_1^* u_1 y_2^* \cdots u_t y_t^* u_{t+1} y_{t+1}^{\omega}]_{\ell}\}$$

Then

$$\#S \cap T \le 2^{s+t+1} \cdot (s+t+2)^{s+t+2} \exp(3(6(s+t+1))^{3(s+t+1)}).$$

*Proof.* By Lemma 5.1 we have that S is contained in a set of the form

$$\left\{c_0k^{-q} + c_1k^{-q+\delta_s n_s} + c_2k^{-q+\delta_s n_s+\delta_{s-1}n_{s-1}} \dots + c_sk^{-q+\delta_s n_s+\dots+\delta_1 n_1} : n_1, \dots, n_s, q \ge 0\right\},\$$

where  $c_0, \ldots, c_s$  are rational numbers. Similarly, T is contained in a set of the form

$$\left\{d_0\ell^{-q'} + d_1\ell^{-q'+\delta_t'm_t} + d_2\ell^{-q'+\delta_t'm_s+\delta_{t-1}'m_{t-1}} \dots + d_t\ell^{-q'+\delta_t'm_t+\dots+\delta_1'm_1} \colon m_1,\dots,m_t,q'\ge 0\right\}$$

where  $d_0, \ldots, d_t$  are rational numbers.

Then an element in  $S \cap T$  corresponds to a solution to the equation

(12) 
$$d_0 X_0 + \dots + d_t X_t + d_{t+1} X_{t+1} + \dots + d_{t+s+1} X_{t+s+1} = 0,$$

where  $X_0 = \ell^{-q'}, X_1 = \ell^{-q'+\delta'_t m_t}, \ldots, X_t = \ell^{-q'+\delta'_t m_t+\cdots+\delta'_1 m_1}, X_{t+1} = k^{-q}, \ldots, X_{t+s+1} = k^{-q+\delta_s n_s+\cdots+\delta_1 n_1}$ , and where we take  $d_{t+j} = -c_{j-1}$  for  $j = 1, \ldots, s+1$ . Moreover, the element in the intersection in this case is given by

$$A := d_0 X_0 + \dots + d_t X_t = -(d_{t+1} X_{t+1} + \dots + d_{t+s+1} X_{t+s+1})$$

Since we are only concerned about the quantity A, we may remove a maximal vanishing subsum of  $d_0X_0 + \cdots + d_tX_t$  and a maximal vanishing subsum of  $d_{t+1}X_{t+1} + \cdots + d_{t+s+1}X_{t+s+1}$  and assume that both subsums are nondegenerate.

Then since there are at most  $2^{t+s+1}$  subsets of  $\{X_1, \ldots, X_{t+s+1}\}$  there are at most  $2^{t+s+1}$  choices for the maximal vanishing subsets we remove. Once we fix these subsets, Lemma 5.3 gives there are at most

$$(s+t+2)^{s+t+2} \exp(3(6(s+t+1))^{3(s+t+1)})$$

solutions to the resulting equation of the desired form. Thus there are at most

$$2^{s+t+1} \cdot (s+t+2)^{s+t+2} \exp(3(6(s+t+1))^{3(s+t+1)})$$

solutions to Equation (12) of the required form.

**Remark 5.5.** In general if S and T are respectively a sparse k-regular subset of [0,1] and a sparse  $\ell$ -regular subset of [0,1], then there are  $p,q \geq 1$  such that S is the union of p simple sparse sets of lengths  $s_1, \ldots, s_p$  and T is the union of q simple sparse sets of lengths  $t_1, \ldots, t_q$ . Moreover, we can determine p, q,  $s_1, \ldots, s_p, t_1, \ldots, t_q$  from the Büchi automata that accept S and T. In fact, if a and b are respectively the number of states in the Büchi automata that accept S and T, we have  $p \leq k^a, q \leq \ell^b, s := \max(s_1, \ldots, s_p) \leq a$ , and  $t := \max(t_1, \ldots, t_q) \leq b$ . Proposition 5.4 then gives a computable bound for  $S \cap T$  purely in terms of the number of states in the Büchi automata the sets S and T.

**Theorem 5.6.** Let k and  $\ell$  be multiplicatively independent positive integers and let S and T be respectively k- and  $\ell$ -regular sparse subsets of  $[0, 1]^d$ , with  $d \ge 1$ . Then  $S \cap T$  is finite.

Proof. Let  $\pi_i : [0,1]^d \to [0,1]$  denote the projection onto the *i*-th coordinate for  $i = 1, \ldots, d$ . Then  $\pi_i(S)$  is a sparse k-regular subset of [0,1] and  $\pi_i(T)$  is a sparse  $\ell$ -regular subset of [0,1]. It follows that  $Z_i := \pi_i(S) \cap \pi_i(T)$  is a finite set by Remark 5.5. Then since  $S \cap T \subseteq Z_1 \times Z_2 \times \cdots \times Z_d$ , we see that  $S \cap T$  is finite.

**Remark 5.7.** From the number of states in trim Büchi automata that accept S and T, we can obtain bounds for the intersection  $\pi_i(S) \cap \pi_i(T)$  for i = 1, ..., d by Remark 5.5 and so we can determine an upper bound for  $S \cap T$  in terms of this data.

#### References

- [AB23] S. Albayrak and J. P. Bell, Quantitative estimates for the size of an intersection of sparse automatic sets. arXiv: 2304.09223, 2023.
- [AS03] J.-P. Allouche and J. Shallit, Automatic Sequences. Theory, applications, generalizations. Cambridge University Press, Cambridge, 2003.
- [BGM20] J. P. Bell, D. Ghioca and R. Moosa, Effective isotrivial Mordell-Lang in positive characteristic. arXiv: 2010.08579, 2020.
- [BHS18] J. P. Bell, K. Hare and J. Shallit, When is an automatic set an additive basis? Proc. Amer. Math. Soc. Ser. B 5 (2018), 50–63.
- [BM19] J. Bell and R. Moosa, F-sets and finite automata. J. Théor. Nombres Bordeaux, 31(1),101–130, 2019.
- [BG et al.20] A. Block Gorman, P. Hieronymi, E. Kaplan, R. Meng, E. Walsberg, Z. Wang, Z. Xiong, and H. Yang, Continuous regular functions, *Log. Methods Comput. Sci.*, 16(1): 17:1–17:24, 2020.
- [BRW98] B. Boigelot, S. Rassart, and P. Wolper, On the expressiveness of real and integer arithmetic automata (extended abstract). Proceedings of the 25th International Colloquium on Automata, Languages and Programming (London, UK), ICALP '98, Springer-Verlag, pp. 152–163, 1998.
- [Büc62] J. Büchi, On a decision method in restricted second order arithmetic. *Logic, Methodology and Philosophy* of Science (Proc. 1960 Internat. Congr.), Stanford Univ. Press, Stanford, Calif., pp. 1–11, 1962.
- [CHM02] C. A. Cabrelli and K. E. Hare and U. M. Molter, Sums of Cantor sets yielding an interval. J. Aust. Math. Soc. 73 (2002), 405–418.
- [CLR15] É. Charlier, J. Leroy, and M. Rigo, An analogue of Cobham's theorem for graph directed iterated function systems. Adv. Math., 280:86–120, 2015.
- [Der07] H. Derksen, A Skolem-Mahler-Lech theorem in positive characteristic and finite automata. Invent. Math. 168 (2007), no. 1. 175–224.
- [vdD85] L. van den Dries, The field of reals with a predicate for the powers of two. Manuscripta Math., 54, 187–195, 1985.
- [EG15] J.-H. Evertse and K. Győry, Unit equations in Diophantine number theory. Cambridge Studies in Advanced Mathematics, 146. Cambridge University Press, Cambridge, 2015.
- [ESS02] J.-H. Evertse, H. P. Schlickewei, and W. M. Schmidt, Linear equations in variables which lie in a multiplicative group. Ann. of Math. 155 (2002), 807–836.
- [Fal03] Kenneth Falconer, Fractal Geometry: Mathematical Foundations and Application. John Wiley & Sons, 2003.
- [GKRS10] P. Gawrychowski, D. Krieger, N. Rampersad, and J. Shallit, Finding the growth rate of a regular or context-free language in polynomial time. *Int. J. Found. Comput. Sci.* **21** (2010), 597–618.

[GS66]	S. Ginsburg and E. Spanier, Bounded regular sets. Proc. Amer. Math. Soc. 17 (1966), 1043–1049.
[GH11]	A. Günaydın and P. Hieronymi, Dependent Pairs. J. Symbolic Logic, 76(2): 377–390, 2011.
[Haw 20]	C. Hawthorne, Automata and tame expansions of $(\mathbb{Z}, +)$ . Isr. J. Math., 249: 651–693, 2022.
[HW19]	P. Hieronymi and E. Walsberg, Fractals and the monadic second order theory of one successor. J. Log.
	Anal., 15(5): 1–25, 2023.
[IR86]	O. H. Ibarra and B. Ravikumar, On sparseness, ambiguity, and other decision problems for acceptors and transducers. STACS 86 (Orsay, 1986), 171–179, Lecture Notes in Comput. Sci., 210, Springer,
	Berlin, 1986.
[Kec95]	A. S. Kechris, Classical descriptive set theory, Grad. Texts in Math., 156 Springer-Verlag, New York, 1995.
[Ked06]	K. S. Kedlaya, Finite automata and algebraic extensions of function fields. J. Théorie Nombres Bordeaux 18 (2006), no. 2, 379–420.
[Ked17]	K. S. Kedlaya, On the algebraicity of generalized power series. <i>Beitr. Algebra Geom.</i> 58 (2017), no. 3, 499–527.
[M02]	D. Marker. Model Theory: An Introduction. Graduate Texts in Mathematics. Springer New York, 2002.
[Mi05]	C. Miller, Tameness in expansions of the real field. In <i>Logic Colloquium '01</i> , volume 20 of <i>Lecture Notes</i> in Logic pages 281–216. Association of Symbolic Logic Urbano, H. 2005.
[MTT10]	<i>in Logic</i> , pages 201–510. Association of symbolic Logic, Orbana, H., 2003.
[101116]	Proc. Amer. Math. Soc., 146: 5169–5179, 2018.
[MT06]	C. Miller and J. Tyne, Expansions of o-Minimal Structures by Iteration Sequences. <i>Notre Dame J. Formal Logic</i> , 47(1):93–99, 2006.
[MS02]	R. Moosa and T. Scanlon. The Mordell-Lang conjecture in positive characteristic revisited. <i>Model</i>
[]	Theory and its applications, 273–296, Quad. Mat. 11, Aracne, Rome, 2002.
[MS04]	R. Moosa and T. Scanlon, <i>F</i> -structures and integral points on semiabelian varieties over finite fields. <i>Amer. J. Math.</i> 126 (3):473–522, 2004.
[Mor66]	P. Moran, Additive functions of intervals and Hausdorff measure. Proc. Cambridge Philos. Soc. 42:15–23, 1046
[Lot02]	1340. M. Lothairo Alashraic Combinatorics on Words Cambridge University Press Cambridge UK 2002
[L0102] [Sch00]	H. P. Schlickowai, S unit equations over number fields. Invent. Math. 102 (1000), 05-107
[Sim15]	In I. Schneckwei, S-unit equations over humber nettes. Internet. Math. 102 (1950), 95 101.
[Sin13]	M Signa Introduction to the Theorem of Commutation 2nd edition Compared Learning Conserved to compare
ျပားစု၊	India Private Limited, Delhi, India, 2013.
[Tro81]	V. I. Trofimov, Growth functions of some classes of languages. (Russian). Kibernetika (Kiev) 1981, no.
	6, i, 9–12, 149.

UNIVERSITY OF WATERLOO, DEPARTMENT OF PURE MATHEMATICS, WATERLOO, ONTARIO, N2L 3G1, CANADA *Email address*: jpbell@uwaterloo.ca

The Ohio State University, Department of Mathematics, Mathematics Tower 100, 231 W 18th Ave, Columbus, OH, 43210

Email address: blockgorman.1@osu.edu