Learning Causal Representations from General Environments: Identifiability and Intrinsic Ambiguity

Jikai Jin*,1 and Vasilis Syrgkanis*,2

 1 Institute for Computational & Mathematical Engineering, Stanford University, CA, USA ²Department of Management Science & Engineering, Stanford University, CA, USA *Corresponding author. Email: [jkjin@stanford.edu;](jkjin@stanford.edu) <vsyrgk@stanford.edu>

Abstract

We study causal representation learning, the task of recovering high-level latent variables and their causal relationships in the form of a causal graph from low-level observed data (such as text and images), assuming access to observations generated from multiple environments. Prior results on the identifiability of causal representations typically assume access to single-node interventions which is rather unrealistic in practice, since the latent variables are unknown in the first place. In this work, we provide the first identifiability results based on data that stem from *general environments*. We show that for linear causal models, while the causal graph can be fully recovered, the latent variables are only identified up to the *surrounded-node ambiguity* (SNA) (Varici et al. [2023\)](#page-13-0). We provide a counterpart of our guarantee, showing that SNA is basically unavoidable in our setting. We also propose an algorithm, LiNGCReL which provably recovers the ground-truth model up to SNA, and we demonstrate its effectiveness via numerical experiments. Finally, we consider general non-parametric causal models and show that the same identification barrier holds when assuming access to groups of soft single-node interventions.

Keywords: Causal Representation Learning

1. Introduction

Artificial intelligence (AI) has achieved tremendous success in various domains in the past decade (Bengio, Courville, and Vincent [2013;](#page-11-0) Silver et al. [2016;](#page-13-1) Bubeck et al. [2023\)](#page-11-1). However, current approaches are largely based on learning the *statistical* structures and relationships in the data that we observe. As a result, it is not surprising that these approaches often capture spurious statistical dependencies between different features, resulting in poor performance in the presence of test distribution shift (Ovadia et al. [2019;](#page-13-2) Koh et al. [2021\)](#page-12-0) or adversarial attacks (Akhtar and Mian [2018;](#page-11-2) Wang et al. [2023\)](#page-13-3).

In view of these pitfalls, a recent line of work has explored the problem of *causal representation learning* (Schölkopf et al. [2021\)](#page-13-4), the task of learning the causal relationships between high-level latent variables underlying our low-level observations. Notably, it is widely believed in cognitive psychology that humans take a causal approach to distill information from the world and make decisions to achieve their goals (Shanks and Dickinson [1988;](#page-13-5) Dunbar and Fugelsang [2004;](#page-12-1) Holyoak and Cheng [2011\)](#page-12-2). As a result, there is reason to believe that learning causal representations has the potential to significantly improve the power of AI, especially on tasks where performance lags far behind human level (Geirhos et al. [2020\)](#page-12-3).

[©] Working Paper 2023. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence [\(http://creativecommons.org/licenses/by/4.0/\)](http://creativecommons.org/licenses/by/4.0/), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

2 Jikai Jin *et al.*

Despite such promise, a crucial challenge in causal representation learning is the *identifiability* of the data generating process; in other words, given the data that we observe, can we uniquely identify the underlying causal model. It has been shown that given observational data (*i.e.*, i.i.d. data generated from a single environment), the model is already non-identifiable in strictly simpler settings where the latent variables are known to be independent (Locatello et al. [2019;](#page-12-4) Locatello, Bauer, Lucic, et al. [2020\)](#page-12-5), or where there is no mixing function and one directly observes the latent variables (Silva et al. [2006\)](#page-13-6). A natural question that arises is what types of data do we need to acquire to make identification possible.

One line of works assumes access to counterfactual data (Locatello, Poole, et al. [2020;](#page-12-6) Von Kügelgen et al. [2021;](#page-13-7) Brehmer et al. [2022\)](#page-11-3), where some form of *weak supervision* is typically required. A common assumption here is that one observes data in *pairs*, where each pair of data is related via sharing part of the latent representation. However, such data is hard to acquire since it requires direct control on the latent representation.

Another line of works (Ahuja et al. [2023;](#page-11-4) Kügelgen et al. [2023;](#page-12-7) Buchholz et al. [2023;](#page-11-5) Varıcı et al. [2023\)](#page-13-8) instead considers an interventional setting, where the learner observes data generated from multiple different environments. This is arguably a much more realistic setup and reflects common practices in robotics (Lippe et al. [2023\)](#page-12-8) and genomics (Lopez et al. [2023;](#page-12-9) Tejada-Lapuerta et al. [2023\)](#page-13-9) applications. However, a vast majority of identifiability guarantees assume that each environment corresponds to *single-node*, *hard* interventions, which is defined as interventions that isolate a single latent variable from its causal parents. Again, this is quite a restrictive assumption because of two reasons. *First*, since the latent variables are unknown and need to be learned from data, it is unclear how to perform interventions that only affect one variable. *Second*, even if one can perform single-node interventions, it may not be feasible to artificially remove causal effects in the data generating processes. This issue is ubiquitous in real-world applications as pointed out in Campbell [\(2007\)](#page-11-6), Eberhardt [\(2014\)](#page-12-10), and Eronen [\(2020\)](#page-12-11). Motivated by these challenges, we instead consider the following two settings:

- Learning from *single-node, soft* interventions, which only change the dependency of each latent variable on its direct causes, but does not remove their causal relationships. This setting is considered in Seigal, Squires, and Uhler [\(2022\)](#page-13-10), H. Zhang et al. [\(2022\)](#page-13-11), Varici et al. [\(2023\)](#page-13-0), and Buchholz et al. [\(2023\)](#page-11-5), which, however, make parametric assumptions on either the causal model or the mixing function. The most related paper is Varici et al. [\(2023\)](#page-13-0), which proves identifiability under an ambiguity induced by the so-called "surrounded set". In this paper, we show that this type of ambiguity is intrinsic in the soft intervention setting.
- Learning from *fully general and diverse environments*. This is a significantly more general and challenging setting, and to the best of our knowledge, no identifiability guarantees are known. Khemakhem et al. [\(2020\)](#page-12-12) and Lu et al. [\(2021\)](#page-12-13) also consider a multi-environment setup without assuming single-node interventions, but they still assume that the distributions of latent variables all come from a certain parametric family with a fixed set of sufficient statistics.

We make the following contributions:

- For linear causal models, with a linear mixing function, we prove identification results assuming access to data from *general and diverse environments* [\(Theorem 1\)](#page-5-0). To the best of our knowledge, this is the first identification guarantee that makes no assumption on the relationship between the environments. Interestingly, while we show that the causal graph can be exactly recovered, the latent variables are only recovered up to a *surrounded-node ambiguity* (SNA) [\(Theorem 3\)](#page-6-0).
- We propose an algorithm, LiNGCReL, in [Section 5](#page-6-1) that provably recovers the ground-truth model up to SNA [\(Theorem 4\)](#page-7-0) in the setting of [Theorem 1.](#page-5-0) To demonstrate the effectiveness of LiNGCReL, we present extensive experimental results in [Section 6](#page-7-1) using it to learn causal representations from randomly generated causal models.

• Going beyond the linear setting, we study the limit of identification for non-parametric causal models and general mixing functions, assuming access to single-node soft interventions. We show that the model is identifiable up to SNA [\(Theorem 5\)](#page-10-0), and then prove that SNA is actually the best achievable guarantee in this setting [\(Theorem 6\)](#page-10-1), thereby highlighting a key difference between soft and hard interventions.

2. Preliminaries

We consider the standard setup of causal representation learning from multiple environments $E \in \mathfrak{E}$. Let $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ be the ground-truth causal graph which is directed and acylic (DAG), where $\mathcal{V} = [d]$ and $\mathcal E$ describes the causal relationship between different nodes. Each node corresponds to a latent variable $z_i \in \mathbb{R}$.

For any node *i*, we let pa $_{\mathcal{G}}(i)$, ch $_{\mathcal{G}}(i)$, ans $_{\mathcal{G}}(i)$ and nd $_{\mathcal{G}}(i)$ to be the set of all parents, children, ancestors and non-descendants of *i* in $\cal G$ respectively. We also define $\overline{\text{pa}}_{\cal G}(i)$ = $\text{pa}_{\cal G}(i) \cup \{i\}$ and similarly for $\overline{\ch}_{G}(i)$, $\overline{\mathrm{ans}}_{G}(i)$ and $\overline{\mathrm{nd}}_{G}(i)$. Assuming that all probability distributions have continuous densities, the joint density of the latent variables *z* can then be written as

$$
p_E(z) = \prod_{i=1}^d p_i^E \left(z_i \mid z_{\text{pa}_G(i)} \right).
$$
 (1)

where p_i^E is the (unknown) latent generating distribution from environment *E* at node *i*. Here for a given vector v , we write $v_i = e_i^\top v$, and let $v_S = (v_i : i \in S) \in \mathbb{R}^{|S|}$.

The causal graph model with density given by [\(1\)](#page-2-0) necessarily enjoys the following property:

 $\bf{Definition 1}$ (Causal Markov Condition). *For any node i, conditioned on* $\boldsymbol{z}_{\text{pa}_{\mathcal{G}}(i)}$ *,* \boldsymbol{z}_i *is independent of* $z_{\mathrm{nd}_{\mathcal{G}}(i)}$. As a consequence, for any node i, $j\in [d]$ and $S\subseteq [d]$, if S d-separates i from j (cf. [Definition 8\)](#page-18-0), ι *hen* $z_i ⊥ z_j ∣ z_S$.

The latent variables *z* are unknown to the learner. Instead, the learner has access to observations $x \in \mathbb{R}^n$ $(n \geqslant d)$ from all environments $E \in \mathfrak{E}$ that are related to the latent z via an injective mixing function *g*:

$$
x = g(z). \tag{2}
$$

The main assumption here that the mixing function is the same across all environments:

 ${\bf Assumption~1.~}$ All environments $E\in \mathfrak{E}$ share the same diffeomorphic mixing function $g:\mathbb{R}^d\mapsto \mathbb{R}^n$ *.*

In causal representation learning, the goal of the learner is to 1) recover the inverse of the mixing function $h = g^{-1}$ (often called the *unmixing* function) which allows recovering the latent variables given any observations, and, 2) recover the underlying causal graph G . In the remaining part of this paper, we refer to (h, \mathcal{G}) as the causal model to be learned. Obviously, there would be some ambiguities in learning (*h*, G). For example, choosing a different permutation of the nodes in the causal graph would lead to a different model, and so does element-wise transformations on each component *hi* of *h*.

A line of recent works show that the ground-truth model can be identified up to these ambiguities in various settings, assuming access to single-node hard interventions (Seigal, Squires, and Uhler [2022;](#page-13-10) Kügelgen et al. [2023;](#page-12-7) Varıcı et al. [2023\)](#page-13-8). On the other hand, some weaker notions of identifiability have also been proposed and studied in the literature (Seigal, Squires, and Uhler [2022;](#page-13-10) Varici et al. [2023;](#page-13-0) Liang et al. [2023\)](#page-12-14) for single-node soft interventions. Here, we provide a generic definition of single-node soft interventions that we will rely on in this paper.

Definition 2. We say that a collection of environments $\hat{\mathfrak{E}}$ is a set of (soft) interventions on a subset of latent variables $\bm{z}_i, i\in S$ if $\forall i\in[d],\forall E_1,E_2\in \hat{\mathfrak{E}},E_1\neq E_2$, we have $p_i^{E_1}=p_i^{E_2}\Leftrightarrow i\notin S$ (the notation p_i^E comes from [\(1\)](#page-2-0)). Equivalently, we write $\mathcal{I}_{z}^{\hat{\mathfrak{E}}}$ = S.

We note that soft interventions are very different from hard interventions, since they do not remove causal relationships between latent variables. The goal of this paper is to address the following question:

What is the best-achievable identification guarantee when hard interventions are not available, and what are the intrinsic ambiguities?

3. The effect domination set and a notion of identifiability

One may expect that identifiability with soft interventions is not much different from hard interventions, since soft interventions can approximate hard interventions with arbitrary accuracy. However, we will show that this is not the case. At a high level, hard intervention is more powerful than soft intervention because it is capable of isolating a latent variable from its direct cause while soft interventions is not, so soft interventions can sometimes fail to identify the true causal relationship from a mixture of causal effects.

To quantify what kind of ambiguities may arise, we can define the surrounding set for each node in a causal graph G as follows:

Definition 3. *(Varici et al.* [2023,](#page-13-0) Definition 3) For two nodes $i, j \in [d]$ in \mathcal{G} , we say that j is effect-dominated b y i, or i \in $\text{sur}_{\mathcal{G}}(j)$ if $i \in \text{pa}_{\mathcal{G}}(j)$, and $\text{ch}_{\mathcal{G}}(j) \subseteq \text{ch}_{\mathcal{G}}(i)$ *. Moreover, we define* $\overline{\text{sur}}_{\mathcal{G}}(j) = \text{sur}_{\mathcal{G}}(j) \cup \{j\}$.

In other words, the effect of *j* on its child set ch_G(*j*) is dominated by the effect of *i*. Intuitively, if there exists some *i* ∈ sur $G(j)$, then ambiguities may arise for the causal variable at node *j*, since any effect of *j* on any of its child *k* can also be interpreted as an effect of *i*. In [Section E](#page-19-0) we discuss a three-node example to further illustrate such ambiguities.

Figure 1. An illustration of [Definition 3;](#page-3-0) here $i \in \text{sur}_G(i)$.

[Definition 3](#page-3-0) naturally induces the following relationship between causal models:

Definition 4. *Using the notations in [Definition 11,](#page-18-1) we write* (*h*, G) ∼sur (ˆ*h*, Gˆ) *if there exists a permutations* π on [d], and a diffeomorphism $\psi:\R^d\mapsto\R^d$ where the j-th component of ψ , denoted by $\psi_j(\bm z)$, is a function *of* $\boldsymbol{z}_{\overline{\text{sur}}_\mathcal{G}(j)}$ *for* $\forall j \in [d]$ *, such that the following holds:*

- *For* $\forall i, j \in [d]$, $i \in pa_{\mathcal{G}}(j) \Leftrightarrow \pi(i) \in pa_{\hat{\mathcal{G}}}(\pi(j))$,
- $P_\pi \circ \hat{h}$ = $\psi \circ h$ *, where* P_π *is a permutation matrix satisfying* $(P_\pi)_{ij}$ = 1 \Leftrightarrow *j* = $\pi(i)$ *.*

In other words, \sim _{sur} requires that the causal graph to be exactly the same up to some permutation of nodes, but allows each latent variable v_i to be entangled with $z_{\textrm{sur}_{\mathcal{G}}(i)}.$ Although not obvious from definition, one can actually check that ∼sur defines an *equivalence relation* (see [Lemma 12\)](#page-22-0). Moreover, we will show later that \sim _{sur} is in general the best that we can hope for in our problem setting.

4. Identifiability of linear causal models from general environments

In this section, we consider learning causal models from *general* environments. Specifically, we assume that the environments $E_k, k \in [K]$ share the same causal graph, but the dependencies between connected nodes (latent variables) are completely unknown, and, in contrast with existing literature on single-node interventions, we impose no similarity constraints on the environments. We begin our investigation of identifiability in this setting in the context of linear causal models with a linear mixing function.

4.1 Problem setup

Formally, we assume the following generative model in *K* distinct environments $\mathfrak{E} = \{E_k : k \in [K]\}$ with data generating process

$$
z = A_k z + \Omega_k^{\frac{1}{2}} \epsilon, \quad x = Gz \quad k \in [K], \tag{3}
$$

where the matrix A_k satisfies $(A_k)_{ij} \neq 0$ if and only if $j \to i$ in $\mathcal G$. We refer to (A_k, Ω_k) as the weight matrices of latent variables z in the environment E_k . It is easy to see that [Assumption 1](#page-2-1) in our general setup translates into the following assumption:

Assumption 2. *The mixing matrix* $\mathbf{G} \in \mathbb{R}^{n \times d}$ has full column rank. Equivalently, the unmixing matrix *H* = *G*† *has full row rank.*

Let $B_k = \Omega_k^{-\frac{1}{2}} (I - A_k)$, $k \in [K]$, then we have $\epsilon = B_k z = B_k H x$. Since in the linear case, there is an easy to see one-to-one correspondence between the matrix *H* and the un-mixing function $x \mapsto Hx$, we abuse the notation and write (H, G) to represent the model instead of (h, G) . Using h_i to denote the *i*-th row of *H*, the following lemma translates [Definition 4](#page-3-1) the the linear setting:

Lemma 1. *According to [Definition 4,](#page-3-1)* $(H, G) \sim_{\text{sur}} (\hat{H}, \hat{G})$ *if and only if there exists a permutation* π *on* $[d]$ *, such that* $i \in pa_{\mathcal{G}}(j) \Leftrightarrow \pi(i) \in pa_{\hat{\mathcal{G}}}(\pi(j))$ *, and for* $\forall i \in [d]$ *,* $\hat{h}_i \in span \langle h_j : \pi(j) \in \overline{\text{sur}}_{\mathcal{G}}(i) \rangle$ *.*

We also need to make the following assumption on noise.

 $\bf{Assumption~3.}$ $\it The$ noise vector ${\bm \epsilon} \in \mathbb{R}^d$ has independent components, at most one component is Gaussian *distributed, and any two components have different distribution.*

The non-gaussianity of the noise vectors is a typical assumption in causal discovery within linear models (Comon [1994;](#page-12-15) Silva et al. [2006\)](#page-13-6) and is always assumed in the LinGAM setting (Shimizu et al. [2006\)](#page-13-12). The assumption that all components have a different distribution is not so standard, but is quite natural in real-world scenarios.

4.2 Identifiability guarantee

For each node $i \in [d]$ of G, we use $w_k(i)$ to be the *weight vector* of environment E_k at node *i*, $i.e.,\, \pmb{w}_k(i)=\left((\pmb{A}_k)_{ij}: j\in\text{pa}_\mathcal{G}(i)\right)\in\mathbb{R}^{\left|\text{pa}_\mathcal{G}(i)\right|}.$ In other words, the structural equation for node i in environment *k* is of the form:

$$
z_i = w_k(i)^{\top} z_{\text{pa}_{\mathcal{G}}(i)} + \sqrt{\omega_{k,i,i}} \varepsilon_i
$$
 (4)

To obtain our identifiability result, the main assumption we need to make is the non-degeneracy of the weights at each node:

 $\bf{Assumption~4.}$ *For each node* $i \in [d]$ *of* \mathcal{G} *, we have* $\operatorname{aff}\left({\bm{w}}_k(i): k \in [K]\right) = \mathbb{R}^{\left|\Pr{\mathcal{G}}(i)\right|}_i$ where $\operatorname{aff}(\cdot)$ denotes the affine hull. Equivalently, the weights $w_k(i)$, $k = 1, 2, \cdots, K$ do not lie in a $\left(\left|\text{pa}_\mathcal{G}(i)\right|-1\right)$ -dimensional h yperplane of $\mathbb{R}^{\left|\mathrm{pa}_{\mathcal{G}}\left(i\right)\right|}$ *.*

This assumption is quite mild since it only requires the weight vectors to be in general positions, and it holds with probability 1 if the weights at each node are sampled from continuous distributions. Moreover, as shown in [Lemma 6,](#page-21-0) it is equivalent to the following assumption.

 $\bf Assumption~5$ (Node-level non-degeneracy). *We say that the matrices* $\{\mathcal{B}_k\}_{k=1}^K$ *are node-level non-* \deg enerate if for all node $i \in [d]$, we have \dim span $\langle (\boldsymbol{B}_k)_i : k \in [K] \rangle$ = $\left| \text{pa}_{\mathcal{G}}(i) \right|$ + 1, where $(\boldsymbol{B}_k)_i$ is the i -th *row of* B_k . *.*

In the following, we state our main result in this section, which shows that $K = d$ non-degenerate environments suffices for the model to be identifiable up to \sim _{sur}.

Theorem 1. *Suppose that* $K \geq d$ *and we have access to observations generated from the linear causal model* (H, \mathcal{G}) across multiple environments $\mathfrak{E} = \{E_k : k \in [K]\}$ with observation distributions $\{\mathbb{P}^E_x\}_{E \in \mathfrak{E}}$, and the *data generating processes are given by* [\(3\)](#page-4-0)*. Let* (*H*ˆ , Gˆ) *be any candidate solution with the hypothetical data generating process*

$$
\mathbf{v} = \hat{\mathbf{A}}_k \mathbf{v} + \hat{\mathbf{\Omega}}_k^{\frac{1}{2}} \hat{\mathbf{\varepsilon}}, \quad \mathbf{x} = \hat{\mathbf{H}}^\dagger \mathbf{v} \quad \text{in the environment } E_k
$$

*where H*ˆ *has full row rank, such that*

- (i) the observation distribution that this hypothetical model generates in E_k is exactly $\mathbb{P}_{\mathbf{x}}^{E_k},$
- *(ii)* all environments share the same causal graph: ∀ $k \in [K]$ and i, $j \in [d]$, $(A_k)_{ij} \neq 0 \Leftrightarrow j \in \text{pa}_{\mathcal{G}}(i)$,
- $(\hat{\bm{A}}_k)_{ij} \neq 0 \Leftrightarrow j \in \text{pa}_{\hat{\mathcal{G}}}(i)$ and $\bm{\Omega}_k, \hat{\bm{\Omega}}_k$ are diagonal matrices with positive entries;
- *(iii)* $\{B_k\}_{k=1}^K$ *and* $\left\{\hat{B}_k = \hat{\Omega}_k^{-\frac{1}{2}}(I \hat{A}_k)\right\}_{k=1}^K$ *k*=1 *are non-degenerate in the sense of [Assumption 5;](#page-5-1)*

(iv) the noise variables ϵ *and* $\hat{\epsilon}$ *satisfy [Assumption 3.](#page-4-1)*

Then we must have $(H, \mathcal{G}) \sim_{\text{sur}} (\hat{H}, \hat{\mathcal{G}})$ *.*

The proof of [Theorem 1](#page-5-0) is given in [Section H.1.](#page-23-0) In the next section, we will introduce an algorithm, LiNGCReL, that provably recovers the ground-truth up to \sim_{sur} .

To the best of our knowledge, this is the first identifiability guarantee in the literature for causal representation learning from general environments. Remarkably, while the fact that existing works (Seigal, Squires, and Uhler [2022;](#page-13-10) J. Zhang et al. [2023\)](#page-13-13) focus on single-node interventions seem to suggest that learning from *diverse* environments is hard, our result indicates that such diversity is actually helpful. Specifically, we show that in the worst case, $\Theta(d^2)$ interventions are required for identifying the ground-truth model under \sim_{sur} :

Theorem 2 (informal version of [Theorem 8\)](#page-27-0)**.** *There exists a causal graph* G *with* Θ(*d* 2) *edges, such* that for any unmixing matrix $\bm{H}\in\mathbb{R}^{d\times n}$ with full row rank, any independent noise variables $\bm{\epsilon}$, and any $0 \le s_i \le \left| \text{pa}_{\mathcal{G}}(i) \right|, i \in [d],$ the ground-truth model (H, \mathcal{G}) is non-identifiable up to ∼ $_{\text{sur}}$ *with s_i* soft *interventions for node i, unless the (ground-truth and intervened) weights of the causal model lie in a null set (w.r.t the Lebesgue measure).*

A formal version and the proof of [Theorem 2](#page-5-2) can be found in [Section H.2.](#page-26-0) On the other hand, by having *d* single-node interventions per node, [Assumption 5](#page-5-1) can be satisfied as long as the weights are in general positions, so in this case we have $(H, \mathcal{G}) \sim_{\text{sur}} (\hat{H}, \hat{\mathcal{G}})$ by [Theorem 1.](#page-5-0) Therefore, [Theorems 1](#page-5-0) and [8](#page-27-0) together imply that $\Theta(d^2)$ single-node interventions are necessary and sufficient for identification up to \sim _{sur}.

Given that [Theorem 1](#page-5-0) only guarantees identification up to \sim_{sur} that is strictly weaker than full identification, one might naturally ask whether [Theorem 1](#page-5-0) can be further improved. Our last theorem in this section indicates that ∼sur is indeed the *best* one can hope for in our setting, even assuming access to *single node, soft intervention*.

Theorem 3 (Counterpart to [Theorem 1,](#page-5-0) informal version of [Theorem 9\)](#page-36-0)**.** *For any linear causal model* (H, \mathcal{G}) and any set of environments \mathfrak{E} = $\{E_k : k \in [K]\}$ such that all conditions in [Theorem 1](#page-5-0) are satisfied, *there must exists a candidate solution* (*H*ˆ , G) *and a hypothetical data generating process that satisfy the same set of conditions, but*

$$
\frac{\partial v_i}{\partial z_j} \neq 0, \quad \forall j \in \overline{\text{sur}}_{\mathcal{G}}(i).
$$

Moreover, if we additionally assume that the environments are groups of single-node soft interventions, then we can guarantee the existence of (H, \mathcal{G}) *and weight matrices which, besides the properties listed above, are also groups of single-node soft interventions.*

5. LinGCReL**: Algorithm for linear non-Gaussian causal representation learning**

In this section, we describe Linear Non-Gaussian Causal Representation Learning (LiNGCReL), an algorithm that provably recovers the underlying causal graph and latent variables up to \sim _{sur} in the infinite-sample limit. At this point, it is instructive to recall the celebrated LiNGAM algorithm (Shimizu et al. [2006\)](#page-13-12) for linear causal graph discovery. Different from their setting, we only observe some unknown linear mixture of the latent variables. Hence, running linear ICA as in LiNGAM only gives us $M_k = B_k H$ rather than the weight matrix B_k itself.

The key idea in our approach is an effect cancellation scheme that allows us to determine the "remaining degree of freedom" (RDF) of any node (*a.k.a.* latent variable) given any subset of its ancestors. This scheme allows us to not only find a topological order of the nodes, but also figure out direct causes by tracking the changes of the RDF. In the following, we present the main steps of LiNGCReL in more details.

Suppose that we are given samples of observations $X^{(k)} = \left\{ x_i^{(k)} \right\}$ $\begin{bmatrix} (k) \\ i \end{bmatrix}$ $\begin{bmatrix} N \\ i \end{bmatrix}$ $\sum_{i=1}^{N}$, $k \in [K]$ where $x_i^{(k)}$ $i_i^{(n)}$ is the *i*-th sample from the *k*-th environment.

Step 1. Recover the matrices $M_k = B_k H$ **Since** $\epsilon = B_k z = B_k H x$ **in the** *k***-th environment, so** we can use any identification algorithm for linear ICA to recover the matrix M_k . Then we properly rearrange the rows of M_k so that all $M_k x, k = 1, 2, \cdots, K$ correspond to the same permutation of noise variables. This step is quite standard and details can be found in [Section B.1.](#page-15-0)

Step 2. Causal representation learning based on M_k **Now we have obtained** M_k **=** B_kH **, but** the unmixing matrix *H* is still unknown. We propose [Algorithm 3](#page-8-0) to learn *H* and the causal graph G. The main part of [Algorithm 3](#page-8-0) contains a loop that maintains a node set *S* which, we will show later, is ancestral, *i.e., i* ∈ *S* \Rightarrow ans $g(i) \subseteq S$. In each round the algorithm finds a new node *i* ∉ *S* such that ans $_G(i) \subseteq S$, and a subroutine Identify-Parents [\(Algorithm 2\)](#page-7-2) is used to find all parents of *i*. After that, we append *i* into *S* and continue until *S* contains all nodes in G. Finally, the rows of the mixing matrix H is obtained by intersections of properly-chosen row spaces of M_k .

Both [Algorithm 2](#page-7-2) and [Algorithm 3](#page-8-0) include a crucial step, which we call it *orthogonal projection*, as described in [Algorithm 1.](#page-7-3) At a high level, it helps determine the minimal RDF for *zi* after fixing the

latent variables z_S , and this exactly corresponds to the number of parents of z_i that are not in z_S . We provide a simple example in [Section E.2](#page-20-0) to illustrate why this approach works.

The following result states that [Algorithm 3](#page-8-0) can recover the ground-truth causal model up to ∼sur:

<code>Theorem 4.</code> *Suppose that* $M_k, k \in [K]$ *are perfectly identified in <code>Step 1.</code> Let* $(\hat{H},\hat{\mathcal{G}})$ *<i>be the solution returned by [Algorithm 3,](#page-8-0) then we must have* $(H, \mathcal{G}) \sim_{\text{sur}} (\hat{H}, \hat{\mathcal{G}})$ *.*

The full proof of [Theorem 4](#page-7-0) is given in [Section H.3.](#page-29-0) It crucially relies on the following two propositions that reveal how [Algorithm 3](#page-8-0) and the subroutine [Algorithm 2](#page-7-2) work.

Algorithm 1 Orthogonal-projections

```
1: Input: Ordered set S = \{s_1, s_2, \cdots, s_m\} \subseteq [d], index i \notin S, matrices M_k \in \mathbb{R}^{d \times n}, k \in [K]2: Output: Set of vectors \{p_k\}_{k=1}^K3: for k \leftarrow 1 to K do
4: W \leftarrow \text{span} \langle (\boldsymbol{M}_k)_s : s \in S \rangle\triangleright (\boldsymbol{M}_k)_s is the s-th row of \boldsymbol{M}_k5: p_k \leftarrow \text{proj}_{W^{\perp}} ((M_k)_i)6: end for
```
Proposition 1. *The following two propositions hold for [Algorithm 3:](#page-8-0)*

- ans $_G(i) \subseteq S \Leftrightarrow$ the *if condition in line 8 of [Algorithm 3](#page-8-0) is fulfilled;*
- the set S maintained in [Algorithm 3](#page-8-0) is always an ancestral set, in the sense that $j \in S \Rightarrow \text{ans}_G(j) \subseteq S$.

 ${\bf Proposition~2.~}$ *Given any ordered ancestral set S that contains* $\mathop{\rm pa}\nolimits_{\mathcal{G}}(i)$ *for some* $i \notin S$ *, [Algorithm 2](#page-7-2) returns a* $\mathit{set\ } P_i \subseteq S\ \mathit{that\ is\ exactly}\ pa_{\mathcal{G}}(i).$

Algorithm 2 Identify-Parents

```
1: Input: An ordered set S = \{s_1, s_2, \dots, s_m\} \subseteq [d], a node i \notin S and matrices M_k, k \in [K]2: Output: The parent set Pi of node i
 3: P_i \leftarrow \emptyset4: for m′ ← 0 to m do
 \{\boldsymbol{p}_k\}_{k=1}^K \leftarrow \texttt{Orthogonal-projections}{s_j : j \le m'}, i, {M_k}_{k \in [K]}6: r_{m'} \leftarrow \dim \text{span} \langle p_k : k \in [K] \rangle7: if m' \ge 1 and r_{m'} = r_{m'-1} - 1 then<br>8: P_i \leftarrow P_i \cup \{m'\}P_i \leftarrow P_i \cup \{m'\}9: end if
10: end for
```
6. Experiments

In this section, we present our experimental setup and results for LiNGCReL. Note that LiNGCReL as described in the previous section only works in the population regime. When the number of samples is limited, two main challenges in implementing LiNGCReL are to accurately compute the dimension of a subspace (line 6 of [Algorithm 2](#page-7-2) and line 8 of [Algorithm 3\)](#page-8-0), and to find a vector in the intersection of multiple subspaces (line 20, [Algorithm 3\)](#page-8-0). Due to space limit, the implementation details are described in [Section B.2.](#page-16-0)

Experimental setup. We generate the independent noise variables from generalized Gaussian distributions $p_{\beta}(x) \propto \exp(-|x|^{\beta})$ with parameters $\beta_k = 0.2k^2, k = 1, 2, \cdots, d$, multiplied by

Algorithm 3 Learn-Causal-Model

1: **Input:** Matrices $M_k, k \in [K]$ 2: **Output:** The edge set $\mathcal E$ on the vertex set $[d]$ and the mixing matrix $\hat H$
3: $S \leftarrow \emptyset$: 3: *S* ← ∅; ▷ *S* is an ordered set 4: $\mathcal{E} \leftarrow \emptyset$ 5: **while** $|S| \le d$ **do**
6: **for** $i \notin S$ **do** 6: **for** $i \notin S$ **do**
7: $\{n_i\}_{i=1}^K$ $\{p_k\}_{k=1}^K \leftarrow \texttt{Orthogonal-projections}$ *S*, *i*, ${M_k}_{k \in [K]}$ 8: **if** dim span $\langle q_k : k \in [K] \rangle = 1$ **then** 9: **break break break b break b b b b b b b we will show that such an** *i* **must exist** 10[.] **end if** 10: **end if** 11: **end for** 12: $P_i \leftarrow \text{Identity-Parents}(S, i)$
13: $S \leftarrow S \cup \{i\}$ 13: $S \leftarrow S \cup \{i\}$
14: $\mathcal{E} \leftarrow \mathcal{E} \cup \{i\}$ 14: $\mathcal{E} \leftarrow \mathcal{E} \cup \{(j, i) : j \in P_i\}$ 15: **end while** 16: **for** *i* = 1 to *d* **do** 17: $E_i \leftarrow \text{span} \langle (\boldsymbol{M}_k)_i : k \in [K] \rangle$ 18: **end for** 19: **for** *i* = 1 to *d* **do** 20: $\quad \hat{\textbf{\textit{h}}}_{i} \leftarrow \text{any non-zero vector in} \left(\cap_{j: (i,j) \in \mathcal{E}} E_{j} \right) \cap E_{i}$ 21: **end for** 22: $\hat{H} \leftarrow \begin{bmatrix} \hat{\textbf{\textit{h}}}^{\top} \ \hat{\textbf{\textit{h}}}^{\prime\top} \end{bmatrix}$ $\tilde{\mathbf{h}}_1^{\top}$, $\hat{\mathbf{h}}_2^{\top}$ $\mathbf{z}_2^{\top}, \cdots, \mathbf{\hat{h}}_d^{\top}$ *d* i⊤

normalization constants to make their variances equal to 1. The ground-truth causal graph is generated by first fixing a total order of the vertices, say 1, 2, \dots , *d*, then add directed edges $i \rightarrow j(i < j)$ according to i.i.d. Bernoulli(*p*) distributions, where $p \in (0, 1)$. The non-zero entries of matrices B_k and *H* are all generated independently from Gaussian distributions. For simplicity, we focus on the case *n* = *d* since recovery of the latent graphs only requires information from *d* components of *x*.

Metrics of estimation error. Since causal representation learning seeks to learn both the causal graphs and the latent variables, for each output of our algorithm we first check if it exactly recovers the ground-truth causal graph. Then, recall that the latent variables and the observations are related by $z = Hx$, given any output unmixing matrix \hat{H} from [Algorithm 3,](#page-8-0) we define the relative estimation error ∆*ⁱ* for *zi* as the solution of the following optimization problem:

$$
\min ||\Delta||_{\infty} \quad s.t. \Delta_i = \frac{\left\| \text{proj}_{\text{span}\langle h_j : j \in \text{sur}_{\mathcal{G}}(i) \rangle}(\hat{h}_i) \right\|_2}{\left\| \hat{h}_i \right\|_2},\tag{5}
$$

 \hat{H} = \hat{PH} for some signed permutation matrix *P*.

where signed permutation is allowed here since the noise distribution in our experiments is symmetric and the order of latent variables z_i , $i = 1, 2, \cdots, d$ does not matter. We refer to the errors Δ_i defined in [\(5\)](#page-8-1) as the *SNA error*. The SNA error measures how much of the row $\hat{\hat{\boldsymbol{h}}}_i$ that we learn is contained in the span of the ground-truth rows $h_j, j \in \overline{\text{sur}_\mathcal{G}}(i).$ Indeed, recall that given any observation $x,$ the ground-truth latent variable is $z = Hx$ while our algorithm outputs $\hat{v}_i = \hat{h}_i^\top x$, so the SNA error essentially captures whether the recovered latent variable is close to some linear mixture of latent variables in the effect-dominating set of *i*. When the SNA error is zero for some node *i*, we know that the recovered latent variable at node *i* is exactly a linear mixture of the ground-truth latent variables in $\overline{\text{sur}_G}(i)$, according to [Lemma 1.](#page-4-2)

Figure 2. Left: plots of SNA Error and graph recovery accuracy achieved by LiNGCReL as functions of sample size (per environment) for different choices of graph size *d* and number of environments *K*. Right: an example of causal graph generated in our experiments, and the estimation error of LiNGCReL for each node.

We also define the *true error* for estimating each latent variable. Formally, let *H* be the unmixing matrix that corresponds to the solution of [\(5\),](#page-8-1) then we define the true estimation error $\tilde{\Delta}_i$ of \boldsymbol{z}_i as

$$
\tilde{\Delta}_i = \left\| \left(\boldsymbol{I} - \boldsymbol{h}_i \boldsymbol{h}_i^\top \right) \hat{\boldsymbol{h}}_i \right\|_2.
$$
 (6)

Results. We randomly sample 100 causal models with size *d* = 5, 30 causal models with size *d* = 8 ad 30 causal models of size *d* = 10. In light of [Theorem 1,](#page-5-0) for each $d \in \{5, 8, 10\}$, we sample data from $K = d$ randomly chosen environments; for $d = 5$ we also consider $K = 20$ to study how different choices of *K* can affect the result. We run LiNGCReL for each model with different sample sizes, compute the SNA error and true error of the obtained solution from [\(5\)](#page-8-1) and [\(6\)](#page-9-1) respectively for each latent variable, and check whether the ground-truth causal graph is exactly recovered.

[Figure 2](#page-9-0) shows how the average SNA error (over all latent variables) and the accuracy of graph recovery changes when sample size grows. We can see LiNGCReL successfully recovers about 80% of all models within each category, and the median of the average SNA error is smaller than 1%. Moreover, by comparing [Figure 2a](#page-9-0) with [Figure 2b,](#page-9-0) one can observe that if we fix the total number of samples but choose a larger *K* (*i.e.*, fewer samples per environment), LiNGCReL can still achieve the same level of performance compared with the choice $K = d$. Intuitively, this is because $K \gg d$ vectors sampled from an $r(r \leq d)$ dimensional subspace are unlikely to approximately lie in an (*r* – 1)-dimensional subspace, so that the calculation of line 6 of [Algorithm 2](#page-7-2) and line 8 of [Algorithm 3](#page-8-0) can be more accurate. We leave a better and quantitative understanding of the trade-off between *d* and *K* to future work.

SNA error v.s. true error. To understand the implication of our theory, we dive deeper by looking into the learning outcome of LiNGCReL on a specific model, of which the causal graph is shown in [Figure 2e.](#page-9-0)

In [Figure 2f,](#page-9-0) we list the surrounding set of each node and the corresponding SNA error and true error. We can see that if sur $G(i) = \emptyset$, the two errors equal and both are small, but if sur $G(i) \neq \emptyset$, the true error is much larger than the SNA error. This indicates that LiNGCReL indeed learns the ground-truth model up to \sim_{sur} , as [Theorem 1](#page-5-0) predicts.

7. Identification limit of general causal models with soft interventions

While [Theorem 1](#page-5-0) guarantees identifiability with general environments, it only applies to linear causal models. In this section, we show that if we have access to single-node soft interventions, then we can identify general non-parametric causal models up to $\sim_{\rm sur}$. To obtain our identifiability result, we also require that the environments are non-degenerate in the following sense:

 $\bf{Definition 5 (Non-degeneracy set of interventions).}$ *Let* $\hat{p}_k\left(z_i\mid z_{\text{pa}_\mathcal{G}(i)}\right), k \in [K_i]$ *be conditional* probability densities at node i, then $\{ \hat{p}_k \}_{k=1}^{K_i}$ is said to be non-degenerate on node i at point $\hat{\boldsymbol{z}} \in \mathbb{R}^d$ if all *these conditional densities are well-defined and positive at z*ˆ*, and the matrix*

$$
\left[\frac{\partial (\hat{p}_1/\hat{p}_k)}{\partial z_j}\right]_{2\leq k\leq K_i, j\in \bar{\text{pa}}_G(i)}\bigg|_{z=\hat{z}} \in \mathbb{R}^{(K_i-1)\times \left(\left|p_1\right(g(i)\right|+1\right)}
$$

has full row rank. Moreover, we say that $\{\hat{p}_k\}_{k=1}^{K_i}$ *is non-degenerate in a point set* O *if for all* $\hat{\bm{z}} \in O$ *, it is non-degenrate at z*ˆ*.*

The following lemma shows how [Definition 5](#page-10-2) is related to [Assumption 5](#page-5-1) in the linear setting:

 $\bf{Lemma~2.~}$ *Suppose that* $\hat{p}_k(z) = \prod_{i=1}^d \hat{p}_k\left(z_i\ |\ z_{\text{pa}_G(i)}\right), k\in[K]$ *be probability distributions of latent* variables z generated from the linear causal models [\(3\)](#page-4-0), such that for $\forall i\in[d]$, $\hat{p}_k\left(z_i\mid z_{\text{pa}_\mathcal{G}(i)}\right)$, $k\in[K]$ are *non-degenerate on node i in the sense of [Definition 5.](#page-10-2) Then the corresponding matrices B^k* , *k* ∈ [*K*] *satisfy [Assumption 5.](#page-5-1)*

Now we are ready to state our main result in this section:

Theorem 5. *Suppose that we have access to observations generated from multiple environments* $\{P^E_X\}_{E\in\mathfrak{E}}$ *.* Let $\hat{(\hat{\textbf{h}},\hat{\mathcal{G}})}$ be any candidate solution with data generated according to [Assumption 1](#page-2-1) with latent variables $\bm{v} = \hat{\bm{h}}(\bm{x})$ and joint distribution q_E with factors q_i^E . Assuming that

- (i) the joint densities $\{p_E(\bm{z})\}_{E\in\mathfrak{E}}$ are continuous differentiable on \mathbb{R}^d with common support $\mathrm{O}_{\bm{z}},$ and $\left\{ q_E(\nu) \right\}_{E \in \mathfrak{E}}$ are continuous differentiable on \mathbb{R}^d with common support \mathbf{O}_ν ;
- *(ii) we have access to multiple single-node soft interventions on each node with unknown targets: there exists a partition* $\mathfrak{E} = \bigcup_{i=1}^d \mathfrak{E}_i$ such that $\mathcal{I}^{\mathfrak{E}_i}_{\mathcal{Z}} = \{\pi(i)\}, \mathcal{I}^{\mathfrak{E}_i}_{\mathcal{V}} = \{\pi'(i)\}, \forall i \in [d]$ for some unknown permutations π *and* π ′ *on* [*d*]*;*
- *(iii) the intervention distributions on each node are non-degenerate in the sense of [Definition 5:](#page-10-2) there exists* $N_z \subseteq O_z$ *and* $N_v \subseteq O_v$ *satisfying* $N_z^{\rm o}$ = $N_v^{\rm o}$ = \emptyset *where S* $^{\rm o}$ *denotes the interior of a set S, such that* f or all $i\in [d]$, $\left\{p_i^E(\cdot): E\in \mathfrak{E}_{\pi^{-1}(i)}\right\}$ (resp. $\left\{q_i^E(\cdot): E\in \mathfrak{E}_{\pi'^{-1}(i)}\right\}$) is non-degenerate on node i in $O_z \setminus N_z$ (resp. $O_v \setminus N_v$).

Then we must have $(h, \mathcal{G}) \sim_{\text{sur}} (\hat{h}, \hat{\mathcal{G}})$ *.*

Previous works on the identifiability of non-parametric causal models typically require that all the joint distributions are supported on the whole space \mathbb{R}^d (Kügelgen et al. [2023;](#page-12-14) Liang et al. 2023; Varıcı et al. [2023\)](#page-13-8). In contrast, we only assume that the densities have common and unknown support across all interventions.

[Theorem 5](#page-10-0) can be regarded as a soft-intervention version of Kügelgen et al. [2023,](#page-12-7) Theorem 4.3, which assumes access to hard interventions and only need two paired interventions per node. While they are able to show full identifiability, we show in the following that identifiability up to \sim_{sur} is the best we can hope for with soft interventions.

Theorem 6 (Counterpart to [Theorem 5,](#page-10-0) informal version of [Theorem 10\)](#page-39-0)**.** *For any causal model* (h, \mathcal{G}) and any set of environments \mathfrak{E} = $\big\{E_k : k \in [K]\big\}$ such that all conditions in [Theorem 5](#page-10-0) are satisfied, *there must exists a candidate solution* (ˆ*h*, G) *and a hypothetical data generating process that satisfy the same set of conditions, but*

$$
\frac{\partial v_i}{\partial z_j} \neq 0, \quad \forall j \in \overline{\text{sur}}_{\mathcal{G}}(i).
$$

Finally, the ambiguity still exists if we additionally assume standard axioms such as causal minimality [\(Assumption 6\)](#page-18-2) and faithfulness [\(Assumption 7\)](#page-18-3) on the causal model.

8. Conclusions

This paper studies the limit of learning identifiable causal representations using data from multiple environments. When hard interventions are not available, we provide theory and algorithm for identification up to SNA, and also show that SNA is an intrinsic ambiguity in our setting.

It is interesting to further investigate the setting where we do not assume that the causal model is linear. Moreover, it is important to understand the concrete form of available interventions in real-world applications. For instance, it is suggested that for single-cell genomics, the intervention is sometimes a "mixture" of hard and soft interventions, and sometimes can even reverse the direction of an edge (Tejada-Lapuerta et al. [2023\)](#page-13-9). Modelling such more complicated interventions appears to be crucial to reveal the underlying causal mechanisms in real-world problems.

9. Broader Impact

This paper presents work whose goal is to advance the field of Machine Learning and in particular the sub-field Causal Representation Learning. There are many potential societal consequences of our work, especially as it pertains to building more reliable machine learning models, none which we feel must be specifically highlighted here.

References

- Ahuja, Kartik, Divyat Mahajan, Yixin Wang, and Yoshua Bengio. 2023. Interventional causal representation learning. In *International conference on machine learning,* 372–407. PMLR.
- Ahuja, Kartik, Amin Mansouri, and Yixin Wang. 2023. Multi-domain causal representation learning via weak distributional invariances. *arXiv preprint arXiv:2310.02854.*
- Akhtar, Naveed, and Ajmal Mian. 2018. Threat of adversarial attacks on deep learning in computer vision: a survey. *Ieee Access* 6:14410–14430.
- Bengio, Yoshua, Aaron Courville, and Pascal Vincent. 2013. Representation learning: a review and new perspectives. *IEEE transactions on pattern analysis and machine intelligence* 35 (8): 1798–1828.
- Brehmer, Johann, Pim De Haan, Phillip Lippe, and Taco S Cohen. 2022. Weakly supervised causal representation learning. *Advances in Neural Information Processing Systems* 35:38319–38331.
- Bubeck, Sébastien, Varun Chandrasekaran, Ronen Eldan, Johannes Gehrke, Eric Horvitz, Ece Kamar, Peter Lee, Yin Tat Lee, Yuanzhi Li, Scott Lundberg, et al. 2023. Sparks of artificial general intelligence: early experiments with gpt-4. *arXiv preprint arXiv:2303.12712.*
- Buchholz, Simon, Goutham Rajendran, Elan Rosenfeld, Bryon Aragam, Bernhard Schölkopf, and Pradeep Ravikumar. 2023. Learning linear causal representations from interventions under general nonlinear mixing. *arXiv preprint arXiv:2306.02235.*
- Campbell, John. 2007. An interventionist approach to causation in psychology. *Causal learning: Psychology, philosophy, and computation,* 58–66.

Comon, Pierre. 1994. Independent component analysis, a new concept? *Signal processing* 36 (3): 287–314.

- Cooper, Gregory F, and Changwon Yoo. 1999. Causal discovery from a mixture of experimental and observational data. In *Proceedings of the fifteenth conference on uncertainty in artificial intelligence,* 116–125.
- Dunbar, Kevin N, and Jonathan A Fugelsang. 2004. Causal thinking in science: how scientists and students interpret the unexpected. In *Scientific and technological thinking,* 57–79. Psychology Press.
- Eberhardt, Frederick. 2008. Almost optimal intervention sets for causal discovery. In *Proceedings of the twenty-fourth conference on uncertainty in artificial intelligence,* 161–168.

. 2014. Direct causes and the trouble with soft interventions. *Erkenntnis* 79:755–777.

- Eronen, Markus I. 2020. Causal discovery and the problem of psychological interventions. *New Ideas in Psychology* 59:100785.
- Fisher, Ronald Aylmer, et al. 1960. The design of experiments. *The design of experiments.,* no. 7th Ed.
- Geirhos, Robert, Jörn-Henrik Jacobsen, Claudio Michaelis, Richard Zemel, Wieland Brendel, Matthias Bethge, and Felix A Wichmann. 2020. Shortcut learning in deep neural networks. *Nature Machine Intelligence* 2 (11): 665–673.
- Hauser, Alain, and Peter Bühlmann. 2014. Two optimal strategies for active learning of causal models from interventional data. *International Journal of Approximate Reasoning* 55 (4): 926–939.
- Holyoak, Keith J, and Patricia W Cheng. 2011. Causal learning and inference as a rational process: the new synthesis. *Annual review of psychology* 62:135–163.
- Hyttinen, Antti, Frederick Eberhardt, and Patrik O Hoyer. 2013. Experiment selection for causal discovery. *Journal of Machine Learning Research* 14:3041–3071.
- Khemakhem, Ilyes, Diederik Kingma, Ricardo Monti, and Aapo Hyvarinen. 2020. Variational autoencoders and nonlinear ica: a unifying framework. In *International conference on artificial intelligence and statistics,* 2207–2217. PMLR.
- Kıcıman, Emre, Robert Ness, Amit Sharma, and Chenhao Tan. 2023. Causal reasoning and large language models: opening a new frontier for causality. *arXiv preprint arXiv:2305.00050.*
- Koh, Pang Wei, Shiori Sagawa, Henrik Marklund, Sang Michael Xie, Marvin Zhang, Akshay Balsubramani, Weihua Hu, Michihiro Yasunaga, Richard Lanas Phillips, Irena Gao, et al. 2021. Wilds: a benchmark of in-the-wild distribution shifts. In *International conference on machine learning,* 5637–5664. PMLR.
- Kügelgen, Julius von, Michel Besserve, Wendong Liang, Luigi Gresele, Armin Kekić, Elias Bareinboim, David M Blei, and Bernhard Schölkopf. 2023. Nonparametric identifiability of causal representations from unknown interventions. *arXiv preprint arXiv:2306.00542.*
- Liang, Wendong, Armin Kekić, Julius von Kügelgen, Simon Buchholz, Michel Besserve, Luigi Gresele, and Bernhard Schölkopf. 2023. Causal component analysis. *arXiv preprint arXiv:2305.17225.*
- Lippe, Phillip, Sara Magliacane, Sindy Löwe, Yuki M Asano, Taco Cohen, and Efstratios Gavves. 2023. Biscuit: causal representation learning from binary interactions. *arXiv preprint arXiv:2306.09643.*
- Locatello, Francesco, Stefan Bauer, Mario Lucic, Gunnar Raetsch, Sylvain Gelly, Bernhard Schölkopf, and Olivier Bachem. 2019. Challenging common assumptions in the unsupervised learning of disentangled representations. In *International conference on machine learning,* 4114–4124. PMLR.
- Locatello, Francesco, Stefan Bauer, Mario Lucic, Gunnar Rätsch, Sylvain Gelly, Bernhard Schölkopf, and Olivier Bachem. 2020. A sober look at the unsupervised learning of disentangled representations and their evaluation. *The Journal of Machine Learning Research* 21 (1): 8629–8690.
- Locatello, Francesco, Ben Poole, Gunnar Rätsch, Bernhard Schölkopf, Olivier Bachem, and Michael Tschannen. 2020. Weakly-supervised disentanglement without compromises. In *International conference on machine learning,* 6348–6359. PMLR.
- Lopez, Romain, Natasa Tagasovska, Stephen Ra, Kyunghyun Cho, Jonathan Pritchard, and Aviv Regev. 2023. Learning causal representations of single cells via sparse mechanism shift modeling. In *Conference on causal learning and reasoning,* 662–691. PMLR.
- Lu, Chaochao, Yuhuai Wu, José Miguel Hernández-Lobato, and Bernhard Schölkopf. 2021. Invariant causal representation learning for out-of-distribution generalization. In *International conference on learning representations.*
- Ovadia, Yaniv, Emily Fertig, Jie Ren, Zachary Nado, David Sculley, Sebastian Nowozin, Joshua Dillon, Balaji Lakshminarayanan, and Jasper Snoek. 2019. Can you trust your model's uncertainty? evaluating predictive uncertainty under dataset shift. *Advances in neural information processing systems* 32.
- Pearl, Judea. 2009. *Causality.* Cambridge university press.
- Prystawski, Ben, and Noah D Goodman. 2023. Why think step-by-step? reasoning emerges from the locality of experience. *arXiv preprint arXiv:2304.03843.*
- Roeder, Geoffrey, Luke Metz, and Durk Kingma. 2021. On linear identifiability of learned representations. In *International conference on machine learning,* 9030–9039. PMLR.
- Rudelson, Mark, and Roman Vershynin. 2009. Smallest singular value of a random rectangular matrix. *Communications on Pure and Applied Mathematics: A Journal Issued by the Courant Institute of Mathematical Sciences* 62 (12): 1707–1739.
- Schölkopf, Bernhard, Francesco Locatello, Stefan Bauer, Nan Rosemary Ke, Nal Kalchbrenner, Anirudh Goyal, and Yoshua Bengio. 2021. Toward causal representation learning. *Proceedings of the IEEE* 109 (5): 612–634.
- Schwartz, J. 1954. The formula for change in variables in a multiple integral. *The American Mathematical Monthly* 61 (2): 81–85.
- Seigal, Anna, Chandler Squires, and Caroline Uhler. 2022. Linear causal disentanglement via interventions. *arXiv preprint arXiv:2211.16467.*
- Shanks, David R, and Anthony Dickinson. 1988. Associative accounts of causality judgment. In *Psychology of learning and motivation,* 21:229–261. Elsevier.
- Shimizu, Shohei, Patrik O Hoyer, Aapo Hyvärinen, Antti Kerminen, and Michael Jordan. 2006. A linear non-gaussian acyclic model for causal discovery. *Journal of Machine Learning Research* 7 (10).
- Silva, Ricardo, Richard Scheines, Clark Glymour, Peter Spirtes, and David Maxwell Chickering. 2006. Learning the structure of linear latent variable models. *Journal of Machine Learning Research* 7 (2).
- Silver, David, Aja Huang, Chris J Maddison, Arthur Guez, Laurent Sifre, George Van Den Driessche, Julian Schrittwieser, Ioannis Antonoglou, Veda Panneershelvam, Marc Lanctot, et al. 2016. Mastering the game of go with deep neural networks and tree search. *nature* 529 (7587): 484–489.
- Spirtes, Peter, Clark N Glymour, and Richard Scheines. 2000. *Causation, prediction, and search.*
- Strevens, Michael. 2007. *Review of woodward," making things happen".*
- Tejada-Lapuerta, Alejandro, Paul Bertin, Stefan Bauer, Hananeh Aliee, Yoshua Bengio, and Fabian J Theis. 2023. Causal machine learning for single-cell genomics. *arXiv preprint arXiv:2310.14935.*
- Tillman, Robert E, and Frederick Eberhardt. 2014. Learning causal structure from multiple datasets with similar variable sets. *Behaviormetrika* 41 (1): 41–64.
- Tong, Simon, and Daphne Koller. 2001. Active learning for structure in bayesian networks. In *International joint conference on artificial intelligence,* 17:863–869. 1. Citeseer.
- Varici, Burak, Emre Acarturk, Karthikeyan Shanmugam, Abhishek Kumar, and Ali Tajer. 2023. Score-based causal representation learning with interventions. *arXiv preprint arXiv:2301.08230.*
- Varıcı, Burak, Emre Acartürk, Karthikeyan Shanmugam, and Ali Tajer. 2023. General identifiability and achievability for causal representation learning. *arXiv preprint arXiv:2310.15450.*
- Von Kügelgen, Julius, Yash Sharma, Luigi Gresele, Wieland Brendel, Bernhard Schölkopf, Michel Besserve, and Francesco Locatello. 2021. Self-supervised learning with data augmentations provably isolates content from style. *Advances in neural information processing systems* 34:16451–16467.
- Wang, Tony Tong, Adam Gleave, Tom Tseng, Kellin Pelrine, Nora Belrose, Joseph Miller, Michael D Dennis, Yawen Duan, Viktor Pogrebniak, Sergey Levine, et al. 2023. Adversarial policies beat superhuman go ais.
- Woodward, James. 2005. *Making things happen: a theory of causal explanation.* Oxford university press.
- Zhang, Hanlin, Yi-Fan Zhang, Weiyang Liu, Adrian Weller, Bernhard Schölkopf, and Eric P Xing. 2022. Towards principled disentanglement for domain generalization. In *Proceedings of the ieee/cvf conference on computer vision and pattern recognition,* 8024–8034.
- Zhang, Jiaqi, Chandler Squires, Kristjan Greenewald, Akash Srivastava, Karthikeyan Shanmugam, and Caroline Uhler. 2023. Identifiability guarantees for causal disentanglement from soft interventions. *arXiv preprint arXiv:2307.06250.*

Appendices

A. Related works

The interventionist approach to causation For the problem of causal graph discovery, it is well-known that the underlying causal structure is non-identifiable given only "passively observed" (equivalently, *i.i.d.*) data alone. As a result, randomized controlled experiments (Fisher et al. [1960\)](#page-12-16) is often used to infer causality. These experiments typically take the form of interventions (Spirtes, Glymour, and Scheines [2000;](#page-13-14) Pearl [2009\)](#page-13-15), *i.e.*, manipulations on the "natural state" of the system of interest. Early works (Woodward [2005;](#page-13-16) [Strevens](#page-13-17) [2007\)](#page-13-17) define the "hard" (also called "surgical" or "arrow-breaking") interventions in which the value of the intervened variable is entirely determined by the experimenter, thereby removing the dependence of this variable on its direct causes. This type of intervention is arguably the most natural one to consider, and following this definition, a line of works explore sufficient conditions for designing experiments that guarantee identifiability of the causal model in various settings (Cooper and Yoo [1999;](#page-12-17) Tong and Koller [2001;](#page-13-18) Eberhardt [2008;](#page-12-18) Hyttinen, Eberhardt, and Hoyer [2013;](#page-12-19) Hauser and Bühlmann [2014\)](#page-12-20).

Intervention *v.s.* **passive observation** While extensive works demonstrate the success of the interventionist approach, it faces several key challenges that significantly limit its applicability. First, Eberhardt [\(2014\)](#page-12-10) finds that in the presence of unobserved variables, certain causal structures are indistinguishable if we only perform hard interventions. This issue can be resolved by performing soft interventions *i.e.*, interventions that do not remove the dependency on direct causes but only changes the conditional distribution. Second, as pointed out in (Tillman and Eberhardt [2014\)](#page-13-19), interventions — whether hard or soft — are often expensive or even infeasible to perform in practice. For example, a psychological intervention is likely to affect multiple psychological variables simultaneously Eronen [\(2020\)](#page-12-11). As a result, (Tillman and Eberhardt [2014\)](#page-13-19) returns to the "passive observation" setting but with multiple datasets with overlapping latent variables.

Interventional causal representation learning Motivated by the interventionist literature in causal graph discovery, a recent line of works (Ahuja et al. [2023;](#page-11-4) Seigal, Squires, and Uhler [2022;](#page-13-10) Varici et al. [2023;](#page-13-0) Kügelgen et al. [2023;](#page-12-7) Buchholz et al. [2023;](#page-11-5) J. Zhang et al. [2023;](#page-13-13) Varıcı et al. [2023\)](#page-13-8) consider performing interventions to resolve the non-identifiability issue in causal representation learning (Locatello et al. [2019\)](#page-12-4). Roughly speaking, these result indicate that identification (possibly with some ambiguities) is possible if one can perform intervention on every latent variable. However, it is unclear how to perform such interventions in practice, given that the underlying latent variables are unknown. Khemakhem et al. [\(2020\)](#page-12-12), Lu et al. [\(2021\)](#page-12-13), and Roeder, Metz, and Kingma [\(2021\)](#page-13-20) do not require single-node interventions to achieve identifiability, but assumes that the joint distribution of latent variables in each environment lie in a certain exponential family. This assumption can be understood as a prior on the latent variables, but it is unclear when or why it is reasonable to make in reality. Recently, Ahuja, Mansouri, and Wang [\(2023\)](#page-11-7) considers learning causal representations from multiple domains that relate to each other via an invariance constraint on the subset S of *stable* latent variables, and they prove identification up to affine mixtures within S.

Causal reasoning capacity of LLMs In view of the tremendous success of large language models (LLMs), several works aim to understand the causal reasoning ability of LLMs. Kıcıman et al. [\(2023\)](#page-12-21) conducts an an extensive experimental study and finds that LLMs outperforms all existing causal discovery methods on multiple datasets, but also have simple and mysterious failure modes. Prystawski and Goodman [\(2023\)](#page-13-21) provide theoretical evidence that the chain-of-thought (CoT) prompt allows LLMs to reduce their uncertainty when answering questions related to causal variables that are far apart.

B. Experiment details for [Section 6](#page-7-1)

B.1 Details for step 1 in [Section 5](#page-6-1)

Since $\epsilon = B_k z = B_k H x$ in the *k*-th environment, so we can use any identification algorithm for linear ICA to recover the matrix \pmb{M}_k . Note that while standard linear ICA algorithms only apply to the case where $n = d$, for $n > d$ we can arbitrarily choose *d* principal components of x to reduce it to the *n* = *d* case. This is without loss of generality, since when *n* > *d* there is redundant information in *x*.

After recovering \boldsymbol{M}_k for each k by running linear ICA, we still do not know whether each $\boldsymbol{M}_k \boldsymbol{x}$ corresponds to the same permutation of the ground-truth noise variables ϵ. To resolve this issue, we choose test function Ψ mapping any distribution on $\mathbb R$ to a deterministic real value, which we expect to take different values for different ϵ_i 's. We choose $\Psi(\mathbb{P}) = \mathbb{P}[|X| \leq 1]$ in our experiments. For all *k* ⩾ 2, we calculate the Ψ value of each component of the *d*-dimensional empirical distribution $\hat{\mathbb{P}}_k = \frac{1}{N} \sum_{i=1}^N \mathbb{1}_{\mathbf{M}_k \mathbf{x}_i^{(k)}}$, and choose a permutation π_k to rearrange them in increasing order. Then, we rearrange the columns of \pmb{M}_k using the same permutation $\pi_k.$ This procedure would asymptotically produce correct alignments as long as $\Psi(\epsilon_i), i\in [d]$ are different, and we find that it empirically works well.

Alternatively, this alignment step can be done as follows: for each pair of environments (*E*1, *Et*), and for each pair of nodes (i, j) , we calculate the distribution distance between ϵ_i in environment E_1 and ϵ_j in environment E_t , based on some notion of distribution distance (*e.g.* kernel maximum mean discrepancy). Then we find the min-cost perfect matching, where the cost of an edge is the distribution distance.

B.2 Details for the implementation of LiNGCReL **in the finite-sample regime**

Although LiNGCReL provably works in the population regime, it faces several challenges when there is only a finite number of samples:

- First, since rank is not a continuous function, it is sensitive to finite-sample estimation errors. In our implementation of [Algorithm 3,](#page-8-0) in each iteration we instead choose $i \notin S$ that has the largest ratio between the first and second singular values of $[q_1, q_2, \cdots, q_K]$. And in line 6 of [Algorithm 2,](#page-7-2) we introduce a hyper-parameter tl such that the matrix $[q_1, q_2, \cdots, q_K]$ is considered to have rank $r_{m'-1}$ if its $r_{m'}$ -th singular value is smaller than t 1. Since the smallest singular value of a random matrix $A \in \mathbb{R}^{K \times m}$ ($K \ge m$) is at the order of \sqrt{K} – √ *m* – 1 with high probability (Rudelson and Vershynin [2009\)](#page-13-22), when *K* = *d* one shall choose tl \sim √ *d* – √ $\overline{d-1} = \mathcal{O}\left(\frac{1}{\sqrt{2}}\right)$ *d* $\big)$. On the other hand, for larger *K* we can correspondingly choose a larger t1. Note that a small t1 potentially has the risk of being dominated the noise in the estimation, which means that we need more samples per environment to reduce the noise. In contrast, for larger tl the estimation is more robust to noise and we can use fewer samples.
- Second, finite-sample estimation errors of M_k make it harder to obtain h_i in [Algorithm 3](#page-8-0) of [Algorithm 3.](#page-8-0) We implement this step in the following way: first let *Q^j* be the orthogonal projection matrix onto E_j^\perp *i.e.*, $\mathbf{Q}_j^\top\boldsymbol{x}$ = $\text{proj}_{E_j^\perp}(\boldsymbol{x})$, then choose h_i to be the singular vector of P *^j*:(*j*,*i*)∈E or *^j*=*ⁱ ^Q*[⊤] *^j Q^j* that corresponds to the smallest singular value (including zero). Indeed, in the noiseless case we would have $\left(\sum_{j:(j,i)\in\mathcal{E}}\,_{\text{or}\,j=i}\mathbf{Q}_j^\top\mathbf{Q}_j\right)\pmb{h}_i$ = 0 if and only if $\pmb{h}_i\in\left(\cap_{j:(i,j)\in\mathcal{E}}E_j\right)\cap$ *Ei* .

C. Further experiment results

SNA error v.s. true error We plot the SNA error v.s. true error achieved by LiNGCReL in [Figure 3.](#page-17-0) We observe that

- For most nodes, SNA error is exactly equal to the true error and both errors are small, indicating that the corresponding latent variables have been successfully learned by LiNGCReL.
- The remaining nodes typically have true error much larger than SNA error. This indicates that there exists some ambiguities at these nodes in the sense that sur $_G(i) \neq \emptyset$. Note that the true error for many nodes are close to 1; one possible reason is that one selects the wrong singular vector in the second part of [Section B.2,](#page-16-0) so that it is orthogonal to the ground-truth vector.

Sensitivity of LiNGCReL **to the hyperparameter** tl We examine how different choices of tl would affect the performance of LiNGCReL. Specifically, we run LiNGCReL on the 100 models with size *d* = 5 and number of environments $K = 5$ sampled in [Section 6](#page-7-1) with $\texttt{tl} \in \{0.1, 0.15, 0.2, 0.25, 0.3\}$ and the results are reported in [Figure 4.](#page-17-1) We can see that the permance is actually quite sensitive to tl.

Figure 3. Comparing SNA error with true error for the 500 latent variables in the 100 graphs of size *d* = 5 that we sample in [Section 6.](#page-7-1)

Figure 4. Performance of LiNGCReL as a function of tl. tl= 0.15 achieves the best performance in terms of both SNA error and graph recovery accuracy.

D. Background on causal representation learning

It is common to assume some axioms on what kind of (conditional) dependency information is encoded in a causal graph (see Spirtes, Glymour, and Scheines [2000,](#page-13-14) Section 3.4 for a detailed discussion). The most natural one is the Causal Markov Condition introduced in [Definition 1](#page-2-2) that gives sufficient conditions for conditional independence via *d*-separation. We introduce the formal definition of *d*-separation below:

Definition 6 (paths and colliders)**.** *Let i*, *j be two nodes of a DAG* G*, a path is a sequence of nodes* i_0 = i, i_1, \cdots, i_k = j such that there is an edge (in either direction) between i_j and i_{j+1}, j = $0, 1, \cdots, k$ – 1*. A node i_j is called a collider on this path if i* j *∈* $\mathrm{ch}_{\mathcal{G}}(i_{j-1})$ *∩* $\mathrm{ch}_{\mathcal{G}}(i_{j+1})$ *.*

Definition 7 (blocked path)**.** *A path in a DAG* G *between node i and node j is said to be blocked by a node set S if either of the following holds:*

- *there exists a node v on the path that is in S but not a collider, or*
- *there exists a node v on the path that is a collider, but none of its descendants (including itself) are in S.*

Definition 8 (d-separation). *For a DAG G with node set* [d], any two nodes $i \neq j$ are said to be d-separated *by a set* $S \subset [d] \setminus \{i, j\}$ *if all paths from i to j are blocked by S.*

The *minimality* condition states that there is no redundant edges in the causal graph, and is a natural consequence of the Occam's Razor Principle.

Assumption 6 (Causal minimality, Spirtes, Glymour, and Scheines [2000,](#page-13-14) Section 3.4.2)**.** *For latent variables z, removing any edge from* G *would render violation of the causal Markov condition [Definition 1.](#page-2-2) In other words, let* G_1 *be the graph obtained by removing any single edge from* G, then there must exist $i \in [d]$ $\mathcal{Z}_{\text{std}_{\mathcal{G}_1}(i)} \mid \mathcal{Z}_{\text{pa}_{\mathcal{G}_1}(i)}$ *.*

The *faithfulness* condition states that the Causal Markov Condition actually entails all (conditional) independence in the latent variables.

Assumption 7 (Faithfulness, Spirtes, Glymour, and Scheines [2000,](#page-13-14) Section 3.4.3)**.** *Every (conditional) independence in the latent variables z is entailed by the Causal Markov Condition applied to* G*. In other words, zⁱ* ⊥⊥ *z^j* | *z^S* ⇔ *i*, *j are d-separated by S.*

Existing works have explored different notions of identifiability. For observational data, it is well known that Markov equivalence of graphs is an intrinsic ambiguity that one cannot resolve:

Definition 9 (Markov equivalence/Faithful Indistinguishability, Spirtes, Glymour, and Scheines [2000,](#page-13-14) Section 4.2)**.** *If two DAGs encodes the same set of dependency relations, we say that they are Markov equivalent.*

Any DAG G induces a partial order on its nodes which we denote by \prec_G . In the special case when for all *i* \neq *j*, either *i* \prec G *j* or *j* \prec G *i* holds, we say that \prec G is a total order. This partial order is equivalent to the transitional closure of the graph, as defined below:

Definition 10 (Transitional closure). *Given any DAG G, its transitional closure* \bar{G} *is defined to be the graph obtained by connecting all edges i* \rightarrow *j where i is an ancestor of j in G.*

When \prec_G is a total order, each pair of nodes are connected by a directed edge in its transitive closure G. Such G is often called a *tournament* in graph theory.

In the following, we list different forms of identifiability that appear in the literature:

 $\bf{Definition~11}$ (different notions of identifiability). $\it{Let}~ {\cal H} : \mathbb{R}^n \supseteq {\cal X} \mapsto \mathbb{R}^d$ be the space of diffeomorphic m appings from observation to latent, and $\mathfrak G$ be the space of all DAGs with d nodes, then for $h, \hat h \in \mathcal H$ and $\mathcal{G}, \hat{\mathcal{G}} \in \mathfrak{G}$, we write

- (i) (Seigal, Squires, and Uhler [2022;](#page-13-10) Liang et al. [2023\)](#page-12-14) (h, G) $\frac{T}{C}$ (ĥ, Ĝ) if there exists a permutation π on $[d]$ *such that* $\pi(\mathcal{G})$ *and* $\hat{\mathcal{G}}$ *have the same transitional closure*;
- *(ii) (Kügelgen et al. [2023;](#page-12-7) Varıcı et al. [2023\)](#page-13-8)* (*h*, G) ∼CRL (ˆ*h*, Gˆ) *if we actually have* G ⁼ Gˆ *for the* ϕ *defined above.*

Given an equivalence relation \sim on $\mathcal{H} \times \mathfrak{G}$, we say that a causal model (*h*, *G*) is identifiable under ∼ if any candidate solution (*h*, Ĝ) satisfies (*h*, Ĝ) ∼ (*h*, Ĝ). The notion of identification up to *^T*∼*G*, as shown in Seigal, Squires, and Uhler [\(2022\)](#page-13-10) with single-node soft interventions on linear causal models, is highly related to this paper. Compared with their result, our ∼sur guarantee is must stronger, since not only the causal graph can be fully recovered, but the latent variables can be identified up to mixtures of the effect-dominating sets as well.

E. Illustrating examples for our theory and algorithm

E.1 An example for understanding the SNA ambiguity

We provide a simple example below to illustrate the SNA ambiguity discussed in [Section 3.](#page-3-2)

Example 1. Let G be a causal graph with $d = 3$ nodes and edges $1 \rightarrow 2$ and $2 \rightarrow 3$. We have access to *observations from a set of environments* E*. It turns out that there is no way to distinguish between the following two structural equation models:*

$$
z_{1} = \epsilon_{1}^{E}
$$

\n
$$
z_{2} = f_{2}^{E}(z_{1}, \epsilon_{2}^{E})
$$

\n
$$
z_{3} = f_{3}^{E}(z_{2}, \epsilon_{3}^{E})
$$

\n
$$
x = z = (z_{1}, z_{2}, z_{3})^{T}
$$

\n
$$
w_{1} = \epsilon_{1}^{E}
$$

\n
$$
v_{2} = f_{2}^{E}(\nu_{1}, \epsilon_{2}^{E})
$$

\n
$$
v_{3} = \nu_{2} + f_{3}^{E}(\nu_{2}, \epsilon_{3}^{E})
$$

\n
$$
x = (\nu_{1}, \nu_{2}, \nu_{3} - \nu_{2})^{T}
$$

where ϵ_i^E , $i = 1, 2, 3$ are independent noise variables, if we do not change the causal graph $\mathcal G$, no matter what *environment E that we have.*

This issue does not exist when we assume access to hard interventions on node 3*, which effectively removes the edge* 2 \rightarrow 3*. Specifically, with hard intervention on* z_3 *, the variables* z_2 *and* z_3 *become independent. But by definition,* $v_2 = z_2$ *and* $v_3 = z_2 + z_3$ *must be dependent, so this intervention cannot be realized by any hard intervention on v*3*, thereby providing a way to distinguish between the above models.*

Without node 3*, the same ambiguity would arise on node* 2*. However, node* 3 *can help us to overcome this ambiguity, thanks to the fact that node* 2 *is the only causal parent of node* 3*. Suppose for example that* $v_2 = m(z_1, z_2)$ is some mixture of z_1 and z_2 , then $v_3 = \hat{f}_3^E(v_2, \epsilon_3^E) = \hat{f}_3^E(m(z_1, z_2), \epsilon_3^E)$. Since all *environments share the same mixing function,* v_3 *must be some deterministic function* $\psi_3(z)$ *of* z *, where* ψ_3 *is the same across all environment E. Hence, we have*

$$
\hat{f}_3^E\left(m(z_1,z_2),\epsilon_3^E\right)=\psi_3\left(z_1,z_2,f_3^E(z_2,\epsilon_3^E)\right) \tag{7}
$$

*Now we note that the dependencies of LHS on z*¹ *and z*² *are through a single scalar-valued function m, but since we would have different f ^E* 3 *'s in different environments, this in general does not hold for the RHS. Therefore, any causal model with latent variable* v_2 *as a mixture of* z_1 *and* z_2 *<i>cannot be equivalent to the ground-truth model.*

According to [Definition 3,](#page-3-0) in [Example 1](#page-19-2) we have $\text{sur}_G(1) = \text{sur}_G(2) = \emptyset$ but $\text{sur}_G(3) = \{2\}$.

E.2 An example for the main idea behind LiNGCReL

To illustrate our main algorithm on how we can recover the graph G and the matrix *H*, we first provide some intuition using a simple three-node example:

Example 2. Let G be the graph with $d = 3$ nodes and edges $1 \rightarrow 2, 1 \rightarrow 3$ and $2 \rightarrow 3$, so that each \mathbf{B}_k is of *form*

$$
\boldsymbol{B}_{k} = \begin{pmatrix} \times & 0 & 0 \\ \times & \times & 0 \\ \times & \times & \times \end{pmatrix} \stackrel{\rightsquigarrow} \boldsymbol{b}_{k1} \rightsquigarrow \boldsymbol{b}_{k2} \tag{8}
$$

We can identify the graph as follows: first, for $i \in \{1,2,3\}$ *, look at the space* W_i *spanned by the rows* $(M_k)_i, k \in [K]$. If $\dim W_i$ = 1, we know that i is a source node (i.e., $\text{pa}_\mathcal{G}(i)$ = \emptyset) in $\mathcal G$. Otherwise it is not, *due to [Assumption 5.](#page-5-1) Hence we can know that node* 1 *is a source node.*

In our example, there is no other node that satisfies this requirement. We then proceed to search for some i ≠ 1 such that the projection of W ^{*i*} onto W_1^{\perp} has dimension 1. If this holds, then one can show that $\operatorname{pa}_{\mathcal{G}}(i)$ = {1}. Otherwise, *i must have parents other than* 1.

It turns this requirement is satisfied for node 2 since $\dim\left(\text{proj}_{\pmb{h}_1},\text{span}\,\langle \pmb{h}_1,\pmb{h}_2\rangle\right)$ = 1, but is not satisfied

for node 3 since $\dim (\operatorname{proj}_{h_1}$ span $\langle h_1, h_2, h_3 \rangle \geqslant 2$ (by *[Lemma 5\)](#page-21-2). Hence we know that* $\operatorname{pa}_{\mathcal{G}}(2)$ = {1}*.*

Finally, it remains to determine $\rm{pa}_{\mathcal{G}}(3)$ *. To do this, we first note that* $\dim W_3$ = 3*. Then we project W*₃ onto W_1^{\perp} and W_2^{\perp} respectively, and the resulting dimensions are 2 and 1. As we rigorously show in [Proposition 2,](#page-7-4) a decrease of the dimension exactly indicates finding a new parent. Thus we have $\text{pa}_\mathcal{G}(3)$ = {1,2}, *completing the recovery of the graph.*

Finally, we recover the unmixing matrix H (and thus the latent variables) by noticing that $h_1 \in W_1$, *h*² ∈ *W*² ∩ *W*³ *and h*³ ∈ *W*3*. Ambiguities would arise at nodes* 2 *and* 3*, which are exactly the nodes that have non-empty effect-dominating sets.*

F. Auxiliary lemmas

Lemma 3. For any family of m-dimensional vectors $\{\boldsymbol{v}_k\}_{k=1}^K$ and $\{\boldsymbol{z}_k\}_{k=1}^K$ if $\boldsymbol{v}_k = \boldsymbol{z}_k \boldsymbol{T}$ and $\boldsymbol{T} \in \mathbb{R}^{m \times m}$ is *invertible, then*

$$
\dim \operatorname{span} \langle v_k : k \in [K] \rangle = \dim \operatorname{span} \langle z_k : k \in [K] \rangle
$$

Theorem 7 (Darmois-Skitovic Theorem). Let ϵ_i , $i \in [d]$ be independent random variables and X = $\sum_{i=1}^d \alpha_i \epsilon_i$, $Y = \sum_{i=1}^d \beta_i \epsilon_i$. If $X \perp Y$, then for $\forall i \in [d]$, $\alpha_i \beta_i \neq 0 \Rightarrow \epsilon_i$ is Gaussian distributed.

Lemma 4. *Suppose that* $\epsilon = (\epsilon_1, \cdots, \epsilon_d)$ *is a d-dimensional random vector with independent components* s uch that $\text{Var}(\epsilon_i)$ = 1, $\forall i \in [d]$, and there exists an invertible and non-diagonal matrix \bm{M} such that $\bm{M}\epsilon \overset{d}{=} \epsilon_i$ *then at least one of the following statements must hold:*

(1) there exists at least two Gaussian variables in $\epsilon_1, \cdots, \epsilon_d$;

(2) ${\bm M}$ *is a permutation matrix and there exists* $1 \leqslant i < j \leqslant d$ *such that* ${\bm \epsilon}_i \overset{d}{=} {\bm \epsilon}_j$ *.*

Proof. Suppose that (1) does not hold, then there is at most one Gaussian variable in $\epsilon_1, \dots, \epsilon_d$. We assume WLOG that $\epsilon_1, \cdots, \epsilon_{d-1}$ are all non-Gaussian. Then by the Darmois-Skitovic Theorem, we know that for ∀1 ≤ *j* < k ≤ [*d*] and i ∈ [*d* – 1], $M_{ii} \cdot M_{ki}$ = 0 \Rightarrow there is at most one non-zero entry in each of the first *d* – 1 columns of *M*.

Assume that $M_{k_i,i} \neq 0$, $i \in [d-1]$. Since M is invertible, we know that $k_i, i \in [d-1]$ must be different. Let k_d be the remaining element in [*d*] that does not appear in k_i , $i < d$, then $(\boldsymbol{M}\boldsymbol{\epsilon})_{k_d}$ = $\bm{M}_{k_d,d} \bm{\epsilon}_d$, while $(\bm{M} \bm{\epsilon})_{k_i} = \bm{M}_{k_i,i} \bm{\epsilon}_i + \bm{M}_{k_i,d} \bm{\epsilon}_d$. Since the components of $\bm{M} \bm{\epsilon}$ are independent, it is easy to see that $M_{id} \neq 0, \forall i \neq k_d$. In other words, M only has non-zero entries at (k_i, i) , $i \in [d]$.

Since Var(ϵ_i) = 1, we know that M must be a signed permutation matrix. Finally, let π be the permutation on [*d*] such that $\pmb{M}_{i,\pi(i)}\neq 0.$ Since \pmb{M} is not diagonal, π must have a cycle (i_1,i_2,\cdots,i_k) with length $k \geq 2$, so that $\epsilon_{i_1}, \cdots, \epsilon_{i_k}$ all have the same distribution, which implies that (2) holds, as desired. \Box

Lemma 5. Let V_1, V_2 be two subspaces of \mathbb{R}^d such that $V_1 \cap V_2 = \{0\}$, and $P_{V_1^\perp}$ be the orthogonal *projection onto* V_1^{\perp} *, then we have that* $\dim(V_2) = \dim\left(P_{V_1^{\perp}}V_2\right)$ *.*

Proof. Obviously we have $\dim(V_2)\geqslant \dim\left(P_{V_1^\perp}V_2\right).$ On the other hand, let u_1,u_2,\cdots,u_m be a basis of V_2 , then $w_i = P_{V_1^{\perp}} u_i$, $i = 1, 2, \cdots, m$ are also independent. Indeed, suppose that λ_i , $i =$ 1, 2, ..., *m* satisfy $\sum_{i=1}^{m} \lambda_i w_i = 0$, then $P_{V_1^{\perp}}(\sum_{i=1}^{m} \lambda_i u_i) = 0$, implying that $\sum_{i=1}^{m} \lambda_i u_i \in V_1$. However, we know that $V_1 \cap V_2 = \{0\}$, so $\lambda_1 = \cdots = \lambda_m = 0$. This concludes the proof. \Box

Lemma 6. *[Assumption 4](#page-4-4) is equivalent to [Assumption 5.](#page-5-1)*

Proof. The main observation is that for each $k \in [K]$, $(B_k)_i$ only has non-zero entries at the *j*-th coordinate where $j \in \overline{\text{pa}}_{\mathcal{G}}(i)$. Moreover, let $\tilde{\textbf{w}}_k(i)$ be the vector consisting of these entries, then $\tilde{\bm{w}}_k(i) = (\mathbf{\Omega}_k)_{ii}^{-\frac{1}{2}}$ (– $\bm{w}_k(i)$, 1). Hence,

$$
\dim \mathrm{span}\left\langle (\mathbf{B}_k)_i : k \in [K] \right\rangle = \dim \mathrm{span}\left\langle (-\mathbf{w}_k(i), 1) : k \in [K] \right\rangle.
$$

Suppose that [Assumption 4](#page-4-4) holds, then for $\forall x \in \mathbb{R}^{\left| pa_{\mathcal{G}}(i)\right|}$, there exists $\lambda_k \in \mathbb{R}, 1 \leqslant k \leqslant \left| pa_{\mathcal{G}}(i)\right|$ such that $\sum_{k} \lambda_{k} = 1$ and $\sum_{k} \lambda_{k} w_{k}(i) = x$. Hence,

$$
(\mathbf{x},1)=\sum_{k}\lambda_{k}\tilde{\boldsymbol{\psi}}_{k}(i)\in \mathrm{span}\left\langle (\boldsymbol{B}_{k})_{i}:k\in[K]\right\rangle .
$$

This immediately implies that span $\langle (\pmb B_k)_i: k\in[K]\rangle$ = $\mathbb{R}^{\left|{\rm pa}_{\cal G}(i)\right|+1},$ so that [Assumption 5](#page-5-1) holds.

 $\text{Conversely, suppose that Assumption 5 holds, then for \ }\forall \pmb{x}\in\mathbb{R}^{\Big|\text{pa}_{\mathcal{G}}(i)\Big|},$ $\text{Conversely, suppose that Assumption 5 holds, then for \ }\forall \pmb{x}\in\mathbb{R}^{\Big|\text{pa}_{\mathcal{G}}(i)\Big|},$ $\text{Conversely, suppose that Assumption 5 holds, then for \ }\forall \pmb{x}\in\mathbb{R}^{\Big|\text{pa}_{\mathcal{G}}(i)\Big|},$ there exists $\lambda_k\in\mathbb{R},1\leqslant k$ $k \leq \left| \text{pa}_{\mathcal{G}}(i) \right|$ such that $\sum_{k} \lambda_{k} \tilde{\psi}_{k}(i) = (\mathbf{x}, 1)$. Hence we have $\sum_{k} \lambda_{k} \psi_{k}(i) = \mathbf{x}$ and $\sum_{k} \lambda_{k} = 1$, implying [Assumption 4.](#page-4-4) П

G. Properties of effect-domination sets

Lemma 7. • $j \in \text{sur}_G(i)$ *if and only if* $\text{ch}_G(i) \subseteq \text{ch}_G(j)$; • *when* $i \neq j$, $j \in \text{sur}_G(i)$ *if and only if* $\overline{\text{ch}}_G(i) \subseteq \overline{\text{ch}}_G(j)$.

Proof. If $j \in \text{sur}_G(i)$, by definition $i \in \text{ch}_G(i)$ and $\text{ch}_G(i) \subseteq \text{ch}_G(i)$, so that $\overline{\text{ch}_G(i)} \subseteq \text{ch}_G(i)$. Conversely, $ch_G(i) ⊆ ch_G(j)$ implies that $i ∈ ch_G(j)$ and $ch_G(i) ⊆ ch_G(j)$, so $j ∈ sur_G(i)$. This proves the first claim.

To prove the second claim, assume that ch_G(*i*) \subseteq ch_G(*j*) holds but ch_G(*i*) \subseteq ch_G(*j*) does not hold, then we must have $j \in \overline{ch}_G(i)$. since $j \neq i$, we have $j \in ch_G(i)$, but then $i \notin \overline{ch}_G(j)$, which is a contradiction. Hence $ch_G(i) \subseteq ch_G(j)$ and the conclusion follows from the first claim. \Box

 ${\bf Lemma~8.~}$ *Let* ${\cal G}$ *be a DAG and i be its node, then for* $\forall j \in \text{pa}_{\cal G}(i)$ *, we have* $\text{sur}_{\cal G}(j) \subseteq \text{pa}_{\cal G}(i)$ *.*

Proof. Let $k \in \text{sur}_G(i)$, then by definition we have $\text{ch}_G(i) \subseteq \text{ch}_G(k)$. In particular, we have $i \in$ $ch_{\mathcal{G}}(k) \Rightarrow k \in pa_{\mathcal{G}}(i).$ \Box

Lemma 9. Let G be a DAG and i be its node, then for $\forall j \in \text{sur}_{G}(i)$, we have $\text{sur}_{G}(j) \subseteq \text{sur}_{G}(i)$.

Proof. Let $k \in \text{sur}_G(j)$, then by definition we have $\overline{\text{ch}}_G(j) \subset \overline{\text{ch}}_G(k)$. We also know that $\overline{\text{ch}}_G(i) \subset$ $ch_G(i)$, so $ch_G(i) \subset ch_G(k)$, implying that $k \in \text{sur}_G(i)$. \Box

Lemma 10. *If* $M \in \mathcal{M}_{\text{sur}}^0(\mathcal{G})$, then $M^{-1} \in \mathcal{M}_{\text{sur}}^0(\mathcal{G})$.

Proof. Assume WLOG that the nodes of G satisfy $i \in pa_{\mathcal{G}}(j) \Rightarrow i < j$ (otherwise we can choose a different index of the nodes and correspondingly swap some rows and columns of *M*). Since $i \in \text{sur}_{\mathcal{G}}(j) \Rightarrow i \in \text{pa}_{\mathcal{G}}(j)$, it follows that M must be lower triangular and the diagonal entries are nonzero.

Let $N = M^{-1}$, then for $\forall i \in [d]$, we have

$$
\sum_{j=1}^{d} \mathbf{N}_{ij} \mathbf{M}_{j\ell} = 0, \quad \forall \ell \notin \overline{\text{sur}}_{\mathcal{G}}(i). \tag{9}
$$

Since $M\in \mathcal{M}_{\text{sur}}^0(\mathcal{G})$, we have $M_{j\ell}$ = 0 for $\forall j$ such that $\ell\notin\overline{\text{sur}}_\mathcal{G}(j)$. By [Lemma 9,](#page-22-1) if $j\in\overline{\text{sur}}_\mathcal{G}(i)$, then $\ell \notin \overline{\text{sur}}_G(i)$ necessarily implies that $\ell \notin \overline{\text{sur}}_G(i)$. Hence the left hand side of [\(9\)](#page-22-2) is essentially a sum over $j \notin \overline{\sup}_{G}(i)$, *i.e.*,

$$
\sum_{j \notin \overline{\text{sur}}_{\mathcal{G}}(i)} \mathbf{N}_{ij} \mathbf{M}_{j\ell} = 0, \quad \forall \ell \notin \overline{\text{sur}}_{\mathcal{G}}(i).
$$

Viewing the above as a system of linear equations in $\bm{N}_{ij}, j \notin \overline{\text{sur}}_{\mathcal{G}}(i)$, the coefficient matrix $\left(\bm{M}_{j\ell}\right)$ $j, l \in \notin$ sur $_{\mathcal{G}}(i)$

must be invertible since it is a sub-matrix of the invertible lower-triangular matrix *M*. As a result, we necessary have N_{ij} = 0, $\forall j \notin \overline{\text{sur}}_{\mathcal{G}}(i)$. Finally, $N = M^{-1}$ must be invertible, so $N \in \mathcal{M}^0_{\text{sur}}(\mathcal{G})$ as desired.

Lemma 11. *Suppose that* ψ : R *d* 7→ R *d is a diffeomorphism and* G *be a DAG, such that for* ∀*i* ∈ [*d*]*,* $\psi_i(z)$ *is a function of* $z_{\overline{\text{sur}}_\mathcal{G}(i)}$ *. Then for* $\forall j\in[d]$, $(\psi^{-1})_j(\pmb{\nu})$ *is a function of* $v_{\overline{\text{sur}}_\mathcal{G}(j)}$ *.*

Proof. Let J_z = $J_\psi(z)$ be the Jacobian matrix of ψ. Since ψ is a diffeomorphism, J_z is invertible for any $\boldsymbol{z}\in\mathbb{R}^d$. Moreover, our assumption implies that $(\boldsymbol{J}_{\boldsymbol{z}})_{ij}$ = 0, $\forall j\notin\overline{\mathrm{sur}}_{\mathcal{G}}(i)$, so $\boldsymbol{J}_{\boldsymbol{z}}\in\mathcal{M}^0_{\mathrm{sur}}(\mathcal{G})$. By Lemma 10 , $J_z^{-1} \in \mathcal{M}_{\text{sur}}^0(\mathcal{G})$. But J_z^{-1} is exactly the Jacobian matrix of ψ^{-1} at $\nu = \psi(z)$, hence it follows that $(\uppsi^{-1})_j(\pmb{\nu})$ is only a function of $\nu_{\overline{\text{sur}}_\mathcal{G}(j)},$ as desired. \Box

Lemma 12. *The binary relation* ∼sur *defined in [Definition 4](#page-3-1) is an equivalence relation.*

Proof. It is obvious that $(h, \mathcal{G}) \sim_{\text{sur}} (h, \mathcal{G})$ holds for any model (h, \mathcal{G}) .

Suppose that $(h_1, G_1) \sim_{sur} (h_2, G_2)$, then there exists a permutation π on [*d*] and a diffeomorphism $\psi:\mathbb{R}^d\mapsto\mathbb{R}^d$ where $\psi_i(z)$ is a function of $z_{\overline{\text{sur}}_{\mathcal{G}_1}(i)},$ such that $i\in\text{pa}_{\mathcal{G}_1}(j)\Leftrightarrow\pi(i)\in\text{pa}_{\mathcal{G}_2}(\pi(j))$ and $P_π ∘ h₂ = √ ∘ h₁$. Then we can write $P_π^{-1} ∘ h₁ = √ ∘ h₂$ where $\hat{ψ} = P_π^{-1} ∘ ψ^{-1} ∘ P_π$. By [Lemma 11,](#page-22-4) we know that $(\psi^{-1})_j(\nu)$ is a function of $\nu_{\overline{\text{sur}}_{\mathcal{G}_1}(j)}$, so $(\hat{\psi})_j$ is a function of $\nu_{\pi(\overline{\text{sur}}_{\mathcal{G}_1}(j))} = \nu_{\overline{\text{sur}}_{\mathcal{G}_2}(j)}$, implying that $(h_2, \mathcal{G}_2) \sim_{\text{sur}} (h_1, \mathcal{G}_1)$.

Finally, let $(h_1, G_1) \sim_{\text{sur}} (h_2, G_2)$ and $(h_2, G_2) \sim_{\text{sur}} (h_3, G_3)$, then we can write

$$
\boldsymbol{P}_{\boldsymbol{\pi}} \circ \boldsymbol{h}_2 = \boldsymbol{\psi} \circ \boldsymbol{h}_1 \quad \text{and} \quad \boldsymbol{P}_{\hat{\boldsymbol{\pi}}} \circ \boldsymbol{h}_3 = \hat{\boldsymbol{\psi}} \circ \boldsymbol{h}_2
$$

where: for $\forall i \in [d]$, $\psi_i(z)$ is a function of $z_{\overline{\text{sur}}_{\mathcal{G}_1}(i)}$, $\hat{\psi}_i(z)$ is a function of $z_{\overline{\text{sur}}_{\mathcal{G}_2}(i)}$, $i \in \text{pa}_{\mathcal{G}_1}(j) \Leftrightarrow$ $\pi(i) \in \text{pa}_{\mathcal{G}_2}(\pi(j))$ and $i \in \text{pa}_{\mathcal{G}_2}(j) \Leftrightarrow \hat{\pi}(i) \in \text{pa}_{\mathcal{G}_2}(\hat{\pi}(j)).$ Then, we can write

$$
P_{\pi}\circ P_{\hat{\pi}}\circ h_3=P_{\pi}\circ \hat{\psi}\circ P_{\pi}^{-1}\circ \psi\circ h_1.
$$

Since $\hat{\psi}_i(z)$ is a function of $z_{\overline{\text{sur}}_{\mathcal{G}_2}(i)}$, we deduce that $\left(P_\pi \circ \hat{\psi} \circ P_\pi^{-1}\right)_i(z)$ is a function of $z_{\overline{\text{sur}}_{\mathcal{G}_1}(i)}$. Hence, $\left(P_\pi \circ \hat{\psi} \circ P_\pi^{-1} \circ \psi\right)_i (z) = \left(P_\pi \circ \hat{\psi} \circ P_\pi^{-1}\right)_i (\psi(z))$ is a function of $\psi_{\overline{\text{sur}}_{\mathcal{G}_1}(i)}(z)$. The definition of ψ implies that for each $j \in \overline{\text{sur}}_{\mathcal{G}_1}(i)$, $\psi_j(z)$ is a function of $z_{\overline{\text{sur}}_{\mathcal{G}_1}(j)}$. By [Lemma 9,](#page-22-1) we have $\cup_{j\in\overline{\text{sur}}_{\mathcal{G}_1}(j)}\overline{\text{sur}}_{\mathcal{G}_1}(j)\subseteq\overline{\text{sur}}_{\mathcal{G}_1}(i).$ Hence $\left(P_\pi\circ\hat{\psi}\circ P_\pi^{-1}\circ\psi\right)$ $\mathcal{Z}_{\text{sur}}(i)$. Moreover, we also have $i \in pa_{\mathcal{G}_1}(j) \Leftrightarrow \pi(i) \in pa_{\mathcal{G}_2}(\pi(j)) \Leftrightarrow \hat{\pi} \circ \pi(i) \in pa_{\mathcal{G}_2}(\hat{\pi} \circ \pi(j))$, so by definition, (*h*1, G1) ∼sur (*h*3, G3), as desired. \Box

H. Omitted proofs from [Section 4](#page-4-3) and [Section 5](#page-6-1)

H.1 Proof of [Theorem 1](#page-5-0)

According to the assumption, we have that $\epsilon = B_k H x$ and $\hat{\epsilon} = \hat{B}_k \hat{H} x$, so that $\epsilon = B_k H (\hat{B}_k \hat{H})^{\dagger} \hat{\epsilon}, \forall k \in$ [*K*]. By [Lemma 4,](#page-20-2) we know that for each k , $P_k := B_k H(\hat{B}_k \hat{H})^{\dagger}$ is a signed permutation matrix, so that $\epsilon = P_k \hat{\epsilon}$. Since for any $i \neq j$, $\hat{\epsilon}_i \neq \hat{\epsilon}_j$, we must have $|P|_1 = |P|_2 = \cdots = |P|_K =: P$ and $\epsilon = P \hat{\epsilon}$, where $|M|$ denotes the resulting matrix by taking the absolute value of all entries in *M*. Thus, we can WLOG assume that $\epsilon = \hat{\epsilon}$, since otherwise we can permute the noise variables $\hat{\epsilon}$, and also permute the rows of B_k correspondingly. In other words, suppose that the permutation matrix $|P|$ has $|P|_{k_i,i} = 1$, $i \in [d]$, then we can assign to each node *i* in \hat{G} a new index k_i and work with the new indices.

In this case, by [Lemma 4](#page-20-2) we have $B_k H = \Sigma_k \hat{B}_k \hat{H}$, $\forall k \in [K]$ or equivalently $\Sigma_k \hat{B}_k = B_k T$, where $\bm{T} = \bm{H}\hat{\bm{H}}^{\dagger} \in \mathbb{R}^{d \times d}$, and $\bm{\Sigma}_k$ is a diagonal matrix with diagonal entries in {+1,-1}. Let $\hat{\hat{\bm{B}}}_k = \bm{\Sigma}_k \hat{\bm{B}}_k$, then the rows of $\hat{\hat{\textbf{\textit{B}}}}_k$ equals (up to sign) to the rows of $\hat{\textbf{\textit{B}}}_k$.

To summarize, we now know that i) $\hat{\hat{\bm{B}}}_k = \bm{B}_k \bm{T}, k \in [K],$ ii) $(\bm{B}_k)_{ij} \neq 0 \Leftrightarrow j \in \overline{\text{pa}}_{\mathcal{G}}(i)$, and similarly, $(\hat{\hat{\bm{B}}}_k)_{ij} \neq 0 \, \Leftrightarrow j \in \overline{\mathrm{pa}}_{\hat{\mathcal{G}}}(i)$, and iii) Both $\{\bm{B}_k\}$ and $\left\{\hat{\hat{\bm{B}}}_k\right\}$ satisfy the node-level non-degeneracy assumption [Assumption 5.](#page-5-1) For any two such matrices that satisfy such a set of conditions, it must necessarily be true that $G = \tilde{G}$.

 $\bf{Lemma \ 13}$ (Graph Identifiability). *Consider any two sets matrices* $\{ \hat{\hat{\bm{B}}}_k \}_{k \in [K]}$ and $\{ \bm{B}_k \}_{k \in [K]}$ and *associated graphs* G, Gˆ*. If these sets and graphs satisfy that:*

- *1.* $\hat{B}_k = B_k T, k \in [K]$;
- 2. $(\mathbf{B}_k)_{ij} \neq 0 \Leftrightarrow j \in \overline{\mathrm{pa}}_{\mathcal{G}}(i)$, and similarly, $(\hat{\hat{\mathbf{B}}}_k)_{ij} \neq 0 \Leftrightarrow j \in \overline{\mathrm{pa}}_{\hat{\mathcal{G}}}(i)$.
- *3. Both* $\{B_k\}$ *and* $\{\hat{\hat{B}}_k\}$ *satisfy the node-level non-degeneracy assumption [Assumption 5.](#page-5-1)*

then it must hold that $G = \hat{G}$ *.*

Proof. We prove this via induction on the size of the graph *d*. Note that here $G = \hat{G}$ is not up to permutation and our statement is equivalent to pa $_{\mathcal{G}}(i)$ = pa $_{\hat{\mathcal{G}}}(i), \forall i \in [\mathcal{d}].$

If $d = 1$, *i.e.*, $\mathcal{G} = \hat{\mathcal{G}}$ obviously holds since both are graphs with only 1 node.

Suppose that for all graphs G of size $d-1$, the graph \hat{G} satisfying all given assumptions must necessarily be equal to G. Now, we consider the case that G has *d* nodes. WLOG we can assume that the nodes of G are properly indexed such that $i \in \text{pa}_{\mathcal{G}}(j) \Rightarrow i < j$, so $\mathcal{B}_k, k \in [K]$ are lower-triangular

matrices. (However, it is currently unknown whether $\hat{\hat{\bm{B}}}_k$ are also lower-triangular.)

By our assumption that $i \in \text{pa}_{\mathcal{G}}(j) \Rightarrow i \leq j$, the node d in $\mathcal G$ has no child. Thus we can write

$$
\boldsymbol{B}_k = \begin{pmatrix} \boldsymbol{B}_k^- & \boldsymbol{0} \\ \boldsymbol{b}_k & c_k \end{pmatrix}, \boldsymbol{T} = \begin{pmatrix} \boldsymbol{T}^- & \times \\ \times & \times \end{pmatrix} \text{ and } \hat{\boldsymbol{B}}_k = \boldsymbol{B}_k \boldsymbol{T} = \begin{pmatrix} \hat{\boldsymbol{B}}_k^- & \times \\ \times & \times \end{pmatrix}
$$

where \bm{B}_{k}^{-} , $\bm{T}, \hat{\hat{\bm{B}}}_{k}^{-} = \bm{B}_{k}^{-} \bm{T}^{-} \in \mathbb{R}^{(d-1)\times(d-1)}, \bm{b}_{k} \in \mathbb{R}^{d-1}, c_{k} \in \mathbb{R}$ and \times denotes irrelevant entries.

Let $\vec{A}_{k}^-, \hat{A}_{k}^ \sum_{k}^{T} \Omega_{k}^{-}$ and $\hat{\Omega}_{k}^{-}$ be the top-left $(d-1) \times (d-1)$ sub-matrices of A_{k} , \hat{A} , Ω_{k} and $\hat{\Omega}_{k}^{-}$ respectively, and ${\cal G}^-$ and $\hat{\cal G}^-$ are graphs obtained by deleting node d and all related edges from ${\cal G}$ and \hat{G} . Then it is easy to see that

$$
\left(A_k^-\right)_{ij} \neq 0 \Leftrightarrow j \in \text{pa}_{\mathcal{G}^-}(i) \quad \text{and} \quad \left(\hat{A}_k^-\right)_{ij} \neq 0 \Leftrightarrow j \in \text{pa}_{\hat{\mathcal{G}}^-}(i). \tag{10}
$$

Moreover,

$$
\begin{pmatrix}\n\mathbf{B}_k^- & \mathbf{0} \\
\mathbf{b}_k^+ & c_k\n\end{pmatrix} = \mathbf{B}_k = \Omega_k^{-\frac{1}{2}} \left(\mathbf{I} - \mathbf{A}_k \right) = \begin{pmatrix}\n\left(\Omega_k^- \right)^{-\frac{1}{2}} & \mathbf{0} \\
\mathbf{0} & \times\n\end{pmatrix} \begin{pmatrix}\n\mathbf{I} - \mathbf{A}_k^- & \times \\
\times & \times\n\end{pmatrix} = \begin{pmatrix}\n\left(\Omega_k^- \right)^{-\frac{1}{2}} \left(\mathbf{I} - \mathbf{A}_k^- \right) & \times \\
\times & \times\n\end{pmatrix}
$$

so that $\bm{B}_k^- = \left(\bm{\Omega}_k^-\right)^{-\frac{1}{2}} \left(\bm{I} - \bm{A}_k^-\right)$. Similarly, we have $\hat{\hat{\bm{B}}}_k^- = \left(\hat{\bm{\Omega}}_k^-\right)$ $\left(\frac{1}{k}\right)^{-\frac{1}{2}}$ $(I - \hat{A}_k)$ *k*).

We can also verify that $\left\{\textit{B}_{\overline{k}}\right\}_{\overline{k}^{\circ}}^{K}$ $\left\{\hat{\hat{\mathbf{B}}}_k^{\!\!\top}\right\}_{k=1}^K$ are node-level independent in the sense of [As](#page-5-1)[sumption 5.](#page-5-1) We only prove this for $\left\{\hat{\hat{\bm{B}}}_k\right\}_{k=1}^K$; the arguments used for $\{\bm{B}_k\}_{k=1}^K$ are exactly the same as the first case considered below. Now for each $i \in [d-1]$, let $R_i \in \mathbb{R}^{K \times d}$ be the matrix whose k -th row is the *i*-th row of \hat{B}_k , and $R_i^- \in \mathbb{R}^{K \times (d-1)}$ be the matrix whose *k*-th row is the *i*-th row of \hat{B}_k^- . then obviously R_i is of form $[R_i^-, r_i]$. We consider two cases:

- **Case 1.** $d \notin \text{pa}_{\hat{\mathcal{G}}}(i)$ This means that the last entry of the *i*-th row of $\hat{\hat{\textbf{B}}}_k$ is zero. Thus $\textbf{\textit{r}}_i$ = **0**, and $\text{rank}(\mathcal{R}_i^-) = \text{rank}(\mathcal{R}_i) = \left| \overline{\text{pa}}_{\hat{\mathcal{G}}}(i) \right| = \left| \overline{\text{pa}}_{\hat{\mathcal{G}}^-}(i) \right|$, where the second equality follows from [Assumption 5.](#page-5-1)
- Case $2.d \in pa_{\hat{\mathcal{G}}}(i)$ In this case we have rank $(R_i^-) \geqslant$ rank $(R_i) 1 = \left|\overline{\text{pa}}_{\hat{\mathcal{G}}}(i)\right| 1 = \left|\overline{\text{pa}}_{\hat{\mathcal{G}}^+}(i)\right|$. Due to our assumption on $\hat{\bm{A}}_k$ and the relationship $\hat{\hat{\bm{B}}}_k^- = \left(\hat{\bm{\Omega}}_k^- \right)$ $\left(\frac{1}{k}\right)^{-\frac{1}{2}}$ $(I - \hat{A}_k)$ *k*), we know that each row of R_i^- , namely the *i*-th row of some \hat{B}_k only has $\left| \overline{\mathrm{pa}}_{\hat{\mathcal{G}}}(i) \right| - 1 = \left| \overline{\mathrm{pa}}_{\hat{\mathcal{G}}^-}(i) \right|$ non-zero entries, so that rank $(R_i^-) = \left| \overline{\text{pa}}_{\hat{\mathcal{G}}^{-}}(i) \right|$ holds.

Since we have shown that the matrices $B_{\vec{k}}^-$ and $\hat{B}_{\vec{k}}^-$ satisfy the three properties that we assume for induction with T replaced by T^- and $\mathcal{G},\hat{\mathcal{G}}$ replaced by $\mathcal{G}^-,\hat{\mathcal{G}}^-$ respectively, by induction hypothesis, we can thus deduce that G^- = \hat{G}^- . To prove G = \hat{G} it remains to show that the dependency of node *d* on the remaining nodes are the same in G and \hat{G} .

First, we show that $ch_{\hat{G}}(d) = \emptyset$. Suppose in contrary that there is some $i \in ch_{\hat{G}}(d)$, then $\left|\text{pa}_{\mathcal{G}}(i)\right| = \left|\text{pa}_{\mathcal{G}^-}(i)\right| = \left|\text{pa}_{\hat{\mathcal{G}}^-}(i)\right| = \left|\text{pa}_{\hat{\mathcal{G}}}(i)\right| - 1$. Recalling that $\left(\mathcal{B}\right)_i$ denotes the *i*-th row of matrix *B*, we have

$$
\dim \left(\text{span}\left\langle \left(\hat{\hat{\mathbf{B}}}_k\right)_i : 1 \leq k \leq K \right\rangle \right) = \dim \left(\text{span}\left\langle \left(\mathbf{B}_k\right)_i : 1 \leq k \leq K \right\rangle \right) \leq \left| \text{pa}_{\hat{\mathcal{G}}}(i) \right| + 1 < \left| \text{pa}_{\hat{\mathcal{G}}}(i) \right| + 1,\n\tag{11}
$$

where the first inequality follows from $\left(\hat{\hat{B}}_k\right)_i$ = $\left(B_k\right)_i$ T and [Lemma 3,](#page-20-3) the second holds since each $(B_k)_i$ has nonzero elements only at coordinates in $j \in \overline{\text{pa}}_G(i)$, and the last one holds since $\left|\text{pa}_{\mathcal{G}}(i)\right| = \left|\text{pa}_{\hat{\mathcal{G}}}(i)\right| - 1$. However, [\(11\)](#page-24-0) contradicts the non-degeneracy condition [Assumption 5](#page-5-1) that we assume for matrices $\hat{B}_k, k \in [K]$ in the statement of the theorem. Therefore we have $ch_{\hat{G}}(d) = \emptyset = ch_{G}(d).$

Second, by a similar argument comparing the number of nonzero elements in the last row of *B^k* and $\hat{\hat{\textbf{\textit{B}}}}_k$, we can also deduce that

$$
\left|\mathrm{pa}_{\mathcal{G}}(d)\right|=\left|\mathrm{pa}_{\hat{\mathcal{G}}}(d)\right|.
$$

Indeed, since $\left(\hat{\hat{\textbf{\textit{B}}}}_k\right)$ $d = (\mathcal{B}_k)_d$ **T**, by [Lemma 3](#page-20-3) we have

$$
\dim\left(\text{span}\left\langle\left(\hat{\hat{\mathbf{B}}}_k\right)_d:1\leqslant k\leqslant K\right\rangle\right)=\dim\left(\text{span}\left\langle\left(\mathbf{B}_k\right)_d:1\leqslant k\leqslant K\right\rangle\right)
$$

However, since we assume that [Assumption 5](#page-5-1) is satisfied for $\{\bm B_k\}_{k=1}^K$ and $\left\{\hat{\bm B}_k\right\}_{k=1}^K$ $_{k=1}$, we know that the LHS and RHS of the above equation are equal to $\left|\text{pa}_{\mathcal{G}}(d)\right|$ + 1 and $\left|\text{pa}_{\hat{\mathcal{G}}}(d)\right|$ + 1 respectively, implying [\(12\).](#page-25-0)

Third, we show that pa $_{\cal G}(d)$ = pa $_{\hat {\cal G}}(d)$. Suppose the contrary, let ℓ be the smallest element in pa $_G(d)$ ∆pa $_{\hat{\cal G}}(d)$, where $A\Delta B$:= $(A\setminus B)\cup (B\setminus A)$. Recall that while ${\cal G}$ and $\hat{\cal G}$ are originally not symmetric as nodes are topologically sorted according to $\mathcal G$, now we have shown that $\mathcal G^-\equiv\hat{\mathcal G}^-$ and that $\ch_{\mathcal{G}}(d)$ = $\ch_{\hat{\mathcal{G}}}(d)$ = \emptyset , so we can assume WLOG that $\ell \in$ pa $_{\mathcal{G}}(d)$ and $\ell \notin$ pa $_{\hat{\mathcal{G}}}(d)$, and the other case can be handled symmetrically. Since \bm{B}_k is lower triangular and $\left(\bm{B}_k\right)_{jj}=\left(\Omega_k\right)_{jj}^{-\frac{1}{2}}\neq 0, \forall j\in [d],$ the top-left $\ell \times \ell$ sub-matrix of \mathcal{B}_k , which we denote by $[\mathcal{B}_k]_{\ell,\ell}$, must be invertible. This implies ${\rm that}\, \left\{ [{\bm B}_k]_{\ell,\ell}^\top\, \bm{\lambda}: \bm{\lambda} \in \mathbb{R}^\ell\right\} = \mathbb{R}^\ell,$ so we can always find coefficients $\lambda_{kj}, j \in [\ell]$ such that the first ℓ entries of the vector $(\bm{B}_k)_{d}$ – $\sum_{i=1}^{\ell} \lambda_{ki} (\bm{B}_k)_{i} \in \mathbb{R}^{d}$ are all zero. Since $\hat{\hat{\bm{B}}}_k$ = $\bm{B}_k \bm{T}$ and \bm{T} is invertible, we have $\left(\hat{\hat{\textbf{\textit{B}}}}_k\right)$ \mathcal{L}_{d} – $\sum_{j=1}^{\ell} \lambda_{kj} \left(\hat{\hat{\mathbf{B}}}_{k}\right)$ j = $\left(\left(\bm{B}_{k}\right)_{d}-\sum_{j=1}^{\ell}\lambda_{kj}\left(\bm{B}_{k}\right)_{j}\right)\bm{T},\forall k\in[K]$ and

$$
\dim \left(\operatorname{span} \left\langle \left(\hat{\hat{\mathbf{B}}}_{k} \right)_{d} - \sum_{j=1}^{\ell} \lambda_{kj} \left(\hat{\hat{\mathbf{B}}}_{k} \right)_{j} : k \in [K] \right\rangle \right) = \dim \left(\operatorname{span} \left\langle \left(\mathbf{B}_{k} \right)_{d} - \sum_{j=1}^{\ell} \lambda_{kj} \left(\mathbf{B}_{k} \right)_{j} : k \in [K] \right\rangle \right)
$$

$$
\leqslant \left| \operatorname{pa}_{\mathcal{G}}(d) \setminus [\ell] \right| + 1.
$$

Here, the inequality holds because for any coordinate $t \in [d]$,

$$
\left(\left(\mathbf{B}_k \right)_d - \sum_{j=1}^{\ell} \lambda_{kj} \left(\mathbf{B}_k \right)_j \right)_t = \begin{cases} 0 & \text{if } t \leq \ell \\ \left(\mathbf{B}_k \right)_{d,t} & \text{otherwise} \end{cases} \tag{12}
$$

where we note that B_k is lower-triangular and thus $(B_k)_{j,t} = 0, \forall j \leq \ell, t > \ell$. This implies that $\left(\left(\mathbf{B}_{k} \right)_{d} - \sum_{j=1}^{\ell} \lambda_{kj} \left(\mathbf{B}_{k} \right)_{j} \right)$ *t* is nonzero only if $t > l$ and $t \in pa_G(d)$.

On the other hand, let $S = \left(\text{pa}_{\hat{\mathcal{G}}}(d) \cap [\ell]^c\right) \cup \{d\}$, then

$$
\dim \left(\text{span}\left\langle \left(\hat{\hat{\mathbf{B}}}_{k}\right)_{d}-\sum_{j=1}^{\ell} \lambda_{kj} \left(\hat{\hat{\mathbf{B}}}_{k}\right)_{j}: k \in [K] \right\rangle\right) \geq \dim \left(\text{span}\left\langle \left(\left(\hat{\hat{\mathbf{B}}}_{k}\right)_{d}-\sum_{j=1}^{\ell} \lambda_{kj} \left(\hat{\hat{\mathbf{B}}}_{k}\right)_{j}\right)_{S}: k \in [K] \right\rangle\right) \n= \dim \left(\text{span}\left\langle \left(\left(\hat{\hat{\mathbf{B}}}_{k}\right)_{d}\right)_{S}: k \in [K] \right\rangle\right) = |S|.
$$

where we recall that u_S denotes the vector $(u_i : i \in S) \in \mathbb{R}^{|S|}$. Here the first equality holds due to the same reason as [\(12\),](#page-25-0) and the second follows from [Assumption 5.](#page-5-1) To see why this is the case, note that [Assumption 5](#page-5-1) implies that the $K \times (\left|\text{pa}_{\mathcal{G}}(d)\right|+1)$ having $((\mathcal{B}_k)_d)_{\overline{\text{pa}}_{\mathcal{G}}(d)}$ as the *k*-th row has full column rank, so that the sub-matrix obtained by extracting columns corresponding to the node set *S* also has full column rank.

We have shown that $\left|\overline{\text{pa}}_{\hat{\mathcal{G}}}(d) \cap [\ell]^c\right| = |S| \leqslant \left|\text{pa}_{\mathcal{G}}(d) \cap [\ell]^c\right| + 1 = \left|\overline{\text{pa}}_{\mathcal{G}}(d) \cap [\ell]^c\right|$. On the other hand, recall that by our choice of ℓ , we have $\left|\overline{\mathrm{pa}}_{\mathcal{G}}(d) \cap [\ell-1]\right| = \left|\overline{\mathrm{pa}}_{\hat{\mathcal{G}}}(d) \cap [\ell-1]\right|$ and $\ell \in \overline{\mathrm{pa}}_{\mathcal{G}}(d)$ $\overline{\text{pa}}_{\hat{\mathcal{G}}}(d)$. Putting these together, we have $\left|\overline{\text{pa}}_{\hat{\mathcal{G}}}(d)\right| > \left|\overline{\text{pa}}_{\hat{\mathcal{G}}}(d)\right|$. However, we know from [\(12\)](#page-25-0) that $\left|\text{pa}_{\mathcal{G}}(d)\right| = \left|\text{pa}_{\hat{\mathcal{G}}}(d)\right|$, leading to a contradiction. Hence, such ℓ shouldn't exist and we must have $\text{pa}_{\mathcal{G}}(d)$ = $\text{pa}_{\hat{\mathcal{G}}}(d)$, completing the induction step for graphs of size *d*.

By the principle of induction, we have shown that $G = \hat{G}$ holds for any graphs under given assumptions. \Box

Now that we have established that $G = \hat{G}$, we prove the remaining part of the theorem. Note that for any $i,j\in [d]$ such that $i\notin \overline{\mathrm{pa}}_{\mathcal{G}}(j)$, we have $(\boldsymbol{B}_k)_{ji}$ = $(\hat{\hat{\boldsymbol{B}}}_k)_{ji}$ = $0,\forall k\in [K].$ Since $\hat{\hat{\boldsymbol{B}}}_k$ = $\boldsymbol{B}_k\boldsymbol{T},$ we have

$$
\sum_{\ell\in\overline{\text{pa}}_{\mathcal{G}}(j)}(\pmb{B}_k)_{j\ell}\,\pmb{T}_{\ell\,i}=0.
$$

By [Assumption 5,](#page-5-1) the above implies that $\bm{T}_{\ell i}$ = 0 for $\forall \ell \in \overline{\bm{\mathsf{pa}}}_{\mathcal{G}}(j).$ In short, we have argued that if there exists *j* such that $i \notin \overline{\text{pa}}_{\mathcal{G}}(j)$ and $\ell \in \overline{\text{pa}}_{\mathcal{G}}(j)$, then $\bm{T}_{\ell i}$ = 0.

This implies that $T_{\ell i}$ is non-zero only if $\bar{\ch}_\mathcal{G}(\ell) \subseteq \bar{\ch}_\mathcal{G}(i)$. Since v = \bm{Tz} , we have v_ℓ = \sum_i^α This implies that $T_{\ell i}$ is non-zero only if $\ch_{\mathcal{G}}(\ell) \subseteq \ch_{\mathcal{G}}(i)$. Since $v = Tz$, we have $v_{\ell} = \sum_{i=1}^{u} T_{\ell i} z_i = \sum_{i \in [d]: \bar{\ch}_{\mathcal{G}}(\ell) \subseteq \bar{\ch}_{\mathcal{G}}(i)} T_{\ell i} z_i$. Note that when $i \neq \ell$, $\bar{\ch}_{\mathcal{G}}(\ell) \subseteq \bar{\ch$ only depends on $z_{\overline{\mathrm{sur}}_\mathcal{G}(\ell)}$ by [Lemma 7,](#page-21-3) as desired.

H.2 Formal version and proof of [Theorem 2](#page-5-2)

In previous works (Seigal, Squires, and Uhler [2022;](#page-13-10) J. Zhang et al. [2023\)](#page-13-13), it is common to consider single-node soft interventions in the following sense:

Assumption 8. *For* $\forall 2 \leq k \leq K$ *, there exists* $i_k \in [d]$ *, such that the structural equation in environment* k *satisfies* [\(4\)](#page-4-5) *satisfies* $w_k(i) = w_1(i)$ *and* $\omega_{k,i,i} = \omega_{1,i,i}$ for $\forall i \neq i_k$ *.*

Let $S_i = \{k : 2 \le k \le K, i_k = i\}$, $i \in [d]$ and $s_i = |S_i|$. Suppose that G has $e = \sum_{i=1}^d |pa_G(i)|$ edges, then we can view the weight vectors $\{(\mathbf{w}_k(i), \mathbf{w}_{k,i,i}): k = 1 \text{ or } i = i_k\}$ as elements of the Euclidean $\exp \exp \left(\frac{e^{i\pi \sum_{k=2}^K \left| \mathrm{pa}_{\mathcal{G}}(i_k) \right|}}{\sum_{k=1}^K \mathbb{R}^{d+k-1}} \right)$. Under [Assumption 8,](#page-26-1) the models can be fully determined by these weight vectors. The following result states that if we restrict ourselves to single-node interventions, then in the worst case, $\Theta(d^2)$ interventions are required.

 $\bf{Theorem 8.}$ $\it There \; exists \; a \; causal \; graph \; \mathcal{G} \; with \; \Theta(d^2) \; edges, such \; that \; for \; any \; unmixing \; matrix \; H \in \mathbb{R}^{d \times n}$ with full row rank, any independent noise variables ϵ , and any $s_i > 0, i \in [d]$ such that $s_i \leqslant \left|\text{pa}_{\mathcal{G}}(i)\right|$ for some i, the following holds: except from a null set of the weight space $\mathbb{R}^{e+\sum_{k=2}^K\left|\text{pa}_{\mathcal{G}}(\mathbf{i}_k)\right|}\times\mathbb{R}_+^{d+K-1}$ (w.r.t the *Lebesgue measure), there must exist a candidate solution* (*H*ˆ , Gˆ) *and a hypothetical data generating process*

$$
\forall k \in [K], \quad \mathbf{v} = \hat{A}_k \mathbf{v} + \hat{\Omega}_k^{\frac{1}{2}} \mathbf{\epsilon}, \quad \mathbf{x} = \hat{H}^{\dagger} \mathbf{v}
$$

such that

 (i') the unmixing matrix $\hat{\boldsymbol{H}} \in \mathbb{R}^{d \times n}$ has full row rank;

(ii') $\forall k \in [K]$ and $i, j \in [d], (\hat{A}_k)_{ij} \neq 0 \Leftrightarrow j \in \text{pa}_{\hat{\mathcal{G}}}(i)$ and $\hat{\mathbf{\Omega}}_k$ is a diagonal matrix with positive entries;

(iii') for $\forall 2\leqslant k\leqslant K,$ the weight matrices $\hat A_k,\hat{\bf\Omega}_k$ of environment E_k are from a single-node soft intervention *on E*¹ *on node i^k , in the sense of [Assumption 8,](#page-26-1)*

but G *is non-isomorphic to* \hat{G} *.*

In this subsection we give the full proof of [Theorem 8.](#page-27-0) We say that $S \subseteq \mathbb{R}^m$ is a null set if it has zero Lebesgue measure. Obviously, any hyperplanes in R *^m* are null sets. We will also need the following simple lemma:

Lemma 14. Suppose that $m \in \mathbb{Z}_+$ and V is a subspace of \mathbb{R}^m . Then for any set of vectors $u_i \in \mathbb{R}^m$, $i =$ 1, 2, \cdots , *n* that does not lie in V , there must exists $v \in \mathbb{R}^m$ such that $u_i^{\top} v \neq 0, \forall i \in [n]$ but $v \in V^{\perp}$, where V^{\perp} *is the orthogonal space of* V *.*

Proof. Let w_i be the orthogonal projection of u_i onto V^{\perp} . Since $u_i \notin V$, we know that $w_i \neq 0$. The solution space of each equation $w_i^{\top} v = 0$ in V^{\perp} must then be a proper subspace of V^{\perp} . Equipped with the Lebesgue measure, all these spaces are null sets in V^{\perp} , so one can always choose a $v \in V^{\perp}$ that does not lie in any of these solution spaces. Such *v* satisfies all the requirements. \Box

We choose G to be the graph with $i \to j$ for $\forall 1 \leqslant i < j \leqslant d$, so that G has $\frac{d(d-1)}{2}$ edges. Suppose that $i_0 \in [d]$ satisfies $s_i \leqslant \left| \text{pa}_{\mathcal{G}}(i) \right| - 1$, then we must have $i_0 \geqslant 2$, so there is an edge $1 \to i_0$ in $\mathcal{G},$ Let \hat{G} be the resulting graph obtained via removing the edge $1 \to i_0$ in G , then G and \hat{G} are clearly non-isomorphic.

Note that the *i*-th row of B_k can be written as $\omega_{k,i,i}^{-\frac{1}{2}}(e_i-(A_k)_i)$. Let's choose an lower-triangular matrix \bm{T} = $(t_{ij})_{i,j=1}^d \in \mathbb{R}^{d \times d}$ with columns $t_i, i \in [d]$ such that the following holds:

$$
(e_i - (A_k)_i)^{\top} t_j = \begin{cases} = 0, & \forall k \in \{1\} \cup S_{i_0}, j = 1 \text{ and } i = i_0 \\ > 0, & \forall i = j \text{ and } k \in \{1\} \cup S_i \\ \neq 0, & \forall \text{ remaining } (i, j, k) \in \{k = 1, j < i\} \cup \{k \ge 2, i = i_k, j < i\} \end{cases} \tag{13}
$$

and

$$
t_{ii} \neq 0, \quad \forall i \in [d]. \tag{14}
$$

We now show that: except from a null set in the weight space, such *T* can always be chosen. To see why this is the case, we first consider all the constraints on t_1 :

$$
(e_i - (A_k)_i)^{\top} t_1 =\begin{cases} = 0, & \forall k \in \{1\} \cup S_{i_0} \text{ and } i = i_0 \\ > 0, & \forall i = 1 \text{ and } k \in \{1\} \cup S_i \\ \neq 0, & \forall \text{ remaining } (i, k) \in \{k = 1, i > 1\} \cup \{k \ge 2, i = i_k > 1\} \end{cases}
$$
(15)

Now let $V = \text{span}\langle e_i - (A_k)_i : k \in \{1\} \cup S_{i_0} \text{ and } i = i_0 \rangle$ and R be the set of pairs (i, k) specified in the second and third row of [\(15\).](#page-28-0) For $\forall (i, k)$, let $w_k(i)$ be the weight vector of node *i* in the environment *k*, *i.e.*, the vector of nonzero entries in $(A_k)_i$. Then for ∀(*i*, *k*) ∈ *R*, the following set (as a subset of the weight space)

$$
\bigcup_{k^* \in \{1\} \cup S_{i_0}} \{e_{i_0} - w_{k^*}(i_0) \in \text{span}\,\langle e_i - w_k(i), e_{i_0} - w_{k'}(i_0) : k' \in \{1\} \cup S_{i_0} \setminus \{k^*\}\rangle\}
$$
(16)

must be a null set. Thus

$$
E = \bigcup_{(i,k)\in R} \bigcup_{k^* \in \{1\} \cup S_{i_0}} \{e_{i_0} - w_{k^*}(i_0) \in \text{span}\, \langle e_i - w_{k^*}(i), e_{i_0} - w_{k'}(i_0) : k' \in \{1\} \cup S_{i_0} \setminus \{k^*\}\rangle\} \tag{17}
$$

is also a null set. For any weights that are not in *E*, we necessarily have

$$
e_i - w_k(i) \notin \text{span}\langle e_i - (A_k)_i : k \in \{1\} \cup S_{i_0} \text{ and } i = i_0 \rangle = V, \quad (i, k) \in R.
$$

Let $U = \{e_i - w_k(i) : (i, k) \in R\}$, then we can apply [Lemma 14](#page-27-1) to deduce that there exists t_1 such that

$$
(\mathbf{e}_i - (\mathbf{A}_k)_i)^{\top} \mathbf{t}_1 = \begin{cases} = 0, & \forall k \in \{1\} \cup S_{i_0} \text{ and } i = i_0 \\ \neq 0, & \forall \text{ remaining } (i, k) \in \{k = 1\} \cup \{k \ge 2, i = i_k\} \end{cases}
$$
(18)

Note that the only difference between [\(18\)](#page-28-1) and [\(15\)](#page-28-0) is that the latter one further requires that

$$
\left(e_1 - (A_k)_1\right)^\top t_1 > 0, \quad \forall k \in \{1\} \cup S_i.
$$

while the former only guarantees that these terms are nonzero. However, recall that $(A_k)_{ij} \neq 0 \Rightarrow j \in$ $pa_{\mathcal{G}}(i) \Rightarrow j \leq i$, so the above essentially says that $t_{11} > 0$. This can be easily guaranteed by replacing the solution t_1 we obtained satisfying (18) with $-t_1$ if needed.

Assuming that the weights do not lie in the null set E we have shown that t_1 can always be chosen to satisfy all constraints imposed on it. We now proceed to choose the remaining entries of *T*. The remaining entries in t_1 can be chosen arbitrarily. For $t_j, j \geq 1,$ we note that the remaining constraints in [\(13\)](#page-26-2) that need to be satisfied consist of the "nonzero" part and the "positivity" part. The positivity constrains can always be satisfied by choosing a sufficiently large *tjj* for *j* > 1.

After choosing the *tj* 's satisfying the positivity constraints, the nonzero constraints along with [\(14\)](#page-28-2) are easy to fulfill by slightly perturbing t_j if they are violated; since each of these constraints are only violated in a zero-measure set of the weight space. Hence, we have shown that except a null set *E* in the weight space, there always exists some *T* satisfying [\(13\).](#page-26-2) Such *T* must be invertible since it is lower-triangular and its diagonal entries are nonzero. Now let \hat{H} = $T^{-1}H$ and $\hat{\mathbf{\Omega}}_k$ be the diagonal matrix with entries $\hat{\omega}_{k,i,i} = t_{ii}^{-2} \cdot \omega_{k,i,i}, i \in [d]$ and

$$
\hat{A}_k = \mathbf{I} - \hat{\Omega}_k^{\frac{1}{2}} \Omega_k^{-\frac{1}{2}} (\mathbf{I} - A_k) \mathbf{T}.
$$
\n(19)

First since T is invertible and H has full rank, \hat{H} must also have full row rank. Second,

$$
(\hat{\boldsymbol{A}}_k)_{ij} = \begin{cases} 1 - \hat{\omega}_{k,i,i}^{\frac{1}{2}} \omega_{k,i,i}^{-\frac{1}{2}} t_{ii} = 0 & \text{if } j = i \\ -\hat{\omega}_{k,i,i}^{\frac{1}{2}} \omega_{k,i,i}^{-\frac{1}{2}} (e_i - (\boldsymbol{A}_k)_i)^\top \boldsymbol{t}_j = 0 & \text{if } j > i \\ -\hat{\omega}_{k,i,i}^{\frac{1}{2}} \omega_{k,i,i}^{-\frac{1}{2}} (e_i - (\boldsymbol{A}_k)_i)^\top \boldsymbol{t}_j & \text{if } j < i. \end{cases}
$$

where we again recall that both A_k and \pmb{T} are lower-triangular. From [\(13\)](#page-26-2) we can see that

- When $i = i_0$ and $j = 1$, we have
	- **−** $(\hat{A}_k)_{i_0,1}$ = 0 if $k \in \{1\} \cup S_{i_0}$, and **−** $(\hat{A}_k)_{i_0,1} = (\hat{A}_1)_{i_0,1} = 0$ if $k \notin \{1\} \cup S_{i_0}$, by definition of S_{i_0} and [Assumption 8.](#page-26-1)
- When $i > j$ and $(i, j) \neq (i_0, 1)$, we have
	- $(\hat{A}_k)_{ij} \neq 0$ if $k = 1$ or $i = i_k$, which directly follows from [\(13\),](#page-26-2) and
	- $(\hat{\boldsymbol{A}}_k)_{ij} = (\hat{\boldsymbol{A}}_1)_{ij} \neq 0$, by [Assumption 8.](#page-26-1)

To summarize, for each k , $(A_k)_{ij} \neq 0 \Leftrightarrow j \in \text{pa}_{\mathcal{G}}(i)$ and $(i, j) \neq (i_0, 1)$.

Finally, let $\hat{\boldsymbol{w}}_k(i)$ be the weight vector of node *i* in environment *k* in the hypothetical model *i.e.*, the vector of nonzero entries in $(A_k)_i$, and T_S be the submatrix of T by selecting the rows and columns in the index set *S*, then by [\(19\)](#page-28-3) we have that

$$
\hat{\omega}_{k,i,i} = t_{ii}^{-2} \cdot \omega_{k,i,i}, \quad \hat{\omega}_{k,i,i}^{\frac{1}{2}} \omega_{k,i,i}^{-\frac{1}{2}} \boldsymbol{w}_k(i) \boldsymbol{T}_{\text{pa}_{\mathcal{G}}(i)} = \begin{cases} \hat{\boldsymbol{w}}_k(i) & \text{if } i \neq i_0 \\ [0, \hat{\boldsymbol{w}}_k(i)] & \text{if } i = i_0 \end{cases}
$$
(20)

By our assumption, for $\forall k \geqslant 2$, $i \neq i_k \Rightarrow w_k(i) = w_1(i)$ and $\omega_{k,i,i} = \omega_{1,i,i}$. Thus [\(20\)](#page-29-1) imply that $∀k ≥ 2$, $i ≠ i_k ⇒ ω_k(i) = ψ_1(i)$ and $ω_{k,i,i} = ω_{1,i,i}$. In other words, a single-node intervention on node *i^k* in environment *k* in the ground-truth model corresponds to a single-node intervention on node *i^k* in environment *k* in the hypothetical model, thereby completing the proof.

H.3 Proof of [Theorem 4](#page-7-0)

We first prove two lemmas.

Lemma 15.
$$
\forall i \in [d]
$$
, we have span $\langle (\mathbf{M}_k)_i : k \in [K] \rangle = \text{span} \langle \mathbf{h}_j : j \in \overline{\text{pa}}_G(i) \rangle$.

Proof. Since $(\bm{M}_k)_i$ = $(\bm{B}_k)_i\bm{H}$, and $(\bm{B}_k)_{ij}\neq 0\Leftrightarrow j\in \overline{\mathrm{pa}}_{\mathcal{G}}(i)$, we can see that $(\bm{M}_k)_i\in\mathrm{span}\left<\bm{h}_j:j\in\overline{\mathrm{pa}}_{\mathcal{G}}(i)\right>$. On the other hand, since H is invertible, by [Assumption 5](#page-5-1) we have $\dim \mathrm{span}\bra{({\bm M}_k)_i:k\in[K]}$ = dim span $\langle (\mathcal{B}_k)_i : k \in [K] \rangle = \Big|\overline{\text{pa}}_{\mathcal{G}}(i)\Big|$. Thus we must have span $\big\langle (\mathcal{M}_k)_i : k \in [K] \big\rangle$ = span $\Big\langle h_j : j \in \overline{\text{pa}}_{\mathcal{G}}(i) \Big\rangle$.

 $\bf{Lemma 16.}$ $\,$ $\it Let$ $\hat{\cal S}$ $\it be$ an ancestral set of graph ${\cal G}$ and $\hat{\bm V}_k$ = ${\rm span}\left\langle ({\bm M}_k)_s: s\in \hat{\cal S} \right\rangle, k\in [K].$ $\,$ $\it Then$ we have $V_1 = V_2 = \cdots = V_K = \text{span}\left\langle h_s : s \in \hat{S} \right\rangle$ *.*

Proof. Recall that M_k = B_kH , so for $\forall s \in \hat{S}$, the *s*-th row of M_k can be written as

$$
(\boldsymbol{M}_{k})_{s} = \sum_{t=1}^{d} (\boldsymbol{B}_{k})_{st} \boldsymbol{h}_{t} = \sum_{t \in \overline{\text{pa}}_{\mathcal{G}}(s)} (\boldsymbol{B}_{k})_{st} \boldsymbol{h}_{t} \in \text{span} \left\langle \boldsymbol{h}_{s} : s \in \hat{S} \right\rangle
$$
 (21)

where the last equation is because \hat{S} is ancestral $\Rightarrow \overline{pa}_G(s) \subseteq \hat{S}$. Thus, for $\forall k \in [K]$, $\hat{V}_k =$ $\big\langle (\bm{M}_k)_s : s \in \hat{S} \big\rangle \ \subseteq \ \mathrm{span} \ \Big\langle \bm{h}_s : s \in \hat{S} \Big\rangle. \ \ \mathrm{On \ \ the \ \ other \ \ hand, \ \ recall \ that \ both \ \bm{B}_k \ \ and \ \bm{H} \ \ have \ \ }$ full rank, so \pmb{M}_k has full row rank as well, which implies that $\dim \pmb{V}_k$ = |S| = $\dim \text{span}\left\langle \pmb{h}_s: s \in \hat{S}\right\rangle$. Hence, \boldsymbol{V}_k = span $\langle \boldsymbol{h}_s : s \in \hat{S} \rangle$, $\forall k \in [K]$. \Box

The following two propositions show that our algorithm always maintain an ancestral set, recursively adds a new node into the set and correctly identifies its parents.

Proposition 3 [\(Proposition 1](#page-7-5) restated)**.** *The following two propositions hold for [Algorithm 3:](#page-8-0)*

- $\text{ans}_G(i) \subseteq S \Leftrightarrow$ the *i* f condition in line 8 of [Algorithm 3](#page-8-0) is fulfilled;
- the set S maintained in *[Algorithm 3](#page-8-0)* is always an ancestral set, in the sense that $j \in S \Rightarrow \text{ans}_G(j) \subseteq S$.

Proof. At the starting point, we have *S* = ∅ which is obviously an ancestral set. Now suppose that after the ℓ -th iteration, $S = \{s_1, s_2, \cdots, s_\ell\}$ is an ancestral set. In the following, we show that ans $g(i) \subseteq S \Leftrightarrow$ the if condition in line 8 is fulfilled. This would immediately imply that there always exists a node *i* that can be added into *S* in the $(\ell + 1)$ -th iteration, and that after adding *i*, *S* is still an ancestral set.

Suppose that $\mathrm{ans}_{\mathcal{G}}(i) \subseteq S$ for some $i \notin S$, by [Lemma 15](#page-29-2) we know that $(\bm{M}_k)_i \in \mathrm{span}\left\langle \bm{h}_j : j \in \overline{\mathrm{pa}}_{\mathcal{G}}(i) \right\rangle$, so there exists $\alpha_k \in \mathbb{R}$ such that $(\bm{M}_k)_i - \alpha_k \bm{h}_i \in \text{span}\left\langle \bm{h}_j : j \in \text{pa}_\mathcal{G}(i)\right\rangle$. Moreover, since $(\bm{M}_k)_i =$ $\sum_{j\in\overline{\text{pa}}_G(i)}(\mathcal{B}_k)_{jj}h_j,$ ($\mathcal{B}_k)_{ii}$ = $\omega_{k,i,i}^{-\frac{1}{2}}$ ≠ 0 and *H* has full row rank by assumption, we must have $(\bm{M}_k)_i \,\notin\, \mathrm{span}\left\langle \bm{h}_j : j \in \mathrm{pa}_\mathcal{G}(i) \right\rangle$ and so $\alpha_k \neq 0.$ Thus, we have by the linearity of the projection operator

$$
q_k := \text{proj}_{V^\perp_k} \left((\pmb{M}_k)_i \right) = \text{proj}_{V^\perp_k} \left((\pmb{M}_k)_i - \alpha_k \pmb{h}_i \right) + \text{proj}_{V^\perp_k} \left(\alpha_k \pmb{h}_i \right) = \alpha_k \text{proj}_{V^\perp_k} \left(\pmb{h}_i \right).
$$

Recall that all the V_k 's are the same and equal span $\langle h_s: s \in S \rangle$ by [Lemma 16.](#page-29-3) So dim span $\langle q_k: k \in [K] \rangle \leq$ 1. Since *H* has full row rank, we have $h_i \notin \text{span } \langle h_s : s \in S \rangle = V_k$, so that dim span $\langle q_k : k \in [K] \rangle = 1$ holds, which is exactly the if condition in line 8.

Conversely, suppose that there is an *i* ∉ *S* such that ans $g(i) \nsubseteq S$ but dim span $\langle q_k : k \in [K] \rangle = 1$ holds. Since *S* is ancestral, we know that there must be some $j \in pa_G(i)$ such that $j \notin S$. Since e_i and e_j both have support on the coordinates in $\overline{pa}_G(i)$, by [Assumption 5](#page-5-1) we know that span $\langle e_i, e_j \rangle \subseteq$ $\mathrm{span}\langle (\boldsymbol{B}_k)_i : k \in [K]\rangle,$ so that $\mathrm{span}\langle \boldsymbol{h}_i, \boldsymbol{h}_j\rangle$ = $\mathrm{span}\langle \boldsymbol{e}_i, \boldsymbol{e}_j\rangle \boldsymbol{H} \subseteq \mathrm{span}\langle (\boldsymbol{B}_k)_i : k \in [K]\rangle \boldsymbol{H}$ = $\mathrm{span}\langle (\boldsymbol{M}_k)_i : k \in [K]\rangle$ $k \in [K]$). Since dim span $\langle q_k : k \in [K] \rangle = 1$, there must exist some vector $u \in \mathbb{R}^n$ and $\alpha_i, \alpha_j \in \mathbb{R}$ such that $h_i - \alpha_i u, h_j - \alpha_j u \in V_k$ = $\text{span}\langle h_s : s \in S \rangle$. Since $i, j \notin S$ and H has full row rank, we can deduce that $h_i, h_j \notin \text{span}\langle h_s: s \in S\rangle$, and so both of α_i and α_j are non-zero. Hence $\alpha_j h_i - \alpha_i h_j \in \text{span}\langle h_s : s \in S \rangle$, which is impossible since we know that H has full row-rank. \Box

 ${\bf Proposition 4}$ [\(Proposition 2](#page-7-4) restated). *Given any ordered ancestral set S that contains* $\operatorname{pa}_{\mathcal G}(i)$ *for some* $i \notin S$ *, [Algorithm 2](#page-7-2) returns a set* $P_i \subseteq S$ *that is exactly* $\text{pa}_{\mathcal{G}}(i)$ *.*

Proof. As we have shown in [Proposition 1,](#page-7-5) for each possible input (*S*, *i*) to [Algorithm 2,](#page-7-2) both *S* and *S*∪{*i*} are ancestral sets, so that ans $_G(i) \subseteq S$. Similarly one can see that inside the set *S* := ${s_1, s_2, \cdots, s_m}$, all the ancestors of s_j are contained in ${s_1, s_2, \cdots, s_{j-1}}$. In the following, we show that $\forall m' \in \{0, \ldots, m\}, r_{m'} = \left| \overline{\text{pa}}_{\mathcal{G}}(i) - \left\{s_j : j \leq m' \right\} \right|$ (*).

By [Lemma 16](#page-29-3) we have W_1 = W_2 = \cdots = W_K = span $\left\langle h_{s_j}:j\leqslant m'\right\rangle$. Let t_1,t_2,\cdots,t_ℓ be elements of $\overline{\text{pa}}_{\mathcal{G}}(i)$ that are not in $\{s_j : j \leqslant m'\}$, then

$$
r_{m'} = \dim \operatorname{span} \langle p_k : k \in [K] \rangle = \dim \left(\operatorname{proj}_{\operatorname{span} \langle h_{j} : j \leq m'} \rangle^{\perp \operatorname{span} \langle (M_k)_i : k \in [K] \rangle} \right)
$$

=
$$
\dim \left(\operatorname{proj}_{\operatorname{span} \langle h_{j} : j \leq m'} \rangle^{\perp \operatorname{span} \langle h_j : j \in \overline{\text{pa}}_G(i) \rangle} \right) \quad \text{(by Lemma 15)}
$$

=
$$
\dim \left(\operatorname{proj}_{\operatorname{span} \langle h_{j} : j \leq m'} \rangle^{\perp \operatorname{span} \langle h_{l_1}, h_{l_2}, \cdots, h_{l_\ell} \rangle} \right)
$$

=
$$
\ell \quad \text{(by Lemma 5 and non-degeneracy of } H)
$$

which proves (*). From (*) it is easy to see that $m' \in \overline{pa}_G(i)$ (and thus in $pa_G(i)$ since $i \notin S$) if and only if $r_{m'} = r_{m'-1} - 1$. \Box

Now we conclude the proof of [Theorem 4.](#page-7-0) [Propositions 1](#page-7-5) and [2](#page-7-4) directly imply that [Algorithm 3](#page-8-0) is able to exactly recover the ground-truth causal graph G. It remains to show that Line 20 in [Algorithm 3](#page-8-0) produces the correct $\hat h_i$'s. By [Lemma 15](#page-29-2) we know that E_j = span $\big\langle h_\ell:\ell\in\overline{\mathrm{pa}}_{\mathcal{G}}(j)\big\rangle,$ so

$$
\bigcap_{j\in\overline{\text{ch}}_{\mathcal{G}}(i)} E_j = \bigcap_{j\in\overline{\text{ch}}_{\mathcal{G}}(i)} \text{span}\left\langle h_\ell : \ell \in \overline{\text{pa}}_{\mathcal{G}}(j)\right\rangle = \text{span}\left\langle h_\ell : \ell \in \overline{\text{sur}}_{\mathcal{G}}(i)\right\rangle.
$$

where the last step holds because H has full row rank and $\cap_{j\in\overline{\ch}_G(i)}\overline{\mathrm{pa}}_\mathcal{G}(j)$ = $\overline{\mathrm{sur}}_\mathcal{G}(i)$ by definition. Hence, each \hat{h}_i is a linear combination of $h_\ell, \ell \in \overline{\mathrm{sur}}_\mathcal{G}(i)$, completing the proof.

I. Omitted Proofs from [Section 7](#page-10-3)

I.1 Proof of [Lemma 2](#page-10-4)

Let $w_k(i) \in \mathbb{R}^{\big|pa_{\mathcal{G}}(i)\big|}$ be the vector obtained by removing all zero entries in the *i*-th row of A_k and $\omega_{k,i,i}$ be the *i*-th diagonal entry in Ω_k , then for the *k*-th environment we have $z_i = w_k(i)^\top z_{\text{pa}_{\mathcal{G}}(i)}$ + $\omega_{k,i,i}^{\frac{1}{2}}$ ε_i , so that

$$
\hat{p}_k\left(\boldsymbol{z}_i\mid \boldsymbol{z}_{\text{pa}_{\mathcal{G}}(i)}\right) = \boldsymbol{\omega}^{-\frac{1}{2}}_{k,i,i} p_{\boldsymbol{\epsilon}_i}\left(\boldsymbol{\omega}^{-\frac{1}{2}}_{k,i,i}(\boldsymbol{z}_i-\langle \boldsymbol{w}_k(i), \boldsymbol{z}_{\text{pa}_{\mathcal{G}}(i)}\rangle)\right)
$$

where $p_{\boldsymbol{\varepsilon}_i}(\cdot)$ is the density of $\boldsymbol{\varepsilon}_i$. As a result, we have

$$
\nabla \frac{\hat{p}_1}{\hat{p}_k} \left(\mathbf{z}_i \mid \mathbf{z}_{\text{pa}_{\mathcal{G}}(i)} \right) = \frac{\hat{p}_1}{\hat{p}_k} \left(\mathbf{z}_i \mid \mathbf{z}_{\text{pa}_{\mathcal{G}}(i)} \right) \cdot \nabla \log \frac{\hat{p}_1}{\hat{p}_k} \left(\mathbf{z}_i \mid \mathbf{z}_{\text{pa}_{\mathcal{G}}(i)} \right)
$$

$$
= \frac{\hat{p}_1}{\hat{p}_k} \left(\mathbf{z}_i \mid \mathbf{z}_{\text{pa}_{\mathcal{G}}(i)} \right) \cdot \left[c_{i1} (1, -\mathbf{w}_1(i)) - c_{ik} (1, -\mathbf{w}_k(i)) \right]
$$

where for convenience we use ∇ to denote the gradient with respect to all variables $z_{\overline{\mathrm{pa}}_{\mathcal{G}}(i)}$, and $c_{ik} = \omega_{k,i,i}^{-\frac{1}{2}} \cdot \frac{p'_{\epsilon_i}}{p_{\epsilon_i}}$ $\left(\omega^{-\frac{1}{2}}_{k,i,i}(z_i - \langle \boldsymbol{w}_k(i), z_{\text{pa}_{\mathcal{G}}(i)} \rangle \right)$ (we omit the dependency on \boldsymbol{z} for simplicity).

[Definition 5](#page-10-2) implies that span $\langle c_{i1}(1,-{\bm{\nu}}_1(i))-c_{ik}(1,-{\bm{\nu}}_k(i)):2\leqslant k\leqslant K\rangle\ =\ {\mathbb R}^{\left|{\mathrm{pa}}_{\mathcal{G}}(i)\right|+1},$ thus it $\text{holds that span}\langle (1,-\pmb{\nu}_k(i)): k\in [K]\rangle=\mathbb{R}^{\left\vert \mathbf{p}a_{\mathcal{G}}(i)\right\vert+1}$ as well. By definition of \pmb{B}_k , this immediately implies that dim $(\text{span}\langle (\mathcal{B}_k)_i : k \in [K] \rangle) = |\text{pa}_{\mathcal{G}}(i)| + 1$ as desired.

I.2 Proof of [Theorem 5](#page-10-0)

Define $\bm{\tau}:=\hat{\bm{h}}\circ\bm{h}^{-1}:\mathbb{R}^d\mapsto\mathbb{R}^d,$ then we have that $\pmb{\nu}=\bm{\tau}(\bm{z}).$ Since both \bm{h} and $\hat{\bm{h}}$ are diffeomorphisms by assumption, so is τ. To avoid confusion, in this section we use *z* (resp. *v*) to denote random variables while using $\hat{\boldsymbol{z}}$ (resp. $\hat{\boldsymbol{v}}$) to denote (deterministic) vectors.

Let $\mathfrak{E}_j = \left\{ E_k^{(j)} \right\}$ $\left\{ k^{(j)}_k : k \in [K_j]\right\}$ be the *j*-th collection of environments according to our assumption. We first prove the following lemma:

Lemma 17. $O_{\nu} = \tau(O_{z})$.

Proof. By the change of variable formula (Schwartz [1954\)](#page-13-23), for $\forall \hat{z} \in \mathbb{R}^d$ and $\forall E \in \mathfrak{E}$ we have $p_E(\hat{z}) = q_E(\hat{\nu}) |\det J_{\tau}(\hat{z})|$, where $\hat{\nu} = \tau(\hat{z})$. Since τ is a diffeomorphism, we must have $|\det J_{\tau}(\hat{z})| \neq 0$ so $\hat{z} \in \mathcal{O}_z \Leftrightarrow \hat{\mathbf{v}} = \tau(\hat{z}) \in \mathcal{O}_{\mathbf{v}}$, concluding the proof.

Lemma 18. Let $\hat{\mathbf{z}} \in \mathbf{O}_{\mathbf{z}}$. For $\forall j \in [d]$ and $2 \leqslant k \leqslant K_j$, we have

$$
\frac{F_{j}^{(j)}}{F_{j}^{(j)}}\left(\hat{\mathbf{z}}_{j}\mid\hat{\mathbf{z}}_{\text{pa}_{\mathcal{G}}(j)}\right)=\frac{q_{j}^{(j)}}{q_{j}^{(j)}}\left(\hat{\mathbf{v}}_{j}\mid\hat{\mathbf{v}}_{\text{pa}_{\hat{\mathcal{G}}}(j)}\right),\tag{22}
$$

where $\hat{v} = \tau(\hat{z}) \in O_{\nu}$ *.*

Proof. Since $v = \tau(z)$, by the change-of-measure formula (Schwartz [1954\)](#page-13-23) we have that for $\forall \hat{z} \in O_z$,

$$
\prod_{i=1}^{d} p_i^{E} \left(\hat{\mathbf{z}}_i \mid \hat{\mathbf{z}}_{\text{pa}_{\mathcal{G}}(i)} \right) = p_E(\hat{\mathbf{z}}) = q_E(\hat{\mathbf{v}}) \left| \det \mathbf{J}_{\boldsymbol{\tau}}(\hat{\mathbf{z}}) \right| = \prod_{i=1}^{d} q_i^{E} \left(\boldsymbol{\tau}_i(\hat{\mathbf{z}}) \mid \boldsymbol{\tau}_{\text{pa}_{\hat{\mathcal{G}}}(i)}(\hat{\mathbf{z}}) \right) \left| \det \mathbf{J}_{\boldsymbol{\tau}}(\hat{\mathbf{z}}) \right| \tag{23}
$$

for all $E \in \mathfrak{E}_j$, where $\hat{\mathbf{v}} = \boldsymbol{\tau}(\hat{\mathbf{z}})$. By Assumption (*ii*) and [Definition 2,](#page-2-3) we know that $p_i^{E^1_k} = p_i^{E^{(1)}_1} \Leftrightarrow i \neq 1$ and $q_i^{E^1_k} = q_i^{E^{(1)}_1} \Leftrightarrow i \neq 1$ for all $k > 1$. Thus, we have that

$$
\prod_{i=1}^d \frac{p_i^{E_k^{(j)}} \left(\hat{\mathbf{z}}_i \mid \hat{\mathbf{z}}_{\text{pa}_G(i)}\right)}{p_i^{E_i^{(j)}} \left(\hat{\mathbf{z}}_i \mid \hat{\mathbf{z}}_{\text{pa}_G(i)}\right)} = \frac{p_j^{E_k^{(j)}}}{p_j^{E_1^{(j)}}} (\hat{\mathbf{z}}_j \mid \hat{\mathbf{z}}_{\overline{\text{pa}}_G(j)})
$$

and

$$
\prod_{i=1}^d\frac{q_i^{E_{\hat{h}}^{(j)}}\left(\hat{\nu}_i\mid\hat{\nu}_{\text{pa}_{\hat{\mathcal{G}}}(i)}\right)}{q_i^{E_{\hat{h}}^{(j)}}\left(\hat{\nu}_i\mid\hat{\nu}_{\text{pa}_{\hat{\mathcal{G}}}(i)}\right)}=\frac{q_j^{E_{\hat{h}}^{(j)}}}{q_j^{E_{\hat{h}}^{(j)}}}\left(\hat{\nu}_j\mid\hat{\nu}_{\overline{\text{pa}}_G(j)}\right).
$$

Since the LHS of the above two equations are the same by [\(23\),](#page-32-1) the RHS must also be the same, concluding the proof. \Box

We assume WLOG that the vertices of G are labelled such that $i \rightarrow j \Rightarrow i \leq j$, and that $\pi(i) = i, \forall i \in [d]$. Also we can assume the nodes are fixed and only consider how they are connected, $i.e., \pi'(i) = i, \forall i \in [d].$ ^{[1](#page-33-0)}

Lemma 19. We have $(\tau(N_z))^{\circ} = (\tau^{-1}(N_v))^{\circ} = \emptyset$.

Proof. The result immediately follows from the assumption that $N_z^{\circ} = N_v^{\circ} = \emptyset$ and that τ is a diffeomorphism.

For any vertex set V, we use \mathcal{G}_V to denote its corresponding induced subgraph of \mathcal{G} . We first prove the following statements by induction on *j*:

- (1) $\forall i \neq j$, $i \in pa_{\mathcal{G}}(j) \Leftrightarrow i \in pa_{\mathcal{G}'}(j);$
- (2) $\forall j \in [d]$, there exists a continuously differentiable function ϕ_i such that $v_j = \phi_j\left(z_{\overline{\text{pa}}_G(j)}\right)$. Moreover, $\frac{\partial \Phi_j}{\partial z_j} \not\equiv 0$ (*i.e.*, not always zero).
- (3) $\forall j \in [d]$, there exists a continuously differentiable function γ_j such that $\nu_{\overline{\mathrm{pa}}_\mathcal{G}(j)} = \gamma_j(\bm{z}_{\overline{\mathrm{pa}}_\mathcal{G}(j)}).$

For $j = 1$, by assumption pa $_G(j) = \emptyset$. [Lemma 18](#page-32-2) implies that for any $\hat{\boldsymbol{z}} \in \mathbf{O_z}$,

$$
\frac{p_1^{(1)}}{p_1^{(1)}}(\hat{\mathbf{z}}_1) = \frac{q_1^{(1)}}{q_1^{(1)}}\left(\hat{\mathbf{\nu}}_1 \mid \hat{\mathbf{\nu}}_{\text{pa}_{\hat{\mathcal{G}}}(1)}\right), \forall 2 \leq k \leq K_1.
$$
\n(24)

Then for $\forall i \in \overline{\text{pa}}_{\hat{G}}(1)$, taking the partial derivative w.r.t v_i gives

$$
\frac{\partial}{\partial \hat{\mathbf{v}}_i}\frac{q_1^{(1)}}{E_1^{(1)}}\left(\hat{\mathbf{v}}_1 \mid \hat{\mathbf{v}}_{\text{pa}_{\hat{\mathcal{G}}}(1)}\right)=\left(\frac{p_1^{(1)}}{p_1^{(1)}}\right)'(\hat{\mathbf{z}}_1)\cdot\frac{\partial \hat{\mathbf{z}}_1}{\partial \hat{\mathbf{v}}_i}\Rightarrow \nabla_{\mathbf{v}_{\overline{\text{pa}}_{\hat{\mathcal{G}}}(1)}}\frac{q_1^{(1)}}{q_1^{(1)}}\left(\hat{\mathbf{v}}_1 \mid \hat{\mathbf{v}}_{\text{pa}_{\hat{\mathcal{G}}}(1)}\right)=\left(\frac{p_1^{(1)}}{p_1^{(1)}}\right)'(\hat{\mathbf{z}}_1)\cdot \nabla_{\mathbf{v}_{\overline{\text{pa}}_{\hat{\mathcal{G}}}(1)}}\hat{\mathbf{z}}_1.
$$

Thus,

$$
\operatorname{rank}\left[\nabla_{\nu_{\overline{\text{pa}}}_{\hat{\mathcal{G}}}(1)}\frac{q_1^{(1)}}{q_1^{(1)}}\left(\hat{\nu}_1\mid\hat{\nu}_{\text{pa}_{\hat{\mathcal{G}}}(1)}\right):2\leq k\leq K_1\right]\leq 1.
$$

Note that the above inequality holds for $\forall \hat{v} \in O_{\nu}$. If $pa_{\hat{G}}(1) \neq \emptyset$, then this would contradict the non-degeneracy assumption *(iii)* which implies that the above matrix should have rank ≥ 2 at some point $\hat{v} \in O_{\nu}$. Hence we must have pa_{$\hat{c}(1) = \emptyset$, implying that (1) holds for $j = 1$.}

Taking the derivative of both sides of [\(24\)](#page-32-3) w.r.t z_i , $i \geqslant 2$ implies that $\sqrt{ }$ \mathcal{L} $q_1^{E_k^{(1)}}$ $q_1^{E_1^{(1)}}$ \setminus $\overline{1}$ ′ $(\hat{\mathbf{v}}_1) \cdot \frac{\partial \hat{\mathbf{v}}_1}{\partial \hat{\mathbf{z}}_i} = 0.$ By our assumption *(iii)*, for $\forall \hat{\mathbf{v}} \in \mathbf{O}_{\mathbf{v}} \setminus \mathbf{N}_{\mathbf{v}}$, there exists $2 \leqslant k \leqslant K_1$ such that $\sqrt{ }$ \mathcal{L} $q_1^{E_k^{(1)}}$ $q_1^{E_1^{(1)}}$ \setminus $\overline{1}$ ′ $(\hat{\boldsymbol{\nu}}_1) \neq 0$,

and thus we have $\frac{\partial \hat{v}_1}{\partial \hat{z}_i} = 0, \forall \hat{z} \in \tau^{-1} \left(O_\nu \setminus N_\nu \right)$. Since τ is a diffeomorphism, we can deduce that $\tau^{-1}(O_\nu\setminus N_\nu)=O_z\setminus\tau^{-1}(N_\nu)$ and $(\tau^{-1}(N_\nu))^{\circ}=\emptyset$ by [Lemma 19.](#page-33-1) As a result, we actually have

^{1.} This is also WLOG because we now have groups of soft interventions where each group corresponds to a single node, so we can just relabel the node in \hat{G} that corresponds to the *i*-th group as node *i*.

 $\frac{\partial \hat{v}_1}{\partial \hat{z}_i}$ = 0, $\forall \hat{z} \in \mathbf{O_z}$. Hence in $\mathbf{O_z}$ there exists a continuous differentiable function ϕ_1 such that $v_1 = \phi_1(z_1)$, proving (2). Finally, (3) directly follows from (2) since $pa_G(1) = \emptyset$, concluding the proof for $j = 1$.

Now suppose that the statement holds up to $j - 1$, and we need to prove it for *j*. Again by [Lemma 18](#page-32-2) we have for $\forall \hat{z} \in O_z$ that

$$
\frac{p_j^{E_k^{\{j\}}}}{p_j^{E_1^{\{j\}}}}\left(\hat{\mathbf{z}}_j \mid \hat{\mathbf{z}}_{\text{pa}_{\mathcal{G}}(j)}\right) = \frac{q_j^{E_k^{\{j\}}}}{q_j^{E_1^{\{j\}}}}\left(\hat{\mathbf{v}}_j \mid \hat{\mathbf{v}}_{\text{pa}_{\mathcal{\hat{G}}}(j)}\right), \quad \forall 2 \leq k \leq K_j.
$$
\n(25)

For all $i \notin \overline{\text{pa}}_{\mathcal{G}}(j)$, taking partial derivative w.r.t. \boldsymbol{z}_i gives

$$
0=\sum_{\ell\in\overline{\text{pa}}_{\hat{\mathcal{G}}}(j)}\frac{\partial}{\partial \hat{\nu}_{\ell}}\frac{q_j^{E_k^{(j)}}}{E_1^{E_1^{(j)}}}\left(\hat{\nu}_j\mid\hat{\nu}_{\text{pa}_{\hat{\mathcal{G}}}(j)}\right)\cdot\frac{\partial \hat{\nu}_{\ell}}{\partial \hat{z}_i},\quad\forall 2\leq k\leq K_j,
$$

i.e.,

$$
\left[\nabla_{\nu_{\overline{\text{pa}}}_{\hat{\mathcal{G}}}}(\hat{y})\frac{q_j^{E_k^{(j)}}}{q_j^{E_1^{(j)}}}\left(\hat{\nu}_j\mid\hat{\nu}_{\text{pa}_{\hat{\mathcal{G}}}(j)}\right):2\leq k\leq K_j\right]^\top\frac{\partial\hat{\nu}_{\overline{\text{pa}}_{\hat{\mathcal{G}}}}(j)}{\partial\hat{z}_i}=0.
$$

Similar to the *j* = 1 case, by assumption *(iii)*, we know that the above corfficient matrix has full $\text{row rank for }\forall \hat{\nu} \in \mathbf{O}_{\mathbf{\nu}} \setminus \mathbf{N}_{\mathbf{\nu}}, \text{ so for }\forall \boldsymbol{z} \in \boldsymbol{\tau}^{-1}\left(\mathbf{O}_{\mathbf{\nu}} \setminus \mathbf{N}_{\mathbf{\nu}}\right) = \mathbf{O}_{\boldsymbol{z}} \setminus \boldsymbol{\tau}^{-1}\left(\mathbf{N}_{\mathbf{\nu}}\right), \text{ we have } \frac{\mathfrak{d} \hat{\nu}_{\overline{\text{pa}}_{\hat{\mathcal{G}}}(\hat{\nu})}}{\mathfrak{d} \hat{z}_{\hat{\nu}}}$ $\frac{1}{\partial \hat{z}_i}$ = 0. Since $(\tau^{-1}(N_{\nu}))^{\circ} = \emptyset$ by [Lemma 19,](#page-33-1) for all $\hat{\varkappa} \in N_{\varkappa}$ we can choose a sequence of points $\hat{\varkappa}^{(i)}$, $i = 1, 2, \cdots$ in O_z such that $\hat{z}^{(i)} \to \hat{z}$. Since τ is a diffeomorphism, its derivatives are continuous and we can deduce that $\frac{\partial \hat{v}_{\overline{\text{pa}}_{\hat{\mathcal{G}}}(j)}}{\partial \hat{z}_i}$ $rac{1}{\partial \hat{z}_i}$ ^{*i*} = lim_{$\ell \rightarrow +\infty$} $\partial \hat{v}_{\overline{p_2}}^{(\ell)}$ $\overline{\mathrm{pa}}_{\hat{\mathcal{G}}}$ (j) $\partial \hat{z}_i^{(\ell)}$ = 0. As a result, $\frac{\partial \hat{v}_{\overline{\text{pa}}_{\hat{\mathcal{G}}}(j)}}{\partial \hat{z}_i}$ $rac{F^2\mathcal{G}^{yy}}{\partial \hat{z}_i} = 0$ actually holds for all $z \in \mathbf{O}_z$. Hence,

there exists a continuous differentiable function γ_j such that $\nu_{\overline{\text{pa}}_{\hat{\mathcal{G}}}(j)} = \gamma_j \left(z_{\overline{\text{pa}}_{\mathcal{G}}(j)} \right)$.

By our assumption, $\text{pa}_{\mathcal{G}}(j) \subseteq [j-1]$. Suppose that $\text{pa}_{\hat{\mathcal{G}}}(j) \nsubseteq \{i : i \leq j\}$, let $\ell \in \text{pa}_{\hat{\mathcal{G}}}(j) \setminus \{i : i \leq j\}$, then by induction hypothesis, $\hat{\pmb{\nu}}_t = \pmb{\tau}_t(\hat{\pmb{z}}), \hat{\pmb{z}} \in \mathbf{O}_{\pmb{z}}, \, t=1,2,\cdots,j, \ell$ are all functions of $\hat{\pmb{z}}_1,\cdots,\hat{\pmb{z}}_j.$ Since τ is a diffeomorphism and O_z is the support of the distributions $p_E, E \in \mathfrak{E}$, we can deduce that the support of the latent variables $(v_t : t = 1, 2, \cdots, j, \ell)$ lie on a submanifold with dimension $\leq j$, which is impossible since ν is supported on the open set $O_\nu\subseteq \mathbb{R}^d$ by assumption *(i)*.

Hence, we must have $\text{pa}_{\hat{\mathcal{G}}}(j) \subseteq \{i : i < j\}$. Furthermore, if there exists $i \in \text{pa}_{\hat{\mathcal{G}}}(j)$ such that *i* ∉ pa_G(*j*), then the induction hypothesis implies that $\frac{\partial v_i}{\partial z_i}$ ≢ 0, but v_i is a function of $z_{\overline{\text{pa}}_G(j)}$ as previously derived, which is also a contradiction. Thus we actually have pa $_{\hat{G}}(j) \subseteq$ pa $_{G}(j)$.

In a completely symmetric manner, we can take the derivatives of [\(25\)](#page-33-2) w.r.t. $v_i, \forall i \in \overline{\text{pa}}_{\hat{\mathcal{G}}}(j)$ and obtain that pa $_{\mathcal{G}}(j)\subseteq$ pa $_{\hat{\mathcal{G}}}(j)$. Hence, pa $_{\hat{\mathcal{G}}}(j)$ = pa $_{\mathcal{G}}(j)$, completing the proof of (1) and (3) for the *j* case.

Finally, if $\frac{\partial v_j}{\partial z_j} \equiv 0$, then by (3) and the induction hypothesis, v_1, \cdots, v_j are all functions of $z_{[j-1]}$, which implies that (v_1, \dots, v_j) lies on a submanifold with dimension $\leq j-1$, again contradicting assumption *(i)*. Thus $\frac{\partial v_j}{\partial z_j} \neq 0$. This completes the proof of our inductive step.

To recap, we now know that

•
$$
\hat{G} = \hat{G}
$$
, and

• For $\forall i \in [d]$, there exists a function γ_i such that $\nu_{\overline{\text{pa}}_G(i)} = \gamma_i \left(z_{\overline{\text{pa}}_G(i)} \right)$.

It remains to show that for $\forall k \in pa_G(i) \setminus \text{sur}_G(i)$, Υ_i doesn't depend on z_k .

By definition, if $k \in pa_G(i) \setminus \text{sur}_G(i)$, we know that there exists $j \in ch_G(i)$ such that $j \notin ch_G(k)$. We have shown that v_i , as a component of $v_{\overline{pa}_G(j)}$, is a function of $z_{\overline{pa}_G(j)}$. By the choice of *k*, we have $k \notin \overline{\text{pa}}_{\mathcal{G}}(j)$, so that v_i does not depend on \boldsymbol{z}_k . The conclusion follows.

J. Omitted Proofs for [Theorem 3](#page-6-0) and [Theorem 6](#page-10-1)

In this section we provide detailed proofs of main ambiguity results.

 $\bf{Definition~12.}$ $\emph{We say that a matrix}\ \bm{M}\in\mathbb{R}^{d\times d}$ is effect-respecting for a causal graph \mathcal{G} , or $\bm{M}\in\mathcal{M}_{\rm sur}(\mathcal{G})_d$ if $M_{ij} \neq 0 \Leftrightarrow j \in \overline{\sup}_{\mathcal{G}}(i)$. We also write $\boldsymbol{M} \in \mathcal{M}_{\text{sur}}^0(\mathcal{G})$ if \boldsymbol{M} is invertible and $\boldsymbol{M}_{ij} \neq 0 \Rightarrow j \in \overline{\sup}_{\mathcal{G}}(i)$. *Finally, we write* $\mathbf{M} \in \overline{\mathcal{M}}_{\text{sur}}(\mathcal{G})$ *if* $\mathbf{M}_{ij} \neq 0 \Rightarrow j \in \overline{\text{sur}}_{\mathcal{G}}(i)$ *.*

 $\bf{Remark\ 1.}$ *By definition* $\mathcal{M}^0_{\rm sur}(G)$ *is the set of all matrices* \bf{M} *where* $\bf{M}_{ij}\neq 0, \forall j \notin \overline{\rm sur}_G(i)$ *, so it* can be identified as \mathbb{R}^{d+d_G} where $d_G = \sum_{i=1}^d |\text{sur}_G(i)|$. Equipped with the Lebesgue measure, we have $\mathcal{M}_{\text{sur}}(\mathcal{G})\subset\mathcal{M}_{\text{sur}}^0(\mathcal{G})\subset\overline{\mathcal{M}}_{\text{sur}}(\mathcal{G})$ and $\overline{\mathcal{M}}_{\text{sur}}(\mathcal{G})\backslash\mathcal{M}_{\text{sur}}(\mathcal{G})$ is a null set. In the remaining part of this section, *we will use measure-theoretic statement for* $\boldsymbol{M} \in \mathcal{M}_{\text{sur}}(\mathcal{G})$ *in the above sense.*

We first present a result that serves as a good starting point to understand why this is the case. It states that latent representations that are equivalent under \sim_sur are essentially generated from the same causal graph.

Proposition 5. Let *M* be an invertible matrix such that $M_{ij} \neq 0 \Rightarrow j \in \overline{\text{sur}}_G(i)$. Suppose that the l *atent variables* $\bm{z}\in\mathbb{R}^d$ are generated from any distributions $p_i\left(\bm{z}_i\mid\bm{z}_{\text{pa}_\mathcal{G}(i)}\right), i\in[d]$ with joint density $p(\boldsymbol{z})$ = $\prod_{i=1}^{d} p_i\left(\boldsymbol{z}_i \mid \boldsymbol{z}_{\text{pa}_{\mathcal{G}}(i)}\right)$, then the joint density of $\boldsymbol{\nu}$ = $\boldsymbol{M}\boldsymbol{z}$ can be written as $q(\boldsymbol{\nu})$ = $\prod_{i=1}^{d} q_i\left(\boldsymbol{\nu}_i \mid \boldsymbol{\nu}_{\text{pa}_{\mathcal{G}}(i)}\right)$ *for some density functions* $q_i, i \in [d].$

J.1 Proof of [Proposition 5](#page-35-2)

We first prove the following lemma:

Lemma 20. Let $M \in \mathcal{M}_{\text{sur}}^0(\mathcal{G})$ and latent variables $v = Mz$, then for $\forall i \in [d]$, there exists invertible m *atrices* \bm{M}_i and \bm{M}_i^- such that $\nu_{\text{pa}_G(i)} = \bm{M}_i^- \bm{z}_{\text{pa}_G(i)}$ and $\nu_{\overline{\text{pa}}_G(i)} = \bm{M}_i \bm{z}_{\overline{\text{pa}}_G(i)}$ *.*

Proof. $\forall j \in \overline{\text{pa}}_{\mathcal{G}}(i)$, we know that ν_j is a linear function of $\bm{z}_\ell, \ell \in \overline{\text{sur}}_{\mathcal{G}}(j)$. By [Lemma 8,](#page-21-4) we know that $\overline{\sup}_\mathcal{G}(j)\subseteq\overline{\mathrm{pa}}_\mathcal{G}(i)$, so each $v_j, j\in\overline{\mathrm{pa}}_\mathcal{G}(i)$ is a linear function of $z_{\overline{\mathrm{pa}}_\mathcal{G}(i)}.$ Thus we can write $v_{\overline{\text{pa}}_\mathcal{G}(i)}$ = $\bm{M}_i \bm{z}_{\overline{\text{pa}}_\mathcal{G}(i)}.$ In the following we argue that \bm{M}_i is invertible. Let π be a permutation on $\overline{\text{pa}}_\mathcal{G}(i)$ $\text{such that } k \in \text{pa}_\mathcal{G}(\ell) \Rightarrow \pi(k) \lt \pi(\ell)$ (such π can always be chosen since $\mathcal G$ is acyclic), then we can write

$$
\left(\hat{\boldsymbol{\nu}}_{\pi(j)}:j\in\overline{\text{pa}}_{\mathcal{G}}(i)\right)^{\top}=\tilde{\boldsymbol{M}}_{i}\left(\hat{\boldsymbol{z}}_{\pi(j)}:j\in\overline{\text{pa}}_{\mathcal{G}}(i)\right)^{\top}
$$
\n(26)

where $\tilde{\bm{M}}_i$ is an upper triangular matrix with non-zero diagonal entries by our choice of $\bm{M}.$ Since \bm{M}_i can be obtained from $\tilde{\bm{M}}_i$ be exchanging a few rows and columns, \bm{M}_i is invertible as well.

Similarly, using the fact that $\forall j\in{\rm pa}_{\cal G}(i)$, $\overline{\rm sur}_{\cal G}(j)\subseteq{\rm pa}_{\cal G}(i)$, we can prove the existence of an invertible matrix \boldsymbol{M}_{i}^{-} such that $\nu_{\text{pa}_{\mathcal{G}}(i)} = \boldsymbol{M}_{i}^{-} \boldsymbol{z}_{\text{pa}_{\mathcal{G}}(i)}$. \Box

Returning to the proof of [Proposition 5.](#page-35-2) Assume WLOG that the nodes of G are ordered in a way such that $i \in pa_{\mathcal{G}}(j) \Rightarrow i < j$, so that M is a lower-triangular matrix. The joint density of ν can be written as

$$
q(\mathbf{v}) = \prod_{i=1}^d q\left(\mathbf{v}_i \mid \mathbf{v}_1, \cdots, \mathbf{v}_{i-1}\right).
$$

Since $v = Mz$ and M is lower triangular and invertible (hence, with non-zero diagonals), we know that $(v_1, v_2, \dots, v_{i-1})$ is an invertible linear function of $(z_1, z_2, \dots, z_{i-1})$ and (v_1, v_2, \dots, v_i) is an invertible linear function of (z_1, z_2, \dots, z_i) . Let $\hat{\mathbf{v}} = \mathbf{M}\hat{\mathbf{z}} \in \mathbb{R}^d$, then we have

$$
q(\hat{\mathbf{v}}_i | \hat{\mathbf{v}}_1, \cdots, \hat{\mathbf{v}}_{i-1}) = \frac{q(\hat{\mathbf{v}}_1, \hat{\mathbf{v}}_2, \cdots, \hat{\mathbf{v}}_i)}{q(\hat{\mathbf{v}}_1, \hat{\mathbf{v}}_2, \cdots, \hat{\mathbf{v}}_{i-1})} = \frac{p(\hat{z}_1, \hat{z}_2, \cdots, \hat{z}_i) \det \hat{\mathbf{M}}_{1:i,1:i}}{p(\hat{z}_1, \hat{z}_2, \cdots, \hat{z}_{i-1}) \det \hat{\mathbf{M}}_{1:i-1,1:i-1}} \propto \frac{p(\hat{z}_1, \hat{z}_2, \cdots, \hat{z}_i)}{p(\hat{z}_1, \hat{z}_2, \cdots, \hat{z}_{i-1})} = p(\hat{z}_i | \hat{z}_1, \cdots, \hat{z}_{i-1}) = p_i(\hat{z}_i | \hat{z}_{\text{pa}_\mathcal{G}(i)}),
$$

where $\hat{\bm{M}}_{1:i, i:i}$ denotes that top-left submatrix of $\hat{\bm{M}}$ of size $i\times i,$ and the last step follows from the causal Markov condition [\(Definition 1\)](#page-2-2). On the other hand, let $q_i\left(\hat{\bm{\nu}}_i\mid \hat{\bm{\nu}}_{\text{pa}_{\mathcal{G}}(i)}\right)$ be the conditional density of v_i on its parents at $\hat{v} \in \mathbb{R}^d$. For $\forall j \in \text{pa}_\mathcal{G}(i)$, from v = Mz we know that v_j is a linear function of $z_{\overline{\sup}}_G(j)$. By [Lemma 20](#page-35-3) we know that $\hat{\nu}_{\text{pa}_G(i)}$ is a linear function of $\hat{z}_{\text{pa}_G(i)}$ and $\hat{\nu}_{\overline{\text{pa}}_G(i)}$ is a linear function of $\hat{\boldsymbol{z}}_{\overline{\text{pa}}_{\mathcal{G}}(i)},$ so that

$$
q\left(\hat{\nu}_{\text{pa}_{\mathcal{G}}(i)}\right) \propto p\left(\hat{z}_{\text{pa}_{\mathcal{G}}(i)}\right) \quad \text{and} \quad q\left(\hat{\nu}_{\overline{\text{pa}}_{\mathcal{G}}(i)}\right) \propto p\left(\hat{z}_{\overline{\text{pa}}_{\mathcal{G}}(i)}\right)
$$

and

$$
q_i\left(\hat{\boldsymbol{\nu}}_i\mid\hat{\boldsymbol{\nu}}_{\text{pa}_{\mathcal{G}}}(i)\right)\propto\frac{p\left(\hat{\boldsymbol{z}}_{\overline{\text{pa}}_{\mathcal{G}}}(i)\right)}{p\left(\hat{\boldsymbol{z}}_{\text{pa}_{\mathcal{G}}}(i)\right)}=p_i\left(\hat{\boldsymbol{z}}_i\mid\hat{\boldsymbol{z}}_{\text{pa}_{\mathcal{G}}}(i)\right).
$$

Hence, we have $q_i\left(\hat{\nu}_i\mid \hat{\nu}_{\text{pa}_{\mathcal{G}}(i)}\right) \propto q\left(\hat{\nu}_i\mid \hat{\nu}_1,\cdots,\hat{\nu}_{i-1}\right)$, so that

$$
q(\hat{\boldsymbol{\nu}}) = \prod_{i=1}^d q_i \left(\hat{\boldsymbol{\nu}}_i \mid \hat{\boldsymbol{\nu}}_{\text{pa}_{\mathcal{G}}(i)} \right) \propto \prod_{i=1}^d q_i \left(\hat{\boldsymbol{\nu}}_i \mid \hat{\boldsymbol{\nu}}_{\text{pa}_{\mathcal{G}}(i)} \right).
$$

Since both sides integrate to 1, it turns out that they are equal, as desired.

J.2 Formal version and proof of [Theorem 3:](#page-6-0) the linear case

Theorem 9 (Counterpart to [Theorem 1\)](#page-5-0)**.** *For any causal model* (*H*, G) *and any set of environments* **€** = ${E_k : k ∈ [K]}$, suppose that we have observations ${P_X^E}_{E ∈ \mathfrak{E}}$ satisfying [Assumption 1:](#page-2-1)

$$
\forall k \in [K], \quad z = A_k z + \Omega_k^{\frac{1}{2}} \epsilon, \quad x = H^{\dagger} z
$$

such that

- *(i)* the unmixing matrix $\boldsymbol{H} \in \mathbb{R}^{d \times n}$ has full row rank;
- (ii) ∀ $k \in [K]$ and $i,j \in [d]$, $(A_k)_{ij} \neq 0 \Leftrightarrow j \in \text{pa}_{\mathcal{G}}(i)$ and $\bm{\Omega}_k$ is a diagonal matrix with positive entries;

(iii)
$$
\left\{ \mathbf{B}_k = \mathbf{\Omega}_k^{-\frac{1}{2}} (\mathbf{I} - \mathbf{A}_k) \right\}_{k=1}^K
$$
 are node level non-degenerate in the sense of Assumption 5,

then there must exist a candidate solution (*H*ˆ , G) *and a hypothetical data generating process*

$$
\forall k \in [K], \quad \mathbf{v} = \hat{A}_k \mathbf{v} + \hat{\Omega}_k^{\frac{1}{2}} \mathbf{\varepsilon}, \quad \mathbf{x} = \hat{H}^{\dagger} \mathbf{v}
$$

such that

- (i') the unmixing matrix $\hat{\boldsymbol{H}} \in \mathbb{R}^{d \times n}$ has full row rank;
- (ii') $\forall k \in [K]$ and $i, j \in [d]$, $(\hat{A}_k)_{ij} \neq 0 \Leftrightarrow j \in \text{pa}_{\mathcal{G}}(i)$ and $\hat{\Omega}_k$ is a diagonal matrix with positive entries; $(iii') \left\{ \hat{B}_k = \hat{\Omega}_k^{-\frac{1}{2}} (I - \hat{A}_k) \right\}^K$ *k*=1 *are node level non-degenerate in the sense of [Assumption 5,](#page-5-1)*

but

$$
\frac{\partial v_i}{\partial z_j} \neq 0, \quad \forall j \in \overline{\text{sur}}_{\mathcal{G}}(i).
$$

Finally, if we additionally assume that

(iii) the environments are groups of single-node interventions: there exists a partition $\mathfrak{E} = \bigcup_{i=1}^d \mathfrak{E}_i$ such that $\mathcal{I}^{\mathfrak{E}_i}_{\boldsymbol{z}}$ = {*i*} *(see [Definition 2\)](#page-2-3)*,

then we can guarantee the existence of (\hat{H} , \hat{G}) *and weight matrices which, besides the properties listed above, also satisfy*

(*iii'*) *for the same partition* $\mathfrak{E} = \bigcup_{i=1}^{d} \mathfrak{E}_i$ *, we have* $\mathcal{I}_{\mathfrak{v}}^{\mathfrak{E}_i} = \{i\}$ *.*

In other words, additionally assuming that the environments are from single-node interventions does not resolve the ambiguity.

Remark 2. *Compared with our identifiability guarantee [Theorem 1,](#page-5-0) [Theorem 9](#page-36-0) actually demonstrates a stronger form of impossibility. Specifically, it states that the SNA cannot be resolved even if both the ground-truth causal graph and the noise variables are known.*

We define

$$
v = Mz \tag{27}
$$

where *M* is an effect-respecting matrix. At this point we do not make any other restrictions on *M*, but we will specify the appropriate choise of *M* later.

By assumption, the latent variables in the *k*-th environment are generated by

$$
z = A_k z + \Omega_k^{\frac{1}{2}} \epsilon,
$$

then $\nu = \bm{M}(\bm{I}-\bm{A}_k)^{-1}\bm{\Omega}_k^{\frac{1}{2}}$ e. Let $\hat{\bm{\Omega}}_k$ be the diagonal matrix with entries $\bm{M}_{ii}^2\cdot(\bm{\Omega}_k)_{ii}, i\in [d]$ and $\hat{\bm{A}}_k = \bm{I} - \hat{\bm{\Omega}}_k^{\frac{1}{2}} \bm{\Omega}_k^{-\frac{1}{2}} (\bm{I} - \bm{A}_k) \bm{M}^{-1}$, then $\bm{\nu} = \hat{\bm{A}}_k \bm{\nu} + \hat{\bm{\Omega}}_k^{\frac{1}{2}} \bm{\epsilon}$. Note that the choice of $\hat{\bm{\Omega}}_k$ here is to that the diagonal entries of $\hat{\bm{A}}_k$ are zero, as we show below. It remains to show that: for almost all *M* ∈ $\mathcal{M}^0_{\text{sur}}(\mathcal{G})$, it holds for $\forall k \in [K]$ that $(\hat{A}_k)_{ij}$ = 0 ⇔ $j \notin$ pa $_{\mathcal{G}}(i)$.

For the \Leftarrow direction, since $M \in \mathcal{M}^0_{\text{sur}}(\mathcal{G}), M^{-1} \in \mathcal{M}^0_{\text{sur}}(\mathcal{G})$ as well. Thus, $\forall j \notin \text{pa}_{\mathcal{G}}(i)$ we have

$$
\left[(I - A_k) \mathbf{M}^{-1} \right]_{ij} = \sum_{\ell=1}^d (I - A_k)_{i\ell} \cdot (\mathbf{M}^{-1})_{\ell j} = \sum_{\ell \in \overline{\text{pa}}_G(i) \cap \{\ell': j \in \overline{\text{sur}}_G(\ell')\}} (I - A_k)_{i\ell} \cdot (\mathbf{M}^{-1})_{\ell j}
$$

$$
= \begin{cases} 0 & \text{if } j \notin \overline{\text{pa}}_G(i) \\ (\mathbf{M}^{-1})_{ii} & \text{if } j = i \end{cases}
$$

where the last step holds because $\forall l \in [d]$, $l \in \overline{\text{pa}}_{\mathcal{G}}(i)$, $j \in \overline{\text{sur}}_{\mathcal{G}}(\ell) \Rightarrow j \in \overline{\text{pa}}_i$, and when $j = i$, the only such ℓ is $\ell = i$. Hence, we can see that our choice of \hat{A}_k satisfies

$$
\left(\hat{\boldsymbol{A}}_k\right)_{ij} = \begin{cases} 0 - 0 = 0 & \text{if } j \notin \overline{\text{pa}}_{\mathcal{G}}(i) \\ 1 - \hat{\omega}_{k,i,i}^{\frac{1}{2}} \omega_{k,i,i}^{-\frac{1}{2}} (\boldsymbol{M}^{-1})_{ii} = 0 & \text{if } j = i, \end{cases}
$$

so $\left(\hat{A}_k\right)$ $j_j \neq 0 \Rightarrow j \in \text{pa}_{\mathcal{G}}(i).$ Conversely, for $\forall j \in pa_{\mathcal{G}}(i)$,

$$
(\hat{\boldsymbol{A}}_k)_{ij} = 0 \Leftrightarrow \sum_{s \in \overline{\text{pa}}_{\mathcal{G}}(i)} (\boldsymbol{I} - \boldsymbol{A}_k)_{is} (\boldsymbol{M}^{-1})_{sj} = 0 \Leftrightarrow \sum_{s \in \overline{\text{pa}}_{\mathcal{G}}(i)} (-1)^s (\boldsymbol{I} - \boldsymbol{A}_k)_{is} \det \boldsymbol{M}_{sj}^{-} = 0
$$
 (28)

where \bm{M}^-_{sj} is the $(d-1)\times(d-1)$ matrix obtained by removing the *s*-th row and *j*-th column of $\bm{M},$ and the second step in the equation above follows from the fact that \bm{M}^{-1} = $\det(\bm{M})^{-1}\mathrm{adj}(\bm{M}),$ where adj (M) denotes the adjugate matrix of M whose (i,j) -th entry is $(-1)^{i+j}$ det $M^-_{ij}.$

[\(28\)](#page-37-0) holds if only if M takes values on a lower-dimensional algebraic manifold of its embedded space $\mathbb{R}^{d+d_\mathcal{G}}$ (see [Remark 1\)](#page-35-4). As a result, for almost every $\bm{M}\in\mathcal{M}^0_{\mathrm{sur}}(\mathcal{G})$, ν is generated from a linear causal model with graph G as defined in [\(3\).](#page-4-0) Moreover, let $\hat{\bm{B}}_k = \bm{B}_k \bm{M}^{-1}$, $k \in [K]$, so that $\bm{\epsilon} = \hat{\bm{B}}_k \bm{\nu}$ in the *k*-th environment. Then for all nodes $i \in [d]$ and $S ⊆ pa(i) ∪ {i}$, we have

$$
\dim \mathrm{span}\left\langle \left(\hat{\boldsymbol{B}_{k}}^{\top}\boldsymbol{e}_{i}\right)_{S} : k \in [K]\right\rangle = \dim \mathrm{span}\left\langle \boldsymbol{M}^{\top}\left(\left(\boldsymbol{B}_{k}^{\top}\boldsymbol{e}_{i}\right)_{S} : k \in [K]\right) \right\rangle
$$
\n
$$
= \dim \mathrm{span}\left\langle \left(\boldsymbol{B}_{k}^{\top}\boldsymbol{e}_{i}\right)_{S} : k \in [K]\right\rangle = \left|\mathrm{pa}_{\mathcal{G}}(i)\right| + 1,
$$

implying that $\hat{\textbf{\textit{B}}}_{k}, k \in [K]$ satisfy [Assumption 5.](#page-5-1)

Now we have shown that for almost every $M \in \mathcal{M}^0_{\text{sur}}(\mathcal{G}),$ we can construct a hypothetical data generating process with latent variables $v = Mz$ that satisfies all requirements in [Theorem 9.](#page-36-0) Choose an arbitrary M that is in $\mathcal{M}_{\text{sur}}(\mathcal{G})$, then we have that

$$
\frac{\partial v_i}{\partial z_j} \neq 0, \quad j \notin \overline{\text{sur}}_{\mathcal{G}}(i).
$$

Finally, if we additionally assume single-node interventions, $\forall k, \ell \in \mathfrak{E}_i$, we have that $(\bm{B}_k)_j$ \neq $(\bm B_\ell)_j \Leftrightarrow j$ = *i*. For any $\bm M\in\mathcal{M}^0_{\rm sur}(\mathcal{G})$ (and specifically the $\bm M$ that we have already chosen above), we have $(\hat{\bm{B}}_k)_j = (\bm{B}_k)_j \bm{M}^{-1}$ and $(\hat{\bm{B}}_\ell)_j = (\bm{B}_\ell)_j \bm{M}^{-1}, \forall j \in [\![d]\!].$ Thus, $(\hat{\bm{B}}_k)_j \neq (\hat{\bm{B}}_\ell)_j \Leftrightarrow j = i$ as well, implying that \mathfrak{E}_i is also a group of single-node interventions on ν , concluding the proof.

J.3 Formal statement and proof of [Theorem 10:](#page-39-0) the non-parametric case

Theorem 10 (Counterpart to [Theorem 5\)](#page-10-0)**.** *For any causal model* (*h*, G) *and any set of environments* E*,* $suppose that we have observations ${P^E_{\boldsymbol{X}}}\}_{E \in \mathfrak{E}}$ satisfying Assumption 1:$ $suppose that we have observations ${P^E_{\boldsymbol{X}}}\}_{E \in \mathfrak{E}}$ satisfying Assumption 1:$

$$
\forall E \in \mathfrak{E}, \mathbf{z} \sim p_E(\hat{\mathbf{z}}) = \prod_{i=1}^d p_i^E\left(\hat{\mathbf{z}}_i \mid \hat{\mathbf{z}}_{\text{pa}_{\mathcal{G}}(i)}\right), \mathbf{x} = \mathbf{h}^{-1}(\mathbf{z})
$$

such that

- (i) all densities p^E_i are continuously differentiable and the joint density p_E is positive everywhere;
- *(ii)* the environments are groups of single-node interventions: there exists a partition $\mathfrak{E} = \bigcup_{i=1}^d \mathfrak{E}_i$ such that $\mathcal{I}_{z}^{\mathfrak{E}_{i}} = \{i\};$
- *(iii) the intervention distributions on each node are non-degenerate:* ∀*i* ∈ [*d*]*, the set of distributions* $\{p_i^E : E \in \mathfrak{E}_i\}$ satisfy [Definition 5](#page-10-2) at any point $\hat{\boldsymbol{z}} \in \mathbb{R}^d$ *,*

then there must exist a candidate solution (ˆ*h*, G) *and a hypothetical data generating process*

$$
\forall E \in \mathfrak{E}, \nu \sim q_E(\hat{\nu}) = \prod_{i=1}^d q_i^E \left(\hat{\nu}_i \mid \hat{\nu}_{pa_{\mathcal{G}}(i)} \right), \mathbf{x} = \hat{\boldsymbol{h}}^{-1}(\nu)
$$

such that

(i') all densities q^E_i are continuously differentiable and the joint density q_E is positive everywhere; (*ii*^{\prime}) for the same partition $\mathfrak{E} = \bigcup_{i=1}^{d} \mathfrak{E}_i$, we have $\mathcal{I}_{\mathfrak{v}}^{\mathfrak{E}_i} = \{i\}$; (*iii*^{\prime}) $\forall i \in [d]$, the set of distributions $\{q_i^E : E \in \mathfrak{E}_i\}$ satisfy [Definition 5](#page-10-2) at any point $\hat{\mathbf{v}} \in \mathbb{R}^d$ *,*

but

$$
\frac{\partial v_i}{\partial z_j} \neq 0, \quad \forall j \in \overline{\text{sur}}_{\mathcal{G}}(i).
$$

Remark 3. *Similar to the case of [Theorem 9,](#page-36-0) [Section J.3](#page-39-1) also establishes a stronger form of identifiability. First, it is assumed that the causal graph* G *is known. Second, we only focus on a special case of the setting of [Theorem 5](#page-10-0) by assuming that the support is the whole space, and the non-degeneracy condition [Definition 5](#page-10-2) holds at any point. Even in this case, we show that our identification guarantee up to SNA cannot be improved.*

We state and prove a stronger version of [Theorem 10:](#page-39-0)

Theorem 11. *For any causal model* (*h*, G) *and any set of environments* E*, suppose that we have observations* $\{P_{\textbf{X}}^E\}_{E\in\mathfrak{E}}$ *satisfying Assumption* 1:

$$
\forall E \in \mathfrak{E}, \quad \mathfrak{z} \sim p_E(z) = \prod_{i=1}^d p_i^E \left(z_i \mid z_{\text{pa}_{\mathcal{G}}(i)} \right), \quad \mathfrak{x} = h^{-1}(z)
$$

such that

- (i) all densities p^E_i are continuously differentiable and the joint density p_E is positive everywhere;
- *(ii)* the environments are groups of single-node interventions: there exists a partition $\mathfrak{E} = \bigcup_{i=1}^d \mathfrak{E}_i$ such that $\mathcal{I}_{z}^{\mathfrak{E}_{i}} = \{i\};$

(iii) the intervention distributions on each node are non-degenerate: ∀*i* ∈ [*d*]*, the set of distributions* $\{p_i^E : E \in \mathfrak{E}_i\}$ *satisfy [Definition 5,](#page-10-2)*

then there must exist a candidate solution (ˆ*h*, G) *and a hypothetical data generating process*

$$
\forall E \in \mathfrak{E}, \quad \mathbf{\nu} \sim q_E(\mathbf{\nu}) = \prod_{i=1}^d q_i^E \left(\nu_i \mid \nu_{\text{pa}_G(i)} \right), \quad \mathbf{x} = \hat{\boldsymbol{h}}^{-1}(\mathbf{\nu})
$$

such that

(i') all densities q_i^E are continuously differentiable and the joint density q_E is positive everywhere; (*ii*^{\prime}) *for the same partition* $\mathfrak{E} = \bigcup_{i=1}^{d} \mathfrak{E}_i$ *, we have* $\mathcal{I}_{\mathfrak{v}}^{\mathfrak{E}_i} = \{i\}$; *(iii'*) ∀*i* ∈ [*d*], the set of distributions $\{q_i^E : E \in \mathfrak{E}_i\}$ satisfy [Definition 5,](#page-10-2)

but

$$
\frac{\partial v_i}{\partial z_j} \neq 0, \quad \forall j \in \overline{\text{sur}}_{\mathcal{G}}(i).
$$

Finally, if we additionally assume minimality [\(Assumption 6\)](#page-18-2) and/or faithfulness [\(Assumption 7\)](#page-18-3) of all pE's, we can guarantee the existence of (^ˆ*h*, ^G) *and qE's satisfying minimality and/or faithfulness in addition to the properties listed above. In other words, assuming minimality and/or faithfulness does not resolve the ambiguity.*

Proof. We define

$$
v = Mz \tag{29}
$$

where *M* is an effect-respecting matrix. At this point we do not make any other restrictions on *M*, and we will choose appropriate M later. By [Lemma 20,](#page-35-3) there exists invertible matrices M_i and $M_i^ \sup_{\text{pa}_{\mathcal{G}}(i)}$ = $\bm{M}_i^- \bm{z}_{\text{pa}_{\mathcal{G}}(i)}$ and $\bm{\nu}_{\overline{\text{pa}}_{\mathcal{G}}(i)}$ = $\bm{M}_i \bm{z}_{\overline{\text{pa}}_{\mathcal{G}}(i)},$ so for all environment $E \in \mathfrak{E}$ we have

$$
q_i^E(\nu_{\text{pa}_{\mathcal{G}}(i)}) = p_i^E(z_{\text{pa}_{\mathcal{G}}(i)}) \cdot \left| \det(\boldsymbol{M}_i^-)^{-1} \right|, \quad q_i^E(\nu_{\overline{\text{pa}}_{\mathcal{G}}(i)}) = p_i^E(z_{\overline{\text{pa}}_{\mathcal{G}}(i)}) \cdot \left| \det(\boldsymbol{M}_i)^{-1} \right|
$$

so that

$$
q_i^E\left(\boldsymbol{v}_i \mid \boldsymbol{v}_{\text{pa}_{\mathcal{G}}(i)}\right) = p_i^E\left(\boldsymbol{z}_i \mid \boldsymbol{z}_{\text{pa}_{\mathcal{G}}(i)}\right) \frac{|\det \boldsymbol{M}_i^{-1}|}{|\det(\boldsymbol{M}_i^{-})^{-1}|}, \quad \forall i \in [d]. \tag{30}
$$

In the following, assuming that $(p_i^E : E \in \mathfrak{E})$ satisfies any of the listed assumptions, we show that $(q_i^E : E \in \mathfrak{E})$ satisfies the same assumption as well.

Firstly, (30) immediately implies that the density of ν is continuous differentiable and positive everywhere. Secondly, ∀*k*,ℓ ∈ E*ⁱ* , we have that

$$
p_j^{E_k}\left(z_j\mid z_{\text{pa}_{\mathcal{G}}(j)}\right)=p_j^{E_\ell}\left(z_j\mid z_{\text{pa}_{\mathcal{G}}(j)}\right)\Leftrightarrow j=i.
$$

By [\(30\)](#page-40-0) it is easy to see that

$$
q_j^{E_k}\left(\boldsymbol{v}_j\mid\boldsymbol{v}_{\text{pa}_{\mathcal{G}}(j)}\right)=q_j^{E_\ell}\left(\boldsymbol{v}_j\mid\boldsymbol{v}_{\text{pa}_{\mathcal{G}}(j)}\right)\Leftrightarrow j=i
$$

as well, *i.e.,* q^k *,* $k \in \mathfrak{E}_i$ are single-node interventions on $\pmb{\nu}_i$ according to [Definition 2.](#page-2-3)

Thirdly, we verify the non-degeneracy condition for q_i^{E} 's. Indeed we have for $\forall k \geq 2$ that

$$
\nabla_{\nu_{\overline{\text{pa}}}_{\mathcal{G}}(i)} \frac{q_i^{E_1}}{q_i^{E_k}} \left(\nu_i \mid \nu_{\text{pa}_{\mathcal{G}}(i)} \right) = \frac{\partial z_{\overline{\text{pa}}_{\mathcal{G}}}(i)}{\partial \nu_{\overline{\text{pa}}_{\mathcal{G}}}(i)} \nabla_{z_{\overline{\text{pa}}_{\mathcal{G}}}(i)} \frac{q_i^{E_1}}{q_i^{E_k}} \left(z_i \mid z_{\text{pa}_{\mathcal{G}}(i)} \right) = M_i^{-1} \nabla_{z_{\overline{\text{pa}}_{\mathcal{G}}}(i)} \frac{q_i^{E_1}}{q_i^{E_k}} \left(z_i \mid z_{\text{pa}_{\mathcal{G}}(i)} \right).
$$

Since M_i is invertible, the above equation and the non-degeneracy of $p^{E_k}, k \in [K]$ immediately implies that non-degeneracy of $q^{E_k}, k \in [K].$

Thus, for arbitrary $M \in \mathcal{M}_{\text{sur}}(\mathcal{G})$, we have constructed a hypothetical data generating process with latent variable $v = Mz$ that satisfies all given conditions. It remains to show that such construction is still possible under additional minimality and faithfulness conditions.

Claim 1. There exists a neighbourhood \odot of the identity matrix I in $\overline{\mathcal{M}}_{\text{sur}}(\mathcal{G})$ (in the **sense of [Remark 1\)](#page-35-4)** such that for $\forall M \in O \cap \mathcal{M}_{\text{sur}}^0(\mathcal{G})$, $p^{E_k}, k \in [K]$ satisfy [Assumption 7](#page-18-3) ⇒ *q ^E^k* , *k* ∈ [*K*] **satisfy [Assumption 7.](#page-18-3)**

For $\forall i,j$ not *d*-separated by $S \subseteq [d]$, for all $k \in [K]$ there exists $\hat{\boldsymbol{z}} \in \mathbb{R}^d$ such that $\Delta_k^{(i,j,S)}$ $\binom{n}{k}$ $p^{E_k}\left(\hat{\bm{z}}_i,\hat{\bm{z}}_j\mid\hat{\bm{z}}_S\right)-p^{E_k}\left(\hat{\bm{z}}_i\mid\hat{\bm{z}}_S\right)p^{E_k}\left(\hat{\bm{z}}_j\mid\hat{\bm{z}}_S\right)\neq 0.$ By continuous differentiability of p^{E_k} , we know that there exists $\delta_k^{(i,j,S)}$ $\frac{f(i,j,S)}{k} > 0$ such that for all $M \in \overline{\mathcal{M}}_{\text{sur}}(\mathcal{G})$ such that $\|M - I\|_F \leqslant \delta_k^{(i,j,S)}$ $\hat{k}^{(y,0)}$, the density of the variable \bm{v} = \bm{Mz} satisfies $q^{E_k}\left(\hat{\bm{v}}_i,\hat{\bm{\nu}}_j\mid\hat{\bm{\nu}}_S\right)\neq q^{E_k}\left(\hat{\bm{\nu}}_i\mid\hat{\bm{\nu}}_S\right)q^k\left(\hat{\bm{\nu}}_j\mid\hat{\bm{\nu}}_S\right)$ for $\hat{\bm{\nu}}$ = $\bm{M\hat{z}},$ which implies that $\bm{\nu}_i$ and v_j are dependent given v_S . Now choose $\delta = \min_{k,i,j,S} \delta_k^{(i,j,S)}$ $\mathcal{M}_k^{(i,j),\sigma} > 0$, then for all $\boldsymbol{M} \in \mathcal{M}_{\text{sur}}(\mathcal{G})$ such that $\|\boldsymbol{M}-\boldsymbol{I}\|_F$ ≤ δ, the resulting distributions $q^{E_k}, k ∈ [K]$ satisfy assumption [Assumption 6.](#page-18-2)

Claim 2. There exists a neighbourhood \bigcirc of *I* in $\overline{\mathcal{M}}_{\text{sur}}(\mathcal{G})$ (in the sense of [Remark 1\)](#page-35-4) such ${\bf t}$ hat for almost all $M\in O\cap {\cal M}_{\rm sur}^0({\cal G}),$ $p^{E_k}, k\in [K]$ satisfies [Assumption 6](#page-18-2) $\Rightarrow p^{E_k}, k\in [K]$ satisfies **[Assumption 6.](#page-18-2)**

The proof is similar to the previous statement. Since [Assumption 6](#page-18-2) causal minimality is satisfied for *z*, for $\forall k \in [K]$, $i \in [d]$, let G_{ij} be the resulting graph obtained by removing the edge $j \to i$ from $\mathcal{G},$ then there must exists some $\alpha_{ijk}\in [d]$ such that $z_{\alpha_{ijk}}\not\perp z_{\mathrm{nd}_{\mathcal{G}_{ij}}(\alpha_{ijk})}\mid z_{\mathrm{pa}_{\mathcal{G}_{ij}}(\alpha_{ijk})}.$ Hence, there exists $\hat{\boldsymbol{z}}^{ijk} \in \mathbb{R}^d$ such that

$$
p^{E_k}\left(\hat{\boldsymbol{z}}_{\alpha_{ijk}}^{ijk}\mid\hat{\boldsymbol{z}}_{\text{pa}_{\mathcal{G}_{ij}}(\alpha_{ijk})}^{ijk}\right)p^{E_k}\left(\hat{\boldsymbol{z}}_{\text{nd}_{\mathcal{G}_{ij}}(\alpha_{ijk})}^{ijk}\mid\hat{\boldsymbol{z}}_{\text{pa}_{\mathcal{G}_{ij}}(\alpha_{ijk})}^{ijk}\right)\neq p^{E_k}\left(\hat{\boldsymbol{z}}_{\text{nd}_{\mathcal{G}_{ij}}(\alpha_{ijk})}^{ijk}\mid\hat{\boldsymbol{z}}_{\text{pa}_{\mathcal{G}_{ij}}(\alpha_{ijk})}^{ijk}\right).
$$

By continuous differentiability of p^{E_k} , there exists $\delta_k^{(i,j)}$ $\mathcal{M}_k^{(t,j)} > 0$ such that for all $\boldsymbol{M} \in \bar{\mathcal{M}}_{\text{sur}}(\mathcal{G})$ such that $\|\boldsymbol{M}-\boldsymbol{I}\|_F \leqslant \delta_k^{(i,j)}$ $\hat{\bm{h}}_k^{(i,j)}$, the density $q_{ij}^{E_k}$ of the variable $\hat{\bm{\nu}}^{ijk} = \bm{M}\hat{\bm{z}}^{ijk}$ satisfies

$$
q^{E_k}\left(\hat{\boldsymbol{\nu}}_{\boldsymbol{\alpha}_{ijk}}^{ijk}\mid\hat{\boldsymbol{\nu}}_{\text{pa}_{\mathcal{G}_{ij}}(\boldsymbol{\alpha}_{ijk})}^{ijk}\right)q^{E_k}\left(\hat{\boldsymbol{\nu}}_{\text{nd}_{\mathcal{G}_{ij}}(\boldsymbol{\alpha}_{ijk})}^{ijk}\mid\hat{\boldsymbol{\nu}}_{\text{pa}_{\mathcal{G}_{ij}}(\boldsymbol{\alpha}_{ijk})}^{ijk}\right)\neq q^{E_k}\left(\hat{\boldsymbol{\nu}}_{\text{nd}_{\mathcal{G}_{ij}}(\boldsymbol{\alpha}_{ijk})}^{ijk}\mid\hat{\boldsymbol{\nu}}_{\text{pa}_{\mathcal{G}_{ij}}(\boldsymbol{\alpha}_{ijk})}^{ijk}\right).
$$

for $\hat{\bm{\nu}}^{ijk}$ = $\bm{M}\hat{\bm{z}}^{ijk}.$ This implies that removing the edge $j\to i$ in ${\cal G}$ would break the causal Markov condition for q^{E_k} . Now let δ = min_{k,i,j} δ^(i,j) $\mathcal{L}_{k}^{(i,j)}$ > 0, then for all $\boldsymbol{M} \in \bar{\mathcal{M}}_{\text{sur}}(\mathcal{G})$ such that $\|\boldsymbol{M}-\boldsymbol{I}\|_{F} \leqslant \delta$, the resulting distributions $q^{E_k}, k \in [K]$ satisfy assumption [Assumption 1.](#page-2-1)

Combining the above two statements and what we have proven before, it is straightfoward to see that one can choose some $M \in \mathcal{M}_{\text{sur}}(\mathcal{G})$ in a small neighbourhood of *I* that satisfies all the requirements, completing the proof. \Box