

A NEW APPROACH TO WEIGHTED SOBOLEV SPACES

DJAMEL EDDINE KEBICHE

ABSTRACT. We present in this paper a new way to define weighted Sobolev spaces when the weight functions are arbitrary small. This new approach can replace the old one consisting in modifying the domain by removing the set of points where at least one of the weight functions is very small. The basic idea is to replace the distributional derivative with a new notion of weak derivative. In this way, non-locally integrable functions can be considered in these spaces. Indeed, assumptions under which a degenerate elliptic partial differential equation has a unique non-locally integrable solution are given. Tools like a Poincaré inequality and a trace operator are developed, and density results of smooth functions are established.

1. INTRODUCTION

By a weight function $u : \Omega \rightarrow \mathbb{R}$ we mean a locally integrable function which is non-negative almost everywhere in Ω . For $p \in [1, \infty)$, $L^p(\Omega, u)$ denotes the space of all functions $f : \Omega \rightarrow \mathbb{R}$ satisfying $fu^{1/p} \in L^p(\Omega)$. Let $m \in \mathbb{N}$ and consider a collection $S := \{u_\alpha\}_{\alpha \in \pi_m}$ ($\pi_m := \{\alpha \in \mathbb{N}^d \mid |\alpha| \leq m\}$) of weight functions on Ω . The weighted Sobolev space $W^{m,p}(\Omega, S)$ is defined as the space of all functions $f \in L^p(\Omega, u) \cap L^1_{\text{loc}}(\Omega)$ such that their distributional derivative are elements of $L^p(\Omega, u_\alpha) \cap L^1_{\text{loc}}(\Omega)$. The space $W^{m,p}(\Omega, S)$ is equipped with the norm

$$\forall f \in W^{m,p}(\Omega, S) : \|f\|_{W^{m,p}(\Omega, S)} := \left(\sum_{|\alpha| \leq m} \|u_\alpha^{1/p} D^\alpha f\|_{L^p}^p \right)^{1/p}.$$

A sufficient condition for the space $W^{m,p}(\Omega, S)$ to be a Banach space is

$$\forall \alpha \in \pi_m : u_\alpha^{-1} \in L^{\frac{1}{p-1}}_{\text{loc}}(\Omega) \quad (1.1)$$

(see Thm. 1.11 of [5]). Indeed, the assumption (1.1) implies that for all $\alpha \in \pi_m$, the space $L^p(\Omega, u_\alpha)$ is continuously embedded into $L^1_{\text{loc}}(\Omega)$, and the functional $W^{m,p}(\Omega, S) \rightarrow \mathbb{R}, f \rightarrow \int_\Omega f \partial^\alpha \varphi dx$ ($\varphi \in \mathcal{D}(\Omega)$) is continuous, a crucial property used in the proof of the completeness of the space $W^{m,p}(\Omega, S)$ (see Remark 1.8 of [5]). However, when (1.1) is violated, the weighted Sobolev space $W^{m,p}(\Omega, S)$ is not necessary a Banach space (see Example 1.12 of [5]). This is because, when (1.1) is violated, the latter functionals are, in general, no longer continuous. The existing remedy consists in replacing the set Ω with a smaller one so that the

2020 *Mathematics Subject Classification.* 46E35, 35J70, 35A23.

Key words and phrases. Weighted Sobolev spaces, degenerate elliptic PDEs, Poincaré inequality.

This research was funded in whole or in part by the Austrian Science Fund (FWF) 10.55776/P33538

For open access purposes, the author has applied a CC BY public copyright license to any author-accepted manuscript version arising from this submission.

property (1.1) holds. More precisely, the new set, which will be denoted by Ω_S^* , is defined by $\Omega_S^* := \Omega \setminus \cup_{\alpha \in \pi_m} P(u_\alpha)$ where $P(u_\alpha)$ is the set of points $x \in \Omega$ for which $u_\alpha^{-1} \notin L^{\frac{1}{p-1}}(\Omega \cap U)$ for all neighborhood U of x . The set $B := \cup_{\alpha \in \pi_m} P(u_\alpha)$ is closed (and hence Ω_S^* is open) and Lebesgue negligible. Moreover, property (1.1) holds when Ω is replaced with Ω_S^* and hence the weighted Sobolev space $W^{m,p}(\Omega_S^*, S)$ is a Banach space (see Sec. 3 of [5] for more details). However, the disadvantage of this remedy is a modification of the domain. As a consequence, the convexity might be lost and the boundary of Ω is changed.

The alternative we present in this paper consists in replacing the notion of distributional derivative with a new notion of weak derivative, called weak derivative in the sense of $L^1_{v,\text{loc}}(\Omega)$ (v being a $C^m(\Omega)$ function). It consists in replacing the space of test functions, which is $C_c^m(\Omega_S^*)$ in the case of $W^{m,p}(\Omega_S^*, S)$, with a larger one whose elements do not necessary vanish near the set B (see Def. 4). When the function v is well chosen, i.e. $\Omega_S^* = \Omega_{|v|^{m+p}}^* := \Omega \setminus P(|v|^{m+p})$, the two notions of derivative coincide (see Prop. 7).

In Sec. 5, several density results of smooth functions in weighted Sobolev spaces are proved. In Sec. 6, we give sufficient conditions under which (continuous) trace operators can be defined. A Poincaré inequality is established in Sec. 8. In the last section, we deal with degenerate elliptic linear partial differential equations, where in particular we give assumptions under which a solution is not locally integrable.

2. THE WEIGHTED LEBESGUE SPACE $L_w^p(\Omega)$

Throughout this paper, a weight function $u : \Omega \longrightarrow \mathbb{R}$ is a locally integrable function, which is positive a.e. in Ω . The set of all the weight functions on Ω is denoted by $W(\Omega)$.

We now give a definition of weighted Lebesgue spaces (using notations slightly different from the classical ones)

Definition 1. Let $\Omega \subseteq \mathbb{R}^d$ be an open set, $p \in [1, \infty)$, and let w be such that $|w|^p \in W(\Omega)$ (in the sequel we write simply $w \in E_p(\Omega)$). We define the weighted Lebesgue space $L_w^p(\Omega)$ as the space of measurable functions f on Ω such that $fw \in L^p(\Omega)$. This space will be equipped with the norm $\|\cdot\|_{L_w^p}$ defined by

$$\forall f \in L_w^p(\Omega) : \|f\|_{L_w^p(\Omega)} := \|fw\|_{L^p(\Omega)}. \quad (2.1)$$

The space $L_w^2(\Omega)$ is equipped with the scalar product $(\cdot, \cdot)_{L_w^2}$ defined by

$$\forall f, g \in L_w^2(\Omega) : (f, g)_{L_w^2} := (wf, wg)_{L^2} \quad (2.2)$$

where $(\cdot, \cdot)_{L^2}$ is the usual scalar product of the space $L^2(\Omega)$.

Moreover, we write $f \in L_{w,\text{loc}}^p(\Omega)$ if $f \in L_w^p(K)$ for all $K \Subset \Omega$.

If we denote by $L^p(\Omega, u)$ the weighted Lebesgue space with the weight function u , then we clearly have

$$L_w^p(\Omega) = L^p(\Omega, |w|^p).$$

The reason for using w instead of the weight function $|w|^p$ is purely for notational reason.

The following theorem is well known (see e.g. Thm. III. 6.6 of [2]).

Theorem 2. Let $\Omega \subseteq \mathbb{R}^d$ be an open set, $w \in E_p(\Omega)$, and $p \in [1, \infty)$. Then, the space $L_w^p(\Omega)$ equipped with the norm (2.1) is a Banach space (Hilbert space when $p = 2$).

Let \mathcal{S} be the Schwartz space. For a fixed non-null polynomial P we set

$$\mathcal{S}_P = \{\psi \in \mathcal{S} : \psi = P\varphi, \varphi \in \mathcal{S}\}$$

which is a linear subspace of \mathcal{S} equipped with the sequence $(\|\cdot\|_k)$ of semi norms of \mathcal{S} . Having denoted by \mathcal{S}'_P the dual space of \mathcal{S}_P , we have the following proposition.

Proposition 3. *Let P be a non-null polynomial function and let $p \in [1, \infty)$. Then, any element $f \in L^p_P(\mathbb{R}^d)$ defines an element of \mathcal{S}'_P in following way*

$$\forall \psi \in \mathcal{S}_P : \langle f, \psi \rangle_{\mathcal{S}'_P, \mathcal{S}_P} := \int_{\mathbb{R}^d} f(x) \psi(x) dx.$$

Proof. Let $f \in L^p_P(\mathbb{R}^d)$, $\psi \in \mathcal{S}_P$ and pick φ so that $\psi = P\varphi$. By Hölder's inequality we have

$$|\langle f, \psi \rangle_{\mathcal{S}'_P, \mathcal{S}_P}| \leq \int_{\mathbb{R}^d} |f(x)P(x)\varphi(x)| dx \leq \|f\|_{L^p_P} \|\varphi\|_{L^{p'}} \leq C \|f\|_{L^p_P} \|\varphi\|_k$$

for some $k \in \mathbb{N}$ and $C > 0$ that depend only on d and p , where p' is the conjugate exponent of p . By Thm. 1 of [4], the linear continuous multiplication map $M_P : \mathcal{S} \longrightarrow \mathcal{S}_P$ defined by $M_P\varphi = P\varphi$ has a continuous inverse $M_P^{-1} : \mathcal{S}_P \longrightarrow \mathcal{S}$. It follows that

$$\exists C' > 0 \exists k' \in \mathbb{N} \forall \psi \in \mathcal{S}_P : |\langle f, \psi \rangle_{\mathcal{S}'_P, \mathcal{S}_P}| \leq C' \|f\|_{L^p_P} \|\psi\|_{k'}$$

which completes the proof. \square

3. THE WEAK DERIVATIVE IN THE SENSE OF $L^1_{v, \text{loc}}(\Omega)$

Let $w \in E_p(\Omega)$. When w is arbitrary small, elements of $L^p_w(\Omega)$ are not necessary locally integrable over Ω , and hence the notion of distributional derivative to elements of $L^p_w(\Omega)$ is not well defined in general. The existing remedy is to replace Ω with $\Omega^*_{|w|^p} := \Omega \setminus P(|w|^p)$ where $P(|w|^p)$ is the set of points $x \in \Omega$ for which

$$|w|^{-p} \notin L^{\frac{1}{p-1}}(\Omega \cap U)$$

for all neighborhood U of x . The set $P(|w|^p)$ is closed and Lebesgue negligible. Moreover, $L^p_w(\Omega) \subseteq L^1_{\text{loc}}(\Omega^*_{|w|^p})$ with continuous embedding (see Sec. 3 of [5] for more details). Hence, the notion of weak derivative in the sense of $L^1_{\text{loc}}(\Omega^*_{|w|^p})$ is well defined on the space $L^p_w(\Omega)$. However, note that with this weak derivative, we are allowed to take only test functions having a compact support in $\Omega^*_{|w|^p}$.

The main aim of this section is to present a new notion of weak derivative where the test functions are not necessary equal to 0 near the set $P(|w|^p)$ as it is the case for the weak derivative in the sense of $L^1_{\text{loc}}(\Omega^*_{|w|^p})$.

Definition 4. Let $\Omega \subseteq \mathbb{R}^d$ be an open set, $\alpha \in \mathbb{N}^d$ with $|\alpha| \neq 0$, $p \in [1, \infty)$, $w \in E_p(\Omega)$, and $v \in \mathcal{C}^{|\alpha|, *}_w(\Omega)$ (i.e. $v \in \mathcal{C}^{|\alpha|}_w(\Omega)$ and $v \neq 0$ a.e. in Ω) be a map satisfying

$$\forall K \Subset \Omega \exists C > 0 : |v| \leq C|w| \text{ a.e. in } K. \quad (3.1)$$

We say that $f \in L^1_{w, \text{loc}}(\Omega)$ is α -weakly differentiable in the sense of $L^1_{v, \text{loc}}(\Omega)$ if there exists an element of $L^1_{v^{|\alpha|+1}, \text{loc}}(\Omega)$ denoted by $D^\alpha_v f$ satisfying

$$\forall \varphi \in \mathcal{D}(\Omega) : \int_{\Omega} f \partial^\alpha (v^{|\alpha|+1} \varphi) dx = (-1)^{|\alpha|} \int_{\Omega} v^{|\alpha|+1} \varphi D^\alpha_v f dx. \quad (3.2)$$

In the sequel, for some given $f \in L^1_{w,\text{loc}}(\Omega)$, we will frequently write $\exists D^\alpha_v f$ to say that the α -weak derivative of f in the sense of $L^1_{v,\text{loc}}(\Omega)$ exists (and hence belongs to $L^1_{v^{|\alpha|+1},\text{loc}}(\Omega)$).

Remark 5. Let Ω , α , p , w , v be as in the previous definition.

- (i) The weak derivative in the sense of $L^1_{v,\text{loc}}(\Omega)$ (when it exists) is unique. The proof is identical to the proof of the uniqueness of the weak derivative in the sense of $L^1_{\text{loc}}(\Omega)$.
- (ii) Take α with $|\alpha| = 1$ and let $f \in L^p_w(\Omega)$. Assume that $\exists D^\alpha_v f \in L^p_{v^2}(\Omega)$, and that $f\partial^\alpha(v^2)$, $v^2 f \in L^p(\Omega)$, e.g. Ω is bounded and $\nabla v \in L^\infty(\Omega)$, then (3.2) yields

$$\forall \varphi \in \mathcal{D}(\Omega) : \int_{\Omega} (f\partial^\alpha(v^2) + v^2 D^\alpha_v f) \varphi \, dx = - \int_{\Omega} v^2 f \partial^\alpha \varphi \, dx$$

which implies that $v^2 f$ is α -weakly differentiable in the sense of $L^1_{\text{loc}}(\Omega)$ (or simply $\exists D^\alpha(v^2 f)$) and $D^\alpha(v^2 f) = f\partial^\alpha(v^2) + v^2 D^\alpha_v f \in L^p(\Omega)$. Consequently, if we assume that the above conditions hold for all $\alpha \in \mathbb{N}^d$ with $|\alpha| = 1$ we get that $v^2 f \in W^{1,p}(\Omega)$.

- (iii) One can show that for any $\varphi \in \mathcal{D}(\Omega)$, there exists a compactly supported continuous function h_α on Ω such that $\partial^\alpha(v^{|\alpha|+1}\varphi) = v h_\alpha$. By (3.1) we have $|\partial^\alpha(v^{|\alpha|+1}\varphi)| \leq C|w|h_\alpha$ a.e. on $K := \text{supp}(\varphi)$. It follows that

$$\left| \int_{\Omega} f \partial^\alpha(v^{|\alpha|+1}\varphi) \, dx \right| \leq C \|f\|_{L^p_w(\Omega)} \|h_\alpha\|_{L^\infty(K)}.$$

Hence, the left hand side of (3.2) is always finite. Moreover, for any test function φ , the functional $L^p_w(\Omega) \rightarrow \mathbb{C}$, $f \mapsto \int_{\Omega} f \partial^\alpha(v^{|\alpha|+1}\varphi) \, dx$ is continuous. This property will be used to prove the completeness of the spaces $W^{m,p}_{V,v}(\Omega)$ defined below.

- (iv) By density of $\mathcal{D}(\Omega)$ in $\mathcal{C}^{|\alpha|}_c(\Omega)$, we can equally well use the space $\mathcal{C}^{|\alpha|}_c(\Omega)$ instead of the space $\mathcal{D}(\Omega)$ in (3.2). This is because using (3.1) we have

$$\left| \int_{\Omega} f \partial^\alpha(v^{|\alpha|+1}\varphi) \, dx \right| \leq C \|fw\|_{L^1(K)} \sum_{0 \leq \beta \leq \alpha} C_\beta \|g_\beta\|_{L^\infty(K)} \|\partial^\beta \varphi\|_{L^\infty(K)},$$

where K is the support of φ , g_β is a continuous function that depends on v , α and β .

- (v) In case $w \in \mathcal{C}^{|\alpha|}(\Omega)$ we can simply choose $v = w$.
- (vi) The case $w \equiv 1$ can be considered since (3.1) trivially holds.

Example 6. Let $f(x) = 1/x^2 \in L^2_{x^2}(-1, 1)$. Then, $\exists D_{x^2} f = -2x^{-3} \in L^2_{x^3}(-1, 1)$.

Now, we compare the notion of weak derivative in the sense of $L^1_{v,\text{loc}}(\Omega)$ with the notion of the weak derivative in the sense of L^1_{loc} .

If $m, k \in \mathbb{N}$ are such that $k < m$, then $\pi_m^k := \pi_m \setminus \pi_k$. More generally, if $\alpha \in \mathbb{N}^d$, then π_α is the set of multi-indices $\beta \in \mathbb{N}^d$ such that $\beta \leq \alpha$ (i.e. $\beta_i \leq \alpha_i$ for all i). If $\lambda \in \mathbb{N}^d$ is such that $\lambda < \alpha$ (i.e. $\lambda \leq \alpha$ with $\lambda_j < \alpha_j$ for some j), then π_α^λ denotes the set $\pi_\alpha \setminus \pi_\lambda$.

The relation between the weak derivative in the sense of L^1_{loc} and the weak derivative in the sense of $L^1_{v,\text{loc}}(\Omega)$ is given in the next proposition.

Proposition 7. Let $\Omega \subseteq \mathbb{R}^d$ be an open set, $\alpha \in \mathbb{N}^d$ with $|\alpha| \neq 0$, $p \in [1, \infty)$, $w \in E_p(\Omega)$, $v \in \mathcal{C}^{|\alpha|,*}(\Omega)$ be a map satisfying (3.1).

- (i) Let $f \in L_{\text{loc}}^1(\Omega) (= L_{w,\text{loc}}^1(\Omega)$ with $w \equiv 1$). If $\exists D^\alpha f \in L_{\text{loc}}^1(\Omega)$ then $\exists D_v^\alpha f$ and $D^\alpha f = D_v^\alpha f$.
- (ii) Let $f \in L_{w,\text{loc}}^p(\Omega)$. Then, f is α -weakly differentiable in the sense of $L_{v,\text{loc}}^1(\Omega)$ with $D_v^\alpha f \in L_{v^{|\alpha|+1},\text{loc}}^p(\Omega)$ if and only if its α -weak derivative in the sense of $L_{\text{loc}}^1(\Omega_{|v|^{p|\alpha|+p}}^*)$, denoted by $D^\alpha f$, exists and belongs to $L_{v^{|\alpha|+1},\text{loc}}^p(\Omega)$. In particular, we have $D_v^\alpha f = D^\alpha f$ a.e. in $\Omega_{|v|^{p|\alpha|+p}}^*$.

Proof. (i): Recall that, by density of $\mathcal{D}(\Omega)$ in the space $\mathcal{C}_c^{|\alpha|}(\Omega)$, we can replace the space $\mathcal{D}(\Omega)$ in the definition of the weak derivative in the sense of $L_{\text{loc}}^1(\Omega)$ with the space $\mathcal{C}_c^{|\alpha|}(\Omega)$. Hence, it suffices to use the definition of the weak derivative in the sense of $L_{\text{loc}}^1(\Omega)$ with the test function $\varphi v^{|\alpha|+1} \in \mathcal{C}_c^{|\alpha|}(\Omega)$ instead of $\varphi \in \mathcal{D}(\Omega)$.

(ii): Assume that f is α -weakly differentiable in the sense of $L_{v,\text{loc}}^1(\Omega)$ with $D_v^\alpha f \in L_{v^{|\alpha|+1},\text{loc}}^p(\Omega)$. Let $\eta \in \mathcal{C}^\infty(\mathbb{R})$ be such that

$$\eta \equiv 1 \text{ on } [-1, 1]^c, \eta \equiv 0 \text{ on } \left[-\frac{1}{2}, \frac{1}{2}\right], 0 \leq \eta \leq 1,$$

and for all $n \in \mathbb{N}$, set $\chi_n := \eta_n \circ v$, where $\eta_n(\cdot) := \eta(n \cdot)$. Setting

$$\forall n \in \mathbb{N} : M_n := \{x \in \Omega \mid |v(x)| \leq 1/n\}$$

Clearly $\chi_n \in \mathcal{C}^{|\alpha|}(\Omega)$, and

$$\forall n \in \mathbb{N} : \chi_n \equiv 1 \text{ on } M_n^c, \chi_n \equiv 0 \text{ on } M_{2n}, \text{ and } 0 \leq \chi_n \leq 1 \quad (3.3)$$

where M_n^c is the complementary (within Ω) of M_n . For any $\varphi \in \mathcal{D}(\Omega_{|v|^{p|\alpha|+p}}^*)$, we are allowed to use (3.2) with $\chi_n \varphi v^{-|\alpha|-1} \in \mathcal{C}_c^{|\alpha|}(\Omega_{|v|^{p|\alpha|+p}}^*)$ (see (iv) of Rem. 5). Thus, we obtain

$$\sum_{0 \leq \beta \leq \alpha} \binom{\alpha}{\beta} \int_{\Omega} f \partial^\beta \chi_n \partial^{\alpha-\beta} \varphi \, dx = (-1)^{|\alpha|} \int_{\Omega} \chi_n \varphi D_v^\alpha f \, dx.$$

By the continuous inclusions $L_{w,\text{loc}}^p(\Omega) \subseteq L_{v,\text{loc}}^p(\Omega) \subseteq L_{v^{|\alpha|+1},\text{loc}}^p(\Omega) \subseteq L_{\text{loc}}^1(\Omega_{|v|^{p|\alpha|+p}}^*)$ (the first inclusion follows from (3.1), and the latter inclusion follows from Thm. 1.5 of [5]) $f, D_v^\alpha f \in L_{\text{loc}}^1(\Omega_{|v|^{p|\alpha|+p}}^*)$. Hence, by the dominated convergence theorem, we have that

$$\int_{\Omega} \chi_n \varphi D_v^\alpha f \, dx \longrightarrow \int_{\Omega} \varphi D_v^\alpha f \, dx \text{ and } \int_{\Omega} f \chi_n \partial^\alpha \varphi \, dx \longrightarrow \int_{\Omega} f \partial^\alpha \varphi \, dx.$$

Left to show that

$$\forall \beta \in \pi_\alpha^0 : \int_{\Omega} f \partial^\beta (\chi_n) \partial^{\alpha-\beta} \varphi \, dx \longrightarrow 0. \quad (3.4)$$

Fix $\beta \in \pi_\alpha^0$. Note that $\partial^\beta \chi_n \rightarrow 0$ pointwise a.e. in Ω . On the other hand, the Faà di Bruno formula ([7]) gives

$$\partial^\beta \chi_n(x) = \sum_{k=1}^{|\beta|} n^k |\{\pi \in \Pi \mid |\pi| = k\}| \eta^{(k)}(nv(x)) \sum_{\pi \in \Pi, |\pi|=k} \prod_{B \in \pi} \frac{\partial^{|B|} v}{\prod_{j \in B} \partial x_j}$$

where

- π runs through the set Π of all partitions of the set \mathcal{A} of non-null indices of β with multiplicity e.g. if $\beta = (2, 3, 0, 1)$ then $\mathcal{A} := \{1, 1, 2, 2, 2, 4\}$

- “ $B \in \pi$ ” means the variable B runs through the list of all of the “blocks” of the partition π , and
- $|A|$ denotes the cardinality of the set A (so that $|\pi|$ is the number of blocks in the partition π and $|B|$ is the size of the block B).

By construction, $\partial^\beta \chi_n(x) = 0$ for all $x \in M_n^c \cup M_{2n}$. Let now $x \in M_n \cap M_{2n}^c$. Since $v \in \mathcal{C}^{|\alpha|}(\Omega)$, we have that

$$\exists C > 0 \forall n \in \mathbb{N} \forall x \in \text{supp}(\varphi) \cap M_n \cap M_{2n}^c : |\partial^\beta \chi_n(x)| \leq Cn^{|\beta|},$$

and hence

$$\exists C > 0 \forall n \in \mathbb{N} \forall x \in \text{supp}(\varphi) \cap M_n \cap M_{2n}^c : |v^{|\alpha|} \partial^\beta \chi_n(x)| \leq Cn^{|\beta| - |\alpha|} \leq C.$$

It follows that

$$\exists C > 0 \forall n \in \mathbb{N} : |f \partial^\beta (\chi_n) \partial^{\alpha-\beta} \varphi| \leq C |v^{-|\alpha|} f \partial^{\alpha-\beta} \varphi| \quad \text{a.e. in } \Omega.$$

Since $f \in L_{v, \text{loc}}^p(\Omega)$ (by (3.1)) and $v^{-|\alpha|-1} \in L_{\text{loc}}^{\frac{p}{p-1}}(\Omega_{|v|^{p|\alpha|+p}}^*)$ (which follows from the definition of the set $\Omega_{|v|^{p|\alpha|+p}}^*$), Hölder’s inequality implies that $v^{-|\alpha|} f \partial^{\alpha-\beta} \varphi \in L^1(\Omega)$. Therefore, (3.4) is now a consequence of the dominated convergence theorem.

Assume now that f is α -weakly differentiable in the sense of $L_{\text{loc}}^1(\Omega_{|v|^{p|\alpha|+p}}^*)$ with $D^\alpha f \in L_{v^{|\alpha|+1}, \text{loc}}^p(\Omega)$. Let χ_n be defined as above. Since $\chi_n v^{|\alpha|+1} \varphi \in \mathcal{C}_c^{|\alpha|}(\Omega_{v^{p|\alpha|+p}}^*)$, we have

$$\begin{aligned} \sum_{0 \leq \beta \leq \alpha} \binom{\alpha}{\beta} \int_{\Omega} f \partial^\beta (\chi_n) \partial^{\alpha-\beta} (v^{|\alpha|+1} \varphi) \, dx = \\ \int_{\Omega} f \partial^\alpha (v^{|\alpha|+1} \chi_n \varphi) \, dx = (-1)^{|\alpha|} \int_{\Omega} v^{|\alpha|+1} \chi_n \varphi D^\alpha f \, dx. \end{aligned} \quad (3.5)$$

One can easily show by induction that

$$\forall \beta \in \pi_\alpha \exists g_\beta \in \mathcal{C}_c^{|\beta|}(\Omega) : \partial^{\alpha-\beta} (v^{|\alpha|+1} \varphi) = v^{|\beta|+1} g_\beta. \quad (3.6)$$

As shown above, it is easy to see that

$$\forall \beta \in \pi_\alpha \exists C > 0 \forall n \in \mathbb{N} : |f \partial^{\alpha-\beta} (v^{|\alpha|+1} \varphi) \partial^\beta \chi_n| \leq C |f v g_\beta|$$

By the dominated convergence theorem (taking the limit in (3.5)) we get

$$\int_{\Omega} f \partial^\alpha (v^{|\alpha|+1} \varphi) \, dx = (-1)^{|\alpha|} \int_{\Omega} v^{|\alpha|+1} \varphi D^\alpha f \, dx,$$

which completes the proof. \square

4. THE WEIGHTED SOBOLEV SPACE $W_{V,v}^{m,p}(\Omega)$

If $V := \{w_\alpha\}_{\alpha \in \pi_m}$ is a collection of elements of $E_p(\Omega)$, then the function w_0 ($0 \in \mathbb{N}^d$) will be simply denoted by w .

Definition 8. Let $\Omega \subseteq \mathbb{R}^d$ be an open set, $p \in [1, \infty)$, $m \in \mathbb{N}$, and $V := \{w_\alpha\}_{\alpha \in \pi_m}$ be a collection of elements of $E_p(\Omega)$, $v \in \mathcal{C}^{m,*}(\Omega)$ be a map satisfying (3.1). We set

$$W_{V,v}^{m,p}(\Omega) := \{f \in L_w^p(\Omega) \mid \forall \alpha \in \pi_m^0 : \exists D_v^\alpha f \in L_{w_\alpha}^p(\Omega)\}.$$

This space is equipped with the norm

$$\forall f \in W_{V,v}^{m,p}(\Omega) : \|f\|_{W_{V,v}^{m,p}} := \left(\sum_{|\alpha| \leq m} \|D_v^\alpha f\|_{L_{w_\alpha}^p}^p \right)^{1/p}.$$

When $p = 2$, the space $W_{V,v}^{m,2}(\Omega)$ will be denoted by $H_{V,v}^m(\Omega)$ and its norm is generated by the scalar product

$$\forall f, g \in H_{V,v}^m(\Omega) : (f, g)_{H_{V,v}^m} := \sum_{|\alpha| \leq m} (D_v^\alpha f, D_v^\alpha g)_{L_{w_\alpha}^2}.$$

We now prove that the space $W_{V,v}^{m,p}(\Omega)$ is Cauchy complete.

Theorem 9. *Let $\Omega \subseteq \mathbb{R}^d$ be an open set, $p \in [1, \infty)$, $m \in \mathbb{N}$, and let $V := \{w_\alpha\}_{\alpha \in \pi_m}$ be a collection of elements of $E_p(\Omega)$, and $v \in \mathcal{C}^{m,*}(\Omega)$. Assume that*

$$\forall \alpha \in \pi_m \forall K \Subset \Omega \exists C_K : |v(x)|^{|\alpha|+1} \leq C_K |w_\alpha(x)| \text{ a.e. in } K. \quad (4.1)$$

Then, the space $W_{V,v}^{m,p}(\Omega)$ is Cauchy complete.

Proof. Let (f_n) be a Cauchy sequence of $W_{V,v}^{m,p}(\Omega)$. By definition, for all $\alpha \in \pi_m$, the sequence $(D_v^\alpha f_n)$ is a Cauchy sequence of $L_{w_\alpha}^p(\Omega)$. By completeness of $L_{w_\alpha}^p(\Omega)$ (see Thm. 2) there exists a unique $f_\alpha \in L_{w_\alpha}^p(\Omega)$ limit of the sequence $(D_v^\alpha f_n)$ in $L_{w_\alpha}^p(\Omega)$. Set $f := f_0$. We shall prove that f is weakly differentiable in the sense of $L_{v, \text{loc}}^1(\Omega)$ with $D_v^\alpha f = f_\alpha$ for all $\alpha \in \pi_m$. Indeed, by assumption we have

$$\forall \varphi \in \mathcal{D}(\Omega) : \int_\Omega f_n \partial^\alpha (v^{|\alpha|+1} \varphi) \, dx = (-1)^{|\alpha|} \int_\Omega \varphi v^{|\alpha|+1} D_v^\alpha f_n \, dx. \quad (4.2)$$

We have $\lim_{n \rightarrow \infty} \int_\Omega f_n \partial^\alpha (v^{|\alpha|+1} \varphi) \, dx = \int_\Omega f \partial^\alpha (v^{|\alpha|+1} \varphi) \, dx$ by (iii) of Rem. 5. On the other hand, (4.1) implies that

$$\left| \int_\Omega (D_v^\alpha f_n) v^{|\alpha|+1} \varphi \, dx \right| \leq C_{\text{supp}(\varphi)} \int_\Omega |(D_v^\alpha f_n) w_\alpha \varphi| \, dx \leq C_{\text{supp}(\varphi)} \|D_v^\alpha f_n\|_{L_{w_\alpha}^p} \|\varphi\|_{L^q}$$

where q is the conjugate exponent of p , which implies that the right hand side of (4.2) converges to $(-1)^{|\alpha|} \int_\Omega \varphi v^{|\alpha|+1} f_\alpha \, dx$ when $n \rightarrow \infty$. It follows that

$$\forall \varphi \in \mathcal{D}(\Omega) : \int_\Omega f \partial^\alpha (v^{|\alpha|+1} \varphi) \, dx = (-1)^{|\alpha|} \int_\Omega \varphi v^{|\alpha|+1} f_\alpha \, dx$$

which proves that $\exists D_v^\alpha f = f_\alpha \in L_{w_\alpha}^p(\Omega) \subseteq L_{v^{|\alpha|+1}, \text{loc}}^1(\Omega)$ (where the last inclusion follows from (4.1)), and hence $f \in W_{V,v}^{m,p}(\Omega)$. Therefore, the sequence (f_n) converges to f in $W_{V,v}^{m,p}(\Omega)$, and the proof is completed. \square

Remark 10. If $w_\alpha = |v|^\sigma$ for some $\sigma \in (0, |\alpha| + 1)$ then (4.1) holds. This is because $|v|^{|\alpha|+1-\sigma}$ is bounded on any compact subset of Ω .

We now compare the new approach of weighted Sobolev spaces presented in this paper with the classical one.

Let $V := \{w_\alpha\}_{\alpha \in \pi_m}$ be a collection of elements of $E_p(\Omega)$, and let $S := \{u_\alpha\}_{\alpha \in \pi_m}$ where $u_\alpha = |w_\alpha|^p \in W(\Omega)$. We recall that the weighted Sobolev space $W^{m,p}(\Omega, S)$ is defined as the space of all functions $f \in L^p(\Omega, u) \cap L_{\text{loc}}^1(\Omega)$ such that their

distributional derivative are elements of $L^p(\Omega, u_\alpha) \cap L^1_{\text{loc}}(\Omega)$. The space $W^{m,p}(\Omega, S)$ is equipped with the norm

$$\forall f \in W^{m,p}(\Omega, S) : \|f\|_{W^{m,p}(\Omega, S)} := \left(\sum_{|\alpha| \leq m} \|u_\alpha^{1/p} D^\alpha f\|_{L^p}^p \right)^{1/p}.$$

In order to guarantee the Cauchy completeness of $W^{m,p}(\Omega, S)$ we assume that

$$\forall \alpha \in \pi_m : u_\alpha^{-1} \in L^{\frac{1}{p-1}}_{\text{loc}}(\Omega) \quad (4.3)$$

(see Thm. 1.11 of [5]). However, when (4.3) is violated, the weighted Sobolev space $W^{m,p}(\Omega, S)$ is not necessarily a Banach space (see Example 1.12 of [5]). Indeed, the “bad” set which may cause the non-completeness of $W^{m,p}(\Omega, S)$ is the set $B := \cup_{\alpha \in \pi_m} P(u_\alpha)$ where $P(u_\alpha)$ is the set of points $x \in \Omega$ for which $u_\alpha^{-1} \notin L^{\frac{1}{p-1}}(\Omega \cap U)$ for all neighborhood U of x . The existing remedy is to define the space $W^{m,p}(\Omega, S)$ as the space $W^{m,p}(\Omega_S^*, S)$ where $\Omega_S^* := \Omega \setminus B$. It is shown that the set B is closed (and hence Ω_S^* is open) and Lebesgue negligible. Moreover, assumption (4.3) holds if we replace Ω with Ω_S^* . Therefore, the weighted Sobolev space $W^{m,p}(\Omega, S) (= W^{m,p}(\Omega_S^*, S))$ is now Cauchy complete (see Sec. 3 of [5] for more details). However, the space of test functions used in the definition of the distributional derivative is, in general, limited to the space $\mathcal{C}_c^m(\Omega_S^*)$.

On the other hand, if there exists a $\mathcal{C}^{m,*}(\Omega)$ function v that satisfies (4.1), then, by assertion (ii) of Prop. 7, we have that

$$W^{m,p}(\Omega_S^*, S) \subseteq W^{m,p}(\Omega_{|v|^{pm+p}}^*, S) = W_{V,v}^{m,p}(\Omega). \quad (4.4)$$

In case $\Omega_S^* = \Omega_{v^{pm+p}}^*$ e.g. $w_\alpha := v^{|\alpha|+1}$, the inclusion in (4.4) becomes an equality.

In conclusion, we have the following remark

Remark 11. For a given collection $S := \{|w_\alpha|^p\}_{\alpha \in \pi_m}$ of arbitrarily small weight functions, the weighted Sobolev space $W^{m,p}(\Omega_S^*, S)$ is a Banach space while the space $W^{m,p}(\Omega, S)$ in general is not. The price we pay is a modification of the domain and hence the convexity property might be lost and the boundary of Ω is modified. If there exists a $\mathcal{C}^{m,*}(\Omega)$ function v satisfying (4.1), then we can consider the space $W_{V,v}^{m,p}(\Omega)$ (where $V := \{w_\alpha\}_{\alpha \in \pi_m}$) instead of $W^{m,p}(\Omega_S^*, S)$, which is in general larger. In this case, no modification of the domain is needed and elements of $W_{V,v}^{m,p}(\Omega)$ are weakly differentiable in the sense of $L^1_{\text{loc}}(\Omega_{|v|^{pm+p}}^*)$ and in the sense of $L^1_{v,\text{loc}}(\Omega)$ but not necessary in the sense of $L^1_{\text{loc}}(\Omega_S^*)$. If furthermore, the function v is well chosen so that $\Omega_S^* = \Omega_{|v|^{pm+p}}^*$, then $W^{m,p}(\Omega_S^*, S) = W_{V,v}^{m,p}(\Omega)$.

Let $m \in \mathbb{N}$ and let $s : \pi_m \rightarrow \mathbb{R}_{\geq 0}$ be a map having the following properties

$$\forall \alpha \in \pi_m \forall \beta \in \pi_\alpha : s(\alpha) \leq |\alpha|, \text{ and } s(\alpha - \beta) + s(\beta) \leq s(\alpha) \quad (4.5)$$

A typical example of s is the map $|\cdot|$. Let $v \in \mathcal{C}^{m,*}(\Omega)$, and consider the collection $V = \{w_\alpha\}_{\alpha \in \pi_m}$ of elements of $E_p(\Omega)$ defined by

$$\forall \alpha \in \pi_m : w_\alpha := |v|^{s(\alpha)+1}.$$

Then, clearly (4.1) holds since $|\alpha| - s(\alpha) \geq 0$. It follows that the space $W_{s,v}^{m,p}(\Omega) := W_{V,v}^{m,p}(\Omega)$ is a Banach space (see Thm. 9).

Proposition 12. *Let $\Omega \subseteq \mathbb{R}^d$ be an open set, $p \in [1, \infty)$, $m \in \mathbb{N}$, $v \in \mathcal{C}^{m,*}(\Omega)$, $s : \pi_m \rightarrow \mathbb{R}_{\geq 0}$ be a map satisfying (4.5). Then, the operator $W_{s,v}^{m,p}(\Omega) \rightarrow W_{\text{loc}}^{m,p}(\Omega)$, $f \mapsto v^{m+1}f$ is continuous. Moreover*

$$\forall f \in W_{s,v}^{m,p}(\Omega) \forall \alpha \in \pi_m : D^\alpha(v^{m+1}f) = \sum_{0 \leq \beta \leq \alpha} \binom{\alpha}{\beta} D_v^\beta f \partial^{\alpha-\beta} v^{m+1}. \quad (4.6)$$

In particular, one can replace $W_{\text{loc}}^{m,p}(\Omega)$ with $W^{m,p}(\Omega)$ in case $v \in \mathcal{C}_b^m(\Omega)$.

Proof. Let $f \in W_{s,v}^{m,p}(\Omega)$ and $\alpha \in \pi_m$. For any $\sigma \in \pi_\alpha$, there exists $g_\sigma \in \mathcal{C}^{m-|\sigma|}(\Omega)$ such that $\partial^\sigma v^{m+1} = g_\sigma v^{m+1-|\sigma|}$, which implies that

$$\forall K \Subset \Omega : \|D_v^\beta f \partial^{\alpha-\beta} v^{m+1}\|_{L^p(K)} \leq C \|v^{m-|\alpha|} g_{\alpha-\beta}\|_{L^\infty(K)} \left\| |v|^{s(\beta)+1} D_v^\beta f \right\|_{L^p(\Omega)} \quad (4.7)$$

where we used (4.1) (with $w_\beta = |v|^{s(\beta)+1}$) in the last inequality, which shows that the right hand side of (4.6) belongs to $L_{\text{loc}}^p(\Omega)$ for all $\alpha \in \pi_m$. In case $v \in \mathcal{C}_b^m(\Omega)$, $v^{m-|\alpha|} g_{\alpha-\beta} \in L^\infty(\Omega)$ and the constant C does not depend on the compact set K . Thus, (4.7) holds with Ω instead of K . Therefore, the right hand side of (4.6) belongs to $L^p(\Omega)$ for all $\alpha \in \pi_m$. We show now that $v^{m+1}f$ is weakly differentiable in the sense of $L_{\text{loc}}^1(\Omega)$ and that the Leibniz formula (4.6) holds. Indeed, we will prove this by induction. Suppose first that $\alpha \in \pi_1^0$. Choose $\varphi \in \mathcal{D}(\Omega)$. We have

$$\begin{aligned} \int_{\Omega} f v^{m+1} \partial^\alpha \varphi \, dx &= \int_{\Omega} f (\partial^\alpha (v^{m+1} \varphi) - \varphi \partial^\alpha v^{m+1}) \, dx \\ &= - \int_{\Omega} (v^{m+1} D_v^\alpha f + f \partial^\alpha v^{m+1}) \varphi \, dx, \end{aligned}$$

where we used (3.2) with the test function $\varphi v^{m-|\alpha|} \in \mathcal{C}_c^m(\Omega)$, which proves (4.6) when $\alpha \in \pi_1^0$. We assume now that (4.6) holds for all $\alpha \in \pi_l^0$ for some $l < m$. Choose now $\alpha \in \pi_{l+1}^l$. Then $\alpha = \beta + \gamma$ for some $|\beta| = l$ and $|\gamma| = 1$. Then for φ as above,

$$\begin{aligned} \int_{\Omega} f v^{m+1} \partial^\alpha \varphi \, dx &= (-1)^{|\beta|} \int_{\Omega} D^\beta (f v^{m+1}) \partial^\gamma \varphi \, dx \\ &= (-1)^{|\beta|} \sum_{0 \leq \sigma \leq \beta} \binom{\beta}{\sigma} \int_{\Omega} D_v^\sigma f \partial^{\beta-\sigma} v^{m+1} \partial^\gamma \varphi \, dx \end{aligned}$$

where we used the induction assumption. Using the equality

$$\partial^{\beta-\sigma} v^{m+1} \partial^\gamma \varphi = \partial^\gamma (\varphi \partial^{\beta-\sigma} v^{m+1}) - \varphi \partial^{\alpha-\sigma} v^{m+1}$$

we get

$$\begin{aligned} \int_{\Omega} f v^{m+1} \partial^\alpha \varphi \, dx &= (-1)^{|\beta|} \sum_{0 \leq \sigma \leq \beta} \binom{\beta}{\sigma} \int_{\Omega} D_v^\sigma f \partial^\gamma (\varphi \partial^{\beta-\sigma} v^{m+1}) \, dx \\ &\quad + (-1)^{|\alpha|} \sum_{0 \leq \sigma \leq \beta} \binom{\beta}{\sigma} \int_{\Omega} \varphi D_v^\sigma f \partial^{\alpha-\sigma} v^{m+1} \, dx. \end{aligned}$$

Again, using (3.2) we obtain

$$\int_{\Omega} D_v^\sigma f \partial^\gamma (\varphi \partial^{\beta-\sigma} v^{m+1}) \, dx = - \int_{\Omega} \varphi \partial^{\beta-\sigma} v^{m+1} D_v^{\gamma+\sigma} f \, dx$$

Thus

$$\int_{\Omega} f v^{m+1} \partial^{\alpha} \varphi \, dx = (-1)^{|\alpha|} \sum_{0 \leq \sigma \leq \beta} \binom{\beta}{\sigma} \int_{\Omega} \varphi (D_v^{\rho} f \partial^{\alpha-\rho} v^{m+1} + D_v^{\sigma} f \partial^{\alpha-\sigma} v^{m+1}) \, dx,$$

where $\rho = \sigma + \gamma$. It follows that

$$\int_{\Omega} f v^{m+1} \partial^{\alpha} \varphi \, dx = (-1)^{|\alpha|} \sum_{0 \leq \sigma \leq \alpha} \binom{\alpha}{\sigma} \int_{\Omega} \varphi (D_v^{\sigma} f \partial^{\alpha-\sigma} v^{m+1}) \, dx,$$

since

$$\binom{\beta}{\sigma-\gamma} + \binom{\beta}{\sigma} = \binom{\alpha}{\sigma},$$

Therefore, the map $W_{s,v}^{m,p}(\Omega) \rightarrow W_{\text{loc}}^{m,p}(\Omega)$, $f \rightarrow v^{m+1} f$ is well defined and (4.6) holds. Moreover, the continuity follows from (4.6) and (4.7), which completes the proof. \square

5. DENSITY OF SMOOTH FUNCTIONS

In this section, we are interested in finding sufficient conditions under which smooth functions are dense in weighted Sobolev spaces $W_{s,v}^{m,p}(\Omega)$.

Let $v \in \mathcal{C}^{m,*}(\Omega)$, and let $s : \pi_m \rightarrow \mathbb{R}_{\geq 0}$ be a map satisfying (4.5). In order to prove the density of smooth functions in $W_{s,v}^{m,p}(\Omega)$, we show first that

$$\forall f \in W_{s,v}^{m,p}(\Omega) \, \forall n \in \mathbb{N} : f \chi_n \in W^{m,p}(\Omega), \quad (5.1)$$

where (χ_n) is the sequence constructed in Prop. 7. After that, we show that the sequence $(f \chi_n)$ converges to f in $W_{s,v}^{m,p}(\Omega)$. Then, we use the density results of smooth functions in Sobolev spaces $W^{m,p}(\Omega)$ to conclude.

Theorem 13. *Let $\Omega \subseteq \mathbb{R}^d$ be an open set, $p \in [1, \infty)$, $m \in \mathbb{N}$, $v \in \mathcal{C}^{m,*}(\Omega)$, $s : \pi_m \rightarrow \mathbb{R}_{\geq 0}$ be a map satisfying (4.5). Assume that*

$$\forall \alpha \in \pi_m^0 \, \exists C > 0 \, \forall x \in M_{2n}^c \cap M_n : |\partial^{\alpha} v(x)| \leq C. \quad (5.2)$$

Then, the mapping $W_{s,v}^{m,p}(\Omega) \rightarrow W^{m,p}(\Omega)$, $f \rightarrow \chi_n f$ is continuous. In particular, for any $f \in W_{s,v}^{m,p}(\Omega)$ and for all $\alpha \in \pi_m^0$ we have

$$D^{\alpha}(f \chi_n) = \sum_{0 \leq \beta \leq \alpha} \binom{\alpha}{\beta} D_v^{\beta} f \partial^{\alpha-\beta} \chi_n. \quad (5.3)$$

Proof. First note that $\chi_n \in \mathcal{C}_b^m(\Omega)$ i.e. $\partial^{\alpha} \chi_n$ is bounded on Ω for all $\alpha \in \pi_m$. This is because $\eta_n^{(k)} \equiv 0$ on $M_{2n} \cup M_n^c$ for all $k \neq 0$, and $\partial^{\alpha} v$ is bounded on $M_{2n}^c \cap M_n$. Let $f \in W_{s,v}^{m,p}(\Omega)$. We show now that $D_v^{\beta} f \partial^{\alpha-\beta} \chi_n \in L^p(\Omega)$ for all $\alpha \in \pi_m$ and for all $\beta \in \pi_{\alpha}$. Indeed, since $\text{supp}(\chi_n) \subseteq M_{2n}^c$ we have

$$\begin{aligned} \|D_v^{\beta} f \partial^{\alpha-\beta} \chi_n\|_{L^p(\Omega)} &= \| |v|^{-s(\beta)-1} \partial^{\alpha-\beta} \chi_n |v|^{s(\beta)+1} D_v^{\beta} f \|_{L^p(\Omega)} \\ &\leq (2n)^{s(\beta)+1} \|\partial^{\alpha-\beta} \chi_n\|_{L^{\infty}(\Omega)} \| |v|^{s(\beta)+1} D_v^{\beta} f \|_{L^p(\Omega)}. \end{aligned} \quad (5.4)$$

We claim now that $\chi_n f$ is α -weakly differentiable in the sense of $L_{\text{loc}}^1(\Omega)$ for all $\alpha \in \pi_m^0$ and that (5.3) holds. Indeed, we will prove this by induction as we did for (4.6). Suppose first that $\alpha \in \pi_1^0$. Choose $\varphi \in \mathcal{D}(\Omega)$. Using (3.2) with the test function $\varphi \chi_n v^{-|\alpha|-1} \in \mathcal{C}_c^m(\Omega)$ (because $\chi_n \equiv 0$ on M_{2n}) we get

$$\int_{\Omega} f \partial^{\alpha}(\chi_n \varphi) \, dx = - \int_{\Omega} \chi_n \varphi D_v^{\alpha} f \, dx.$$

It follows that

$$\int_{\Omega} f \chi_n \partial^{\alpha} \varphi \, dx = \int_{\Omega} f (\partial^{\alpha} (\chi_n \varphi) - \varphi \partial^{\alpha} \chi_n) \, dx = - \int_{\Omega} (\chi_n D_v^{\alpha} f + f \partial^{\alpha} \chi_n) \varphi \, dx,$$

which proves our claim when $\alpha \in \pi_1^0$. We assume now that (5.3) holds for all $\alpha \in \pi_l^0$ for some $l < m$. Choose now $\alpha \in \pi_{l+1}^l$. Then $\alpha = \beta + \gamma$ for some $|\beta| = l$ and $|\gamma| = 1$. Then for φ as above,

$$\begin{aligned} \int_{\Omega} f \chi_n \partial^{\alpha} \varphi \, dx &= (-1)^{|\beta|} \int_{\Omega} D^{\beta} (f \chi_n) \partial^{\gamma} \varphi \, dx \\ &= (-1)^{|\beta|} \sum_{0 \leq \sigma \leq \beta} \binom{\beta}{\sigma} \int_{\Omega} D_v^{\sigma} f \partial^{\beta-\sigma} \chi_n \partial^{\gamma} \varphi \, dx \end{aligned}$$

where we used the induction assumption. Using the equality

$$\partial^{\beta-\sigma} \chi_n \partial^{\gamma} \varphi = \partial^{\gamma} (\varphi \partial^{\beta-\sigma} \chi_n) - \varphi \partial^{\alpha-\sigma} \chi_n$$

we get

$$\begin{aligned} \int_{\Omega} f \chi_n \partial^{\alpha} \varphi \, dx &= (-1)^{|\beta|} \sum_{0 \leq \sigma \leq \beta} \binom{\beta}{\sigma} \int_{\Omega} D_v^{\sigma} f \partial^{\gamma} (\varphi \partial^{\beta-\sigma} \chi_n) \, dx \\ &\quad + (-1)^{|\alpha|} \sum_{0 \leq \sigma \leq \beta} \binom{\beta}{\sigma} \int_{\Omega} \varphi D_v^{\sigma} f \partial^{\alpha-\sigma} \chi_n \, dx. \end{aligned}$$

Again, using (3.2) we obtain

$$\int_{\Omega} D_v^{\sigma} f \partial^{\gamma} (\varphi \partial^{\beta-\sigma} \chi_n) \, dx = - \int_{\Omega} \varphi \partial^{\beta-\sigma} \chi_n D_v^{\gamma+\sigma} f \, dx$$

Thus

$$\int_{\Omega} f \chi_n \partial^{\alpha} \varphi \, dx = (-1)^{|\alpha|} \sum_{0 \leq \sigma \leq \beta} \binom{\beta}{\sigma} \int_{\Omega} \varphi (D_v^{\rho} f \partial^{\alpha-\rho} \chi_n + D_v^{\sigma} f \partial^{\alpha-\sigma} \chi_n) \, dx,$$

where $\rho = \sigma + \gamma$. It follows that

$$\int_{\Omega} f \chi_n \partial^{\alpha} \varphi \, dx = (-1)^{|\alpha|} \sum_{0 \leq \sigma \leq \alpha} \binom{\alpha}{\sigma} \int_{\Omega} \varphi (D_v^{\sigma} f \partial^{\alpha-\sigma} \chi_n) \, dx,$$

since

$$\binom{\beta}{\sigma-\gamma} + \binom{\beta}{\sigma} = \binom{\alpha}{\sigma},$$

Therefore, the map $W_{s,v}^{m,p}(\Omega) \rightarrow W^{m,p}(\Omega)$, $f \rightarrow \chi_n f$ is well defined and (5.3) holds. Moreover, the continuity follows from (5.3) and (5.4), completing the proof. \square

Remark 14. Using Thm. 13 with $m = 0$ we get that $\chi_n f \in L^p(\Omega)$. By the dominated convergence theorem we conclude that $\chi_n f \rightarrow f$ in $L_v^p(\Omega)$ and hence $L^p(\Omega)$ is dense in $L_v^p(\Omega)$. If v is bounded on Ω , then $\mathcal{D}(\Omega)$ is dense in $L_v^p(\Omega)$.

Another way to prove the density of $\mathcal{D}(\Omega)$ in $L_w^p(\Omega)$ ($w \in E_p(\Omega)$) is to assume that $w \in C^{\infty}(\Omega)$. In this setting, given a sequence (φ_n) of $\mathcal{D}(\Omega)$ that converges to wf in $L^2(\Omega)$, the sequence $(\varphi_n \chi_n w^{-1})$ (χ_n defined with $v = w$) of $\mathcal{D}(\Omega)$ converges to f in $L_w^p(\Omega)$.

The aim now is to show that the sequence $(\chi_n f)$ of $W^{m,p}(\Omega)$ (provided assumption (5.2) holds) converges to f in $W_{s,v}^{m,p}(\Omega)$ when v is bounded. More precisely, we plan to show that

$$\forall \alpha \in \pi_m : |v|^{s(\alpha)+1} D^\alpha(f \chi_n) = |v|^{s(\alpha)+1} \sum_{0 \leq \beta \leq \alpha} \binom{\alpha}{\beta} D_v^\beta f \partial^{\alpha-\beta} \chi_n \rightarrow |v|^{s(\alpha)+1} D_v^\alpha f$$

with convergence in $L^p(\Omega)$, where we used (5.3) in the first equality. Since $\chi_n |v|^{s(\beta)+1} D_v^\beta f \rightarrow |v|^{s(\beta)+1} D_v^\beta f$ in $L^p(\Omega)$ for all $\beta \in \pi_m$ by the dominated convergence theorem, it suffices to show that

$$\forall \alpha \in \pi_m \forall \beta \in \pi_\alpha \setminus \{\alpha\} : |v|^{s(\beta)+1} (D_v^\beta f) |v|^{s(\alpha)-s(\beta)} \partial^{\alpha-\beta} \chi_n \rightarrow 0 \text{ in } L^p(\Omega).$$

However, since

- (i) $\partial^\sigma \chi_n \rightarrow 0$ pointwise a.e. in Ω for all $\sigma \in \pi_m^0$,
- (ii) $|v|^{s(\beta)+1} (D_v^\beta f) \in L^p(\Omega)$ for all $\beta \in \pi_m$,
- (iii) $\partial^\sigma \chi_n = 0$ on $M_n^c \cup M_{2n}$ and $|v|^{s(\sigma)} \leq n^{-s(\sigma)}$ on $M_n \cap M_{2n}^c$ for all $\sigma \in \pi_m^0$,
- (iv) $s(\alpha) - s(\beta) \geq s(\alpha - \beta)$ and hence $|v|^{s(\alpha)-s(\beta)} \leq C |v|^{s(\alpha-\beta)}$ for all $x \in \Omega$ for some C independent of x (since v is bounded on Ω)

it suffices to have the following property

$$\exists C > 0 \forall n \in \mathbb{N} \forall \sigma \in \pi_m^0 \forall x \in M_n \cap M_{2n}^c : |\partial^\sigma \chi_n| \leq C n^{s(\sigma)} \quad (5.5)$$

and conclude then by the dominated convergence theorem.

In the next lemma we give an assumption on v that yields (5.5).

Lemma 15. *Let $\Omega \subseteq \mathbb{R}^d$ be an open set, $p \in [1, \infty)$, $m \in \mathbb{N}$, $v \in \mathcal{C}^{m,*}(\Omega)$, $s : \pi_m \rightarrow \mathbb{R}_{\geq 0}$ be a map satisfying (4.5). Assume that*

$$\exists C > 0 \forall n \in \mathbb{N} \forall \sigma \in \pi_m^0 \forall x \in M_n \cap M_{2n}^c : |\partial^\sigma v(x)| \leq C n^{s(\sigma)-1}. \quad (5.6)$$

Then, (5.5) holds.

Proof. Fix $n \in \mathbb{N}$ and let $\sigma \in \pi_m^0$. By the Faà di Bruno formula ([7]), for any $x \in M_n \cap M_{2n}^c$ we have

$$\partial^\sigma \chi_n(x) = \sum_{k=1}^{|\sigma|} |\{\pi \in \Pi \mid |\pi| = k\}| \eta_n^{(k)}(v(x)) \sum_{\pi \in \Pi, |\pi|=k} \prod_{B \in \pi} \frac{\partial^{|B|} v}{\prod_{j \in B} \partial x_j}.$$

(See the proof of (ii) of Prop. 7 for more details). By construction of the sequence (η_n) we have

$$\forall k \in \mathbb{N} \exists C > 0 \forall n \in \mathbb{N} \forall x \in \mathbb{R} : |\eta_n^{(k)}(x)| \leq C n^k.$$

For any $B \in \pi$, we associate $\beta \in \mathbb{N}^d$ by setting $\beta_j := |\{j \in B\}|$ for all j . It follows that

$$|\partial^\sigma \chi_n(x)| \leq C \sum_{k=1}^{|\sigma|} C_k n^k \sum_{\pi \in \Pi, |\pi|=k} \prod_{B \in \pi} n^{s(\beta)-1} \quad (5.7)$$

where we used assumption (5.6). It is easy to see that

$$\forall \pi \in \Pi : \prod_{B \in \pi} n^{s(\beta)-1} = n^{\sum_{B \in \pi} s(\beta) - |\pi|} \leq n^{s(\sigma) - |\pi|}$$

where we used (4.5), which gives (5.5) when used in (5.7). \square

Theorem 16. *Let $\Omega \subseteq \mathbb{R}^d$ be an open set, $p \in [1, \infty)$, $m \in \mathbb{N}$, and $v \in \mathcal{C}^{m,*}(\Omega) \cap L^\infty(\Omega)$, $s : \pi_m \longrightarrow \mathbb{R}_{\geq 0}$ be a map satisfying (4.5). Assume that (5.6) holds. Then, for any $f \in W_{s,v}^{m,p}(\Omega)$, the sequence $(\chi_n f)$ of $W^{m,p}(\Omega)$ (by Thm. 13) converges to f in $W_{s,v}^{m,p}(\Omega)$.*

Proof. It follows from Lem. 15 and the paragraph preceding it. \square

We deduce from this theorem the following density results

Corollary 17. *Under the assumptions of Thm. 16, we have*

- (i) *The space $\mathcal{D}(\Omega)$ is dense in $W_{s,v}^{m,p}(\Omega)$ when $\Omega = \mathbb{R}^d$.*
- (ii) *The space $\mathcal{C}^\infty(\Omega) \cap W^{m,p}(\Omega)$ is dense in $W_{s,v}^{m,p}(\Omega)$ in case Ω is bounded.*

Proof. Let $f \in W_{s,v}^{m,p}(\mathbb{R}^d)$. By Thm. 16, the sequence $(\chi_n f)$ of $W^{m,p}(\mathbb{R}^d)$ converges to f in $W_{s,v}^{m,p}(\mathbb{R}^d)$. By density of $\mathcal{D}(\mathbb{R}^d)$ in $W^{m,p}(\mathbb{R}^d)$, for all n there exists a sequence $(\varphi_{n,k})_k$ of $\mathcal{D}(\mathbb{R}^d)$ that converges to $\chi_n f$ in $W^{m,p}(\mathbb{R}^d)$, and hence in $W_{s,v}^{m,p}(\mathbb{R}^d)$ since $v \in L^\infty(\mathbb{R}^d)$, which proves assertion (i).

Similarly, to prove assertion (ii), we use Thm. 16 and the fact that the space $\mathcal{C}^\infty(\Omega) \cap W^{m,p}(\Omega)$ is dense in $W^{m,p}(\Omega)$ when Ω is bounded (see Thm. 2 of Subsection 5.3.2 of [3]). \square

Remark 18. The boundedness assumption of v in Thm. 16 can be dropped in case $s(\alpha) = |\alpha|$ since the assumption was used to guarantee that the constant C in (4.1) is independent of K .

6. THE TRACE OPERATOR

Now, we discuss the different possibilities to define a trace operator on $W_{s,v}^{1,p}(\Omega)$

Remark 19. Let $\Omega \subseteq \mathbb{R}^d$ be a bounded open set with a sufficiently smooth boundary¹, e.g. $\partial\Omega$ is \mathcal{C}^1 , $p \in [1, \infty)$, $v \in \mathcal{C}^{1,*}(\Omega)$, $s : \pi_1 \longrightarrow \mathbb{R}_{\geq 0}$ be a map satisfying (4.5).

- (i) Assume that

$$\exists K \Subset \Omega \exists \delta > 0 \forall x \in \Omega \setminus K : |v(x)| > \delta \quad (6.1)$$

(or equivalently $M_{n_0} \Subset \Omega$ for some $n_0 \in \mathbb{N}$). Fix $n \geq n_0$. Then, (5.2) holds and $\partial\Omega \subseteq \partial M_n^c$. By Thm. 13 we have

$$\chi_n f \in W^{1,p}(\Omega) \text{ and } \forall \alpha \in \pi_1^0 : D^\alpha(\chi_n f) = f \partial^\alpha \chi_n + \chi_n D_v^\alpha f. \quad (6.2)$$

If we denote by $|\partial\Omega : W^{1,p}(\Omega) \longrightarrow L^p(\partial\Omega)$ the trace operator, then $(f \chi_n)|_{\partial\Omega}$ is well defined. Since $\chi_n \equiv 1$ on $\Omega \setminus K \subseteq M_n^c$, we have that $(f \chi_n)|_{\partial\Omega} = f|_{\partial\Omega}$ which implies that $|\partial\Omega$ can be extended to the space $W_{s,v}^{1,p}(\Omega)$. Furthermore, by Thm. 13 and the trace theorem in $W^{1,p}(\Omega)$ we get

$$\|f|_{\partial\Omega}\|_{L^p(\partial\Omega)} \leq C \|\chi_n f\|_{W^{1,p}(\Omega)} \leq C_n \|f\|_{W_{s,v}^{1,p}(\Omega)} \quad (6.3)$$

where C_n is a constant that depends on n but not in f , which proves that $|\partial\Omega$ is continuous on $W_{s,v}^{1,p}(\Omega)$.

¹so that the trace operator on $W^{1,p}(\Omega)$ is well defined.

- (ii) If we assume that $v \in \mathcal{C}_b^1(\Omega)$, then by Prop. 12, we have that the map $W_{s,v}^{1,p}(\Omega) \longrightarrow W^{1,p}(\Omega)$, $f \longrightarrow v^2 f$ is continuous. In particular $(v^2 f)|_{\partial\Omega}$ is well defined and belongs to $L^p(\partial\Omega)$. Moreover, by the trace theorem in $W^{1,p}(\Omega)$ we have

$$\|(v^2 f)|_{\partial\Omega}\|_{L^p(\partial\Omega)} \leq C \|v^2 f\|_{W^{1,p}(\Omega)} \leq C \|f\|_{W_{s,v}^{1,p}(\Omega)} \quad (6.4)$$

Therefore, the operator $W_{s,v}^{1,p}(\Omega) \longrightarrow L^p(\partial\Omega)$, $f \longrightarrow (v^2 f)|_{\partial\Omega}$ is continuous.

Therefore, we can define the trace operator on the space $W_{s,v}^{1,p}(\Omega)$ either by setting $\text{Tr}(f) = f|_{\partial\Omega} \in L^p(\partial\Omega)$, or by setting $\text{Tr}(f) = (v^2 f)|_{\partial\Omega} \in L^p(\partial\Omega)$ depending on the assumptions satisfied by the function v .

Remark 20. Note that in the favorite situation in which assumption (6.1) holds and $v \in \mathcal{C}_b^1(\Omega)$, both $f|_{\partial\Omega}$ and $(v^2 f)|_{\partial\Omega}$ are well defined and belong to $L^p(\partial\Omega)$ for all $f \in W_{s,v}^{1,p}(\Omega)$. Moreover, both v and v^{-1} are bounded on $\partial\Omega$, which implies that the operators $W_{s,v}^{1,p}(\Omega) \longrightarrow L^p(\partial\Omega)$, $f \longrightarrow f|_{\partial\Omega}$, $f \longrightarrow (f v^2)|_{\partial\Omega}$ are equivalent, and hence $f|_{\partial\Omega} = 0$ if and only if $(v^2 f)|_{\partial\Omega} = 0$.

These different possibilities motivate the following definition

Definition 21. Let $\Omega \subseteq \mathbb{R}^d$ be a bounded open set and $\partial\Omega$ is \mathcal{C}^1 , $p \in [1, \infty)$, $v \in \mathcal{C}^{1,*}(\Omega)$, $s : \pi_1 \longrightarrow \mathbb{R}_{\geq 0}$ be a map satisfying (4.5).

- (i) If assumption (6.1) holds, then

$$\forall f \in W_{s,v}^{1,p}(\Omega) : \text{Tr}_1(f) := f|_{\partial\Omega} \in L^p(\partial\Omega) \quad (6.5)$$

- (ii) If $v \in \mathcal{C}_b^1(\Omega)$, then

$$\forall f \in W_{s,v}^{1,p}(\Omega) : \text{Tr}_2(f) := (v^2 f)|_{\partial\Omega} \in L^p(\partial\Omega). \quad (6.6)$$

As we have seen in the beginning of this section, under the assumptions given in the latter definition, the trace operators are continuous (see (6.3) and (6.4)).

7. SPACES OF NULL TRACE

Definition 22. Let $\Omega \subseteq \mathbb{R}^d$ be an open set, $p \in [1, \infty)$, $V := \{w_\alpha\}_{\alpha \in \pi_1}$ be a collection of elements of $E_p(\Omega)$, and let $v \in \mathcal{C}^{1,*}(\Omega)$ be such that (4.1) holds; $X_{V,v,0}^{1,p}(\Omega)$ denotes the closure of $\mathcal{D}(\Omega)$ in $W_{V,v}^{1,p}(\Omega)$.

The assumption (4.1) guarantees that the weighted Sobolev space $W_{V,v}^{1,p}(\Omega)$ is Cauchy complete, and hence $X_{V,v,0}^{1,p}(\Omega)$ is well defined.

We now define the spaces $W_{s,v,0}^{1,p,i}$ ($i = 1, 2$) using the trace operators

Definition 23. Let Ω be a bounded open set and $\partial\Omega$ is \mathcal{C}^1 , $p \in [1, \infty)$, $v \in \mathcal{C}^{1,*}(\Omega)$, $s : \pi_1 \longrightarrow \mathbb{R}_{\geq 0}$ be a map satisfying (4.5).

- (i) Assume that (6.1) holds. Then we set

$$W_{s,v,0}^{1,p,1}(\Omega) := \{f \in W_{s,v}^{1,p}(\Omega) \mid \text{Tr}_1(f) = 0\}.$$

- (ii) Assume that $v \in \mathcal{C}_b^1(\Omega)$. Then we set

$$W_{s,v,0}^{1,p,2}(\Omega) := \{f \in W_{s,v}^{1,p}(\Omega) \mid \text{Tr}_2(f) = 0\}.$$

Remark 24. Recall from Rem. 20 that the operators Tr_1 , Tr_2 are equivalent when $v \in \mathcal{C}_b^1(\Omega)$ and (6.1) holds, which implies that

$$W_{s,v,0}^{1,p,1}(\Omega) = W_{s,v,0}^{1,p,2}(\Omega).$$

Proposition 25. *Let Ω be a bounded open set and $\partial\Omega$ is \mathcal{C}^1 , $p \in [1, \infty)$, $v \in \mathcal{C}^{1,*}(\Omega)$, $s : \pi_1 \longrightarrow \mathbb{R}_{\geq 0}$ be a map satisfying (4.5).*

- (i) *If (6.1) holds, then the space $W_{s,v,0}^{1,p,1}(\Omega)$ is Cauchy complete.*
- (ii) *If $v \in \mathcal{C}_b^1(\Omega)$, then the space $W_{s,v,0}^{1,p,2}(\Omega)$ is Cauchy complete.*

Proof. It is sufficient to prove that the space $W_{s,v,0}^{1,p,i}(\Omega)$ ($i = 1, 2$) is closed since it is a subspace of a Banach space $W_{s,v}^{1,p}(\Omega)$. However, the closedness follows from the fact that $W_{s,v,0}^{1,p,i}(\Omega) = \text{Ker}(\text{Tr}_i)$ and Tr_i is continuous. \square

Theorem 26. *Let Ω be a bounded open set and $\partial\Omega$ is \mathcal{C}^1 , $p \in [1, \infty)$, $v \in \mathcal{C}^{1,*}(\Omega)$, $s : \pi_1 \longrightarrow \mathbb{R}_{\geq 0}$ be a map satisfying (4.5).*

- (i) *If v is bounded and (6.1) holds, then $X_{s,v,0}^{1,p}(\Omega) = W_{s,v,0}^{1,p,1}(\Omega)$.*
- (ii) *If $v \in \mathcal{C}_b^1(\Omega)$ then $X_{s,v,0}^{1,p}(\Omega) = W_{s,v,0}^{1,p,2}(\Omega)$.*

Proof. For $i = 1, 2$, the inclusion $X_{s,v,0}^{1,p}(\Omega) \subseteq W_{s,v,0}^{1,p,i}(\Omega)$ follows from the fact that $\text{Tr}_i(\varphi) = 0$ for all $\varphi \in \mathcal{D}(\Omega)$, and Tr_i is continuous.

We deal now with the opposite inclusion. Claim first that $\chi_n f \in W_0^{1,p}(\Omega)$ and that the sequence $(\chi_n f)$ converges to f in $W_{s,v}^{1,p}(\Omega)$ for all $f \in W_{s,v,0}^{1,p,i}(\Omega)$ ($i = 1, 2$). Indeed, in case (6.1) holds, the first part of the claim follows from the definition of the space $W_{s,v,0}^{1,p,1}(\Omega)$ (see also (i) of Rem. 19). In case $v \in \mathcal{C}_b^1(\Omega)$, $\chi_n f \in W^{1,p}(\Omega)$ by Thm. (13). Moreover, we have that $(f\chi_n)|_{\partial M_{2n} \cap \partial\Omega} = 0$ because $\chi_n \equiv 0$ on M_{2n} , and $(f\chi_n)|_{\partial M_{2n}^c \cap \partial\Omega} = (fv^2\chi_n v^{-2})|_{\partial M_{2n}^c \cap \partial\Omega} = 0$ because $|v^{-2}| \leq (2n)^2$ on M_{2n}^c , which proves that $f\chi_n \in W_0^{1,p}(\Omega)$, and hence the first part of the claim is proved. The second part of the claim is a consequence of Thm. 16. Finally, by density of $\mathcal{D}(\Omega)$ in $W_0^{1,p}(\Omega)$, and since v is bounded, we conclude that $\mathcal{D}(\Omega)$ is dense $W_{s,v,0}^{1,p,i}(\Omega)$. \square

In the next lemma, we consider a functional of the form $X_{s,v,0}^{1,p}(\Omega) \longrightarrow \mathbb{R}$, $f \longrightarrow \int_{\Omega} g D^{\alpha}(v^3 f) dx$ where $v \in \mathcal{C}^{1,*}(\Omega)$, $s : \pi_1 \longrightarrow \mathbb{R}_{\geq 0}$ is a map satisfying (4.5), and $g \in W_{s,v}^{1,p'}(\Omega)$ (p' is the exponent conjugate to p). We show in particular that

$$\forall f \in X_{s,v,0}^{1,p}(\Omega) \forall \alpha \in \pi_1^0 : \int_{\Omega} g D^{\alpha}(v^3 f) dx = - \int_{\Omega} v^3 f D_v^{\alpha} g dx \quad (7.1)$$

which implies that the functional $f \longrightarrow \int_{\Omega} g D^{\alpha}(v^3 f) dx$ is continuous. As far as we know, the latter functional and the integration by part formula (7.1) was not considered in the literature.

Lemma 27. *Let $\Omega \subseteq \mathbb{R}^d$ be a bounded open set, $v \in \mathcal{C}^{1,*}(\Omega)$, $s : \pi_1 \longrightarrow \mathbb{R}_{\geq 0}$ be a map satisfying (4.5). Assume that*

- (i) *$v \in L^{\infty}(\Omega)$*
- (ii) *$a \in L_{v^{-3}}^{\infty}(\Omega)$ with $av^{-2} \in \mathcal{C}^1(\Omega)$ and $\partial^{\alpha} a \in L_{v^{-2}}^{\infty}(\Omega)$ for all $\alpha \in \pi_1^0$.*

Then

$$\forall q \in [1, \infty) \forall g \in W_{s,v}^{1,q}(\Omega) : ag \in W^{1,q}(\Omega) \text{ and } D^{\alpha}(ag) = g\partial^{\alpha}a + aD_v^{\alpha}g.$$

Moreover,

$$\forall h \in W_{s,v}^{1,p}(\Omega) \forall f \in X_{s,v,0}^{1,p'}(\Omega) : \int_{\Omega} h D^{\alpha}(af) dx = - \int_{\Omega} af D_v^{\alpha} h dx \quad (7.2)$$

for all $p \in (1, \infty)$, where p' is the exponent conjugate to p .

Proof. The proof of the first part of the lemma is very similar to the proof of Thm. 12. We prove now (7.2). By definition of $X_{s,v,0}^{1,p'}$, there exists a sequence (φ_n) of $\mathcal{D}(\Omega)$ that converges to f in $W_{s,v}^{1,p'}(\Omega)$. Since $h \in W_{s,v}^{1,p}(\Omega)$, we have

$$\int_{\Omega} h \varphi_n \partial^{\alpha} a \, dx + \int_{\Omega} h a \partial^{\alpha} \varphi_n \, dx = \int_{\Omega} h \partial^{\alpha} (a \varphi_n) \, dx = - \int_{\Omega} a \varphi_n D_v^{\alpha} h \, dx$$

for all n and for all $\alpha \in \pi_1^0$, where we simply used the definition of the weak derivative in the sense of $L_{v,\text{loc}}^1(\Omega)$ with the test function $av^{-2}\varphi_n \in \mathcal{C}_0^1(\Omega)$. By letting n tends to ∞ we get

$$\int_{\Omega} h f \partial^{\alpha} a \, dx + \int_{\Omega} h a D_v^{\alpha} f \, dx = - \int_{\Omega} a f D_v^{\alpha} h \, dx \quad (7.3)$$

since

$$\begin{aligned} \left| \int_{\Omega} h \partial^{\alpha} a (\varphi_n - f) \, dx \right| &\leq C \|\partial^{\alpha} a\|_{L_{v^{-2}}^{\infty}(\Omega)} \|h v\|_{L^p(\Omega)} \|v(\varphi_n - f)\|_{L^{p'}(\Omega)}, \\ \left| \int_{\Omega} h a (\partial^{\alpha} \varphi_n - D_v^{\alpha} f) \, dx \right| &\leq C \|a\|_{L_{v^{-3}}^{\infty}(\Omega)} \|h v\|_{L^p(\Omega)} \|(\partial^{\alpha} \varphi_n - D_v^{\alpha} f) v\|_{L^{p'}(\Omega)}^{s(\alpha)+1}, \\ \text{and} \\ \left| \int_{\Omega} a (\varphi_n - f) D_v^{\alpha} h \, dx \right| &\leq C \|a\|_{L_{v^{-3}}^{\infty}(\Omega)} \|v\|_{L^p(\Omega)}^{s(\alpha)+1} \|D_v^{\alpha} h\|_{L^p(\Omega)} \|v(\varphi_n - f)\|_{L^{p'}(\Omega)}. \end{aligned}$$

Using now (7.3) and the first part of the lemma with $g = f$ we obtain (7.2). \square

Remark 28.

- (i) When $v \in \mathcal{C}_b^1(\Omega)$, the function a in the previous lemma can be any function of the form $a = \tilde{a}v^n$ where $\tilde{a} \in \mathcal{C}_b^1(\Omega)$ and $n \in \mathbb{N}$ with $n \geq 3$.
- (ii) Clearly, the first part of the lemma can be proved with less assumptions on Ω , v and a .

8. POINCARÉ INEQUALITY

In this section we are interesting in proving a Poincaré inequality for the weighted Sobolev space $X_{s,v,0}^{1,p}(\Omega)$.

We first derive an interesting inequality where we prove that the term $\|f \nabla(v^2)\|_{L^p}$ is bounded by $\|v^2 D_v f\|_{L^p}$. We use then this inequality to prove an interesting Poincaré inequality under the assumption $v \in \mathcal{C}_b^{1,*}(\Omega) := \mathcal{C}_b^1(\Omega) \cap \mathcal{C}^{1,*}(\Omega)$.

Theorem 29. *Let $\Omega \subseteq \mathbb{R}^d$ ($d \geq 2$) be a bounded open set, $p \in [1, d)$, $v \in \mathcal{C}_b^{1,*}(\Omega)$, $s : \pi_1 \rightarrow \mathbb{R}_{\geq 0}$ be a map satisfying (4.5). Assume that*

$$\exists \sigma > 0 \forall x \neq 0 : |\nabla v(x)|_p := \left(\sum_{i=1}^d |\partial_i v(x)|^p \right)^{1/p} \leq \sigma \frac{|v(x)|}{|x|} \quad (8.1)$$

where $|x|$ is the euclidean norm of \mathbb{R}^d , and that

$$0 < \frac{2\sigma p}{d-p} < 1. \quad (8.2)$$

Then,

$$\forall f \in X_{s,v,0}^{1,p}(\Omega) : \|f \nabla(v^2)\|_{L^p} \leq \frac{2\sigma p}{d-p-2\sigma p} \|v^2 D_v f\|_{L^p}. \quad (8.3)$$

Proof. Let $f \in X_{s,v,0}^{1,p}(\Omega)$. By Thm. 26 we have

$$fv^2 \in W_0^{1,p}(\Omega) \text{ with } D^\alpha(fv^2) = v^2 D_v^\alpha f + f \partial^\alpha v^2, \quad \forall \alpha \in \pi_1^0.$$

(See also Prop. 12.) Using (8.1) and Hardy inequality we get

$$\begin{aligned} \|f \nabla(v^2)\|_{L^p(\Omega)} &:= 2 \left(\sum_{i=1}^d \|fv \partial_i v\|_{L^p(\Omega)}^p \right)^{\frac{1}{p}} \\ &\leq 2\sigma \left\| \frac{fv^2}{x} \right\|_{L^p(\Omega)} \\ &\leq 2\sigma \frac{p}{d-p} \|D(fv^2)\|_{L^p(\Omega)} \\ &\leq 2\sigma \frac{p}{d-p} \|v^2 D_v f\|_{L^p(\Omega)} + 2\sigma \frac{p}{d-p} \|f \nabla(v^2)\|_{L^p(\Omega)} \end{aligned}$$

This inequality together with (8.2) gives (8.3). \square

Example 30. The typical example of a function satisfying (8.1) when $p \leq 2$ is the function $v(x) = |x|^\beta$ with $\beta \geq 2$, where $\sigma = \beta d^{\frac{2-p}{2}}$.

Similarly, we have

Theorem 31. Let $\Omega \subseteq \mathbb{R}$ be an open set, $p \in (1, \infty)$, $v \in \mathcal{C}^{1,*}(\Omega)$ with $v' \in L^\infty(\Omega)$, and let $s : \pi_1 \rightarrow \mathbb{R}_{\geq 0}$ be a map satisfying (4.5). Assume that

$$\exists \sigma > 0 \forall x \neq 0 : |v'(x)| \leq \sigma \frac{|v(x)|}{|x|}, \quad 0 < \frac{2\sigma p}{p-1} < 1.$$

Then

$$\forall f \in X_{s,v,0}^{1,p}(\Omega) : \|f(v^2)'\|_{L^p(\Omega)} \leq \frac{2\sigma p}{p-1-2\sigma p} \|v^2 D_v f\|_{L^p(\Omega)}.$$

The proof is identical to the proof of Thm. 29.

Using Poincaré inequality and Thm. 29 we get

Corollary 32. Let $\Omega \subseteq \mathbb{R}^d$ ($d \geq 2$) be a bounded open set, $p \in [1, d)$, $v \in \mathcal{C}_b^{1,*}(\Omega)$, $s : \pi_1 \rightarrow \mathbb{R}_{\geq 0}$ be a map satisfying (4.5). Assume that (8.1) and (8.2) hold. Then,

$$\exists C_\Omega > 0 \forall f \in X_{s,v,0}^{1,p} : \|v^2 f\|_{L^p} \leq C_\Omega \|v^2 D_v f\|_{L^p}$$

Proof. This is a direct consequence of Poincaré inequality in $W_0^{1,p}(\Omega)$ and Thm. 29. \square

9. DEGENERATE ELLIPTIC PDE

In the sequel, if $\alpha \in \pi_1^0$, we write D_v^i instead of D_v^α where i is the index of the non-null component of α . We also write D_v for the operator (D_v^i) .

We consider in this section, a boundary-value problem of the form

$$\begin{cases} -\sum_{i,j=1}^d D_j(a_{ij} D_i f) + (b \cdot D)f + cf = h & \text{in } \Omega \\ f = 0 & \text{on } \partial\Omega \end{cases} \quad (9.1)$$

where the coefficients (a_{ij}) , $b = (b_i)$, c are given functions, and h is some given functional. In particular, we consider that the bilinear form $\sum_{i,j=1}^d a_{ij}(x) \xi_i \xi_j$ degenerates i.e. is not uniformly elliptic. Let $V := \{w_\alpha\}_{\alpha \in \pi_1}$ be a collection of elements of $E_p(\Omega)$, and $v \in \mathcal{C}^{1,*}(\Omega)$ that satisfy (4.1), and h be a continuous functional

on $X_{V,v,0}^1(\Omega) (:= X_{V,v,0}^{1,2}(\Omega))$. We say that $f \in X_{V,v,0}^1(\Omega)$ is a weak solution to the boundary-value problem (9.1) if

$$\forall g \in X_{V,v,0}^1(\Omega) : B_v(f, g) = h(g) \quad (9.2)$$

where $B_v(\cdot, \cdot) : X_{V,v,0}^1(\Omega) \times X_{V,v,0}^1(\Omega) \rightarrow \mathbb{R}$ is the bilinear form given by

$$B_v(f, g) := \sum_{i,j=1}^d \int_{\Omega} a_{ij} D_v^i f D_v^j g \, dx + \int_{\Omega} g(b \cdot D_v) f \, dx + \int_{\Omega} c f g \, dx. \quad (9.3)$$

Assume that $f \in X_{V,v,0}^1(\Omega)$ is a weak solution to (9.1) and assume that

$$\forall i, j = 1, \dots, d : a_{ij}, b_i, c \in L_{\text{loc}}^{\infty}(\Omega_{|v|^{2p}}^*) \text{ and } h \in L_{\text{loc}}^2(\Omega_{|v|^{2p}}^*). \quad (9.4)$$

Then the differential equation of (9.1) holds in the sense of $\mathcal{D}'(\Omega_{|v|^{2p}}^*)$. Indeed, we know that $X_{V,v,0}^1(\Omega) \subseteq W^{1,2}(\Omega_{|v|^{2p}}^*, V)$. Hence,

$$\forall i, j = 1, \dots, d : a_{ij} D_i f, b_i D_i f, c f, h \in L_{\text{loc}}^1(\Omega_{|v|^{2p}}^*) \subseteq \mathcal{D}'(\Omega_{|v|^{2p}}^*).$$

The remaining part of the proof is trivial.

For an in-depth study of weak solution existence to boundary value problems of the form (9.1), we refer to [6] and to reference therein. In the next theorem, we give sufficient assumptions that guarantee the existence of a unique weak solution in $X_{s,v,0}^1(\Omega)$ to the boundary-value problem (9.1). We use in particular the Poincaré inequality proved in Cor. 32. Moreover, it is possible to take $h(g) = \int_{\Omega} k \nabla(v^3 g) \, dx$ for all $g \in X_{s,v,0}^1(\Omega)$ where $k \in W_{s,v}^{1,2}(\Omega)$ (see Lem. 27).

For simplicity, we will limit ourselves to the case where $s(\cdot) = |\cdot|$ (in particular $w_{\alpha} = v^{|\alpha|+1} \in \mathcal{C}^1(\Omega)$ for all α). We will then write $X_{|\cdot|,v,0}^1(\Omega)$ instead of $X_{s,v,0}^1(\Omega)$. A generalization to an arbitrary map s satisfying (4.5), is possible with some small modification of the assumptions.

Theorem 33. *Let $\Omega \subseteq \mathbb{R}^d$ be a bounded open set, $v \in \mathcal{C}^{1,*}(\Omega)$, (a_{ij}) be such that $a_{ij} \in L_{v^{-4}}^{\infty}(\Omega)$ and satisfies*

$$\exists \mu > 0 \forall \xi \in \mathbb{R}^d : \mu v^4(x) |\xi|^2 \leq \sum_{i,j=1}^d a_{ij}(x) \xi_i \xi_j \text{ a.e. in } \Omega. \quad (9.5)$$

Let c be such that $v^{-2}c \in L^{\infty}(\Omega)$, and h a continuous functional on $X_{|\cdot|,v,0}^1(\Omega)$. Suppose furthermore that one of the following assumptions holds

- (i) $v^{-3}b_i \in L^{\infty}(\Omega)$ for all $i = 1, \dots, d$, $cv^{-2} \geq \sigma > 0$ a.e. in Ω , and $d^{\frac{1}{2}} \|v^{-3}b\|_{L^{\infty}} < 2\sqrt{\mu\sigma}$, where $\|v^{-3}b\|_{L^{\infty}} := \max_i \|v^{-3}b_i\|_{L^{\infty}}$.
- (ii) $v \in \mathcal{C}_b^1(\Omega)$ satisfies (8.1), (8.2) with $p = 2$ (and hence we should have $d \geq 3$), $v^{-4}b_i \in L^{\infty}(\Omega)$ for all $i = 1, \dots, d$, $cv^{-2} \geq \sigma > 0$ a.e. in Ω , and

$$0 < \mu - C_{\Omega} d^{\frac{1}{2}} \|v^{-4}b\|_{L^{\infty}} \quad (9.6)$$

- (iii) $v^{-3}b_i \in L^{\infty}(\Omega) \cap \mathcal{C}^1(\Omega)$ for all $i = 1, \dots, d$, and one of the following assertions holds

- (a) $\text{div}(b) \leq 0$ a.e. in Ω and $cv^{-2} \geq \sigma > 0$ a.e. in Ω
- (b) $v^{-2} \text{div}(b) \in L^{\infty}(\Omega)$ and $(c - \frac{1}{2} \text{div}(b))v^{-2} \geq \sigma > 0$ a.e. in Ω .

Then, the boundary-value problem (9.1) has a unique weak solution $f \in X_{|\cdot|,v,0}^1(\Omega)$. In addition, we have the estimate

$$\exists \gamma > 0 : \|f\|_{X_{|\cdot|,v}^1} \leq \frac{1}{\gamma} |h| \quad (9.7)$$

where $|h|$ is the norm operator of h .

Proof. We check that the bilinear form B_v defined in (9.3) satisfies the assumptions of the Lax-Milgram theorem in the Hilbert space $X_{|\cdot|,v,0}^1(\Omega)$. Note that in any cases, we have $b_i v^{-3} \in L^\infty(\Omega)$. Hence, for any $f, g \in X_{|\cdot|,v,0}^1(\Omega)$ we have

$$\begin{aligned} |B_v(f, g)| &\leq C \sum_{i,j=1}^d \|v^2 D_v^i f\|_{L^2} \|v^2 D_v^j g\|_{L^2} + C \sum_{i=1}^d \|v^2 D_v^i f\|_{L^2} \|v g\|_{L^2} \\ &\quad + C \|v f\|_{L^2} \|v g\|_{L^2} \leq C \|f\|_{X_{|\cdot|,v}^1} \|g\|_{X_{|\cdot|,v}^1} \end{aligned}$$

for some non-negative constant C , which proves the continuity of the bilinear form B_v . We prove now that the bilinear form B_v is coercive, that is,

$$\exists \gamma > 0 \forall f \in X_{|\cdot|,v,0}^1(\Omega) : \gamma \|f\|_{X_{|\cdot|,v}^1}^2 \leq B_v(f, f). \quad (9.8)$$

Fix $f \in X_{|\cdot|,v,0}^1$. Under the assumptions on (a_{ij}) we have

$$\mu \|v^2 D_v f\|_{L^2}^2 + \int_{\Omega} c f^2 dx + \int_{\Omega} f(b \cdot D_v) f dx \leq B_v(f, f).$$

Suppose that assertion (i) holds. Then by Hölder's inequality, we have

$$\mu \|v^2 D_v f\|_{L^2}^2 + \sigma \|v f\|_{L^2}^2 - d^{\frac{1}{2}} \|v^{-3} b\|_{L^\infty} \|v^2 D_v f\|_{L^2} \|v f\|_{L^2} \leq B_v(f, f) \quad (9.9)$$

Thanks to the assumptions of the assertion (i), it is possible to choose γ sufficiently small so that $0 < \gamma \leq \min(\mu, \sigma)$ and $d^{\frac{1}{2}} \|b\|_{L_{v^{-3}}^\infty} < 2\sqrt{(\mu - \gamma)(\sigma - \gamma)}$. Thus, it follows that left hand side of (9.9) is bounded from below by $\gamma(\|v^2 D_v f\|_{L^2}^2 + \|v f\|_{L^2}^2)$ which gives (9.8).

In case assertion (ii) holds, we use Hölder's inequality to obtain

$$\mu \|v^2 D_v f\|_{L^2}^2 + \sigma \|v f\|_{L^2}^2 - d^{\frac{1}{2}} \|v^{-4} b\|_{L^\infty} \|v^2 D_v f\|_{L^2} \|v^2 f\|_{L^2} \leq B_v(f, f)$$

By Cor. 32, we have

$$(\mu - C_\Omega d^{\frac{1}{2}} \|v^{-4} b\|_{L^\infty}) \|v^2 D_v f\|_{L^2}^2 + \sigma \|v f\|_{L^2}^2 \leq B_v(f, f).$$

Therefore, (9.8) holds with the constant $\gamma := \min((\mu - C_\Omega d^{\frac{1}{2}} \|v^{-4} b\|_{L^\infty}), \sigma)$ which is non-negative thanks to (9.6).

Assume now that (iii) holds. By (9.5) we have

$$\mu \|v^2 D_v f\|_{L^2}^2 + \int_{\Omega} f(b \cdot D_v) f dx + \int_{\Omega} c f^2 dx \leq B_v(f, f). \quad (9.10)$$

Let (φ_n) be a sequence of $\mathcal{D}(\Omega)$ that converges to f in $X_{|\cdot|,v}^1(\Omega)$. Since $v^{-3} b_i \in L^\infty(\Omega)$ we have

$$\begin{aligned} \left| \int_{\Omega} b_i (f D_v^i f - \varphi_n \partial_i \varphi_n) dx \right| &\leq \\ \|b_i v^{-3}\|_{L^\infty} \|f v\|_{L^2} \|v^2 (D_v^i f - \partial_i \varphi_n)\|_{L^2} &+ \|b_i v^{-3}\|_{L^\infty} \|v(f - \varphi_n)\|_{L^2} \|v^2 \partial_i \varphi_n\|_{L^2}. \end{aligned}$$

Thus,

$$\int_{\Omega} f(b \cdot D_v) f \, dx = \lim_{n \rightarrow \infty} \int_{\Omega} \varphi_n(b \cdot \nabla) \varphi_n \, dx = -\frac{1}{2} \lim_{n \rightarrow \infty} \int_{\Omega} \operatorname{div}(b) \varphi_n^2 \, dx. \quad (9.11)$$

Hence, in case (a) of (iii) holds, the right hand side of the latter equality is positive. Thus, the left hand side of (9.10) is bounded from below by $\gamma \|f\|_{X_{|\cdot|,v}^1}^2$ where $\gamma = \min(\mu, \sigma) > 0$, and hence (9.8) is proved, and in case (b) of (iii) holds, the right hand side of (9.11) is equal to $-\frac{1}{2} \int_{\Omega} \operatorname{div}(b) f^2 \, dx$. Replacing this in (9.10) we get

$$\gamma \|f\|_{X_{|\cdot|,v}^1}^2 \leq \mu \|v^2 D_v f\|_{L^2}^2 + \sigma \|vf\|_{L^2}^2 \leq B_v(f, f)$$

where $\gamma := \min(\mu, \sigma) > 0$. Thus, (9.8) holds.

Therefore, if one of the assertions (i)-(iii) holds, the bilinear form B_v is coercive. On the other hand, h is a continuous functional on $X_{|\cdot|,v,0}^1$. We conclude then by the Lax-Milgram theorem.

For the estimate (9.7), it follows easily from (9.2), (9.8), and the continuity of h . Indeed, if we denote by f the unique weak solution of (9.1), then

$$\gamma \|f\|_{X_{|\cdot|,v}^1}^2 \leq B_v(f, f) = h(f) \leq |h| \|f\|_{X_{|\cdot|,v}^1}.$$

□

Example 34. Let $v \in \mathcal{C}^{1,*}(\Omega)$, and let (\tilde{a}_{ij}) , $(\tilde{b}) = (\tilde{b}_i)$, \tilde{c} be such that

- (i) $\forall i, j = 1, \dots, d : \tilde{a}_{ij}, \tilde{b}_i, \tilde{c} \in L^\infty(\Omega)$
- (ii) $\exists \mu > 0$ such that $\sum_{i,j} \tilde{a}_{ij} \xi_i \xi_j \geq \mu |\xi|^2$ a.e. in Ω and for all $\xi \in \mathbb{R}^d$
- (iii) $\exists \sigma > 0$ such that $\tilde{c} \geq \sigma$ a.e. in Ω , and $d^{\frac{1}{2}} \|\tilde{b}\|_{L^\infty(\Omega)} < 2\sqrt{\mu\sigma}$.

Set

$$\forall i, j = 1, \dots, d : a_{ij} = v^4 \tilde{a}_{ij}, b_i = v^3 \tilde{b}_i, c = v^2 \tilde{c}.$$

Then, the functions (a_{ij}) satisfy (9.5), and the functions (b_i) , c satisfy (i) of Thm. 33.

In the next proposition we give sufficient conditions under which the solution is not locally integrable.

Proposition 35. Let $\Omega \subseteq \mathbb{R}^d$ be a bounded open set, $v \in \mathcal{C}^{2,*}(\Omega)$ be a map satisfying (6.1). Let $f \in X_{|\cdot|,v,0}^1(\Omega)$ be such that (9.2) holds where

- (i) $\forall i, j = 1, \dots, d : a_{ij} \in \mathcal{C}^1(\Omega) \cap L_{v^{-4}}^\infty(\Omega)$ and $\forall \alpha \in \pi_1^0 : \partial^\alpha a_{ij} \in L_{v^{-3},\text{loc}}^\infty(\Omega)$
- (ii) $\forall i = 1, \dots, d : b_i \in \mathcal{C}^1(\Omega) \cap L_{v^{-3}}^\infty(\Omega)$ and $\forall \alpha \in \pi_1^0 : \partial^\alpha b_i \in L_{v^{-2},\text{loc}}^\infty(\Omega)$
- (iii) $c \in L_{v^{-2}}^\infty(\Omega)$
- (iv) the map h is defined by $h(g) := \int_{\Omega} kg \, dx$, where $k \in L_{v^{-1}}^2(\Omega)$, $k \geq 0$ a.e. in Ω and $kv^{-2} \notin L_{\text{loc}}^1(\Omega)$

Then, $f \notin L_{\text{loc}}^1(\Omega)$.

Proof. Use (9.2) with $g = \varphi \chi_n v^{-2} \in \mathcal{C}_c^2(\Omega)$ (because $\operatorname{supp}(\chi_n) \subseteq M_{2n}^c$) where $\varphi \in \mathcal{D}(\Omega)$ and (χ_n) is the sequence of $\mathcal{C}^2(\Omega)$ functions constructed in the proof of

Prop. 7 we get

$$\begin{aligned} \sum_{i,j=1}^d \int_{\Omega} a_{ij} D_v^i f \partial_j (\varphi \chi_n v^{-2}) dx + \sum_{i=1}^d \int_{\Omega} b_i \varphi \chi_n v^{-2} D_v^i f dx \\ + \int_{\Omega} c f \varphi \chi_n v^{-2} dx = \int_{\Omega} k \varphi \chi_n v^{-2} dx \end{aligned} \quad (9.12)$$

Using the weak derivative in the sense of $L_{v,\text{loc}}^1(\Omega)$ with $a_{ij} v^{-2} \partial_j (\varphi \chi_n v^{-2}) \in \mathcal{C}_c^1(\Omega)$ we get

$$\int_{\Omega} a_{ij} D_v^i f \partial_j (\varphi \chi_n v^{-2}) dx = - \int_{\Omega} f \partial_i (a_{ij} \partial_j (\varphi \chi_n v^{-2})) dx \quad (9.13)$$

Similarly, using the weak derivative in the sense of $L_{v,\text{loc}}^1(\Omega)$ with the test function $b_i \varphi \chi_n v^{-4} \in \mathcal{C}_c^1(\Omega)$ we get

$$\int_{\Omega} b_i \varphi \chi_n v^{-2} D_v^i f dx = - \int_{\Omega} f \partial_i (b_i \varphi \chi_n v^{-2}) dx \quad (9.14)$$

Using (9.13) and (9.14) in (9.12) we get

$$\begin{aligned} - \sum_{i,j=1}^d \int_{\Omega} f \partial_i (a_{ij} \partial_j (\varphi \chi_n v^{-2})) dx - \sum_{i=1}^d \int_{\Omega} f \partial_i (b_i \varphi \chi_n v^{-2}) dx \\ + \int_{\Omega} c f \varphi \chi_n v^{-2} dx = \int_{\Omega} k \varphi \chi_n v^{-2} dx. \end{aligned} \quad (9.15)$$

A simple calculation gives

$$\begin{aligned} \partial_i (a_{ij} \partial_j (\varphi \chi_n v^{-2})) &= \chi_n (\partial_i a_{ij} \partial_j (v^{-2} \varphi) + a_{ij} \partial_{ij} (v^{-2} \varphi)) \\ &+ \partial_j \chi_n (v^{-2} \varphi \partial_i a_{ij} + a_{ij} \partial_i (v^{-2} \varphi)) + \partial_i \chi_n (a_{ij} \partial_j (v^{-2} \varphi)) + a_{ij} v^{-2} \varphi \partial_{ij} \chi_n \end{aligned} \quad (9.16)$$

Developing the expression $\partial_i a_{ij} \partial_j (v^{-2} \varphi) + a_{ij} \partial_{ij} (v^{-2} \varphi)$ we get

$$\begin{aligned} \partial_i a_{ij} (-2(\partial_j v) v^{-3} \varphi + v^{-2} \partial_j \varphi) + \\ a_{ij} (v^{-2} \partial_{ij} \varphi - 2(\partial_j v) v^{-3} \partial_i \varphi - 2(\partial_i v) v^{-3} \partial_j \varphi - 2(\partial_{ij} v) v^{-3} \varphi + 6(\partial_i v)(\partial_j v) v^{-4} \varphi) \end{aligned}$$

By assumption (i)

$$\exists C > 0 \forall n \in \mathbb{N} : |\chi_n (\partial_i a_{ij} \partial_j (v^{-2} \varphi) + a_{ij} \partial_{ij} (v^{-2} \varphi))| \leq C 1_V \quad (9.17)$$

where 1_V is the characteristic function of the set $V := \text{supp}(\varphi)$. Note now that, for all $\alpha \in \pi_2^0$ we have $\partial^\alpha \chi_n \equiv 0$ on M_n^c (because by $\chi_n \equiv 1$ on M_n^c), and that $|\partial^\alpha \chi_n| \leq C n^{|\alpha|}$ on M_n for all n sufficiently large, where $C > 0$ is independent of n . This is because $M_n \Subset \Omega$ (by assumption (6.1)) (and hence $\partial^\alpha v$ is bounded on M_n), and because $|\eta_n^{(k)}| \leq C n^{|k|}$ where the constant $C > 0$ depends on k but not on n . It follows that

$$\exists C > 0 \exists n_0 \in \mathbb{N} \forall n \geq n_0 \forall \alpha \in \pi_2^0 : |v^{|\alpha|} \partial^\alpha \chi_n| \leq C \quad (9.18)$$

since $|v| \leq 1/n$ on M_n . Thus, one can easily prove (as above), using assumption (i) that

$$|\partial_j \chi_n (v^{-2} \varphi \partial_i a_{ij} + a_{ij} \partial_i (v^{-2} \varphi)) + \partial_i \chi_n (a_{ij} \partial_j (v^{-2} \varphi)) + a_{ij} v^{-2} \varphi \partial_{ij} \chi_n| \leq C 1_V. \quad (9.19)$$

for all $n \geq n_0$. Assume by contradiction that $f \in L^1_{\text{loc}}(\Omega)$. Then, by (9.16), (9.17) and (9.19) we have

$$\forall n \geq n_0 : \left| \sum_{i,j=1}^d \int_{\Omega} f(\partial_i(a_{ij}\partial_j(\varphi\chi_n v^{-2}))) dx \right| \leq C\|f\|_{L^1(V)}. \quad (9.20)$$

By (ii) and (iii), and using similar arguments as above one can prove that

$$\forall n \geq n_0 : \left| \sum_{i=1}^d \int_{\Omega} f\partial_i(b_i\varphi\chi_n v^{-2}) + fc\varphi\chi_n v^{-2} dx \right| \leq C\|f\|_{L^1(V)} \quad (9.21)$$

We deduce from (9.15), (9.20), (9.21) that

$$\exists C > 0 \forall n \geq n_0 : \int_{\Omega} k\varphi\chi_n v^{-2} dx \leq C\|f\|_{L^1(V)}$$

Let now $K \Subset \Omega$ be such that $kv^{-2} \notin L^1(K)$ and choose φ so that $\varphi \geq 0$ on Ω and $\varphi \equiv 1$ on K . It follows that $\int_K k\chi_n v^{-2} dx \leq C$ for all $n \geq n_0$, which leads to a contradiction when choosing n sufficiently large since

$$\lim_{n \rightarrow \infty} \int_K k\chi_n v^{-2} dx = \int_K kv^{-2} dx = \infty$$

by the monotone convergence theorem. \square

Remark 36. If

$$\forall i, j = 1, \dots, d : a_{ij} = \tilde{a}_{ij}v^4, b_i = \tilde{b}_i v^3$$

where

$$\forall i, j = 1, \dots, d : \tilde{a}_{ij} \in \mathcal{C}^1(\Omega), \tilde{b}_i \in \mathcal{C}^1(\Omega),$$

then assumptions (i), (ii) of the latter proposition hold.

Example 37. For any integer $m \geq 1$, and for any $\beta \in (d/2 - 2m, d/2)$, the boundary-value problem

$$\begin{cases} -\sum_{i=1}^d D_i(|x|^{8m} D_i f) + |x|^{4m} f = |x|^{-\beta+2m} & \text{in } \{x \in \mathbb{R}^d \mid |x| < 1\} \\ f = 0 & \text{on } \{x \in \mathbb{R}^d \mid |x| = 1\} \end{cases}$$

has a unique non-locally integrable solution $f \in X^1_{|\cdot|, |x|^{2m}, 0}(\Omega)$ thanks to Thm. 33 and to Prop. 35.

REFERENCES

- [1] Cavalheiro, A., Weighted Sobolev Spaces and Degenerate Elliptic Equations. Boletim da Sociedade Paranaense de Matemática. 26.- 10.5269/bspm.v26i1-2.7415. (2008).
- [2] Dunford, N., Schwartz, J. T., Linear Operators. Part I : General Theory. Interscience Publishers, New York, London 1958.
- [3] Evans, L., Partial differential equations. Providence, R.I.: American Mathematical Society, 2010.
- [4] Hörmander, L., On the division of distributions by polynomials, Arkiv för Mat., t. 3, 1958, p. 555-568.
- [5] Kufner, A., Opic, B., How to define reasonably weighted Sobolev spaces. (English). Commentationes Mathematicae Universitatis Carolinae, vol. 25 (1984), issue 3, pp. 537-554
- [6] Kufner, A., Sändig, A., Some applications of weighted Sobolev spaces." (1987).
- [7] Wikipedia contributors. Faà di Bruno's formula. In Wikipedia, The Free Encyclopedia. en.wikipedia.org/w/index.php?title=Fa%C3%A0_di_Bruno%27s_formula&oldid=1145953219

FACULTY OF MATHEMATICS, UNIVERSITY OF VIENNA, AUSTRIA, OSKAR-MORGENSTERN-PLATZ
1, 1090 WIEN, AUSTRIA
Email address: `djameleddine.kebiche@univie.ac.at`