A NONLOCAL EQUATION DESCRIBING TUMOR GROWTH

RAFAEL GRANERO-BELINCHÓN AND MARTINA MAGLIOCCA

Abstract. Cancer is a very complex phenomenon that involves many different scales and situations. In this paper we consider a free boundary problem describing the evolution of a tumor colony and we derive a new asymptotic model for tumor growth. We focus on the case of a single phase tumor colony taking into account chemotactic effects in an early stage where there is no necrotic inner region. Thus, our model is valid for the case of multilayer avascular tumors with very little access to both nutrients and inhibitors or the case where the amount of nutrients and inhibitors is very similar to the amount consumed by the multilayer tumor cells. Our model takes the form of a single nonlocal and nonlinear partial differential equation. In its simplest dimensionless form, our model reads

$$
\partial_t g(x_1, t)
$$

= $-\eta \Lambda^3 g(x_1, t) + \varepsilon \eta ([H, g(x_1, t)] H g_{1,11}(x_1, t)),$

$$
-\varepsilon \left(2\theta e^{-Nt} \alpha(t) (g(x_1, t) - g(x_1, 0)) + e^{-Nt} \alpha(t) (2\theta - \rho)t + \rho e^{-Nt} t c_{\mathcal{B}}\right)
$$

$$
+ \theta \varepsilon e^{-Nt} 2\alpha(t) ((g(x_1, t) - g(x_1, 0)) + t) - \varepsilon \frac{\rho}{2} \alpha(t) e^{-Nt} g_{111}(x_1, t) - \frac{e^{-Nt}}{2} \rho (\alpha(t) - c_{\mathcal{B}}),
$$

where g denote the interface of the multilayer tumor colony, H denotes the Hilbert transform, $\varepsilon, \theta, \eta$ are dimensionless parameters measuring different ratios related to amplitude of the tumor, stength of chemotactic and cohesion effects, respectively. Similarly, N, ρ , c_B and $\alpha(t)$ capture the effect of nutrients, inhibitors and vasculation of the tumor colony. Besides deriving the model, we also prove a well-posedness result.

CONTENTS

Key words and phrases. Tumor growth, Free boundary problem, Nonlocal PDE, Well-posedness.

Date: December 25, 2023.

1. INTRODUCTION

Free boundary problems for partial differential equations are a very hot research area in Mathematical Analysis nowadays. These problems are mathematically challenging and physically interesting. Moreover, their applications are really spread, from geothermal reservoirs to tumour growth, passing through weather forecasting. Many free boundary problems arise during the interaction between two fluids or between a fluid and an elastic solid, *i.e.* when the dynamics of both fluids or the fluid and the elastic solid are connected. The study of fluid $\&$ fluid and fluid & solid interactions are classical problems in Applied Sciences and Mathematics. In this paper, we will consider tumor growth as a free boundary problem for appropriate partial differential equations.

Cancer is a very complex phenomenon that involves many different scales and situations. Cancer starts at a primary site in the body with the mutation of a small number of cells. Some time after this, proliferating cancer cells start their cell division and the tumor grows. Some of the cells situated in the inner part of the tumor die due to absence of enough nutrients such as oxygen or glucose and a central necrotic core is formed. Usually, the closest cells to this necrotic core are viable non-proliferating cells while the outer part of the carcinoma contains the proliferating cells [\[9,](#page-22-0) [7\]](#page-22-1). Without access to the vasculature, the tumor colony may grow up to the order of 10^6 cells. If these cancer cells manage to grow successfully they may acquire a vascular network. Once they have access to a constant source of nutrients, the tumor can grow beyond that limit and at a faster rate [\[46\]](#page-23-0).

The importance of this illness together with the complexity of the phenomenon lead to a large number of research studies by many different research groups in Biology and Medicine but also in Physics, Mathematics or Engineering. In fact, the first mathematical models related to cancer goes back to the fifties (see [\[3,](#page-21-2) [8\]](#page-22-2) for very complete reviews). However, these first attempts do not provide any mechanical insight in the growth of tumors. Due to the large number of cells involved, a continuum modelling approach seems appropriate [\[8\]](#page-22-2).

Although cancer research has improved the available treatments and our understanding of this disease, there is still a lack of a good mechanical description of certain phenomena. Such mathematical approach to oncology has many benefits as evidenced for instance by the work of Gatenby & Gawlinski [\[25\]](#page-22-3) (see also [\[2,](#page-21-3) [8\]](#page-22-2)). There, the predictions of the mathematical models where verified experimentally later. Furthermore, as pointed out in [\[2\]](#page-21-3), mathematical models could provide unexpected insight into the underlying mechanisms and generate novel hypotheses for experimentation. The purpose of this work is to study a mechanical model of tumor growth and derive a new asymptotic model that accurately describe the evolution of the colony. In particular, we will focus on the case of a single phase tumor colony where only the tumor (without healthy surrounding tissue) is considered (see figure [1\)](#page-1-0). The tumor is assumed to be in an early stage where there is no necrotic inner region but we allow the tumor colony to possibly have access to the vascular network. The problem that we consider is very similar to the growth of a so called multilayer tumor, i.e. a colony of tumor cells cultivated in laboratory [\[34,](#page-22-4) [38\]](#page-23-1).

Figure 1. The one-phase case with only the tumor colony and infinite depth

In that regards, when studying tumor growth there are different approaches present in the literature, each of them focusing on a specific feature of such a complex phenomenon.

First, one has the approach based on systems of nonlinear differential equations posed on a fixed domain. We could possibly call this approach the reaction-diffusion approach. Examples of such an approach are the works $\begin{bmatrix} 4 \\ 12 \\ 26 \\ 27 \\ 33 \\ 47 \\ 45 \\ 48 \\ 51 \end{bmatrix}$. Typically, the system of differential equations takes the form of a coupled parabolic partial differential equation(s) and a ordinary differential equation to model haptotaxis. In general, such systems of differential equations are able to exhibit very complex and even spatio-temporal chaotic dynamics [\[6,](#page-21-5) [42\]](#page-23-6).

Second, one can consider *hybrid multiscale* and also *agent based* models such as [\[39,](#page-23-7) [41,](#page-23-8) [43,](#page-23-9) [44,](#page-23-10) [50,](#page-23-11) [49\]](#page-23-12). In this approach, partial differential equations of reaction-diffusion type are considered at the tissue level. However, at the cell level, agent based models are used to track each cell together with its state individually. In particular, these models allow to differentiate cancer cells into quiescent, proliferative, apoptotic, hypoxic, and even necrotic states. Finally, ordinary differential equations are used at the subcellular scale.

A third approach could possibly be called the phase-field models. In this approach, the tumor microenvironment is considered as a mixture of tumor and healthy tissue. For this twoconstituent medium, a function ϕ describes the tumor and healthy tissue volume fractions, respectively and a diffuse, typically thin, interface is considered. In this approach, the tumor is mathematically described by Cahn-Hilliard-type equations. The Cahn-Hilliar equation is a fourth order parabolic equation that was originally developed as a model phase separation and now is widely applied in tumor growth. Examples of such an approach are [\[13,](#page-22-9) [15,](#page-22-10) [14,](#page-22-11) [24,](#page-22-12) [23,](#page-22-13) [36,](#page-23-13) [40,](#page-23-14) [41\]](#page-23-8).

There is however another approach which is focused on the boundary of the tumor colony. In this approach, the tumor colony is described mathematically as a *free boundary problem*. These approach started with the pioneer works of Greenspan [\[31,](#page-22-14) [32\]](#page-22-15) (see also Byrne & Chaplain [\[10,](#page-22-16) [9,](#page-22-0) [7\]](#page-22-1)) and continues nowadays (see [\[5,](#page-21-6) [16\]](#page-22-17) and the references therein).

In particular, these early models by Greenspan study the growth of *avascular, i.e.* the carcinoma is not linked to the blood vessels, multicellular spheroids. Typically this type of tumor grows in vitro. This situation is then described by several reaction diffusion equations for the different nutrients and inhibitors, and an ordinary differential equation for the radius of the tumor (which is assumed to be radially symmetric). This is the approach in the works of many authors such as those in [\[18,](#page-22-18) [19,](#page-22-19) [22,](#page-22-20) [20,](#page-22-21) [21\]](#page-22-22).

Although this setting was succesful in describing certain phenomena related to cancer growth it is also very limited. In particular, such a setting can describe how the tumor grows or how the colony diminishes, but it cannot describe observed features of certain tumors such as invasive fingers or tumor boundaries with large and variable curvature.

This paper adscribes himself to the so called free boundary approach. In this regards, we consider the problem of tumor growth as the dynamics of a free boundary. The main purpose of this paper is to derive and study a single simplified nonlinear and nonlocal partial differential equation that describe (up to certain degree of accuracy) the full free boundary dynamics in certain biological regime. A similar research program has been widely applied in many different areas in Applied Sciences. That is for instance the case of Coastal Engineering where asymptotic models of water waves are used to accurately describe the motion of the surface of the sea in the appropriate physical regime. While there are many results for free boundary problems arising when studying the motion of incompressible fluids (see for instance [\[28,](#page-22-23) [35,](#page-22-24) [1,](#page-21-7) [11,](#page-22-25) [17\]](#page-22-26) and the references therein), much less is known for free boundary problems modelling the growth of different sorts of tumors.

In this regards, the asymptotic model that we derive and study in this paper reads

$$
\partial_t g(x_1, t) = -\eta \Lambda^3 g(x_1, t) + \varepsilon \eta \left([H, g(x_1, t)] H g_{1111}(x_1, t) \right),_1 - \varepsilon \theta g(x_1, t) e^{-N_2 t} \mathcal{L}_{11}(x_1, 0, t) + \varepsilon \widetilde{K}^{(1)}(x_1, 0, t) + K^{(0)}(x_1, 0, t).
$$

for appropriate, explicit (see below) functions $\widetilde{K}^{(1)}, K^{(0)}$ and **L**. Such a model es valid up to certain explicit error (see below) in the case of an avascular tumor with very little access to both nutrients and inhibitors. Similarly, the same model is also valid in the case of vascular tumors where the amount of nutrients and inhibitors is very similar to the amount consumed by the tumor cells.

To the best of our knowledge, not only the model of tumor growth that we find is new and interesting, but also the derivation technique is different to the more heuristic derivations present in the literature. Furthermore, besides deriving a new asymptotic model for multilayer tumor growth, we also establish a well-posedness result for a specific case.

Notation. We denote the one dimensional torus by $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$. Alternatively, this domain can be thought as the interval $[-\pi, \pi]$ with periodic boundary conditions.

Given a matrix $A \in \mathbb{R}^{n \times m}$ we denote with A_j^i the entry of A at row i and column j, and we adopt Einstein convention for the summation of repeated indexes. We write

$$
f_{,j} = \partial_{x_j} f
$$

for the space derivative in the j−th direction. Similarly, H denotes the Hilbert transform. Let $v(x_1)$ denote a L^2 function on $\mathbb T$. We recall its Fourier series representation:

$$
\hat{v}(n) = \frac{1}{2\pi} \int_{\mathbb{T}} v(x_1) e^{-inx_1} dx_1,
$$

where $n \in \mathbb{Z}$. Then we have that

$$
v(x_1) = \sum_{n \in \mathbb{Z}} \hat{v}(n) e^{inx_1}.
$$

Finally,

$$
[A, B] = AB - BA \tag{1.1}
$$

denotes the commutator.

2. Free boundary problems describing tumor growth

We consider the case in which the growth of cancer regions is affected by the action of certain inhibitors, whose aim is preventing the tumor expansion. These inhibitors can be a blood-borne anticancer drug or any other substance secreted by neighboring cells in response to the anomalous tissue. Then, the growth of the tumor is related to the interactions between the nutrients such as oxygen or glucose and the previously mentioned inhibitors. However, we consider that every nutrient acts via the function σ and every inhibitor products its effect through the function β . Due to these assumptions, the system of PDE will contain two reaction diffusion equations, one for β and one for σ .

As we said, we allow for the tumor colony to have access to the vasculature. The capillary network can supply nutrients to the tumor at a rate

$$
\delta_n(\sigma_B-\sigma),
$$

where σ_B is the constant nutrient concentration in the vasculature, and δ_n is the constant rate of blood-tissue transfer per unit length. We observe that the avascular case reduces then to $\delta_n=0$

The nutrient then diffuses in the tumor tissue with diffusion constant D_n^T and it is consumed at a rate $\lambda_n \sigma$. Furthermore, the inhibitor may interact with the nutrient with a function $G_n(\sigma, \beta)$. Similarly, the inhibitor diffuses with constant D_i^T , it is consumed with constant λ_i and it is supplied from the capillary network with rate δ_i . In the case of blood-borne anticancer drug, a blood-tissue exchange rate of the form

$$
\delta_i(\beta_B-\beta)
$$

has to be taken into account. Here, β_B is the constant inhibitor concentration in the vasculature. In addition to this, the nutrient can potentially have an effect on the inhibitors via the function $G_i(\sigma, \beta)$. We can use the same reasoning with respect to the inhibitor β . Then, writing $\Omega_T(t)$ for the tumor region, these reaction diffusion equations take the generic form (see for instance [\[9,](#page-22-0) [37\]](#page-23-15))

$$
\frac{\partial \sigma}{\partial t} - D_n \Delta \sigma = \delta_n (\sigma_B - \sigma) - \lambda_n \sigma + G_n(\sigma, \beta) \quad \text{in } \Omega_T(t) \times (0, T),
$$

$$
\frac{\partial \beta}{\partial t} - D_i \Delta \beta = \delta_i (\beta_B - \beta) - \lambda_i \beta + G_i(\sigma, \beta) \quad \text{in } \Omega_T(t) \times (0, T).
$$

The sources G_i depend on the hypotheses of the model [\[9\]](#page-22-0). In general, it is assumed that the inhibitor affects both cell proliferation and the nutrient concetration.

Following [\[9\]](#page-22-0), we are going to choose

$$
G_n(\sigma,\beta)=-\gamma_n\beta,
$$

and

$$
G_i(\sigma, b) = 0,
$$

i.e. the inhibitor affects the nutrient concentration with some constant rate γ_n , but not the cell proliferation rate.

For more examples of choices of G_n and G_i , we refer to [\[9\]](#page-22-0).

The cells are assumed to move according to *pressure* gradients created by the birth and death of cells $[32]$. The introduction of this oncotic pressure function, p, make a conceptual change from a mechanical viewpoint. As a consequence of this hypothesis, the velocity vector of the tumor, u , is assumed to follow Darcy's law

$$
u = -\nabla p + \chi \nabla \sigma \qquad \text{in } \Omega_T(t) \times (0, T),
$$

where

$$
\chi \nabla \sigma \quad \text{with} \quad \chi \ge 0
$$

is a chemotaxis term modelling the motion towards high concentration of nutrients. Being μ a constant threshold reflecting the expansion by mitosis and $\tilde{\sigma}$ being a threshold concentration, the local rate of volume change is given by

$$
\nabla \cdot u = \mu(\sigma - \tilde{\sigma} - \tau \beta) \qquad \text{in } \Omega_T(t) \times (0, T),
$$

where the term $\tau\beta$ is the death rate of tumor cells due to the effect of the inhibitor. The role played by this threshold concentration $\tilde{\sigma}$ can be explained as follows. On the one hand, if the amount of nutrients is below this value in a certain region, the cells die and the tumor colony diminishes locally. On the other hand, if the amount of nutrients is above this value in certain area of the tumor tissue, the cells replicate themselves and the tumor colony grows locally.

There are a number of boundary conditions that need to be imposed for this system. First, let us denote the tumor surface by $\Gamma(t)$. We assume that the tumor colony has boundary given by

$$
\Gamma(t) = (x_1, h(x_1, t)),
$$

for certain function h. We write K for the curvature of $\Gamma(t)$, *i.e.*

$$
\mathcal{K} = \frac{\partial_{x_1}^2 h}{(1 + (\partial_{x_1} h)^2)^{3/2}}.
$$

Then, although there are other available models [\[10\]](#page-22-16), in this work we consider that the compactness of the tumor is mantained by surface tension via a Young-Laplace equation [\[32\]](#page-22-15)

$$
p = -\nu \mathcal{R} \qquad \text{on } \Gamma(t) \times (0, T).
$$

Then, as the surface of the tumor colony is transported by the cell velocity, we have that

$$
\frac{\partial h}{\partial t} = u \cdot (-\partial_{x_1} h(x_1, t), 1) \qquad \text{on } \Gamma(t) \times (0, T).
$$

The appropriate boundary conditions are

$$
\sigma = \sigma_D, \quad \beta = \beta_D \qquad \text{on } \Gamma(t) \times (0, T).
$$

As a consequence, under the assumption that the inhibitor acts on the nutrient concentration and the cell proliferation, the free boundary problem can be written as follows

$$
\partial_t \sigma - D_n \Delta \sigma = \delta_n (\sigma_B - \sigma) - \lambda_n \sigma - \gamma_n \beta \quad \text{in } \Omega_T(t) \times (0, T),
$$

\n
$$
\partial_t \beta - D_i \Delta \beta = \delta_i (\beta_B - \beta) - \lambda_i \beta \quad \text{in } \Omega_T(t) \times (0, T),
$$

\n
$$
\nabla \cdot u = \mu(\sigma - \tilde{\sigma} - \tau \beta) \quad \text{in } \Omega_T(t) \times (0, T),
$$

\n
$$
u = -\nabla p + \chi \nabla \sigma \quad \text{in } \Omega_T(t) \times (0, T),
$$

\n
$$
p = -\nu \frac{h_{11}}{(1 + h_{11})^{3/2}} \quad \text{on } \Gamma(t) \times (0, T),
$$

\n
$$
\partial_t h = u \cdot (h_{11}, 1) \quad \text{on } \Gamma(t) \times (0, T),
$$

\n
$$
\beta = \beta_D \quad \text{on } \Gamma(t) \times (0, T),
$$

\n
$$
\beta = \beta_D \quad \text{on } \Gamma(t) \times (0, T),
$$

with the initial data

$$
\sigma(x,0) = \sigma_0(x) \quad \text{in } \Omega(0),
$$

\n
$$
\beta(x,0) = \sigma_0(x) \quad \text{in } \Omega(0),
$$

\n
$$
h(x_1,0) = g(x_1) \quad \text{on } \Gamma(0),
$$

with

$$
\Omega_T(t) = \left\{ x \in \mathbb{R}^2, x_1 \in L\mathbb{S}^1, -\infty < x_2 < h(x_1, t) \right\},\
$$

where L is the typical length of the tumor colony and $L\mathbb{S}^1$ denotes the circle with length 2L (or equivalently the interval $[-L\pi, L\pi]$ with periodic boundary conditions). Similarly, the boundary Γ(*t*) of this domain is

$$
\Gamma(t) = \{ x \in \mathbb{R}^2, x_1 \in L\mathbb{S}^1, x_2 = h(x_1, t) \}.
$$

In the previous free boundary problem, the units of the quantities and parameters are as follows.

Concentrations σ , β , σ_B , β_B , σ_D , β_D , $\tilde{\sigma}$, : g · cm⁻² Diffusion coefficients D_i , D_n and pressure $p:$ cm² \cdot s⁻¹ Rates δ_i , δ_n , λ_i , λ_n , γ_n : s⁻¹ Velocities $u, \frac{\partial h}{\partial t} : \text{ cm} \cdot \text{s}^{-1}$ Curvature $\mathcal{K}: \quad \mathrm{cm}^{-1}$ Parameter $\nu: \text{ cm}^3 \cdot \text{s}^{-1}$ Parameter $\chi: \quad g^{-1} \cdot \text{cm}^4 \cdot \text{s}^{-1}$ Parameter $\mu: \text{ g}^{-1} \cdot \text{cm}^2 \cdot \text{s}^{-1}$

With these units in mind, we consider the dimensionless variables

$$
\overline{x} = \left(\frac{x_1}{L}, \frac{x_2}{L}\right), \qquad \overline{t} = \frac{D_n}{LH}t,
$$

and functions

$$
\overline{\sigma}(\overline{x},\overline{t}) = \frac{\sigma(x,t) - \sigma_D}{\tilde{\sigma}}, \qquad \overline{u}(\overline{x},\overline{t}) = \frac{L}{D_n}u(x,t), \qquad \overline{p}(\overline{x},\overline{t}) = \frac{p(x,t)}{D_n},
$$

A NONLOCAL EQUATION DESCRIBING TUMOR GROWTH $7\,$

$$
\overline{h}(\overline{x},\overline{t}) = \frac{h(x,t)}{H}, \qquad \overline{\beta}(\overline{x},\overline{t}) = \frac{\beta(x,t) - \beta_D}{\tilde{\sigma}},
$$

being L and H typical lengths associated to the tumor colony. Furthermore, we define the dimensionless parameters \overline{H} \overline{D}

$$
\varepsilon = \frac{H}{L}, \quad \alpha = \frac{D_i}{D_n},\tag{2.1}
$$

$$
N_1 = LH \frac{\delta_i + \lambda_i}{D_n}, \quad N_2 = LH \frac{\delta_n + \lambda_n}{D_n}, \quad N_3 = \frac{\gamma_n LH}{D_n}, \tag{2.2}
$$

$$
M_1 = \frac{LH}{D_n\tilde{\sigma}} \left[-(\delta_i + \lambda_i)\beta_D + \delta_i\beta_B \right], \quad M_2 = \frac{LH}{D_n\tilde{\sigma}} \left[-\sigma_D(\delta_n + \lambda_n) + \sigma_B\delta_n - \gamma_n\beta_D \right], \tag{2.3}
$$

$$
\theta = \frac{\chi \tilde{\sigma}}{D_n}, \quad \rho = \frac{\mu \tilde{\sigma} L^2}{D_n}, \quad \omega = \frac{\mu L^2}{D_n} (\sigma_D - \tau \beta_D - \tilde{\sigma}), \quad \eta = \frac{\nu H}{L^2 D_n}.
$$
 (2.4)

Then, dropping the $\overline{\cdot}$ from the notation and substituting u by its value, the problem in dimensionless form reads as follows

$$
\partial_t \beta - \varepsilon \alpha \Delta \beta = -N_1 \beta + M_1 \qquad \text{in } \Omega_T(t) \times (0, T), \qquad (2.5a)
$$

$$
\partial_t \sigma - \varepsilon \Delta \sigma = -N_2 \sigma - N_3 \beta + M_2 \qquad \text{in } \Omega_T(t) \times (0, T), \tag{2.5b}
$$

$$
-\Delta p + \theta \Delta \sigma = \rho (\sigma - \tau \beta) + \omega \qquad \text{in } \Omega_T(t) \times (0, T), \qquad (2.5c)
$$

$$
p = -\eta \frac{\partial_{x_1}^2 h}{(1 + (\varepsilon \partial_{x_1} h)^2)^{3/2}} \qquad \text{on } \Gamma(t) \times (0, T), \tag{2.5d}
$$

$$
\partial_t h = (-\nabla p + \theta \nabla \sigma) \cdot (-\partial_{x_1} h, 1) \qquad \text{on } \Gamma(t) \times (0, T), \tag{2.5e}
$$

$$
\sigma = 0 \qquad \text{on } \Gamma(t) \times (0, T), \tag{2.5f}
$$

$$
\beta = 0 \qquad \text{on } \Gamma(t) \times (0, T), \tag{2.5g}
$$

where the dimensionless domain and boundary are

$$
\Omega_T(t) = \{ x \in \mathbb{R}^2, x_1 \in \mathbb{S}^1, -\infty < x_2 < \varepsilon h(x_1, t) \},
$$
\n
$$
\Gamma(t) = \{ x \in \mathbb{R}^2, x_1 \in \mathbb{S}^1, x_2 = \varepsilon h(x_1, t) \}.
$$

3. The free boundary problem in Arbitrary Lagrangian-Eulerian coordinates

In order to fix the domain, we now pass to the ALE formulation. The reference domain (see figure [2\)](#page-6-1) and boundary are

and

$$
\Omega_T = \left\{ x \in \mathbb{R}^2, x_1 \in \mathbb{S}^1, -\infty < x_2 < 0 \right\},\
$$

$$
\Gamma = \{ x \in \mathbb{R}^2, x_1 \in \mathbb{S}^1, x_2 = 0 \}.
$$

Vacuum

FIGURE 2. The reference domain.

As our current goal is transforming the free boundary problem [\(2.5\)](#page-6-2) into a fixed boundary one, we introduce the time dependent diffeomorphism

$$
\psi(\cdot,t):\Omega_T\to\Omega_T(t)
$$

defined as

$$
\psi(x_1, x_2, t) = (x_1, x_2 + \varepsilon h(x_1, t)).
$$

Then, we are going to compose the equations in problem (2.5) with ψ to find its equivalent formulation over Ω_T . Let us define the concentrations of nutrient, inhibitor and pressure in the fixed boundary frame as

$$
S = \sigma \circ \psi, \qquad B = \beta \circ \psi, \qquad P = p \circ \psi.
$$

In order to write (2.5) in terms of S, B , and P , we will need

$$
\nabla \psi = \begin{pmatrix} 1 & 0 \\ \varepsilon h, 1 & 1 \end{pmatrix}, \qquad \psi_t = \varepsilon \begin{pmatrix} 0 \\ \partial_t h \end{pmatrix} \qquad \text{and} \qquad A = (\nabla \psi)^{-1} = \begin{pmatrix} 1 & 0 \\ -\varepsilon h, 1 & 1 \end{pmatrix}.
$$

Since we are interested in the asymptotic behavior in ε , we split these equivalent equations in S, B, and P as follows: we collect all the terms which depend on ε in the r.h.s., leaving the rest to the left.

In the following computations, we will need to compute

$$
\partial_t f \circ \psi
$$
 and $\Delta f \circ \psi$

in terms of $F = f \circ \psi$. We use the chain rule obtaining that

$$
\partial_t f \circ \psi = \partial_t F - A_i^k F_{,k} \, \partial_t \psi^i = \partial_t F - \varepsilon F_{,2} \, \partial_t h,
$$

$$
(\Delta f) \circ \psi = A_j^i \left(A_j^k F_{,k}^T \right)_{,i} = \Delta F + \varepsilon \mathbf{R}_1[h] \left(F \right) + \varepsilon^2 \mathbf{R}_2[h] \left(F \right),
$$

where we have used Einstein convention for summation over repeated indexes, and

$$
\mathbf{R}_{1}[\ell(x_{1},t)]\left(F(x,t)\right) = -(\ell_{11}(x_{1},t)F(x,t) + 2\ell_{11}(x_{1},t)F_{11}(x,t)),_{2},
$$
\n
$$
\mathbf{R}_{2}[\ell(x_{1},t)]\left(F(x,t)\right) = \ell_{11}^{2}(x_{1},t)F_{22}(x,t).
$$
\n(3.1)

Reasoning in this way, we obtain the system

$$
\partial_t B + N_1 B - M_1 = \varepsilon (B, 2 \partial_t h + \alpha \Delta B) + \varepsilon^2 \alpha \mathbf{R}_1[h](B) \quad \text{in } \Omega_T \times (0, T), \tag{3.2a}
$$

$$
\partial_t S + N_2 S + N_3 B - M_2 = \varepsilon \left(S_{2} \partial_t h + \Delta S \right) + \varepsilon^2 \mathbf{R}_1[h](S) \qquad \text{in } \Omega_T \times (0, T), \tag{3.2b}
$$

$$
-\Delta P + \theta \Delta S - \rho (S - \tau B) - \omega = \varepsilon (\mathbf{R}_1[h](P) - \theta \mathbf{R}_1[h](S))
$$

+
$$
\varepsilon^2
$$
 ($\mathbf{R}_2[h](P) - \theta \mathbf{R}_2[h](P)$) in $\Omega_T \times (0,T)$, (3.2c)

$$
P = -\eta \frac{h_{,11}}{\left(1 + \varepsilon^2 h_{,1}^2\right)^{3/2}} \quad \text{on } \Gamma \times (0, T), \quad (3.2d)
$$

$$
\partial_t h = -P_{,2} + \theta S_{,2} + \varepsilon h_{,1} (P_{,1} - \theta S_{,1})
$$

$$
-\varepsilon^2 h_{11}^2 (P_{2}-\theta S_{2}) \qquad \text{on } \Gamma \times (0,T), \quad (3.2e)
$$

$$
S = 0 \qquad \text{on } \Gamma \times (0, T), \quad (3.2f)
$$

$$
B = 0 \qquad \text{on } \Gamma \times (0, T). \quad (3.2g)
$$

4. DERIVATION OF THE ASYMPTOTIC MODEL

We introduce the ansatz

$$
h(x_1, t) = \sum_{n=0}^{\infty} \varepsilon^n h^{(n)}(x_1, t), \qquad B(x, t) = \sum_{n=0}^{\infty} \varepsilon^n B^{(n)}(x, t),
$$

$$
S(x, t) = \sum_{n=0}^{\infty} \varepsilon^n S^{(n)}(x, t), \qquad P(x, t) = \sum_{n=0}^{\infty} \varepsilon^n P^{(n)}(x, t),
$$

where

$$
B(x,0) = B^{(0)}(x,0), \qquad S(x,0) = S^{(0)}(x,0), \qquad h(x_1,0) = g(x_1,0),
$$

and

$$
S^{(n)}(x,0) = B^{(n)}(x,0) = 0, \qquad h^{(n)}(x_1,0) = 0 \qquad \text{for } n \ge 1.
$$

We consider now the case where

$$
M_1 = M_2 = \omega = O(\varepsilon^2).
$$

This is, for instance, the case of an avascular tumor with very little access to both nutrients and inhibitors. Similarly, it is the case of vascular tumors where the amount of nutrients and inhibitors is very similar to the amount consumed by the tumor cells.

4.1. The case $n = 0$. The first term in the asymptotic series is then

$$
\partial_t B^{(0)} + N_1 B^{(0)} = 0 \qquad \text{in } \Omega_T \times (0, T), \qquad (4.1a)
$$

$$
\partial_t S^{(0)} + N_2 S^{(0)} + N_3 B^{(0)} = 0 \qquad \text{in } \Omega_T \times (0, T), \qquad (4.1b)
$$

$$
-\Delta P^{(0)} + \theta \Delta S^{(0)} - \rho \left(S^{(0)} - \tau B^{(0)} \right) = 0 \qquad \text{in } \Omega_T \times (0, T), \qquad (4.1c)
$$

$$
P^{(0)} = -\eta h^{(0)},11 \qquad \text{on } \Gamma \times (0, T), \qquad (4.1d)
$$

$$
\partial_t h^{(0)} = -P^{(0)},_2 + \theta S^{(0)},_2 \qquad \text{on } \Gamma \times (0, T), \tag{4.1e}
$$

$$
S^{(0)} = 0 \qquad \text{on } \Gamma \times (0, T), \tag{4.1f}
$$

$$
B^{(0)} = 0 \qquad \text{on } \Gamma \times (0, T). \tag{4.1g}
$$

In order to simplify the notation, we set

$$
\mathcal{B}(x) = B^{(0)}(x,0)
$$
 and $\mathcal{S}(x) = S^{(0)}(x,0)$.

We can solve for B and S using, respectively, $(4.1a) - (4.1g)$ $(4.1a) - (4.1g)$ and $(4.1b) - (4.1f)$ $(4.1b) - (4.1f)$, finding that

$$
B^{(0)}(x,t) = e^{-N_1 t} \mathcal{B}(x),\tag{4.2}
$$

$$
S^{(0)}(x,t) = e^{-N_2 t} \mathcal{L}(x,t),
$$
\n(4.3)

where

$$
\mathcal{L}(x,t) = \mathcal{S}(x) - \frac{N_3}{N_2 - N_1} \left(e^{(N_2 - N_1)t} - 1 \right) \mathcal{B}(x).
$$
 (4.4)

The expression of the pressure $P^{(0)}$ can be found applying [\[29,](#page-22-27) Lemma A.1] to [\(4.1c\)](#page-8-5)-[\(4.1d\)](#page-8-6):

$$
P^{(0)}(x,t) = -\frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} \left[\frac{1}{2|k|} \left(\int_{-\infty}^{0} \hat{w}^{(0)}(k, y_2, t) e^{|k|y_2} dy_2 \right) \left(e^{|k|x_2} - e^{-|k|x_2} \right) \right.
$$

$$
+ \frac{1}{2|k|} \left(\int_{0}^{x_2} \hat{w}^{(0)}(k, y_2, t) \left(e^{-|k|(x_2 - y_2)} - e^{|k|(x_2 - y_2)} \right) dy_2 \right)
$$

$$
+ -\widehat{g^{(0)}}(k, t) e^{|k|x_2|} e^{ikx_1}, \tag{4.5}
$$

with $\hat{b} = \hat{w}^{(0)}$, $\hat{g} = g^{(0)}$, and

$$
\widehat{w}^{(0)} = \widehat{\theta \Delta S^{(0)}} - \rho \left(\widehat{S^{(0)}} - \tau \widehat{B^{(0)}} \right),
$$

$$
\widehat{g^{(0)}} = \eta |k|^2 \widehat{h^{(0)}}.
$$

We gather the above computations, together with [\(4.1e\)](#page-8-7), to say that

$$
\partial_t h^{(0)}(x_1, t) = -P^{(0)}, \quad \quad \ (x_1, 0, t) + \theta e^{-N_2 t} \mathcal{L}, \quad \quad \ (x_1, 0, t)
$$
\n
$$
= -\eta \Lambda^3 h^{(0)}(x_1, t) + K^{(0)}(x_1, 0, t), \tag{4.6}
$$

where

$$
K^{(0)}(x_1, 0, t) = -\theta \int_{-\infty}^{0} e^{y_2 \Lambda} e^{-N_2 t} \Delta \mathcal{L}(x_1, y_2, t) dy_2
$$

+ $\rho \int_{-\infty}^{0} e^{y_2 \Lambda} e^{-N_2 t} \left(\mathcal{L}(x_1, y_2, t) - \tau e^{(N_2 - N_1)t} \mathcal{B}(x_1, y_2) \right) dy_2$
+ $\theta e^{-N_2 t} \mathcal{L}_{,2}(x_1, 0, t).$ (4.7)

4.2. The case $n = 1$. We now consider the case $n = 1$.

The expressions of $B^{(1)}$ and $S^{(1)}$. The equations $(3.2a)$ - $(3.2g)$ for the second term of the asymptotic expansion reads

$$
\partial_t B^{(1)} + N_1 B^{(1)} = B^{(0)}, \quad \partial_t h^{(0)} + \alpha \Delta B^{(0)} = e^{-N_1 t} \left(\mathcal{B}, \quad \partial_t h^{(0)} + \alpha \Delta \mathcal{B} \right) \quad \text{in } \Omega_T \times (0, T),
$$

$$
B^{(1)} = 0 \qquad \text{on } \Gamma \times (0, T),
$$

thanks also to the expression of $B^{(0)}(x,t)$ in [\(4.2\)](#page-8-8). This means that

$$
B^{(1)}(x,t) = e^{-N_1t} \left[\mathcal{B}_{2}(x) \left(h^{(0)}(x_1,t) - \mathcal{g}(x_1) \right) + \alpha t \Delta \mathcal{B}(x) \right].
$$

We now set

$$
\mathbf{Q}_{\alpha}[\ell(x_1, t)](f(x, t)) = (\ell(x_1, t) - \mathbf{g}(x_1)) f_{1,2}(x, t) + \alpha t \Delta f(x, t), \qquad (4.8)
$$

and rewrite $B^{(1)}$ as

$$
B^{(1)}(x,t) = e^{-N_1 t} Q_{\alpha}[h^{(0)}(x_1,t)](\mathcal{B}(x)).
$$
\n(4.9)

Note that, even if **B** does not depend on t, $Q_{\alpha}[h^{(0)}(x_1, t)](\cdot)$ always depends on time.

As far as (3.2b) - (3.2f) is concerned, the computations above and (4.3) lead to
\n
$$
\partial_t S^{(1)}(x,t) + N_2 S^{(1)}(x,t)
$$
\n
$$
= -N_3 B^{(1)}(x,t) + S^{(0)},_2 (x,t) \partial_t h^{(0)}(x_1,t) + \Delta S^{(0)}(x,t)
$$
\n
$$
= -N_3 e^{-N_1 t} \left[\mathcal{B},_2 (x) \left(h^{(0)}(x_1,t) - \mathcal{g}(x_1) \right) + \alpha t \Delta \mathcal{B}(x) \right]
$$
\n
$$
+ e^{-N_2 t} \left(\mathcal{S},_2 (x) - \frac{N_3}{N_2 - N_1} \left(e^{(N_2 - N_1)t} - 1 \right) \mathcal{B},_2 (x) \right) \partial_t h^{(0)}(x_1,t)
$$
\n
$$
+ e^{-N_2 t} \left(\Delta \mathcal{S}(x) - \frac{N_3}{N_2 - N_1} \left(e^{(N_2 - N_1)t} - 1 \right) \Delta \mathcal{B}(x) \right) \quad \text{in } \Omega_T \times (0, T),
$$
\n
$$
S^{(1)} = 0 \quad \text{on } \Gamma \times (0, T),
$$

so that $S^{(1)}$ has the form

$$
S^{(1)}(x,t) = -N_3 e^{-N_2 t} \int_0^t e^{(N_2 - N_1)s} \left[\mathcal{B}_{2}(x) \left(h^{(0)}(x_1, s) - \mathcal{g}(x_1) \right) + \alpha s \Delta \mathcal{B}(x) \right] ds
$$

+ $e^{-N_2 t} \mathcal{S}_{2}(x) \left(h^{(0)}(x_1, t) - \mathcal{g}(x_1) \right)$
- $\frac{N_3}{N_2 - N_1} e^{-N_2 t} \mathcal{B}_{2}(x) \int_0^t \left(e^{(N_2 - N_1)s} - 1 \right) \partial_s h^{(0)}(x_1, s) ds$

$$
+ e^{-N_2 t} \left[t \Delta \mathcal{S}(x) - \frac{N_3}{N_2 - N_1} \Delta \mathcal{B}(x) \int_0^t e^{(N_2 - N_1)s} - 1 ds \right].
$$

We explicitly compute the integrals appearing in the above formulation:

$$
-N_{3}e^{-N_{2}t}\mathcal{B}_{2}(x)\int_{0}^{t}e^{(N_{2}-N_{1})s}\left(h^{(0)}(x_{1},s)-g(x_{1})\right)ds
$$

\n
$$
= -N_{3}e^{-N_{2}t}\mathcal{B}_{2}(x)\int_{0}^{t}e^{(N_{2}-N_{1})s}h^{(0)}(x_{1},s)ds
$$

\n
$$
+\frac{N_{3}}{N_{2}-N_{1}}e^{-N_{2}t}g(x_{1})\mathcal{B}_{2}(x)\left(e^{(N_{2}-N_{1})t}-1\right),
$$

\n
$$
-\alpha N_{3}e^{-N_{2}t}\Delta\mathcal{B}(x)\int_{0}^{t}e^{(N_{2}-N_{1})s}sds
$$

\n
$$
=\alpha \frac{N_{3}}{N_{2}-N_{1}}\Delta\mathcal{B}(x)e^{-N_{2}t}\left(e^{(N_{2}-N_{1})t}t-\frac{1}{N_{2}-N_{1}}\left(e^{(N_{2}-N_{1})t}-1\right)\right),
$$

\n
$$
-\frac{N_{3}}{N_{2}-N_{1}}e^{-N_{2}t}\mathcal{B}_{2}(x)\int_{0}^{t}\left(e^{(N_{2}-N_{1})s}-1\right)\partial_{s}h^{(0)}ds
$$

\n
$$
=\frac{N_{3}}{N_{2}-N_{1}}e^{-N_{2}t}\mathcal{B}_{2}(x)\left(e^{(N_{2}-N_{1})t}-1\right)h^{(0)}(x_{1},t)
$$

\n
$$
+N_{3}e^{-N_{2}t}\mathcal{B}_{3}(x)\int_{0}^{t}e^{(N_{2}-N_{1})s}h^{(0)}(x_{1},s)ds,
$$

\n
$$
-\frac{N_{3}}{N_{2}-N_{1}}e^{-N_{2}t}\Delta\mathcal{B}(x)\int_{0}^{t}e^{(N_{2}-N_{1})s}-1ds
$$

$$
= e^{-N_2 t} \frac{N_3}{N_2 - N_1} \Delta \mathcal{B}(x) \left(t - \frac{1}{N_2 - N_1} \left(e^{(N_2 - N_1)t} - 1 \right) \right).
$$

Then, $S^{(1)}$ becomes

−

$$
S^{(1)}(x,t) = e^{-N_2 t} \left(h^{(0)}(x_1,t) - g(x_1) \right) \left(\mathcal{S}_{2}(x) - \frac{N_3}{N_2 - N_1} \left(e^{(N_2 - N_1)t} - 1 \right) \mathcal{B}_{2}(x) \right)
$$

+ $e^{-N_2 t} t \Delta \left[\mathcal{S}(x) + \frac{N_3}{N_2 - N_1} \left(1 - \alpha e^{(N_2 - N_1)t} \right) \mathcal{B}(x) \right]$
- $\frac{N_3 (1 - \alpha)}{(N_2 - N_1)^2} e^{-N_2 t} \left(e^{(N_2 - N_1)t} - 1 \right) \Delta \mathcal{B}(x)$
= $e^{-N_2 t} \left(h^{(0)}(x_1,t) - g(x_1) \right) \mathcal{L}_{2}(x,t) + e^{-N_2 t} t \Delta \mathcal{L}(x,t)$
+ $\frac{N_3 (1 - \alpha)}{N_2 - N_1} e^{-N_2 t} \Delta \mathcal{B}(x) \left[t e^{(N_2 - N_1)t} - \frac{1}{N_2 - N_1} \left(e^{(N_2 - N_1)t} - 1 \right) \right].$

We set

$$
M(t) = \frac{N_3(1-\alpha)}{N_2 - N_1} e^{-N_2 t} \left[t e^{(N_2 - N_1)t} - \frac{1}{N_2 - N_1} \left(e^{(N_2 - N_1)t} - 1 \right) \right],
$$
 (4.10)

and recall the expression of $S^{(1)}$ in (4.8) so that we can rewrite $S^{(1)}$ in compact form as

$$
S^{(1)}(x,t) = e^{-N_2 t} Q_1[h^{(0)}(x_1,t)](\mathcal{L}(x,t)) + M(t) \Delta \mathcal{B}(x).
$$
\n(4.11)

The expression of $P^{(1)}$. We now focus on $P^{(1)}$. Equations $(3.2c) - (3.2d)$ $(3.2c) - (3.2d)$ for the second term of the asymptotic expansion read

$$
\Delta P^{(1)} = \theta \Delta S^{(1)} - \rho \left(S^{(1)} - \tau B^{(1)} \right) - \mathbf{R}_1[h^{(0)}](P^{(0)}) + \theta \mathbf{R}_1[h^{(0)}](S^{(0)}) \quad \text{in } \Omega_T \times (0,T),
$$

$$
P^{(1)} = \eta h^{(1)},
$$
 on $\Gamma \times (0, T)$,

where $\boldsymbol{R}_1[h^{(0)}]$ is given by (3.1) .

To find $P^{(1)}$, we apply again [\[29,](#page-22-27) Lemma A.1] with $\hat{b} = \widehat{w_{(1)}}, \hat{g} = \widehat{g_{(1)}},$ for $\widehat{w_{(1)}} = \widehat{\theta \Delta S^{(1)}} - \widehat{\rho \left(\widehat{S^{(1)}} - \tau \widehat{B^{(1)}} \right)} - \widehat{\mathbf{R_1}[h^{(0)}]}(P^{(0)}) + \widehat{\mathbf{R_1}[h^{(0)}]}(S^{(0)}),$

$$
w_{(1)} = \theta \Delta S^{(1)} - \rho \left(S^{(1)} - \tau B^{(1)} \right) - \mathbf{R}_1 [h^{(0)}](P^{(0)}) + \mathbf{R}_1 [h^{(0)}](S^{(0)}),
$$

$$
\widehat{g_{(1)}} = \eta |k|^2 \widehat{h^{(1)}},
$$

obtaining that

$$
P^{(1)}(x,t) = -\frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} \left[\frac{1}{2|k|} \left(\int_{-\infty}^{0} \widehat{w_{(1)}}(k, y_2, t) e^{|k|y_2} dy_2 \right) \left(e^{|k|x_2} - e^{-|k|x_2} \right) \right.
$$

$$
+ \frac{1}{2|k|} \left(\int_{0}^{x_2} \widehat{w_{(1)}}(k, y_2, t) \left(e^{-|k|(x_2 - y_2)} - e^{|k|(x_2 - y_2)} \right) dy_2 \right)
$$

$$
- \widehat{g_{(1)}}(k, t) e^{|k|x_2|} e^{ikx_1}.
$$

The evolution equation of $\partial_t h^{(1)}$. Finally, the second term of the asymptotic expansion of [\(3.2e\)](#page-7-8) is

$$
\partial_t h^{(1)} = -P^{(1)},_2 + \theta S^{(1)},_2 + h^{(0)},_1 \left(P^{(0)},_1 - \theta S^{(0)},_1 \right) \qquad \text{on } \Gamma \times (0, T).
$$

We want to compute the explicit expression of $\partial_t h^{(1)}$ in terms of **B** and **S**. To this aim, we have to develop each term in the above equation.

We begin focusing on $P^{(1)}$, 2. [\[29,](#page-22-27) Lemma A.1] gives us

$$
P^{(1)},_2 = \eta \Lambda^3 h^{(1)} + F_1 + F_2 + F_3 + F_4 + F_5 \qquad \text{on } \Gamma \times (0, T),
$$

where

$$
F_1(x_1, 0, t) = \theta \int_{-\infty}^0 e^{y_2 \Lambda} \Delta S^{(1)}(x_1, y_2, t) dy_2,
$$

\n
$$
F_2(x_1, 0, t) = -\rho \int_{-\infty}^0 e^{y_2 \Lambda} S^{(1)}(x_1, y_2, t) dy_2,
$$

\n
$$
F_3(x_1, 0, t) = \rho \tau \int_{-\infty}^0 e^{y_2 \Lambda} B^{(1)}(x_1, y_2, t) dy_2,
$$

\n
$$
F_4(x_1, 0, t) = -\int_{-\infty}^0 e^{y_2 \Lambda} \mathbf{R}_1[h^{(0)}](P^{(0)}(x_1, y_2, t)) dy_2,
$$

\n
$$
F_5(x_1, 0, t) = \int_{-\infty}^0 e^{y_2 \Lambda} \mathbf{R}_1[h^{(0)}](S^{(0)}(x_1, y_2, t)) dy_2.
$$

We are going to write the F_i terms putting in evidence their dependence on the data. We recall the expressions of $S^{(1)}$ and $B^{(1)}$ in (4.11) and (4.9) , and we write F_2 and F_3 as

$$
F_2(x_1, 0, t) = -\rho e^{-N_2 t} \int_{-\infty}^0 e^{y_2 \Lambda} \mathbf{Q}_1[h^{(0)}(x_1, t)](\mathcal{L}(x_1, y_2, t)) dy_2
$$

$$
-\rho M(t) \int_{-\infty}^0 e^{y_2 \Lambda} \Delta \mathbf{B}(x_1, y_2), \qquad (4.12)
$$

$$
F_3(x_1, 0, t) = \rho \tau e^{-N_1 t} \int_{-\infty}^0 e^{y_2 \Lambda} \mathbf{Q}_{\alpha}[h^{(0)}(x_1, t)] (\mathbf{B}(x_1, y_2)) dy_2.
$$
 (4.13)

Since

$$
\Delta(fg) = g\Delta f + f\Delta g + 2(f, g, g_1 + f, g, g_2),
$$

we develop F_1 as

$$
F_1(x_1, 0, t) = \theta e^{-N_2 t} \int_{-\infty}^0 e^{y_2 \Lambda} \left(\ell(x_1, t) - g(x_1) \right)_{1,11} \mathcal{L}_{1,2}(x_1, y_2, t) dy_2
$$

+ $\theta e^{-N_2 t} \int_{-\infty}^0 e^{y_2 \Lambda} \left(\ell(x_1, t) - g(x_1) \right) \Delta \mathcal{L}_{1,2}(x_1, y_2, t) dy_2$
+ $2\theta e^{-N_2 t} \int_{-\infty}^0 e^{y_2 \Lambda} \left(\ell(x_1, t) - g(x_1) \right)_{11} \mathcal{L}_{1,2}(x_1, y_2, t) dy_2$
+ $\theta M(t) \int_{-\infty}^0 e^{y_2 \Lambda} \Delta^2 \mathcal{B}(x_1, y_2) dy_2.$ (4.14)

Furthermore, by (3.1) , F_5 becomes

$$
F_5(x_1, 0, t) = -\int_{-\infty}^0 e^{y_2 \Lambda} \left(h^{(0)},_{11} (x_1, t) S^{(0)},_2 (x_1, y_2, t) + 2h^{(0)},_1 (x_1, t) S^{(0)},_{12} (x_1, y_2, t) \right) dy_2
$$

$$
= -e^{-N_2 t} \int_{-\infty}^0 e^{y_2 \Lambda} \left(h^{(0)},_{11} (x_1, t) \mathcal{L}(x_1, y_2, t) + 2h^{(0)},_1 (x_1, t) \mathcal{L},_1 (x_1, y_2, t) \right),_2 dy_2
$$

$$
= -e^{-N_2 t} \int_{-\infty}^0 e^{y_2 \Lambda} \mathbf{R}_1[h^{(0)}(x_1, t)] (\mathcal{L}(x_1, y_2, t)) dy_2.
$$
 (4.15)

We define the sum of F_1 , F_2 , F_3 , F_5 as

$$
K_1^{(1)} = F_1 + F_2 + F_3 + F_5, \tag{4.16}
$$

with F_i in (4.14) , (4.12) , (4.13) , (4.15) .

We are left with \mathbb{F}_4 which can be written as

$$
F_4(x_1, 0, t) = \int_{-\infty}^0 e^{y_2 \Lambda} \left(h^{(0)},_{11} (x_1, t) P^{(0)}(x_1, y_2, t) + 2h^{(0)},_{1} (x_1, t) P^{(0)},_{1} (x_1, y_2, t) \right),_2 dy_2
$$

\n
$$
= \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} e^{ikx_1} \int_{-\infty}^0 e^{y_2 |k|} \overline{h^{(0)},_{11} (k, t) P^{(0)},_2 (k, y_2, t)} dy_2
$$

\n
$$
+ \frac{2}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} e^{ikx_1} \int_{-\infty}^0 e^{y_2 |k|} \overline{h^{(0)},_{1} (k, t) P^{(0)},_{12} (k, y_2, t)} dy_2.
$$

\norder to develop the Fourier terms, we recall (4.5) and we compute
\n
$$
\overline{0, 0},_2 (k, x_2, t) = -\frac{1}{2} \left(\int_{-\infty}^0 \widehat{w}^{(0)}(k, y_2, t) e^{|k|y_2} dy_2 \right) \left(e^{|k|x_2} + e^{-|k|x_2} \right)
$$

In order to develop the Fourier terms, we recall (4.5) and we compute

$$
= \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} e^{ikx_1} \int_{-\infty} e^{iy_2|k|} h^{(0)}{}_{,11}(k, t) P^{(0)}{}_{,2}(k, y_2, t) dy_2
$$

+
$$
\frac{2}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} e^{ikx_1} \int_{-\infty}^0 e^{y_2|k|} \overline{h^{(0)}{}_{,1}(k, t) P^{(0)}{}_{,12}(k, y_2, t) dy_2}.
$$

In order to develop the Fourier terms, we recall (4.5) and we compute

$$
\widehat{P^{(0)}}{}_{,2}(k, x_2, t) = -\frac{1}{2} \left(\int_{-\infty}^0 \widehat{w}^{(0)}(k, y_2, t) e^{|k|y_2} dy_2 \right) \left(e^{|k|x_2} + e^{-|k|x_2} \right)
$$

$$
+ \frac{1}{2} \left(\int_0^{x_2} \widehat{w}^{(0)}(k, y_2, t) \left(e^{-|k|(x_2 - y_2)} + e^{|k|(x_2 - y_2)} \right) dy_2 \right) + |k| \widehat{g^{(0)}}(k, t) e^{|k|x_2},
$$

$$
\widehat{P^{(0)}}{}_{,12}(k, x_2, t) = -\frac{ik}{2} \left(\int_{-\infty}^0 \widehat{w}^{(0)}(k, y_2, t) e^{|k|y_2} dy_2 \right) \left(e^{|k|x_2} + e^{-|k|x_2} \right)
$$

$$
+ \frac{ik}{2} \left(\int_0^{x_2} \widehat{w}^{(0)}(k, y_2, t) \left(e^{-|k|(x_2 - y_2)} + e^{|k|(x_2 - y_2)} \right) dy_2 \right) + ik|k| \widehat{g^{(0)}}(k, t) e^{|k|x_2}.
$$

Then, F_4 becomes

$$
F_4(x_1, 0, t) = \eta \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} e^{ikx_1} \sum_{m \in \mathbb{Z}} \int_{-\infty}^0 e^{y_2(|k| + |m|)} |m|^3 \widehat{h^{(0)}}(m, t)(k - m)^2 \widehat{h^{(0)}}(k - m, t) dy_2
$$

$$
+ 2\eta \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} e^{ikx_1} \sum_{m \in \mathbb{Z}} \int_{-\infty}^0 e^{y_2(|k|+|m|)} (k-m) \widehat{h^{(0)}}(k-m,t)m|m|^3 \widehat{h^{(0)}}(m,t) dy_2
$$

+
$$
\frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} e^{ikx_1} \left(\widehat{I}_1(k,0,t) + \widehat{I}_2(k,0,t) \right),
$$

where

$$
\widehat{I}_{1}(k,0,t) = -\frac{1}{2} \sum_{m \in \mathbb{Z}} \left(\int_{-\infty}^{0} \widehat{w}^{(0)}(m, y_{2}, t) e^{|m|y_{2}} dy_{2} \right) \times
$$
\n
$$
\times \int_{-\infty}^{0} e^{y_{2}|k|}(k-m)^{2} \widehat{h^{(0)}}(k-m) \left(e^{|m|y_{2}} + e^{-|m|y_{2}} \right) dy_{2}
$$
\n
$$
+ \frac{1}{2} \sum_{m \in \mathbb{Z}} \int_{-\infty}^{0} e^{y_{2}|k|}(k-m)^{2} \widehat{h^{(0)}}(k-m) \times
$$
\n
$$
\times \left(\int_{0}^{y_{2}} \widehat{w}^{(0)}(m, y_{2}', t) \left(e^{-|m|(y_{2} - y_{2}') } + e^{|m|(y_{2} - y_{2}')} \right) dy_{2}' \right) dy_{2}, \qquad (4.17)
$$
\n
$$
\widehat{I}_{2}(k,0,t) = \sum_{m \in \mathbb{Z}} \left(\int_{-\infty}^{0} \widehat{w}^{(0)}(m, y_{2}, t) e^{|m|y_{2}} dy_{2} \right) \times
$$
\n
$$
\times \int_{-\infty}^{0} e^{y_{2}|k|}(k-m) \widehat{h^{(0)}}(k-m) \left(e^{|m|y_{2}} + e^{-|m|y_{2}} \right) dy_{2}
$$
\n
$$
- \sum_{m \in \mathbb{Z}} \int_{-\infty}^{0} e^{y_{2}|k|}(k-m) \widehat{h^{(0)}}(k-m) \times
$$
\n
$$
\times \left(\int_{0}^{y_{2}} \widehat{w}^{(0)}(m, y_{2}', t) \left(e^{-|m|(y_{2} - y_{2}') } + e^{|m|(y_{2} - y_{2}')} \right) dy_{2}' \right) dy_{2}, \qquad (4.18)
$$

and with

$$
\widehat{w}^{(0)} = \widehat{\theta \Delta S^{(0)}} - \rho \left(\widehat{S^{(0)}} - \tau \widehat{B^{(0)}} \right).
$$

We can simplify the expression of F_4 through the following steps:

$$
F_4(x_1, 0, t) = \eta \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} e^{ikx_1} \sum_{m \in \mathbb{Z}} \int_{-\infty}^0 e^{y_2(|k| + |m|)} |m|^3 (k - m)(k + m) \widehat{h^{(0)}}(m, t) \widehat{h^{(0)}}(k - m, t) dy_2
$$

+
$$
\frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} e^{ikx_1} \left(\widehat{I}_1(k, 0, t) + \widehat{I}_2(k, 0, t) \right)
$$

=
$$
\eta \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} e^{ikx_1} \sum_{m \in \mathbb{Z}} |m|^3 (|k| - |m|) \widehat{h^{(0)}}(m, t) \widehat{h^{(0)}}(k - m, t)
$$

+
$$
\frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} e^{ikx_1} \left(\widehat{I}_1(k, 0, t) + \widehat{I}_2(k, 0, t) \right)
$$

=
$$
\eta \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} e^{ikx_1} \sum_{m \in \mathbb{Z}} |m|^3 (|k| - |m|) \widehat{h^{(0)}}(m, t) \widehat{h^{(0)}}(k - m, t)
$$

+
$$
\frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} e^{ikx_1} \left(\widehat{I}_1(k, 0, t) + \widehat{I}_2(k, 0, t) \right).
$$
(4.19)

Finally, setting

$$
I(x_1, 0, t) = \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} e^{ikx_1} \left(\widehat{I}_1(k, 0, t) + \widehat{I}_2(k, 0, t) \right), \tag{4.20}
$$

we have found the compact form of F_4

$$
F_4 = \eta \left(\Lambda \left(h^{(0)} \Lambda^3 h^{(0)} \right) - h^{(0)} \Lambda^4 h^{(0)} \right) + I = \eta [\Lambda, h^{(0)}] \Lambda^3 h^{(0)} + I.
$$

We thus rewrite $P^{(1)}$, as

$$
P^{(1)},2 = \eta \Lambda^3 h^{(1)}(x_1, t) + \sum_{i=1}^5 F_i
$$

= $\eta \Lambda^3 h^{(1)}(x_1, t) + \eta [\Lambda, h^{(0)}] \Lambda^3 h^{(0)} + I + K_1^{(1)},$

with $K_1^{(1)}$ $1^{(1)}$ as in (4.16) .

To complete the computations for $\partial_t h^{(1)}$, we still need the sum of the following terms:

$$
\theta S^{(1)},_2(x_1,0,t) = \theta e^{-N_2 t} \mathbf{Q}_1[h^{(0)}(x_1,t)](\mathcal{L}_{,2}(x_1,0,t)) + \theta M(t) \Delta \mathcal{B}_{,2}(x_1,0), \tag{4.21}
$$

$$
h^{(0)},_{1}(x_{1},t)P^{(0)},_{1}(x_{1},0,t)=-\eta h^{(0)},_{1}(x_{1},t)\left(\Lambda^{2}h^{(0)}(x_{1},t)\right),_{1},
$$
\n(4.22)

$$
-\theta h^{(0)}, \mathbf{1}(x_1, t)S^{(0)}, \mathbf{1}(x_1, 0) = -\theta h^{(0)}, \mathbf{1}(x_1, t)e^{-N_2 t}\mathcal{L}, \mathbf{1}(x_1, 0, t).
$$
\n(4.23)

We rewrite the sum of (4.21) , (4.22) , (4.23) as

$$
\theta S^{(1)},_2 + h^{(0)},_1 \left(P^{(0)},_1 - \theta S^{(0)},_1 \right) = -\eta h^{(0)},_1 (x_1, t) \left(\Lambda^2 h^{(0)}(x_1, t) \right),_1 + K_2^{(1)},
$$

being

$$
K_2^{(1)} = \theta e^{-N_2 t} \mathbf{Q}_1[h^{(0)}](\mathcal{L}_{,2}) + \theta M(t) \Delta \mathbf{B}_{,2} - \theta h^{(0)}, e^{-N_2 t} \mathcal{L}_{,1}.
$$
 (4.24)

We can finally gather the previous computations to obtain the following expression of $\partial_t h^{(1)}$:

$$
\partial_t h^{(1)}(x_1, t) = -\eta \Lambda^3 h^{(1)}(x_1, t) - \eta [\![\Lambda, h^{(0)}]\!] \Lambda^3 h^{(0)} - \eta h^{(0)}, \quad (x_1, t) \left(\Lambda^2 h^{(0)}(x_1, t) \right),
$$

+ $K^{(1)}(x_1, 0, t).$

with

$$
K^{(1)}(x_1,0,t) = -I(x_1,0,t) - K_1^{(1)}(x_1,0,t) + K_2^{(1)}(x_1,0,t),
$$
\n(4.25)

for I, $K_1^{(1)}$, $K_2^{(1)}$ as in [\(4.20\)](#page-14-3), [\(4.16\)](#page-12-2), and [\(4.24\)](#page-14-4) respectively. We now perform some manipulations on

$$
-\eta[\![\Lambda, h^{(0)}]\!] \Lambda^3 h^{(0)}
$$
 and $-\eta h^{(0)}, \left(\Lambda^2 h^{(0)}\right),$

in order to rewrite them in terms of the commutator in (1.1) . We have that

$$
- \eta [\Lambda, h^{(0)}] \Lambda^3 h^{(0)} - \eta h^{(0)}, \quad (\Lambda^2 h^{(0)}) ,_1
$$

= $-\eta \left(\Lambda \left(h^{(0)} \Lambda^3 h^{(0)} \right) - h^{(0)} \Lambda^4 h^{(0)} \right) - \eta h^{(0)}, \quad (\Lambda^2 h^{(0)}) ,_1$
= $\eta \left([H, h^{(0)}] H h^{(0)},_{111} \right),_1$,

and then $\partial_t h^{(1)}$ reads

$$
\partial_t h^{(1)}(x_1, t) = -\eta \Lambda^3 h^{(1)}(x_1, t) + \eta \left([H, h^{(0)}(x_1, t)] H h^{(0)}, 111}(x_1, t) \right), \quad 1 + K^{(1)}(x_1, 0, t) \tag{4.26}
$$

for $K^{(1)}$ as in [\(4.25\)](#page-14-5).

4.3. The asymptotic model. We use the equations of $\partial_t h^{(0)}$ in [\(4.6\)](#page-9-4) and of $\partial_t h^{(1)}$ in [\(4.26\)](#page-14-6) to write the equation of $\partial_t h$:

$$
\partial_t h^{(0)} + \varepsilon \partial_t h^{(1)} = -\eta \Lambda^3 h^{(0)} - \varepsilon \eta \Lambda^3 h^{(1)} + \varepsilon \eta \left([H, h^{(0)}] H h^{(0)}, 111 \right), \mathbf{1} + K^{(0)} - \varepsilon K^{(1)}, \quad (4.27)
$$

for $K^{(0)}$ as in [\(4.7\)](#page-9-5) and for $K^{(1)}$ as in [\(4.25\)](#page-14-5). Now, let

$$
g = h^{(0)} + \varepsilon h^{(1)}.
$$

We want to write (4.27) in terms of g. The term with the commutator becomes

$$
\varepsilon\eta\left(\llbracket H, h^{(0)} \rrbracket H h^{(0)}, 111 \right),_1 = \varepsilon\eta\left(\llbracket H, g - \varepsilon h^{(1)} \rrbracket H \left(g - \varepsilon h^{(1)}\right), 111 \right),_1
$$

$$
= \varepsilon\eta\left(\llbracket H, g \rrbracket H g, 111 \right),_1 + O(\varepsilon^2).
$$

We now want to rewrite

$$
I = \widetilde{I} + O(\varepsilon),
$$
 $K_1^{(1)} = \widetilde{K}_1^{(1)} + O(\varepsilon),$ $K_2^{(1)} = \widetilde{K}_2^{(1)} + O(\varepsilon),$

so that $K^{(1)}$ as

$$
K^{(1)} = \widetilde{K}^{(1)} + O(\varepsilon) = -\widetilde{I} - \widetilde{K}_1^{(1)} + \widetilde{K}_2^{(1)} + O(\varepsilon).
$$

To this aim, we first observe that the functionals $\mathbf{Q}_{\alpha}[\ell(x_1, t)]$ and $\mathbf{R}_1[\ell(x_1, t)]$ defined in [\(4.8\)](#page-9-2) and (3.1) respectively can be written in terms of g as

$$
Q_{\alpha}[h^{(0)}(x_1,t)](f(x,t))
$$

= $(h^{(0)}(x_1,t) - g(x_1)) f_{2}(x,t) + \alpha t \Delta f(x,t)$
= $(g(x_1,t) - g(x_1)) f_{2}(x,t) + \alpha t \Delta f(x,t) - \varepsilon h^{(1)}(x_1,t) f_{2}(x,t)$
= $Q_{\alpha}[g(x_1,t)](f(x,t)) + O(\varepsilon),$
 $R_1[h^{(0)}(x_1,t)](f(x,t))$
= $-(h^{(0)},11)} (x_1,t) f(x,t) + 2h^{(0)},1) (x_1,t) f_{11}(x,t) ,$
= $-(\left(g(x_1,t) - \varepsilon h^{(1)}(x_1,t)\right),11} f(x,t) + 2\left(g(x_1,t) - \varepsilon h^{(1)}(x_1,t)\right),1 f_{11}(x,t) ,$
= $R_1[g(x_1,t)](f(x,t)) + O(\varepsilon).$

Then, the terms F_i in [\(4.14\)](#page-12-0), [\(4.12\)](#page-11-1), [\(4.13\)](#page-11-2), [\(4.15\)](#page-12-1) composing $K_1^{(1)}$ $1^{(1)}$ (see (4.16)) become

$$
F_1(x_1, 0, t) = \theta e^{-N_2 t} \int_{-\infty}^0 e^{y_2 \Lambda} Q_1[h^{(0)}(x_1, t)] (\Delta \mathcal{L}(x_1, y_2, t)) dy_2
$$

+ $\theta M(t) \int_{-\infty}^0 e^{y_2 \Lambda} \Delta^2 \mathcal{B}(x_1, y_2) dy_2$
= $\theta e^{-N_2 t} \int_{-\infty}^0 e^{y_2 \Lambda} Q_1[g(x_1, t)] (\Delta \mathcal{L}(x_1, y_2, t)) dy_2$
+ $\theta M(t) \int_{-\infty}^0 e^{y_2 \Lambda} \Delta^2 \mathcal{B}(x_1, y_2) dy_2 + O(\varepsilon)$

$$
F_2(x_1, 0, t) = -\rho e^{-N_2 t} \int_{-\infty}^0 e^{y_2 \Lambda} Q_1[h^{(0)}(x_1, t)] (\mathcal{L}(x_1, y_2, t)) dy_2
$$

- $\rho M(t) \int_{-\infty}^0 e^{y_2 \Lambda} \Delta \mathcal{B}(x_1, y_2)$

A NONLOCAL EQUATION DESCRIBING TUMOR GROWTH $\hfill 17$

$$
= -\rho e^{-N_2 t} \int_{-\infty}^{0} e^{y_2 \Lambda} \mathbf{Q}_1[g(x_1, t)] (\mathcal{L}(x_1, y_2, t)) dy_2
$$

\n
$$
- \rho M(t) \int_{-\infty}^{0} e^{y_2 \Lambda} \Delta \mathbf{B}(x_1, y_2) + O(\varepsilon)
$$

\n
$$
F_3(x_1, 0, t) = \rho \tau e^{-N_1 t} \int_{-\infty}^{0} e^{y_2 \Lambda} \mathbf{Q}_{\alpha}[h^{(0)}(x_1, t)] (\mathbf{B}(x_1, y_2)) dy_2
$$

\n
$$
= \rho \tau e^{-N_1 t} \int_{-\infty}^{0} e^{y_2 \Lambda} \mathbf{Q}_{\alpha}[g(x_1, t)] (\mathbf{B}(x_1, y_2)) dy_2 + O(\varepsilon)
$$

\n
$$
F_5(x_1, 0, t) = -e^{-N_2 t} \int_{-\infty}^{0} e^{y_2 \Lambda} \mathbf{R}_1[h^{(0)}(x_1, t)] (\mathcal{L}(x_1, y_2, t)) dy_2
$$

\n
$$
= -e^{-N_2 t} \int_{-\infty}^{0} e^{y_2 \Lambda} \mathbf{R}_1[g(x_1, t)] (\mathcal{L}(x_1, y_2, t)) dy_2 + O(\varepsilon).
$$

We rewrite their sum, which appears in the definition of $K^{(1)}$, as

$$
K_1^{(1)}(x_1,0,t) = \widetilde{K}_1^{(1)}(x_1,0,t) + O(\varepsilon),
$$

with

$$
\widetilde{K}_{1}^{(1)}(x_{1},0,t) = \theta e^{-N_{2}t} \int_{-\infty}^{0} e^{y_{2}\Lambda} \mathbf{Q}_{1}[g(x_{1},t)](\Delta \mathcal{L}(x_{1},y_{2},t)) dy_{2} + \theta M(t) \int_{-\infty}^{0} e^{y_{2}\Lambda} \Delta^{2} \mathbf{B}(x_{1},y_{2}) dy_{2}
$$
\n
$$
- \rho e^{-N_{2}t} \int_{-\infty}^{0} e^{y_{2}\Lambda} \mathbf{Q}_{1}[g(x_{1},t)](\mathcal{L}(x_{1},y_{2},t)) dy_{2} - \rho M(t) \int_{-\infty}^{0} e^{y_{2}\Lambda} \Delta \mathbf{B}(x_{1},y_{2})
$$
\n
$$
+ \rho \tau e^{-N_{1}t} \int_{-\infty}^{0} e^{y_{2}\Lambda} \mathbf{Q}_{\alpha}[g(x_{1},t)](\mathbf{B}(x_{1},y_{2})) dy_{2}
$$
\n
$$
- e^{-N_{2}t} \int_{-\infty}^{0} e^{y_{2}\Lambda} \mathbf{R}_{1}[g(x_{1},t)](\mathcal{L}(x_{1},y_{2},t)) dy_{2}.
$$
\n(4.28)

Similarly,

$$
\widetilde{K}_2^{(1)}(x_1,0,t) = \theta e^{-N_2 t} \mathbf{Q}_1[g(x_1,t)](\mathcal{L}_{,2}) + \theta M(t) \Delta \mathbf{B}_{,2} - \theta g_{,1}(x_1,t) e^{-N_2 t} \mathcal{L}_{,1}. \tag{4.29}
$$

We now consider the components I_i^k in [\(4.17\)](#page-13-0) and [\(4.18\)](#page-13-1) of the term I defined in [\(4.20\)](#page-14-3):

$$
\begin{split}\n\widehat{I}_{1}(k,0,t) &= -\frac{1}{2} \sum_{m \in \mathbb{Z}} \left[\left(\int_{-\infty}^{0} \widehat{w}^{(0)}(m,y_{2},t) e^{|m|y_{2}} \, dy_{2} \right) \times \\
&\quad \times \int_{-\infty}^{0} e^{y_{2}|k|}(k-m)^{2} \left(\widehat{g}(k-m) - \varepsilon \widehat{h^{(1)}}(k-m) \right) \left(e^{|m|y_{2}} + e^{-|m|y_{2}} \right) \, dy_{2} \right] \\
&\quad + \frac{1}{2} \sum_{m \in \mathbb{Z}} \left[\int_{-\infty}^{0} e^{y_{2}|k|}(k-m)^{2} \left(\widehat{g}(k-m) - \varepsilon \widehat{h^{(1)}}(k-m) \right) \times \\
&\quad \times \left(\int_{0}^{y_{2}} \widehat{w}^{(0)}(m,y_{2}',t) \left(e^{-|m|(y_{2}-y_{2}')} + e^{|m|(y_{2}-y_{2}')} \right) \, dy_{2} \right) \, dy_{2} \right], \\
&= \widehat{I}_{1}(k,0,t) + O(\varepsilon), \\
\widehat{I}_{2}(k,0,t) &= \sum_{m \in \mathbb{Z}} \left[\left(\int_{-\infty}^{0} \widehat{w}^{(0)}(m,y_{2},t) e^{|m|y_{2}} \, dy_{2} \right) \times \\
\end{split}
$$

A NONLOCAL EQUATION DESCRIBING TUMOR GROWTH $\hfill 18$

$$
\times \int_{-\infty}^{0} e^{y_2|k|} (k - m) \left(\hat{g}(k - m) - \varepsilon \widehat{h^{(1)}}(k - m) \right) \left(e^{|m|y_2} + e^{-|m|y_2} \right) dy_2 \Big]
$$

$$
- \sum_{m \in \mathbb{Z}} \left[\int_{-\infty}^{0} e^{y_2|k|} (k - m) \left(\hat{g}(k - m) - \varepsilon \widehat{h^{(1)}}(k - m) \right) \times \right.
$$

$$
\times \left(\int_{0}^{y_2} \hat{w}^{(0)}(m, y_2', t) \left(e^{-|m|(y_2 - y_2')} + e^{|m|(y_2 - y_2')} \right) dy_2' \right) dy_2 \Big]
$$

$$
= \widehat{I}_2(k, 0, t) + O(\varepsilon),
$$

being

$$
I_{1}(k,0,t)
$$
\n
$$
= -\frac{1}{2} \sum_{m \in \mathbb{Z}} \left(\int_{-\infty}^{0} \widehat{w}^{(0)}(m,y_{2},t) e^{|m|y_{2}} dy_{2} \right) \times
$$
\n
$$
\times \int_{-\infty}^{0} e^{y_{2}|k|}(k-m)^{2} \widehat{g}(k-m) \left(e^{|m|y_{2}} + e^{-|m|y_{2}} \right) dy_{2}
$$
\n
$$
+ \frac{1}{2} \sum_{m \in \mathbb{Z}} \int_{-\infty}^{0} e^{y_{2}|k|}(k-m)^{2} \widehat{g}(k-m) \times
$$
\n
$$
\times \left(\int_{0}^{y_{2}} \widehat{w}^{(0)}(m,y_{2}',t) \left(e^{-|m|(y_{2}-y_{2}')} + e^{|m|(y_{2}-y_{2}')} \right) dy_{2}' \right) dy_{2}, \qquad (4.30)
$$
\n
$$
\widehat{I}_{2}(k,0,t)
$$

$$
= \sum_{m\in\mathbb{Z}} \left(\int_{-\infty}^{0} \widehat{w}^{(0)}(m, y_2, t) e^{|m|y_2} dy_2 \right) \int_{-\infty}^{0} e^{y_2|k|} (k - m) \widehat{g}(k - m) \left(e^{|m|y_2} + e^{-|m|y_2} \right) dy_2
$$

$$
- \sum_{m\in\mathbb{Z}} \int_{-\infty}^{0} e^{y_2|k|} (k - m) \widehat{g}(k - m) \times
$$

$$
\times \left(\int_{0}^{y_2} \widehat{w}^{(0)}(m, y_2', t) \left(e^{-|m|(y_2 - y_2')} + e^{|m|(y_2 - y_2')} \right) dy_2 \right) dy_2.
$$
(4.31)

Then, we can rewrite the term I as

$$
I(x_1,0,t) = \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} e^{ikx_1} \left(\widehat{\tilde{I}}_1(k,0,t) + \widehat{\tilde{I}}_2(k,0,t) \right) + O(\varepsilon) = \widetilde{I}(x_1,0,t) + O(\varepsilon).
$$
 (4.32)

Finally, the equation (4.27) in terms of g is

$$
\partial_t g(x_1, t) = -\eta \Lambda^3 g(x_1, t) + \varepsilon \eta \left([H, g(x_1, t)] H g_{111}(x_1, t) \right),_1 - \varepsilon \theta g(x_1, t) e^{-N_2 t} \mathcal{L}_{,1}(x_1, 0, t) + \varepsilon \widetilde{K}^{(1)}(x_1, 0, t) + K^{(0)}(x_1, 0, t) + O(\varepsilon^2).
$$
\n(4.33)

As a consequence, our asymptotic model reads

$$
\partial_t g(x_1, t) = -\eta \Lambda^3 g(x_1, t) + \varepsilon \eta \left([H, g(x_1, t)] H g_{1,111}(x_1, t) \right),_1 - \varepsilon \theta g(x_1, t) e^{-N_2 t} \mathcal{L}_{11}(x_1, 0, t) + \varepsilon \widetilde{K}^{(1)}(x_1, 0, t) + K^{(0)}(x_1, 0, t).
$$

5. A particular case

We focus our attention on the particular cases where both the inhibitor and the nutrient depend only on the depth x_2 .

To simplify the computations below, we assume that inhibitor and nutrient have the same parameters, and we use the following notation: $D_i = D_n = D$, $\delta_i = \delta_n = \delta$, and so on. We also set $\tau = 1$. In the previous sections, we supposed that $M_1 = M_2 = \omega = O(\varepsilon^2)$. Then, equations $(2.1), (2.2), (2.3), (2.4)$ $(2.1), (2.2), (2.3), (2.4)$ $(2.1), (2.2), (2.3), (2.4)$ $(2.1), (2.2), (2.3), (2.4)$ $(2.1), (2.2), (2.3), (2.4)$ $(2.1), (2.2), (2.3), (2.4)$ $(2.1), (2.2), (2.3), (2.4)$ become

$$
\varepsilon = \frac{H}{L}, \qquad \alpha = 1,
$$

\n
$$
N_1 = N_2 = N = L H \frac{\delta + \lambda}{D}, \qquad N_3 = \gamma \frac{L H}{D},
$$

\n
$$
M_1 = 0, \qquad M_2 = 0, \qquad \theta = \frac{\chi \tilde{\sigma}}{D}, \qquad \rho = \frac{\mu \tilde{\sigma} L^2}{D}, \qquad \omega = 0, \qquad \eta = \frac{\nu H}{L^2 D}.
$$
\n(5.1)

Note that, with this choice of parameters, we have that the function $M(t)$ defined in [\(4.10\)](#page-10-2) is zero.

If the data do not depend on x_1 , then the [\[29,](#page-22-27) Lemma A1] and the computations we made before apply with $k = 0$. Indeed, suppose that $b = b(x_2)$ and $g \equiv 0$ in [\[29,](#page-22-27) Lemma A1, Eq. (66)]. Hence, passing to Fourier, the equation $\Delta u = b$ becomes

$$
-k^{2}\widehat{u}(k,x_{2})+\widehat{u}_{22}(k,x_{2})=\widehat{b}(x_{2}).
$$

The lack of dependence on k of \hat{b} implies that $k = 0$ in the whole equation, hence \hat{u} verifies

$$
\widehat{u}_{22}(x_2) = \widehat{b}(x_2).
$$

Remark 5.1. This is not in contrast with $[29, \text{Lemma A1}, \text{Eqs (71)} - (72)]$ $[29, \text{Lemma A1}, \text{Eqs (71)} - (72)]$. Indeed, we first use [\[29,](#page-22-27) Lemma A1, Eq (72)] in [29, Lemma A1, Eq (71)], and then we take the limit on $k \to 0$ and we derive twice in x_2 , we obtain the same expression of \widehat{u}_{22} .

We consider the following initial data:

$$
\mathcal{B}(x) = c_{\mathcal{B}}e^{x_2}\sin x_2 \quad \text{and} \quad \mathcal{S}(x) = c_{\mathcal{S}}e^{x_2}\sin x_2 \quad \text{for } c_{\mathcal{B}}, c_{\mathcal{S}} > 0.
$$

We want to simplify (4.33) with this choice. This means that we have to find the expressions of $K^{(0)}$ and $\widetilde{K}^{(1)}$ according to this data.

The definition of $\mathcal L$ in [\(4.4\)](#page-8-11) becomes

$$
\mathcal{L}(x_2, t) = \alpha(t)e^{x_2}\sin x_2
$$
 where $\alpha(t) = c_8 - N_3c_8t$.

Then, the third term in the r.h.s. of [\(4.33\)](#page-17-1) is zero.

We now focus on the term $K^{(0)}$, whose general expression is given in [\(4.7\)](#page-9-5). When no dependence on x_1 is considered and with the choice of parameters given in (5.1) , it reduces to

$$
K^{(0)}(0,t) = -\theta e^{-Nt} \int_{-\infty}^{0} \mathcal{L}_{22}(y_2,t) dy_2 + \rho e^{-Nt} \int_{-\infty}^{0} (\mathcal{L}(y_2,t) - \mathcal{B}(y_2)) dy_2
$$

+ $\theta e^{-Nt} \mathcal{L}_{22}(0,t).$

Since

$$
\mathcal{L}_{2}(x_2,t) = \alpha(t)e^{x_2}(\sin x_2 + \cos x_2) \quad \text{and} \quad \mathcal{L}_{2}(x_2,t) = 2\alpha(t)e^{x_2}\cos x_2,
$$

we simplify $K^{(0)}$ as

$$
K^{(0)}(0,t) = e^{-Nt} \left[-2\theta \alpha(t) \int_{-\infty}^{0} e^{y_2} \cos y_2 \, dy_2 + \rho \left(\alpha(t) - c_{\mathcal{B}} \right) \int_{-\infty}^{0} e^{y_2} \sin y_2 \, dy_2 + \theta \alpha(t) \right].
$$

This expression can be further simplified using that

$$
\int_{-\infty}^{0} e^{y_2} \sin y_2 dy_2 = -\int_{-\infty}^{0} e^{y_2} \cos y_2 dy_2 = -\frac{1}{2},
$$

then $K^{(0)}(0, t)$ is given by

$$
K^{(0)}(0,t) = -\frac{e^{-Nt}}{2}\rho\left(\alpha(t) - c_{\mathcal{B}}\right). \tag{5.2}
$$

We now consider $\widetilde{K}_1^{(1)}(x_1,0,t) \equiv \widetilde{K}_1^{(1)}(0,t)$ (see [\(4.28\)](#page-16-0) for its general expression):

$$
\widetilde{K}_1^{(1)}(0,t) = \theta e^{-Nt} \int_{-\infty}^0 \mathbf{Q}_1[g(x_1,t)](\mathcal{L}_{,22}(y_2,t)) dy_2 - \rho e^{-Nt} \int_{-\infty}^0 \mathbf{Q}_1[g(x_1,t)](\mathcal{L}(y_2,t)) dy_2
$$

$$
+ \rho e^{-Nt} \int_{-\infty}^0 \mathbf{Q}_1[g(x_1,t)](\mathcal{B}(y_2)) dy_2 - e^{-Nt} \int_{-\infty}^0 \mathbf{R}_1[g(x_1,t)](\mathcal{L}(y_2,t)) dy_2.
$$

We have

$$
Q_1[g(x_1,t)](\mathcal{L}(x_2,t)) = \alpha(t) [(g(x_1,t) - g(x_1)) (\sin x_2 + \cos x_2) + 2t \cos x_2] e^{x_2},
$$

\n
$$
Q_1[g(x_1,t)](\mathcal{L}_{22}(x_2,t)) = 2\alpha(t) [(g(x_1,t) - g(x_1)) (\cos x_2 - \sin x_2) - 2t \sin x_2] e^{x_2},
$$

\n
$$
Q_1[g(x_1,t)](\mathcal{B}(x_2)) = (g(x_1,t) - g(x_1)) c_2 e^{x_2} (\sin x_2 + \cos x_2) + 2t c_2 e^{x_2} \cos x_2,
$$

\n
$$
R_1[g(x_1,t)](\mathcal{L}(x_2,t)) = -g(x_1,t)_{11} \alpha(t) e^{x_2} (\sin x_2 + \cos x_2).
$$

The integrals of the above quantities are the following:

$$
\int_{-\infty}^{0} \mathbf{Q}_1[g(x_1,t)](\mathcal{L}(y_2,t)) dy_2 = \alpha(t)t,
$$
\n
$$
\int_{-\infty}^{0} \mathbf{Q}_1[g(x_1,t)](\Delta \mathcal{L}(y_2,t)) dy_2 = 2\alpha(t) (g(x_1,t) - g(x_1)) + 2\alpha(t)t,
$$
\n
$$
\int_{-\infty}^{0} \mathbf{Q}_1[g(x_1,t)](\mathcal{B}(y_2)) dy_2 = t c_{\mathcal{B}},
$$
\n
$$
\int_{-\infty}^{0} \mathbf{R}_1[g(x_1,t)](\mathcal{L}(y_2,t)) dy_2 = 0.
$$

Then, gathering the above computations, we find that

$$
\widetilde{K}_1^{(1)}(0,t) = 2\theta e^{-Nt} \alpha(t) \left(g(x_1,t) - g(x_1) \right) + e^{-Nt} \alpha(t) (2\theta - \rho)t + \rho e^{-Nt} t c_{\mathcal{B}}.
$$
\n(5.3)

We now deal with $\widetilde{K}_2^{(1)}(x_1, 0, t) \equiv \widetilde{K}_2^{(1)}(0, t)$:

$$
\widetilde{K}_2^{(1)}(0,t) = \theta e^{-Nt} \mathbf{Q}_1[g(x_1,t)](\mathbf{L}_{2}(0,t)) - \theta g_{1}(x_1,t)e^{-Nt}\mathbf{L}_{1}(0,t),
$$

(see (4.29)). Since

$$
Q_1[g(x_1,t)](\mathcal{L}_{,2}(0,t)) = (g(x_1,t) - g(x_1))\mathcal{L}_{,22}(0,t) + t\Delta \mathcal{L}_{,2}(0,t)
$$

= 2\alpha(t)((g(x_1,t) - g(x_1)) + t),

$$
\Delta \mathcal{B}_{,2}(0) = 2c_{\mathcal{B}},
$$

then

$$
\widetilde{K}_2^{(1)}(0,t) = \theta e^{-Nt} 2\alpha(t) \left(\left(g(x_1,t) - g(x_1) \right) + t \right). \tag{5.4}
$$

We now compute $\widetilde{I}(x_1, 0, t) = \widetilde{I}(0, t)$. This term has been defined in [\(4.32\)](#page-17-2), and its expression can be deduced from (4.30) and (4.31) . The quantities in (4.30) and (4.31) follows from an application of $[29, \text{Lemma A.1}]$ (see also $(4.19), (4.17),$ $(4.19), (4.17),$ $(4.19), (4.17),$ and (4.18)). In this particular case, $[29, \text{$ Lemma A.1] applies with $b = b(x_2)$ and $g = g(x_1)$, so that $\overline{I}(0,t)$ simplifies as

$$
\widetilde{I}(0,t) = \frac{1}{\sqrt{2\pi}} \frac{\rho}{2} \alpha(t) e^{-Nt} \sum_{n \in \mathbb{Z}} n^2 \widehat{g}(n,t) = -\frac{\rho}{2} \alpha(t) e^{-Nt} g_{111}(x_1,t).
$$
\n(5.5)

Finally, we are ready to simplify (4.33) using (5.2) , (5.3) , (5.4) , and (5.5) as follows: $\partial_t g(x_1, t)$ = $-\eta \Lambda^{3} g(x_1, t) + \varepsilon \eta \left([H, g(x_1, t)] H g_{1111}(x_1, t) \right)$,

$$
-\varepsilon \left(2\theta e^{-Nt}\alpha(t)\left(g(x_1,t) - g(x_1)\right) + e^{-Nt}\alpha(t)\left(2\theta - \rho\right)t + \rho e^{-Nt}t c_{\mathcal{B}}\right) + \theta \varepsilon e^{-Nt} 2\alpha(t)\left(\left(g(x_1,t) - g(x_1)\right) + t\right) - \varepsilon \frac{\rho}{2}\alpha(t)e^{-Nt}g_{11}(x_1,t) - \frac{e^{-Nt}}{2}\rho\left(\alpha(t) - c_{\mathcal{B}}\right),
$$
(5.6)

being $g(x_1) = g(x_1, 0)$.

We now prove the local well-posedness of (5.6) . We here recall the definition of Wiener spaces $A^s(\mathbb{T})$

$$
A^{s}(\mathbb{T}) = \left\{ u \in L^{1}(\mathbb{T}), \quad \|u\|_{A^{s}(\mathbb{T})} := \sum_{k \in \mathbb{Z}} |k|^{s} |\widehat{u}(k)| < \infty \right\}.
$$

In the following, we use the notation

$$
||f||_{A^s} = ||f||_{A^s(\mathbb{T})}.
$$

Then, the main result reads

Theorem 5.1 (Well-posedness). Let $g \in A^1(\mathbb{T})$ be such that

$$
\|\boldsymbol{g}\|_{A^1}\ll 1.
$$

Then, there exists a time T and a unique solution

$$
g \in C([0, T], A^1(\mathbb{T})) \cap L^1(0, T; A^4(\mathbb{T}))
$$

of [\(5.6\)](#page-20-0).

Proof. Sketch of the proof: The proof follows the ideas in [\[30\]](#page-22-28). Finally, we need to impose a smallness asumption on the initial data to close the estimates. Furthermore, due to the forcing induced by the nutrients and inhibitors, our estimates allow to a controlled norm growth which eventually imply that the smallness assumption on the initial data does not propagate in time. As a consequence, we have a local in time existence for small initial data.

This goal is achieved in different steps.

A priori estimate on the mean of q : We compute

$$
\frac{d}{dt} \int_{\mathbb{T}} g(x,t) dx \leq e^{-Nt} \gamma(t) \left| \int_{\mathbb{T}} g(x,t) dx \right| + \gamma(t) e^{-Nt},
$$

for certain (explicit) $\gamma(t)$ depending on the initial data and the parameters of the problem that grows linearly in time and may change from line to line.

A priori estimate on the derivative of g : We will make use of the following Poincaré type inequalities

$$
||g||_{A^1} \le \left| \int_{\mathbb{T}} g(x, t) dx \right| + ||g||_{\dot{A}^1},
$$

$$
||g||_{\dot{A}^0} \le ||g||_{\dot{A}^1}.
$$

We have that

$$
\sum_{k \in \mathbb{Z}} |k| \partial_t |\hat{g}(k, t)| = \frac{d}{dt} ||g(t)||_{\dot{A}^1},
$$

$$
-\eta \sum_{k \in \mathbb{Z}} |k|^4 |\hat{g}(k, t)| = -\eta ||g(t)||_{\dot{A}^4},
$$

$$
2\eta \sum_{k \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} |k||m||k - m|^3 |\hat{g}(k - m, t)| |\hat{g}(m, t)| \le 2\eta ||g(t)||_{A^1} ||g(t)||_{A^4}.
$$

Thus,

$$
\frac{d}{dt}\left\|g(t)\right\|_{\dot{A}^1} \leq -\eta \left(1-2\left\|g(t)\right\|_{\dot{A}^1}\right) \left\|g(t)\right\|_{\dot{A}^4} + e^{-Nt} \left(\left\|g(t)\right\|_{\dot{A}^3} + \left\|{\bm g}\right\|_{\dot{A}^1} + 1 \right) \gamma(t).
$$

Using interpolation, we find that

$$
\frac{d}{dt} \|g(t)\|_{\dot{A}^1} \leq -\eta \left(1-2\|g(t)\|_{\dot{A}^1}\right) \|g(t)\|_{\dot{A}^4} + e^{-Nt} \left(\|g(t)\|_{\dot{A}^1}^{1/3} \|g(t)\|_{\dot{A}^4}^{2/3} + \|g\|_{\dot{A}^1} + 1 \right) \gamma(t),
$$

and, using Young inequality, we conclude that

$$
\frac{d}{dt}\left\|g(t)\right\|_{\dot{A}^1} \leq -\eta\left(1-\gamma(t)\delta e^{-Nt}-2\left\|g(t)\right\|_{\dot{A}^1}\right)\left\|g(t)\right\|_{\dot{A}^4}+e^{-Nt}\left(\frac{1}{\delta}\left\|g(t)\right\|_{\dot{A}^1}+\left\|{\pmb g}\right\|_{\dot{A}^1}+1\right)\gamma(t),
$$

for $\delta > 0$ artbitrary. Using the smallness of the initial data together with taking a small enough $\delta > 0$ we obtain that

$$
\frac{d}{dt} ||g(t)||_{\dot{A}^1} + c\eta ||g(t)||_{\dot{A}^4} \leq e^{-Nt} \gamma(t) (||g(t)||_{\dot{A}^1} + 1).
$$

Closing the estimates: Collecing the previous estimates we find the inequality

$$
\frac{d}{dt} ||g(t)||_{A^1} \le e^{-Nt} \gamma(t) (||g(t)||_{A^1} + 1).
$$

From where a uniform time of existence $0 < T \ll 1$ can be concluded. This time cannot, in general, be taken arbitrarily large because the smallness condition may not be satisfied for large time intervals.

Uniqueness: The uniqueness follows from a standard contradiction argument together with the parabolic gain of regularity.

 \Box

ACKNOWLEDGMENTS

R.G-B was supported by the project "Mathematical Analysis of Fluids and Applications" Grant PID2019-109348GA-I00 funded by MCIN/AEI/ 10.13039/501100011033 and acronym "MAFyA". This publication is part of the project PID2019-109348GA-I00 funded by MCIN/ AEI /10.13039/501100011033. R.G-B is also supported by a 2021 Leonardo Grant for Researchers and Cultural Creators, BBVA Foundation. The BBVA Foundation accepts no responsibility for the opinions, statements, and contents included in the project and/or the results thereof, which are entirely the responsibility of the authors.

The work of M.M. was partially supported by a public grant as part of the Investissement d'avenir project, FMJH, and by Grant RYC2021-033698-I, funded by the Ministry of Science and Innovation/State Research Agency/10.13039/501100011033 and by the European Union "NextGenerationEU/Recovery, Transformation and Resilience Plan". Both authors are funded by the project "Análisis Matemático Aplicado y Ecuaciones Diferenciales" Grant PID2022-141187NB-I00 funded by MCIN /AEI /10.13039/501100011033 / FEDER, UE and acronym "AMAED". This publication is part of the project PID2022-141187NB-I00 funded by MCIN/ AEI /10.13039/501100011033.

REFERENCES

- [1] Thomas Alazard. Stabilization of gravity water waves. *Journal de Mathématiques Pures et Appliquées*, 114:51– 84, 2018.
- [2] Alexander RA Anderson and Vito Quaranta. Integrative mathematical oncology. *Nature Reviews Cancer*, 8(3):227, 2008.
- [3] Robyn P Araujo and DL Sean McElwain. A history of the study of solid tumour growth: the contribution of mathematical modelling. *Bulletin of mathematical biology*, 66(5):1039–1091, 2004.
- [4] Nicola Bellomo, Abdelghani Bellouquid, Youshan Tao, and Michael Winkler. Toward a mathematical theory of keller–segel models of pattern formation in biological tissues. *Mathematical Models and Methods in Applied Sciences*, 25(09):1663–1763, 2015.
- [5] Beatriz Blanco, Juan Campos, Juan Melchor, and Juan Soler. Modeling interactions among migration, growth and pressure in tumor dynamics. *Mathematics*, 9(12):1376, 2021.
- [6] Jan Burczak and Rafael Granero-Belinchón. On a generalized doubly parabolic keller–segel system in one spatial dimension. *Mathematical Models and Methods in Applied Sciences*, 26(01):111–160, 2016.
- [7] H M Byrne and MAJ Chaplain. Growth of necrotic tumors in the presence and absence of inhibitors. *Mathematical biosciences*, 135(2):187–216, 1996.
- [8] Helen M Byrne. Dissecting cancer through mathematics: from the cell to the animal model. *Nature Reviews Cancer*, 10(3):221, 2010.
- [9] HM Byrne and M A J Chaplain. Growth of nonnecrotic tumors in the presence and absence of inhibitors. *Mathematical biosciences*, 130(2):151–181, 1995.
- [10] HM Byrne and Mark AJ Chaplain. Modelling the role of cell-cell adhesion in the growth and development of carcinomas. *Mathematical and Computer Modelling*, 24(12):1–17, 1996.
- [11] Ángel Castro, Diego Córdoba, Charles Fefferman, Francisco Gancedo, and María López-Fernández. Rayleightaylor breakdown for the muskat problem with applications to water waves. *Annals of Mathematics*, pages 909–948, 2012.
- [12] MAJ Chaplain and Georgios Lolas. Mathematical modelling of cancer cell invasion of tissue: The role of the urokinase plasminogen activation system. *Mathematical Models and Methods in Applied Sciences*, 15(11):1685–1734, 2005.
- [13] Pierluigi Colli, Gianni Gilardi, and Danielle Hilhorst. On a cahn-hilliard type phase field system related to tumor growth. *arXiv preprint arXiv:1401.5943*, 2014.
- [14] Pierluigi Colli, Gianni Gilardi, Elisabetta Rocca, and Jürgen Sprekels. Asymptotic analyses and error estimates for a cahn-hilliard type phase field system modelling tumor growth. *Discrete & Continuous Dynamical Systems-Series S*, 10(1), 2017.
- [15] Pierluigi Colli, Gianni Gilardi, Elisabetta Rocca, and Jürgen Sprekels. Optimal distributed control of a diffuse interface model of tumor growth. *Nonlinearity*, 30(6):2518, 2017.
- [16] Martina Conte, Sergio Casas-Tintò, and Juan Soler. Modeling invasion patterns in the glioblastoma battlefield. *PLoS computational biology*, 17(1):e1008632, 2021.
- [17] Daniel Coutand and Steve Shkoller. On the finite-time splash and splat singularities for the 3-d free-surface euler equations. *Communications in Mathematical Physics*, 325(1):143–183, 2014.
- [18] Shangbin Cui and Avner Friedman. Analysis of a mathematical model of the effect of inhibitors on the growth of tumors. *Mathematical Biosciences*, 164(2):103–137, 2000.
- [19] Shangbin Cui and Avner Friedman. Analysis of a mathematical model of the growth of necrotic tumors. *Journal of Mathematical Analysis and Applications*, 255(2):636–677, 2001.
- [20] Avner Friedman. A hierarchy of cancer models and their mathematical challenges. *Discrete and Continuous Dynamical Systems Series B*, 4(1):147–160, 2004.
- [21] Avner Friedman. Mathematical analysis and challenges arising from models of tumor growth. *Mathematical Models and Methods in Applied Sciences*, 17(supp01):1751–1772, 2007.
- [22] Avner Friedman and Fernando Reitich. Analysis of a mathematical model for the growth of tumors. *Journal of mathematical biology*, 38:262–284, 1999.
- [23] Harald Garcke, Kei Fong Lam, Robert Nürnberg, and Emanuel Sitka. A multiphase cahn–hilliard–darcy model for tumour growth with necrosis. *Mathematical Models and Methods in Applied Sciences*, 28(03):525– 577, 2018.
- [24] Harald Garcke, Kei Fong Lam, Emanuel Sitka, and Vanessa Styles. A cahn–hilliard–darcy model for tumour growth with chemotaxis and active transport. *Mathematical Models and Methods in Applied Sciences*, 26(06):1095–1148, 2016.
- [25] Robert A Gatenby and Edward T Gawlinski. A reaction-diffusion model of cancer invasion. *Cancer research*, 56(24):5745–5753, 1996.
- [26] Alf Gerisch and Mark AJ Chaplain. Mathematical modelling of cancer cell invasion of tissue: local and non-local models and the effect of adhesion. *Journal of Theoretical Biology*, 250(4):684–704, 2008.
- [27] Rafael Granero-Belinch´on. A nonlocal model describing tumor angiogenesis. *Nonlinear Analysis*, 227:113180, 2023.
- [28] Rafael Granero-Belinch´on and Omar Lazar. Growth in the muskat problem. *Mathematical Modelling of Natural Phenomena*, 15:7, 2020.
- [29] Rafael Granero-Belinch´on and Stefano Scrobogna. Models for damped water waves. *SIAM Journal on Applied Mathematics*, 79(6):2530–2550, 2019.
- [30] Rafael Granero-Belinchón and Stefano Scrobogna. On an asymptotic model for free boundary darcy flow in porous media. *SIAM Journal on Mathematical Analysis*, 52(5):4937–4970, 2020.
- [31] HP Greenspan. Models for the growth of a solid tumor by diffusion. *Studies in Applied Mathematics*, 51(4):317–340, 1972.
- [32] HP Greenspan. On the growth and stability of cell cultures and solid tumors. *Journal of theoretical biology*, 56(1):229–242, 1976.
- [33] Thomas Hillen, Kevin J Painter, and Michael Winkler. Convergence of a cancer invasion model to a logistic chemotaxis model. *Mathematical Models and Methods in Applied Sciences*, 23(01):165–198, 2013.
- [34] Jong Bin Kim, Robert Stein, and Mike J O'hare. Three-dimensional in vitro tissue culture models of breast cancer—a review. *Breast cancer research and treatment*, 85(3):281–291, 2004.
- [35] David Lannes. *The water waves problem: mathematical analysis and asymptotics*, volume 188. American Mathematical Soc., 2013.
- [36] EABF Lima, JT Oden, DA Hormuth, TE Yankeelov, and RC Almeida. Selection, calibration, and validation of models of tumor growth. *Mathematical Models and Methods in Applied Sciences*, 26(12):2341–2368, 2016.
- [37] John S Lowengrub, Hermann B Frieboes, Fang Jin, Yao-Li Chuang, Xiangrong Li, Paul Macklin, Steven M Wise, and Vittorio Cristini. Nonlinear modelling of cancer: bridging the gap between cells and tumours. *Nonlinearity*, 23(1):R1, 2009.
- [38] Wolfgang Mueller-Klieser. Three-dimensional cell cultures: from molecular mechanisms to clinical applications. *American Journal of Physiology-Cell Physiology*, 273(4):C1109–C1123, 1997.
- [39] Lev Naumov, Alfons Hoekstra, and Peter Sloot. The influence of mitoses rate on growth dynamics of a cellular automata model of tumour growth. *Procedia Computer Science*, 1(1):971–978, 2010.
- [40] J Tinsley Oden, Andrea Hawkins, and Serge Prudhomme. General diffuse-interface theories and an approach to predictive tumor growth modeling. *Mathematical Models and Methods in Applied Sciences*, 20(03):477–517, 2010.
- [41] J Tinsley Oden, Ernesto ABF Lima, Regina C Almeida, Yusheng Feng, Marissa Nichole Rylander, David Fuentes, Danial Faghihi, Mohammad M Rahman, Matthew DeWitt, Manasa Gadde, et al. Toward predictive multiscale modeling of vascular tumor growth: computational and experimental oncology for tumor prediction. *Archives of Computational Methods in Engineering*, 23:735–779, 2016.
- [42] Kevin J Painter and Thomas Hillen. Spatio-temporal chaos in a chemotaxis model. *Physica D: Nonlinear Phenomena*, 240(4-5):363–375, 2011.
- [43] Mohammad Mamunur Rahman, Yusheng Feng, Thomas E Yankeelov, and J Tinsley Oden. A fully coupled space–time multiscale modeling framework for predicting tumor growth. *Computer methods in applied mechanics and engineering*, 320:261–286, 2017.
- [44] HL Rocha, RC Almeida, EABF Lima, ACM Resende, JT Oden, and TE Yankeelov. A hybrid three-scale model of tumor growth. *Mathematical Models and Methods in Applied Sciences*, 28(01):61–93, 2018.
- [45] Christian Stinner, Christina Surulescu, and Michael Winkler. Global weak solutions in a pde-ode system modeling multiscale cancer cell invasion. *SIAM Journal on Mathematical Analysis*, 46(3):1969–2007, 2014.
- [46] Robert M Sutherland, John A McCredie, and W Rodger Inch. Growth of multicell spheroids in tissue culture as a model of nodular carcinomas. *Journal of the National Cancer Institute*, 46(1):113–120, 1971.
- [47] ZUZANNA SZYMAŃSKA, Cristian Morales Rodrigo, MIROSŁAW LACHOWICZ, and Mark AJ Chaplain. Mathematical modelling of cancer invasion of tissue: the role and effect of nonlocal interactions. *Mathematical Models and Methods in Applied Sciences*, 19(02):257–281, 2009.
- [48] Youshan Tao and Mingjun Wang. Global solution for a chemotactic–haptotactic model of cancer invasion. *Nonlinearity*, 21(10):2221, 2008.
- [49] Zhihui Wang, Joseph D Butner, Romica Kerketta, Vittorio Cristini, and Thomas S Deisboeck. Simulating cancer growth with multiscale agent-based modeling. In *Seminars in cancer biology*, volume 30, pages 70–78. Elsevier, 2015.
- [50] Zhihui Wang, Le Zhang, Jonathan Sagotsky, and Thomas S Deisboeck. Simulating non-small cell lung cancer with a multiscale agent-based model. *Theoretical Biology and Medical Modelling*, 4(1):1–14, 2007.
- [51] Anna Zhigun, Christina Surulescu, and Aydar Uatay. Global existence for a degenerate haptotaxis model of cancer invasion. *Zeitschrift für angewandte Mathematik und Physik*, 67:1–29, 2016.

DEPARTAMENTO DE MATEMÁTICAS, ESTADÍSTICA Y COMPUTACIÓN, UNIVERSIDAD DE CANTABRIA, SPAIN.

Email address: rafael.granero@unican.es

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, UNIVERSIDAD DE SEVILLA, SPAIN.

Email address: mmagliocca@us.es