The analogue of overlap-freeness for the Fibonacci morphism

James D. Currie & Narad Rampersad Department of Mathematics & Statistics The University of Winnipeg j.currie@uwinnipeg.ca, n.rampersad@uwinnipeg.ca

November 23, 2023

Abstract

A 4⁻-power is a non-empty word of the form $XXXX^-$, where X^- is obtained from X by erasing the last letter. A binary word is called *faux-bonacci* if it contains no 4⁻-powers, and no factor 11. We show that faux-bonacci words bear the same relationship to the Fibonacci morphism that overlap-free words bear to the Thue-Morse morphism. We prove the analogue of Fife's Theorem for faux-bonacci words, and characterize the lexicographically least and greatest infinite faux-bonacci words.

Mathematics Subject Classifications: 68R15

1 Introduction

We study binary words, that is words over the alphabet $\mathcal{B} = \{0, 1\}$. We use lower case letters (e.g., w) to denote finite words, and we use bold-face letters to denote words with letters indexed by \mathbb{N} ; e.g.,

 $\boldsymbol{w}=w_1w_2w_3\cdots$.

In the literature, words with letters indexed by \mathbb{N} are variously referred to as ω -words, infinite words, one-sided infinite words, etc. In this article we

refer to them as ω -words. We freely use notions from combinatorics on words and from automata theory. Thus, for example, the set of finite words over \mathcal{B} is denoted by \mathcal{B}^* , and the set of ω -words is denoted by \mathcal{B}^{ω} . We record morphisms inline, i.e., g = [g(0), g(1)].

The binary overlap-free words constitute a classical object of study in combinatorics on words. They are particularly well understood because of their intimate connection to the Thue-Morse morphism $\mu = [01, 10]$.

Theorem 1.1. Let w be binary word. Then w is overlap-free if and only if $\mu(w)$ is overlap-free.

Thue [15] proved that, for two-sided infinite words and for circular words, every overlap-free binary word arises as the μ image of an overlap-free word. The analysis of finite words is more complicated, but these also arise via iterating μ . (See, e.g., Restivo and Salemi [14].)

Theorem 1.2. Let $w \in \mathcal{B}^*$ be overlap-free. Then we can write $w = a\mu(u)b$, where $a, b \in \{\epsilon, 0, 00, 1, 11\}$, and u is overlap-free. If $|w| \ge 7$ this factorization is unique. If $w \in \mathcal{B}^{\omega}$ is overlap-free, then we can write $w = a\mu(u)$, for some overlap-free word $u \in \mathcal{B}^{\omega}$ where $a \in \{\epsilon, 0, 00, 1, 11\}$.

Characterizations of binary overlap-free words in terms of μ have allowed sharp enumerations of these words [14, 12, 5, 6, 11, 10]. These enumerations are closely connected to a classical result known as Fife's Theorem [9].

Theorem 1.3 (Fife's Theorem). Let $u \in \mathcal{B}^{\omega}$. Then u is overlap-free if and only if

$$oldsymbol{u} = w ullet oldsymbol{f}$$

for some $w \in \{01, 001\}$, and some $f \in \{\alpha, \beta, \gamma\}^{\omega}$ containing no factor in

$$I = (\alpha + \beta)(\gamma \gamma)^* (\beta \alpha + \gamma \beta + \alpha \gamma).$$

In this theorem, each of α , β , and γ is an operator that maps a finite word ending in $\mu^n(0)$ or $\mu^n(1)$ to a finite word ending in $\mu^{n+1}(0)$ or $\mu^{n+1}(1)$. The second author's thesis [13] contains a modern exposition of Fife's Theorem.

The Thue-Morse sequence \mathbf{t} is a fixed point of μ , namely,

$$\mathbf{t} = \lim_{n \to \infty} \mu^n(0)$$

Due to Theorem 1.2, \mathbf{t} arises naturally in any study of overlap-free binary words. For example, Berstel [3] proved (See also Allouche *et al.* [1]):

Theorem 1.4. The lexicographically greatest overlap-free binary ω -word starting with θ is **t**.

Our rich understanding of binary overlap-free words comes from the strong connection between these words and the Thue-Morse morphism. The thesis of the present article is that there exist analogous connections between other pairs of languages and morphisms. In a recent paper, the first author [7] showed such a connection between the period-doubling morphism $\delta = [01, 00]$ and good words. A binary word is good if it doesn't contain factors 11 or 1001, and doesn't encounter pattern 0000 or 00010100. He showed that:

- Good words factorize under δ ;
- Word $\delta(w)$ is good if and only if w is good;
- An analog of Fife's Theorem holds for good ω -words;
- One can exhibit the lexicographically least and greatest good ω -words.

Unfortunately, one may object that the period-doubling morphism does not give a proper 'new' example of a language/morphism connection because of the close relationship of the period-doubling sequence d to the Thue-Morse sequence t; it is well-known [8] that the period-doubling sequence can be obtained from the Thue-Morse sequence as follows: Let vtm be given by

$$vtm = g^{-1}(t),$$

where g is the morphism on $\{0, 1, 2\}$ given by g = [011, 01, 0]. Then

$$\boldsymbol{d} = h(\boldsymbol{vtm}),$$

where h is the morphism on $\{0, 1, 2\}$ given by h = [0, 1, 0]. For this reason, in this article we consider another morphism, not connected to μ in the same way.

Another famous binary sequence is the Fibonacci word, which is the fixed point

 $\phi = 0100101001001001001001001001001001001 \cdots$

of the binary morphism φ , where $\varphi = [01, 0]$. We call φ the **Fibonacci morphism**. The word ϕ is central to the study of Sturmian words, and has a large literature. (See [4], for example.)

For a non-empty word X, the word X^- is obtained from X by erasing its last letter. The word ^-X is obtained by erasing its first letter. We define a 4⁻-**power** (said "four minus"-power) to be a word of the form $XXXX^-$, some non-empty word X. Equivalently, a 4⁻-power is a word of period p, length 4p - 1 for some positive p. Extending periodically on the right or left with period p by a single letter gives a fourth power. Thus a 4⁻-power $XXXX^-$ can also be written as ^-YYYY , where $Y = aXa^{-1}$ and a is the last letter of X.

A binary word is called *faux-bonacci* if it contains no factor 11 and no 4^- -power. For the remainder of this paper we abbreviate 'faux-bonacci' as 'fb'. We will show that fb words bear the same relationship to the Fibonacci morphism that overlap-free words bear to the Thue-Morse morphism. We show that:

- The fb words factorize under φ (Theorem 2.3);
- Word $\varphi(w)$ is fb if and only if w is fb (Theorem 2.4);
- An analog of Fife's Theorem holds for fb ω -words (Theorem 3.10);
- One can exhibit the lexicographically least and greatest fb ω -words (Theorem 4.5).

These results raise the question of which well-studied properties of the Thue-Morse word t may generalize to the class of all morphic fixed points.

2 Faux-bonacci words

Unless otherwise specified, our words and morphisms are over the binary alphabet $\mathcal{B} = \{0, 1\}$.

Lemma 2.1. Let $u \in \mathcal{B}^{\omega}$. Suppose $\varphi(u)$ is fb. Then u is fb.

Proof. We prove the contrapositive: Suppose \boldsymbol{u} is not fb; we prove that $\varphi(\boldsymbol{u})$ is not fb.

If 11 is a factor of \boldsymbol{u} , then one of $\varphi(110) = 0001$ and $\varphi(111) = 000$ is a factor of $\varphi(\boldsymbol{u})$. Each of these contains the 4⁻-power 000.

Suppose u contains an 4⁻-power $XXXX^-$ where X = xa for some $a \in \mathcal{B}$. If a = 1, then $\varphi(u)$ contains $\varphi(x1x1x1x) = \varphi(x)0\varphi(x)0\varphi(x)0\varphi(x)$, which is an 4⁻-power. If a = 0, then u contains one of x0x0x0x0 and x0x0x0x1. In either case, $\varphi(u)$ contains the 4⁻-power $\varphi(x)01\varphi(x)01\varphi(x)01\varphi(x)0$. Remark 2.2. Let $w = w_1 w_2 w_3 \cdots w_n$ with $w_i \in \mathcal{B}$ and suppose $|w|_{11} = 0$. There is a unique word u such that $0w = \varphi(u)$ for some u. Let $w \in \mathcal{B}^{\omega}$ and suppose $|w|_{11} = 0$. Then we can write $w = a\varphi(u)$, some $u \in \mathcal{B}^{\omega}$ where $a \in \{\epsilon, 1\}$.

Theorem 2.3. Let $\boldsymbol{w} \in \mathcal{B}^{\omega}$ be fb. Then we can write $\boldsymbol{w} = a\varphi(\boldsymbol{u})$, where \boldsymbol{u} is fb and where $a \in \{\epsilon, 1\}$.

Proof. This is immediate from Lemma 2.1 and Remark 2.2.

Theorem 2.4. Let $w \in \mathcal{B}^{\omega}$. Then $\varphi(w)$ is fb if and only if w is fb.

Proof. The only if direction is Lemma 2.1. Suppose then that \boldsymbol{w} is fb. Certainly $\varphi(\boldsymbol{w})$ cannot contain 11 as a factor. Further, if 000 is a factor of $\varphi(\boldsymbol{w})$, then one of 110 and 111 is a factor of \boldsymbol{w} ; however, \boldsymbol{w} is fb, so this is impossible.

Suppose that $\varphi(\boldsymbol{w})$ has a factor XXXx where X = xa, some $a \in \mathcal{B}$.

If a = 1 then XX = x1x1. Since 11 is not a factor of $\varphi(\boldsymbol{w})$, x must be non-empty and have first letter 0. Write $X = \varphi(Y0)$. Then $XXXx = \varphi(Y0Y0Y0Y)0$, and \boldsymbol{w} contains the overlap Y0Y0Y0Y. This is a contradiction, since \boldsymbol{w} is fb.

Assume then that a = 0. If the first letter of X is 1, then the factor XXXx = x0x0x0x of $\varphi(w)$ must appear in the context 0x0x0x0x. Word x cannot be empty, since 000 is not a factor of $\varphi(w)$. If the last letter of x is 1, then replacing X by 0x reduces to the previous case. Suppose then that the last letter of x is 0. Write x = x'0. Then XXXx = x'00x'00x'00x'0. Since 000 is not a factor of $\varphi(w)$, x' is non-empty and starts and ends with 1. This implies that $\varphi(w)$ contains the factor 0XXXx' = 0x'00x'00x'00x'. Write $0x'0 = \varphi(Y1)$. Then $0x'00x'00x'00x' = \varphi(Y1Y1Y1Y)$, and the 4⁻-power Y1Y1Y1Y is a factor of w. This is impossible.

Corollary 2.5. The Fibonacci word ϕ is fb.

Lemma 2.6. Suppose that $w \in \mathcal{B}^{\omega}$ is fb. Then 10101 is not a factor of -w.

Proof. Suppose 10101 is a factor of -w. Since 11 is not a factor of w, extending 10101 to the left and right we find that 0101010 is a factor of w. But 0101010 is a 4⁻-power.

Lemma 2.7. Suppose $0101 \boldsymbol{w} \in \boldsymbol{\mathcal{B}}^{\omega}$ is fb. Then $10101 \boldsymbol{w}$ is fb.

Proof. If $10101\boldsymbol{w}$ is not fb it must begin with a 4⁻-power with some positive period p and length 4p - 1. If $p \ge 5$, then 10101 is a factor of the fb word $-0101\boldsymbol{w}$. This is impossible by Lemma 2.6, so that $p \le 4$. Thus p is a period of 10101, so that p is 2 or 4. This forces 1010101 to be a prefix of $10101\boldsymbol{w}$, and again 10101 is a factor of the fb word $-0101\boldsymbol{w}$.

3 An analogue of Fife's Theorem

Let U be the set of infinite fb words. For $u \in \mathcal{B}^*$, let $U_u = U \cap u\mathcal{B}^{\omega}$.

Lemma 3.1. Let $v \in \mathcal{B}^{\omega}$.

- (i) Word $\varphi(\boldsymbol{v}) \in U \iff \boldsymbol{v} \in U;$
- (ii) Word $1\varphi(\boldsymbol{v}) \in U \iff 0\boldsymbol{v} \in U \text{ or } \boldsymbol{v} \in U_{00}.$

Proof. Part (i) is Theorem 2.4. For part (ii), first suppose that $0\mathbf{v} \in U$ or $\mathbf{v} \in U_{00}$. If $0\mathbf{v} \in U$, then by Theorem 2.4 it follows that $01\varphi(\mathbf{v}) = \varphi(0\mathbf{v})$ is fb, so in particular $1\varphi(\mathbf{v})$ is fb; if $\mathbf{v} \in U_{00}$, then $\varphi(\mathbf{v})$ is fb by Theorem 2.4 and has prefix 0101, so that $1\varphi(\mathbf{v})$ is fb by Lemma 2.7.

In the other direction, suppose that $1\varphi(\boldsymbol{v}) \in U$. To get a contradiction, suppose that $0\boldsymbol{v} \notin U$ and $\boldsymbol{v} \notin U_{00}$. Since $1\varphi(\boldsymbol{v}) \in U$, we must have $\varphi(\boldsymbol{v}) \in U$, so that $\boldsymbol{v} \in U$ by Theorem 2.4. From $0\boldsymbol{v} \notin U$ we deduce that a prefix of $0\boldsymbol{v}$ is not fb. Since 11 cannot be a prefix of $0\boldsymbol{v}$, we deduce that $0\boldsymbol{v}$ has a prefix of the form $XXXX^-$ for some non-empty word X. It follows that 0 is a prefix of X. Since $\boldsymbol{v} \notin U_{00}$, we conclude that $|X| \geq 2$. Since XX is a factor of the fb word \boldsymbol{v} , and XX has prefix X0, we conclude that 00 is not a suffix of X; otherwise \boldsymbol{v} would have factor $000 = 0000^-$, which is impossible.

Let \boldsymbol{v} have prefix $^{-}XXXX^{-}a$, where $a \in \mathcal{B}$. Letter a cannot be the last letter of X, or \boldsymbol{v} would start with the 4⁻-power $^{-}XXXX$. It follows that X^{-} is followed in $^{-}XXXX^{-}a$ variously by a and by the other letter of \mathcal{B} . This implies that the last letter of X^{-} is 0, since 1 cannot be followed by 1 in \boldsymbol{v} . Since 00 is not a suffix of X, word X must end in 01.

We cannot, however, have X = 01. In this case, word \boldsymbol{v} would have prefix $^{-}XXXX^{-}0 = 1010100$, causing $1\varphi(\boldsymbol{v})$ to have prefix 100100100101, which begins with the 4⁻-power 10010010010. It follows that $|X| \geq 3$. Since X^{-} is sometimes followed by 0 in \boldsymbol{v} , word X^{-} cannot have suffix 00, since 000 is not a factor of \boldsymbol{v} . As the last letter of X^{-} is 0, this implies that X^{-} ends in 10. Now X ends in 101, and XX is a factor of \boldsymbol{v} . Thus Lemma 2.6 implies

that X does not start 01. It follows that X starts 00. Write X = 00Y101and $\boldsymbol{v} = XXXX^{-0}\boldsymbol{u}$. Then

$$1\varphi(\boldsymbol{v}) = 101\varphi(Y)00100101\varphi(Y)00100101\varphi(Y)00100101\varphi(Y)0010010\varphi(Y)$$

which begins with the 4⁻-power $ZZZZ^-$ where $Z = 101\varphi(Y)00100$. This is a contradiction.

Consider the finite Fibonacci words F_n defined for non-negative integers n by

$$F_0 = 0,$$

$$F_1 = 01, \text{ and}$$

$$F_{n+2} = F_{n+1}F_n \text{ for } n \ge 0$$

The following lemma is proved by induction.

Lemma 3.2. Suppose that u is a binary word and p is a prefix of $\varphi(u)$. If p ends in F_{n+1} , some $n \ge 2$, then $p = \varphi(q)$, where q is a prefix of u ending in F_n .

Remark 3.3. The condition $n \ge 2$ is necessary. If u = 00, then F_2 is a factor of $\varphi(u)$, but F_1 is not a factor of u.

We use the notation $\pi(w)$ for the *Parikh vector* of a binary word. Thus

$$\pi(w) = [|w|_0, |w|_1].$$

Suppose that w is a word of the form $y_n F_n$ where $\pi(y_n) \leq \pi(F_n) - \pi(0)$. We define operations on w by

$$\alpha(w) = y_n F_{n+1}$$

$$\beta(w) = y_n F_{n-1} F_{n+1}$$

One checks that F_n is a prefix of $F_{n-1}F_{n+1}$, so that w is always a prefix of $\alpha(w)$ and $\beta(w)$. Because $\pi(y_n) \leq \pi(F_n) - \pi(0)$, we have that $\pi(y_nF_{n-1}) \leq \pi(F_nF_{n-1}) - \pi(0) = \pi(F_{n+1}) - \pi(0)$, so that we can iterate the maps α and β . Let $\mathcal{O} = \{\alpha, \beta\}$. We define operators $w \bullet f$ for $f \in \mathcal{O}^*$ by

$$w \bullet \epsilon = w$$
$$w \bullet (f\gamma) = (w \bullet f) \bullet \gamma \text{ for } \gamma \in \mathcal{O}.$$

Remark 3.4. If g is a prefix of f, then $w \bullet g$ will be a prefix of $w \bullet f$.

Lemma 3.5. Let \boldsymbol{u} be a fb ω -word, and for $n \geq 2$, let $w_n = y_n F_n$ be the shortest prefix of \boldsymbol{u} ending in F_n . For $n \geq 2$, we have $\pi(y_n) \leq \pi(F_n) - \pi(0)$.

Proof. Word w_2 is a fb word containing $F_2 = 010$ exactly once, as a suffix. The candidates are 010, 0010, 1010, and 10010, which would yield $y_2 = \epsilon$, 0, 1, and 10, respectively. Thus the result is true for n = 2.

Suppose the result has been found to be true for n = k, some $k \ge 2$. Write $\boldsymbol{u} = a\varphi(\boldsymbol{u}')$ where \boldsymbol{u}' is fb, and $a \in \{\epsilon, 1\}$. Since the first letter of F_{k+1} is 0 and $a \in \{\epsilon, 1\}$, any occurrence of F_{k+1} in \boldsymbol{u} starts in $\varphi(\boldsymbol{u}')$. By Lemma 3.2, any prefix qF_{k+1} of \boldsymbol{u} has the form $a\varphi(q'F_k)$ where $q'F_k$ is a prefix of \boldsymbol{u}' . Thus $w_{k+1} = a\varphi(w'_k)$ where $w'_k = y'_k F_k$ is the shortest prefix of \boldsymbol{u}' ending in F_k . By the induction hypothesis, $\pi(y'_k) \le \pi(F_k) - \pi(0)$. It follows that

$$\pi(a\varphi(y'_k)) \le \pi(1) + \pi(\varphi(F_k)) - \pi(\varphi(0)) = \pi(1) + \pi(F_{k+1}) - \pi(01) = \pi(F_{k+1}) - \pi(0).$$

Remark 3.6. Again, the condition $n \ge 2$ is necessary. If $w_2 = 1001$, then $y_2 = 10$, and $\pi(y_2) \le \pi(F_2) - \pi(0)$.

Lemma 3.7. Let u be a fb ω -word, and for $n \ge 2$, let $w_n = y_n F_n$ be the shortest prefix of u ending in F_n . For $n \ge 2$, we have $w_{n+1} \in \{w_n \bullet \alpha, w_n \bullet \beta\}$.

Proof. The only fb words starting with F_2 , and containing F_3 exactly once, as a suffix, are $01001 = 010 \bullet \alpha$ and $0101001 = 010 \bullet \beta$. Thus the result is true for n = 2.

Suppose the result has been found to be true for n = k, some $k \ge 2$. Write $\boldsymbol{u} = a\varphi(\boldsymbol{u}')$ where \boldsymbol{u}' is fb, and $a \in \{\epsilon, 1\}$. For each n, let $w'_n = y'_n F_n$ be the shortest prefix of \boldsymbol{u}' ending in F_n . By the induction hypothesis, w'_{k+1} is either $w'_k \bullet \alpha = y'_k F_{k+1}$ or $w'_k \bullet \beta = y'_k F_{k-1} F_{k+1}$. As in the previous proof, $w_{k+2} = a\varphi(w'_{k+1})$ and $y_{k+2} = a\varphi(y'_{k+1})$. Then

$$w_{k+2} = a\varphi(w'_{k+1}) \\ \in \{a\varphi(y'_kF_{k+1}), a\varphi(y'_kF_{k-1}F_{k+1})\} \\ = \{y_{k+1}F_{k+2}, y_{k+1}F_kF_{k+2}\} \\ = \{w_{k+1} \bullet \alpha, w_{k+1} \bullet \beta\}.$$

Corollary 3.8. Let \boldsymbol{u} be a fb ω -word. There is an ω -word $\boldsymbol{f} = \prod_{n=2}^{\infty} f_n$, $f_n \in \mathcal{O}$, and a 'seed' word $w_2 \in \{010, 0010, 1010, 10010\}$, such that

$$\boldsymbol{u} = \lim_{n \to \infty} w_2 \bullet (f_2 f_3 \cdots f_n).$$

Proof. Letting the w_n be as in the previous lemma, choose f_n such that $w_{n+1} = w_n \bullet f_n$.

Suppose that w is a word of the form $y_n F_n$ where $\pi(y_n) \leq \pi(F_n) - \pi(0)$. We note that

$$w \bullet \alpha = wF_n^{-1}F_{n+1} = wF_n^{-1}\varphi^{n-2}(F_2 \bullet \alpha), w \bullet \beta = wF_n^{-1}F_{n-1}F_{n+1} = wF_n^{-1}\varphi^{n-2}(F_2 \bullet \beta).$$

If $f \in \mathcal{O}^k$, write $f = f_1 f_2 \cdots f_k$, where each $f_i \in \mathcal{O}$. Suppose that w is a word of the form $y_2 F_2$ where $\pi(y_2) \leq \pi(F_2) - \pi(0)$. By induction, $w \bullet f$ is a word of the form $y_{k+2}F_{k+2}$ where $\pi(y_{k+2}) \leq \pi(F_{k+2}) - \pi(0)$. We find that

$$w \bullet f$$

= $(w \bullet f_1 f_2 \cdots f_{k-1}) \bullet f_k$
= $(w \bullet f_1 f_2 \cdots f_{k-1}) F_{k+1}^{-1} \varphi^{k-1} (F_2 \bullet f_k)$
= $(w \bullet f_1 f_2 \cdots f_{k-2}) F_k^{-1} \varphi^{k-2} (F_2 \bullet f_{k-1}) F_{k+1}^{-1} \varphi^{k-1} (F_2 \bullet f_k)$
:
= $w \prod_{j=1}^k F_{j+1}^{-1} \varphi^{j-1} (F_2 \bullet f_j)$ (1)

Let $\mathbf{f} \in \mathcal{O}^{\omega}$, $\mathbf{f} = f_1 f_2 f_3 \cdots$, where each $f_i \in \mathcal{O}$. For $w \in \{010, 0010, 1010, 10010\}$, define

$$w \bullet f = \lim_{n \to \infty} w \bullet (f_1 f_2 f_3 \cdots f_n).$$

Lemma 3.9. Suppose $w \in \{010, 0010, 1010, 10010\}$ and $\mathbf{f} \in \mathcal{O}^{\omega}$ and $g \in \mathcal{O}^k$. Suppose $F_2 \bullet \mathbf{f} = \mathbf{x}$. Then

$$w \bullet (gf) = (w \bullet g) F_{k+2}^{-1} \varphi^k(\boldsymbol{x}).$$
⁽²⁾

Proof. Write $\mathbf{f} = f_1 f_2 f_3 \cdots$ and $g = g_1 g_2 \cdots g_k$ where the $f_i, g_i \in \mathcal{O}$. Suppose that $F_2 \bullet f_1 \cdots f_n = x_n$. We will show that $w \bullet (g_1 g_2 \cdots g_k f_1 f_2 \cdots f_n) = (w \bullet g) F_{k+2}^{-1} \varphi^k(x_n)$, and the result follows by taking limits. From (1),

$$x_n = F_2 \bullet f_1 \cdots f_n = F_2 \prod_{j=1}^n F_{j+1}^{-1} \varphi^{j-1} (F_2 \bullet f_j)$$

and

$$\begin{split} & w \bullet (g_1 g_2 \cdots g_k f_1 f_2 \cdots f_n) \\ &= w \Pi_{j=1}^k F_{j+1}^{-1} \varphi^{j-1} (F_2 \bullet g_j) \Pi_{j=1}^n F_{k+j+1}^{-1} \varphi^{k+j-1} (F_2 \bullet f_j) \\ &= (w \bullet g) \Pi_{j=1}^n \varphi^k (F_{j+1})^{-1} \varphi^k (\varphi^{j-1} (F_2 \bullet f_j)) \\ &= (w \bullet g) \varphi^k (F_2^{-1} F_2 \Pi_{j=1}^n F_{j+1}^{-1} \varphi^{j-1} (F_2 \bullet f_j)) \\ &= (w \bullet g) \varphi^k (F_2)^{-1} \varphi^k (F_2 \Pi_{j=1}^n F_{j+1}^{-1} \varphi^{j-1} (F_2 \bullet f_j))) \\ &= (w \bullet g) F_{k+2}^{-1} \varphi^k (x_n), \end{split}$$

as desired.

Define sets F and V by

$$F = \alpha^* \beta (\alpha \alpha + \alpha \beta + \beta \alpha \alpha + \beta \beta \beta^* \alpha \alpha)^* (\beta \alpha \beta + \beta \beta \beta^* \alpha \beta),$$
$$V = \mathcal{O}^{\omega} - \mathcal{O}^* F \mathcal{O}^{\omega}.$$

Theorem 3.10. Let $x \in \mathcal{B}^{\omega}$. If x begins with 010, then x is fb if and only if $x = 010 \bullet f$ for some $f \in V$.

The set F consists of forbidden factors for words of V.

Let $W = \{ \boldsymbol{f} \in \mathcal{O}^{\omega} : 010 \bullet \boldsymbol{f} \in U \}$. To prove Theorem 3.10 it is enough to prove that W = V. Let $L \subseteq \mathcal{O}^{\omega}$ and let $x \in \mathcal{O}^*$. We define the (left) quotient $x^{-1}L$ by $x^{-1}L = \{ \boldsymbol{y} \in \mathcal{O}^{\omega} : x\boldsymbol{y} \in L \}$. The next lemma establishes several identities concerning quotients of the set W. They are proved using (2) and Lemma 3.1. The identities demonstrate that W is precisely the set of infinite labeled paths through the automaton A_{010} given in Figure 1. These are just the labeled paths omitting factors in F, so that W = V. Thus, proving Lemma 3.11 establishes Theorem 3.10.

Lemma 3.11. The following identities hold:

(a) $W = \alpha^{-1}W;$



Figure 1: 'Fife' automaton A_{010} for U.

- (b) $\beta^{-1}W = (\beta\alpha\alpha)^{-1}W = (\beta\alpha\beta)^{-1}W^{-1} = (\beta\beta\alpha\alpha)^{-1}W;$
- (c) $(\beta\beta\alpha)^{-1}W = (\beta\beta\beta\alpha)^{-1}W;$
- (d) $(\beta\beta\beta)^{-1}W = (\beta\beta\beta\beta)^{-1}W;$
- (e) $(\beta\beta\alpha\beta)^{-1}W = \emptyset.$

Each set of identities corresponds to the state of A_{010} with the same label as the identities. There are two further states: (f), corresponding to $(\beta\alpha)^{-1}W$, and (g), corresponding to $(\beta\beta)^{-1}W$ The non-accepting sink (e) is not shown in the figure. The automaton A_{010} is not minimal; for example, (d)can be identified with (g). However, this form highlights parallels between it and the three automata we present later. *Proof.* (a) We have

$$\alpha \boldsymbol{f} \in W$$

$$\iff 010 \bullet \alpha \boldsymbol{f} \in U$$

$$\iff (010 \bullet \alpha)\varphi(F_2)^{-1}\varphi(\boldsymbol{x}) \in U \text{ by } (2)$$

$$\iff 01001(01001)^{-1}\varphi(\boldsymbol{x}) \in U$$

$$\iff \boldsymbol{x} \in U \text{ by Lemma 3.1 (i)}$$

$$\iff 010 \bullet \boldsymbol{f} \in U$$

$$\iff \boldsymbol{f} \in W,$$

so that $\alpha^{-1}W = W$.

(b) Here

$$\beta \boldsymbol{f} \in W$$

$$\iff 010 \bullet \beta \boldsymbol{f} \in U$$

$$\iff (010 \bullet \beta)\varphi(F_2)^{-1}\varphi(\boldsymbol{x}) \in U$$

$$\iff 01 \ \frac{01001(01001)}{}^{-1}\varphi(\boldsymbol{x}) \in U$$

$$\iff \varphi(0\boldsymbol{x}) \in U$$

$$\iff 0\boldsymbol{x} \in U \text{ by Lemma 3.1 (i).}$$

Similarly we find that

$$\beta \alpha \alpha \boldsymbol{f} \in W$$

$$\iff 010 \bullet \beta \alpha \alpha \boldsymbol{f} \in U$$

$$\iff (010 \bullet \beta \alpha \alpha) \varphi^3 (F_2)^{-1} \varphi^3 (\boldsymbol{x}) \in U$$

$$\iff 01 \ \frac{0100101001001(01001001001)^{-1}}{\varphi^3} (\boldsymbol{x}) \in U$$

$$\iff \varphi (0\varphi^2(\boldsymbol{x})) \in U$$

$$\iff 0\varphi^2(\boldsymbol{x}) \in U \text{ by Lemma 3.1 (i)}$$

$$\iff \varphi (1\varphi(\boldsymbol{x})) \in U$$

$$\iff 1\varphi(\boldsymbol{x}) \in U \text{ by Lemma 3.1 (i)}$$

$$\iff 0\boldsymbol{x} \in U \text{ or } \boldsymbol{x} \in U_{00} \text{ by Lemma 3.1 (ii)}$$

$$\iff 0\boldsymbol{x} \in U.$$

Here we use the fact that 01 is a prefix of \boldsymbol{x} , so that $\boldsymbol{x} \notin U_{00}$.

Again,

$$\beta \alpha \beta \boldsymbol{f} \in W$$

$$\iff 010 \bullet \beta \alpha \beta \boldsymbol{f} \in U$$

$$\iff (010 \bullet \beta \alpha \beta) \varphi^{3}(F_{2})^{-1} \varphi^{3}(\boldsymbol{x}) \in U$$

$$\iff 0101001 \frac{0100101001001(01001001)^{-1}}{\varphi^{3}(\boldsymbol{x})} \in U$$

$$\iff \varphi(0010\varphi^{2}(\boldsymbol{x})) \in U$$

$$\iff \varphi(101\varphi(\boldsymbol{x})) \in U$$

$$\iff 101\varphi(\boldsymbol{x}) \in U$$

$$\iff 10\varphi(0\boldsymbol{x}) \in U$$

$$\iff 00\boldsymbol{x} \in U \text{ or } 0\boldsymbol{x} \in U_{00}$$

$$\iff 0\boldsymbol{x} \in U.$$

Here we use the fact that 0 is a prefix of \boldsymbol{x} , so that 00 is a prefix of $0\boldsymbol{x}$. Finally we get

Thus $\beta^{-1}W = (\beta\alpha\alpha)^{-1}W = (\beta\alpha\beta)^{-1}W = (\beta\beta\alpha\alpha)^{-1}W$, as desired.

(c) Here

$$\beta \beta \alpha \boldsymbol{f} \in W$$

$$\iff 010 \bullet \beta \beta \alpha \boldsymbol{f} \in U$$

$$\iff (010 \bullet \beta \beta \alpha) \varphi^{3}(F_{2})^{-1} \varphi^{3}(\boldsymbol{x}) \in U$$

$$\iff 01010 \ \frac{0100101001001(01001001001)^{-1}}{\varphi^{3}(\boldsymbol{x})} \in U$$

$$\iff \varphi(001\varphi^{2}(\boldsymbol{x})) \in U$$

$$\iff 001\varphi^{2}(\boldsymbol{x}) \in U$$

$$\iff 01\varphi(\boldsymbol{x}) \in U$$

$$\iff 01\boldsymbol{x} \in U \text{ or } 1\boldsymbol{x} \in U_{00}$$

$$\iff 01\boldsymbol{x} \in U.$$

Similarly,

Thus $(\beta\beta\alpha)^{-1}W = (\beta\beta\beta\alpha)^{-1}W$, as desired.

(d) We have

$$\beta\beta\beta \boldsymbol{f} \in W$$

$$\iff 010 \bullet \beta\beta\beta \boldsymbol{f} \in U$$

$$\iff (010 \bullet \beta\beta\beta)\varphi^{3}(F_{2})^{-1}\varphi^{3}(\boldsymbol{x}) \in U$$

$$\iff 0101001001 \frac{0100101001(01001001001)^{-1}}{\varphi^{3}(\boldsymbol{x})} \in U$$

$$\iff \varphi(001010\varphi^{2}(\boldsymbol{x})) \in U$$

$$\iff 001010\varphi^{2}(\boldsymbol{x}) \in U$$

$$\iff 00101\varphi^{2}(\boldsymbol{x}) \in U$$

$$\iff 010\boldsymbol{x} \in U \text{ or } 10\boldsymbol{x} \in U_{00}$$

$$\iff 010\boldsymbol{x} \in U.$$

Similarly,

Thus $(\beta\beta\beta)^{-1}W = (\beta\beta\beta\beta)^{-1}W$, as desired.

(e) We have

However, \boldsymbol{x} has prefix 0, so $00\boldsymbol{x}$ has the 4⁻-power 000 as a prefix. Therefore, $00\boldsymbol{x} \notin U$. It follows that $(\beta\beta\alpha\beta)^{-1}W = \emptyset$.

Remark 3.12. We mention without proof that

$$\beta \alpha \bullet f \in W \iff 1f \in V,$$

and

$$\beta\beta \bullet f \in W \iff 10f \in V.$$

We do not need these equivalences to formulate the automaton.

Remark 3.13. As Fife's Theorem features a forbidden factor characterization, we have given such a characterization for V. In fact, however, all information about V is captured in the automaton A_{010} . For a finite string $g \in \mathcal{O}^*$, word $010 \bullet g$ is fb exactly when g can be walked on the automaton A_{010} ; such a string g never encounters the (undepicted) non-acepting sink (e), which can only be reached via (c) on input β . If $\mathbf{f} \in \mathcal{O}^{\omega}$, word $010 \bullet \mathbf{f}$ is fb exactly when $010 \bullet f$ is fb for finite prefix f of \mathbf{f} .

It is routine to write down an expression for the regular language of finite words arriving at the sink, which is $F\mathcal{O}^*$. It happens F will take us from



Figure 2: 'Fife' automaton A_{0010} for W_{0010} .

any given state to the sink. For this reason, $010 \bullet g$ is fb exactly when g has no *factor* in F.

For $u \in \{010, 0010, 1010, 10010\}$, let $W_u = \{\mathbf{f} \in \mathcal{O}^{\omega} : u \bullet \mathbf{f} \in U\}$. Using the same method as in Lemma 3.11, one shows that W_u is the subset of \mathcal{O}^{ω} which can be walked on A_u , where the additional automata are depicted in Figures 2, 3, and 4.

4 Lexicographically extremal fb words

The lexicographic order on binary words is given recursively by

 $u < v \iff v \neq \epsilon \text{ and } ((u = \epsilon) \text{ or } (u^- < v^-) \text{ or } ((u = u^- 0) \text{ and } (v = u^- 1))).$

Note that the morphism φ is order-reversing: Let u and v be non-empty binary words so that u < v. Write u = u'0u'', v = u'1v'' where u' is the



Figure 3: 'Fife' automaton A_{1010} for U_{1010} .



Figure 4: 'Fife' automaton A_{10010} for U_{10010} .

longest common prefix of u and v. Then $\varphi(u')01$ is a prefix of $\varphi(u)$, while $\varphi(u')00$ is a prefix of $\varphi(v)$, so that $\varphi(u) > \varphi(v)$.

For each non-negative integer n, let ℓ_n (resp., m_n) be the lexicographically least (resp., greatest) word of length n such that ℓ_n (resp., m_n) is the prefix of an fb ω -word.

Lemma 4.1. Let n be a non-negative integer. Word ℓ_n is a prefix of ℓ_{n+1} . Word m_n is a prefix of m_{n+1} .

Proof. We prove the result for the ℓ_n ; the proof for the m_n is similar. Let $\ell_n \mathbf{r}$ be an fb ω -word. Let p be the length n + 1 prefix of $\ell_n \mathbf{r}$, and let q be the length n prefix of ℓ_{n+1} . We need to show that $q = \ell_n$. Both p and q are prefixes of fb ω -words. By definition we have $\ell_{n+1} \leq p$ and $\ell_n \leq q$. If $\ell_n < q$, then $p^- = \ell_n < q = \ell_{n+1}^-$, so that $p < \ell_{n+1}$. This is a contradiction. Therefore $\ell_n = q$, as desired.

Let $\boldsymbol{\ell} = \lim_{n \to \infty} \ell_n$, $\mathbf{m} = \lim_{n \to \infty} m_n$.

Lemma 4.2. Word ℓ is the lexicographically least fb ω -word. Word **m** is the lexicographically greatest fb ω -word.

Proof. We show that ℓ is lexicographically least. The proof that **m** is lexicographically greatest is similar. Let **w** be an fb ω -word. For each n let w_n be the length n prefix of **w**, so that $\mathbf{w} = \lim_{n \to \infty} w_n$.

If for some n we have $w_n > \ell_n$, then $\mathbf{w} > \boldsymbol{\ell}$.

Otherwise $w_n \leq \ell_n$ for all n. By the definition of the ℓ_n we have $w_n \geq \ell_n$, so that $w_n = \ell_n$ for all n. Thus $\mathbf{w} = \lim_{n \to \infty} w_n = \lim_{n \to \infty} \ell_n = \boldsymbol{\ell}$.

In all cases we find $\mathbf{w} \geq \boldsymbol{\ell}$.

Lemma 4.3. We have $\ell = \varphi(\mathbf{m})$.

Proof. Since the Fibonacci word ϕ has suffixes beginning with 00, $\ell_2 = 00$, and we can write $\boldsymbol{\ell} = \varphi(\mathbf{m}')$ for some \mathbf{m}' by Theorem 2.3. Since $\boldsymbol{\ell}$ is fb, \mathbf{m}' is fb by Lemma 2.1. It follows that $\mathbf{m}' \leq \mathbf{m}$. However if $\mathbf{m}' < \mathbf{m}$ then $\varphi(\mathbf{m}) < \varphi(\mathbf{m}') = \boldsymbol{\ell}$ since φ is order-reversing. This is impossible, since $\boldsymbol{\ell}$ is least. Therefore $\mathbf{m}' = \mathbf{m}$, and $\boldsymbol{\ell} = \varphi(\mathbf{m})$.

Lemma 4.4. We have $\mathbf{m} = 1\varphi(\boldsymbol{\ell})$.

Proof. Since 00 is a prefix of $\boldsymbol{\ell}$ we see that 0101 is a prefix of $\varphi(\boldsymbol{\ell})$. It follows from Lemma 2.7 that $1\varphi(\boldsymbol{\ell})$ is fb. Since 10101 is the lexicographically greatest fb word of length 5, $m_5 = 10101$. It follows that we can write $\boldsymbol{m} = 1\varphi(\boldsymbol{\ell}')$ for some fb word $\boldsymbol{\ell}'$. However, φ is order reversing, so that if $\boldsymbol{\ell}' > \boldsymbol{\ell}$, then $1\varphi(\boldsymbol{\ell}) > 1\varphi(\boldsymbol{\ell}') = \boldsymbol{m}$, contradicting the maximality of \boldsymbol{m} . Thus $\boldsymbol{\ell}' = \boldsymbol{\ell}$ and $\mathbf{m} = 1\varphi(\boldsymbol{\ell})$.

Theorem 4.5. Word *m* satisfies

$$\boldsymbol{m} = 1\varphi^2(\boldsymbol{m}). \tag{3}$$

Word ℓ satisfies

$$\boldsymbol{\ell} = 0\varphi^2(\boldsymbol{\ell}). \tag{4}$$

Proof. This follows from Lemma 4.3 and Lemma 4.4.

Lemma 4.6. Neither of \boldsymbol{m} and $\boldsymbol{\ell}$ is the fixed point of a binary morphism. Every factor of $\boldsymbol{\phi}$ is a factor of \boldsymbol{m} and $\boldsymbol{\ell}$, but there are infinitely many factors of \boldsymbol{m} (resp., $\boldsymbol{\ell}$) which are not factors of $\boldsymbol{\phi}$ or $\boldsymbol{\ell}$ (resp., \boldsymbol{m}).

Proof. Word \boldsymbol{m} has prefix 10101, but by Lemma 2.6, the word 10101 is not a factor of $-\boldsymbol{m}$. It follows that \boldsymbol{m} cannot be the fixed point of a binary

(1)

morphism. Similarly, $\varphi(10101)$ is a prefix of ℓ , but not a factor of $-\ell$, so that ℓ is not a fixed point of a binary morphism.

Every factor of ϕ is a factor of $\varphi^{2k}(0)$ for some k, and is therefore a factor of

$$\boldsymbol{m} = 1\varphi^2(\boldsymbol{m}) = 1\varphi^2(1\varphi^2(\boldsymbol{m})) = \cdots = 1\varphi^2(1)\varphi^4(1)\cdots\varphi^{2k-2}(1)\varphi^{2k}(\boldsymbol{m})$$

Similarly, every factor of $\boldsymbol{\phi}$ is a factor of $\boldsymbol{\ell}$. However, none of factors $\varphi^{2k}(10101)$ of \boldsymbol{m} (resp., $\varphi^{2k+1}(10101)$ of $\boldsymbol{\ell}$) is a factor of $\boldsymbol{\phi}$ or $\boldsymbol{\ell}$ (resp. \boldsymbol{m}). \Box

5 Acknowledgment

The work of James D. Currie is supported by the Natural Sciences and Engineering Research Council of Canada (NSERC), [funding reference number 2017-03901]. The work of Narad Rampersad is supported by the Natural Sciences and Engineering Research Council of Canada (NSERC), [funding reference number 2019-04111].

References

- J.-P. Allouche, J. D. Currie, J. Shallit, Extremal infinite overlap-free binary words, *Electron. J. Combin.* 5 (1998), #R27.
- [2] J.-P. Allouche and J. Shallit, The ubiquitous Prouhet-Thue-Morse sequence, in C. Ding, T. Helleseth, H. Niederreiter, eds., Sequences and their Applications, Proceedings of SETA '98, Springer-Verlag, 1998, pp. 1–16.
- [3] J. Berstel, A rewriting of Fife's theorem about overlap-free words, in J. Karhumäki, H. Maurer, and G. Rozenberg, eds., *Results and Trends* in *Theoretical Computer Science*, Lect. Notes in Comp. Sci., Vol. 812, Springer-Verlag, 1994, pp. 19–29.
- [4] J. Berstel, Fibonacci words a survey, in G. Rozenberg, A. Salomaa, eds. The Book of L, Springer-Verlag, 1986, pp. 13–27.
- [5] A. Carpi, Overlap-free words and finite automata, *Theoret. Comput. Sci.* 115 (1993), 243–260.

- [6] J. Cassaigne, Counting overlap-free binary words, in P. Enjalbert, A. Finkel, and K. Wagner, eds., STACS 93, Proc. 10th Symp. Theoretical Aspects of Comp. Sci., Lect. Notes in Comp. Sci., Vol. 665, Springer-Verlag, 1993, pp. 216–225.
- [7] J. D. Currie, The analog of overlap-freeness for the period-doubling sequence, J. Int. Seq. 26 (2023), 23.8.2
- [8] D. Damanik, Local symmetries in the period-doubling sequence, *Discrete Appl. Math.* **100** (2000), 115–121.
- [9] E. Fife, Binary sequences which contain no *BBb*, *Trans. Amer. Math.* Soc. **261** (1980), 115–136.
- [10] N. Guglielmi, V. Protasov, Exact Computation of Joint Spectral Characteristics of Linear Operators. Found. Comput. Math. 13 (2013), 37–97.
- [11] R. M. Jungers, V. Y. Protasov, and V. D. Blondel, Overlap-free words and spectra of matrices. em Theoret. Comput. Sci. 410 (38-40), 3670– 3684.
- [12] Y. Kobayashi, Enumeration of irreducible binary words. Discrete Appl. Math. 20 (1988), 221–232.
- [13] N. Rampersad, Overlap-Free Words and Generalizations, Ph.D. thesis, University of Waterloo, 2007.
- [14] A. Restivo and S. Salemi, Overlap-free words on two symbols, in M. Nivat and D. Perrin, eds., Automata on Infinite Words, Vol. 192 of Lect. Notes in Comp. Sci., Springer-Verlag, 1984, pp. 198–206.
- [15] A. Thue, Uber die gegenseitige Lage gleicher Teile gewisser Zeichenreihen. Kra. Vidensk. Selsk. Skrifter. I. Mat. Nat. Kl. 1 (1912), pp. 1–67. Reprinted in Selected Mathematical Papers of Axel Thue, T. Nagell et al., editors, Universitetsforlaget, Oslo, 1977, pp. 413–477.