

Robust Multi-Model Subset Selection

Anthony-Alexander Christidis
Department of Statistics
University of British Columbia
anthony.christidis@stat.ubc.ca

Gabriela Cohen-Freue
Department of Statistics
University of British Columbia
gcohen@stat.ubc.ca

Abstract

Outlying observations can be challenging to handle and adversely affect subsequent analyses, particularly, in complex high-dimensional datasets. Although outliers are not always undesired anomalies in the data and may possess valuable insights, only methods that are robust to outliers are able to accurately identify them and resist their influence. In this paper, we propose a method that generates an ensemble of sparse and diverse predictive models that are resistant to outliers. We show that the ensembles generally outperform single-model sparse and robust methods in high-dimensional prediction tasks. Cross-validation is used to tune model parameters to control levels of sparsity, diversity and resistance to outliers. We establish the finite-sample breakdown point of the ensembles and the models that comprise them, and we develop a tailored computing algorithm to learn the ensembles by leveraging recent developments in ℓ_0 optimization. Our extensive numerical experiments on synthetic and artificially contaminated real datasets from bioinformatics and cheminformatics demonstrate the competitive advantage of our method over state-of-the-art single-model methods.

Keywords: Robust methods; High-dimensional data; Ensemble methods; Multi-model optimization.

1 Introduction

The rapid growth of digital technologies has led to an explosive increase in data, revolutionizing the approaches to modelling and predicting real-world phenomena. For example, advances in genomics allow the simultaneous quantitation of thousands of genes from a patient’s sample, which can be used to predict pathogenic processes or responses to therapies (e.g., [Byron et al., 2016](#)). Large volumes of data of different types, formats, and structures can be rapidly collected, generated and integrated. Thus, modern datasets are often characterized by the presence of a large number of variables (in columns), of which some are irrelevant or redundant, and in general exceeds the number of observations (in rows). Along with more sophisticated processes of obtaining data comes the inclusion of outlying observations, also known as data contamination. In this article, an outlier refers to a case (or row) in the dataset with observed values differing from those of the bulk of the data. These outliers may arise due to errors in data collection, sensitive detection of rare cases, discrepancies in data sources, or data corruption, among other reasons. Although outliers may contain valuable information, only robust methods resistant to outliers can flag or handle them appropriately. In this work, we leverage modern computational tools in ℓ_0 -optimization to generate a collection of sparse and diverse predictive models that are resistant to outlying rows in the dataset, which can be ensembled to improve the performance of single-model sparse and robust methods.

Regularized regression methods have been developed to model datasets with many predictors relative to the number of samples, enabling the selection of an optimal subset of predictors for building interpretable predictive models. Prominent examples include the least absolute shrinkage and selection operator (LASSO, [Tibshirani, 1996](#)) and the smoothly clipped absolute deviation (SCAD, [Fan and Li, 2001](#)) methods. The empirical performance and theoretical properties of these methods have been studied extensively (e.g., [Hastie et al., 2019](#)). However, depending on the loss function used, these methods may be very sensitive to outliers, which may adversely affect their variable selection and prediction performances. To address this problem, robust statistical procedures are usually derived using loss functions that downweight the effect of outlying observations ([Maronna et al., 2019](#)). In recent years, there have been many proposals that combine regularized and robust methods to obtain predictive models that are resistant to outlying rows with different characteristics ([Maronna, 2011](#); [Alfons et al., 2013](#); [Smucler and Yohai, 2017](#); [Cohen Freue et al., 2019](#)). Many of these methods have a breakdown point of 50%, meaning that they can provide a robust prediction under the assumption that the majority of the cases is not contaminated with

outlying observations.

Ensemble methods can be used to generate and aggregate multiple diverse models, and often outperform single-model methods in high-dimensional prediction tasks. Traditionally, ensemble methods rely on randomization or some form of heuristics to generate diverse models, and are thus considered “blackbox” methods. Some notable examples of these ensemble methods include random forests (RF) (Breiman, 2001), random generalized linear models (RGLM) (Song et al., 2013), gradient boosting (Friedman, 2001) and all its variations (e.g., Bühlmann and Yu, 2003; Chen and Guestrin, 2016). In general, these types of ensemble methods generate a large number of uninterpretable and inaccurate models that are only useful when they are pooled together. More recently, Christidis et al. (2020) and Christidis et al. (2024) proposed methods that generate ensembles comprised of a small number of sparse and diverse models learned directly from the data without any form of randomization or heuristics. Each of the models in these ensembles have a high prediction accuracy similar to that achieved by many single-model sparse methods, and the ensembling of these small models have been shown to outperform state-of-the-art blackbox ensemble methods on synthetic as well as complex biological and chemical data. However, both the ensembles and the individual models that comprise them are not robust and are thus very sensitive to outliers.

In this article, we introduce a robust multi-model subset selection (RMSS) method to generate ensembles comprised of a small number of sparse, robust and diverse models in a regression setting. The levels of sparsity, diversity and robustness within each model are driven directly by the data. We establish the finite-sample breakdown point of these robust ensembles and the individual models that comprise them. To fit these models, we harness recent developments in ℓ_0 -optimization to develop a tailored computational algorithm with attractive convergence properties. RMSS is shown to outperform state-of-the-art sparse and robust methods in an extensive simulation study and in biological and chemical datasets with simulated outlying cases. To the best of our knowledge, this is the first robust ensemble method proposed in the literature. In addition, the flexibility offered by our method can be particularly appealing for practitioners collecting and analyzing high-dimensional complex data.

The remainder of this article is organized as follows. In section 2, we provide a literature review. In section 3, we introduce RMSS and some of its special cases. In section 4, we study robustness properties of RMSS. In section 5, we provide a computational algorithm to generate RMSS and establish some of its convergence properties. In section 6, we present a large simulation study. In section 7, we apply RMSS on artificially contaminated datasets from bioinformatics and

cheminformatics. Concluding remarks are given in section 8.

2 Literature Review

In this section, we review a variety of predictive methods proposed in the literature that are related to RMSS and introduce important notation.

We consider the usual linear regression setting where a dataset comprised of n observations and p predictor variables can be used to build a predictive model for a response variable of interest. Let $\mathbf{y} = (y_1, \dots, y_n)^T \in \mathbb{R}^n$, $\mathbf{X} \in \mathbb{R}^{n \times p}$, and $\mathbf{x}_i = (x_{i1}, \dots, x_{ip})^T$ be the rows of \mathbf{X} for $1 \leq i \leq n$. We assume a standard linear model

$$y_i = \mu + \mathbf{x}_i^T \boldsymbol{\beta}_0 + \sigma \varepsilon_i, \quad 1 \leq i \leq n, \quad (1)$$

where $\mu \in \mathbb{R}$ and $\boldsymbol{\beta}_0 \in \mathbb{R}^p$ are the regression coefficients, and the elements of the noise vector $\boldsymbol{\varepsilon} = (\varepsilon_1, \dots, \varepsilon_n)^T \in \mathbb{R}^n$ are independent and identically distributed with mean zero and variance one. We focus our attention on the high-dimensional setting ($p \gg n$) where the underlying model is sparse, i.e., the number of nonzero elements of the true coefficient vector $\|\boldsymbol{\beta}_0\|_0 \ll p$.

2.1 Single-Model Methods

Several regression methods were proposed to generate sparse predictive models based on only a subset of the predictor variables, particularly needed when p is very large compared to the number of observations n . The Best Subset Selection (BSS) estimator proposed by [Garside \(1965\)](#) was one of the first variable selection method, which can be defined as the solution to the non-convex (and non-differentiable) minimization problem given by

$$\min_{\boldsymbol{\beta} \in \mathbb{R}^p} \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|_2^2 \quad \text{subject to} \quad \|\boldsymbol{\beta}\|_0 \leq t, \quad (2)$$

where $t \leq \min(n - 1, p)$ is the number of nonzero coefficients, which may be chosen by a model selection criterion (see e.g., [Mallows, 1973](#); [Akaike, 1974](#)) or by cross-validation (CV).

Since the BSS optimization problem (2) is an NP-hard problem ([Welch, 1982](#)), many sparse regularization methods in the form of convex relaxations of (2) were proposed, such as LASSO, Elastic Net (EN, [Zou and Hastie, 2005](#)), and SCAD ([Fan and Li, 2001](#)) methods. Although convex relaxations have much lower computational cost, BSS enjoys better estimation and variable selection properties compared to sparse regularization methods ([Shen et al., 2013](#)), and often out-

performs regularization methods in high-dimensional prediction tasks (Hastie et al., 2020). In an effort to make BSS computational feasible in high-dimensional settings, Bertsimas et al. (2016) proposed fast and scalable algorithms to generate solutions to the BSS problem (2) directly.

Since most of these methods are based on the squared loss function, they are very sensitive to atypical observations in the data, which may adversely affect their variable selection and prediction performances. Over the last two decades, several robust methods have been proposed that can be used when $p \gg n$, can select only a subset of relevant predictors, and are resistant to outliers. Khan et al. (2007a) and Khan et al. (2007b) were among the first ones to develop robust stepwise and least angle regression (LARS) (Efron et al., 2004) algorithms using robust estimators of pairwise correlations instead of their classical sample counterparts. Alfons et al. (2013) introduced a penalized version of the least trimmed squares estimator (Rousseeuw, 1984), called sparseLTS, by minimizing the LASSO-penalized sum of h smallest squared residuals, with $h \leq \lfloor n/2 \rfloor$. Also inspired by LTS, Thompson (2022) introduced the Robust Best Subset Selection (RBSS) method by combining the LTS loss with the ℓ_0 -penalty for the vector of coefficients. The objective function becomes

$$\min_{\beta \in \mathbb{R}^p} \sum_{i \in I} (y_i - \mathbf{x}_i^T \beta)^2 \quad \text{subject to} \quad \begin{cases} \|\beta\|_0 \leq t, \\ |I| \geq h, \end{cases} \quad (3)$$

where $I = \{1, \dots, n\}$ and $|\cdot|$ is the cardinality operator.

Thompson (2022) established the finite-sample breakdown point of RBSS, and developed a computing algorithm. Other sparse robust regression estimators have been later proposed using other loss and penalty functions, including the MM-LASSO (Smucler and Yohai, 2017), PENSE(M) (Cohen Freue et al., 2019) and their adaptive versions (Kepplinger, 2023).

2.2 Ensemble Methods

Ensemble methods have been proposed to generate and aggregate multiple models with appealing performance in high-dimensional prediction tasks. Ueda and Nakano (1996) decomposed the mean squared prediction error (MSPE) of regression ensembles and showed that the variance of an ensemble is largely determined by how correlated its individual models are. Thus, until recently, most ensemble methods relied on a large number of weak decorrelated models (typically more than 100). For example, decorrelation of the individual trees in RF is achieved by random sampling of the data (i.e., bagging, Breiman, 1996a) and random sampling of the predictors (i.e., the random

predictor subspace method, [Ho, 1998](#)). Similarly, ensembles from large number of diverse linear models are generated in the RGLM method ([Song et al., 2013](#)) and through gradient boosting ([Chen and Guestrin, 2016](#)). However, their individual models are not interpretable and have weak predictive accuracy. In addition, the selection of predictors is unreliable if randomization is used, and in the case of gradient boosting the models are fit on residuals rather than the original data.

To generate ensembles of sparse, accurate and diverse models, [Christidis et al. \(2020\)](#) and [Christidis et al. \(2024\)](#) relied on the principle of the multiplicity of good models ([McCullagh and Nelder, 1989](#)). [Christidis et al. \(2020\)](#) proposed a method called Split-Regularized Regression (SplitReg) that splits the set of predictors into groups and builds a set of sparse models by minimizing an objective function that encourages sparsity within each group and diversity among them. To alleviate using the multi-convex relaxation of SplitReg and control the degrees of sparsity and diversity directly, [Christidis et al. \(2024\)](#) introduced a multi-model subset selection (MSS) as a generalization of BSS in (2). The degree of sparsity of the models and diversity between them are chosen by CV and thus driven directly by the data. Despite the high prediction accuracy and interpretability of these ensembles and the models that comprise them, they are very sensitive outliers in the data.

3 Robust Multi-Model Subset Selection

In this section we introduce our Robust Multi-Model Subset Selection estimator (RMSS) to build an ensemble of strong sparse predictive models from a high-dimensional and complex dataset containing outlying samples. RMSS aims to find $G \geq 2$ robust, sparse and diverse well-performing models that can also be combined into a highly accurate robust ensemble model.

Let β_j^g denote the coefficient for predictor j in model g , for $1 \leq j \leq p$ and $1 \leq g \leq G$. Let $\beta^g = (\beta_1^g, \beta_2^g, \dots, \beta_p^g)^T \in \mathbb{R}^p$ be the vector of coefficients of model g , for $1 \leq g \leq G$, and $\beta_j = (\beta_j^1, \beta_j^2, \dots, \beta_j^G)^T \in \mathbb{R}^G$ be the vector of coefficients of predictor j across the G models. For a fixed number of models $G \geq 2$, RMSS solves the constrained optimization problem

$$\min_{\beta^1, \dots, \beta^G \in \mathbb{R}^p} \sum_{g=1}^G \sum_{i \in I^{(g)}} (y_i - \mathbf{x}_i^T \beta^g)^2 \quad \text{subject to} \quad \begin{cases} \|\beta^g\|_0 \leq t, & 1 \leq g \leq G, \\ \|\beta_j\|_0 \leq u, & 1 \leq j \leq p, \\ |I^{(g)}| \geq h, & 1 \leq g \leq G. \end{cases} \quad (4)$$

where the subset $I^{(g)} \subseteq I = \{1, \dots, n\}$ indicates the subset of at least $h \leq n$ observations used to estimate model g ; t controls the number of predictors selected for each model ($t \leq \min(n-1, p)$);

and u sets maximum number of times a predictor can be shared among models ($u \leq G$).

In our algorithm, the tuning parameters h , t , and u are chosen by CV. The argument h can also be predetermined by the user if there is some a priori knowledge of the degree of data contamination. Since only a subset of the predictor variables may be contaminated for any given sample, a nice feature of RMSS is that the individual models in the ensembles may be fit on different subsamples (i.e., different subsets $I^{(g)}$). Thus, RMSS may potentially use more observations relative to other single-model sparse and robust methods, reducing the loss of information from the training data.

We now address some special cases of RMSS for different configurations of the tuning parameters. If $h = n$ in (12) it follows immediately that RMSS is equivalent to MSS for the same values of t and u . RMSS can also be seen as a generalization of RBSS that leverages the use of multiple models.

Proposition 1 *If $u = G$ in (12), then there is no diversity among the individual models in RMSS and the solution to each of these models is the optimal solution to RBSS in (3) with sparsity and robustness parameters t and h , respectively.*

The proof of Proposition 1 provided in the supplementary material follows directly from the fact that there is no restriction on the sharing of predictors. Thus, the minimum loss for each model is achieved by the RBSS optimal solution with the same tuning parameters t and h . Corollary 1 below follows immediately from Proposition 1.

Corollary 1 *If $u = G$ in (12), then if*

- (I) $h = n$, RMSS is equivalent to BSS, and
- (II) $t = p < n - 1$, RMSS is equivalent to LTS.

Since the tuning parameters t , u and h are chosen by CV, RMSS can easily adapt to data with different characteristics, e.g., data with very few predictors or without any outliers.

In this article, we generate ensembles using the simple model averaging method, where the coefficients of an ensemble $\bar{\beta}$ are the average of the estimated coefficients $\hat{\beta}^g$ of the G models. However, other methods can also be implemented, including the weighted model averaging methods (Breiman, 1996b) or model aggregation methods (Biau et al., 2016).

4 Finite-Sample Breakdown Point

This section establishes the finite-sample breakdown point of RMSS ensembles, a standard robustness measure defined by Donoho and Huber (1983) that indicates the smallest fraction of contam-

inated observations needed to render the estimator meaningless. The mathematical definition of the finite-sample breakdown point is given in Definition 1 below.

Definition 1 Let (\mathbf{X}, \mathbf{y}) be an uncontaminated sample of size n , and denote by $(\mathbf{X}_c, \mathbf{y}_c)$ the sample (\mathbf{X}, \mathbf{y}) with $1 \leq m \leq n$ observations replaced by outliers. Let $T(\mathbf{X}, \mathbf{y})$ be some estimator of data (\mathbf{X}, \mathbf{y}) . The finite-sample breakdown point of $T(\mathbf{X}, \mathbf{y})$ is given by

$$B(T|\mathbf{X}, \mathbf{y}) = \min_{1 \leq m \leq n} \left\{ \frac{m}{n} : \sup_{(\mathbf{X}_c, \mathbf{y}_c)} \|T(\mathbf{X}, \mathbf{y}) - T(\mathbf{X}_c, \mathbf{y}_c)\|_2 = \infty \right\}. \quad (5)$$

In Theorem 1 and Corollary 2, we establish the finite-sample breakdown point of RMSS ensembles and the individual models that comprise them, respectively. The proofs are provided in the supplementary material.

Theorem 1 Let $T(\mathbf{X}, \mathbf{y})$ be the optimal value of the objective function of RMSS in (12). Then, $T(\mathbf{X}, \mathbf{y})$ has finite-sample breakdown point

$$B(T|\mathbf{X}, \mathbf{y}) = \frac{n - h + 1}{n}.$$

Corollary 2 Let $T^g(\mathbf{X}, \mathbf{y})$ be the trimmed sum of squares of model g ,

$$T^g(\mathbf{X}, \mathbf{y}) = \sum_{i \in \hat{I}^{(g)}} \left(y_i - \mathbf{x}_i^T \hat{\boldsymbol{\beta}}^g \right)^2, \quad 1 \leq g \leq G,$$

where $\hat{I}^{(g)}$ and $\hat{\boldsymbol{\beta}}^g$ are the optimal subset of samples and vector of coefficients for model g in RMSS (12), respectively. Then $T^g(\mathbf{X}, \mathbf{y})$ has finite-sample breakdown point

$$B(T^g|\mathbf{X}, \mathbf{y}) = \frac{n - h + 1}{n}, \quad 1 \leq g \leq G.$$

From Theorem 1 and Corollary 2, it follows that RMSS ensembles and the individual models that comprise them are resistant to up to $n - h$ contaminated samples. Since $h = n$ corresponds to the MSS ensemble method of Christidis et al. (2024), the breakdown point of MSS ensembles and the individual models that comprise them is $1/n$ and are thus not resistant to any contamination level. If the trimming parameter h is chosen by CV, the extent to which RMSS ensembles and their individual models are resistant to outliers is data-driven.

5 Computing Algorithm

The evaluation of every possible combination of predictors in RMSS is not feasible, even for a low-dimensional case in which predictors are not shared between models (see a combinatorics result in the supplementary material). Thus, we propose an algorithm to search over a three-dimensional grid of the tuning parameters h , t and u , reducing the computational cost of the CV.

We first center and scale the response \mathbf{y} and the columns of the design matrix \mathbf{X} using their medians and median absolute deviations, respectively. We then generate disjoint subsets of predictors for the particular case $u = 1$ (section 5.1) and generate solutions for any combination of values in the three-dimensional grid of h , t and u (section 5.2). We also developed a three-dimensional neighborhood search to sequentially improve incumbent solutions generated by our algorithm (see supplementary material). However, numerical experiments show only marginal improvements in prediction and variable selection performances. Nevertheless, this option is available in our software. Once all computations are done on the scaled data, the estimated coefficients are returned to their original scale.

5.1 Initial Predictor Subsets

Christidis et al. (2024) proposed a multi-model forward selection algorithm that selects G disjoint subsets of predictors to initialize the algorithm of MSS. However, their approach is sensitive to outliers in the data. Thus, in Algorithm 1 we propose a robust multi-model stepwise selection criteria to select disjoint initial subsets of predictors $J_{\text{init}}^{(g)} \subseteq J = \{1, \dots, p\}$, $1 \leq g \leq G$.

Algorithm 1 generalizes the robust forward stepwise regression algorithm of Khan et al. (2007a) to multiple models. For a single-model framework, Khan et al. (2007a) showed that the stepwise forward search algorithm depends only the sample means, variances and correlations between the variables in the model. Thus, to make the algorithm robust to outlying observations, they proposed replacing these sample estimators by robust counterparts to compute the residual sum of squares and partial F -rules (see section 4 in the supplementary material for further details).

In our implementation, we compute robust correlation estimates via the state-of-the-art Detect Deviating Cells (DDC) method of Rousseeuw and Bossche (2018), which can be scaled to high-dimensional settings using properties of product moments (Raymaekers and Rousseeuw, 2021). We denote with $\hat{\mathbf{r}}_{\mathbf{y}}$ the robust estimates of the correlations between the response and predictors, and $\hat{\mathbf{\Sigma}}$ the robust estimate of the matrix of correlations of the predictors. These robust estimates are used to compute robust residual sum of squares (rRSS) and robust partial F -rules as in Khan

et al. (2007a). Due to post-inference problems and the use of plug-in estimators, the resulting methodology can not be used to perform formal F -tests with valid p -values from the F -distribution. In Algorithm 1 these rules and p -values are only used to develop a selection and a stopping criterion.

Algorithm 1 Robust Multi-Model Stepwise Selection

Input: Robust correlation vector of the response $\hat{\mathbf{r}}_{\mathbf{y}} \in \mathbb{R}^p$, robust correlation matrix of predictors $\hat{\mathbf{\Sigma}} \in \mathbb{R}^{p \times p}$, number of models $G \geq 2$, and saturation threshold $\gamma \in (0, 1)$.

Initialize: The set of candidates $J = \{1, \dots, p\}$, and for each model the set of model predictors $J^{(g)} = \emptyset$ and the model saturation indicator $T^{(g)} = \text{FALSE}$, $1 \leq g \leq G$.

- 1: Repeat the following steps until $T^{(g)} = \text{TRUE}$ for all $1 \leq g \leq G$:
 - 1.1. For each model g satisfying $T^{(g)} = \text{FALSE}$:
 - 1.1.1: Identify candidate predictor $j^{(g)}$ that maximizes the decrease in rRSS when combined with the variables in $J^{(g)}$.
 - 1.1.2: Calculate the p -value $\gamma^{(g)}$ from the robust F -rule once the predictor $j^{(g)}$ is added to the current g model.
 - 1.1.3: If $\gamma^{(g)} \geq \gamma$ set $T^{(g)} = \text{TRUE}$.
 - 1.2. Identify the unsaturated model g^* with the smallest p -value $\gamma^{(g^*)}$.
 - 1.3. If $\gamma^{(g^*)} < \gamma$:
 - 1.3.1: Update the set of predictors for model g^* : $J^{(g^*)} = J^{(g^*)} \cup \{j^{(g^*)}\}$.
 - 1.3.2: Update the set of candidate predictors $J = J \setminus \{j^{(g^*)}\}$.
 - 1.3.3: If $|J^{(g^*)}| = n - 1$, set $T^{(g^*)} = \text{TRUE}$.
 - 2: Return the sets of model predictors $J^{(g)}$, $1 \leq g \leq G$.
-

5.2 Computing RMSS over a Grid of Tuning Constants

To develop an efficient computing algorithm to compute RMSS over a grid of t , u and h values, we recast (12) in its equivalent form using auxiliary variables $\boldsymbol{\eta}^g = (\eta_1^g, \dots, \eta_n^g)^T \in \mathbb{R}^n$, $1 \leq g \leq G$, with 0 entries for the smallest h residuals of each model g , and the corresponding residuals otherwise. For any t , u and h values, the equivalent reformulation of RMSS is given by

$$\min_{\beta^1, \dots, \beta^G \in \mathbb{R}^p} \sum_{g=1}^G \mathcal{L}_n(\beta^g, \boldsymbol{\eta}^g | \mathbf{y}, \mathbf{X}) \quad \text{subject to} \quad \begin{cases} \|\beta^g\|_0 \leq t, & 1 \leq g \leq G, \\ \|\beta_j\|_0 \leq u, & 1 \leq j \leq p, \\ \|\boldsymbol{\eta}^g\|_0 \leq n - h, & 1 \leq g \leq G, \end{cases} \quad (6)$$

where the loss function \mathcal{L}_n used for each model g is given by

$$\mathcal{L}_n(\boldsymbol{\beta}^g, \boldsymbol{\eta}^g | \mathbf{y}, \mathbf{X}) = \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}^g - \boldsymbol{\eta}^g\|_2^2. \quad (7)$$

The gradients $\nabla_{\boldsymbol{\beta}} \mathcal{L}_n(\boldsymbol{\beta}, \boldsymbol{\eta} | \mathbf{y}, \mathbf{X})$ and $\nabla_{\boldsymbol{\eta}} \mathcal{L}_n(\boldsymbol{\beta}, \boldsymbol{\eta} | \mathbf{y}, \mathbf{X})$ are both Lipschitz continuous with Lipschitz constants $\ell_{\boldsymbol{\beta}} = 2\|\mathbf{X}^T \mathbf{X}\|_2^2$ and $\ell_{\boldsymbol{\eta}} = 2$, respectively (see proofs in the supplementary material).

We now introduce some notation needed to outline our computing algorithms.

Definition 2 For any $v \in \mathbb{R}^p$ and $r \in \mathbb{R}$, the projection operator $\mathcal{P}(v; r)$, defined as

$$\mathcal{P}(v; r) \in \arg \min_{w \in \mathbb{R}^p} \|w - v\|_2^2 \quad \text{subject to} \quad \|w\|_0 \leq r \quad (8)$$

retains the r largest elements in absolute value of the vector v .

Definition 3 For any $v \in \mathbb{R}^p$, $S \subseteq J = \{1, \dots, p\}$ and $r \in \mathbb{R}$, the projected subset operator $\mathcal{Q}(v; S, r)$, defined as

$$\mathcal{Q}(v; S, r) \in \arg \min_{w \in \mathbb{R}^p} \|w - v\|_2^2 \quad \text{subject to} \quad \begin{cases} \|w\|_0 \leq r \\ \{j \in J : w_j \neq 0\} \subseteq S \end{cases} \quad (9)$$

retains the r largest elements in absolute value of the vector v that belong to the subset S .

Note that both $\mathcal{P}(v; r)$ and $\mathcal{Q}(v; S, r)$ are set-valued maps since more than one possible permutation of the indices $J = \{1, \dots, p\}$ and $\{j \in J : j \in S\}$ may exist.

For a given set of multi-model coefficient estimates $\hat{\boldsymbol{\beta}}^g = (\hat{\beta}_1^g, \dots, \hat{\beta}_p^g)^T \in \mathbb{R}^p$, $1 \leq g \leq G$, let $J^{(g)} = \{j \in J : \hat{\beta}_j^g \neq 0\}$ be the subsets of predictors' indices included in each model and $S_u^{(g)}$ be the subsets of predictors' indices used in at most $u - 1$ models excluding model g ,

$$S_u^{(g)} = \left\{ j \in J : \sum_{\substack{l=1 \\ l \neq g}}^G \mathbb{I}(j \in J^{(l)}) \leq u - 1 \right\}. \quad (10)$$

Lastly, denote $\mathbf{X}_S \in \mathbb{R}^{n \times |S|}$ the submatrix of \mathbf{X} with column indices $S \subseteq J = \{1, \dots, p\}$.

For fixed tuning constants t , u and h and given some starting values $(\tilde{\boldsymbol{\beta}}^g, \tilde{\boldsymbol{\eta}}^g)$, $1 \leq g \leq G$, Algorithm 2, outlines the steps to perform a projected subset block gradient descent (PSBGD) to generate robust multi-model estimates. For each model at a time, the algorithm alternates between

updates of the coefficient estimates $\hat{\beta}^g$ and updates of the subsamples selected given the size of the updated residuals in $\hat{\eta}^g$ until convergence is achieved.

Algorithm 2 Projected Subset Block Gradient Descent (PSBGD)

Input: Matrix of predictor variables $\mathbf{X} \in \mathbb{R}^{n \times p}$, response vector $\mathbf{y} \in \mathbb{R}^n$, starting values $(\tilde{\beta}^g, \tilde{\eta}^g)$, $1 \leq g \leq G$, tuning parameters t, u and h , and tolerance parameter $\epsilon > 0$.

Initialize: For each model g , the set of model predictors $J^{(g)} = \{j \in J : \tilde{\beta}_j^g \neq 0\}$, $1 \leq g \leq G$.

1: Repeat the following steps for each model g , $1 \leq g \leq G$:

1.1: Create the set of predictors' indices $S_u^{(g)}$ via (10) and compute the Lipschitz constant $\ell_{\beta^g} = 2\|\mathbf{X}_{S_u^{(g)}}^T \mathbf{X}_{S_u^{(g)}}\|_2$.

1.2: Update, sequentially, current estimates $\tilde{\beta}^g$ and $\tilde{\eta}^g$ via

$$\begin{aligned}\hat{\beta}^g &\in \mathcal{Q} \left(\tilde{\beta}^g - \frac{1}{L_{\beta^{(g)}}} \nabla_{\beta} \mathcal{L}_n(\beta, \hat{\eta}^g | \mathbf{y}, \mathbf{X}) \Big|_{\beta = \tilde{\beta}^g; S_u^{(g)}, r} \right) \\ \hat{\eta}^g &\in \mathcal{P} \left(\tilde{\eta}^g - \frac{1}{L_{\eta}} \nabla_{\eta} \mathcal{L}_n(\hat{\beta}^g, \eta | \mathbf{y}, \mathbf{X}) \Big|_{\eta = \tilde{\eta}^g; n-h} \right)\end{aligned}$$

with $L_{\beta^g} \geq \ell_{\beta^g}$ and $L_{\eta} \geq \ell_{\eta}$, and repeat until $\mathcal{L}_n(\tilde{\beta}^g, \tilde{\eta}^g | \mathbf{y}, \mathbf{X}) - \mathcal{L}_n(\hat{\beta}^g, \hat{\eta}^g | \mathbf{y}, \mathbf{X}) \leq \epsilon$.

1.3: Update the model predictors $J^{(g)} = \{j \in J : \hat{\beta}_j^g \neq 0\}$ and compute the set of model subsamples $I^{(g)} = \{i \in I : \hat{\eta}_i^g = 0\}$.

1.4: Compute the final model coefficients

$$\hat{\beta}^g = \arg \min_{\beta \in \mathbb{R}^p} \sum_{i \in I^{(g)}} (y_i - \mathbf{x}_i^T \beta)^2 \quad \text{subject to} \quad \beta_j = 0, j \notin J^{(g)}, \quad 1 \leq g \leq G.$$

2: Return the pairs $(\hat{\beta}^g, \hat{\eta}^g)$, $1 \leq g \leq G$.

Proposition 2 below establishes the convergence of the Algorithm 2 (see the proof in the supplementary material).

Proposition 2 For each model g in (6), step 1 of Algorithm 2 generates a converging sequence for the pair $(\hat{\beta}^g, \hat{\eta}^g)$ and the inequalities

$$\begin{aligned}\left\| \hat{\beta}^g - \mathcal{Q} \left(\hat{\beta}^g - \frac{1}{L_{\beta^{(g)}}} \nabla_{\beta} \mathcal{L}_n(\beta, \hat{\eta}^g | \mathbf{y}, \mathbf{X}) \Big|_{\beta = \hat{\beta}^g; S_u^{(g)}, r} \right) \right\|_2^2 &\leq \epsilon, \\ \left\| \hat{\eta}^g - \mathcal{P} \left(\hat{\eta}^g - \frac{1}{L_{\eta}} \nabla_{\eta} \mathcal{L}_n(\hat{\beta}^g, \eta | \mathbf{y}, \mathbf{X}) \Big|_{\eta = \hat{\eta}^g; n-h} \right) \right\|_2^2 &\leq \epsilon,\end{aligned}$$

can be achieved in $O(1/\epsilon)$ iterations.

In Algorithm 3, we outline the steps to generate the solutions $(\hat{\beta}^g[t, u, h], \hat{\eta}^g[t, u, h])$, $1 \leq g \leq$

G , for any t , u , and h over the grids $T = \{t_1, \dots, t_q\}$, $U = \{1, \dots, G\}$ and $H = \{h_1, \dots, h_r\}$, respectively.

Algorithm 3 Decrementing Diversity PSBGD

Input: Design matrix $\mathbf{X} \in \mathbb{R}^{n \times p}$, response vector $\mathbf{y} \in \mathbb{R}^n$, grids of tuning parameters $T = \{t_1, \dots, t_q\}$, $U = \{1, \dots, G\}$ and $H = \{h_1, \dots, h_r\}$, and tolerance parameter $\epsilon > 0$.

- 1: Use Algorithm 1 to obtain the sets of predictors' indices $J_{\text{init}}^{(g)}$, $1 \leq g \leq G$, and compute the initial model coefficients

$$\hat{\beta}_{\text{init}}^g = \arg \min_{\beta \in \mathbb{R}^p} \|\mathbf{y} - \mathbf{X}\beta\|_2^2 \quad \text{subject to} \quad \beta_j = 0, j \notin J_{\text{init}}^{(g)}.$$

- 2: For each combination of $t \in T$ and $h \in H$:

- 2.1: Compute the pairs $(\hat{\beta}^g[t, 1, h], \hat{\eta}^g[t, 1, h])$, $1 \leq g \leq G$, using Algorithm 2 initialized with $(\hat{\beta}_{\text{init}}^g, \mathbf{0}_n)$ where $\mathbf{0}_n = (0, \dots, 0)^T \in \mathbb{R}^n$, $1 \leq g \leq G$.

- 2.2: For $u = 2, \dots, G$:

- 2.2.1: Compute the pairs $(\hat{\beta}^g[t, u, h], \hat{\eta}^g[t, u, h])$, $1 \leq g \leq G$, using Algorithm 2 initialized with $(\hat{\beta}^g[t, u-1, h], \hat{\eta}^g[t, u-1, h])$, $1 \leq g \leq G$.

- 3: Return the pairs $(\hat{\beta}^g[t, u, h], \hat{\eta}^g[t, u, h])$, $1 \leq g \leq G$, for all combinations of $t \in T$, $u \in U$ and $h \in H$.
-

5.3 Selection of Tuning Parameters

We use 5-fold CV to select the final combination of t , u and h from the grids of candidates T , U and H . Since the test folds may contain outliers, we use the fast and robust τ -estimator of location (Maronna and Zamar, 2002) on the prediction residuals of the test folds for each combination of t , u and h . Our final selection is the combination with the smallest estimate.

The fine grids $T = \{1, \dots, n-1\}$ and $H = \{\lfloor n/2 \rfloor + 1, \dots, n\}$ may be used for the sparsity and robustness parameters, respectively. But less dense grids can also be used to speed up computation. Oftentimes, a priori information is available about the level of contamination of the data which may guide the choice for H . Our method has the flexibility to choose h by CV or set it to a fix value. We illustrate this feature in the simulation section. In general, the grid $U = \{1, \dots, G\}$ should be fixed since it is required for the computation for solutions in Algorithm 3.

5.4 Software

The implementation of robust multi-model stepwise selection outlined in Algorithm 1 is available on CRAN (R Core Team, 2022) in the R package `robStepSplitReg` (Christidis and Cohen-Freue, 2023b). The implementation of Algorithms 2 - 3 to fit RMSS ensembles is also available on CRAN

in the R package RMSS (Christidis and Cohen-Freue, 2023a), which generates RBSS if $G = 1$. The source code of RMSS is written in C++, and multithreading via OpenMP (Chandra et al., 2001) is available in the package to further speed up computations.

6 Simulations

In this section, we investigate the performance of RMSS against robust and sparse methods as well as blackbox ensemble methods in an extensive simulation study where the data is contaminated in both the predictor space and the response. We also use a block correlation structure between predictor variables to mimic as closely as possible the behavior of many modern datasets (Zhang and Coombes, 2012).

6.1 Simulation of Uncontaminated Data

In each setting of our simulation study, we generate the uncontaminated data from the linear model

$$y_i = \mu + \mathbf{x}_i^T \boldsymbol{\beta}_0 + \sigma \varepsilon_i, \quad 1 \leq i \leq n,$$

with independent and identically distributed $x_i \sim N_p(\mathbf{0}, \boldsymbol{\Sigma})$ and $\varepsilon_i \sim N(0, 1)$, sample size $n = 50$ and the number of predictors $p = 500$. Alternative choices for n and p where $p \gg n$ lead to similar conclusions and are omitted for conciseness.

We examine models with different proportions ζ of active predictors, with $\zeta \in \{0.1, 0.2, 0.4\}$. Without loss of generality, we construct the correlation matrix of the active predictors as a block matrix, with each block corresponding to 25 predictors, a within-block correlation $\rho_1 = 0.8$ and a between-block correlation $\rho_2 = 0.2$. The non-active predictors are independent and uncorrelated with the active ones.

For simplicity, the intercept is set to $\mu = 0$. The coefficients of the active variables, $\{\beta_j : \beta_j \neq 0, 1 \leq j \leq p\}$, are randomly generated from the random variable $(-1)^Z \times U$, where Z is Bernoulli distributed with parameter 0.7 and U is uniformly distributed on the interval $(0, 5)$.

The noise parameter σ is computed based on the desired signal to noise ratio, $\text{SNR} = \boldsymbol{\beta}'_0 \boldsymbol{\Sigma} \boldsymbol{\beta}_0 / \sigma^2$. We consider SNRs of 0.5 (low signal), 1 (moderate signal), 2 (high signal), which correspond to proportions of variance explained $\text{PVE} = \text{SNR} / (\text{SNR} + 1)$ of 33.3%, 50% and 66.7%, respectively.

6.2 Data Contamination

We contaminate the first $m = \lfloor \tau n \rfloor$ samples (\mathbf{x}_i, y_i) according to the model proposed in [Maronna \(2011\)](#). The regression outliers are introduced by replacing the predictors \mathbf{x}_i with

$$\tilde{\mathbf{x}}_i = \Theta_i + \frac{k_{\text{lev}}}{\sqrt{\mathbf{a}^T \boldsymbol{\Sigma}^{-1} \mathbf{a}}} \mathbf{a}, \quad 1 \leq i \leq m,$$

where $\Theta_i \sim \mathcal{N}(\mathbf{0}_p, 0.01 \times \mathbf{I}_p)$ and $\mathbf{a} = \tilde{\mathbf{a}} - (1/p)\tilde{\mathbf{a}}^T \mathbf{1}_p$ where \mathbf{I}_p is the p -dimensional identity matrix, $\mathbf{0}_p = (0, \dots, 0)^T \in \mathbb{R}^p$, $\mathbf{1}_p = (1, \dots, 1)^T \in \mathbb{R}^p$, and the entries of \tilde{a}_j of $\tilde{\mathbf{a}}$ follow a uniform distribution on the interval $(-1, 1)$, $1 \leq j \leq p$. The parameter k_{lev} controls the distance in the direction most influential for the estimator.

We also contaminate the observation in the response by altering the regression coefficient

$$\tilde{y}_i = \tilde{\mathbf{x}}_i^T \tilde{\boldsymbol{\beta}}, \quad \tilde{\beta}_j = \begin{cases} \beta_j(1 + k_{\text{slo}}), & \beta_j \neq 0, \\ k_{\text{slo}} \|\boldsymbol{\beta}\|_{\infty}, & \text{otherwise,} \end{cases} \quad 1 \leq i \leq m.$$

The parameters k_{lev} and k_{slo} control the position of the contaminated observations. Our preliminary experiments showed that the effect of k_{lev} on estimators examined was almost the same for any $k_{\text{lev}} > 1$, hence we fixed $k_{\text{lev}} = 2$. We also found that the position the vertical outliers affects the estimators much more, and that the performance of non-robust estimators degraded significantly for any $k_{\text{slo}} \geq 100$, hence we fixed $k_{\text{slo}} = 100$. We consider the contamination proportions of $\tau = 0$ (no contamination), $\tau = 0.15$ (moderate contamination) and $\tau = 0.3$ (high contamination).

6.3 Methods

Our simulation study compares the prediction and variable selection accuracy of eight methods. All computations were carried out in **R** using the implementations listed below

1. Elastic Net (**EN**, [Zou and Hastie, 2005](#)), with `glmnet` package ([Friedman et al., 2010](#)).
2. Adaptive **PENSE** ([Kepplinger, 2023](#)), with `pense` package ([Kepplinger et al., 2023](#)).
3. EN Penalized Huber (**HuberEN**, [Yi and Huang, 2017](#)), with `hqreg` package ([Yi, 2017](#)).
4. Sparse LTS (**SparseLTS**, [Alfons et al., 2013](#)), with `robustHD` package ([Alfons, 2021](#)).
5. Robust Best Subset Selection (**RBSS**, [Thompson, 2022](#)), with `RMSS` package.
6. Random GLM (**RGLM**, [Song et al., 2013](#)), with `RGLM` package ([Song and Langfelder, 2013](#)).

7. Random Forest (**RF**, Breiman, 2001), with `randomForest` package (Liaw and Wiener, 2002).
8. Robust Multi-Model Subset Selection (**RMSS**) with $G = 10$ models, proposed in this paper, with `RMSS` package.

To reduce the computational cost of RMSS in our extensive simulation study, we use the candidate grid $T = \{0.3n, 0.4n, 0.5n\} = \{15, 20, 25\}$ for the sparsity tuning parameter. We use a grid of values $H = \{(1 - (\tau + 0.1))n, (1 - (\tau + 0.05))n, (1 - \tau)n\}$ where $\tau \in \{0, 0.15, 0.3\}$ and determine the robustness tuning parameter h by CV. We use the default grid $U = \{1, \dots, G\}$ with $G = 10$ for the diversity parameter as required by our computing algorithm. RBSS is computed at the same time as RMSS in a single function call of the `RMSS` package by fixing $u = G$ (see Proposition 1). While better empirical results may be obtained with RMSS when combined with a larger number of models and more refined grids for the tuning parameters, we find that even with our suboptimal settings RMSS is competitive with state-of-the-art sparse and robust methods. To select tuning parameters we use 10-fold CV for EN and 5-fold CV for the robust methods. Additional details about the selection of tuning parameters of each estimator is given in the supplementary material.

6.4 Performance Measures

For each combination of the sparsity parameter ζ , SNR and contamination level τ , we randomly generate $N = 50$ training sets and a large (uncontaminated) independent test set of size 2,000. In each replication of a particular configuration, we fit the methods on the training sets and we compute the MSPE using the independent test set. The reported MSPEs are relative to the variance of the irreducible error σ^2 , i.e., the best possible result is 1. We also compute the recall (RC) and precision (PR) which are defined as

$$\text{RC} = \frac{\sum_{j=1}^p \mathbb{I}(\beta_j \neq 0, \hat{\beta}_j \neq 0)}{\sum_{j=1}^p \mathbb{I}(\beta_j \neq 0)}, \quad \text{PR} = \frac{\sum_{j=1}^p \mathbb{I}(\beta_j \neq 0, \hat{\beta}_j \neq 0)}{\sum_{j=1}^p \mathbb{I}(\hat{\beta}_j \neq 0)},$$

where β and $\hat{\beta}$ are the true and estimated regression coefficients, respectively. We use the ensemble fit $\bar{\beta}$ for RMSS. We do not report RC and PR of RGLM and RF since their recall tends to be always close to 1 while their precision equals approximately the proportion of active variables ζ . RC and PR range between 0 and 1 and large values are desirable.

6.5 Results

In Table 1, we report the average and lowest MSPE rank of the eight methods over the nine possible combinations of the SNR and sparsity level for each contamination proportion. The top performance in each column of a table are highlighted in bold fonts.

In the $\tau = 0$ (no contamination) case, RGLM achieved the best overall rank followed by RMSS and RF. RMSS consistently outperformed the non-robust EN and never ranked worse than fourth place among the eight methods. This demonstrates the ability of RMSS to adapt to the level of contamination of the data. In the $\tau = 0.15$ (moderate contamination) case PENSE and RMSS traded the top rank over every possible combination of SNR and sparsity level, with PENSE achieving the highest rank more often. However in the $\tau = 0.30$ (high contamination) case, PENSE’s performance deteriorated while RMSS achieved the top rank in every possible configuration of the simulation. RBSS was not competitive with RMSS and PENSE, except in the high contamination case where it achieved the second best average rank behind RMSS. The non-robust ensemble methods’ predictive performance deteriorated completely for both the moderate and high contamination cases, achieving the lowest two ranks for both the moderate and high contamination scenarios.

To better understand the relevance of these summary results, in Figure 1, we plot the MSPE of the top robust competing methods in the contaminated scenarios, PENSE, RBSS and RMSS, estimated from $N = 50$ random training sets over different contamination and sparsity levels for $\text{SNR} = 1$ (moderate signal). It is evident that in the moderate contamination case PENSE and RMSS perform very similarly for all sparsity level, while for the high contamination case RMSS significantly outperforms PENSE and RBSS for any sparsity level. Similar conclusion are obtained for all three SNRs considered.

In Table 2, we report the average and lowest RC and PR rank of the six methods for which we compute the RC and PR, over the nine possible combinations of the SNR and sparsity level for each contamination proportion. RMSS achieved the best average RC rank over all contamination levels, while PENSE was the second best performing method over all contamination levels. In terms of PR, RBSS was the best performing method, followed by RMSS when the data was contaminated.

In Figure 2, we plot the RC and PR of PENSE, RBSS and RMSS over the $N = 50$ random training sets over different contamination and sparsity levels for a moderate SNR. RMSS with $G = 10$ models outperformed PENSE and RBSS in terms of RC over all contamination levels when $\zeta = 0.1$ and 0.2 , and was competitive with PENSE when $\zeta = 0.4$. In our numerical experiments, RMSS achieved even superior RC when we used more than $G = 10$ models. RBSS generally had the best performance in terms of PR but at the cost of a poor RC for any combination of

Table 1: Average and lowest MSPE rank of the eight methods over the nine SNR and sparsity level combinations for each contamination proportion.

Method	MSPE Rank					
	$\tau = 0$		$\tau = 0.15$		$\tau = 0.3$	
	Avg	Low	Avg	Low	Avg	Low
EN	5.2	6	6.0	6	4.7	5
PENSE	4.1	7	1.2	2	4.1	5
HuberEN	5.0	6	4.4	5	2.8	4
SparseLTS	7.7	8	3.3	4	7.0	8
RBSS	7.1	8	4.2	5	2.4	4
RGLM	1.1	2	8.0	8	7.7	8
RF	2.9	5	7.0	7	6.3	7
RMSS	2.9	4	1.8	2	1.0	1




Figure 1: MSPE of PENSE, RBSS and RMSS over $N = 50$ random training sets over different contamination and sparsity levels for SNR = 1.

sparsity, contamination and SNR level. RMSS combined a high RC with a high PR consistently across all configurations of our simulation study. Overall, the results of this study demonstrate a steady desirable performance in complex data with high-dimensional block correlation structure and containing regression outliers.

Table 2: Average and lowest RC and PR rank of the six methods over the nine SNR and sparsity level combinations for each contamination proportion.

Method	RC Rank						PR Rank					
	$\tau = 0$		$\tau = 0.15$		$\tau = 0.3$		$\tau = 0$		$\tau = 0.15$		$\tau = 0.3$	
	Avg	Low	Avg	Low	Avg	Low	Avg	Low	Avg	Low	Avg	Low
EN	5.4	6	5.7	6	5.3	6	2.1	4	5.7	6	5.3	6
PENSE	2.3	3	1.8	2	1.9	2	5.0	5	3.0	3	3.0	3
HuberEN	2.0	3	5.1	6	5.7	6	3.7	4	5.3	6	5.7	6
SparseLTS	5.4	6	4.1	5	4.0	4	6.0	6	4.0	4	4.0	4
RBSS	4.1	5	3.1	4	3.0	3	1.2	2	1.0	1	1.0	1
RMSS	1.7	3	1.2	2	1.1	2	3.0	4	2.0	2	2.0	2




Figure 2: RC and PR of PENSE, RBSS and RMSS over $N = 50$ random training sets over different contamination and sparsity levels for SNR = 1.

6.6 Computing Times

The average computing times of the R function calls over all configurations of our simulation is given in Table 3. The time of RMSS also includes the computation of RBSS since the latter is simultaneously computed by setting $u = G$ in (12). Our implementation of RMSS generates $G = 10$ sparse and robust models simultaneously and still achieved a lower average computing time than PENSE, which generates a single sparse and robust model, with a smaller number of initial estimates than the default in its R implementation. Moreover, RMSS must perform CV over three tuning parameters compared to only one for all the single-model sparse and robust methods.

The computing time of RMSS would increase significantly if it uses a neighborhood search strategy. In general, we find that the neighborhood search improves the solutions $\hat{\beta}^g$ ($1 \leq g \leq G$) in terms of variable selection and minimizing the objective function (12) but not in terms of prediction.

Table 3: Computation time of R function calls for the methods in CPU seconds. CPU seconds are on a 2.7 GHz Intel Xeon processor in a machine running Linux 7.8 with 125 GB of RAM.

Method	EN	PENSE	HuberEN	SparseLTS	RMSS
Time	0.1	281.8	0.1	169.5	239.7

7 Contamination of Bioinformatics and Cheminformatics Data

In biomedical sciences new deoxyribonucleic acid (DNA) microarray and ribonucleic acid (RNA) sequencing technologies allow for an increase in the type and volume of the genomics data collected (e.g., [Byron et al., 2016](#)). In chemistry, innovative microscopic technologies allow for the collection of data on the composition of chemical compounds and molecules. Outliers are not rare in these datasets. For example, in genomics, many datasets contain atypical observations obtained from samples with poor measurement quality or incorrect reads (e.g., [Sangiovanni et al., 2019](#)). In cheminformatics the topic of high-dimensionality and robustness has gained a lot of attention in recent years due to the emerging fields of computer-aided drug design and computational toxicology among others (see e.g. [Basak and Vracko, 2022](#)).

In this section we artificially contaminate real bioinformatic and cheminformatic datasets to evaluate the performance of RMSS and the other methods in situations that mimic real applications. We also show that RMSS can uncover some predictors that may be relevant to predict the outcome

of interest but that may not be picked up by single-model sparse robust methods.

7.1 Bioinformatics Data

In a study analyzing the genetic basis of Bardet-Biedl syndrome (BBS), [Scheetz et al. \(2006\)](#) performed mutation and functional studies and identified TRIM32 (tripartite motif-containing protein 32) as a gene whose expression highly correlates with the incidence of BBS. The R package `abess` ([Zhu et al., 2022](#)) contains a dataset with the expression of TRIM32 and $p = 500$ genes for 120 mammalian-eye tissue samples, which is a subset of the original dataset analyzed by [Scheetz et al. \(2006\)](#). The $p = 500$ genes were selected from the 18,976 available genes based on their marginal correlation with TRIM32 and are used to predict the expression of TRIM32.

The normalized gene expression levels in the uncontaminated dataset are all below 10 in magnitude. We randomly split the samples $N = 50$ times in a training set of size $n = 50$ and a test set of size 70. We contaminate 25% of the samples of each training set by replacing the expression of TRIM32 and 100 randomly selected predictor genes with a normal random variable with mean 25 and standard deviation 1. We evaluate the MSPE of EN and the five robust methods used in section 6 on the uncontaminated test set. For RMSS we fix $h = \lfloor 0.75n \rfloor = 37$ but keep the grid $T = \{0.3n, 0.4n, 0.5n\} = \{15, 20, 25\}$ for the sparsity parameter. For the other methods we use the same configurations as in section 6.

The MSPE and standard deviation (SD) of the MSPE reported in Table 5 are relative to the lowest value attained by the six methods, thus the best possible value is 1. The best performance for each measure is highlighted in bold fonts. RMSS achieved the best performance in terms of MSPE with the lowest MSPE variability. SparseLTS was the closest competitor but its MSPE was still 7% higher than the MSPE of RMSS. The individual $G = 10$ models of RMSS also achieved a high prediction accuracy with an average MSPE only 10% higher than the MSPE of RBSS. This observation indicates that each individual model correctly identified the outlying samples (see Corollary 2). As expected, the EN completely deteriorated with the addition outliers to the data.

Table 4: MSPE and SD of the six methods relative to the best performance for the artificially contaminated BBS bioinformatic dataset.

Method	EN	PENSE	HuberEN	SparseLTS	RBSS	RMSS
MSPE	> 25	1.11	1.26	1.07	1.77	1.00
SD	> 100	1.40	1.64	1.26	4.11	1.00

Beyond the good predictive performance of RMSS, the ensembles can potentially uncover genes that are not identified by the other methods. In particular, in the presence of high-dimensional data there are multiple models comprised of different subsets of predictors that can each achieve a high prediction accuracy. This phenomenon is known as the “the multiplicity of good models” in the statistical literature (see relevant discussions in [McCullagh and Nelder, 1989](#)). Thus, single-model sparse and robust methods may potentially discard important genes from the decision-making process. On the BBS dataset, no gene was selected more than 50% of the time by PENSE or SparseLTS over the $N = 50$ random training sets, while 30 genes were selected more than 50% of the time by RMSS. Moreover, the genes most often selected by PENSE and SparseLTS were often selected by RMSS. For example, PENSE selected gene at probe 13704291_at most often, and this gene was selected the same number of times by RMSS. Conversely, the genes most often selected by RMSS were seldom selected by PENSE or SparseLTS. For example, the gene at probe 1374809_at was selected 72% of the time by RMSS and only 2% and 4% of the time by PENSE and SparseLTS, respectively.

7.2 Cheminformatics Data

We analyze the glass dataset from [Lemberge et al. \(2000\)](#) for which the goal is to predict the concentration of the chemical compound Na₂O based on its frequency measurements obtained from an electron probe X-ray microanalysis (EPXMA). After removing variables with little variation, the dataset is comprised of $p = 486$ frequency measurements for $n = 180$ samples. We split the data into $N = 50$ training sets of size $n = 50$ and test sets of size $m = 130$. We contaminate the training sets in the same way as we did for the BBS data, and compute the MSPE of the same methods using the uncontaminated test sets.

In Table 4, we report the MSPE and SD of the MSPE of the methods relative to the best performing method. RMSS again achieved the best MSPE by far as SparseLTS coming was the second method with an MSPE 54% larger. RMSS was also the most stable method in terms of prediction accuracy. The $G = 10$ individual models of RMSS achieved an MSPE similar to RBSS (only 3% larger). The good predictive performance of RMSS is not restricted to the chemical compound Na₂O and may be observed in more compounds available in [Lemberge et al. \(2000\)](#).

RMSS also uncovered frequency measurements that may be relevant to predict the concentration of Na₂O that were missed by PENSE and SparseLTS. The frequency measurement most often selected by RMSS (92% of the time over the $N = 50$ random training sets) was only selected 2% of the time by both PENSE and SparseLTS. On the other hand, the frequency measurements

Table 5: MSPE and SD of the six methods relative to the best performance for the artificially contaminated glass cheminformatic dataset.

Method	EN	PENSE	HuberEN	SparseLTS	RBSS	RMSS
MSPE	> 150	1.90	10.91	1.54	2.58	1.00
SD	> 300	2.62	7.04	1.28	3.86	1.00

most often selected by PENSE and SparseLTS were often selected by RMSS. In fact, the frequency measurement most often selected by PENSE was selected only 30% of the time, and this same frequency measurement was selected 90% of the time by RMSS.

8 Summary and Future Works

In this article, we introduce RMSS, a data-driven method to build an ensemble of sparse and robust models to predict a response of interest and select important predictors from high-dimensional datasets possibly containing outlying observations. To the best of our knowledge, this is the first method proposed for this aim. The levels of sparsity, diversity and robustness of each model are driven directly by the data based on a CV criterion. We established the finite-sample breakdown point of the ensembles and the individual models within the ensembles. To bypass the NP-hard computational complexity of RMSS, we developed a tailored computing algorithm with a local convergence property by leveraging recent developments in the ℓ_0 -optimization literature. Our extensive numerical experiments on synthetic and real data demonstrate the excellent performance of RMSS relative to state-of-the-art sparse and robust methods in high-dimensional prediction tasks when the data is also contaminated. We also showed how RMSS can potentially uncover important predictor variables that may be discarded by single-model sparse and robust methods.

Since RMSS can potentially uncover predictor variables that are not picked up by single-model methods, the addition of interaction terms may potentially further increase the competitive advantage of RMSS over single-model sparse and robust methods. For example, in the -omics sciences where interactions between genes or proteins may drive the outcome of interest. The empirical performance of RMSS can be improved further by considering alternative ways to combine the models in the ensembles other than the simple model average we used in this article. Our work can be extended by considering other robust loss functions to build sparse robust models.

With the growing emphasis on interpretable statistical and machine learning algorithms in the literature and in real data applications, our proposal will potentially pave the way for the

development of other robust ensemble methods. A potential bottleneck in this area of research is the high computational cost of such methods, thus new optimization tools will be needed to render such ensemble methods feasible in practice.

Acknowledgement

Most of the numerical results were generated using a computational infrastructure funded by the Canada Foundation for Innovation (CFI). Authors were supported by NSERC Discovery grant and the Data Science Institute of the authors' affiliation.

Code and Data

The code and data to reproduce the numerical results of this manuscript is submitted with this article are available at <https://doi.org/10.5281/zenodo.14202073>.

Appendix A: Special Case of RMSS

In the proof below we show that when $u = G$, each individual model in RMSS is the solution to RBSS for a fixed sparsity parameter $t \leq \min(n - 1, p)$ and trimming parameter $h \leq n$.

If $u = G$, then there is no restriction in the sharing of predictor variables. In particular, a predictor variable can appear in as many as G models.

$$\begin{aligned} & \min_{\beta^1, \dots, \beta^G \in \mathbb{R}^p} \sum_{g=1}^G \sum_{i \in I^{(g)}} (y_i - \mathbf{x}_i^T \beta^g)^2 \quad \text{subject to} \quad \begin{cases} \|\beta^g\|_0 \leq t, & 1 \leq g \leq G, \\ \|\beta_j\|_0 \leq G, & 1 \leq j \leq p, \\ |I^{(g)}| \geq h, & 1 \leq g \leq G. \end{cases} \\ & = \sum_{g=1}^G \min_{\beta^g \in \mathbb{R}^p} \sum_{i \in I^{(g)}} (y_i - \mathbf{x}_i^T \beta^g)^2 \quad \text{subject to} \quad \begin{cases} \|\beta^g\|_0 \leq t, & 1 \leq g \leq G, \\ |I^{(g)}| \geq h, & 1 \leq g \leq G. \end{cases} \end{aligned}$$

The RMSS objective function is thus separable and can be written as the sum of $G \geq 2$ independent objective functions. Also, each of these objective functions is the RBSS objective function. The pairs of solutions $(\hat{I}^{(g)}, \hat{\beta}^g)$ are all given by the RBSS solution for fixed t and h . \square

Appendix B: Finite-Sample Breakdown Point Proofs

We adapt the proof of the finite sample breakdown point of RBSS by [Thompson \(2022\)](#) to the finite-sample breakdown point of RMSS ensembles and the individual models that comprise them.

Proof of Theorem 1

Denote the matrix of coefficients for G models in an ensemble by

$$\boldsymbol{\beta}_{1:G} = \begin{pmatrix} \beta_1^1 & \beta_1^2 & \dots & \beta_1^G \\ \beta_2^1 & \beta_2^2 & \dots & \beta_2^G \\ \vdots & \vdots & \ddots & \vdots \\ \beta_p^1 & \beta_p^2 & \dots & \beta_p^G \end{pmatrix}, \quad (11)$$

where $\boldsymbol{\beta}_{1:G} \in \mathbb{R}^{p \times G}$ and β_j^g is the coefficient for predictor j in model g , $1 \leq g \leq G$. Again denote $\boldsymbol{\beta}^g = (\beta_1^g, \beta_2^g, \dots, \beta_p^g)^T \in \mathbb{R}^p$ the coefficients of model g , and also let $\boldsymbol{\beta}_j = (\beta_j^1, \beta_j^2, \dots, \beta_j^G)^T \in \mathbb{R}^G$ be the coefficients of predictor j across the G models. RMSS solves for a fixed number of models $G \geq 2$ the constrained optimization problem

$$\min_{\boldsymbol{\beta}^1, \dots, \boldsymbol{\beta}^G \in \mathbb{R}^p} \sum_{g=1}^G \sum_{i \in I^{(g)}} (y_i - \mathbf{x}_i^T \boldsymbol{\beta}^g)^2 \quad \text{subject to} \quad \begin{cases} \|\boldsymbol{\beta}^g\|_0 \leq t, & 1 \leq g \leq G, \\ \|\boldsymbol{\beta}_j\|_0 \leq u, & 1 \leq j \leq p, \\ |I^{(g)}| \geq h, & 1 \leq g \leq G. \end{cases} \quad (12)$$

where the subsets $I^{(g)} \subseteq \{1, \dots, n\}$, $1 \leq g \leq G$ indicate the subset of samples used to estimate the g -th model.

Let $T(\mathbf{X}, \mathbf{y})$ be the optimal value of the objective function of RMSS in (12). Then $T(\mathbf{X}, \mathbf{y})$ satisfies the equality

$$T(\mathbf{X}, \mathbf{y}) = \min_{I^{(1)} \subseteq \mathcal{I}} \dots \min_{I^{(G)} \subseteq \mathcal{I}} \min_{\boldsymbol{\beta}_{1:G} \in M_{1:G}} \sum_{g=1}^G \sum_{i \in I^{(g)}} (y_i - \mathbf{x}_i^T \boldsymbol{\beta}^g)^2,$$

where

$$\mathcal{I} = \{I \subseteq \{1, \dots, n\} : |I| \geq h\}, \quad \text{and} \\ M_{1:G} = \{\boldsymbol{\beta}_{1:G} \in \mathbb{R}^{p \times G} : \|\boldsymbol{\beta}^g\|_0 \leq t, \|\boldsymbol{\beta}_j\|_0 \leq u, 1 \leq g \leq G, 1 \leq j \leq p\}.$$

First assume that $n - h$ arbitrarily chosen rows of the original data (\mathbf{X}, \mathbf{y}) are replaced by contaminated observations, and denote the contaminated sample $(\mathbf{X}_c, \mathbf{y}_c)$, where the rows of $\mathbf{X}_c \in \mathbb{R}^{n \times p}$ are given by $\mathbf{x}_{i,c} \in \mathbb{R}^p$, $1 \leq i \leq n$, and $\mathbf{y}_c = (y_{1,c}, \dots, y_{n,c})^T \in \mathbb{R}^n$. Also denote I^0 the indices of the uncontaminated samples. Since $I^0 \in \mathcal{I}$, it follows that

$$\begin{aligned} T(\mathbf{X}_c, \mathbf{y}_c) &= \min_{I^{(1)} \subseteq \mathcal{I}} \dots \min_{I^{(G)} \subseteq \mathcal{I}} \min_{\beta_{1:G} \in M_{1:G}} \sum_{g=1}^G \sum_{i \in I^{(g)}} (y_{i,c} - \mathbf{x}_{i,c}^T \beta^g)^2 \\ &\leq \min_{\beta_{1:G} \in M_{1:G}} \sum_{g=1}^G \sum_{i \in I^0} (y_{i,c} - \mathbf{x}_{i,c}^T \beta^g)^2 \end{aligned}$$

Since there are no contaminated samples in the sum of the right-hand side, the latter is finite. Thus, it follows that $B(T|\mathbf{X}, \mathbf{y}) > (n - h)/n$. Now suppose that $n - h + 1$ arbitrarily chosen rows of the original data (\mathbf{X}, \mathbf{y}) are replaced by contaminated observations. Then every possible subset $I \subseteq \mathcal{I}$ includes a contaminated data point. Let this contaminated sample for model g^* be indexed by $i^* \in \{1, \dots, n\}$. Also let the pairs $(\hat{I}^{(g)}, \hat{\beta}^g)$ denote the optimal solutions to the RMSS objective function (12). Then, the optimal value of the objective function has the lower bound

$$T(\mathbf{X}_c, \mathbf{y}_c) = \sum_{g=1}^G \sum_{i \in \hat{I}^{(g)}} (y_{i,c} - \mathbf{x}_{i,c}^T \hat{\beta}^g)^2 \geq (y_{i^*,c} - \mathbf{x}_{i^*,c}^T \hat{\beta}^{g^*})^2$$

The right-hand side can be made arbitrarily large since $y_{i^*,c}$ can be made arbitrarily large, thus it follows that $B(T|\mathbf{X}, \mathbf{y}) \leq (n - h + 1)/n$. Thus, it follows that $B(T|\mathbf{X}, \mathbf{y}) = (n - h + 1)/n$. \square .

Proof of Corollary 2

Suppose again that $n - h$ arbitrarily chosen rows of the original data (\mathbf{X}, \mathbf{y}) are replaced by contaminated observations, and denote again the pairs $(\hat{I}^{(g)}, \hat{\beta}^g)$ the optimal solutions to the RMSS objective function (12). For any model g in RMSS, it follows that

$$\begin{aligned} T^g(\mathbf{X}_c, \mathbf{y}_c) &= \sum_{i \in \hat{I}^{(g)}} (y_{i,c} - \mathbf{x}_{i,c}^T \hat{\beta}^g)^2 \\ &\leq \min_{I^{(1)} \subseteq \mathcal{I}} \dots \min_{I^{(G)} \subseteq \mathcal{I}} \min_{\beta_{1:G} \in M_{1:G}} \sum_{g=1}^G \sum_{i \in I^{(g)}} (y_{i,c} - \mathbf{x}_{i,c}^T \beta^g)^2 \\ &\leq \min_{\beta_{1:G} \in M_{1:G}} \sum_{g=1}^G \sum_{i \in I^0} (y_{i,c} - \mathbf{x}_{i,c}^T \beta^g)^2 \end{aligned}$$

by Theorem 1. Thus $T^{(g)}(\mathbf{X}_c, \mathbf{y}_c)$ is finite and $B(T^{(g)}|\mathbf{X}, \mathbf{y}) > (n - h)/n$. If $n - h + 1$ arbitrarily chosen rows are contaminated, every possible subset $I \subseteq \mathcal{I}$ includes a contaminated sample. For some model g^* in RMSS, index this contaminated sample by $i^* \in \{1, \dots, n\}$. Also let $(\hat{I}^{(g^*)}, \hat{\beta}^{g^*})$ denote the pair of optimal solutions for model g^* . Then it follows that

$$T^{(g^*)}(\mathbf{X}_c, \mathbf{y}_c) = \sum_{i \in \hat{I}^{(g^*)}} \left(y_{i,c} - \mathbf{x}_{i,c}^T \hat{\beta}^{g^*} \right)^2 \geq \left(y_{i^*,c} - \mathbf{x}_{i^*,c}^T \hat{\beta}^{g^*} \right)^2$$

The right-hand side can be made arbitrarily large since $y_{i^*,c}$ can be made arbitrarily large, thus it follows that $B(T^{(g)}|\mathbf{X}, \mathbf{y}) \leq (n - h + 1)/n$, $1 \leq g \leq G$. Finally, it immediately follows that $B(T^{(g)}|\mathbf{X}, \mathbf{y}) = (n - h + 1)/n$. \square .

Appendix C: Combinatorics of RMSS

Let $h \leq n$ be the smallest sample size allowed for each model and $p_g \leq t$ the number of variables in model g , $1 \leq g \leq G$. Also let $q = \sum_{g=1}^G p_g$ and $h_j(p_1, \dots, p_G)$ be the number of elements in the sequence $\{p_1, \dots, p_G\}$ that are equal to j , $1 \leq j \leq t$. Then the total number of possible sample and predictor combinations for RMSS when $u = 1$ is given by

$$\left[\sum_{i=h}^n \binom{n}{i} \right]^G \times \sum_{p_1 \leq \dots \leq p_G \leq t} \binom{p}{q} \left[\frac{q!}{p_1! \dots p_G!} \prod_{j=1}^t \frac{1}{h_j(p_1, \dots, p_G)!} \right]. \quad (13)$$

The combinatorics result above for the particular case of $u = 1$ demonstrates that the evaluation of every possible combination of predictors in RMSS is not feasible, even if predictor variables are not allowed to be shared between models.

To illustrate the significant computational obstacle that RMSS poses, we consider the simple low-dimensional case with $n = 10$ samples, $p = 10$ predictor variables and $G = 2$ models. Setting the tuning parameters $t = 5$, $h = 5$ and $u = 1$, there are over 9 trillion unique combinations for the two models comprised of at least 5 samples and at most 5 predictor variables. Thus even when the sample size and number of predictor variables are small the evaluation of every possible combination in RMSS is not feasible. The computational obstacle of RMSS is magnified further when the tuning parameters t , h and u are chosen by CV.

Appendix D: Robust Stepwise Algorithm

In this section, we give the two main lemmas from [Khan et al. \(2007a\)](#) to show that the stepwise search algorithm can be written as a function of correlations only. Our derivations of the forward stepwise algorithm based on correlations will lead to the two lemmas originally proved by [Khan et al. \(2007a\)](#).

The Lemmas

In each step of the classical (non-robust) single-model stepwise regression algorithm, one must identify the predictor variable that reduces the residual sum of squares (RSS) by the largest amount when combined with variables already in the model, and test whether this reduction is statistically significant with respect to some threshold $\gamma \in (0, 1)$ via a partial F -test (see e.g. [Pope and Webster, 1972](#)). Let $\mathbf{X} \in \mathbb{R}^{n \times p}$ be the matrix for n samples of p predictor variables and let $\mathbf{y} \in \mathbb{R}^n$ be the vector of response variables. Denote $\hat{\mathbf{r}}_{\mathbf{y}} = (\hat{r}_{1y}, \dots, \hat{r}_{py})^T \in \mathbb{R}^p$ the vector of sample correlations between the response and predictor variables, and $\hat{\Sigma} \in \mathbb{R}^{p \times p}$ the sample correlation matrix of the design matrix \mathbf{X} . The following lemmas proved by induction in [Khan et al. \(2007a\)](#) form the basis for the robustification of stepwise regression. As a reference for our code, we include a complete proof below.

Lemma 1 *Only the original correlations $\hat{\mathbf{r}}_{\mathbf{y}}$ and $\hat{\Sigma}$ are needed to generate the sequence of variables from a stepwise regression search algorithm.*

Lemma 2 *The partial F -test at each step of a stepwise regression search algorithm for nested model comparison can be written as a function of $\hat{\mathbf{r}}_{\mathbf{y}}$ and $\hat{\Sigma}$ only.*

The stepwise algorithm can thus be robustified by replacing the classical estimates of $\hat{\mathbf{r}}_{\mathbf{y}}$ and $\hat{\Sigma}$ by their robust counterparts, which we obtain via the DDC (detecting deviating cells) method of [Rousseeuw and Bossche \(2018\)](#). The latter method can be scaled efficiently to high-dimensional settings ([Raymaekers and Rousseeuw, 2021](#)).

Derivations and Proofs

For notational convenience we denote the set $J = \{1, \dots, p\}$ and \mathbf{X}_j the j -th column of the design matrix. We assume that the variables have been centered to have mean zero and scaled to have standard deviation one. In a robust setting, the variables can be centered using the median and scaled using the MAD (median absolute deviation from the median).

Proof of Lemma 1

We first derive the first three steps of the forward stepwise algorithm to add variables to the null model using partial correlations between variables, and generalize the computations for any step of the algorithm.

- Variable $j_1 \in J$ maximizes the correlation r_{jy} ,

$$j_1 = \arg \max_{j \in J} \hat{r}_{jy},$$

so j_1 enters the model first.

- Denote the residual vector $\mathbf{Z}_{j \cdot j_1} = X_j - \beta_{jj_1} X_{j_1}$ where $\beta_{jj_1} = \hat{\Sigma}_{jj_1}$, $j \in J \setminus \{j_1\}$. Variable j_2 maximizes the (scaled) partial correlation,

$$j_2 = \arg \max_{j \in J \setminus \{j_1\}} \frac{\mathbf{Z}_{j \cdot j_1}^T \mathbf{y} / n}{\sqrt{\mathbf{Z}_{j \cdot j_1}^T \mathbf{Z}_{j \cdot j_1} / n}},$$

so j_2 enters the model second.

- Denote the residual vector $\mathbf{Z}_{j \cdot j_1 j_2} = \mathbf{Z}_{j \cdot j_1} - \beta_{jj_2 \cdot j_1} \mathbf{Z}_{j_2 \cdot j_1}$, where

$$\beta_{jj_2 \cdot j_1} = \frac{\mathbf{Z}_{j_2 \cdot j_1}^T \mathbf{Z}_{j \cdot j_1}}{\mathbf{Z}_{j_2 \cdot j_1}^T \mathbf{Z}_{j_2 \cdot j_1}}, \quad j \in J \setminus \{j_1, j_2\}.$$

Variable j_3 maximizes the (scaled) partial correlation,

$$j_3 = \arg \max_{j \in J \setminus \{j_1, j_2\}} \frac{\mathbf{Z}_{j \cdot j_1 j_2}^T \mathbf{y} / n}{\sqrt{\mathbf{Z}_{j \cdot j_1 j_2}^T \mathbf{Z}_{j \cdot j_1 j_2} / n}},$$

so j_3 enters the model third.

For steps $k \geq 3$, we can generalize the forward stepwise in the following way. Denote the residual vector $\mathbf{Z}_{j \cdot j_1 \dots j_{(k-1)}} = \mathbf{Z}_{j \cdot j_1 \dots j_{(k-2)}} - \beta_{jj_{(k-1)} \cdot j_1 \dots j_{(k-2)}} \mathbf{Z}_{j_{(k-1)} \cdot j_1 \dots j_{(k-2)}}$, where

$$\beta_{jj_{(k-1)} \cdot j_1 \dots j_{(k-2)}} = \frac{\mathbf{Z}_{j_{(k-1)} \cdot j_1 \dots j_{(k-2)}}^T \mathbf{Z}_{j \cdot j_1 \dots j_{(k-2)}}}{\mathbf{Z}_{j_{(k-1)} \cdot j_1 \dots j_{(k-2)}}^T \mathbf{Z}_{j_{(k-1)} \cdot j_1 \dots j_{(k-2)}}}, \quad j \in J \setminus \{j_1, \dots, j_{(k-1)}\}.$$

At step $k \geq 3$, variable j_k maximizes the (scaled) partial correlation,

$$j_k = \arg \max_{j \in J \setminus \{j_1, \dots, j_{(k-1)}\}} \frac{\mathbf{Z}_{j \cdot j_1 \dots j_{(k-2)}}^T \mathbf{y} / n}{\sqrt{\mathbf{Z}_{j \cdot j_1 \dots j_{(k-2)}}^T \mathbf{Z}_{j \cdot j_1 \dots j_{(k-2)}} / n}},$$

so j_k enters the model third. Thus we only need to show that for any $k \geq 2$:

1. $\mathbf{Z}_{j_k \cdot j_1 \dots j_{(k-1)}}^T \mathbf{Z}_{j_k \cdot j_1 \dots j_{(k-1)}}$, and
2. $\mathbf{Z}_{j \cdot j_1 \dots j_k}^T \mathbf{y}$

can be written as a function of $\hat{\Sigma}$ and $\hat{\mathbf{r}}_{\mathbf{y}}$ only. First, note that

$$\mathbf{Z}_{j_2 \cdot j_1}^T \mathbf{Z}_{j_2 \cdot j_1} = \left[\mathbf{X}_{j_2} - \hat{\Sigma}_{j_1 j_2} \mathbf{X}_{j_1} \right]^T \left[\mathbf{X}_j - \hat{\Sigma}_{j_1 j} \mathbf{X}_{j_1} \right] = \hat{\Sigma}_{j j_2} - \hat{\Sigma}_{j j_1} \hat{\Sigma}_{j_1 j_2}.$$

For $k \geq 2$:

$$\begin{aligned} \mathbf{Z}_{j_k \cdot j_1 \dots j_{(k-1)}}^T \mathbf{Z}_{j \cdot j_1 \dots j_{(k-1)}} &= \left[\mathbf{Z}_{j_k \cdot j_1 \dots j_{(k-2)}} - \beta_{j_k j_{(k-1)} \cdot j_1 \dots j_{(k-2)}} \mathbf{Z}_{j_{(k-1)} \cdot j_1 \dots j_{(k-2)}} \right]^T \\ &\quad \left[\mathbf{Z}_{j \cdot j_1 \dots j_{(k-2)}} - \beta_{j j_{(k-1)} \cdot j_1 \dots j_{(k-2)}} \mathbf{Z}_{j_{(k-1)} \cdot j_1 \dots j_{(k-2)}} \right] \end{aligned}$$

where

$$\beta_{j j_{(k-1)} \cdot j_1 \dots j_{(k-2)}} = \frac{\mathbf{Z}_{j_{(k-1)} \cdot j_1 \dots j_{(k-2)}}^T \mathbf{Z}_{j \cdot j_1 \dots j_{(k-2)}}}{\mathbf{Z}_{j_{(k-1)} \cdot j_1 \dots j_{(k-2)}}^T \mathbf{Z}_{j_{(k-1)} \cdot j_1 \dots j_{(k-2)}}}, \quad j \in J \setminus \{j_1, \dots, j_{(k-1)}\}.$$

If the inner products $\mathbf{Z}_{j_{(k-1)} \cdot j_1 \dots j_{(k-2)}}^T \mathbf{Z}_{j \cdot j_1 \dots j_{(k-2)}}$ can be written as a function of $\hat{\Sigma}$ and $\hat{\mathbf{r}}_{\mathbf{y}}$, then the inner products $\mathbf{Z}_{j_k \cdot j_1 \dots j_{(k-1)}}^T \mathbf{Z}_{j \cdot j_1 \dots j_{(k-1)}}$ can be written as a function of $\hat{\Sigma}$ and $\hat{\mathbf{r}}_{\mathbf{y}}$. Since $\mathbf{Z}_{j_2 \cdot j_1}^T \mathbf{Z}_{j_2 \cdot j_1} = \hat{\Sigma}_{j j_2} - \hat{\Sigma}_{j j_1} \hat{\Sigma}_{j_1 j_2}$, this proves the first part of Lemma 1. Second, note that

$$\begin{aligned} \mathbf{X}_{j_1}^T \mathbf{y} &= \hat{r}_{y j_1}, \\ \mathbf{Z}_{j \cdot j_1}^T \mathbf{y} &= \left[\mathbf{X}_j - \hat{\Sigma}_{j j_1} \mathbf{X}_{j_1} \right]^T \mathbf{y} = \hat{r}_{y j} - \hat{\Sigma}_{j j_1} \hat{r}_{y j_1} \end{aligned}$$

For $k \geq 2$,

$$\mathbf{Z}_{j \cdot j_1 \dots j_k}^T \mathbf{y} = \mathbf{Z}_{j \cdot j_1 \dots j_{(k-1)}}^T \mathbf{y} - \beta_{j j_k \cdot j_1 \dots j_{(k-1)}} \mathbf{Z}_{j_k \cdot j_1 \dots j_{(k-1)}}^T \mathbf{y}.$$

Since it was already proved that $\beta_{j j_k \cdot j_1 \dots j_{(k-1)}}$ can be written as a function of $\hat{\Sigma}$ and $\hat{\mathbf{r}}_{\mathbf{y}}$ for any $k \geq 2$, it follows that $\mathbf{Z}_{j \cdot j_1 \dots j_k}^T \mathbf{y}$ can be written as a function of $\hat{\Sigma}$ and $\hat{\mathbf{r}}_{\mathbf{y}}$ for any $k \geq 1$. This

completes the proof of Lemma 1. □

Proof of Lemma 2

The F -statistic at step $k \geq 2$ of the forward stepwise search is given by

$$F = \frac{\text{RSS}_{(k-1)} - \text{RSS}_k}{\text{RSS}_k} \times (n - k - 1),$$

where F has $\text{df}_1 = 1$ and $\text{df}_2 = k$ degrees of freedom, respectively. Note that

$$\begin{aligned} \text{RSS}_1 &= \|\mathbf{y} - \hat{r}_{yj_1} \mathbf{X}_{j_1}\|_2^2 = 1 - \hat{r}_{yj_1}, \quad \text{and} \\ \text{RSS}_k &= \left\| \mathbf{y} - \hat{r}_{yj_1} \mathbf{X}_{j_1} - \sum_{r=2}^k \beta_{yj_r \cdot j_1 \dots j_{(r-1)}} \mathbf{Z}_{j_r \cdot j_1 \dots j_{(r-1)}} \right\|_2^2, \end{aligned}$$

where

$$\beta_{yj_r \cdot j_1 \dots j_{(r-1)}} = \frac{\mathbf{Z}_{j_r \cdot j_1 \dots j_{(r-1)}}^T \mathbf{y}}{\mathbf{Z}_{j_r \cdot j_1 \dots j_{(r-1)}}^T \mathbf{Z}_{j_r \cdot j_1 \dots j_{(r-1)}}}.$$

By the orthogonality of \mathbf{X}_{j_1} and the residual vectors, it follows that

$$\begin{aligned} \text{RSS}_k &= \text{RSS}_{k-1} - \beta_{yj_k \cdot j_1 \dots j_{(k-1)}} \mathbf{Z}_{j_k \cdot j_1 \dots j_{(k-1)}}^T \mathbf{y} \\ &= \text{RSS}_{k-1} - \frac{\left(\mathbf{Z}_{j_k \cdot j_1 \dots j_{(k-1)}}^T \mathbf{y} \right)^2}{\mathbf{Z}_{j_k \cdot j_1 \dots j_{(k-1)}}^T \mathbf{Z}_{j_k \cdot j_1 \dots j_{(k-1)}}}. \end{aligned}$$

It follows from the proof of Lemma 1 that the numerator and denominator on the right-hand side can be written as a function of $\hat{\Sigma}$ and $\hat{\mathbf{r}}_{\mathbf{y}}$ only for any $k \geq 2$. This completes the proof of Lemma 2. □

Appendix E: Projected Subset Block Gradient Descent Algorithm

In this section we provide the proofs relating to the projected subset block gradient descent (PS-BGD) algorithm. First, we derive the Lipschitz constants of the gradients of the loss function \mathcal{L}_n used for each model,

$$\mathcal{L}_n(\boldsymbol{\beta}, \boldsymbol{\eta} | \mathbf{y}, \mathbf{X}) = \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta} - \boldsymbol{\eta}\|_2^2.$$

The gradients of \mathcal{L}_n with respect to $\boldsymbol{\beta}$ and $\boldsymbol{\eta}$ are given by

$$\nabla_{\boldsymbol{\beta}} \mathcal{L}_n(\boldsymbol{\beta}, \boldsymbol{\eta} | \mathbf{y}, \mathbf{X}) = 2\mathbf{X}^T (\mathbf{X}\boldsymbol{\beta} + \boldsymbol{\eta} - \mathbf{y}), \quad (14)$$

$$\nabla_{\boldsymbol{\eta}} \mathcal{L}_n(\boldsymbol{\beta}, \boldsymbol{\eta} | \mathbf{y}, \mathbf{X}) = 2(\mathbf{X}\boldsymbol{\beta} + \boldsymbol{\eta} - \mathbf{y}). \quad (15)$$

Since

$$\begin{aligned} & \left\| \nabla_{\boldsymbol{\beta}} \mathcal{L}_n(\boldsymbol{\beta}, \boldsymbol{\eta} | \mathbf{y}, \mathbf{X}) - \nabla_{\boldsymbol{\beta}} \mathcal{L}_n(\tilde{\boldsymbol{\beta}}, \boldsymbol{\eta} | \mathbf{y}, \mathbf{X}) \right\|_2 \\ &= \left\| 2\mathbf{X}^T (\mathbf{X}\boldsymbol{\beta} + \boldsymbol{\eta} - \mathbf{y}) - 2\mathbf{X}^T (\mathbf{X}\tilde{\boldsymbol{\beta}} + \boldsymbol{\eta} - \mathbf{y}) \right\|_2 \\ &= \left\| 2\mathbf{X}^T \mathbf{X} (\boldsymbol{\beta} - \tilde{\boldsymbol{\beta}}) \right\|_2 \\ &\leq \|2\mathbf{X}^T \mathbf{X}\|_2 \left\| \boldsymbol{\beta} - \tilde{\boldsymbol{\beta}} \right\|_2 \\ &= 2 \|\mathbf{X}^T \mathbf{X}\|_2 \left\| \boldsymbol{\beta} - \tilde{\boldsymbol{\beta}} \right\|_2, \end{aligned}$$

it follows that (14) is Lipschitz continuous with Lipschitz constant $\ell_{\boldsymbol{\beta}} = 2\|\mathbf{X}^T \mathbf{X}\|_2$, where $\|\mathbf{X}^T \mathbf{X}\|_2$ is the spectral norm of $\mathbf{X}^T \mathbf{X}$. Also, since

$$\begin{aligned} & \left\| \nabla_{\boldsymbol{\eta}} \mathcal{L}_n(\boldsymbol{\beta}, \boldsymbol{\eta} | \mathbf{y}, \mathbf{X}) - \nabla_{\boldsymbol{\eta}} \mathcal{L}_n(\boldsymbol{\beta}, \tilde{\boldsymbol{\eta}} | \mathbf{y}, \mathbf{X}) \right\|_2 \\ &= \left\| 2(\mathbf{X}\boldsymbol{\beta} + \boldsymbol{\eta} - \mathbf{y}) - 2(\mathbf{X}\boldsymbol{\beta} + \tilde{\boldsymbol{\eta}} - \mathbf{y}) \right\|_2 \\ &= \left\| 2(\boldsymbol{\eta} - \tilde{\boldsymbol{\eta}}) \right\|_2 \\ &\leq 2 \|\boldsymbol{\eta} - \tilde{\boldsymbol{\eta}}\|_2 \end{aligned}$$

it follows that (15) is Lipschitz continuous with Lipschitz constant $\ell_{\boldsymbol{\eta}} = 2$.

Proof of Proposition 2

We adapt the proof of Proposition 1 in [Thompson \(2022\)](#) for RBSS to RMSS and our notation. First fix any model g in RMSS and consider the update

$$\hat{\boldsymbol{\beta}}^g \in \mathcal{Q} \left(\tilde{\boldsymbol{\beta}}^g - \frac{1}{L_{\boldsymbol{\beta}}(g)} \nabla_{\boldsymbol{\beta}} \mathcal{L}_n(\boldsymbol{\beta}, \hat{\boldsymbol{\eta}}^g | \mathbf{y}, \mathbf{X}) \Big|_{\boldsymbol{\beta}=\tilde{\boldsymbol{\beta}}^g; S^{(g)}, t} \right)$$

where $L_{\boldsymbol{\beta}}(g) \geq \ell_{\boldsymbol{\beta}}(g)$ is an upper bound to the partial Lipschitz constant for model g , $\ell_{\boldsymbol{\beta}}(g) = 2\|\mathbf{X}_{S^{(g)}}^T \mathbf{X}_{S^{(g)}}\|_2$. It can be easily shown that the loss function is bounded from above by its quadratic

approximation with Lipschitz constant $\ell_{\beta^{(g)}}$ (Boyd et al., 2004),

$$\begin{aligned}\mathcal{L}_n(\boldsymbol{\beta}, \boldsymbol{\eta} | \mathbf{y}, \mathbf{X}) &\leq \mathcal{L}_n(\tilde{\boldsymbol{\beta}}, \boldsymbol{\eta} | \mathbf{y}, \mathbf{X}) + \nabla_{\boldsymbol{\beta}} \mathcal{L}_n(\boldsymbol{\beta}, \boldsymbol{\eta} | \mathbf{y}, \mathbf{X}) \Big|_{\boldsymbol{\beta}=\tilde{\boldsymbol{\beta}}}^T (\boldsymbol{\beta} - \tilde{\boldsymbol{\beta}}) + \frac{1}{2} \ell_{\boldsymbol{\beta}} \|\boldsymbol{\beta} - \tilde{\boldsymbol{\beta}}\|_2^2 \\ &= \mathcal{L}_n^Q(\boldsymbol{\beta}, \boldsymbol{\eta} | \mathbf{y}, \mathbf{X}, \tilde{\boldsymbol{\beta}}).\end{aligned}$$

Then it can be shown that

$$\begin{aligned}\mathcal{L}_n(\boldsymbol{\beta}^g, \boldsymbol{\eta}^g) &= \mathcal{L}_n^Q(\boldsymbol{\beta}^g, \boldsymbol{\eta}^g | \mathbf{y}, \mathbf{X}, \boldsymbol{\beta}^g) \\ &\geq \min_{\tilde{\boldsymbol{\beta}}^g \in \mathbb{R}^p} \mathcal{L}_n^Q(\boldsymbol{\beta}^g, \boldsymbol{\eta}^g | \mathbf{y}, \mathbf{X}, \tilde{\boldsymbol{\beta}}^g) \quad \text{subject to} \quad \begin{cases} \|\tilde{\boldsymbol{\beta}}^g\|_0 \leq t \\ \{j \in J : \tilde{\boldsymbol{\beta}}_j^g \neq 0\} \subseteq S^{(g)} \end{cases} \\ &= \min_{\tilde{\boldsymbol{\beta}}^g \in \mathbb{R}^p} \left(\mathcal{L}_n(\tilde{\boldsymbol{\beta}}^g, \boldsymbol{\eta}^g | \mathbf{y}, \mathbf{X}) - \frac{1}{2L_{\boldsymbol{\beta}^{(g)}}} \|\nabla_{\boldsymbol{\beta}} \mathcal{L}_n(\boldsymbol{\beta}, \boldsymbol{\eta}^g | \mathbf{y}, \mathbf{X})\|_2^2 + \frac{1}{2} \|\tilde{\boldsymbol{\beta}}^g - \boldsymbol{\beta}^g\|_2^2 \right) \\ &\quad \text{subject to} \quad \begin{cases} \|\tilde{\boldsymbol{\beta}}^g\|_0 \leq t \\ \{j \in J : \tilde{\boldsymbol{\beta}}_j^g \neq 0\} \subseteq S^{(g)} \end{cases} \\ &\geq \mathcal{L}_n(\boldsymbol{\beta}^g, \boldsymbol{\eta}^g) + \frac{1}{2} (L_{\boldsymbol{\beta}^{(g)}} - \ell_{\boldsymbol{\beta}^{(g)}}) \|\hat{\boldsymbol{\beta}}^g - \boldsymbol{\beta}^g\|_2^2\end{aligned}$$

Denoting and taking the estimates at step k to be $\boldsymbol{\beta}^g = \boldsymbol{\beta}^{g,(k)}$, $\hat{\boldsymbol{\beta}}^g = \hat{\boldsymbol{\beta}}^{g,(k+1)}$ and $\boldsymbol{\eta} = \boldsymbol{\eta}^{g,(k)}$ yields the inequality

$$\mathcal{L}_n(\boldsymbol{\beta}^{g,(k)}, \boldsymbol{\eta}^{g,(k)}) - \mathcal{L}_n(\boldsymbol{\beta}^{g,(k+1)}, \boldsymbol{\eta}^{g,(k)}) \geq \frac{1}{2} (L_{\boldsymbol{\beta}^{(g)}} - \ell_{\boldsymbol{\beta}^{(g)}}) \|\boldsymbol{\beta}^{g,(k+1)} - \boldsymbol{\beta}^{g,(k)}\|_2^2 \quad (16)$$

A similar derivation for the trimming parameter leads to the inequality

$$\mathcal{L}_n(\boldsymbol{\beta}^{g,(k+1)}, \boldsymbol{\eta}^{g,(k)}) - \mathcal{L}_n(\boldsymbol{\beta}^{g,(k+1)}, \boldsymbol{\eta}^{g,(k+1)}) \geq \frac{1}{2} (L_{\boldsymbol{\eta}^{(g)}} - \ell_{\boldsymbol{\eta}^{(g)}}) \|\hat{\boldsymbol{\eta}}^{g,(k+1)} - \hat{\boldsymbol{\eta}}^{g,(k)}\|_2^2 \quad (17)$$

Adding inequalities (16) and (17) together yield the new inequality

$$\begin{aligned}\mathcal{L}_n(\boldsymbol{\beta}^{g,(k)}, \boldsymbol{\eta}^{g,(k)}) - \mathcal{L}_n(\boldsymbol{\beta}^{g,(k+1)}, \boldsymbol{\eta}^{g,(k+1)}) \\ \geq \frac{1}{2} \left((L_{\boldsymbol{\beta}^{(g)}} - \ell_{\boldsymbol{\beta}^{(g)}}) \|\boldsymbol{\beta}^{g,(k+1)} - \boldsymbol{\beta}^{g,(k)}\|_2^2 + (L_{\boldsymbol{\eta}^{(g)}} - \ell_{\boldsymbol{\eta}^{(g)}}) \|\hat{\boldsymbol{\eta}}^{g,(k+1)} - \hat{\boldsymbol{\eta}}^{g,(k)}\|_2^2 \right).\end{aligned}$$

Since $\{\mathcal{L}_n(\boldsymbol{\beta}^{g,(k)}, \boldsymbol{\eta}^{g,(k)})\}$ is decreasing and $\mathcal{L}_n(\boldsymbol{\beta}^g, \boldsymbol{\eta}^g)$ is bounded from below by zero, the sequence converges. Now notice that

$$\begin{aligned} & \sum_{k=1}^K \left(\mathcal{L}_n \left(\boldsymbol{\beta}^{g,(k)}, \boldsymbol{\eta}^{g,(k)} \right) - \mathcal{L}_n \left(\boldsymbol{\beta}^{g,(k+1)}, \boldsymbol{\eta}^{g,(k+1)} \right) \right) \\ & \geq \frac{1}{2} \sum_{k=1}^K \left(\left(L_{\boldsymbol{\beta}^{(g)}} - \ell_{\boldsymbol{\beta}^{(g)}} \right) \left\| \boldsymbol{\beta}^{g,(k+1)} - \boldsymbol{\beta}^{g,(k)} \right\|_2^2 + \left(L_{\boldsymbol{\eta}^{(g)}} - \ell_{\boldsymbol{\eta}^{(g)}} \right) \left\| \hat{\boldsymbol{\eta}}^{g,(k+1)} - \hat{\boldsymbol{\eta}}^{g,(k)} \right\|_2^2 \right). \end{aligned}$$

Thus it follows that

$$\begin{aligned} & \mathcal{L}_n \left(\boldsymbol{\beta}^{g,(1)}, \boldsymbol{\eta}^{g,(1)} \right) - \mathcal{L}_n \left(\boldsymbol{\beta}^{g,(K+1)}, \boldsymbol{\eta}^{g,(K+1)} \right) \\ & \geq \frac{K}{2} \min \left(L_{\boldsymbol{\beta}^{(g)}} - \ell_{\boldsymbol{\beta}^{(g)}}, L_{\boldsymbol{\eta}^{(g)}} - \ell_{\boldsymbol{\eta}^{(g)}} \right) \min_{1 \leq k \leq K} \left(\left\| \boldsymbol{\beta}^{g,(k+1)} - \boldsymbol{\beta}^{g,(k)} \right\|_2^2 + \left\| \hat{\boldsymbol{\eta}}^{g,(k+1)} - \hat{\boldsymbol{\eta}}^{g,(k)} \right\|_2^2 \right). \end{aligned}$$

Finally, since $\{\mathcal{L}_n(\boldsymbol{\beta}^{g,(k)}, \boldsymbol{\eta}^{g,(k)})\}$ is decreasing and converges to $\mathcal{L}_n(\boldsymbol{\beta}^{g,(\infty)}, \boldsymbol{\eta}^{g,(\infty)})$,

$$\min_{1 \leq k \leq K} \left(\left\| \boldsymbol{\beta}^{g,(k+1)} - \boldsymbol{\beta}^{g,(k)} \right\|_2^2 + \left\| \hat{\boldsymbol{\eta}}^{g,(k+1)} - \hat{\boldsymbol{\eta}}^{g,(k)} \right\|_2^2 \right) \leq 2 \frac{\mathcal{L}_n(\boldsymbol{\beta}^{g,(1)}, \boldsymbol{\eta}^{g,(1)}) - \mathcal{L}_n(\boldsymbol{\beta}^{g,(\infty)}, \boldsymbol{\eta}^{g,(\infty)})}{K \min \left(L_{\boldsymbol{\beta}^{(g)}} - \ell_{\boldsymbol{\beta}^{(g)}}, L_{\boldsymbol{\eta}^{(g)}} - \ell_{\boldsymbol{\eta}^{(g)}} \right)}.$$

This completes the proof of Proposition 2 of the main article. \square

Appendix F: Neighborhood Search

In Algorithm 4 we introduce a three-dimensional neighborhood search to refine solutions generated by Algorithms 1 - 3 of the main article. For each combination of $t \in T$, $u \in U$ and $h \in H$, neighboring solutions for alternative combinations of t , u and h are used as warm starts. This process is iterated until the solutions in the grid have stabilized up to some tolerance level $\epsilon > 0$.

While Algorithm 4 yields considerable improvements in terms of minimizing the objective function, it increases the computational cost of RMSS significantly. Furthermore based on our numerical experiments, the improvements Algorithm 4 provides in terms of prediction accuracy and variable selection are marginal compared to only using Algorithms 1 - 3 of the main article.

Appendix G: Simulation Tuning Parameters

We use the $\ell_1 - \ell_2$ mixing parameter $\alpha = 3/4$ for EN, PENSE and HuberEN. To decrease the long computing time of PENSE we reduce the number of initial candidates in its computing algorithm.

Algorithm 4 Three-Dimensional Neighborhood Search

Input: Pairs $(\hat{\beta}^g[t, u, h], \hat{\eta}^g[t, u, h])$, $1 \leq g \leq G$, for all combinations of $t \in T = \{t_1, \dots, t_q\}$, $u \in U = \{1, \dots, G\}$ and $h \in H = \{h_1, \dots, h_r\}$, and tolerance parameter $\epsilon > 0$.

1: For all possible (i, j, k) where $i \in \{1, \dots, q\}$, $j \in \{1, \dots, G\}$ and $k \in \{1, \dots, r\}$:

1.1 Generate the neighborhood

$$\mathcal{N}(i, j, k) = \{a \in \{1, \dots, q\}, b \in \{1, \dots, G\}, c \in \{1, \dots, r\} : |i - a| + |j - b| + |k - c| \leq 1\}.$$

1.2 For all $(a, b, c) \in \mathcal{N}(i, j, k)$:

1.2.1 Run Algorithm 2 of the main article initialized with $(\hat{\beta}^g[t_a, u_b, h_c], \hat{\eta}^g[t_a, u_b, h_c])$, $1 \leq g \leq G$.

1.2.2 Update $(\hat{\beta}^g[t_i, u_j, h_k], \hat{\eta}^g[t_i, u_j, h_k])$, $1 \leq g \leq G$, with incumbent solution if it achieves a lower value of the objective function.

2: Repeat step 1 until ϵ -small change is achieved for the sum

$$\sum_{t \in T} \sum_{u \in U} \sum_{h \in H} \sum_{g=1}^G \left\| \mathbf{y} - \mathbf{X} \hat{\beta}^g[t, u, h] - \hat{\eta}^g[t, u, h] \right\|_2^2$$

3: Return the updated grid of solutions $\hat{\beta}^g[t, u, h]$, $1 \leq g \leq G$, for all combinations of $t \in T$, $u \in U$ and $h \in H$.

To select the level of penalty λ a candidate grid of size 100 is used for EN and a candidate grid of size 50 is used for PENSE, HuberEN and SparseLTS. The candidate grids are generated internally by the respective implementations of EN, PENSE and HuberEN. We use the `lambda0` function of the `robustHD` package to generate the log-equispaced candidate grid of size 50 for SparseLTS.

References

- Akaike, H. (1974). A new look at the statistical model identification. *IEEE Transactions on Automatic Control* 19(6), 716–723.
- Alfons, A. (2021). `robustHD`: An R package for robust regression with high-dimensional data. *Journal of Open Source Software* 6(67), 3786.
- Alfons, A., C. Croux, and S. Gelper (2013). Sparse least trimmed squares regression for analyzing high-dimensional large data sets. *The Annals of Applied Statistics*, 226–248.
- Basak, S. C. and M. Vracko (2022). *Big Data Analytics in Chemoinformatics and Bioinformatics*:

With Applications to Computer-Aided Drug Design, Cancer Biology, Emerging Pathogens and Computational Toxicology. Elsevier.

Bertsimas, D., A. King, and R. Mazumder (2016). Best subset selection via a modern optimization lens. *The Annals of Statistics* 44(2), 813–852.

Biau, G., A. Fischer, B. Guedj, and J. D. Malley (2016). Cobra: A combined regression strategy. *Journal of Multivariate Analysis* 146, 18–28.

Boyd, S., S. P. Boyd, and L. Vandenberghe (2004). *Convex optimization.* Cambridge university press.

Breiman, L. (1996a). Bagging predictors. *Machine Learning* 24(2), 123–140.

Breiman, L. (1996b). Stacked regressions. *Machine Learning* 24(1), 49–64.

Breiman, L. (2001, October). Random forests. *Machine Learning* 45(1), 5–32.

Bühlmann, P. and B. Yu (2003). Boosting with the l_2 loss: regression and classification. *Journal of the American Statistical Association* 98(462), 324–339.

Byron, S. A., K. R. Van Keuren-Jensen, D. M. Engelthaler, J. D. Carpten, and D. W. Craig (2016). Translating rna sequencing into clinical diagnostics: opportunities and challenges. *Nature Reviews Genetics* 17(5), 257–271.

Chandra, R., L. Dagum, D. Kohr, R. Menon, D. Maydan, and J. McDonald (2001). *Parallel programming in OpenMP.* Morgan Kaufmann.

Chen, T. and C. Guestrin (2016). Xgboost: A scalable tree boosting system. In *Proceedings of the 22nd acm sigkdd international conference on knowledge discovery and data mining*, pp. 785–794.

Christidis, A. and G. Cohen-Freue (2023a). *RMSS: Robust Multi-Model Subset Selection.* R package version 1.1.1.

Christidis, A. and G. Cohen-Freue (2023b). *robStepSplitReg: Robust Stepwise Split Regularized Regression.* R package version 1.1.0.

Christidis, A.-A., S. V. Aelst, and R. Zamar (2024). Multi-model subset selection. *Computational Statistics and Data Analysis.* In press.

- Christidis, A.-A., L. Lakshmanan, E. Smucler, and R. Zamar (2020). Split regularized regression. *Technometrics* 62(3), 330–338.
- Cohen Freue, G. V., D. Kepplinger, M. Salibián-Barrera, and E. Smucler (2019). Robust elastic net estimators for variable selection and identification of proteomic biomarkers.
- Donoho, D. L. and P. J. Huber (1983). The notion of breakdown point. *A festschrift for Erich L. Lehmann 157184*.
- Efron, B., T. Hastie, I. Johnstone, and R. Tibshirani (2004). Least angle regression. *The Annals of Statistics* 32(2), 407–499.
- Fan, J. and R. Li (2001). Variable selection via nonconcave penalized likelihood and its oracle properties. *Journal of the American Statistical Association* 96(456), 1348–1360.
- Friedman, J. H. (2001). Greedy function approximation: A gradient boosting machine. *Ann. Statist.* 29(5), 1189–1232.
- Friedman, J. H., T. Hastie, and R. Tibshirani (2010). Regularization paths for generalized linear models via coordinate descent. *Journal of Statistical Software* 33(1), 1.
- Garside, M. (1965). The best sub-set in multiple regression analysis. *Journal of the Royal Statistical Society: Series C (Applied Statistics)* 14(2-3), 196–200.
- Hastie, T., R. Tibshirani, and R. Tibshirani (2020). Best subset, forward stepwise or lasso? analysis and recommendations based on extensive comparisons. *Statistical Science* 35(4), 579–592.
- Hastie, T., R. Tibshirani, and M. Wainwright (2019). *Statistical learning with sparsity: the lasso and generalizations*. Chapman and Hall/CRC.
- Ho, T. K. (1998). The random subspace method for constructing decision forests. *IEEE Transactions on Pattern Analysis and Machine Intelligence* 20(8), 832–844.
- Kepplinger, D. (2023). Robust variable selection and estimation via adaptive elastic net s-estimators for linear regression. *Computational Statistics & Data Analysis* 183, 107730.
- Kepplinger, D., M. Salibián-Barrera, and G. Cohen Freue (2023). *pense: Penalized Elastic Net S/MM-Estimator of Regression*. R package version 2.2.0.
- Khan, J. A., S. Van Aelst, and R. H. Zamar (2007a). Building a robust linear model with forward selection and stepwise procedures. *Computational Statistics & Data Analysis* 52(1), 239–248.

- Khan, J. A., S. Van Aelst, and R. H. Zamar (2007b). Robust linear model selection based on least angle regression. *Journal of the American Statistical Association* 102(480), 1289–1299.
- Lemberge, P., I. De Raedt, K. H. Janssens, F. Wei, and P. J. Van Espen (2000). Quantitative analysis of 16–17th century archaeological glass vessels using pls regression of epxma and μ -xrf data. *Journal of Chemometrics: A Journal of the Chemometrics Society* 14(5-6), 751–763.
- Liaw, A. and M. Wiener (2002). Classification and regression by randomforest. *R News* 2(3), 18–22.
- Mallows, C. L. (1973). Some comments on cp. *Technometrics* 15(4), 661–675.
- Maronna, R. A. (2011). Robust ridge regression for high-dimensional data. *Technometrics* 53(1), 44–53.
- Maronna, R. A., R. D. Martin, V. J. Yohai, and M. Salibián-Barrera (2019). *Robust statistics: theory and methods (with R)*. John Wiley & Sons.
- Maronna, R. A. and R. H. Zamar (2002). Robust estimates of location and dispersion for high-dimensional datasets. *Technometrics* 44(4), 307–317.
- McCullagh, P. and J. A. Nelder (1989). Monographs on statistics and applied probability. *Generalized Linear Models* 37.
- Pope, P. and J. Webster (1972). The use of an f-statistic in stepwise regression procedures. *Technometrics* 14(2), 327–340.
- R Core Team (2022). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Raymaekers, J. and P. J. Rousseeuw (2021). Fast robust correlation for high-dimensional data. *Technometrics* 63(2), 184–198.
- Rousseeuw, P. J. (1984). Least median of squares regression. *Journal of the American statistical association* 79(388), 871–880.
- Rousseeuw, P. J. and W. V. D. Bossche (2018). Detecting deviating data cells. *Technometrics* 60(2), 135–145.
- Sangiovanni, M., I. Granata, A. S. Thind, and M. R. Guarracino (2019). From trash to treasure: detecting unexpected contamination in unmapped ngs data. *BMC bioinformatics* 20(4), 1–12.

- Scheetz, T. E., K.-Y. A. Kim, R. E. Swiderski, A. R. Philp, T. A. Braun, K. L. Knudtson, A. M. Dorrance, G. F. DiBona, J. Huang, T. L. Casavant, et al. (2006). Regulation of gene expression in the mammalian eye and its relevance to eye disease. *Proceedings of the National Academy of Sciences* 103(39), 14429–14434.
- Shen, X., W. Pan, Y. Zhu, and H. Zhou (2013). On constrained and regularized high-dimensional regression. *Annals of the Institute of Statistical Mathematics* 65(5), 807–832.
- Smucler, E. and V. J. Yohai (2017). Robust and sparse estimators for linear regression models. *Computational Statistics & Data Analysis* 111, 116–130.
- Song, L. and P. Langfelder (2013). *randomGLM: Random General Linear Model Prediction*. R package version 1.02-1.
- Song, L., P. Langfelder, and S. Horvath (2013). Random generalized linear model: a highly accurate and interpretable ensemble predictor. *BMC Bioinformatics* 14(1), 5.
- Thompson, R. (2022). Robust subset selection. *Computational Statistics & Data Analysis*, 107415.
- Tibshirani, R. (1996). Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society: Series B (Statistical Methodological)* 58(1), 267–288.
- Ueda, N. and R. Nakano (1996). Generalization error of ensemble estimators. In *Proceedings of International Conference on Neural Networks (ICNN'96)*, Volume 1, pp. 90–95. IEEE.
- Welch, W. J. (1982). Algorithmic complexity: three np-hard problems in computational statistics. *Journal of Statistical Computation and Simulation* 15(1), 17–25.
- Yi, C. (2017). *hqreg: Regularization Paths for Lasso or Elastic-Net Penalized Huber Loss Regression and Quantile Regression*. R package version 1.4.
- Yi, C. and J. Huang (2017). Semismooth newton coordinate descent algorithm for elastic-net penalized huber loss regression and quantile regression. *Journal of Computational and Graphical Statistics* 26(3), 547–557.
- Zhang, J. and K. R. Coombes (2012). Sources of variation in false discovery rate estimation include sample size, correlation, and inherent differences between groups. *BMC Bioinformatics* 13(S13), S1.

Zhu, J., X. Wang, L. Hu, J. Huang, K. Jiang, Y. Zhang, S. Lin, and J. Zhu (2022). abess: a fast best-subset selection library in python and r. *The Journal of Machine Learning Research* 23(1), 9206–9212.

Zou, H. and T. Hastie (2005). Regularization and variable selection via the elastic net. *Journal of the Royal Statistical Society: Series B (Statistical Methodological)* 67(2), 301–320.