

LIOUVILLE THEOREM FOR QUASILINEAR ELLIPTIC EQUATIONS IN \mathbb{R}^N

WANGZHE WU AND QIQI ZHANG

ABSTRACT. We prove Liouville theorem for the equation $\Delta_m v + v^p + M|\nabla v|^q = 0$ in a domain $\Omega \subset \mathbb{R}^n$, with $M \in \mathbb{R}$ in the critical and subcritical case. As a natural extension of our recent work [14], the proof is based on an integral identity and Young's inequality.

1. INTRODUCTION

In this paper, we are concerned with a Liouville type theorem for the global solutions of the following equations in \mathbb{R}^n :

$$\Delta_m v + v^p + M|\nabla v|^q = 0, \quad \text{in } \Omega, \quad (1)$$

where $q = \frac{mp}{p+1}$, $\Delta_m v = \operatorname{div}(|\nabla v|^{m-2}\nabla v)$, p and q are exponents larger than 1, $M \in \mathbb{R}$. This kind of equations are widely concerned and studied depending on the sign and value of M . The most important motivation of the present study is to extend the results obtained for the semilinear equation

$$\Delta v + v^p + M|\nabla v|^q = 0, \quad \text{in } \Omega, \quad (2)$$

in [4, 5]. By using a delicate combination of refined Bernstein techniques and Keller-Osserman estimate, they obtained a series of a priori estimates for any positive solution of (2) in arbitrary domain of \mathbb{R}^n .

Theorem 1.1 (Theorem E in [4]). *Let $n \geq 3$, $1 < p < \frac{n+2}{n-2}$, $q = \frac{2p}{p+1}$. Then there exists $\epsilon_0 > 0$ depending on n and p such that for any $|M| \leq \epsilon_0$, there exists no nontrivial nonnegative solution of (2) in \mathbb{R}^n .*

It's obviously to see that the result in [4] did not cover all the cases that $m = 2$, $1 < p \leq \frac{n+2}{n-2}$ for $q = \frac{2p}{p+1}$ when $M > 0$. Thus, our recent work [14] came out, which complete the corresponding results of the range of p with $1 < p \leq \frac{n+2}{n-2}$ when $M > 0$, based on an integral identity and Young's inequality. For $m = 2$, $M < 0$, in [5] they gave some Liouville theorems for the equation (2) when v is radial. In addition, for general case $1 < m < p$, there are also parts of results varies as M changes. If $M > 0$, Filippucci-Sun-Zheng [10] derived various a priori estimate concerning $|\nabla v|$ for positive solutions of (1) in the cases q is less, greater or equal to $\frac{mp}{p+1}$, and consequently obtain Liouville type theorems. For the case $q = \frac{mp}{p+1}$ and M large enough, they got the following non-existence result in \mathbb{R}^n .

Theorem 1.2 (Theorem 1.4 in [10]). *Let $\Omega \subset \mathbb{R}^n$, $p > \max\{m - 1, 1\}$ and $q = \frac{mp}{p+1}$. Then for any*

$$M > \frac{\sqrt{n}(p+1)}{(4p)^{\frac{p}{p+1}}} \left(\frac{p-1}{\sqrt{a}} \right)^{\frac{p-1}{p+1}},$$

where $0 < a \leq 1/n$, (1) does not admit positive solutions in \mathbb{R}^n .

If $M < 0$, especially $M = -1$, which has been introduced by Chipot-Weissler [8] in 1989 and obtained the existence of ground states when the parameter values p and q satisfy $1 < q < \frac{2p}{p+1}$, $p > 1$. we recall that a ground states for (1) or (2) is a non-negative non-trivial entire solution. It's known that radial ground states for (2) always exist for the supercritical parameter range $p > (n+2)/(n-2)$ and may or may not exist when p is critical or subcritical, depending on the value of q . Turning to the case of ground states for equation (1) with $M = -1$, Serrin-Zou [20] showed that existence always holds when p is supercritical, that is $p > \frac{(m-1)n+m}{n-m}$. For critical p , existence of radial ground states holds if and only if

$$\begin{cases} 0 < q < (m-1)\frac{(m-1)n+m}{(m-1)n-m}, & \text{if } m > \frac{n}{n-1}; \\ q > 0, & \text{if } m < \frac{n}{n-1}. \end{cases}$$

For subcritical p , existence holds when $q < \frac{mp}{p+1}$. Besides, when M is allowed to be other negative constant, Filippucci-Sun-Zheng [10] derived a non-existence result for supersolutions of (1) in an exterior domain.

Theorem 1.3 (Theorem 1.5 in [10]). *Let $p > m - 1$ if $n = m$ or $m - 1 < p < \frac{n(m-1)}{n-m}$ if $n > m$, $q = \frac{mp}{p+1}$ and $M > -\nu^*(n)$ where*

$$\nu^*(n) := (p+1) \left(\frac{n(m-1) - p(n-m)}{mp} \right)^{\frac{p}{p+1}}.$$

Then there exist no nontrivial nonnegative supersolutions of (1) in $\mathbb{R}^n \setminus \bar{B}_R$ for any $R > 0$.

If $M = 0$, then (1) reduces to the generalized Lane-Emden equation

$$\Delta_m v + v^p = 0, \quad \text{in } \Omega, \tag{3}$$

which has been widely studied in the literature [1, 2, 6, 7, 9, 11, 12, 16, 17, 18, 21, 22], both when Ω is bounded and unbounded. Especially, in the semilinear case $m = 2$, one of the celebrated results is given by Gidas and Spruck [11]: if $n > 2$, and $p \in \left[1, \frac{n+2}{n-2}\right)$, then any nonnegative solution of (3) is identically zero and the result is sharp. For the case of $m > 1$, radially symmetric positive solutions were studied by Ni and Serrin [15, 16, 17], and further results in this direction were obtained by Guedda and Véron [12] and Bidaut-Véron [1]. When one studies the so called Liouville property of (3), namely whether all positive C^1 solutions of (3) in \mathbb{R}^n are constant, two critical exponents appear

$$m_* = \frac{n(m-1)}{n-m}, \quad m^* = \frac{n(m-1) + m}{n-m},$$

where $n > m$, known as the Serrin exponent and the Sobolev exponent, respectively.

For the case only with gradient terms ($M \neq 0$), we first recall the Hamilton-Jacobi equation

$$\Delta_m v + |\nabla v|^q = 0, \quad \text{in } \Omega, \quad (4)$$

The Liouville property of (4) was studied by Lions in [13] for special case $m = 2$, who proved that any C^2 solution to (4) with $q > 1$ has to be a constant by using the Bernstein technique. In [8] and [19], most of the study deals with the case $q \neq \frac{2p}{p+1}$. In the critical case, then not only the sign of M but also its value plays a fundamental role, with a delicate interaction with the exponent p . Other work on this kind of equation can be seen in [3]. For general case, Bidaut-Véron, Garcia-Huidobro and Véron [3] proved that any C^1 solution of (4) in an arbitrary domain Ω with $n \geq m > 1$ and $q > m - 1$ satisfies

$$|\nabla v(x)| \leq c_{n,m,q} (\text{dist}(x, \partial\Omega))^{-\frac{1}{q-m+1}}.$$

Estimates of this type, not only for the gradient but also for the solutions are called by Serrin and Zou "universal a priori estimates", because they are independent of the solutions and do not need any boundary conditions.

In this paper, we continue the idea used in our recent work [14] to derive Liouville type theorems for positive solutions of (1) in the case $q = \frac{mp}{p+1}$ varies in the sign of M , that produces the following results.

Theorem 1.4. *Let $n \geq 2$, $m - 1 < p \leq \frac{(m-1)n+m}{n-m}$, $q = \frac{mp}{p+1}$, then there exists $M_1 > 0$, which depends on n, m, p , such that for any $0 < M < M_1$, all the nonnegative solutions of (1) are $v \equiv 0$.*

Theorem 1.5. *Let $n \geq 2$, $m - 1 < p < \frac{(m-1)n+m}{n-m}$, $q = \frac{mp}{p+1}$, then there exists $M_2 > 0$, which depends on n, m, p , such that for any $-M_2 < M < 0$, all the nonnegative solutions of (1) are $v \equiv 0$.*

The second theorem extends the result in [10] from $m - 1 < p < \frac{n(m-1)}{n-m}$ to $m - 1 < p < \frac{(m-1)n+m}{n-m}$. But when $m - 1 < p < \frac{n(m-1)}{n-m}$, we don't know whether $M_2 > \nu^*(n)$.

2. INTEGRAL IDENTITY

Multiplying (1) by $v^\alpha \Delta_m v$,

$$\begin{aligned} v^\alpha (\Delta_m v)^2 &= -v^{\alpha+p} \underset{I}{\Delta_m v} - M v^\alpha \underset{II}{|\nabla v|^q} \Delta_m v. \\ \textcircled{1} \end{aligned} \quad (5)$$

We observed that the term ① is important. Denote

$$\begin{aligned} X^i &= |\nabla v|^{m-2} v_i, \\ \Delta_m v &= (|\nabla v|^{m-2} v_i)_i, \\ X_j^i &= (|\nabla v|^{m-2} v_i)_j, \\ E_j^i &= X_j^i - \frac{\Delta_m v}{n} \delta_{ij}, \end{aligned}$$

then we will do some computations on some important terms as follows.

2.1. The term $v^{\alpha-1}|\nabla v|^m \Delta_m v$.

$$\begin{aligned}
 v^{\alpha-1}|\nabla v|^m \Delta_m v &= v^{\alpha-1}|\nabla v|^m (|\nabla v|^{m-2} v_i)_i \\
 &= (v^{\alpha-1}|\nabla v|^{2m-2} v_i)_i - (\alpha-1)v^{\alpha-2}|\nabla v|^{2m} - m v^{\alpha-1}|\nabla v|^{2m-4} v_i v_j v_{ij}, \\
 &\quad \text{III} \tag{6}
 \end{aligned}$$

here

$$\begin{aligned}
 \text{III} &= v^{\alpha-1}|\nabla v|^{2m-4} v_i v_j v_{ij} \\
 &= v^{\alpha-1}|\nabla v|^{m-2} v_i v_j (|\nabla v|^{m-2} v_{ij}) \\
 &= v^{\alpha-1}|\nabla v|^{m-2} v_i v_j \left[(|\nabla v|^{m-2} v_i)_j - (m-2)|\nabla v|^{m-4} v_i v_\ell v_{\ell j} \right] \\
 &= v^{\alpha-1}|\nabla v|^{m-2} v_i v_j X_j^i - (m-2)v^{\alpha-1}|\nabla v|^{2m-4} v_i v_j v_{ij}, \tag{7}
 \end{aligned}$$

therefore,

$$\begin{aligned}
 \text{III} &= \frac{1}{m-1} v^{\alpha-1}|\nabla v|^{m-2} v_i v_j X_j^i \\
 &= \frac{1}{m-1} v^{\alpha-1}|\nabla v|^{m-2} v_i v_j (E_j^i + \frac{\Delta_m v}{n} \delta_{ij}) \\
 &= \frac{1}{m-1} v^{\alpha-1}|\nabla v|^{m-2} v_i v_j E_j^i + \frac{1}{n(m-1)} v^{\alpha-1}|\nabla v|^m \Delta_m v. \tag{8}
 \end{aligned}$$

Substitute (8) into (6), we obtain

$$\begin{aligned}
 &v^{\alpha-1}|\nabla v|^m \Delta_m v \\
 &= \frac{n(m-1)}{n(m-1)+m} \cdot \left[(v^{\alpha-1}|\nabla v|^{2m-2} v_i)_i - (\alpha-1)v^{\alpha-2}|\nabla v|^{2m} - \frac{m}{m-1} v^{\alpha-1}|\nabla v|^{m-2} v_i v_j E_j^i \right]. \tag{9}
 \end{aligned}$$

2.2. The term ①.

In fact, we have

$$\begin{aligned}
 v^\alpha (\Delta_m v)^2 &= v^\alpha X_i^i X_j^j \\
 &= (v^\alpha X_i^i X^j)_j - \alpha v^{\alpha-1} X_i^i X^j v_j - v^\alpha X^j (X_i^i)_j \\
 &= (v^\alpha X_i^i X^j)_j - \alpha v^{\alpha-1} |\nabla v|^m \Delta_m v - \left[(v^\alpha X^j X_i^i)_i - (v^\alpha X^j)_i X_j^i \right] \\
 &= (v^\alpha X_i^i X^j)_j - \alpha v^{\alpha-1} |\nabla v|^m \Delta_m v - \left[(v^\alpha X^j X_i^i)_i - \alpha v^{\alpha-1} v_i X^j X_j^i - v^\alpha X_j^i X_i^j \right] \\
 &= \left(v^\alpha \Delta_m v X^i - v^\alpha X^j X_j^i \right)_i - \alpha v^{\alpha-1} |\nabla v|^m \Delta_m v + \alpha v^{\alpha-1} v_i X^j X_j^i + v^\alpha X_j^i X_i^j.
 \end{aligned}$$

Since $X_j^i = E_j^i + \frac{\Delta_m v}{n} \delta_{ij}$, then

$$\begin{aligned}
 v^\alpha (\Delta_m v)^2 &= (1 - \frac{1}{n})(v^\alpha \Delta_m v X^i)_i - (v^\alpha E_j^i X^j)_i - \alpha v^{\alpha-1} |\nabla v|^m \Delta_m v \\
 &\quad + \alpha v^{\alpha-1} v_i X^j E_j^i + \frac{\alpha}{n} v^{\alpha-1} |\nabla v|^m \Delta_m v + v^\alpha E_j^i E_i^j + \frac{1}{n} v^\alpha (\Delta_m v)^2,
 \end{aligned}$$

\Rightarrow

$$(1 - \frac{1}{n})v^\alpha(\Delta_m v)^2 = (1 - \frac{1}{n})(v^\alpha \Delta_m v X^i)_i - (v^\alpha E_j^i X^j)_i - (1 - \frac{1}{n})\alpha v^{\alpha-1} |\nabla v|^m \Delta_m v + \alpha v^{\alpha-1} v_i X^j E_j^i + v^\alpha E_j^i E_i^j.$$

As a result, we obtain

$$\begin{aligned} ① &= v^\alpha(\Delta_m v)^2 \\ &= (v^\alpha \Delta_m v X^i)_i - \frac{n}{n-1}(v^\alpha E_j^i X^j)_i - \alpha v^{\alpha-1} |\nabla v|^m \Delta_m v + \frac{n}{n-1} \alpha v^{\alpha-1} |\nabla v|^{m-2} E_j^i v_i v_j + \frac{n}{n-1} v^\alpha E_j^i E_i^j \\ &= (v^\alpha \Delta_m v X^i)_i - \frac{n}{n-1}(v^\alpha E_j^i X^j)_i + \frac{n}{n-1} v^\alpha E_j^i E_i^j + \frac{n}{n-1} \alpha v^{\alpha-1} |\nabla v|^{m-2} E_j^i v_i v_j \\ &\quad - \alpha \frac{n(m-1)}{n(m-1)+m} \left[(v^{\alpha-1} |\nabla v|^{2m-2} v_i)_i - (\alpha-1) v^{\alpha-2} |\nabla v|^{2m} - \frac{m}{m-1} v^{\alpha-1} |\nabla v|^{m-2} v_i v_j E_j^i \right] \\ &= (v^\alpha \Delta_m v X^i)_i - \frac{n}{n-1} (v^\alpha E_j^i X^j)_i - \alpha \frac{n(m-1)}{n(m-1)+m} (v^{\alpha-1} |\nabla v|^{2m-2} v_i)_i + \frac{n}{n-1} v^\alpha E_j^i E_i^j \\ &\quad + \alpha(\alpha-1) \frac{n(m-1)}{n(m-1)+m} v^{\alpha-2} |\nabla v|^{2m} \\ &\quad + \alpha \left[\frac{n}{n-1} + \frac{nm}{n(m-1)+m} \right] v^{\alpha-1} |\nabla v|^{m-2} E_j^i v_i v_j. \end{aligned} \tag{10}$$

2.3. The term A .

$$\begin{aligned} A &= (v^\alpha X^i \Delta_m v)_i \\ &= \alpha v^{\alpha-1} |\nabla v|^m \Delta_m v + v^\alpha (\Delta_m v)^2 + v^\alpha X^i \underset{A_1}{(\Delta_m v)_i}. \end{aligned}$$

Substitute (1) into A_1 ,

$$\begin{aligned} A_1 &= v^\alpha X^i \left(-v^p - M |\nabla v|^q \right)_i \\ &= -p v^{\alpha+p-1} |\nabla v|^m - q M v^\alpha |\nabla v|^{m+q-4} v_i v_j v_{ij}. \end{aligned}$$

Multiply (1) by $v^{\alpha-1} |\nabla v|^m$

$$-v^{\alpha+p-1} |\nabla v|^m = v^{\alpha-1} |\nabla v|^m \Delta_m v + M v^{\alpha-1} |\nabla v|^{m+q}. \tag{11}$$

Therefore, we have

$$A_1 = p v^{\alpha-1} |\nabla v|^m \Delta_m v + p M v^{\alpha-1} |\nabla v|^{m+q} - q M v^\alpha |\nabla v|^{m+q-4} v_i v_j v_{ij}. \tag{12}$$

On the other hand, we know

$$\begin{aligned}
 -qMv^\alpha|\nabla v|^{m+q-4}v_i v_j v_{ij} &= -Mv^\alpha(|\nabla v|^q)_i |\nabla v|^{m-2} v_i \\
 &= -Mv^\alpha(|\nabla v|^q)_i X^i \\
 &= -M \left[(v^\alpha |\nabla v|^q X^i)_i - |\nabla v|^q (v^\alpha X^i)_i \right] \\
 &= -M(v^\alpha |\nabla v|^q X^i)_i + \alpha Mv^{\alpha-1} |\nabla v|^{m+q} + Mv^\alpha |\nabla v|^q \Delta_m v \\
 &= -M(v^\alpha |\nabla v|^q X^i)_i + \alpha Mv^{\alpha-1} |\nabla v|^{m+q} + Mv^\alpha |\nabla v|^q \left(-v^p - M|\nabla v|^q \right) \\
 &= -M(v^\alpha |\nabla v|^q X^i)_i + \alpha Mv^{\alpha-1} |\nabla v|^{m+q} - Mv^{\alpha+p} |\nabla v|^q - M^2 v^\alpha |\nabla v|^{2q}.
 \end{aligned} \tag{13}$$

\Rightarrow the last two terms of (12) is

$$\begin{aligned}
 &pMv^{\alpha-1}|\nabla v|^{m+q} - qMv^\alpha|\nabla v|^{m+q-4}v_i v_j v_{ij} \\
 &= -M(v^\alpha |\nabla v|^q X^i)_i + (p + \alpha) Mv^{\alpha-1} |\nabla v|^{m+q} - Mv^{\alpha+p} |\nabla v|^q - M^2 v^\alpha |\nabla v|^{2q}.
 \end{aligned} \tag{14}$$

So we get that A_1 and

$$\begin{aligned}
 A &= (v^\alpha X^i \Delta_m v)_i \\
 &= (p + \alpha)v^{\alpha-1} |\nabla v|^m \Delta_m v + v^\alpha (\Delta_m v)^2 \\
 &\quad - M(v^\alpha |\nabla v|^q X^i)_i + (p + \alpha) Mv^{\alpha-1} |\nabla v|^{m+q} - Mv^{\alpha+p} |\nabla v|^q - M^2 v^\alpha |\nabla v|^{2q}.
 \end{aligned}$$

Substitute (9) into it:

$$\begin{aligned}
 &-v^\alpha (\Delta_m v)^2 \\
 &= -A + (\alpha + p) \cdot \frac{n(m-1)}{n(m-1) + m} \left[(v^{\alpha-1} |\nabla v|^{2m-2} v_i)_i - (\alpha - 1)v^{\alpha-2} |\nabla v|^{2m} \right. \\
 &\quad \left. - \frac{m}{m-1} v^{\alpha-1} |\nabla v|^{m-2} v_i v_j E_j^i \right] \\
 &\quad - M(v^\alpha |\nabla v|^q X^i)_i + (p + \alpha) Mv^{\alpha-1} |\nabla v|^{m+q} - Mv^{\alpha+p} |\nabla v|^q - M^2 v^\alpha |\nabla v|^{2q}.
 \end{aligned} \tag{15}$$

2.4. The final equation. Combine (10) and (15) to delete $v^\alpha(\Delta_m v)^2$, we have

$$\begin{aligned}
0 = & -A + (\alpha + p) \cdot \frac{n(m-1)}{n(m-1)+m} \left[(v^{\alpha-1} |\nabla v|^{2m-2} v_i)_i - (\alpha - 1) v^{\alpha-2} |\nabla v|^{2m} \right. \\
& - \frac{m}{m-1} v^{\alpha-1} |\nabla v|^{m-2} v_i v_j E_j^i \Big] \\
& - M(v^\alpha |\nabla v|^q X^i)_i + (p + \alpha) M v^{\alpha-1} |\nabla v|^{m+q} - M v^{\alpha+p} |\nabla v|^q - M^2 v^\alpha |\nabla v|^{2q} \\
& + (v^\alpha \Delta_m v X^i)_i - \frac{n}{n-1} (v^\alpha E_j^i X^j)_i - \alpha \frac{n(m-1)}{n(m-1)+m} (v^{\alpha-1} |\nabla v|^{2m-2} v_i)_i + \frac{n}{n-1} v^\alpha E_j^i E_i^j \\
& + \alpha(\alpha - 1) \frac{n(m-1)}{n(m-1)+m} v^{\alpha-2} |\nabla v|^{2m} \\
& + \alpha \left[\frac{n}{n-1} + \frac{nm}{n(m-1)+m} \right] v^{\alpha-1} |\nabla v|^{m-2} E_j^i v_i v_j,
\end{aligned}$$

that is

$$\begin{aligned}
0 = & -A + (\alpha + p) \cdot \frac{n(m-1)}{n(m-1)+m} (v^{\alpha-1} |\nabla v|^{2m-2} v_i)_i - M(v^\alpha |\nabla v|^q X^i)_i \\
& + (v^\alpha \Delta_m v X^i)_i - \frac{n}{n-1} (v^\alpha E_j^i X^j)_i - \alpha \frac{n(m-1)}{n(m-1)+m} (v^{\alpha-1} |\nabla v|^{2m-2} v_i)_i \\
& + \frac{n}{n-1} v^\alpha E_j^i E_i^j - p(\alpha - 1) \frac{n(m-1)}{n(m-1)+m} v^{\alpha-2} |\nabla v|^{2m} \\
& + \left[\frac{n\alpha}{n-1} - \frac{npm}{n(m-1)+m} \right] v^{\alpha-1} |\nabla v|^{m-2} v_i v_j E_j^i \\
& + (p + \alpha) M v^{\alpha-1} |\nabla v|^{m+q} - M v^{\alpha+p} |\nabla v|^q - M^2 v^\alpha |\nabla v|^{2q}.
\end{aligned}$$

Now we can integrate it into the following form:

$$\begin{aligned}
0 = & W + \frac{n}{n-1} v^\alpha E_j^i E_i^j - p(\alpha - 1) \frac{n(m-1)}{n(m-1)+m} v^{\alpha-2} |\nabla v|^{2m} \\
& + \left[\frac{n\alpha}{n-1} - \frac{npm}{n(m-1)+m} \right] v^{\alpha-1} |\nabla v|^{m-2} v_i v_j E_j^i + (p + \alpha) M v^{\alpha-1} |\nabla v|^{m+q} \\
& - M v^{\alpha+p} |\nabla v|^q - M^2 v^\alpha |\nabla v|^{2q},
\end{aligned} \tag{16}$$

where W consists of all the divergence terms.

3. CONDITIONS

In order to obtain the Liouville theorem, we need some extra conditions and we also insert coefficients T, U, P for more detailed computations.

Multiply (1) by $v^{\alpha+p}$:

$$\begin{aligned}
v^{\alpha+2p} + M v^{\alpha+p} |\nabla v|^q &= -v^{\alpha+p} \Delta_m v \\
&= -v^{\alpha+p} (|\nabla v|^{m-2} v_i)_i \\
&= -(v^{\alpha+p} |\nabla v|^{m-2} v_i)_i + (\alpha + p) v^{\alpha+p-1} |\nabla v|^m.
\end{aligned} \tag{17}$$

Multiply (1) by $v^{\alpha-1}|\nabla v|^m$:

$$\begin{aligned} & -Mv^{\alpha-1}|\nabla v|^{m+q} - v^{\alpha+p-1}|\nabla v|^m \\ & = v^{\alpha-1}|\nabla v|^m \Delta_m v \\ & = \frac{n(m-1)}{n(m-1)+m} \left[(v^{\alpha-1}|\nabla v|^{2m-2}v_i)_i - (\alpha-1)v^{\alpha-2}|\nabla v|^{2m} - \frac{m}{m-1}v^{\alpha-1}|\nabla v|^{m-2}v_i v_j E_j^i \right]. \end{aligned}$$

Multiply (1) by $Mv^\alpha|\nabla v|^q$:

$$Mv^{\alpha+p}|\nabla v|^q + M^2v^\alpha|\nabla v|^{2q} = -Mv^\alpha|\nabla v|^q \Delta_m v, \quad (18)$$

as we can see that

$$\begin{aligned} X_j^i v_i v_j &= (|\nabla v|^{m-2}v_i)_j v_i v_j \\ &= (m-2)|\nabla v|^{m-4}v_{\ell j}v_\ell v_i v_i v_j + |\nabla v|^{m-2}v_{ij}v_i v_j \\ &= (m-1)|\nabla v|^{m-2}v_{ij}v_i v_j, \end{aligned} \quad (19)$$

if we insert (19) into (18), we obtain

$$\begin{aligned} v^\alpha|\nabla v|^q \Delta_m v &= (v^\alpha|\nabla v|^q X^i)_i - \alpha v^{\alpha-1}|\nabla v|^{m+q} - q v^\alpha|\nabla v|^{m+q-4}v_{ij}v_i v_j \\ &= (v^\alpha|\nabla v|^q X^i)_i - \alpha v^{\alpha-1}|\nabla v|^{m+q} - \frac{q}{m-1}v^\alpha|\nabla v|^{q-2}X_j^i v_i v_j \\ &= (v^\alpha|\nabla v|^q X^i)_i - \alpha v^{\alpha-1}|\nabla v|^{m+q} - \frac{q}{m-1}v^\alpha|\nabla v|^{q-2}E_j^i v_i v_j - \frac{q}{n(m-1)}v^\alpha|\nabla v|^q \Delta_m v, \end{aligned}$$

$$\begin{aligned} \Rightarrow v^\alpha|\nabla v|^q \Delta_m v &= \frac{n(m-1)}{n(m-1)+q}(v^\alpha|\nabla v|^q X^i)_i - \frac{n(m-1)\alpha}{n(m-1)+q}v^{\alpha-1}|\nabla v|^{m+q} \\ &\quad - \frac{nq}{n(m-1)+q}v^\alpha|\nabla v|^{q-2}E_j^i v_i v_j, \end{aligned} \quad (20)$$

so (18) turns to be

$$\begin{aligned} & Mv^{\alpha+p}|\nabla v|^q + M^2v^\alpha|\nabla v|^{2q} \\ &= -M \frac{n(m-1)}{n(m-1)+q}(v^\alpha|\nabla v|^q X^i)_i + \frac{n(m-1)\alpha}{n(m-1)+q}Mv^{\alpha-1}|\nabla v|^{m+q} \\ &\quad + \frac{nq}{n(m-1)+q}Mv^\alpha|\nabla v|^{q-2}E_j^i v_i v_j. \end{aligned} \quad (21)$$

After that, we now rewrite (16) as follows

$$\begin{aligned}
0 = & W + \frac{n}{n-1} v^\alpha E_j^i E_i^j - p(\alpha-1) \frac{n(m-1)}{n(m-1)+m} v^{\alpha-2} |\nabla v|^{2m} \\
& + \left[\frac{n\alpha}{n-1} - \frac{npm}{n(m-1)+m} \right] v^{\alpha-1} |\nabla v|^{m-2} v_i v_j E_j^i \\
& + (p+\alpha) M v^{\alpha-1} |\nabla v|^{m+q} - M v^{\alpha+p} |\nabla v|^q - M^2 v^\alpha |\nabla v|^{2q} \\
& + T \left[v^{\alpha+2p} + M v^{\alpha+p} |\nabla v|^q - (\alpha+p) v^{\alpha+p-1} |\nabla v|^m \right] \\
& + U \left[M v^{\alpha+p} |\nabla v|^q + M^2 v^\alpha |\nabla v|^{2q} - \frac{n(m-1)\alpha}{n(m-1)+q} M v^{\alpha-1} |\nabla v|^{m+q} \right. \\
& \quad \left. - \frac{nq}{n(m-1)+q} M v^\alpha |\nabla v|^{q-2} E_j^i v_i v_j \right] \\
& + P \left[-M v^{\alpha-1} |\nabla v|^{m+q} - v^{\alpha+p-1} |\nabla v|^m + \frac{n(m-1)(\alpha-1)}{n(m-1)+m} v^{\alpha-2} |\nabla v|^{2m} \right. \\
& \quad \left. + \frac{nm}{n(m-1)+m} v^{\alpha-1} |\nabla v|^{m-2} v_i v_j E_j^i \right], \tag{22}
\end{aligned}$$

that is

$$\begin{aligned}
0 = & W + \frac{n}{n-1} v^\alpha E_j^i E_i^j + \left[\frac{n\alpha}{n-1} - \frac{nm(p-P)}{n(m-1)+m} \right] v^{\alpha-1} |\nabla v|^{m-2} v_i v_j E_j^i \\
& - \frac{nqU}{n(m-1)+q} M v^\alpha |\nabla v|^{q-2} E_j^i v_i v_j - (p-P)(\alpha-1) \frac{n(m-1)}{n(m-1)+m} v^{\alpha-2} |\nabla v|^{2m} \\
& + T v^{\alpha+2p} + (U-1) M^2 v^\alpha |\nabla v|^{2q} + \left[-(\alpha+p)T - P \right] v^{\alpha+p-1} |\nabla v|^m \\
& + \left[\alpha + p - P - \frac{n(m-1)\alpha U}{n(m-1)+q} \right] M v^{\alpha-1} |\nabla v|^{m+q} + (T+U-1) M v^{\alpha+p} |\nabla v|^q.
\end{aligned}$$

Define

$$L_{ij} := v_i v_j - \frac{1}{n} |\nabla v|^2 \delta_{ij},$$

then

$$|L_{ij}|^2 = \frac{n-1}{9^n} |\nabla v|^4,$$

and

$$\begin{aligned}
 0 &\geq W - \frac{n-1}{4n}v^\alpha \left\{ \left[\frac{n\alpha}{n-1} - \frac{nm(p-P)}{n(m-1)+m} \right] v^{-1} |\nabla v|^{m-2} L_{ij} - \frac{nqU}{n(m-1)+q} M |\nabla v|^{q-2} L_{ij} \right\}^2 \\
 &\quad - (p-P)(\alpha-1) \frac{n(m-1)}{n(m-1)+m} v^{\alpha-2} |\nabla v|^{2m} \\
 &\quad + T v^{\alpha+2p} + (U-1) M^2 v^\alpha |\nabla v|^{2q} + \left[-(\alpha+p)T - P \right] v^{\alpha+p-1} |\nabla v|^m \\
 &\quad + \left[\alpha + p - P - \frac{n(m-1)\alpha U}{n(m-1)+q} \right] M v^{\alpha-1} |\nabla v|^{m+q} + (T+U-1) M v^{\alpha+p} |\nabla v|^q \\
 &= W - \frac{(n-1)^2}{4n^2} v^\alpha |\nabla v|^4 \left\{ \left[\frac{n\alpha}{n-1} - \frac{nm(p-P)}{n(m-1)+m} \right] v^{-1} |\nabla v|^{m-2} - \frac{nqU}{n(m-1)+q} M |\nabla v|^{q-2} \right\}^2 \\
 &\quad - (p-P)(\alpha-1) \frac{n(m-1)}{n(m-1)+m} v^{\alpha-2} |\nabla v|^{2m} \\
 &\quad + T v^{\alpha+2p} + (U-1) M^2 v^\alpha |\nabla v|^{2q} + \left[-(\alpha+p)T - P \right] v^{\alpha+p-1} |\nabla v|^m \\
 &\quad + \left[\alpha + p - P - \frac{n(m-1)\alpha U}{n(m-1)+q} \right] M v^{\alpha-1} |\nabla v|^{m+q} + (T+U-1) M v^{\alpha+p} |\nabla v|^q.
 \end{aligned}$$

After integration,

$$\begin{aligned}
 RHS &= W + \left\{ - (p-P)(\alpha-1) \frac{n(m-1)}{n(m-1)+m} - \frac{(n-1)^2}{4n^2} \left[\frac{n\alpha}{n-1} - \frac{nm(p-P)}{n(m-1)+m} \right]^2 \right\} v^{\alpha-2} |\nabla v|^{2m} \\
 &\quad + T v^{\alpha+2p} + \left\{ U-1 - \frac{(n-1)^2}{4n^2} \left[\frac{nqU}{n(m-1)+q} \right]^2 \right\} M^2 v^\alpha |\nabla v|^{2q} \\
 &\quad + \left[-(\alpha+p)T - P \right] v^{\alpha+p-1} |\nabla v|^m \\
 &\quad + \left\{ \alpha + p - P - \frac{n(m-1)\alpha U}{n(m-1)+q} + \frac{(n-1)^2}{2n^2} \left[\frac{n\alpha}{n-1} - \frac{nm(p-P)}{n(m-1)+m} \right] \frac{nqU}{n(m-1)+q} \right\} M v^{\alpha-1} |\nabla v|^{m+q} \\
 &\quad + (T+U-1) M v^{\alpha+p} |\nabla v|^q.
 \end{aligned} \tag{23}$$

If the following conditions hold at the same time:

$$\begin{cases} -(p-P)(\alpha-1)\frac{n(m-1)}{n(m-1)+m} - \frac{(n-1)^2}{4n^2} \left[\frac{n\alpha}{n-1} - \frac{nm(p-P)}{n(m-1)+m} \right]^2 > 0, \end{cases} \quad (24)$$

$$T > 0, \quad (25)$$

$$\begin{cases} U - 1 - \frac{(n-1)^2}{4n^2} \left[\frac{nqU}{n(m-1)+q} \right]^2 > 0, \end{cases} \quad (26)$$

$$\begin{cases} -(\alpha+p)T - P \geq 0, \end{cases} \quad (27)$$

$$\begin{cases} \left\{ \alpha + p - P - \frac{n(m-1)\alpha U}{n(m-1)+q} + \frac{(n-1)^2}{2n^2} \left[\frac{n\alpha}{n-1} - \frac{nm(p-P)}{n(m-1)+m} \right] \cdot \frac{nqU}{n(m-1)+q} \right\} M \geq 0, \end{cases} \quad (28)$$

$$\begin{cases} (T+U-1)M \geq 0, \end{cases} \quad (29)$$

then there always exists $\varepsilon > 0$, such that

$$\begin{aligned} & \varepsilon v^{\alpha-2} |\nabla v|^{2m} + \varepsilon v^\alpha |\nabla v|^{2m-6} \left(\sum_j v_j v_{ij} \right)^2 + \varepsilon v^{\alpha+2p} + \varepsilon v^\alpha |\nabla v|^{2q} \\ & \leq B_1 (v^{\alpha-1} |\nabla v|^{2m-2} v_i)_i + B_2 (v^\alpha |\nabla v|^q X^i)_i + B_3 (v^{\alpha+p} X^i)_i + B_4 (v^\alpha X_j^i X^j)_i. \end{aligned} \quad (30)$$

To make sure (24) hold, we can choose

$$\alpha = \frac{-mn - m + 2n}{n(m-1) + m} (p - P),$$

and

$$m - 1 < p - P < \frac{(m-1)n + m}{n - m}.$$

4. YOUNG'S INEQUALITY AND CUT-OFF FUNCTIONS

In this part, we assume (30) holds and prove that $|\nabla v| \equiv 0$. Define η is a smooth cut-off function, satisfying that

$$\begin{aligned} \eta &\equiv 1 \quad \text{in } B_R, \\ \eta &\equiv 0 \quad \text{in } \mathbb{R}^n \setminus B_{2R}. \end{aligned}$$

Multiply (30) by η^δ and integrate over \mathbb{R}^n ,

$$\begin{aligned} & \varepsilon \int_{\mathbb{R}^n} v^{\alpha-2} |\nabla v|^{2m} \eta^\delta + \varepsilon \int_{\mathbb{R}^n} v^\alpha |\nabla v|^{2m-6} \left(\sum_j v_j v_{ij} \right)^2 \eta^\delta + \varepsilon \int_{\mathbb{R}^n} v^{\alpha+2p} \eta^\delta + \varepsilon \int_{\mathbb{R}^n} v^\alpha |\nabla v|^{2q} \eta^\delta \\ & \leq B_1 \delta \int_{B_{2R} \setminus B_R} v^{\alpha-1} |\nabla v|^{2m-2} v_i \eta^{\delta-1} \eta_i + B_2 \delta \int_{B_{2R} \setminus B_R} v^\alpha |\nabla v|^{m+q-2} v_i \eta^{\delta-1} \eta_i \\ & + B_3 \delta \int_{B_{2R} \setminus B_R} v^{\alpha+p} |\nabla v|^{m-2} v_i \eta^{\delta-1} \eta_i + B_4 \delta \int_{B_{2R} \setminus B_R} v^\alpha X_j^i X^j \eta^{\delta-1} \eta_i. \end{aligned}$$

Since we know that

$$\begin{aligned}
 & \int_{B_{2R} \setminus B_R} v^\alpha X_j^i X^j \eta^{\delta-1} \eta_i \\
 &= - \int_{B_{2R} \setminus B_R} (v^\alpha X^j \eta^{\delta-1} \eta_i)_j X^i \\
 &= -\alpha \int_{B_{2R} \setminus B_R} v^{\alpha-1} |\nabla v|^{2m-2} v_i \eta^{\delta-1} \eta_i - \int_{B_{2R} \setminus B_R} v^\alpha \Delta_m v \eta^{\delta-1} \eta_i X^i - \int_{B_{2R} \setminus B_R} v^\alpha X^j (\eta^{\delta-1} \eta_i)_j X^i \\
 &= -\alpha \int_{B_{2R} \setminus B_R} v^{\alpha-1} |\nabla v|^{2m-2} v_i \eta^{\delta-1} \eta_i + \int_{B_{2R} \setminus B_R} v^\alpha (v^p + M |\nabla v|^q) \eta^{\delta-1} \eta_i X^i \\
 &\quad - \int_{B_{2R} \setminus B_R} v^\alpha |\nabla v|^{2m-4} v_i v_j (\eta^{\delta-1} \eta_i)_j \\
 &= -\alpha \int_{B_{2R} \setminus B_R} v^{\alpha-1} |\nabla v|^{2m-2} v_i \eta^{\delta-1} \eta_i + \int_{B_{2R} \setminus B_R} v^{\alpha+p} |\nabla v|^{m-2} v_i \eta^{\delta-1} \eta_i + M \int_{B_{2R} \setminus B_R} v^\alpha |\nabla v|^{m+q-2} v_i \eta^{\delta-1} \eta_i \\
 &\quad - \int_{B_{2R} \setminus B_R} v^\alpha |\nabla v|^{2m-4} v_i v_j (\eta^{\delta-1} \eta_i)_j,
 \end{aligned}$$

then we get

$$\begin{aligned}
 & \varepsilon \int_{R^n} v^{\alpha-2} |\nabla v|^{2m} \eta^\delta + \varepsilon \int_{R^n} v^\alpha |\nabla v|^{2m-6} \left(\sum_j v_j v_{ij} \right)^2 \eta^\delta + \varepsilon \int_{R^n} v^{\alpha+2p} \eta^\delta + \varepsilon \int_{R^n} v^\alpha |\nabla v|^{2q} \eta^\delta \\
 &\leq \frac{\varepsilon}{2} \int_{B_{2R} \setminus B_R} v^{\alpha-2} |\nabla v|^{2m} \eta^\delta + \frac{C}{R^2} \int_{B_{2R} \setminus B_R} v^\alpha |\nabla v|^{2m-2} \eta^{\delta-2} \\
 &\quad + \frac{\varepsilon}{2} \int_{B_{2R} \setminus B_R} v^\alpha |\nabla v|^{2q} \eta^\delta + \frac{C}{R^2} \int_{B_{2R} \setminus B_R} v^\alpha |\nabla v|^{2m-2} \eta^{\delta-2} \\
 &\quad + \frac{\varepsilon}{2} \int_{B_{2R} \setminus B_R} v^{\alpha+2p} \eta^\delta + \frac{C}{R^2} \int_{B_{2R} \setminus B_R} v^\alpha |\nabla v|^{2m-2} \eta^{\delta-2}.
 \end{aligned}$$

Therefore, we get that

$$\int_{R^n} v^{\alpha-2} |\nabla v|^{2m} \eta^\delta + \int_{R^n} v^{\alpha+2p} \eta^\delta \leq \frac{C}{R^2} \int_{B_{2R} \setminus B_R} v^\alpha |\nabla v|^{2m-2} \eta^{\delta-2}. \quad (31)$$

Define $p_1, q_1, \sigma_1 > 0$, such that

$$\frac{1}{p_1} + \frac{1}{q_1} + \frac{1}{\sigma_1} = 1, \text{ and } p_1, q_1, \sigma_1 > 0. \quad (32)$$

So by Young's inequality, we know

$$\begin{aligned}
& R^{-2} \int_{B_{2R} \setminus B_R} v^\alpha |\nabla v|^{2m-2} \eta^{\delta-2} \\
&= R^{-2} \int_{B_{2R} \setminus B_R} v^{\alpha-A} |\nabla v|^{2m-2} \cdot v^A \eta^{\delta-2} \\
&\leq \varepsilon_0 \int_{B_{2R} \setminus B_R} v^{\alpha-2} |\nabla v|^{2m} \eta^\delta + \varepsilon_0 \int_{B_{2R} \setminus B_R} v^{\alpha+2p} \eta^\delta + CR^{-2\sigma_1} \int_{B_{2R} \setminus B_R} \eta^{\delta-2\sigma_1}.
\end{aligned} \tag{33}$$

Combine (31) and (33), it follows that

$$\begin{aligned}
& \int_{R^n} v^{\alpha+2p} \eta^\delta + \int_{R^n} v^{\alpha-2} |\nabla v|^{2m} \eta^\delta \\
&\leq CR^{-2\sigma_1} \int_{B_{2R} \setminus B_R} \eta^{\delta-2\sigma_1} \\
&\leq CR^{n-2\sigma_1} \rightarrow 0, \text{ as } R \text{ tends to infinity.}
\end{aligned} \tag{34}$$

The last two steps come from the condition that

$$\begin{aligned}
& \delta - 2\sigma_1 > 0, \\
& n - 2\sigma_1 < 0.
\end{aligned} \tag{35}$$

So we always take δ large enough. In conclusion, we need

$$\begin{aligned}
& \frac{\alpha+2p}{A} = p_1 > 0, \\
& \frac{\alpha-2}{\alpha-A} = \frac{2m}{2m-2} = q_1 > 0, \\
& 1 - \frac{2}{n} < \frac{1}{p_1} + \frac{1}{q_1} < 1.
\end{aligned} \tag{36}$$

Therefore, as long as the conditions (24) - (29) are satisfied, and there exists such p_1, q_1, σ_1 , then $v \equiv 0$.

By (36), we know

$$\begin{aligned}
A &= \frac{\alpha+2m-2}{m}, \\
\frac{1}{p_1} + \frac{1}{q_1} &= \frac{m-1}{m} + \frac{\alpha+2m-2}{m(\alpha+2p)}.
\end{aligned} \tag{37}$$

Next, we set out to prove

$$1 - \frac{2}{n} < \frac{1}{p_1} + \frac{1}{q_1} < 1.$$

Firstly, for the left hand:

$$\begin{aligned} 1 - \frac{2}{n} &< \frac{1}{p_1} + \frac{1}{q_1}, \\ \Leftrightarrow \frac{2}{n} - \frac{1}{m} + \frac{\alpha + 2m - 2}{m(\alpha + 2p)} &> 0, \\ \Leftrightarrow \alpha &> -2p - n + \frac{n(p+1)}{m}. \end{aligned}$$

As we choose in the previous section,

$$\alpha = \frac{-mn - m + 2n}{n(m-1) + m}(p - P),$$

then it means to show

$$\frac{-mn - m + 2n}{n(m-1) + m}(p - P) > -2p - n + \frac{n(p+1)}{m}.$$

Since $m - 1 < p - P < \frac{(m-1)n+m}{n-m}$, we only need to show that

$$\left\{ \begin{array}{l} \frac{-mn - m + 2n}{n(m-1) + m}(m - 1 - P) > -2(m - 1), \end{array} \right. \quad (38)$$

$$\left\{ \begin{array}{l} \frac{-mn - m + 2n}{n(m-1) + m} \cdot \frac{(m-1)n+m}{n-m} - \frac{-mn - m + 2n}{n(m-1) + m}P > \frac{n-2m}{m} \frac{(m-1)n+m}{n-m} - n + \frac{n}{m}. \end{array} \right. \quad (39)$$

It is equivalent to be

$$\left\{ \begin{array}{l} (m-1)(mn+m) - (-mn - m + 2n)P > 0, \end{array} \right. \quad (40)$$

$$\left\{ \begin{array}{l} \frac{m^2}{m(n-m)} - \frac{-mn - m + 2n}{n(m-1) + m}P > 0. \end{array} \right. \quad (41)$$

Next, for the right hand:

by (37), it's namely to prove

$$\frac{\alpha + 2m - 2 - \alpha - 2p}{m(\alpha + 2p)} < 0.$$

We only need to show that

$$\alpha + 2m - 2 > 0,$$

then

$$\alpha + 2p > \alpha + 2m - 2 > 0,$$

$$p_1 > 0.$$

There are three different cases:

- If $\alpha + 2m - 2 > 0$, then we can get desired $p_1, q_1 > 0$.
- If $\alpha + 2m - 2 \leq 0$ and $M > 0$, then we have the following Lemma:

Lemma 4.1. *If we choose the same α as before, then*

$$\int_{R^n} v^{\alpha+2p} \eta^\delta \leq CR^{-2m} \int_{B_{2R} \setminus B_R} v^{\alpha+2m-2} \eta^{\delta-2m}. \quad (42)$$

Proof. By (31) and Young's inequality , we have

$$\begin{aligned}
& \int_{R^n} v^{\alpha-2} |\nabla v|^{2m} \eta^\delta + \int_{R^n} v^{\alpha+2p} \eta^\delta \\
& \leq \frac{C}{R^2} \int_{B_{2R} \setminus B_R} v^\alpha |\nabla v|^{2m-2} \eta^{\delta-2} \\
& \leq CR^{-2m} \int_{B_{2R} \setminus B_R} v^{\alpha+2m-2} \eta^{\delta-2m} + \frac{1}{2} \int_{B_{2R} \setminus B_R} v^{\alpha-2} |\nabla v|^{2m} \eta^\delta
\end{aligned}$$

with $(m, \frac{m}{m-1})$. So we get that

$$\int_{R^n} v^{\alpha+2p} \eta^\delta \leq CR^{-2m} \int_{B_{2R} \setminus B_R} v^{\alpha+2m-2} \eta^{\delta-2m}.$$

□

Besides, we need the following result which comes from Lemma 2.3 in [21] by Serrin-Zou.

Lemma 4.2. Suppose $\{|x| > R > 0\} \subset \Omega$ and $1 < m < n$. Let v be a positive solution of the inequality

$$\Delta_m v \leq 0, x \in \Omega.$$

Then there exists a constant $C = C(m, n, \min_{|x|=2R} v, R)$ such that

$$v(x) \geq C|x|^{-\frac{n-m}{m-1}}, \text{ when } |x| > 2R.$$

Therefore, if $M > 0$, we know that $\Delta_m v \leq 0$, then

$$v(x) \geq CR^{-\frac{n-m}{m-1}}.$$

Thus we get that

$$\begin{aligned}
\int_{R^n} v^{\alpha+2p} \eta^\delta & \leq CR^{-2m} \int_{B_{2R} \setminus B_R} v^{\alpha+2m-2} \eta^{\delta-2m} \\
& \leq CR^{-2m+n-(\alpha+2m-2)\cdot\frac{n-m}{m-1}}.
\end{aligned}$$

We hope that

$$-2m + n - (\alpha + 2m - 2) \cdot \frac{n-m}{m-1} < 0.$$

If $-mn - m + 2n \geq 0$, then $\alpha \geq 0$ and $\alpha + 2m - 2 > 0$. This is the first case.
So we only need to consider the case that $-mn - m + 2n < 0$:

Since

$$\begin{aligned}
 & \alpha + 2m - 2 \\
 &= \frac{-mn - m + 2n}{n(m-1) + m} p - \frac{-mn - m + 2n}{n(m-1) + m} P + 2m - 2 \\
 &\geq \frac{-mn - m + 2n}{n-m} - \frac{-mn - m + 2n}{n(m-1) + m} P + 2m - 2 \\
 &= \frac{-mn - m + 2n + 2mn - 2m^2 - 2n + 2m}{n-m} - \frac{-mn - m + 2n}{n(m-1) + m} P \\
 &= \frac{m(n+1-2m)}{n-m} - \frac{-mn - m + 2n}{n(m-1) + m} P,
 \end{aligned}$$

then

$$\begin{aligned}
 & -2m + n - (\alpha + 2m - 2) \cdot \frac{n-m}{m-1} \\
 &\leq -2m + n - \frac{m(-2m+n+1)}{n-m} \cdot \frac{n-m}{m-1} + \frac{-mn - m + 2n}{n(m-1) + m} P \cdot \frac{n-m}{m-1} \\
 &= -2m + n - \frac{m(-2m+n+1)}{m-1} + \frac{-mn - m + 2n}{n(m-1) + m} P \cdot \frac{n-m}{m-1} \\
 &= \frac{-2m^2 + 2m + mn - n + 2m^2 - mn - m}{m-1} + \frac{-mn - m + 2n}{n(m-1) + m} P \cdot \frac{n-m}{m-1} \\
 &= \frac{m-n}{m-1} + \frac{-mn - m + 2n}{n(m-1) + m} P \cdot \frac{n-m}{m-1} \\
 &= \frac{m-n}{m-1} \left[1 - \frac{-mn - m + 2n}{n(m-1) + m} P \right]. \tag{43}
 \end{aligned}$$

In conclusion, it follows by (40), (41) and (43) that if

$$M > 0, 1 < m < n, m-1 < p-P < \frac{(m-1)n+m}{n-m}, -P_m < P < P_m,$$

where

$$P_m = \min \left\{ \frac{(mn+m)(m-1)}{|-mn-m+2n|}, \frac{m^2}{m(n-m)} \left[\frac{|-mn-m+2n|}{n(m-1)+m} \right]^{-1}, \frac{n(m-1)+m}{|-mn-m+2n|} \right\},$$

then we can deduce by (30) that

$$v \equiv 0.$$

- If $\alpha + 2m - 2 \leq 0$ and $M < 0$. The condition $\Delta_m v \leq 0$ does not hold any more and we can not use Lemma 4.2 directly. But inspired by [10], if we define $v = w^\sigma$ with $\sigma > 1$, we can get that $\Delta_m w \leq 0$ under some conditions for

p, P, m, M .

$$\begin{aligned} -\Delta_m w &= (\sigma - 1)(m - 1)w^{-1}|\nabla w|^m + \sigma^{1-m}w^{m+\sigma(p-m+1)-1} \\ &\quad + M\sigma^{q-m+1}w^{(\sigma-1)(q-m+1)}|\nabla w|^q, \end{aligned}$$

and then setting $z = |\nabla v|^m$ yields

$$-\Delta_m w = \sigma^{1-m}w^{-1}\Psi(z),$$

where

$$\Psi(z) = \sigma^{m-1}(\sigma - 1)(m - 1)z + M\sigma^q w^{(\sigma-1)(q-m+1)+1}z^{\frac{q}{m}} + w^{m+\sigma(p-m+1)}.$$

Since $q = \frac{mp}{p+1}$, it is easy to see that $\Psi(z)$ achieves its minimum at

$$z_0 = \left[\frac{|M| p \sigma^{1-\frac{m}{p+1}}}{(\sigma - 1)(m - 1)(p + 1)} \right]^{p+1} v^{m+\sigma(p-m+1)},$$

and

$$\Psi(z_0) = \left[1 - \left(\frac{|M|}{p+1} \right)^{p+1} \frac{(\sigma p)^p}{(\sigma - 1)^p(m - 1)^p} \right] v^{m+\sigma(p-m+1)}.$$

If

$$1 - \left(\frac{|M|}{p+1} \right)^{p+1} \frac{(\sigma p)^p}{(\sigma - 1)^p(m - 1)^p} \geq 0, \quad (44)$$

then we know that $\Delta_m w \geq 0$ and by Lemma 4.2 we get that

$$v = w^\sigma \geq CR^{-\frac{n-m}{m-1}\sigma}.$$

Thus we hope the follow conditions will hold at the same time:

$$\begin{cases} -2m + n - (\alpha + 2m - 2) \cdot \frac{n-m}{m-1}\sigma < 0, \end{cases} \quad (45)$$

$$\begin{cases} 1 - \left(\frac{|M|}{p+1} \right)^{p+1} \cdot \frac{(\sigma p)^p}{(\sigma - 1)^p(m - 1)^p} \geq 0. \end{cases} \quad (46)$$

Observe that

$$\begin{aligned} &-2m + n - (\alpha + 2m - 2) \cdot \frac{n-m}{m-1}\sigma \\ &= \frac{m-n}{m-1} \left[1 + \frac{m(-2m+n+1)}{n-m}(\sigma - 1) \right] - \frac{m-n}{m-1} \cdot \frac{-mn-m+2n}{n(m-1)+m} P\sigma. \end{aligned}$$

If $-2m + n + 1 \geq 0$, we choose $\sigma = 2$; if $-2m + n + 1 < 0$, we choose

$$\frac{m(-2m+n+1)}{n-m}(\sigma - 1) = -\frac{1}{2}.$$

Then there exists $M_3 = M_3(n, m, p) > 0$, such that if $-M_3 < M < 0$, then the condition (46) hold. Besides, for condition (45) we have

$$\begin{aligned}
 & -2m + n - (\alpha + 2m - 2) \cdot \frac{n-m}{m-1} \sigma \\
 & \leq \frac{m-n}{m-1} \cdot \frac{1}{2} - \frac{m-n}{m-1} \cdot \frac{-mn-m+2n}{n(m-1)+m} P\sigma \\
 & = \frac{m-n}{m-1} \cdot \left[\frac{1}{2} - \frac{-mn-m+2n}{n(m-1)+m} P\sigma \right].
 \end{aligned} \tag{47}$$

In conclusion, combined with (40), (41) and (47), if

$$-M_3 < M < 0, 1 < m < n, m-1 < p-P < \frac{(m-1)n+m}{n-m}, -P_m < P < P_m,$$

where

$$P_m = \min \left\{ \frac{(mn+m)(m-1)}{|-mn-m+2n|}, \frac{m^2}{m(n-m)} \left[\frac{|-mn-m+2n|}{n(m-1)+m} \right]^{-1}, \frac{n(m-1)+m}{2\sigma|-mn-m+2n|} \right\},$$

then we can deduce by (30) that

$$v \equiv 0.$$

5. CASE $M > 0$ AND $M < 0$

In this section, we are working on proving (24) - (29) hold, and talk about them separately according to the sign of M .

Case 1: $M > 0$

Theorem 5.1. Let $n \geq 2$, $m-1 < p \leq \frac{(m-1)n+m}{n-m}$, $q = \frac{mp}{p+1}$, then there exists $M_1 > 0$, which depends on n, m, p , such that for any $0 < M < M_1$, all the nonnegative solutions of (1) are $v \equiv 0$.

Proof of Theorem 5.1. In this proof, we choose $P = T = 0$ at first and then

$$\begin{aligned}
 \alpha &= \frac{-mn-m+2n}{n(m-1)+m} p, \\
 \alpha + p &= \frac{np}{n(m-1)+m} > 0,
 \end{aligned}$$

then the condition (24) holds for $m - 1 < p < \frac{(m-1)n+m}{n-m}$. And the condition (28) becomes that:

$$\begin{aligned}
0 &\leq \alpha + p - \frac{n(m-1)\alpha U}{n(m-1)+q} + \frac{(n-1)^2}{2n^2} \left[\frac{n\alpha}{n-1} - \frac{nmp}{n(m-1)+m} \right] \cdot \frac{nqU}{n(m-1)+q} \\
&= \frac{np}{n(m-1)+m} - \frac{-mn-m+2n}{n(m-1)+m} p \cdot \frac{n(m-1)U}{n(m-1)+q} \\
&\quad + \frac{(n-1)^2}{2n^2} \left[\frac{n}{n-1} \cdot \frac{-mn-m+2n}{n(m-1)+m} p - \frac{nmp}{n(m-1)+m} \right] \cdot \frac{nqU}{n(m-1)+q} \\
&= \frac{np}{n(m-1)+m} + \frac{npU}{(nm-n+q)(nm-n+m)} \left[(mn+m-2n)(m-1) \right. \\
&\quad \left. + \frac{(n-1)^2 q}{2n} \left(\frac{-mn-m+2n}{n-1} - m \right) \right] \\
&= \frac{np}{nm-n+m} + \frac{np(m-1)U}{(nm-n+q)(nm-n+m)} \left[mn+m-2n-(n-1)q \right] \\
&= \frac{np}{(nm-n+q)(nm-n+m)} \left[nm-n+q+(m-1)U(mn+m-2n-(n-1)q) \right].
\end{aligned}$$

Define

$$f_1(m, q) := nm - n + q + (m-1) \left(1 + \frac{1}{mn} \right) \left[mn + m - 2n - (n-1)q \right].$$

claim 1. For any $m - 1 < q \leq m - 1 + \frac{m}{n}$, $1 < m < n$, we have

$$f_1(m, q) > 0.$$

Proof of claim 1. We only need to show that

$$f_1(m, m-1) \quad \text{and} \quad f_1 \left(m, m-1 + \frac{m}{n} \right) > 0.$$

In fact, we have

$$\begin{aligned}
f_1(m, m-1) &= nm - n + m - 1 + \left(1 + \frac{1}{mn} \right) (m-1) \left[mn + m - 2n - (n-1)(m-1) \right] \\
&= (m-1) \left(1 + \frac{1}{mn} \right) \left[n + 1 - \frac{n+1}{mn+1} + mn + m - 2n - (n-1)(m-1) \right] \\
&= (m-1) \left(1 + \frac{1}{mn} \right) \left(2m - \frac{n+1}{mn+1} \right) > 0,
\end{aligned}$$

and

$$\begin{aligned}
 & f_1 \left(m, m - 1 + \frac{m}{n} \right) \\
 &= nm - n + m - 1 + \frac{m}{n} + (m - 1) \left(1 + \frac{1}{mn} \right) \left[mn + m - 2n - (n - 1) \left(m - 1 + \frac{m}{n} \right) \right] \\
 &= \frac{m^2 n^3 - mn^3 + m^2 n^2 - mn^2 + m^2 n + (m - 1)(mn + 1) \left[mn^2 + mn - 2n^2 - (n - 1)(mn - n + m) \right]}{mn^2} \\
 &= \frac{m^2 n^3 - mn^3 + m^2 n^2 - mn^2 + m^2 n + (m^2 n + m - mn - 1) \left[mn - n^2 - n + m \right]}{mn^2} > 0.
 \end{aligned}$$

Define

$$\begin{aligned}
 f(n) &:= m^2 n^3 - mn^3 + m^2 n^2 - mn^2 + m^2 n + (m^2 n + m - mn - 1) \left[mn - n^2 - n + m \right] \\
 &= m^2 n^3 - mn^3 + m^2 n^2 - mn^2 + m^2 n + m^3 n^2 - m^2 n^3 - m^2 n^2 + m^3 n + m^2 n - mn^2 - mn + m^2 \\
 &\quad - m^2 n^2 + mn^3 + mn^2 - m^2 n - mn + n^2 + n - m \\
 &= m^3 n^2 + m^3 n + m^2 n - mn^2 - 2mn + m^2 - m^2 n^2 + n^2 + n - m \\
 &= (m^3 - m^2 - m + 1)n^2 + (m^3 + m^2 - 2m + 1)n + m^2 - m,
 \end{aligned}$$

it can be seen that

$$\begin{aligned}
 f'(n) &= 2n(m^3 - m^2 - m + 1) + m^3 + m^2 - 2m + 1 \\
 &> 2m(m^3 - m^2 - m + 1) + m^3 + m^2 - 2m + 1 \\
 &= 2m^4 - m^3 - m^2 + 1 \\
 &= m^2(m - 1)(2m + 1) + 1 > 0.
 \end{aligned}$$

so $f(n) > f(m) = m^5 > 0$.

□

Since $m - 1 < p \leq \frac{(m-1)n+m}{n-m}$, and $q = \frac{mp}{p+1}$, then we obtain the range of q , that is

$$m - 1 < q \leq m - 1 + \frac{m}{n}.$$

Therefore, if we choose $U = 1 + \frac{1}{mn}$, then the condition (28) and (29) hold. That is if we choose

$$\begin{aligned}
 P &= T = 0, \\
 \alpha &= \frac{-mn - m + 2n}{n(m - 1) + m}(p - P), \\
 U &= 1 + \frac{1}{mn},
 \end{aligned}$$

then for any $m - 1 < p < \frac{(m-1)n+m}{n-m}$, we have

$$\alpha + p - \frac{n(m-1)\alpha U}{n(m-1)+q} + \frac{(n-1)^2}{2n^2} \left[\frac{n\alpha}{n-1} - \frac{nmp}{n(m-1)+m} \right] \cdot \frac{nqU}{n(m-1)+q} > 0,$$

and the conditions (24) and (29) hold. It remains to show the condition (26) hold, while if

$$U - 1 - \frac{(n-1)^2}{4n^2} \left[\frac{nqU}{n(m-1)+q} \right]^2 < 0,$$

then by Young's inequality, it follows that

$$M^2 v^\alpha |\nabla v|^{2q} \leq K |M| v^{\alpha-1} |\nabla v|^{m+q} + \frac{\left(K \cdot \frac{p+1}{p} \right)^{-p}}{p+1} |M|^{p+2} v^{\alpha+p} |\nabla v|^q, \quad (48)$$

with $(\frac{p+1}{p}, p+1)$, by (23) we only need

$$\left\{ \alpha + p - \frac{n(m-1)\alpha U}{n(m-1)+q} + \frac{(n-1)^2}{2n^2} \left[\frac{n\alpha}{n-1} - \frac{nmp}{n(m-1)+m} \right] \cdot \frac{nqU}{n(m-1)+q} \right. \quad (49)$$

$$\left. + K \left(U - 1 - \frac{(n-1)^2}{4n^2} \left[\frac{nqU}{n(m-1)+q} \right]^2 \right) = 0, \right. \quad (50)$$

$$\left. U - 1 + \frac{\left(K \frac{p+1}{p} \right)^{-p}}{p+1} |M|^{p+1} \left(U - 1 - \frac{(n-1)^2}{4n^2} \left[\frac{nqU}{n(m-1)+q} \right]^2 \right) > 0, \right. \quad (51)$$

then

$$U - 1 - \frac{p^p}{(p+1)^{p+1}} |M|^{p+1} \left(-U + 1 + \frac{(n-1)^2}{4n^2} \left[\frac{nqU}{n(m-1)+q} \right]^2 \right)^{p+1} \\ \cdot \left[\alpha + p - \frac{n(m-1)\alpha U}{n(m-1)+q} + \frac{(n-1)^2}{2n^2} \left[\frac{n\alpha}{n-1} - \frac{nmp}{n(m-1)+m} \right] \cdot \frac{nqU}{n(m-1)+q} \right]^{-p} > 0.$$

Since U has been chosen as before, by the inequality above, we can get $M < M_1$. Thus we can say there exists $M_1 > 0$, such that for any $0 < M < M_1$, all the inequality conditions hold. Therefore, by continuity, when we choose

$$-\varepsilon < P < 0, \quad (\alpha + p)T + P = 0,$$

such that

$$p - P < \frac{(m-1)n+m}{n-m},$$

then conditions (24) - (29) all hold. Finally we explain the critical case $p = \frac{(m-1)n+m}{n-m}$. For fixed $M > 0$, we can still choose $0 < P < \varepsilon$, where $\varepsilon > 0$ is much smaller than M . Thus using the following Young's inequality, we can get the same result as $p < \frac{(m-1)n+m}{n-m}$:

$$v^{\alpha+p-1}|\nabla v|^m \leq \frac{q}{m} \left(J|M| \cdot \frac{m}{m-q} \right)^{-\frac{m-q}{q}} v^{\alpha+p} |\nabla v|^q + J|M|v^{\alpha-1} |\nabla v|^{m+q} \quad (52)$$

with $(\frac{m}{q}, \frac{m}{m-q})$.

□

Case 2: $M < 0$

Theorem 5.2. Let $n \geq 2$, $m-1 < p < \frac{(m-1)n+m}{n-m}$, $q = \frac{mp}{p+1}$, then there exists $M_2 > 0$, which depends on n, m, p , such that for any $-M_2 < M < 0$, all the nonnegative solutions of (1) are $v \equiv 0$.

Proof of Theorem 5.2. We still choose

$$\alpha = \frac{-mn - m + 2n}{n(m-1) + m} (p - P).$$

There are two different cases:

- $1 < m \leq \frac{n}{2} + \frac{1}{4}$:

In this case, we know

$$\begin{aligned} \alpha + p - P - \frac{n(m-1)\alpha U}{n(m-1) + q} + \frac{(n-1)^2}{2n^2} \left[\frac{n\alpha}{n-1} - \frac{nm(p-P)}{n(m-1) + m} \right] \cdot \frac{nqU}{n(m-1) + q} \\ = \frac{n(p-P)}{nm - n + m} + \frac{n(p-P)(m-1)U}{(nm - n + q)(nm - n + m)} \left[mn + m - 2n - (n-1)q \right], \end{aligned}$$

and

$$\begin{aligned} mn + m - 2n - (n-1)q \\ < mn + m - 2n - (n-1)(m-1) \\ = 2m - n - 1 \leq -\frac{1}{2}. \end{aligned}$$

So we can choose $U > 0$, such that

$$\alpha + p - \frac{n(m-1)\alpha U}{n(m-1) + q} + \frac{(n-1)^2}{2n^2} \left[\frac{n\alpha}{n-1} - \frac{nm(p-P)}{n(m-1) + m} \right] \cdot \frac{nqU}{n(m-1) + q} < 0.$$

Besides, for any $m-1 < p-P < \frac{nm-n+m}{n-m}$, we have

$$\begin{aligned} & - (p-P)(\alpha-1) \frac{n(m-1)}{n(m-1) + m} - \frac{(n-1)^2}{4n^2} \left[\frac{n\alpha}{n-1} - \frac{nm(p-P)}{n(m-1) + m} \right]^2 \\ &= \frac{p-P}{(nm - n + m)^2} (nm - n) \left[nm - n + m - (n-m)(p-P) \right] > 0. \end{aligned}$$

So for any $m-1 < p < \frac{nm-n+m}{n-m}$, there always exists P_0 , such that $-P_m < P_0 < 0$ and

$$p - P_0 < \frac{nm - n + m}{n - m},$$

$$\alpha + p > 0.$$

Choose

$$T = \frac{-0.5P_0}{\alpha + p} > 0,$$

then the conditions (24), (25), (27) and (28) all hold. It remains to show (26) and (29). If they don't hold, we can use Young's inequality,

$$|M|v^{\alpha+p}|\nabla v|^q \leq Kv^{\alpha+2p} + \frac{q}{m} \left(K \cdot \frac{m}{m-q} \right)^{-\frac{m-q}{q}} |M|^{\frac{m}{q}} v^{\alpha+p-1} |\nabla v|^m, \quad (53)$$

with $(\frac{m}{m-q}, \frac{m}{q})$.

Using (48) and (53), there exists $0 < M_2 < M_3$ such that for any $-M_2 < M < 0$, the result (30) hold.

- $\frac{n}{2} + \frac{1}{4} \leq m < n$:

In this case, we consider the condition (26):

$$U - 1 - \frac{(n-1)^2}{4n^2} \left[\frac{nqU}{n(m-1)+q} \right]^2 > 0.$$

claim 2. When $\frac{n}{2} + \frac{1}{4} \leq m < n$, we have

$$\Delta = 1 - \frac{(n-1)^2}{n^2} \left[\frac{nq}{n(m-1)+q} \right]^2 > 0.$$

Proof of claim 2.

$$\begin{aligned} \Delta &= 1 - (n-1)^2 \left[\frac{q}{n(m-1)+q} \right]^2 \\ &= \frac{(mn-n+q)^2 - (n-1)^2 q^2}{(mn-n+q)^2} \\ &= \frac{-(n^2-2n)q^2 + 2(mn-n)q + (mn-n)^2}{(mn-n+q)^2}. \end{aligned}$$

Since $m-1 < q < m-1 + \frac{m}{n}$, we only need to check that

$$\left\{ -(n^2-2n)(m-1)^2 + 2(mn-n)(m-1) + (mn-n)^2 > 0, \quad (54) \right.$$

$$\left. -(n^2-2n) \left(m-1 + \frac{m}{n} \right)^2 + 2(mn-n) \left(m-1 + \frac{m}{n} \right) + (mn-n)^2 > 0. \quad (55) \right.$$

In fact, we have

$$(54) \Leftrightarrow n(m-1) \left[-(n-2)(m-1) + 2(m-1) + (mn-n) \right] > 0$$

$$\Leftrightarrow 4n(m-1)^2 > 0.$$

On the other hand, we know

$$(55) \Leftrightarrow -(n-2)(mn-n+m)^2 + 2(mn-n)(mn-n+m) + n(mn-n)^2 > 0,$$

$$\Leftrightarrow 2m^2n^2 + 4n^2 + 2m^2 + 5m^2n - 6mn(n+1) > 0.$$
(56)

Define

$$f(m) := 2m^2n^2 + 4n^2 + 2m^2 + 5m^2n - 6mn(n+1)$$

$$= m^2(2n^2 + 5n + 2) - m(6n^2 + 6n) + 4n^2.$$

It is equivalent to prove

$$f(m) > 0.$$

Since $m \geq \frac{n}{2} + \frac{1}{4} > \frac{3n^2+3n}{2n^2+5n+2}$, we get that

$$f(m) \geq f\left(\frac{n}{2} + \frac{1}{4}\right)$$

$$= \left(\frac{n}{2} + \frac{1}{4}\right)^2 (2n^2 + 5n + 2) - \left(\frac{n}{2} + \frac{1}{4}\right) (6n^2 + 6n) + 4n^2$$

$$= 0.5n^4 - 1.25n^3 + 1.375n^2 - 0.6875n + 0.125$$

$$= 0.5(n-0.5)^2(n^2 - 1.5n + 1).$$

So $f(m) > 0$, i.e. (56) holds. \square

So we know there exists $U_0 > 0$ such that condition (26) holds. At this time, we still choose the same P, T as the first case, then (24) - (27) hold, but

$$\begin{cases} \alpha + p - \frac{n(m-1)\alpha U}{n(m-1)+q} + \frac{(n-1)^2}{2n^2} \left[\frac{n\alpha}{n-1} - \frac{nmp}{n(m-1)+m} \right] \cdot \frac{nqU}{n(m-1)+q} > 0, \\ T + U - 1 > 0. \end{cases}$$

By Young's inequality,

$$|M|v^{\alpha-1}|\nabla v|^{m+q} \leq Jv^{\alpha-2}|\nabla v|^{2m} + \frac{1}{p+2} \left(J \cdot \frac{p+2}{p+1} \right)^{-p-1} |M|^{p+2}v^{\alpha+p}|\nabla v|^q, \quad (57)$$

with $(\frac{p+2}{p+1}, p+2)$, and

$$|M|v^{\alpha+p}|\nabla v|^q \leq Kv^{\alpha+2p} + \frac{q}{m} \left(K \cdot \frac{m}{m-q} \right)^{-\frac{m-q}{q}} |M|^{\frac{m}{q}} v^{\alpha+p-1} |\nabla v|^m,$$

with $(\frac{m}{m-q}, \frac{m}{q})$. Therefore there exists $0 < M_2 < M_3$ such that for any $-M_2 < M < 0$, the result (30) holds. \square

REFERENCES

- [1] M.F. Bidaut-Véron, Local and global behavior of solutions of quasilinear equations of Emden-Fowler type, *Arch. Rational Mech. Anal.* 107 (1989), 293-324.
- [2] M.F. Bidaut-Véron, Liouville results and asymptotics of solutions of a quasilinear elliptic equation with supercritical source gradient term, *Adv. Nonlinear Stud.* 21 (2021), 57-76.
- [3] M.F. Bidaut-Véron, M. Garcia-Huidobro and L. Véron, Local and global properties of solutions of quasilinear Hamilton-Jacobi equations. *J. Funct. Anal.* 267(9) (2014), 3294-3331.
- [4] M.F. Bidaut-Véron, M. Garcia-Huidobro and L. Véron, A priori estimates for elliptic equations with reaction terms involving the function and its gradient. *Math. Ann.* 378(1-2) (2020), 13-56.
- [5] M.F. Bidaut-Véron, M. Garcia-Huidobro and L. Véron, Radial solutions of scaling invariant nonlinear elliptic equations with mixed reaction terms, *Discrete Contin. Dyn. Syst.* 40 (2020), 933-982.
- [6] M.F. Bidaut-Véron, L. Véron, Nonlinear elliptic equations on compact Riemannian manifolds and asymptotics of Emden equations, *Invent. Math.* 106 (1991), 489-539.
- [7] W.X. Chen, C. Li, Classification of solutions of some nonlinear elliptic equations, *Duke Math. J.* 63 (1991), 615-622.
- [8] M. Chipot, F. Weissler, Some blow up results for nonlinear evolution equations with a gradient term. *SIAM J. Math. Anal.* 20 (1989), 886-907.
- [9] L. Damascelli, S. Merchán, L. Montoro and B. Sciunzi, Radial symmetry and applications for a problem involving the $-\Delta_p(\cdot)$ operator and critical nonlinearity in \mathbb{R}^n , *Adv. Math.* 265 (2014), 313-335.
- [10] R. Filippucci, Y. Sun and Y. Zheng, A priori estimates and Liouville type results for quasilinear elliptic equations involving gradient terms, arXiv:2205.07484, (2022).
- [11] B. Gidas, J. Spruck, Global and local behavior of positive solutions of nonlinear elliptic equations, *Comm. Pure Appl. Math.* 34 (1981), 525-598.
- [12] M. Guedda, L.Véron, Local and global properties of solutions of quasilinear elliptic equations, *J. Differential Equations*, 76 (1988), 159-189.
- [13] P.L. Lions, Quelques remarques sur les problés elliptiques quasilinéaires du second ordre. (French) [Some remarks on second order elliptic quasilinear problems] *J. Analyse Math.* 45 (1985), 234-254.
- [14] X.N. Ma, W.Z. Wu and Q.Q. Zhang, Liouville theorem for elliptic equations involving the sum of the function and its gradient in \mathbb{R}^n , arXiv:2311.04641.
- [15] W.M. Ni, J. Serrin, Non-existence theorems for quasilinear partial differential equations, *Rend. Circ. Mat. Palermo*, suppl. 8 (1985), 171-185.
- [16] W.M. Ni, J. Serrin, Existence and nonexistence theorems for ground states of quasilinear partial differential equations: The anomalous case, *Accad. Naz. dei Lincei*, 77 (1986), 231-257.

- [17] W.M. Ni, J. Serrin, Nonexistence theorems for singular solutions of quasilinear partial differential equations, *Comm. Pure Appl. Math.* (1986), 379-399.
- [18] J. Serrin, Local behavior of solutions of quasilinear equations, *Acta Math.* 111 (1964), 247-302.
- [19] J. Serrin, H. Zou, Existence and non-existence results for ground states of quasilinear elliptic equations. *Arch. Rat. Mech. Anal.* 121 (1992), 101-130.
- [20] J. Serrin, H. Zou, Ground States of Degenerate Quasilinear Equations. *Mathematics in Science and Engineering*, 192 (1993), 287-305.
- [21] J. Serrin, H. Zou, Cauchy Liouville and universal boundedness theorems for quasilinear elliptic equations and inequalities, *Acta Math.* 189 (2002), 791-42.
- [22] J. Vétois, A priori estimates and application to the symmetry of solutions for critical p-Laplace equations, *J. Differential Equations*, 260 (2016), 149-161.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA, HEFEI, CHINA

Email address: wuwz18@mail.ustc.edu.cn

COLLEGE OF MATHEMATICS AND INFORMATION SCIENCE, NANCHANG HANGKONG UNIVERSITY, NANCHANG, CHINA

Email address: qiqizmath@126.com