A NOTE ON ŁOŚ'S THEOREM WITHOUT THE AXIOM OF CHOICE

TOSHIMICHI USUBA

ABSTRACT. We study some topics about Los's theorem without assuming the Axiom of Choice. We prove that Los's fundamental theorem of ultraproducts is equivalent to a weak form that every ultrapower is elementarily equivalent to its source structure. On the other hand, it is consistent that there is a structure M and an ultrafilter U such that the ultrapower of M by U is elementarily equivalent to M, but the fundamental theorem for the ultrapower of M by U fails. We also show that weak fragments of the Axiom of Choice, such as the Countable Choice, do not follow from Los's theorem, even assuming the existence of non-principal ultrafilters.

1. INTRODUCTION

Ultraproduct and ultrapower are basic tools in model theory, set theory, and many other fields. For an indexed family of first order structures $\{M_i \mid i \in I\}$ with same language and an ultrafilter U over I, let $\prod_{i \in I} M_i/U$ denote the ultraproduct of the family $\{M_i \mid i \in I\}$ by U. For ultraproducts, the following Loś's theorem is very important:

Theorem 1.1 (Loś's fundamental theorem of ultraproducts [10]). Let U be an ultrafilter over a set I, and $\{M_i \mid i \in I\}$ an indexed family of structures with language \mathcal{L} . For $f \in \prod_{i \in I} M_i$, let [f] denote the equivalence class of f modulo U. Then for every \mathcal{L} -formula $\varphi(v_0, \ldots, v_n)$ and $f_0, \ldots, f_n \in \prod_{i \in I} M_i$,

$$\prod_{i \in I} M_i / U \models \varphi([f_0], \dots, [f_n])$$
$$\iff \{i \in I \mid M_i \models \varphi(f_0(i), \dots, f_n(i))\} \in U.$$

This theorem provides considerable information about ultraproducts and ultrapowers; For instance, ${}^{I}M/U$, the ultrapower of a first order structure M by an ultrafilter U over a set I, is elementarily equivalent to M.

Loś's theorem is a theorem of ZFC, and it turned out that Loś's theorem is not provable from ZF alone: Let BPI be the statement that every Boolean algebra has a prime ideal. Howard [5] showed that Loś's theorem + BPI is equivalent to AC in ZF. It is known that ZF + BPI does not imply AC, so Loś's theorem is not provable from ZF.

Moreover, it was appeared that Los's theorem is strictly weaker than AC. Blass [1] constructed a model of ZF in which there is no non-principal

²⁰²⁰ Mathematics Subject Classification. Primary 03C20, 03E25, 03E35, 03E55.

Key words and phrases. Axiom of Choice, Loś's theorem, Symmetric extension, Ultrapower, Ultraproduct.

ultrafilter. In his model, AC fails but Łoś's theorem holds (in the trivial sense). Hence in ZF, Łoś's theorem does not imply AC.

Here we focus on the statement that "every ultrapower is elementarily equivalent to its source structure". As stated before, this is just a consequence of Loś's theorem, and it seems weaker than Loś's theorem. However, some papers and books (e.g., Howard [5], Howard-Rubin [6]) present this weak statement as Loś's theorem. Actually Howard's proof was carried out with this weak statement, and many consequences of Loś's theorem follow from this weak statement (e.g., see Tachtsis [13]). This situation arises the following natural question, which was also asked in [13].

Question 1.2 (In ZF). Is the weak statement that "every ultrapower is elementarily equivalent to its source structure" really weaker than Loś's fundamental theorem of ultraproducts?

To distinguish various forms of Los's theorem, let us introduce the following definitions: Let U be an ultrafilter over a set I. Let M be a first order structure, and $\{M_i \mid i \in I\}$ an indexed family of first order structures with same language.

- (1) LT1(M, U): M is elementarily equivalent to ${}^{I}M/U$.
- (2) LT2(M, U): The fundamental theorem holds for the ultrapower ${}^{I}M/U$.
- (3) LT3($\{M_i \mid i \in I\}, U$): If $\prod_{i \in I} M_i \neq \emptyset$, then the fundamental theorem holds for the ultraproduct $\prod_{i \in I} M_i/U$.

Let $\mathsf{LT}n$ for n = 1, 2 be the statement that $\mathsf{LT}n(M, U)$ holds for every structure M and ultrafilter U over I, and $\mathsf{LT3}$ be that $\mathsf{LT3}(\{M_i \mid i \in I\}, U)$ holds for every family $\{M_i \mid i \in I\}$ of structures with same language and every ultrafilter U over I. $\mathsf{LT3}$ is Loś's fundamental theorem of ultraproducts, and $\mathsf{LT1}$ is the weak statement in question. The following implications are clear from the definitions:

$$LT2(M,U) \Rightarrow LT1(M,U)$$

and

$$LT3 \Rightarrow LT2 \Rightarrow LT1.$$

In this paper, first we show the following theorem in Section 3:

Theorem 1.3 (In ZF). LT1, LT2, and LT3 are equivalent.

Because of this theorem, we can let LT denote LT1, LT2, and/or LT3.

We now know that LT1 implies LT2, but the difference between LT1(M, U) and LT2(M, U) is worthy of attention, because Spector [12] proved that if Mis a model of ZF¹, then LT1(M, U) is equivalent to LT2(M, U). This result suggests that an implication LT1 \Rightarrow LT2 is in fact *uniform*, that is, the implication LT1(M, U) \Rightarrow LT2(M, U) always holds for every M and U. In Section 4, however, we prove that an implication LT1 \Rightarrow LT2 is not uniform, in fact this uniform implication is equivalent to LT.

Theorem 1.4 (In ZF). (1) If there is a non-principal ultrafilter U over ω but the Countable Choice fails, then there is a first order structure M such that LT1(M,U) holds but LT2(M,U) fails.

¹Not necessary an \in -model.

(2) LT is equivalent to the following statement: For every first order structure M and every ultrafilter U, if LT1(M, U) holds then LT2(M, U) holds as well.

In Sections 5 and 6, we reexamine the fact that LT does not imply AC. Blass's model satisfies LT but the Countable Choice fails in his model. However there is no non-principal ultrafilter in the model, and LT only holds in the trivial sense. This observation lead us to ask the following question.

Question 1.5 (In ZF). Does LT + WUF or LT + WUF(?) deduce (some fragment of) AC? Where WUF is the weak ultrafilter theorem that every infinite set carries a non-principal ultrafilter, and WUF(?) is the statement that there is a non-principal ultrafilter over some set².

Concerning this question, Tachtsis [13] proved that LT + WUF (in fact LT+"there is a non-principal ultrafilter over ω ") implies the Countable Choice, and he asked if LT + WUF implies AC_{fin} , where AC_{fin} is the statement that every family of non-empty finite sets has a choice function.

For this question, in Sections 5 and 6 we provide two answers. First answer is the following. For a cardinal κ and a natural number n, let $AC_{\kappa,n}$ be the statement that every indexed family $\{A_{\alpha} \mid \alpha < \kappa\}$ of non-empty sets with $|A_{\alpha}| \leq n$ has a choice function.

Theorem 1.6. If ZFC is consistent, then so is $ZF + LT + WUF + \neg AC_{\omega_1,2}$.

Hence LT + WUF does not imply $AC_{\omega_1,2}$. This gives an answer to Tachtsis's question.

We give a second answer by using large cardinals:

Theorem 1.7. If ZFC+ "there is a measurable cardinal" is consistent, then so is ZF + LT + $\neg AC_{\omega,2}$ + "there is a measurable cardinal"+ "every ultrafilter is σ -complete".

Thus LT + WUF(?) does not imply $AC_{\omega,2}$, under the consistency of large cardinals.

In Section 7, we prove the following result, which contrasts with Theorem 1.6 in the view point of the consistency strength. It also shows that the large cardinal assumption in Theorem 1.7 cannot be eliminated. Let AC_{ω} denote the Countable Choice.

Theorem 1.8. The following theories are equiconsistent:

- (1) $\mathsf{ZF} + \mathsf{LT} + \mathsf{WUF}(?) + \neg \mathsf{AC}_{\omega,2}$.
- (2) $ZF + LT + WUF(?) + \neg AC_{\omega}$.
- (3) ZFC+ "there is a measurable cardinal".

We also consider the statement that WUF(?)+"every ultrafilter is σ complete". We prove that this statement is equiconsistent with ZFC, but
the statement with LT has a large cardinal strength.

²This notation is due to Herrlich [4].

2. Preliminaries

From now on, our base theory is ZF unless otherwise specified. Throughout this paper, a *filter over a set I* (ultrafilter, respectively) will be a proper filter (proper ultrafilter, respectively) on the Boolean algebra $\mathcal{P}(I)$. An element of an ultrafilter is called *a measure one set*.

A structure is a first order structure with some language \mathcal{L} , and we sometimes identify each symbol in \mathcal{L} with its interpretation in M. In Sections 6 and 7, a set (or class) X will be identified with \in -structure $\langle X; \in \rangle$.

Here we recall ultraproducts and ultrapowers. Let I be a set and U an ultrafilter over I. Let $\{M_i \mid i \in I\}$ be an indexed family of structures with same language \mathcal{L} . For $f, g \in \prod_{i \in I} M_i$, define $f \simeq g$ if $\{i \in I \mid f(i) = g(i)\} \in U$, this is an equivalence relation on $\prod_{i \in I} M_i$. For $f \in \prod_{i \in I} M_i$, let [f] be the equivalence class of f. The ultraproduct of $\{M_i \mid i \in I\}$ by U, denoted by $\prod_{i \in I} M_i/U$, is the structure with language \mathcal{L} defined as in the following manner. The base set of the ultraproduct is the set $\{[f] \mid f \in \prod_{i \in I} M_i\}$. For an *n*-array relation symbol $R \in \mathcal{L}$, its interpretation R^* in $\prod_{i \in I} M_i/U$ is defined as:

$$R^*([f_0],\ldots,[f_n]) \iff \{i \in I \mid M_i \models R(f_0(i),\ldots,f_n(i))\} \in U.$$

We define the interpretations of function symbols and constant symbols by a similar manner.

If every M_i is the same structure M, let us denote $\prod_{i \in I} M_i/U$ by ${}^I M/U$, it is called the *ultrapower of* M by U.

Here we state various forms of Łoś's theorem explicitly.

Definition 2.1. Let U be an ultrafilter over a set I. Let M be a first order structure with language \mathcal{L} , and $\{M_i \mid i \in I\}$ an indexed family of first order structures with language \mathcal{L} . Let $\mathsf{LT1}(M,U)$, $\mathsf{LT2}(M,U)$, and $\mathsf{LT3}(\{M_i \mid i \in I\}, U)$ be the following statements:

- (1) LT1(M, U): M is elementarily equivalent to $^{I}M/U$.
- (2) LT2(M,U): For every \mathcal{L} -formula $\varphi(v_0,\ldots,v_n)$ and $f_0,\ldots,f_n \in {}^{I}M$,

$${}^{I}M/U \models \varphi([f_0], \dots, [f_n]) \\ \iff \{i \in I \mid M \models \varphi(f_0(i), \dots, f_n(i))\} \in U.$$

(3) LT3($\{M_i \mid i \in I\}, U$): If $\prod_{i \in I} M_i \neq \emptyset$, then for every \mathcal{L} -formula $\varphi(v_0, \ldots, v_n)$ and $f_0, \ldots, f_n \in \prod_{i \in I} M_i$,

$$\prod_{i \in I} M_i / U \models \varphi([f_0], \dots, [f_n])$$
$$\iff \{i \in I \mid M_i \models \varphi(f_0(i), \dots, f_n(i))\} \in U.$$

Let $\mathsf{LT}n$ for n = 1, 2 be the statement that $\mathsf{LT}n(M, U)$ holds for every structure M and ultrafilter U, and $\mathsf{LT}3$ be that $\mathsf{LT}3(\{M_i \mid i \in I\}, U)$ holds for every family $\{M_i \mid i \in I\}$ of structures with same language and every ultrafilter U.

We will show that LT1, LT2, and LT3 are equivalent.

A cardinal is a well-orderable cardinal. An ultrafilter U over an infinite cardinal κ is uniform if every measure one set of U has cardinality κ .

For an infinite cardinal κ , an ultrafilter U is κ -complete if for every $\alpha < \kappa$ and $f : \alpha \to U$, we have $\bigcap_{\beta < \alpha} f(\beta) \in U$. U is σ -complete if it is ω_1 -complete. Every ultrafilter is ω -complete, and if U is a principal ultrafilter, then U is κ -complete for every cardinal κ . For an ultrafilter U, if κ is the least cardinal such that U is not κ -complete, then κ must be a successor cardinal.

Let S, T be sets, and U an ultrafilter over a set S. For a map $f: S \to T$, put $f_*(U) = \{X \subseteq T \mid f^{-1}(X) \in U\}$. $f_*(U)$ is an ultrafilter over T, and if U is κ -complete for some cardinal κ , then $f_*(U)$ is also κ -complete.

Lemma 2.2. Let U be an ultrafilter over a set S and κ a cardinal. Suppose κ is the least cardinal such that there is $f : \kappa \to U$ with $\bigcap_{\alpha < \kappa} f(\alpha) \notin U$, that is, U is κ -complete but not κ^+ -complete. Then κ is regular and there is a function $g : S \to \kappa$ such that $g^{-1}(\alpha) \notin U$ for every $\alpha < \kappa$.

Proof. It is easy to show that κ is regular. To find a required function g, we may assume $\bigcap_{\alpha < \kappa} f(\alpha) = \emptyset$. For $\gamma < \kappa$, let $X_{\gamma} = \bigcap_{\alpha < \gamma} f(\alpha)$. By the minimality of κ , we have that $X_{\gamma} \in U$, and $\bigcap_{\gamma < \kappa} X_{\gamma} = \emptyset$. Now define $g: S \to \kappa$ as in the following manner. For $s \in S$, let γ_s be the least ordinal γ with $s \notin X_{\gamma}$. Let $g(s) = \gamma_s$. It is easy to check that this g works. \Box

An uncountable cardinal κ is a *measurable cardinal* if κ carries a κ complete non-principal ultrafilter. Every measurable cardinal is regular.
In ZFC, there is a measurable cardinal if and only if there is a non-principal σ -complete ultrafilter, but this equivalence can fail in ZF, see Section 7.

Fact 2.3. If there is a measurable cardinal, then there is an inner model of ZFC+ "there is a measurable cardinal".

We recall weak forms of AC. For a cardinal κ and a natural number n, let $\mathsf{AC}_{\kappa,n}$ be the statement that every indexed family $\{A_{\alpha} \mid \alpha < \kappa\}$ of nonempty sets with $|A_{\alpha}| \leq n$ has a choice function. AC_{κ} be the statement that every indexed family $\{A_{\alpha} \mid \alpha < \kappa\}$ of non-empty sets has a choice function. AC_{ω} is called *the Countable Choice*. The weak ultrafilter theorem, denoted by WUF, is the statement that every infinite set carries a non-principal ultrafilter. $\mathsf{WUF}(?)$ is that there is a non-principal ultrafilter over some set. Clearly WUF implies $\mathsf{WUF}(?)$.

For a set X and an inner model M of ZF, let M(X) be the inner model of ZF containing X and all elements of M, namely, $M(X) = \bigcup_{\alpha} L(M_{\alpha} \cup \operatorname{trcl}(\{X\}))$ where M_{α} is the set of all sets in M with rank $< \alpha$. The following follows from the construction of M(X).

Fact 2.4. Let M be an inner model of ZF and X a set. If M satisfies AC, then for every set $S \in M(X)$, there is an ordinal γ and a surjection $f \in M$ from ${}^{<\omega}\gamma \times {}^{<\omega}\operatorname{trcl}({X})$ onto S.

3. Łoś's Theorem

In this section we prove that LT1, LT2, and LT3 are equivalent. First we prove the following useful lemma, a similar result was proved by Spector [12].

Lemma 3.1. Let U be an ultrafilter over a set I. Then the following are equivalent:

- (1) LT3($\{M_i \mid i \in I\}, U$) holds for every indexed family $\{M_i \mid i \in I\}$ of structures with same language.
- (2) For every indexed family $\{A_i \mid i \in I\}$ of non-empty sets, there is a function g on I such that $\{i \in I \mid g(i) \in A_i\} \in U$.

Proof. $(2) \Rightarrow (1)$ follows from a standard proof of Loś's fundamental theorem of ultraproducts; To carry out a proof of Loś's theorem without AC, a problematic part is the direction that

$$\{i \in I \mid M_i \models \exists v \varphi(f_0(i), \dots, f_n(i), v)\} \in U$$
$$\Rightarrow \prod_{i \in I} M_i / U \models \exists v \varphi([f_0], \dots, [f_n], v)$$

for $f_0, \ldots, f_n \in \prod_{i \in I} M_i$ and a formula $\varphi(v_0, \ldots, v_n, v)$. Now suppose $\{i \in I \mid M_i \models \exists v \varphi(f_0(i), \ldots, f_n(i), v)\} \in U$. Define A_i as: If $M_i \models \exists v \varphi(f_0(i), \ldots, f_n(i), v)$, then let $A_i = \{x \in M_i \mid M_i \models \varphi(f_0(i), \ldots, f_n(i), x)\}$. Otherwise, let A_i be M_i . By (2), we can find a function g on I such that $\{i \in I \mid g(i) \in A_i\} \in U$. Clearly [g] witnesses that $\prod_{i \in I} M_i/U \models \exists v \varphi([f_0], \ldots, [f_n], v)$.

(1) \Rightarrow (2). For a given family $\{A_i \mid i \in I\}$, fix a point $p \notin \bigcup_{i \in I} A_i$, and let $B_i = A_i \cup \{p\}$. We identify each B_i with some structure. Since $\prod_{i \in I} B_i \neq \emptyset$, we can apply (1) to the ultraproduct $\prod_{i \in I} B_i/U$. For each B_i , obviously $\exists v(v \neq p)$ holds. Let $c_p : I \rightarrow \{p\}$ be the constant function. By (1), we know $\prod_{i \in I} B_i/U \models \exists v(v \neq [c_p])$, hence there is $[g] \in \prod_{i \in I} B_i/U$ such that $\prod_{i \in I} B_i/U \models [g] \neq [c_p]$. Then $\{i \in I \mid g(i) \neq p\} \in U$, so $\{i \in I \mid g(i) \in A_i\} \in U$.

Now we prove Theorem 1.3.

Theorem 3.2. LT1 implies LT3.

Proof. Suppose LT1. Fix an ultrafilter U over I. By Lemma 3.1, it is sufficient to show that for every indexed family $\{A_i \mid i \in I\}$ of non-empty sets, there is a function g on I with $\{i \in I \mid g(i) \in A_i\} \in U$.

Let $S = I \cup \bigcup_{i \in I} A_i$. Define the unary relation $R_1 \subseteq S$ and the binary relation $R_2 \subseteq S^2$ as follows: $R_1(x) \iff x \in I$, and $R_2(x,y) \iff x \in I$ and $y \in A_x$. Let M be the structure $\langle S; R_1, R_2 \rangle$. Let $M^* = \langle S^*; R_1^*, R_2^* \rangle$ be the ultrapower of M by U, where $R_1^* \subseteq S^*$ and $R_2^* \subseteq (S^*)^2$ are relations corresponding to R_1 and R_2 respectively.

By the definition of R_1 and R_2 , we have $M \models \forall x(R_1(x) \to \exists yR_2(x,y))$. By LT1, we also have $M^* \models \forall x(R_1^*(x) \to \exists yR_2^*(x,y))$. Let $\mathrm{id} : I \to I$ be the identity function on I. Since $M \models R_1(\mathrm{id}(i))$ for every $i \in I$, we have $M^* \models R_1^*([\mathrm{id}])$. Hence there is $[g] \in M^*$ such that $M^* \models R_2^*([\mathrm{id}], [g])$. By the definition of the ultrapower, we have $\{i \in I \mid M \models R_2(i, g(i))\} \in U$. Then $\{i \in I \mid g(i) \in A_i\} \in U$ by the definition of R_2 .

From now on, we let LT denote LT1, LT2, and/or LT3. This does not cause any problem by the theorem.

The following are immediate from the proof of Theorem 3.2.

Corollary 3.3. LT is equivalent to the following weak form of LT1: For every ultrafilter U over I and structure M with relational language, LT1(M, U) holds.

4. LT1(M, U) AND LT2(M, U)

In this section, we show that, under some assumption, there are M and U such that LT1(M, U) holds but LT2(M, U) fails. First we prove it under the assumption that ω carries a non-principal ultrafilter but the Countable Choice fails. This is (1) of Theorem 1.4.

Theorem 4.1. Suppose there is a non-principal ultrafilter U over ω but the Countable Choice fails. Then there is a structure M such that LT1(M, U) holds but LT2(M, U) fails.

Note that the assumption of this theorem is consistent; In the first Cohen model, the Countable Choice fails but BPI holds (see [6]). We also note that by this theorem, if LT holds and ω carries a non-principal ultrafilter then the Countable Choice holds, and this result already proved by Tachtsis ([13]).

Proof. Take an indexed family $\{E_n \mid n < \omega\}$ of non-empty sets which has no choice function. We may assume that E_0 is infinite and $\omega \cap \bigcup_{n < \omega} E_n = \emptyset$.

Claim 4.2. There is an infinite set $S, A \subseteq S$, and $B_x \subseteq S$ for $x \in A$ satisfying the following:

- (1) A is countably infinite.
- (2) For each $x \in A$, the set B_x is infinite.
- (3) For distinct $x, x' \in A$, $B_x \cap B_{x'} = \emptyset$.
- (4) $A \cap \bigcup_{x \in A} B_x = \emptyset.$
- (5) $S \setminus (A \cup \bigcup_{x \in A} B_x)$ is infinite.
- (6) For every infinite $A' \subseteq A$, the subfamily $\{B_x \mid x \in A'\}$ has no choice function.

Proof. Let $A = \omega$, $B_x = \{x\} \times \prod_{n \le x} E_n$ for $x \in A$, and $S = (\omega + \omega) \cup \bigcup_{x \in A} B_x$. We check that (6) holds. Let A' be an infinite subset of $A (= \omega)$ and suppose to the contrary that $\{B_x \mid x \in A'\}$ has a choice function f. Define g(n) as follows: Pick the minimal $x \in A'$ with $x \ge n$, and let $f(x) = \langle x, \langle e_0, \ldots, e_x \rangle \rangle$. Then let $g(n) = e_n$. It is clear that g is a choice function of $\{E_n \mid n < \omega\}$, this is a contradiction.

Let S, A, B_x be from the claim. Define the binary relation $R \subseteq S^2$ by $R(x, y) \iff x \in A$ and $y \in B_x$. Let M be the structure $M = \langle S; R \rangle$, and $M^* = \langle S^*; R^* \rangle$ the ultrapower of M by U.

Claim 4.3. There is a function $f : \omega \to S$ and a formula $\varphi(x)$ such that $M \models \varphi(f(n))$ for every $n < \omega$, but $M^* \models \neg \varphi([f])$. Hence LT2(M, U) fails.

Proof. Since A is countably infinite, we can take a bijection $f : \omega \to A$. Consider the formula $\varphi(v)$ that $\exists w R(v, w)$. For each $n < \omega$, we have that $\exists w R(f(n), w)$ holds in M. If $M^* \models \exists w \varphi([f], w)$, then there is $[g] \in S^*$ with $M^* \models R^*([f], [g])$. So $X = \{n \mid R(f(n), g(n))\} = \{n \mid g(n) \in B_{f(n)}\} \in U$, and X is infinite. Then the set $A' = \{f(n) \mid n \in X\}$ is infinite and g induces a choice function of $\{B_x \mid x \in A'\}$, this contradicts (6) of Claim 4.2.

Let $C^* = \{x \in S^* \mid \exists w R^*(x, w)\}$, and for $x \in C^*$, let $D_x^* = \{y \in S^* \mid R^*(x, y)\}$.

Claim 4.4. (1) For $[f] \in S^*$, if $[f] \in C^*$ then $\{n \mid f(n) \in A\} \in U$.

- (2) For distinct $x, x' \in C^*$, we have $D_x^* \cap D_{x'}^* = \emptyset$.
- (3) $C^* \cap \bigcup_{x \in C^*} D_x^* = \emptyset.$

Proof. (1) Since $[f] \in C^*$, there is $[g] \in S^*$ with $R^*([f], [g])$. Then $X = \{n \mid R(f(n), g(n))\} \in U$. For $n \in X$, we know R(f(n), g(n)), thus $f(n) \in A$. Then $X \subseteq \{n \mid f(n) \in A\}$, and $\{n \mid f(n) \in A\} \in U$.

(2) Let [f] = x and [f'] = x'. If there is $[g] \in D_x^* \cap D_{x'}^*$, then $Y = \{n \mid R(f(n), g(n)), R(f'(n), g(n))\} \in U$. For $n \in Y$, we know $g(n) \in B_{f(n)} \cap B_{f'(n)}$, so f(n) = f'(n) must hold. Hence $Y \subseteq \{n \mid f(n) = f'(n)\} \in U$, and [f] = [f']. This is a contradiction.

(3) is similar.

Note that the converse of (1) does not hold; If $f : \omega \to A$ is a bijection, then $\{n \mid f(n) \in A\} \in U$ but $[f] \notin C^*$ by the proof of Claim 4.3.

For $z \in S$, let $c_z : \omega \to \{z\}$ be the constant function. The following is easy to check.

Claim 4.5. If $z \in A$ then $[c_z] \in C^*$. In particular the assignment $z \mapsto [c_z]$ is an injection from A into C^* .

Claim 4.6. $S^* \setminus (C^* \cup \bigcup_{x \in C^*} D_x^*)$ is infinite.

Proof. If $z \in S \setminus (A \cup \bigcup_{x \in A} B_x)$ then it is easy to show that $[c_z] \notin (C^* \cup \bigcup_{x \in C^*} D_x^*)$. Since $S \setminus (A \cup \bigcup_{x \in A} B_x)$ is infinite, we have that $S^* \setminus (C^* \cup \bigcup_{x \in C^*} D_x^*)$ is infinite as well. \Box

Claim 4.7. For every $x \in C^*$, there is $a \in A$ with $[c_a] = x$.

Proof. Let x = [f] and suppose to the contrary that there is no $a \in A$ with $\{n \mid f(n) = c_a(n) = a\} \in U$. Pick $[g] \in S^*$ with $R^*([f], [g])$. We know $Y = \{n \mid f(n) \in A, R(f(n), g(n))\} \in U$. Then we can take a sequence $n_0 < n_1 < \cdots$ from Y such that $f(n_k) \neq f(n_l)$ for every $k < l < \omega$. The map g induces a choice function of the family $\{B_x \mid x \in \{f(n_k) \mid k < \omega\}\}$, this is a contradiction.

Claim 4.8. Let $a \in A$ and $x = [c_a] \in C^*$. Then D_x^* is infinite.

Proof. Take an arbitrary $m < \omega$. Pick m many elements $b_0, \ldots, b_{m-1} \in B_a$. Then $R(a, b_k)$ holds for every k < m, hence $R^*([c_a], [c_{b_k}])$ holds, and $[c_{b_0}], \ldots, [c_{b_{m-1}}] \in D_x^*$.

Finally we prove LT1(M, U) holds.

Claim 4.9. M is elementarily equivalent to M^* .

Proof. Take a generic extension V[G] in which both S and S^* are countable. In V[G], we shall construct an isomorphism from M onto M^* . Then M is elementarily equivalent to M^* in V[G]. On the other hand, the satisfaction relations on M and M^* are absolute between V and V[G], hence we conclude that M is elementarily equivalent to M^* in V.

We work in V[G]. Note the following observations:

- (1) The assignment $a \mapsto [c_a]$ gives a bijection from A onto C^* by Claims 4.5 and 4.7.
- (2) Since $S \setminus (A \cup \bigcup_{x \in A} B_x)$ and $S^* \setminus (C^* \cup \bigcup_{x \in C^*} D_x^*)$ are countably infinite in V[G], there is a bijection between them.

(3) By Claim 4.8, we have both B_a and $D^*_{[c_a]}$ are countably infinite for every $a \in A$. Because S and S^* are countable in V[G], for each $a \in A$, we can take a bijection between B_a and $D^*_{[c_a]}$ in some uniform way.

By these observations, we can take a bijection $j: S \to S^*$ such that:

(1) $j``A = C^*$.

(2) $j^{*}B_x = D_{j(x)}^*$ for every $x \in A$.

(3) $j^{"}(S \setminus (A \cup \bigcup_{x \in A} B_x)) = S^* \setminus (C^* \cup \bigcup_{x \in C^*} D_x^*).$

Then for every $x, y \in S$,

$$R(x,y) \iff y \in B_x \iff j(y) \in D^*_{j(x)} \iff R^*(j(x), j(y)).$$

Thus j is an isomorphism from M onto M^* .

Note 4.10. Under LT, the Countable Choice follows from the existence of a non-principal ultrafilter over ω , but the converse does not hold: Pincus-Solovay [11] constructed a model of ZF in which the Dependent Choice holds but there is no non-principal ultrafilter. In their model, LT holds trivially.

Next, we prove that if LT fails then LT1(M, U) holds but LT2(M, U) fails for some M and U. This immediately shows (2) of Theorem 1.4.

Theorem 4.11. Suppose LT fails. Then there is a structure M and an ultrafilter U over a set I such that LT1(M, U) holds but LT2(M, U) fails.

Proof. By Lemma 3.1, there is an indexed family $\{E_i \mid i \in I\}$ of non-empty sets such that there is no function g on I with $\{i \in I \mid g(i) \in E_i\} \in U$. By using this family, we shall construct a structure M and an ultrafilter U such that $\mathsf{LT1}(M, U)$ holds but $\mathsf{LT2}(M, U)$ fails as in the proof of Theorem 4.1.

First, we may assume that every $i \in I$ is neither an ordinal nor an ordered pair. Let $A = I \cup \omega$.

Claim 4.12. There is an infinite set S and non-empty $B_x \subseteq S$ for $x \in A$ satisfying the following:

(1) $A \subseteq S$.

(2) The set $\{x \in A \mid B_x \text{ is infinite}\}$ is infinite.

- (3) For every positive $n \in \omega$, the set $\{x \in A \mid |B_x| = n\}$ is infinite.
- (4) For distinct $x, x' \in A$, $B_x \cap B_{x'} = \emptyset$.
- (5) $A \cap \bigcup_{x \in A} B_x = \emptyset.$
- (6) $S \setminus (A \cup \bigcup_{x \in A} B_i)$ is infinite.
- (7) There is no function g with dom(g) = I and $\{i \in I \mid g(i) \in B_i\} \in U$.

Proof. Fix an indexed family $\{E_k \mid k < \omega\}$ of non-empty sets such that:

• The set $\{k < \omega \mid E_k \text{ is infinite}\}$ is infinite.

• For every positive $n \in \omega$, the set $\{k < \omega \mid |E_k| = n\}$ is infinite.

For $x \in A = I \cup \omega$, let $B_x = \{x\} \times E_x$. Then take a large set $S \supseteq A \cup \bigcup_{x \in A} B_x$ such that $S \setminus (A \cup \bigcup_{x \in A} B_x)$ is infinite. It is easy to check that S and the B_x 's are as required.

Define $R \subseteq S^2$ by $R(x,y) \iff x \in A$ and $y \in B_x$. Let M be the structure $\langle S; R \rangle$, and $M^* = \langle S^*; R^* \rangle$ the ultrapower of M by U.

Claim 4.13. There is a function $f : I \to S$ and a formula $\varphi(v)$ such that $M \models \varphi(f(i))$ for every $i \in I$, but $M^* \models \neg \varphi([f])$. Hence $\mathsf{LT2}(M, U)$ fails.

Proof. As in the proof of Claim 4.3, consider the formula $\varphi(v)$ that $\exists w R(v, w)$. Let id be the identity function on I. For each $i \in I$, we have that $\exists w R(\operatorname{id}(i), w)$ holds in M. If $M^* \models \exists w \varphi([\operatorname{id}], w)$, then there is $[g] \in S^*$ with $M^* \models R^*([\operatorname{id}], [g])$. Then $\{i \in I \mid R(\operatorname{id}(i), g(i))\} = \{i \in I \mid g(i) \in B_i\} \in U$, this contradicts to (7) of Claim 4.12. \Box

Let $C^* = \{x \in S^* \mid \exists w R^*(x, w)\}$, and $D^*_x = \{y \in S^* \mid R^*(x, y)\}$ for $x \in C^*$. The following claims can be verified easily.

Claim 4.14. (1) For $[f] \in S^*$, if $[f] \in C^*$ then $\{i \in I \mid f(i) \in A\} \in U$. (2) For distinct $x, x' \in C^*$, we have $D_x^* \cap D_{x'}^* = \emptyset$. (3) $C^* \cap \bigcup_{x \in C^*} D_x^* = \emptyset$.

Claim 4.15. $S^* \setminus (C^* \cup \bigcup_{x \in C^*} D^*_x)$ is infinite.

Recall that $c_z: I \to \{z\}$ is the constant function.

Claim 4.16. If $z \in A$ then $[c_z] \in C^*$. In particular the assignment $z \mapsto [c_z]$ is an injection from A into C^* , and C^* is infinite.

Claim 4.17. Let $a \in A$ and $x = [c_a] \in C^*$.

- (1) If B_a is infinite, then D_x^* is infinite as well.
- (2) If B_a is finite, then D_x^* is finite and $|B_x| = |D_x^*|$.

By the claims and the choice of the B_x 's, we have:

Claim 4.18. (1) The set $\{x \in C^* \mid D_x^* \text{ is infinite}\}$ is infinite. (2) For every positive $n \in \omega$, the set $\{x \in C^* \mid |D_x^*| = n\}$ is infinite.

Claim 4.19. M is elementarily equivalent to M^* . Hence LT1(M, U) holds.

Proof. We work in a generic extension V[G] of V in which S and S^* are countably infinite. By the claims above, we have:

- (1) A and C^* are countably infinite.
- (2) The sets $\{x \in A \mid B_x \text{ is countably infinite}\}$ and $\{x \in C^* \mid D_x^* \text{ is countably infinite}\}$ are countably infinite.
- (3) For every positive $n < \omega$, the sets $\{x \in A \mid |B_x| = n\}$ and $\{x \in C^* \mid |D_x^*| = n\}$ are countably infinite.

(4) $S \setminus (A \cup \bigcup_{x \in A} B_x)$ and $S^* \setminus (C^* \cup \bigcup_{x \in C^*} D_x^*)$ are countably infinite. Hence we can construct a bijection $j: S \to S^*$ such that

- (1) $j``A = C^*$.
- (2) $j "B_x = D_{j(x)}^*$ for $x \in A$.
- (3) $j^{"}(S \setminus (A \cup \bigcup_{x \in A} B_x)) = S^* \setminus (C^* \cup \bigcup_{x \in C^*} D_x^*).$

Then j is an isomorphism between M and M^* , hence M is elementarily equivalent to M^* .

To conclude this section, we see another weak form of Loś's theorem. Let M be a structure and U an ultrafilter over a set I. The map $i: M \to {}^{I}M/U$ defined by $i(x) = [c_x]$ is called the *ultrapower embedding* where c_x is the

constant function with value x. Under LT, every ultrapower embedding is an elementary embedding. Now we let LT1.5(M, U) be the statement that the ultrapower embedding $i: M \to {}^{I}M/U$ is an elementary embedding.

It is clear that $LT2(M, U) \Rightarrow LT1.5(M, U) \Rightarrow LT1(M, U)$, and Spector [12] showed that LT2(M, U), LT1.5(M, U), and LT1(M, U) are equivalent if M is a model of ZF. Beside this result, we point out that, if M and U are from the proof of Theorem 4.11, then LT1.5(M, U) holds but LT2(M, U) fails. Here we sketch the proof. Let $i: M \to {}^{I}M/U (= M^*)$ be the ultrapower embedding. Take a formula $\varphi(v_0, \ldots, v_n)$ and $x_0, \ldots, x_n \in M$. In a generic extension V[G] in which S and S^* are countable, we can construct an isomorphism $j: M \to M^*$ such that $i(x_k) = j(x_k)$ for every $k \leq n$. Then

$$M \models \varphi(x_0, \dots, x_n) \iff M^* \models \varphi(j(x_0), \dots, j(x_n))$$
$$\iff M^* \models \varphi(i(x_0), \dots, i(x_n)).$$

So the ultrapower embedding i is an elementary embedding. This result also shows that LT is equivalent to: For every M and U, if LT1.5(M, U) holds then LT2(M, U) holds as well.

However we do not know the answer to the following question:

Question 4.20. (1) Is it consistent that there are M and U such that LT1(M, U) holds but LT1.5(M, U) fails?

(2) Is LT equivalent to the following statement?: For every M and U, if LT1(M, U) holds then LT1.5(M, U) holds as well.

5. Choiceless models of LT (I)

To prove Theorems 1.6 and 1.7, we will use *symmetric extensions* (or *symmetric submodels*). Here we review it. See Jech [7] for details, and we follow Karagila's notations in [9].

Let \mathbb{P} be a poset with maximum element $\mathbb{1}$. For a set x, let \check{x} be a canonical name for x, namely, $\check{x} = \{\langle \check{y}, \mathbb{1} \rangle \mid y \in x\}.$

Every automorphism π on \mathbb{P} induces the isomorphism π on the \mathbb{P} -names, namely, $\pi(\dot{x}) = \{ \langle \pi(\dot{y}), \pi(p) \rangle \mid \langle \dot{y}, p \rangle \in \dot{x} \}$ for a \mathbb{P} -name \dot{x} .

Fact 5.1. Let $p \in \mathbb{P}$, φ be a formula of set theory, and $\dot{x}_0, \ldots, \dot{x}_n \mathbb{P}$ -names. Let π be an automorphism on \mathbb{P} . Then $p \Vdash \varphi(\dot{x}_0, \ldots, \dot{x}_n)$ if and only if $\pi(p) \Vdash \varphi(\pi(\dot{x}_0), \ldots, \pi(\dot{x}_n))$.

Let \mathcal{G} be a subgroup of the automorphism group on \mathbb{P} . A non-empty family \mathcal{F} of subgroups of \mathcal{G} is a *normal filter on* \mathcal{G} if the following hold:

- (1) If $H \in \mathcal{F}$ and H' is a subgroup of \mathcal{G} with $H \subseteq H'$, then $H' \in \mathcal{F}$.
- (2) For $H, H' \in \mathcal{F}$, we have $H \cap H' \in \mathcal{F}$.
- (3) For every $H \in \mathcal{G}$ and $\pi \in \mathcal{G}$, the set $\pi^{-1}H\pi = \{\pi^{-1} \circ \sigma \circ \pi \mid \sigma \in H\}$ is in \mathcal{F} .

A triple $\langle \mathbb{P}, \mathcal{G}, \mathcal{F} \rangle$ is called a *symmetric system*.

For a \mathbb{P} -name \dot{x} , let sym $(\dot{x}) = \{\pi \in \mathcal{G} \mid \pi(\dot{x}) = \dot{x}\}$, which is a subgroup of \mathcal{G} . A name \dot{x} is symmetric if sym $(\dot{x}) \in \mathcal{F}$, and hereditarily symmetric if \dot{x} is symmetric and for every $\langle \dot{y}, p \rangle \in \dot{x}, \dot{y}$ is hereditarily symmetric.

Fact 5.2. If \dot{x} is a hereditarily symmetric name and $\pi \in \mathcal{G}$, then $\pi(\dot{x})$ is also hereditarily symmetric.

For \mathbb{P} -names $\dot{x}_0, \ldots, \dot{x}_n$, let $\{\dot{x}_0, \ldots, \dot{x}_n\}^{\bullet}$ be the \mathbb{P} -name $\{\langle \dot{x}_0, \mathbb{1} \rangle, \ldots, \langle \dot{x}_n, \mathbb{1} \rangle\}$, and $\langle \dot{x}_0, \dot{x}_1 \rangle^{\bullet} = \{\{\dot{x}_0\}^{\bullet}, \{\dot{x}_0, \dot{x}_1\}^{\bullet}\}^{\bullet}$. The name $\{\dot{x}_0, \ldots, \dot{x}_n\}^{\bullet}$ is a canonical name for the family of the interpretations of the \dot{x}_i 's, and $\langle \dot{x}_0, \dot{x}_1 \rangle^{\bullet}$ is for the ordered pair. If $\dot{x}_0, \ldots, \dot{x}_n$ are hereditarily symmetric, then so are $\{\dot{x}_0, \ldots, \dot{x}_n\}^{\bullet}$ and $\langle \dot{x}_0, \dot{x}_1 \rangle^{\bullet}$.

Let HS be the class of all hereditarily symmetric names. For a (V, \mathbb{P}) generic G, let HS^G be the class of all interpretations of hereditarily symmetric names. HS^G is a transitive model of ZF with $V \subseteq \mathrm{HS}^G \subseteq V[G]$. HS^G is
called a symmetric extension of V.

Here we construct our symmetric extension. Starting from a model of ZFC, fix a regular cardinal κ . In a κ -closed forcing extension of V, we shall construct a symmetric extension satisfying the following:

- (1) AC_{λ} holds for every $\lambda < \kappa$.
- (2) Every infinite cardinal $\lambda < \kappa$ carries a uniform ultrafilter.
- (3) Every ultrafilter over κ has a measure one set with size $< \kappa$.
- (4) If $0^{\#}$ does not exist, then for every cardinal $\lambda > \kappa$ and ultrafilter U over λ , U has a measure one set with size $< \kappa$.
- (5) $AC_{\kappa,2}$ fails.

 $0^{\#}$ is a set of natural numbers which code true formulas about indiscernibles in the Gödel's constructible universe L. The existence of $0^{\#}$ is a large cardinal property, e.g., if $0^{\#}$ exists, then every uncountable cardinal is inaccessible in L. In particular $0^{\#}$ does not exist in L. It is also known that if a measurable cardinal exists, then $0^{\#}$ exists. See Kanamori [8] for details of $0^{\#}$.

Our symmetric extension is a variant of Feferman's model (see [6] or [7]), and essentially the same construction is used in Hayut-Karagila [3].

We notice that our symmetric extension may not be a model of Theorems 1.6 and 1.7. After constructing symmetric extension, we will take an intermediate model between the ground model and a symmetric extension, which is a required model. This argument is based on Blass's one in [1].

For a set X, let $\operatorname{Fn}(X, 2, < \kappa)$ be the poset of all partial functions $p : X \to 2$ with size $< \kappa$. The order is given by the reverse inclusion. It is a κ -closed poset. We define a symmetric system $\langle \mathbb{P}, \mathcal{G}, \mathcal{F} \rangle$ as follows. Let $\mathbb{P} = \operatorname{Fn}(\kappa \times \kappa, 2, < \kappa)$. Let $\vec{X} = \langle X_{\alpha} \mid \alpha < \kappa \rangle$ be a sequence of subsets of κ . Define the automorphism $\pi_{\vec{X}}$ on \mathbb{P} by : dom $(\pi_{\vec{X}}(p)) = \operatorname{dom}(p)$, and

$$\pi_{\vec{X}}(p)(\alpha,\beta) = \begin{cases} 1 - p(\alpha,\beta) & \text{if } \beta \in X_{\alpha}, \\ p(\alpha,\beta) & \text{if } \beta \notin X_{\alpha}. \end{cases}$$

We note that for \vec{X} and \vec{Y} ,

$$\pi_{\vec{Y}}(\pi_{\vec{X}}(p))(\alpha,\beta) = \begin{cases} 1 - p(\alpha,\beta) & \text{if } \beta \in X_{\alpha} \triangle Y_{\alpha}, \\ p(\alpha,\beta) & \text{if } \beta \notin X_{\alpha} \triangle Y_{\alpha}. \end{cases}$$

Where $X_{\alpha} \triangle Y_{\alpha}$ is the symmetric difference of X_{α} and Y_{α} . Let \mathcal{G} be the group of automorphisms σ which is of the form $\pi_{\vec{X}}$ for some sequence \vec{X} of subsets of κ . For a set $e \in [\kappa]^{<\kappa}$, let $\operatorname{fix}(e) = \{\pi_{\vec{X}} \mid X_{\alpha} = \emptyset$ for every $\alpha \in e\}$. fix(e) is a subgroup of \mathcal{G} . Let \mathcal{F} be the filter on \mathcal{G} generated by

{fix(e) $| e \in [\kappa]^{<\kappa}$ }, that is, for a subgroup $H \subseteq \mathcal{G}, H \in \mathcal{F} \iff \text{fix}(e) \subseteq H$ for some set $e \in [\kappa]^{<\kappa}$. It is routine to check that \mathcal{F} is a normal filter on \mathcal{G} . For $e \subseteq \kappa$ and $p \in \mathbb{P}$, let $p \upharpoonright e$ be the function $p \cap (e \times \kappa \times 2)$, which is in

 $\operatorname{Fn}(e \times \kappa, 2, < \kappa).$

Lemma 5.3. Let $\dot{x}_0, \ldots, \dot{x}_n$ be hereditarily symmetric names and $e \in [\kappa]^{<\kappa}$ with fix $(e) \subseteq \bigcap_{i \leq n} \operatorname{sym}(\dot{x}_i)$. Then for every $p \in \mathbb{P}$ and formula $\varphi(v_0, \ldots, v_n)$ of set theory, if p forces $\varphi(\dot{x}_0, \ldots, \dot{x}_n)$, then $p \upharpoonright e$ also forces $\varphi(\dot{x}_0, \ldots, \dot{x}_n)$.

Proof. Suppose $p \Vdash \varphi(\dot{x}_0, \ldots, \dot{x}_n)$, but there is $q \leq p \upharpoonright e$ with $q \Vdash \neg \varphi(\dot{x}_0, \ldots, \dot{x}_n)$. Let $d = \{\alpha \in \kappa \setminus e \mid \exists \beta(\langle \alpha, \beta \rangle \in \operatorname{dom}(p))\}$. Take a sequence $\langle X_\alpha \mid \alpha < \kappa \rangle$ of subsets of κ such that $X_\alpha = \emptyset$ for $\alpha \notin d$, and for $\alpha \in d$, $X_\alpha = \{\beta < \kappa : p(\alpha, \beta) \neq q(\alpha, \beta)\}$. It is easy to check that $\pi_{\vec{X}}(q)$ is compatible with $p, \pi_{\vec{X}}(\dot{x}_i) = \dot{x}_i$ for $i \leq n$, but $\pi_{\vec{X}}(q) \Vdash \neg \varphi(\pi_{\vec{X}}(\dot{x}_0), \ldots, \pi_{\vec{X}}(\dot{x}_n))$, this is a contradiction. \Box

Take a (V, \mathbb{P}) -generic G and work in V[G]. Let HS^G be the class of all interpretations of hereditarily symmetric names.

Since \mathbb{P} is κ -closed, we have $([\kappa]^{<\kappa})^V = ([\kappa]^{<\kappa})^{V[G]}$. Hence if $\mu < \kappa$ is a cardinal and $U \in V$ is an ultrafilter over μ in V, then U remains an ultrafilter in V[G]. Because $V \subseteq \mathrm{HS}^G \subseteq V[G]$, we have:

Lemma 5.4. In HS^G, every infinite cardinal $\mu < \kappa$ carries a uniform ultrafilter.

For $\alpha < \kappa$, let $G_{\alpha} = G \cap \operatorname{Fn}(\alpha \times \kappa, 2, < \kappa)$. G_{α} is $(V, \operatorname{Fn}(\alpha \times \kappa, 2, < \kappa))$ -generic.

Lemma 5.5. (1) $V[G_{\alpha}] \subseteq HS^G$ for every $\alpha < \kappa$.

(2) For every ordinal γ and $x \in \mathcal{P}(\gamma)^{\mathrm{HS}^G}$, there is $\alpha < \kappa$ with $x \in V[G_{\alpha}]$.

Proof. (1) G_{α} can be coded in some element of $\mathcal{P}(\kappa)^{V[G_{\alpha}]}$, and every element x of $\mathcal{P}(\kappa)^{V[G_{\alpha}]}$ has a $\operatorname{Fn}(\alpha \times \kappa, 2, < \kappa)$ -name \dot{x} , which is hereditarily symmetric with $\operatorname{fix}(\alpha) \subseteq \operatorname{sym}(\dot{x})$. Hence $G_{\alpha} \in \operatorname{HS}^{G}$, so $V[G_{\alpha}] \subseteq \operatorname{HS}^{G}$.

(2) Take a hereditarily symmetric name \dot{x} for x. Take $e \in [\kappa]^{<\kappa}$ with fix $(e) \subseteq \operatorname{sym}(\dot{x})$, and take a large $\alpha < \kappa$ with $\alpha > \sup(e)$. By Lemma 5.3, we have that for every $\xi < \gamma$,

$$\xi \in x \iff p \Vdash \check{\xi} \in \dot{x} \text{ for some } p \in G_{\alpha}.$$

Hence x is definable in $V[G_{\alpha}]$, and $x \in V[G_{\alpha}]$.

Let $x_{\alpha} = \{\beta \mid \exists p \in G(p(\alpha, \beta) = 1)\}$. The name $\dot{x}_{\alpha} = \{\langle \check{\beta}, p \rangle \mid p \in \mathbb{P}, p(\alpha, \beta) = 1\}$ is a canonical hereditarily symmetric name for x_{α} with $\operatorname{fix}(\{\alpha\}) \subseteq \operatorname{sym}(\dot{x}_{\alpha})$, in particular $x_{\alpha} \in \operatorname{HS}^{G}$. For a sequence $\vec{X} = \langle X_{\alpha} \mid \alpha < \kappa \rangle$ of subsets of κ and $\alpha < \kappa$, if $\kappa \setminus X_{\alpha} \subseteq \xi$ for some $\xi < \kappa$, then $\Vdash \pi_{\vec{X}}(\dot{x}_{\alpha}) \cap \dot{x}_{\alpha} \subseteq \check{\xi}$. Similarly, if $X_{\alpha} \subseteq \xi$ for some $\xi < \kappa$, then $\Vdash \pi_{\vec{X}}(\kappa \setminus \dot{x}_{\alpha}) \cap (\kappa \setminus \dot{x}_{\alpha}) \subseteq \check{\xi}$.

Lemma 5.6. Let $U \in HS^G$ be an ultrafilter over κ in HS^G . Then U has a measure one set with size $< \kappa$.

Proof. Take a hereditarily symmetric name U for U, and take $e \in [\kappa]^{<\kappa}$ with fix $(e) \subseteq \operatorname{sym}(\dot{U})$. Fix a large $\alpha < \kappa$ with $\alpha > \sup(e)$. If $x_{\alpha} \in U$, pick $p \in G$ such that $p \Vdash \dot{x}_{\alpha} \in \dot{U}$. Take a large $\xi < \kappa$ such that $\{\beta \mid \langle \alpha, \beta \rangle \in f\}$

dom(p)} $\subseteq \xi$. Fix a sequence $\vec{X} = \langle X_{\delta} | \delta < \kappa \rangle$ such that $X_{\delta} = \emptyset$ if $\delta \neq \alpha$, and $X_{\alpha} = \kappa \setminus \xi$. Then $\pi_{\vec{X}}(p) = p$ and $p \Vdash \pi_{\vec{X}}(\dot{x}_{\alpha}) \in \pi_{\vec{X}}(\dot{U}) = \dot{U}$ and $\pi_{\vec{X}}(\dot{x}_{\alpha}) \cap \dot{x}_{\alpha} \subseteq \check{\xi}$." Hence $p \Vdash \check{\xi} \in \dot{U}$. The case that $x_{\alpha} \notin U$ is similar. \Box

To prove Lemma 5.9 below, we need the notion of *regular ultrafilters*.

Definition 5.7. Let U be an ultrafilter over a set S, and λ, μ infinite cardinals. U is said to be (λ, μ) -regular if there is an indexed family $\{X_{\alpha} \mid \alpha < \mu\} \subseteq U$ such that $\bigcap_{\alpha \in A} X_{\alpha} = \emptyset$ for every $A \in [\mu]^{\lambda}$.

Fact 5.8 (Donder [2], in ZFC). Let ν be a cardinal $\geq \omega_1$. If there is a uniform ultrafilter over ν which is not (ω, μ) -regular for some cardinal $\mu < \nu$, then there is an inner model of a measurable cardinal. In particular $0^{\#}$ exists.

Lemma 5.9. Suppose $0^{\#}$ does not exist. In HS^G , let $\lambda > \kappa$ be an infinite cardinal and U an ultrafilter over λ . Then U has a measure one set with size $< \kappa$.

Proof. We work in HS^G . We prove it by induction on $\lambda \ge \kappa$. The base step $\lambda = \kappa$ is Lemma 5.6.

Let $\lambda > \kappa$ and U an ultrafilter over λ . We shall prove that there is $X \in U$ with $|X| < \lambda$. Then, by applying the induction hypothesis, we can take a measure one set of U with size $< \kappa$.

Suppose to the contrary that $|X| = \lambda$ for every $X \in U$, that is, U is a uniform ultrafilter.

Case 1: There is $f : \lambda \to \kappa$ such that $f^{-1}(Y) \notin U$ for every set $Y \in [\kappa]^{<\kappa}$. In this case, consider the ultrafilter $f_*(U) = \{Y \subseteq \kappa \mid f^{-1}(Y) \in U\}$. By the assumption, $f_*(U)$ forms a uniform ultrafilter over κ , this contradicts to Lemma 5.6.

Case 2: For every $f : \lambda \to \kappa$, there is a set $Y \in [\kappa]^{<\kappa}$ such that $f^{-1}(Y) \in U$. Take a hereditarily symmetric name \dot{U} for U, and an ordinal $\alpha < \kappa$ with $\operatorname{fix}(\alpha) \subseteq \operatorname{sym}(\dot{U})$. For every $X \in \mathcal{P}(\lambda)^{V[G_{\alpha}]}$, we can find a $\operatorname{Fn}(\alpha \times \kappa, 2, <\kappa)$ -name \dot{X} for X. \dot{X} is hereditarily symmetric with $\operatorname{fix}(\alpha) \subseteq \operatorname{sym}(\dot{X})$. By Lemma 5.3, we have that for every $X \in \mathcal{P}(\lambda)^{V[G_{\alpha}]}$,

 $X \in U \iff p \Vdash \dot{X} \in \dot{U}$ for some $p \in G_{\alpha}$.

Thus we have $U \cap V[G_{\alpha}] \in V[G_{\alpha}]$. Let $U' = U \cap V[G_{\alpha}]$, which is a uniform ultrafilter over λ in $V[G_{\alpha}]$.

Subcase 1: $\kappa = \omega$. In this case, it is easy to check that U' is σ -complete in $V[G_{\alpha}]$, hence U' is a σ -complete uniform ultrafilter over λ . This means, in $V[G_{\alpha}]$, there is a measurable cardinal $\leq \lambda$, hence $0^{\#}$ exists in $V[G_{\alpha}]$. This contradicts to the assumption.

Subcase 2: $\kappa > \omega$. We prove that U' is not (ω, κ) -regular in $V[G_{\alpha}]$. Since $V[G_{\alpha}]$ is a model of ZFC, we conclude that $0^{\#}$ exists by Fact 5.9, this is also impossible.

We work in $V[G_{\alpha}]$. Suppose to the contrary that U' is (ω, κ) -regular, and take $\{X_{\alpha} \mid \alpha < \kappa\} \subseteq U'$ such that $\bigcap_{\alpha \in A} X_{\alpha} = \emptyset$ for every $A \in [\kappa]^{\omega}$. Define $g: \lambda \to \kappa$ as follows. For $\beta < \lambda$, by the choice of the A_{α} 's, the set $\{\alpha < \kappa \mid \beta \in A_{\alpha}\}$ is bounded in κ . Let $g(\beta) = \sup\{\alpha < \kappa \mid \beta \in A_{\alpha}\} < \kappa$. By the assumption, in HS^{G} , there is a set $Y \in [\kappa]^{<\kappa}$ such that $g^{-1}(Y) \in U$. We know $Y \in V[G_{\alpha}]$ since Y has cardinality $< \kappa$, so $g^{-1}(Y) \in V[G_{\alpha}]$ and $g^{-1}(Y) \in U'$. Pick $\gamma < \kappa$ with $\gamma > \sup(Y)$. By the definition of g, we know that $\beta \notin A_{\gamma}$ for every $\beta \in g^{-1}(Y)$, hence $A_{\gamma} \cap g^{-1}(Y) = \emptyset$. This is a contradiction.

To check that $\mathsf{AC}_{\kappa,2}$ fails in HS^G , we need more. For $\alpha < \kappa$, the name $\dot{x}_{\alpha} = \{\langle \check{\beta}, p \rangle \mid p \in \mathbb{P}, p(\alpha, \beta) = 1\}$ is a canonical hereditarily symmetric name for \dot{x}_{α} with fix $(\{\alpha\}) \subseteq \operatorname{sym}(\dot{x}_{\alpha})$. Then for $\pi_{\vec{X}} \in \mathcal{G}$, the name $\pi_{\vec{X}}(\dot{x}_{\alpha})$ is a hereditarily symmetric name for the symmetric difference $x_{\alpha} \bigtriangleup X_{\alpha}$ with fix $(\{\alpha\}) \subseteq \operatorname{sym}(\pi_{\vec{X}}(\dot{x}_{\alpha}))$.

Now fix a uniform ultrafilter W over κ in V. Let d_{α} be the set $\{x_{\alpha} \bigtriangleup Y \mid Y \in \mathcal{P}(\kappa)^{V}, Y \notin W\}$, and \dot{d}_{α} the name $\{\langle \pi_{\vec{X}}(\dot{x}_{\alpha}), \mathbb{1} \rangle \mid \pi_{\vec{X}} \in \mathcal{G}, X_{\alpha} \notin W\}$. \dot{d}_{α} is a hereditarily symmetric name for d_{α} . Similarly, let e_{α} be the set $\{x_{\alpha} \bigtriangleup Y \mid Y \in \mathcal{P}(\kappa)^{V}, Y \in W\}$, and \dot{e}_{α} the name $\{\langle \pi_{\vec{X}}(\dot{x}_{\alpha}), \mathbb{1} \rangle \mid \pi_{\vec{X}} \in \mathcal{G}, X_{\alpha} \in W\}$. \dot{e}_{α} is a hereditarily symmetric name for e_{α} . We note that $d_{\alpha}, e_{\alpha} \in V[G_{\alpha+1}]$ and $d_{\alpha} \cap e_{\alpha} = \emptyset$.

The following is straightforward:

Lemma 5.10. (1)
$$\pi_{\vec{X}}(\dot{d}_{\alpha}) = \begin{cases} d_{\alpha} & \text{if } X_{\alpha} \notin W, \\ \dot{e}_{\alpha} & \text{if } X_{\alpha} \in W. \end{cases}$$

(2) $\pi_{\vec{X}}(\dot{e}_{\alpha}) = \begin{cases} \dot{e}_{\alpha} & \text{if } X_{\alpha} \notin W, \\ \dot{d}_{\alpha} & \text{if } X_{\alpha} \in W. \end{cases}$

Define the function F_0 on κ by $F_0(\alpha) = \{d_\alpha, e_\alpha\}$ for $\alpha < \kappa$.

Lemma 5.11. $F_0 \in \mathrm{HS}^G$.

Proof. By Lemma 5.10, the name $\{\dot{d}_{\alpha}, \dot{e}_{\alpha}\}^{\bullet}$ is hereditarily symmetric with $\operatorname{sym}(\{\dot{d}_{\alpha}, \dot{e}_{\alpha}\}^{\bullet}) = \mathcal{G}$. Hence the name $\dot{F}_{0} = \{\langle \langle \check{\alpha}, \{\dot{d}_{\alpha}, \dot{e}_{\alpha}\}^{\bullet} \rangle^{\bullet}, \mathbb{1} \rangle \mid \alpha < \kappa\}$, which is a name for F_{0} , is hereditarily symmetric. So we have $F_{0} \in \operatorname{HS}^{G}$. \Box

Lemma 5.12. $AC_{\kappa,2}$ fails in HS^G .

Proof. Let $E_{\alpha} = \{d_{\alpha}, e_{\alpha}\}$ for $\alpha < \kappa$, and consider the indexed family $\{E_{\alpha} \mid \alpha < \kappa\}$, which is in HS^{G} because $F_{0} \in \mathrm{HS}^{G}$. We show that this family has no choice function in HS^{G} .

Suppose to the contrary that the family has a choice function. Take a choice function $f \in \mathrm{HS}^G$, and let \dot{f} be a hereditarily symmetric name for f. Take $p \in G$ and $\alpha < \kappa$ such that $\mathrm{fix}(\alpha) \subseteq \mathrm{sym}(\dot{f})$ and p decides the value of $\dot{f}(\check{\alpha})$, say $p \Vdash \dot{f}(\check{\alpha}) = \dot{d}_{\alpha}$. Fix a large $\beta < \kappa$ with $\mathrm{dom}(p) \cap (\{\alpha\} \times \kappa) \subseteq \{\alpha\} \times \beta$. In V, take a sequence \vec{X} such that $X_{\gamma} = \emptyset$ if $\gamma \neq \alpha$, and $X_{\alpha} = \kappa \setminus \beta$. We have $\pi_{\vec{X}}(p) = p$, and since $X_{\alpha} \in W$, we have $p \Vdash \dot{f}(\check{\alpha}) = \pi_{\vec{X}}(\dot{f})(\check{\alpha}) = \pi_{\vec{X}}(\dot{d}_{\alpha}) = \dot{e}_{\alpha}$ by Lemma 5.10. Hence $p \Vdash \dot{d}_{\alpha} = \dot{e}_{\alpha}$, this is a contradiction. \Box

As mentioned before, our symmetric extension HS^G may not be a model of Theorem 1.6 and 1.7, because we do not know if LT holds in HS^G . Instead, we take an intermediate model M between V and HS^G such that M satisfies not only (1)–(5) but also the following (6) and (7):

- (6) Every ultrafilter has a well-orderable measure one set.
- (7) If $0^{\#}$ does not exist, then every ultrafilter has a well-orderable measure one set with size $< \kappa$.

For $\alpha < \kappa$, if \dot{x} is a \mathbb{P} -name which is of the form $\{\langle \dot{\beta}, p \rangle \mid \beta < \kappa, p \in A_{\beta}\}$ for some $A_{\beta} \subseteq \operatorname{Fn}(\alpha \times \kappa, 2, < \kappa)$, then \dot{x} is a hereditarily symmetric name with fix $(\alpha) \subseteq \operatorname{sym}(\dot{x})$. Let Q_{α} be the set of such names, and $\dot{P}_{\alpha} = \{\langle \dot{x}, 1 \rangle \mid \dot{x} \in Q_{\alpha}\}$. \dot{P}_{α} is a hereditarily symmetric name for $\mathcal{P}(\kappa)^{V[G_{\alpha}]}$ with $\operatorname{sym}(\dot{P}_{\alpha}) = \mathcal{G}$. Using this, we have:

Lemma 5.13. The function F_1 on κ defined by $F_1(\alpha) = \mathcal{P}(\kappa)^{V[G_\alpha]}$ is in HS^G .

Our final model M is $V(\{F_0, F_1\})$, which is the minimal inner model of ZF containing F_0, F_1 and all elements of V. M is an intermediate model between V and HS^G . We note that $V[G_\alpha] \subseteq M$ for every $\alpha < \kappa$ since $G_\alpha \in M$. We show that M satisfies the properties (1)–(5), and (6), (7).

For every ordinal γ , we have $\mathcal{P}(\gamma)^{\hat{M}} = \mathcal{P}(\gamma)^{\text{HS}^{\hat{G}}}$ by Lemma 5.5. By Lemmas 5.4, 5.9, and this observation, we have:

Lemma 5.14. The following hold in M:

- (1) For every infinite cardinal $\lambda < \kappa$, λ carries a uniform ultrafilter.
- (2) Every ultrafilter over κ has a measure one set with size $< \kappa$.
- (3) If $0^{\#}$ does not exist, then for every cardinal $\lambda > \kappa$ and ultrafilter U over λ , U has a measure one set with size $< \kappa$.

Since $F_0 \in M \subseteq \mathrm{HS}^G$, we have:

Lemma 5.15. $AC_{\kappa,2}$ fails in M.

Define the function H on κ by $H(\alpha) = F_1(\alpha) \cup \bigcup_{\beta < \alpha} F_0(\beta) = \mathcal{P}(\kappa)^{V[G_\alpha]} \cup \{d_\beta, e_\beta \mid \beta < \alpha\}$. Note that $H \in M$, $H(\alpha) \in V[G_\alpha]$, and whenever $\alpha < \beta < \kappa$ we have $H(\alpha) \subseteq H(\beta)$.

Lemma 5.16. In M, for every set $S \in M$, there is an ordinal γ and a surjection from $\gamma \times {}^{<\omega}(\bigcup_{\alpha < \kappa} H(\alpha))$ onto S.

Proof. First we show that, in M, there is a surjection from $\kappa \times \bigcup_{\alpha < \kappa} H(\alpha)$ onto $\operatorname{trcl}(\{F_0, F_1\})$. Since $\operatorname{trcl}(\{F_0, F_1\}) = \{F_0, F_1\} \cup \operatorname{trcl}(F_0) \cup \operatorname{trcl}(F_1)$, it is sufficient to show that there is a surjection from $\kappa \times \omega \times \bigcup_{\alpha < \kappa} H(\alpha)$ onto $\operatorname{trcl}(F_0)$ and from $\kappa \times \omega \times \bigcup_{\alpha < \kappa} H(\alpha)$ onto $\operatorname{trcl}(F_1)$. We show only for $\operatorname{trcl}(F_0)$, we can take a surjection onto $\operatorname{trcl}(F_1)$ by a similar way.

Recall that for a set X, the transitive closure $\operatorname{trcl}(X)$ is defined as follows: $X_0 = X, X_{n+1} = \bigcup X_n$, and $\operatorname{trcl}(X) = \bigcup_n X_n$. Hence the set $\operatorname{trcl}(F_0)$ is the union:

$$\begin{aligned} \left\{ \langle \alpha, \{d_{\alpha}, e_{\alpha}\} \rangle \mid \alpha < \kappa \right\} \cup \left\{ \{\alpha\}, \{\alpha, \{d_{\alpha}, e_{\alpha}\} \} \mid \alpha < \kappa \right\} \\ \cup \left\{ \alpha \mid \alpha < \kappa \right\} \cup \left\{ \{d_{\alpha}, e_{\alpha}\} \mid \alpha < \kappa \right\} \\ \cup \left\{ d_{\alpha}, e_{\alpha} \mid \alpha < \kappa \right\} \cup \bigcup_{\alpha < \kappa} (d_{\alpha} \cup e_{\alpha}). \end{aligned}$$

Define $f : \kappa \times \omega \times \bigcup_{\alpha < \kappa} H(\alpha) \to \operatorname{trcl}(F_0)$ in M as follows.

- $f(\alpha, 0, x) = \langle \alpha, \{d_{\alpha}, e_{\alpha}\} \rangle.$
- $f(\alpha, 1, x) = \{\alpha\}$ and $f(\alpha, 2, x) = \{\alpha, \{d_{\alpha}, e_{\alpha}\}\}.$
- $f(\alpha, 3, x) = \alpha$ and $f(\alpha, 4, x) = \{d_\alpha, e_\alpha\}.$
- $f(\alpha, n, x) = x$ for $n \ge 5$.

It is straightforward to check that this f works.

Now fix a set $S \in M$. By Fact 2.4, there is a function $f' \in M$ and an ordinal γ' such that f' is a surjection from ${}^{<\omega}\gamma' \times {}^{<\omega}(\operatorname{trcl}(\{F_0, F_1\}))$ onto S. We have known that there is a surjection from $\kappa \times \bigcup_{\alpha < \kappa} H(\alpha)$ onto $\operatorname{trcl}(\{F_0, F_1\})$. Hence, by modifying f' in M, we can take a map f and an ordinal γ such that f is a surjection from $\gamma \times {}^{<\omega}(\bigcup_{\alpha < \kappa} H(\alpha))$ onto S. \Box

We now check that M satisfies the condition (1).

Lemma 5.17. In V[G], M is closed under $< \kappa$ -sequences. In particular AC_{λ} holds for every $\lambda < \kappa$ in M.

Proof. Take $\lambda < \kappa$ and a set $\{y_{\alpha} \mid \alpha < \lambda\} \subseteq M$. In M, we can take a large set S, a map f, and an ordinal γ such that $\{y_{\alpha} \mid \alpha < \kappa\} \subseteq S$ and f is a surjection from $\gamma \times^{<\omega} (\bigcup_{\alpha < \kappa} H(\alpha))$ onto S. Then there is $\delta < \kappa$ such that $\{y_{\alpha} \mid \alpha < \lambda\} \subseteq f^{*}(\gamma \times {}^{<\omega}H(\delta))$. In M, $H(\delta)$ is well-orderable; $H(\delta) \in V[G_{\delta}] \subseteq M$ and $H(\delta)$ is well-orderable in $V[G_{\delta}]$. In particular $\gamma \times^{<\omega} H(\delta)$ is also well-orderable in M. So, by modifying f in M, we can take an ordinal θ and a map g such that $\operatorname{dom}(g) = \theta$ and $\{y_{\alpha} \mid \alpha < \lambda\} \subseteq \operatorname{range}(g)$. In V[G], pick $\xi_{\alpha} < \theta$ with $g(\xi_{\alpha}) = y_{\alpha}$. Then $\langle \xi_{\alpha} \mid \alpha < \lambda \rangle \in V \subseteq M$, hence $\{y_{\alpha} \mid \alpha < \mu\} = g^{*}\{\xi_{\alpha} \mid \alpha < \lambda\} \in M$.

Next we show that M satisfies the conditions (6) and (7).

Lemma 5.18. In M, let S be an infinite set, and U an ultrafilter over S. Then U has a well-orderable measure one set. In addition, if $0^{\#}$ does not exist, then U has a well-orderable measure one set with size $< \kappa$.

Proof. We work in M. Take a map f and an ordinal γ such that f is a surjection from $\gamma \times {}^{<\omega}(\bigcup_{\alpha < \kappa} H(\alpha))$ onto S. For $\alpha < \kappa$, let $S_{\alpha} = f^{"}(\gamma \times {}^{<\omega}H(\alpha))$. We know $S = \bigcup_{\alpha < \kappa} S_{\alpha}$, and

For $\alpha < \kappa$, let $S_{\alpha} = f''(\gamma \times {}^{<\omega}H(\alpha))$. We know $S = \bigcup_{\alpha < \kappa} S_{\alpha}$, and $\langle S_{\alpha} \mid \alpha < \kappa \rangle \in M$. Define $g: S \to \kappa$ as g(x) is the least $\alpha < \kappa$ with $x \in S_{\alpha}$. Consider the ultrafilter $g_*(U)$ over κ . We can find an ordinal $\alpha < \kappa$ with $\alpha \in g_*(U)$ by Lemma 5.14, so we have $g^{-1}(\alpha) \subseteq S_{\alpha} = f''(\gamma \times {}^{<\omega}H(\alpha)) \in U$. Hence, we may assume that U is an ultrafilter over S_{α} . In $M, H(\alpha)$ is well-orderable. Hence, in $M, S_{\alpha} = f''(\gamma \times {}^{<\omega}H(\alpha))$ is a well-orderable measure one set of U. Moreover, if $0^{\#}$ does not exist, then by Lemma 5.9, U has a well-orderable measure one set with size $< \kappa$.

We are ready to prove Theorem 1.6. We use the well-known fact that if the Countable Choice holds then every infinite set has a countably infinite subset.

Theorem 5.19. If ZFC is consistent, then so is $ZF + LT + WUF + \neg AC_{\omega_{1},2}$.

Proof. Assuming $AC + 0^{\#}$ does not exist" (e.g., suppose V = L), take the model M as the above with the case $\kappa = \omega_1$. The following hold in M:

- (1) The Countable Choice holds (Lemma 5.17).
- (2) ω carries a non-principal ultrafilter (Lemma 5.14).
- (3) Every ultrafilter has a countable measure one set (Lemma 5.18).
- (4) $AC_{\omega_{1},2}$ fails (Lemma 5.15).

By (1) and (3) with Lemma 3.1, LT holds. By (1), every infinite set S has a countably infinite subset S', and (2) guarantees that there is a non-principal ultrafilter U' over S'. Then we can extend U' to a non-principal ultrafilter U over S.

The resulting model of Theorem 5.19 has non-principal ultrafilters, but no uniform ultrafilter over uncountable cardinals.

Question 5.20. Is $ZF+LT+\neg AC+$ "every infinite cardinal carries a uniform ultrafilter" relatively consistent with ZFC?

We do know the answer to this question, but we prove that the statement LT+ "every infinite cardinal carries a uniform ultrafilter" deduces more fragments of AC than the Countable Choice.

Proposition 5.21. Suppose LT holds. Let κ be an infinite cardinal. If every infinite cardinal $\lambda \leq \kappa$ carries a uniform ultrafilter, then AC_{λ} holds for every $\lambda \leq \kappa$.

Proof. By induction on $\lambda \leq \kappa$. The base step $\lambda = \omega$ follows from Theorem 4.1. If λ is singular, then AC_{λ} holds by the induction hypothesis as follows: Take a family $\{A_{\alpha} \mid \alpha < \lambda\}$ of non-empty sets. Take also an increasing sequence $\langle \lambda_i \mid i < \operatorname{cf}(\lambda) \rangle$ with limit λ . For each $i < \operatorname{cf}(\lambda)$, by the induction hypothesis, there is a choice function of $\{A_{\alpha} \mid \alpha < \lambda_i\}$. Again, by $\mathsf{AC}_{\operatorname{cf}(\lambda)}$, we can choose a family $\{f_i \mid i < \operatorname{cf}(\lambda)\}$ such that each f_i is a choice function of $\{A_{\alpha} \mid \alpha < \lambda_i\}$. Then the function f on λ defined by $f(\alpha) = f_i(\alpha)$, where i is the least $i < \operatorname{cf}(\lambda)$ with $\alpha < \lambda_i$, is a choice function of $\{A_{\alpha} \mid \alpha < \lambda\}$.

Now suppose λ is regular. Fix a uniform ultrafilter U over λ . Let $\{B_{\alpha} \mid \alpha < \lambda\}$ be a family of non-empty sets.

For $\alpha < \lambda$, let $A_{\alpha} = \prod_{\beta \leq \alpha} B_{\beta}$. By the induction hypothesis, A_{α} is nonempty. By LT, there is a function f on λ such that $X = \{\alpha < \kappa \mid f(\alpha) \in A_{\alpha}\} \in U$. Since U is uniform, we can define the function g on λ as follows: $g(\alpha) = f(\beta)(\alpha)$, where β is the least element of $X \setminus (\alpha + 1)$. It is easy to check that $g(\alpha) \in B_{\alpha}$ for every $\alpha < \lambda$.

It is known that if AC_{κ} holds for every cardinal κ , then the Dependent Choice DC holds (see Jech [7]).

Proposition 5.22. Suppose LT. If every infinite cardinal carries a uniform ultrafilter, then AC_{κ} holds for every infinite cardinal κ , and DC holds.

Question 5.23. Suppose LT. Does DC hold provided that ω carries a non-principal ultrafilter?

6. CHOICELESS MODELS OF LT (II)

By using a model M in Section 5, we will give a proof of Theorem 1.7. For this sake, we need the following lemma and theorem.

Lemma 6.1. The following are equivalent:

- (1) There is an ultrafilter which is not σ -complete.
- (2) ω carries a non-principal ultrafilter.

Proof. $(2) \Rightarrow (1)$ is trivial.

For (1) \Rightarrow (2), fix an ultrafilter U over a set S which is not σ -complete. We can find a map $g: S \to \omega$ such that $g^{-1}(n) \notin U$ for every $n < \omega$. Then $g_*(U)$ is a non-principal ultrafilter over ω .

Here we recall some basic definitions for filters. Let S be a set and F a filter over S. A set $X \subseteq S$ is F-positive if $X \cap Y \neq \emptyset$ for every $Y \in F$. For an F-positive set X, let $F \upharpoonright X = \{Y \subseteq S \mid Y \cup (S \setminus X) \in F\}$. $F \upharpoonright X$ is a filter over S with $X \in F \upharpoonright X$. Let κ be a cardinal. A filter F is κ -saturated if for every indexed family $\{X_{\alpha} \mid \alpha < \kappa\}$ of positive sets, there are $\alpha < \beta < \kappa$ such that $X_{\alpha} \cap X_{\beta}$ is F-positive.

See Kanamori [8] for the proof of the following Tarski's theorem.

Fact 6.2 (Tarski, in ZFC). If F is a $(2^{\omega})^+$ -complete ω_1 -saturated filter, then there is an F-positive set X such that $F \upharpoonright X$ is an ultrafilter.

Theorem 6.3. Suppose V satisfies AC. Then there is a c.c.c. forcing extension V[G] and an intermediate model M between V and V[G] satisfying the following:

- (1) M is a model of ZF + LT.
- (2) If $S \in V$ is a set, κ is an uncountable cardinal, and $U \in V$ is a κ -complete ultrafilter over S in V, then the filter generated by U, $\{X \in \mathcal{P}(S)^M \mid \exists Y \in U(Y \subseteq X)\}$, is a κ -complete ultrafilter in M.
- (3) In M, every ultrafilter is σ -complete.
- (4) $AC_{\omega,2}$ fails in M.

Proof. Our model M is given in Section 5 with the case $\kappa = \omega$. Note that $\operatorname{Fn}(\omega \times \omega, 2, < \omega)$ has the c.c.c.

We already know that the model M satisfies the following:

- (1) There is no non-principal ultrafilter over ω .
- (2) $AC_{\omega,2}$ fails.
- (3) Every ultrafilter has a well-orderable measure one set.

By (1) and Lemma 6.1, we also have:

(4) Every ultrafilter is σ -complete.

Claim 6.4. If $U \in V$ is a κ -complete ultrafilter in V for some uncountable cardinal κ , then U generates a κ -complete ultrafilter in M.

Proof. Since \mathbb{P} is countable, it is known that U generates a κ -complete ultrafilter in V[G] (e.g., see Kanamori [8]). Hence it also generates a κ -complete ultrafilter in M.

Claim 6.5. In M, let κ be an infinite cardinal, and U an ultrafilter over κ . Then $U \cap V \in V$.

Proof. Let $\lambda = (2^{\omega})^V$. First we prove that U is λ^+ -complete. If not, then we can find a cardinal $\mu \leq \lambda$ and $f : \kappa \to \mu$ such that $f^{-1}(\gamma) \notin U$ for every $\gamma < \mu$. Take μ -many reals $\{r_{\xi} \mid \xi < \mu\} \subseteq \mathcal{P}(\omega)^V$. Define $X_n \in U$ as follows. For $n < \omega$, we know $\{\alpha \mid n \in r_{f(\alpha)}\} \in U$ or $\{\alpha \mid n \notin r_{f(\alpha)}\} \in U$. If $\{\alpha \mid n \in r_{f(\alpha)}\} \in U$ then let $X_n = \{\alpha \mid n \in r_{f(\alpha)}\}$, otherwise X_n is $\{\alpha \mid n \notin r_{f(\alpha)}\}$. Since U is σ -complete, we have $X = \bigcap_{n < \omega} X_n \in U$. By the assumption, we can pick $\alpha, \beta \in X$ with $f(\alpha) \neq f(\beta)$. Then for every $n < \omega$, we have $n \in r_{f(\alpha)} \iff n \in r_{f(\beta)}$, hence $r_{f(\alpha)} = r_{f(\beta)}$. This is a contradiction.

To show that $U \cap V \in V$, fix a name \dot{U} for U and $p \in \mathbb{P}$ such that $p \Vdash ``\dot{U}$ is an ultrafilter over κ in M." In V, let $F = \{X \subseteq \kappa \mid p \Vdash \check{X} \in \dot{U}\}$. Since U is a λ^+ -complete ultrafilter and \mathbb{P} satisfies the c.c.c., we have that, in V, F is a $(2^{\omega})^+$ -complete ω_1 -saturated filter over κ . By Fact 6.2 there must be an F-positive set $X \subseteq \kappa$ such that $F \upharpoonright X$ is an ultrafilter. Pick $q \leq p$ with $q \Vdash \check{X} \in \dot{U}$. Then $q \Vdash \check{F} \upharpoonright \check{X} = \dot{U} \cap V$, as required. \Box

The next claim together with Lemma 3.1 yields LT in M.

Claim 6.6. In M, let U be an ultrafilter over S. Then for every indexed family $\{A_s \mid s \in S\}$ of non-empty sets, there is a function g on S such that $\{s \mid g(x) \in A_s\} \in U$.

Proof. Since U has a well-orderable measure one set, we may assume that U is a non-principal σ -complete ultrafilter over an infinite cardinal κ . Let $U' = U \cap V$, which is a σ -complete ultrafilter over κ in V by Claim 6.5. In V, let $j: V \to N \approx {}^{I}V/U'$ be the ultrapower elementary embedding induced by U' where we identify the ultrapower ${}^{I}V/U'$ with its transitive collapse N. Since \mathbb{P} is countable, in V[G], we can extend j to $j: V[G] \to N[G]$ by the canonical way. Moreover we have that $j(F_0) = F_0, j(F_1) = F_1$, and $j \upharpoonright V(\{F_0, F_1\})$ is an elementary embedding from $V(\{F_0, F_1\})$ to $N(\{F_0, F_1\})$. We also note $j(\bigcup_n H(n)) = \bigcup_n H(n)$. Let id be the identity map on κ . We know that $U = \{X \in \mathcal{P}(\kappa)^M \mid [\mathrm{id}] \in j(X)\}$, and for every ordinal γ , there is a function f on κ in V such that $j(f)([\mathrm{id}]) = \gamma$.

Now fix an indexed family $\{A_{\alpha} \mid \alpha < \kappa\} \in M$ of non-empty sets. Let $Y = \bigcup_{\alpha < \kappa} A_{\alpha}$, and take an ordinal γ and a map $g \in M$ such that g is a surjection from $\gamma \times {}^{<\omega}(\bigcup_{n < \omega} H(n))$ onto Y. By the elementarity, j(g) is a surjection from $j(\gamma) \times {}^{<\omega}(\bigcup_{n < \omega} H(n))$ onto j(Y) and $j(\bigcup_{\alpha < \kappa} A_{\alpha}) = j(Y)$. Consider the family $j(\{A_{\alpha} \mid \alpha < \kappa\}) = \{A_{\alpha}^* \mid \alpha < j(\kappa)\}$. We know that $A_{[id]}^*$ is non-empty, so we can take $\delta < j(\gamma)$ and $\langle r_0, \ldots, r_k \rangle \in {}^{<\omega}(\bigcup_n H(n))$ such that $j(g)(\delta, \langle r_0, \ldots, r_k \rangle) \in A_{[id]}^*$. Pick a function $h \in V$ with $j(h)([id]) = \delta$. By the elementarity of j again, we have that $\{\alpha < \kappa \mid g(h(\alpha), \langle r_0, \ldots, r_k \rangle) \in A_{\alpha}\} \in U$. Hence $\alpha \mapsto g(h(\alpha), \langle r_0, \ldots, r_k \rangle)$ is a required choice function in M.

Now we have Theorem 1.7.

Theorem 6.7. If ZFC+ "there is a measurable cardinal" is consistent, then so is ZF + LT + $\neg AC_{\omega,2}$ + "there is a measurable cardinal"+ "every ultrafilter is σ -complete."

Proof. Suppose V satisfies AC and there is a measurable cardinal κ . In the resulting model M in Theorem 6.3, κ remains a measurable cardinal.

If we suppose the existence of a strongly compact cardinal, we can give a partial answer to Question 5.20. Here we recall strongly compact cardinals. An uncountable cardinal κ is a *strongly compact cardinal* if for every cardinal $\lambda \geq \kappa$, there is a κ -complete fine ultrafilter U over $[\lambda]^{<\kappa}$, that is, $\{x \in [\lambda]^{<\kappa} \mid \lambda \leq \kappa\}$

 $\alpha \in x \} \in U$ for every $\alpha < \lambda$. Every strongly compact cardinal is measurable. If κ is strongly compact, then for every regular $\lambda \geq \kappa$, there is a κ -complete uniform ultrafilter over λ ; Fix a κ -complete fine ultrafilter over λ . Define $f : [\lambda]^{<\kappa} \to \lambda$ by $f(x) = \sup(x)$. Then one can check that $f_*(U)$ is a κ complete uniform ultrafilter over λ . In ZFC, the converse of this fact hold: κ is strongly compact if every regular $\lambda \geq \kappa$ carries a κ -complete uniform ultrafilter. We do not know if the converse still holds in ZF.

Theorem 6.8. If ZFC+ "there is a strongly compact cardinal" is consistent, then so is ZF + LT + $\neg AC_{\omega,2}$ + "every ultrafilter is σ -complete"+ "there is a measurable cardinal κ such that every regular cardinal $\lambda \geq \kappa$ carries a uniform ultrafilter."

Proof. Suppose V satisfies AC and there is a strongly compact cardinal κ . In V, for every regular $\lambda \geq \kappa$, there is a κ -complete uniform ultrafilter over λ . Then, in the resulting model M in Theorem 6.3, every regular cardinal $\lambda \geq \kappa$ carries a κ -complete uniform ultrafilter.

7. Consistency strengths of some statements

We proved that LT + WUF(?) does not imply $AC_{\omega,2}$. However our proof needed a large cardinal assumption. In this section, we show that this large cardinal assumption is necessary.

Proposition 7.1. Suppose LT. Let U be a non-principal ultrafilter over a set S. Then there is a cardinal κ such that U is not κ -complete.

Note that, in ZF, it is possible that there is a non-principal ultrafilter which is κ -complete for every cardinal κ . We will return to this topic in the end of this section.

Proof. Suppose to the contrary that U is κ -complete for every cardinal κ . First we shall prove:

Claim 7.2. For every ordinal α and $f : S \to \alpha$, there is $\beta < \alpha$ with $f^{-1}(\{\beta\}) \in U$.

Proof. Suppose not. For $\beta < \alpha$, let $S_{\beta} = \{x \in S \mid f(x) \neq \beta\}$, which is in U. By the assumption, we know $\bigcap_{\beta < \alpha} S_{\beta} \in U$, in particular, there is $x \in \bigcap_{\beta < \alpha} S_{\beta} \in U$. Then $f(x) \neq \beta$ for every $\beta < \alpha$, this is impossible. \Box

Next, fix a large limit ordinal θ such that V_{θ} is a model of a sufficiently large fragment of ZF. Let M be the structure $\langle V_{\theta}; \in \rangle$, and $M^* = \langle M^*; E \rangle$ the ultrapower ${}^{S}M/U$. By LT, M^* is a model of (a large fragment of) ZF. Let $j: V_{\alpha} \to M^*$ be the ultrapower embedding, that is, $j(a) = [c_a]$ for $a \in V_{\theta}$, where c_a is the constant function with value a. By LT, the ultrapower embedding j is an elementary embedding. By the claim, for every $[f] \in M^*$, if [f] is an ordinal of M^* , then $[f] = j(\alpha)$ for some ordinal α . Thus, we have that the set of ordinals of M^* is isomorphic to some set of ordinals. Hence the relation E is well-founded on the ordinals of M^* . Because M^* is a model of (a large fragment of) ZF, every set in M^* has a rank in M^* . Thus we conclude that E is a well-founded relation on M^* , and we can take the transitive collapse N^* of M^* and the collapsing map $\pi: M^* \to N^*$. Let $j^* = \pi \circ j$, which is an elementary embedding from V_{θ} into N^* . We have that $j^*(\alpha) = \alpha$ for every $\alpha < \theta$.

Now we prove that $j^*(a) = a$ for every $a \in V_{\theta}$, however this is impossible; We know $S \in V_{\theta}$ but $[id] \in j^*(S) \setminus S$ since U is non-principal.

Claim 7.3. $j^*(a) = a$ for every $a \in V_{\theta}$.

Proof. By induction on the rank of a. Let $\alpha < \theta$, and suppose $j^*(b) = b$ for every $b \in V_{\alpha}$. Take a set $a \subseteq V_{\alpha}$ with rank α . By the induction hypothesis, we have $a \subseteq j^*(a)$. For the converse that $j^*(a) \subseteq a$, because rank $(a) = j^*(\operatorname{rank}(a)) = \operatorname{rank}(j^*(a))$, we have that $j^*(a) \subseteq V_{\alpha}$. Pick $b \in j^*(a)$. Then rank $(b) < \alpha$, so $j^*(b) = b \in j^*(a)$ and $b \in a$.

Lemma 7.4. Suppose LT holds but there is no non-principal ultrafilter over ω . Then for every set S, S carries a non-principal ultrafilter if and only if there is a measurable cardinal κ and an injection $f : \kappa \to S$.

Proof. If there is an injection from a measurable cardinal κ to S, then it is clear that S carries a non-principal ultrafilter. For the converse, suppose S carries a non-principal ultrafilter U. By LT and Proposition 7.1, there is a cardinal κ and a map $g: S \to \kappa$ such that U is κ -complete but not κ^+ -complete, and $g^{-1}(\alpha) \notin U$ for every $\alpha < \kappa$. By Lemma 6.1 and the assumption that no non-principal ultrafilter over ω , κ must be uncountable. $g_*(U)$ forms a non-principal κ -complete ultrafilter, hence κ is measurable. Now we may assume that $g^{-1}(\alpha + 1) \setminus g^{-1}(\alpha) \neq \emptyset$ for every $\alpha < \kappa$. Let $S_{\alpha} = g^{-1}(\alpha + 1) \setminus g^{-1}(\alpha)$. By LT, we can find a function h on κ such that $X = \{\alpha < \kappa \mid h(\alpha) \in S_{\alpha}\} \in g_*(U)$. Then $h \upharpoonright X$ is an injection from X into S. By arranging this function, we can take an injection from κ to S.

Proposition 7.5. Suppose LT holds but there is no non-principal ultrafilter over ω . Then the following are equivalent:

- (1) There is a measurable cardinal.
- (2) There is a non-principal σ -complete ultrafilter.
- (3) There is a non-principal ultrafilter.
- (4) There is a non-principal ultrafilter, and every ultrafilter is σ -complete.

Proof. $(1) \Rightarrow (2) \Rightarrow (3)$ is trivial, and $(3) \Rightarrow (1)$ is Lemma 7.4. (4) \Rightarrow (3) is obvious, and (3) \Rightarrow (4) follows form Lemma 6.1.

By this proposition, we have Theorem 1.8.

Theorem 7.6. The following theories are equiconsistent:

(1) $ZF + LT + WUF(?) + \neg AC_{\omega,2}$.

- (2) $ZF + LT + WUF(?) + \neg AC_{\omega}$.
- (3) ZFC+ "there is a measurable cardinal".

Proof. $(1) \Rightarrow (2)$ is trivial, and $(3) \Rightarrow (1)$ is Theorem 6.7.

For $(2) \Rightarrow (3)$, suppose V is a model of $\mathsf{ZF} + \mathsf{LT} + \mathsf{WUF}(?) + \neg \mathsf{AC}_{\omega}$. Then by Theorem 4.1, there is no non-principal ultrafilter over ω . Hence there is a measurable cardinal by Proposition 7.5, and then we can take an inner model of ZFC +"there is a measurable cardinal". We also have:

Corollary 7.7. The following theories are equiconsistent:

- (1) ZFC+ "there is a measurable cardinal".
- (2) ZF + LT + "there is a non-principal σ -complete ultrafilter".
- (3) ZF + LT + WUF(?) + "every ultrafilter is σ -complete".

Finally, unlike Corollary 7.7, we prove that the statement that WUF(?)+ "every ultrafilter is σ -complete" does not have a large cardinal strength.

An infinite set S is said to be *amorphous* if every subset of S is finite or co-finite. If S is amorphous, the set U of all co-finite subsets of S forms a σ -complete non-principal ultrafilter over S; It is easy to check that U is a non-principal ultrafilter. If U is not σ -complete, we can find a partition $\langle S_n | n < \omega \rangle$ of S such that each S_n is non-empty finite subset of S. Then the set $\bigcup_n S_{2n}$ is an infinite co-infinite subset of S, this is a contradiction. In fact, one can check that U is κ -complete for every cardinal κ by the same argument.

Fact 7.8 (E.g., see Jech [7]). If ZFC is consistent, then so is ZF+ "there is an amorphous set".

Theorem 7.9. If ZFC is consistent, then so is ZF + WUF(?) + "every ultrafilter is σ -complete".

Proof. By Lemma 6.1, it is enough to construct a model of ZF in which there is a non-principal ultrafilter but there is no non-principal ultrafilter over ω .

Since the theory ZF +"there is an amorphous set" is equiconsistent with ZFC , we can assume that there is an amorphous set S in V. Let U be the set of all co-finite subsets of S, it is a non-principal σ -complete ultrafilter over S. Let $\mathbb{P} = \operatorname{Fn}(\omega \times \omega, 2, <\omega)$. Note that \mathbb{P} is countable in V. Take a (V, \mathbb{P}) -generic G.

Claim 7.10. In V[G], U generates an ultrafilter over S.

Proof. Take $p \in \mathbb{P}$ and a \mathbb{P} -name \dot{X} such that $p \Vdash \dot{X} \subseteq \check{S}$. For $q \leq p$, let $S_q = \{s \in S \mid q \Vdash \check{s} \in \dot{X}\}$. If $S_q \in U$ for some $q \in \mathbb{P}$, then $q \Vdash \check{S}_q \subseteq \dot{X}$, and q forces that " \dot{X} is in the filter generated by U". If there is no $q \leq p$ with $S_q \in U$, since \mathbb{P} is countable and U is σ -complete, we have $T = \bigcup_{q \leq p} (S \setminus S_q) \in U$ and $p \Vdash \check{T} \cap \dot{X} = \emptyset$, so $p \Vdash \check{S} \setminus \dot{X} \in \dot{U}$.

Now, let HS^G be the symmetric extension of V in Section 5 with the case $\kappa = \omega$. We can carry out the proof of Lemma 5.6 without AC in V, hence HS^G is a model of ZF +"there is no non-principal ultrafilter over ω ". On the other hand, since HS^G is an intermediate model between V and V[G], U generates a non-principal ultrafilter over S in HS^G by Claim 7.10. \Box

References

- A. Blass, A model without ultrafilters. Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.25, no.4 (1977), 329–331.
- [2] H.-D. Donder, Regularity of ultrafilters and the core model, Isr. J. Math. 63, No. 3 (1988), 289–322.
- [3] Y. Hayut, A. Karagila, Spectra of uniformity, Comment. Math. Univ. Carolin. 60 (2019), no.2, 285–298.

- [4] H. Herrlich, Axiom of Choice. Lecture Notes in Mathematics 1876, Springer (2006).
- [5] P. E. Howard, Loś' theorem and the Boolean prime ideal theorem imply the axiom of choice, Proc. Amer. Math. Soc. 49 (1975), 426–428.
- [6] P. E. Howard, J. E. Rubin, Consequences of the axiom of choice, Math. Surveys Monogr., 59, American Mathematical Society, 1998.
- [7] T. Jech, The axiom of choice. Stud. Logic Found. Math., Vol. 75, North-Holland, 1973.
- [8] A. Kanamori, *The higher infinite, Large cardinals in set theory from their beginnings*, Second edition. Springer-Verlag, 2009.
- [9] A. Karagila, Iterating symmetric extensions, J. Symb. Log. 84 (2019), no.1, 123–159.
- [10] J. Loś, Quelques remarques, theoremes et problemes sur les classes definissables d'algebres, Mathematical interpretation of formal systems, 98–113 (1955).
- [11] D. Pincus, R. M. Solovay, Definability of measures and ultrafilters, J. Symbolic Logic 42 (1977), no.2, 179–190.
- [12] M. Spector, Ultrapowers without the axiom of choice, J. Symbolic Logic 53 (1988), no.4, 1208–1219.
- [13] E. Tachtsis, Loś's theorem and the axiom of choice, Math. Log. Quart. 65, No. 3 (2019), 280–292.

(T. Usuba) FACULTY OF SCIENCE AND ENGINEERING, WASEDA UNIVERSITY, OKUBO 3-4-1, SHINJYUKU, TOKYO, 169-8555 JAPAN

Email address: usuba@waseda.jp