

# INEQUALITIES AND ASYMPTOTICS FOR HOOK NUMBERS IN RESTRICTED PARTITIONS

WILLIAM CRAIG, MADELINE LOCUS DAWSEY, AND GUO-NIU HAN

**ABSTRACT.** In this paper, we consider the asymptotic properties of hook numbers of partitions in restricted classes. More specifically, we compare the frequency with which partitions into odd parts and partitions into distinct parts have hook numbers equal to  $h \geq 1$  by deriving an asymptotic formula for the total number of hooks equal to  $h$  that appear among partitions into odd and distinct parts, respectively. We use these asymptotic formulas to prove a recent conjecture of the first author and collaborators that for  $h \geq 2$  and  $n \gg 0$ , partitions into odd parts have, on average, more hooks equal to  $h$  than do partitions into distinct parts. We also use our asymptotics to prove certain probabilistic statements about how hooks distribute in the rows of partitions.

## 1. INTRODUCTION

A *partition*  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_\ell)$  of an integer  $n \geq 0$  is a representation of  $n$  in the form  $n = \lambda_1 + \lambda_2 + \dots + \lambda_\ell$ , where reorderings of this sum are considered identical. We use the standard notations  $\lambda \vdash n$  or  $|\lambda| = n$  to say that  $\lambda$  has *size*  $n$ , and we let  $\ell = \ell(\lambda)$  denote the number of parts, or *length*, of  $\lambda$ . We also let  $p(n)$  be the number of partitions of  $n$ . The theory of partitions, and the closely related theory of  $q$ -series, have a vast literature and intersect regularly with combinatorics, harmonic analysis, number theory, mathematical physics, and many other areas of mathematics. For an overview of the theory of partitions, see Andrews' seminal book [1].

In this paper, we focus on the hook numbers of partitions. These are defined in terms of the *Ferrers diagram* of the partition  $\lambda$ , whereby  $\lambda = (\lambda_1, \dots, \lambda_\ell)$  is represented as a diagram of left-adjusted rows of boxes in which the  $i$ th row contains  $\lambda_i$  boxes. The *hook number*  $h_{i,j}(\lambda)$  of the cell in the  $i$ th row and  $j$ th column of  $\lambda$  is defined as the length of the  $L$ -shape formed by the boxes below and to the right of this box, including the box itself. We represent the multiset of all hook numbers of  $\lambda$  as  $\mathcal{H}(\lambda)$ . We give an example below for the partition  $4 + 3 + 2$ .

6	5	3	1
4	3	1	
2	1		

FIGURE 1. Hook numbers of the partition  $\lambda = (4, 3, 2)$

---

2020 *Mathematics Subject Classification.* 11P82, 05A17, 05A15, 05A19, 11P83.

*Key words and phrases.* hook length, distinct partitions, odd partitions, partition asymptotics, partition inequalities.

Hook numbers of partitions have very important implications for the representation theory of symmetric groups; the irreducible representations of  $S_n$  are in bijection with partitions  $\lambda \vdash n$ , and the dimensions of these representations are a function of the hook numbers of  $\lambda$ . Hook numbers have in recent decades also been the subject of many interesting arithmetic studies. This has arisen in large part due to the Nekrasov–Okounkov hook length formula, which connects hook numbers in a deep way to modular forms and  $q$ -series. This formula [17, 20] says that for any complex number  $z$ , we have

$$\sum_{\lambda \in \mathcal{P}} x^{|\lambda|} \prod_{h \in \mathcal{H}(\lambda)} \left(1 - \frac{z}{h^2}\right) = \prod_{k=1}^{\infty} (1 - x^k)^{z-1}.$$

These connections have led to many interesting studies in the number theory and combinatorics literature regarding the asymptotic, combinatorial, and arithmetic properties of the hook numbers of partitions, especially studies into  $t$ -core partitions and  $t$ -hooks of partitions [3, 8, 9, 13, 15, 17, 22].

The object of this paper is to prove a recent conjecture of the first author and collaborators [4] on hook numbers in *restricted* classes of partitions, where analogues of the Nekrasov–Okounkov formula are not known. To motivate the question, recall Euler’s famous theorem that the number of partitions of  $n$  with only odd parts is exactly equal to the number of partitions of  $n$  all of whose parts are distinct integers. It is natural to wonder whether various partition statistics behave differently on these families of partitions. For instance, it is easy to show that on average, partitions into odd parts have more parts than do partitions into distinct parts. This is expected, as partitions into odd parts might have many repetitions of very small parts. Another result from a paper of Andrews [2] can be interpreted in terms of hook numbers<sup>1</sup>, which we now explain. Letting  $\mathcal{O}(n), \mathcal{D}(n)$  be the collections of partitions of  $n$  into odd parts and distinct parts, respectively, define for any integer  $h \geq 1$  the functions

$$a_h(n) := \sum_{\lambda \in \mathcal{O}(n)} \#\{x \in \mathcal{H}(\lambda) : x = h\}, \quad b_h(n) := \sum_{\lambda \in \mathcal{D}(n)} \#\{x \in \mathcal{H}(\lambda) : x = h\}.$$

Because of Euler’s result that  $\mathcal{O}(n)$  and  $\mathcal{D}(n)$  are in bijection, we can see that for each  $n \geq 0$ , we have

$$np_{\mathcal{O}}(n) = \sum_{h \geq 1} a_h(n) = \sum_{h \geq 1} b_h(n) = np_{\mathcal{D}}(n), \tag{1.1}$$

where we let  $p_{\mathcal{O}}(n) = |\mathcal{O}(n)|$  and  $p_{\mathcal{D}}(n) = |\mathcal{D}(n)|$ . Andrews proves in [2] that for all  $n \geq 0$ , we have  $b_1(n) \geq a_1(n)$ ; that is, partitions into distinct parts have, on average, more hook numbers equal to 1 than do partitions into odd parts. In light of (1.1), it would be natural to suspect that the inequality of Andrews must be balanced out in some way by inequalities between  $a_h(n)$  and  $b_h(n)$  for some  $h > 1$  which are in the other direction. In [4], this concept was formulated as a much more concrete conjecture:

**Conjecture 1.1.** *Let  $h \geq 2$  be fixed. Then the following are true:*

---

<sup>1</sup>Andrews states his results in terms of distinct part sizes, which are easily seen to be equivalent to hook numbers equal to 1.

- (1) *There exists some integer  $N_h > 0$  such that for all  $n > N_h$ , we have  $a_h(n) \geq b_h(n)$ .*
- (2) *There exists some constant  $\gamma_h > 1$  such that  $a_h(n)/b_h(n) \rightarrow \gamma_h$  as  $n \rightarrow \infty$ .*

Note that Conjecture 1.1 (2) is a much stronger statement than Conjecture 1.1 (1). In [4], Conjecture 1.1 is proved only in the cases  $h = 2$  and  $h = 3$ , but no progress is made for any cases  $h \geq 4$ . We also note that the authors of [4] prove that  $\gamma_1 = \frac{1}{2\log(2)}$ ,  $\gamma_2 = \frac{3}{2}$ , and  $\gamma_3 = \frac{2}{3(\log(2)-1/8)}$ . In this paper, we improve on the methods of [4], which enable us to prove Conjecture 1.1 in its entirety.

**Theorem 1.2.** *Conjecture 1.1 (2), and therefore also Conjecture 1.1 (1), is true for all  $h \geq 2$ .*

**Remark 1.3.** In [4], the exact values of  $N_2$  and  $N_3$  are computed. In our case, although Theorem 1.2 establishes the existence of  $N_h$ , our proof is not effective. It would be possible to make the proof effective following the basic outline of [10] if the main results of Section 3 could be made effective. Such an effective proof would not, however, give an optimal value for  $N_h$ . Based on computational data from [4],  $N_h$  appears to grow roughly as  $0.6h^2$ .

In fact, the theorems we prove give much more detail about the behavior of  $a_h(n)$  and  $b_h(n)$ , which we now summarize. In order to prove that  $a_h(n)/b_h(n) \rightarrow \gamma_h$ , we prove separate asymptotic formulas for  $a_h(n)$  and  $b_h(n)$ , which we state below.

**Theorem 1.4.** *Let  $h \geq 1$  be an integer. Then there exists a constant  $\alpha_h \in \mathbb{Q}$  such that*

$$a_h(n) \sim \alpha_h \frac{3^{1/4}}{2\pi n^{1/4}} e^{\pi\sqrt{n/3}}$$

as  $n \rightarrow \infty$ .

**Theorem 1.5.** *Let  $h \geq 1$  be an integer. Then there exists a constant  $\beta_h \in \mathbb{Q}(\log(2))$  such that*

$$b_h(n) \sim \beta_h \frac{3^{1/4}}{2\pi n^{1/4}} e^{\pi\sqrt{n/3}}$$

as  $n \rightarrow \infty$ . Furthermore,  $\beta_h \in \mathbb{Q}$  if and only if  $h$  is even.

A key ingredient to these asymptotic formulas is the construction of the generating functions for  $a_h(n)$  and  $b_h(n)$ , which previously had not been known apart from the cases  $h = 2$  and  $h = 3$ . We make use of two representations of these generating functions, found in Theorems 2.1 and 2.2. In particular, we show in Theorem 2.2 that these generating functions are essentially rational functions of  $q$  multiplied by the generating function for partitions into odd parts, which is a modular form.

From Theorems 1.4 and 1.5, it is immediately clear that  $a_h(n)/b_h(n) \rightarrow \alpha_h/\beta_h$  as  $h \rightarrow \infty$ , so this establishes the existence of  $\gamma_h$ . By careful evaluations of the constants  $\alpha_h$  and  $\beta_h$ , we are able to prove the following result for  $\gamma_h$ .

**Theorem 1.6.** *For each  $h \geq 2$ , we have  $\gamma_h > 1$ . Furthermore, we have*

$$\gamma_h \rightarrow \frac{\log(4)}{\log(3)} \approx 1.2618\dots$$

as  $h \rightarrow \infty$ .

It is now clear that to prove Theorem 1.2, it will suffice to prove Theorems 1.4, 1.5, and 1.6. Therefore, the rest of the paper is dedicated to proving these central results. In Section 2, we construct the generating functions for the sequences  $a_h(n)$  and  $b_h(n)$ . In Section 3, we prove Theorems 1.4 and 1.5 using the circle method. In Section 4 we then closely study the constants  $\alpha_h$  and  $\beta_h$  in order to prove Theorem 1.6. We close with further conjectures and interesting probabilistic corollaries of our results in Section 5; for example, we will show that for  $n \gg 0$ , most rows in partitions into distinct parts contain a hook of any given length  $h \geq 1$ .

## ACKNOWLEDGEMENTS

The first author has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 101001179).

## 2. GENERATING FUNCTIONS

Recall the usual notation of the  $q$ -ascending factorial

$$\begin{aligned} (x; q)_0 &:= 1; \\ (x; q)_n &:= (1-x)(1-xq) \cdots (1-xq^{n-1}); \\ (x; q)_\infty &:= \lim_{n \rightarrow \infty} (x; q)_n. \end{aligned}$$

For  $0 \leq k \leq n$ , let

$$\binom{n}{k}_q := \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}}$$

be the usual  $q$ -binomial coefficient. In this section, we establish the following explicit generating functions for  $a_h(n)$  and  $b_h(n)$  by using a method described in [5]. Let

$$\tilde{a}_h(q) = \sum_{n=0}^{\infty} a_h(n) q^n \quad \text{and} \quad \tilde{b}_h(q) = \sum_{n=0}^{\infty} b_h(n) q^n.$$

**Theorem 2.1.** *For each  $h \geq 1$ , we have*

$$\begin{aligned} \tilde{a}_h(q) &= (-q; q)_\infty \sum_{j=0}^{\lceil h/2 \rceil - 1} q^h \binom{h-j-1}{j}_q \sum_{q^2, m \geq 0} (q^{2m+1}; q^2)_j q^{(2h-4j)m} \\ &\quad + (-q; q)_\infty \sum_{j=0}^{\lfloor h/2 \rfloor - 1} q^{3h-4j-3} \binom{h-j-2}{j}_q \sum_{q^2, m \geq 0} (q^{2m+3}; q^2)_j q^{(2h-4j-2)m}; \\ \tilde{b}_h(q) &= (-q; q)_\infty \sum_{j=0}^{\lceil h/2 \rceil - 1} q^{h+j(j-1)/2} \binom{h-j-1}{j}_q \sum_{q, m \geq 0} \frac{q^{(j+1)m}}{(-q^{m+1}; q)_{h-j}}. \end{aligned}$$

*Proof.* Each partition  $\lambda$  can be represented by its Ferrers diagram (or Young diagram). For each box  $v$  in the Ferrers diagram of a partition  $\lambda$ , or for each box  $v$  in  $\lambda$ , for short, define the *arm length* (resp. *leg length*, *coarm length*, *coleg length*) of  $v$ , denoted by  $\text{arm}(\lambda, v)$  (resp.  $\text{leg}(\lambda, v)$ ,  $\text{coarm}(\lambda, v)$ ,  $\text{coleg}(\lambda, v)$ ), to be the number of boxes  $u$  such that  $u$  lies in the same row as  $v$  and to the right of  $v$  (resp. in the same column

as  $v$  and below  $v$ , in the same row as  $v$  and to the left of  $v$ , in the same column as  $v$  and above  $v$ ). See Figure 2.




FIGURE 2. Arm, leg, coarm, and coleg lengths:  $\text{arm}(\lambda, v) = j$ ,  $\text{leg}(\lambda, v) = l$ ,  $\text{coarm}(\lambda, v) = m$ ,  $\text{coleg}(\lambda, v) = g$

Consider a set  $\mathcal{L}$  of partitions. For each given triplet  $(j, l, m)$  of integers, let  $f_{\mathcal{L}}(j, l, m; n)$  denote the number of ordered pairs  $(\lambda, v)$  such that  $\lambda \in \mathcal{L}$ ,  $v \in \lambda$ ,  $\lambda \vdash n$ ,  $\text{arm}(\lambda, v) = j$ ,  $\text{leg}(\lambda, v) = l$ , and  $\text{coarm}(\lambda, v) = m$ . For a fixed partition  $\lambda$ , it is easy to see that for each box  $v \in \lambda$  and  $\lambda \vdash n$ , the triplets  $(\text{arm}(\lambda, v), \text{leg}(\lambda, v), \text{coarm}(\lambda, v))$  are different. Now, let the triplet  $(j, l, m)$  be fixed. The generating function for those partitions is equal to the product of several “small” generating functions for the different regions of the partitions, as shown in Figure 3.




FIGURE 3. A partition and its different regions

(i) Case  $\tilde{b}_h(q)$  for the distinct partitions  $\mathcal{D}$ . Let  $F_{\mathcal{D}}(j, l, m; q) := \sum_n f_{\mathcal{D}}(j, l, m; n)q^n$  be the generating function for partitions into distinct parts with arm length  $j$ , leg length  $l$ , and coarm length  $m$ , as displayed in the above diagram. It is quite routine (see, e.g., [1, Chap. 3]) to prove that

$$F_{\mathcal{D}}(j, l, m; q) = F_{\mathcal{D}}(A, q)F_{\mathcal{D}}(B, q)F_{\mathcal{D}}(C, q)F_{\mathcal{D}}(D, q),$$

where the generating functions for the regions  $A, B, C, D$  are respectively

$$\begin{aligned} F_{\mathcal{D}}(A, q) &= (-q; q)_m; \\ F_{\mathcal{D}}(B, q) &= \binom{j}{l}_q q^{l(l-1)/2}; \\ F_{\mathcal{D}}(C, q) &= (1 + q^{m+j+2})(1 + q^{m+j+3}) \dots = \frac{(-q; q)_{\infty}}{(-q; q)_{m+j+1}}; \\ F_{\mathcal{D}}(D, q) &= q^{(m+1)(l+1)+j}. \end{aligned}$$

Hence,

$$\begin{aligned} F_{\mathcal{D}}(j, l, m; q) &= F_{\mathcal{D}}(A, q)F_{\mathcal{D}}(B, q)F_{\mathcal{D}}(C, q)F_{\mathcal{D}}(D, q) \\ &= (-q; q)_m \binom{j}{l}_q q^{l(l-1)/2} \frac{(-q; q)_{\infty}}{(-q; q)_{m+j+1}} q^{(m+1)(l+1)+j} \\ &= q^{(m+1)(l+1)+j+l(l-1)/2} \binom{j}{l}_q \frac{(-q; q)_{\infty}}{(-q^{m+1}; q)_{j+1}}. \end{aligned}$$

We see that  $F_{\mathcal{D}}(j, l, m; q) = 0$  if  $j < l$ , since then  $\binom{j}{l}_q = 0$ . Since  $h = j + l + 1$ , the condition  $j \geq l$  implies that  $h - l - 1 \geq l$ , and thus we have  $l \leq (h - 1)/2$ . We therefore have

$$\begin{aligned} \tilde{b}_h(q) &= \sum_{l=0}^{\lceil h/2 \rceil - 1} \sum_{m \geq 0} F_{\mathcal{D}}(h - l - 1, l, m; q) \\ &= \sum_{l=0}^{\lceil h/2 \rceil - 1} \sum_{m \geq 0} q^{(m+1)(l+1)+(h-l-1)+l(l-1)/2} \binom{h-l-1}{l}_q \frac{(-q; q)_{\infty}}{(-q^{m+1}; q)_{h-l-1+1}}, \end{aligned}$$

which is equal to the expression given in the theorem.

(ii) *Case  $\tilde{a}_h(q)$  for the odd partitions  $\mathcal{O}$ .* Let  $F_{\mathcal{O}}(j, l, m; q) := \sum_n f_{\mathcal{O}}(j, l, m; n)q^n$  be the generating function for partitions into odd parts with arm length  $j$ , leg length  $l$ , and coarm length  $m$ . If  $m = 2m'$  is even, then  $j = 2j'$  is also even. With  $j + l + 1 = h$ , we have  $2j' + l + 1 = h$  and  $j' \leq (h - 1)/2$ . The generating functions for the regions  $A, B, C, D$  are respectively

$$\begin{aligned} F_{\mathcal{O}}^e(A, q) &= \frac{1}{(1 - q)(1 - q^3)(1 - q^5) \dots (1 - q^{2m'-1})} = \frac{1}{(q; q^2)_{m'}}, \\ F_{\mathcal{O}}^e(B, q) &= \binom{j' + l}{j'}_{q^2}, \\ F_{\mathcal{O}}^e(C, q) &= \frac{1}{(1 - q^{2m'+2j'+1})(1 - q^{2m'+2j'+3}) \dots} = \frac{1}{(q^{2m'+2j'+1}; q^2)_{\infty}}, \\ F_{\mathcal{O}}^e(D, q) &= q^{(2m'+1)(l+1)+2j'}, \end{aligned}$$

where to obtain  $F_{\mathcal{O}}^e(B, q)$ , we made the following calculation:

$$\sum_{l \geq 0} F_{\mathcal{O}}^e(B, q)t^l = \frac{1}{(t; q^2)_{j'+1}} = \sum_{l \geq 0} \binom{j' + l}{l}_{q^2} t^l.$$

Hence, with  $2j' + l + 1 = h$ , we have

$$\begin{aligned} F_{\mathcal{O}}^e(2j', l, 2m'; q) &= F_{\mathcal{O}}^e(A, q)F_{\mathcal{O}}^e(B, q)F_{\mathcal{O}}^e(C, q)F_{\mathcal{O}}^e(D, q) \\ &= \frac{1}{(q; q^2)_{m'}} \binom{j' + l}{j'}_{q^2} \frac{1}{(q^{2m'+2j'+1}; q^2)_{\infty}} q^{(2m'+1)(l+1)+2j'} \\ &= \frac{1}{(q; q^2)_{\infty}} \binom{h - j' - 1}{j'}_{q^2} (q^{2m'+1}; q^2)_{j'} q^{h+(2h-4j)m} \end{aligned}$$

If  $m = 2m' + 1$  is odd, then  $j = 2j' + 1$  is also odd. Since  $j + l + 1 = h$ , we have  $2j' + 1 + l + 1 = h$  and  $j' \leq (h - 2)/2$ . The generating functions for the regions  $A, B, C, D$  are respectively

$$\begin{aligned} F_{\mathcal{O}}^o(A, q) &= \frac{1}{(1 - q)(1 - q^3)(1 - q^5) \cdots 1 - q^{2m'+1}} = \frac{1}{(q; q^2)_{m'+1}}, \\ F_{\mathcal{O}}^o(B, q) &= \binom{j' + l}{j'}_{q^2} q^l, \\ F_{\mathcal{O}}^o(C, q) &= \frac{1}{(1 - q^{2m'+2j'+3})(1 - q^{2m'+2j'+5}) \cdots} = \frac{1}{(q^{2m'+2j'+3}; q^2)_{\infty}}, \\ F_{\mathcal{O}}^o(D, q) &= q^{(2m'+2)(l+1)+2j'+1}, \end{aligned}$$

where where to obtain  $F_{\mathcal{O}}^o(B, q)$ , we made the following calculation:

$$\sum_{l \geq 0} F_{\mathcal{O}}^o(B, q) t^l = \frac{1}{(tq; q^2)_{j'+1}} = \sum_{l \geq 0} \binom{j' + l}{l}_{q^2} (tq)^l.$$

Hence, for  $2j' + 1 + l + 1 = h$ , we have

$$\begin{aligned} F_{\mathcal{O}}^o(2j' + 1, l, 2m' + 1; q) &= F_{\mathcal{O}}^o(A, q)F_{\mathcal{O}}^o(B, q)F_{\mathcal{O}}^o(C, q)F_{\mathcal{O}}^o(D, q) \\ &= \frac{1}{(q; q^2)_{m'+1}} \binom{j' + l}{j'}_{q^2} q^l \frac{1}{(q^{2m'+2j'+3}; q^2)_{\infty}} q^{(2m'+2)(l+1)+2j'+1} \\ &= \frac{1}{(q; q^2)_{\infty}} \binom{h - j' - 2}{j'}_{q^2} (q^{2m'+3}; q^2)_{j'} q^{3h-4j-3+(2h-4j-2)m}. \end{aligned}$$

Using Euler's well-known identity  $(-q; q)_{\infty}(q; q^2)_{\infty} = 1$ , combining the above two cases yields

$$\begin{aligned} \tilde{a}_h(q) &= \sum_{j'=0}^{\lfloor h/2 \rfloor - 1} \sum_{m' \geq 0} F_{\mathcal{O}}^e(2j', h - 2j' - 1, 2m'; q) \\ &\quad + \sum_{j'=0}^{\lfloor h/2 \rfloor - 1} \sum_{m' \geq 0} F_{\mathcal{O}}^o(2j' + 1, h - 2j' - 2, 2m' + 1; q), \end{aligned}$$

which is exactly the expression for  $\tilde{a}_h(q)$  given in the theorem.  $\square$

From Theorem 2.1, we can derive the following result, which shows the relatively elementary nature of the generating functions for  $a_h(n)$  and  $b_h(n)$ .

**Theorem 2.2.** *We have*

$$\begin{aligned}\tilde{a}_h(q) &= (-q; q)_\infty \tilde{A}_h(q), \\ \tilde{b}_{2h}(q) &= (-q; q)_\infty \tilde{B}_{2h}(q), \\ \tilde{b}_{2h+1}(q) &= (-q; q)_\infty \left( \tilde{B}_{2h+1}(q) + \sum_{k=1}^{\infty} \frac{q^k}{1+q^k} \right),\end{aligned}$$

where  $\tilde{A}_h(q)$  and  $\tilde{B}_h(q)$  are rational functions in  $q$ .

Here are the first few values of  $\tilde{A}_h(q)$  and  $\tilde{B}_h(q)$ :

$$\begin{aligned}\tilde{A}_1(q) &= \frac{q}{1-q^2}, & \tilde{A}_2(q) &= \frac{q^5 + q^3 + q^2}{1-q^4}, \\ \tilde{A}_3(q) &= \frac{q^{10} + q^9 + 2q^7 + 3q^5 - q^4 + 2q^3}{(1-q^6)(1+q^2)}, \\ \tilde{B}_1(q) &= 0, & \tilde{B}_2(q) &= \frac{q^2}{1-q^2}, & \tilde{B}_3(q) &= \frac{q^5 - q^2 - q}{(1-q^4)(1+q)}.\end{aligned}$$

*Proof.* The basic idea of the proof is to simplify the expressions given in Theorem 2.1 by removing  $\sum_{m \geq 0}$  via the geometric summation formula. Recall the following useful identities, which are consequences of well-known identities for  $q$ -binomial coefficients (for example, see [1, Ch. 3]) or are easy to prove:

$$(q^{2m+1}; q^2)_n = \sum_{k=0}^n \binom{n}{k}_{q^2} (-1)^k q^{2mk+k^2}, \quad (2.1)$$

$$\frac{1}{(-q^{m+1}; q)_n} = \sum_{k=0}^{\infty} \binom{n+k-1}{k}_q (-1)^k q^{(m+1)k}, \quad (2.2)$$

$$\sum_{k=1}^{\infty} \frac{(-q)^k}{1-q^k} = - \sum_{k=1}^{\infty} \frac{q^k}{1+q^k}. \quad (2.3)$$

From Theorem 2.1, we have

$$\frac{\tilde{a}_h(q)}{(-q; q)_\infty} = \sum_{j=0}^{\lceil h/2 \rceil - 1} \binom{h-j-1}{j}_{q^2} A_1(h, j; q) + \sum_{j=0}^{\lfloor h/2 \rfloor - 1} \binom{h-j-2}{j}_q A_2(h, j; q), \quad (2.4)$$

where, with the help of (2.1) we have

$$\begin{aligned}A_1(h, j; q) &:= q^h \sum_{m \geq 0} (q^{2m+1}; q^2)_j q^{(2h-4j)m} \\ &= q^h \sum_{m \geq 0} \sum_{k=0}^j \binom{j}{k}_{q^2} (-1)^k q^{2mk+k^2} q^{(2h-4j)m} \\ &= q^h \sum_{k=0}^j \binom{j}{k}_{q^2} (-1)^k q^{k^2} \sum_{m \geq 0} (q^{2h+2k-4j})^m\end{aligned}$$



$$= q^h \sum_{k=0}^j \binom{j}{k}_{q^2} \frac{(-1)^k q^{k^2}}{1 - q^{2h+2k-4j}},$$

and

$$\begin{aligned} A_2(h, j; q) &:= q^{3h-4j-3} \sum_{m \geq 0} (q^{2m+3}; q^2)_j q^{(2h-4j-2)m} \\ &= q^{3h-4j-3} \sum_{m \geq 0} \sum_{k=0}^j \binom{j}{k}_{q^2} (-1)^k q^{2(m+1)k+k^2} q^{(2h-4j-2)m} \\ &= q^{3h-4j-3} \sum_{k=0}^j \binom{j}{k}_{q^2} (-1)^k q^{k^2+2k} \sum_{m \geq 0} (q^{2h-4j+2k-2})^m \\ &= q^{3h-4j-3} \sum_{k=0}^j \binom{j}{k}_{q^2} \frac{(-1)^k q^{k^2+2k}}{1 - q^{2h-4j+2k-2}}. \end{aligned}$$

Thus, it is clear that  $\tilde{A}_h(q)$  is a rational function in  $q$ .

We now consider  $\tilde{b}_h(q)$ . From Theorem 2.1, we write

$$\tilde{b}_h(q) = (-q; q)_\infty \sum_{j=0}^{\lceil h/2 \rceil - 1} q^{h+j(j-1)/2} \binom{h-j-1}{j}_q Y(h, j; q), \quad (2.5)$$

where, with the help of (2.2),

$$\begin{aligned} Y(h, j; q) &:= \sum_{m \geq 0} \frac{q^{m(j+1)}}{(-q^{m+1}; q)_{h-j}} \\ &= \sum_{m \geq 0} q^{m(j+1)} \sum_{k=0}^{\infty} \binom{h-j+k-1}{k}_q (-1)^k q^{(m+1)k} \\ &= \sum_{k=0}^{\infty} (-q)^k \binom{h-j+k-1}{k}_q \sum_{m \geq 0} q^{m(j+k+1)} \\ &= \sum_{k=0}^{\infty} \binom{h-j+k-1}{k}_q \frac{(-q)^k}{1 - q^{j+k+1}}. \end{aligned}$$

Since

$$\binom{h-j+k-1}{k}_q = \frac{(1 - q^{k+1})(1 - q^{k+2}) \cdots (1 - q^{k+h-j-1})}{(q; q)_{h-j-1}},$$

there are two cases to be considered.

Case 1: Suppose  $k + h - j - 1 \geq j + k + 1$ ; i.e.,  $j \leq h/2 - 1$ . Then the denominator term  $1 - q^{j+k+1}$  is cancelled with a factor in the numerator. Hence,

$$\binom{h-j+k-1}{k}_q \frac{(-q)^k}{1 - q^{j+k+1}} = \frac{1}{(q; q)_{h-j-1}} \sum_{s=1}^{h-j-1} C_s(h, j; q) q^{sk},$$

where  $C_s(h, j; q)$  are polynomials in  $q$ . Notice that  $C_s(h, j; q)$  are independent of  $k$ . Thus,

$$Y(h, j; q) = \sum_{k=0}^{\infty} \frac{1}{(q; q)_{h-j-1}} \sum_{s=0}^{h-j-1} C_s(h, j; q) q^{sk} = \frac{1}{(q; q)_{h-j-1}} \sum_{s=0}^{h-j-1} \frac{C_s(h, j; q)}{1 - q^s}.$$

Hence,  $Y(h, j; q)$  is a rational function in  $q$ , so that  $\tilde{b}_h(q)/(-q; q)_{\infty}$  is also a rational function in  $q$ .

Case 2: Suppose on the other hand that  $k + h - j - 1 < j + k + 1$ ; i.e.,  $j > h/2 - 1$ . Since  $j \leq \lceil h/2 \rceil - 1$ , the only possible choice is that  $h = 2j + 1$  is odd. We then have

$$\begin{aligned} Y(2j + 1, j; q) &= \sum_{k=0}^{\infty} \binom{2j + 1 - j + k - 1}{k}_q \frac{(-q)^k}{1 - q^{j+k+1}} \\ &= \sum_{k=0}^{\infty} \frac{(1 - q^{k+1})(1 - q^{k+2}) \cdots (1 - q^{k+j})}{(1 - q)(1 - q^2) \cdots (1 - q^j)} \cdot \frac{(-q)^k}{1 - q^{j+k+1}} \\ &= \frac{1}{(q; q)_j} \sum_{k=0}^{\infty} (-1)^k (1 - q^{k+1})(1 - q^{k+2}) \cdots (1 - q^{k+j}) \frac{q^k}{1 - q^{j+k+1}} \\ &= \frac{1}{(q; q)_j} \sum_{k=0}^{\infty} Z_k(j, q^k; q), \end{aligned}$$

with

$$Z_k(j, x; q) := (-1)^k \frac{x(1 - xq)(1 - xq^2) \cdots (1 - xq^j)}{1 - xq^{j+1}}.$$

We write

$$Z_k(j, x; q) = (-1)^k \left( F(x, q) + \frac{G(q)}{1 - xq^{j+1}} \right),$$

where  $F(x, q), G(q)$ , viewed as polynomials of  $x$ , are determined by

$$F(x, q) + \frac{G(q)}{1 - xq^{j+1}} = \frac{x(1 - xq)(1 - xq^2) \cdots (1 - xq^j)}{1 - xq^{j+1}}.$$

Taking  $x \rightarrow 0$ , we see that  $G(q) = -F(0, q)$  and

$$G(q) = x(1 - xq)(1 - xq^2) \cdots (1 - xq^j) \Big|_{x=q^{-j-1}} = \frac{(-1)^j}{q^{(j+1)(j+2)/2}} (q; q)_j.$$

On the other hand, with the help of (2.3) we have

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{(-1)^k q^{k+j+1}}{1 - q^{k+j+1}} &= (-1)^{j+1} \sum_{k=0}^{\infty} \frac{(-q)^{k+j+1}}{1 - q^{k+j+1}} \\ &= (-1)^{j+1} \sum_{k=1}^{\infty} \frac{(-q)^k}{1 - q^k} + (-1)^j \sum_{k=1}^j \frac{(-q)^k}{1 - q^k} \\ &= (-1)^j \sum_{k=1}^{\infty} \frac{q^k}{1 + q^k} + (-1)^j \sum_{k=1}^j \frac{(-q)^k}{1 - q^k}. \end{aligned}$$

Therefore, we obtain

$$\begin{aligned}
Y(2j+1, j; q) &= \frac{1}{(q; q)_j} \sum_{k=0}^{\infty} Z_k(j, q^k; q) \\
&= \frac{1}{(q; q)_j} \sum_{k=0}^{\infty} (-1)^k \left( F(q^k, q) + G(q) + \frac{G(q)q^{k+j+1}}{1-q^{k+j+1}} \right) \\
&= \frac{1}{(q; q)_j} \sum_{k=0}^{\infty} (-1)^k (F(q^k, q) - F(0, q)) \\
&\quad + \frac{G(q)}{(q; q)_j} \sum_{k=0}^{\infty} (-1)^k \left( \frac{q^{k+j+1}}{1-q^{k+j+1}} \right) \\
&= \text{Rational}(q) + \frac{(-1)^j}{q^{(j+1)(j+2)/2}} \sum_{k=0}^{\infty} (-1)^k \left( \frac{q^{k+j+1}}{1-q^{k+j+1}} \right) \\
&= \text{Rational}(q) + \frac{(-1)^j}{q^{(j+1)(j+2)/2}} \sum_{k=0}^{\infty} (-1)^k \left( \frac{q^{k+j+1}}{1-q^{k+j+1}} \right) \\
&= \text{Rational}(q) + \frac{1}{q^{(j+1)(j+2)/2}} \sum_{k=1}^{\infty} \frac{q^k}{1+q^k}.
\end{aligned}$$

Finally, when  $h = 2m + 1$  is odd, we therefore have

$$\begin{aligned}
\frac{\tilde{b}_h(q)}{(-q; q)_{\infty}} &= \sum_{j=0}^{m-1} q^{h+j(j-1)/2} \binom{h-j-1}{j}_q Y(h, j; q) + q^{2m+1+m(m-1)/2} Y(2m+1, m; q) \\
&= \text{Rational}(q) + \sum_{k=1}^{\infty} \frac{q^k}{1+q^k}.
\end{aligned}$$

This completes the proof.  $\square$

### 3. ASYMPTOTICS FOR $a_h(n)$ AND $b_h(n)$

In this section, we use Wright's circle method to compute the first terms of the asymptotic expansions of the  $q$ -series  $a_h(n)$  and  $b_h(n)$  for integers  $h \geq 1$ , and we use this to prove Theorems 1.4 and 1.5. In order to accomplish this, we will use Euler–Maclaurin summation to compute asymptotics for  $\tilde{a}_h(q)$  and  $\tilde{b}_h(q)$  with  $q = e^{-z}$ , as  $z \rightarrow 0$  in any conical region (i.e. if  $z = x + iy$ , we may restrict  $z$  to the region  $0 \leq x < \delta y$  for any  $\delta > 0$ ).

Although we showed in Section 2 that the generating functions of  $\tilde{a}_h(q)$  and  $\tilde{b}_h(q)$  are rational (or nearly rational) functions multiplied by the modular form  $(-q; q)_{\infty}$ , we use different forms of the generating functions here that are more convenient for Euler–Maclaurin summation. For  $\tilde{a}_h(q)$ , we shall use the representation from (2.4):

$$\tilde{a}_h(q) = (-q; q) \left[ \sum_{j=0}^{\lceil h/2 \rceil - 1} \binom{h-j-1}{j}_q A_1(h, j; q) + \sum_{j=0}^{\lfloor h/2 \rfloor - 1} \binom{h-j-2}{j}_q A_2(h, j; q) \right]$$

where we recall

$$A_1(h, j; q) := q^h \sum_{m \geq 0} (q^{2m+1}; q^2)_j q^{(2h-4j)m},$$

$$A_2(h, j; q) := q^{3h-4j-3} \sum_{m \geq 0} (q^{2m+3}; q^2)_j q^{(2h-4j-2)m}.$$

In both cases, the asymptotic analysis will be carried out using the more general functions

$$F_{j,k,l}(q) := \sum_{m \geq 0} (q^{2m+k}; q^2)_j q^{lm}.$$

Observe that  $A_1(h, j; q) = q^h F_{j,1,2h-4j}(q)$  and  $A_2(h, j; q) = q^{3h-4j-2} F_{j,3,2h-4j-2}(q)$ , and therefore if  $q = e^{-z}$ , then as  $z \rightarrow 0$  we have  $A_1(h, j; q) \sim F_{j,1,2h-4j}(e^{-z})$  and  $A_2(h, j; q) \sim F_{j,3,2h-4j-2}(e^{-z})$ . Thus if  $q = e^{-z}$ , then as  $z \rightarrow 0$  we have the asymptotic formula

$$\begin{aligned} \tilde{a}_h(q) \sim (-q; q)_\infty & \left[ \sum_{j=0}^{\lceil h/2 \rceil - 1} \binom{h-j-1}{j} F_{j,1,2h-4j}(e^{-z}) \right. \\ & \left. + \sum_{j=0}^{\lfloor h/2 \rfloor - 1} \binom{h-j-2}{j} F_{j,3,2h-4j-2}(e^{-z}) \right]. \end{aligned} \quad (3.1)$$

For the asymptotic analysis of  $\tilde{b}_h(q)$ , we will use the representation from (2.5):

$$\tilde{b}_h(q) = (-q; q)_\infty B_h(q), \quad B_h(q) := \sum_{j=0}^{\lceil h/2 \rceil - 1} q^{h + \frac{j(j-1)}{2}} \binom{h-j-1}{j} \sum_{m \geq 0} \frac{q^{(j+1)m}}{(-q^{m+1}; q)_{h-j}}.$$

In order to understand the asymptotic behavior of  $B_h(e^{-z})$  as  $z \rightarrow 0$ , we study the more general functions

$$G_{j,k}(q) := \sum_{m \geq 0} \frac{q^{jm}}{(-q^{m+1}; q)_k}.$$

Observe that if  $q = e^{-z}$ , then as  $z \rightarrow 0$  we have the asymptotic formula

$$\tilde{b}_h(q) \sim (-q; q)_\infty \sum_{j=0}^{\lceil h/2 \rceil - 1} \binom{h-j-1}{j} G_{j+1, h-j}(e^{-z}). \quad (3.2)$$

We now estimate the functions  $F_{j,k,l}(q)$  and  $G_{j,k}(q)$ , and we apply these estimates to the asymptotic formulas in (3.1) and (3.2).

**3.1. Euler–Maclaurin Summation.** Euler–Maclaurin summation gives a method for estimating, and even obtaining exact formulas for, sums involving terms of the form  $f(mz)$ . There are many formulations, some asymptotic and some exact. We will use the following formulation, which comes from [7]. We actually use Euler–Maclaurin summation in a slightly unusual manner to analyze sums whose terms do not quite fit the form  $f(mz)$ , inspired by the methodology applied in [6].

Suppose that  $D_\theta := \{re^{i\alpha} : r \geq 0, |\alpha| \leq \theta\}$  for some  $0 \leq \theta < \frac{\pi}{2}$ . Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be holomorphic in the domain  $D_\theta$ , and assume that  $f$  and all of its derivatives

decay at infinity at least as fast as  $|z|^{-1-\varepsilon}$  for some  $\varepsilon > 0$ . (This property is often called *sufficient decay*.) Finally, let  $B_n(x)$  denote the Bernoulli polynomials, and let  $\tilde{B}_n(x) := B_n(\{x\})$  be the periodic Bernoulli function. Then from [7, Equation (5.7)] in the special case  $a = 1$ , we have the following proposition.

**Proposition 3.1.** *For each  $N \geq 1$ , as  $z \rightarrow 0$  in the region  $D_\theta$ , we have*

$$\begin{aligned} \sum_{m \geq 0} f((m+1)z) &= \frac{1}{z} \int_0^\infty f(x) dx - \sum_{k \geq 0} \frac{f^{(k)}(0) z^k}{(k+1)!} - \sum_{n=0}^{N-1} \frac{B_{n+1}(0) f^{(n)}(z)}{(n+1)!} z^n \\ &\quad - \frac{(-1)^N z^{N-1}}{N!} \int_z^{z\infty} f^{(N)}(w) \tilde{B}_N\left(\frac{w}{z} - 1\right) dw, \end{aligned}$$

when  $f$  and all its derivatives have sufficient decay at infinity, where the last integral is taken along a path of fixed argument.

We now apply this proposition to compute asymptotic expansions of the functions  $F_{j,k,l}(q)$  and  $G_{j,k}(q)$ .

**Proposition 3.2.** *As  $z \rightarrow 0$ , we have*

$$F_{j,k,l}(e^{-z}) \sim \frac{1}{z} \int_0^\infty (1 - e^{-2x})^j e^{-lx} dx.$$

*Proof.* We use Proposition 3.1 to prove Proposition 3.2. It is not quite obvious how to put  $F_{j,k,l}(q)$  into the appropriate form, but this can be fixed by introducing a second auxiliary variable as in [6].

Let  $t$  be an arbitrary complex number in some region  $D_\theta$  as defined in Proposition 3.1. Define the function

$$F_{j,k,l}(t; z) := \sum_{m \geq 0} (e^{-2mz-kt}; e^{-2t})_j e^{-lmz}.$$

Observe that  $F_{j,k,l}(z; z) = F_{j,k,l}(e^{-z})$ . Furthermore, if we define

$$f_{j,k,l}(t; z) := (e^{-2z-kt}; e^{-2t})_j e^{-lz},$$

we can see that

$$F_{j,k,l}(t; z) = \sum_{m \geq 0} f_{j,k,l}(t; mz) = f_{j,k,l}(t; 0) + \sum_{m \geq 0} f_{j,k,l}(t; (m+1)z).$$

Observe now that for any fixed  $t$  in  $D_\theta$ ,  $f_{j,k,l}(t; z)$  is a holomorphic function of  $z$  in  $D_\theta$  and can be written as a linear combination of exponential decay functions. Therefore, it is clear that  $f_{j,k,l}(t; z)$  satisfies the criterion of Proposition 3.1 for all fixed  $t \in D_\theta$ , and therefore

$$\begin{aligned} \sum_{m \geq 0} f_{j,k,l}(t; (m+1)z) &= \frac{1}{z} \int_0^\infty f_{j,k,l}(t; x) dx \\ &\quad - \sum_{m \geq 0} \frac{f_{j,k,l}^{(m)}(t; 0) z^m}{(m+1)!} - \sum_{n=0}^{N-1} \frac{B_{n+1}(0) f_{j,k,l}^{(n)}(t; z)}{(n+1)!} z^n \\ &\quad - \frac{(-1)^N z^{N-1}}{N!} \int_z^{z\infty} f_{j,k,l}^{(N)}(t; w) \tilde{B}_N\left(\frac{w}{z} - 1\right) dw. \end{aligned}$$

Since we restrict to  $z \rightarrow 0$  in the region  $D_\theta$ , we obtain

$$\begin{aligned} F_{j,k,l}(e^{-z}) &= F_{j,k,l}(z; z) = f_{j,k,l}(z; 0) + \frac{1}{z} \int_0^\infty f_{j,k,l}(z; x) dx \\ &\quad - \sum_{m \geq 0} \frac{f_{j,k,l}^{(m)}(z; 0) z^m}{(m+1)!} - \sum_{n=0}^{N-1} \frac{B_{n+1}(0) f_{j,k,l}^{(n)}(z; z)}{(n+1)!} z^n \\ &\quad - \frac{(-1)^N z^{N-1}}{N!} \int_z^{z\infty} f_{j,k,l}^{(N)}(z; w) \tilde{B}_N \left( \frac{w}{z} - 1 \right) dw. \end{aligned}$$

Now, observe that  $f_{j,k,l}(t; z)$  is holomorphic at  $t = 0$  and therefore the identification  $t = z$  does not introduce any additional singularities at  $z = 0$ . Furthermore, because  $f_{j,k,l}(t; z)$  is holomorphic at zero for both the  $t$  and  $z$  variables, it is easy to see that the only term in this expression which contributes to the principal part as  $z \rightarrow 0$  is  $\frac{1}{z} \int_0^\infty f_{j,k,l}(z; x) dx$ , and therefore as  $z \rightarrow 0$  in  $D_\theta$  we obtain

$$F_{j,k,l}(e^{-z}) \sim \frac{1}{z} \int_0^\infty f_{j,k,l}(z; x) dx.$$

It is straightforward to see analytically that  $\lim_{z \rightarrow 0} \int_0^\infty f_{j,k,l}(z; x) dx = \int_0^\infty f_{j,k,l}(0; x) dx$ , and therefore as  $z \rightarrow 0$  in  $D_\theta$  we obtain

$$F_{j,k,l}(e^{-z}) \sim \frac{1}{z} \int_0^\infty f_{j,k,l}(0; x) dx.$$

The result follows.  $\square$

We define for convenience the integrals

$$I_A(j, l) := \int_0^\infty f_{j,k,l}(0; x) dx = \int_0^\infty (1 - e^{-2x})^j e^{-lx} dx. \quad (3.3)$$

Observe that the notation  $I_A(j, l)$  is sufficient because the resulting integral does not depend on  $k$ . Combining these calculations with (3.1), it follows that for  $q = e^{-z}$ , as  $z \rightarrow 0$  in  $D_\theta$ , we have

$$\begin{aligned} \tilde{a}_h(q) &\sim \frac{(-q; q)_\infty}{z} \left[ \sum_{j=0}^{\lceil h/2 \rceil - 1} \binom{h-j-1}{j} I_A(j, 2h-4j) \right. \\ &\quad \left. + \sum_{j=0}^{\lfloor h/2 \rfloor - 1} \binom{h-j-2}{j} I_A(j, 2h-4j-2) \right]. \quad (3.4) \end{aligned}$$

We follow a very similar process to estimate  $G_{j,k}(q)$ , and we obtain the following asymptotic formula.

**Proposition 3.3.** *As  $z \rightarrow 0$ , we have*

$$G_{j,k}(e^{-z}) \sim \frac{1}{z} \int_0^\infty \frac{e^{-jx}}{(1 + e^{-x})^k} dx.$$

*Proof.* As before, we restrict  $t, z$  to a region  $D_\theta$ ,  $q = e^{-z}$ . Consider

$$g_{j,k}(t; z) := \frac{e^{-jz}}{(e^{-z-t}; e^{-t})_k}.$$

Then we have

$$G_{j,k}(e^{-z}) = \sum_{m \geq 0} g_{j,k}(z; mz) = g_{j,k}(z; 0) + \sum_{m \geq 0} g_{j,k}(z; (m+1)z).$$

Using Proposition 3.1 with  $t$  fixed as in the previous case, we obtain for

$$G_{j,k}(t; z) := \sum_{m \geq 0} g_{j,k}(t; mz)$$

the identity

$$\begin{aligned} G_{j,k}(t; z) &= g_{j,k}(t; 0) + \frac{1}{z} \int_0^\infty g_{j,k}(t; x) dx - \sum_{m \geq 0} \frac{g_{j,k}^{(m)}(t; 0) z^m}{(m+1)!} \\ &\quad - \sum_{n=0}^{N-1} \frac{B_{n+1}(0) g_{j,k}^{(n)}(t; z)}{(n+1)!} z^n - \frac{(-1)^N z^{N-1}}{N!} \int_z^{z\infty} g_{j,j}^{(N)}(t; w) \tilde{B}_N \left( \frac{w}{z} - 1 \right) dw, \end{aligned}$$

valid for any  $N \geq 1$ . As before, the holomorphicity properties of  $g_{j,k}(t; z)$  in the  $z$  and  $t$  variables imply that as  $z \rightarrow 0$  in  $D_\theta$ , we have

$$G_{j,k}(e^{-z}) \sim G_{j,k}(z; z) \sim \frac{1}{z} \int g_{j,k}(z; x) dx \sim \frac{1}{z} \int_0^\infty g_{j,k}(0; x) dx.$$

The result follows.  $\square$

If we define the integrals

$$I_B(j, k) := \int_0^\infty g_{j,k}(0; x) dx = \int_0^\infty \frac{e^{-jx}}{(1+e^{-x})^k} dx, \quad (3.5)$$

then for  $q = e^{-z}$ , as  $z \rightarrow 0$  in  $D_\theta$ , we obtain

$$\tilde{b}_h(q) \sim \frac{(-q; q)_\infty}{z} \sum_{j=0}^{\lceil h/2 \rceil - 1} \binom{h-j-1}{j} I_B(j+1, h-j). \quad (3.6)$$

To summarize our asymptotic analysis so far, we have the following asymptotic formulas for  $\tilde{a}_h(q)$  and  $\tilde{b}_h(q)$  from Propositions 3.2 and 3.3.

**Theorem 3.4.** *For  $q = e^{-z}$ , as  $z \rightarrow 0$  we have*

$$\tilde{a}_h(q) \sim \frac{\alpha_h}{z} (-q; q)_\infty \quad \text{and} \quad \tilde{b}_h(q) \sim \frac{\beta_h}{z} (-q; q)_\infty,$$

where  $\alpha_h$  and  $\beta_h$  are constants defined by

$$\alpha_h := \sum_{j=0}^{\lceil h/2 \rceil - 1} \binom{h-j-1}{j} I_A(j, 2h-4j) + \sum_{j=0}^{\lfloor h/2 \rfloor - 1} \binom{h-j-2}{j} I_A(j, 2h-4j-2)$$

and

$$\beta_h := \sum_{j=0}^{\lceil h/2 \rceil - 1} \binom{h-j-1}{j} I_B(j+1, h-j),$$

and  $I_A(j, k)$  and  $I_B(j, k)$  are defined in (3.3) and (3.5), respectively.

Since we have shown in Section 2 that  $\tilde{a}_h(q)/(-q; q)_\infty$  and  $\tilde{b}_h(q)/(-q; q)_\infty$  are essentially rational functions (in particular, when  $q = e^{-z}$ , they only have polar singularities as  $z \rightarrow 0$ ), we can now apply Wright's circle method [21] to obtain an asymptotic expansion for the coefficients  $a_h(n)$  and  $b_h(n)$ .

**3.2. Wright's circle method.** We now apply Wright's circle method to obtain asymptotics for the sequences  $a_h(n)$  and  $b_h(n)$  as  $n \rightarrow \infty$ . We use the following special case of the formulation given by Ngo and Rhoades [21, Proposition 1.8].

**Proposition 3.5.** *Suppose that  $c(n)$  are integers defined by*

$$\sum_{n \geq 0} c(n)q^n = L(q)\xi(q)$$

for analytic functions  $L, \xi$  satisfying the following hypotheses:

(H1) As  $|z| \rightarrow 0$  in the cone  $D_\theta$ , we have

$$L(e^{-z}) \sim \frac{1}{z} \sum_{k \geq 0} a_k z^k$$

for  $a_k \in \mathbb{C}$ ,

(H2) As  $|z| \rightarrow 0$  in the cone  $D_\theta$ , we have

$$\xi(e^{-z}) = K e^{\frac{A}{z}} \left(1 + O_\theta \left(e^{-\frac{B}{z}}\right)\right)$$

for  $K, A \geq 0$  and  $B > 0^2$ ,

(H3) As  $|z| \rightarrow 0$  outside  $D_\theta$  and satisfying  $\operatorname{Re}(z) > 0$ , we have

$$|L(e^{-z})| \ll_\theta |z|^{-C}$$

for some  $C > 0$ ,

(H4) As  $|z| \rightarrow 0$  outside  $D_\theta$  and satisfying  $\operatorname{Re}(z) > 0$ , we have

$$|\xi(e^{-z})| \ll_\theta \xi(|e^{-z}|) e^{-\frac{\delta'}{\operatorname{Re}(z)}}$$

for some  $\delta' > 0$ .

Then as  $n \rightarrow \infty$ , we have for any  $N \in \mathbb{Z}^+$  that

$$c(n) = K e^{2\sqrt{A}n} n^{-\frac{1}{4}} \left( \sum_{r=0}^{N-1} p_r n^{-\frac{r}{2}} + O\left(n^{-\frac{N}{2}}\right) \right),$$

where  $p_r := \sum_{j=0}^r t_j c_{j,r-j}$  with  $t_j \in \mathbb{C}$  and  $c_{j,r} := \frac{\left(-\frac{1}{4\sqrt{A}}\right)^r \sqrt{A}^{j-\frac{1}{2}} \Gamma(j + \frac{1}{2} + r)}{2\sqrt{\pi} r! \Gamma(j + \frac{1}{2} - r)}$ .

**Remark 3.6.** We note that because our generating functions are given as the generating function for partitions into odd parts multiplied by a rational or nearly rational function, there are methods which in principle can give better error terms. See for example [14] for the case of unrestricted partitions.

---

<sup>2</sup>Ngo and Rhoades require that  $B > A$ , which arises naturally in the case that  $\xi$  is modular but is not strictly necessary when proving Proposition 3.5.



Note that hypotheses (H1) and (H2) require asymptotics for  $L$  and  $\xi$  on the major arc, for  $q$  near 1, while hypotheses (H3) and (H4) require that  $L$  and  $\xi$  are small on the minor arc, for  $q$  away from 1.

We are now in a position to prove Theorems 1.4 and 1.5.

*Proofs of Theorems 1.4 and 1.5.* From the modularity of the Dedekind eta function  $\eta(z) := q^{1/24} \prod_{n \geq 1} (1 - q^n)$ , we obtain the transformation law  $\eta(-1/z) = \sqrt{-iz} \eta(z)$ , from which it is easy to show using Proposition 3.5 (see [4] for more details) that for  $q = e^{-z}$  we have

$$(-q; q)_\infty = \frac{1}{\sqrt{2}} \exp\left(\frac{\pi^2}{12z}\right) \left(1 + O_\delta\left(e^{-\frac{B}{z}}\right)\right) \quad (3.7)$$

as  $z \rightarrow 0$  inside  $D_\theta$ . If  $z = x + iy$ , then outside of  $D_\theta$  we have that

$$|\xi(q)| \ll_\theta \xi(e^{-x}) \cdot \exp\left(-\frac{\delta'}{x}\right)$$

for some  $\delta' > 0$ . More specifically, [4, Lemmas 5.9 and 5.10] show that  $(-q; q)_\infty$  satisfies hypotheses (H2) and (H4) with  $K = \frac{1}{\sqrt{2}}$ ,  $A = \frac{\pi^2}{12}$  for both  $\tilde{a}_h(q)$  and  $\tilde{b}_h(q)$ . We calculate that

$$p_0 = \alpha_h c_{0,0} = \alpha_h \frac{3^{1/4}}{\pi\sqrt{2}} \quad \text{and} \quad p_0 = \beta_h c_{0,0} = \beta_h \frac{3^{1/4}}{\pi\sqrt{2}}$$

for  $\tilde{a}_h(q)$  and  $\tilde{b}_h(q)$ , respectively. Therefore, we have

$$a_h(n) \sim \alpha_h \frac{3^{1/4}}{2\pi n^{1/4}} e^{\pi\sqrt{\frac{n}{3}}} \quad \text{and} \quad b_h(n) \sim \beta_h \frac{3^{1/4}}{2\pi n^{1/4}} e^{\pi\sqrt{\frac{n}{3}}},$$

which completes the proofs of the theorems.  $\square$

**Remark 3.7.** Note that in order to compute the constants  $N_h$  in Theorem 1.2, one could use the explicit bounds in [18]. To do this, one must first make the asymptotics for the rational functions  $\tilde{A}_h(q)$  and  $\tilde{B}_h(q)$  from Theorem 2.2 effective, which could potentially be done by either an effective Taylor theorem or effective Euler–Maclaurin summation as used in [10, 18].

We now see that the inequality  $a_h(n) > b_h(n)$  will follow for  $n \gg 0$  if  $\alpha_h > \beta_h$ .

#### 4. EVALUATING $\alpha_h$ AND $\beta_h$

In this section, we complete the proof of Theorem 1.2 by showing that  $\alpha_h > \beta_h$  for all  $h \geq 2$ . In order to do this, recall that

$$\alpha_h := \sum_{j=0}^{\lceil h/2 \rceil - 1} \binom{h-j-1}{j} I_A(j, 2h-4j) + \sum_{j=0}^{\lfloor h/2 \rfloor - 1} \binom{h-j-2}{j} I_A(j, 2h-4j-2)$$

and

$$\beta_h := \sum_{j=0}^{\lceil h/2 \rceil - 1} \binom{h-j-1}{j} I_B(j+1, h-j),$$

where the integrals  $I_A(j, k)$  and  $I_B(j, k)$  are defined by (3.3) and (3.5). In order to prove that  $\alpha_h > \beta_h$  for all  $h \geq 2$ , we proceed in stages. First, we produce simpler

formulas for  $\alpha_h$  and  $\beta_h$  which involve harmonic numbers. We then leverage these simpler formulas to prove that  $\alpha_h > \beta_h$  for  $h \geq 2$ . In fact, we will prove something much stronger:

**Theorem 4.1.** *The following are true:*

- (1) We have  $\alpha_h \rightarrow \log(2)$  and  $\beta_h \rightarrow \frac{\log(3)}{2}$  as  $h \rightarrow \infty$ .
- (2) We have  $\alpha_h > \beta_h$  for all  $h \geq 2$ .

Observe that Theorems 1.4, 1.5, and 4.1 together imply Theorem 1.6 in the introduction, which then completes the proof of Theorem 1.2. Therefore, we focus the remainder of this section on the proof of Theorem 4.1.

To prove this theorem, we prove a sequence of lemmas that give successively simpler values for  $\alpha_h$  and  $\beta_h$ . To help with this, we define

$$\tilde{\alpha}_h := \sum_{j=0}^{\lfloor \frac{h-1}{2} \rfloor} \sum_{k=0}^j \binom{h-j-1}{j} \binom{j}{k} \frac{(-1)^k}{h-2j+k}. \quad (4.1)$$

**Lemma 4.2.** *For any integer  $h \geq 1$ , we have*

$$\alpha_h = \frac{\tilde{\alpha}_h + \tilde{\alpha}_{h-1}}{2}.$$

*Proof.* By checking the changes induced by taking  $h \mapsto h+1$ , the result will follow from the definition of  $\alpha_h$  if we can show that

$$\frac{\tilde{\alpha}_h}{2} = \sum_{j=0}^{\lfloor \frac{h-1}{2} \rfloor} \binom{h-j-1}{j} I_A(j, 2h-4j),$$

which in turn follows if we prove that

$$I_A(j, 2h-4j) = \sum_{k=0}^j \binom{j}{k} \frac{(-1)^k}{h-2j+k}.$$

Now, from the definition of  $I_A(j, l)$  and by the substitution  $u = e^{-2x}$ , we have

$$I_A(j, 2h-4j) = \int_0^\infty (1 - e^{-2x})^a e^{-2(h-2j)x} dx = \frac{(-1)^j}{2} \int_0^1 (u-1)^j u^{h-2j-1} du.$$

The result then follows by expanding  $(u-1)^j$  with the binomial theorem and integrating.  $\square$

**Lemma 4.3.** *Let  $h \geq 1$  be a positive integer. Then we have*

$$\beta_h = \sum_{j=0}^{\frac{h-2}{2}} \binom{h-j-1}{j} \sum_{k=0}^j \binom{j}{k} \frac{(-1)^k}{h-2j+k-1} \left(1 - \frac{1}{2^{h-2j+k-1}}\right)$$

whenever  $h$  is even and

$$\beta_h = \log(2) + \sum_{j=0}^{\frac{h-1}{2}} \binom{h-j-1}{j} \sum_{k=1}^j \binom{j}{k} \frac{(-1)^k}{h-2j+k-1} \left(1 - \frac{1}{2^{h-2j+k-1}}\right)$$

$$+ \sum_{j=0}^{\frac{h-3}{2}} \binom{h-j-1}{j} \frac{1}{h-2j-1} \left(1 - \frac{1}{2^{h-2j-1}}\right)$$

whenever  $h$  is odd.

*Proof.* From the definition of  $I_B(j, k)$ , we have

$$\beta_h = \sum_{j=0}^{\lfloor \frac{h-1}{2} \rfloor} \binom{h-j-1}{j} \int_0^\infty \frac{e^{-(j+1)x}}{(1+e^{-x})^{h-j}} dx.$$

We focus on the integral. Using the substitution  $u = 1 + e^{-x}$  and expanding the resulting power of  $(u-1)$  with the binomial theorem, we have

$$\int_0^\infty \frac{e^{-(j+1)x}}{(1+e^{-x})^{h-j}} dx = \int_1^2 \frac{(u-1)^j}{u^{h-j}} du = \sum_{k=0}^j \binom{j}{k} \int_1^2 u^{2j-h-k} du.$$

Now, considering the limitations  $0 \leq k \leq j$  and  $0 \leq j \leq \lfloor \frac{h-1}{2} \rfloor$ , we have  $-h \leq 2j - h - k \leq -1$ , with equality to  $-1$  if and only if  $j = \frac{h-1}{2}$  and  $k = 0$ . Note that this scenario is only possible if  $h$  is odd. Thus, we have

$$\int_1^2 u^{2j-h-k} du = \begin{cases} \frac{1}{h-2j+k-1} \left(1 - \frac{1}{2^{h-2j+k-1}}\right) & \text{if } (j, k) \neq \left(\frac{h-1}{2}, 0\right), \\ \log(2) & \text{if } (j, k) = \left(\frac{h-1}{2}, 0\right). \end{cases}$$

It is therefore convenient to split into cases. Firstly, if  $h$  is even, then the  $\log(2)$  term does not emerge and we obtain

$$\beta_h = \sum_{j=0}^{\frac{h-2}{2}} \binom{h-j-1}{j} \sum_{k=0}^j \binom{j}{k} \frac{(-1)^k}{h-2j+k-1} \left(1 - \frac{1}{2^{h-2j+k-1}}\right).$$

If  $h$  is odd on the other hand, we must isolate the term  $(j, k) = \left(\frac{h-1}{2}, 0\right)$ , and so we obtain

$$\begin{aligned} \beta_h &= \log(2) + \sum_{j=0}^{\frac{h-1}{2}} \binom{h-j-1}{j} \sum_{k=1}^j \binom{j}{k} \frac{(-1)^k}{h-2j+k-1} \left(1 - \frac{1}{2^{h-2j+k-1}}\right) \\ &\quad + \sum_{j=0}^{\frac{h-3}{2}} \binom{h-j-1}{j} \frac{1}{h-2j-1} \left(1 - \frac{1}{2^{h-2j-1}}\right). \end{aligned}$$

This completes the proof.  $\square$

Lemmas 4.2 and 4.3 simplify our considerations to the evaluation of linear combinations of merely rational numbers. We will analyze these finite sums using the following lemmas studying related polynomials.

**Lemma 4.4.** *Define the polynomials*

$$F_{n,m}(x) := \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k x^{m+k}}{m+k}, \quad R_n(x) := \sum_{k=1}^n \binom{n}{k} \frac{(-1)^k x^k}{k}.$$

*Then the following are true:*

(1) We have

$$F_{n,m}(x) = \frac{x^m \sum_{j=0}^{n+m-1} (1-x)^{n-j} \binom{n+m-1-j}{m-1}}{m \binom{n+m}{m}}.$$

(2) We have

$$R_n(x) = \int_0^x \frac{(1-t)^n - 1}{t} dt.$$

*Proof.* We first observe that

$$F'_{n,m}(x) = \sum_{k=0}^n \binom{n}{k} (-1)^k x^{m+k-1} = x^{m-1} \sum_{k=0}^n \binom{n}{k} (-1)^k x^k = x^{m-1} (1-x)^n. \quad (4.2)$$

To prove (1), we compute the generating function for  $F_{n,m}(x)$ . We have by (4.2) that

$$\sum_{m \geq 1} \sum_{n \geq 0} F_{n,m}(x) \frac{u^n}{n!} \frac{v^{m-1}}{(m-1)!} = \sum_{m \geq 1} \sum_{n \geq 0} \frac{u^n}{n!} \frac{v^{m-1}}{(m-1)!} \int_0^x t^{m-1} (1-t)^n dt.$$

By swapping the order of summation and integration and using the series for  $e^t$ , we may simplify:

$$\begin{aligned} \sum_{m \geq 1} \sum_{n \geq 0} F_{n,m}(x) \frac{u^n}{n!} \frac{v^{m-1}}{(m-1)!} &= \int_0^x e^{vt} e^{(1-t)u} dt \\ &= \int_0^x e^{u+t(v-u)} dt \\ &= \frac{1}{v-u} e^{u+t(v-u)} \Big|_0^x \\ &= \frac{1}{v-u} (e^{xv-xu+u} - e^u) \\ &= \sum_{k \geq 1} \frac{(xv-xu+u)^k - u^k}{(v-u)k!} \\ &= x \sum_{k \geq 1} \frac{\sum_{j=0}^{k-1} (xv-xu+u)^{k-1-j} u^j}{k!}. \end{aligned}$$

Therefore, using the notation  $[x^n]F(x)$  to denote the coefficient of  $x^n$  in the expression  $F(x)$ , we obtain

$$\begin{aligned} F_{n,m}(x) &= [u^n v^{m-1}] n!(m-1)! x \sum_{k \geq 1} \frac{\sum_{j=0}^{k-1} (xv-xu+u)^{k-1-j} u^j}{k!} \\ &= [u^n v^{m-1}] n!(m-1)! x \sum_{k \geq 1} \frac{\sum_{j=0}^{k-1} (xv)^{m-1} ((1-x)u)^{k-j-m} \binom{k-1-j}{m-1} u^j}{k!} \\ &= [u^n] n!(m-1)! t^m \sum_{k \geq 1} \frac{\sum_{j=0}^{k-1} (1-x)^{k-j-m} u^{k-m} \binom{k-1-j}{m-1}}{k!} \end{aligned}$$

$$\begin{aligned}
&= n!(m-1)!x^m \frac{\sum_{j=0}^{n+m-1} (1-x)^{n-j} \binom{n+m-1-j}{m-1}}{(n+m)!} \\
&= \frac{x^m \sum_{j=0}^{n+m-1} (1-x)^{n-j} \binom{n+m-1-j}{m-1}}{m \binom{n+m}{m}}
\end{aligned}$$

This completes the proof of (1). To prove (2), it is enough to observe that

$$R'_n(x) = \sum_{k=1}^n \binom{n}{k} (-1)^k x^{k-1} = \frac{(1-x)^n - 1}{x}. \quad \square$$

We also require the following lemma for evaluating a different type of summation.

**Lemma 4.5.** *For  $n = 2m$  even, we have*

$$\sum_{k=0}^m \frac{1}{n-2k+1} \binom{n-k}{k} 2^k = \frac{1}{n+1} (2^{n+1} - 1).$$

*For  $n = 2m + 1$  odd, we have*

$$\sum_{k=0}^m \frac{1}{n-2k+1} \binom{n-k}{k} 2^k = \frac{1}{n+1} (2^{m+1} - 1)^2.$$

*Proof.* Firstly, we observe the fact that

$$\begin{aligned}
\sum_{n \geq 0} \sum_{k=0}^n \binom{n-k}{k} 2^k x^n &= \sum_{k \geq 0} (2x^2)^k \sum_{n \geq 2k} \binom{n-k}{k} x^{n-2k} = \sum_{k \geq 0} (2x^2)^k \frac{1}{(1-x)^{k+1}} \\
&= \frac{1}{3} \left( \frac{2}{1-2x} + \frac{1}{1+x} \right).
\end{aligned}$$

This generating function identity implies that

$$\sum_{n \geq 0} \sum_{k=0}^n \binom{n-k}{k} 2^k = \frac{2^{n+1} - (-1)^{n+1}}{3}.$$

To continue the proof, consider

$$S(m) := \sum_{k=0}^m \frac{1}{n-2k+1} \binom{n-k}{k} 2^k.$$

Using the previous identity, we have

$$\begin{aligned}
S(n) &= \frac{1}{n+1} \sum_{k=0}^m \binom{n-k}{k} 2^k + \frac{1}{n+1} \sum_{k=0}^m \frac{2k}{n-2k+1} \binom{n-k}{k} 2^k \\
&= \frac{2^{n+1} - (-1)^{n+1}}{3(n+1)} + \frac{2T(n)}{n+1},
\end{aligned}$$

where (for  $n = 2m$  or  $2m + 1$ )

$$T(n) = \sum_{k=0}^m \frac{k}{n-2k+1} \binom{n-k}{k} 2^k = \sum_{k=1}^m \binom{n-k}{k-1} 2^k$$

$$= 2 \sum_{k=1}^m \binom{n-1-(k-1)}{k-1} 2^{k-1} = 2 \sum_{k=0}^{m-1} \binom{n-1-k}{k} 2^k.$$

If  $n = 2m$  is even, we have

$$T(n) = 2 \left( \frac{2^n - (-1)^n}{3} \right),$$

and if  $n = 2m + 1$  is odd, we have

$$T(n) = 2 \left( \sum_{k=0}^m \binom{n-1-k}{k} 2^k - 2^m \right) = 2 \left( \frac{2^n - (-1)^n}{3} \right) - 2^{m+1}.$$

Thus, if  $n = 2m$  is even, we have

$$\begin{aligned} S(n) &= \frac{2^{n+1} - (-1)^{n+1}}{3(n+1)} + \frac{2T(n)}{n+1} \\ &= \frac{2^{n+1} - (-1)^{n+1}}{3(n+1)} + \frac{4}{n+1} \frac{2^n - (-1)^n}{3} = \frac{1}{n+1} (2^{n+1} - 1), \end{aligned}$$

and if  $n = 2m + 1$  is odd, we have

$$\begin{aligned} S(n) &= \frac{2^{n+1} - (-1)^{n+1}}{3(n+1)} + \frac{2T(n)}{n+1} = S(n-1) - \frac{2 \cdot 2^{m+1}}{n+1} \\ &= \frac{1}{2m+2} (2^{2m+2} + 1 - 2^{m+2}) = \frac{1}{2m+2} (2^{m+1} - 1)^2. \end{aligned}$$

This completes the proof.  $\square$

With the aid of Lemma 4.4, we now further simplify the formulas for  $\alpha_h$  and  $\beta_h$  given in Lemmas 4.2 and 4.3. For the following result, we need the well-known *harmonic numbers*, defined by  $H_0 = 0$  and

$$H_n := \sum_{k=1}^n \frac{1}{k}$$

for  $n \geq 1$ .

**Theorem 4.6.** *We have for all  $h \geq 1$  that*

$$\alpha_h = \frac{H_h - H_{\lceil \frac{h-1}{2} \rceil} + H_{h-1} - H_{\lceil \frac{h-2}{2} \rceil}}{2}.$$

*Proof.* By Lemma 4.4 (1), we have the identity

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{m+k} = F_{n,m}(1) = \frac{1}{m \binom{m+n}{n}}.$$

Using this identity with  $n = j$  and  $m = h - 2j$  we obtain from (4.1) that

$$\tilde{\alpha}_h = \sum_{j=0}^{\lfloor \frac{h-1}{2} \rfloor} \binom{h-j-1}{j} \sum_{k=0}^j \binom{j}{k} \frac{(-1)^k}{h-2j+k} = \sum_{j=0}^{\lfloor \frac{h-1}{2} \rfloor} \binom{h-j-1}{j} \frac{1}{(h-2j) \binom{h-j}{j}}$$

$$= \sum_{j=0}^{\lfloor \frac{h-1}{2} \rfloor} \frac{1}{h-2j} \left(1 - \frac{j}{h-j}\right) = \sum_{j=0}^{\lfloor \frac{h-1}{2} \rfloor} \frac{1}{h-j}.$$

Thus,

$$\tilde{\alpha}_h = \begin{cases} H_h - H_{h/2} & \text{if } 2|h, \\ H_h - H_{\frac{h-1}{2}} & \text{if } 2 \nmid h. \end{cases}$$

As it has already been established that  $\alpha = \frac{\tilde{\alpha}_h + \tilde{\alpha}_{h-1}}{2}$ , we obtain the desired formula for  $\alpha_h$ .  $\square$

We also note briefly that this theorem implies part of Theorem 4.1.

**Corollary 4.7.** *We have  $\alpha_h \rightarrow \log(2)$  as  $h \rightarrow \infty$ .*

*Proof.* Recall the well-known fact that

$$\lim_{n \rightarrow \infty} (H_n - \log(n)) = \gamma,$$

where  $\gamma$  is the Euler–Mascheroni constant. Thus,  $\tilde{\alpha}_h \rightarrow \log(h) - \log(h/2) = \log(2)$  and  $\alpha_h = \frac{\tilde{\alpha}_h + \tilde{\alpha}_{h-1}}{2} \rightarrow \log(2)$ .  $\square$

We now prove the limiting theorems for  $\beta_h$ , which is the last major step before we complete the proof of Theorem 4.1, and thus also of Theorem 1.2.

**Theorem 4.8.** *We have for all  $h \geq 1$  that*

$$\beta_h = \begin{cases} \sum_{c=0}^{\frac{h-2}{2}} \frac{1}{(2c+1)2^{2c+1}} & \text{if } 2|h, \\ \sum_{c=0}^{\frac{h-3}{2}} \frac{1}{(c+1)2^{c+1}} - \sum_{c=0}^{\frac{h-3}{2}} \frac{1}{(2c+2)2^{2c+2}} + \int_{1/2}^1 \frac{(1-x)^{\frac{h-1}{2}}}{x} dx & \text{if } 2 \nmid h. \end{cases}$$

*Proof.* We begin first with  $h = 2n$  even, in which case we have

$$\beta_h = \sum_{j=0}^{n-1} \sum_{k=0}^j \binom{h-j-1}{j} \binom{j}{k} \frac{(-1)^k}{h-2j+k-1} \left(1 - \frac{1}{2^{h-2j+k-1}}\right).$$

Using Lemma 4.4 (1), we have

$$\sum_{j=0}^{n-1} \sum_{k=0}^j \binom{h-j-1}{j} \binom{j}{k} \frac{(-1)^k}{h-2j+k-1} = \sum_{j=0}^{n-1} \frac{1}{h-2j-1},$$

and so we may write

$$\beta_h = \sum_{j=0}^{n-1} \frac{1}{h-2j-1} - U(h), \tag{4.3}$$

where

$$U(h) := \sum_{j=0}^{n-1} \frac{1}{h-2j-1} \sum_{k=0}^{h-j-2} \binom{h-j-k-2}{h-2j-2} \frac{1}{2^{h-j-k-1}}.$$

We first simplify  $U(h)$  and swap the order of summation:

$$\begin{aligned} U(h) &= \sum_{j=0}^{n-1} \frac{1}{h-2j-1} \sum_{k=0}^j \binom{h-2j-2+k}{k} \frac{1}{2^{h-2j-1+k}} \\ &= \sum_{k=0}^{n-1} \sum_{j=k}^{n-1} \frac{1}{h-2j-1} \binom{h-2j-2+k}{k} \frac{1}{2^{h-2j-1+k}}. \end{aligned}$$

Since  $h = 2n$ , we write  $b = n - j$  and obtain

$$U(h) = \sum_{k=0}^{n-1} \sum_{b=0}^{n-k-1} \frac{1}{2b+1} \binom{2b+k}{k} \frac{1}{2^{2b+1+k}} = \sum_{k=0}^{n-1} \frac{1}{2^k} \sum_{b=0}^{n-k-1} \frac{1}{2b+1} \binom{2b+k}{k} \frac{1}{2^{2b+1}}.$$

If we further reindex the double sum with  $c = b + k$ , we have by Lemma 4.4 (3) that

$$U(h) = \sum_{c=0}^{n-1} \sum_{k=0}^c \frac{1}{2c-2k+1} \binom{2c-k}{k} \frac{1}{2^{2c-k+1}} = \sum_{c=0}^{n-1} \frac{1}{2c+1} \left(1 - \frac{1}{2^{2c+1}}\right).$$

Thus, we may simplify (4.3) for  $\beta_h$  to the form

$$\beta_h = \sum_{j=0}^{n-1} \frac{1}{h-2j-1} - \sum_{c=0}^{n-1} \frac{1}{2c+1} \left(1 - \frac{1}{2^{2c+1}}\right) = \sum_{c=0}^{n-1} \frac{1}{(2c+1)2^{2c+1}}.$$

This proves the theorem in the case where  $h$  is even.

Now, assume  $h = 2n + 1$  is odd. By Lemma 4.3, we have

$$\begin{aligned} \beta_h &= \log(2) + \sum_{j=0}^n \binom{h-j-1}{j} \sum_{k=1}^j \binom{j}{k} \frac{(-1)^k}{h-2j+k-1} \left(1 - \frac{1}{2^{h-2j+k-1}}\right) \\ &\quad + \sum_{j=0}^{n-1} \binom{h-j-1}{j} \frac{1}{h-2j-1} \left(1 - \frac{1}{2^{h-2j-1}}\right). \end{aligned}$$

From Lemma 4.4 (1), we obtain as in the previous case that

$$\beta_h = \log(2) + \sum_{j=0}^{n-1} \frac{1}{h-2j-1} - U(h) + \sum_{k=1}^n \frac{(-1)^k}{k} \left(1 - \frac{1}{2^k}\right).$$

Now, Lemma 4.4 (2) implies that

$$\log(2) + \sum_{k=1}^n \frac{(-1)^k}{k} \left(1 - \frac{1}{2^k}\right) = \log(2) + \int_{1/2}^1 \frac{(1-x)^n - 1}{x} dx = \int_{1/2}^1 \frac{(1-x)^n}{x} dx,$$

and therefore

$$\beta_h = \sum_{j=0}^{n-1} \frac{1}{h-2j-1} - U(h) + \int_{1/2}^1 \frac{(1-x)^n}{x} dx.$$

Now, using Lemma 4.5, we therefore have

$$\beta_h = \sum_{j=0}^{n-1} \frac{1}{h-2j-1} - \sum_{c=0}^{n-1} \frac{1}{2c+2} \left(1 - \frac{2}{2^{c+1}} + \frac{1}{2^{2c+2}}\right) + \int_{1/2}^1 \frac{(1-x)^n}{x} dx$$



$$= \sum_{c=0}^{n-1} \frac{1}{(c+1)2^{c+1}} - \sum_{c=0}^{n-1} \frac{1}{(2c+2)2^{2c+2}} + \int_{1/2}^1 \frac{(1-x)^n}{x} dx.$$

This completes the proof for  $h$  odd.  $\square$

We also note briefly that this theorem implies the remaining part of Theorem 4.1 (1).

**Corollary 4.9.** *We have  $\beta_h \rightarrow \frac{\log(3)}{2}$  as  $h \rightarrow \infty$ .*

*Proof.* As  $n \rightarrow \infty$  it is clear that

$$\int_{1/2}^1 \frac{(1-x)^n}{x} dx \rightarrow 0$$

and therefore by Theorem 4.8 we have as  $n \rightarrow \infty$  that

$$\beta_{2n+1} \sim \sum_{c=0}^{\infty} \frac{1}{(c+1)2^{c+1}} - \sum_{c=0}^{\infty} \frac{1}{(2c+2)2^{2c+2}} = \sum_{c=1}^{\infty} \frac{1}{(2c+1)2^{2c+1}} = \frac{\log(3)}{2}.$$

Taking into consideration even  $h$ , it is clear that  $\beta_h \rightarrow \frac{\log(3)}{2}$  as  $h \rightarrow \infty$ .  $\square$

We are now ready to prove the remaining part of Theorem 4.1, and therefore also of 1.2.

*Proofs of Theorems 4.1 and 1.2.* By Corollaries 4.7 and 4.9 along with Theorems 1.4 and 1.5, Theorem 4.1 (1) is already proven, and so it only remains to show that  $\alpha_h > \beta_h$  for all  $h \geq 2$ .

To show this inequality, we first consider  $\beta_h$ . For  $h = 2n \geq 2$  even, it is clear from Theorem 4.8 that

$$\beta_{2n} < \sum_{c=1}^{\infty} \frac{1}{(2c+1)2^{2c+1}} = \frac{\log(3)}{2}.$$

To study  $\beta_{2n+1}$  for  $n \geq 1$ , we begin by observing that  $\beta_3 = \log(2) - \frac{1}{8} > \frac{\log(3)}{2}$ . For  $n \geq 2$ , we have by Theorem 4.8 that

$$\begin{aligned} \beta_{2n+1} - \beta_{2n-1} &= \frac{1}{n2^n} - \frac{1}{n2^{2n+1}} + \int_{1/2}^1 \frac{(1-x)^n - (1-x)^{n-1}}{x} dx \\ &= \frac{1}{n2^n} - \frac{1}{n2^{2n+1}} - \int_{1/2}^1 (1-x)^{n-1} dx \\ &= -\frac{1}{n2^{2n+1}}. \end{aligned}$$

Thus,  $\beta_h \leq \log(2) - \frac{1}{8}$  for all  $h \geq 2$ . We now consider a similar study of  $\alpha_h$ . Recall that  $\alpha_h = \frac{\tilde{\alpha}_h + \tilde{\alpha}_{h-1}}{2}$  for

$$\tilde{\alpha}_h = \begin{cases} H_h - H_{\frac{h}{2}} & \text{if } 2|h, \\ H_h - H_{\frac{h-1}{2}} & \text{if } 2 \nmid h. \end{cases}$$

Therefore, we have

$$\alpha_{h+1} - \alpha_h = \frac{\tilde{\alpha}_{h+1} - \tilde{\alpha}_{h-1}}{2}.$$

Now, for  $h$  odd we have

$$\begin{aligned}\tilde{\alpha}_{h+1} - \tilde{\alpha}_{h-1} &= \left(H_{h+1} - H_{\frac{h+1}{2}}\right) - \left(H_{h-1} - H_{\frac{h-1}{2}}\right) \\ &= (H_{h+1} - H_{h-1}) - \left(H_{\frac{h+1}{2}} - H_{\frac{h-1}{2}}\right) = \frac{1}{h} - \frac{1}{h+1} > 0,\end{aligned}$$

and likewise  $\tilde{\alpha}_{h+1} - \tilde{\alpha}_{h-1} > 0$  for  $h$  even as well. Thus,  $\alpha_h$  is an increasing function of  $h$ . Now, we have from [4] that  $\alpha_2 = \frac{3}{4} > \frac{1}{2} = \beta_2$  and  $\alpha_3 = \frac{2}{3} > \log(2) - \frac{1}{8} = \beta_3$ . Therefore, for  $h \geq 4$  we have  $\alpha_h > \alpha_3 > \beta_3 > \beta_h$ , which completes the proof.  $\square$

## 5. FINAL REMARKS

**5.1. Frequency of hooks in the rows of partitions.** Probabilistic features of partitions are of great interest, and asymptotic formulas derived from the circle method are very useful for studying such questions [11, 12, 21]. We briefly give an overview of some statistical corollaries which can be derived from our results when combined with other known asymptotics in the literature. To state these results, we let  $d(n)$  be the number of partitions into odd parts (or into distinct parts). We first give the average number of hooks equal to any  $h \geq 1$  among the partitions of  $n$  into odd parts.

**Corollary 5.1.** *Let  $\text{avg}_{\mathcal{L}}(h; n)$  be the average number of hooks of length  $h$  among the partitions of size  $n$  in the collection  $\mathcal{L}$ . Then we have*

$$\text{avg}_{\mathcal{O}}(h; n) \sim \frac{6\alpha_h}{\pi} \sqrt{\frac{n}{3}}, \quad \text{avg}_{\mathcal{D}}(h; n) \sim \frac{6\beta_h}{\pi} \sqrt{\frac{n}{3}}.$$

as  $n \rightarrow \infty$ .

*Proof.* It is a well-known consequence of the circle method (and indeed can be proven from (3.7) and Proposition 3.5) that

$$p_{\mathcal{O}}(n) = p_{\mathcal{D}}(n) \sim \frac{3^{3/4}}{12n^{3/4}} e^{\pi\sqrt{\frac{n}{3}}}. \quad (5.1)$$

Note that the average number of hooks of length  $h$  in a partition of  $n$  into odd parts or distinct parts is given by  $a_h(n)/p_{\mathcal{O}}(n)$ ,  $b_h(n)/p_{\mathcal{D}}(n)$ , respectively. This completes the proof along with Theorems 1.4 and 1.5.  $\square$

There is another natural probabilistic question which can now be answered about hooks in these restricted classes of partitions, namely the probability that a uniformly selected part from this class of partitions has a hook of that length in the corresponding row of the Ferrers diagram.

**Corollary 5.2.** *Let  $\text{prob}_{\mathcal{L}}(h; n)$  denote the probability that a randomly selected row from among the partitions in  $\mathcal{L}$  of size  $n$  has a hook of length  $h$ . Then we have*

$$\text{prob}_{\mathcal{O}}(h; n) \sim \frac{4\alpha_h}{\log(n)}, \quad \text{prob}_{\mathcal{D}}(h; n) \sim \frac{\beta_h}{\log(2)}$$

as  $n \rightarrow \infty$ .

*Proof.* Note that by construction of the diagram, each row can have at most one hook of length  $h$  in each row. Therefore, if we let  $d_{\mathcal{O}}(n)$  and  $d_{\mathcal{D}}(n)$  denote the number of parts among all partitions into odd parts or distinct parts, respectively,

it follows that the desired probabilities are given by  $a_h(n)/d_{\mathcal{O}}(n)$  and  $b_h(n)/d_{\mathcal{D}}(n)$ , respectively. From the main results of [10, 19] we obtain

$$d_{\mathcal{O}}(n) \sim \frac{3^{\frac{1}{4}} \log(n)}{8\pi n^{\frac{1}{4}}} e^{\pi\sqrt{\frac{n}{3}}}, \quad d_{\mathcal{D}}(n) \sim \frac{3^{\frac{1}{4}} \log(2)}{2\pi n^{\frac{1}{4}}} e^{\pi\sqrt{\frac{n}{3}}}$$

as  $n \rightarrow \infty$ . The result follows from Theorems 1.4 and 1.5.  $\square$

The results of this section together give a detailed comparison between the properties of hook numbers in these two classes of partitions. From Corollary 5.1, we reaffirm that the total number of hooks of length  $h \geq 2$  is larger for  $\mathcal{O}$  than for  $\mathcal{D}$ , and this inequality flips for  $h = 1$ . However, the perspective of individual parts tells a different story. As  $n \rightarrow \infty$ , we see that the “average row” of a partition into odd parts has a vanishing probability of containing a hook of length  $h$ , while that same probability for the number of partitions into distinct parts is the positive number  $\frac{\beta_h}{\log(2)}$ . By Corollary 4.9, this constant approaches  $\frac{\log(3)}{\log(4)} \approx 0.7924$ . Thus, most rows of partitions into distinct parts have a hook of any given length  $h \geq 1$ . This discrepancy is reflected by the fact that partitions into odd parts have many more parts than partitions into distinct parts do on average, as is seen from the asymptotics for  $d_{\mathcal{O}}(n)$  and  $d_{\mathcal{D}}(n)$ . It would be very interesting to more deeply study these probabilistic features of hooks in partitions, as is done for example in [16] for hook numbers which are divisible by  $h$  in unrestricted partitions.

**5.2. Open problems and questions.** The motivating question of this study can be greatly generalized; the underlying concept is to understand using the circle method (or other methods) how combinatorial statistics on partitions behave on different sub-families of partitions. Such studies can be immediately generalized to other related combinatorial objects, such as unimodal sequences or more general integer compositions.

It would also be natural to study such questions for other combinatorial statistics on standard integer partitions, such as ranks and cranks of partitions. Our result does not immediately give an asymptotic count or even inequality for these counting functions, but a sieving argument such as that in [11] might be useful here. Alternatively, one might add together generating functions  $\tilde{a}_{th}(q), \tilde{b}_{th}(q)$  over the values  $t \geq 1$  and perform an analysis parallel to that of this paper.

Finally, it would be interesting to pursue variations of this problem for other families of partitions which lie in bijection with one another. As there remain conjectures of [4] which are not resolved in this aspect, we focus on this case. The primary discussions not addressed here involve the functions  $a_h^*(n)$  and  $b_h^*(n)$ , which count the number of hooks of length  $h$  among self-conjugate partitions and partitions into distinct odd parts, respectively. In light of our main theorems and the data presented in [4], we present the following conjecture in this setting.

**Conjecture 5.3.** *For  $h \geq 2$ , there is a constant  $\gamma_h^* > 1$  such that  $a_h^*(n)/b_h^*(n) \rightarrow \gamma_h^*$  as  $n \rightarrow \infty$ .*

It would also be interesting if  $\gamma_h^*$  itself had a limit as  $h \rightarrow \infty$ , but we do not speculate on this here. The function  $a_h^*(n)$  is of additional interest, as it seems to have significant nontrivial divisibilities. We therefore restate here a conjecture of [4].

**Conjecture 5.4.** *For each  $n \geq 0$  and  $m \geq 1$ , we have*

$$a_{2m}^*(n) \equiv 0 \pmod{2m}.$$

## REFERENCES

- [1] G. Andrews. *The Theory of Partitions*, Cambridge University Press, no. 2 (1998).
- [2] G. Andrews. *Euler’s partition identity and two problems of George Beck*. Math. Student, 86(1-2):115–119, 2017.
- [3] A. Ayyer and S. Sinha. *The size of  $t$ -cores and hook lengths of random cells in partitions*. Ann. Appl. Probab. **33** (1), 85–106 (2023).
- [4] C. Ballantine, H. Burson, W. Craig, A. Folsom, and B. Wen. *Hook length biases and general linear partition inequalities*. Res. Math. Sci. **10**, 41 (2023).
- [5] C. Bessenrodt and G.-N. Han. *Symmetry distribution between hook length and part length for partitions*. Discrete Math. **309** (2009), pp. 6070–6073.
- [6] K. Bringmann, W. Craig, and J. Males. *Asymptotics for  $d$ -fold partition diamonds and related infinite products*. Preprint, arXiv:2311.11805.
- [7] K. Bringmann, C. Jennings-Shaffer, and K. Mahlburg. *On a Tauberian theorem of Ingham and Euler–Maclaurin summation*. Ramanujan J. **61**, 55–86 (2023).
- [8] K. Bringmann, K. Ono, and I. Wagner. *Eichler integrals of Eisenstein series as  $q$ -brackets of weighted  $t$ -hook functions on partitions*. Ramanujan J. **61**, 279–293 (2023).
- [9] H. Cho, B. Kim, H. Nam, and J. Sohn. *A survey on  $t$ -cores partitions*. Hardy-Ramanujan J. **44**, 81–101 (2021).
- [10] W. Craig. *On the number of parts in congruence classes for partitions into distinct parts*. Res. Number Theory, **8** (3):52, 2022.
- [11] P. Erdős and J. Lehner. *The distribution of the number of summands in the partitions of a positive integer*. Duke Math. J. **8**, 335–345 (1941).
- [12] B. Fristedt, *The structure of random large partitions of integers*, Trans. Am. Math. Soc. **337**, 703–735 (1993).
- [13] F. Garvan, D. Kim, and D. Stanton. *Cranks and  $t$ -cores*. Invent. math. **101** 1, 1–18 (1990).
- [14] P.J. Grabner, A. Knopfmacher, and S. Wagner. *A General Asymptotic Scheme for the Analysis of Partition Statistics*. Comb. Probab. Comput., Volume 23, Issue 6: Honouring the Memory of Philippe Flajolet - Part 2, November 2014, pp. 1057–1086.
- [15] A. Granville, K. Ono. *Defect zero blocks for finite simple groups*. Trans. Am. Math. Soc. **348**, 1 (1996).
- [16] M. Griffin, K. Ono, and W.-L. Tsai. *Distributions of hook lengths in integer partitions*. Proc. Am. Math. Soc., Series B. In Press.
- [17] G.-N. Han. *The Nekrasov-Okounkov hook length formula: refinement, elementary proof, extension and applications*. Ann. Inst. Fourier (Grenoble), 60(1):1–29, 2010.
- [18] F. Jackson and M. Otgonbayar. *Biases among congruence classes for parts in  $k$ -regular partitions*. arXiv preprint, arXiv:2207.04352, 2022.
- [19] F. Jackson and M. Otgonbayar. *Unexpected biases between congruence classes for parts in  $k$ -indivisible partitions*. J. Number Theory **248**, 310–342 (2023).
- [20] N. A. Nekrasov and A. Okounkov. *Seiberg-Witten theory and random partitions*. In *The unity of mathematics*, Volume 244 of Progr. Math., pages 525–596. Birkhäuser Boston, Boston, MA, 2006.
- [21] H.T. Ngo and R. Rhoades. *Integer partitions, probabilities and quantum modular forms*. Res. Math. Sci. **4**, Article number: 17 (2017).
- [22] K. Ono and L. Sze. *4-core partitions and class numbers*. Acta Arith. **80** 3, 249–272 (1997).

MATHEMATICAL INSTITUTE, UNIVERSITY OF COLOGNE, GYRHOFSTR. 8B, 50931 COLOGNE,  
GERMANY

*Email address:* `wcraig@uni-koeln.de`

UNIVERSITY OF TEXAS AT TYLER, 3900 UNIVERSITY BOULEVARD, TYLER, TX 75799, USA

*Email address:* `mdawsey@uttyler.edu`

IRMA, UNIVERSITÉ DE STRASBOURG ET CNRS, 7 RUE RENÉ DESCARTES, 67084 STRAS-  
BOURG, FRANCE

*Email address:* `guoniu.han@unistra.fr`