2-covers of wide Young diagrams

Ron Aharoni,* Eli Berger,† He Guo‡ and Daniel Kotlar§

August 2023

Abstract

A Young diagram Y is called wide if every sub-diagram Z formed by a subset of the rows of Y dominates Z' , the conjugate of Z. A Young diagram Y is called Latin if its squares can be assigned numbers so that for each i, the ith row is filled injectively with the numbers $1, \ldots, a_i$, where a_i is the length of ith row of Y , and every column is also filled injectively. A conjecture of Chow and Taylor, publicized by Chow, Fan, Goemans, and Vondrak [\[3\]](#page-16-0) is that a wide Young diagram is Latin. We prove a dual version of the conjecture.

1 Introduction

Young diagrams are a way of representing partitions of natural numbers. Given a partition $n = a_1 + a_2 + a_3$ $\cdots + a_m$ of a number n, where $a_1 \ge a_2 \ge \cdots \ge a_m \ge 0$, the corresponding Young diagram Y is formed by m rows of squares that are left-aligned and the *i*th row consists of a_i squares. The size n of Y is denoted by |Y|. The *i*th row (resp. column) of Y are denoted by $r_i(Y)$ (resp. $c_i(Y)$). We write $a_i(Y)$ for $|r_i(Y)|$ and $b_i(Y)$ for $|c_i(Y)|$. When the identity of Y is clear from the context we omit its mention, and write r_i, c_i, a_i, b_i .

The conjugate Y' of a Young diagram Y is obtained by transposing the diagram. In terms of partitions, this is the partition $n = b_1 + b_2 + \cdots + b_{a_1}$ with $b_1 \geq b_2 \geq \cdots \geq b_{a_1} \geq 0$, where $b_j = |c_j|$, the length of the jth column of Y . The left-aligned-ness of the diagram implies

$$
b_k = |\{i : a_i \ge k\}|,\tag{1}
$$

$$
\sum_{1 \le j \le k} b_j = \sum_{1 \le i \le m} \min(k, a_i) \quad \text{and} \quad \sum_{1 \le i \le k} a_i = \sum_{1 \le j \le a_1} \min(k, b_j). \tag{2}
$$

A Young diagram X dominates a diagram Y partitioning the same n if

$$
\sum_{i=1}^{k} a_i(X) \ge \sum_{i=1}^{k} a_i(Y)
$$

for every $k \geq 1$, where partitions are extended by appending zero parts at the end as necessary.

Definition 1.1. A Young diagram is said to be wide if for every subset S of its rows, the diagram Z formed by S dominates Z' .

Definition 1.2. A filling of a Young diagram Y is an assignment of a number $y(i, j)$ to each square (i, j) that is in the ith row and jth column of Y, satisfying that for each i, the assignment is an injection from the a_i many squares in row i to $[a_i] = \{1, \ldots, a_i\}$. A filling is Latin if it is injective also in each column. A Young diagram is Latin if it has a Latin filling.

In [\[3\]](#page-16-0) the following is ascribed to Victor Reiner:

[∗]Faculty of Mathematics, Technion, Haifa 32000, Israel. E-mail: raharoni@gmail.com

[†]Department of Mathematics, University of Haifa, Haifa 31905, Israel. Email: berger.haifa@gmail.com

[‡]Faculty of Mathematics, Technion, Haifa 32000, Israel. Email: hguo@campus.technion.ac.il

[§]Computer Science Department, Tel-Hai College, Upper Galilee 12210, Israel. Email: dannykotlar@gmail.com

Theorem 1.3. A Latin Young diagram is wide.

Proof. Let Y be a Latin Young diagram with a Latin filling $(y(i, j))_{i,j}$. Let

$$
T_k = |\{(i, j) : y(i, j) \le k\}|.
$$

Clearly, $a_i(Y) \geq k$ if and only if $\{y(i, j) : 1 \leq j \leq a_i\}$ contains [k]. So, for every row i,

$$
|\{j : y(i,j) \le k\}| = \min(k, a_i).
$$

Summing over all rows and using [\(2\)](#page-0-0) gives

$$
T_k = \sum_{1 \le i \le m} \min(k, a_i) = \sum_{1 \le j \le k} b_j.
$$
 (3)

On the other hand, the injectivity of y in every column implies that for every fixed j,

$$
|\{i : y(i,j) \le k\}| \le k,
$$

implying

$$
|\{i : y(i,j) \le k\}| \le \min(k, b_j).
$$

Summing over all columns gives

$$
T_k \le \sum_{1 \le j \le a_i} \min(k, b_j) = \sum_{1 \le i \le k} a_i,
$$
\n(4)

where the last equality is again by (2) . Combining (3) and (4) , we have

$$
\sum_{1 \le i \le k} a_i \ge \sum_{1 \le j \le k} b_j.
$$

This argument works also for sub-diagrams formed by a subset of rows of Y, which verifies that Y is wide. \Box

A conjecture attributed in [\[3\]](#page-16-0) to Chow and Taylor is that the converse is also true:

Conjecture 1.4. If a Young diagram Y is wide then it is Latin.

In [\[3\]](#page-16-0) this is called the Wide Partition Conjecture, or WPC. The original motivation for the conjecture comes from Rota's basis conjecture [\[4\]](#page-16-1). Replacing the symbols with elements of a matroid and the condition of injectivity by matroid independence yields a generalization of Rota's conjecture.

1.1 A re-formulation in matching terminology

Following notation from [\[2\]](#page-16-2), a k-matching in a hypergraph H is an edge subset of H in which every two edges share fewer than k vertices (so, a 1-matching is a classical matching, i.e., a set of disjoint edges). The k th matching number $\nu^{(k)}(H)$ of H is the maximal size of a k-matching in H. Let $H^{(k)}$ be the hypergraph whose vertex set is $\{K \subseteq V(H) : |K| = k, K \subseteq e \text{ for some } e \in E(H)\}$ and whose edge set is $\{\binom{e}{k} : e \in E(H)\}$. Then a k-matching of H corresponds to a matching in $H^{(k)}$, so $\nu^{(k)}(H) = \nu(H^{(k)}),$ where $\nu(F)$ is the classical matching number of a hypergraph F.

A k-cover of a hypergraph H is a subset P of $V(H^{(k)}) = \{K \subseteq V(H) : |K| = k, K \subseteq e \text{ for some } e \in E(H)\}$ covering all edges of H, namely for every edge e of H, there exists some $K \in P$ such that $K \subseteq e$. By $\tau^{(k)}(H)$ we denote the integral LP-dual parameter of $\nu^{(k)}(H)$, namely the minimal size of a k-cover of H. Again, $\tau^{(k)}(H) = \tau(H^{(k)}),$ where $\tau(F)$ is the classical covering number of a hypergraph F. Obviously

$$
\tau^{(k)}(H) \ge \nu^{(k)}(H).
$$

Below we shall only consider the case $k = 2$. We say that two sets e_1, e_2 are almost disjoint if $|e_1 \cap e_2| \leq 1$.

Notation 1.5. Pairs and triples will be often written without delineating parentheses, so we write ab for ${a, b}$ and abc for ${a, b, c}$.

To a Young diagram Y we assign a tripartite hypergraph $H(Y)$, as follows. Its three respective sides are $R = R(Y) = \{r_1, \ldots, r_m\}$ (the set of rows), $C = C(Y) = \{c_1, \ldots, c_{a_1}\}$ (the set of columns), and $S = S(Y) = \{s_1, \ldots, s_{a_1}\}\$ (the set of numerical symbols).

For $1 \leq i \leq m$ and $r_i \in R$, let

$$
H_i(Y) = \{r_i c_j s_k \mid c_j \in C, s_k \in S, 1 \le j, k \le a_i\}.
$$

Let

$$
H(Y) = \bigcup_{1 \le i \le m} H_i(Y).
$$

Observation 1.6. $\tau^{(2)}(H(Y)) \leq |Y|$.

This follows from the fact that $\{r_ic_j \mid 1 \leq i \leq m, 1 \leq j \leq a_i\}$ is a 2-cover for $H(Y)$. An equivalent formulation of the WPC is:

Conjecture 1.7. If a Young diagram Y is wide, then $\nu^{(2)}(H(Y)) = |Y|$.

To see the equivalence, assume there is an assignment $y(i, j)$ satisfying the requirement of Conjecture [1.4.](#page-1-2) Then $M := \{r_i c_j s_{y(i,j)} \mid 1 \leq i \leq m, 1 \leq j \leq a_i\}$ is a 2-matching in $H(Y)$ of size $|Y|$. To see the almost disjointness, note that $r_i c_j$ is contained in one edge of M by the construction; since $y(i, j)$ is injective for fixed i, then $r_i s_k$ is contained in at most one edge of M; since the assignment is injective in each column, $c_i s_k$ is contained in at most one edge of M. For the other direction, let M be a 2-matching in $H(Y)$ of size $|Y| = \sum_{i=1}^{m} a_i$. Then for each $1 \leq i \leq m$ and $1 \leq j \leq a_i$, the pair $r_i c_j$ is contained in exactly one edge $r_i c_j s_k$ of M. Then we set $y(i, j) := k$. Note that $1 \leq y(i, j) \leq a_i$. Then y is a Latin filling of Y, where the row injectivity and column injectivity requirements follow from the fact that $r_i s_k$ and $c_j s_k$ can each be contained in at most one edge of M, respectively.

In this terminology Theorem [1.3](#page-1-3) says that if $\nu^{(2)}(H(Y)) = |Y|$ then Y is wide.

We prove a weaker version of Conjecture [1.7:](#page-2-0)

Theorem 1.8. If a Young diagram Y is wide, then $\tau^{(2)}(H(Y)) = |Y|$.

The converse is also true:

Theorem 1.9. If $\tau^{(2)}(H(Y)) = |Y|$, then the Young diagram Y is wide.

This is a strengthening of Theorem [1.3,](#page-1-3) since if $\nu^{(2)}(H(Y)) = |Y|$ then $\tau^{(2)}(H(Y)) = |Y|$. By these theorems, Y is wide if and only if $\tau^{(2)}(H(Y)) = |Y|$.

Theorem [1.9](#page-2-1) is proved in Section [3](#page-3-0) and Theorem [1.8](#page-2-2) in Section [4.](#page-5-0) In Section [5](#page-13-0) we discuss fractional versions. In Section [6](#page-14-0) we post some questions strengthening Conjecture [1.7.](#page-2-0)

2 Preliminaries

Notation 2.1. For two disjoint sets A and B, we denote $\{ab \mid a \in A, b \in B\}$ by $A \times B$.

This notation is usually reserved for the set of ordered pairs, but the assumption $A \cap B = \emptyset$ removes the risk of ambiguity.

Given a tripartite hypergraph F with sides A, B, C, we denote $\{bc \in B \times C \mid abc \in F \text{ for some } a \in A\}$ by $F[B \times C].$

Notation 2.2. We abbreviate $\{c_1, \ldots, c_i\} \times \{s_1, \ldots, s_j\} \subseteq C \times S$ by $[i] \times [j]$.

Notation 2.3. Given an integer ℓ and a subset Q of $C \times S$, let

$$
\nu(\ell, Q) := \nu([\ell] \times [\ell] - Q).
$$

Lemma 2.4. For any $Q \subseteq C \times S$ and $1 \leq i \leq m$, the minimal number of pairs in $\{r_i\} \times (C \cup S)$ needed to cover all the edges of H_i that are not covered by Q is $\nu(a_i, Q)$.

Proof. Let T_i be a minimum collection of pairs in $\{r_i\} \times (C \cup S)$ that cover all the edges of H_i that are not covered by Q. Then $\cup_{t\in T_i} t\cap (C\cup S)$ is a cover of the bipartite graph $[a_i]\times [a_i]-Q$, otherwise suppose $c_j s_k$ is not covered by $\cup_{t\in T_i} t\cap (C\cup S)$, then $r_ic_js_k\in H_i$ is not covered by T_i or Q , a contradiction. Hence

 $|T_i| \geq \nu(a_i, Q),$

by König's theorem.

On the other hand, let Z_i be a minimum cover of the bipartite graph $[a_i] \times [a_i] - Q$, it is routine to check that ${r_i} \times Z_i$ forms a 2-cover that covers all the edges of H_i that are not covered by Q. Therefore

$$
|T_i| \leq \nu(a_i, Q).
$$

This proves that $\nu(a_i, Q)$ is the desired covering number.

Corollary 2.5. For any $Q \subseteq C \times S$, there exists a 2-cover P of $H(Y)$ satisfying

$$
\tau^{(2)}(H) \le |P| = |Q| + \sum_{i=1}^{m} \nu(a_i, Q)
$$

and $Q = P \cap (C \times S)$.

Proof. Let $P = Q \cup \bigcup_{i=1}^{m} T_i$, where T_i is as in the proof of Lemma [2.4.](#page-2-3)

Observation 2.6. A pair in $\{r_i\} \times (C \cup S)$ does not cover any edge in H_j for $i \neq j$.

Notation 2.7. For a set of pairs $P \subseteq (R \times C) \cup (R \times S) \cup (C \times S)$ let $Q(P) = P \cap (C \times S)$.

Lemma [2.4](#page-2-3) and Observation [2.6](#page-3-1) imply the following.

Corollary 2.8. Let P be a 2-cover of $H(Y)$ and let $Q = Q(P)$. Then

$$
|P| \ge |Q| + \sum_{i=1}^{m} \nu(a_i, Q).
$$

Lemma 2.9. For $0 \leq p \leq q$, we have

$$
\nu(\ell,[p] \times [q]) = \begin{cases} 0, & \text{if } \ell \leq p. \\ \ell - p, & \text{if } p < \ell \leq q. \\ 2\ell - p - q, & \text{if } q < \ell \leq p + q. \\ \ell & \text{if } \ell > p + q. \end{cases}
$$

Proof. For $\ell \leq p$, $[\ell] \times [\ell] - [p] \times [q] = \emptyset$ so that $\nu(\ell, [p] \times [q]) = 0$.

For $p < \ell \leq q$, in $[\ell] \times [\ell] - [p] \times [q]$ we can find a matching $\{c_{p+1}s_{p+1}, \ldots, c_{\ell}s_{\ell}\}\$ of size $\ell - p$ and a vertex-cover $\{c_{p+1}, \ldots, c_{\ell}\}\$ of the same size, implying $\nu(\ell, [p] \times [q]) = \ell - p$.

For $q < \ell \leq p+q$, in $[\ell] \times [\ell]-[p] \times [q]$ we can find a matching $\{c_{p+1}s_1, \ldots, c_\ell s_{\ell-p}, s_{q+1}c_1, \ldots, s_\ell c_{\ell-q}\}$ of size $(\ell-p)+(\ell-q)$ and a vertex-cover $\{c_{p+1},\ldots,c_{\ell},s_{q+1},\ldots,s_{\ell}\}\$ of the same size, therefore $\nu(\ell,[p]\times[q])=2\ell-p-q.$

By the third line of Lemma [2.9,](#page-3-2) $\nu(p+q, [p] \times [q]) = p+q$, which means that $[p+q] \times [p+q] - [p] \times [q]$ has a perfect matching. Then for $\ell > p+q$, this matching together with $\{c_{\ell'}s_{\ell'} | p+q < \ell' \leq \ell\}$ forms a perfect matching in $[\ell] \times [\ell] - [p] \times [q]$. Therefore $\nu(\ell, [p] \times [q]) = \ell$ for $\ell > p + q$. \Box

3 $\tau^{(2)}(H(Y)) = |Y|$ implies wideness of Y

We prove Theorem [1.9](#page-2-1) in its contrapositive form.

Theorem 3.1. If a Young diagram Z is not wide, then $\tau^{(2)}(H(Z)) < |Z|$.

 \Box

Proof of Theorem [3.1.](#page-3-3) By the assumption that Z is not wide, there exists a subpartition Y formed by a set of rows of Z , such that Y does not dominate Y'. Let k be the minimum number that witnesses this fact, namely

$$
\sum_{i=1}^{k} a_i(Y) < \sum_{i=1}^{k} b_i(Y). \tag{5}
$$

Write $a_i = a_i(Y)$ and $b_j = b_j(Y)$.

Claim 3.1.1. $k \leq a_k$.

Proof. Assume the contrary and let $\ell = a_k < k$. We have

$$
\sum_{i=1}^{\ell} a_i = \sum_{i=1}^{k} a_i - \sum_{\ell < i \le k} a_i \quad \text{and} \quad \sum_{j=1}^{\ell} b_j = \sum_{j=1}^{k} b_j - \sum_{\ell < j \le k} b_j. \tag{6}
$$

If we show

$$
\sum_{\ell < i \le k} a_i \ge \sum_{\ell < i \le k} b_j,\tag{7}
$$

then combining (6) , (7) , and the assumption (5) , we have

$$
\sum_{i=1}^{\ell} a_i < \sum_{j=1}^{\ell} b_j,
$$

contradicting the minimality of k.

To prove [\(7\)](#page-4-1), on the one hand,

$$
\sum_{\ell < i \le k} a_i \ge \left| \left\{ (i,j) \in Y : \ell < i \le k, 1 \le j \le \ell \right\} \right| + \underbrace{\left| \left\{ (i,j) \in Y : \ell < i \le k, \ell < j \le k \right\} \right|}_{=:L_{\ell,k}} \tag{8}
$$

where the equality is because $a_i \ge a_k = \ell$ for each $1 \le i \le k$.

On the other hand,

$$
\sum_{\ell < j \le k} b_j \le \left| \left\{ (i,j) \in Y : 1 \le i \le \ell, \ell < j \le k \right\} \right| + \left| \left\{ (i,j) \in Y : \ell < i \le k, \ell < j \le k \right\} \right|
$$
\n
$$
\le (k - \ell)\ell + L_{\ell,k},\tag{9}
$$

where the first inequality is because $a_k = \ell$, then for each $j > \ell$, $b_j = |\{i : a_i \geq j\}| \leq k$ and then the squares in the $\ell + 1, ..., k$ th columns of Y are contained in $\{(i, j) \in Y \mid 1 \le i \le k, \ell < j \le k\}$. Combining [\(8\)](#page-4-3) and [\(9\)](#page-4-4) vields (7), and thus the claim. yields [\(7\)](#page-4-1), and thus the claim.

Claim 3.1.2. $\tau^{(2)}(H(Y)) < \sum_{i=1}^{m} a_i = |Y|.$

Proof. Let $Q = [k] \times [a_k]$, $T_0 = \{i : a_i \le k, i \ge k+1\}$, $T_1 = \{i : k < a_i \le a_k, i \ge k+1\}$, $T_2 = \{i : a_k \le a_i \le k+1\}$ $k + a_k, i \leq k$, and $T_3 = \{i : a_i > k + a_k\}.$

By Corollary [2.5](#page-3-4) and Lemma [2.9](#page-3-2) (that can be applied here in view of Claim [2.9,](#page-3-2))

$$
\tau^{(2)}(H(Y)) \le |Q| + \sum_{i} \nu(a_i, Q)
$$

= $ka_k + \sum_{i \in T_1} (a_i - k) + \sum_{i \in T_2} (2a_i - k - a_k) + \sum_{i \in T_3} a_i,$ (10)

and our aim is to show [\(10\)](#page-4-5) is less than $|Y| = \sum_i a_i$, i.e.,

$$
ka_k + \sum_{i \in T_1} (a_i - k) + \sum_{i \in T_2} (2a_i - k - a_k) + \sum_{i \in T_3} a_i < \sum_{i \in T_0} a_i + \sum_{i \in T_1} a_i + \sum_{i \in T_2} a_i + \sum_{i \in T_3} a_i.
$$

This is equivalent to showing

$$
ka_k + \sum_{i \in T_2} (a_i - a_k) + \sum_{i \in T_3} (a_i - a_k) < \sum_{i \in T_0} a_i + k|T_1| + k|T_2| + \sum_{i \in T_3} (a_i - a_k). \tag{11}
$$

Noting that $|T_2 \cup T_3| = k$, the left-hand side of [\(11\)](#page-5-1) is $\sum_{i \in T_2} a_i + \sum_{i \in T_3} a_i$, i.e., the sum of the lengths of the first k rows of Y. Since $a_i - a_k \ge k$ for $i \in T_3$, the right-hand side is at least $\sum_{i \in T_0} a_i + k|T_1| + k|T_2| + k|T_3|$, which is the sum of the lengths of the first k columns of Y. By our assumption (5) ,

$$
\sum_{i \in T_2} a_i + \sum_{i \in T_3} a_i < \sum_{i \in T_0} a_i + k|T_1| + k|T_2| + k|T_3|.
$$

Therefore , [\(11\)](#page-5-1) holds, completing the proof of Claim [3.1.2.](#page-4-6)

Now, let P be the minimum 2-cover of $H(Y)$. Then $|P| < |Y|$. Let $W = \{r_ic_j \mid (i,j) \in Z \setminus Y\}$. Then $P \cup W$ is a 2-cover of $H(Z)$, of size $|P| + |Z| - |Y|$, which is less than $|Z|$. We prove the theorem. \Box

4 Wideness of Y implies $\tau^{(2)}(H(Y)) = |Y|$

In this section we prove our main result, Theorem [1.8.](#page-2-2) Given a 2-cover P of $H(Y)$ for a wide diagram Y, we want to show that $|P| > |Y|$. As a first step, we replace P by a 2-cover having the same size, but more structure. We "shift" it, which in our context means that we "push left" the edges in $P \cap (C \times S)$ (the explicit definition is given below).

4.1 Shifting

Shifting is defined for an arbitrary set $P \subseteq (C \times S) \cup (R \times S) \cup (R \times C)$ with respect to either the C or S side of $H(Y)$ — we choose the first, namely C. Let $Q = Q(P) = P \cap (C \times S)$ and $W = W(P) = P \cap (C \times R)$.

For an integer t let $L_t = \{r_\ell \in R : a_\ell < t\}.$

Let c_i, c_j be two elements of C with $i < j$. The shifting $P' = \phi_{c_i,c_j}(P)$ of P is obtained from P by replacing all edges of P incident with $\{c_i, c_j\}$ by

$$
\{c_i\} \times \left(N_Q(c_i) \cup N_Q(c_j)\right) \bigcup \{c_j\} \times \left(N_Q(c_j) \cap N_Q(c_i)\right)
$$

$$
\bigcup \{c_i\} \times \left(N_W(c_i) \cap \left(N_W(c_j) \cup L_j\right)\right) \bigcup \{c_j\} \times \left(N_W(c_j) \cup \left(N_W(c_i) \setminus L_j\right)\right)
$$

Lemma 4.1. Let $i < j$ and let $P' = \phi_{c_i, c_j}(P)$. Then

- 1. $|P'| = |P|$, and
- 2. If P is a 2-cover of $H(Y)$, so is P'.

Proof. The first part follows from

$$
|N_Q(c_i) \cup N_Q(c_j)| + |N_Q(c_i) \cap N_Q(c_j)| = |N_Q(c_i)| + |N_Q(c_j)|,
$$

and for $T_1 = N_W(c_i) \cap (N_W(c_j) \cup L_j)$ and $T_2 = N_W(c_i) \cup (N_W(c_i) \setminus L_j)$,

$$
|T_1| + |T_2| = |T_1 \cup T_2| + |T_1 \cap T_2|
$$

=|N_W(c_i) \cup N_W(c_j)| + |N_W(c_i) \cap N_W(c_j)|
=|N_W(c_i)| + |N_W(c_j)|.

For the second part, since the pairs in P that are not incident to c_i or c_j are kept in P', we only need to show that edges of the form $e = r_{\ell} c_i s_k$ or $e = r_{\ell} c_j s_k$ are still covered by P' .

Case 1: $e = r_{\ell} c_i s_k$.

If $r_{\ell} s_k$ or $c_i s_k$ is in P, then they are also in P' and then e is covered by P'. So, we may assume that $r_{\ell}c_i \in P$. If $r_{\ell}c_i \notin P'$, then by the definition of $P', r_{\ell} \notin N_W(c_j) \cup L_j$. The fact that $r_{\ell} \notin L_j$ means that $j \leq a_{\ell}$. Hence $r_{\ell}c_{j} s_{k} \in H(Y)$ $(k \leq a_{\ell}$ follows from $e \in H(Y)$). The fact that $r_{\ell} \notin N_{W}(c_{j})$ means that $r_{\ell}c_{j}s_{k}$ is covered by P via $r_{\ell} s_k$ or $c_j s_k$. If $r_{\ell} s_k \in P$, then it is kept in P'; If $c_j s_k \in P$, then by definition $c_i s_k \in P'$. In both cases, $e = r_{\ell} c_i s_k$ is covered by P' and we are done.

Case II: $e = r_{\ell} c_i s_k$.

If e is covered by $r_\ell s_k$ or by $r_\ell c_j$ in P, then since these pairs are kept in P', e is covered in P'. Thus we may assume that $c_j s_k \in P$. On the other hand, the edge $r_{\ell} c_i s_k$ of $H(Y)$ is covered by P. If $r_{\ell} s_k \in P$, then $r_{\ell} s_k \in P'$ and e is covered by P'. If $c_i s_k \in P$, then $s_k \in N_Q(c_i) \cap N_Q(c_j)$, therefore by definition $c_j s_k \in P'$ and e is covered by P'. If $r_{\ell}c_i \in P$, since $e \in H(Y)$, then $r_{\ell} \notin L_j$ ($r_{\ell}c_js_k \in H(Y)$ means that $a_{\ell} \geq j$). Therefore $r_{\ell} \in N_W(c_i) \setminus L_j$ so that by definition $r_{\ell}c_j \in P'$ and e is covered by P' . \Box

By interchanging the roles of C and S, we can also define the shifting operation $\phi_{s_i,s_j}(P)$ for $s_i,s_j \in S$ with $i < j$, which satisfies similar property as Lemma [4.1.](#page-5-2)

Definition 4.2. A set $\Gamma \subseteq C \times S$ is called closed down (in $C \times S$) if $c_p s_q \in \Gamma$ implies $[p] \times [q] \subseteq \Gamma$.

Lemma 4.3. For every set $\Gamma \subseteq C \times S$ there exists a closed down Γ' that can be obtained from Γ by a finite sequence of shifting operations.

Proof. We first show that after a finite sequence of shifting operations, we reach at a set that is stable under shifting. And then we prove that a stable set is close down.

Claim 4.3.1. After finitely many shifting operations on Γ , we can reach $\Gamma' \subseteq [a_1] \times [a_1]$ such that $\phi_{c_i,c_j}(\Gamma') =$ $\phi_{s_i,s_j}(\Gamma') = \Gamma'$ for all $1 \leq i < j \leq a_1$.

Proof of the claim. Set $f(\Gamma') = \sum_{i=1}^{a_1} i(|N_{\Gamma'}(c_i)|+|N_{\Gamma'}(s_i)|)$. If ϕ_{c_i,c_j} or ϕ_{s_i,s_j} shifts Γ' to Γ'' that is different from Γ' , then $f(\Gamma'') < f(\Gamma')$. But f is non-negative, which means after finitely many shifting operations, the resulting set becomes stable, which completes the proof of the claim. \Box

Claim 4.3.2. If $\phi_{c_i,c_j}(\Gamma') = \phi_{s_i,s_j}(\Gamma') = \Gamma'$ for all $1 \leq i < j \leq a_1$, then Γ' is closed down.

Proof of the claim. Assume Γ' satisfies the shifting stable property. If $c_p s_q \in \Gamma'$, then for all $p' \leq p$, $q' \leq q$, we have $c_{p'}s_{q'} \in \Gamma'$: If $p' < p$, the shifting-stability applied to $c_p, c_{p'}$ yields $c_{p'}s_q \in \Gamma'$. If $q' < q$, then another application of the shifting-stability of s_q , $s_{q'}$ yields $c_{p'}s_{q'} \in \Gamma'$. \Box

Combining the claims, we prove that after finitely many shifting operations on Γ, we can get a closed down Γ′ . \Box

Combining Lemma [4.1](#page-5-2) and Lemma [4.3,](#page-6-0) we have the following:

Lemma 4.4. For any 2-cover P of $H(Y)$, there exists a 2-cover P' of $H(Y)$ such that $|P'| = |P|$ and $P' \cap (C \times S)$ is closed down.

4.2 More facts about $|Q|$ and $\nu(\ell, Q)$

Lemma [2.9](#page-3-2) provides the values of $\nu(\ell, Q)$ when $Q = [p] \times [q]$. In this section, we study $\nu(\ell, Q)$ and $|Q|$ for general closed down Q. Together with Corollary [2.8,](#page-3-5) this will be the key in the proof of Theorem [1.8.](#page-2-2)

Throughout this section $Q \subseteq C \times S$ and is assumed to be closed down. Let $p \geq 0$ be the maximum integer such that $[p] \times [p] \subseteq Q$. This is well-defined as $|Q|$ is finite.

Observation 4.5.

(1)
$$
\nu(\ell, Q) = 0
$$
 for $\ell \leq p$.

(2) $\nu(\ell, Q) - \nu(\ell - 1, Q) \in \{1, 2\}$ for every $\ell > p$.

Proof. The first part is obvious by the definition of p. For the second part, for $\ell > p$, by maximality of p and the closed down property of Q, we have $c_{\ell} s_{\ell} \notin Q$. Then $\nu(\ell, Q) - \nu(\ell - 1, Q) \geq 1$ is because for every matching in $|\ell-1| \times |\ell-1|$ avoiding Q, we can add the edge $c_{\ell} s_{\ell}$ to obtain a matching in $|\ell| \times |\ell|$ avoiding $Q. \nu(\ell, Q) - \nu(\ell - 1, Q) \leq 2$, since adding two vertices to a bipartite graph cannot increase $\tau = \nu$ by more than 2. 口

Let $q \ge 0$ be the minimum integer such that $\nu(\ell, Q) = \ell$ for all $\ell \ge p + q$. This is well-defined since it is easy to check for all $\ell \geq 2|Q|$, $\nu(\ell,Q) = \ell$. (See [\[1,](#page-16-3) Example 1.6 (4)] for a stronger bound.) In particular, we have $q \ge p$, since when $p \ge 1$, for any $p \le \ell < p+p$, by Observation [4.5,](#page-6-1) $\nu(\ell, Q) \le 2(\ell - p) < \ell$.

By the second part of Observation [4.5,](#page-6-1) we can divide the interval $(p, p+q]$ into (left-open and right-closed with integer endpoints) sub-intervals $O_0, T_1, O_1, T_2, \ldots, O_{k-1}, T_k$ in order such that

$$
\nu(\ell, Q) - \nu(\ell - 1, Q) = \begin{cases} 1, & \text{if } \ell \in O_j \text{ for some } 0 \le j \le k - 1, \\ 2, & \text{if } \ell \in T_j \text{ for some } 1 \le j \le k. \end{cases}
$$

Note that all these sub-intervals, except possibly O_0 , are non-empty. Furthermore, by the minimality of q, the last sub-interval must be T_k (rather than some O_i).

Let I_j be the length of O_j for $0 \le j \le k-1$ and II_j be the length of T_j for $1 \le j \le k$.

We can express $\nu(\ell, Q)$ explicitly in terms of $\mathcal{Q} := \{p, q, O_0, T_1, \ldots, O_{k-1}, T_k, I_0, II_1, \ldots, I_{k-1}, II_k\}$ of Q , as follows.

Lemma 4.6. For any closed down $Q \subseteq C \times S$ with Q defined as above, we have

$$
O_j = (p + \sum_{\ell=0}^{j-1} I_\ell + \sum_{t=1}^j II_t, \ p + \sum_{\ell=0}^j I_\ell + \sum_{t=1}^j II_t], \tag{12}
$$

$$
T_j = (p + \sum_{\ell=0}^{j-1} I_\ell + \sum_{t=1}^{j-1} II_t, \ p + \sum_{\ell=0}^{j-1} I_\ell + \sum_{t=1}^{j} II_t]. \tag{13}
$$

And

$$
\nu(\ell, Q) = \begin{cases}\n0, & \text{if } \ell \leq p. \\
\ell - p + \sum_{1 \leq t \leq j} II_t, & \text{if } \ell \in O_j \text{ for some } 0 \leq j \leq k - 1. \\
2\ell - 2p - \sum_{0 \leq t < j} I_t, & \text{if } \ell \in T_j \text{ for some } 1 \leq j \leq k. \\
\ell, & \text{if } \ell > p + q.\n\end{cases} \tag{14}
$$

Proof. The endpoints in [\(12\)](#page-7-0) and [\(13\)](#page-7-1) follow from the definition of the intervals O_i and T_i . The first and last items in (14) follow from the definitions of p and q .

To compute $\nu(\ell, Q)$, when $\ell \in O_j$, let ℓ_0 be the left endpoint of O_j , which is $p + \sum_{\ell=0}^{j-1} I_{\ell} + \sum_{t=1}^{j} II_t$ by [\(12\)](#page-7-0). And $\nu(\ell_0, Q) = \sum_{\ell=0}^{j-1} I_{\ell} + 2 \sum_{t=1}^{j} II_t$. Therefore

$$
\nu(\ell, Q) = (\ell - \ell_0) + \nu(\ell_0, Q)
$$

= $\left(\ell - (p + \sum_{\ell=0}^{j-1} I_{\ell} + \sum_{t=1}^{j} II_{t})\right) + \sum_{\ell=0}^{j-1} I_{\ell} + 2 \sum_{t=1}^{j} II_{t} = \ell - p + \sum_{1 \le t \le j} II_{t}.$

When $\ell \in T_j$, let $\ell_0 = p + \sum_{\ell=0}^{j-1} I_{\ell} + \sum_{t=1}^{j-1} I_{t}$ be the left endpoint of T_j by [\(13\)](#page-7-1). And $\nu(\ell_0, Q) =$ $\sum_{\ell=0}^{j-1} I_{\ell} + 2 \sum_{t=1}^{j-1} II_t$. Therefore

$$
\nu(\ell, Q) = 2(\ell - \ell_0) + \nu(\ell_0, Q) = 2\ell - 2p - \sum_{0 \le t < j} I_t.
$$

Hence we complete the proof.

We have the following results about the sums of II_j and of I_j , and a lower bound on the size of Q with respect to I_{ℓ}, II_j .

Corollary 4.7. For any closed down $Q \subseteq C \times S$ with Q defined as above, we have

$$
\sum_{j=1}^{k} II_j = p, \quad \sum_{j=0}^{k-1} I_j = q - p,\tag{15}
$$

Proof. Note that by the definition of O_j, T_j, I_j, II_j ,

$$
p + \sum_{j=0}^{k-1} I_j + \sum_{j=1}^{k} II_j = p + q = \nu(p + q, Q) = \sum_{j=0}^{k-1} I_j + 2 \sum_{j=1}^{k} II_j,
$$

from which the identities in [\(15\)](#page-8-0) immediately follow.

Lemma 4.8. For any closed down $Q \subseteq C \times S$ with Q defined as above,

$$
|Q| \ge \sum_{j=1}^{k} II_j \left(p + I_0 + \sum_{\ell=1}^{j-1} (II_\ell + I_\ell) \right). \tag{16}
$$

Proof. For $0 \le j \le k-1$ and $\ell \in O_j$, we take a maximum matching $M_{\ell-1}$ in $([\ell-1] \times [\ell-1]) \setminus Q$. The constraint $\nu(\ell,Q)-\nu(\ell-1,Q)=1$ implies that we cannot match both c_{ℓ} and s_{ℓ} to $\{c_1,\ldots,c_{\ell-1}\}\cup\{s_1,\ldots,s_{\ell-1}\}\$ without using edges in Q or intersecting with edges in $M_{\ell-1}$, otherwise we get a matching in $([\ell] \times [\ell]) \setminus Q$ of size $|M_{\ell-1}|+2$. Therefore for one of c_{ℓ} and s_{ℓ} , which we denote by z_{ℓ} , say $z_{\ell}=s_{\ell}$, all the edges $\{c_1, \ldots, c_{\ell-1}\}\times \{s_{\ell}\}\$ are either in Q or incident with edges in $M_{\ell-1}$. Since by [\(14\)](#page-7-2)

$$
|M_{\ell-1}| = \nu(\ell-1, Q) = \nu(\ell, Q) - 1 = \ell - 1 - p + \sum_{1 \le t \le j} II_t,
$$

then

$$
|N_{Q \cap ([\ell-1] \times [\ell-1])}(z_{\ell})| \ge (\ell-1) - |M_{\ell-1}| = p - \sum_{1 \le t \le j} II_t.
$$
\n(17)

For any $1 \le j \le k-1$, let ℓ_j be the largest integer in T_j . For all $\ell' \in T_j$, since $\ell' < \ell_j + 1 \in O_j$, by the closed down property of Q,

$$
N_{Q \cap ([\ell_j] \times [\ell_j])}(z_{\ell_j+1}) \subseteq N_{Q \cap ([\ell_j] \times [\ell_j])}(z_{\ell'}),
$$

where $z_{\ell'}$ is set to be $s_{\ell'}$ if $z_{\ell_j+1} = s_{\ell_j+1}$, and to be $c_{\ell'}$ if $z_{\ell_j+1} = c_{\ell_j+1}$. Therefore

$$
|N_{Q \cap ([\ell_j] \times [\ell_j])}(z_{\ell'})| \ge |N_{Q \cap ([\ell_j] \times [\ell_j])}(z_{\ell_j+1})| \ge p - \sum_{1 \le t \le j} II_t.
$$
\n(18)

Remark 4.9. Since Q is closed down and p is maximal, the neighbors counted in (17) and (18) are the first vertices among the first p vertices on the side opposite to the side of z_{ℓ} and $z_{\ell'}$. Counting the number of edges of Q from that side also leads to the lower bound on $|Q|$ as in Corollary [4.7](#page-8-3) (for example, we can count $|Q|$ from the C-side of the Q constructed in the proof of Corollary [4.10\)](#page-9-0) — a fact not used in the proof.

Hence by the choice of p , combining [\(17\)](#page-8-1) with [\(18\)](#page-8-2), and applying [\(15\)](#page-8-0), we have

$$
|Q| \ge |[p] \times [p] + \sum_{\ell \in O_0} |N_{Q \cap ([\ell-1] \times [\ell-1])}(z_{\ell})| + \sum_{j=1}^{k-1} \Big(\sum_{\ell' \in T_j} |N_{Q \cap ([\ell_j] \times [\ell_j])}(z_{\ell'})| + \sum_{\ell \in O_j} |N_{Q \cap ([\ell-1] \times [\ell-1])}(z_{\ell})|\Big)
$$

\n
$$
\ge p(p+I_0) + \sum_{j=1}^{k-1} (II_j + I_j)(p - \sum_{1 \le t \le j} II_t)
$$

\n
$$
= \sum_{j=1}^k II_j(p+I_0) + \sum_{j=1}^{k-1} (II_j + I_j) \sum_{j < t \le k} II_t
$$

$$
= \sum_{j=1}^{k} II_j(p+I_0) + \sum_{t=2}^{k} II_t \Big(\sum_{j=1}^{t-1} (II_j + I_j) \Big)
$$

=
$$
\sum_{j=1}^{k} II_j \Big(p + I_0 + \sum_{\ell=1}^{j-1} (II_\ell + I_\ell) \Big),
$$

which completes the proof.

The next result will not be used, but it is of independent interest.

Corollary 4.10. For any integers $0 \le p \le q$ and division of the interval $(p, p + q)$ into (left-open and rightclosed with integer endpoints) sub-intervals $O_0, T_1, O_1, T_2, \ldots, O_{k-1}, T_k$ in order such that except possibly O_0 , all the sub-intervals are non-empty, for $I_j = |O_j|$ and $II_j = |T_j|$, there exists a closed down $Q \subseteq C \times S$ (assuming $|C| = |S| > p + q$) satisfying [\(12\)](#page-7-0)–[\(15\)](#page-8-0) and attaining the lower bound in [\(16\)](#page-8-4), i.e.,

$$
|Q| = \sum_{j=1}^{k} II_j \left(p + I_0 + \sum_{\ell=1}^{j-1} (II_{\ell} + I_{\ell}) \right).
$$

Proof. We set

$$
Q := \bigcup_{j=1}^k \left(\{c_{p-\sum_{\ell=1}^j I I_\ell + 1}, \ldots, c_{p-\sum_{\ell=1}^{j-1} I I_\ell}\} \times \left(\{s_\ell : \ell \in [p] \cup O_0 \} \cup \bigcup_{\ell=1}^{j-1} \{s_\ell : \ell \in T_\ell \cup O_\ell\} \right) \right).
$$

It is routine to verify Q satisfies all the requirements.

4.3 Proof of Theorem [1.8](#page-2-2)

We are now set to prove Theorem [1.8.](#page-2-2) We prove the contrapositive statement.

Theorem 4.11. If $\tau^{(2)}(H(Y)) < |Y|$, then the Young diagram Y is not wide.

4.3.1 A toy example

By Lemma [4.4,](#page-6-2) there exists a minimum 2-cover P of $H(Y)$ satisfying $P \cap (C \times S)$ is closed down and $|P| = \tau^{(2)}(H(Y)) < |Y|$. Furthermore, by Corollary [2.8](#page-3-5) and the assumption $|P| < |Y|$, $Q = P \cap (C \times S)$ is non-empty. In a toy case, we assume that $Q = [p] \times [q]$ for some $1 \leq p \leq q$. For this Q, we have $O_0 = (p, q]$, $T_1 = (q, p + q], I_0 = q - p$, and $II_1 = p$. Let Z be the partition formed by the rows of Y whose lengths are at most $p + q$. We shall show that the sum of the lengths of the first $II_1 = p$ rows of Z is less than that of the first p columns of Z . Therefore Z does not dominate Z' , hence Y is not wide.

Indeed, let

$$
D = \{i : a_i \le p\}, \ A_{O_0} = \{i : a_i \in O_0\}, \ A_{T_1} = \{i : a_i \in T_1\}, \text{ and } U = \{i : a_i > p + q\}.
$$

Let $A_{T_1}[I_1]$ be the set of the first $II_1 = p$ elements of A_{T_1} if $II_1 \leq |A_{T_1}|$, and to be A_{T_1} if $II_1 > |A_{T_1}|$.

The first observation is

the sum of the lengths of the first
$$
II_1
$$
 rows of $Z \leq pq + \sum_{i \in A_{T_1}[II_1]} (a_i - q).$ (19)

To see this, note that if $|A_{T_1}| \geq II_1 = p$, then the RHS is equal to the LHS, which is $II_1 \cdot q + \sum_{i \in A_{T_1}[II_1]} (a_i - a_i)$ $q) = \sum_{i \in A_{T_1}[II_1]} a_i$. If $|A_{T_1}| < II_1$, then $|A_{T_1}|q + \sum_{i \in A_{T_1}[II_1]} (a_i - q)$ is the sum of the lengths of the first $|A_{T_1}|$ rows of Z, and the sum of the lengths of the remaining $II_1 - |A_{T_1}|$ rows is at most $(p - |A_{T_1}|)q$, because each of these remaining rows is below those with indices in A_{T_1} and has length at most q.

By Corollary [2.8,](#page-3-5) Lemma [2.9,](#page-3-2) and the assumption $|P| = \tau^{(2)}(H(Y)) < |Y|$, we have

$$
pq + \sum_{i \in A_{O_0}} (a_i - p) + \sum_{i \in A_{T_1}[II_1]} (2a_i - p - q) + \sum_{i \in A_{T_1} \setminus A_{T_1}[II_1]} (2a_i - p - q) + \sum_{i \in U} a_i < \sum_{i \in D \cup A_{O_0} \cup A_{T_1} \cup U} a_i. \tag{20}
$$

Note that

LHS of
$$
(20)
$$

$$
= \sum_{i \in A_{O_0}} a_i + \sum_{i \in A_{T_1}} a_i + pq + \sum_{i \in A_{T_1}[II_1]} (a_i - q) + \sum_{i \in A_{T_1} \setminus A_{T_1}[II_1]} (a_i - q) + \sum_{i \in U} a_i - p(|A_{O_0}| + |A_{T_1}|)
$$

 \geq the sum of the lengths of the first II_1 rows of $Z + \sum$ $i∈A_{O_0} ∪ A_{T_1} ∪ U$ $a_i - p(|A_{O_0}| + |A_{T_1}|)$ (21)

where the last inequality is by [\(19\)](#page-9-2) and the fact that for each $i \in A_{T_1} \setminus A_{T_1}[I_1], a_i \ge q$. Then comparing [\(21\)](#page-10-0) with the RHS of [\(20\)](#page-9-1), after cancellation, we have

the sum of the lengths of the first
$$
II_1
$$
 rows of $Z < \sum_{i \in D} a_i + p(|A_{O_0}| + |A_{T_1}|),$

where the RHS is the sum of the lengths of the first $II_1 = p$ columns of Z, since the rows with indices in $A_{O_0} \cup A_{T_1}$ have lengths greater than p. We are done.

4.3.2 Proof of Theorem [4.11](#page-9-3)

Proof of Theorem [4.11.](#page-9-3) By Lemma [4.4,](#page-6-2) there exists a minimum 2-cover P of $H(Y)$ satisfying $P \cap (C \times S)$ is closed down and $|P| = \tau^{(2)}(H(Y)) < |Y|$. Let $Q = P \cap (C \times S)$. By Corollary [2.8](#page-3-5) and the assumption $|P| < |Y|$, we have $Q \neq \emptyset$. By Corollary [2.8](#page-3-5) and Lemma [4.8,](#page-8-5) and using notations there, where the nonemptiness of Q implies $1 \le p \le q$, we have

$$
f(Q) := \sum_{j=1}^{k} II_j \left(p + I_0 + \sum_{\ell=1}^{j-1} (II_\ell + I_\ell) \right) + \sum_i \nu(a_i, Q)
$$

$$
\leq |Q| + \sum_i \nu(a_i, Q) \leq |P| < \sum_i a_i,
$$
 (22)

where, as above, a_i is the length of the *i*th row of Y.

Using the notation in Lemma [4.6,](#page-7-3) let

$$
D = \{i : a_i \le p\},
$$

\n
$$
A_{O_j} = \{i : a_i \in O_j\} \text{ for } 0 \le j \le k - 1,
$$

\n
$$
A_{T_j} = \{i : a_i \in T_j\} \text{ for } 1 \le j \le k,
$$

\n
$$
U = \{i : a_i > p + q\}.
$$

For $1 \leq j \leq k$, let Z_j be the subpartition of Y that consists of the rows with index in $D \cup_{t=0}^{j-1} A_{O_t} \cup_{\ell=1}^{j} A_{T_{\ell}}$. We define $A_{T_j}[I I_j]$ to be the set of the first II_j elements of A_{T_j} if $II_j \leq |A_{T_j}|$, and to be A_{T_j} if $II_j > |A_{T_j}|$.

We aim to show that the sum of the lengths of the first II_k rows of Z_k is less than that of the first II_k columns of Z_k , which implies that Z_k does not dominate Z'_k and then Y is not wide.

(Note that in the toy example in Section [4.3.1,](#page-9-4) $k = 1$ and we set $Z = Z_k = Z_1$.)

As before, the first observation is an upper bound on the sum of the lengths of the first II_i rows of Z_i .

Claim 4.11.1. For every $1 \leq j \leq k$, the sum of the lengths of the first II_j rows of Z_j is at most

$$
h_j := II_j\Big(p + I_0 + \sum_{\ell=1}^{j-1} (II_\ell + I_\ell)\Big) + \sum_{i \in A_{T_j}[II_j]} (a_i - 2p - \sum_{0 \le t < j} I_t + \sum_{\ell=j}^k II_\ell).
$$

Note that the first item is part of $f(Q)$ in [\(22\)](#page-10-1).

Proof of Claim [4.11.1.](#page-10-2) To verify this, first observe that since $p - \sum_{\ell=j}^{k} II_{\ell} = \sum_{\ell=1}^{j-1} II_{\ell}$ by [\(15\)](#page-8-0), then the term in the second summand of h_j satisfies

$$
a_i - 2p - \sum_{0 \le t < j} I_t + \sum_{\ell=j}^k II_\ell
$$

$$
= a_i - p - \sum_{0 \le t < j} I_t - (p - \sum_{\ell=j}^k II_\ell)
$$
\n
$$
= a_i - p - \sum_{0 \le t < j} I_t - \sum_{\ell=1}^{j-1} II_\ell = a_i - \left(p + I_0 + \sum_{\ell=1}^{j-1} (II_\ell + I_\ell) \right).
$$

Note that $w_j := p + I_0 + \sum_{\ell=1}^{j-1} (II_\ell + I_\ell)$ appears in the first part of the definition of h_j .

If $II_j \leq |A_{T_j}|$, then $h_j = II_j w_j + \sum_{i \in A_{T_j}[II_j]} (a_i - w_j) = \sum_{i \in A_{T_j}[II_j]} a_i$, which is the sum of the lengths of the first II_j rows of $\mathbb{Z}_j.$

If $II_j > |A_{T_j}|$, then

$$
h_j = |A_{T_j}| \cdot w_j + \sum_{i \in A_{T_j}} (a_i - w_j) + (II_j - |A_{T_j}|) \cdot w_j
$$

= $\left(\sum_{i \in A_{T_j}} a_i\right) + (II_j - |A_{T_j}|) \cdot w_j.$

Since the $|A_{T_j}|+1,\ldots, I I_j$ th rows of Z_j are below the rows with indices in A_{T_j} , by [\(13\)](#page-7-1) in Lemma [4.6,](#page-7-3) each of these $II_j - |A_{T_j}|$ rows has size at most $w_j = p + I_0 + \sum_{\ell=1}^{j-1} (II_\ell + I_\ell)$. Therefore in this case, the sum of the lengths of the first II_j rows of Z_j is at most h_j .

Together with [\(14\)](#page-7-2) in Lemma [4.6,](#page-7-3) we can rewrite the representation of $f(Q)$ in [\(22\)](#page-10-1) as follows.

$$
f(Q) = \sum_{j=1}^{k} II_j (p + I_0 + \sum_{\ell=1}^{j-1} (II_{\ell} + I_{\ell}))
$$

+
$$
\sum_{j=0}^{k-1} \sum_{i \in A_{O_j}} (a_i - p + \sum_{1 \leq t \leq j} II_t) + \sum_{j=1}^{k} \sum_{i \in A_{T_j}} (2a_i - 2p - \sum_{0 \leq t < j} I_t) + \sum_{i \in U} a_i
$$

=
$$
\sum_{j=1}^{k} II_j (p + I_0 + \sum_{\ell=1}^{j-1} (II_{\ell} + I_{\ell}))
$$

+
$$
\sum_{j=0}^{k-1} \sum_{i \in A_{O_j}} (a_i - p + \sum_{1 \leq t \leq j} II_t) + \sum_{j=1}^{k} \sum_{i \in A_{T_j}} a_i
$$

+
$$
\sum_{j=1}^{k} \Big(\sum_{i \in A_{T_j} [II_t]} + \sum_{i \in A_{T_j} \setminus A_{T_j} [II_j]} (a_i - 2p - \sum_{0 \leq t < j} I_t + \sum_{\ell=j}^{k} II_{\ell})
$$

-
$$
\sum_{j=1}^{k} |A_{T_j}| \sum_{\ell=j}^{k} II_{\ell} + \sum_{i \in U} a_i
$$

=
$$
\sum_{j=1}^{k} h_j + \sum_{j=0}^{k-1} \sum_{i \in A_{O_j}} (a_i - (p - \sum_{1 \leq t \leq j} II_t) + \sum_{j=1}^{k} \sum_{i \in A_{T_j}} a_i
$$

+
$$
\sum_{j=1}^{k} \sum_{i \in A_{T_j} \setminus A_{T_j} [II_j]} (a_i - p - \sum_{0 \leq t < j} I_t - (p - \sum_{\ell=j}^{k} II_{\ell}))
$$

-
$$
\sum_{j=1}^{k} |A_{T_j}| \sum_{\ell=j}^{k} II_{\ell} + \sum_{i \in U} a_i
$$

=
$$
\sum_{j=1}^{k} h_j + \sum_{i \in Z_k \cup U \setminus D} a_i - \sum_{j=1}^{k} |A_{T_j}| \sum_{\ell=j}^{k} II_{\ell} - \sum_{j=0}^{k
$$

$$
+\sum_{j=1}^k \sum_{i \in A_{T_j} \setminus A_{T_j}[II_j]} (a_i - p - \sum_{0 \le t < j} I_t - \sum_{\ell=1}^{j-1} II_\ell).
$$

For the last term, for $i \in A_{T_j}$, by [\(13\)](#page-7-1) we have $a_i - p - \sum_{0 \leq t < j} I_t - \sum_{\ell=1}^{j-1} II_j \geq 0$. Therefore

$$
f(Q) \ge \sum_{j=1}^{k} h_j + \sum_{i \in Y_k \cup U \setminus D} a_i - \sum_{j=1}^{k} |A_{T_j}| \sum_{\ell=j}^{k} II_{\ell} - \sum_{j=0}^{k-1} |A_{O_j}| \sum_{\ell=j+1}^{k} II_{\ell}.
$$
 (23)

On the other hand, by [\(22\)](#page-10-1) we have

$$
f(Q) < \sum_{i \in Z_k \cup U} a_i. \tag{24}
$$

Combining [\(23\)](#page-12-0) with [\(24\)](#page-12-1), we have

$$
\sum_{j=1}^{k} h_j < \sum_{i \in D} a_i + \sum_{j=1}^{k} |A_{T_j}| \sum_{\ell=j}^{k} II_{\ell} + \sum_{j=0}^{k-1} |A_{O_j}| \sum_{\ell=j+1}^{k} II_{\ell}
$$
\n
$$
= \sum_{i \in D} a_i + \sum_{j=1}^{k} II_j \left(\sum_{1 \le t \le j} |A_{T_t}| + \sum_{0 \le \ell < j} |A_{O_\ell}| \right). \tag{25}
$$

We may assume that for each $1 \leq j \leq k-1$, the sum of the lengths of the first II_j rows of Z_j is at least that of the first II_j columns of Z_j , which is

$$
II_j(\sum_{1 \leq t \leq j} |A_{T_t}| + \sum_{0 \leq \ell < j} |A_{O_\ell}|) + \sum_{i: \ a_i \leq II_j} a_i + \sum_{i: \ a_i > II_j, a_i \in D} II_j,
$$

otherwise Z_j does not dominate Z'_j and we already prove Y is not wide. (Note that the first two summands arise from the fact that the lengths of the corresponding rows are greater than p and $p \geq II_j$ by [\(15\)](#page-8-0).) Hence by Claim [4.11.1,](#page-10-2) we have for every $1 \leq j \leq k - 1$,

$$
h_j \ge II_j \left(\sum_{1 \le t \le j} |A_{T_t}| + \sum_{0 \le \ell < j} |A_{O_\ell}| \right) + \sum_{i: \ a_i \le II_j} a_i + \sum_{i: \ a_i > II_j, a_i \in D} II_j. \tag{26}
$$

Combining [\(25\)](#page-12-2) with [\(26\)](#page-12-3) and subtracting $\sum_{j=1}^{k-1} h_j$ from $\sum_{j=1}^{k} h_j$, we have

$$
h_k < II_k(\sum_{t=1}^k |A_{T_t}| + \sum_{\ell=0}^{k-1} |A_{O_\ell}|) + \sum_{i \in D} a_i - \sum_{j=1}^{k-1} \sum_{i: \ a_i \le II_j} a_i + \sum_{i: \ a_i > II_j, a_i \in D} II_j). \tag{27}
$$

If we can show

$$
\sum_{i \in D} a_i - \sum_{j=1}^{k-1} (\sum_{i: a_i \le II_j} a_i + \sum_{i: a_i > II_j, a_i \in D} II_j) \le \sum_{i: a_i \le II_k} a_i + \sum_{i: a_i > II_k, a_i \in D} II_k, \tag{28}
$$

then by Claim [4.11.1,](#page-10-2) we have the sum of the lengths of the first II_k rows of Z_k is at most

$$
h_k < II_k(\sum_{t=1}^k |A_{T_t}| + \sum_{\ell=0}^{k-1} |A_{O_\ell}|) + \sum_{i: \ a_i \le II_k} a_i + \sum_{i: \ a_i > II_k, a_i \in D} II_k,
$$

where the right-hand side is the sum of the lengths of the first II_k columns of Z_k and we complete the proof.

To see that [\(28\)](#page-12-4) is true, we consider an equivalent form

$$
\sum_{i \in D} a_i \le \sum_{j=1}^k \sum_{i: \ a_i \le II_j} a_i + \sum_{j=1}^k II_j \sum_{i: \ a_i > II_j, a_i \in D} 1.
$$
 (29)

Indeed, for each a_i counted by the left-hand side, if $a_i \leq \max(I_1, \ldots, I_k)$, then it appears at least once in the first summand on the right-hand side. If $p \ge a_i > \max(I_1, \ldots, I_k)$, then it is counted k times in the second summand of the right-hand side, which by [\(15\)](#page-8-0) contribute $\sum_{j=1}^{k} II_j = p \ge a_i$ to the right-hand side. Therefore we prove [\(29\)](#page-12-5) and then [\(28\)](#page-12-4), which completes the proof of the theorem.

5 Fractional versions

An intermediate version between Theorem [1.8](#page-2-2) and Conjecture [1.7](#page-2-0) is a fractional version of the conjecture.

A fractional 2-matching in a hypergraph H is a fractional matching of $H^{(2)}$, i.e., a function $y : E(H) \rightarrow$ $\mathbb{R}_{\geq 0}$ subject to the constraint

$$
\sum_{e:\ p \subseteq e} y(e) \le 1 \text{ for every } p \in V(H^{(2)}), \text{ i.e., for every } p \in \binom{e}{2} \text{ with } e \in E(H).
$$

A fractional 2-cover of a hypergraph H is a fractional cover of $H^{(2)}$, i.e., a function $f: V(H^{(2)}) \to \mathbb{R}_{\geq 0}$ subject to the constraint

$$
\sum_{p \in {e \choose 2}} f(p) \ge 1
$$
 for every $e \in E(H)$.

We denote by $\nu^{(2)*}(H)$ the maximal value $\sum_{e \in E(H)} y(e)$ over all fractional 2-matchings y in H, and by $\tau^{(2)*}(H)$ the minimal value $\sum_{p\in V(H^{(2)})} f(p)$ over all fractional 2-covers f of H. Again we have $\nu^{(2)*}(H)$ = $\nu^*(H^{(2)})$ and $\tau^{(2)*}(H) = \tau^*(H^{(2)})$. By the definition and LP duality,

$$
\nu^{(2)}(H) \le \nu^{(2)*}(H) = \tau^{(2)*}(H) \le \tau^{(2)}(H).
$$

Definition 5.1. A Young diagram Y of size n is said to be fractionally Latin if $\nu^{(2)*}(H(Y)) = n$.

Observation 5.2.

- 1. By the definition of n, there are n pairs in $H[R \times C]$.
- 2. In each row i there are a_i symbols, so there are n pairs in $H[R \times S]$.
- 3. $|H[C \times S]| = (a_1)^2$ can be larger than n.

Corollary 5.3. Let y be a fractional 2-matching in $H(Y)$. Then $\sum_{e \in E(H)} y(e) = n$ if and only if $\sum_{e: p \subseteq e} y(e) =$ 1 for every pair $p \in H[R \times Z]$, where Z is C or S.

Proof. We have

$$
\sum_{e \in E(H)} y(e) \le \sum_{p \in H[R \times Z]} \sum_{e: p \subseteq e} f(e) \le \sum_{p \in H[R \times Z]} 1 = n,
$$

where the equality is by [\(1\)](#page-13-1) and [\(2\)](#page-13-2) in Observation [5.2.](#page-13-3) Therefore $\sum_{e \in E(H)} y(e) = n$ if and only if $\sum_{e: p \subseteq e} y(e) = 1$ for every pair $p \in H[R \times Z].$ □

Conjecture 5.4. If a Young diagram Y is wide, then $\nu^{(2)*}(H(Y)) = |Y|$.

The converse, generalizing Theorem [1.3,](#page-1-3) is true.

Theorem 5.5. If Z is a fractionally Latin Young diagram, i.e., $\nu^{(2)*}(H(Z)) = |Z|$, then Z is wide.

This follows from Theorem [1.9](#page-2-1) and the fact that $\nu^{(2)*}(H) = \tau^{(2)*}(H) \leq \tau^{(2)}(H)$. The direct proof below follows the footsteps of the proof of Theorem [1.3,](#page-1-3) but is still worth noting.

Proof of Theorem [5.5.](#page-13-4) Suppose not. Let Y be a subpartition formed by a subset of rows of Z that does not dominate Y'. Then we have $\nu^{(2)*}(H(Y)) = |Y|$, otherwise an optimal fractional 2-cover of Y together with putting weight 1 on $r_i c_j$ for $(i, j) \in Y \setminus Z$ forms a fractional 2-cover of Z with weight $\nu^{(2)*}(H(Y)) + |Z| - |Y|$, which is less than $|Z|$, a contradiction.

Assume the sum of the lengths of the first k rows of Y is less than that of the first k columns of Y, i.e.,

$$
\sum_{1 \le i \le k} a_i < \sum_{1 \le i \le k} b_i. \tag{30}
$$

Let y be a fractional 2-matching of Y of size $|Y| = n$. We abbreviate $y(i, j, \ell)$ for $y(r_i c_i s_\ell)$ and without loss of generality, we set $y(i, j, \ell) = 0$ if $r_i c_j s_\ell \notin H(Y)$. Let

$$
T_k := \sum_{1 \le i \le m, 1 \le j \le a_i, 1 \le \ell \le k} y(i, j, \ell).
$$

Claim 5.5.1. We have

- 1. $\sum_{1 \leq i \leq k} b_i = T_k$.
- 2. $\sum_{1 \leq i \leq k} a_i \geq T_k$.

Remark 5.6. The source of the difference, why the first part is an equality and the second part is only an inequality, is in the difference between $(1)-(2)$ $(1)-(2)$ in Observation [5.2](#page-13-3) and [\(3\)](#page-13-5) of that observation.

Proof of Claim [5.5.1.](#page-14-1) To prove the first part, let i be fixed. By Corollary [5.3](#page-13-6) for every $1 \leq \ell \leq \min(k, a_i)$ we have $\sum_{1 \leq j \leq a_i} y(i, j, \ell) = 1$. And for $\ell > a_i$, we have $\sum_{1 \leq j \leq a_i} y(i, j, \ell) = 0$ since no edge of H contains such $r_i s_\ell$. Therefore $\sum_{1 \leq j \leq a_i, 1 \leq \ell \leq k} y(i, j, \ell) = \min(k, a_i)$. Hence

$$
T_k = \sum_{1 \le i \le m} \min(k, a_i) = \sum_{1 \le j \le k} b_j,
$$

where the last equality is by [\(2\)](#page-0-0).

To prove the second part, fix j. For each $1 \leq \ell \leq k$, we have $\sum_{1 \leq i \leq m} y(i, j, \ell) \leq 1$, hence

$$
\sum_{1\leq i\leq m, 1\leq \ell\leq k}y(i,j,\ell)\leq k.
$$

Similarly, for any $1 \leq i \leq b_j$, we have $\sum_{1 \leq \ell \leq k} y(i, j, \ell) \leq 1$. Since there is no edge of the form $r_i c_j s_\ell$ for $b_j < i \leq m$, we have

$$
\sum_{1 \le i \le m, 1 \le \ell \le k} y(i, j, \ell) = \sum_{1 \le i \le b_j, 1 \le \ell \le k} y(i, j, \ell) \le b_j.
$$

Therefore

$$
\sum_{1 \le i \le m, 1 \le \ell \le k} y(i, j, \ell) \le \min(k, b_j).
$$

Summing over all $1 \leq j \leq a_i$ yields

$$
T_k \le \sum_{1 \le j \le a_1} \min(k, b_j) = \sum_{1 \le i \le k} a_i,
$$

where the last equality is by [\(2\)](#page-0-0).

By Claim [5.5.1](#page-14-1) $\sum_{1 \leq i \leq k} a_i \geq T_k = \sum_{1 \leq i \leq k} b_i$, a contradiction to [\(30\)](#page-13-7).

 \Box \Box

 6 $\tau^{(2)}$ vs. $\nu^{(2)}$ and some stronger questions

Observation 6.1. $\tau(H(Y)) = \nu(H(Y)).$

Proof. Let $G = H[R \times C]$ (as defined in Section [2\)](#page-2-4). We claim that $\tau(H(Y)) = \nu(H(Y)) = \tau(G)$. Clearly, $\tau(H(Y)) \leq \tau(G)$, and since $\nu(H(Y)) \leq \tau(H(Y))$, by König's theorem it suffices to show that $\nu(H(Y)) =$ $\nu(G)$. As noted above, $\nu(H(Y)) \leq \nu(G)$. To show the converse, for any matching M in G, let $\tilde{M} = \{rc_j s_j \mid s_j \in G\}$ $rc_j \in M$, where $C = \{c_1, \ldots, c_{a_1}\}\$ and $S = \{s_1, \ldots, s_{a_1}\}\$. Then, by the definition of $H(Y)$, \tilde{M} is a matching in $H(Y)$, and $|M| = |M|$. 口

If the wide diagram conjecture is true, then $\tau^{(2)}(H(Y)) = \nu^{(2)}(H(Y))$ whenever Y is wide. We do not know an example, wide or not, in which this integral duality relation fails. We post it as a question.

Question 6.2. Is it true that $\tau^{(2)}(H(Y)) = \nu^{(2)}(H(Y))$ for every Young diagram Y?

Based on Theorem [1.8,](#page-2-2) solving Conjecture [1.4](#page-1-2) is equivalent to solving this question, restricted to wide diagrams.

We dare go even further. For bipartite graphs G_1, \ldots, G_m sharing the same bipartition define the following hypergraph:

$$
H = H(G_1, ..., G_m) = \{ixy \mid xy \in G_i, 1 \le i \le m\}.
$$

Question 6.3. Is it true that if $G_1 \supseteq G_2 \supseteq \ldots \supseteq G_m$, then $\tau^{(2)}(H) = \nu^{(2)}(H)$?

Question [6.2](#page-14-2) is the case in which $G_i = [a_i] \times [a_i]$ for some number a_i , for every i.

Theorem 6.4. If all G_i are the same graph G , then the answer to Question [6.3](#page-15-0) is positive.

Proof. Recall that $\nu^{(2)}(H)$ is the maximal size of the union of m disjoint matchings in G. By König's edge-coloring theorem, this is the maximum of $|E(F)|$, over all subgraphs F of G with $\Delta(F) \leq m$. This is the integral object of an LP program which is the maximal value of $\sum_{e \in E(G)} f(e)$ over all functions $f: E(G) \to \mathbb{R}_{\geq 0}$ satisfying $f(e) \leq 1$ for every edge e of G and $\sum_{e:v \in e} f(e) \leq m$ for every vertex v of G. The polytope defined by these inequalities is obtained from the matching polytope of the bipartite graph G , by adding constraints of the form $f(e) \leq 1$ for all the $e \in E(G)$. The matrix defining the polytope is well-known to be totally unimodular, so the polytope is integral. Hence there exists an integral solution.

In the dual program there is a variable x_v for every $v \in V(G)$ (corresponding to the degree constraint at v in the primal) and a variable y_e for every $e \in E(G)$ (corresponding to the constraint $f(e) \leq 1$), satisfying the conditions $x_v, y_e \ge 0$ and $x_u + x_v + y_e \ge 1$ for every edge $e = uv$. The object function of the dual is the minimum of $\sum_{v \in V(G)} mx_v + \sum_{e \in E(G)} y_e$. By the duality theorem, it follows that $\nu^{(2)}(H)$ is the minimal value of $\sum_{v\in V(G)} mx_v + \sum_{e\in E(G)} y_e$. By minimality, we may assume that in a solution $y_e \leq 1$ for every $e \in E(G)$.

The dual, too, has an integral solution. Taking X to be the set of vertices v with $x_v = 1$ in a solution of the dual, by the minimality of the object function, $y_e = 1$ precisely for those edges e that are not covered by X, hence the value of the dual is the minimum, taken over all $X \subseteq V(G)$, of $m|X| + |E(G - X)|$.

Let $X \subseteq V(G)$ attain the minimum and let $Q = E(G - X)$. As in Corollary [2.5](#page-3-4) we have $\tau^{(2)}(H) \leq$ $|Q| + \sum_{i=1}^{m} \tau(G - Q) \leq |Q| + m|X| = \nu^{(2)}(H)$. Since $\tau^{(2)}(H) \geq \nu^{(2)}(H)$, equality holds. \Box

Question [6.3](#page-15-0) has a nice reformulation that relates to perfect graphs. For a graph $G = (V, E)$ and a chain of sets $V_m \subseteq V_{m-1} \subseteq \ldots \subseteq V_2 \subseteq V_1 \subseteq V$ for some positive integer m, we obtain a new graph $Lev(G; V_1, \ldots, V_m)$ by taking vertex-disjoint copies of $G[V_1], \ldots, G[V_m]$, and then for every $v \in V_1$ we add a clique on the set of copies of v. We say that $Lev(G; V_1, \ldots, V_m)$ is an *m-level* graph of G.

We say that G is m-perfect if for every $V_m \subseteq V_{m-1} \subseteq \ldots \subseteq V_2 \subseteq V_1 \subseteq V(G)$, the m-level graph L of G satisfies

$$
\alpha(L) = \kappa(L),
$$

where α is the independence number and κ is the coloring number of the complement. So clearly (up to the Weak Perfect Graph Theorem) a graph is 1-perfect if and only if it is perfect (according to the original definition of Berge).

We say that a graph is ∞ -perfect if it is m-perfect for every m. For example, a complete graph is ∞ perfect, because every level graph of the complete graph is the line graph of some bipartite graph, which is known to be perfect. To see the isomorphism between the m-level graph L and the line graph of some bipartite graph F, assume the vertex set of L is $\cup_{i=1}^m \{v_{i,1},\ldots,v_{i,p_i}\}$, where $\{v_{i,1},\ldots,v_{i,p_i}\}$ is the vertex set of the copy of $G[V_i]$ and $\{v_{1,j},\ldots,v_{q_j,j}\}\$ is the collection of the copies of v_j for $q_j = \{i \mid p_i \geq j\}$. Then the bipartite graph F has two sides $\{x_1, \ldots, x_m\}$ and $\{y_1, \ldots, y_{p_1}\}$ such that x_i is connected to y_1, \ldots, y_{p_i} . Then the vertex $v_{i,j}$ corresponds to $x_i y_j \in E(F)$ for $1 \leq i \leq m$ and $1 \leq j \leq p_i$, and the level graph L is isomorphic to the line graph of F.

Question [6.3](#page-15-0) can now be reformulated as follows.

Question 6.5. Is the line graph of every bipartite graph ∞ -perfect?

Proof that Questions [6.3](#page-15-0) and [6.5](#page-15-1) are equivalent. Let $G = (V, E)$ be the line graph of some bipartite graph G_1 . Then we note that for every subgraph G_i of $G_1,$ the line graph of G_i is $G[E(G_i)]$. Therefore, if $G_m \subseteq \cdots \subseteq G_1,$ we can take $V_i = E(G_i)$ and define the m-level graph $L = Lev(G; V_1, \ldots, V_m)$, where the vertex set of L is $\bigcup_{i=1}^m \{xy_i \mid xy \in E(G_i)\}.$ Then the vertices of L are in one-to-one correspondence with the edges of $H = H(G_1, \ldots, G_m)$, where the vertex xy_i of L, i.e., the copy of xy in the graph $G[V_i]$, corresponds to $ixy \in E(H)$. We also note that for every pair of vertices of H, the set of edges of H containing this pair correspond to the vertices of some clique in L. Conversely (using the fact that G_1 has no triangles), the vertices of any clique in L correspond to a set of edges in H with two vertices in their intersection. In addition, every independent set in L corresponds to a 2-matching in H and vice versa. From all these we get

$$
\tau^{(2)}(H) = \kappa(L)
$$
 and $\nu^{(2)}(H) = \alpha(L)$.

If we assume that the statement in Question [6.3](#page-15-0) is true, and $G = L(G_1)$ and $V_m \subseteq \ldots \subseteq V_1 = E(G_1)$ are given, then we can define $G_i = (V(G_1), V_i)$ for $i = 2, \ldots, m$ and get $\kappa(L) = \tau^{(2)}(H) = \nu^{(2)}(H) = \alpha(L)$ proving that the statement in Question [6.5](#page-15-1) is true.

Conversely, if we assume that the statement in Question [6.5](#page-15-1) is true, and we are given $G_m \subseteq \cdots \subseteq G_1$, then we can define $G = L(G_1)$ and $V_i = E(G_i)$ for $i = 1, ..., m$ and get $\tau^{(2)}(H) = \kappa(L) = \alpha(L) = \nu^{(2)}(H)$, proving that the statement in Question [6.3](#page-15-0) is true. \Box

An even stronger hypothesis is

Question 6.6. Is every perfect graph ∞ -perfect?

References

- [1] R. Aharoni, E. Berger, J. Briggs, and H. Guo. Looms. Preprint (2023). arXiv:2309.03735.
- [2] R. Aharoni and S. Zerbib. A generalization of Tuza's conjecture. J. Graph Theory 94 (2020), 445–462.
- [3] T. Y. Chow, C. K. Fan, M. X. Goemans, and J. Vondrak. Wide partitions, Latin tableaux, and Rota's basis conjecture. Adv. in Appl. Math. 31 (2003), 334–358.
- [4] R. Huang and G.-C. Rota. On the relations of various conjectures on Latin squares and straightening coefficients. Discrete Math. 128 (1994), 225–236.