SOME CONGRUENCES FOR 3-CORE CUBIC BIPARTITIONS

RUSSELLE GUADALUPE

ABSTRACT. In this paper, we derive certain congruences for the number of 3-core cubic bipartitions using elementary q-series manipulations and dissection formulas.

1. Introduction

Throughout this paper, we denote $(q^m;q^m)_{\infty} = \prod_{n\geq 1} (1-q^{mn})$ for |q|<1 and positive integer m. Recall that a partition of a positive integer n is a nonincreasing sequence of positive integers, known as its parts, whose sum is n. The study of the arithmetic properties of the number p(n) of partitions of n was initiated by Ramanujan [14, 15], who proved the following remarkable congruences

$$p(5n + 4) \equiv 0 \pmod{5},$$

 $p(7n + 5) \equiv 0 \pmod{7},$
 $p(11n + 6) \equiv 0 \pmod{11}.$

Ahlgren [1] generalized this result by showing that infinitely many congruences of the form $p(An + B) \equiv 0 \pmod{M}$ exist when M is coprime to 6. Since the inception of p(n), analogous congruences for the other types of partitions have been established by various authors [2, 8, 9, 13, 16, 17], including those obtained by Chan [4, 5] for the cubic partition function a(n), with generating function given by

$$\sum_{n=0}^{\infty} a(n)q^n = \frac{1}{(q;q)_{\infty}(q^2;q^2)_{\infty}},$$

and by Chen [6, 7] for the t-core partition function $c_t(n)$, with generating function given by

$$\sum_{n=0}^{\infty} c_t(n) q^n = \frac{(q^t; q^t)_{\infty}^t}{(q; q)_{\infty}}$$

for $t \geq 1$. We note that a(n) is the number of partitions in two colors such that the second color only appears in even parts, and $a_t(n)$ is the number of partitions whose Ferrers graph has no hook numbers divisible by t. In 2017, Gireesh [11] examined the arithmetic properties of the function $C_3(n)$, which is the number of cubic partitions of n which are also 3-cores, whose generating function is given by

$$\sum_{n=0}^{\infty} C_3(n)q^n = \frac{(q^3; q^3)_{\infty}^3 (q^6; q^6)_{\infty}^3}{(q; q)_{\infty} (q^2; q^2)_{\infty}}.$$

Recently, da Silva and Sellers [10] extended Gireesh's results by obtaining dissections and congruences for $C_3(n)$, including $C_3(24n + 21) \equiv 0 \pmod{4}$ and $C_3(18n + 14) \equiv 0$

(mod 9). Motivated by their results, we derive in this paper some congruences for the number $CP_3(n)$ of 3-core cubic bipartitions, whose generating function is given by

$$\sum_{n=0}^{\infty} CP_3(n)q^n = \frac{(q^3; q^3)_{\infty}^6 (q^6; q^6)_{\infty}^6}{(q; q)_{\infty}^2 (q^2; q^2)_{\infty}^2}.$$

More precisely, $CP_3(n)$ is the number of ordered pairs (μ, κ) such that both μ and κ are 3-core cubic partitions such that the sum of all of the parts of μ and κ is n. We apply elementary q-series manipulations and dissection formulas to present our results.

We organize our paper as follows. In Section 2, we list some identities that are needed to prove our results. Section 3 contains certain dissection formulas and resulting congruences for $CP_3(n)$.

2. Preliminaries

In this section, we present several 2- and 3-dissection formulas needed to derive our congruences involving $CP_3(n)$. We denote $f_m := (q^m; q^m)_{\infty}$ for integer $m \ge 1$ throughout the rest of the paper.

Lemma 2.1. The following 2-dissections hold:

(1)
$$f_1^2 = \frac{f_2 f_8^5}{f_4^2 f_{16}^2} - 2q \frac{f_2 f_{16}^2}{f_8},$$

(2)
$$\frac{1}{f_1^2} = \frac{f_8^5}{f_2^5 f_{16}^2} + 2q \frac{f_4^2 f_{16}^2}{f_2^5 f_8},$$

(3)
$$f_1^4 = \frac{f_4^{10}}{f_2^2 f_8^4} - 4q \frac{f_2^2 f_8^4}{f_4^2},$$

(4)
$$\frac{f_3^3}{f_1} = \frac{f_4^3 f_6^2}{f_2^2 f_{12}} + q \frac{f_{12}^3}{f_4},$$

(5)
$$\frac{f_3^2}{f_1^2} = \frac{f_4^4 f_6 f_{12}^2}{f_5^5 f_8 f_{24}} + 2q \frac{f_4 f_6^2 f_8 f_{24}}{f_2^4 f_{12}},$$

(6)
$$\frac{f_3}{f_1^3} = \frac{f_4^6 f_6^3}{f_2^9 f_{12}^2} + 3q \frac{f_4^2 f_6 f_{12}^2}{f_2^7},$$

(7)
$$\frac{1}{f_1 f_3} = \frac{f_8^2 f_{12}^5}{f_2^2 f_4 f_6^4 f_{24}^2} + q \frac{f_4^5 f_{24}^2}{f_2^4 f_6^2 f_8^2 f_{12}}.$$

Proof. We refer to [10, Lem. 1] for a proof of (3) and [3, p. 40, Entry 25] for proofs of (1) and (2). Identities (4), (5) and (7) follow from [12, (22.7.5)], [12, (30.9.9)] and [12, (30.12.3)], respectively. Identity (6) follows after replacing q with -q in [12, (27.7.3)]. \square

Lemma 2.2. The following 3-dissections hold:

(8)
$$\frac{1}{f_1 f_2} = \frac{f_9^9}{f_3^6 f_6^2 f_{18}^3} + q \frac{f_9^6}{f_3^5 f_6^3} + 3q^2 \frac{f_9^3 f_{18}^3}{f_3^4 f_6^4} - 2q^3 \frac{f_{18}^6}{f_3^3 f_6^5} + 4q^4 \frac{f_{18}^9}{f_3^2 f_6^6 f_9^3},$$

(9)
$$\frac{f_2}{f_1^2} = \frac{f_6^4 f_9^6}{f_3^8 f_{18}^3} + 2q \frac{f_6^3 f_9^3}{f_3^7} + 4q^2 \frac{f_6^2 f_{18}^3}{f_3^6}.$$

Proof. Identity (8) follows from [4, (13)]. To obtain (9), replace q with $q^{1/3}$ in the q-expansion of $\frac{f_2}{f_1^2}$ and use [2, Lem. 2.1, 2.6-2.8]. The desired identity now follows from replacing q with q^3 .

Lemma 2.3. We have

(10)
$$\frac{f_1 f_2}{f_9 f_{18}} = \frac{1}{x(q^3)} - q - 2q^2 x(q^3),$$

(11)
$$\frac{f_3^4 f_6^4}{f_9^4 f_{18}^4} = \frac{1}{x(q^3)^3} - 7q^3 - 8q^6 x(q^3)^3,$$

where $x(q) = q^{-1/3}c(q)$, and

$$c(q) = \frac{q^{1/3}}{1 + \frac{q + q^2}{1 + \frac{q^2 + q^4}{1 + \dots}}}$$

is the Ramanujan's cubic continued fraction.

Proof. See
$$[11, Lem. 2.2-2.3]$$
.

3. Dissections and congruences for $CP_3(n)$

We start presenting our results by giving a 2-dissection of the generating function

(12)
$$\sum_{n=0}^{\infty} CP_3(n)q^n = \frac{f_3^6 f_6^6}{f_1^2 f_2^2}.$$

Theorem 3.1. We have

(13)
$$\sum_{n=0}^{\infty} CP_3(2n)q^n = \frac{f_2^6 f_3^{10}}{f_1^6 f_6^2} + q \frac{f_3^6 f_6^6}{f_1^2 f_2^2},$$

(14)
$$\sum_{n=0}^{\infty} CP_3(2n+1)q^n = 2\frac{f_2^2 f_3^8 f_6^2}{f_1^4}.$$

Proof. Substituting (4) into (12) yields

$$\sum_{n=0}^{\infty} CP_3(n)q^n = \left(\frac{f_3^3}{f_1}\right)^2 \frac{f_6^6}{f_2^2} = \left(\frac{f_4^3 f_6^2}{f_2^2 f_{12}} + q \frac{f_{12}^3}{f_4}\right)^2 \frac{f_6^6}{f_2^2}.$$

Extracting the terms with odd and even exponents gives

$$\sum_{n=0}^{\infty} CP_3(2n)q^{2n} = \frac{f_4^6 f_6^{10}}{f_2^6 f_{12}^2} + q^2 \frac{f_6^6 f_{12}^6}{f_2^2 f_4^2},$$
$$\sum_{n=0}^{\infty} CP_3(2n+1)q^{2n+1} = 2q \frac{f_4^2 f_6^8 f_{12}^2}{f_2^4}.$$

We then divide the last identity by q and replace q^2 to q, arriving at identities (13) and (14).

We next give congruences modulo 8 and 16 for $CP_3(n)$.

Theorem 3.2. We have

$$CP_3(8n+3) \equiv 0 \pmod{8},$$

 $CP_3(8n+7) \equiv 0 \pmod{16}.$

Proof. Substituting (5) and (3) into (14), we have

$$\sum_{n=0}^{\infty} CP_3(2n+1)q^n = 2\frac{f_2^2 f_3^8 f_6^2}{f_1^4}$$

$$= 2\left(\frac{f_4^4 f_6 f_{12}^2}{f_2^5 f_8 f_{24}} + 2q \frac{f_4 f_6^2 f_8 f_{24}}{f_2^4 f_{12}}\right)^4 \left(\frac{f_4^{16}}{f_2^2 f_8^4} - 4q \frac{f_2^2 f_8^4}{f_4^2}\right) f_2^2 f_6^2.$$

We extract the terms with odd exponents, divide by q, and then replace q^2 with q so that

(15)
$$\sum_{n=0}^{\infty} CP_3(4n+3)q^n \equiv -8\frac{f_2^{14}f_3^6f_6^8}{f_1^{16}f_{12}^4} + 16\frac{f_2^{23}f_3^7f_6^5}{f_1^{19}f_4^6f_{12}^2} \pmod{32}.$$

We next plug (1) and (6) into (15) to get

$$\sum_{n=0}^{\infty} CP_3(4n+3)q^n \equiv -8\left(\frac{f_4^6 f_6^3}{f_2^9 f_{12}^2} + 3q \frac{f_4^2 f_6 f_{12}^2}{f_2^7}\right)^6 \left(\frac{f_2 f_8^5}{f_4^2 f_{16}^2} - 2q \frac{f_2 f_{16}^2}{f_8}\right) \frac{f_2^{14} f_6^8}{f_{12}^4}$$

$$(16) \qquad + 16\left(\frac{f_4^6 f_6^3}{f_2^9 f_{12}^2} + 3q \frac{f_4^2 f_6 f_{12}^2}{f_7^7}\right)^7 \left(\frac{f_2 f_8^5}{f_4^2 f_{16}^2} - 2q \frac{f_2 f_{16}^2}{f_8}\right) \frac{f_2^{23} f_6^5}{f_4^6 f_{12}^2} \pmod{32}.$$

Extracting the terms in (16) with even exponents and replacing q^2 with q yields

$$\sum_{n=0}^{\infty} CP_3(8n+3)q^n \equiv 8 \frac{f_2^{34} f_3^{26} f_4^5}{f_1^{39} f_6^{16} f_8^2} - 8q \frac{f_2^{26} f_6^{22} f_4^5}{f_1^{35} f_6^8 f_8^2} - 8q^2 \frac{f_2^{18} f_3^{18} f_4^5}{f_1^{31} f_8^2} + 8q^3 \frac{f_2^{10} f_3^{14} f_4^5 f_6^8}{f_1^{27} f_8^2} \pmod{32},$$

which shows that $CP_3(8n+3) \equiv 0 \pmod{8}$. On the other hand, choosing the terms in (16) with odd exponents, dividing by q, and replacing q^2 with q leads us to

$$\sum_{n=0}^{\infty} CP_3(8n+7)q^n \equiv -16 \frac{f_2^{36} f_3^{26} f_8^2}{f_1^{39} f_4 f_6^{16}} + 16q \left(\frac{f_2^{22} f_3^{20} f_4^5}{f_1^{33} f_6^4 f_8^2} - \frac{f_2^{28} f_3^{22} f_8^2}{f_1^{35} f_4 f_6^8} \right) - 16q^2 \frac{f_2^{20} f_3^{18} f_8^2}{f_1^{31} f_4} + 16q^3 \left(\frac{f_2^6 f_3^{12} f_4^5 f_6^{12}}{f_1^{25} f_8^2} - \frac{f_2^{12} f_3^{14} f_6^8 f_8^2}{f_1^{27} f_4} \right), \pmod{32}$$

which proves the congruence $CP_3(8n+7) \equiv 0 \pmod{16}$.

The following theorem provides the 3-dissection of $CP_3(n)$.

Theorem 3.3. We have

(17)
$$\sum_{n=0}^{\infty} CP_3(3n)q^n = \frac{f_2^2 f_3^{18}}{f_1^6 f_6^6} + 2q \frac{f_3^9 f_6^3}{f_1^3 f_2} + 28q^2 \frac{f_6^{12}}{f_2^4},$$

(18)
$$\sum_{n=0}^{\infty} CP_3(3n+1)q^n = 2\frac{f_2f_3^{15}}{f_1^5f_6^3} + 13q\frac{f_3^6f_6^6}{f_1^2f_2^2} - 16q^2\frac{f_1f_6^{15}}{f_2^5f_3^3},$$

(19)
$$\sum_{n=0}^{\infty} CP_3(3n+2)q^n = 7\frac{f_3^{12}}{f_1^4} - 4q\frac{f_3^3 f_6^9}{f_1 f_2^3} + 16q^2 \frac{f_1^2 f_6^{18}}{f_2^6 f_3^6}.$$

Proof. Substituting (7) into (12) yields

$$\sum_{n=0}^{\infty} CP_3(n)q^n = \left(\frac{f_9^9}{f_3^6 f_6^2 f_{18}^3} + q \frac{f_9^6}{f_3^5 f_6^3} + 3q^2 \frac{f_9^3 f_{18}^3}{f_3^4 f_6^4} - 2q^3 \frac{f_{18}^6}{f_3^5 f_6^5} + 4q^4 \frac{f_{18}^9}{f_3^2 f_6^6 f_9^3}\right)^2 \frac{f_9^6}{f_6^6}.$$

Extracting the terms with exponents congruent to 0,1 and 2 modulo 3 gives

$$\sum_{n=0}^{\infty} CP_3(3n)q^{3n} = \frac{f_6^2 f_9^{18}}{f_3^6 f_{18}^6} + 2q^3 \frac{f_9^9 f_{18}^3}{f_3^3 f_6} + 28q^6 \frac{f_{18}^{12}}{f_6^4},$$

$$\sum_{n=0}^{\infty} CP_3(3n+1)q^{3n+1} = 2q \frac{f_6 f_9^{15}}{f_3^5 f_{18}^3} + 13q^4 \frac{f_9^6 f_{18}^6}{f_3^2 f_6^2} - 16q^7 \frac{f_3 f_{18}^{15}}{f_6^5 f_9^3},$$

$$\sum_{n=0}^{\infty} CP_3(3n+2)q^{3n+2} = 7q^2 \frac{f_9^{12}}{f_3^4} - 4q^5 \frac{f_9^3 f_{18}^9}{f_3 f_6^3} + 16q^8 \frac{f_3^2 f_{18}^{18}}{f_6^6 f_{18}^6}.$$

We then divide these three identities by 1, q and q^2 , respectively, and replace q^3 with q to arrive at (17), (18) and (19).

As a byproduct of Theorem 3.3, we find an infinite family of internal congruences modulo 2.

Corollary 3.4. We have

(20)
$$CP_3(3^k n + 3^k - 2) \equiv CP_3(n - 1) \pmod{2}$$

for all $k, n \geq 1$.

Proof. Reducing (18) modulo 2 yields

$$\sum_{n=0}^{\infty} CP_3(3n+1)q^n \equiv q \frac{f_3^6 f_6^6}{f_1^2 f_2^2} \equiv \sum_{n=0}^{\infty} CP_3(n)q^{n+1} \pmod{2}$$

Since $CP_3(1) = 2$, we have

$$\sum_{n=1}^{\infty} CP_3(3n+1)q^n \equiv \sum_{n=1}^{\infty} CP_3(n-1)q^n \pmod{2}$$

so that $CP_3(3n+1) \equiv CP_3(n-1) \pmod{2}$, which is (20) for k = 1. Hence, the congruence (20) for $k \geq 2$ now follows by induction.

We next provide the 3-dissection formulas for $CP_3(2n+1)$.

Theorem 3.5. We have

(21)
$$\sum_{n=0}^{\infty} CP_3(6n+1)q^n = 2\frac{f_2^6 f_3^{12}}{f_1^8 f_6^6} + 32q \frac{f_2^7 f_3^3 f_6^3}{f_1^5},$$

(22)
$$\sum_{n=0}^{\infty} CP_3(6n+3)q^n = 8\frac{f_2^9 f_3^9}{f_1^7 f_6^3} + 32q \frac{f_2^6 f_6^6}{f_1^4},$$

(23)
$$\sum_{n=0}^{\infty} CP_3(6n+5)q^n = 24\frac{f_2^8 f_3^6}{f_1^6}.$$

Proof. Substituting (8) into (14) yields

$$\sum_{n=0}^{\infty} CP_3(2n+1)q^n = 2\left(\frac{f_6^4 f_9^6}{f_3^8 f_{18}^3} + 2q \frac{f_6^3 f_9^3}{f_3^7} + 4q^2 \frac{f_6^2 f_{18}^3}{f_3^6}\right)^2 f_3^8 f_6^2.$$

Extracting the terms with exponents congruent to 0,1 and 2 modulo 3 gives

$$\sum_{n=0}^{\infty} CP_3(6n+1)q^{3n} = 2\frac{f_6^6 f_9^{12}}{f_3^8 f_{18}^6} + 32q^3 \frac{f_6^7 f_9^3 f_{18}^3}{f_3^5},$$

$$\sum_{n=0}^{\infty} CP_3(6n+3)q^{3n+1} = 8q \frac{f_6^9 f_9^9}{f_3^7 f_{18}^3} + 32q^4 \frac{f_6^6 f_{18}^6}{f_3^4},$$

$$\sum_{n=0}^{\infty} CP_3(6n+5)q^{3n+2} = 24q^2 \frac{f_6^8 f_9^6}{f_3^6}.$$

We then divide these three identities by 1, q and q^2 , respectively, and replace q^3 with q to arrive at (21), (22) and (23).

We now enumerate congruences for $CP_3(24n+r)$ for some values of nonnegative integers $r \leq 23$.

Theorem 3.6. We have

$$CP_3(24n + 7) \equiv 0 \pmod{16}$$
,
 $CP_3(24n + 13) \equiv 0 \pmod{4}$,
 $CP_3(24n + 19) \equiv 0 \pmod{8}$.

Proof. Substituting (3) and (5) into (21) and reducing modulo 32, we get

(24)

$$\sum_{n=0}^{\infty} CP_3(6n+1)q^n \equiv 2\left(\frac{f_4^4 f_6 f_{12}^2}{f_2^5 f_8 f_{24}} + 2q \frac{f_4 f_6^2 f_8 f_{24}}{f_2^4 f_{12}}\right)^6 \left(\frac{f_4^{10}}{f_2^2 f_8^4} - 4q \frac{f_2^2 f_8^4}{f_4^2}\right) \frac{f_2^6}{f_6^6} \pmod{32}.$$

Extracting the terms in (24) with even exponents and replacing q^2 with q yields

(25)
$$\sum_{n=0}^{\infty} CP_3(12n+1)q^n \equiv 2\frac{f_2^{34}f_6^{12}}{f_1^{26}f_4^{10}f_{12}^6} \pmod{8}.$$

We next put (2) in (25), extract the terms with odd exponents, divide by q, and replace q^2 with q, obtaining

$$\sum_{n=0}^{\infty} CP_3(24n+13)q^n \equiv 4 \frac{f_3^{12} f_4^{59}}{f_1^{31} f_2^8 f_6^6 f_8^{22}} \pmod{8},$$

so that $CP_3(24n+13) \equiv 0 \pmod{4}$. On the other hand, we consider the terms in (24) with odd exponents, divide by q, and replace q^2 with q to get

(26)
$$\sum_{n=0}^{\infty} CP_3(12n+7)q^n \equiv -8\frac{f_2^{22}f_6^{12}}{f_1^{22}f_4^2f_{12}^6} - 8\frac{f_2^{31}f_3f_6^9}{f_1^{25}f_4^8f_{12}^4} \pmod{32}.$$

Substituting (2) and (6) into (26) yields

$$\sum_{n=0}^{\infty} CP_3(12n+7)q^n \equiv -8\left(\frac{f_8^5}{f_2^5 f_{16}^2} + 2q \frac{f_4^2 f_{16}^2}{f_2^5 f_8}\right)^{11} \frac{f_2^{22} f_6^{12}}{f_4^2 f_{12}^6}$$

$$(27) \qquad \qquad -8\left(\frac{f_4^6 f_6^3}{f_2^9 f_{12}^2} + 3q \frac{f_4^2 f_6 f_{12}^2}{f_2^7}\right) \left(\frac{f_8^5}{f_2^5 f_{16}^2} + 2q \frac{f_4^2 f_{16}^2}{f_2^5 f_8}\right)^{11} \frac{f_2^{31} f_6^9}{f_4^8 f_{12}^4} \pmod{32}.$$

We consider the terms in (27) with even exponents and replace q^2 with q to arrive at

$$\sum_{n=0}^{\infty} CP_3(24n+7)q^n \equiv -16 \frac{f_3^{12} f_4^{55}}{f_1^{33} f_2^2 f_6^6 f_8^{22}} \pmod{32},$$

which follows that $CP_3(24n + 7) \equiv 0 \pmod{16}$. On the other hand, we consider the terms in (27) with odd exponents, divide by q, and replace q^2 with q to get

$$\sum_{n=0}^{\infty} CP_3(24n+19)q^n \equiv 8 \frac{f_3^{10} f_4^{55}}{f_1^{31} f_2^6 f_6^2 f_8^{22}} - 8 \frac{f_3^{12} f_4^{49}}{f_1^{33} f_6^6 f_8^{18}} \pmod{32},$$

so that $CP_3(24n + 19) \equiv 0 \pmod{8}$.

Theorem 3.7. We have

$$CP_3(24n + 15) \equiv 0 \pmod{16},$$

 $CP_3(24n + 21) \equiv 0 \pmod{16}.$

Proof. Putting (3), (5) and (7) in (22) and reducing modulo 32, we see that the right-hand side of (22) is congruent to

(28)

$$8\left(\frac{f_4^4f_6f_{12}^2}{f_2^5f_8f_{24}} + 2q\frac{f_4f_6^2f_8f_{24}}{f_2^4f_{12}}\right)^5\left(\frac{f_8^2f_{12}^5}{f_2^2f_4f_6^4f_{24}^2} + q\frac{f_4^5f_{24}^2}{f_2^4f_6^2f_8^2f_{12}}\right)\left(\frac{f_4^{10}}{f_2^2f_8^4} - 4q\frac{f_2^2f_8^4}{f_4^2}\right)\frac{f_2^9}{f_6^3} \pmod{32}.$$

Extracting the terms in (28) with even exponents and replacing q^2 with q yields

(29)
$$\sum_{n=0}^{\infty} CP_3(12n+3)q^n \equiv 8 \frac{f_2^{29} f_6^{15}}{f_1^{20} f_3^2 f_8^7 f_{12}^7} - 16q \frac{f_2^{32} f_3 f_6^6}{f_1^{21} f_4^9 f_{12}} \pmod{32}.$$

We next apply (2), (6) and (7) into (29) to arrive at

$$\sum_{n=0}^{\infty} CP_3(12n+3)q^n \equiv 8 \left(\frac{f_8^5}{f_2^5 f_{16}^2} + 2q \frac{f_4^2 f_{16}^2}{f_2^5 f_8} \right)^9 \left(\frac{f_8^2 f_{12}^5}{f_2^2 f_4 f_6^4 f_{24}^2} + q \frac{f_4^5 f_{24}^2}{f_2^4 f_6^2 f_8^2 f_{12}} \right) \frac{f_2^{29} f_6^{15}}{f_4^7 f_{12}^7}$$

$$(30) \qquad -16q \left(\frac{f_8^5}{f_2^5 f_{16}^2} + 2q \frac{f_4^2 f_{16}^2}{f_2^5 f_8} \right)^9 \left(\frac{f_6^6 f_3^3}{f_2^9 f_{12}^2} + 3q \frac{f_4^2 f_6 f_{12}^2}{f_2^7} \right) \frac{f_2^{32} f_6^6}{f_2^9 f_{12}} \quad (\text{mod } 32).$$

We consider the terms in (30) with odd exponents, divide by q, and replace q^2 with q to get

$$\sum_{n=0}^{\infty} CP_3(24n+15)q^n \equiv 16 \frac{f_3^7 f_4^{43} f_6^3}{f_1^{20} f_7^7 f_8^{14} f_{12}^4} + 16q \frac{f_2^5 f_3^{11} f_4^{35} f_{12}^4}{f_1^{24} f_6^9 f_8^{14}} \pmod{32},$$

so that $CP_3(24n + 15) \equiv 0 \pmod{16}$. On the other hand, we look for the terms in (28) with odd exponents, divide by q, and replace q^2 with q so that

(31)
$$\sum_{n=0}^{\infty} CP_3(12n+9)q^n \equiv 8 \frac{f_2^{35} f_6^9}{f_1^{22} f_4^{11} f_{12}^3} - 16 \frac{f_2^{26} f_6^{12}}{f_1^{19} f_3 f_4^5 f_{12}^5} \pmod{32}.$$

We now apply (2) and (7) into (31) to get

$$\sum_{n=0}^{\infty} CP_3(12n+9)q^n \equiv 8 \left(\frac{f_8^5}{f_2^5 f_{16}^2} + 2q \frac{f_4^2 f_{16}^2}{f_2^5 f_8} \right)^{11} \frac{f_2^{35} f_6^9}{f_4^{11} f_{12}^3}$$

$$-16 \left(\frac{f_8^5}{f_2^5 f_{16}^2} + 2q \frac{f_4^2 f_{16}^2}{f_2^5 f_8} \right)^9 \left(\frac{f_8^2 f_{12}^5}{f_2^2 f_4 f_6^4 f_{24}^2} + q \frac{f_4^5 f_{24}^2}{f_2^4 f_6^2 f_8^2 f_{12}} \right) \frac{f_2^{26} f_6^{12}}{f_2^5 f_{12}^5} \pmod{32}$$

We extract the terms in (32) with odd exponents, divide by q, and replace q^2 with q so that

$$\sum_{n=0}^{\infty} CP_3(24n+21)q^n \equiv 16 \frac{f_3^9 f_4^{49}}{f_1^{20} f_2^9 f_6^3 f_8^{18}} - 16 \frac{f_3^{10} f_4^{43} f_{12}^2}{f_1^{23} f_6^6 f_8^{18}} \pmod{32},$$

which yields $CP_3(24n + 21) \equiv 0 \pmod{16}$.

Theorem 3.8. We have

$$CP_3(24n + 11) \equiv 0 \pmod{48},$$

 $CP_3(24n + 23) \equiv 0 \pmod{96}.$

Proof. We start with putting (5) into (23), obtaining

(33)
$$\sum_{n=0}^{\infty} CP_3(6n+5)q^n = 24\left(\frac{f_4^4 f_6 f_{12}^2}{f_2^5 f_8 f_{24}} + 2q \frac{f_4 f_6^2 f_8 f_{24}}{f_2^4 f_{12}}\right)^3 f_2^8.$$

Considering the terms in (33) with odd exponents, we divide them by q and replace q^2 with q so that

(34)
$$\sum_{n=0}^{\infty} CP_3(12n+11)q^n = 144 \frac{f_2^9 f_3^4 f_6^3}{f_1^6 f_4 f_{12}} + 192q \frac{f_2^3 f_3^6 f_4^3 f_{12}^3}{f_1^4 f_6^3}.$$

We now apply (2), (4) and (5) into (34) to get

$$\sum_{n=0}^{\infty} CP_3(12n+11)q^n = 144 \left(\frac{f_4^4 f_6 f_{12}^2}{f_2^5 f_8 f_{24}} + 2q \frac{f_4 f_6^2 f_8 f_{24}}{f_2^4 f_{12}} \right)^2 \left(\frac{f_8^5}{f_2^5 f_{16}^2} + 2q \frac{f_4^2 f_{16}^2}{f_2^5 f_8} \right) \frac{f_2^9 f_6^3}{f_4 f_{12}}$$

$$+ 192q \left(\frac{f_4^3 f_6^2}{f_2^2 f_{12}} + q \frac{f_{12}^3}{f_4} \right)^2 \left(\frac{f_8^5}{f_2^5 f_{16}^2} + 2q \frac{f_4^2 f_{16}^2}{f_2^5 f_8} \right) \frac{f_2^3 f_4^3 f_{12}^3}{f_6^3}.$$

$$(35)$$

Extracting the terms in (35) with even exponents and replacing q^2 with q gives

$$\sum_{n=0}^{\infty} CP_3(24n+11)q^n \equiv 48 \frac{f_2^7 f_3^5 f_4^3 f_6^3}{f_1^6 f_8^2 f_{12}^2} \pmod{96},$$

which follows that $CP_3(24n + 11) \equiv 0 \pmod{48}$. On the other hand, we look for the terms in (35) with odd exponents, divide by q, and replace q^2 with q so that

$$\sum_{n=0}^{\infty} CP_3(24n+23)q^n \equiv 96 \frac{f_2^9 f_3^5 f_6^3 f_8^2}{f_1^6 f_4^3 f_{12}^2} \pmod{192},$$

so that $CP_3(24n + 23) \equiv 0 \pmod{96}$.

We end this section by presenting an infinite family of congruences modulo twice times powers of 3, as the following result says.

Theorem 3.9. We have

$$CP_3(2 \cdot 3^k n + 3^k - 2) \equiv 0 \pmod{2 \cdot 3^{k-1}}$$

for all $k, n \geq 1$.

Proof. In view of (14) and Corollary 3.4, it suffices to prove that

(36)
$$CP_3(3^k n + 3^k - 2) \equiv 0 \pmod{3^{k-1}}$$

for all $k, n \ge 1$. To show (36), we follow the argument used to prove [11, Thm. 3.1]. Let $a = q^{-1}f_1f_2ff_9^{-1}f_{18}^{-1}$, $b = q^{-1}x(q^3)^{-1}$ and $c = q^{-3}f_3^4f_6^4f_9^{-4}f_{18}^{-4}$. Then by (10) and (11) of Lemma 2.3, we see that

(37)
$$a = b - 1 - \frac{2}{b},$$

$$(38) c = b^3 - 7 - \frac{8}{b^3}$$

From (37) and (38), we note that

$$c+7 = \left(b - \frac{2}{b}\right)^3 + 6\left(b - \frac{2}{b}\right) = (a+1)^3 + 6(a+1),$$

which shows that $c = a^3 + 3a^2 + 9a$ and

(39)
$$\frac{1}{a^2} = \frac{1}{c^2}(a^4 + 6a^3 + 27a^2 + 54a + 81).$$

We rewrite (12) as

(40)
$$\sum_{n=0}^{\infty} CP_3(n)q^{n-1} = \frac{f_3^6 f_6^6}{q^3 f_9^2 f_{18}^2} \cdot \frac{1}{a^2},$$

and define the "huffing" operator H_3 modulo 3 (which is a linear map) by

$$H_3\left(\sum_{n=0}^{\infty} a(n)q^n\right) = \sum_{n=0}^{\infty} a(3n)q^{3n}.$$

Using (37), we compute

$$H_3(1) = 1,$$

$$H_3(a) = -1,$$

$$H_3(a^2) = -3,$$

$$H_3(a^3) = -\frac{8}{b^3} + 11 + b^3,$$

$$H_3(a^4) = \frac{32}{b^3} + 1 - 4b^3,$$

so that from (38) and (39), we get

$$H_3\left(\frac{1}{a^2}\right) = \frac{1}{c^2} \left(\frac{32}{b^3} + 1 - 4b^3 - \frac{48}{b^3} + 66 + 6b^3 - 81 - 54 + 81\right)$$

$$= \frac{1}{c^2} \left(-\frac{16}{b^3} + 13 + 2b^3\right) = \frac{2}{c} + \frac{27}{c^2}.$$
(41)

Thus, applying H_3 to (40), using (41), and replacing q^3 with q lead to

(42)
$$\sum_{n=0}^{\infty} CP_3(3n+1)q^n = 2(f_1f_2f_3f_6)^2 + 27q\frac{f_3^6f_6^6}{f_1^2f_2^2}.$$

We now define

(43)
$$(f_1 f_2 f_3 f_6)^2 = q^2 (f_3 f_6 f_9 f_{18})^2 a^2 := \sum_{n=0}^{\infty} d(n) q^n,$$

so that, in view of (42), we have

$$(44) CP_3(3n+1) = 2d(n) + 27CP_3(n-1)$$

for all $n \geq 1$. Applying H_3 to (43) and replacing q^3 with q lead to

$$\sum_{n=0}^{\infty} d(3n+2)q^n = -3(f_1f_2f_3f_6)^2,$$

which means that

$$(45) d(3n+2) = -3d(n)$$

for all $n \geq 0$. Thus, from (44) and (45), we get

$$CP_3(9n+7) = 2d(3n+2) + 27CP_3(3n+1) = -6d(n) + 27(2d(n) + 27CP_3(n-1))$$

= $48d(n) + 729CP_3(n-1)$

and by induction, we finally arrive at

$$CP_3(3^k n + 3^k - 2) = 3^{k-1} \cdot \frac{9^k - (-1)^k}{5} \cdot d(n) + 3^{3k}CP_3(n-1)$$

for all $k, n \ge 1$. Since $9^k - (-1)^k$ is divisible by 5 for all $k \ge 1$, we immediately obtain (36).

References

- [1] S. Ahlgren, Distribution of the partition function modulo composite integers M, Math. Ann. 318 (2000), 795–803.
- [2] G. E. Andrews, M. D. Hirschhorn and J. A. Sellers, Arithmetic properties of partitions with even parts distinct, Ramanujan J. 23 (2010), 169–181.
- [3] B. C. Berndt, Ramanujan's Notebooks, Part III, Springer, New York, 1991.
- [4] H.-C. Chan, Ramanujan's cubic continued fraction and an analog of his 'Most beautiful identity', Int. J. Number Theory 6 (2010), 673–680.
- [5] H.-C. Chan, Ramanujan's cubic continued fraction and Ramanujan type congruences for a certain partition function, Int. J. Number Theory 6 (2010), 819–834.
- [6] S. Chen, Arithmetical properties of the number of t-core partitions, Ramanujan J. 18 (2009), 103–112.
- [7] S. Chen, Congruences for t-core partition functions, J. Number Theory 133 (2013), 4036–4046.
- [8] S. Chern and M. G. Dastidar, Congruences and recursions for the cubic partition, Ramanujan J. 44 (2017), 559–556.
- [9] S. P. Cui and N. S. S. Gu, Arithmetic properties of l-regular partitions, Adv. Appl. Math. 51 (2013), 507–523.
- [10] R. da Silva and J. A. Sellers, Congruences for 3-core cubic partitions, Indian J. Pure Appl. Math. 54 (2023), 404–420.
- [11] D. S. Gireesh, Formulas for cubic partition with 3-cores, J. Math. Anal. Appl. 453 (2017), 20–31.
- [12] M. D. Hirschhorn, *The Power of q, A Personal Journey*, Developments in Mathematics vol. 49, Springer, New York, 2017.
- [13] M. S. Mahadeva Naika and B. Hemanthkumar, Arithmetic properties of 5-regular bipartitions, Int. J. Number Theory 13 (2017), 939–956.
- [14] S. Ramanujan, Some properties of p(n), the number of partitions of n, Proc. Cambridge Philos. Soc. 19 (1919), 207–210.
- [15] S. Ramanujan, Congruence properties of partitions, Math. Z. 9 (1921), 147–153.
- [16] L. Wang, Arithmetic identities and congruence for partition triples with 3-cores, Int. J. Number Theory 12 (2016), 995–1010.
- [17] L. Wang, Congruences modulo powers of 5 for two restricted bipartitions, Ramanujan J. 44 (2017), 471–4491.

Institute of Mathematics, University of the Philippines-Diliman, Quezon City 1101, Philippines

Email address: rguadalupe@math.upd.edu.ph