

Powers of the operator $u(z)\frac{d}{dz}$ and their connection with the combinatorial numbers

Ioana Petkova

Faculty of Mathematics and Informatics, Sofia University

Sofia, Bulgaria

e-mail: ioana_petkovaa@abv.bg

Abstract: In this paper the operator $A = u(z)\frac{d}{dz}$ is considered, where u is an arbitrary entire function. The expansion of A^k ($k \geq 1$) with the help of powers of differential operators is obtained, and it is shown that this expansion depends on special numbers. A connection between these numbers and known combinatorial numbers is given.

Keywords: Combinatorial numbers, Special numbers, Operators, Euler-Cauchy operator, Bessel functions, Entire functions.

2020 Mathematics Subject Classification: 33C10, 11B73, 11F25, 30D10.

1. Introduction

Let $u : \mathbb{C} \rightarrow \mathbb{C}$ be an entire function. We denote by A the operator

$$A = u(z)\frac{d}{dz}$$

A particular case of A (when $u(z) = z$) is the operator $z\frac{d}{dz}$, considered by L. Euler. Also we may consider the operators $e^z\frac{d}{dz}$, $\frac{1}{z}\frac{d}{dz}$ and many others. Just as every infinite sequence of numbers a_n is connected with a so-called generating function $f(z) = \sum_{n=0}^{\infty} a_n z^n$, it is interesting to know if the powers A^k of A are generating operators for some special sequences of numbers. The mentioned above may motivate one to find an explicit representation for the operators A^k , $k = 1, 2, 3, \dots$

In this paper we show that the powers A^k of A are generating operators for many kinds of special (combinatorial) numbers (some of them are Stirling numbers of the first and second kind).

Also, several new identities for combinatorial numbers are obtained.

2. The main result.

First we want to express the operator A^k with the help of $\left(\frac{d}{dz}\right)^s$.

Theorem 1. There exist numbers $C_{\alpha_1, \dots, \alpha_s}^{s, m, k+1}$, for which the following representation:

$$A^{k+1} = \sum_{s=1}^{k+1} P_{k+1}^s(u) \left(\frac{d}{dz}\right)^s, \quad (1)$$

is valid, where:

$$P_{k+1}^{k+1}(u) := u^{k+1}; \quad (2)$$

$$P_{k+1}^{k+1-s}(u) := \sum_{m=1}^s u^{k+1-m} F_m^{k+1}(u), \quad s = 1, \dots, k, \quad (3)$$

$$F_m^{k+1}(u) := \sum_{m, s} C_{\alpha_1, \dots, \alpha_s}^{s, m, k+1} \left(\frac{du}{dz}\right)^{\alpha_1} \cdots \left(\frac{d^s u}{dz^s}\right)^{\alpha_s} \quad (4)$$

and $\sum_{m, s}$ in (4) is over all nonnegative integers $\alpha_1, \dots, \alpha_s$ for which:

$$\alpha_1 + \alpha_2 + \dots + \alpha_s = m; \quad 1 \cdot \alpha_1 + \dots + s \cdot \alpha_s = s.$$

The numbers $C_{\alpha_1, \dots, \alpha_s}^{s, m, k+1}$ satisfy the following recurrence relations:

$$C_{\gamma_1, \dots, \gamma_s, \gamma_{s+1}}^{s+1, m, k+1} = \sum_{t=s+1}^k [(t+1-m) C_{\gamma_1-1, \gamma_2, \dots, \gamma_s}^{s, m-1, t} + \sum_{j=1}^{s+1} (\gamma_j+1) C_{\gamma_1, \dots, \gamma_{j-1}, \gamma_j+1, \gamma_{j+1}-1, \gamma_{j+2}, \dots, \gamma_s}^{s, m, t}], \quad (5)$$

where: $1 < m < s+1$; $\gamma_{s+1} = 0$; $\sum_{i=1}^{s+1} \gamma_i = m$; $\sum_{i=1}^{s+1} i \gamma_i = s+1$; $s = 1, \dots, k-1$;

$$C_{\gamma_1, \dots, \gamma_s, \gamma_{s+1}}^{s+1, s+1, k+1} = \sum_{t=s+1}^k (t-s) C_{\gamma_1-1, \gamma_2, \dots, \gamma_s}^{s, s, t}, \quad (6)$$

where: $\gamma_{s+1} = 0$; $\sum_{i=1}^{s+1} \gamma_i = s+1$; $\sum_{i=1}^{s+1} i \gamma_i = s+1$; $s = 1, \dots, k-1$

$$C_{\underbrace{0, \dots, 0}_s, 1}^{s+1, 1, k+1} = \sum_{t=s+1}^k C_{\underbrace{0, \dots, 0}_{s-1}, 1}^{s, 1, t}, \quad (7)$$

where $s = 1, 2, \dots, k-1$. ■

Remark. A particular case of (6) is the relation:

$$C_{\underbrace{s+1, 0, \dots, 0}_s}^{s+1, s+1, k+1} = \sum_{t=s+1}^k (t-s) C_{\underbrace{s, 0, \dots, 0}_{s-1}}^{s, s, t}, \quad s = 1, 2, \dots, k-1. \quad (8)$$

Proof of Theorem 1. Let $f : \mathbb{C} \rightarrow \mathbb{C}$ is an entire function and

$$A^k f(z) = \sum_{s=1}^k P_k^s(u) f^{(s)}(z).$$

Then

$$\begin{aligned} \sum_{s=1}^{k+1} P_{k+1}^s(u) f^{(s)}(z) &= A^{k+1} f(z) = A(A^k f(z)) = A\left(\sum_{s=1}^k P_k^s(u) f^{(s)}(z)\right) = \\ &= u\left(\frac{d}{dz} P_k^1(u)\right) f^{(1)}(z) + \sum_{s=2}^k \left(u P_k^{s-1}(u) + u \frac{d}{dz} P_k^s(u)\right) f^{(s)}(z) + u P_k^k(u). \end{aligned}$$

Comparing the left-hand side and the right-hand side, we obtain the relations:

$$P_{k+1}^1(u) = u \frac{d}{dz} P_k^1(u); \quad (9)$$

$$P_{k+1}^s(u) = u P_k^{s-1}(u) + u \frac{d}{dz} P_k^s(u), \quad 1 < s < k + 1; \quad (10)$$

$$P_{k+1}^{k+1}(u) = u P_k^k(u). \quad (11)$$

Since $P_1^1(u) = u$, then from (11) we obtain

$$P_{k+1}^{k+1} = u^{k+1}. \quad (12)$$

From (10) we obtain the equalities:

$$\begin{aligned} P_{k+1}^k(u) &= u P_k^{k-1}(u) + k u^k \frac{d}{dz} u; \\ P_k^{k-1}(u) &= u P_{k-1}^{k-2}(u) + (k-1) u^{k-1} \frac{d}{dz} u; \\ &\vdots \\ P_3^2(u) &= u P_2^1(u) + 2 u^2 \frac{d}{dz} u \end{aligned}$$

We multiply the first equality by 1, the second - by u , the third - by u^2 , ..., the last - by u^{k-2} , then adding the received equalities, we obtain

$$P_{k+1}^k(u) = u^{k-1} P_2^1(u) + u^k \frac{d}{dz} u \sum_{t=2}^k t$$

Since $P_2^1(u) = u \frac{d}{dz} u$, we obtain

$$P_{k+1}^k(u) = \left(\sum_{t=1}^k t\right) u^k \frac{d}{dz} u = \frac{k(k+1)}{2} u^k \frac{d}{dz} u \quad (13)$$

To calculate $P_{k+1}^{k-1}(u)$ we use the same approach, using the above representation for $P_{k+1}^k(u)$.

Continuing the same way, we see that (3) seems true. Let us prove (3). Substituting in (10) s with $k+1-s$, we obtain:

$$P_{k+1}^{k+1-s}(u) = u P_k^{k-s}(u) + u \frac{d}{dz} P_k^{k+s+1}(u), \quad s = 1, 2, \dots, k. \quad (14)$$

With the help of (14), we shall repeat the approach, realized for $P_{k+1}^k(u)$, but this time to obtain $P_{k+1}^{(k+1)-(s+1)}(u)$. Let us assume that $P_{k+1}^{k+1-s}(u)$ is given by (3). Then

$$\frac{d}{dz}P_{k+1}^{k+1-s}(u) = \sum_{m=1}^s (k+1-m)u^{k-m}F_m^{k+1}(u)\frac{d}{dz}u + \sum_{m=1}^s u^{k+1-m}\frac{d}{dz}F_m^{k+1}(u).$$

From (14), using the above equality, we obtain

$$\begin{aligned} \frac{d}{dz}P_{k+1}^{k+1-s}(u) &= u^k \frac{d}{dz}F_1^{k+1}(u) + \\ &+ \sum_{m=2}^s \left((k+2-m)F_{m-1}^{k+1}(u)\frac{d}{dz}u + \frac{d}{dz}F_m^{k+1}(u) \right) u^{k+1-m} + \\ &+ (k+1-s)u^{k-s}F_s^{k+1}(u)\frac{d}{dz}u \end{aligned} \quad (15)$$

Substituting in (15) $k+1$ with $k, k-1, k-2, \dots$, and after that multiplying the first equality by u , the second - by u^2 , the third - by u^3 , and so on, then adding the received equalities, we obtain

$$\begin{aligned} P_{k+1}^{(k+1)-(s+1)}(u) &= \left(\sum_{t=s+1}^k \frac{d}{dz}F_1^t(u) \right) u^k + \\ &+ \sum_{m=2}^s \left(\sum_{t=s+1}^k (t+1-m)F_{m-1}^t(u)\frac{d}{dz}u + \frac{d}{dz}F_m^t(u) \right) u^{k+1-m} + \\ &+ \left(\sum_{t=s+1}^k (t-s)F_s^t(u)\frac{d}{dz}u \right) u^{k-s} \end{aligned} \quad (16)$$

Using the representation of $P_{k+1}^{(k+1)-(s+1)}(u)$, given in (3), with the help of (4), we may conclude that the relations (5), (6) and (7) hold. Indeed, from (4), on one side we have:

$$F_{m-1}^t(u)\frac{d}{dz}u = \sum_{m-1, s} C_{\alpha_1, \dots, \alpha_s}^{s, m-1, t} \left(\frac{du}{dz} \right)^{\alpha_1+1} \left(\frac{d^2u}{dz^2} \right)^{\alpha_2} \dots \left(\frac{d^s u}{dz^s} \right)^{\alpha_s}, \quad (17)$$

and on the other side we have

$$\begin{aligned} \frac{d}{dz}F_m^t(u) &= \\ \sum_{m, s} \sum_{j=1}^s \delta_j C_{\delta_1, \dots, \delta_s}^{s, m, t} \left(\frac{du}{dz} \right)^{\delta_1} \dots \left(\frac{d^{j-1}u}{dz^{j-1}} \right)^{\delta_{j-1}} \left(\frac{d^j u}{dz^j} \right)^{\delta_j-1} \left(\frac{d^{j+1}u}{dz^{j+1}} \right)^{\delta_{j+1}+1} \left(\frac{d^{j+2}u}{dz^{j+2}} \right)^{\delta_{j+2}} \dots \left(\frac{d^s u}{dz^s} \right)^{\delta_s} \end{aligned} \quad (18)$$

Comparing (1), (3), (4) with (16) - (18), we obtain:

$$\delta_1 = \alpha_1 + 1, \dots, \delta_j - 1 = \alpha_j, \delta_{j+1} + 1 = \alpha_{j+1}, \dots, \delta_s = \alpha_s$$

and thus we get (5). The relations (6) and (7) could be received analogically. To finish the proof of Theorem 1, it remains to prove that each monomial in \sum , for example

$$\left(\frac{du}{dz} \right)^{\gamma_1} \left(\frac{d^2u}{dz^2} \right)^{\gamma_2} \dots \left(\frac{d^s u}{dz^s} \right)^{\gamma_s} \left(\frac{d^{s+1}u}{dz^{s+1}} \right)^{\gamma_{s+1}}$$

with $\gamma_{s+1} = 0$, is contained in the right-hand side of (16). This fact can be proved in the same way as in the proof of Lemma 3.2 (see [1]). The theorem is proved. ■

3. Corollaries

Corollary 1. It is valid

$$C_{0,\dots,0,1}^{s,1,k+1} = \binom{k+1}{s+1}, s = 1, \dots, k.$$

Corollary 2. It is valid

$$C_{s+1,0,\dots,0}^{s+1,s+1,k+1} = \sigma(k+1, k-s),$$

where $\sigma(n, m)$ are the Stirling numbers of the second kind (for Stirling numbers $S(n, m)$ of first and second kind, see [2]).

4. Some results for the special choices of the function $u(z)$

Replacing the function $u = u(z)$ in A by different entire functions, we could receive various combinatorial numbers. For example:

- for $u(z) = z$, we receive

$$A^k = \left(u(z) \frac{d}{dz}\right)^k = \left(z \frac{d}{dz}\right)^k = \sum_{s=1}^k \sigma(k, s) x^s \frac{d^s}{dx^s}.$$

where $\sigma(k, s)$ are Stirling numbers of the second kind.

- for $u(z) = e^z$, we receive

$$A^k = \left(u(z) \frac{d}{dz}\right)^k = e^{kz} \sum_{s=1}^k (-1)^{k-s} S(k, s) x^s \frac{d^s}{dx^s},$$

where $S(k, s)$ are Stirling numbers of the first kind.

- for $u(z) = \frac{1}{z}$, we receive

$$A^k = \left(\frac{1}{z} \frac{d}{dz}\right)^k = \sum_{s=0}^{k-1} (-1)^s (2s-1)!! \binom{k-1+s}{2s} \frac{1}{z^{k-s}} \left(\frac{d}{dz}\right)^{k-s},$$

where we define $(-1)!!=1$.

Some combinatorial identities

From the previous section and Theorem 1, one may deduce the following relations:

- $$\sum_{m=1}^s \sum_{m,s} C_{\alpha_1, \dots, \alpha_s}^{s,m,k} = (-1)^s S(k, k-s)$$
- $$\sum_{s,k} \frac{k!}{1^{\alpha_1} \alpha_1! \dots k^{\alpha_k} \alpha_k!} = \sum_{m=1}^{k-s} \sum_{m,k-s} C_{\alpha_1, \dots, \alpha_{k-s}}^{k-s,m,k}$$
- $$\sum_{m=1}^s \sum_{m,s} (1!)^{\alpha_1} \dots (s!)^{\alpha_s} C_{\alpha_1, \dots, \alpha_s}^{s,m,k+1} = (2s-1)!! \binom{k+s}{2s}.$$

A particular case of the operator A

If we choose $u(z) = \frac{1}{z}$, then $A = \frac{1}{z} \frac{d}{dz}$. In this case we have

Theorem 2. For $k = 1, 2, 3, \dots$, the following representation is valid:

$$\left(\frac{1}{z} \frac{d}{dz}\right)^{k+1} = \sum_{s=1}^k (-1)^{k+1-s} (2k - 2s + 1)!! \binom{2k+1-s}{s-1} z^{s-2(k+1)} \frac{d^s}{dz^s} + z^{-k+1} \frac{d^{k+1}}{dz^{k+1}},$$

where $(2n+1)!! = 1.3.5 \dots (2n+1)$ and $\binom{a}{b}$ are the binomial coefficients.

Proof. With the help of mathematical induction it is seen that, if $f(z)$ is an entire function, then

$$\left(\frac{1}{z} \frac{d}{dz}\right)^k f(z) = \sum_{s=1}^k \frac{a_k^s}{z^{2k-s}} f^{(s)}(z),$$

with $a_1^1 = 1$, $a_k^k = 1$.

Furthermore, we have

$$\begin{aligned} \left(\frac{1}{z} \frac{d}{dz}\right)^{k+1} f(z) &= \\ &= \left(\frac{1}{z} \frac{d}{dz}\right) \left[\left(\frac{1}{z} \frac{d}{dz}\right)^k f(z) \right] = \\ &= \frac{(-1)(2k-1)a_k^1}{z^{2(k+1)-1}} f^{(1)}(z) + \\ &+ \sum_{s=2}^k \frac{a_k^{s-1} - (2k-s)a_k^s}{z^{2(k+1)-s}} f^{(s)}(z) + \\ &+ \frac{a_k^k}{z^{k+1}} f^{z+1} = \\ &= \sum_{s=1}^{k+1} \frac{a_{k+1}^s}{z^{2(k+1)-s}} f^{(s)}(z) \end{aligned}$$

Then the following relations hold:

- 1) $a_{k+1}^1 = (-1)(2k-1)a_k^1$; a_1^1 , $k = 1, 2$,
- 2) $a_{k+1}^{k+1} = a_k^k = 1$, $k = 1, 2, \dots$
- 3) $a_{k+1}^s = a_k^{s-1} - (2k-s)a_k^s$, $s = 2, 3, \dots, k$

With the help of 2) for $s = k$ from 3) we receive

$$a_{k+1}^k = a_2^1 - (2+3+\dots+k) = -\frac{k(k+1)}{2}$$

For $s = k-1$ we receive

$$a_{k+1}^{k-1} = a_3 - (4a_3^2 + 5a_4^3 + \dots + (k+1)a_k^{k-1})$$

i.e.

$$a_{k+1}^{k-1} = \frac{1}{2} \sum_{s=2}^k (s-1)s(s+1) = \frac{(k-1)k(k+1)(k+2)}{2.4}$$

For $s = k - 2$ we receive

$$\begin{aligned} a_{k+1}^{k-2} &= a_4^1 - \frac{1}{2} \frac{1}{4} \sum_{s=4}^k (s-2)(s)(s+1)(s+2) = \\ &= -\frac{1}{2.4} \sum_{s=3}^k (s-2)(s-1)(s)(s+1)(s+2) = \\ &= -\frac{(k-2)(k-1)k(k+1)(k+2)(k+3)}{2.4.6} \end{aligned}$$

The previous calculations lead us to the conjecture:

$$a_{k+1}^{(k+1)-s} = (-1)^s \frac{(k-s+1)\dots(k-1)k\dots(k+s)}{2.4.6\dots(2s)}$$

This expression we rewrite in the form:

$$a_{k+1}^{(k+1)-s} = (-1)^s (2s-1)!! \binom{k+s}{2s}$$

After substituting s with $k+1-s$ in the above equality, we receive

$$a_{k+1}^s = (-1)^{k+1-s} (2k-2s+1)!! \binom{2k+1-s}{s-1}, s = 1, \dots, k; k = 1, 2, 3, \dots$$

Now, to prove strictly that the above representation of a_{k+1}^s holds, it is enough to verify with its help that 1) and 3) are satisfied.

The proof of 1) is trivial.

3) follows after substitution $2k-s = n$ and checking the equality

$$(n+1-s) \binom{n+1}{n+2-s} = (n+1-s) \binom{n}{n+2-s} + n \binom{n-1}{n-s},$$

which follows from the binomial equation

$$(n+1-s) \binom{n}{n+1-s} = n \binom{n-1}{n-s}$$

□

Acknowledgements

The work has been (financially) supported by Grant KP-06-N 62/5 of Bulgarian National Science Fund.

References

- [1] I. Petkova, *Series expansion of the Gamma function and its reciprocal*, Notes on Number Theory and Discrete Mathematics, 27(4), 104-115, doi: 10.7546/nntdm.2021.27.4.104-115, 2021, <https://nntdm.net/papers/nntdm-27/NNTDM-27-4-104-115.pdf>
- [2] Abramowitz, M. & Stegun, I. A. (1972). *Handbook of Mathematical Functions, Tenth Printing*, Washington.
- [3] Luke, Y. L. (1980). *Special Mathematical Functions and Their Approximations*, Mir Publishing House, Moscow (in Russian).
- [4] JANKE - EMDE - LÖSH, *Tafeln Höherer Funktionen*, Moscow, 1977, (in Russian)