

ABOUT UNIVERSALITY OF LARGE DEVIATION PRINCIPLES FOR CONJUGACY INVARIANT PERMUTATIONS

ALICE GUIONNET AND MOHAMMED SLIM KAMMOUN

ABSTRACT. We prove the universality of the large deviations for conjugacy invariant permutations with few cycles. As an application, we establish the universality of large deviations principles at speeds n and \sqrt{n} for the length of monotone subsequences in conjugacy invariant permutations, with a sharp control over the total number of cycles. This universality class includes the well-known Ewens measures.

CONTENTS

1. Introduction and main results	1
1.1. Monotone subsequences	3
1.2. Main Technical trick	5
1.3. Applications	6
1.3.1. Applications to the longest increasing subsequence.	6
1.3.2. Applications to large deviations for the Eulerian statistics	7
1.3.3. Edge of RSK	7
1.4. Some comments	7
2. Proof of the results	8
2.1. Proof of Property 8	8
2.2. Proof of Property $(CI_{\alpha,\alpha})$ for Ewens distributions	10
2.3. Proof of Theorem 6	10
2.4. Proof of Theorem 4	13
2.5. Proof of Proposition 5	15
2.6. Proof of Proposition 13	16
3. Further discussions	17
References	19

1. INTRODUCTION AND MAIN RESULTS

De Moivre was the first to show in 1738 that errors from the law of large numbers were described by the bell curve, also known as the Gaussian or the Normal curve. However, his approach was in essence combinatorial and his result only concerned the sum of independent Bernoulli variables. In a tour de force, Laplace [36] generalized this result and showed that the Gaussian law characterizes the fluctuations of many more random variables. This was the first instance of universality, and a key result in probability theory. Other distributions such as Poisson processes or the Gumbel distribution appear as limits of different models with weakly correlated structure. It was more recently showed that models with strongly correlated structures also have universal fluctuations, which are characterized by

limiting distributions such as the semicircular law [52, 18], Tracy-Widom distribution [5, 29], Airy process [48, 19], sine process [47, 19], etc., emerge. On the contrary to fluctuations and the central limit theorem, large deviations theory [16] rarely displays universal feature, the probability to take an unlikely state being in general much more dependent on the underlying model. However, recent advances in random matrix theory have shown a surprising universality feature in large deviations principles [8, 39, 25, 11, 24, 2, 23]. It was shown in [24] that the large deviations for the largest eigenvalues of Wigner matrices are universal, and the same as those of matrices with Gaussian entries derived in [8], within a class of matrices with subGaussian entries whose distribution is called sharp subGaussian because their Laplace transform is uniformly bounded by the Gaussian one with the same variance. [3, 15] show that the probability of deviating towards a sufficiently small value stay universal for all subGaussian entries, but are different for deviations towards large enough value if the entries are not sharp subGaussian. The goal of this article is to study a similar universality phenomenon for random permutations.

In the last few years, universality for random permutations have been investigated thoroughly, including various aspects like global convergence and fluctuations. In particular, for conjugacy invariant permutations (see Definition 1), many results suggest that for a large family of functions on permutations, the asymptotic behavior depends only on the number of fixed points [22, 34], or only on the number of fixed points and two cycles [13, 26, 33]. For general functions, some results have been proven for example in [21, 32].

Nevertheless, relatively few results are known for large deviations for general conjugacy invariant permutations. In some sense, this question is related with large deviations for random matrices since [28] shows that the law of the longest subsequence in a permutation chosen at random has the same limiting distribution as the largest eigenvalue of a Gaussian Wishart matrices. We will show that large deviations for random permutations can easily be seen to be universal in a rather wide class of conjugacy invariant permutations with a sharp control over the total number of cycles. The main technique that we will introduce in this article is based on coupling of random permutations and exponential approximations [16, Section 4.2.2], see section 1.2.

Hereafter, we will denote by \mathfrak{S}_n the set of permutations of $\{1, \dots, n\} = [n]$. To state more precisely our result, let us remind the definition of conjugacy invariant permutations.

Definition 1. *A random permutation σ_n of size n is said to be conjugacy invariant if for every $\rho \in \mathfrak{S}_n$,*

$$\rho \sigma_n \rho^{-1} \stackrel{d}{=} \sigma_n.$$

In other words, σ_n is conjugacy invariant if and only if the map $\sigma \in \mathfrak{S}_n \mapsto \mathbb{P}(\sigma_n = \sigma)$ depends only on the cycle structure of σ . This class of permutations first emerged in Biology. It includes well-known measures like the Ewens measures (see Definition 1), along with various generalizations, such as the Kingman virtual permutations [20, 35, 49]. In this article, we are interested in the following class of conjugacy-invariant permutations with a sharp control over the total number of cycles described as follows.

Definition 2. *Let $0 < \alpha, \beta \leq 1$. We say that a sequence of random permutations $(\sigma_n)_{n \geq 1}$ satisfies $(\mathcal{CI}_{\alpha, \beta})$ if*

- *for any n , σ_n is a conjugacy invariant permutation of size n ,*

- and, for any $\varepsilon > 0$,

$$(\mathcal{CI}_{\alpha,\beta}) \quad \lim_{n \rightarrow \infty} \frac{\ln \mathbb{P}\left(\frac{\#\sigma_n}{n^\alpha} > \varepsilon\right)}{n^\beta} = -\infty,$$

where $\#(\sigma)$ is the total number of cycles of σ .

The main goal of this article will be to show that the sets $(\mathcal{CI}_{\alpha,\beta})$ provide natural universality classes for the large deviation of the uniform law on permutations for some appropriate choices of α and β .

We observe that for any $\beta \leq \alpha$, $\mathcal{CI}_{\alpha,\beta} \subset \mathcal{CI}_{\alpha,\alpha}$. A classical example of random permutations satisfying $(\mathcal{CI}_{\alpha,\alpha})$ (for any α) is the uniform permutation. More generally, we prove in section 2.2, see Corollary 16, that the Ewens measures satisfy $(\mathcal{CI}_{\alpha,\alpha})$ for any positive real number α . Let θ be a non-negative real number. We say that a random permutation $\sigma_{\theta,n}^{\text{Ew}}$ follows the Ewens distribution with parameter θ if for every $\sigma \in \mathfrak{S}_n$,

$$(1) \quad \mathbb{P}\left(\sigma_{\theta,n}^{\text{Ew}} = \sigma\right) = \frac{\theta^{\#(\sigma)-1}}{\prod_{i=1}^{n-1} (\theta + i)}.$$

In particular, $\sigma_{1,n}^{\text{Ew}} = \sigma_n^{\text{Unif}}$ is the uniform permutation and $\sigma_{0,n}^{\text{Ew}}$ is a uniform cyclic permutation. Clearly, Ewens measures are conjugacy invariant since by definition $\mathbb{P}(\sigma_{\theta,n}^{\text{Ew}} = \sigma)$ depends only on the number of cycles of σ .

Our primary emphasis is the application of our universality results to monotone subsequences of random permutations.

1.1. Monotone subsequences. Let \mathfrak{S}_n be the set of permutations of size n . Given $\sigma \in \mathfrak{S}_n$, a subsequence $(\sigma(i_1), \dots, \sigma(i_k))$ is an increasing (resp. decreasing) subsequence of σ of length k if $i_1 < \dots < i_k$ and $\sigma(i_1) < \dots < \sigma(i_k)$ (resp. $\sigma(i_1) > \dots > \sigma(i_k)$). We denote by $\text{LIS}(\sigma)$ (resp. $\text{LDS}(\sigma)$) the length of the longest increasing (resp. decreasing) subsequence of σ . For example,

$$\text{if } \sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 5 & 3 & 2 & 4 & 1 \end{pmatrix}, \text{ LIS}(\sigma) = 2 \text{ and } \text{LDS}(\sigma) = 4.$$

The study of the asymptotic behavior of monotone subsequences within random permutations is famously known as Ulam's problem. In his seminal work [50], Ulam posed the conjecture that the limit, as n tends towards infinity, of $\frac{\mathbb{E}(\text{LIS}(\sigma_n^{\text{Unif}}))}{\sqrt{n}}$ exists. Vershik, Kerov, Logan, and Shepp [51, 37] proved that this limit is equal to 2. The limiting fluctuations are the same as that of some random matrices models [5, 6]. For historical details and full proofs, we refer to [44]. Large deviations of speed \sqrt{n} and n were also proved when $\sigma_n = \sigma_n^{\text{Unif}}$ follows the uniform law on \mathfrak{S}_n :

Theorem 3. [46, Theorem 2][17]

For any $x \geq 2$,

$$\lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} \ln \mathbb{P}\left(\text{LIS}\left(\sigma_n^{\text{Unif}}\right) \geq x\sqrt{n}\right) = -I_{\text{LIS}, \frac{1}{2}}(x)$$

and for any $0 < x < 2$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P}\left(\text{LIS}\left(\sigma_n^{\text{Unif}}\right) \leq x\sqrt{n}\right) = -I_{\text{LIS}, 1}(x)$$

where

$$I_{\text{LIS},\frac{1}{2}}(x) = 2x \cosh^{-1}\left(\frac{x}{2}\right) \quad \text{and} \quad I_{\text{LIS},1}(x) = -1 + \frac{x^2}{4} + 2 \ln\left(\frac{x}{2}\right) - \left(2 + \frac{x^2}{2}\right) \ln\left(\frac{2x^2}{4+x^2}\right).$$

Note that the large deviations to the right (for $x \geq 2$) and to the left (for $x < 2$) have different speed. This is very similar to the large deviations principles for the largest eigenvalue of the Gaussian matrix ensembles [40, 8, 9]. The large deviations to the left necessitate to move all the spectrum and in fact are related to the large deviations of the empirical measure of the eigenvalues : their speed is n^2 , typically, the number of random entries of the matrices [9]. Whereas large deviations to the right are easier as they only require to move one eigenvalue, namely the largest, which can be achieved for instance by making one very large entry, which corresponds to a speed of order n . Therefore, these two results are, in fact, rather different in nature.

Our first result, proven in Section 2.4, is the following.

Theorem 4. *If the sequence of random permutations $(\sigma_n)_{n \geq 1}$ satisfies $(\mathcal{CI}_{\frac{1}{2},\frac{1}{2}})$, then the law of $\frac{\text{LIS}(\sigma_n)}{\sqrt{n}}$ satisfies a large deviation principle with speed \sqrt{n} and good rate function $J_{\text{LIS},\frac{1}{2}}$ which is equal to $I_{\text{LIS},\frac{1}{2}}$ on $[2, +\infty)$ and $+\infty$ on $(-\infty, 2)$. In other words, for any measurable subset E of \mathbb{R} , we have :*

$$(2) \quad - \inf_{x \in E^\circ} J_{\text{LIS},\frac{1}{2}}(x) \leq \liminf_{n \rightarrow \infty} \frac{\ln \mathbb{P}\left(\frac{\text{LIS}(\sigma_n)}{\sqrt{n}} \in E\right)}{\sqrt{n}} \leq \limsup_{n \rightarrow \infty} \frac{\ln \mathbb{P}\left(\frac{\text{LIS}(\sigma_n)}{\sqrt{n}} \in E\right)}{\sqrt{n}} \leq - \inf_{x \in \bar{E}} J_{\text{LIS},\frac{1}{2}}(x).$$

Moreover, if the sequence of random permutations $(\sigma_n)_{n \geq 1}$ satisfies $(\mathcal{CI}_{\frac{1}{2},1})$, then the law of $\frac{\text{LIS}(\sigma_n)}{\sqrt{n}}$ satisfies a large deviation principle with speed n and good rate function $J_{\text{LIS},1}$ where

$$J_{\text{LIS},1}(x) = \begin{cases} I_{\text{LIS},1}(x) & 0 < x \leq 2 \\ 0 & x > 2 \\ +\infty & x \leq 0 \end{cases}.$$

In other words, for every measurable subset E of \mathbb{R} ,

$$(3) \quad - \inf_{x \in E^\circ} J_{\text{LIS},1}(x) \leq \liminf_{n \rightarrow \infty} \frac{\ln \mathbb{P}\left(\frac{\text{LIS}(\sigma_n)}{\sqrt{n}} \in E\right)}{n} \leq \limsup_{n \rightarrow \infty} \frac{\ln \mathbb{P}\left(\frac{\text{LIS}(\sigma_n)}{\sqrt{n}} \in E\right)}{n} \leq - \inf_{x \in \bar{E}} J_{\text{LIS},1}(x),$$

The same result holds if we replace LIS par LDS, the length of the longest decreasing subsequence.

Even if Ewens measures do not satisfy $(\mathcal{CI}_{\frac{1}{2},1})$ (except for $\theta = 0$) so that (3) is not guaranteed, we will show by a separate argument that universality still holds in the sense that the Ewens distribution still satisfies the same large deviation principle, see section 2.2:

Proposition 5. *For any $\theta \geq 0$, the sequence of random permutations following the Ewens distribution $\sigma_n \stackrel{d}{=} \sigma_{\theta,n}^{\text{Ew}}$, $n \geq 1$, satisfy the large deviation principle (3).*

1.2. Main Technical trick. In this section, we introduce our main technical trick, namely Theorem 6, which can be thought as an exponential approximation argument as presented in [16, Section 4.2.2]. To this end, we see the statistic under study, such as LIS, as a function f of the permutation. Our point is then that if this function f depends sufficiently smoothly on the permutation in the sense of Hypothesis \mathcal{H}_1 and the law of $n^{-\alpha} f(\sigma_n^{\text{Unif}})$ satisfies a large deviation principle with speed n^β , then this large deviation principle remains true for all the random permutations satisfying $(\mathcal{CI}_{\alpha,\beta})$. We then show that many well known statistics verify hypothesis \mathcal{H}_1 and then apply our result to various examples.

The goal of this section is to introduce a tool to prove universality for the large and moderate deviations bounds for some statistics. We first define for a permutation $\sigma \in \mathfrak{S}_n$, the set

$$A_\sigma = \begin{cases} \{\sigma\} & \text{if } \#(\sigma) = 1 \\ \{\rho \in \mathfrak{S}_n, \rho = \sigma \circ (i_1, i_2) \circ (i_1, i_3) \cdots \circ (i_1, i_{\#(\sigma)}) \text{ and } \#(\rho) = 1\} & \text{if } \#(\sigma) > 1 \end{cases}.$$

Above, $\{i_1, \dots, i_{\#(\sigma)}\}$ are distinct elements of $\{1, \dots, n\}$.

For example: $A_{(3,4)} = \{(1, 2, 3, 4), (1, 2, 4, 3), (1, 3, 4, 2), (1, 4, 3, 2)\}$. Indeed, $(1, 2, 3, 4) = (3, 4)(4, 2)(4, 1)$, $(1, 2, 4, 3) = (3, 4)(3, 2)(3, 1)$, etc.

Our main theorem states that large deviations are universal within the class $(\mathcal{CI}_{\alpha,\beta})$.

Theorem 6. *Let f be a function defined on $\bigcup_{n \geq 1} \mathfrak{S}_n$ and having values in \mathbb{R}^d for some fixed integer number d . We denote $|\cdot|$ the Euclidean distance in \mathbb{R}^d . Suppose that*

- *the following holds true*

$$(\mathcal{H}_1) \quad \sup_{\sigma \in \bigcup_{n \geq 1} \mathfrak{S}_n} \sup_{\rho \in A_\sigma} \frac{|f(\sigma) - f(\rho)|}{\#(\sigma)} < +\infty.$$

- *there exists some function J **continuous**, some $0 < \beta \leq \alpha < 1$ and \mathcal{I} an open set of \mathbb{R}^d such that for any $x \in \mathcal{I}$*

$$(\mathcal{H}_2) \quad \lim_{n \rightarrow \infty} \frac{1}{n^\beta} \ln \mathbb{P} \left(f \left(\sigma_n^{\text{Unif}} \right) \geq xn^\alpha \right) = -J(x).$$

The relation $a = (a_1, \dots, a_d) \geq b = (b_1, \dots, b_d)$ means $a_i \geq b_i$ for all $i \in [d]$.

- *Moreover, the sequence $(\sigma_n)_{n \geq 1}$ of random permutations satisfies $(\mathcal{CI}_{\alpha,\beta})$*

Then, for any $x \in \mathcal{I}$

$$\lim_{n \rightarrow \infty} \frac{1}{n^\beta} \ln \mathbb{P}(f(\sigma_n) \geq xn^\alpha) = -J(x).$$

It may appear challenging to verify (\mathcal{H}_1) for an arbitrary statistic. However, in some cases, it is more straightforward to prove the following condition:

$$(\mathcal{H}'1) \quad \sup_{\sigma \in \bigcup_{n \geq 1} \mathfrak{S}_n} \sup_{i,j} |f(\sigma) - f(\sigma \circ (i, j))| < +\infty.$$

A noteworthy observation is that $(\mathcal{H}'1)$ implies (\mathcal{H}_1) , as it directly follows from the triangle inequality.

Remark 7. The set of functions satisfying (\mathcal{H}_1) (resp. $(\mathcal{H}'1)$) is a vector space.

We next show that Hypothesis $(\mathcal{H}'1)$ is satisfied for a large class of well-known statistics. Hence, Theorem 6 implies the universality of the large deviations for these statistics as soon as they are known for the uniform measure. We prove the following Property in subsection 2.1.

Property 8. *The following functions satisfy $(\mathcal{H}'1)$:*

- i) *The longest increasing subsequence LIS.*
- ii) *The longest decreasing subsequence LDS.*
- iii) *The vector $(\lambda_1, \dots, \lambda_d)$ if λ_i denotes the length of the i^{th} row of the RSK image for $i \in [d]$. d is a fixed integer number.*
- iv) *The normalized Inversions count : $\frac{\text{Inv}(\sigma)}{n} = \frac{\text{card}(\{(i,j):i<j,\sigma(i)>\sigma(j)\})}{n}$.*
- v) *The descents count : $D(\sigma) := \text{card}\{i : \sigma(i+1) < \sigma(i)\}$.*
- vi) *The ascents count : $A(\sigma) := \text{card}\{i : \sigma(i+1) > \sigma(i)\}$.*
- vii) *The peaks count : $\text{Peaks}(\sigma) = \text{card}\{i : \sigma(i-1) < \sigma(i) > \sigma(i+1)\}$.*
- viii) *The valleys count : $\text{Valleys}(\sigma) = \text{card}\{i : \sigma(i-1) > \sigma(i) < \sigma(i+1)\}$.*
- ix) *The exceedance count : $\text{Exc}(\sigma) := \text{card}\{i : \sigma(i) > i\}$ ¹*
- x) *The normalized major index: $\frac{\text{Maj}(\sigma)}{n} = \frac{\sum_{\sigma(i+1) > \sigma(i)} i}{n}$.*
- xi) *The longest alternating subsequence LAS: Let $1 \leq i_1 < i_2 < \dots < i_k \leq n$. We say that $\sigma(i_1), \dots, \sigma(i_k)$ is an alternating subsequence of length k if $\sigma(i_1) < \sigma(i_2) > \sigma(i_3) < \sigma(i_4) \dots$ and let LAS be the length of the longest alternating subsequence.*

In (iii), $\lambda(\sigma) = (\lambda_i(\sigma))_{i \geq 1}$ denotes the RSK shape associated with the permutation σ , see a definition in [44, Section 1.6]

1.3. Applications.

1.3.1. *Applications to the longest increasing subsequence.* A direct application of Theorem 6 gives the upper large deviations (2) of Theorem 4.

Corollary 9. *If the sequence $(\sigma_n)_{n \geq 1}$ of random permutations satisfies $(CI_{\frac{1}{2}, \frac{1}{2}})$, then for any $x > 2$*

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} \ln \mathbb{P}(\text{LIS}(\sigma_n) \geq x\sqrt{n}) &= -I_{\text{LIS}, \frac{1}{2}}(x) \\ \lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} \ln \mathbb{P}(\text{LDS}(\sigma_n) \geq x\sqrt{n}) &= -I_{\text{LIS}, \frac{1}{2}}(x). \end{aligned}$$

Proof. The first point is a direct application of Theorem 6 by taking $f = \text{LIS}$ and $\alpha = \beta = \frac{1}{2}$. The hypotheses (\mathcal{H}_1) is satisfied thanks to Property 8 and (\mathcal{H}_2) is satisfied thanks to Theorem 3. Moreover, when σ_n follows the uniform law, the LDS has the same distribution as the LIS. This can be seen for instance by replacing the collection $\{i\}$ into $\{n-i\}$. The proof of the large deviation principle for the function LDS is therefore the same as for the function LIS. □

For the upper moderate deviation, we find the following universal result:

Corollary 10. *Let $\frac{1}{6} < \nu < \frac{1}{2}$. If the sequence $(\sigma_n)_{n \geq 1}$ of random permutations satisfies $(CI_{\nu, \frac{3\nu}{2} - \frac{1}{4}})$, then for any $x > 0$,*

$$\lim_{n \rightarrow \infty} \frac{1}{n^{\frac{3\nu}{2} - \frac{1}{4}}} \ln \mathbb{P}(\text{LIS}(\sigma_n) \geq 2\sqrt{n} + xn^\nu) = -\frac{4}{3}x^{\frac{3}{2}}.$$

¹For the specialists, in general, the normalized number of occurrences of any set of classical (inversions for example), consecutive (descents, ascents, double descents, peaks, valleys, etc.) or more generally (bi-)vincular pattern.

This result was proven in the uniform case in [38]. Our universality result therefore follows from Theorem 6 since the coefficients $(\alpha, \beta) = (\nu, \frac{3\nu}{2} - \frac{1}{4})$ satisfy $\beta = \frac{3\nu}{2} - \frac{1}{4} \leq \nu = \alpha$ for every $\nu \leq 1/2$ so that (\mathcal{H}_2) is satisfied. (\mathcal{H}_1) follows from Property 8.

1.3.2. *Applications to large deviations for the Eulerian statistics.* Let D be the number of descents in σ .

Definition 11. A function f is called an Eulerian statistic if $f(\sigma_n^{\text{Unif}}) \stackrel{d}{=} D(\sigma_n^{\text{Unif}})$ for every integer number n .

It is known, for example that the exceedances count Exc and the ascents count A are Eulerian. Note that the equality in distribution is not true for general conjugacy invariant permutations.

Corollary 12. If the sequence of random permutations $(\sigma_n)_{n \geq 1}$ satisfies $(\mathcal{CI}_{1,1})$, then for $f \in \{D, A, \text{Exc}\}$, for any $\frac{1}{2} < x < 1$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P}(f(\sigma_n) \geq xn) = -I_D(x).$$

Where $I_D(x) = \sup_t \{xt - \ln(\frac{\exp(t)-1}{t})\}$.

Proof. For the descents, this LDP is already known in the uniform case [10] which implies the same large deviation principle for the other Eulerian statistics readily as they have the same distribution. Hence, (\mathcal{H}_2) is satisfied with $\alpha = \beta = 1$. Universality follows again from Theorem 6, since Property 8 implies that D, A, Exc satisfy (\mathcal{H}_1) . \square

1.3.3. *Edge of RSK.* In this subsection, we give an additional result, which is not a direct application of Theorem 6 but which proof uses the same techniques.

We are interested in the lower tail of LIS, and more precisely the length of the first rows of the RSK image.

Proposition 13. If the sequence $(\sigma_n)_{n \geq 1}$ of random permutations satisfies $(\mathcal{CI}_{\frac{1}{2},1})$, then for any $0 < x_d < \dots < x_1 < 2$,

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P}(\forall i \in [d], \lambda_i(\sigma_n) \leq x_i \sqrt{n}) = -I_{\text{LIS},1}(x_d).$$

To the best of our knowledge, this result is not stated in the uniform case for general integer number d but it is immediate to adapt the proof of [17] to get the uniform case, see Section 2.6.

1.4. **Some comments.** Except for the special case of the uniform permutation, there is a lack of existing results regarding the large deviations of general conjugacy-invariant permutations. Even for the uniform case, the large deviation theory of many statistics has not been studied yet. For example, we are not aware of existing results for the large deviations for LAS, the permutations patterns counts, and the upper edge of RSK in the uniform case. Even if the hypotheses $(\mathcal{H}'1)$ is satisfied for a large family of functions, since we are using comparison techniques, the application of Theorem 6 is not possible without knowing the large deviations in the case of uniform permutations.

In the context of random matrices, two distinct classes of large deviation results emerge: those concerning the edge of the spectrum and those concerning the the global spectral measure. The methodologies employed in proving these results typically differ significantly.

Our results primarily align with edge results. The analogue to the spectral measure in the context of random permutations is the shape of the RSK image. Using our techniques, one could investigate the universality of the shape. However, choosing the appropriate topology may be challenging.

In the literature, some non-universality results have been established for certain statistics but for other families of random permutations. For instance, [17] studied the longest increasing subsequence of i.i.d. points sampled from a measure on the unit square. Additionally, [42] explores the large deviations of LIS and LAS for permutations uniformly chosen among those that avoid a pattern of length 3. Moreover, in [41, Theorem B], a large deviations principle for the major index has been proven for a distinct family of random permutations.

A permutation can be viewed from two perspectives: as a matrix or as a word. In the context of Ewens permutations, their interpretation as random matrices reveals that many spectral properties are non-universal, i.e. depends on θ as the size of the permutation, goes to infinity [7, 4]. However, when considered as random words, numerous statistics become universal in the limit [21, 32]. This work can be seen as an extension of the latter results to large deviations.

While our proof uses a key argument of [32], the context of large deviations necessitates new ideas, particularly concentration inequalities, to ensure exponential approximations. Moreover, for the specific case of the large deviations of the longest increasing subsequence at speed n , we introduce novel probabilistic arguments, notably Lemma 18 and (19).

2. PROOF OF THE RESULTS

2.1. Proof of Property 8. In this section, we prove that many natural statistics satisfy $(\mathcal{H}'1)$.

- *Monotone subsequences* (LIS, LDS, and λ_i): The cases of LIS and LDS have already been demonstrated in [30, Lemma 3.1], and the proof for λ_i is presented in [30, Lemma 3.4].
- *Inversions*:

Let $\sigma \in \mathfrak{S}_n$ be a fixed permutation, and let $1 \leq i_1 < j_1 \leq n$. Define $\rho = \sigma \circ (i_1, j_1)$.

The key observation is that for all $i \notin \{i_1, j_1\}$, we have $\rho(i) = \sigma(i)$. The remaining of the proof varies depending on the specific statistic considered, but the underlying idea remains the same. We will provide detailed explanations for a few of these statistics.

Inversions can be expressed as follows:

$$\text{Inv}(\sigma) = |\{(i, j) : i < j, \sigma(i) > \sigma(j)\}|.$$

We make the following decomposition:

$$\underbrace{\{(i, j) : i < j, \sigma(i) > \sigma(j)\}}_s = \underbrace{\{(i, j) : i < j, \sigma(i) > \sigma(j), \{i, j\} \cap \{i_1, j_1\} \neq \emptyset\}}_{s_1} \cup \underbrace{\{(i, j) : i < j, \sigma(i) > \sigma(j), \{i, j\} \cap \{i_1, j_1\} = \emptyset\}}_{s_2}.$$

Furthermore, for inversions of ρ , we have:

$$\underbrace{\{(i, j) : i < j, \rho(i) > \rho(j)\}}_{s'} = \underbrace{\{(i, j) : i < j, \rho(i) > \rho(j), \{i, j\} \cap \{i_1, j_1\} \neq \emptyset\}}_{s'_1} \cup \underbrace{\{(i, j) : i < j, \rho(i) > \rho(j), \{i, j\} \cap \{i_1, j_1\} = \emptyset\}}_{s'_2}.$$

Using the key observation, one can see that $s'_2 = s_2$, which allows us to write:

$$\begin{aligned} |\text{Inv}(\sigma) - \text{Inv}(\rho)| &= |\text{card}(s) - \text{card}(s')| = |\text{card}(s'_1) - \text{card}(s_1)| \\ &\leq \max(\text{card}(s'_1), \text{card}(s_1)) \\ &\leq 2n. \end{aligned}$$

As a consequence,

$$\frac{1}{n} |\text{Inv}(\sigma) - \text{Inv}(\rho)| \leq 2$$

which proves that $n^{-1}\text{Inv}$ satisfies $(\mathcal{H}'1)$.

- *Descents and Major index:* The same idea applies here.

Recall that $D(\sigma) = |\{i : \sigma(i+1) < \sigma(i)\}|$. Let

$$\underbrace{\{i : \sigma(i+1) < \sigma(i)\}}_s = \underbrace{\{i : \sigma(i+1) < \sigma(i), \{i, i+1\} \cap \{i_1, j_1\} \neq \emptyset\}}_{s_1} \cup \underbrace{\{i : \sigma(i+1) < \sigma(i), \{i, i+1\} \cap \{i_1, j_1\} = \emptyset\}}_{s_2}$$

and

$$\underbrace{\{i : \rho(i+1) < \rho(i)\}}_{s'} = \underbrace{\{i : \rho(i+1) < \rho(i), \{i, i+1\} \cap \{i_1, j_1\} \neq \emptyset\}}_{s'_1} \cup \underbrace{\{i : \rho(i+1) < \rho(i), \{i, i+1\} \cap \{i_1, j_1\} = \emptyset\}}_{s'_2}.$$

Then, we have

$$|D(\sigma) - D(\rho)| \leq \max(\text{card}(s_1), \text{card}(s'_1)) \leq 4.$$

Similarly, with $\text{Maj}(\sigma) = \sum_{\sigma(i+1) > \sigma(i)} i$, we find

$$|\text{Maj}(\sigma) - \text{Maj}(\rho)| = \left| \sum_{i \in s_1} i - \sum_{i \in s'_1} i \right| \leq \max \left(\sum_{i \in s_1} i, \sum_{i \in s'_1} i \right) \leq n \max(\text{card}(s_1), \text{card}(s'_1)) \leq 4n.$$

- For *Peaks and Valleys*, the proof is similar to that of descents.
- For longest alternating subsequence LAS, we can use the following characterization (see [27] and [43, Corollary 2])

$$\text{LAS}(\sigma) = 1 + \sum_{i=1}^{n-1} M_k(\sigma),$$

where $M_1(\sigma) = \mathbb{1}_{\sigma(1) > \sigma(2)}$ and for $1 < k < n$,

$$M_k(\sigma) = \mathbb{1}_{\sigma(k-1) > \sigma(k) < \sigma(k+1)} + \mathbb{1}_{\sigma(k-1) < \sigma(k) > \sigma(k+1)}.$$

We have that $\text{LAS} = M_1 + \text{Valleys} + \text{Peaks}$. M_1 satisfies $(\mathcal{H}'1)$ since it is a bounded function. Therefore, by using Remark 7, LAS also satisfies $(\mathcal{H}'1)$.

2.2. Proof of Property $(\mathcal{CI}_{\alpha,\alpha})$ for Ewens distributions. For the Ewens distribution, it is known that the law of the total number of cycles has a nice description as a sum of independent Bernoulli variables.

Proposition 14. *The number of cycles of $\sigma_{\theta,n}^{\text{Ew}}$ is equal in distribution to $\sum_{i=1}^n a_{\theta,i}$ where $(a_{\theta,i})_i$ are independent Bernoulli variables with $\mathbb{P}(a_{\theta,i} = 1) = \frac{\theta}{i+\theta-1}$.*

This property can be proved independently by the Chinese restaurant process description of Ewens permutations or by the Feller coupling. It is a classical result, we can cite for example [14, equation (1.1)] and [1]. Many concentration inequalities are known for the sum of independent variables. For our purpose, we use a special form of the Bennett's inequality.

Proposition 15. [12, Theorem 2.9]²

Let X_1, X_2, \dots, X_n be independent random variables such that almost surely $X_i \leq 1$ then

$$(4) \quad \ln \left(\mathbb{P} \left(\sum_{i=1}^n X_i - \mathbb{E}(X_i) > t \right) \right) \leq -(v+t) \ln \left(1 + \frac{t}{v} \right) + t.$$

where $v = \sum_{i=1}^n \mathbb{E}X_i^2$.

Corollary 16. *The sequence $(\sigma_{\theta,n}^{\text{Ew}})_{n \geq 1}$ satisfies $(\mathcal{CI}_{\alpha,\alpha})$, for every $\theta \geq 0$, and every $\alpha > 0$,*

Proof. Using Proposition 14, we have $\#(\sigma_{\theta,n}^{\text{Ew}}) \stackrel{d}{=} \sum_{i=1}^n a_{\theta,i}$, and

$$\sum_{i=1}^n \mathbb{E}(a_{\theta,i}) = \sum_{i=1}^n \mathbb{E}(a_{\theta,i}^2) = \theta \ln(n) + O(1).$$

We want to prove that for every $\varepsilon > 0$, every $\alpha > 0$,

$$(5) \quad \lim_{n \rightarrow \infty} \frac{\ln \mathbb{P} \left(\frac{\# \sigma_{\theta,n}^{\text{Ew}}}{n^\alpha} > \varepsilon \right)}{n^\alpha} = -\infty,$$

This is a direct consequence of Proposition 14 by setting $X_i = a_{\theta,i}$, $t = \varepsilon n^\alpha - v$ in Proposition 15. \square

2.3. Proof of Theorem 6. In order to prove Theorem 6, we need to introduce a one step Markov chain T . It is the same as in [31]. It maps a conjugacy invariant random permutations σ_n to a permutation having the same law as $\sigma_{0,n}^{\text{Ew}}$.

This Markov chain does not change a lot statistics satisfying (\mathcal{H}_1) . Let T be the Markov chain defined on \mathfrak{S}_n and associated to the stochastic matrix $\left[\frac{1_{A_\sigma(\rho)}}{\text{card}(A_\sigma)} \right]_{\sigma, \rho \in \mathfrak{S}_n}$ where we recall that

$$A_\sigma = \begin{cases} \{\sigma\} & \text{if } \#(\sigma) = 1 \\ \{\rho \in \mathfrak{S}_n, \sigma^{-1} \circ \rho = (i_1, i_2) \circ (i_1, i_3) \cdots \circ (i_1, i_{\#(\sigma)}) \text{ and } \#(\rho) = 1\} & \text{if } \#(\sigma) > 1 \end{cases}.$$

T is then the Markov operator mapping a permutation σ to a permutation uniformly chosen at random among the permutations obtained by merging the cycles of σ using transpositions having all a common point.

²The version we use is [12, Theorem 2.9] by setting $b = 1$.

Lemma 17. [31, Lemma 6] *For any conjugacy invariant random permutation σ_n on \mathfrak{S}_n ,*

$$T(\sigma_n) \stackrel{d}{=} \sigma_{0,n}^{\text{Ew}}.$$

The proof is detailed in [31, Lemma 6] but the idea is rather simple. By construction, $T(\sigma_n)$ has almost surely one cycle. Since permutations with one cycle belong to the same conjugacy class, it is sufficient to prove that $T(\sigma_n)$ is conjugacy invariant as soon as σ_n is conjugacy invariant.

We move now to the proof of Theorem 6

Proof of Theorem 6. We consider a function f on $\cup_{n \geq 1} \mathfrak{S}_n$ with values in \mathbb{R}^d . One can suppose for simplicity that

$$(6) \quad \sup_{\sigma \in \cup_{n \geq 1} \mathfrak{S}_n} \sup_{\rho \in A_\sigma} \frac{|f(\sigma) - f(\rho)|}{\#(\sigma)} \leq 1.$$

Otherwise one can apply the theorem to $\frac{f}{\sup_{\sigma \in \cup_{n \geq 1} \mathfrak{S}_n} \sup_{\rho \in A_\sigma} \frac{|f(\sigma) - f(\rho)|}{\#(\sigma)}}$ since by hypothesis the denominator is finite. Let σ_n be conjugacy invariant. By Lemma 17, we know that

$$(7) \quad \mathbb{P}(f(\sigma_{0,n}^{\text{Ew}}) \geq xn^\alpha) = \mathbb{P}(f(T(\sigma_n)) \geq xn^\alpha).$$

Moreover, by (6), we have

$$\mathbb{P}(f(\sigma_n) \geq xn^\alpha + \#\sigma_n \mathbf{1}) \leq \mathbb{P}(f(T(\sigma_n)) \geq xn^\alpha) \leq \mathbb{P}(f(\sigma_n) \geq xn^\alpha - \#\sigma_n \mathbf{1}).$$

Here, $\mathbf{1}$ is the vector of \mathbb{R}^d with all components equal to 1. Let $\varepsilon > 0$. We write the following decomposition

$$\begin{aligned} p^\pm := \mathbb{P}(f(\sigma_n) \geq xn^\alpha \pm \#\sigma_n \mathbf{1}) &= \underbrace{\mathbb{P}(f(\sigma_n) \geq xn^\alpha \pm \#\sigma_n \mathbf{1} | \#(\sigma_n) < \varepsilon n^\alpha)}_{p_1^\pm} \underbrace{\mathbb{P}(\#(\sigma_n) < \varepsilon n^\alpha)}_{p_2} \\ &+ \underbrace{\mathbb{P}(f(\sigma_n) \geq xn^\alpha \pm \#\sigma_n \mathbf{1} | \#(\sigma_n) \geq \varepsilon n^\alpha)}_{p_3^\pm} \underbrace{\mathbb{P}(\#(\sigma_n) \geq \varepsilon n^\alpha)}_{p_4}. \end{aligned}$$

Moreover,

$$p_1^- \leq \mathbb{P}(f(\sigma_n) \geq (x - \varepsilon \mathbf{1})n^\alpha | \#(\sigma_n) < \varepsilon n^\alpha)$$

gives readily that

$$p_1^- p_2 \leq \mathbb{P}(f(\sigma_n) \geq (x - \varepsilon \mathbf{1})n^\alpha \text{ and } \#(\sigma_n) < \varepsilon n^\alpha) \leq \mathbb{P}(f(\sigma_n) \geq (x - \varepsilon \mathbf{1})n^\alpha).$$

Consequently, we find that

$$(8) \quad p^- \leq \mathbb{P}(f(\sigma_n) \geq (x - \varepsilon \mathbf{1})n^\alpha) + p_4.$$

Similarly,

$$p_1^+ \geq \mathbb{P}(f(\sigma_n) \geq (x + \varepsilon \mathbf{1})n^\alpha | \#(\sigma_n) < \varepsilon n^\alpha)$$

and then

$$\begin{aligned} (9) \quad p^+ &\geq p_1^+ p_2 \geq \mathbb{P}(f(\sigma_n) \geq (x + \varepsilon \mathbf{1})n^\alpha \text{ and } \#(\sigma_n) < \varepsilon n^\alpha) \\ &= \mathbb{P}(f(\sigma_n) \geq (x + \varepsilon \mathbf{1})n^\alpha) - \mathbb{P}(f(\sigma_n) \geq (x + \varepsilon \mathbf{1})n^\alpha \text{ and } \#(\sigma_n) \geq \varepsilon n^\alpha) \\ &\geq \mathbb{P}(f(\sigma_n) \geq (x + \varepsilon \mathbf{1})n^\alpha) - p_4. \end{aligned}$$

To sum-up, for any conjugacy invariant permutation σ_n , for any $\varepsilon > 0$, (6),(7),(8),(9) imply

$$(10) \quad \mathbb{P}(f(\sigma_n) \geq (x + \varepsilon \mathbf{1})n^\alpha) - \mathbb{P}(\#(\sigma_n) \geq \varepsilon n^\alpha) \leq \mathbb{P}(f(\sigma_{0,n}^{\text{Ew}}) \geq xn^\alpha) \\ \leq \mathbb{P}(f(\sigma_n) \geq (x - \varepsilon \mathbf{1})n^\alpha) + \mathbb{P}(\#(\sigma_n) \geq \varepsilon n^\alpha).$$

We next choose $\sigma_n = \sigma_n^{\text{Unif}}$ to be the uniform permutation. Because the Ewens distribution with $\theta = 1$ is the uniform distribution, Corollary 16 implies that σ_n^{Unif} is $(\mathcal{CI}_{\alpha,\alpha})$. Therefore, for any $\varepsilon > 0$, for n large enough we have under the hypothesis of Proposition 6 and because $\beta \leq \alpha$, we find for all $M > 0$

$$\mathbb{P}(f(\sigma_n^{\text{Unif}}) \geq (x + \varepsilon \mathbf{1})n^\alpha) - \mathbb{P}(\#(\sigma_n^{\text{Unif}}) \geq \varepsilon n^\alpha) = \exp(-n^\beta(J(x + \varepsilon \mathbf{1}) + o(1))) - o(\exp(-n^\beta M))$$

If $J(x + \varepsilon \mathbf{1})$ is infinite, then the right hand side will also be smaller than $\exp(-n^\beta M)$ for M as large as wished, whereas if it is finite, taking $M > J(x + \varepsilon \mathbf{1})$ also gives

$$\mathbb{P}(f(\sigma_n^{\text{Unif}}) \geq (x + \varepsilon \mathbf{1})n^\alpha) - \mathbb{P}(\#(\sigma_n^{\text{Unif}}) \geq \varepsilon n^\alpha) = \exp(-n^\beta(J(x + \varepsilon \mathbf{1}) + o(1))).$$

Similarly

$$\mathbb{P}(f(\sigma_n^{\text{Unif}}) \geq (x - \varepsilon \mathbf{1})n^\alpha) + \mathbb{P}(\#(\sigma_n^{\text{Unif}}) \geq \varepsilon n^\alpha) = \exp(-n^\beta J(x - \varepsilon \mathbf{1}) + o(n^\beta))(1 + o(1)).$$

We therefore conclude that, for every $\varepsilon > 0$,

$$-J(x + \varepsilon \mathbf{1}) \leq \liminf_{n \rightarrow \infty} \frac{1}{n^\beta} \ln \mathbb{P}(f(\sigma_{0,n}^{\text{Ew}}) \geq xn^\alpha) \leq \limsup_{n \rightarrow \infty} \frac{1}{n^\beta} \ln \mathbb{P}(f(\sigma_{0,n}^{\text{Ew}}) \geq xn^\alpha) \leq -J(x - \varepsilon \mathbf{1}).$$

Consequently, since we assumed that J is continuous, we find by letting ε going to zero

$$(11) \quad \liminf_{n \rightarrow \infty} \frac{1}{n^\beta} \ln \mathbb{P}(f(\sigma_{0,n}^{\text{Ew}}) \geq xn^\alpha) = \limsup_{n \rightarrow \infty} \frac{1}{n^\beta} \ln \mathbb{P}(f(\sigma_{0,n}^{\text{Ew}}) \geq xn^\alpha) = -J(x).$$

Now let σ_n be a conjugacy invariant permutation. Equation (10) implies (by choosing first to replace x by $x + \varepsilon \mathbf{1}$ then by $x - \varepsilon \mathbf{1}$) that

$$(12) \quad \mathbb{P}(f(\sigma_{0,n}^{\text{Ew}}) \geq (x + \varepsilon \mathbf{1})n^\alpha) - \mathbb{P}(\#(\sigma_n) \geq \varepsilon n^\alpha) \leq \mathbb{P}(f(\sigma_n) \geq xn^\alpha) \\ \leq \mathbb{P}(f(\sigma_{0,n}^{\text{Ew}}) \geq (x - \varepsilon \mathbf{1})n^\alpha) + \mathbb{P}(\#(\sigma_n) \geq \varepsilon n^\alpha).$$

Under hypothesis $(\mathcal{CI}_{\alpha,\beta})$, (11) implies that

$$(13) \quad \mathbb{P}(f(\sigma_{0,n}^{\text{Ew}}) \geq (x - \varepsilon \mathbf{1})n^\alpha) + \mathbb{P}(\#(\sigma_n) \geq \varepsilon n^\alpha) = \exp(-n^\beta J(x - \varepsilon \mathbf{1}) + o(n^\beta))(1 + o(1)),$$

and

$$(14) \quad \mathbb{P}(f(\sigma_{0,n}^{\text{Ew}}) \geq (x + \varepsilon \mathbf{1})n^\alpha) - \mathbb{P}(\#(\sigma_n) \geq \varepsilon n^\alpha) = \exp(-n^\beta J(x + \varepsilon \mathbf{1}) + o(n^\beta))(1 + o(1)).$$

Plugging these estimates in (12), letting n going to infinity and then ε going to zero (while using the continuity of J) completes the proof of Theorem 6 for permutations satisfying $(\mathcal{CI}_{\alpha,\beta})$. \square

2.4. Proof of Theorem 4. We first remark that it is enough to prove the Theorem with $E = [x, \infty)$ for $x \geq 2$ or $E = (0, x)$ if $x \leq 2$. Indeed, for $x > 2$, it is not hard to see that $I_{\text{LIS}, \frac{1}{2}}$ is strictly increasing so that the probability that LIS belongs to $[x, +\infty)$ is equivalent to the probability that it belongs to $[x, x + \delta]$ for any $\delta > 0$. Hence, (2) for $E = [x, \infty)$ and every $x \geq 2$ yields the weak large deviation principle. Exponential tightness is as well clear as $I_{\text{LIS}, \frac{1}{2}}$ goes to $+\infty$ at infinity. Hence, Corollary 9 implies the full large deviation principle above 2. Similarly, it is easy to see that $I_{\text{LIS}, 1}$ is strictly decreasing on $(0, 2)$ so that proving (3) for $E = (0, x]$ yields the full large deviation principle below 2. However, proving (3) for $E = (0, x]$ for $x < 2$ is more complicated because it is not possible to use Theorem 6 since $1 = \beta > \alpha = 1/2$. We need a proof specific to the longest increasing subsequence that we detail below. Before proving this result we prove a key lemma.

Lemma 18. *For every given permutation σ , for every integer number k ,*

$$\mathbb{P}(\text{LIS}(T(\sigma)) \leq \text{LIS}(\sigma) + k) \geq \inf_{\tau \in \mathcal{S}_{\#(\sigma)}} \mathbb{P}(\text{LIS}(\tau \circ \sigma_{0, \#(\sigma)}^{\text{Ew}}) < k)$$

Proof. The idea is to observe first that $\rho \in A_\sigma$ if and only if $\sigma^{-1} \circ \rho = (i_1, i_2, \dots, i_{\#(\sigma)})$ where i_j and i_k are in different cycles of σ as soon as $j \neq k$. This implies that one way to construct $T(\sigma)$ is to choose first uniformly, $j_1 < j_2 < \dots < j_{\#(\sigma)}$ each from one cycle of σ , π a uniform permutation of size $\#(\sigma)$. Then, it is easy to see that $T(\sigma)$ has the same law as

$$\sigma \circ (j_1, j_{\pi(1)}, j_{\pi^2(1)}, \dots, j_{\pi^{\#(\sigma)-1}(1)}).$$

Fix now $j_1, j_2, \dots, j_{\#(\sigma)}$ each on a cycle of σ and π a cyclic permutation and let $\ell_1 < \dots < \ell_{\text{LIS}(T(\sigma))}$ be such that $T(\sigma)(\ell_1) < \dots < T(\sigma)(\ell_{\text{LIS}(T(\sigma))})$ be a maximal increasing subsequence of $T(\sigma)$. Let

$$E = \{j_1, j_2, \dots, j_{\#(\sigma)}\} \cap \{\ell_1, \ell_2, \dots, \ell_{\text{LIS}(T(\sigma))}\} = \{j_{a_1}, j_{a_2}, j_{a_{\text{card}(E)}}\}$$

with $a_1 < a_2 < \dots < a_{\text{card}(E)}$ and let $F = \{\ell_1, \ell_2, \dots, \ell_{\text{LIS}(T(\sigma))}\} \setminus E$. For any $\ell_k \in F$, $T(\sigma)(\ell_k) = \sigma(\ell_k)$ and then $\text{card}(F) \leq \text{LIS}(\sigma)$. Let τ be the unique permutation of $\{1, \dots, \#(\sigma)\}$ such that

$$\sigma(j_{\tau^{-1}(1)}) < \sigma(j_{\tau^{-1}(2)}) < \dots < \sigma(j_{\tau^{-1}(\#(\sigma))}).$$

Moreover, $T(\sigma)(j_{a_k}) = \sigma(j_{\pi(a_k)})$ and then

$$\sigma(j_{\pi(a_1)}) < \dots < T(\sigma)(j_{\pi(a_{\text{card}(E)})})$$

In particular, the following holds true:

$$\text{card}(E) \leq \text{LIS}(\tau\pi).$$

Consequently, we find that

$$\text{LIS}(T(\sigma)) = \text{card}(E) + \text{card}(F) \leq \text{LIS}(\sigma) + \text{LIS}(\tau \circ \pi).$$

Now, applying this inequality we find that for any integer number k ,

$$\begin{aligned} \mathbb{P}(\text{LIS}(T(\sigma)) - \text{LIS}(\sigma) \leq k | \tau) &\geq \mathbb{P}(\text{LIS}(\tau \circ \pi) < k) \\ &= \mathbb{P}(\text{LIS}(\tau \circ \sigma_{0, \#(\sigma)}^{\text{Ew}}) < k) \geq \inf_{\tau' \in \mathcal{S}_{\#(\sigma)}} \mathbb{P}(\text{LIS}(\tau' \circ \sigma_{0, \#(\sigma)}^{\text{Ew}}) < k) \end{aligned}$$

where in the second line we used that π follows the uniform cyclic permutation of length $\#(\sigma)$. \square

To prove the second part of Theorem 4, we first show the result for the Ewens distribution with $\theta = 0$, namely that for every $x < 2$,

$$(15) \quad \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P}(\text{LIS}(\sigma_{0,n}^{\text{Ew}}) \leq x\sqrt{n}) = \limsup_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P}(\text{LIS}(\sigma_{0,n}^{\text{Ew}}) \leq x\sqrt{n}) = -I_{\text{LIS},1}(x).$$

It is then straightforward to generalize this result to $\sigma_n \in (\mathcal{CI}_{\frac{1}{2},1})$ as in the proof of Theorem 6 by taking $f = -\text{LIS}$, $\alpha = \frac{1}{2}$ and $\beta = 1$.

The upper bound is trivial since for any $k \in \mathbb{N}$,

$$\mathbb{P}(\text{LIS}(\sigma_{0,n}^{\text{Ew}}) = k) = \frac{\text{card}\{\sigma : \text{LIS}(\sigma) = k, \#(\sigma) = 1\}}{(n-1)!} \leq \frac{\text{card}\{\sigma : \text{LIS}(\sigma) = k\}}{(n-1)!} = n\mathbb{P}(\text{LIS}(\sigma_n^{\text{Unif}}) = k).$$

Consequently, by Theorem 3, we deduce

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P}(\text{LIS}(\sigma_{0,n}^{\text{Ew}}) \leq x\sqrt{n}) \leq \limsup_{n \rightarrow \infty} \frac{1}{n} (\ln \mathbb{P}(\text{LIS}(\sigma_n^{\text{Unif}}) \leq x\sqrt{n}) + \ln(n)) = -I_{\text{LIS},1}(x).$$

The lower bound is more sophisticated. Fix $x < 2$, $0 < \varepsilon < x$, and $\sigma \in \mathfrak{S}_n$. We assume that σ is such that $\text{LIS}(\sigma) < (x - 3\varepsilon)\sqrt{n}$ and $\#(\sigma) < \varepsilon^2 n$. Then, we find that

$$(16) \quad \begin{aligned} \mathbb{P}(\text{LIS}(T(\sigma)) \leq x\sqrt{n}) &= \mathbb{P}(\text{LIS}(T(\sigma)) \leq (x - 3\varepsilon)\sqrt{n} + 3\varepsilon\sqrt{n}) \\ &\geq \mathbb{P}(\text{LIS}(T(\sigma)) \leq \text{LIS}(\sigma) + 3\varepsilon\sqrt{n}) \end{aligned}$$

By Lemma 18, (16) gives

$$(17) \quad \mathbb{P}(\text{LIS}(T(\sigma)) \leq x\sqrt{n}) \geq \inf_{\tau \in \mathfrak{S}_{\#(\sigma)}} \mathbb{P}(\text{LIS}(\tau \circ \sigma_{0,\#(\sigma)}^{\text{Ew}}) < 3\varepsilon\sqrt{n}).$$

To conclude, we need the following easy lemma.

Lemma 19. *For any $x > 2$,*

$$\lim_{n \rightarrow \infty} \inf_{\tau \in \mathfrak{S}_n} \mathbb{P}(\text{LIS}(\tau \circ \sigma_{0,n}^{\text{Ew}}) < x\sqrt{n}) = 1$$

Proof. The distribution of $\tau \circ \sigma_{0,n}^{\text{Ew}}$ is uniform over a subset of size exactly $(n-1)!$ of \mathfrak{S}_n (the image S_τ by τ of the cyclic permutations). This set has probability $1/n$ for σ_n^{Unif} . We have then

$$\mathbb{P}(\text{LIS}(\tau \circ \sigma_{0,n}^{\text{Ew}}) \geq x\sqrt{n}) = n\mathbb{P}(\{\text{LIS}(\sigma_n^{\text{Unif}}) \geq x\sqrt{n}\} \cap \{\sigma_n^{\text{Unif}} \in S_\tau\}) \leq n\mathbb{P}(\{\text{LIS}(\sigma_n^{\text{Unif}}) \geq x\sqrt{n}\}) = o(1)$$

because of Theorem 3 and $x > 2$. Note that these bounds do not depend on τ . Therefore

$$\mathbb{P}(\text{LIS}(\tau \circ \sigma_{0,n}^{\text{Ew}}) < x\sqrt{n}) = 1 - \mathbb{P}(\text{LIS}(\tau \circ \sigma_{0,n}^{\text{Ew}}) \geq x\sqrt{n}) = 1 - o(1).$$

□

Consequently, for any $x < 2$ and $\varepsilon > 0$, and σ_n conjugacy invariant, we have

$$\begin{aligned}
\mathbb{P}(\text{LIS}(\sigma_{0,n}^{\text{Ew}}) \leq x\sqrt{n}) &= \sum_{\sigma \in \mathfrak{S}_n} \mathbb{P}(\text{LIS}(T(\sigma)) \leq x\sqrt{n})\mathbb{P}(\sigma_n = \sigma) \\
&\geq \sum_{\text{LIS}(\sigma) \leq (x-3\varepsilon)\sqrt{n}, \#(\sigma) < \varepsilon^2 n} \inf_{\tau \in \mathfrak{S}_{\#(\sigma)}} \mathbb{P}(\text{LIS}(\tau \circ \sigma_{0,k}^{\text{Ew}}) < 3\varepsilon\sqrt{n})\mathbb{P}(\sigma_n = \sigma) \\
&\geq \inf_{k \leq \varepsilon^2 n} \inf_{\tau \in \mathfrak{S}_k} \mathbb{P}(\text{LIS}(\tau \circ \sigma_{0,k}^{\text{Ew}}) < 3\varepsilon\sqrt{n})\mathbb{P}(\text{LIS}(\sigma_n) \leq (x-3\varepsilon)\sqrt{n}, \#(\sigma_n) < \varepsilon^2 n) \\
(18) \quad &\geq (1+o(1))\mathbb{P}(\text{LIS}(\sigma_n) \leq (x-3\varepsilon)\sqrt{n}, \#(\sigma_n) < \varepsilon^2 n)
\end{aligned}$$

In the first line we used Lemma 17, in the second we restricted the summation over σ and bounded uniformly from below the first probability by using (17), in the third line we summed the second probability over the remaining permutations, and in the last line we used Lemma 19 when k goes to infinity (noticing that $\varepsilon\sqrt{n}/\sqrt{k} \geq 1$ for n large enough), while the bound is clear when k is finite. Now choose σ_n to be uniform. We deduce from (16) that

$$\begin{aligned}
\mathbb{P}(\text{LIS}(\sigma_{0,n}^{\text{Ew}}) \leq x\sqrt{n}) &\geq (1+o(1))\mathbb{P}(\text{LIS}(\sigma_n^{\text{Unif}}) \leq (x-3\varepsilon)\sqrt{n}, \#(\sigma_n^{\text{Unif}}) < \varepsilon^2 n) \\
&= (1+o(1))(\mathbb{P}(\text{LIS}(\sigma_n^{\text{Unif}}) \leq (x-3\varepsilon)\sqrt{n}) \\
&\quad - \mathbb{P}(\text{LIS}(\sigma_n^{\text{Unif}}) \leq (x-3\varepsilon)\sqrt{n}, \#(\sigma_n^{\text{Unif}}) \geq \varepsilon^2 n)) \\
&\geq (1+o(1))\mathbb{P}(\text{LIS}(\sigma_n^{\text{Unif}}) \leq (x-3\varepsilon)\sqrt{n}) = e^{-nI_{\text{LIS}, \frac{1}{2}}(x-\varepsilon)}(1+o(1))
\end{aligned}$$

where we used Theorem 3 and that σ_n^{Unif} is $(\mathcal{CI}_{\alpha,\alpha})$. We finally can let n going to infinity and ε going to zero to get (15). To complete the proof of Theorem 4, one only needs to check that

- If the sequence $(\sigma_n)_{n \geq 1}$ of random permutations satisfies $(\mathcal{CI}_{\frac{1}{2},1})$, for any $\varepsilon > 0$, for any $x \geq 2$.

$$\lim_{n \rightarrow \infty} \frac{1}{n} \ln \left(\mathbb{P} \left(\text{LIS}(\sigma_n) \in \left((x-\varepsilon)\sqrt{n}, (x+\varepsilon)\sqrt{n} \right) \right) \right) = 0.$$

Since $(\mathcal{CI}_{\frac{1}{2},1})$ implies $(\mathcal{CI}_{\frac{1}{2},\frac{1}{2}})$, this is a direct consequence of Corollary 9.

- If the sequence $(\sigma_n)_{n \geq 1}$ satisfies $(\mathcal{CI}_{\frac{1}{2},\frac{1}{2}})$, for any $0 < y < 2$.

$$\lim_{n \rightarrow \infty} \frac{\ln(\mathbb{P}(\text{LIS}(\sigma_n) < y\sqrt{n}))}{\sqrt{n}} = -\infty$$

Indeed, by taking $f = -\text{LIS}$, $\varepsilon = (2-y)/2$, $x = -(y+\varepsilon)$ and $\alpha = \beta = \frac{1}{2}$ in (10), the first inequality becomes

$$\mathbb{P}(\text{LIS}(\sigma_n) \leq y\sqrt{n}) \leq \mathbb{P}(\#(\sigma_n) \geq \varepsilon\sqrt{n}) + \mathbb{P}(\text{LIS}(\sigma_{0,n}^{\text{Ew}}) \leq x\sqrt{n}).$$

The first term goes to zero faster than $e^{-M\sqrt{n}}$ for any $M > 0$, since the sequence $(\sigma_n)_{n \geq 1}$ satisfies $(\mathcal{CI}_{\frac{1}{2},\frac{1}{2}})$, whereas the second term goes to zero

2.5. Proof of Proposition 5. For the upper tail, one can apply directly Corollary 9.

For the lower bound of the lower tail, let $\theta > 0$.

$$\begin{aligned} \mathbb{P}(\text{LIS}(\sigma_{\theta,n}^{\text{Ew}}) = k) &\geq \mathbb{P}(\text{LIS}(\sigma_{\theta,n}^{\text{Ew}}) = k, \#\sigma_{\theta,n}^{\text{Ew}} = 1) \\ &= \frac{\text{card}\{\sigma : \text{LIS}(\sigma) = k, \#(\sigma) = 1\} \Gamma(\theta)}{\Gamma(n + \theta)} \\ &= \frac{\Gamma(\theta) \Gamma(n)}{\Gamma(n + \theta)} \mathbb{P}(\text{LIS}(\sigma_{0,n}^{\text{Ew}}) = k) \geq \frac{\Gamma(\theta)}{(n + \theta)^\theta} \mathbb{P}(\text{LIS}(\sigma_{0,n}^{\text{Ew}}) = k). \end{aligned}$$

This implies by taking $k \leq x\sqrt{n}$,

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P}(\text{LIS}(\sigma_{\theta,n}^{\text{Ew}}) \leq x\sqrt{n}) \geq \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P}(\text{LIS}(\sigma_{0,n}^{\text{Ew}}) \leq x\sqrt{n}) = -I_{\text{LIS}, \frac{1}{2}}(x).$$

For the upper bound, one can conclude directly by (18), by choosing σ_n to be $\sigma_{\theta,n}^{\text{Ew}}$.

2.6. Proof of Proposition 13. We will adapt the proof of the lower tail of Theorem 4. We choose to give two different separate proofs for readability reasons.

First remark that Proposition 13 is equivalent to that

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P}\left(\forall j \in [d], \sum_{i=1}^j \lambda_i(\sigma_n) \leq \sum_{i=1}^j x_i \sqrt{n}\right) = -I_{\text{LIS}, 1}(x_d).$$

Indeed, because the right hand side depends only on x_d , we see that mostly the deviations of $\lambda_d(\sigma_n)$ matters. The same phenomenon appears for random matrices: the probability fix the d largest eigenvalues to make a deviation below 2 is equivalent to the probability that the spectrum stays below the smallest one, namely x_d .

We recall that according to Green's theorem ([45, Theorem 3.5.3]) $\lambda_1 + \dots, \lambda_k$ is the maximum sum of lengths of k disjoint increasing subsequences.

The counterpart of Lemma 18 is that for every σ deterministic, for every integer numbers d, k ,

$$(19) \quad \mathbb{P}\left(\sum_{i=1}^d \lambda_i(T(\sigma)) \leq \left(\sum_{i=1}^d \lambda_i(\sigma)\right) + k\right) \geq \inf_{\tau \in \mathfrak{S}_{\#(\sigma)}} \mathbb{P}\left(\sum_{i=1}^d \lambda_i(\tau \circ \sigma_{0, \#(\sigma)}^{\text{Ew}}) < k\right).$$

As in the proof of Theorem 4, we want to prove that

$$(20) \quad \begin{aligned} \liminf_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P}\left(\forall j \in [d], \sum_{i=1}^j \lambda_i(\sigma_{\theta,n}^{\text{Ew}}) \leq \sum_{i=1}^j x_i \sqrt{n}\right) &= \limsup_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P}\left(\forall j \in [d], \sum_{i=1}^j \lambda_i(\sigma_{0,n}^{\text{Ew}}) \leq \sum_{i=1}^j x_i \sqrt{n}\right) \\ &= -I_{\text{LIS}, 1}(x_d). \end{aligned}$$

It is then straightforward to generalize this result to $\sigma_n \in (\mathcal{CI}_{\frac{1}{2}, 1})$ as in the proof of Theorem 6 by taking $f = (\lambda_1, \lambda_1 + \lambda_2, \dots, \sum_{j=1}^d \lambda_j)$, $\alpha = \frac{1}{2}$ and $\beta = 1$.

The upper bound is trivial for the same reason that in the proof of Theorem 4 (the probability of any event under Ewens with parameter 0 is at most n times its probability under the uniform permutation).

The lower bound is more sophisticated. Fix $0 < x_d < \dots < x_1 < 2$, $0 < \varepsilon < x_d$, and $\sigma \in S_n$. We assume that σ is such that for any j , $\sum_{i=1}^j \lambda_i(\sigma) < ((\sum_{i=1}^j x_i) - 3\varepsilon)\sqrt{n}$ and $\#(\sigma) < \frac{\varepsilon^2 n}{d^2}$. Then, we find that

$$(21) \quad \begin{aligned} \mathbb{P}\left(\forall j \in [d], \sum_{i=1}^j \lambda_i(T(\sigma)) \leq \sum_{i=1}^j x_i \sqrt{n}\right) &= \mathbb{P}\left(\forall i \in [d], \sum_{i=1}^j \lambda_i(T(\sigma)) \leq \left(\left(\sum_{i=1}^j x_i\right) - 3\varepsilon\right)\sqrt{n} + 3\varepsilon\sqrt{n}\right) \\ &\geq \mathbb{P}\left(\forall j \in [d], \sum_{i=1}^j \lambda_i(T(\sigma)) \leq \left(\sum_{i=1}^j \lambda_i(\sigma)\right) + 3\varepsilon\sqrt{n}\right) \end{aligned}$$

By (19), (21) gives

$$(22) \quad \mathbb{P}\left(\forall j \in [d], \sum_{i=1}^j \lambda_i(T(\sigma)) \leq \sum_{i=1}^j x_i \sqrt{n}\right) \geq \inf_{\tau \in \mathfrak{S}_{\#(\sigma)}} \mathbb{P}\left(\forall j \in [d], \sum_{i=1}^j \lambda_i(\tau \circ \sigma_{0, \#(\sigma)}^{\text{Ew}}) < 3\varepsilon\sqrt{n}\right).$$

To conclude, we need the following easy lemma.

Lemma 20. *For any $x > 2$ and any integer number j ,*

$$\lim_{n \rightarrow \infty} \inf_{\tau \in \mathfrak{S}_n} \mathbb{P}\left(\sum_{i=1}^j \lambda_i(\tau \circ \sigma_{0, n}^{\text{Ew}}) < jx\sqrt{n}\right) = 1$$

Proof. First we have,

$$\lim_{n \rightarrow \infty} \inf_{\tau \in \mathfrak{S}_n} \mathbb{P}\left(\sum_{i=1}^j \lambda_i(\tau \circ \sigma_{0, n}^{\text{Ew}}) < jx\sqrt{n}\right) \geq \lim_{n \rightarrow \infty} \inf_{\tau \in \mathfrak{S}_n} \mathbb{P}(j \text{LIS}(\tau \circ \sigma_{0, n}^{\text{Ew}}) < jx\sqrt{n})$$

because of the non-increasing of λ_i . Therefore, one can conclude by Lemma 19 that

$$\lim_{n \rightarrow \infty} \inf_{\tau \in \mathfrak{S}_n} \mathbb{P}(j \text{LIS}(\tau \circ \sigma_{0, n}^{\text{Ew}}) < jx\sqrt{n}) = \mathbb{P}(\text{LIS}(\tau \circ \sigma_{0, n}^{\text{Ew}}) < x\sqrt{n}) = 1 + o(1).$$

□

Consequently, for any σ_n conjugacy invariant, similarly to (18), we obtain

$$(23) \quad \begin{aligned} \mathbb{P}\left(\forall j \in [d], \sum_{i=1}^j \lambda_i(\sigma_{0, n}^{\text{Ew}}) \leq \sum_{i=1}^j x_i \sqrt{n}\right) \\ \geq (1 + o(1)) \mathbb{P}\left(\forall j \in [d], \sum_{i=1}^j \lambda_i(\sigma_n) \leq \left(\left(\sum_{i=1}^j x_i\right) - 3\varepsilon\right)\sqrt{n}, \#(\sigma_n) < \frac{\varepsilon^2 n}{d^2}\right). \end{aligned}$$

The remaining of the proof is identical to that of the lower tail of Theorem 4.

3. FURTHER DISCUSSIONS

We start by discussing the optimality of our conditions. The condition $(\mathcal{CI}_{\frac{1}{2}, \frac{1}{2}})$ for LIS is optimal as a condition on cycles. Indeed, let us construct a sequence of random permutations σ_n such that $\#(\sigma_n)/\sqrt{n}$ is of order $x \neq 0$ with probability of order $e^{\sqrt{n}C}$ for some C finite and show that our result does not apply to σ_n . In fact, let σ_n be a permutation constructed

as follows. With probability $e^{-(I_{\text{LIS},1/2}(x)-\varepsilon)\sqrt{n}}$, σ_n has $\lfloor x\sqrt{n} \rfloor$ fixed points, the other points belonging to a cycle of length $n - \lfloor x\sqrt{n} \rfloor$. With probability $1 - e^{-(I_{\text{LIS},1/2}(x)-\varepsilon)\sqrt{n}}$, σ_n is a uniform cyclic permutation. In this case, it is not difficult to check that since the fixed points furnish an increasing subsequence

$$\lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} \ln \mathbb{P}(\text{LIS}(\sigma_n) \geq x\sqrt{n}) \geq -I_{\text{LIS},1/2}(x) + \varepsilon > -I_{\text{LIS},1/2}(x),$$

showing that $\text{LIS}(\sigma_n)$ can not follow the same large deviation principle than $\text{LIS}(\sigma_n^{\text{Unif}})$ as stated in Theorem 3.

When the number of cycles is not sufficiently controlled, it is only possible to obtain one bound.

Proposition 21. *If $(\sigma_n)_{n \geq 1}$ satisfies $(\mathcal{CI}_{1,1})$, then for any $0 < x < 2$,*

$$(24) \quad \limsup_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P}(\text{LIS}(\sigma_n) \leq x\sqrt{n}) \leq -I_{\text{LIS},1}(x).$$

Moreover, if $(\sigma_n)_{n \geq 1}$ satisfies that for any $\varepsilon > 0$

$$\liminf_{n \rightarrow \infty} \mathbb{P}\left(\frac{\#(\sigma_n)}{n} < \varepsilon\right) > 0,$$

then for any $x > 2$.

$$\liminf_{n \rightarrow \infty} \frac{1}{\sqrt{n}} \ln \mathbb{P}(\text{LIS}(\sigma_n) \geq x\sqrt{n}) \geq -I_{\text{LIS},\frac{1}{2}}(x).$$

Sketch of the proof of Proposition 21. We will compare directly $\text{LIS}(\sigma_n)$ to $\text{LIS}(\sigma_{0,n}^{\text{Ew}})$. For the first inequality, (18) and (3) imply that

$$(25) \quad e^{-nI_{\text{LIS},1}(x)(1+o(1))} \geq \mathbb{P}(\text{LIS}(\sigma_{0,n}^{\text{Ew}}) \leq x\sqrt{n}) \\ \geq (1+o(1))\mathbb{P}(\text{LIS}(\sigma_n) \leq (x-3\varepsilon)\sqrt{n}, \#(\sigma_n) < \varepsilon^2 n).$$

Moreover, if σ_n satisfies $(\mathcal{CI}_{1,1})$, for every $M > 0$ and n large enough,

$$\mathbb{P}(\text{LIS}(\sigma_n) \leq (x-3\varepsilon)\sqrt{n}, \#(\sigma_n) < \varepsilon^2 n) \geq \mathbb{P}(\text{LIS}(\sigma_n) \leq (x-3\varepsilon)\sqrt{n}) - e^{-nM}.$$

which gives (24) with (25).

The second inequality is an application of [32, Lemma 21]. First remark that a conjugacy invariant permutation conditioned on $\#(\sigma_n) \leq k$ is still conjugacy invariant. One needs only to prove this result for permutations where the number of cycles is less than εn for $n \geq n_0$ almost surely.

By choosing ρ to be $\sigma_{0,n}^{\text{Ew}}$ and $k = 1$ in [32, Lemma 21], one can reformulate the lemma to obtain that for any conjugacy invariant permutation σ_n , there exists $\widehat{\sigma}_n \stackrel{d}{=} \sigma_n$ such that, for any $1 \leq a < b$ and for any $c > 0$

$$\mathbb{E}\left(\left(\text{LIS}(\widehat{\sigma}_n) - \text{LIS}(\sigma_{0,n}^{\text{Ew}})\right)_- \mid \#\widehat{\sigma}_n < c, \text{LIS}(\sigma_{0,n}^{\text{Ew}}) \in [a, b]\right) \leq \frac{cb}{n}.$$

By Markov inequality, and by taking $a = x\sqrt{n} + 4x\varepsilon\sqrt{n}$ and $b = 2x\sqrt{n}$ and $c = \varepsilon n$ we obtain that

$$\begin{aligned}
\mathbb{P}\left(\text{LIS}(\sigma_n) \geq x\sqrt{n}\right) &= \mathbb{P}\left(\text{LIS}(\widehat{\sigma}_n) \geq x\sqrt{n}\right) \\
&\geq \mathbb{P}\left(\text{LIS}(\sigma_{0,n}^{\text{Ew}} \in [a, b], \left(\text{LIS}(\widehat{\sigma}_n) - \text{LIS}(\sigma_{0,n}^{\text{Ew}})\right)_- \leq 4x\varepsilon\sqrt{n}\right) \\
&= \mathbb{P}\left(\left(\text{LIS}(\widehat{\sigma}_n) - \text{LIS}(\sigma_{0,n}^{\text{Ew}})\right)_- \leq 4x\varepsilon\sqrt{n} \mid \text{LIS}(\sigma_{0,n}^{\text{Ew}} \in [a, b])\right) \mathbb{P}(\text{LIS}(\sigma_{0,n}^{\text{Ew}} \in [a, b]) \\
&\geq \left(1 - \frac{\frac{cb}{n}}{4x\varepsilon\sqrt{n}}\right) I_{\text{LIS}, \frac{1}{2}} \mathbb{P}(\text{LIS}(\sigma_{0,n}^{\text{Ew}} \in [a, b]) = \frac{1}{2} \mathbb{P}(\text{LIS}(\sigma_{0,n}^{\text{Ew}} \in [a, b])
\end{aligned}$$

which concludes the proof. \square

Acknowledgement Both authors are supported by ERC Project LDRAM (ERC-2019-ADG Project: 884584). M.S.K. is partially supported by LabEx CIMI. He would also acknowledge a useful discussion with Christian Houdré and Valentin Féray.

REFERENCES

- [1] R. Arratia, A. D. Barbour, and S. Tavaré. Exploiting the Feller coupling for the Ewens sampling formula. *Statist. Sci.*, 31(1):27–29, 2016.
- [2] F. Augeri. Large deviations principle for the largest eigenvalue of Wigner matrices without Gaussian tails. *Electron. J. Probab.*, 21:Paper No. 32, 49, 2016.
- [3] F. Augeri, A. Guionnet, and J. Husson. Large deviations for the largest eigenvalue of sub-Gaussian matrices. *Comm. Math. Phys.*, 383(2):997–1050, 2021.
- [4] V. Bahier and J. Najnudel. On smooth mesoscopic linear statistics of the eigenvalues of random permutation matrices. *J. Theor. Probab.*, 35(3):1640–1661, 2022.
- [5] J. Baik, P. Deift, and K. Johansson. On the distribution of the length of the longest increasing subsequence of random permutations. *J. Amer. Math. Soc.*, 12(4):1119–1178, 1999.
- [6] J. Baik and E. M. Rains. The asymptotics of monotone subsequences of involutions. *Duke Math. J.*, 109(2):205–281, 2001.
- [7] G. Ben Arous and K. Dang. On fluctuations of eigenvalues of random permutation matrices. *Ann. Inst. Henri Poincaré, Probab. Stat.*, 51(2):620–647, 2015.
- [8] G. Ben Arous, A. Dembo, and A. Guionnet. Aging of spherical spin glasses. *Probab. Theory Related Fields*, 120(1):1–67, 2001.
- [9] G. Ben Arous and A. Guionnet. Large deviations for Wigner’s law and Voiculescu’s non-commutative entropy. *Probab. Theory Relat. Fields*, 108(4):517–542, 1997.
- [10] B. Bercu, M. Bonnefont, and A. Richou. Sharp large deviations and concentration inequalities for the number of descents in a random permutation, 2022.
- [11] G. Biroli and A. Guionnet. Large deviations for the largest eigenvalues and eigenvectors of spiked Gaussian random matrices. *Electron. Commun. Probab.*, 25:Paper No. 70, 13, 2020.
- [12] S. Boucheron, G. Lugosi, and P. Massart. *Concentration Inequalities: A Nonasymptotic Theory of Independence*. Oxford University Press, 02 2013.
- [13] T. Budzinski, N. Curien, and B. Petri. Universality for random surfaces in unconstrained genus. *Electron. J. Comb.*, 26(4):research paper p4.2, 35, 2019.
- [14] D. Chafaï, Y. Doumerc, and F. Malrieu. Processus des restaurants chinois et loi d’Ewens. *Revue de Mathématiques Spéciales (RMS)*, 123(3):56–74, Mar. 2013.
- [15] N. Cook, R. Ducatez, and A. Guionnet. Full large deviation principles for the largest eigenvalue of sub-gaussian wigner matrices. *arXiv:2302.14823*, 2023.
- [16] A. Dembo and O. Zeitouni. *Large deviations techniques and applications*, volume 38 of *Applications of Mathematics (New York)*. Springer-Verlag, New York, second edition, 1998.
- [17] J.-D. DEUSCHEL and O. ZEITOUNI. On increasing subsequences of i.i.d. samples. *Combinatorics, Probability and Computing*, 8(3):247–263, 1999.
- [18] I. Dumitriu and S. Pal. Sparse regular random graphs: spectral density and eigenvectors. *Ann. Probab.*, 40(5):2197–2235, 2012.

- [19] L. Erdős and H.-T. Yau. Universality of local spectral statistics of random matrices. *Bull. Amer. Math. Soc. (N.S.)*, 49(3):377–414, 2012.
- [20] W. J. Ewens. The sampling theory of selectively neutral alleles. *Theoret. Population Biology*, 3:87–112; erratum, *ibid.* 3 (1972), 240; erratum, *ibid.* 3 (1972), 376, 1972.
- [21] V. Féray. Asymptotic behavior of some statistics in Ewens random permutations. *Electron. J. Probab.*, 18(76):1–32, 2013.
- [22] J. Fulman, G. B. Kim, and S. Lee. Central limit theorem for peaks of a random permutation in a fixed conjugacy class of S_n . *Annals of Combinatorics*, 26(1):97–123, Dec. 2021.
- [23] S. Ganguly, E. Hiesmayr, and K. Nam. Spectral large deviations of sparse random matrices. *arXiv 2206.06954*, 2022.
- [24] A. Guionnet and J. Husson. Large deviations for the largest eigenvalue of Rademacher matrices. *Ann. Probab.*, 48(3):1436–1465, 2020.
- [25] A. Guionnet and M. Maïda. Large deviations for the largest eigenvalue of the sum of two random matrices. *Electron. J. Probab.*, 25:Paper No. 14, 24, 2020.
- [26] Z. Hamaker and B. Rhoades. Characters of local and regular permutation statistics, 2022.
- [27] C. Houdré and R. Restrepo. A probabilistic approach to the asymptotics of the length of the longest alternating subsequence. *Electron. J. Combin.*, 17(1):Research Paper 168, 19, 2010.
- [28] K. Johansson. Shape fluctuations and random matrices. *Comm. Math. Phys.*, 209(2):437–476, 2000.
- [29] K. Johansson. Discrete orthogonal polynomial ensembles and the Plancherel measure. *Ann. Math. (2)*, 153(1):259–296, 2001.
- [30] M. S. Kammoun. Monotonous subsequences and the descent process of invariant random permutations. *Electron. J. Probab.*, 23:31 pp., 2018.
- [31] M. S. Kammoun. On the Longest Common Subsequence of Conjugation Invariant Random Permutations. *The Electronic Journal of Combinatorics*, 27, 2020.
- [32] M. S. Kammoun. Universality for random permutations and some other groups. *Stochastic Processes Appl.*, 147:76–106, 2022.
- [33] M. S. Kammoun and M. Maïda. A product of invariant random permutations has the same small cycle structure as uniform. *Electron. Commun. Probab.*, 25:14, 2020. Id/No 57.
- [34] G. Kim and S. Lee. Central limit theorem for descents in conjugacy classes of S_n . *J. Combin. Th., Series A*, 169:105–123, 2020.
- [35] J. F. C. Kingman. Random partitions in population genetics. *Proc. Roy. Soc. London Ser. A*, 361(1704):1–20, 1978.
- [36] P.-S. Laplace. *Théorie analytique des probabilités. Vol. II. Livre II: Théorie générale des probabilités. Suppléments.* Paris: Éditions Jacques Gabay, reprint of the 1820 third edition (Book II) and of the 1816, 1818, 1820 and 1825 originals edition, 1995.
- [37] B. F. Logan and L. A. Shepp. A variational problem for random Young tableaux. *Advances in Math.*, 26(2):206–222, 1977.
- [38] M. Löwe and F. Merkl. Moderate deviations for longest increasing subsequences: the upper tail. *Comm. Pure Appl. Math.*, 54(12):1488–1520, 2001.
- [39] M. Maïda. Large deviations for the largest eigenvalue of rank one deformations of Gaussian ensembles. *Electron. J. Probab.*, 12:1131–1150, 2007.
- [40] S. N. Majumdar and G. Schehr. Top eigenvalue of a random matrix: large deviations and third order phase transition. *J. Stat. Mech. Theory Exp.*, (1):P01012, 31, 2014.
- [41] P.-L. Méliot and A. Nikeghbali. Asymptotics of the major index of a random standard tableau, 2022.
- [42] R. G. Pinsky. Large deviations for the longest alternating and the longest increasing subsequence in a random permutation avoiding a pattern of length three, 2023.
- [43] D. Romik. Local extrema in random permutations and the structure of longest alternating subsequences. In *23rd International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2011)*, Discrete Math. Theor. Comput. Sci. Proc., AO, pages 825–834. Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2011.
- [44] D. Romik. *The surprising mathematics of longest increasing subsequences*, volume 4 of *Institute of Mathematical Statistics Textbooks*. Cambridge University Press, New York, 2015.
- [45] B. E. Sagan. *The symmetric group*, volume 203 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 2001. Representations, combinatorial algorithms, and symmetric functions.

- [46] T. Seppäläinen. Large deviations for increasing sequences on the plane. *Probab. Theory Related Fields*, 112(2):221–244, 1998.
- [47] T. Tao and V. Vu. Random matrices: The universality phenomenon for Wigner ensembles. In *Modern aspects of random matrix theory. Based on lectures delivered at the 2013 AMS short course on random matrices, San Diego, CA, USA, January 6–7, 2013*, pages 121–172. Providence, RI: American Mathematical Society (AMS), 2014.
- [48] C. A. Tracy and H. Widom. Level-spacing distributions and the Airy kernel. *Commun. Math. Phys.*, 159(1):151–174, 1994.
- [49] N. Tsilevich. Stationary measures on the space of virtual permutations for an action of the infinite symmetric group. 1998.
- [50] S. M. Ulam. Monte Carlo calculations in problems of mathematical physics. In *Modern mathematics for the engineer: Second series*, pages 261–281. McGraw-Hill, New York, 1961.
- [51] A. M. Vershik and S. V. Kerov. Asymptotic behavior of the Plancherel measure of the symmetric group and the limit form of Young tableaux. *Dokl. Akad. Nauk SSSR*, 233(6):1024–1027, 1977.
- [52] E. P. Wigner. Characteristic vectors of bordered matrices with infinite dimensions. *Ann. Math. (2)*, 62:548–564, 1955.

Email address: `alice.guionnet@ens-lyon.fr`

Email address: `slim.kammoun@ens-lyon.fr`