# Lih Wang and Dittert's Conjectures on **Permanents**

Divya.K.U and K. Somasundaram

Department of Mathematics, Amrita School of Physical Sciences, Coimbatore Amrita Vishwa Vidyapeetham, India. divyaku93@gmail.com, s sundaram@cb.amrita.edu

#### Abstract

Let  $\Omega_n$  denote the set of all doubly stochastic matrices of order n. Lih and Wang conjectured that for  $n \geq 3$ , per $(tJ_n + (1-t)A) \leq t$  per $J_n + (1-t)$  per A, for all  $A \in \Omega_n$ and all  $t \in [0.5, 1]$ , where  $J_n$  is the  $n \times n$  matrix with each entry equal to  $\frac{1}{n}$ . This conjecture was proved partially for  $n \leq 5$ .

Let  $K_n$  denote the set of non-negative  $n \times n$  matrices whose elements have sum n. Let  $\phi$  be a real valued function defined on  $K_n$  by  $\phi(X) = \prod_{i=1}^n r_i + \prod_{j=1}^n c_j$  - per X for  $X \in K_n$  with row sum vector  $(r_1, r_2, ... r_n)$  and column sum vector  $(c_1, c_2, ... c_n)$ . A matrix  $A \in K_n$  is called a  $\phi$ -maximizing matrix if  $\phi(A) \geq \phi(X)$  for all  $X \in K_n$ . Dittert conjectured that  $J_n$  is the unique  $\phi$ -maximizing matrix on  $K_n$ . Sinkhorn proved the conjecture for  $n = 2$  and Hwang proved it for  $n = 3$ .

In this paper, we prove the Lih and Wang conjecture for  $n = 6$  and Dittert conjecture for  $n = 4$ .

**Keywords:**Permanents, Doubly Stochastic Matrices, Lih-Wang Conjecture,  $\phi$ -maximizing matrix, Dittert's conjecture.

AMS Subject Classification: 15A15.

## 1 Introduction

The permanent of an  $n \times n$  matrix  $A = (a_{ij})$  is defined by

$$
\text{per}(A) = \sum_{\sigma \in S_n} a_{1\sigma(1)} a_{2\sigma(2)} \dots a_{n\sigma(n)},
$$

where  $S_n$  is the symmetric group of degree n.

Let  $\Omega_n$  denote the set of all doubly stochastic matrices of order n and let  $J_n$  be an  $n \times n$ matrix with each entry equal to  $\frac{1}{n}$ .

For positive integers n and k with  $(1 \le k \le n)$ ,  $Q_{k,n}$  denotes the set  $\{(i_1, ..., i_k)/1 \leq i_1 < ... < i_k \leq n\}$ . For  $\alpha, \beta \in Q_{k,n}$ , let  $A(\alpha/\beta)$  be the submatrix of A obtained by deleting the rows indexed by  $\alpha$  and columns indexed by  $\beta$  and  $A[\alpha/\beta]$  be the submatrix of A with rows and columns indexed by  $\alpha$  and  $\beta$  respectively.

For  $1 \leq k \leq n$ , the  $k^{th}$  order subpermanent of A is defined by

$$
\sigma_k(A) = \sum_{\alpha,\beta \in Q_{k,n}} \text{per}\left(A[\alpha/\beta]\right)
$$

In this paper, we use the following results quoted by Minc [\[25\]](#page-14-0): If A and B are two  $n \times n$  matrices and  $1 \leq k \leq n$ , then

$$
\text{per}(A) = \sum_{\beta \in Q_{k,n}} \text{per}(A[\alpha/\beta]) \text{ per}(A(\alpha/\beta)), \text{ for } \alpha \in Q_{k,n},
$$

and

$$
per(A + B) = \sum_{k=0}^{n} S_k (A, B),
$$

where  $S_k(A, B) = \sum$  $\alpha,\beta \in Q_{k,n}$  $\text{per}(A[\alpha/\beta])\text{per}(B(\alpha/\beta)), \text{per}(A[\alpha/\beta]) = 1$  when  $k = 0$ and per $(B(\alpha/\beta)) = 1$  when  $k = n$ .

Several authors have considered the problem of finding an upper bound for the permanent of a convex combination of  $J_n$  and A, where  $A \in \Omega_n$ . In particular, Marcus and Minc [\[13\]](#page-13-0) have proposed a conjecture, which states that if  $A \in \Omega_n, n \geq 2$ , then  $\text{per}(\frac{nJ_n-A}{n-1})$  $\frac{J_n-A}{n-1}\leq \text{per}(A),$ with equality holding if and only if  $A = J_n$ ,  $n \geq 3$ . They established that the conjecture is true for  $n = 2$ , or if A is a positive semi-definite symmetric matrix, or if A is in a sufficiently small neighbourhood of  $J_n$ . Wang [\[16\]](#page-13-1) proved the Marcus - Minc conjecture for  $n = 3$  with a revised statement of the case of equality and he proposed a conjecture that  $\text{per}(\frac{nJ_n+A}{n+1}) \leq \text{per}(A)$ .

Rather than consider a particular convex combination of A with  $J_n$ , Foregger [\[4\]](#page-13-2) raised a question whether

$$
\operatorname{per}\left(tJ_n + (1-t)A\right) \le \operatorname{per}(A) \tag{1}
$$

holds for all  $A \in \Omega_n$  and  $0 < t \leq \frac{n}{n-1}$  $\frac{n}{n-1}$ ? He proved that the inequality (1) holds for  $0 < t \leq \frac{3}{2}$ 2 and  $A \in \Omega_3$  with equality if and only if  $A = J_3$  or  $t = \frac{3}{2}$  $\frac{3}{2}$  and  $A = \frac{1}{2}$  $rac{1}{2}(I+P)$ 

(up to permutations of rows and columns), where  $P$  is a full cycle permutation matrix. Later, Foregger himself proved in [\[5\]](#page-13-3) that the inequality (1) holds for  $n = 4$  and  $t_0 < t \leq \frac{4}{3}$  $\frac{4}{3}$ , where  $t_0$ is approximately equal to 0.28095.

Another conjecture in permanents is due to Holens [\[8\]](#page-13-4) and Dokovic [\[2\]](#page-13-5), namely the Holens - Dokovic conjecture, which states that if  $A \in \Omega_n$  and k is an integer,  $1 \leq k \leq n$ , then

$$
\sigma_k(A) \ge \frac{(n-k+1)^2}{kn} \sigma_{k-1}(A). \tag{2}
$$

They proved this conjecture for  $k \leq 3$ . Kopotun [\[10\]](#page-13-6) proved the inequality (2) for  $k = 4$  and  $n \geq 5$ .

By Minc [\[25\]](#page-14-0), the inequality (2) is equivalent to the Monotonicity conjecture, which states that if A is any matrix in  $\Omega_n$  and  $2 \leq k \leq n$ , then  $\sigma_k(X)$  is monotonically increasing on the line segment from  $J_n$  to A. For  $k = n$ , the monotonicity conjecture on permanents is equivalent to

$$
per((1-t)J_n + tA) \leq per(A)
$$

for all  $A \in \Omega_n$  and  $0 \le t \le 1$  [\[9,](#page-13-7) [11\]](#page-13-8). Most interestingly, Wanless [\[19\]](#page-14-1) disproved the Holens -Dokovic inequality (2).

Brualdi and Newman [\[1\]](#page-13-9) considered the inequality

$$
\operatorname{per}(\alpha J_n + (1 - \alpha) A) \le \alpha \operatorname{per}(J_n) + (1 - \alpha) \operatorname{per}(A) \tag{3}
$$

and showed that it did not always hold for all  $A \in \Omega_n$  and for all  $\alpha \in [0,1]$ .

For  $n = 3$ , Lih and Wang [\[12\]](#page-13-10) proved that (3) is true for all  $A \in \Omega_n$  and for all  $\alpha \in \left[\frac{1}{2}\right]$  $\frac{1}{2}$ , 1] and they proposed the following conjecture (quoted by Cheon and Wanless [\[7\]](#page-13-11)): The inequality (3) is true for all  $A \in \Omega_n$  and for all  $\alpha \in \left[\frac{1}{2}\right]$  $\frac{1}{2}$ , 1]. For  $n = 4$ , Foregger [\[5\]](#page-13-3) proved that the inequality (3) holds for all  $A \in \Omega_4$  and for all  $\alpha \in [0.63, 1]$ . Kopotun [\[11\]](#page-13-8) extended this inequality (3) to subpermanents. Subramanian and Somasundaram [\[15\]](#page-13-12) proved that the inequality (3) holds for  $n = 4$  and  $n = 5$  with some conditions on A. A detailed survey of conjectures on permanent was given by Gi-Sang Cheon and Wanless [\[7\]](#page-13-11) and Fuzhen Zhang [\[6\]](#page-13-13). In the next section, we prove that the Lih Wang conjecture is true for  $n = 4$  and the inequality (3) is valid in  $[\beta, 1]$ , where the value of  $\beta$  is about 0.486. Also, we prove the Lih Wang Conjecture for  $n = 6$  with some conditions on A in the interval [0.7836, 1]. The following theorem is due to Foregger [\[5\]](#page-13-3).

**Theorem 1.1.** Let  $A \in \Omega_4$  and  $t_2 < t \leq 1$ , where  $t_2$  is the unique real root of the polynomial  $106t^3 - 418t^2 + 465t - 153$ . *Then* 

$$
per(tJ_4 + (1-t)A) \leq tper(J_4) + (1-t)perA
$$

*with equality iff*  $A = J_4$ *.* 

He found that the value of  $t_2$  is about 0.6216986477375.

The following theorem is due to Subramanian and Somasundaram [\[15\]](#page-13-12)

**Theorem 1.2.** Let  $A \in \Omega_n$  and  $2 \leq k \leq n$ . If the polynomial  $\sum_{r=2}^k r \frac{(k-r)!}{n^{k-r}} {n-r \choose k-r}$  $\binom{n-r}{k-r}^2 \sigma_r(A J_n) t^{r-2}$  has no root in  $(0,1)$  then  $\sigma_k(A) \ge \frac{(n-k+1)^2}{nk} \sigma_{k-1}(A), k = 2, 3, ..., n$ .

Another conjecture on permanents is Dittert's conjecture [[\[25\]](#page-14-0), conjecture 28]. Let  $K_n$ denote the set of all  $n \times n$  real nonnegative matrices whose entries have sum n. Let  $\phi$  denote a real valued function defined on  $K_n$  by  $\phi(X) = \prod_{i=1}^n r_i + \prod_{j=1}^n c_j - per(X)$  for  $X \in K_n$  with row sum vector  $R_X = (r_1, r_2, ... r_n)$  and column sum vector  $C_X = (c_1, c_2, ... c_n)$ . For the same  $X$ , let  $\phi_{ij}(X) = \prod_{k\neq i} r_k + \prod_{l\neq j} c_l - perX(i|j).$ 

A matrix  $A \in K_n$  is called a  $\phi$ -maximizing matrix on  $K_n$  if  $\phi(A) \ge \phi(X)$  for all  $X \in K_n$ . An  $n \times n$  is called fully indecomposable if it does not contain an  $s \times t$  zero submatrix with  $s + t = n.$ 

E. Dittert conjectured that [[\[25\]](#page-14-0), conjecture 28]  $J_n$  is the unique  $\phi$ -maximizing matrix on  $K_n$ . R.Sinkhorn[\[20\]](#page-14-2) proved that every  $\phi$ -maximizing matrix on  $K_n$  has a positive permanent and also that the conjecture is true for  $n = 2$ .

Suk Geun Hwang[\[21,](#page-14-3) [22\]](#page-14-4) investigated some properties of  $\phi$ -maximizing matrices. The following Lemmas and theorems are due to SG Hwang [\[22\]](#page-14-4).

**Lemma 1.1.** Let A be a  $\phi$ -maximizing matrix on  $K_n$ , and let  $1 \leq s < t \leq n$ . If columns s and t of A have either the same sums or the same  $(0, 1)$  patterns, then the matrix obtained from A by replacing each of columns s and t by its average is also a  $\phi$ -maximizing matrix on  $K_n$ . A similar statement holds for rows.

He also proved that Dittert's conjecture is true for positive semidefinite symmetric matrices in  $K_n$  and for matrices in a sufficiently small neighbourhood of  $J_n$  in  $K_n$ .

**Lemma 1.2.** Let  $A = [a_{ij}]$  be a  $\phi$ -maximizing matrix on  $K_n$ . Then

- $\phi_{ii}(A) = \phi_{kl}(A)$  if  $a_{ij} > 0$  and  $a_{kl} > 0$ .
- $\phi_{ii}(A) \leq \phi_{kl}(A)$  if  $a_{ii} = 0$  and  $a_{kl} > 0$ .

**Theorem 1.3.** *If*  $A = [a_{ij}]$  *is a*  $\phi$ *-maximizing matrix on*  $K_n$ *, then*  $\phi_{ij}(A) = \phi(A)$  *if*  $a_{ij} > 0$  *and*  $\phi_{ij}(A) \leq \phi(A)$  *if*  $a_{ij} = 0$ .

**Theorem 1.4.**  $J_3$  *is the unique*  $\phi$ *-maximizing matrix on*  $K_3$ *.* 

He [\[22\]](#page-14-4) stated that one can at lest prove that if A is a  $\phi$  maximizing matrix on  $K_n$  then A is fully indecomposable. Based on his remark we prove that if A is a  $\phi$ -maximizing matrix on  $K_4$  then A is fully indecomposable.

Cheon and Wanless [\[23\]](#page-14-5) proved that if  $A \in K_n$  is partly decomposable then  $\phi(A) < \phi(J_n)$  and that if the zeroes in  $A \in K_n$  form a block then A is not a  $\phi$ -maximizing matrix. Cheon and Yoon [\[24\]](#page-14-6) obtained some sufficient conditions for which the Dittert conjecture holds.

# 2 Lih Wang conjecture

In this section, first we prove the Lih Wang conjecture completely for  $n = 4$ . We improved the results of Theorem 1.1 and Theorem 1.2. Later in Theorem 2.2, we prove this conjecture partially for  $n = 6$ .

**Theorem 2.1.** *If*  $A \in \Omega_4$  *then*  $per(\alpha J_4 + (1-\alpha)A) \leq \alpha per(J_4) + (1-\alpha) perA$  *for all*  $\alpha \in [t,1]$ *where the value of* t *is about* 0.485.

*Proof.* Let  $f(\alpha) = \alpha \text{ per}(J_4) + (1 - \alpha) \text{ per}A - \text{ per}(\alpha J_4 + (1 - \alpha)A)$ . As we know that  $\text{per}(J_4) = \frac{3}{32}$ ,

$$
f(\alpha) = 3\alpha + 32(1 - \alpha) \text{per}(A) - 32\text{per}(\alpha J_4 + (1 - \alpha)A) \ge 0.
$$
 (4)

Let  $a_i$  denote the  $i^{th}$  column of A and  $e_r(\mathbf{x}) = e_r(x_1, x_2, ... x_n)$  denote the  $r^{th}$  elementary symmetric function of  $\mathbf{x} = (x_1, x_2...x_n)$ . That is, the sum of products of components taken r at a time,  $r = 1, 2, ...n$ . Let  $T_r(A)$  denote the set of  $\binom{n}{r}$  $r \choose r$  sums of columns of A taken r at a time. That is, let  $T_r(A) = {\mathbf{x} = a_{i_1} + a_{i_2} + ... + a_{i_r} | (i_1, i_2, ..., i_r) \text{ is a } r\text{-subset of } (1, 2, ... n)},$ then

$$
\text{per}A = \sum_{T_n(A)} e_n(\mathbf{x}) - \sum_{T_{n-1}(A)} e_n(\mathbf{x}) + \dots + (-1)^{n-1} \sum_{T_1(A)} e_n(\mathbf{x}).
$$

When  $A$  is doubly stochastic we have

$$
\sum_{T_n(A)} e_n(\mathbf{x}) = 1.
$$

Let  $B = \alpha J_4 + (1 - \alpha)A$ . We use the formula of Eberlein and Mudholkar ([\[3\]](#page-13-14), page 392) to calculate permanent of B.

$$
per(B) = \frac{-1}{3} + \frac{1}{9} \sum_{T_1(B)} (-4e_2 + 9e_3 - 18e_4)(x) + \frac{1}{18} \sum_{T_2(B)} (4e_2 - 9e_3 + 18e_4)(\mathbf{x}).
$$

The elements of *T*<sub>1</sub>(*B*) are the columns of *B*. Let *b<sub>i</sub>* denote the *i*<sup>th</sup> column of *B*. Then *b<sub>i</sub>* is of the form 
$$
b_i = \alpha \begin{bmatrix} \frac{1}{4} \\ \frac{1}{4} \\ \frac{1}{4} \\ \frac{1}{4} \end{bmatrix} + (1 - \alpha) \begin{bmatrix} a_{1i} \\ a_{2i} \\ a_{3i} \\ a_{4i} \end{bmatrix}, \text{ where } \begin{bmatrix} a_{2i} \\ a_{3i} \\ a_{4i} \\ a_{4i} \end{bmatrix} \text{ is the } i^{th}
$$
 column of *A*.  
\nNow, 
$$
e_2(b_i) = e_2 \begin{bmatrix} \frac{1}{4} + (1 - \alpha)a_{1i} \\ \frac{1}{4} + (1 - \alpha)a_{1i} \\ \frac{1}{4} + (1 - \alpha)a_{3i} \\ \frac{1}{4} + (1 - \alpha)a_{4i} \end{bmatrix}
$$
\nNow, 
$$
e_2(b_i) = e_2 \begin{bmatrix} \frac{1}{4} + (1 - \alpha)a_{1i} \\ \frac{1}{4} + (1 - \alpha)a_{2i} \end{bmatrix} + \frac{16}{16} + (1 - \alpha)a_{1i} \Big| \frac{16}{4} + (1 - \alpha)a_{3i} \Big| + \dots
$$
\n
$$
= \frac{3}{8}\alpha^2 + \frac{3}{4}\alpha(1 - \alpha) + (1 - \alpha)^2 e_2(a_i).
$$
\n
$$
e_3(b_i) = \Big[\frac{\alpha}{4} + (1 - \alpha)a_{1i}\Big] \Big[\frac{\alpha}{4} + (1 - \alpha)a_{2i}\Big] \Big[\frac{\alpha}{4} + (1 - \alpha)a_{3i}\Big] + \frac{\alpha}{4} + (1 - \alpha)a_{1i} \Big] \Big[\frac{\alpha}{4} + (1 - \alpha)a_{1i} \Big] \Big[\frac{\alpha}{4
$$

Therefore,

$$
\sum_{i=1}^{4} e_2(b_i) = \frac{3}{2}\alpha^2 + 3\alpha(1-\alpha) + (1-\alpha)^2 \sum_{i=1}^{4} e_2(a_i),
$$

$$
\sum_{i=1}^{4} e_3(b_i) = \frac{\alpha^3}{4} + \frac{3}{4}\alpha^2(1-\alpha) + \frac{1}{2}\alpha(1-\alpha)^2 \sum_{i=1}^{4} e_2(a_i) + (1-\alpha)^3 \sum_{i=1}^{4} e_3(a_i)
$$

and

$$
\sum_{i=1}^{4} e_4(b_i) = \frac{1}{64} \alpha^4 + \frac{1}{16} \alpha^3 (1-\alpha) + \frac{1}{16} \alpha^2 (1-\alpha)^2 \sum_{i=1}^{4} e_2(a_i) + \frac{1}{4} \alpha (1-\alpha)^3 \sum_{i=1}^{4} e_3(a_i) + (1-\alpha)^4 \sum_{i=1}^{4} e_4(a_i).
$$

$$
\sum_{T_2(B)} e_2(\mathbf{x}) = 18\alpha - 9\alpha^2 + (1 - \alpha)^2 \sum_{T_2(A)} e_2(\mathbf{y}),
$$

$$
\sum_{T_2(B)} e_3(\mathbf{x}) = -6\alpha^3 + 9\alpha^2 + \alpha(1-\alpha)^2 \sum_{T_2(A)} e_2(\mathbf{y}) + (1-\alpha)^3 \sum_{T_2(A)} e_3(\mathbf{y})
$$

and

$$
\sum_{T_2(B)} e_4(\mathbf{x}) = \frac{3\alpha^3}{2} - \frac{9\alpha^4}{8} + \alpha(1-\alpha)^2 \sum_{T_2(A)} e_2(\mathbf{y}) + (1-\alpha)^3 \sum_{T_2(A)} e_3(\mathbf{y}).
$$

Now, we use the formula of Eberlein and Mudholkar ([\[3\]](#page-13-14), equations (3.4) and (3.5)) for any  $A \in \Omega_4$ ,

$$
\sum_{T_2(A)} e_2(\mathbf{x}) = 2 \sum_{T_1(A)} e_2(\mathbf{x}) + 6,
$$

$$
\sum_{T_2(A)} e_3(\mathbf{x}) = 2 \sum_{T_1(A)} e_2(\mathbf{x}).
$$

Using the above two equations, we get

$$
\sum_{T_2(B)} e_2(\mathbf{x}) = 18\alpha - 9\alpha^2 + (2 - 4\alpha + 2\alpha^2) \sum_{i=1}^4 e_2(a_i) + 6 - 12\alpha + 6\alpha^2,
$$

$$
\sum_{T_2(B)} e_3(\mathbf{x}) = -6\alpha^3 + 9\alpha^2 + (\alpha - 2\alpha^2 + \alpha^3)(2\sum_{i=1}^4 e_2(a_i) + 6) + (2 - 6\alpha + 6\alpha^2 - 2\alpha^3)\sum_{i=1}^4 e_2(a_i)
$$

and

$$
\sum_{T_2(B)} e_4(\mathbf{x}) = \frac{3\alpha^3}{2} - \frac{9\alpha^4}{8} + \left(\frac{\alpha^2 - 2\alpha^3 + \alpha^4}{4}\right)(2\sum_{i=1}^4 e_2(a_i) + 6) +
$$

$$
(\alpha - 3\alpha^2 + 3\alpha^3 - \alpha^4) \sum_{i=1}^4 e_2(a_i) + (1 - \alpha)^4 \sum_{T_2(A)} e_4(\mathbf{y}).
$$

Therefore

$$
per(B) = \frac{-1}{3} + \frac{1}{9}(-4(\frac{3\alpha^2}{2} + 3\alpha(1 - \alpha) + (1 - \alpha)^2 \sum_{i=1}^4 e_2(a_i))
$$

$$
+9\left(\frac{\alpha^3}{4} + \frac{3\alpha^2(1-\alpha)}{4} + \frac{\alpha(1-\alpha)^2}{2}\sum_{i=1}^4 e_2(a_i) + (1-\alpha)^3 \sum_{i=1}^4 e_3(a_i)\right)
$$

$$
-18\left(\frac{\alpha^4}{64} + \frac{\alpha^3(1-\alpha)}{16} + \frac{\alpha^2(1-\alpha)^2}{16}\right) \sum_{i=1}^4 e_2(a_i) + \frac{\alpha(1-\alpha)^3}{4} \sum_{i=1}^4 e_3(a_i) + (1-\alpha)^4 \sum_{i=1}^4 e_4(a_i))
$$

$$
+\frac{1}{18}(4(18\alpha - 9\alpha^2 + (2 - 4\alpha + 2\alpha^2))\sum_{i=1}^{4}e_2(a_i) + 6 - 12\alpha + 6\alpha^2)
$$

$$
-9(-6\alpha^3 + 9\alpha^2 + (2\alpha - 4\alpha^2 + 2\alpha^3) \sum_{i=1}^4 e_2(a_i) + 6\alpha - 12\alpha^2 + 6\alpha^3 + (2 - 6\alpha + 6\alpha^2)
$$

$$
-2\alpha^3\left(\sum_{i=1}^4 e_2(a_i)\right) + 18\left(\frac{3\alpha^3}{2} - \frac{9\alpha^4}{8} + \frac{\alpha^2 - 2\alpha^3 + \alpha^4}{2}\sum_{i=1}^4 e_2(a_i) + \frac{3(\alpha^2 - 2\alpha^3 + \alpha^4)}{2}
$$

$$
+(\alpha - 3\alpha^2 + 3\alpha^3 - \alpha^4) \sum_{i=1}^4 e_2(a_i) + (1 - \alpha)^4 \sum_{T_2(A)} e_4(\mathbf{y})).
$$

Now,

$$
f(\alpha) = 3\alpha + 32(1 - \alpha)\left(\frac{-1}{3} + \frac{1}{9}\sum_{T_1(A)} (-4e_2 + 9e_3 - 18e_4)(\mathbf{x})\right)
$$

$$
+\frac{1}{18}\sum_{T_2(A)}(4e_2-9e_3+18e_4)(\mathbf{x}))
$$

$$
-32\left(\frac{-1}{3} + \frac{1}{9}\left(-4\left(\frac{3\alpha^2}{2} + 3\alpha(1-\alpha) + (1-\alpha)^2\right)\right)\right)_{i=1}^4 e_2(a_i))
$$

$$
+9\left(\frac{\alpha^3}{4} + \frac{3\alpha^2(1-\alpha)}{4} + \frac{\alpha(1-\alpha)^2}{2}\sum_{i=1}^4 e_2(a_i) + (1-\alpha)^3 \sum_{i=1}^4 e_3(a_i)\right)
$$

$$
-18\left(\frac{\alpha^4}{64} + \frac{\alpha^3(1-\alpha)}{16} + \frac{\alpha^2(1-\alpha)^2}{16}\sum_{i=1}^4 e_2(a_i) + \frac{\alpha(1-\alpha)^3}{4}\sum_{i=1}^4 e_3(a_i) + (1-\alpha)^4\sum_{i=1}^4 e_4(a_i))\right)
$$

$$
+\frac{1}{18}(4(18\alpha - 9\alpha^2 + (2 - 4\alpha + 2\alpha^2))\sum_{i=1}^{4}e_2(a_i) + 6 - 12\alpha + 6\alpha^2)
$$

$$
-9(-6\alpha^3 + 9\alpha^2 + (2\alpha - 4\alpha^2 + 2\alpha^3) \sum_{i=1}^4 e_2(a_i) + 6\alpha - 12\alpha^2 + 6\alpha^3 + (2 - 6\alpha + 6\alpha^2)
$$

$$
-2\alpha^3\left(\sum_{i=1}^4 e_2(a_i)\right) + 18\left(\frac{3\alpha^3}{2} - \frac{9\alpha^4}{8} + \frac{\alpha^2 - 2\alpha^3 + \alpha^4}{2}\sum_{i=1}^4 e_2(a_i) + \frac{3(\alpha^2 - 2\alpha^3 + \alpha^4)}{2}
$$

$$
+(\alpha - 3\alpha^2 + 3\alpha^3 - \alpha^4) \sum_{i=1}^4 e_2(a_i) + (1 - \alpha)^4 \sum_{T_2(A)} e_4(\mathbf{y}))).
$$

After simplifying we get a 4<sup>th</sup> degree polynomial in  $\alpha$ ,

$$
f(\alpha) = \frac{128}{3} + \alpha[67 - 80 \sum_{i=1}^{4} e_2(a_i) + 80 \sum_{i=1}^{4} e_3(a_i) - 192 \sum_{i=1}^{4} e_4(a_i) + 96 \sum_{T_2(A)} e_4(x)]
$$
  

$$
+ \alpha^2[-120 + 148 \sum_{i=1}^{4} e_2(a_i) - 144 \sum_{i=1}^{4} e_3(a_i) + 384 \sum_{i=1}^{4} e_4(a_i) - 192 \sum_{T_2(A)} e_4(x)]
$$
  

$$
+ \alpha^3[68 - 88 \sum_{i=1}^{4} e_2(a_i) + 80 \sum_{i=1}^{4} e_3(a_i) - 256 \sum_{i=1}^{4} e_4(a_i) + 128 \sum_{T_2(A)} e_4(x)]
$$
  

$$
+ \alpha^4[-15 + 24 \sum_{i=1}^{4} e_2(a_i) - 16 \sum_{i=1}^{4} e_3(a_i) + 64 \sum_{i=1}^{4} e_4(a_i) - 32 \sum_{T_2(A)} e_4(x)].
$$

Using the MATLAB, we obtained the minimum values of the function  $f(\alpha)$  for different values of  $\alpha$ . We obtained minimum values of  $f(\alpha)$  in different sub intervals of [0, 1] and the values are shown in table 1.

Therefore, the function  $f(\alpha) \ge 0$  in [0.5, 1]. Hence, Lih-Wang conjecture is true for  $n = 4$  and in particular the inequality  $(3)$  is true in [0.486, 1].  $\Box$ 

Interval	Minimum at $\alpha$	Minumum value $f(\alpha)$
[0.5, 1]	0.5	0.1250
[0.4, 1]	0.4	$-0.5472$
[0.45, 1]	0.45	$-0.2654$
[0.48, 1]	0.48	$-0.0452$
[0.485, 1]	0.45	$-0.0044$
[0.486, 1]	0.486	0.0039
[0, 1]	0.2545	$-0.8675$

Table 1: Minimum value

**Theorem 2.2.** *If* A *is a matrix in*  $\Omega_6$  *such that the polynomials* 

 $\sum_{r=2}^{5} r \frac{(5-r)!}{6^{5-r}}$  $\frac{5-r)!}{6^{5-r}} \binom{6-r}{5-r}$  $\int_{5-r}^{6-r} \int_{0}^{2\pi} \sigma_r(A-J_6) t^{r-2}$  and  $\sum_{r=2}^{6} r \frac{(6-r)!}{6^{6-r}}$  $\frac{6-r)!}{6^{6-r}}\sigma_r(A-J_6)t^{r-2}$  have no roots in  $(0,1)$  then  $per(tJ_6 + (1-t)A) \le t$   $perJ_6 + (1-t)perA$  *for all*  $t \in [0.7836, 1]$ .

*Proof.*  $t$  per $J_6 + (1 - t)$  per  $A$ -per $(tJ_6 + (1 - t)A)$  $=\frac{5}{324}t + (1-t)\text{per}A - \sum_{r=0}^{6} t^r(1-t)^{6-r}\frac{r!}{6r}$  $\frac{r_!}{6^r}\sigma_{6-r}(A)$  $=\frac{5}{324}t+(1-t)$  per  $A-[(1-t)^6$  per  $A+\frac{1}{6}$  $=\frac{5}{324}t+(1-t)$  per  $A-[(1-t)^6$  per  $A+\frac{1}{6}t(1-t)^5\sigma_5(A)+\frac{2}{62}t^2(1-t)^4\sigma_4(A)+\frac{6}{63}t^3(1-t)^3\sigma_3(A)+\frac{24}{64}t^4(1-t)^2\sigma_5(A)+\frac{120}{63}t^3(1-t)\sigma_6(A)$  $\frac{24}{6^4}\tilde{t}^4(1-t)^2\sigma_2(A)+\frac{120}{6^5}t^5(1-t)\sigma_1(A)+\frac{720}{6^6}t^6]$  $=\frac{5}{324}t+\text{per}A(1-t)[1-(1-t)^5]-\frac{1}{6}$  $\frac{1}{6}t(1-t)^5\sigma_5(A)-\frac{1}{18}t^2(1-t)^4\sigma_4(A)-\frac{1}{36}t^3(1-t)^3\sigma_3(A) \frac{1}{54}t^4(1-t)^2\sigma_2(A)-\frac{5}{54}t^5(1-t)-\frac{5}{324}t^6$  $=\frac{5}{324}t(1-t)(1+t+t^2+t^3+t^4)-\frac{5}{54}t^5(1-t)+\text{per}A(1-t)(5t-10t^2+10t^3-5t^4+t^5) \frac{1}{6}t(1-t)^5\sigma_5(A) - \frac{1}{18}t^2(1-t)^4\sigma_4(A) - \frac{1}{36}t^3(1-t)^3\sigma_3(A) - \frac{1}{54}t^4(1-t)^2\sigma_2(A)$  $=\frac{5}{324}t(1-t)(1+t+t^2+t^3-5t^4)$ +per $At(1-t)(5-10t+10t^2-5t^3+t^4)-\frac{1}{6}$  $=\frac{5}{324}t(1-t)(1+t+t^2+t^3-5t^4)$ +per $At(1-t)(5-10t+10t^2-5t^3+t^4)-\frac{1}{6}t(1-t)^5σ_5(A)-\frac{1}{18}t^2(1-t)^4σ_4(A)-\frac{1}{36}t^3(1-t)^3σ_3(A)-\frac{1}{54}t^4(1-t)^2σ_2(A)$ 

 $=t(1-t)F_A(t)$ , where

$$
F_A(t) = \frac{5}{324}(1 + t + t^2 + t^3 - 5t^4) + \text{per}A(5 - 10t + 10t^2 - 5t^3 + t^4) - \frac{1}{6}(1 - t)^4 \sigma_5(A) - \frac{1}{18}t(1 - t)^3 \sigma_4(A) - \frac{1}{36}t^2(1 - t)^2 \sigma_3(A) - \frac{1}{54}t^3(1 - t)\sigma_2(A).
$$

It is enough to prove that  $F_A(t) \geq 0$  for all  $t \in [0.7836, 1]$ . Since the theorem holds good for  $t = 1$ , it is sufficient to consider  $t < 1$ .

$$
F_A(t) = (5 - 10t + 10t^2 - 5t^3 + t^4)(per A - \frac{1}{36}\sigma_5(A)) + \frac{1}{36}(5 - 10t + 10t^2 - 5t^3 + t^4)\sigma_5(A) - \frac{1}{6}(1 - t)^4\sigma_5(A) - \frac{1}{18}t(1 - t)^3\sigma_4(A) - \frac{1}{36}t^2(1 - t)^2\sigma_3(A) - \frac{1}{54}t^3(1 - t)\sigma_2(A) + \frac{5}{324}(1 + t + t^2 + t^3 - 5t^4).
$$
  
\nBy Theorem 1.2,  $F_A(t) \ge \frac{1}{36}(-1 + 14t - 26t^2 + 19t^3 - 5t^4)\sigma_5(A) - \frac{1}{18}t(1 - t)^3\sigma_4(A) - \frac{1}{36}t^2(1 - t)^2\sigma_3(A) - \frac{1}{54}t^3(1 - t)\sigma_2(A) + \frac{5}{324}(1 + t + t^2 + t^3 - 5t^4)$   
\n
$$
= \frac{1}{36}(-1 + 14t - 26t^2 + 19t^3 - 5t^4)(\sigma_5(A) - \frac{2}{15}\sigma_4(A)) + \frac{1}{270}\sigma_4(A)(-1 + 14t - 26t^2 + 19t^3 - 5t^4) - \frac{1}{18}t(1 - t)^3\sigma_4(A) - \frac{1}{36}t^2(1 - t)^2\sigma_3(A) - \frac{1}{54}t^3(1 - t)\sigma_2(A) + \frac{5}{324}(1 + t + t^2 + t^3 - 5t^4).
$$

By Theorem 1.2, 
$$
F_A(t) \geq \frac{1}{270} \sigma_4(A)(-1 + 14t - 26t^2 + 19t^3 - 5t^4) - \frac{1}{18}t(1 - t)^3 \sigma_4(A) - \frac{1}{36}t^2(1 - t)^2 \sigma_3(A) - \frac{1}{54}t^3(1 - t)\sigma_2(A) + \frac{5}{324}(1 + t + t^2 + t^3 - 5t^4).
$$

This implies that 
$$
F_A(t) \ge \frac{1}{270} \sigma_4(A) \{-1 + 14t - 26t^2 + 19t^3 - 5t^4 - 15t(1 - 3t + 3t^2 - t^3)\} - \frac{1}{36}t^2(1-t)^2\sigma_3(A) - \frac{1}{54}t^3(1-t)\sigma_2(A) + \frac{5}{324}(1+t+t^2+t^3 - 5t^4)
$$
  
\n $= \frac{1}{270} \sigma_4(A)(-1-t+19t^2 - 26t^3 + 10t^4) - \frac{1}{36}t^2(1-t)^2\sigma_3(A) - \frac{1}{54}t^3(1-t)\sigma_2(A) + \frac{5}{324}(1+t+t^2+t^3 - 5t^4)$ 

$$
=\frac{1}{270}(-1-t+19t^2-26t^3+10t^4)(\sigma_4(A)-\frac{3}{8}\sigma_3(A))+\frac{3}{2160}\sigma_3(A)(-1-t+19t^2-26t^3+10t^4)-\frac{1}{36}t^2(1-t)^2\sigma_3(A)-\frac{1}{54}t^3(1-t)\sigma_2(A)+\frac{5}{324}(1+t+t^2+t^3-5t^4)
$$

Again by Theorem 1.2,  $F_A(t) \ge \frac{3}{2160} \sigma_3(A)(-1-t+19t^2-26t^3+10t^4)-\frac{1}{36}t^2(1-t)^2\sigma_3(A)-\frac{1}{54}t^3(1-t)\sigma_2(A)+\frac{5}{324}(1+t+t^2+t^3-5t^4)$ 

$$
=\sigma_3(A)\left(\frac{-3}{2160} - \frac{3}{2160}t - \frac{3}{2160}t^2 + \frac{42}{2160}t^3 - \frac{30}{2160}t^4\right) - \frac{1}{54}t^3(1-t)\sigma_2(A) + \frac{5}{324}(1+t+t^2+t^3 - 5t^4)
$$

From Dokovic[\[2\]](#page-13-5) and Tverberg[\[17\]](#page-14-7) we know that for  $A \in \Omega_n$ ,  $\sigma_2(A) = \frac{1}{2} \sum_{i,j=1}^n a_{ij}^2 + \frac{1}{2}$  $\frac{1}{2}n(n-2)$  and

$$
\sigma_3(A) = \frac{2}{3} \sum_{i,j=1}^n a_{ij}^3 + \frac{1}{2}(n-4) \sum_{i,j=1}^n a_{ij}^2 + \frac{1}{6}n(n^2 - 6n + 10).
$$

Therefore  $F_A(t) \ge \frac{1}{2160}(-3-3t-3t^2+42t^3-30t^4)(\frac{2}{3}\sum_{i,j=1}^6 a_{ij}^3 + \sum_{i,j=1}^6 a_{ij}^2 + 10) - \frac{1}{54}t^3(1$  $t$ )( $\frac{1}{2}\sum_{i,j=1}^{6} a_{ij}^2 + 12$ ) +  $\frac{5}{324}(1 + t + t^2 + t^3 - 5t^4)$ 

$$
\frac{1}{3240}(-3-3t-3t^2+42t^3-30t^4)\sum_{i,j=1}^6 a_{ij}^3+\frac{1}{2160}\sum_{i,=1}^6 a_{ij}^2(-3-3t-3t^2+22t^3-10t^4)+\frac{1}{648}(1+t+t^2-8t^3+4t^4)
$$

 $=\sum_{i=1}^{6} \left\{\frac{1}{3240}(-3 - 3t - 3t^2 + 42t^3 - 30t^4)\sum_{j=1}^{6} a_{ij}^3 + \frac{1}{216}t^2\right\}$  $\frac{1}{2160}\sum_{j=1}^{6} a_{ij}^2(-3-3t-3t^2+22t^3 10t^4$ } +  $\frac{1}{648}(1 + t + t^2 - 8t^3 + 4t^4)$ .

It is easy to see that the coefficient of  $\sum_{j=1}^{6} a_{ij}^3$  is nonnegative for all  $t \in [0.6530, 1]$  and  $(\sum_{j=1}^{6} a_{ij}^2)^2 \leq \sum_{j=1}^{6} a_{ij}^3$  for  $i = 1, 2, ...6$  (Kopotun[\[10\]](#page-13-6)).

Therefore, for all  $t \in [0.6530, 1]$ ,  $F_A(t) \ge \sum_{i=1}^6 \left[\frac{1}{3240}(-3 - 3t - 3t^2 + 42t^3 - 30t^4)(\sum_{j=1}^6 a_{ij}^2)^2 + \frac{1}{2160}(-3 - 3t - 3t^2 + 22t^3 10t^4$ )  $\sum_{j=1}^6 a_{ij}^2$  +  $\frac{1}{648}(1 + t + t^2 - 8t^3 + 4t^4)$ . Put  $x = \sum_{j=1}^{6} a_{ij}^2$  and  $f(x) = \frac{1}{3240}(-3 - 3t - 3t^2 + 42t^3 - 30t^4)x^2 + \frac{1}{2160}(-3 - 3t - 3t^2 + 22t^3 - 10t^4)x.$ Now,  $f'(x) = \frac{2}{3240}x(-3 - 3t - 3t^2 + 42t^3 - 30t^4) + \frac{1}{2160}(-3 - 3t - 3t^2 + 22t^3 - 10t^4)$ It it easy to see that for  $t \in [0.7836, 1], f'(x) \ge 0.$ 

London and Minc(Lemma1,[\[18\]](#page-14-8)) showed that  $x = \sum_{j=1}^{6} a_{ij}^2 \ge \frac{1}{6}$  $\frac{1}{6}$ . Hence  $f(x)$  is an increasing function on  $\left[\frac{1}{6}\right]$  $\frac{1}{6}, \infty$ ).

That is,  $f(x) \ge f(\frac{1}{6})$  $\frac{1}{6}$ ) =  $\frac{-1}{3888} - \frac{1}{3888}t - \frac{1}{3888}t^2 + \frac{1}{486}t^3 - \frac{1}{972}t^4$ 

Therefore  $t \in [0.7836, 1]$  $F_A(t) \ge 6\left(\frac{-1}{3888} - \frac{1}{3888}t - \frac{1}{3888}t^2 + \frac{1}{486}t^3 - \frac{1}{972}t^4\right) + \frac{1}{648}(1 + t + t^2 - 8t^3 + 4t^4) = 0.$ 

Hence the theorem.

The following two corollaries are consequence of the above theorem and theorems in [\[15\]](#page-13-12).

 $\Box$ 

**Corollary 2.1.** If A is a normal matrix in  $\Omega_6$  whose eigenvalues lie in the sector  $\left[\frac{-\pi}{10}, \frac{\pi}{10}\right]$  and the polynomial  $\sum_{r=2}^{6} r \frac{(6-r)!}{6^{6-r}}$  $\frac{6-r)!}{6^{6-r}}\sigma_r(A-J_6)t^{r-2}$  has no root in  $(0, 1)$  then  $per(tJ_6 + (1-t)A) \le t$  per  $J_6 + (1-t)$ perA for all  $t \in [0.7836, 1]$ .

*Proof.* The proof follows since if A is a normal matrix in  $\Omega_6$  whose eigenvalues lie in the sector  $\int_{5-r}^{6-r} \int_{0}^{2} \sigma_r (A - J_6) t^{r-2}$  has no roots in (0, 1).  $\left[\frac{-\pi}{10}, \frac{\pi}{10}\right]$  then the polynomial  $\sum_{r=2}^{5} r \frac{(5-r)!}{6^{5-r}}$  $\frac{5-r)!}{6^{5-r}} \binom{6-r}{5-r}$ 口

**Corollary 2.2.** If A is a normal matrix in  $\Omega_6$  whose eigenvalues lie in the sector  $\left[\frac{-\pi}{12}, \frac{\pi}{12}\right]$  then  $per(tJ_6 + (1-t)A) \leq t$  per $J_6 + (1-t)$  per A for all  $t \in [0.7836, 1]$ .

*Proof.* The proof follows since if A is a normal matrix in  $\Omega_6$  whose eigenvalues lie in the  $\int_{5-r}^{6-r} \int_{0}^{2\pi} \sigma_r(A-J_6)t^{r-2}$  and  $\sum_{r=2}^{6} r \frac{(6-r)!}{6^{6-r}}$ sector  $\left[\frac{-\pi}{12}, \frac{\pi}{12}\right]$  then the polynomials  $\sum_{r=2}^{5} r \frac{(5-r)!}{6^{5-r}}$  $\frac{5-r)!}{6^{5-r}} \binom{6-r}{5-r}$  $\frac{6-r}{6^{6-r}}\sigma_r(A J_6$ ) $t^{r-2}$  have no roots in (0, 1).  $\Box$ 

# 3 Dittert's Conjecture for  $n = 4$

SG Hwang [\[22\]](#page-14-4) stated that if A is a  $\phi$  maximizing matrix on  $K_n$  then A is fully indecomposable. Based on this we prove that if A is a  $\phi$ -maximizing matrix on  $K_4$  then A is fully indecomposable. We denote  $R_A = (r_1, ..., r_n)$  and  $C_A = (c_1, ..., c_n)$  are respectively the row and column sum vectors of A.

**Theorem 3.1.** If A is a  $\phi$ -maximizing matrix on  $K_4$  then A is fully indecomposble.

*Proof.* Let  $A = [a_{ij}]$  be a  $\phi$ -maximizing matrix on  $K_4$ . First, we assume that A is not fully indecomposable. Without loss of generality we may assume that  $a_{12} = a_{13} = a_{14} = 0$ . Now we intend to show that  $(a_{21}, a_{31}, a_{41}) \neq (0, 0, 0)$ . Suppose  $a_{21} = a_{31} = a_{41} = 0$ . Let  $B = A(1|1)$ and  $a_{11} = a$ . Then  $A = aI_1 \oplus B$ ,  $\phi(A) = a\phi(B)$  and  $\phi_{11}(A) = \phi(B)$ . By Theorem 1.3,  $a = 1$ and hence  $B \in K_3$  and by Theorem 1.4,  $B = J_3$  and hence  $\phi(A) = \phi(B) = \phi(J_3) < \phi(J_4)$ , which is a contradiction. Thus  $(a_{21}, a_{31}, a_{41}) \neq (0, 0, 0)$ .

If  $a_{21} > 0, a_{31} > 0$  and  $a_{41} > 0$ , by Lemma 1.2,  $\phi_{21}(A) = \phi_{31}(A) = \phi_{41}(A), \phi_{21}(A) =$  $r_1r_3r_4 + c_2c_3c_4$ -per $A(2|1)$  and  $\phi_{31}(A) = r_1r_2r_4 + c_2c_3c_4$ -per $A(3|1)$ . Therefore  $\phi_{21}(A) =$  $\phi_{31}(A) = o$  implies  $ar_4(r_3 - r_2) = 0$  and hence  $r_2 = r_3$ . Then by Lemma 1.1 (the matrix  $A_1$  obtained from A by replacing the 2nd and 3rd rows by its average and  $A_1$  is also a  $\phi$ maximizing matrix on  $K_4$ ), we assume that  $A_1$  is of the form

$$
A_1 = \begin{bmatrix} a & 0 & 0 & 0 \\ x & y & z & t \\ x & y & z & t \\ p & q & r & s \end{bmatrix},
$$

where  $axyztpqrs > 0$ . Since the second, third and fourth columns have the same (0, 1) patterns, by Lemma 1.1 we assume that  $A_1$  is of the form

$$
A_2 = \begin{bmatrix} a & 0 & 0 & 0 \\ x & y & y & y \\ x & y & y & y \\ p & q & q & q \end{bmatrix}
$$

,

where  $axypq > 0$ . Now since the second, third and fourth rows have the same (0, 1) patterns, by Lemma 1.1 we can assume that  $A_2$  is of the form

$$
A_3 = \begin{bmatrix} a & 0 & 0 & 0 \\ x & y & y & y \\ x & y & y & y \\ x & y & y & y \end{bmatrix},
$$

where  $axy > 0$ . Suppose  $x \ge y$ . We construct  $A_4$  from  $A_3$  such that  $R_{A_4} = R_{A_3}, C_{A_4} = C_{A_3}$ and  $perA_4 = perA_3$ ,

$$
A_4 = \begin{bmatrix} a & 0 & 0 & 0 \\ x & y & y & y \\ x+y & 0 & y & y \\ x-y & 2y & y & y \end{bmatrix}.
$$

The matrix  $A_4$  is also a  $\phi$ -maximizing matrix on  $K_4$  since  $R_{A_4} = R_{A_3}, C_{A_4} = C_{A_3}$  and per $A_4$  =per $A_3$ . Now  $a_{33} > 0$  and  $a_{42} > 0$ , but  $\phi_{42}(A_4) - \phi_{33}(A_4) = ay^2 > 0$  which is a contradiction to Lemma 1.2. Hence  $x < y$ . We construct  $A_5$  from  $A_3$  such that  $R_{A_5}$  =  $R_{A_3}, C_{A_5} = C_{A_3}$  and per $A_5 =$ per $A_3$ ,

$$
A_5 = \begin{bmatrix} a & 0 & 0 & 0 \\ x & y & y & y \\ 2x & y & y & y - x \\ 0 & y & y & y + x \end{bmatrix}.
$$

The matrix  $A_5$  is a  $\phi$ -maximizing matrix on  $K_4$  since  $R_{A_5} = R_{A_3}$ ,  $C_{A_5} = C_{A_3}$  and per $A_5 =$ per $A_3$ . So, by Lemma 1.1, we construct the matrix  $A_6$  from  $A_5$  by taking average of 3rd and 4th columns of  $A_5$ .

$$
A_6 = \begin{bmatrix} a & 0 & 0 & 0 \\ x & y & y & y \\ 2x & y & y - \frac{x}{2} & y - \frac{x}{2} \\ 0 & y & y + \frac{x}{2} & y + \frac{x}{2} \end{bmatrix}
$$

is also a  $\phi$ -maximizing matrix on  $K_4$ .  $R_{A_6} = R_{A_5}$ ,  $C_{A_6} = C_{A_5}$  but per $A_6 = a(6y^3 - \frac{yx^2}{2})$  $\frac{x^2}{2}$ ) <  $6ay^3 = perA_5$ . This implies that  $\phi(A_6) > \phi(A_5) = \phi(A_3)$ , which is a contradiction. Hence at least one of  $a_{21}, a_{31}, a_{41}$  is zero.

Without loss of generality we may assume that  $a_{21} > 0$  and  $a_{31} = a_{41} = 0$ . Hence A has the form

$$
A = \begin{bmatrix} a & 0 & 0 & 0 \\ b & B & \\ 0 & & \\ 0 & & \end{bmatrix}
$$

with  $ab > 0$ . We claim that B is not a positive matrix. Because if  $B > 0$  then we may say

$$
A = \begin{bmatrix} a & 0 & 0 & 0 \\ b & x & x & x \\ 0 & y & y & y \\ 0 & z & z & z \end{bmatrix}.
$$

Since the third and fourth rows have the same  $(0, 1)$  patterns, we construct the matrix  $\tilde{A}$  from A by taking the average of 3rd and 4th rows of A.

$$
\tilde{A} = \begin{bmatrix} a & 0 & 0 & 0 \\ b & x & x & x \\ 0 & y & y & y \\ 0 & y & y & y \end{bmatrix}
$$

with  $abxy > 0$ .  $\phi_{11}(\tilde{A}) - \phi_{21}(\tilde{A}) = 3y^2(3b - 3a + 7x)$ . From Lemma 1.2,  $\phi_{11}(\tilde{A}) = \phi_{21}(\tilde{A})$ , this implies  $3a = 3b + 7x$ . For a sufficiently small  $\epsilon > 0$ , consider the perturbation matrix  $\tilde{A}_{\epsilon}$ ,

$$
\tilde{A}_{\epsilon} = \begin{bmatrix} a - \epsilon & \epsilon & 0 & 0 \\ b + \epsilon & x - \epsilon & x & x \\ 0 & y & y & y \\ 0 & y & y & y \end{bmatrix}.
$$

Then  $R_{\tilde{A}_{\epsilon}} = R_{\tilde{A}}, C_{\tilde{A}_{\epsilon}} = C_{\tilde{A}}$  but per $\tilde{A}_{\epsilon} = 6axy^2 - \frac{32}{3}$  $\frac{32}{3}\epsilon xy^2 + O(\epsilon^2) < 6axy^2 = \text{per}(\tilde{A})$  which gives us  $\phi(\tilde{A}_{\epsilon}) > \phi(\tilde{A})$  which is a contradiction. Therefore B is not a positive matrix.

Since aper  $B = per A > 0$ , we have per  $B > 0$  and hence without loss of generality we assume that

$$
B = \begin{bmatrix} x & l & m \\ 0 & y & v \\ 0 & 0 & z \end{bmatrix},
$$

where  $xyzlmv > 0$ . Consider another perturbation matrix of A

$$
\hat{A} = \begin{bmatrix} a & 0 & 0 & 0 \\ b - \epsilon & x & l & m + \epsilon \\ 0 & 0 & y & v \\ \epsilon & 0 & 0 & z - \epsilon \end{bmatrix}
$$

.

A is a matrix in  $K_4$  whose permanent value is  $axy(z - \epsilon)$  which is strictly less than perA = axyz. So  $\phi(\hat{A})$  is strictly greater than  $\phi(A)$  which is a contradiction to our assumption that A is  $\phi$  – maximizing matrix. So A is fully indecomposable.  $\Box$ 

### <span id="page-13-9"></span>References

- <span id="page-13-5"></span>[1] R. A. Brualdi and M. Newman (1965), Inequality for permanents and permanental minors, *Proc. Cambridge Philos. Soc.*, 61: 741 - 746.
- <span id="page-13-14"></span>[2] D.Z. Dokovic, On a conjecture by van der Waerden, Mat. Vesnik, 19 (1967), 566-569.
- <span id="page-13-2"></span>[3] Eberlein, P. J., and Govind S. Mudholkar.,Some remarks on the van der Waerden conjecture, Journal of Combinatorial Theory, 5.4, 386-396, (1968).
- <span id="page-13-3"></span>[4] T.H. Foregger, Remarks on a conjecture of M.Marcus and H.Minc, Linear and Multilinear Algebra, 7 (1979), 123 - 126.
- <span id="page-13-13"></span>[5] T.H. Foregger, Permanents of convex combinations of doubly stochastic matrices, Linear and Multilinear Algebra, 23 (1988), 79 - 90.
- <span id="page-13-11"></span>[6] Fuzhen Zhang, An update on a few permanent conjectures, Special Matrices 4.1, 305-316 (2016).
- [7] Gi-Sang Cheon and I.M. Wanless, An update on Minc's survey of open problems involving permanents, Linear Algebra Appl., 403. (2005), 314-342.
- <span id="page-13-7"></span><span id="page-13-4"></span>[8] F. Holens, Two aspects of doubly stochastic matrices: Permutation matrices and the minimum permanent function, Ph.D. Thesis, University of Manitoba, 1964.
- <span id="page-13-6"></span>[9] S.G Hwang, A Note on The Monotonicity of Permanents function, Discrete Mathematics, 91 (1991), 99-104.
- [10] Kirill.A.Kopotun, On some permanental conjectures, Linear and Multilinear Algebra, 36 (1994), 205-216.
- <span id="page-13-8"></span>[11] Kirill A. Kopotun, A Note on the Convexity of the Sum of Subpermanents, Linear Algebra and its Applications, 245:157-169(1996).
- <span id="page-13-10"></span>[12] Ko-wei Lih and E.T.H.Wang, A convexity inequality on the permanent of doubly stochastic matrices, Congressus Numerantium, vol. 36, 189 -198 (1982).
- <span id="page-13-0"></span>[13] M.Marcus and H.Minc, On a conjecture of B.L.van der Waerden, Proc. Cambridge Philos. Soc., 63 (1967), 305-309.
- <span id="page-13-12"></span>[14] H.Minc, Permanents, Encyclopedia Math. Appl., 6 (1978), Addison-Wesley, Reading, Mass.
- [15] Subramanian.P and Somasundaram.K, Some Conjectures on Permanents of Doubly Stochastic Matrices, Journal of Discrete Mathematical Sciences and Cryptography, 19(5- 6), 997-1011, (2016).
- <span id="page-13-1"></span>[16] E.T.H.Wang, On a conjecture of Marcus and Minc, Linear and multilinear Algebra, 5 (1977), 145-148.
- <span id="page-14-8"></span><span id="page-14-7"></span>[17] H.Tverberg, On the permanent of a bistochastic matrix, Math Scand,12, (1963), 25-35
- <span id="page-14-1"></span>[18] D.London and H.Minc, On the permanent of doubly stochastic matrices with zero diagonal, Linear and Multilinear Algebra,24 (1989), 289-300
- <span id="page-14-2"></span>[19] I.M. Wanless, The Holens-Dokovic conjecture on permanents fails, Linear Algebra Appl., 286 (1999), 273 - 285.
- <span id="page-14-3"></span>[20] Richard Sinkhorn, A problem related to the van der waerden permanent theorem, Linear and Multilinear Algebra, 16:1-4,167-173(1984)
- <span id="page-14-4"></span>[21] Suk Geun Hwang, A note on a conjecture on Permanents, Linear Algebra and its Applications, 76:31-44(1986)
- <span id="page-14-5"></span>[22] Suk Geun Hwang, On a conjecture of E. Dittert, Linear Algebra and its Applications, 95:161-169(1987)
- <span id="page-14-6"></span>[23] G.S. Cheon, I.M. Wanless, Some results towards the Dittert conjecture on permanents, Linear Algebra and its Applications, 436: 791–801 (2012)
- <span id="page-14-0"></span>[24] G.S.Cheon, H.W.Yoon, A note on the Dittert conjecture for permanents, International Mathematical Forum, 1, 2006, no. 39, 1943-1949
- [25] H.Minc, Permanents, Encyclopedia Math. Appl., 6 (1978), Addison-Wesley, Reading, Mass.
- [26] Suk Geun Hwang, Mun Gu Sohn, Si Ju Kim, The Dittert's function on a set of nonnegative matrices, International Journal of Mathematics and Mathematical Sciences, 13(4), 709- 716(1990)
- [27] G.S.Cheon, On the monotonicity of the Dittert function on classes of nonnegative matrices, Bulletin of the Korean Mathematical Society, 30(2), 265-275(1993)