# Extremal graphs without long paths and a given graph<sup>\*</sup>

Yichong Liu, Liying Kang<sup>†</sup>

Department of Mathematics, Shanghai University, Shanghai 200444, P.R. China

#### Abstract

For a family of graphs  $\mathcal{F}$ , the Turán number  $ex(n, \mathcal{F})$  is the maximum number of edges in an *n*-vertex graph containing no member of  $\mathcal{F}$  as a subgraph. The maximum number of edges in an *n*-vertex connected graph containing no member of  $\mathcal{F}$  as a subgraph is denoted by  $ex_{conn}(n, \mathcal{F})$ . Let  $P_k$  be the path on k vertices and H be a graph with chromatic number more than 2. Katona and Xiao [Extremal graphs without long paths and large cliques, European J. Combin., 2023 103807] posed the following conjecture: Suppose that the chromatic number of H is more than 2. Then  $ex(n, \{H, P_k\}) = n \max\{\lfloor \frac{k}{2} \rfloor -1, \frac{ex(k-1,H)}{k-1}\} + O_k(1)$ . In this paper, we determine the exact value of  $ex_{conn}(n, \{P_k, H\})$  for sufficiently large n. Moreover, we obtain asymptotical result for  $ex(n, \{P_k, H\})$ , which solves the conjecture proposed by Katona and Xiao.

Keywords: Turán number, extremal graph, chromatic number, path

AMS (2000) subject classification: 05C35

#### 1 Introduction

Let G = (V, E) be a simple graph with vertex set V = V(G) and edge set E = E(G). The number of edges in G is denoted by e(G). For  $S \subseteq V(G)$ , we denote by  $N_G^{co}(S)$  the common neighborhood of vertices of S in V(G). Denote by G[S] the graph induced by S, and denote by  $G \setminus S$  the graph obtained from G by deleting all vertices of S and all edges incident with S. For  $V_1, V_2 \subseteq V(G)$ ,  $E(V_1, V_2)$  denotes the set of edges between  $V_1$  and  $V_2$  in G, and  $e(V_1, V_2) = |E(V_1, V_2)|$ . The chromatic number of G is denoted by  $\chi(G)$ .

 $<sup>^{*}</sup>$ Research was partially supported by the National Nature Science Foundation of China (grant number 12331012

<sup>&</sup>lt;sup>†</sup>Corresponding author. Email address: lykang@shu.edu.cn (L. Kang), lyc328az@163.com (Y. Liu)

Let  $\mathcal{F}$  be a family of graphs. A graph G is called  $\mathcal{F}$ -free if G does not contain any member of the graphs in  $\mathcal{F}$  as a subgraph. The Turán number, denoted by  $ex(n, \mathcal{F})$ , is the maximum number of edges in an *n*-vertex graph containing no member of  $\mathcal{F}$  as a subgraph. We call an *n*-vertex  $\mathcal{F}$ -free graph attaining  $ex(n, \mathcal{F})$  edges an *extremal graph* for  $\mathcal{F}$ . The family of extremal graphs for  $\mathcal{F}$  is denoted by  $EX(n, \mathcal{F})$ . The maximum number of edges in an *n*-vertex connected graph containing no member of  $\mathcal{F}$  as a subgraph is denoted by  $ex_{conn}(n, \mathcal{F})$ , and the family of *n*-vertex  $\mathcal{F}$ -free connected graphs attaining  $ex_{conn}(n, \mathcal{F})$  edges is denoted by  $EX_{conn}(n, \mathcal{F})$ .

Let  $K_n$  denote the complete graph on n vertices. Let  $P_k$  and  $C_k$  denote the path and the cycle on k vertices respectively. The n-vertex independent set is denoted by  $I_n$ . The complete k-partite graph  $K_{n_1,n_2,\dots,n_k}$  is a graph formed by partitioning the set of n vertices into k subsets with  $n_i(1 \le i \le k)$  vertices in each subset, and connecting two vertices by an edge if and only if they belong to different subsets. The Turán graph T(n,k) is a complete multipartite graph formed by partitioning the set of n vertices into k subsets with size as equal as possible. For two graphs G and H, the disjoint union of G and H is denoted by  $G \cup H$ . The *join* of G and H, denoted by  $G \vee H$ , is the graph obtained from  $G \cup H$  by adding all possible edges between G and H.

In 1959, Erdős and Gallai [4] determined the Turán number for  $P_k$ .

**Theorem 1.1** ([4]). Let G be an n-vertex graph. If  $e(G) > \frac{k-2}{2}n$  where  $k \ge 2$ , then G contains a copy of  $P_k$ .

The extremal problem for  $P_k$  under the condition that G is connected was considered by Kopylov [6]. He determined the Turán number for  $P_k$ . After 30 years, Balister, Győri, Lehel and Schelp [2] found all the extremal graphs for  $P_k$ .

**Theorem 1.2** ([2]). Let G be a connected graph on n vertices containing no path on k vertices where  $n > k \ge 4$ . Then

$$e(G) \le \max\left\{ \binom{k-2}{2} + (n-k+2), \binom{\left\lceil \frac{k}{2} \right\rceil}{2} + \left\lfloor \frac{k-2}{2} \right\rfloor \left(n - \left\lceil \frac{k}{2} \right\rceil \right) \right\}.$$

The equality holds when G is either  $(K_{k-3} \cup I_{n-k+2}) \vee K_1$  or  $(K_{k-2\lfloor \frac{k}{2} \rfloor+1} \cup I_{n-\lceil \frac{k}{2} \rceil}) \vee K_{\lfloor \frac{k}{2} \rfloor-1}$ .

Recently, Katona and Xiao [5] determined the exact value of  $ex(n, \{P_k, K_m\})$  if k > 2m - 1and  $ex_{conn}(n, \{P_k, K_m\})$  if k > m for sufficiently large n.

**Theorem 1.3** ([5]). Let G be a connected n-vertex  $\{K_m, P_k\}$ -free graph where m < k. For sufficiently large n (> N(k)),

$$ex_{conn}(n, \{K_m, P_k\}) = \left(\left\lfloor \frac{k}{2} \right\rfloor - 1\right) \left(n - \left\lfloor \frac{k}{2} \right\rfloor + 1\right) + e\left(T\left(\left\lfloor \frac{k}{2} \right\rfloor - 1, m - 2\right)\right).$$

 $T\left(\lfloor \frac{k}{2} \rfloor - 1, m - 2\right) \vee I_{n - \lfloor \frac{k}{2} \rfloor + 1} \text{ is an extremal graph.}$ 

**Theorem 1.4** ([5]). Let G be an n-vertex  $\{K_m, P_k\}$ -free graph where 2m - 1 < k. For sufficiently large n (> N'(k)),

$$ex(n, \{K_m, P_k\}) = \left(\left\lfloor \frac{k}{2} \right\rfloor - 1\right) \left(n - \left\lfloor \frac{k}{2} \right\rfloor + 1\right) + e\left(T\left(\left\lfloor \frac{k}{2} \right\rfloor - 1, m - 2\right)\right).$$

 $T\left(\lfloor \frac{k}{2} \rfloor - 1, m - 2\right) \vee I_{n - \lfloor \frac{k}{2} \rfloor + 1} \text{ is an extremal graph.}$ 

Katona and Xiao [5] proposed to study  $ex(n, \{P_k, H\})$  for an H with  $\chi(H) > 2$  and posed the following conjecture.

Conjecture 1.5 ([5]). Suppose  $\chi(H) > 2$ . Then

$$ex(n, \{H, P_k\}) = n \max\left\{ \left\lfloor \frac{k}{2} \right\rfloor - 1, \frac{ex(k-1, H)}{k-1} \right\} + O_k(1).$$

In this paper, we determine the exact value of  $ex_{conn}(n, \{P_k, H\})$  for sufficiently large n and obtain asymptotical result of  $ex(n, \{P_k, H\})$  which confirms Conjecture 1.5.

Let  $\mathcal{H}$  be the family of graphs obtained from H by deleting a color class of H. Our main results are the following.

**Theorem 1.6.** If H is a graph with  $\chi(H) > 2$  and k is even, then

$$ex_{conn}(n, \{P_k, H\}) = ex\left(\frac{k}{2} - 1, \mathcal{H}\right) + \left(\frac{k}{2} - 1\right)\left(n - \frac{k}{2} + 1\right)$$

for sufficiently large n. The extremal graph is  $T \vee I_{n-k/2+1}$ , where  $T \in EX(\frac{k}{2}-1,\mathcal{H})$ .

**Theorem 1.7.** If H is a graph with  $\chi(H) > 2$  and k is odd, then

$$ex_{conn}(n, \{P_k, H\}) = ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\left(n - \frac{k-3}{2}\right) + c$$

for sufficiently large n, where c = 0 or 1.

The following theorem gives a sufficient condition for c = 0 in Theorem 1.7. Let  $\mathcal{H}'$  be the family of graphs obtained from H by deleting two adjacent vertices in V(H).

**Theorem 1.8.** If H is a graph with  $\chi(H) > 2$ , k is odd, and every graph in  $EX(\frac{k-3}{2}, \mathcal{H})$  contains at least a member in  $\mathcal{H}'$ , then

$$ex_{conn}(n, \{P_k, H\}) = ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\left(n - \frac{k-3}{2}\right)$$

for sufficiently large n. The extremal graph is  $T \vee I_{n-(k-3)/2}$ , where T is a graph in  $EX(\frac{k-3}{2},\mathcal{H})$ .

We will show that Theorem 1.3 can be deduced from Theorem 1.8 in Section 3.

**Theorem 1.9.** Suppose n is sufficiently large and  $\chi(H) > 2$ . Then

$$ex(n, \{H, P_k\}) = n \max\left\{ \left\lfloor \frac{k}{2} \right\rfloor - 1, \frac{ex(k-1, H)}{k-1} \right\} + O_k(1).$$

The rest of the paper is organized as follows. In Section 2, we present some preliminaries and lemmas. The proofs of main results will be given in Sections 3 and 4. We give some discussion in the last section.

#### 2 Preliminaries

The following results can be found in [5].

**Lemma 2.1** ([5]). Let G be a connected graph on n vertices with a path P on k-1 vertices but no path on k vertices. Let  $u \in V(G) \setminus V(P)$  be a vertex adjacent to  $s \ge 1$  vertices of P and assume a longest path Q in  $G \setminus V(P)$  starting at u has  $j (j \ge 0)$  vertices. Then,  $s+j \le \lfloor \frac{k}{2} \rfloor$ .

**Lemma 2.2** ([5]). Let G be a connected graph on k vertices with no Hamiltonian path but with a path  $P = (v_1, v_2, \dots, v_{k-1})$  on k-1 vertices. Suppose the vertex  $u \in V(G) \setminus V(P)$  is adjacent to s vertices of P, that is  $N_P(u) = \{v_{i_1}, v_{i_2}, \dots, v_{i_s}\}, i_1 < i_2 < \dots < i_s$ . Then (1)  $s \leq \lfloor \frac{k}{2} \rfloor - 1$ ; (2) there are no edges of the form  $v_{i_j+1}v_{i_r+1}, v_{i_j-1}v_{i_r-1}, v_1v_{i_t+1}$  or  $v_{i_t-1}v_{k-1}, 1 \leq j < r \leq s, 1 \leq t \leq s$ .

By Lemma 2.2, if G is an n-vertex  $P_k$ -free connected graph with a path  $P_{k-1}$ , then  $e(u, V(P_{k-1})) \le \lfloor \frac{k}{2} \rfloor - 1$  for each  $u \in V(G) \setminus V(P_{k-1})$ . We partition the vertices of  $V(G) \setminus V(P_{k-1})$  into mutually disjoint sets in two different ways.

(1) Let  $u \in A_i$ , if  $e(u, V(P_{k-1})) = i, 0 \le i \le \lfloor \frac{k}{2} \rfloor - 1$ .

(2) Let  $u \in B_i$ , if  $u \in A_{i_1}$  and u is connected to vertices in  $A_{i_2}, A_{i_3}, \dots, A_{i_r}$  by a path, then  $i = \max\{i_1, i_2, \dots, i_r\}$ .

One can easily verify that  $A_i \cap A_j = \emptyset$ ,  $B_i \cap B_j = \emptyset$  for  $0 \le i \ne j \le \lfloor \frac{k}{2} \rfloor - 1$  and

$$\sum_{i=0}^{\lfloor k/2 \rfloor - 1} A_i = \sum_{i=0}^{\lfloor k/2 \rfloor - 1} B_i = V(G) \setminus V(P_{k-1}).$$

Moreover,  $B_0$  is empty since G is connected, and there are no edges between  $B_i$  and  $B_j$  for  $1 \le i \ne j \le \lfloor \frac{k}{2} \rfloor - 1$ .

**Lemma 2.3** ([5]). Let G be an n-vertex  $P_k$ -free connected graph with a path on k-1 vertices. Let  $A_i$  and  $B_i$  are the sets of vertices defined above. Then

$$e(G) - e(G[V(P_{k-1})]) \le \left(\left\lfloor \frac{k}{2} \right\rfloor - 1\right) \left| A_{\lfloor k/2 \rfloor - 1} \right| + \sum_{l=0}^{\lfloor k/2 \rfloor - 2} \left(\left\lfloor \frac{k}{2} \right\rfloor - \frac{3}{2}\right) |A_l|.$$

Now suppose G is an n-vertex  $\{P_k, H\}$ -free connected graph, and G contains a path on k-1 vertices. The following lemma estimates the number of edges in  $G[V(P_{k-1})]$ .

**Lemma 2.4.** For a graph H with  $\chi(H) > 2$  and sufficiently large n, let G be an n-vertex  $\{P_k, H\}$ -free connected graph containing a  $P_{k-1}$ . Let  $\mathcal{H}$  be the family of graphs obtained from H by deleting a color class of H. If  $|A_{\lfloor k/2 \rfloor - 1}| = \Theta(n)$ , then

$$e(G[V(P_{k-1})]) \le ex\left(\left\lfloor \frac{k}{2} \right\rfloor - 1, \mathcal{H}\right) + \left(\left\lfloor \frac{k}{2} \right\rfloor - 1\right) \left\lceil \frac{k}{2} \right\rceil + c,$$

where c = 0 if k is even and c = 1 otherwise.

*Proof.* Let  $P_{k-1} := (v_1, v_2, \dots, v_{k-1})$ . We distinguish two different cases depending on the parity of k.

Case 1: k is even. Since for any vertex u in  $A_{k/2-1}$ , u cannot be adjacent to  $v_1$ ,  $v_{k-1}$ , or consecutive vertices on  $P_{k-1}$ ,  $N_{P_{k-1}}(u) = \{v_2, v_4, \cdots, v_{k-2}\}$ . Then  $A_{k/2-1}$  is the common neighborhood of vertices in  $\{v_2, v_4, \cdots, v_{k-2}\}$ . The assumption  $|A_{k/2-1}| = \Theta(n)$  implies that  $G[\{v_2, v_4, \cdots, v_{k-2}\}]$  is  $\mathcal{H}$ -free. Otherwise,  $G[\{v_2, v_4, \cdots, v_{k-2}\}]$  contains a graph in  $\mathcal{H}$ , we can always find the missing color class in  $A_{k/2-1}$  which forms an H, a contradiction. Thus  $G[\{v_2, v_4, \cdots, v_{k-2}\}]$  must be  $\mathcal{H}$ -free. Moreover, by Lemma 2.2, there are no edges in  $G[\{v_1, v_3, \cdots, v_{k-1}\}]$ . Then  $e(G[V(P_{k-1})]) \leq ex(\frac{k}{2}-1,\mathcal{H}) + (\frac{k}{2}-1)\frac{k}{2}$ .

Case 2: k is odd. Since for any vertex u in  $A_{(k-3)/2}$ , u cannot be adjacent to  $v_1, v_{k-1}$ , or consecutive vertices on  $P_{k-1}, N_{P_{k-1}}(u) = \{v_2, v_4, \cdots, v_{k-3}\}$  or  $N_{P_{k-1}}(u) = \{v_2, v_4, \cdots, v_{2l}, v_{2l+3}, v_{2l+5}, \cdots, v_{k-2}\}$  where  $1 \le l \le \frac{k-5}{2}$ . Let  $G_1 = G \setminus V(P_{k-1})$ . Then

$$A_{(k-3)/2} = \bigcup_{l=1}^{(k-5)/2} N_{G_1}^{co}(\{v_2, v_4, \cdots, v_{2l}, v_{2l+3}, v_{2l+5}, \cdots, v_{k-2}\}) \cup N_{G_1}^{co}(\{v_2, v_4, \cdots, v_{k-3}\}).$$

Since  $|A_{(k-3)/2}| = \Theta(n)$ , we must have  $|N_{G_1}^{co}(\{v_2, v_4, \cdots, v_{k-3}\})| = \Theta(n)$  or  $|N_{G_1}^{co}(\{v_2, v_4, \cdots, v_{2l}, v_{2l+3}, v_{2l+5}, \cdots, v_{k-2}\})| = \Theta(n)$  for some l.

For the former case,  $G[\{v_2, v_4, \cdots, v_{k-3}\}]$  is  $\mathcal{H}$ -free. Moreover, by Lemma 2.2, there are no

edges in  $G[\{v_1, v_3, \cdots, v_{k-2}\}]$  and  $v_1v_{k-1}, v_3v_{k-1}, \cdots, v_{k-4}v_{k-1}$  are not edges of G. Then

$$e(G[V(P_{k-1})]) \leq ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \left(\frac{k-1}{2}\right)^2 - \frac{k-3}{2} + \frac{k-3}{2}$$
$$= ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \left(\frac{k-1}{2}\right)^2$$
$$= ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\frac{k+1}{2} + 1.$$

For the latter case,  $G[\{v_2, v_4, \cdots, v_{2l}, v_{2l+3}, v_{2l+5}, \cdots, v_{k-2}\}]$  is  $\mathcal{H}$ -free. Moreover, by Lemma 2.2, there are no edges in  $G[\{v_1, v_3, \cdots, v_{2l+1}, v_{2l+4}, v_{2l+6}, \cdots, v_{k-1}\}]$  and  $v_1v_{2l+2}, v_3v_{2l+2}, \cdots, v_{2l-1}v_{2l+2}, v_{2l+2}v_{2l+4}, v_{2l+2}v_{2l+6}, \cdots, v_{2l+2}v_{k-1}]$  are not edges of G. Then

$$e(G[V(P_{k-1})]) \leq ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \left(\frac{k-1}{2}\right)^2 - \frac{k-3}{2} + \frac{k-3}{2}$$
$$= ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \left(\frac{k-1}{2}\right)^2$$
$$= ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\frac{k+1}{2} + 1.$$

**Lemma 2.5.** Suppose  $V(G) = M \cup \{u, v\}$ , where |M| = l,  $\mathcal{H}'$  is the family of graphs obtained from H by deleting two adjacent vertices in V(H). If G[M] is  $\mathcal{H}$ -free, G is H-free and every graph in  $EX(l, \mathcal{H})$  contains at least a member in  $\mathcal{H}'$ . Then  $e(G) \leq ex(l, \mathcal{H}) + 2l$ .

Proof. Since G[M] is  $\mathcal{H}$ -free,  $e(G[M]) \leq ex(l, \mathcal{H})$ . If  $e(G) = ex(l, \mathcal{H}) + 2l + 1$ , then  $G \cong T \bigvee K_2$ where T is a graph in  $EX(l, \mathcal{H})$ . However, since every graph in  $EX(l, \mathcal{H})$  contains at least a member in  $\mathcal{H}'$ , G must contains H as a subgraph, a contradiction.

**Lemma 2.6.** For odd k, let G be the graph in Lemma 2.4 and  $|A_{(k-3)/2}| = \Theta(n)$ ,  $\mathcal{H}'$  be the family of graphs obtained from H by deleting two adjacent vertices in V(H). If every graph in  $EX(\frac{k-3}{2},\mathcal{H})$  contains at least a member in  $\mathcal{H}'$ , then

$$e(G[V(P_{k-1})]) \le ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\frac{k+1}{2}$$

for sufficiently large n.

*Proof.* As in the proof of Lemma 2.4, we have

$$A_{(k-3)/2} = \bigcup_{l=1}^{(k-5)/2} N_{G_1}^{co} \left( \{ v_2, v_4, \cdots, v_{2l}, v_{2l+3}, v_{2l+5}, \cdots, v_{k-2} \} \right) \cup N_{G_1}^{co} \left( \{ v_2, v_4, \cdots, v_{k-3} \} \right).$$

The assumption  $|A_{(k-3)/2}| = \Theta(n)$  implies that  $|N_{G_1}^{co}(\{v_2, v_4, \cdots, v_{k-3}\})| = \Theta(n)$  or there exists an l with  $1 \le l \le \frac{k-5}{2}$  such that  $|N_{G_1}^{co}(\{v_2, v_4, \cdots, v_{2l}, v_{2l+3}, v_{2l+5}, \cdots, v_{k-2}\})| = \Theta(n)$ .

If  $|N_{G_1}^{co}(\{v_2, v_4, \cdots, v_{k-3}\})| = \Theta(n)$ , then  $G[\{v_2, v_4, \cdots, v_{k-3}\}]$  is  $\mathcal{H}$ -free, and  $G[\{v_2, v_4, \cdots, v_{k-3}, v_{k-2}, v_{k-1}\}]$  is H-free. By Lemma 2.5,

$$e(G[\{v_2, v_4, \cdots, v_{k-3}, v_{k-2}, v_{k-1}\}]) \le ex(\frac{k-3}{2}, \mathcal{H}) + k - 3$$

Moreover, by Lemma 2.2, there are no edges in  $G[\{v_1, v_3, \cdots, v_{k-2}\}]$  and  $v_1v_{k-1}, v_3v_{k-1}, \cdots, v_{k-4}v_{k-1}$  are not edges of G. Then

$$e(G[V(P_{k-1})]) \leq ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \left(\frac{k-3}{2}\right)^2 + k - 3$$
$$= ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\frac{k+1}{2}.$$

If  $|N_{G_1}^{co}(\{v_2, v_4, \cdots, v_{2l}, v_{2l+3}, v_{2l+5}, \cdots, v_{k-2}\})| = \Theta(n)$  for some l, then  $G[\{v_2, v_4, \cdots, v_{2l}, v_{2l+3}, v_{2l+5}, \cdots, v_{k-2}\}]$  is  $\mathcal{H}$ -free, and  $G[\{v_2, v_4, \cdots, v_{2l}, v_{2l+3}, v_{2l+5}, \cdots, v_{k-2}, v_{2l+1}, v_{2l+2}\}]$  is  $\mathcal{H}$ -free. By Lemma 2.5,

$$e(G[\{v_2, v_4, \cdots, v_{2l}, v_{2l+3}, v_{2l+5}, \cdots, v_{k-2}, v_{2l+1}, v_{2l+2}\}]) \le ex\left(\frac{k-3}{2}, \mathcal{H}\right) + k - 3.$$

Moreover, by Lemma 2.2, there are no edges in  $G[\{v_1, v_3, \cdots, v_{2l+1}, v_{2l+4}, v_{2l+6}, \cdots, v_{k-1}\}]$  and  $v_1v_{2l+2}, v_3v_{2l+2}, \cdots, v_{2l-1}v_{2l+2}, v_{2l+2}v_{2l+4}, v_{2l+2}v_{2l+6}, \cdots, v_{2l+2}v_{k-1}]$  are not edges of G. Then

$$e(G[V(P_{k-1})]) \leq ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \left(\frac{k-3}{2}\right)^2 + k - 3$$
$$= ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\frac{k+1}{2}.$$

Let  $M_t$  be a matching on t edges and  $S_t$  be a star on t+1 vertices. In 1972, Abbott, Hanson and Sauer[1] determined  $ex(n, \{M_t, S_t\})$ .

**Theorem 2.1** ([1]). We omit isolated vertices in extremal graphs. If k is odd and  $n \ge 2t$ ,  $ex(n, \{M_t, S_t\}) = t^2 - t$  and the extremal graph is  $K_t \cup K_t$ . When k is even and  $n \ge 2t - 1$ ,  $ex(n, \{M_t, S_t\}) = t^2 - \frac{3}{2}t$  and the extremal graphs are all the graphs with 2t - 1 vertices,  $t^2 - \frac{3}{2}t$ edges and maximum degree t - 1.

We also need the stronger version proved by Chvátal and Hanson [3].

**Theorem 2.2** ([3]). For every  $k \ge 1$  and  $t \ge 1$ ,

$$ex(n, \{M_{k+1}, S_{t+1}\}) = kt + \left\lfloor \frac{t}{2} \right\rfloor \cdot \left\lfloor \frac{k}{\lceil t/2 \rceil} \right\rfloor \le kt + k.$$

### 3 Proofs of Theorems 1.6, 1.7 and 1.8

**Proof of Theorems 1.6.** For any graph T in  $EX(\frac{k}{2}-1,\mathcal{H})$ , obviously  $T \vee I_{n-k/2+1}$  is a  $\{P_k, H\}$ -free connected graph. Then

$$ex_{conn}(n, \{P_k, H\}) \ge ex\left(\frac{k}{2} - 1, \mathcal{H}\right) + \left(\frac{k}{2} - 1\right)\left(n - \frac{k}{2} + 1\right).$$
 (3.1)

Next we show that  $ex_{conn}(n, \{P_k, H\}) \leq ex(\frac{k}{2} - 1, \mathcal{H}) + (\frac{k}{2} - 1)(n - \frac{k}{2} + 1)$ . Let  $G \in EX_{conn}(n, \{P_k, H\})$ .

If  $P_{k-1} \nsubseteq G$ , by Theorem 1.1,

$$e(G) \le \frac{k-3}{2}n < ex\left(\frac{k}{2}-1,\mathcal{H}\right) + \left(\frac{k}{2}-1\right)\left(n-\frac{k}{2}+1\right)$$

for sufficiently large n.

If  $P_{k-1} \subseteq G$ , we claim that  $|A_{k/2-1}| = \Theta(n)$ . Otherwise,  $|A_{k/2-1}| = o(n)$ . By Lemma 2.3,

$$e(G) - e(G[V(P_{k-1})]) \le \left(\frac{k}{2} - 1\right) \left| A_{k/2-1} \right| + \sum_{l=0}^{k/2-2} \left(\frac{k}{2} - \frac{3}{2}\right) |A_l|$$
  
$$= \left(\frac{k}{2} - 1\right) \left| A_{k/2-1} \right| + \left(\frac{k}{2} - \frac{3}{2}\right) \left(n - k + 1 - \left| A_{k/2-1} \right| \right)$$
  
$$= \left(\frac{k}{2} - \frac{3}{2}\right) n + o(n).$$

Combining with  $e(G[V(P_{k-1})]) \leq ex(k-1, H)$ , we have

$$e(G) \leq ex(k-1,H) + \left(\frac{k}{2} - \frac{3}{2}\right)n + o(n)$$
  
$$< ex\left(\frac{k}{2} - 1, \mathcal{H}\right) + \left(\frac{k}{2} - 1\right)\left(n - \frac{k}{2} + 1\right)$$

for sufficiently large n, which is a contradiction to (3.1). So  $|A_{k/2-1}| = \Theta(n)$ . By Lemma 2.4,

$$e(G[V(P_{k-1})]) \le ex\left(\frac{k}{2} - 1, \mathcal{H}\right) + \left(\frac{k}{2} - 1\right)\frac{k}{2}.$$

According to Lemma 2.3,

$$e(G) - e(G[V(P_{k-1})]) \leq \left(\frac{k}{2} - 1\right) \left| A_{k/2-1} \right| + \sum_{l=0}^{k/2-2} \left(\frac{k}{2} - \frac{3}{2}\right) |A_l|$$
$$\leq \left(\frac{k}{2} - 1\right) (n - k + 1).$$

Then

$$e(G) \leq ex\left(\frac{k}{2}-1,\mathcal{H}\right) + \left(\frac{k}{2}-1\right)\frac{k}{2} + \left(\frac{k}{2}-1\right)(n-k+1)$$
$$= ex\left(\frac{k}{2}-1,\mathcal{H}\right) + \left(\frac{k}{2}-1\right)\left(n-\frac{k}{2}+1\right).$$

So  $ex_{conn}(n, \{P_k, H\}) = ex(\frac{k}{2}-1, \mathcal{H}) + (\frac{k}{2}-1)(n-\frac{k}{2}+1)$ . The extremal graph is  $T \vee I_{n-k/2+1}$ , where  $T \in EX(\frac{k}{2}-1, \mathcal{H})$ .

**Proof of Theorem 1.7.** For any graph T in  $EX(\frac{k-3}{2}, \mathcal{H})$ , obviously  $T \vee I_{n-(k-3)/2}$  is a  $\{P_k, H\}$ -free connected graph. Then

$$ex_{conn}(n, \{P_k, H\}) \ge ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\left(n - \frac{k-3}{2}\right).$$
 (3.2)

Next we will show that  $ex_{conn}(n, \{P_k, H\}) \leq ex(\frac{k-3}{2}, \mathcal{H}) + \frac{k-3}{2}(n - \frac{k-3}{2}) + 1$ . Let  $G \in EX_{conn}(n, \{P_k, H\})$ . If  $P_{k-2} \not\subseteq G$ , by Theorem 1.1,  $e(G) \leq \frac{k-4}{2}n < ex(\frac{k-3}{2}, \mathcal{H}) + \frac{k-3}{2}(n - \frac{k-3}{2})$  for sufficiently large n.

If  $P_{k-1} \nsubseteq G$ ,  $P_{k-2} \subseteq G$ , we claim that  $|A_{(k-3)/2}| = \Theta(n)$ . Otherwise,  $|A_{(k-3)/2}| = o(n)$ . By Lemma 2.3,

$$e(G) - e(G[V(P_{k-2})]) \le \frac{k-3}{2} |A_{(k-3)/2}| + \sum_{l=0}^{(k-5)/2} \frac{k-4}{2} |A_l|$$
  
$$= \frac{k-3}{2} |A_{(k-3)/2}| + \frac{k-4}{2} (n-k+2 - |A_{(k-3)/2}|)$$
  
$$= \frac{k-4}{2} n + o(n).$$

Combining with  $e(G[V(P_{k-2})]) \leq ex(k-2, H)$ , we have

$$e(G) \leq ex(k-2,H) + \frac{k-4}{2}n + o(n)$$
  
<  $ex\left(\frac{k-3}{2},\mathcal{H}\right) + \frac{k-3}{2}\left(n - \frac{k-3}{2}\right)$ 

for sufficiently large n, which is a contradiction to (3.2). So  $|A_{(k-3)/2}| = \Theta(n)$ .

By Lemmas 2.4 and 2.3, we have

$$e(G[V(P_{k-2})]) \le ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\frac{k-1}{2},$$

and

$$e(G) - e(G[V(P_{k-2})]) \leq \frac{k-3}{2} |A_{(k-3)/2}| + \sum_{l=0}^{(k-5)/2} \frac{k-4}{2} |A_l|$$
$$\leq \frac{k-3}{2} (n-k+2).$$

Then

$$e(G) \leq ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\frac{k-1}{2} + \frac{k-3}{2}(n-k+2)$$
  
=  $ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\left(n - \frac{k-3}{2}\right).$ 

If  $P_{k-1} \subseteq G$ , we claim that  $|A_{(k-3)/2}| = \Theta(n)$ . Otherwise,  $|A_{(k-3)/2}| = o(n)$ . By Lemma 2.3,

$$e(G) - e(G[V(P_{k-1})]) \le \frac{k-3}{2} |A_{(k-3)/2}| + \sum_{l=0}^{(k-5)/2} \frac{k-4}{2} |A_l|$$
  
=  $\frac{k-3}{2} |A_{(k-3)/2}| + \frac{k-4}{2} (n-k+1-|A_{(k-3)/2}|)$   
=  $\frac{k-4}{2} n + o(n).$ 

Combining with  $e(G[V(P_{k-1})]) \leq ex(k-1, H)$ , we have

$$e(G) \leq ex(k-1,H) + \frac{k-4}{2}n + o(n)$$
  
<  $ex\left(\frac{k-3}{2},\mathcal{H}\right) + \frac{k-3}{2}\left(n - \frac{k-3}{2}\right),$ 

for sufficiently large n, which is a contradiction to (3.2). So  $|A_{(k-3)/2}| = \Theta(n)$ . By Lemmas 2.4 and 2.3, we have

$$e(G[V(P_{k-1})]) \le ex(\frac{k-3}{2}, \mathcal{H}) + \frac{k-3}{2}\frac{k+1}{2} + 1,$$

and

$$e(G) - e(G[V(P_{k-1})]) \leq \frac{k-3}{2} |A_{(k-3)/2}| + \sum_{l=0}^{(k-5)/2} \frac{k-4}{2} |A_l|$$
$$\leq \frac{k-3}{2} (n-k+1).$$

Then

$$e(G) \leq ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\frac{k+1}{2} + 1 + \frac{k-3}{2}(n-k+1)$$
  
=  $ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\left(n - \frac{k-3}{2}\right) + 1.$ 

Consequently,  $ex_{conn}(n, \{P_k, H\}) = ex(\frac{k-3}{2}, \mathcal{H}) + \frac{k-3}{2}(n - \frac{k-3}{2}) + c$ , where c = 0 or 1.  $\square$ **Proof of Theorem 1.8.** Let  $G \in EX_{conn}(n, \{P_k, H\})$ . According to the proof of Theorem

1.7, we only need to consider the case  $P_{k-1} \subseteq G$ . As in the proof of Theorem 1.7, we can show that  $|A_{(k-3)/2}| = \Theta(n)$ . By Lemma 2.6,

$$e(G[V(P_{k-1})]) \le ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\frac{k+1}{2}$$

Then

$$e(G) \leq ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\frac{k+1}{2} + \frac{k-3}{2}(n-k+1)$$
  
=  $ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\left(n - \frac{k-3}{2}\right).$ 

 $\operatorname{So}$ 

$$ex_{conn}(n, \{P_k, H\}) \le ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\left(n - \frac{k-3}{2}\right).$$

Combining with the lower bound

$$ex_{conn}(n, \{P_k, H\}) \ge ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\left(n - \frac{k-3}{2}\right),$$

we have

$$ex_{conn}(n, \{P_k, H\}) = ex\left(\frac{k-3}{2}, \mathcal{H}\right) + \frac{k-3}{2}\left(n - \frac{k-3}{2}\right).$$

Moreover,  $T \vee I_{n-(k-3)/2}$  is an extremal graph, where T is a graph in  $EX\left(\frac{k-3}{2}, \mathcal{H}\right)$ .

We show that the above results imply Theorem 1.3. When k is even, Theorem 1.6 yields that

$$ex_{conn}(n, \{P_k, K_m\}) = ex\left(\frac{k}{2} - 1, K_{m-1}\right) + \left(\frac{k}{2} - 1\right)\left(n - \frac{k}{2} + 1\right)$$
$$= e\left(T\left(\frac{k}{2} - 1, m - 2\right)\right) + \left(\frac{k}{2} - 1\right)\left(n - \frac{k}{2} + 1\right),$$

and the extremal graph is  $T(\frac{k}{2}-1, m-2) \vee I_{n-k/2+1}$ . When k is odd, we have the following corollary.

**Corollary 1.** For k is odd, if  $k \ge 2m - 1$ , then  $ex_{conn}(n, \{K_m, P_k\}) = e(T(\frac{k-3}{2}, m-2)) + \frac{k-3}{2}(n-\frac{k-3}{2})$  for sufficiently large n, and the extremal graph is  $T(\frac{k-3}{2}, m-2) \lor I_{n-(k-3)/2}$ . If m < k < 2m - 1,  $ex_{conn}(n, \{K_m, P_k\}) = \frac{(k-3)(k-5)}{8} + \frac{k-3}{2}(n-\frac{k-3}{2}) + 1$  and the extremal graph is  $K_{(k-3)/2} \lor (I_{n-(k-1)/2} \cup K_2)$ .

Proof. Take  $H = K_m$ , then  $\mathcal{H} = \{K_{m-1}\}$ ,  $EX(\frac{k-3}{2}, K_{m-1}) = \{T(\frac{k-3}{2}, m-2)\}$  and  $\mathcal{H}' = \{K_{m-2}\}$ . If  $k \ge 2m-1$ , since  $T(\frac{k-3}{2}, m-2)$  contains a  $K_{m-2}$ , by Theorem 1.8,  $ex_{conn}(n, \{P_k, K_m\}) = e(T(\frac{k-3}{2}, m-2)) + \frac{k-3}{2}(n-\frac{k-3}{2})$  and the extremal graph is  $T(\frac{k-3}{2}, m-2) \lor I_{n-(k-3)/2}$ .

If m < k < 2m - 1, it is easily seen that  $K_{(k-3)/2} \vee (I_{n-(k-1)/2} \cup K_2)$  is  $\{P_k, K_m\}$ -free, by Theorem 1.7, we have

$$ex_{conn}(n, \{K_m, P_k\}) = e\left(T\left(\frac{k-3}{2}, m-2\right)\right) + \frac{k-3}{2}\left(n-\frac{k-3}{2}\right) + 1$$
$$= \frac{(k-3)(k-5)}{8} + \frac{k-3}{2}\left(n-\frac{k-3}{2}\right) + 1.$$

Take  $H = F_t$ , where  $F_t$  is the graph formed by t triangles intersecting at a vertex. Then  $\mathcal{H} = \{M_t, S_t\}$  and  $\mathcal{H}' = \{F_{t-1}, M_{t-1} \cup K_1\}$ . By Theorem 2.1, when k is odd, the extremal graph for  $\{M_t, S_t\}$  is  $K_t \cup K_t$ . When k is even, the extremal graphs for  $\{M_t, S_t\}$  are all the graphs with 2t - 1 vertices,  $t^2 - \frac{3}{2}t$  edges and maximum degree t - 1.

**Corollary 2.** For sufficiently large n,  $ex_{conn}(n, \{F_t, P_k\}) = ex(\lfloor \frac{k}{2} \rfloor - 1, \{M_t, S_t\}) + (\lfloor \frac{k}{2} \rfloor - 1)(n - \lfloor \frac{k}{2} \rfloor + 1)$  and the extremal graph is  $T \vee I_{n - \lfloor k/2 \rfloor + 1}$ , where  $T \in EX(\lfloor \frac{k}{2} \rfloor - 1, \{M_t, S_t\})$ .

*Proof.* Take  $H = F_t$ . When k is even, Theorem 1.6 yields

$$ex_{conn}(n, \{P_k, F_t\}) = ex\left(\frac{k}{2} - 1, \{M_t, S_t\}\right) + \left(\frac{k}{2} - 1\right)\left(n - \frac{k}{2} + 1\right),$$

and the extremal graph is  $T \vee I_{n-k/2+1}$ , where  $T \in EX(\frac{k}{2}-1, \{M_t, S_t\})$ .

If k is odd, then  $\mathcal{H} = \{M_t, S_t\}$  and  $\mathcal{H}' = \{F_{t-1}, M_{t-1} \cup K_1\}$ . We first show that every graph in  $EX(\frac{k-3}{2}, \{M_t, S_t\})$  contains a copy of  $M_{t-1} \cup K_1$ . By Theorem 2.6,  $ex(\frac{k-3}{2}, \{M_t, S_t\}) = t^2 - t$ if t is odd and  $ex(\frac{k-3}{2}, \{M_t, S_t\}) = t^2 - \frac{3}{2}t$  otherwise. By Theorem 2.1,  $ex(\frac{k-3}{2}, \{M_{t-1}, S_t\}) \leq (t-1)^2$ . Then  $ex(\frac{k-3}{2}, \{M_t, S_t\}) > ex(\frac{k-3}{2}, \{M_{t-1}, S_t\})$ . So every graph in  $EX(\frac{k-3}{2}, \{M_t, S_t\})$ must contain a copy of  $M_{t-1} \cup K_1$ . Hence, by Theorem 1.8,

$$ex_{conn}(n, \{P_k, F_t\}) = ex\left(\frac{k-3}{2}, \{M_t, S_t\}\right) + \frac{k-3}{2}\left(n - \frac{k-3}{2}\right),$$

and the extremal graph is  $T \vee I_{n-(k-3)/2}$ , where  $T \in EX(\frac{k-3}{2}, \{M_t, S_t\})$ .

**Remark.** We give an example for c = 1 in Theorem 1.7. Take  $H = K_{2,2,2}$ . We will show that  $ex_{conn}(n, \{P_k, H\}) = ex(\frac{k-3}{2}, \mathcal{H}) + \frac{k-3}{2}(n-\frac{k-3}{2}) + 1.$ 

For  $H = K_{2,2,2}$ , then  $\mathcal{H} = C_4$ . It is sufficient to prove that for every  $T \in EX(\frac{k-3}{2}, C_4)$ ,  $G = T \vee (I_{n-(k-1)/2} \cup \{uv\})$  is  $\{P_k, K_{2,2,2}\}$ -free. Obviously G is  $P_k$ -free. If G contains a copy of  $K_{2,2,2}$ , say W, then W must contain uv. Note that every edge of  $K_{2,2,2}$  belongs to a  $C_4$ in  $K_{2,2,2}$ , then uv belongs to a  $C_4$  in W. Let the  $C_4$  containing uv in W be  $\{x, y, u, v\}$ , then  $x, y \in V(T)$ . Suppose  $\{a, b\} = V(W) \setminus \{x, y, u, v\}$ , then vertices a, b must be in T. However,  $W[\{a, b, x, y\}]$  is a  $C_4$  in T, which contradicts to that T is  $C_4$ -free. Then by Theorem 1.7, we have

$$ex_{conn}(n, \{P_k, K_{2,2,2}\}) = ex\left(\frac{k-3}{2}, C_4\right) + \frac{k-3}{2}\left(n - \frac{k-3}{2}\right) + 1.$$

#### 4 Proof of Theorem 1.9

For the lower bound, it is easily seen that  $ex(n, \{P_k, H\}) \ge ex_{conn}(n, \{P_k, H\}) = (\lfloor \frac{k}{2} \rfloor - 1)n + O_k(1)$  by Theorems 1.6 and 1.7. Moreover, suppose n = m(k-1) + r, where  $0 \le r \le k-2$ . Then the disjoint union of m copies from a graph in EX(k-1, H) and a copy from a graph in EX(r, H) is also  $\{P_k, H\}$ -free. Hence

$$ex(n, \{P_k, H\}) \ge m \cdot ex(k-1, H) + ex(r, H)$$
  
=  $\frac{ex(k-1, H)}{k-1}n + O_k(1).$ 

So  $ex(n, \{H, P_k\}) \ge n \max\left\{\lfloor \frac{k}{2} \rfloor - 1, \frac{ex(k-1,H)}{k-1}\right\} + O_k(1).$ 

Suppose G is an n-vertex  $\{P_k, H\}$ -free graph. Let  $G_i (1 \le i \le t)$  be the components of G and  $l_i = |V(G_i)|$ .

Case 1. k is even. For the upper bound, we distinguish two cases for  $l_i$ . Case 1.1.  $l_i < k$ . Then  $e(G_i) \le ex(l_i, H) \le \frac{l_i(l_i-1)}{2} \le \frac{k-2}{2}l_i$ .

Case 1.2.  $l_i \geq k$ . Let the longest path in  $G_i$  be  $P_m$ , where  $m \leq k - 1$ . Then

$$e(G_i) \le ex(m, H) + \left(\frac{k}{2} - 1\right)(l_i - m)$$
  
$$\le \frac{m(m-1)}{2} + \left(\frac{k}{2} - 1\right)l_i - \frac{m(k-2)}{2}$$
  
$$= \left(\frac{k}{2} - 1\right)l_i - \frac{m}{2}(m-k+1)$$
  
$$\le \left(\frac{k}{2} - 1\right)l_i.$$

The first inequality holds by  $e(G_i[V(P_m)]) \leq ex(m, H)$  and Lemma 2.3. Therefore,

$$e(G) = \sum_{i=1}^{t} e(G_i) \le \sum_{i=1}^{t} \left(\frac{k}{2} - 1\right) l_i = \left(\frac{k}{2} - 1\right) n.$$

Then  $ex(n, \{H, P_k\}) \le n \cdot \max\left\{ \lfloor \frac{k}{2} \rfloor - 1, \frac{ex(k-1,H)}{k-1} \right\}$ . Combining with the lower bound, we have

$$ex(n, \{H, P_k\}) = n \cdot \max\left\{ \left\lfloor \frac{k}{2} \right\rfloor - 1, \frac{ex(k-1, H)}{k-1} \right\} + O_k(1)$$

Case 2. k is odd. For the upper bound, we distinguish three different cases for  $l_i$ .

Case 2.1.  $l_i < k - 1$ . Then  $e(G_i) \le ex(l_i, H) \le \frac{l_i(l_i - 1)}{2} \le \frac{k - 3}{2}l_i$ . Case 2.2.  $l_i = k - 1$ . Then  $e(G_i) \le ex(k - 1, H)$ .

Case 2.3.  $l_i \ge k$ . Let the longest path in  $G_i$  be  $P_m$  where  $m \le k-1$ .

If  $P_{k-1} \not\subseteq G_i$ , then  $m \leq k-2$  and

$$e(G_i) \le ex(m, H) + \frac{k-3}{2}(l_i - m)$$
  
$$\le \frac{m(m-1)}{2} + \frac{k-3}{2}l_i - \frac{m(k-3)}{2}$$
  
$$= \frac{k-3}{2}l_i - \frac{m}{2}(m-k+2)$$
  
$$\le \frac{k-3}{2}l_i.$$

The first inequality holds by  $e(G_i[V(P_m)]) \leq ex(m, H)$  and Lemma 2.3. If  $P_{k-1} \subseteq G_i$ , then

$$e(G_i) \leq ex(k-1,H) + \frac{k-3}{2}(l_i - k + 1)$$
  
=  $\frac{k-3}{2}l_i + ex(k-1,H) - \frac{(k-1)(k-3)}{2}.$ 

The first inequality holds by  $e(G_i[V(P_{k-1})]) \leq ex(k-1, H)$  and Lemma 2.3.

Let  $n_1 := \sum_{l_j < k-1} l_j$ ,  $n_2 := \sum_{l_j = k-1} l_j$ ,  $n_3 := \sum_{l_j \ge k, P_{k-1} \notin G_j} l_j$  and  $n_4 := \sum_{l_j \ge k, P_{k-1} \subseteq G_j} l_j$ .

Then we have

$$\begin{split} e(G) &= \frac{k-3}{2}n_1 + \frac{n_2}{k-1}ex(k-1,H) + \sum_{l_i \ge k, P_{k-1} \nsubseteq G_i} e(G_i) + \sum_{l_j \ge k, P_{k-1} \subseteq G_j} e(G_j) \\ &\leq \frac{k-3}{2}n_1 + \frac{n_2}{k-1}ex(k-1,H) + \frac{k-3}{2}n_3 + \frac{k-3}{2}n_4 + \frac{n_4}{k} \Big(ex(k-1,H) - \frac{(k-1)(k-3)}{2}\Big) \\ &\leq \frac{k-3}{2}n_1 + \frac{n_2}{k-1}ex(k-1,H) + \frac{k-3}{2}n_3 + \frac{k-3}{2}n_4 + \frac{n_4}{k-1} \Big(ex(k-1,H) - \frac{(k-1)(k-3)}{2}\Big) \\ &= \frac{k-3}{2}(n_1+n_3+n_4) + \frac{n_2+n_4}{k-1}ex(k-1,H) - \frac{k-3}{2}n_4 \\ &= \frac{k-3}{2}(n_1+n_3) + \frac{ex(k-1,H)}{k-1}(n_2+n_4). \end{split}$$

Let  $x = n_1 + n_3$  and  $y = n_2 + n_4$ . Then x + y = n and  $e(G) \le \frac{k-3}{2}x + \frac{cx(n-1,n_1)}{k-1}y$ . Hence

$$e(G) \le n \cdot \max\left\{\frac{k-3}{2}, \frac{ex(k-1,H)}{k-1}\right\}.$$

Then  $ex(n, \{H, P_k\}) \le n \cdot \max\left\{ \lfloor \frac{k}{2} \rfloor - 1, \frac{ex(k-1,H)}{k-1} \right\}$ . Combining with the lower bound, we have  $ex(n, \{H, P_k\}) = n \cdot \max\left\{ \left\lfloor \frac{k}{2} \right\rfloor - 1, \frac{ex(k-1, H)}{k-1} \right\} + O_k(1).$ The proof is finished.

#### $\mathbf{5}$ **Concluding Remarks**

Theorem 1.8 gives a sufficient condition for c = 0 in Theorem 1.7 and we also give an example to show c can be 1 in Theorem 1.7. It is natural to ask whether we can determine when c is 0 or 1 exactly in Theorem 1.7.

We obtain asymptotical result of  $ex(n, \{P_k, H\})$  with the constant term  $O_k(1)$  left, it is an interesting question to give the exact value of  $ex(n, \{P_k, H\})$ .

#### **Declaration of interests**

The authors declare that there is no conflict of interest.

## References

[1] H. L. Abbott, D. Hanson, N. Sauer, Intersection theorems for systems of sets, Journal of Combinatorial Theory, Series A, 12 (1972) 381–389.

- [2] P. N. Balister, E. Győri, J. Lehel, R. H. Schelp, Connected graphs without long paths, Discrete Mathematics, 308 (2008) 4487–4494.
- [3] V. Chvátal, D. Hanson, Degrees and matchings, Journal of Combinatorial Theory, Series B, 20 (1976) 128–138.
- [4] P. Erdős, T. Gallai, On maximal paths and circuits of graphs, Acta Mathematica Academiae Scientiarum Hungarica, 10 (1959) 337–356.
- [5] G. O. H. Katona, C. Q. Xiao, Extremal graphs without long paths and large cliques, European Journal of Combinatorics, (2023) 103807.
- [6] G. N. Kopylov, On maximal paths and cycles in a graph, Doklady Akademii Nauk, 234 (1977) 19–21.