A structure theorem for pseudo-segments and its applications

Jacob Fox^{*} János Pach[†]

Andrew Suk[‡]

Abstract

We prove a far-reaching strengthening of Szemerédi's regularity lemma for intersection graphs of pseudo-segments. It shows that the vertex set of such a graph can be partitioned into a bounded number of parts of roughly the same size such that almost all bipartite graphs between different pairs of parts are *complete* or *empty*. We use this to get an improved bound on disjoint edges in simple topological graphs, showing that every *n*-vertex simple topological graph with no *k* pairwise disjoint edges has at most $n(\log n)^{O(\log k)}$ edges.

1 Introduction

Given a set of curves C in the plane, we say that C is a collection of *pseudo-segments* if any two members in C have at most one point in common, and no three members in C have a point in common. The intersection graph of a collection C of sets has vertex set C and two sets in C are adjacent if and only if they have a nonempty intersection.

A partition of a set is an *equipartition* if each pair of parts in the partition differ in size by at most one. Szemerédi's celebrated regularity lemma roughly says that the vertex set of any graph has an equipartition such that the bipartite graph between almost all pairs of parts is random-like.

Our main result is a strengthening of Szemerédi's regularity lemma for intersection graphs of pseudo-segments. It replaces the condition that the bipartite graphs between almost all pairs of parts is random-like to being homogeneous, either complete or empty.

Theorem 1.1. For each $\varepsilon > 0$ there is $K = K(\varepsilon)$ such that for every finite collection C of pseudosegments in the plane, there is an equipartition of C into K parts C_1, \ldots, C_K such that for all but at most εK^2 pairs C_i , C_j of parts, either every curve in C_i crosses every curve in C_j , or every curve in C_i is disjoint from every curve in C_j .

Pach and Solymosi [35] proved the special case of Theorem 1.1 where C is a collection of segments in the plane, and this result was later extended to semi-algebraic graphs [3] and hypergraphs [15] of bounded description complexity. However, the techniques used to prove these results heavily rely on the algebraic structure. In fact, while it follows from the Milnor-Thom theorem that there are only $2^{O(n \log n)}$ graphs on n vertices which are semialgebraic of bounded description complexity

^{*}Stanford University, Stanford, CA. Supported by a Packard Fellowship and by NSF award DMS-1855635. Email: jacobfox@stanford.edu.

[†]Rényi Institute of Mathematics, H-1364 Budapest, POB 127, Hungary. Supported by NKFIH grants K-131529, Austrian Science Fund Z 342-N31, and ERC Advanced Grant 882971 "GeoScape." Email: pach@cims.nyu.edu.

¹Department of Mathematics, University of California at San Diego, La Jolla, CA, 92093 USA. Supported an NSF CAREER award and NSF award DMS-1952786. Email: asuk@ucsd.edu.

(see [36, 3, 42]) there are many more (namely $2^{\Omega(n^{4/3})}$) graphs on *n* vertices which are intersection graphs of pseudo-segments [17].

Theorem 1.1 does not hold if we do not allow to have exceptional pairs of parts. Indeed, the so-called half-graph (i.e., the graph on the vertex set $\{u_i, v_i : 1 \le i \le n\}$ with $u_i v_j \in E(G)$ if and only if i < j) can be represented as the intersection graph of segments, and it is easy to see that it has no equipartition without exceptional pairs [30].

Next, we discuss an application of Theorem 1.1 in graph drawing.

Disjoint edges in simple topological graphs. A *topological graph* is a graph drawn in the plane such that its vertices are represented by points and its edges are represented by nonself-intersecting arcs connecting the corresponding points. The edges are allowed to intersect, but they may not intersect vertices apart from their endpoints. Furthermore, no two edges are tangent, i.e., if two edges share an interior point, then they must properly cross at that point in common. A topological graph is *simple* if every pair of its edges intersect at most once. Two edges of a topological graph *cross* if their interiors share a point, and are *disjoint* if they neither share a common vertex nor cross.

A thrackle is a simple topological graph with no two disjoint edges. A famous conjecture due to Conway states that every *n*-vertex thrackle has at most *n* edges. In fact, Conway offered a \$1000 reward for a proof or disproof of this conjecture. The first linear upper bound was established by Lovász, Pach, and Szegedy in [29], who showed that every *n*-vertex thrackle has at most 2n edges. This bound was subsequently improved in [5, 22, 23].

Determining the maximum number of edges in a simple topological graph with no k pairwise disjoint edges seems to be a difficult task. In [38], Pach and Tóth showed that every n-vertex simple topological graph with no $k \geq 3$ pairwise disjoint edges has at most $O(n \log^{4k-8} n)$ edges. They conjectured that for every fixed k, the number of edges in such graphs is at most $O_k(n)$. Our next result substantially improves the upper bound for large k.

Theorem 1.2. If G = (V, E) is an n-vertex simple topological graph with no k pairwise disjoint edges, then $|E(G)| \leq n(\log n)^{O(\log k)}$.

In [21], Fox and Sudakov showed that every dense *n*-vertex simple topological graph contains $\Omega(\log^{1+\delta} n)$ pairwise disjoint edges, where $\delta \approx 1/40$. As an immediate Corollary to Theorem 1.2, we improve this bound to nearly polynomial under a much weaker assumption.

Corollary 1.3. Let $\varepsilon > 0$, and let G = (V, E) be an *n*-vertex simple topological graph with at least $2n^{1+\varepsilon}$ edges. Then G has $n^{\Omega(\varepsilon/\log \log n)}$ pairwise disjoint edges.

For complete *n*-vertex simple topological graphs, Aichholzer et al. [2] showed that one can always find $\Omega(n^{1/2})$ pairwise disjoint edges. Whether or not this can be improved to $\Omega(n)$ is an open problem.

The proofs of the above theorems heavily rely on a bipartite Ramsey-type result for intersection graphs of pseudo-segments (Theorem 2.8), as will be explained in Section 2. At the end of Section 2, using a variant of Szemerédi's regularity lemma originally proposed by Komlós and Sós [27, 31, 33], we show how this Ramsey-type result implies two "density-type" theorems (Theorems 2.12 and 2.11). Roughly speaking, we prove that if there are many (resp., few) crossings between two families of pseudo-segments, then they have two large subfamilies such that every member of the first subfamily crosses every member of the second (resp., no member of the first subfamily crosses any member of the second). In Subsection 2.3, we show that any finite collection of pseudo-segments

in the plane contains a linear-sized subset with the property that only a small fraction of pairs in the subset are crossing, or nearly all of them cross. In Section 3, we prove Theorem 2.8 in the special case where one of the families is double grounded. Building on these results, in Section 4, we establish our bipartite Ramsey-type theorem (Theorem 2.8) for any two families of pseudosegments with the property that for each family, only a small fraction of pairs are crossing, or nearly all of them cross. Finally, in Section 5, we prove Theorem 2.8 in its full generality. We conclude our paper with a number of remarks and with an application of our results to an old problem in graph drawing (Theorem 1.2).

2 Ramsey-type properties of hereditary families of graphs

A family of graph is *hereditary* if it is closed under taking induced subgraphs. In this section, we show that the property that in a hereditary family of graphs, a homogeneous regularity lemma holds, is equivalent to some seemingly weaker Ramsey-type properties. See Theorem 2.2, below. It will imply that in order to prove Theorem 1.1, it is enough to establish that the family of intersection graphs of pseudo-segments satisfies some bipartite Ramsey-type condition (Theorem 2.8).

We need the following definitions.

Definition 2.1. Let \mathcal{F} be a family of graphs.

- 1. \mathcal{F} has the *Erdős-Hajnal property* if there is $\varepsilon = \varepsilon(\mathcal{F}) > 0$ such that every graph in \mathcal{F} with n vertices contains a clique or independent set of size n^{ε} .
- 2. \mathcal{F} has the *polynomial Rödl property* if there is $C = C(\mathcal{F})$ such that for every $\varepsilon > 0$, every *n*-vertex graph in \mathcal{F} has an induced subgraph on at least $\epsilon^{C(\mathcal{F})}n$ vertices that is ε -homogeneous.
- 3. \mathcal{F} has the strong Erdős-Hajnal property if there is a constant $\varepsilon = \varepsilon(\mathcal{F}) > 0$ such that the vertex set of every *n*-vertex graph in \mathcal{F} with $n \geq 2$ has two disjoint subsets, each of size at least εn , such that the bipartite graph between them is complete or empty.
- 4. \mathcal{F} has the mighty Erdős-Hajnal property if there is a constant $\varepsilon = \varepsilon(\mathcal{F}) > 0$ such that for every graph G in \mathcal{F} and every pair of disjoint subsets $A, B \subset V(G)$ with |A| = |B|, there are $A' \subset A$ and $B' \subset B$ with $|A'| \ge \varepsilon |A|$ and $|B'| \ge \varepsilon |B|$ such that the bipartite graph between A and B in G is complete or empty.
- 5. \mathcal{F} has the homogeneous regularity property if for every $\varepsilon > 0$ there is $K = K_{\mathcal{F}}(\varepsilon)$ such that the vertex set of every graph $G \in \mathcal{F}$ has an equipartition $V(G) = V_1 \cup \cdots \cup V_K$ into K parts such that for all but at most εK^2 pairs of parts (V_i, V_j) , the bipartite graph between them is complete or empty (i.e., $V_i \times V_j \subset E(G)$ or $V_i \times V_j \cap E(G) = \emptyset$).
- 6. \mathcal{F} has the homogeneous density property if for every $\varepsilon > 0$ there is $C = C(\varepsilon)$ such that for every $G \in \mathcal{F}$ and every pair $A, B \subset V(G)$ of disjoint subsets of the same size with at least $\varepsilon |A||B|$ edges between them, there are $A' \subset A$ and $B' \subset B$ with $|A'| \ge \varepsilon^C |A|$ and $|B'| \ge \varepsilon^C |B|$ with the property that $A' \times B' \subset E(G)$.

For any family of graphs, \mathcal{F} , let $\overline{\mathcal{F}}$ be the family consisting of the complements of the elements of \mathcal{F} . In the sequel, we will heavily use the following.

Theorem 2.2. Let \mathcal{F} be a hereditary family of graphs. Then the following are equivalent

1. \mathcal{F} has the mighty Erdős-Hajnal property.

2. \mathcal{F} has the homogeneous regularity property.

3. \mathcal{F} and $\overline{\mathcal{F}}$ both have the homogeneous density property.

Before proving Theorem 2.2, we make some remarks about the relative strengths of the above properties.

A classical result of Erdős and Szekeres [10] states that every graph on n vertices has a clique or independent set on $\frac{1}{2} \log_2 n$ vertices. The first improvement on the constant $\frac{1}{2}$ was found recently by Campos, Griffiths, Morris, and Sahasrabudhe [6]. From the other direction, Erdős [8] showed that for every integer n > 2, there is a graph on n vertices that contains no clique or independent set on $2 \log_2 n$ vertices.

It is known that this result can be substantially improved in several restricted settings; see, e.g., [7, 14]. According to the famous *Erdős-Hajnal conjecture* [9], which has been established only in some special cases, every hereditary family of graphs that is not the family of all graphs, satisfies the Erdős-Hajnal property.

Rödl [41] proved that for any hereditary family \mathcal{F} of graphs that is missing a graph and any $\varepsilon > 0$, there is $\delta = \delta_{\mathcal{F}}(\varepsilon) > 0$ such that any graph in \mathcal{F} on n vertices contains an induced subgraph on at least δn vertices which is ε -homogeneous. Rödl's proof used Szemerédi's regularity lemma and consequently gives a weak quantitative bound. Better quantitative bounds were obtained more recently in [20, 4].

Fox and Sudakov [20] conjectured that a polynomial dependence holds in Rödl's theorem. That is, every hereditary family of graphs that is not the family of all graphs, satisfies the polynomial Rödl property. It is a simple exercise that any family having the polynomial Rödl property also has the Erdős-Hajnal property. In particular, the conjecture of Fox and Sudakov is a strengthening of the Erdős-Hajnal conjecture. Many cases of this strengthening of the Erdős-Hajnal conjecture were recently verified in [11, 32].

One way to prove that a family of graphs satisfies the Erdős-Hajnal property is to show that it has the *strong* Erdős-Hajnal property. However, this technique is limited, as the strong Erdős-Hajnal property is strictly stronger than the Erdős-Hajnal property. For example, it was proved by Tomon [43] that the family of intersection graphs of curves in the plane (the family of "*string* graphs") has the Erdős-Hajnal property, yet, as was shown by Pach and G. Tóth [39], it does not have the strong Erdős-Hajnal property. Interestingly, the family of intersection graphs of curves in the plane, each pair of which have a *bounded number of intersections*, has the strong Erdős-Hajnal property [19]. The same is true for intersection graphs of x-monotone curves or convex sets in the plane [18].

In Lemma 2.13 below, we prove that every hereditary family of graphs that has the strong Erdős-Hajnal property must also have the polynomial Rödl property. The strong Erdős-Hajnal property is strictly stronger than the polynomial Rödl property, as it is a simple exercise to show that the family of triangle-free graphs has the polynomial Rödl property but does not have the strong Erdős-Hajnal property.

However, for some applications, even the strong Erdős-Hajnal property is not sufficient, and one needs to introduce a stronger property: the *mighty* Erdős-Hajnal property, which is a bipartite version of the strong Erdős-Hajnal property. The simplest example of a class of graphs that has the strong property, but not the mighty one, is the family of bipartite graphs. A more interesting example is the family of intersection graphs of *convex sets* in the plane, which has the strong Erdős-Hajnal property by [18], but not he mighty Erdős-Hajnal property as it contains the complement of every bipartite graph. On the other hand, it was proved by Pach and Solymosi [36, 35] that intersection graphs of segments, or, more generally, intersection graphs of semialgebraic sets of bounded description complexity, have even the mighty Erdős-Hajnal property [3].

2.1 Proof of Theorem 2.2

We split the proof of the equivalences in Theorem 2.2 into four smaller lemmas.

Lemma 2.3. Suppose that a hereditary family \mathcal{F} of graphs has the homogeneous regularity property. Then \mathcal{F} also has the mighty Erdős-Hajnal property.

Proof. Let $G \in \mathcal{F}$ and $A, B \subset V(G)$ be disjoint with |A| = |B| = n. Let G' be the induced subgraph of G with vertex set $A \cup B$. Since \mathcal{F} is hereditary and G' is an induced subgraph of $G \in \mathcal{F}$, then $G' \in \mathcal{F}$. By the homogeneous regularity property of G' applied with $\varepsilon = 1/65$ there is an absolute constant K > 100 and an equipartition of $A \cup B$ into K parts V_1, \ldots, V_K such that all but εK^2 pairs of parts (V_i, V_j) are complete or empty. Note that the parts V_i that have at most n/(2K) elements in A together have at most $K \cdot n/(2K) = n/2$ elements from A. Each part V_i has at most $\lceil 2n/K \rceil \leq 4n/K$ elements. So there at least (n/2)/(4n/K) = K/8 parts V_i that have at least n/(2K) elements from A. Similarly, at least K/8 parts have at least n/(2K) elements from A. Similarly, at least K/8 parts have at least n/(2K) elements from A. Similarly, at least K/8 parts have at least n/(2K) elements from A. Similarly, at least K/8 parts have at least n/(2K) elements from A. Since we chose $\varepsilon = 1/200$, there are at least $K^2/64 - K^2/65 > 0$ pairs of parts (V_i, V_j) that are complete or empty to each other, V_i contains at least n/(2K) elements from A and V_j contains at least n/(2K) elements of B. Fixing such a pair (V_i, V_j) and letting $A' = V_i \cap A$ and $B' = V_j \cap B$, we have A' and B' are complete or empty to each other and are of the desired size. So \mathcal{F} has the mighty Erdős-Hajnal property.

Lemma 2.4. Suppose that a hereditary family \mathcal{F} of graphs has the mighty Erdős-Hajnal property. Then \mathcal{F} also has the homogeneous regularity property.

Proof. Since \mathcal{F} has the mighty Erdős-Hajnal property, there is c > 0 such that for every graph G_0 in \mathcal{F} and for all disjoint vertex subsets $A, B \subset V(G_0)$ with |A| = |B|, there is $A' \subset A$ and $B' \subset B$ with $|A'| \ge c|A|$ and $|B'| \ge c|B|$ such that the pair (A', B') is homogeneous.

We will show that \mathcal{F} also has the homogeneous regularity property. Let $G \in \mathcal{F}$ have *n* vertices. We will show that *G* has an equipartition into *K* parts such that the bipartite graph between all but at most εK^2 pairs of parts are complete or empty. If $n \leq K$, then partition V(G) into *K* sets which have size at most one. Then every pair of parts is complete or empty between them. So we may assume n > K.

Arbitrarily equipartition $V(G) = V_1 \cup \ldots \cup V_t$ into $t = 4/\varepsilon$ sets, so at most a fraction $\varepsilon/4$ of the pairs of vertices are in the same part in the partition. Delete one vertex from each part V_i that has size $\lfloor n/t \rfloor + 1$ so all parts have the same size. Still, all but at least an $\varepsilon/3$ fraction of the pairs of vertices go between these pairs of sets of the same. Let $\mathcal{B}_0 = \{(V_i, V_j)\}_{1 \le i < j \le t}$.

Let $\varepsilon' = c^2 \varepsilon / 100$ At each step *i* of a process we will have a family \mathcal{B}_i of pairs of disjoint vertex subsets all of the same size satisfying the following four properties:

- 1. $|\mathcal{B}_i| \leq t^2 (1/\varepsilon')^{2i}$.
- 2. Each pair of vertices of G go between at most one pair of vertex subsets of \mathcal{B}_i .

- 3. All but a fraction $(1 2\varepsilon')^{2i}$ of the pairs of vertices that go between a pair of vertex subsets in \mathcal{B}_0 go between a pair of vertex subsets in \mathcal{B}_i .
- 4. At most a $(1 c^2)^i$ fraction of the pairs of vertices of the graph go between a pair of vertex subsets in \mathcal{B}_i that are not homogeneous.

Note that the above properties hold for i = 0. Suppose we have already found \mathcal{B}_i . For each pair $(A, B) \in \mathcal{B}_i$, by the mighty Erdős-Hajnal property, there are subsets $A' \subset A$ and $B' \subset B$ with $|A'| \geq c|A|$ and $|B'| \geq c|B|$ and the bipartite graph between A and B is complete or empty. Partition each of the four sets A', $A \setminus A'$, B' and $B \setminus B'$ into parts of size $\varepsilon'|A|$ with one possible remaining set. For each pair of these subsets of A and B of size $\varepsilon'|A|$ such that one is a subset of A and the other is a subset of B, we place the pair of subsets in \mathcal{B}_{i+1} .

We next observe that each of the four properties listed hold. For each pair $(A, B) \in \mathcal{B}_i$, we get at most $(1/\varepsilon')^2$ pairs in \mathcal{B}_{i+1} , so $|\mathcal{B}_{i+1}| \leq (1/\varepsilon')^2 |\mathcal{B}_i|$, which inductively gives the first claimed property. The second claimed property holds since every pair of vertices that go between a pair of vertex subsets in \mathcal{B}_{i+1} also go between a pair of vertex subsets in \mathcal{B}_i . At least a $(1-2\varepsilon')^2$ -fraction of the pairs of vertices in $A \times B$ go between pairs of vertices in \mathcal{B}_{i+1} , which inductively gives the third claimed property. Finally, for each $(A, B) \in \mathcal{B}_i$, since at least a $(1-c)^2$ fraction of the pairs of vertices go between the homogeneous pair (A', B'), the last claimed property holds.

After $i_0 = c^{-2} \log(4/\varepsilon)$ steps, all but a fraction $(1 - c^2)^{i_0} < e^{-c^2 i_0} = \varepsilon/4$ of the pairs of vertices in G go between a pair of vertex subsets in \mathcal{B}_{i_0} that are not homogeneous. The fraction of pairs of vertices in G that do no go between a pair of vertex subsets in \mathcal{B}_{i_0} is at most $\varepsilon/3 + 1 - (1 - 2\varepsilon')^{2i_0} \le \varepsilon/2$.

For each pair $(A, B) \in \mathcal{B}_{i_0}$, we obtain two bipartitions $V(G) = A \cup (V(G) \setminus A)$ and $V(G) = B \cup (V(G) \setminus B)$. Let \mathcal{Q} be the common refinement of all of these $2|\mathcal{B}_{i_0}|$ bipartitions, so \mathcal{Q} is a partition of V(G) into $2^{2|\mathcal{B}_{i_0}|}$ parts.

We next obtain an equipartition \mathcal{P} of V(G) into K parts. For every equipartition into K parts, we have parts of size $\lfloor n/K \rfloor$ or $\lceil n/K \rceil$, with the number of parts of size $\lceil n/K \rceil$ being the remainder when n is divided by K. Partition each part of \mathcal{Q} into parts of size $\lfloor n/K \rfloor$ or $\lceil n/K \rceil$, with additional one part of size at most n/K. We partition the union of the $|\mathcal{Q}|$ additional parts into sets of size $\lfloor n/K \rfloor$ or $\lceil n/K \rceil$ so that to obtain an equipartition P of V(G) into K parts. The fraction of pairs of vertices in G not going between a homogeneous pair in partition P is at most $\varepsilon/2 + \varepsilon/4 + 2|\mathcal{Q}|/K \leq \varepsilon$. Thus, P is the desired equipartition.

Lemma 2.5. Suppose that \mathcal{F} and $\overline{\mathcal{F}}$ are hereditary families that both have the homogeneous density property. Then \mathcal{F} also has the mighty Erdős-Hajnal property.

Proof. Let $G \in \mathcal{F}$ and A, B be disjoint vertex subsets of G of the same size. The bipartite graph between A and B have edge density at least 1/2 or at most 1/2. In the first case, since \mathcal{F} has the homogeneous density property, there is a constant c > 0 and subsets $A' \subset A$ with $|A'| \ge c|A|$ and $B' \subset B$ with $|B'| \ge c|B|$ such that A' is complete to B'. In the second case, since $\overline{\mathcal{F}}$ has the homogeneous density property, there is a constant c > 0 and subsets $A' \subset A$ with $|A'| \ge c|A|$ and $B' \subset B$ with $|B'| \ge c|B|$ such that A' is empty to B'. So we get \mathcal{F} has the mighty Erdős-Hajnal property.

Lemma 2.6. Suppose that a hereditary family \mathcal{F} has the mighty Erdős-Hajnal property. Then \mathcal{F} and $\overline{\mathcal{F}}$ both have the homogeneous density property.

The proof of Lemma 2.6 is now a standard application of a weak bipartite version of Szemerédi's regularity lemma. The version we use is due to Komlós and Sós (see [27, 33] or Theorem 9.4.1 on page 223 of [31]).

Lemma 2.7 ([27, 33]). Let G = (A, B, E) be a bipartite graph with parts A and B such that |A| = |B| = n. Let $0 < c \le 1/2$. If $|E| \ge \varepsilon n^2$, then there exists subsets $A_0 \subset A, B_0 \subset B$ such that

1.
$$|A_0| = |B_0| = \varepsilon^{1/c^2} n$$
, and

2. $|E(A_0, B_0)| \ge \varepsilon |A_0| |B_0|$, and

3. |E(A', B')| > 0 for any $A' \subset A_0$ and $B' \subset B_0$ with $|A'|, |B'| \ge c|A_0|$.

Proof of Lemma 2.6. Since \mathcal{F} has the mighty Erdős-Hajnal property, there is a constant $0 < c \leq 1/2$ such that for every $G \in \mathcal{F}$ and disjoint vertex subsets A, B of G of the same size, there are $A' \subset A$ and $B' \subset B$ with $|A'|, |B'| \geq c|A|$ such that the bipartite graph between A' and B' is either complete or empty.

By symmetry, it suffices to show that \mathcal{F} has the homogeneous density property. Let $G \in \mathcal{F}$ and A, B be disjoint vertex subsets of equal size such that the edge density between A and B is at least ε . By Lemma 2.7, there are $A_0 \subset A$ and $B_0 \subset B$ such that $|A_0| = |B_0| = \varepsilon^{1/c^2} |A|$, the edge density between A_0 and B_0 is at least ε , and |E(A', B')| > 0 for any $A' \subset A_0$ and $B' \subset B_0$ with $|A'|, |B'| \ge c|A_0|$. Applying the mighty Erdős-Hajnal property to A_0 and B_0 , we get subsets A' and B' with $|A'|, |B'| \ge c|A_0|$ that are complete or empty to each other. However, the above condition that there is at least one edge between A' and B' implies that A' and B' are complete to each other, completing the proof.

2.2 Deductions for intersection graphs of pseudo-segments

By Theorem 2.2, the main result in this paper (Theorem 1.1) is equivalent to saying that the family of intersection graphs of pseudo-segments has the mighty Erdős-Hajnal property. Therefore, to prove Theorem 1.1, it suffices to prove the following result.

Theorem 2.8. Let \mathcal{R} be a set of n red curves, and \mathcal{B} be a set of n blue curves in the plane such that $\mathcal{R} \cup \mathcal{B}$ is a collection of pseudo-segments.

Then there are subsets $\mathcal{R}' \subset \mathcal{R}$ and $\mathcal{B}' \subset \mathcal{B}$, where $|\mathcal{R}'|, |\mathcal{B}'| \geq \Omega(n)$, such that either every curve in \mathcal{R}' crosses every curves in \mathcal{B}' , or every curve in \mathcal{R}' is disjoint from every curves in \mathcal{B}' .

As mentioned earlier, a weaker result, that the family of intersection graphs of pseudo-segments has the strong Erdős-Hajnal property, was proved earlier by Fox, Pach, and Tóth [19]. We repeat it below for convenience.

Theorem 2.9 ([19]). Let C be a collection on n pseudo-segments in the plane. Then there are subsets $C_1, C_2 \subset C$, where $|C_1|, |C_2| \geq \Omega(n)$, such that either every curve in C_1 crosses every curve in C_2 , or every curve in C_1 is disjoint from every curve in C_2

By Theorem 2.2, Theorem 2.8 is equivalent to the following two theorems holding.

Theorem 2.10. There is an absolute constant c > 0 such that the following holds. Let \mathcal{R} be a collection of n red curves, and \mathcal{B} be a collection of n blue curves in the plane such that $\mathcal{R} \cup \mathcal{B}$ is a collection of pseudo-segments.

If there are at least εn^2 crossing pairs in $\mathcal{R} \times \mathcal{B}$, then there are subsets $\mathcal{R}' \subset \mathcal{R}, \mathcal{B}' \subset \mathcal{B}$, where $|\mathcal{R}'|, |\mathcal{B}'| \geq \varepsilon^c n$, such that every curve in \mathcal{R}' crosses every curve in \mathcal{B}' .

Theorem 2.11. There is an absolute constant c > 0 such that the following holds. Let \mathcal{R} be a collection of n red curves, and \mathcal{B} be a collection of n blue curves in the plane such that $\mathcal{R} \cup \mathcal{B}$ is a collection of pseudo-sequents.

If there are at least εn^2 disjoint pairs in $\mathcal{R} \times \mathcal{B}$, then there are subsets $\mathcal{R}' \subset \mathcal{R}, \mathcal{B}' \subset \mathcal{B}$, where $|\mathcal{R}'|, |\mathcal{B}'| \geq \varepsilon^c n$, such that every curve in \mathcal{R}' is disjoint from every curve in \mathcal{B}' .

Theorem 2.11 is the first density-type result for *disjointness graphs* of pseudo-segments, and is the key ingredient in the proof of Theorem 1.2. For the proof of Theorem 2.8, we need the following non-bipartite version of Theorem 2.10 established by the first two authors in [13].

Theorem 2.12 ([13]). There is an absolute constant c' > 0 such that the following holds. Let C be a collection of n pseudo-segments in the plane with at least εn^2 crossing pairs. Then there are subsets $C_1, C_2 \subset C$, each of size $c'\varepsilon n$, such that every curve in C_1 crosses every curve in C_2 .

2.3 The strong Erdős-Hajnal property implies the polynomial Rödl property

The goal of this subsection is to prove the following lemma, and deduce that the family of intersection graphs of pseudo-segments has the polynomial Rödl property.

Lemma 2.13. If a hereditary family \mathcal{F} of graphs has the strong Erdős-Hajnal property, then it also has the polynomial Rödl property.

In proving Lemma 2.13, given a graph G in a hereditary family of graphs satisfying the strong Erdős-Hajnal property, we find a large subgraph of G or its complement that is a balanced complete multipartite graph with many parts. Such a subgraph must have density close to one, giving the desired ε -homogeneous induced subgraph.

Proof of Lemma 2.13. Since \mathcal{F} has the strong Erdős-Hajnal property, there is a constant c > 0such that every $G \in \mathcal{F}$ on $n \geq 2$ vertices has disjoint vertex subsets V_1, V_2 with $|V_1| = |V_2| = cn$ such that the graph between V_1 and V_2 is complete or empty. We repeatedly apply this to the induced subgraph of each of the vertex subsets of G we obtain. After t levels of iteration, we obtain 2^t disjoint vertex subsets, each of size $c^t n$, that are complete or empty between each pair of them, and the induced subgraph H of G formed by taking one vertex from each of these 2^t subsets is a cograph. Since cographs are perfect graphs, and any perfect graph with m vertices contains a clique or independent set of size at least $m^{1/2}$, H contains a clique or independent set with at least $2^{t/2}$ vertices. Let U_1, \ldots, U_s be the vertex subsets that the vertices from this clique or independent set belong to. Then the edge density of the induced subgraph on the union $\bigcup_{i=1}^s U_i$ is either more than 1-1/s or less than 1/s. Let $t = 2\log_2(1/\varepsilon)$. Since $s \geq 2^{t/2} = 1/\varepsilon$ and $c^t = \varepsilon^C$ with $C = 2\log_2(1/c)$, this completes the proof.

Given a collection C of curves in the plane, let G(C) denote the intersection graph of C. According to Theorem 2.9, the family of intersection graphs of pseudo-segments has the strong Erdős-Hajnal property. Applying Lemma 2.13, we obtain

Corollary 2.14. The family of intersection graphs of pseudo-segments has the polynomial Rödl property. That is, there is an absolute constant $c_1 > 0$ such that the following holds. Let $\varepsilon > 0$ and C be a collection of n pseudo-segments in the plane. Then there is a subset $C' \subset C$ of size $\varepsilon^{c_1} n$ whose intersection graph G(C') is ε -homogeneous.

We will frequently use the following simple lemma in this paper.

Lemma 2.15. Let G = (V, E) be a graph on n vertices. If the edge density of G is at most ε , then any induced subgraph on δn vertices has edge density at most $2\varepsilon/\delta^2$. Likewise, if the edge density of G is at least $1 - \varepsilon$, then any induced subgraph on δn vertices has edge density at least $1 - 2\varepsilon/\delta^2$.

Proof. Suppose the edge density of G is at most εn^2 edges. Hence, any induced subgraph on δn vertices will have at most $(2\varepsilon/\delta^2)(\delta n)^2/2$ edges, which implies that the edge density of the induced subgraph is at most $2\varepsilon/\delta^2$. A symmetric argument follows in the case when the edge density of G is at least $1 - \varepsilon$.

3 Proof of Theorem 2.8 – for double grounded red curves

Given a collection of curves C in the plane, we say that C is *double grounded* if there are two distinct curves γ_1 and γ_2 such that for each curve $\alpha \in C$, α has one endpoint on γ_1 and the other on γ_2 , and the interior α is disjoint from γ_1 and γ_2 . Throughout this paper, for simplicity, we will always assume that both endpoints of each of our curves have distinct *x*-coordinates. We refer to the endpoint of a curve with the smaller (larger) *x*-coordinate as its *left (right) endpoint*. The aim of this section is to prove Theorem 2.8 in the special case where one of the color classes (the red one, say) consists of double grounded curves.

A curve in the plane is called x-monotone if every vertical line intersects it in at most one point. We start by considering double grounded x-monotone curves, and at the end of this section, we will remove the x-monotone condition. We will need the following result, known as the cutting-lemma for x-monotone curves. See, for example, Proposition 2.11 in [26].

Lemma 3.1 (The Cutting Lemma). Let C be a collection of n double grounded x-monotone curves, whose grounds that are disjoint vertical segments γ_1 and γ_2 , and let r > 1 be a parameter. Then $\mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2)$ can be subdivided into t connected regions $\Delta_1, \ldots, \Delta_t$, such that the interior of each Δ_i is intersected by at most n/r curves from C, and we have $t = O(r^2)$.

In the proof of the following lemma and throughout the paper, we will implicitly use the Jordan curve theorem.

Lemma 3.2. Let \mathcal{R} be a set of n red double grounded x-monotone curves, whose grounds are disjoint vertical segments γ_1 and γ_2 . Let \mathcal{B} be a set of n blue curves (not necessarily x-monotone) such that every blue curve in \mathcal{B} is disjoint from grounds γ_1 and γ_2 , and suppose that $\mathcal{R} \cup \mathcal{B}$ is a collection of pseudo-segments.

Then there are subsets $\mathcal{R}' \subset \mathcal{R}$ and $\mathcal{B}' \subset \mathcal{B}$ such that $|\mathcal{R}'|, |\mathcal{B}'| \geq \Omega(n)$, and either every curve in \mathcal{R}' crosses every curve in \mathcal{B}' , or every curve in \mathcal{R}' is disjoint from every curve in \mathcal{B}' .

Proof. Let P be the set of left-endpoints of the curves in \mathcal{B} . We apply Lemma 3.1 to \mathcal{R} with parameter r = 4 to obtain a subdivision

$$\mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2) = \Delta_1 \cup \cdots \cup \Delta_t,$$

such that for each Δ_i , the interior of Δ_i intersects at most n/4 members in \mathcal{R} , and $t \leq c_0 4^2$ where c_0 is an absolute constant from Lemma 3.1. By the pigeonhole principle, there is a region Δ_i such that Δ_i contains at least $n/c_0 4^2$ points from P. Let $\mathcal{B}_0 \subset \mathcal{B}$ be the set of blue curves whose left endpoints are in Δ_i . Hence $|\mathcal{B}_0| = \Omega(n)$.

Let Q be the right endpoints of the curves in \mathcal{B}_0 . Using the same subdivision described above, there is a region Δ_i such that Δ_i contains at least $|Q|/(c_0 4^2) \ge n/(c_0 4^2)^2$ points from Q. Let



Figure 1: Case 1, α_0 and α are disjoint.

 $\mathcal{B}_1 \subset \mathcal{B}_0$ be the set of blue curves with their left endpoint in Δ_i and right endpoint in Δ_j . Let $\mathcal{R}_1 \subset \mathcal{R}$ consists of all red curves that do not intersect the interior of Δ_i and Δ_j . Lemma 3.1 implies that

$$|\mathcal{R}_1| \ge n - \frac{2n}{4} = \frac{n}{2},$$

and $|\mathcal{B}_1| = \Omega(n)$. Recall that each blue curve in \mathcal{B}_1 does not intersect the grounds γ_1 nor γ_2 . Fix an arbitrary curve $\alpha_0 \in \mathcal{R}_1$. The proof now falls into the following cases.

Case 1. Suppose at least $|\mathcal{R}_1|/2$ curves in \mathcal{R}_1 are disjoint from α_0 . Let $\mathcal{R}_2 \subset \mathcal{R}_1$ be the set of red curves disjoint from α_0 . For each $\alpha \in \mathcal{R}_2 \setminus \{\alpha_0\}$,

$$\mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2 \cup \alpha_0 \cup \alpha),$$

consists of two connected components, one bounded and the other unbounded.

Case 1.a. Suppose for at least $|\mathcal{R}_2|/2$ red curves $\alpha \in \mathcal{R}_2$, both Δ_i and Δ_j lie in the same connected component of $\mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2 \cup \alpha_0 \cup \alpha)$. See Figure 1(a). Let $\mathcal{R}_3 \subset \mathcal{R}_2$ be the collection of such red curves. Then for each $\alpha \in \mathcal{R}_3$, each blue curve $\beta \in \mathcal{B}_1$ crosses α if and only if β crosses α_0 . Hence, there is a subset $\mathcal{B}_2 \subset \mathcal{B}_1$ of size at least $\Omega(n)$, such that either every blue curve in \mathcal{B}_2 crosses every red curve in \mathcal{R}_3 , or every blue curve in \mathcal{B}_2 is disjoint from every red curve in \mathcal{R}_3 . Moreover, $|\mathcal{R}_3| = \Omega(n)$ and we are done.

Case 1.b. Suppose for at least $|\mathcal{R}_2|/2$ red curves $\alpha \in \mathcal{R}_2$, regions Δ_i and Δ_j lie in different connected component of $\mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2 \cup \alpha_0 \cup \alpha)$. See Figure 1(b). Similar to above, let $\mathcal{R}_3 \subset \mathcal{R}_2$ be the collection of such red curves. By the pseudo-segment condition, for each $\alpha \in \mathcal{R}_3$, each blue curve $\beta \in \mathcal{B}_1$ crosses α if and only if β is disjoint from α_0 . Hence, there is a subset $\mathcal{B}_2 \subset \mathcal{B}_1$ of size $\Omega_r(n)$, such that either every blue curve in \mathcal{B}_2 crosses every red curve in \mathcal{R}_3 , or every blue curve in \mathcal{B}_2 is disjoint from every red curve in \mathcal{R}_3 . Moreover, $|\mathcal{R}_3| = \Omega(n)$ and we are done.

Case 2. Suppose at least $|\mathcal{R}_1|/2$ curves in \mathcal{R}_1 cross α_0 . Let $\mathcal{R}_2 \subset \mathcal{R}_1$ be the set of red curves that crosses α_0 . For each $\alpha \in \mathcal{R}_2 \setminus \{\alpha_0\}$,

$$\mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2 \cup \alpha_0 \cup \alpha),$$

consists of three connected components, two of which are bounded and the other unbounded.

Case 2.a. Suppose for at least $|\mathcal{R}_2|/3$ red curves $\alpha \in \mathcal{R}_2$, Both Δ_i and Δ_j lie in the same connected component of $\mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2 \cup \alpha_0 \cup \alpha)$. See Figure 2(a). Let $\mathcal{R}_3 \subset \mathcal{R}_2$ be the collection of such red curves. By the pseudo-segment condition, for each $\alpha \in \mathcal{R}_3$, each blue curve $\beta \in \mathcal{B}_1$ crosses α if and



Figure 2: Case 2, α_0 and α cross.

only if β crosses α_0 . Hence, there is a subset $\mathcal{B}_2 \subset \mathcal{B}_1$ of size at least $\Omega(n)$, such that either every blue curve in \mathcal{B}_2 crosses every red curve in \mathcal{R}_3 , or every blue curve in \mathcal{B}_2 is disjoint from every red curve in \mathcal{R}_3 . Moreover, $|\mathcal{R}_3| = \Omega(n)$.

Case 2.b. Suppose for at least $|\mathcal{R}_2|/3$ red curves $\alpha \in \mathcal{R}_2$, regions Δ_i and Δ_j lie in different bounded connected components of $\mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2 \cup \alpha_0 \cup \alpha)$. See Figure 2(b). Let $\mathcal{R}_3 \subset \mathcal{R}_2$ be the collection of such red curves. Then for each $\alpha \in \mathcal{R}_3$, every blue curve $\beta \in \mathcal{B}_1$ crosses α . Since $|\mathcal{R}_3| = \Omega(n)$ and $|\mathcal{B}_1| = \Omega(n)$.

Case 2.c. Suppose for at least $|\mathcal{R}_2|/3$ red curves $\alpha \in \mathcal{R}_2$, regions Δ_i and Δ_j lie in different connected components of $\mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2 \cup \alpha_0 \cup \alpha)$, one of which is bounded and the other unbounded. See Figure 2(c). Let $\mathcal{R}_3 \subset \mathcal{R}_2$ be the collection of such red curves. By the pseudo-segment condition, for each $\alpha \in \mathcal{R}_3$, each blue curve $\beta \in \mathcal{B}_1$ crosses α if and only if β is disjoint from α_0 . Hence, there is a subset $\mathcal{B}_2 \subset \mathcal{B}_1$ of size $\Omega(n)$, such that either every blue curve in \mathcal{B}_2 crosses every red curve in \mathcal{R}_3 , or every blue curve in \mathcal{B}_2 is disjoint from every red curve in \mathcal{R}_3 . Moreover, $|\mathcal{R}_3| = \Omega(n)$, and we are done.

Recall that a *pseudoline* is an unbounded arc in \mathbb{R}^2 , whose complement is disconnected. An *arrangement of pseudolines* is a set of pseudolines such that every pair meets exactly once, and no three members have a point in common. A classic result of Goodman [24] states that every arrangement of pseudolines is isomorphic to an arrangement of *wiring diagrams* (bi-infinite *x*-monotone curves). Moreover, Goodman and Pollack showed the following.

Theorem 3.3 ([25]). Every arrangement of pseudolines can be continuously deformed (through isomorphic arrangements) to a wiring diagram.

We also need the following simple lemma.

Lemma 3.4. Given a finite linearly ordered set whose elements are colored red or blue, we can select half of the red elements and half of the blue elements such that all of the selected elements of one color come before all of the selected elements of the other color.

We are now ready to establish the main result of this section:

Theorem 3.5. Let \mathcal{R} be a set of n red double grounded curves with grounds γ_1 and γ_2 , where γ_1 and γ_2 cross each other. Let \mathcal{B} be a set of n blue curves such that $\mathcal{R} \cup \mathcal{B} \cup \{\gamma_1, \gamma_2\}$ is a collection of pseudo-segments.

Then there are subsets $\mathcal{R}' \subset \mathcal{R}$ and $\mathcal{B}' \subset \mathcal{B}$ such that $|\mathcal{R}'|, |\mathcal{B}'| \geq \Omega(n)$, and either every curve in \mathcal{R}' crosses every curve in \mathcal{B}' , or every curve in \mathcal{R}' is disjoint from every curve in \mathcal{B}' .

Proof. By passing to linear-sized subsets of \mathcal{R} and \mathcal{B} and subcurves of γ_1 and γ_2 , we will reduce the problem to the setting of Lemma 3.2. Let us assume that γ_1 and γ_2 cross at point p. Hence, $(\gamma_1 \setminus \gamma_2) \cup (\gamma_2 \setminus \gamma_1)$ consists of four connected components. By the pigeonhole principle, there is a subset $\mathcal{R}_1 \subset \mathcal{R}$ of size n/4 such that every curve in \mathcal{R}_1 has an endpoint on one of the connected components of $\gamma_1 \setminus \gamma_2$, and all of the other endpoints lie on one of the connected components of $\gamma_2 \setminus \gamma_1$. Let $\gamma'_i \subset \gamma_i$, for i = 1, 2, be these connected components so that they have a common endpoint at p and their interiors are disjoint.

For each $\alpha \in \mathcal{R}_1$, the sequence of curves $(\gamma'_1, \gamma'_2, \alpha)$ appear either in *clockwise* or *counterclockwise* order along the unique simple closed curve that lies in $\gamma'_1 \cup \gamma'_2 \cup \alpha$. Without loss of generality, we can assume that there is a subset $\mathcal{R}_2 \subset \mathcal{R}_1$, where $|\mathcal{R}_2| = \Omega(n)$, such that for every curve $\alpha \in \mathcal{R}_2$, the sequence $(\gamma'_1, \gamma'_2, \alpha)$ appears in clockwise order, since a symmetric argument would follow otherwise.

We define the orientation of each curve $\alpha \in \mathcal{R}_2$ as the sequence of turns, either left-left, leftright, right-left, or right-right, made by starting at p and moving along γ'_1 in the arrangement $\gamma'_1 \cup \gamma'_2 \cup \alpha$, until we return back to p. More precisely, starting at p we move along γ'_1 until we reach the endpoint of α . We then turn either left or right to move along α towards γ'_2 . Once we've reached γ'_2 , we either turn left or right in order to move along γ'_2 and reach p again. By the pigeonhole principle, there is a subset $\mathcal{R}_3 \subset \mathcal{R}_2$ of size at least $\Omega(n)$ such that all curves in \mathcal{R}_3 have the same orientation. Without loss of generality, we can assume that the orientation is left-left, since a symmetric argument would follow otherwise.

Starting at p and moving along γ'_1 towards its other endpoint, let us consider the sequence of curves from $\mathcal{R}_3 \cup \mathcal{B}$ intersecting γ'_1 . Then, by Lemma 3.4, there are subsets $\mathcal{R}_4 \subset \mathcal{R}_3$ and $\mathcal{B}_1 \subset \mathcal{B}$, where $|\mathcal{R}_4| \geq |\mathcal{R}_3|/2$ and $|\mathcal{B}_1| \geq |\mathcal{B}|/2$, such that either all of the curves in \mathcal{R}_4 appear before all of the curves in \mathcal{B}_1 that intersect γ'_1 in this sequence, or all of the curves in \mathcal{R}_4 appear after all of the curves in \mathcal{B}_1 in this sequence. Note that \mathcal{B}_1 consists of the blue curves in \mathcal{B} that are disjoint to γ'_1 and at least half of the curves in \mathcal{B} that intersect γ'_1 found by the application of Lemma 3.4. Hence, there is a subcurve $\gamma''_1 \subset \gamma'_1$ such that γ''_1 is one of the grounds for \mathcal{R}_4 , and is disjoint from every curve in \mathcal{B}_1 . We apply the same argument to $\mathcal{R}_4 \cup \mathcal{B}_1$ and γ'_2 , and obtain subsets $\mathcal{R}_5 \subset \mathcal{R}_4$, $\mathcal{B}_2 \subset \mathcal{B}_1$, and a subcurve $\gamma''_2 \subset \gamma'_2$, such that $|\mathcal{R}_5|, |\mathcal{B}_2| = \Omega(n)$, and \mathcal{R}_5 is double grounded with disjoint grounds γ''_1 and γ''_2 , and every curve in \mathcal{B}_2 is disjoint from γ''_1 and γ''_2 .

For $i \in \{1, 2\}$, let p_i be the endpoint of γ_i'' that lies closest to p along γ_i' . Starting at p_i and moving along γ_i'' , let π_i be the sequence of curves in \mathcal{R}_5 that appear on γ_i'' . Since every curve in \mathcal{R}_5 has the same left-left orientation, and appears clockwise order with respect to γ_1' and γ_2' , two curves $\alpha, \alpha' \in \mathcal{R}_5$ cross if and only if the order in which they appear in π_1 and π_2 changes. Let γ_3'' be a curve very close to γ_2'' such that γ_3'' has the same endpoints as γ_2'' , and is disjoint from all curves in $\mathcal{R}_5 \cup \mathcal{B}_2$. Hence, $\gamma_2'' \cup \gamma_3''$ makes an empty lens in the arrangement $\mathcal{R}_5 \cup \mathcal{B}_2$. We slightly extend each curve $\alpha \in \mathcal{R}_5$ through this lens to γ_3'' so that the resulting curve, α' properly crosses γ_2'' and has its new endpoint on γ_3'' . Moreover, the extension will be made in such a way that the sequence π_3 of curves in \mathcal{R}_5 appearing along γ_3'' starting from p_2 will appear in the opposite order of π_1 . Let

$$\mathcal{R}_5' = \{ \alpha' : \alpha \in \mathcal{R}_5 \}.$$

Thus, every pair of curves in \mathcal{R}'_5 will cross exactly once.

For each curve $\alpha' \in \mathcal{R}'_5$, we further extend α' by moving both endpoints towards p along γ_1 and γ_2 , so that we do not create any additional crossings within \mathcal{R}'_5 . Let $\hat{\alpha}$ be the resulting extension, where both endpoints of $\hat{\alpha}$ lie arbitrarily close to p. Set $\hat{\mathcal{R}}_5 = \{\hat{\alpha} : \alpha' \in \mathcal{R}'_5\}$. See Figure 3/ Furthermore, we can assume that p lies in the unbounded face of the arrangement $\hat{\mathcal{R}}_5$, since



Figure 3: The resulting extension $\hat{\mathcal{R}}_5$.

otherwise we could project the arrangement $\hat{\mathcal{R}}_5$ onto a sphere, and then project it back to the plane so that p lies in the unbounded face, without creating or removing any crossing. Therefore, $\hat{\mathcal{R}}_5$ can be extended to a family of pseudolines. By Theorem 3.3, we can apply a continuous deformation of the plane so that $\hat{\mathcal{R}}_5$ becomes a collection of unbounded x-monotone curves. Hence, after the deformation, the original set \mathcal{R}_5 becomes a collection of double grounded x-monotone curves, with grounds γ_1'', γ_2'' , such that every curve in \mathcal{B}_2 is disjoint from the grounds γ_1'' and γ_2'' , the crossing pattern in the arrangement $\mathcal{R}_5 \cup \mathcal{B}_2$ is the same as before. Moreover, γ_1'' and γ_2'' will be disjoint vertical segments. We apply Lemma 3.2 to \mathcal{R}_5 and \mathcal{B}_2 and obtain subsets $\mathcal{R}_6 \subset \mathcal{R}_5$ and $\mathcal{B}_3 \subset \mathcal{B}_2$, each of size $\Omega(n)$, such that either every curve in \mathcal{R}_6 crosses every curve in \mathcal{B}_3 , or every curve in \mathcal{R}_6 is disjoint from every curve in \mathcal{B}_3 . This completes the proof.

By combining Theorem 3.5 with Lemma 2.7, as shown in Section 2, we obtain the following density theorems.

Theorem 3.6. There is an absolute constant c' > 0 such that the following holds. Let \mathcal{R} be a collection of n red double grounded curves with grounds γ_1 and γ_2 , such that γ_1 and γ_2 cross. Let \mathcal{B} be a collection of n blue curves such that $\mathcal{R} \cup \mathcal{B} \cup \{\gamma_1, \gamma_2\}$ is a collection of pseudo-segments.

If there are at least εn^2 crossing pairs in $\mathcal{R} \times \mathcal{B}$, then there are subsets $\mathcal{R}' \subset \mathcal{R}, \mathcal{B}' \subset \mathcal{B}$, where $|\mathcal{R}'|, |\mathcal{B}'| \geq \varepsilon^{c'}n$, such that every curve in \mathcal{R}' crosses every curves in \mathcal{B}' .

Theorem 3.7. There is an absolute constant c' > 0 such that the following holds. Let \mathcal{R} be a collection of n red double grounded curves with grounds γ_1 and γ_2 , such that γ_1 and γ_2 cross. Let \mathcal{B} be a collection of n blue curves such that $\mathcal{R} \cup \mathcal{B}$ is a collection of pseudo-segments.

If there are at least εn^2 disjoint pairs in $\mathcal{R} \times \mathcal{B}$, then there are subsets $\mathcal{R}' \subset \mathcal{R}, \mathcal{B}' \subset \mathcal{B}$, where $|\mathcal{R}'|, |\mathcal{B}'| \geq \varepsilon^{c'}n$, such that every curve in \mathcal{R}' is disjoint from every curves in \mathcal{B}' .

We will apply Theorems 3.5 and 3.6 in the next section.

4 Proof of Theorem 2.8 – for ε -homogeneous families

The aim of this section is to prove Theorem 2.8, the main result of this paper, in the special case where the edge density of the intersection graph of the red curves is nearly 0 or nearly 1, and the same is true for the intersection graph of the blue curves. This will easily imply Theorem 2.8 in its full generality, as shown in the next section.

4.1 Low versus low density

By Corollary 2.14, it is sufficient to establish Theorem 2.8 in the special case where the intersection graphs $G(\mathcal{R})$ and $G(\mathcal{B})$ are both ε -homogeneous. Below, we first consider the cases when the edge

densities of both $G(\mathcal{R})ndG(\mathcal{B})$ are smaller than ε . Recall that a separator for a graph G = (V, E)is a subset $V_0 \subset V$ such that there is a partition $V = V_1 \cup V_1 \cup V_2$ with $|V_1|, |V_2| \leq \frac{2}{3}|V|$, and no vertex in V_1 is adjacent to any vertex in V_2 . The celebrated Lipton-Tarjan separator theorem states that every planar graph with n vertices has a separator of size $O(\sqrt{n})$. We will need the following result due to the first two authors. For a strengthening, see [28].

Lemma 4.1 ([12]). Let C be a collection of curves in the plane with a total of m crossings. Then the intersection graph G(C) has a separator of size $O(\sqrt{m})$.

We now prove the following.

Lemma 4.2. There is an absolute constant $\varepsilon_0 > 0$ such that the following holds. Let \mathcal{R} be a set of n red curves in the plane and let \mathcal{B} be a set of n blue curves such that $\mathcal{R} \cup \mathcal{B}$ is a collection of pseudo-segments.

If the edge densities of the intersection graphs $G(\mathcal{R})$ and $G(\mathcal{B})$ are both less than ε_0 , and there are at most $\varepsilon_0 n^2$ crossing pairs in $\mathcal{R} \times \mathcal{B}$, then there are subsets $\mathcal{R}' \subset \mathcal{R}$ and $\mathcal{B}' \subset \mathcal{B}$, each of size $\Omega(n)$, such that every red curve in \mathcal{R}' is disjoint from every blue curve in \mathcal{B}' .

Proof. Let ε_0 be a small constant that will be determined later. Set $C = \mathcal{R} \cup \mathcal{B}$. We apply Lemma 4.1 to C and obtain a partition $C = C_0 \cup C_1 \cup C_2$, such that $|C_1|, |C_2| \leq (2/3)(2n)$, and every curve in C_1 is disjoint from every curve in C_2 . Let c be the constant from Lemma 4.1. By setting $\varepsilon_0 < \frac{1}{(2c)^{2}16}$, we have

$$|\mathcal{C}_0| \le c\sqrt{\varepsilon_0(2n)^2} \le 2c\sqrt{\varepsilon_0}n \le n/4.$$

Without loss of generality, we can assume that at least half of the curves in C_1 are red, since a symmetric argument would follow otherwise. Since $|C_1| \ge (2n)/3 - |C_0|$, there are at least

$$\frac{2n/3 - n/4}{2} = 5n/24$$

red curves in C_1 . Moreover, there are at most 2n/3 blue curves in C_1 , which implies that there are at least

$$n - \frac{2n}{3} - \frac{n}{4} \ge \frac{n}{12},$$

blue curves in C_2 . By setting \mathcal{R}' to be the red curves in C_1 and \mathcal{B}' to be the blue curves in C_2 , we obtain the desired subsets.

Theorem 4.3. There is an absolute constant $\varepsilon_1 > 0$ such that the following holds. Let \mathcal{R} be a set of n red curves and \mathcal{B} be a set of n blue curves in the plane such that $\mathcal{R} \cup \mathcal{B}$ is a collection of pseudo-segments.

If the edge densities of the intersection graphs $G(\mathcal{R})$ and $G(\mathcal{B})$ are both less than ε_1 , then there are subsets $\mathcal{R}' \subset \mathcal{R}$ and $\mathcal{B}' \subset \mathcal{B}$, each of size $\Omega(n)$, such that every red curve in \mathcal{R}' crosses every blue curve in \mathcal{B}' , or every red curve in \mathcal{R}' is disjoint from every blue curve in \mathcal{B}' .

Proof. Let ε_1 be a small constant that will be determined later. Suppose there are at most $\varepsilon_0 n^2$ crossing pairs in $\mathcal{R} \times \mathcal{B}$, where ε_0 is defined in Lemma 4.2. Since the edge density of the intersection graphs $G(\mathcal{R})$ and $G(\mathcal{B})$ are both less than ε_1 , by setting $\varepsilon_1 < \varepsilon_0$, Lemma 4.2 implies that there



Figure 4: Partitioning of the red curve $\alpha = \alpha_u \cup \alpha_\ell$.

are subsets $\mathcal{R}' \subset \mathcal{R}$ and $\mathcal{B}' \subset \mathcal{B}$, each of size $\Omega(n)$, such that every red curve in \mathcal{R}' is disjoint from every blue curve in \mathcal{B}' and we are done.

If there are at at least $\varepsilon_0 n^2$ crossing pairs in $\mathcal{R} \times \mathcal{B}$, then, by Theorem 2.12, there are subsets $\mathcal{C}_1, \mathcal{C}_2 \subset \mathcal{R} \cup \mathcal{B}$, each of size $\varepsilon_0^{c'} n$, where c' > 1 is an absolute constant from Theorem 2.12, such that each curve in \mathcal{C}_1 crosses each curve in \mathcal{C}_2 . Without loss of generality, we can assume that at least $\varepsilon_0^{c'} n/2$ curves in \mathcal{C}_1 are red. Since $G(\mathcal{R})$ has edge density at most ε_1 , by setting $\varepsilon_1 = \varepsilon_0^{2c'}/4 < \varepsilon_0$, at most $\varepsilon_0^{c'} n/2$ curves in \mathcal{C}_2 are red. Hence, at least $\varepsilon_0^{c'} n/2$ curves in \mathcal{C}_2 are blue and we are done. \Box

4.2 High versus low edge density

In this subsection, we consider the case when the intersection graph $G(\mathcal{R})$ has edge density at least $1 - \varepsilon$, and $G(\mathcal{B})$ has edge density less than ε . Since the edge density in the intersection graph $G(\mathcal{R})$ is at least $1 - \varepsilon$, we can further reduce to the case when there is a red curve γ_1 that crosses every member in \mathcal{R} exactly once.

Lemma 4.4. For each integer $t \geq 1$, there is a constant $\varepsilon'_t > 0$ such that the following holds. Let \mathcal{R} be a set of n red curves in the plane, all crossed by a curve γ_1 exactly once, and \mathcal{B} be a set of n blue curves in the plane such that $\mathcal{R} \cup \mathcal{B} \cup \{\gamma_1\}$ is a collection of pseudo-segments. Suppose that the intersection graph $G(\mathcal{B})$ has edge density less than ε'_t , and $G(\mathcal{R})$ has edge density at least $1 - \varepsilon'_t$.

Then there are subsets $\hat{\mathcal{R}} \subset \mathcal{R}$, $\hat{\mathcal{B}} \subset \mathcal{B}$, each of size $\Omega_{\varepsilon'_t}(n)$, such that either every red curve in $\hat{\mathcal{R}}$ crosses every blue curve in $\hat{\mathcal{B}}$, or every red curve in $\hat{\mathcal{R}}$ is disjoint from every blue curve in $\hat{\mathcal{B}}$, or each curve $\alpha \in \hat{\mathcal{R}}$ has a partition into two connected parts $\alpha = \hat{\alpha}_u \cup \hat{\alpha}_\ell$, such that for

$$\hat{\mathcal{U}} = \{ \hat{\alpha}_u : \alpha \in \hat{\mathcal{R}}, \alpha = \hat{\alpha}_u \cup \hat{\alpha}_\ell \} \quad \text{and} \quad \hat{\mathcal{L}} = \{ \hat{\alpha}_\ell : \alpha \in \hat{\mathcal{R}}, \alpha = \hat{\alpha}_u \cup \hat{\alpha}_\ell \},$$

every curve in $\hat{\mathcal{L}}$ is disjoint to every curve in $\hat{\mathcal{B}}$, and the edge density of $G(\hat{\mathcal{U}})$ is less that 2^{-t} .

Proof. Each curve $\alpha \in \mathcal{R}$ is partitioned into two connected parts by γ_1 , say an upper and lower part. More precisely, we have the partition $\alpha = \alpha_u \cup \alpha_\ell$, where the parts α_u and α_ℓ are defined, as follows. We start at the left endpoint of γ_1 and move along γ_1 until we reach $\alpha \cap \gamma_1$. At this point, we turn left along α to obtain α_u and right to obtain α_ℓ . See Figure 4. Let $\mathcal{U}(\mathcal{L})$ be the upper (lower) part of each curve in \mathcal{R} , that is,

$$\mathcal{U} = \{ \alpha_u : \alpha \in \mathcal{R}, \alpha = \alpha_\ell \cup \alpha_u \} \quad \text{and} \quad \mathcal{L} = \{ \alpha_\ell : \alpha \in \mathcal{R}, \alpha = \alpha_\ell \cup \alpha_u \}.$$

In what follows, for every integer $t \ge 1$, we will obtain subsets $\mathcal{R}^{(t)} \subset \mathcal{R}$, $\mathcal{B}^{(t)} \subset \mathcal{B}$, each of size $\Omega_{\varepsilon'_{t}}(n)$, such that either every red curve in $\mathcal{R}^{(t)}$ crosses every blue curve in $\mathcal{B}^{(t)}$, or every red curve

in $\mathcal{R}^{(t)}$ is disjoint from every blue curve in $\mathcal{B}^{(t)}$, or each curve $\alpha \in \mathcal{R}^{(t)}$ has a new partition into two connected parts $\alpha = \alpha'_{\mu} \cup \alpha'_{\rho}$, upper and lower, such that the following holds.

- 1. We have $\alpha'_u \subset \alpha_u$, that is, the upper part α'_u is a subcurve of the previous upper part α_u .
- 2. The lower part α'_{ℓ} of each curve in $\mathcal{R}^{(t)}$ is disjoint from each blue curve in $\mathcal{B}^{(t)}$.
- 3. There is an equipartition $\mathcal{R}^{(t)} = \mathcal{R}_1^{(t)} \cup \cdots \cup \mathcal{R}_{2^t}^{(t)}$ into 2^t parts such that for $1 \le i < j \le 2^{t-1}$, the upper part α'_u of each curve $\alpha \in \mathcal{R}_i^{(t)}$ is disjoint from the upper part β'_u of each curve $\beta \in \mathcal{R}_j^{(t)}$.

Hence, the lemma follows from the statement above by setting $\hat{\mathcal{B}} = \mathcal{B}^{(t)}, \hat{\mathcal{R}} = \mathcal{R}^{(t)}$.

We proceed by induction on t. The bulk of the argument below is actually for the base case t = 1, since we will just repeat the entire argument for the inductive step with parameter ε'_t . Let ε'_1 be a small positive constant that will be determined later such that $\varepsilon'_1 < \varepsilon_1$, where ε_1 is from Theorem 4.3. Thus, $G(\mathcal{R})$ has edge density at least $1 - \varepsilon'_1$ and $G(\mathcal{B})$ has edge density less than ε'_1 .

Let $\delta > 0$ also be a sufficiently small constant determined later, such that $\varepsilon'_1 < \delta < \varepsilon_1$. We apply Corollary 2.14 to \mathcal{L} with parameter δ and obtain a subset $\mathcal{L}_1 \subset \mathcal{L}$ such that \mathcal{L}_1 is δ -homogeneous and $|\mathcal{L}_1| = \Omega_{\delta}(n)$. Let $\mathcal{R}_1 \subset \mathcal{R}$ be the red curves in \mathcal{R} corresponding to the curves in \mathcal{L}_1 , and let $\mathcal{U}_1 \subset \mathcal{U}$ be the curves in \mathcal{U} that corresponds to the red curves in \mathcal{R}_1 .

Without loss of generality, we can assume that the intersection graph $G(\mathcal{L}_1)$ has edge density less than δ . Indeed, otherwise if $G(\mathcal{L}_1)$ has edge density greater than $1 - \delta$, by the pseudosegment condition, the intersection graph $G(\mathcal{U}_1)$ must have edge density less than δ and a symmetric argument would follow. In order to apply Theorem 4.3, we need two subsets of equal size. By averaging, there is a subset $\mathcal{B}' \subset \mathcal{B}$ with $|\mathcal{B}'| = |\mathcal{L}_1|$ such that the edge density of $G(\mathcal{B}')$ is at most that of $G(\mathcal{B})$. Since $G(\mathcal{L}_1)$ has edge density less than δ and $G(\mathcal{B}')$ has edge density less than ε'_1 , by setting $\varepsilon'_1 < \delta < \varepsilon_1$, we can apply Theorem 4.3 to \mathcal{L}_1 and \mathcal{B}' and obtain subsets $\mathcal{L}_2 \subset \mathcal{L}_1$ and $\mathcal{B}_1 \subset \mathcal{B}'$, each of size $\Omega_{\delta}(n)$, such that every curve in \mathcal{L}_2 crosses every blue curve in \mathcal{B}_1 , or every curve in \mathcal{L}_2 is disjoint from every blue curve in \mathcal{B}_1 . If we are in the former case, then we are done. Hence, we can assume that we are in the latter case. Let $\mathcal{R}_2 \subset \mathcal{R}_1$ be the red curves that corresponds to \mathcal{L}_2 , and let $\mathcal{U}_2 \subset \mathcal{U}_1$ be the curves in \mathcal{U}_1 that corresponds to \mathcal{R}_2 . We apply Corollary 2.14 to \mathcal{U}_2 with parameter δ and obtain a subset $\mathcal{U}_3 \subset \mathcal{U}_2$ such that \mathcal{U}_3 is δ -homogeneous and $|\mathcal{U}_3| = \Omega_{\delta}(n)$. Let \mathcal{R}_3 be the red curves in \mathcal{R} corresponding to \mathcal{U}_3 , and let \mathcal{L}_3 be the curves in \mathcal{L}_2 that corresponds to \mathcal{R}_3 .

Suppose that the intersection graph $G(\mathcal{U}_3)$ has edge density less than δ . Since $|\mathcal{B}_1| = \delta_0 n$, where $\delta_0 = \delta_0(\delta, \varepsilon_1)$, by Lemma 2.15, the intersection graph $G(\mathcal{B}_1)$ has edge density at most $2\varepsilon'_1/\delta_0^2$. Thus, we set δ and ε'_1 sufficiently small so that $\delta < \varepsilon_1$ and $2\varepsilon'_1/\delta_0^2 < \varepsilon_1$. By averaging, we can find subsets of \mathcal{U}_3 and \mathcal{B}_1 , each of size min $(|\mathcal{U}_3|, |\mathcal{B}_1|)$ and with densities less than ε_1 , and apply Theorem 4.3 to these subsets and obtain subsets $\mathcal{U}_4 \subset \mathcal{U}_3$ and $\mathcal{B}_2 \subset \mathcal{B}_1$, each of size $\Omega_{\delta}(n)$, such that every curve in \mathcal{U}_4 crosses every blue curve in \mathcal{B}_2 , or every curve in \mathcal{U}_4 is disjoint from every blue curve in \mathcal{B}_2 . In both cases, we are done since every curve in \mathcal{L}_3 is disjoint from every curve in \mathcal{B}_2 . Therefore, we can assume that $G(\mathcal{U}_3)$ has edge density greater than $1 - \delta$.

For each curve $\alpha \in \mathcal{U}_3$, let $N(\alpha)$ denote the set of curves in \mathcal{U}_3 that intersects α , and let $d(\alpha) = |N(\alpha)|$. We label the curves $\beta \in N(\alpha)$ with integers 0 to $d(\alpha) - 1$ according to their closest intersection point to the ground γ_1 along α , that is, the label $f_{\alpha}(\beta)$ of $\beta \in N(\alpha)$ is the number of curves in \mathcal{U}_3 that intersects the portion of α strictly between γ_1 and $\alpha \cap \beta$. Since

$$\sum_{\alpha \in \mathcal{U}_3} d(\alpha) - 1 \ge 2(1-\delta) \binom{|\mathcal{U}_3|}{2} - |\mathcal{U}_3|,$$

by Jensen's inequality, we have

$$\sum_{\alpha \in \mathcal{U}_3} \sum_{\beta \in N(\alpha)} f_{\alpha}(\beta) = \sum_{\alpha \in \mathcal{U}_3} \binom{d(\alpha)}{2} \ge |\mathcal{U}_3| \binom{\frac{\sum_{\alpha \in \mathcal{U}_3} d(\alpha)}{|\mathcal{U}_3|}}{2} \ge \frac{|\mathcal{U}_3|^3}{4}.$$

Let the weight $w(\beta)$ of a curve $\beta \in \mathcal{U}_3$ be the sum of its labels, that is,

$$w(\beta) = \sum_{\alpha:\beta \in N(\alpha)} f_{\alpha}(\beta).$$

Hence, the weight $w(\beta)$ is the total number of crossing points along curves α strictly between γ_1 and β , where α crosses both γ_1 and β . By averaging, there is a curve $\gamma_2 \in \mathcal{U}_3$ whose weight is at least $|\mathcal{U}_3|^2/4$.

Using γ_2 , we partition each curve $\alpha \in \mathcal{U}_3 \setminus \{\gamma_2\}$ that crosses γ_2 into two connected parts, $\alpha = \alpha_w \cup \alpha_m$, where α_m is the connected subcurve with endpoints on γ_1 and γ_2 , and α_w is the other connected part. Set

$$\mathcal{W}_3 = \{ \alpha_w : \alpha \in \mathcal{U}_3 \setminus \{\gamma_2\}, \alpha \cap \gamma_2 \neq \emptyset \} \text{ and } \mathcal{M}_3 = \{ \alpha_m : \alpha \in \mathcal{U}_3 \setminus \{\gamma_2\}, \alpha \cap \gamma_2 \neq \emptyset \}.$$

Since γ_2 has weight at least $|\mathcal{U}_3|^2/4$, by the pigeonhole principle, there are at least $|\mathcal{U}_3|^2/8$ intersecting pairs in $\mathcal{M}_3 \times \mathcal{M}_3$, or at least $|\mathcal{U}_3|^2/8$ intersecting pairs in $\mathcal{M}_3 \times \mathcal{W}_3$.

Case 1. Suppose there are at least $|\mathcal{U}_3|^2/8$ pairs in $\mathcal{M}_3 \times \mathcal{W}_3$ that cross. The set \mathcal{M}_3 is double grounded with grounds γ_1 and γ_2 that cross exactly once, and every curve in \mathcal{W}_3 is disjoint from γ_1 and γ_2 . As $|\mathcal{M}_3|, |\mathcal{W}_3| \leq |\mathcal{U}_3|$, the density of edges in the bipartite intersection graph of \mathcal{M}_3 and \mathcal{W}_3 is at least 1/8. By averaging, we can find subsets of \mathcal{M}_3 and \mathcal{W}_3 each of size min $(|\mathcal{M}_3|, |\mathcal{W}_3|)$ such that the density of edges in the bipartite intersection graph of these subsets is at least 1/8. By setting $\delta > 0$ sufficiently small, we can apply Theorem 3.6 to these subsets of \mathcal{M}_3 and \mathcal{W}_3 and obtain subsets $\mathcal{M}_4 \subset \mathcal{M}_3$ and $\mathcal{W}'_4 \subset \mathcal{W}_3$, each of size $\Omega_{\delta}(n)$, such that each curve in \mathcal{M}_4 crosses each curve in \mathcal{W}'_4 . Moreover, by the pseudo-segment condition, each curve in $\mathcal{M}_4 \cup \mathcal{W}'_4$ corresponds to a unique curve in \mathcal{U}_3 . Let $\mathcal{U}_4 \subset \mathcal{U}_3$ be the curves that corresponds to \mathcal{M}_4 and let $\mathcal{U}'_4 \subset \mathcal{U}_3$ be the curves that corresponds to \mathcal{W}'_4 . Hence, we set

$$\mathcal{W}_4 = \{ \alpha_m : \alpha \in \mathcal{U}_4, \alpha = \alpha_w \cup \alpha_m \} \text{ and } \mathcal{M}'_4 = \{ \alpha_m : \alpha \in \mathcal{U}'_4, \alpha = \alpha_w \cup \alpha_m \}.$$

See Figure 5(a).

We apply Theorem 3.5 to arbitrary subsets of \mathcal{M}_4 and \mathcal{B}_2 , each of size $\min(|\mathcal{M}_4|, |\mathcal{B}_2|)$, and obtain subsets $\mathcal{M}_5 \subset \mathcal{M}_4$ and $\mathcal{B}_3 \subset \mathcal{B}_2$, each of size $\Omega_{\delta}(n)$, such that either every red curve in \mathcal{M}_5 crosses every blue curve in \mathcal{B}_3 , or every red curve in \mathcal{M}_5 is disjoint from every blue curve in \mathcal{B}_3 . In the former case, we are done. Hence, we can assume that we are in the latter case.

We again apply Theorem 3.5 to arbitrary subsets of \mathcal{M}'_4 and \mathcal{B}_3 , each of size $\min(|\mathcal{M}'_4|, |\mathcal{B}_3|)$, to obtain subsets $\mathcal{M}'_5 \subset \mathcal{M}'_4$ and $\mathcal{B}_4 \subset \mathcal{B}_3$, each of size $\Omega_\delta(n)$, such that either every red curve in \mathcal{M}'_5 crosses every blue curve in \mathcal{B}_4 , or every red curve in \mathcal{M}'_5 is disjoint from every blue curve in



Figure 5: In both cases, \mathcal{W}_4 is disjoint to \mathcal{W}'_4 .

 \mathcal{B}_4 . Again, if we are in the former case, we are done. Hence, we can assume that we are in the latter case. Let

$$\mathcal{W}_5 = \{ \alpha_w : \alpha = \alpha_w \cup \alpha_m, \alpha_m \in \mathcal{M}_5 \} \text{ and } \mathcal{W}'_5 = \{ \alpha_w : \alpha = \alpha_w \cup \alpha_m, \alpha_m \in \mathcal{M}'_5 \},\$$

and recall that every element in \mathcal{M}_5 crosses every element in \mathcal{W}'_5 . By the pseudo-segment condition, every element in \mathcal{W}_5 is disjoint from every element in \mathcal{W}'_5 .

Let \mathcal{R}_5 be the red curves in \mathcal{R} that corresponds to \mathcal{W}_5 , and let \mathcal{R}'_5 be the red curves in \mathcal{R} that corresponds to \mathcal{W}'_5 . We have $|\mathcal{R}_5|, |\mathcal{R}'_5| = \Omega_\delta(n)$, and moreover, we can assume that $|\mathcal{R}_5| = |\mathcal{R}'_5|$. For each curve $\alpha \in \mathcal{R}_5 \cup \mathcal{R}'_5$, and its original partition $\alpha = \alpha_u \cup \alpha_\ell$ defined by γ_1 , we have a new partition $\alpha = \alpha'_u \cup \alpha'_\ell$ defined by γ_2 , where $\alpha'_u = \alpha_w$ and $\alpha'_\ell = \alpha_m \cup \alpha_\ell$. By setting $\mathcal{R}^{(1)} = \mathcal{R}_5 \cup \mathcal{R}'_5$, and $\mathcal{B}^{(1)} = \mathcal{B}_4$, where each curve $\alpha \in \mathcal{R}^{(1)}$ is equipped with the partition $\alpha = \alpha'_u \cup \alpha'_\ell$, we satisfy the base case of the statement.

Case 2. The argument is essentially the same as Case 1. Suppose we have at least $|\mathcal{U}_3|^2/8$ crossing pairs in $\mathcal{M}_3 \times \mathcal{M}_3$. Then by Theorem 2.9, there are subsets $\mathcal{M}_4, \mathcal{M}'_4 \subset \mathcal{M}_3$, each of size $\Omega_{\delta}(n)$, such that every curve in \mathcal{M}_4 crosses every curve in \mathcal{M}'_4 . Let $\mathcal{U}_4 \subset \mathcal{U}$ be the curves that corresponds to \mathcal{M}_4 and let $\mathcal{U}'_4 \subset \mathcal{U}$ be the curves that corresponds to \mathcal{M}'_4 . Set

$$\mathcal{W}_4 = \{ \alpha_w : \alpha \in \mathcal{U}_4, \alpha = \alpha_w \cup \alpha_m \} \text{ and } \mathcal{W}_4' = \{ \alpha_w : \alpha \in \mathcal{U}_4', \alpha = \alpha_w \cup \alpha_m \}.$$

See Figure 5(b).

Hence, by the pseudo-segment condition, every curve in \mathcal{W}_4 is disjoint from every curve in \mathcal{W}'_4 . By taking arbitrary subsets of \mathcal{M}_4 and \mathcal{B}_2 of size $\min(|\mathcal{M}_4|, |\mathcal{B}_2|)$, we can apply Theorem 3.5 to these subsets and obtain subsets $\mathcal{M}_5 \subset \mathcal{M}_4$ and $\mathcal{B}_3 \subset \mathcal{B}_2$, each of size $\Omega_\delta(n)$, such that either every red curve in \mathcal{M}_5 crosses every blue curve in \mathcal{B}_3 , or every red curve in \mathcal{M}_5 is disjoint from every blue curve in \mathcal{B}_3 . In the former case, we are done. Hence, we can assume that we are in the latter case.

Again, we take an arbitrary subset of \mathcal{M}'_4 and \mathcal{B}_3 of size $\min(|\mathcal{M}'_4|, |\mathcal{B}_3|)$ and apply Theorem 3.5 to \mathcal{M}'_4 and \mathcal{B}_3 , to obtain subsets $\mathcal{M}'_5 \subset \mathcal{M}'_4$ and $\mathcal{B}_4 \subset \mathcal{B}_3$, each of size $\Omega_\delta(n)$, such that either every red curve in \mathcal{M}'_5 crosses every blue curve in \mathcal{B}_4 , or every red curve in \mathcal{M}'_5 is disjoint from every blue curve in \mathcal{B}_4 . Again, if we are in the former case, we are done. Hence, we can assume that we are in the latter case. Set \mathcal{R}_5 be the red curves in \mathcal{R} that corresponds to \mathcal{M}_5 , and let \mathcal{R}'_5 be the red curves in \mathcal{R} that corresponds to \mathcal{M}'_5 .

We have $|\mathcal{R}_5|, |\mathcal{R}'_5| = \Omega_{\delta}(n)$, and moreover, we can assume that $|\mathcal{R}_5| = |\mathcal{R}'_5|$. For each curve $\alpha \in \mathcal{R}_5 \cup \mathcal{R}'_5$, and its original partition $\alpha = \alpha_u \cup \alpha_\ell$ defined by γ_1 , we have a new partition

 $\alpha = \alpha'_u \cup \alpha'_\ell$ defined by γ_2 , where $\alpha'_u = \alpha_w$ and $\alpha'_\ell = \alpha_m \cup \alpha_\ell$. By setting $\mathcal{R}^{(1)} = \mathcal{R}_5 \cup \mathcal{R}'_5$, and $\mathcal{B}^{(1)} = \mathcal{B}_4$, where each curve $\alpha \in \mathcal{R}^{(1)}$ is equipped with the partition $\alpha = \alpha'_u \cup \alpha'_\ell$, we satisfy the base case of the statement.

For the inductive step, suppose we have obtained constants $\varepsilon'_{t-1} < \cdots < \varepsilon'_1$ such that the statement follows. Let ε'_t be a small constant that will be determined later such that $\varepsilon'_t < \varepsilon'_{t-1}$. Let \mathcal{R} be a set of n red curves in the plane, all crossed by a curve γ_1 exactly once, and \mathcal{B} be a set of n blue curves in the plane such that $\mathcal{R} \cup \mathcal{B} \cup \{\gamma_1\}$ is a collection of pseudo-segments. Moreover, $G(\mathcal{R})$ has edge density at least $1 - \varepsilon'_t$ and $G(\mathcal{B})$ has edge density less than ε'_t . We set $\delta' < 0$ to be a small constant such that $\varepsilon'_t < \delta' < \varepsilon_{t-1}$. We repeat the entire argument above, replacing ε'_1 with ε'_t and δ with δ' , to obtain subsets $\mathcal{R}_5, \mathcal{R}'_5 \subset \mathcal{R}$ and $\mathcal{B}_4 \subset \mathcal{B}$, each of size $\Omega_{\delta'}(n)$, such that each $\alpha \in \mathcal{R}_5 \cup \mathcal{R}'_5$ is equipped with the partition $\alpha = \alpha'_u \cup \alpha'_\ell$, and α'_ℓ is disjoint to β'_u .

Since $|\mathcal{R}_5|, |\mathcal{B}_4| \geq \delta_1 n$, where δ_1 depends only on δ' , by Theorem 2.15, $G(\mathcal{R}_5)$ has edge density at least $1 - 2\varepsilon'_t/\delta_1^2$ and $G(\mathcal{B}_4)$ has edge density less than $2\varepsilon'_t/\delta_1^2$. By setting ε'_t sufficiently small, $G(\mathcal{R}_5)$ has edge density at least $1 - \varepsilon'_{t-1}$, and $G(\mathcal{B}_4)$ has edge density less than ε'_{t-1} . By averaging, we can find subsets of \mathcal{R}_5 and \mathcal{B}_4 , each of size $\min(|\mathcal{R}_5|, |\mathcal{B}_4|)$ and with densities at least $1 - \varepsilon'_{t-1}$ and less than ε'_{t-1} respectively, and apply induction to these subsets parameter t' = t - 1, and obtain subsets $\mathcal{R}^{(t-1)} \subset \mathcal{R}_5$, $\mathcal{B}^{(t-1)} \subset \mathcal{B}_4$, each of size $\Omega_{\varepsilon'_{t-1}}(n)$, with the desired properties. If every red curve in $\mathcal{R}^{(t-1)}$ is disjoint from every blue curve in $\mathcal{B}^{(t-1)}$, or if every red curve in $\mathcal{R}^{(t-1)}$ crosses every blue curve in $\mathcal{B}^{(t-1)}$, then we are done. Hence, we can assume that each curve $\alpha \in \mathcal{R}^{(t-1)}$ has a partition $\alpha = \alpha''_u \cup \alpha''_\ell$ such that α''_u is a subcurve of α'_u , α''_ℓ is disjoint from every blue curve in $\mathcal{B}^{(t-1)}$, and there is an equipartition

$$\mathcal{R}^{(t-1)} = \mathcal{R}_1^{(t-1)} \cup \cdots \cup \mathcal{R}_{2^{t-1}}^{(t-1)},$$

such that for $1 \leq i < j \leq 2^{t-1}$, the upper part α''_u of each curve $\alpha \in \mathcal{R}_i^{(t-1)}$ is disjoint the upper part β''_u of each curve $\beta \in \mathcal{R}_i^{(t-1)}$.

Finally, since $|\mathcal{R}'_5|, |\mathcal{B}^{(t-1)}| \geq \delta_2 n$, where δ_2 depends only on δ' , by Theorem 2.15, $G(\mathcal{R}'_5)$ has edge density at least $1 - 2\varepsilon'_t/\delta_2^2$ and $G(\mathcal{B}^{(t-1)})$ has edge density less than $2\varepsilon'_t/\delta_2^2$. By setting ε'_t sufficiently small, $G(\mathcal{R}'_5)$ has edge density at least $1 - \varepsilon'_{t-1}$, and $G(\mathcal{B}^{(t-1)})$ has edge density less than ε'_{t-1} . By averaging, we can find subsets of \mathcal{R}'_5 and $\mathcal{B}^{(t-1)}$, each of size $\min(|\mathcal{R}'_5|, |\mathcal{B}^{(t-1)}|)$ and with densities at least $1 - \varepsilon'_{t-1}$ and less than ε'_{t-1} respectively, and apply induction to these subsets parameter t' = t - 1, and obtain subsets $\mathcal{S}^{(t-1)} \subset \mathcal{R}'_5$, $\mathcal{B}^{(t)} \subset \mathcal{B}^{(t-1)}$, each of size $\Omega_{\varepsilon'_{t-1}}(n)$, with the desired properties. If every red curve in $\mathcal{S}^{(t-1)}$ is disjoint from every blue curve in $\mathcal{B}^{(t)}$, or if every red curve in $\mathcal{S}^{(t-1)}$ crosses every blue curve in $\mathcal{B}^{(t)}$, then we are done. Hence, we can assume that each curve $\alpha \in \mathcal{S}^{(t-1)}$ has a partition $\alpha = \alpha''_u \cup \alpha''_\ell$ such that α''_u is a subcurve of α'_u, α''_ℓ is disjoint from every blue curve in $\mathcal{B}^{(t-1)}$, and there is an equipartition

$$\mathcal{S}^{(t-1)} = \mathcal{S}_1^{(t-1)} \cup \dots \cup \mathcal{S}_{2^{t-1}}^{(t-1)},$$

such that for $1 \leq i < j \leq 2^{t-1}$, the upper part α''_u of each curve $\alpha \in \mathcal{S}_i^{(t-1)}$ is disjoint the upper part β''_u of each curve $\beta \in \mathcal{S}_j^{(t-1)}$. We then (arbitrarily) remove curves from each part in $\mathcal{R}_i^{(t-1)}$ and $\mathcal{S}_j^{(t-1)}$ such that the resulting parts all have the same size and for

$$\mathcal{R}^{(t)} = \mathcal{R}_1^{(t-1)} \cup \dots \cup \mathcal{R}_{2^{t-1}}^{(t-1)} \cup \mathcal{S}_1^{(t-1)} \cup \dots \cup \mathcal{S}_{2^{t-1}}^{(t-1)},$$

we have $|\mathcal{R}^{(t)}| = \Omega_{\varepsilon'_{t-1}}(n)$. Then $\mathcal{R}^{(t)}$ and $\mathcal{B}^{(t)}$ has the desired properties.

We now prove the following.

Theorem 4.5. There is an absolute constant $\varepsilon_3 > 0$ such that the following holds. Let \mathcal{R} be a set of n red curves in the plane and \mathcal{B} be a set of n blue curves in the plane such that $\mathcal{R} \cup \mathcal{B}$ is a collection of pseudo-segments, and the intersection graph $G(\mathcal{B})$ has edge density less than ε_3 , and $G(\mathcal{R})$ has edge density at least $1 - \varepsilon_3$.

Then there are subsets $\mathcal{R}' \subset \mathcal{R}$, $\mathcal{B}' \subset \mathcal{B}$, each of size $\Omega(n)$, such that either every red curve in \mathcal{R} crosses every blue curve in \mathcal{B} , or every red curve in \mathcal{R} is disjoint from every blue curve in \mathcal{B} .

Proof. Let t be a fixed large integer such that $2^{-t} < \varepsilon_1$, where ε_1 is defined in Theorem 4.3. Let ε_3 be a small constant determined later such that $\varepsilon_3 < \varepsilon'_t$, where ε'_t is defined in Lemma 4.4. Recall that $\varepsilon'_t < \varepsilon_1$. Since $G(\mathcal{R})$ has edge density at least $1 - \varepsilon_3$, there is a curve $\gamma_1 \in \mathcal{R}$ such that γ_1 crosses at least n/2 red curves in \mathcal{R} . Let $\mathcal{R}_0 \subset \mathcal{R}$ be the red curves that crosses γ_1 . By Lemma 2.15, $G(\mathcal{R}_0)$ has edge density at least $1 - \varepsilon_3$. By averaging, we can find a subset $\mathcal{B}' \subset \mathcal{B}$ of size $|\mathcal{R}_0|$ whose edge density less than ε_3 . By setting ε_3 sufficiently small so that $\varepsilon_3 < \varepsilon'_t$, we can apply Lemma 4.4 to \mathcal{R}_0 and \mathcal{B}' with parameter t, and obtain subsets $\hat{\mathcal{R}} \subset \mathcal{R}_0$, $\hat{\mathcal{B}} \subset \mathcal{B}$, each of size $\Omega_{\varepsilon'_t}(n)$, with the desired properties. If every red curve in $\hat{\mathcal{R}}$ then we are done. Therefore, we can assume that each curve $\alpha \in \hat{\mathcal{R}}$ has a partition into two parts $\alpha = \alpha'_u \cup \alpha'_\ell$ with the properties described in Lemma 4.4. Set

$$\mathcal{U} = \{ \alpha'_u : \alpha \in \hat{\mathcal{R}}, \alpha = \alpha'_u \cup \alpha'_\ell \} \quad \text{ and } \quad \mathcal{L} = \{ \alpha'_\ell : \alpha \in \hat{\mathcal{R}}, \alpha = \alpha'_u \cup \alpha'_\ell \}.$$

Hence, every curve in \mathcal{L} is disjoint from every curve in $\hat{\mathcal{B}}$, and $G(\mathcal{U})$ has edge density at most $2^{-t} < \varepsilon_1$. Since $|\hat{\mathcal{B}}| \ge \delta n$, where δ depends only on ε'_t , by Lemma 2.15, $G(\hat{\mathcal{B}})$ has edge density at most $2\varepsilon_3/\delta^2$. By setting ε_3 sufficiently small so that $2\varepsilon_3/\delta_0^2 < \varepsilon_1$, $G(\hat{\mathcal{B}})$ has edge density at most ε_1 . By averaging, we can find subsets of \mathcal{U} and $\hat{\mathcal{B}}$, each of size $\min(|\mathcal{U}|, |\hat{\mathcal{B}}|)$ and with densities at most ε_1 , and apply Theorem 4.3 to these subsets to obtain subsets $\mathcal{U}' \subset \mathcal{U}$ and $\mathcal{B}' \subset \hat{\mathcal{B}}$, each of size $\Omega_{\varepsilon_3}(n)$, such that every curve in \mathcal{U}' is disjoint from every curve in \mathcal{B}' , or every curve in \mathcal{U}' , every red curve in \mathcal{R}' is disjoint from every blue curve in \mathcal{B}' , or every red curve in \mathcal{B}' , and each subset has size $\Omega_{\varepsilon_3}(n)$.

4.3 High versus high edge density

Finally, we consider the case when the intersection graphs $G(\mathcal{R})$ and $G(\mathcal{B})$ both have edge density at least $1 - \varepsilon$. By copying the proof of Theorem 4.5, except using Theorem 4.5 (high versus low density) instead of Theorem 4.3 (low versus low density) in the argument, we obtain the following.

Theorem 4.6. There is an absolute constant $\varepsilon_4 > 0$ such that the following holds. Let \mathcal{R} be a set of n red curves in the plane and \mathcal{B} be a set of n blue curves in the plane such that $\mathcal{R} \cup \mathcal{B}$ is a collection of pseudo-segments, and the intersection graphs $G(\mathcal{B})$ and $G(\mathcal{R})$ both have edge density at least $1 - \varepsilon_4$.

Then there are subsets $\mathcal{R}' \subset \mathcal{R}$, $\mathcal{B}' \subset \mathcal{B}$, each of size $\Omega(n)$, such that either every red curve in \mathcal{R} crosses every blue curve in \mathcal{B} , or every red curve in \mathcal{R} is disjoint from every blue curve in \mathcal{B} .

5 Proof of Theorem 2.8

Let \mathcal{R} be a set of n red curves in the plane, and \mathcal{B} be a set of n blue curves in the plane such that $\mathcal{R}\cup\mathcal{B}$ is a collection of pseudo-segments. Let ε be a sufficiently small constant such that $\varepsilon < \varepsilon_4 < \varepsilon_3 < \varepsilon_1$, where ε_1 is from Theorem 4.3, ε_3 is from Theorem 4.5, and ε_4 is from Theorem 4.6. We apply Corollary 2.14 to both \mathcal{R} and \mathcal{B} and obtain subsets $\mathcal{R}_1 \subset \mathcal{R}$ and $\mathcal{B}_1 \subset \mathcal{B}$ such that both $G(\mathcal{R}_1)$ and $G(\mathcal{B}_1)$ are ε -homogeneous. Moreover, we can assume that $|\mathcal{R}_1| = |\mathcal{B}_1|$.

If both $G(\mathcal{R}_1)$ and $G(\mathcal{B}_1)$ have edge densities less than ε , then, since ε is sufficiently small, we can apply Theorem 4.3 to obtain subsets $\mathcal{R}_2 \subset \mathcal{R}_1$ and $\mathcal{B}_2 \subset \mathcal{B}_1$, each of size $\Omega_{\varepsilon}(n)$, such that either every red curve in \mathcal{R}_2 is disjoint from every blue curve in \mathcal{B}_2 , or every red curve in \mathcal{R}_2 crosses every blue curve in \mathcal{B}_2 .

If one of the graphs of $G(\mathcal{R}_1)$ and $G(\mathcal{B}_1)$ has edge density less than ε , and the other has edge density greater than $1 - \varepsilon$, then we apply Theorem 4.5 to \mathcal{R}_1 and \mathcal{B}_1 to obtain subsets $\mathcal{R}_2 \subset \mathcal{R}_1$ and $\mathcal{B}_2 \subset \mathcal{B}_1$, each of size $\Omega_{\varepsilon}(n)$, such that either every red curve in \mathcal{R}_2 is disjoint from every blue curve in \mathcal{B}_2 , or every red curve in \mathcal{R}_2 crosses every blue curve in \mathcal{B}_2 .

Finally, if both $G(\mathcal{R}_1)$ and $G(\mathcal{B}_1)$ have edge densities at least $1 - \varepsilon$, then, since ε is sufficiently small, we can apply Theorem 4.6 to obtain subsets $\mathcal{R}_2 \subset \mathcal{R}_1$ and $\mathcal{B}_2 \subset \mathcal{B}_1$, each of size $\Omega_{\varepsilon}(n)$, such that either every red curve in \mathcal{R}_2 is disjoint from every blue curve in \mathcal{B}_2 , or every red curve in \mathcal{R}_2 crosses every blue curve in \mathcal{B}_2 .

6 Topological graphs with no k pairwise disjoint edges

In this section, we prove Theorem 1.2. Recall that the *odd-crossing number* of G, denoted by odd-cr(G), is the minimum possible number of pairs of edges that cross an odd number of times, over all drawings of G. The *bisection width* of a graph G is defined as

$$b(G) = \min_{|V_1|, |V_2| \le 2n/3} |E(V_1, V_2)|,$$

where the minimum is take over all partitions $V(G) = V_1 \cup V_2$, such that $|V_1|, |V_2| \leq 2n/3$. The following lemma, due to Pach and Tóth [37], relates the odd-crossing number of a graph to its bisection width.

Lemma 6.1 ([38]). If G is a graph with n vertices of degrees d_1, \ldots, d_n , then

$$b(G) \le O\left(\log n \sqrt{\operatorname{odd-cr}(G) + \sum_{i=1}^{n} d_i^2}\right).$$

Since all graphs contain a bipartite subgraph with at least half of its edges, Theorem 1.2 immediately follows from the following theorem.

Theorem 6.2. If G = (V, E) is an n-vertex simple topological bipartite graph with no k pairwise disjoint edges, then $|E(G)| \leq n(\log n)^{O(\log k)}$.

Proof. Let c > 0 and $c_1 > 0$ be the absolute constants from Theorem 2.11 and Lemma 6.1 respectively, and let $c_2 > 0$ be a sufficiently large constant that will be determined later. We define f(n,k) to be the maximum number of edges in an *n*-vertex simple topological bipartite graph with no k pairwise disjoint edges. We will prove by induction on n and k that



Figure 6: Redrawing procedure

 $f(n,k) \le n(\log n)^{c_2 \log k}.$

Clearly, we have $f(2,k) \leq 1$, and by the results on thrackles imply that $f(n,2) \leq 1.4n$. Assume the statement holds for n' < n and k' < k, and let G be an n-vertex simple topological bipartite graph with no k pairwise disjoint edges. The proof falls into two cases.

Case 1. Suppose there are at least $|E(G)|^2/((2c_1)^2 \log^6 n)$ disjoint pairs of edges in G. Let us partition $E(G) = E_1 \cup E_2$ into two parts, such that there are at least $|E(G)|^2/(2(2c_1)^2 \log^6 n)$ disjoint pairs in $E_1 \times E_2$. Since E(G) is a collection of pseudo-segments, by Theorem 2.11, there exist subsets $E'_1 \subset E_1, E'_2 \subset E_2$, each of size $|E(G)|/(2c_1 \log n)^{6c}$, such that every edge in E'_1 is disjoint from every edge in E'_2 . Since G does not contain k pairwise disjoint edges, this implies that either E'_1 or E'_2 does not contain k/2 pairwise disjoint edges. Without loss of generality, suppose E'_1 does not contain k/2 pairwise disjoint edge. Hence

$$\frac{|E(G)|}{(2c_1 \log n)^{6c}} \le |E_1'| \le f(n, k/2).$$

By the induction hypothesis, we have

$$f(n, k/2) \le n(\log n)^{c_2 \log(k/2)} \le n(\log n)^{c_2 \log(k) - c_2}$$

Hence, for c_2 sufficiently large, we have $|E(G)| \le n(\log n)^{c_2 \log(k)}$.

Case 2. Suppose there are at most $|E(G)|^2/((2c_1)^2 \log^6 n)$ disjoint pairs of edges in G. In what follows, we will apply a redrawing technique due to Pach and Tóth in [38]. Since G is bipartite, let $V(G) = V_1 \cup V_2$. We can redraw G such that the vertices in V_1 lie above the line y = 1, the vertices in V_2 lie below the line y = 0, the edges in the strip $0 \le y \le 1$ are vertical segments, and we have neither created nor removed any crossings. We then "flip" the half-plane bounded by the y = 1 line from left to right about the y-axis, and replace the edges in the strip $0 \le y \le 1$ with straight line segments that reconnect the corresponding pairs on the line y = 0 and y = 1. See Figure 6.

Notice that if any two edges crossed in the original drawing, then they must cross an even number of times in the new drawing. Indeed, suppose the edges e_1 and e_2 crossed in the original drawing. By the simple condition, they share exactly 1 point in common. Let k_i denote the number of times edge e_i crosses the strip for $i \in \{1, 2\}$, and note that k_i must be odd since V_1 lies above the line y = 1 and v_2 lies below the line y = 0. After we have redrawn our graph, these $k_1 + k_2$ segments inside the strip will now pairwise cross, creating $\binom{k_1+k_2}{2}$ crossing points. Since edge e_i will now cross itself $\binom{k_i}{2}$ times, this implies that there are now

$$\binom{k_1+k_2}{2} - \binom{k_1}{2} - \binom{k_2}{2} \tag{1}$$

crossing points between edges e_1 and e_2 inside the strip $0 \le y \le 1$. One can easily check that (1) is odd when k_1 and k_2 are odd. Since e_1 and e_2 had 1 point in common outside of the strip, this implies that e_1 and e_2 cross each other an even number of times in the new drawing. Moreover, we can easily get rid of self-intersections by making modifications in a small ball at these crossing points.

Hence, the odd-crossing number in our new drawing is at most the number of disjoint pair of edges in the original drawing of G, plus the number of pair of edges that share a common vertex. Since there are at most

$$\sum_{v \in V(G)} d^2(v) \le 2|E(G)|n$$

pairs of edges that share a vertex in G, this implies

odd-cr(G)
$$\leq \frac{|E(G)|^2}{(2c_1)^2 \log^6 n} + 2|E(G)|n.$$

By Lemma 6.1, there is a partition of the vertex set $V = V_1 \cup V_2$ with $|V|/3 \le |V_i| \le 2|V|/3$, where i = 1, 2, and

$$|E(V_1, V_2)| \le b(G) \le c_1 \log n \sqrt{\frac{|E(G)|^2}{(2c_1)^2 \log^6 n} + 4n|E(G)|}.$$

If $|E(G)|^2/((2c_1)^2 \log^6 n) \le 4n|E(G)|$, then for c_2 sufficiently large we have $|E(G)| \le n(\log n)^{c_2 \log k}$ and we are done. Therefore, we can assume

$$b(G) \le c_1 \log n \sqrt{\frac{2|E(G)|^2}{(2c_1)^2 \log^6 n}} \le \frac{|E(G)|}{\log^2 n}.$$

Let $|V_1| = n_1$ and $|V_2| = n_2$. Since $n_1, n_2 < n$, by the induction hypothesis, we have

$$|E(G)| \leq b(G) + n_1(\log n_1)^{c_2 \log k} + n_2(\log n_2)^{c_2 \log k}$$

$$\leq \frac{|E(G)|}{\log^2 n} + n(\log(2n/3))^{c_2 \log k}$$

$$\leq \frac{|E(G)|}{\log^2 n} + n(\log n - \log(3/2))^{c_2 \log k},$$

which implies

$$|E(G)| \le n(\log n)^{c_2 \log k} \frac{(1 - \log(3/2)/\log n)^{c_2 \log k}}{1 - 1/\log^2 n} \le n(\log n)^{c_2 \log k}$$

7 Concluding remarks

A graph is said to be a *cograph* if it consists of a single-vertex or if it can be obtained from two smaller cographs by either taking their disjoint union or their join. We next define the depth of a cograph. The graph on one vertex has depth 0. The depth of a cograph which is the disjoint union or join of two smaller cographs is one more than the maximum depth of the two smaller cographs. For example, the depth of a complete or empty graph on k vertices is $\lceil \log_2 k \rceil$.

The proof of Theorem 1.2 can be generalized as follows.

Theorem 7.1. Let F be a cograph with depth d. If G = (V, E) is an n-vertex simple topological graph with no matching M whose intersection graph is isomorphic to F, then $|E(G)| \leq n(\log n)^{O(d)}$.

Proof sketch. The proof is very similar to the proof of Theorem 6.2. We proceed by induction on the depth d and n. When d = 0 or n = 1, the statement is trivial. For the inductive step, let $F = F_1 \cup F_2$, where F_1 and F_2 are two smaller cographs. If F is the disjoint union of F_1 and F_2 , then we follow the proof of Theorem 6.2.

If F is the join of F_1 and F_2 , then the proof is again very similar to the proof of Theorem 6.2. We consider the two cases when G has at least $|E(G)|^2/(c\log^6 n)$ crossing edges, or less than $|E(G)|^2/(c\log^6 n)$ crossing edges, where c is a large constant. In the former case, we apply Theorem 2.12 to the edges of G and obtain two subsets $E_1, E_2 \subset E(G)$, each of size $|E(G)|/(2c\log n)^{6c'}$ such that every edge in E_1 crosses every edge in E_1 and c' is an absolute constant. Hence we can apply induction to each E_i and we are done. In the latter case, we apply a bisection width formula due to Pach, Shahrokhi, and Szegedy [34] and apply induction on n.

A natural k-grid in a topological graph is a pair of edge sets E_1, E_2 , such that E_i consists of k pairwise disjoint edges, and every edge in E_1 crosses every edge in E_2 . In [1], Ackerman and the authors showed that every n-vertex simple topological graph with no natural k-grid has at most $O(n \log^{4k-6} n)$ edges. As a corollary to Theorem 7.1, noting that the balanced complete bipartite graph with k vertices in each part is a cograph with depth $\lceil \log_2 k \rceil + 1$, we have the following.

Theorem 7.2. If G = (V, E) is an *n*-vertex simple topological graph with no natural k-grid, then $|E(G)| \leq 3n(\log n)^{O(\log k)}$.

7.1 Further remarks on properties of hereditary families of graphs

In Section 2, we proved several properties of hereditary families of graphs are equivalent. Here we extend on these equivalences with some natural variants of these properties.

It is natural to drop the assumption that the vertex subsets have the same size in the definition of the mighty Erdős-Hajnal property. So we say a family \mathcal{F} of graphs has the unbalanced mighty Erdős-Hajnal property if there is a constant c > 0 such that for every graph $G \in \mathcal{F}$ and for all disjoint vertex subsets A and B of G, there are subsets $A' \subset A$ and $B' \subset B$ with $|A'| \ge c|A|$ and $|B'| \ge c|B|$ and the bipartite graph between A' and B' is complete or empty. Trivially, if \mathcal{F} has the unbalanced mighty Erdős-Hajnal property, then it also has the mighty Erdős-Hajnal property. The following proposition shows that something close to a converse also holds.

Lemma 7.3. If a hereditary family \mathcal{F} of graphs has the mighty Erdős-Hajnal property and is closed under cloning vertices, then \mathcal{F} also has the unbalanced mighty Erdős-Hajnal property.

Proof. Since \mathcal{F} has the mighty Erdős-Hajnal property, there is a constant c > 0 such that if $G \in \mathcal{F}$ and A, B are disjoint vertex subsets of G of equal size, then there are $A' \subset A$ and $B' \subset B$ each of size at least c|A| that are complete or empty to each other.

Let $G \in \mathcal{F}$ and A, B be disjoint vertex subsets of G. Every vertex in A we clone |B| times and every vertex in B we clone |A| times to get another graph $G' \in \mathcal{F}$. Let A^* be the vertex subset of G' of clones of vertices in A, and B^* be the vertex subset of G' of clones of vertices in B, so $|A^*| = |B^*| = |A||B|$. Applying the mighty Erdős-Hajnal property to G' and vertex subsets A^* and B^* , we get subsets $A' \subset A^*$ and $B^* \subset B$ that are complete or empty to each other and $|A'|, |B'| \ge c|A||B|$. Let $A'' \subset A$ be those vertices that contain which has at least one clone in A^* and $B'' \subset B$ be those vertices that contain which has at least one clone in B^* . Then $|A''| \ge c|A|$, $|B''| \ge c|B|$, and A'' and B'' are complete or empty to each other. So \mathcal{F} also has the unbalanced mighty Erdős-Hajnal property.

Noting that by drawing another pseudo-segment very close to an original pseudo-segment, we see that the family of intersection graphs of pseudo-segments is closed under cloning. So we have the following corollary of Lemma 7.3 and Theorem 2.8.

Corollary 7.4. The family of intersection graphs of pseudo-segments has the unbalanced mighty Erdős-Hajnal property.

One can similarly define an unbalanced homogeneous density property. A family \mathcal{F} of graphs has the unbalanced homogeneous density property if there is $C = C(\varepsilon)$ such that for every $G \in \mathcal{F}$ and every pair $A, B \subset V$ of disjoint vertex subsets of G with at least $\varepsilon |A||B|$ edges, there are $A' \subset A$ and $B' \subset B$ with $|A'| \ge \varepsilon^C |A|$ and $|B'| \ge \varepsilon^C |B|$ that are complete between them. Trivially, if a family of graphs has the unbalanced homogeneous density property, then it also has the homogeneous density property. In the other direction, just as in the proof of Lemma 7.3, if a hereditary family of graphs has the homogeneous density property and is closed under cloning vertices, then it also has the unbalanced homogeneous density property. In particular, the family of intersection graphs of pseudo-segments has the unbalanced homogeneous density property.

In the definition of the homogeneous density property, we require a polynomial dependence on the density ε . It turns out that for a hereditary family \mathcal{F} and the family $\overline{\mathcal{F}}$ of complements of graphs in \mathcal{F} , having this property is equivalent to the seemingly weaker property that the there is just some dependence on the density ε . We formalize this now. A family \mathcal{F} of graphs has the *weak homogeneous density property* if

• there is a function $\delta : (0,1] \to (0,1]$ such that for every $G \in \mathcal{F}$ and every pair $A, B \subset V$ of disjoint vertex subsets of G of the same size with at least $\varepsilon |A||B|$ edges between them, there are $A' \subset A$ and $B' \subset B$ with $|A'| \ge \delta(\varepsilon)|A|$ and $|B'| \ge \delta(\varepsilon)|B|$ that are complete between them.

We can add to the list of equivalent properties in Theorem 2.2 that \mathcal{F} and $\overline{\mathcal{F}}$ have the *weak* homogeneous density property.

Theorem 7.5. A hereditary family of \mathcal{F} of graphs satisfies \mathcal{F} and $\overline{\mathcal{F}}$ have the homogeneous density property if and only if \mathcal{F} and $\overline{\mathcal{F}}$ have the weak homogeneous density property.

The "only if" direction of Theorem 7.5 is trivial. To see the "if" direction, observe that the proof of Lemma 2.5 that \mathcal{F} and $\overline{\mathcal{F}}$ have the homogeneous density property implies that \mathcal{F} has the mighty Erdős-Hajnal property extends to show that \mathcal{F} and $\overline{\mathcal{F}}$ have the weak homogeneous density property

implies that \mathcal{F} has the mighty Erdős-Hajnal property. Further, Lemma 2.6 implies hereditary \mathcal{F} has the mighty Erdős-Hajnal property implies that \mathcal{F} and $\overline{\mathcal{F}}$ have the homogeneous density property. Thus, the "if" direction holds as well.

References

- E. Ackerman, J. Fox, J. Pach, and A. Suk, On grids in topological graphs, Comput. Geom. 47 (2014), 710–723.
- [2] O. Aichholzer, A. García, J. Tejel, B. Vogtenhuber, and A. Weinberger, Twisted ways to find plane structures in simple drawings of complete graphs, In *Proc. 38th Symp. Comput. Geometry*, LIPIcs, Dagstuhl, Germany, 2022, pages 5:1–5:18.
- [3] N. Alon, J. Pach, R. Pinchasi, R. Radoičić, and M. Sharir, Crossing patterns of semi-algebraic sets, J. Comb. Theory Ser. A 111 (2005), 310–326.
- [4] M. Bucić, T. Nguyen, A. Scott, and P. Seymour, Induced subgraph density. I. A loglog step towards Erdős-Hajnal, preprint arXiv:2301.10147 (2023).
- [5] G. Cairns and Y. Nikolayevsky, Bounds for generalized thrackles, *Discrete Comput. Geom.* 23 (2000), 191–206.
- [6] M. Campos, S. Griffiths, R. Morris, and J. Sahasrabudhe, An exponential improvement for diagonal Ramsey, arXiv preprint arXiv:2303.09521, 2023.
- [7] M. Chudnovsky, The Erdős–Hajnal Conjecture–A Survey, J. Graph Theory 75 (2014), 178– 190.
- [8] P. Erdős, Some remarks on the theory of graphs, Bull. Amer. Math. Soc. 53 (1947), 292–294.
- [9] P. Erdős and A. Hajnal, Ramsey-type theorems, Discrete Appl. Math. 25 (1989), 37–52.
- [10] P. Erdős and G. Szekeres, A combinatorial problem in geometry, Compositio Mathematica 2 (1935), 463–470.
- [11] J. Fox, T. Nguyen, A. Scott, and P. Seymour, Induced subgraph density. II. Sparse and dense sets in cographs, preprint arXiv:2307.00801 (2023).
- [12] J. Fox and J. Pach, Separator theorems and Turán-type results for planar intersection graphs, Adv. Math. 219 (2008), 1070–1080.
- [13] J. Fox, J. Pach, A Separator Theorem for String Graphs and its Applications, Combin. Probab. Comput. (2010) 19, 371–390.
- [14] J. Fox and J. Pach, Erdős-Hajnal-type results on intersection patterns of geometric objects, G.O.H. Katona, et al. (Eds.), Horizon of Combinatorics, Bol. Soc. Stud. Math., Springer (2008), pp. 79–103.
- [15] J. Fox, J. Pach, and A. Suk, A polynomial regularity lemma for semialgebraic hypergraphs and its applications in geometry and property testing, SIAM J. Comput. 45 (2016), 2199–2223.

- [16] J. Fox, J. Pach, and A. Suk, Erdős-Hajnal conjecture for graphs with bounded VC-dimension, Disc. Comput. Geom. 61 (2019), 809–829.
- [17] J. Fox, J. Pach, and A. Suk, Enumeration of intersection graphs of x-monotone curves, in preparation.
- [18] J. Fox, J. Pach, and C. D. Tóth, Turán-type results for partial orders and intersection graphs of convex sets, *Israel J. Math.* **178** (2010), 29–50.
- [19] J. Fox, J. Pach, and C. D. Tóth, Intersection patterns of curves, J. Lond. Math. Soc. 83 (2011), 389–406.
- [20] J. Fox and B. Sudakov, Induced Ramsey-type theorems, Adv. Math. 219 (2008), 1771–1800.
- [21] J. Fox and B. Sudakov, Density theorems for bipartite graphs and related Ramsey-type results, Combinatorica 29 (2009), 153–196.
- [22] R. Fulek and J. Pach, A computational approach to Conway's thrackle conjecture, Comput. Geom. 44 (2011), 345–355.
- [23] R. Fulek and J. Pach, Thrackles: An improved upper bound, Discrete Appl. Math. 259 (2019), 226–231.
- [24] J.E. Goodman. Proof of a conjecture of Burr, Grünbaum, and Sloane, Discrete Math. 32 (1980), 27–35.
- [25] J.E. Goodman and R. Pollack, Semispaces of configurations, cell complexes of arrangements. J. Combin. Theory, Ser. A 37 (1984), 257–293.
- [26] S. Har-Peled, Constructing planar cuttings in theory and practice, SIAM J. Comput. 29 (2000), 2016–2039
- [27] J. Komlós and M. Simonovits, Szemerédi's regularity lemma and its applications in graph theory. In: *Combinatorics, Paul Erdős is Eighty, Vol. 2. Bolyai Soc. Math. Studies*, Bolyai Math. Soc., 2002, 84–112.
- [28] J. R. Lee, Separators in region intersection graphs, in: 8th Innovations in Theoretical Comp. Sci. Conf. (ITCS 2017), LIPIcs 67 (2017), 1–8.
- [29] L. Lovász, J. Pach, and M. Szegedy, On Conway's thrackle conjecture, Discrete Comput. Geom. 18 (1997), 369–376.
- [30] M. Malliaris and S. Shelah, Regularity lemma for stable graphs, Trans. Amer. Math. Soc. 366 (2014), 1551–1585.
- [31] J. Matoušek, Lectures on Discrete Geometry. Springer-Verlag New York, Inc., 2002.
- [32] T. Nguyen, A. Scott, and P. Seymour, Induced subgraph density. III. The pentagon and the bull, preprint arXiv:2307.06379 (2023).
- [33] J. Pach, A Tverberg-type result on multicolored simplices, Comput. Geom.: Theor. Appl. 10, (1998), 71–76.

- [34] J. Pach, F. Shahrokhi, M. Szegedy, Applications of the crossing number, Algorithmica 16 (1996), 111–117.
- [35] J. Pach and J. Solymosi, Structure theorems for systems of segments, in: Discrete and computational geometry (Tokyo 2000), Lect. Notes Comput. Sci. 2098, Springer-Verlag, Berlin (2001), 308–317.
- [36] J. Pach and J. Solymosi, Crossing patterns of segments. J. Comb. Theory Ser. A 96 (2001), 316–325.
- [37] J. Pach and G. Tóth, Which crossing number is it anyway?, J. Comb. Theory Ser. B 80 (2000), 225–246.
- [38] J. Pach and G. Tóth, Disjoint edges in topological graphs. In Proceedings of the 2003 Indonesia-Japan joint conference on Combinatorial Geometry and Graph Theory (IJCCGGT'03), Jin Akiyama, Edy Tri Baskoro, and Mikio Kano (Eds.). Springer-Verlag, Berlin, Heidelberg, 133– 140.
- [39] J. Pach and G. Tóth, How many ways can one draw a graph?, *Combinatorica* **26** (2006), 559–576.
- [40] J. Pach and G. Tóth, Comment on Fox news, Geombinatorics 15 (2006), 150–154.
- [41] V. Rödl, On universality of graphs with uniformly distributed edges, Discrete Math. 59 (1986), 125–134.
- [42] L. Sauermann, On the speed of algebraically defined graph classes, Adv. Math. 380 (2021), Paper No. 107593, 55 pp.
- [43] I. Tomon, String graphs have the Erdős–Hajnal property. J. Eur. Math. Soc. (2023).