

ON THE MOSER'S BERNSTEIN THEOREM

JOSÉ EDSON SAMPAIO AND EURÍPEDES CARVALHO DA SILVA

ABSTRACT. In this paper, we prove the following version of the famous Bernstein's theorem: Let $X \subset \mathbb{R}^{n+k}$ be a closed and connected set with Hausdorff dimension n . Assume that X satisfies the monotonicity formula at $p \in X$. Then, the following statements are equivalent:

- (1) X is an affine linear subspace;
- (2) X is a definable set that is Lipschitz regular at infinity and its geometric tangent cone at infinity, $C(X, \infty)$, is a linear subspace;
- (3) X is a definable set, blow-spherical regular at infinity and $C(X, \infty)$ is a linear subspace;
- (4) X is a definable set that is Lipschitz normally embedded at infinity and $C(X, \infty)$ is a linear subspace;
- (5) the density of X at infinity is 1.

Consequently, we prove the following generalization of Bernstein's theorem: Let $X \subset \mathbb{R}^{n+1}$ be a closed and connected set with Hausdorff dimension n . Assume that X satisfies the monotonicity formula at $p \in X$ and there are compact sets $K \subset \mathbb{R}^n$ and $\tilde{K} \subset \mathbb{R}^{n+1}$ such that $X \setminus \tilde{K}$ is a minimal hypersurface that is the graph of a C^2 -smooth function $u: \mathbb{R}^n \setminus K \rightarrow \mathbb{R}$. Assume that u has bounded derivative whenever $n > 7$. Then X is a hyperplane. Several other results are also presented. For example, we generalize the o-minimal Chow's theorem, we prove that any entire complex analytic set that is bi-Lipschitz homeomorphic to a definable set in an o-minimal structure must be an algebraic set. We also obtain that Yau's Bernstein Problem, which says that an oriented stable complete minimal hypersurface in \mathbb{R}^{n+1} with $n \leq 6$ must be a hyperplane, holds true whether the hypersurface is a definable set in an o-minimal structure.

CONTENTS

1. Introduction 2

2010 *Mathematics Subject Classification*. 53A10; 53A07 (primary); 14J17; 53C42; 14P10 (secondary).

Key words and phrases. Bernstein Theorem, Moser's Bernstein Theorem, Minimal surfaces, monotonicity formula, Lipschitz regularity.

The first named author was partially supported by CNPq-Brazil grant 310438/2021-7. This work was supported by the Serrapilheira Institute (grant number Serra – R-2110-39576).

2. Preliminaries	5
2.1. O-minimal structures	6
2.2. Dimension of definable sets	6
2.3. Lipschitz regularity at infinity	7
2.4. Lipschitz normal embedding at infinity	7
2.5. Tangent cones	8
2.6. Relative multiplicities at infinity	9
2.7. Blow-spherical invariance of the relative multiplicities at infinity	9
2.8. Bi-Lipschitz invariance of the relative multiplicities at infinity	10
2.9. Density at infinity	12
3. Kurdyka-Raby's formula at infinity	13
3.1. Proof of the Kurdyka-Raby's formula at infinity	13
3.2. Some examples	19
3.3. First consequences	20
4. Parametric versions of the Bernstein Theorem	22
4.1. Some examples	24
4.2. Some direct consequences	25
5. Generalization of the Moser's Bernstein Theorem	27
5.1. Examples	29
5.2. Consequences	29
References	31

1. INTRODUCTION

We start this article by reminding the famous Bernstein's problem:

Bernstein's problem. *If the graph of a function on \mathbb{R}^n is a minimal hypersurface in \mathbb{R}^{n+1} , does this imply that the function is linear?*

The answer is negative when n is at least 8 (see [10]), but the answer was proved to be positive the cases $n = 2$ by Bernstein (see [4]), $n = 3$ by DeGiorgi (see [13]), $n = 4$ by Almgren (see [2]) and $n \leq 7$ by Simons (see [50]). The positive answers can be summarized as follows:

Theorem 1.1. *Let $M \subset \mathbb{R}^{n+1}$ be a minimal hypersurface which is a graph of C^2 -smooth function $u: \mathbb{R}^n \rightarrow \mathbb{R}$ with $2 \leq n \leq 7$. Then u is a linear function and M must be a hyperplane.*

Furthermore, the celebrated theorem due to J. Moser in [31], called Moser's Bernstein Theorem, says the following:

Theorem 1.2 (Moser's Bernstein Theorem). *Let $M \subset \mathbb{R}^{n+1}$ be a minimal hypersurface which is a graph of an entire Lipschitz function $u: \mathbb{R}^n \rightarrow \mathbb{R}$. Then u is a linear function and M must be a hyperplane.*

Let us remark that, in general, Moser's Bernstein theorem does not hold in higher codimension. Lawson and Osserman in [28] presented a minimal cone which is the graph of a Lipschitz mapping, but is not an affine linear subspace, more precisely, they presented the following:

Example 1.3 (Theorem 7.1 in [28]). *The graph of the Lipschitz mapping $f: \mathbb{R}^4 \rightarrow \mathbb{R}^3$ given by $f(0) = 0$ and*

$$f(x) = \frac{\sqrt{5}}{2} \|x\| \eta \left(\frac{x}{\|x\|} \right), \quad \forall x \neq 0,$$

is a minimal cone, where $\eta: \mathbb{S}^3 \rightarrow \mathbb{S}^2$ is the Hopf mapping given by

$$\eta(z_1, z_2) = (|z_1|^2 - |z_2|^2, 2z_1\bar{z}_2).$$

Several other mathematicians approached Moser's Bernstein Theorem in higher codimension (for example, see [3], [23], [24], [25] and [54]).

In this direction, we prove the following result:

Theorem 5.1. *Let $X \subset \mathbb{R}^{n+k}$ be a connected closed set with Hausdorff dimension n . Then, X is an n -dimensional affine linear subspace if and only if we have the following:*

- (1) X satisfies the monotonicity formula at some $p \in X$;
- (2) there are compact sets $K \subset \mathbb{R}^n$ and $\tilde{K} \subset \mathbb{R}^{n+k}$ such that $X \setminus \tilde{K}$ is the graph of a C^1 -smooth function $u: \mathbb{R}^n \setminus K \rightarrow \mathbb{R}^k$ with bounded derivative;
- (3) $\mathcal{N}(X, \infty)$ is a linear subspace.

In order to know, for a subset $X \subset \mathbb{R}^{n+k}$, with Hausdorff dimension n and $p \in X$, we say that X **satisfies the monotonicity formula at p** if the function $\theta^n(X, p, \cdot): (0, +\infty) \rightarrow [1, +\infty)$ given by

$$\theta^n(X, p, r) = \frac{\mathcal{H}^n(X \cap B_r^n(p))}{\mu_n r^n}$$

is well-defined and nondecreasing, and satisfies the following expression:

- (i) X is C^1 -smooth at p if and only if $\theta^n(X, p) := \lim_{r \rightarrow 0^+} \theta^n(X, p, r) = 1$;

- (ii) $\theta^n(X, p, \cdot)$ is a constant function if and only if X is a cone with vertex at p ,

where μ_n is the volume of the n -dimensional Euclidean unit ball, $\mathcal{H}^n(A)$ is the n -dimensional Hausdorff measure of A , and $B_r^m(p) \subset \mathbb{R}^m$ is the open Euclidean ball centered at p of radius $r > 0$. Moreover, $\mathcal{N}(X, \infty)$ is the union of all hyperplanes T such that there is a sequence $\{x_j\}_j \subset X \setminus \text{Sing}_1(X)$ such that $\lim \|x_j\| = +\infty$ and $T_{x_j}X$ converges to T , where $\text{Sing}_1(X)$ is the set of points $x \in X$ such that $X \cap U$ is not a C^1 smooth submanifold of \mathbb{R}^{n+k} , for any open neighborhood $U \subset \mathbb{R}^{n+k}$ of x .

Remark 1.4. *If $X \subset \mathbb{R}^{n+k}$ is a complete minimal submanifold or a closed area-minimizing set, then X satisfies the monotonicity formula at any $p \in X$. In particular, any entire-pure dimensional complex analytic set $X \subset \mathbb{C}^n$ satisfies the monotonicity formula at any $p \in X$.*

In Section 5, we present several consequences of Theorem 5.1. Moreover, in subsection 5.1, by presenting several examples, we show that Theorem 5.1 is sharp such that no hypothesis can be dropped.

Another natural question is whether Moser's Bernstein Theorem can be generalized to the parametric case. Thus, the following questions arise:

Question 1. *Given a smooth minimal hypersurface $M \subset \mathbb{R}^{n+1}$ which is bi-Lipschitz homeomorphic to \mathbb{R}^n , is M an affine linear subspace?*

Recently, [18] and [42] approached Question 1 in higher codimension for complex analytic sets. For instance, they proved that a pure dimensional complex analytic set that is Lipschitz regular at infinity (see Definition 2.6) must be an affine linear subset.

In this article, we use the concepts of [18] and [42] to provide partial positive answers to Question 1, even for high codimensions. We prove the following:

Theorem 4.1. *Let $X \subset \mathbb{R}^{n+k}$ be a closed and connected set with $n = \dim_H X$. Assume that X satisfies the monotonicity formula at some $p \in X$. Then the following statements are equivalent:*

- (1) X is an affine linear subspace;
- (2) X is a definable set, blow-spherical regular at infinity and $C(X, \infty)$ is a linear subspace;
- (3) X is a definable set that is Lipschitz regular at infinity and $C(X, \infty)$ is a linear subspace;

(4) X is an LNE at infinity definable set and $C(X, \infty)$ is a linear subspace;

$$(5) \theta^n(X) := \lim_{r \rightarrow +\infty} \frac{\mathcal{H}^n(X \cap B_r^n(0))}{\mu_n r^n} = 1.$$

We obtain several consequences, for instance, we recover the results proven in [18] and [45].

The main tool to prove Theorem 4.1 is the Kurdyka-Raby's formula at infinity.

Theorem 3.1. (Kurdyka-Raby's formula at infinity). *Let $X \subset \mathbb{R}^{n+k}$ be a definable set in an o-minimal structure on \mathbb{R} with $n = \dim X$. Let C_1, \dots, C_m be the connected components of $C_{\text{Smp}}(X, \infty)$. Then, $\theta^n(X)$ is defined; moreover for each $x \in \mathbb{R}^{n+k}$,*

$$\theta^n(X) = \lim_{r \rightarrow +\infty} \frac{\mathcal{H}^n(X \cap B_r^{n+k}(x))}{\mu_n r^n} = \sum_{j=1}^m k_{X, \infty}(C_j) \frac{\mathcal{H}^n(C_j \cap B_1^{n+k})}{\mu_n}.$$

We prove this formula in Section 3. Furthermore, we present some consequences of Theorem 3.1. For instance, we obtain the o-minimal Chow's theorem proved in [35, Corollary 4.5] as follows:

Corollary 3.9 (O-minimal Chow's theorem). *Let $X \subset \mathbb{C}^{n+k}$ be a pure-dimensional entire complex analytic set with $\dim_{\mathbb{C}} X = n$. Then, X is a complex algebraic set if and only if X is definable in an o-minimal structure on \mathbb{R} .*

We obtain the following generalization of the o-minimal Chow's theorem:

Corollary 3.10 (Lipschitz o-minimal Chow's theorem). *Let $X \subset \mathbb{C}^{n+k}$ be a pure-dimensional entire complex analytic set with $\dim_{\mathbb{C}} X = n$. Then, X is a complex algebraic set if and only if X is lipeomorphic at infinity to a definable set in an o-minimal structure on \mathbb{R} .*

We obtain also that the Yau's Bernstein Problem (see Problem 102 in [55]), which says that *an oriented stable complete minimal hypersurface in \mathbb{R}^{n+1} with $n \leq 6$ must be a hyperplane*, holds true whether the hypersurface is a definable set in an o-minimal structure on \mathbb{R} (see Corollary 3.11).

2. PRELIMINARIES

All the sets herein are assumed to be equipped with the induced Euclidean metric.

2.1. O-minimal structures.

Definition 2.1. A structure on \mathbb{R} is a collection $\mathcal{S} = \{\mathcal{S}_n\}_{n \in \mathbb{N}}$ where each \mathcal{S}_n is a set of subsets of \mathbb{R}^n , satisfying the following axioms:

- 1) All algebraic subsets of \mathbb{R}^n are in \mathcal{S}_n ;
- 2) For every n , \mathcal{S}_n is a Boolean subalgebra of the powerset of \mathbb{R}^n ;
- 3) If $A \in \mathcal{S}_m$ and $B \in \mathcal{S}_n$, then $A \times B \in \mathcal{S}_{m+n}$.
- 4) If $\pi: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ is the projection on the first n coordinates and $A \in \mathcal{S}_{n+1}$, then $\pi(A) \in \mathcal{S}_n$.

The structure \mathcal{S} is said to be **o-minimal** if it satisfies the following condition:

- 5) The elements of \mathcal{S}_1 are precisely finite unions of points and intervals.

A element of \mathcal{S}_n is called **definable in \mathcal{S}** .

Throughout this paper, we fix an o-minimal structure \mathcal{S} on \mathbb{R} .

Definition 2.2. A mapping $f: A \subset \mathbb{R}^n \rightarrow \mathbb{R}^m$ is called **definable in \mathcal{S}** if its graph is an element of \mathcal{S}_{n+m} .

In the sequel, the adjective definable denotes definable in \mathcal{S} .

2.2. Dimension of definable sets.

Definition 2.3. Let $X \subset \mathbb{R}^m$ be a subset and k be a positive integer. The C^k **singular set** of X , denoted by $\text{Sing}_k(X)$, is the set of points $x \in X$ such that $U \cap X$ is not a smooth submanifold of \mathbb{R}^m for any open neighbourhood U of x . A point of $\text{Sing}_k(X)$ is called a C^k **singular point** (or a C^k **singularity**) of X . If $p \in \text{Reg}_k(X) := X \setminus \text{Sing}_k(X)$, we say that X is smooth at p .

Thus, if $p \in \text{Reg}_k(X) = X \setminus \text{Sing}_k(X)$, there is open neighbourhood $U \subset \mathbb{R}^m$ of p such that $X \cap U$ is a C^k smooth submanifold of \mathbb{R}^m and, then, we define the **real dimension of X at p** by $\dim_p X = \dim X \cap U$. Thus, we define the **dimension of X** by

$$\dim X = \max_{p \in \text{Reg}_1(X)} \dim_p X.$$

We say that X is a **pure-dimensional set**, if $\dim X = \dim_p X$ for all $p \in \text{Reg}_1(X)$.

In the case that $X \subset \mathbb{R}^m$ is a definable set, we have that $\text{Sing}_k(X)$ is also a definable set with $\dim \text{Sing}_k(X) < \dim X$ and $\dim X$ is equal to the Hausdorff dimension of X , denoted here by $\dim_H X$.

In the case that $X \subset \mathbb{C}^m \cong \mathbb{R}^{2m}$ is a complex analytic set, we define the **complex dimension of X** as $\dim_{\mathbb{C}} X = \frac{1}{2} \dim X$.

2.3. Lipschitz regularity at infinity.

Definition 2.4. Let $X \subset \mathbb{R}^n$ and $Y \subset \mathbb{R}^m$ be two subsets. A mapping $\phi: X \rightarrow Y$ is **Lipschitz** if there is a constant $C \geq 0$ such that

$$\|\phi(x) - \phi(y)\| \leq C\|x - y\|, \quad \forall x, y \in X.$$

In this case, we also say that ϕ is C -Lipschitz. Moreover, a mapping $\phi: X \rightarrow Y$ is a **lipeomorphism** if ϕ is a homeomorphism such that ϕ and ϕ^{-1} are Lipschitz. When there is a lipeomorphism $\phi: X \rightarrow Y$, we say that X and Y are **lipeomorphic**.

Definition 2.5. Let $X \subset \mathbb{R}^n$ and $Y \subset \mathbb{R}^m$ be two subsets. We say that X and Y are **lipeomorphic at infinity**, if there exist compact subsets $K \subset \mathbb{R}^n$ and $\tilde{K} \subset \mathbb{R}^m$ such that $X \setminus K$ and $Y \setminus \tilde{K}$ are lipeomorphic.

Definition 2.6. A subset $X \subset \mathbb{R}^m$ is called **Lipschitz regular at infinity** if X and \mathbb{R}^n are lipeomorphic at infinity for some $n \in \mathbb{N}$.

2.4. Lipschitz normal embedding at infinity. Let us recall the definition of the inner distance. Given a path connected subset $X \subset \mathbb{R}^m$ the *inner distance* on X is defined as follows: given two points $x_1, x_2 \in X$, $d_X(x_1, x_2)$ is the infimum of the lengths of paths on X connecting x_1 to x_2 . As stated in Section 2, all the sets considered herein are supposed to be equipped with the induced Euclidean metric. Whenever we consider the inner distance, we emphasize it clearly.

Definition 2.7 (See [6]). A subset $X \subset \mathbb{R}^m$ is called **Lipschitz normally embedded** (or shortly **LNE**) if there exists $C > 0$ such that

$$d_X(x_1, x_2) \leq C\|x_1 - x_2\|$$

for all $x_1, x_2 \in X$. In this case, we also say that X is **C -LNE**.

In [18], the following definition was presented:

Definition 2.8. A subset $X \subset \mathbb{R}^n$ is **Lipschitz normally embedded at infinity** (or shortly **LNE at infinity**) if there exists a compact subset $K \subset \mathbb{R}^n$ such that $X \setminus K$ is Lipschitz normally embedded.

2.5. Tangent cones.

Definition 2.9. Let $X \subset \mathbb{R}^m$ be an unbounded set (resp. $p \in \overline{X}$ be a non-isolated point). We say that $v \in \mathbb{R}^m$ is a tangent vector of X at infinity (resp. p) if there are a sequence of points $\{x_i\} \subset X$ tending to infinity (resp. p) and a sequence of positive real numbers $\{t_i\}$ such that

$$\lim_{i \rightarrow \infty} \frac{1}{t_i} x_i = v \quad (\text{resp. } \lim_{i \rightarrow \infty} \frac{1}{t_i} (x_i - p) = v).$$

Let $C(X, \infty)$ (resp. $C(X, p)$) denote the set of all tangent vectors of X at infinity (resp. p). We call $C(X, \infty)$ the **geometric tangent cone of X at infinity** (resp. p).

Another way to present the geometric tangent cone at infinity (resp. p) of a subset $X \subset \mathbb{R}^m$ is via the spherical blow-up at infinity (resp. p) of \mathbb{R}^m . Let us consider the **spherical blowing-up at infinity** (resp. p) of \mathbb{R}^m , $\rho_\infty: \mathbb{S}^{m-1} \times (0, +\infty) \rightarrow \mathbb{R}^m$ (resp. $\rho_p: \mathbb{S}^{m-1} \times [0, +\infty) \rightarrow \mathbb{R}^m$), given by $\rho_\infty(x, r) = \frac{1}{r}x$ (resp. $\rho_p(x, r) = rx + p$).

Note that $\rho_\infty: \mathbb{S}^{m-1} \times (0, +\infty) \rightarrow \mathbb{R}^m \setminus \{0\}$ (resp. $\rho_p: \mathbb{S}^{m-1} \times (0, +\infty) \rightarrow \mathbb{R}^m \setminus \{0\}$) is a homeomorphism with inverse mapping $\rho_\infty^{-1}: \mathbb{R}^m \setminus \{0\} \rightarrow \mathbb{S}^{m-1} \times (0, +\infty)$ (resp. $\rho_p^{-1}: \mathbb{R}^m \setminus \{0\} \rightarrow \mathbb{S}^{m-1} \times (0, +\infty)$) given by $\rho_\infty^{-1}(x) = (\frac{x}{\|x\|}, \frac{1}{\|x\|})$ (resp. $\rho_p^{-1}(x) = (\frac{x-p}{\|x-p\|}, \|x-p\|)$).

The **strict transform** of the subset X under the spherical blowing-up ρ_∞ (resp. ρ_p) is $X'_\infty := \overline{\rho_\infty^{-1}(X \setminus \{0\})}$ (resp. $X'_p := \overline{\rho_p^{-1}(X \setminus \{0\})}$). The subset $X'_\infty \cap (\mathbb{S}^{m-1} \times \{0\})$ (resp. $X'_p \cap (\mathbb{S}^{m-1} \times \{0\})$) is called the **boundary** of X'_∞ (resp. X'_p) and is denoted as $\partial X'_\infty$ (resp. $\partial X'_p$).

Remark 2.10. If $X \subset \mathbb{R}^m$ is a definable set in an o-minimal structure, then $\partial X'_\infty = (C(X, \infty) \cap \mathbb{S}^{m-1}) \times \{0\}$ (resp. $\partial X'_p = (C(X, p) \cap \mathbb{S}^{m-1}) \times \{0\}$).

We finish this Subsection by reminding the invariance of the tangent cone at infinity under lipeomorphisms at infinity obtained in the paper [44] (see also [18, Theorem 2.19]).

Theorem 2.11 (Corollary 2.11 in [44]). Let $X \subset \mathbb{R}^m$ and $Y \subset \mathbb{R}^k$ be unbounded definable sets with $n = \dim X = \dim Y$. If there exist compact subsets $K \subset \mathbb{R}^m$ and $\tilde{K} \subset \mathbb{R}^k$, and a lipeomorphism $\varphi: X \setminus K \rightarrow Y \setminus \tilde{K}$, then there is a globally defined lipeomorphism $d\varphi: C(X, \infty) \rightarrow C(Y, \infty)$ with $d\varphi(0) = 0$.

2.6. Relative multiplicities at infinity. Let $X \subset \mathbb{R}^m$ be an n -dimensional definable subset in an o-minimal structure and $p \in \mathbb{R}^m \cup \{\infty\}$. We say that $x \in \partial X'_p$ is **simple point of $\partial X'_p$** , if there is an open subset $U \subset \mathbb{R}^{m+1}$ with $x \in U$ such that:

- a) the connected components X_1, \dots, X_r of $(X'_p \cap U) \setminus \partial X'_p$ are topological submanifolds of \mathbb{R}^{m+1} with $\dim X_i = \dim X$, for all $i = 1, \dots, r$;
- b) $(X_i \cup \partial X'_p) \cap U$ are topological manifolds with boundary $\partial X'_p \cap U$, for all $i = 1, \dots, r$.

Let $\text{Smp}(\partial X'_p)$ be the set of simple points of $\partial X'_p$ and we define $C_{\text{Smp}}(X, p) = \{t \cdot x; t > 0 \text{ and } (x, 0) \in \text{Smp}(\partial X'_p)\}$. Let $k_{X,p}: C_{\text{Smp}}(X, p) \rightarrow \mathbb{N}$ be the function such that $k_{X,p}(x)$ is the number of connected components of the germ $(\rho^{-1}(X \setminus \{p\}), v)$, where $v = (\frac{x}{\|x\|}, 0)$.

Remark 2.12. $\text{Smp}(\partial X'_p)$ is an open dense subset of the $(n-1)$ -dimensional part of $\partial X'_p$ whenever $\partial X'_p$ is an $(n-1)$ -dimensional subset, where $n = \dim X$.

Definition 2.13. It is clear that the function $k_{X,p}$ is locally constant. In fact, $k_{X,p}$ is constant for each connected component X_j of $C_{\text{Smp}}(X, p)$. We define **the relative multiplicity at p of X (along of X_j)** as $k_{X,p}(X_j) := k_{X,p}(x)$ where $x \in X_j$.

Definition 2.14. Let $X \subset \mathbb{R}^m$ and $Y \subset \mathbb{R}^k$ be closed sets. Let $p \in \mathbb{R}^m \cup \{\infty\}$, $q \in \mathbb{R}^k \cup \{\infty\}$ and a homeomorphism $\varphi: X \rightarrow Y$ such that $q = \lim_{x \rightarrow p} \varphi(x)$, is said a **blow-spherical homeomorphism at p** , if the homeomorphism

$$\rho_q^{-1} \circ \varphi \circ \rho_p: \rho_p^{-1}(X \setminus \{p\}) \rightarrow \rho_q^{-1}(Y \setminus \{q\})$$

extends to a homeomorphism $\varphi': X'_p \rightarrow Y'_q$.

2.7. Blow-spherical invariance of the relative multiplicities at infinity. The following result was presented in [45, Proposition 3.5] for semi-algebraic sets, but the proof is the same for definable sets in an o-minimal structure as we can see in [38, Teorema 3.1.7].

Proposition 2.15. Let $X \subset \mathbb{R}^m$ and $Y \subset \mathbb{R}^k$ be definable subsets in an o-minimal structure on \mathbb{R} . Let $\varphi: X \rightarrow Y$ be a blow-spherical homeomorphism at $p \in \mathbb{R}^m \cup \{\infty\}$. Then $\varphi'(\text{Smp}(\partial X'_p)) = \text{Smp}(\partial Y'_q)$ and

$$k_{X,p}(x) = k_{Y,q}(\varphi'(x)),$$

for all $x \in \text{Smp}(\partial X'_p)$, where $q = \lim_{x \rightarrow p} \varphi(x)$.

2.8. Bi-Lipschitz invariance of the relative multiplicities at infinity.

Theorem 2.16. *Let $X \subset \mathbb{R}^m$ and $Y \subset \mathbb{R}^k$ be unbounded definable sets with $n = \dim X = \dim Y$. If there exist compact subsets $K \subset \mathbb{R}^m$ and $\tilde{K} \subset \mathbb{R}^k$, and a lipeomorphism $\varphi: X \setminus K \rightarrow Y \setminus \tilde{K}$, then there exists a lipeomorphism $d\varphi: C(X, \infty) \rightarrow C(Y, \infty)$ that satisfies*

$$k_{X, \infty}(x) = k_{Y, \infty}(d\varphi(x)), \quad \forall x \in C_{\text{Smp}}(X, \infty) \cap d\varphi^{-1}(C_{\text{Smp}}(Y, \infty)).$$

In particular, $d\varphi|_{C_{\text{Smp}}(X, \infty)}: C_{\text{Smp}}(X, \infty) \rightarrow C_{\text{Smp}}(Y, \infty)$ is a lipeomorphism where $C_{\text{Smp}}(X, \infty) \neq \emptyset$.

Proof. We closely follow the proof of Theorem 4.2 in [41], but because its importance in this article, we present the proof here.

By making identifications $X \setminus K \longleftrightarrow X \setminus K \times \{0\}$ and $Y \setminus \tilde{K} \longleftrightarrow \{0\} \times Y \setminus \tilde{K}$, we can assume that $m = k$ and φ is globally defined in \mathbb{R}^m . We know that there exist a sequence of positive real numbers $S = \{t_j\}_{j \in \mathbb{N}}$ and a lipeomorphism $d\varphi: C(X, \infty) \rightarrow C(Y, \infty)$ such that

$$t_j \rightarrow +\infty \quad \text{and} \quad \frac{\varphi(t_j v)}{t_j} \rightarrow d\varphi(v), \quad \forall v \in C(X, \infty).$$

For more details, see [39], [18], [19] and [44].

Thus, $\rho^{-1} \circ \varphi \circ \rho: SX \rightarrow Y'$ is an injective and continuous mapping that continuously extends to a mapping $\varphi': \overline{SX} \rightarrow Y'$, where $\rho = \rho_\infty$ and

$$SX = \{(x, s) \in \mathbb{S}^{m-1} \times (0, +\infty); \frac{1}{s} \cdot x \in X \text{ and } \frac{1}{s} \in S\}.$$

We note that $\text{Smp}(\partial X'_\infty) = \emptyset$ (resp. $\text{Smp}(\partial Y'_\infty) = \emptyset$) if and only if $\dim C(X, \infty) < \dim X$ (resp. $\dim C(Y, \infty) < \dim Y$). Since by Theorem 2.11, $\dim C(X, \infty) = \dim C(Y, \infty)$, then we obtain that $\text{Smp}(\partial X'_\infty) = \emptyset$ if and only if $\text{Smp}(\partial Y'_\infty) = \emptyset$. However, when $\text{Smp}(\partial X'_\infty) = \text{Smp}(\partial Y'_\infty) = \emptyset$, we have nothing to do. Thus, we can assume that $\text{Smp}(\partial X'_\infty) \neq \emptyset$ and, thus, $\text{Smp}(\partial Y'_\infty) \neq \emptyset$ as well. Further, $C_{\text{Smp}}(X, \infty)$ (resp. $C_{\text{Smp}}(Y, \infty)$) is a dense subset in the n -dimensional part of $C(X, \infty)$ (resp. $C(Y, \infty)$). Therefore, $d\varphi(C_{\text{Smp}}(X, \infty)) \cap C_{\text{Smp}}(Y, \infty)$ is a dense subset in the n -dimensional part of $C(Y, \infty)$ and $C_{\text{Smp}}(Y, \infty)$.

Let X_1, \dots, X_r be the connected components of $C_{\text{Smp}}(X, \infty)$ and let Y_1, \dots, Y_s be the connected components of $C_{\text{Smp}}(Y, \infty)$. For each point $x \in X_j$, we know that $k_{X, \infty}(X_j) = k_{X, \infty}(x)$ is the number of connected components of the germ $(\rho_\infty^{-1}(X) \cap B_\delta^{m+1}(x), x)$. Then, $k_{X, \infty}(x)$ can be seen as the number of connected components of the set $(SX \cap \mathbb{S}^{m-1} \times \{t_j\}) \cap B_\delta^{m+1}(x)$ for all sufficiently large k .

Let $\pi: \mathbb{R}^m \rightarrow \mathbb{R}^n$ be a linear projection such that

$$\pi^{-1}(0) \cap (C(X, \infty) \cup C(Y, \infty)) = \{0\}.$$

Let ramification loci of

$$\pi|_X: X \rightarrow \mathbb{R}^n \quad \text{and} \quad \pi|_{C(X, \infty)}: C(X, \infty) \rightarrow \mathbb{R}^n$$

be $\sigma(X)$ and $\sigma(C(X, \infty))$ respectively. Similarly, we define $\sigma(Y)$ and $\sigma(C(Y, \infty))$.

Let $Z = \sigma(X) \cup \sigma(C(X, \infty))$, $W = \sigma(Y) \cup \sigma(C(Y, \infty))$, and

$$\Sigma = Z \cup C(Z, \infty) \cup \pi(d\varphi^{-1}(\pi|_{Y \cup C(Y, \infty)}^{-1}(W \cup C(W, \infty))).$$

Since $\pi|_{Y \cup C(Y, \infty)}$ is a proper mapping, $d\varphi$ is a lipeomorphism, and $\dim Z \cup W \leq n-1$, we obtain that $\dim \Sigma \leq n-1$. Thus, we obtain that if $v' \in \mathbb{R}^n \setminus \Sigma$ then $\pi|_X^{-1}(v') \subset \mathbb{R}^m \setminus \pi|_X^{-1}(Z \cup C(Z, \infty))$ and for any $v_i \in \pi|_X^{-1}(v')$, we have that $w'_i = \pi(d\varphi(v_i)) \in \mathbb{R}^n \setminus (W \cup C(W, \infty))$. For $\eta, R > 0$, we define the following set

$$C_{\eta, R}(v') = \{w \in \mathbb{R}^n \mid \exists t > 0; \|tv' - w\| < \eta t\} \setminus B_R^n(0).$$

Let $\eta, R > 0$ be sufficiently large such that $C_{\eta, R}(v') \subset \mathbb{R}^n \setminus (Z \cup C(Z, \infty))$. Then, we obtain that $\pi|_V: V \rightarrow C_{\eta, R}(v')$ is a lipeomorphism for each connected component V of $\pi^{-1}(C_{\eta, R}(v')) \cap X$. Therefore, for each $j = 1, \dots, r$, there are different connected components $V_{j1}, \dots, V_{jk_X(X_j)}$ of $\pi^{-1}(C_{\eta, R}(v')) \cap X$ such that

$$C(\overline{V_{ji}}, \infty) \setminus \{0\} \subset X_j, \quad \forall i \in \{1, \dots, k_{X, \infty}(X_j)\}.$$

Let $\rho_\infty: \mathbb{S}^{m-1} \times (0, +\infty) \rightarrow \mathbb{R}^m$ be the spherical blow-up at infinity. Fixed $j \in \{1, \dots, r\}$, let us suppose that there is $q \in \{1, \dots, s\}$ such that $d\varphi(X_j) \cap Y_q \neq \emptyset$ and $k_{X, \infty}(X_j) > k_{Y, \infty}(Y_q)$, it means that if we consider a point $x = (v, 0) \in (X_j \cap \mathbb{S}^{m-1}) \times \{0\}$ with $d\varphi(x) \in Y_q$ and, since $\dim \Sigma \leq n-1$, we can assume that $\pi(v) \notin \Sigma$, then there are at least two different connected components V_{ji} and V_{jl} of $\pi^{-1}(C_{\eta, R}(\pi(v))) \cap X$ and sequences $\{(x_j, t_j^{-1})\}_{j \in \mathbb{N}} \subset \rho^{-1}(V_{ji}) \cap SX$ and $\{(y_j, t_j^{-1})\}_{j \in \mathbb{N}} \subset \rho^{-1}(V_{jl}) \cap SX$ such that $\lim(x_j, t_j^{-1}) = \lim(y_j, t_j^{-1}) = x$ and $\varphi'(x_j, t_j^{-1}), \varphi'(y_j, t_j^{-1}) \in \rho^{-1}(\tilde{V}_{q\ell})$ for all $j \in \mathbb{N}$, where $\tilde{V}_{q\ell}$ is a connected component of $\pi^{-1}(C_{\tilde{\eta}, \tilde{R}}(\pi(d\varphi(v)))) \cap Y$ for some $\tilde{\eta}, \tilde{R} > 0$ such that

$$C_{\tilde{\eta}, \tilde{R}}(w') = \{\tilde{w} \in \mathbb{R}^n \mid \exists t > 0; \|tw' - \tilde{w}\| < \tilde{\eta}t\} \setminus B_{\tilde{R}}^n(0)$$

satisfies $C_{\tilde{\eta}, \tilde{R}}(w') \subset \mathbb{R}^n \setminus (W \cup C(W, \infty))$, where $w' = \pi(d\varphi(v))$. In particular, $\pi|_{\tilde{V}}: \tilde{V} \rightarrow C_{\tilde{\eta}, \tilde{R}}(w')$ is a lipeomorphism, where $\tilde{V} = \tilde{V}_{q\ell}$. Since

$\varphi(t_j x_j), \varphi(t_j y_j) \in \tilde{V}$ for all $j \in \mathbb{N}$, we have

$$\|\varphi(t_j x_j) - \varphi(t_j y_j)\| = o_\infty(t_j)$$

and

$$d_Y(\varphi(t_j x_j), \varphi(t_j y_j)) \leq d_{\tilde{V}}(\varphi(t_j x_j), \varphi(t_j y_j)) = o_\infty(t_j),$$

where $g(t_j) = o_\infty(t_j)$ means that $\lim_{j \rightarrow +\infty} \frac{g(t_j)}{t_j} = 0$. Now, since X is lipeomorphic at infinity to Y , we obtain $d_X(t_j x_j, t_j y_j) \leq o_\infty(t_j)$. On the other hand, since $t_j x_j$ and $t_j y_j$ lie in different connected components of $\pi^{-1}(C_{\eta, R}(\pi(v))) \cap X$, there exists a constant $C > 0$ such that $d_X(t_j x_j, t_j y_j) \geq Ct_j$, which is a contradiction.

We have proved that $k_{X, \infty}(X_j) \leq k_{Y, \infty}(Y_q)$. By similar arguments, by using that φ^{-1} is a lipeomorphism, we can also prove $k_{Y, \infty}(Y_q) \leq k_{X, \infty}(X_j)$. Therefore, $k_{X, \infty}(X_j) = k_{Y, \infty}(Y_q)$ for any $q \in \{1, \dots, s\}$ such that $d\varphi(X_j) \cap Y_q \neq \emptyset$. Then

$$k_{X, \infty}(x) = k_{Y, \infty}(d\varphi(x)), \quad \forall x \in C_{\text{Smp}}(X, \infty) \cap d\varphi^{-1}(C_{\text{Smp}}(Y, \infty)).$$

□

Remark 2.17. *Since the inversion mapping $\iota: \mathbb{R}^m \setminus \{0\} \rightarrow \mathbb{R}^m \setminus \{0\}$, given by $\iota(x) = \frac{x}{\|x\|^2}$, is a blow-spherical homeomorphism at infinity, for $Z = \iota(X \setminus \{0\})$, we obtain that $C(Z, 0) = C(X, \infty)$, $C_{\text{Smp}}(X, \infty) = C_{\text{Smp}}(Z, 0)$ and $k_{X, \infty}(v) = k_{Z, 0}(v)$, for all $v \in C_{\text{Smp}}(X, \infty)$, and thus the o -minimal version of [41, Theorem 4.2] follows from Theorem 2.16 and [43, Theorem 4.1].*

2.9. Density at infinity. Let $X \subset \mathbb{R}^m$ be a definable set. Fixed $n \in \mathbb{N}$ and $p \in X$, we define the function $\theta^n(X, p, \cdot): (0, +\infty) \rightarrow X$ as follows:

$$\theta^n(X, p, r) = \frac{\mathcal{H}^n(X \cap B_r^n(p))}{\mu_n r^n},$$

where μ_n is the volume of the n -dimensional Euclidean unit ball, $\mathcal{H}^n(A)$ is the n -dimensional Hausdorff measure of A and $B_r^m(p) \subset \mathbb{R}^m$ is the open Euclidean ball centered at p of radius $r > 0$.

For simplicity, we denote $B_r^m := B_r^m(0)$.

Definition 2.18. *Let $X \subset \mathbb{R}^{n+k}$ be a set. We say that X has n -dimensional density at p and we denote it by $\theta^n(X, p)$, if the limit exists:*

$$\theta^n(X, p) := \lim_{r \rightarrow 0^+} \theta^n(X, p, r).$$

Definition 2.19. Let $X \subset \mathbb{R}^{n+k}$ be a set. We say that X has n -dimensional density at infinity and we denote it by $\theta^n(X)$, if the limit exists:

$$\theta^n(X) := \lim_{r \rightarrow +\infty} \frac{\mathcal{H}^n(X \cap B_r^{n+k})}{\mu_n r^n}.$$

When X has n -dimensional density at infinity and $n = \dim X$, we say that X has density at infinity.

3. KURDYKA-RABY'S FORMULA AT INFINITY

The goal of this section is to bring the Theorem of Kurdyka and Raby in [27] and its ideas to a global perspective. Then, we prove the following Kurdyka-Raby's formula at infinity:

Theorem 3.1 (Kurdyka-Raby's formula at infinity). Let $X \subset \mathbb{R}^{n+k}$ be a definable set in an o -minimal structure on \mathbb{R} with $n = \dim X$. Let C_1, \dots, C_m be the connected components of $C_{\text{Smp}}(X, \infty)$. Then, $\theta^n(X)$ is defined and for each $x \in \mathbb{R}^{n+k}$,

$$\theta^n(X) = \lim_{r \rightarrow +\infty} \frac{\mathcal{H}^n(X \cap B_r^{n+k}(x))}{\mu_n r^n} = \sum_{j=1}^m k_{X, \infty}(C_j) \frac{\mathcal{H}^n(C_j \cap B_1^{n+k})}{\mu_n}.$$

3.1. Proof of the Kurdyka-Raby's formula at infinity.

Proof of Theorem 3.1. We first prove that the result is true for open definable sets.

Claim 3.1.1. Let $\Omega \subset \mathbb{R}^n$ be an open definable set. Then, the density at infinity satisfies

$$\lim_{r \rightarrow +\infty} \frac{\mathcal{H}^n(\Omega \cap B_r^n)}{r^n} = \mathcal{H}^n(B_1^n \cap C(\Omega, \infty)).$$

Proof of the Claim 3.1.1. First, note that $\theta^n(\Omega)$ there exists if and only if there exists $\theta^n(\Omega \setminus K)$, where $K \subset \Omega$ is a compact subset. Moreover, in this case, $\theta^n(\Omega) = \theta^n(\Omega \setminus K)$. In fact,

$$\begin{aligned} \theta^n(\Omega) &= \lim_{r \rightarrow +\infty} \frac{\mathcal{H}^n(\Omega \cap B_r^n)}{\mu_n r^n} \\ &= \lim_{r \rightarrow +\infty} \frac{\mathcal{H}^n(\Omega \cap K \cap B_r^n) + \mathcal{H}^n((\Omega \setminus K) \cap B_r^n)}{\mu_n r^n} \\ &= \lim_{r \rightarrow +\infty} \frac{\mathcal{H}^n((\Omega \setminus K) \cap B_r^n)}{\mu_n r^n} \\ &= \theta^n(\Omega \setminus K). \end{aligned}$$

When C is a cone with the vertex at 0 in \mathbb{R}^n , we have

$$\mathcal{H}^n(C \cap B_r^n) = r^n \mathcal{H}^n(C \cap B_1^n),$$

for every $r > 0$. Now, we consider the cone $C_r := \text{Cone}_0(\Omega \setminus B_r^n)$.

Claim 3.1.2. $\bigcap_{r>0} \overline{C_r} = C(\Omega, \infty)$.

Proof of the Claim 3.1.2. Let $v \in \bigcap_{r>0} \overline{C_r}$. Then, $v \in C_k$ for every $k \in \mathbb{N}$. Therefore, $v = t_l x_l$, where $x_l \in \Omega \setminus B_l^n$, with $\|x_l\| \geq l$, and $t_l \in (0, +\infty)$. Hence, $v \in C(\Omega, \infty)$.

Conversely, if $v \in C(\Omega, \infty)$, it follows from [18, Proposition 2.15] that there exists a continuous and definable curve $\alpha: (\epsilon, +\infty) \rightarrow \Omega$ such that $\alpha(t) = tv + o_\infty(t)$ with $\|\alpha(t)\| > t$, where $g(t) = o_\infty(t)$ means that $\lim_{t \rightarrow +\infty} \frac{g(t)}{t} = 0$. Now, define $\beta(t) = \frac{\alpha(t)}{t} = v + \frac{o_\infty(t)}{t}$, and note that $\|\beta(t) - v\| \rightarrow 0$ as $t \rightarrow +\infty$. Since $\alpha(t) \in C_t$, we have $\frac{\alpha(t)}{t} \in C_t$, and therefore $\beta(t) \in C_t$. Thus,

$$\text{dist}(v, C_t) \leq \|v - \beta(t)\|,$$

for every $t \geq 0$. However, $C_t \subset C_s$ when $s < t$. Then for any $s > 0$ and $s < t$, we have

$$\text{dist}(v, \overline{C_s}) \leq \text{dist}(v, \overline{C_t}) \leq \|v - \beta(t)\| \rightarrow 0,$$

as $t \rightarrow +\infty$. Hence, $v \in \overline{C_s}$ for any $s > 0$ and, consequently, $v \in \bigcap_{r>0} \overline{C_r}$. \square

Now, consider $Z = \bigcap_{r>0} C_r$, and it follows from the claim that $C(\Omega, \infty) \supset Z$.

Moreover, the functions $r \mapsto \mathcal{H}^n(C_r \cap B_1^n)$ and $r \mapsto \mathcal{H}^n(\overline{C_r} \cap B_1^n)$ are nonincreasing; hence, the following limits exist:

$$\begin{aligned} \lim_{r \rightarrow +\infty} \frac{\mathcal{H}^n(C_r \cap B_r^n)}{r^n} &= \lim_{r \rightarrow +\infty} \mathcal{H}^n(C_r \cap B_1^n) \\ &= \mathcal{H}^n\left(B_1^n \cap \bigcap_{r>0} C_r\right) \\ &= \mathcal{H}^n(B_1^n \cap Z) \end{aligned}$$

and

$$\begin{aligned}
\lim_{r \rightarrow +\infty} \frac{\mathcal{H}^n(\overline{C_r} \cap B_r^n)}{r^n} &= \lim_{r \rightarrow +\infty} \mathcal{H}^n(\overline{C_r} \cap B_1^n) \\
&= \mathcal{H}^n\left(B_1^n \cap \bigcap_{r>0} \overline{C_r}\right) \\
&= \mathcal{H}^n(B_1^n \cap C(\Omega, \infty)).
\end{aligned}$$

Since C_r is a definable set, we have

$$\mathcal{H}^n(B_1^n \cap C_r) = \mathcal{H}^n(B_1^n \cap \overline{C_r}).$$

Therefore, $\mathcal{H}^n(B_1^n \cap C(\Omega, \infty)) = \mathcal{H}^n(B_1^n \cap Z)$.

Moreover, for every $R > 0$, we have $C_R \supset \Omega \setminus (\Omega \cap B_R^n)$. Thus,

$$\begin{aligned}
\limsup_{r \rightarrow +\infty} \frac{\mathcal{H}^n(\Omega \cap B_r^n)}{r^n} &= \limsup_{r \rightarrow +\infty} \frac{\mathcal{H}^n((\Omega \setminus (\Omega \cap B_R^n)) \cap B_r^n)}{r^n} \\
&\leq \lim_{r \rightarrow +\infty} \frac{\mathcal{H}^n(C_r \cap B_r^n)}{r^n} \\
&= \mathcal{H}^n(Z \cap B_1^n) = \mathcal{H}^n(B_1^n \cap C(\Omega, \infty)).
\end{aligned}$$

For $w \in \mathbb{R}^n \setminus \{0\}$, we set $(w, +\infty) := \{tw; t > 1\}$. Now, take $z \in Z \setminus \{0\}$. Without any loss of generality, we assume that $0 \notin \Omega$. Then, $(z, +\infty) \cap \Omega$ is a definable set in Ω , and we define $\alpha(z) = \sup\{\lambda \geq 0; \lambda z \notin \Omega\}$, which is nonnegative. If $\alpha(z) \neq 0$, the segment $(\alpha(z)z, +\infty)$ is contained in Ω and if $\alpha(z) = 0$, then $(\epsilon z, +\infty)$ is contained in Ω , for all $\epsilon > 0$.

We define the following set

$$Z_r = \{z \in Z \setminus \{0\} / \|\alpha(z) \cdot z\| \leq r\}.$$

Since $\alpha(\mu z) = \sup\{\lambda \geq 0; \lambda(\mu z) \notin \Omega\} = \lambda_0$ and $\alpha(z) = \sup\{\eta \geq 0; \eta z \notin \Omega\} = \eta_0$, we have $\lambda_0 = \frac{\eta_0}{\mu}$, and Z_r is indeed a cone.

Now, note that $Z_r \setminus \overline{B_r^n} \subset \Omega \setminus \overline{B_r^n}$. Indeed, if $z \in Z_r \setminus \overline{B_r^n}$, then $\|z\| > r$, and additionally, $\|\alpha(z)z\| \leq r$. Therefore, $\alpha(z) \leq \frac{r}{\|z\|} < 1$. Note that for $r < s$, we have $Z_r \subset Z_s$. In fact, if $z \in Z_r$, then $\alpha(z)\|z\| \leq r < s$, so $z \in Z_s$.

Claim 3.1.3. $Z \setminus \{0\} = \bigcup_{r>0} Z_r$.

Proof of the Claim 3.1.3. It follows from the definition that $\bigcup_{r>0} Z_r \subset Z \setminus \{0\}$. Reciprocally, assume that $z \in Z \setminus \{0\}$. Thus for each $r > 0$ there exists a points $z_r \in \Omega \setminus B_r^n$ and $\lambda_r \in (0, +\infty)$ such that $\lambda_r z_r = z$ and since $\Omega \setminus \overline{B_r^n}$ is open set there exists $t_{r_j} \in (0, +\infty)$ such that $t_{r_j} z_j \in \Omega \setminus \overline{B_r^n}$. So, if the semiline $0\vec{z}$ we have $(t_{r_j} z_j, +\infty) := 0\vec{z} \setminus (0\vec{z} \cap B_{t_{r_j}}^n) \subset \Omega$, we have $\alpha(z) < +\infty$. Finally, note that $(z, +\infty) \subset \Omega \setminus B_r^n$.

□

Moreover, for $r > R > 0$, we have

$$\begin{aligned}
\liminf_{r \rightarrow +\infty} \frac{\mathcal{H}^n(\Omega \cap B_r^n)}{r^n} &= \liminf_{r \rightarrow +\infty} \frac{\mathcal{H}^n((\Omega \setminus B_R^n) \cap B_r^n)}{r^n} \\
&\geq \liminf_{r \rightarrow +\infty} \frac{\mathcal{H}^n((Z_r \setminus B_R^n) \cap B_r^n)}{r^n} \\
&= \liminf_{r \rightarrow +\infty} \frac{\mathcal{H}^n(Z_r \cap B_r^n) - \mathcal{H}^n(Z_r \cap B_R^n)}{r^n} \\
&= \liminf_{r \rightarrow +\infty} \left(\mathcal{H}^n(Z_r \cap B_1^n) - \frac{\mathcal{H}^n(Z \cap B_R^n)}{r^n} \right) \\
&= \mathcal{H}^n(Z \cap B_1^n) = \mathcal{H}^n(B_1^n \cap C(\Omega, \infty)).
\end{aligned}$$

Therefore, Ω has density at infinity and

$$\theta^n(\Omega) = \frac{\mathcal{H}^n(B_1^n \cap C(\Omega, \infty))}{\mu_n} = \theta^n(C(\Omega, \infty)).$$

□

Now we prove that X has density at infinity. Recently, this was proved in [33], but here we present a different proof.

Claim 3.1.4. *X has density at infinity.*

Proof of the Claim 3.1.4. Since the tangent mapping $\nu: \mathcal{M} := X \setminus \text{Sing}_1(X) \rightarrow G_n(\mathbb{R}^{n+k})$, given by $\nu(x) = T_x \mathcal{M}$, is a definable mapping (see [26, Lemma 1.14]), likewise it was done in [51], we obtain that for each $\epsilon > 0$, there exist $N(\epsilon)$ and disjoint definable sets $\Gamma_1^\epsilon, \dots, \Gamma_{N(\epsilon)}^\epsilon$ contained in X such that:

- (1) $\dim X \setminus \cup_{i=1}^{N(\epsilon)} \Gamma_i^\epsilon < n$;
- (2) Each Γ_i^ϵ is the graph of a mapping $\varphi_i^\epsilon: U_i^\epsilon \rightarrow (E_i^\epsilon)^\perp$ with derivative bounded by ϵ and U_i^ϵ is a definable open set in $E_i^\epsilon \in G_n(\mathbb{R}^{n+k})$, where $G_n(\mathbb{R}^{n+k})$ is the Grassmannian of all n -dimensional linear subspaces of \mathbb{R}^{n+k} .

By refining the above decomposition, we can assume, if necessary that each U_i^ϵ is M -LNE, where $M = M(n)$ depends only on n (see, e.g., [26, Theorem 1.2]). Therefore, each φ_i^ϵ is $M\epsilon$ -Lipschitz. Thus, the mapping $\psi_i^\epsilon: U_i^\epsilon \rightarrow \Gamma_i^\epsilon$, given by $\psi_i^\epsilon(x) = x + \varphi_i^\epsilon(x)$ is a definable lipeomorphism such that

$$\|x - y\| \leq \|\psi_i^\epsilon(x) - \psi_i^\epsilon(y)\| \leq (1 + M\epsilon)\|x - y\| \quad \forall x, y \in U_i^\epsilon.$$

By Claim 3.1.1, it follows that each U_i^ϵ has a density at infinity and

$$\theta^n(U_i^\epsilon) = \lim_{r \rightarrow +\infty} \frac{\mathcal{H}^n(U_i^\epsilon \cap B_r^n)}{\mu_n r^n} = \frac{\mathcal{H}^n(B_1^n \cap C(U_i^\epsilon, \infty))}{\mu_n} = \theta^n(C(U_i^\epsilon, \infty)).$$

Thus,

$$\frac{\theta^n(U_i^\epsilon)}{(1 + M\epsilon)^n} \leq \liminf_{r \rightarrow +\infty} \frac{\mathcal{H}^n(\Gamma_i^\epsilon \cap B_r^n)}{\mu_n r^n} \leq \limsup_{r \rightarrow +\infty} \frac{\mathcal{H}^n(\Gamma_i^\epsilon \cap B_r^n)}{\mu_n r^n} \leq (1 + M\epsilon)^n \theta^n(U_i^\epsilon).$$

By setting $\lambda(\epsilon) = \sum_{i=1}^{N(\epsilon)} \theta^n(U_i^\epsilon)$ and since for every $r > 0$, we have

$$\mathcal{H}^n(X \cap B_r^{n+k}) = \sum_{i=1}^{N(\epsilon)} \mathcal{H}^n(\Gamma_i^\epsilon \cap B_r^{n+k}),$$

then we obtain

$$\frac{\lambda(\epsilon)}{(1 + M\epsilon)^n} \leq \liminf_{r \rightarrow +\infty} \frac{\mathcal{H}^n(X \cap B_r^n)}{\mu_n r^n} \leq \limsup_{r \rightarrow +\infty} \frac{\mathcal{H}^n(X \cap B_r^n)}{\mu_n r^n} \leq (1 + M\epsilon')^n \lambda(\epsilon'),$$

for all $\epsilon, \epsilon' > 0$.

Note that $\lambda(\epsilon) \leq (1 + M\epsilon)^n 2^n \lambda(1)$. Then

$$\lim_{\epsilon \rightarrow 0} (1 + M\epsilon)^n \lambda(\epsilon) - \frac{\lambda(\epsilon)}{(1 + M\epsilon)^n} = 0.$$

Therefore $\liminf_{r \rightarrow +\infty} \frac{\mathcal{H}^n(X \cap B_r^n)}{\mu_n r^n} = \limsup_{r \rightarrow +\infty} \frac{\mathcal{H}^n(X \cap B_r^n)}{\mu_n r^n}$, and thus $\theta^n(X)$ is defined. □

In this case, it is easy to see that

$$\theta^n(X) = \lim_{r \rightarrow +\infty} \frac{\mathcal{H}^n(X \cap B_r^{n+k}(x))}{\mu_n r^n}$$

for all $x \in \mathbb{R}^{n+k}$.

Finally, we are going to prove the second part, more precisely,

$$\theta^n(X) = \sum_{j=1}^m k_{X, \infty}(C_j) \cdot \mathcal{H}^n(C_j),$$

where C_1, \dots, C_m are the connected components of $C_{\text{Smp}}(X, \infty)$.

It is worth noting that the above formula is not presented in [33].

Fixed $\epsilon > 0$ and $i \in \{1, \dots, N(\epsilon)\}$, let $U = U_i^\epsilon$, $\Gamma = \Gamma_i^\epsilon$, $\varphi = \varphi_i^\epsilon$ and $E = E_i^\epsilon$. Therefore, it follows from Theorems 2.11 and 2.16 that

- (1) $C(\Gamma, \infty)$ is the graph of a definable $M\epsilon$ -Lipschitz mapping $d_\infty \varphi: C(U, \infty) \rightarrow E^\perp$;
- (2) $k_{\Gamma, \infty}(x) = 1$ for all $x \in C_{\text{Smp}}(\Gamma, \infty)$.

Then

$$\frac{1}{(1+M\epsilon)^n} \theta^n(U_i^\epsilon) \leq \theta^n(\Gamma_i^\epsilon) \leq (1+M\epsilon)^n \theta^n(U_i^\epsilon)$$

and

$$\frac{1}{(1+M\epsilon)^n} \theta^n(C(U_i^\epsilon, \infty)) \leq \theta^n(C(\Gamma_i^\epsilon, \infty)) \leq (1+M\epsilon)^n \theta^n(C(U_i^\epsilon, \infty)).$$

Again, according to Claim 3.1.1, we have

$$\theta^n(C_{\text{Smp}}(U_i^\epsilon, \infty)) = \theta^n(C(\Gamma_i^\epsilon, \infty)) = \theta^n(U_i^\epsilon).$$

Therefore,

$$(1) \quad \frac{1}{(1+M\epsilon)^{2n}} \theta^n(C(\Gamma_i^\epsilon, \infty)) \leq \theta^n(\Gamma_i^\epsilon) \leq (1+M\epsilon)^{2n} \theta^n(C(\Gamma_i^\epsilon, \infty)).$$

So, let \mathcal{A} be a stratification of $C_{\text{Smp}}(X, \infty)$ compatible with \mathbb{S}^{n+k-1} , $C_1, \dots, C_m, C_{\text{Smp}}(\Gamma_1^\epsilon, \infty), \dots, C_{\text{Smp}}(\Gamma_{N(\epsilon)}^\epsilon, \infty)$.

For all $1 \leq i \leq N(\epsilon)$, $k_{\Gamma_i^\epsilon, \infty}(x) = 1$ for all $x \in C_{\text{Smp}}(\Gamma_i^\epsilon, \infty)$. Therefore, if $T \in \mathcal{A}$ is a stratum of dimension $n-1$ contained in $C_j \cap \mathbb{S}^{n+k-1}$, it follows from the definition of $k_{X, \infty}(C_j)$ that it is the number of pieces Γ_i^ϵ such that T is contained in $C_{\text{Smp}}(\Gamma_i^\epsilon)$. Then

$$\sum_{i=1}^{N(\epsilon)} \mathcal{H}^n(C_{\text{Smp}}(\Gamma_i^\epsilon) \cap B^{n+k}) = \sum_{j=1}^m k_{X, \infty}(C_j) \mathcal{H}^{n-1}(C_j \cap B^{n+k})$$

and thus

$$(2) \quad \sum_{i=1}^{N(\epsilon)} \theta^n(C_{\text{Smp}}(\Gamma_i^\epsilon)) = \sum_{j=1}^m k_{X, \infty}(C_j) \theta^n(C_j).$$

Therefore, from Eq. (1), (2) and

$$\theta^n(X) = \sum_{i=1}^{N(\epsilon)} \theta^n(\Gamma_i^\epsilon),$$

we obtain

$$\theta^n(X) = \sum_{j=1}^m k_{X, \infty}(C_j) \cdot \theta^n(C_j) = \sum_{j=1}^m k_{X, \infty}(C_j) \cdot \frac{\mathcal{H}^n(C_j \cap B^{n+k})}{\mu_n}.$$

□

3.2. Some examples.

Example 3.2. Let $X = \{(x, y) \in \mathbb{R}^2; y - x^2 = 0\}$. Then, $\theta^1(X) = 1$.

Example 3.3. Let $C = \{(x, y, z) \in \mathbb{R}^3; x^2 + y^2 = \cosh^2(z)\}$ be the catenoid surface. Then, $\theta^2(C) = 2$.




FIGURE 1. The catenoid minimal surface (see [9])

Example 3.4. Let $X = \{(x, y, z) \in \mathbb{R}^3; z^2 = \alpha(x^2 + y^2)\}$, where $\alpha > 0$. Then, $\theta^2(X) = 2(1 + \alpha)^{-\frac{1}{2}}$ (see [27, Example 2.5]).

If we remove the hypothesis that X is a definable set, then density can be infinite or cannot even exist.

Example 3.5. Let $H = \{(x, y, z) \in \mathbb{R}^3; z = \tan^{-1}(\frac{y}{x})\}$ be the helicoid surface. Then, $\theta^2(H) = +\infty$.




FIGURE 2. The helicoid minimal surface (see [9])

Example 3.6. Let $I_1 = (0, a_1)$ be an open interval where $a_1 > 0$. Through recurrence, we define the following sets: $A_1 := I_1 \times \{0\}$, $A_2 := I_2 \times$

$(\{-\frac{1}{2}\} \cup \{\frac{1}{2}\})$, where $I_2 = (a_1, a_1 + 2\mathcal{H}^1(I_1))$, $A_3 := I_3 \times \{0\}$, where $I_3 = (a_2, a_2 + 2\mathcal{H}^1(I_2))$ and in several cases

$$A_{2j+1} = I_{2j} \times \{0\},$$

and

$$A_{2j} = I_{2j-1} \times \left(\left\{ -\frac{1}{2} \right\} \cup \left\{ \frac{1}{2} \right\} \right),$$

where and $I_j = (a_{j-1}, a_{j-1} + 2\mathcal{H}^1(I_{j-1}))$, and $\mathcal{H}^1(I_k)$ is the length of the interval I_k . Thus we define

$$X = \bigcup_{j \in \mathbb{N}} A_j.$$

Then, the following limit does not exist

$$\lim_{r \rightarrow +\infty} \frac{\mathcal{H}^1(X \cap B_r^2)}{\mu_1 r}.$$

3.3. First consequences.

3.3.1. Degree of an algebraic set is its density at infinity.

Corollary 3.7. *Let $X \subset \mathbb{C}^{n+k}$ be a pure-dimensional complex algebraic set with $\dim_{\mathbb{C}} X = n$. Then, $\theta^{2n}(X) = \deg(X)$.*

Proof. Note that when X is a complex algebraic set, there is a complex analytic set σ with $\dim \sigma < \dim X$, such that for each irreducible component X_j of the tangent cone $C(X, \infty)$, $X_j \setminus \sigma$ intersect only one connected component C_i of $C_{\text{Smp}}(X, \infty)$, then we define $k_{X, \infty}(X_j) := k_{X, \infty}(C_i)$. Thus, from [20, §2.3], we obtain

$$\deg(X) = \sum_{j=0}^r k_{X, \infty}(X_j) \cdot \deg(X_j),$$

where X_1, \dots, X_r are all the irreducible components of $C(X, \infty)$. Since each X_j is a complex algebraic cone, we have that $\deg(X_j) = m(X_j, 0)$, where $m(X_j, 0)$ denotes the multiplicity of X_j at 0 (see the definitions of multiplicity and degree in [11]). By [15, Theorem 7.3], $\theta^{2n}(X_j, 0) = m(X_j, 0)$ for each $j \in \{1, \dots, r\}$. Since each X_j is a cone with vertex at 0, we have that $\theta^{2n}(X_j, 0) = \theta^{2n}(X_j)$. Thus,

$$\deg(X) = \sum_{j=0}^r k_{X, \infty}(X_j) \cdot \theta^{2n}(X_j).$$

By Theorem 3.1, we have that

$$\theta^{2n}(X) = \sum_{j=0}^r k_{X, \infty}(X_j) \cdot \theta^{2n}(X_j).$$

Therefore $\theta^{2n}(X) = \deg(X)$. \square

3.3.2. *The o-minimal Chow's theorem.* Let us remember the Stoll-Bishop's Theorem:

Theorem 3.8 (Stoll-Bishop's Theorem (see [52, 53, 8])). *Let $Z \subset \mathbb{C}^m$ be a pure-dimensional entire complex analytic subset with $\dim_{\mathbb{C}} X = n$. Then Z is a complex algebraic set if and only if there exists a constant $R > 0$ such that*

$$\frac{\mathcal{H}^{2n}(Z \cap \overline{B}_r^{2m}(0))}{r^{2n}} \leq R, \quad \text{for all } r > 0.$$

Thus, we obtain the o-minimal Chow's Theorem proved in [35, Corollary 4.5]:

Corollary 3.9 (O-minimal Chow's theorem). *Let $X \subset \mathbb{C}^{n+k}$ be a pure-dimensional entire complex analytic set with $\dim_{\mathbb{C}} X = n$. Then, X is a complex algebraic set if and only if X is definable in an o-minimal structure on \mathbb{R} .*

Proof. By Theorem 3.1, $\theta^{2n}(X)$ is finite. Then there exists a constant $R > 0$ such that

$$\frac{\mathcal{H}^{2n}(X \cap \overline{B}_r^{2(n+k)}(0))}{r^{2n}} \leq R, \quad \text{for all } r > 0.$$

Therefore, by Stoll-Bishop's Theorem, X is a complex algebraic set. \square

3.3.3. *The Lipschitz o-minimal Chow's theorem.*

Corollary 3.10 (Lipschitz o-minimal Chow's theorem). *Let $X \subset \mathbb{C}^{n+k}$ be a pure-dimensional entire complex analytic set with $\dim_{\mathbb{C}} X = n$. Then, X is a complex algebraic set if and only if X is lipeomorphic at infinity to a definable set in an o-minimal structure on \mathbb{R} .*

Proof. We closely follow the proof of Theorem 3.1 in [42].

We assume that $0 \in X$ and that X is lipeomorphic at infinity to a definable set in an o-minimal structure on \mathbb{R} . Let $A \subset \mathbb{R}^m$ be a definable set such that there exist compact subsets $K \subset \mathbb{C}^{n+k}$ and $\tilde{K} \subset \mathbb{R}^m$ and a lipeomorphism $\varphi: A \setminus \tilde{K} \rightarrow X \setminus K$. Let $\lambda \geq 1$ such that

$$\frac{1}{\lambda} \|x - y\| \leq \|\varphi(x) - \varphi(y)\| \leq \lambda \|x - y\|, \quad \forall x, y \in A \setminus \tilde{K}.$$

Fix $x_0 \in A \setminus \tilde{K}$ and set $y_0 = \varphi(x_0)$. Let $\tilde{r}_0 = \|x_0\|$ and $r_0 = \|y_0\|$. Thus, for any $r > 0$, we obtain that

$$(X \setminus K) \cap B_r^{2(n+k)}(0) \subset \varphi((A \setminus \tilde{K}) \cap B_{\lambda(r+r_0)+\tilde{r}_0}^m(0))$$

and

$$\mathcal{H}^{2n}(X \cap B_r^{2(n+k)}(0)) \leq \lambda^{2n} \mathcal{H}^{2n}(A \cap B_{\lambda(r+r_0)+\tilde{r}_0}^m(0)) + \mathcal{H}^{2n}(X \cap K).$$

Since A is a definable set, by Theorem 3.1, $\theta^{2n}(A)$ is finite. Then there exists a constant $C > 0$ such that

$$\frac{\mathcal{H}^{2n}(A \cap \overline{B}_r^m(0))}{r^{2n}} \leq C, \quad \text{for all } r > 0.$$

But X is analytic at 0, then there exist $r_1, R_1 > 0$ such that

$$\frac{\mathcal{H}^{2n}(X \cap B_r^{2(n+k)}(0))}{r^{2n}} \leq R_1,$$

for all $r \leq r_1$. Moreover, $\mathcal{H}^{2n}(X \cap K) < +\infty$. Then, there exists $R_2 > 0$ such that

$$\frac{\mathcal{H}^{2n}(X \cap B_r^{2(n+k)}(0))}{r^{2n}} \leq \frac{\lambda^{2n} C (\lambda(r+r_0) + \tilde{r}_0)^{2n} + \mathcal{H}^{2n}(X \cap K)}{r^{2n}} \leq R_2,$$

for all $r \geq r_1$.

Therefore, by Stoll-Bishop's Theorem, X is a complex algebraic set. \square

3.3.4. *O-minimal Yau's Bernstein Problem.*

Corollary 3.11. *Let $X \subset \mathbb{R}^{n+1}$ be an oriented stable complete minimal hypersurface with $n \leq 6$. If X is a definable set then X is a hyperplane.*

Proof. By Theorem 3.1, we have

$$\lim_{r \rightarrow +\infty} \frac{\mathcal{H}^n(X \cap B_r^{n+1}(0))}{r^n} < +\infty.$$

Thus, by [46, Corollary, p. 104] (see also [47] and [48]), X is a hyperplane. \square

4. PARAMETRIC VERSIONS OF THE BERNSTEIN THEOREM

Theorem 4.1. *Let $X \subset \mathbb{R}^{n+k}$ be a closed and connected set with $n = \dim_H X$. Assume that X satisfies the monotonicity formula at some $p \in X$. Then, the following statements are equivalent.*

- (1) X is an affine linear subspace;
- (2) X is a definable set, blow-spherical regular at infinity and $C(X, \infty)$ is a linear subspace;
- (3) X is a definable set that is Lipschitz regular at infinity and $C(X, \infty)$ is a linear subspace;
- (4) X is an LNE at infinity definable set and $C(X, \infty)$ is a linear subspace;
- (5) $\theta^n(X) = 1$.

Proof. It is obvious that item (1) implies the items (2), (3), (4) and (5).

(2) \implies (5). Assume that X is a definable set, blow-spherical regular at infinity and that $C(X, \infty)$ is a linear subspace.

Since X is blow-spherical regular at infinity, by Proposition 2.14 in [45], we have $k_{X, \infty}(C) = 1$ for all connected component C of $C_{\text{Smp}}(X, \infty)$. It follows from the hypothesis that $C(X, \infty)$ is a hyperplane and Theorem 3.1 that $\theta^n(X) = 1$.

(3) \implies (5). Assume that X is a definable set that is Lipschitz regular at infinity and $C(X, \infty)$ is a linear subspace.

By Theorem 2.16, we obtain that $k_{X, \infty}(v) = 1$, for all $v \in C_{\text{Smp}}(X, \infty)$. Therefore, as $C(X, \infty)$ is a linear subspace, it follows from Claim 4.1.1, Remark 2.12, and Theorem 3.1 that $\theta^n(X) = \mathcal{H}^n(C_1) = 1$.

(4) \implies (5). Assume that X is a definable set that is LNE at infinity and $C(X, \infty)$ is a linear subspace.

Claim 4.1.1. *If $X \subset \mathbb{R}^m$ is a definable set and LNE at infinity, then $k_{X, \infty}(v) = 1$ for all $v \in C_{\text{Smp}}(X, \infty)$.*

Proof of Claim 4.1.1. Indeed, we suppose there exists $x = (x', 0) \in \text{Smp}(\partial X'_\infty) = C(X, \infty) \cap \mathbb{S}^{m-1} \times \{0\}$ such that $k_{X, \infty}(x') \geq 2$, we know that $k_{X, \infty}(X_j) = k_{X, \infty}(x')$ is the number of connected components of the germ $(\rho_\infty^{-1}(X), x)$, where X_j is a connected component of $C_{\text{Smp}}(X, \infty)$ and $x' \in X_j$. Thus, for a sequence of positive real numbers $\{t_j\}_{j \in \mathbb{N}}$ such that $\lim t_j = +\infty$, $k_{X, \infty}(x')$ can be seen as the number of connected components of the set $(SX \cap \mathbb{S}^{m-1} \times \{t_j^{-1}\}) \cap B_\delta^{m+1}(x)$ for all large k and a small enough $\delta > 0$, where

$$SX = \{(x, s) \in \mathbb{S}^{m-1} \times (0, +\infty); \frac{1}{s} \cdot x \in X \text{ and } \frac{1}{s} \in S\}$$

and $S = \{t_j^{-1}\}_{j \in \mathbb{N}}$.

Let X_1 and X_2 be two connected components of $X'_\infty \cap B_\delta^{m+1}(x)$, and for $\eta, R > 0$, we define the following set $C_{\eta, R}(x') = \{v \in \mathbb{R}^n \setminus \{0\}; \exists t > 0; \|v - tx'\| \leq \eta t\} \setminus B_R^n$. For each $j \in \mathbb{N}$, we can take $x_j \in \rho_\infty(X_1)$ and $y_j \in \rho_\infty(X_2)$ such that $\|x_j\| = \|y_j\| = t_j$ and $\lim_{j \rightarrow +\infty} \frac{x_j}{t_j} = \lim_{j \rightarrow +\infty} \frac{y_j}{t_j} = x'$. Now, taking subsequence, if necessary, we assume that $x_j, y_j \in C_{\frac{\eta}{2}, R}$. Thus, let $\beta_j: [0, 1] \rightarrow X$ be a curve connecting x_j to y_j , we have that there exists a $t_0 \in [0, 1]$ such that $\beta_j(t_0) \notin C_{\eta, R}$, since x_j and y_j belong to different

connected components of $X \cap C_{\eta,R}$. Therefore, $\text{length}(\beta_j) \geq \eta t_j$ and then $d_X(x_j, y_j) \geq \frac{\eta}{2} t_j$.

On the other hand, X is a Lipschitz normally embedded at infinity set, then there exists a ball B_r^n and a constant $C > 0$ such that $d_{X \setminus B_r^n}(v, w) \leq C \|v - w\|$, for all $v, w \in X \setminus B_r(0)$. Therefore, $C \left\| \frac{x_j}{t_j} - \frac{y_j}{t_j} \right\| \geq \eta$, for all $j \in \mathbb{N}$ and this is a contradiction, since $\lim \frac{x_j}{t_j} = \lim \frac{y_j}{t_j}$.

Therefore $k_{X,\infty}(v) = 1$ for all $v \in C_{\text{Smp}}(X, \infty)$. \square

Therefore, as $C(X, \infty)$ is a linear subspace, it follows from Claim 4.1.1, Remark 2.12, and Theorem 3.1 that $\theta^n(X) = \mathcal{H}^n(C_1) = 1$.

(5) \implies (1). Suppose that $\theta^n(X) = 1$.

Since X satisfies the monotonicity formula at p , $\theta^n(X, p, \cdot)$ is a nondecreasing function such that $1 \leq \theta^n(X, p, r) \leq \theta^n(X) = 1$ for all $r > 0$. Then $\theta^n(X, p, r) = 1$ for all $r > 0$, and by using again that X satisfies the monotonicity formula at p , we have that X is a cone with vertex at p and X is C^1 -smooth at p . Therefore X is an affine linear subspace. \square

4.1. Some examples.

Example 4.2. *The catenoid is the surface $C = \{(x, y, z) \in \mathbb{R}^3; x^2 + y^2 = \cosh^2(z)\}$. The Catenoid is a complete embedded minimal hypersurface and its geometric tangent cone at infinity is the plane $z = 0$. Moreover, it is definable in the o-minimal structure \mathbb{R}_{exp} (see [16]).*

Example 4.3. *Let $L \subset \mathbb{S}^2$ be a curve lipeomorphic to \mathbb{S}^1 with $2\pi < \mathcal{H}^1(L) < +\infty$. Thus, the surface $X = \text{Cone}_0(L) := \{tx; x \in L \text{ and } t \geq 0\}$ is lipeomorphic to \mathbb{R}^2 and, in particular, it is Lipschitz regular at infinity. Since X is a cone, then $\theta^2(X, 0, r) = \mathcal{H}^1(L)/(2\pi) > 1$ for all $r > 0$. Therefore X satisfies the monotonicity formula at 0. However, it is not a plane.*

Example 4.4. *The surface $P = \{(z, w) \in \mathbb{C}^2; z = w^2\}$ is a smooth complex algebraic curve and, in particular, it is an area-minimizing set. Its geometric tangent cone at infinity is the complex line $w = 0$. However, P is not a plane.*

Example 4.5. *We cannot remove the hypothesis that X satisfies the monotonicity formula at some p . Indeed, $X = \{(x, y, z) \in \mathbb{R}^3; z^3 = x^2 + y^2 + 1\}$ is lipeomorphic to \mathbb{R}^2 and, in particular, it is Lipschitz regular at infinity. In fact, X is the graph of the Lipschitz function $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ given by $f(x, y) = (x^2 + y^2 + 1)^{\frac{1}{3}}$. Moreover, X is a smooth surface, $C(X, \infty)$ is the plane $z = 0$ and $\theta^2(X) = 1$. However, X is not a plane.*

4.2. Some direct consequences. It is a direct consequence of Theorem 3.1 and Claim 4.1.1 the following result:

Corollary 4.6. *Let $X \subset \mathbb{R}^{n+k}$ be an n -dimensional definable set. If X is LNE at infinity, then $\theta^n(X) = \theta^n(C(X, \infty))$.*

In particular, since for a pure dimensional complex algebraic set, $X \subset \mathbb{C}^{n+k}$ with $\dim_{\mathbb{C}} X = n$, we have $\theta^{2n}(X) = \deg(X)$ (see Corollary 3.7), we obtain the main result of [14]:

Corollary 4.7. *Let $X \subset \mathbb{C}^{n+k}$ be pure dimensional complex algebraic set. If X is LNE at infinity, then $\deg(X) = \deg(C(X, \infty))$.*

We also obtain the following:

Corollary 4.8. *Let $X \subset \mathbb{R}^{n+k}$ be a pure-dimensional algebraic set. Suppose that X is LNE at infinity and $C(X, \infty)$ is a linear subspace. If X is a minimal submanifold or an area-minimizing set, then X is an affine linear subspace.*

Proof. By Corollary 4.6, $\theta^n(X) = \theta^n(C(X, \infty))$. Since $C(X, \infty)$ is a hyperplane, we have $\theta^n(X) = 1$. By Theorem 4.1, X is an affine linear subspace. \square

We obtain the main results of [18] (see also [42]):

Corollary 4.9. *Let $X \subset \mathbb{C}^{n+k}$ be a pure-dimensional complex analytic set with $\dim_{\mathbb{C}} X = n$. Suppose that X is LNE at infinity and $C(X, \infty)$ is a dimensional complex linear subspace with $\dim_{\mathbb{C}} C(X, \infty) = n$. Then X is an affine linear subspace.*

Proof. Since $C(X, \infty)$ is a dimensional complex algebraic set with $\dim_{\mathbb{C}} C(X, \infty) = n$, we have that X is a complex algebraic set as well (see, e.g., [42, Theorem 3.1]). In particular, X is a definable set and satisfies the monotonicity formula at any $p \in X$. By Theorem 3.1, X is an affine linear subspace. \square

Corollary 4.10. *Let $X \subset \mathbb{C}^{n+k}$ be a pure-dimensional complex algebraic set with $\dim_{\mathbb{C}} X = n$. Suppose that X is Lipschitz regular at infinity. Then X is an affine linear subspace.*

Proof. X is Lipschitz regular at infinity, then X is LNE at infinity, and by Theorem 2.11, $C(X, \infty)$ is lipeomorphic to \mathbb{R}^{2n} . In particular, $C(X, \infty)$ is a topological manifold, and thus by Prill's theorem ([37, Theorem]), $C(X, \infty)$ is a dimensional complex linear subspace with $\dim_{\mathbb{C}} C(X, \infty) = n$. By Theorem 3.1, X is an affine linear subspace. \square

We obtain also the following version of Moser's result [31, Corollary, p. 591]:

Corollary 4.11. *Let $X \subset \mathbb{R}^{n+1}$ be a complete minimal hypersurface. Suppose that there are compact sets $K \subset \mathbb{R}^n$ and $\tilde{K} \subset \mathbb{R}^{n+1}$ such that $X \setminus \tilde{K}$ is the graph of a definable Lipschitz function $u: \mathbb{R}^n \setminus K \rightarrow \mathbb{R}$. Then u is the restriction of an affine function and, in particular, X is a hyperplane.*

Proof. We need a preliminary result. For a set $A \subset \mathbb{R}^n$, denote by $\mathcal{N}(A, \infty)$ the union of all hyperplanes T such that there is a sequence $\{x_j\}_j \subset A \setminus \text{Sing}_1(A)$ such that $\lim \|x_j\| = +\infty$ and $T_{x_j}A$ converges to T .

Lemma 4.12. *Let $A \subset \mathbb{R}^n$ be an unbounded definable set. Then, $C(A, \infty) \subset \mathcal{N}(A, \infty)$.*

Proof. Firstly, note that $C(A \setminus \text{Sing}_1(A), \infty) = C(A, \infty)$ and $\mathcal{N}(A \setminus \text{Sing}_1(x), \infty) = \mathcal{N}(A, \infty)$. Thus, we may assume that A is C^1 -smooth.

Let $v \in C(A, \infty)$. We are going to show that $v \in \mathcal{N}(A, \infty)$ and we may assume that $\|v\| = 1$. In order to do that, it is enough to find, for some $R > 0$, a definable C^1 -smooth arc $\alpha: (R, +\infty) \rightarrow A$ such that $\alpha(t) = tv + o_\infty(t)$. Indeed, by taking a subsequence, if necessary, we may assume that $T_{\alpha(j)}A$ converges to $T \subset \mathcal{N}(A, \infty)$. Since $\alpha'(j) \in T_{\alpha(j)}A$ for all j , then $\lim \alpha'(j) = v \in T$.

Now, let us prove that there is such an arc α . Let $Y = \rho_\infty^{-1}(A)$. We have that Y is a definable set. Note that the closure of Y in $\mathbb{S}^{n-1} \times [0, \infty)$ satisfies the following:

$$\overline{Y} \cap (\mathbb{S}^{n-1} \times \{0\}) = (C(A, \infty) \cap \mathbb{S}^{n-1}) \times \{0\}.$$

Thus, $(v, 0) \in \overline{Y}$. By the Curve Selection Lemma, there is a definable continuous arc $\beta: [0, \epsilon) \rightarrow \overline{Y}$ such that $\beta((0, \epsilon)) \subset Y$ and $\beta(0) = (v, 0)$. Let us write $\beta(s) = (x(s), h(s))$. Since h is not a constant function around 0, by Monotonicity Lemma, one can suppose that β is C^1 -smooth and that h is C^1 smooth, strictly increasing in the domain $(0, \delta)$ and $h'(s) \neq 0$ for all $s \in (0, \delta)$. Hence, $h: (0, \delta/2) \rightarrow (0, \delta')$ is a C^1 -diffeomorphism, where $\delta' = h(\frac{\delta}{2})$. Let $\alpha: (\frac{1}{\delta'}, +\infty) \rightarrow A$ be the arc $\alpha(t) = \rho_\infty \circ \beta \circ h^{-1}(\frac{1}{t})$.

Clearly, α is C^1 -smooth, and since $\lim_{s \rightarrow 0^+} x(s) = x(0) = v$, we obtain

$$\alpha(t) = tx(h^{-1}(\frac{1}{t})) = tv + o_\infty(t).$$

□

From the above lemma, we obtain that $C(X, \infty) \subset \mathcal{N}(X, \infty)$.

By Lemma 1.18 in [49], we have that the following limit exists:

$$\lim_{x \rightarrow \infty} (Du(x), -1) = w.$$

Therefore, $\mathcal{N}(X, \infty) = w^\perp$. In particular, $C(X, \infty)$ is a subset of an n -dimensional hyperplane, and thus $\mathcal{H}^n(C(X, \infty) \cap B_1^{n+1}) \leq \mu_n$. Since there is a blow-spherical homeomorphism at infinity between X and \mathbb{R}^n , then there is only one relative multiplicity which is 1. Then, by Theorem 3.1, $\theta^n(X) \leq 1$. However, since X is a minimal submanifold, $\theta^n(X) \geq 1$. Therefore, $\theta^n(X) = 1$ and by Theorem 4.1, X is a hyperplane. It is easy to verify that u is the restriction of an affine function. \square

In the next section, we generalize Corollary 4.11 (see Corollary 5.6).

5. GENERALIZATION OF THE MOSER'S BERNSTEIN THEOREM

In this section, we generalize the Moser's Bernstein Theorem.

Theorem 5.1. *Let $X \subset \mathbb{R}^{n+k}$ be a connected closed set with Hausdorff dimension n . Then, X is an n -dimensional affine linear subspace if and only if we have the following:*

- (1) X satisfies the monotonicity formula at some $p \in X$;
- (2) there are compact sets $K \subset \mathbb{R}^n$ and $\tilde{K} \subset \mathbb{R}^{n+k}$ such that $X \setminus \tilde{K}$ is the graph of a C^1 -smooth function $u: \mathbb{R}^n \setminus K \rightarrow \mathbb{R}^k$ with bounded derivative;
- (3) $\mathcal{N}(X, \infty)$ is a linear subspace.

Proof. It is obvious that if X is an n -dimensional affine linear subspace, then X satisfies (1), (2) and (3).

Reciprocally, assume that X satisfies (1), (2) and (3).

As before, for a set $A \subset \mathbb{R}^n$, denote by $\mathcal{N}(A, \infty)$ the union of all hyperplanes T such that there is a sequence $\{x_j\}_j \subset A \setminus \text{Sing}_1(A)$ such that $\lim \|x_j\| = +\infty$ and $T_{x_j}A$ converges to T .

Let P be the hyperplane $\mathcal{N}(X, \infty)$. We choose linear coordinates (y_1, \dots, y_{n+k}) of \mathbb{R}^{n+k} such that $P = \{(y_1, \dots, y_{n+k}) \in \mathbb{R}^{n+k}; y_{n+1} = \dots = y_{n+k} = 0\}$. For a larger enough $R > 0$, $X \setminus (\overline{B_R^n} \times \mathbb{R}^k)$ is the graph of a function $v = (v_1, \dots, v_k): P \setminus \overline{B_R^n} \rightarrow \mathbb{R}^k$. By the Implicit Mapping Theorem, we have that v is a Lipschitz function and that the following limit exists:

$$\lim_{y \rightarrow \infty} Dv_i(y) = \tilde{w}_i, \quad i \in \{1, \dots, k\}.$$

Therefore $\tilde{w} = (\tilde{w}_1, \dots, \tilde{w}_k) = 0$. Then, for each $\varepsilon > 0$, there is $r > 0$ such that $\|Dv(y)\| \leq \varepsilon$ for all $y \in \Omega := P \setminus \overline{B_r^n}$.

Claim 5.1.1. $v|_\Omega$ is a Lipschitz mapping with constant $\pi\varepsilon$.

Proof. Firstly, we define the set of all piecewise C^1 -smooth curves connecting x to y in Ω . We denote such a set by:

$$\Omega(x, y) := \{ \gamma: [0, 1] \rightarrow \Omega; \gamma(0) = x, \gamma(1) = y \text{ and } \gamma \text{ is piecewise } C^1\text{-smooth} \}.$$

Let $x, y \in \Omega$ be any two points. Then

$$\begin{aligned} \|v(x) - v(y)\| &\leq \inf_{\gamma \in \Omega(x, y)} \int_0^1 \|(v \circ \gamma)'(t)\| dt \\ &= \inf_{\gamma \in \Omega(x, y)} \int_0^1 \|Dv(\gamma(t)) \cdot \gamma'(t)\| dt \\ &\leq \inf_{\gamma \in \Omega(x, y)} \int_0^1 \|\nabla(\gamma(t))\| \|\gamma'(t)\| dt \\ &\leq \varepsilon \inf_{\gamma \in \Omega(x, y)} \int_0^1 \|\gamma'(t)\| dt \\ &= \varepsilon d_\Omega(x, y). \end{aligned}$$

However, Ω is a set that is LNE such that $d_\Omega(x, y) \leq \pi\|x - y\|$ for all $x, y \in \Omega$. Therefore,

$$\|v(x) - v(y)\| \leq \pi\varepsilon\|x - y\|.$$

□

Claim 5.1.2. $1 \leq \theta^n(X) \leq (1 + \pi\varepsilon)^n$.

Proof. Let $\varphi: \Omega \rightarrow A = X \setminus (\overline{B_r^n} \times \mathbb{R}^k)$ be the mapping given by $\varphi(x) = (x, v(x))$. We have that φ is a bi-Lipschitz mapping such that

$$\|x - y\| \leq \|\varphi(x) - \varphi(y)\| \leq (1 + \pi\varepsilon)\|x - y\|.$$

Therefore,

$$\mathcal{H}^n(B_t^n \setminus B_r^n) \leq \mathcal{H}^n(A \cap B_t^{n+k}) \leq (1 + \pi\varepsilon)^n \mathcal{H}^n(B_{t+\pi\varepsilon}^n),$$

for all $t > r$. Thus, we obtain the following $\theta^n(\mathbb{R}^n \setminus B_r^n) \leq \theta^n(X) \leq (1 + \pi\varepsilon)^n \theta^n(\mathbb{R}^n)$. Since $\theta^n(\mathbb{R}^n \setminus B_r^n) = \theta^n(\mathbb{R}^n) = 1$ and $\theta^n(A) = \theta^n(X)$, we obtain that $1 \leq \theta^n(X) \leq (1 + \pi\varepsilon)^n$, which finishes the proof of the claim. □

Setting $\varepsilon \rightarrow 0^+$, we obtain that $\theta^n(X) = 1$. By Theorem 4.1, X is an affine linear subspace.

□

5.1. Examples. In this subsection, by presenting several examples, we show that Theorem 5.1 is sharp in the sense that no hypothesis of it can be removed.

We cannot remove the hypothesis that $\mathcal{N}(X, \infty)$ is a linear subspace.

Example 5.2 (Theorem 7.1 in [28]). *There is a connected closed semialgebraic set $X \subset \mathbb{R}^{n+k}$ which is an area-minimizing set and is the graph of a semialgebraic Lipschitz function $u: \mathbb{R}^n \rightarrow \mathbb{R}^k$. Moreover, X is not an affine linear subspace.*

We cannot remove the condition that X is the graph of a function outside of a compact.

Example 5.3. *The catenoid $C = \{(x, y, z) \in \mathbb{R}^3; x^2 + y^2 = \cosh^2(z)\}$ (see Example 4.2) satisfies the monotonicity formula at any point $p \in C$ and $N(C, \infty)$ is the plane $z = 0$; however, C is not a plane.*

We cannot remove the hypothesis that the function u has bounded derivative.

Example 5.4. *The surface $P = \{(z, w) \in \mathbb{C}^2; z = w^2\}$ is a smooth complex algebraic curve and, in particular, it is an area-minimizing set. $N(P, \infty)$ is the complex line $w = 0$. However, P is not a plane.*

We cannot remove the hypothesis that X satisfies the monotonicity formula at some p .

Example 5.5. *$X = \{(x, y, z) \in \mathbb{R}^3; z^3 = x^2 + y^2 + 1\}$ is the graph of the smooth Lipschitz function $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ given by $f(x, y) = (x^2 + y^2 + 1)^{\frac{1}{3}}$. Moreover, X is a smooth surface and $\mathcal{N}(X, \infty)$ is the plane $z = 0$. However, X is not a plane.*

5.2. Consequences.

Corollary 5.6. *Let $X \subset \mathbb{R}^{n+1}$ be a connected closed set with Hausdorff dimension n and that satisfies the monotonicity formula at some $p \in X$. Suppose that there are compact sets $K \subset \mathbb{R}^n$ and $\tilde{K} \subset \mathbb{R}^{n+1}$ such that $X \setminus \tilde{K}$ is a minimal hypersurface that is the graph of a C^2 -smooth function $u: \mathbb{R}^n \setminus K \rightarrow \mathbb{R}$ with bounded derivative. Then u is the restriction of an affine function and, in particular, X is a hyperplane.*

Proof. By Lemma 1.18 in [49] and Theorem X in [5], we have that the following limit exists:

$$\lim_{x \rightarrow \infty} (Du(x), -1) = w.$$

Therefore, $\mathcal{N}(X, \infty)$ is the hyperplane w^\perp .

By Theorem 5.1, X is a hyperplane. It is obvious that u is the restriction of an affine function. \square

Corollary 5.7. *Let $X \subset \mathbb{R}^{n+1}$ be a connected closed set with Hausdorff dimension n and that satisfies the monotonicity formula at some $p \in X$. Suppose that $2 \leq n \leq 7$ and there are compact sets $K \subset \mathbb{R}^n$ and $\tilde{K} \subset \mathbb{R}^{n+1}$ such that $X \setminus \tilde{K}$ is a minimal hypersurface that is the graph of a C^2 -smooth function $u: \mathbb{R}^n \setminus K \rightarrow \mathbb{R}$. Then u is the restriction of an affine function and, in particular, X is a hyperplane.*

Proof. By Theorem 1 in [49] and Theorem X in [5], Du is bounded. By Corollary 5.6, X is a hyperplane and u is the restriction of an affine function. \square

Since a minimal hypersurface satisfies the monotonicity formula at any of its points, we obtain, as direct consequences of Corollaries 5.7 and 5.6, the following results:

Corollary 5.8. *Let $X \subset \mathbb{R}^{n+1}$ be a connected complete minimal hypersurface. Suppose that there are compact sets $K \subset \mathbb{R}^n$ and $\tilde{K} \subset \mathbb{R}^{n+1}$ such that $X \setminus \tilde{K}$ is the graph of a C^2 -smooth function $u: \mathbb{R}^n \setminus K \rightarrow \mathbb{R}$ with bounded derivative. Then X is a hyperplane and u is the restriction of an affine function.*

Corollary 5.9. *Let $X \subset \mathbb{R}^{n+1}$ be a connected complete minimal hypersurface with $2 \leq n \leq 7$. Suppose that there are compact sets $K \subset \mathbb{R}^n$ and $\tilde{K} \subset \mathbb{R}^{n+1}$ such that $X \setminus \tilde{K}$ is the graph of a C^2 -smooth function $u: \mathbb{R}^n \setminus K \rightarrow \mathbb{R}$. Then X is a hyperplane and u is the restriction of an affine function.*

We finish this section by presenting the following example, which shows that the hypothesis that X is a complete hypersurface in the above corollaries cannot be dropped:

Example 5.10. *Outside of a compact set, the upper part of the catenoid is the graph of a Lipschitz function (see Example 4.2). Indeed, we can see the upper part of the catenoid as the graph of the function $f: \mathbb{R}^2 \setminus \overline{B_2^2} \rightarrow \mathbb{R}$ given by $f(x, y) = \operatorname{arccosh}(\sqrt{x^2 + y^2})$. Now, we note that*

$$\left| \frac{\partial f}{\partial x}(x, y) \right| = \frac{x}{\sqrt{x^2 + y^2}} \frac{1}{\sqrt{x^2 + y^2 - 1}} \leq 1,$$

and

$$\left| \frac{\partial f}{\partial y}(x, y) \right| = \frac{y}{\sqrt{x^2 + y^2}} \frac{1}{\sqrt{x^2 + y^2 - 1}} \leq 1.$$

Since $\mathbb{R}^2 \setminus \overline{B_2^2}$ is Lipschitz normally embedded set, we have that f is Lipschitz (see Claim 5.1.1).

Acknowledgements. The authors would like to thank Alexandre Fernandes and Pacelli Bessa for their interest, encouragement, and comments on this research.

REFERENCES

- [1] Allard, W. K. *On the first variation of a varifold*. Ann. of Math. (2), 95:417–491, 1972.
- [2] Almgren, F. J. *Some interior regularity theorems for minimal surfaces and an extension of Bernstein's theorem*. Annals of Mathematics, Second Series, 84 (2), (1966): 277–292.
- [3] Assimos, R. and Jost, J. *The Geometry of Maximum Principles and a Bernstein Theorem in Codimension 2*. Preprint (2018), arXiv:1811.09869 [math.DG].
- [4] Bernstein, S. N. *Sur une théorème de géométrie et ses applications aux équations dérivées partielles du type elliptique*. Comm. Soc. Math. Kharkov, (15), 38–45.
- [5] Bers, L. *Isolated Singularities of Minimal Surfaces*. Ann. Math., vol. 53 (1951), 364–386.
- [6] Birbrair, L. and Mostowski, T. *Normal embeddings of semialgebraic sets*. Michigan Math. J., vol. 47 (2000), 125–132.
- [7] Birbrair, L.; Fernandes, A.; Lê D. T. and Sampaio, J. E. *Lipschitz regular complex algebraic sets are smooth*. Proceedings of the American Mathematical Society, vol. 144 (2016), 983–987.
- [8] Bishop, E. *Condition for the analyticity of certain sets*. Michigan Math. J., vol. 11 (1964), 289–304.
- [9] Classical Surfaces. *Bloomington's Virtual Minimal Surface Museum*. <https://minimal.sitehost.iu.edu/archive/Classical/index.html>. Accessed: November 29, 2023.
- [10] Bombieri, E.; De Giorgi, E.; Giusti, E. *Minimal cones and the Bernstein problem*. Inventiones Mathematicae, vol. 7 (1969), 243–268.
- [11] Chirka, E.M. *Complex analytic sets*. Kluwer Academic Publishers, 1989.
- [12] Collin, P. *Topologie et courbure des surfaces minimales proprement plongées de \mathbb{R}^3* . Ann. of Math., (145)2 (1997), 1-31.
- [13] De Giorgi, E. , *Una estensione del teorema di Bernstein*. Ann. Scuola Norm. Sup. Pisa (3) (1965), 19: 79–85.
- [14] Dias, L.R.G. and Ribeiro, N.R. *Lipschitz normally embedded set and tangent cones at infinity*. The Journal of Geometric Analysis, vol. 32 (2022), article number 51, 1-10.

- [15] Draper, R. N. *Intersection Theory in Analytic Geometry*. Mathematische Annalen, vol. 180 (1969), 175–204.
- [16] van den Dries, L. *Remarks on Tarski's problem concerning $(R, +, \cdot, exp)$* . G. Lolli (ed.) G. Longo (ed.) A. Marcja (ed.), Logic Colloquium '82, North-Holland (1984), 97–121.
- [17] Ephraim, R. *C^1 preservation of multiplicity*. Duke Math., vol. 43 (1976), 797–803.
- [18] Fernandes, A. and J. E. Sampaio. *On Lipschitz rigidity of complex analytic sets*. The Journal of Geometric Analysis, vol. 30 (2020), 706–718.
- [19] Fernandes, A. and Sampaio, J. E. *Bi-Lipschitz invariance of the multiplicity*. Handbook of Geometry and Topology of Singularities IV. Jose Luis Cisneros Molina, José Seade, Dũng Tráng Lê (editors). Springer Nature Switzerland AG, 2023.
- [20] Fernández de Bobadilla, J.; Fernandes, A. and Sampaio, J. E. *Multiplicity and degree as bi-Lipschitz invariants for complex sets*. Journal of Topology, vol. 11, pp. 957–965, 2018.
- [21] Ghomi, M. and Howard, R. *Tangent cones and regularity of real hypersurfaces*. Journal für die reine und angewandte Mathematik (Crelles Journal), vol. 697 (2014), 221–247.
- [22] Harvey, R. and Lawson, B. *Extending Minimal Varieties*. Inventiones math. 28 (1975), 209–226.
- [23] Jost, J. and Xin, Y.L. *Bernstein type theorems for higher codimension*. Calc. Var. Partial Differ. Equ., vol. 9 (1999), 277–296.
- [24] Jost, J.; Xin, Y.L. and Yang, L. *The geometry of Grassmannian manifolds and Bernstein type theorems for higher codimension*. Ann. Del. Scu. Norm. Sup. Di Pisa, vol. XVI (2016), 1–39.
- [25] Jost, J.; Xin, Y. L. and Yang, L. *A spherical Bernstein theorem for minimal submanifolds of higher codimension*. Calc. Var. Partial Differ. Equ., vol. 57 (2018), article no. 166, 1–21.
- [26] Kurdyka, K. and Parsusiński, A. *Quasi-convex decomposition in \mathcal{O} -minimal structures. Application to the gradient conjecture*. Advanced Studies in Pure Mathematics, vol. 43 (2006), Singularity Theory and Its Applications pp. 137-177.
- [27] Kurdyka, K. and Raby, G. *Densité des ensembles sous-analytiques*. Ann. Inst. Fourier (Grenoble), vol. 39 (1989), no. 753–771.
- [28] Lawson Jr., H. B. and Osserman, R. *Non-existence, non-uniqueness and irregularity of solutions to the minimal surface system*. Acta Math., vol. 139 (1977), no. 1-2, 1–17.
- [29] Meeks III, W. H. *Global Problems in Classical Minimal Surface Theory*. Global theory of minimal surfaces, Clay Math. Proc., vol. 2, Amer. Math. Soc., 2005, p. 453-469.
- [30] Moore, H. *Stable Minimal Hypersurfaces and Tangent Cone Singularities*. International Journal of Mathematics 10.03 (1999): 407-413.
- [31] Moser, J. *On Harnack's Theorem for Elliptic Differential Equations*. Communications on Pure and Applied Mathematics, vol. 14 (1961), 577–591.
- [32] Mumford, M. *The topology of normal singularities of an algebraic surface and a criterion for simplicity*. Inst. Hautes Études Sci. Publ. Math., vol. 9 (1961), 5–22.
- [33] Nguyen, N. and Si Tiep, D. *On the density at infinity of definable functions*. Preprint (2023) arXiv:2310.06324 [math.AG].

- [34] Pawłucki, W. *Quasi-regular boundary and Stokes' formula for a sub-analytic leaf*. In: Lawrynowicz J. (eds) Seminar on Deformations. Lecture Notes in Math., vol 1165. Berlin, Heidelberg: Springer. 1985.
- [35] Peterzil, Y. and Starchenko, S. *Complex analytic geometry and analytic-geometric categories*. J. reine angew. Math., vol. 626 (2009), 39–74.
- [36] Pokrovskii, A. V. *Removable singularities of solutions of the minimal surface equation*. (Russian) Funktsional. Anal. i Prilozhen. 39 (2005), no. 4, 62–68, 95–96; translation in Funct. Anal. Appl. 39 (2005), no. 4, 296–300.
- [37] Prill, D. *Cones in complex affine space are topologically singular*. Proc. of AMS, vol. 18 (1967), 178–182.
- [38] Sampaio, J. E. *Regularidade Lipschitz, invariância da multiplicidade e a geometria dos cones tangentes de conjuntos analíticos*. 2015. Tese (Doutorado em Matemática) - Universidade Federal do Ceará, Fortaleza, 2015.
- [39] Sampaio, J. E. *Bi-Lipschitz homeomorphic subanalytic sets have bi-Lipschitz homeomorphic tangent cones*. Selecta Mathematica: New Series, vol. 22 (2016), no. 2, 553–559.
- [40] Sampaio, J. E. *Globally subanalytic CMC surfaces in \mathbb{R}^3 with singularities*. Proceedings A of the Royal Society of Edinburgh, vol. 151 (2021), no. 1, 407–424.
- [41] Sampaio, J. E. *Multiplicity, regularity and Lipschitz Geometry of real analytic hypersurfaces*. Israel Journal of Mathematics, vol. 246 (2021), no. 1, 371–394.
- [42] Sampaio, J. E. *On Lipschitz Geometry at infinity of complex analytic sets*. Calculus of Variations and Partial Differential Equations, vol. 62 (2023), no. 2, article number 69.
- [43] Sampaio, J. E. *Local vs. global Lipschitz geometry*. Preprint (2023), arXiv:2305.11830 [math.MG].
- [44] Sampaio, J. E. and da Silva, E. C. *On bi-Lipschitz invariance and the uniqueness of tangent cones*. Journal of Singularities, vol. 25 (2022), 393–402.
- [45] Sampaio, J. E. and da Silva, E. C. *Classification of complex algebraic curves under blow-spherical equivalence*. Preprint (2023), arXiv:2302.02026 [math.AG].
- [46] Shen, Y.-B. and Zhu, X.-H. *On Stable Complete Minimal Hypersurfaces in $\mathbb{R}^n + 1$* . American Journal of Mathematics, vol. 120 (1998), no. 1, 103–116.
- [47] Shoen, R. and Simon, L. *Regularity of stable minimal hypersurfaces*. Comm. Pure Appl. Math., vol. 34, (1981), 741–797.
- [48] Shoen, R.; Simon, L. and Yau, S.-T. *Curvature estimates for minimal hypersurfaces*. Acta Math., vol. 134, (1975), 275–288.
- [49] Simon, L. *Asymptotic behaviour of minimal graphs over exterior domains*. Annales de l'I.H.P., section C, vol. 4 (1987), no. 3, 231–242.
- [50] Simons, J. *Minimal varieties in riemannian manifolds*. Annals of Mathematics, Second Series, 88 (1) (1968): 62–105.
- [51] Stasica, J. *Whitney property of sub-analytic sets*. Zeszyty Naukowe Uniw. Jag., vol. 23 (1982), 211–221.
- [52] Stoll, W. *The growth of the area of a transcendental analytic set. I*. Math. Ann., 156 (1964), no. 1, 47–78.

- [53] Stoll, W. *The growth of the area of a transcendental analytic set. II.* Math. Ann., vol. 156 (1964), 144–170.
- [54] Xin, Y.L. *Bernstein type theorems without graphic condition.* Asian J. Math., vol. 9 (2005), no. 1, 31–44.
- [55] Yau, S.-T. *Problem section.* In: Seminar on Differential Geometry. Ann. of Math. Stud., vol. 102, Princeton Univ. Press, 1982, 669-706.

(José Edson Sampaio) DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO CEARÁ, RUA CAMPUS DO PICI, S/N, BLOCO 914, PICI, 60440-900, FORTALEZA-CE, BRAZIL. E-MAIL: edsonsampaio@mat.ufc.br

EURIPEDES CARVALHO DA SILVA: DEPARTAMENTO DE MATEMÁTICA, INSTITUTO FEDERAL DE EDUCAÇÃO, CIÊNCIA E TECNOLOGIA DO CEARÁ, AV. PARQUE CENTRAL, 1315, DISTRITO INDUSTRIAL I, 61939-140, MARACANAÚ-CE, BRAZIL.
AND DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO CEARÁ, RUA CAMPUS DO PICI, S/N, BLOCO 914, PICI, 60440-900, FORTALEZA-CE, BRAZIL. E-MAIL: euripedes.carvalho@ifce.edu.br