

The minimum number of detours in graphs*

Xingzhi Zhan

Department of Mathematics, East China Normal University, Shanghai 200241, China

Abstract

A longest path in a graph is called a detour. It is easy to see that a connected graph of minimum degree at least 2 and order at least 4 has at least 4 detours. We prove that if the number of detours in such a graph of order at least 9 is odd, then it is at least 9, and this lower bound can be attained for every order. Thus the possibilities 3, 5 and 7 are excluded. Two open problems are posed.

Key words. Detour; longest path; minimum degree

Mathematics Subject Classification. 05C30, 05C35, 05C38

We consider finite simple graphs and use terminology and notations in [4]. Following Kapoor, Kronk, and Lick [3], we call a longest path in a graph G a *detour* of G . This concise term has now been widely used (e.g. [1] and [2]). The *order* of a graph is its number of vertices. We denote by $V(G)$ and $E(G)$ the vertex set and edge set of a graph G , respectively. For vertices x and y , an (x, y) -*path* is a path with endpoints x and y . We denote by $\delta(G)$ the minimum degree of a graph G , and by $N(x)$ the neighborhood of a vertex x . If u, v are two vertices on a path P , then $P[u, v]$ denotes the subpath of P with endpoints u and v . A basic fact about detours is that a detour of a connected graph G of order n has order at least $\min\{2\delta(G) + 1, n\}$.

It is easy to see that a connected graph of minimum degree at least 2 and order at least 4 has at least 4 detours. We will show that if the number of detours in such a graph of order at least 9 is odd, then it is at least 9, and this lower bound can be attained for

*E-mail address: zhan@math.ecnu.edu.cn

every order. Thus the possibilities 3, 5 and 7 are excluded. At the end we pose two open problems.

Notation. $f(G)$ denotes the number of detours in a graph G .

Theorem 1. *The minimum number of detours in a connected graph of minimum degree at least 2 and order at least 4 is 4.*

Proof. Let G be a connected graph of order at least 4 with $\delta(G) \geq 2$ and let $P : x_1, x_2, \dots, x_k$ be a detour of G . Then $N(x_1) \subseteq V(P)$ and $N(x_k) \subseteq V(P)$. Since $\delta(G) \geq 2$, x_1 has a neighbor x_i with $i \geq 3$ and x_k has a neighbor x_j with $j \leq k - 2$. If $i = k$ or $j = 1$, then G contains a k -cycle and we clearly have $f(G) \geq 4$. Next suppose $3 \leq i \leq k - 1$ and $2 \leq j \leq k - 2$.

Case 1. $i \leq j$.

G has at least the following four detours:

$$P, \quad P[x_1, x_j] \cup x_j x_k \cup P[x_k, x_{j+1}],$$

$$P[x_{i-1}, x_1] \cup x_1 x_i \cup P[x_i, x_k], \quad P[x_{i-1}, x_1] \cup x_1 x_i \cup P[x_i, x_j] \cup x_j x_k \cup P[x_k, x_{j+1}].$$

See Figure 1.




Fig. 1. The case $i \leq j$

Case 2. $i > j$.

G has at least the following six detours:

$$P, \quad P[x_1, x_j] \cup x_j x_k \cup P[x_k, x_{j+1}], \quad P[x_{i-1}, x_1] \cup x_1 x_i \cup P[x_i, x_k],$$

$$P[x_{j-1}, x_1] \cup x_1 x_i \cup P[x_i, x_k] \cup x_k x_j \cup P[x_j, x_{i-1}], \quad P[x_{i+1}, x_k] \cup x_k x_j \cup P[x_j, x_1] \cup x_1 x_i \cup P[x_i, x_{j+1}],$$

$$P[x_{j-1}, x_1] \cup x_1 x_i \cup P[x_i, x_j] \cup x_j x_k \cup P[x_k, x_{i+1}].$$

See Figure 2.




Fig. 2. The case $i > j$

This shows $f(G) \geq 4$. Conversely, for every order $n \geq 4$ we construct a graph G_n of order n with $\delta(G) \geq 2$ satisfying $f(G_n) = 4$. $G_4 = C_4$, the 4-cycle. G_5 is the bowtie, the graph consisting of two triangles sharing one vertex. G_6 consists of a triangle and a 4-cycle sharing one vertex. G_7 , G_8 and G_9 are depicted in (a), (b) and (c) of Figure 3, respectively.




Fig. 3. The graphs G_7 , G_8 and G_9

For $n \geq 10$, G_n is obtained from G_9 in (c) of Figure 3 by replacing the path x, z, y by an (x, y) -path of order $n - 6$. \square

Remark 1. Note that for $n \geq 7$, the graphs G_n in the above proof of Theorem 1 are 2-connected. Thus, if we replace “minimum degree at least 2” by “2-connected” in Theorem 1, we obtain the same conclusion for graphs of order at least 7.

Remark 2. In the above proof of Theorem 1, an edge e of P appears on at least four detours unless (1) $i \leq j$ and $e = x_{i-1}x_i$ or $e = x_jx_{j+1}$ or (2) $i = j + 1$ and $e = x_ix_j$.

Theorem 2. *Let G be a connected graph of minimum degree at least 2 and order at least 9. If $f(G)$ is an odd number, then $f(G) \geq 9$. Furthermore, the lower bound 9 can be attained for every order by both graphs of connectivity 1 and graphs of connectivity 2.*

Proof. We first prove that if $f(G)$ is an odd number, then $f(G) \geq 9$. It suffices to show that either $f(G) \geq 8$ or $f(G) = 4$ or $f(G) = 6$.

We make the following conventions: (1) For a positive integer r , “ r detours” means “ r pair-wise distinct detours”; (2) for an edge e of G and a detour D , we say that e appears on D if $e \in E(D)$.

Let $P : x_1, x_2, \dots, x_k$ be a detour of G . If there is another detour Q with $V(Q) \neq V(P)$, by the proof of Theorem 1, there are at least 4 detours with the same vertex set $V(P)$ and there are at least 4 detours with the same vertex set $V(Q)$. These detours are clearly distinct. Hence we have $f(G) \geq 8$. Next suppose that all detours of G have $V(P)$ as their vertex set.

Recall that an edge e of G is called a *chord* of a path R if the two endpoints of e lie in R but $e \notin E(R)$. A chord e of R is called an *inner chord* if both endpoints of e are internal vertices of R . Otherwise e is called a *boundary chord*. A detour D is called a *basic detour* if no inner chord of P is an edge of D ; otherwise D is called a *non-basic detour*.

Let the order of G be n . If G is hamiltonian, then $f(G) \geq n \geq 9$. Next assume that G is non-hamiltonian.

Since P is a detour, $N(x_1) \subseteq V(P)$ and $N(x_k) \subseteq V(P)$. The condition $\delta(G) \geq 2$ implies that x_1 has a neighbor x_i with $i \geq 3$ and x_k has a neighbor x_j with $j \leq k - 2$. If $i = k$ or $j = 1$, then G has a k -cycle C which contains P . Since P is a detour, C must be a Hamilton cycle, contradicting our assumption that G is non-hamiltonian. Hence $3 \leq i \leq k - 1$ and $2 \leq j \leq k - 2$. We distinguish two cases.

Case 1. Every detour of G is a basic detour.

We need consider only the boundary chords of P .

Subcase 1.1. P contains exactly two boundary chords.

As analyzed in the proof of Theorem 1, in this case $f(G) = 4$ or $f(G) = 6$.

Subcase 1.2. P contains exactly three boundary chords.

Without loss of generality, let x_1x_q be the third chord of P with $q \neq i$. Note that the two boundary chords x_1x_i and x_1x_q are in symmetric positions. If $q > i$ we may interchange the roles of x_1x_i and x_1x_q . Thus we may and do assume that $q < i$.

Suppose $i \leq j$. We have four basic detours not containing the edge x_1x_q . If $3 \leq q \leq i - 2$, we have exactly the following two detours containing the edge x_1x_q :

$$P[x_{q-1}, x_1] \cup x_1x_q \cup P[x_q, x_k], \quad P[x_{q-1}, x_1] \cup x_1x_q \cup P[x_q, x_j] \cup x_jx_k \cup P[x_k, x_{j+1}].$$

Hence $f(G) = 6$. If $q = i - 1$, we have exactly the following four detours containing the edge x_1x_q :

$$P[x_2, x_q] \cup x_qx_1 \cup x_1x_i \cup P[x_i, x_k], \quad P[x_2, x_q] \cup x_qx_1 \cup x_1x_i \cup P[x_i, x_j] \cup x_jx_k \cup P[x_k, x_{j+1}],$$

$$P[x_{q-1}, x_1] \cup x_1x_q \cup P[x_q, x_k], \quad P[x_{q-1}, x_1] \cup x_1x_q \cup P[x_q, x_j] \cup x_jx_k \cup P[x_k, x_{j+1}].$$

Hence $f(G) = 8$.

Suppose $i > j$. We have six basic detours not containing the edge x_1x_q . In this case it is easy to check that there are at least two detours containing the edge x_1x_q by considering the subgraph $P \cup x_1x_q \cup x_kx_j$. Thus $f(G) \geq 8$.

Subcase 1.3. P contains at least four boundary chords.

Based on Subcase 1.2, we deduce that $f(G) \geq 8$ in this case.

Case 2. G contains a non-basic detour.

Claim 1. Every edge in a detour appears on at least two detours.

This claim can be verified by checking the proof of Theorem 1, replacing P there by the detour in question. In fact, except for possible one or two edges, every edge in a detour appears on at least four distinct detours. See Remark 2 above.

Since G contains a non-basic detour, some inner chord e of P is an edge of a detour. By Claim 1, there are at least two detours containing e as an edge. Thus G has at least two non-basic detours. If $i > j$, we have six basic detours, and consequently $f(G) \geq 8$. By Subcases 1.2 and 1.3, if P contains at least three boundary chords, then G contains at least six basic detours. Again we have $f(G) \geq 8$. If an inner chord of P appears on at least four detours, then we have at least four non-basic detours. It follows that $f(G) \geq 8$.

It remains to treat the case when (1) $i \leq j$, (2) P contains exactly two boundary chords and (3) every inner chord of P appears on at most three detours. Next we make these three assumptions.

Let $D : y_1, y_2, \dots, y_k$ be a detour of G . Suppose y_c is a neighbor of y_1 and y_d is a neighbor of y_k . As in the proof of Theorem 1, there are four detours (if $c \leq d$) or six detours (if $c > d$) whose edges belong to $E(D) \cup \{y_1y_c, y_ky_d\}$. We denote by $\Psi(D)$ the set of these four or six detours according as $c \leq d$ or $c > d$. When we write $\Psi(D)$ we assume that the two boundary chords y_1y_c and y_ky_d have been prescribed.

Claim 2. If an inner chord h of P appears on a detour D such that $\Psi(D) \cap \Psi(P) \neq \phi$, then one of the two endpoints of h belongs to the set $\{x_{i-1}, x_{j+1}\}$.

Let $D = y_1, y_2, \dots, y_k$ with $h \in E(D)$. Suppose y_c is a neighbor of y_1 and y_d is a neighbor of y_k . Since h appears on at most three detours, by Remark 2 after the proof of Theorem 1, if $c > d$ we must have $c = d + 1$ and then D is contained in a cycle which must be a Hamilton cycle since D is a detour, contradicting our assumption that G is non-hamiltonian. Thus $c \leq d$ and then either $h = y_{c-1}y_c$ or $h = y_dy_{d+1}$. Note that since h is an inner chord of P , the two endpoints of h cannot be x_1 or x_k . Let $R \in \Psi(D) \cap \Psi(P)$. Then R does not contain h . Each of the two detours in $\Psi(D)$ not containing h has one endpoint which is an endpoint of h . Thus one endpoint v of R is an endpoint of h . Since the endpoints of the four detours in $\Psi(P)$ are $x_1, x_k, x_{i-1}, x_{j+1}$, we

deduce that $v \in \{x_1, x_k, x_{i-1}, x_{j+1}\}$ but $v \notin \{x_1, x_k\}$. Hence $v \in \{x_{i-1}, x_{j+1}\}$.

Subcase 2.1. G has a detour which contains at least two inner chords of P .

Let h and e be two inner chords of P that appear on one common detour. Consider the subgraph $G' = P \cup x_1x_i \cup x_kx_j \cup h \cup e$. The path $T = P[x_{i-1}, x_1] \cup x_1x_i \cup P[x_i, x_j] \cup x_jx_k \cup P[x_k, x_{j+1}]$ is a detour of G' with endpoints x_{i-1} and x_{j+1} , which is also a detour of G . Since we have assumed that G is non-hamiltonian, x_{i-1} and x_{j+1} are non-adjacent. By Claim 2, each of h and e has exactly one endpoint in the set $\{x_{i-1}, x_{j+1}\}$. Now in G' , the detour T has four boundary chords $x_{i-1}x_i$, $x_{j+1}x_j$, h and e . By Subcase 1.3 above (replacing P there by T), we obtain $f(G) \geq f(G') \geq 8$.

Subcase 2.2. Every non-basic detour contains exactly one inner chord of P .

Denote by Ω the set of the inner chords of P that appear on at least one detour. By the above Claim 1, if one inner chord of P appears on a detour, then there are at least two detours containing that chord. Thus, if $|\Omega| \geq 2$ then we have at least four non-basic detours, and consequently we have $f(G) \geq 8$. Next suppose $|\Omega| = 1$ and let $\Omega = \{x_sx_t\}$ with $2 \leq s \leq t-2$. Recall that we have assumed $i \leq j$. Using Claim 2, we deduce that $f(G) = 6$ if (1) $t = i-1$; (2) $s = j+1$; (3) $s = i-1$ and $i+2 \leq t \leq j$; (4) $i \leq s \leq j-2$ and $t = j+1$. In all other cases $f(G) \geq 8$. This completes the proof that if $f(G)$ is an odd number, then $f(G) \geq 9$.

Next for every integer $n \geq 9$ we construct a graph H_n of order n and connectivity 1 which contains exactly 9 detours. Every H_n is traceable. H_9, H_{10} and H_{11} are depicted in Figure 4.




Fig. 4. The graphs H_9, H_{10} and H_{11}

For $n \geq 11$, H_n is obtained from H_{10} by subdividing the edge $(4, 5)$ $n - 10$ times, i.e., replacing the edge $(4, 5)$ by a $(4, 5)$ -path of order $n - 8$. The 9 detours in H_9 are

$$\begin{aligned}
 & (0, 1, 2, 3, 4, 5, 6, 7, 8), \quad (0, 1, 2, 3, 4, 5, 6, 8, 7), \quad (0, 1, 2, 3, 4, 7, 8, 6, 5), \\
 & (1, 0, 2, 3, 4, 5, 6, 7, 8), \quad (1, 0, 2, 3, 4, 5, 6, 8, 7), \quad (1, 0, 2, 3, 4, 7, 8, 6, 5), \\
 & (3, 2, 0, 1, 4, 5, 6, 7, 8), \quad (3, 2, 0, 1, 4, 5, 6, 8, 7), \quad (3, 2, 0, 1, 4, 7, 8, 6, 5).
 \end{aligned}$$

Finally for every integer $n \geq 9$ we construct a graph M_n of order n and connectivity 2 which contains exactly 9 detours. Every M_n is traceable. M_9 and M_{10} are depicted in Figure 5.




Fig. 5. The graphs M_9 and M_{10}

For $n \geq 10$, M_n is obtained from M_9 by subdividing the edge $(7, 8)$ $n-9$ times. Observe that M_9 is obtained from H_9 in Figure 4 (a) by adding the edge $(2, 6)$, and any detour of M_9 cannot contain the edge $(2, 6)$. Hence M_9 and H_9 have the same set of detours, in particular, the same number of detours, i.e., 9. Note that each detour of M_9 contains the edge $(7, 8)$. Thus for every $n \geq 10$, M_n has the same number of detours as M_9 . \square

Finally we pose two problems. Recall that $f(G)$ denotes the number of detours in a graph G .

Problem 1. Let k and n be integers with $3 \leq k \leq n - 2$. Denote by $\Gamma(k, n)$ the set of connected graphs with minimum degree k and order n . Define

$$a(k, n) = \min\{f(G) \mid G \in \Gamma(k, n)\}.$$

Determine $a(k, n)$.

Problem 2. Let k, n and $\Gamma(k, n)$ be as in Problem 1. Define

$$b(k, n) = \min\{f(G) \mid G \in \Gamma(k, n) \text{ and } f(G) \text{ is odd}\}.$$

Determine $b(k, n)$.

Perhaps for sufficiently large orders n , $a(k, n)$ and $b(k, n)$ are independent of n .

Acknowledgement. This research was supported by the NSFC grant 12271170 and Science and Technology Commission of Shanghai Municipality grant 22DZ2229014.

References

- [1] L.W. Beineke, J.E. Dunbar and M. Frick, Detour-saturated graphs, *J. Graph Theory* 49(2005), 116–134.
- [2] G. Chartrand, G.L. Johns and S.L. Tian, Detour distance in graphs, *Quo vadis, graph theory?*, 127–136, *Ann. Discrete Math.*, 55, North-Holland, Amsterdam, 1993.
- [3] S.F. Kapoor, H.V. Kronk and D.R. Lick, On detours in graphs, *Canad. Math. Bull.* 11(1968), 195–201.
- [4] D.B. West, *Introduction to Graph Theory*, Prentice Hall, Inc., 1996.