

ON MONOMIAL DEFORMATIONS OF GENERALIZED DELSARTE POLYNOMIALS

ALAN ADOLPHSON AND STEVEN SPERBER

ABSTRACT. By a generalized Delsarte polynomial we mean a Laurent polynomial whose exponent vectors are linearly independent. We consider certain monomial deformations of generalized Delsarte polynomials and study their associated differential modules. We determine the solutions at the origin, which are all classical ${}_kF_{k-1}$ -hypergeometric functions with the variable raised to a power. Using standard changes of variable, we also obtain the solutions at infinity.

1. INTRODUCTION

Let $A = \{\mathbf{a}_j\}_{j=1}^m \subseteq \mathbb{Z}^n$ be a linearly independent set in \mathbb{R}^n . We assume there is a point $\mathbf{a}_0 \in \mathbb{Z}^n$ lying in the interior of the convex hull of the set A . This is equivalent to the existence of a relation

$$(1.1) \quad \ell_0 \mathbf{a}_0 = \sum_{j=1}^m \ell_j \mathbf{a}_j$$

where $\ell_0, \ell_1, \dots, \ell_m$ are positive integers having greatest common divisor equal to one satisfying

$$(1.2) \quad \ell_0 = \sum_{j=1}^m \ell_j.$$

Set $A_+ = A \cup \{\mathbf{a}_0\}$. We associate a differential module to the polynomial

$$(1.3) \quad f_\lambda = \sum_{j=1}^m \ell_j x^{\mathbf{a}_j} - \ell_0 \lambda x^{\mathbf{a}_0} \in \mathbb{Z}[\lambda][x_1^{\pm 1}, \dots, x_n^{\pm 1}],$$

where λ is an indeterminate. (In the special case of invertible polynomials, the ℓ_j , $j = 1, \dots, m$, are the dual weights and ℓ_0 is the dual degree.)

Let $V \subseteq \mathbb{R}^n$ be the subspace generated by A , so $\dim_{\mathbb{R}} V = m$. Let $V_{\mathbb{Z}} = V \cap \mathbb{Z}^n$ and let $C(A) \subseteq V$ be the real cone generated by A . Put $M = V_{\mathbb{Z}} \cap C(A)$ and let $S \subseteq \mathbb{C}[\lambda][x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ be the $\mathbb{C}[\lambda]$ -algebra generated by $\{x^u \mid u \in M\}$. The differential operators

$$D_i := x_i \frac{\partial}{\partial x_i} + x_i \frac{\partial f_\lambda}{\partial x_i} \left(= \frac{1}{\exp f_\lambda} \circ x_i \frac{\partial}{\partial x_i} \circ \exp f_\lambda \right)$$

for $i = 1, \dots, n$ operate on S and commute with the operator

$$D_\lambda := \frac{\partial}{\partial \lambda} - \ell_0 x^{\mathbf{a}_0} \left(= \frac{1}{\exp f_\lambda} \circ \frac{\partial}{\partial \lambda} \circ \exp f_\lambda \right),$$

Date: December 5, 2023.

which also operates on S . Let $\mathcal{D} := \mathbb{C}[\lambda]\langle\partial_\lambda\rangle$ be the ring of differential operators with coefficients in $\mathbb{C}[\lambda]$. Then both S and the $\mathbb{C}[\lambda]$ -module $\mathcal{W} := S/\sum_{i=1}^n D_i S$ are left \mathcal{D} -modules, where ∂_λ acts on S as D_λ . For $\xi \in S$, we also denote by ξ its image in the quotient \mathcal{W} . The context should prevent this abuse of notation from causing confusion.

Let $S' \subseteq \mathbb{C}(\lambda)[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ be the $\mathbb{C}(\lambda)$ -algebra generated by $\{x^u \mid u \in M\}$ and let $\mathcal{W}' = S'/\sum_{i=1}^n D_i S'$. Both S' and \mathcal{W}' are left \mathcal{D}' -modules, where $\mathcal{D}' = \mathbb{C}(\lambda)\langle\partial_\lambda\rangle$. For $\xi \in S'$, we also denote by ξ its image in the quotient \mathcal{W}' .

Our main result is an explicit construction of a basis for the finite-dimensional \mathbb{C} -vector space $\text{Hom}_{\mathcal{D}}(\mathcal{W}, \mathbb{C}[[\lambda]])$, the formal solutions of \mathcal{W} at the origin, in terms of classical ${}_kF_{k-1}$ -hypergeometric series for various k (Corollary 3.18 and Theorem 4.7). We apply this result to two related questions. First of all, the quotient \mathcal{W}' is a finite-dimensional $\mathbb{C}(\lambda)$ -vector space, so for any basis there is an associated Picard-Fuchs equation. We find the fundamental solution matrix for one such basis, again in terms of classical ${}_kF_{k-1}$ -hypergeometric series. As a second application, we compute the hypergeometric equations satisfied by elements $x^{\mathbf{b}}$ of this basis for \mathcal{W}' . Using standard changes of variables, we also explain how to find the solutions of \mathcal{W}' at infinity, which may involve $\log \lambda$.

Special cases of \mathcal{W} and \mathcal{W}' have been studied by various authors. The relation between these spaces and de Rham cohomology has been known since Dwork [8] and Katz [10] (see also Batyrev [6] and Adolphson-Sperber [3]). The problem of expressing solutions of the Picard-Fuchs equation for certain hypersurfaces in terms of ${}_kF_{k-1}$ -hypergeometric functions or for determining the differential operator satisfied by a differential form has also been studied by Kloosterman [12], Gähns [9], Miyatani [14], Doran et al. [7], and Negishi [15]. We give an example of the application of our results to such a computation in Section 11.

In the case of certain monomial deformations of diagonal hypersurfaces, Katz [11] makes a normalization of the equation $f_\lambda = 0$ that puts the singular points at roots of unity and leads to an expression of cohomology in terms of hypergeometric sheaves. Our normalization (1.1), a special case of [5, Equation (1.1)], generalizes Katz's normalization. This observation motivated us to combine Dwork's original method [8] with some ideas from [5] and [4, Section 3] to calculate the solutions.

We describe the type of hypergeometric equations that arise here. Let $\delta_x = x \frac{d}{dx}$ and consider the differential operator

$$(1.4) \quad (\delta_x + \beta_1 - 1) \cdots (\delta_x + \beta_k - 1) - x(\delta_x + \alpha_1) \cdots (\delta_x + \alpha_k).$$

Suppose that the β_j are distinct elements in $(0, 1]$ with $\beta_1 = 1$. One solution of the operator (1.4) at the origin is then

$$(1.5) \quad {}_kF_{k-1} \left(\begin{matrix} \alpha_1, \dots, \alpha_k \\ \beta_2, \dots, \beta_k \end{matrix}; x \right) := \sum_{s=0}^{\infty} \frac{(\alpha_1)_s \cdots (\alpha_k)_s}{(\beta_2)_s \cdots (\beta_k)_s (1)_s} x^s.$$

The other $k-1$ solutions are obtained by multiplying the series on the right-hand side by $x^{1-\beta_j}$, $j = 2, \dots, k$, and adding $1 - \beta_j$ to the quantity inside each Pochhammer symbol:

$$(1.6) \quad x^{1-\beta_j} {}_kF_{k-1} \left(\begin{matrix} \alpha_1 + 1 - \beta_j, \dots, \alpha_k + 1 - \beta_j \\ \beta_1 + 1 - \beta_j, \hat{}, \beta_k + 1 - \beta_j \end{matrix}; x \right),$$

where the symbol $\hat{}$ indicates that the term $\beta_j + 1 - \beta_j = 1$ has been omitted (as is usual in hypergeometric notation). In particular, one can read off the differential

operator (1.4) and the remaining solutions (1.6) from the single solution (1.5). The hypergeometric equations we encounter will be of this type with the variable x replaced by λ^{ℓ_0} .

We illustrate how the series (1.5) appear in our work. Let $P(A) \subseteq V$ be the parallelepiped defined by A :

$$P(A) = \left\{ \sum_{j=1}^m c_j \mathbf{a}_j \mid 0 \leq c_j < 1 \text{ for } j = 1, \dots, m \right\}.$$

Let $\mathbb{Z}A \subseteq V_{\mathbb{Z}}$ be the abelian group generated by A . The set $\mathcal{B} := V_{\mathbb{Z}} \cap P(A)$ is a complete set of coset representatives for $\mathbb{Z}A$ in $V_{\mathbb{Z}}$. Put $d = |\mathcal{B}| = [V_{\mathbb{Z}} : \mathbb{Z}A]$. Let $\mathbb{Z}A_+ \subseteq V_{\mathbb{Z}}$ be the subgroup generated by A_+ . Equation (1.2) implies that $[\mathbb{Z}A_+ : \mathbb{Z}A] = \ell_0$.

We show (Proposition 5.34) that the set $X^{\mathcal{B}} := \{x^{\mathbf{b}} \mid \mathbf{b} \in \mathcal{B}\}$ is a basis for \mathcal{W}' . In Section 6, we construct a fundamental solution matrix for the Picard-Fuchs equation associated to this basis. Its rows and columns are indexed by the set \mathcal{B} , and series of the type (1.5) with x replaced by λ^{ℓ_0} appear as the diagonal entries. Constant multiples of the series (1.6) with x replaced by λ^{ℓ_0} are the remaining nonzero entries in that row.

These diagonal entries are easy to describe. Let $\mathbf{b} \in \mathcal{B}$ and write $\mathbf{b} = \sum_{j=1}^m v_j \mathbf{a}_j$ with $v_j \in [0, 1) \cap \mathbb{Q}$ for $j = 1, \dots, m$. The (\mathbf{b}, \mathbf{b}) -entry in our solution matrix is

$$(1.7) \quad \sum_{s=0}^{\infty} \frac{\prod_{j=1}^m \binom{v_j}{\ell_j}_s \binom{v_j+1}{\ell_j}_s \cdots \binom{v_j+\ell_j-1}{\ell_j}_s}{\binom{1}{\ell_0}_s \binom{2}{\ell_0}_s \cdots \binom{\ell_0}{\ell_0}_s} \lambda^{s\ell_0}.$$

In general there is cancellation in this quotient. Let $\mathcal{B}_1 \subseteq \mathcal{B}$ be the subset consisting of those points that are congruent to \mathbf{b} modulo $\mathbb{Z}A_+$, a set of cardinality ℓ_0 , and let R be the number of elements of \mathcal{B}_1 that are interior to $P(A)$. If \mathbf{b} is an interior point of $P(A)$, then (1.7) will be an ${}_R F_{R-1}$ -hypergeometric function. If \mathbf{b} is a boundary point of $P(A)$, then (1.7) will be an ${}_{R+1} F_R$ -hypergeometric function. (If \mathbf{b} is a boundary point, then $v_j = 0$ for some j , so (1.7) equals the constant function 1.) This follows from the fact (Proposition 6.14) that in either case, if $\mathbf{b}' \in \mathcal{B}_1$ is a boundary point of $P(A)$ and $\mathbf{b} + t\mathbf{a}_0 \equiv \mathbf{b}' \pmod{\mathbb{Z}A}$ with $t \in \{1, 2, \dots, \ell_0 - 1\}$, then the factor $(1 - \frac{t}{\ell_0})_s$ cancels in (1.7). Using (1.6), one can then read off the other solutions of the corresponding Equation (1.4). Constant multiples of these solutions with the variable x replaced by λ^{ℓ_0} are the nonzero entries in the row corresponding to \mathbf{b} .

Proposition 6.14 describes the cancellation algorithm. Theorems 6.17 and 6.18 describe the entries in row \mathbf{b} . In Section 8 we show that the element $x^{\mathbf{b}} \in \mathcal{W}'$ satisfies the differential equation whose solutions are the nonzero entries in row \mathbf{b} .

Example. Take $f_{\lambda} = 3x_1^2 + 2x_2^3 + x_3^6 - 6\lambda x_1 x_2 x_3$. We have $\mathbf{a}_1 = (2, 0, 0)$, $\mathbf{a}_2 = (0, 3, 0)$, $\mathbf{a}_3 = (0, 0, 6)$, and $\mathbf{a}_0 = (1, 1, 1)$, giving the relation

$$6\mathbf{a}_0 = 3\mathbf{a}_1 + 2\mathbf{a}_2 + \mathbf{a}_3.$$

Thus $\ell_1 = 3$, $\ell_2 = 2$, $\ell_3 = 1$, $\ell_0 = 6$, and $|\mathcal{B}| = 36$. Let \mathcal{B}_1 be the subset of \mathcal{B} consisting of elements congruent to $(0, 0, 0)$ modulo $\mathbb{Z}A_+$:

$$\mathcal{B}_1 = \{(0, 0, 0), (1, 1, 1), (0, 2, 2), (1, 0, 3), (0, 1, 4), (1, 2, 5)\}.$$

In this case $R = 2$, so the row corresponding to $\mathbf{b} = (0, 0, 0)$ has three nonzero entries. We have $v_j = 0$ for $j = 1, 2, 3$, so (1.7) becomes

$$\sum_{s=0}^{\infty} \frac{\left(\frac{0}{3}\right)_s \left(\frac{1}{3}\right)_s \left(\frac{2}{3}\right)_s \left(\frac{0}{2}\right)_s \left(\frac{1}{2}\right)_s \left(\frac{0}{6}\right)_s}{\left(\frac{1}{6}\right)_s \left(\frac{2}{6}\right)_s \left(\frac{3}{6}\right)_s \left(\frac{4}{6}\right)_s \left(\frac{5}{6}\right)_s \left(\frac{6}{6}\right)_s} \lambda^{6s} = {}_3F_2(0, 0, 0; 1/6, 5/6; \lambda^6) = 1.$$

The cancellation in this quotient is determined by the boundary points (other than $(0, 0, 0)$) in \mathcal{B}_1 . We have

$$(0, 0, 0) + \begin{cases} 2(1, 1, 1) \\ 3(1, 1, 1) \\ 4(1, 1, 1) \end{cases} \equiv \begin{cases} (0, 2, 2) \\ (1, 0, 3) \\ (0, 1, 4) \end{cases} \pmod{\mathbb{Z}A}.$$

By Proposition 6.14(b), the terms that cancel are $(1 - \frac{t}{6})_s$ for $t = 2, 3, 4$.

The other two nonzero entries in this row are given, up to a constant multiple, by (1.6) with x replaced by λ^6 :

$$\lambda {}_3F_2(1/6, 1/6, 1/6; 1/3, 7/6; \lambda^6) \quad \text{and} \quad \lambda^5 {}_3F_2(5/6, 5/6, 5/6; 5/3, 11/6; \lambda^6).$$

These entries appear in columns $(1, 1, 1)$ and $(1, 2, 5)$, respectively.

Now consider the row corresponding to $\mathbf{b} = (1, 1, 1)$. We have $v_1 = 1/2$, $v_2 = 1/3$, and $v_3 = 1/6$, so by (1.7) the diagonal entry is

$$\sum_{s=0}^{\infty} \frac{\left(\frac{1}{6}\right)_s \left(\frac{3}{6}\right)_s \left(\frac{5}{6}\right)_s \left(\frac{1}{6}\right)_s \left(\frac{4}{6}\right)_s \left(\frac{1}{6}\right)_s}{\left(\frac{1}{6}\right)_s \left(\frac{2}{6}\right)_s \left(\frac{3}{6}\right)_s \left(\frac{4}{6}\right)_s \left(\frac{5}{6}\right)_s \left(\frac{6}{6}\right)_s} \lambda^{6s} = {}_2F_1(1/6, 1/6; 1/3; \lambda^6).$$

For the boundary points of \mathcal{B}_1 we have

$$(1, 1, 1) + \begin{cases} 1(1, 1, 1) \\ 2(1, 1, 1) \\ 3(1, 1, 1) \\ 5(1, 1, 1) \end{cases} \equiv \begin{cases} (0, 2, 2) \\ (1, 0, 3) \\ (0, 1, 4) \\ (0, 0, 0) \end{cases} \pmod{\mathbb{Z}A}.$$

By Proposition 6.14(a), the terms that cancel are $(1 - \frac{t}{6})_s$ for $t = 1, 2, 3, 5$.

From (1.6), the entry in the column corresponding to $(1, 2, 5)$ is, up to a constant multiple, $\lambda^4 {}_2F_1(5/6, 5/6; 5/3; \lambda^6)$.

2. SOLUTIONS AT THE ORIGIN

Let $S_0 \subseteq \mathbb{C}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ be the \mathbb{C} -algebra generated by $\{x^u \mid u \in M\}$. Let f_0 be the Laurent polynomial obtained from f_λ by setting $\lambda = 0$:

$$f_0 = \sum_{j=1}^m \ell_j x^{\mathbf{a}_j} \in S_0.$$

Put for $i = 1, \dots, n$

$$D_{i,0} = x_i \frac{\partial}{\partial x_i} + x_i \frac{\partial f_0}{\partial x_i},$$

operators on S_0 . Finally, let $\mathcal{W}_0 = S_0 / \sum_{i=1}^n D_{i,0} S_0$.

Theorem 2.1. *There is an isomorphism of \mathbb{C} -vector spaces*

$$(2.2) \quad \text{Hom}_{\mathcal{D}}(\mathcal{W}, \mathbb{C}[[\lambda]]) \cong \text{Hom}_{\mathbb{C}}(\mathcal{W}_0, \mathbb{C}).$$

The isomorphism (2.2) is a special case of well-known, more general results (see, for example, [1, Section 4]). We follow here an idea of Dwork [8, Section 5(c)] to make this isomorphism explicit, so that by computing a basis of the simpler right-hand side we obtain explicit formulas for a basis of the left-hand side. We begin with some general observations.

Let \mathcal{F} be a left \mathcal{D} -module and consider the $\mathbb{C}[\lambda]$ -module

$$R(\mathcal{F}) = \left\{ \sum_{u \in M} A_u x^{-u} \mid A_u \in \mathcal{F} \right\}.$$

Consider the pairing $R(\mathcal{F}) \times S \rightarrow \mathcal{F}$ defined by

$$\left\langle \sum_{u \in M} A_u x^{-u}, \sum_{u \in M} B_u x^u \right\rangle = \sum_{u \in M} B_u A_u,$$

where the B_u lie in $\mathbb{C}[\lambda]$ and the sum on the right-hand side is finite because the second sum on the left-hand side is. This pairing defines an isomorphism

$$(2.3) \quad \text{Hom}_{\mathbb{C}[\lambda]}(S, \mathcal{F}) \cong R(\mathcal{F}).$$

We next determine which elements of $R(\mathcal{F})$ correspond to elements of $\text{Hom}_{\mathcal{D}}(S, \mathcal{F})$. The condition to be satisfied is that for all $u \in M$ we have

$$\begin{aligned} \left\langle \sum_{v \in M} A_v x^{-v}, D_\lambda(x^u) \right\rangle &= \partial_\lambda \left\langle \sum_{v \in M} A_v x^{-v}, x^u \right\rangle \\ &= \partial_\lambda(A_u). \end{aligned}$$

But $D_\lambda(x^u) = -\ell_0 x^{u+\mathbf{a}_0}$, so the left-hand side is just $-\ell_0 A_{u+\mathbf{a}_0}$. Thus if we put

$$R^*(\mathcal{F}) = \left\{ \sum_{u \in M} A_u x^{-u} \mid A_u \in \mathcal{F} \text{ and } \partial_\lambda(A_u) = -\ell_0 A_{u+\mathbf{a}_0} \text{ for all } u \in M \right\},$$

then

$$(2.4) \quad \text{Hom}_{\mathcal{D}}(S, \mathcal{F}) \cong R^*(\mathcal{F}).$$

We want to compute $\text{Hom}_{\mathcal{D}}(\mathcal{W}, \mathcal{F})$, which is the intersection of $\text{Hom}_{\mathcal{D}}(S, \mathcal{F})$ and $\text{Hom}_{\mathbb{C}[\lambda]}(\mathcal{W}, \mathcal{F}) (\subseteq \text{Hom}_{\mathbb{C}[\lambda]}(S, \mathcal{F}))$. We compute $\text{Hom}_{\mathbb{C}[\lambda]}(\mathcal{W}, \mathcal{F})$. By (2.3), these are the elements of $R(\mathcal{F})$ that vanish on $\sum_{i=1}^n D_i S$. Let

$$\xi = \sum_{u \in M} A_u x^{-u} \in R(\mathcal{F}).$$

Then ξ vanishes on $\sum_{i=1}^n D_i S$ if and only if it vanishes on $D_i(x^v)$ for all $i = 1, \dots, n$ and all $v \in M$. So the condition to be satisfied is (where $\mathbf{a}_j = (a_{1j}, \dots, a_{nj})$)

$$(2.5) \quad \begin{aligned} \langle \xi, D_i(x^v) \rangle &= \left\langle \xi, v_i x^v - \ell_0 a_{i0} \lambda x^{v+\mathbf{a}_0} + \sum_{j=1}^m \ell_j a_{ij} x^{v+\mathbf{a}_j} \right\rangle \\ &= v_i A_v - \ell_0 a_{i0} \lambda A_{v+\mathbf{a}_0} + \sum_{j=1}^m \ell_j a_{ij} A_{v+\mathbf{a}_j} \\ &= 0. \end{aligned}$$

Let γ_- be defined by $\mathbb{C}[\lambda]$ -linearity and the condition

$$\gamma_-(x^{-u}) = \begin{cases} x^{-u} & \text{if } u \in M, \\ 0 & \text{if } u \notin M. \end{cases}$$

Let D_i^* be the operator on $R(\mathcal{F})$ defined for $i = 1, \dots, n$ by

$$D_i^* = \gamma_- \circ \left(-x_i \frac{\partial}{\partial x_i} - \ell_0 a_{i0} \lambda x^{\mathbf{a}_0} + \sum_{j=1}^m \ell_j a_{ij} x^{\mathbf{a}_j} \right).$$

For $v \in M$ the coefficient of x^{-v} in $D_i^*(\xi)$ is

$$v_i A_v - \ell_0 a_{i0} \lambda A_{v+\mathbf{a}_0} + \sum_{j=1}^m \ell_j a_{ij} A_{v+\mathbf{a}_j}.$$

It now follows from (2.5) that

$$\mathrm{Hom}_{\mathbb{C}[\lambda]}(\mathcal{W}, \mathcal{F}) = \{ \xi \in R(\mathcal{F}) \mid D_i^*(\xi) = 0 \text{ for } i = 1, \dots, n \}.$$

Define

$$\mathcal{K}(\mathcal{F}) = \left\{ \xi = \sum_{u \in M} A_u x^{-u} \mid A_u \in \mathcal{F} \text{ for } u \in M, D_i^*(\xi) = 0 \text{ for } i = 1, \dots, n, \right. \\ \left. \text{and } \partial_\lambda(A_u) = -\ell_0 A_{u+\mathbf{a}_0} \text{ for all } u \in M \right\}.$$

We have proved the following result.

Proposition 2.6. $\mathcal{K}(\mathcal{F}) \cong \mathrm{Hom}_{\mathcal{D}}(\mathcal{W}, \mathcal{F})$.

We analyze similarly $\mathrm{Hom}_{\mathbb{C}}(\mathcal{W}_0, Y)$ for an arbitrary \mathbb{C} -vector space Y . Let

$$R_0(Y) = \left\{ \sum_{u \in M} A_u x^{-u} \mid A_u \in Y \right\}.$$

The pairing $R_0(Y) \times S_0 \rightarrow Y$ defined by

$$\left\langle \sum_{u \in M} A_u x^{-u}, \sum_{u \in M} B_u x^u \right\rangle = \sum_{u \in M} B_u A_u,$$

where the B_u lie in \mathbb{C} , defines an isomorphism

$$(2.7) \quad \mathrm{Hom}_{\mathbb{C}}(S_0, Y) \cong R_0(Y).$$

Let $\xi = \sum_{u \in M} A_u x^{-u} \in R_0(Y)$. Then ξ vanishes on $\sum_{i=1}^n D_{i,0} S_0$ if and only if it vanishes on $D_{i,0}(x^v)$ for all $i = 1, \dots, n$ and all $v \in M$. The condition to be satisfied is

$$(2.8) \quad \langle \xi, D_{i,0}(x^v) \rangle = \left\langle \xi, v_i x^v + \sum_{j=1}^m \ell_j a_{ij} x^{v+\mathbf{a}_j} \right\rangle \\ = v_i A_v + \sum_{j=1}^m \ell_j a_{ij} A_{v+\mathbf{a}_j} \\ = 0.$$

Let $D_{i,0}^*$ be the operator on $R_0(Y)$ defined for $i = 1, \dots, n$ by

$$D_{i,0}^* = \gamma_- \circ \left(-x_i \frac{\partial}{\partial x_i} + \sum_{j=1}^m \ell_j a_{ij} x^{\mathbf{a}_j} \right).$$

For $v \in M$, the coefficient of x^{-v} in $D_{i,0}^*(\xi)$ is

$$v_i A_v + \sum_{j=1}^m \ell_j a_{ij} A_{v+\mathbf{a}_j}.$$

If we put

$$\mathcal{K}_0(Y) = \left\{ \xi = \sum_{u \in M} A_u x^{-u} \mid A_u \in Y \text{ for } u \in M \text{ and } D_{i,0}^*(\xi) = 0 \text{ for } i = 1, \dots, n \right\},$$

then from Equation (2.8) we get the following result.

Proposition 2.9. $\mathcal{K}_0(Y) \cong \text{Hom}_{\mathbb{C}}(\mathcal{W}_0, Y)$.

Proof of Theorem 2.1. By Propositions 2.6 and 2.9 we need to prove that

$$(2.10) \quad \mathcal{K}(\mathbb{C}[[\lambda]]) \cong \mathcal{K}_0(\mathbb{C}).$$

For $\xi = \sum_{u \in M} c_u x^{-u} \in \mathcal{K}_0(\mathbb{C})$, define

$$(2.11) \quad \phi(\xi) = \gamma_-(\exp(-\ell_0 \lambda x^{\mathbf{a}_0}) \xi) =: \sum_{u \in M} A_u(\lambda) x^{-u},$$

where the $A_u(\lambda)$ lie in $\mathbb{C}[[\lambda]]$. We claim that ϕ is an isomorphism from $\mathcal{K}_0(\mathbb{C})$ onto $\mathcal{K}(\mathbb{C}[[\lambda]])$.

We first show that $\phi(\xi) \in \mathcal{K}(\mathbb{C}[[\lambda]])$. We have

$$\partial_\lambda(\phi(\xi)) = \sum_{u \in M} (\partial_\lambda A_u)(\lambda) x^{-u}.$$

Since ∂_λ commutes with γ_- we also have

$$\begin{aligned} \partial_\lambda(\phi(\xi)) &= \gamma_-(-\ell_0 x^{\mathbf{a}_0} \exp(-\ell_0 \lambda x^{\mathbf{a}_0}) \xi) \\ &= \gamma_-(-\ell_0 x^{\mathbf{a}_0} \phi(\xi)) \\ &= -\ell_0 \sum_{u \in M} A_{u+\mathbf{a}_0}(\lambda) x^{-u}. \end{aligned}$$

It follows that $(\partial_\lambda A_u)(\lambda) = -\ell_0 A_{u+\mathbf{a}_0}(\lambda)$ for all $u \in M$.

We also need to check that $D_i^*(\phi(\xi)) = 0$ for $i = 1, \dots, n$. First note that

$$x_i \frac{\partial}{\partial x_i} \left(\exp(-\ell_0 \lambda x^{\mathbf{a}_0}) \xi \right) = \exp(-\ell_0 \lambda x^{\mathbf{a}_0}) \left(x_i \frac{\partial \xi}{\partial x_i} - \ell_0 \lambda a_{i0} x^{\mathbf{a}_0} \right).$$

We thus have

$$\begin{aligned} D_i^*(\phi(\xi)) &= \\ &= \gamma_- \left(\exp(-\ell_0 \lambda x^{\mathbf{a}_0}) \left(\left(-x_i \frac{\partial \xi}{\partial x_i} + \ell_0 \lambda a_{i0} x^{\mathbf{a}_0} \right) - \ell_0 a_{i0} \lambda x^{\mathbf{a}_0} \xi + \sum_{j=1}^m \ell_j a_{ij} x^{\mathbf{a}_j} \xi \right) \right). \end{aligned}$$

Hence

$$\begin{aligned} D_i^*(\phi(\xi)) &= \gamma_- \left(\exp(-\ell_0 \lambda x^{\mathbf{a}_0}) \left(-x_i \frac{\partial \xi}{\partial x_i} + \sum_{j=1}^m \ell_j a_{ij} x^{\mathbf{a}_j} \xi \right) \right) \\ &= \gamma_- \left(\exp(-\ell_0 \lambda x^{\mathbf{a}_0}) D_{i,0}^*(\xi) \right) \\ &= 0 \end{aligned}$$

since $\xi \in \mathcal{K}_0(\mathbb{C})$.

For $\sum_{u \in M} A_u(\lambda)x^{-u} \in \mathcal{K}(\mathbb{C}[[\lambda]])$ define

$$\phi' \left(\sum_{u \in M} A_u(\lambda)x^{-u} \right) = \sum_{u \in M} A_u(0)x^{-u},$$

i. e., we evaluate all the $A_u(\lambda)$ at $\lambda = 0$. We claim that ϕ' is the inverse of ϕ . It is straightforward to check that

$$D_{i,0}^* \left(\sum_{u \in M} A_u(0)x^{-u} \right) = 0$$

for $i = 1, \dots, n$, so ϕ' maps $\mathcal{K}(\mathbb{C}[[\lambda]])$ into $\mathcal{K}_0(\mathbb{C})$. It follows from the definition of ϕ that if $\xi = \sum_{u \in M} c_u x^{-u} \in \mathcal{K}_0(\mathbb{C})$, then $\phi'(\phi(\xi)) = \xi$, so ϕ' is onto. The map ϕ' is also injective because $\sum_{u \in M} A_u(\lambda)x^{-u} \in \mathcal{K}(\mathbb{C}[[\lambda]])$ is determined by its constant terms $\sum_{u \in M} A_u(0)x^{-u}$. To see this, suppose that c_k is the coefficient of λ^k in the series $A_u(\lambda)$. The condition $(\partial_\lambda A_u)(\lambda) = -\ell_0 A_{u+\mathbf{a}_0}(\lambda)$ for all $u \in M$ implies that the constant term of $(-\ell_0)^k A_{u+k\mathbf{a}_0}(\lambda)$ equals $k!c_k$. So if the constant terms $A_u(0)$ vanish for all $u \in M$ then the series $A_u(\lambda)$ vanish for all $u \in M$. \square

3. BASIS FOR $\text{Hom}_{\mathcal{D}}(\mathcal{W}, \mathbb{C}[[\lambda]])$

In this section we find a basis for \mathcal{W}_0 , compute the dual basis of $\text{Hom}_{\mathbb{C}}(\mathcal{W}_0, \mathbb{C})$, and use the isomorphism ϕ of the previous section to get a basis for $\text{Hom}_{\mathcal{D}}(\mathcal{W}, \mathbb{C}[[\lambda]])$. Let \mathcal{B} be as in Section 1 and let $X^{\mathcal{B}} = \{x^{\mathbf{b}} \mid \mathbf{b} \in \mathcal{B}\}$.

Let $\alpha = (a_{ij})$ ($i = 1, \dots, n$, $j = 1, \dots, m$) be the $n \times m$ matrix whose columns are the \mathbf{a}_j . Since the \mathbf{a}_j are linearly independent, there exists an $n \times n$ matrix $\beta = (\beta_{ij})$ of rational numbers such that

$$(3.1) \quad \beta\alpha = \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \dots & \vdots & \vdots \\ 0 & 0 & \dots & 0 & 1 \\ 0 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & 0 & 0 \end{pmatrix}.$$

We define differential operators $\tilde{D}_{i,0}$, $i = 1, \dots, n$, on S_0 by the formula

$$(3.2) \quad \begin{pmatrix} \tilde{D}_{1,0} \\ \vdots \\ \tilde{D}_{n,0} \end{pmatrix} = \beta \begin{pmatrix} D_{1,0} \\ \vdots \\ D_{n,0} \end{pmatrix} = \begin{pmatrix} \sum_{j=1}^n \beta_{1j} D_{j,0} \\ \vdots \\ \sum_{j=1}^n \beta_{nj} D_{j,0} \end{pmatrix}.$$

Lemma 3.3. *Let $u \in M$ and write $u = (u_1, \dots, u_n) = \sum_{j=1}^m u'_j \mathbf{a}_j$ with $u'_j \in \mathbb{Q}$ for $j = 1, \dots, m$. We have*

$$\tilde{D}_{j,0}(x^u) = \begin{cases} u'_j x^u + \ell_j x^{u+\mathbf{a}_j} & \text{for } j = 1, \dots, m, \\ 0 & \text{for } j = m+1, \dots, n. \end{cases}$$

Proof. We note that

$$(3.4) \quad \begin{pmatrix} x_1 \frac{\partial f_0}{\partial x_1} \\ \vdots \\ x_n \frac{\partial f_0}{\partial x_n} \end{pmatrix} = \boldsymbol{\alpha} \begin{pmatrix} \ell_1 x^{\mathbf{a}_1} \\ \vdots \\ \ell_m x^{\mathbf{a}_m} \end{pmatrix}$$

and

$$(3.5) \quad \boldsymbol{\alpha} \begin{pmatrix} u'_1 \\ \vdots \\ u'_m \end{pmatrix} = \begin{pmatrix} u_1 \\ \vdots \\ u_n \end{pmatrix}.$$

From the definition we have

$$(3.6) \quad \begin{aligned} \begin{pmatrix} \tilde{D}_{1,0} \\ \vdots \\ \tilde{D}_{n,0} \end{pmatrix} (x^u) &= \boldsymbol{\beta} \begin{pmatrix} x_1 \frac{\partial}{\partial x_1} + x_1 \frac{\partial f_0}{\partial x_1} \\ \vdots \\ x_n \frac{\partial}{\partial x_n} + x_n \frac{\partial f_0}{\partial x_n} \end{pmatrix} (x^u) \\ &= \boldsymbol{\beta} \begin{pmatrix} u_1 \\ \vdots \\ u_n \end{pmatrix} x^u + \boldsymbol{\beta} \begin{pmatrix} x_1 \frac{\partial f_0}{\partial x_1} \\ \vdots \\ x_n \frac{\partial f_0}{\partial x_n} \end{pmatrix} x^u \\ &= \boldsymbol{\beta} \boldsymbol{\alpha} \begin{pmatrix} u'_1 \\ \vdots \\ u'_m \end{pmatrix} x^u + \boldsymbol{\beta} \boldsymbol{\alpha} \begin{pmatrix} \ell_1 x^{\mathbf{a}_1} \\ \vdots \\ \ell_m x^{\mathbf{a}_m} \end{pmatrix} x^u \\ &= \begin{pmatrix} u'_1 x^u + \ell_1 x^{u+\mathbf{a}_1} \\ \vdots \\ u'_m x^u + \ell_m x^{u+\mathbf{a}_m} \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \end{aligned}$$

where the third equality follows from (3.4) and (3.5) and the fourth equality follows from (3.1). \square

Since $\boldsymbol{\beta}$ is invertible, Equation (3.2) and Lemma 3.3 imply the following result.

Corollary 3.7. *We have $\mathcal{W}_0 = S_0 / \sum_{j=1}^m \tilde{D}_{j,0} S_0$.*

The following result is clear.

Lemma 3.8. *The set $X^{\mathcal{B}}$ is a basis for $S_0 / \sum_{j=1}^m x^{\mathbf{a}_j} S_0$.*

The set A_+ lies on a hyperplane in V , which allows us to define a grading on the spaces S , S' , and S_0 . For $u = (u_1, \dots, u_n) \in M$ we write $u = \sum_{j=1}^m u'_j \mathbf{a}_j$ and define $\deg x^u = \sum_{j=1}^m u'_j$. Equivalently, if $\sum_{i=1}^n h_i y_i = 1$ is the equation of a hyperplane containing A_+ , then $\deg x^u = \sum_{i=1}^n h_i u_i$. The degree is a surjective map from $\{x^u \mid u \in M\}$ onto the set $(q_0)^{-1} \mathbb{Z}_{\geq 0}$ for some positive integer q_0 . In the study of invertible polynomials, the integer q_0 is the degree of the polynomial and the $q_0 h_i$, $i = 1, \dots, n$, are the weights.

Notation: Let L be a module (over some ring) and let $\{\phi_j\}_{j=1}^m$ be commuting endomorphisms of L . We denote by $\text{Kos}^\bullet(L, \{\phi_j\}_{j=1}^m)$ the associated (cohomological)

Koszul complex. More precisely, let the e_i , $i = 1, \dots, m$ be formal symbols and define

$$\mathrm{Kos}^k(L, \{\phi\}_{j=1}^m) = \bigoplus_{1 \leq j_1 < \dots < j_k \leq m} L e_{j_1} \wedge \dots \wedge e_{j_k}.$$

The boundary map $\nabla : \mathrm{Kos}^k(L, \{\phi\}_{j=1}^m) \rightarrow \mathrm{Kos}^{k+1}(L, \{\phi\}_{j=1}^m)$ is given for $\xi \in L$ by

$$\nabla(\xi e_{j_1} \wedge \dots \wedge e_{j_k}) = \left(\sum_{j=1}^m \phi_j(\xi) e_j \right) \wedge e_{j_1} \wedge \dots \wedge e_{j_k}.$$

Proposition 3.9. *The set $X^{\mathcal{B}}$ is a basis for \mathcal{W}_0 .*

Proof. Consider the Koszul complex $\mathrm{Kos}^\bullet(S_0, \{\tilde{D}_{j,0}\}_{j=1}^m)$ defined by the $\tilde{D}_{j,0}$. The above grading makes it into a filtered complex for which

$$H^m(\mathrm{Kos}^\bullet(S_0, \{\tilde{D}_{j,0}\}_{j=1}^m)) = \mathcal{W}_0.$$

By Lemma 3.3 the associated graded complex is $\mathrm{Kos}^\bullet(S_0, \{\ell_j x^{\mathbf{a}_j}\}_{j=1}^m)$. Since the set $\{x^{\mathbf{a}_j}\}_{j=1}^m$ is a regular sequence on S_0 , we have $H^i(\mathrm{Kos}^\bullet(S_0, \{\ell_j x^{\mathbf{a}_j}\}_{j=1}^m)) = 0$ for $0 \leq i < m$. It follows that $\mathcal{W}_0 \cong H^m(\mathrm{Kos}^\bullet(S_0, \{\ell_j x^{\mathbf{a}_j}\}_{j=1}^m))$. But

$$H^m(\mathrm{Kos}^\bullet(S_0, \{\ell_j x^{\mathbf{a}_j}\}_{j=1}^m)) = S_0 / \sum_{j=1}^m x^{\mathbf{a}_j} S_0,$$

so Lemma 3.8 implies the proposition. \square

We use the following notation, which we find simpler for some formulas than the classical Pochhammer notation. Later we switch to Pochhammer notation to make the connection with hypergeometric series. Define for $z \in \mathbb{C}$

$$[z]_l = \begin{cases} 1 & \text{if } l = 0, \\ \frac{1}{(z+1)(z+2)\dots(z+l)} & \text{if } l > 0 \text{ and } z \notin \{-1, -2, \dots, -l\}, \\ z(z-1)\dots(z+l-1) & \text{if } l < 0. \end{cases}$$

The relation to the Pochhammer symbol $(z)_l$ is that $[z]_l$ is defined if and only if $(-z)_{-l}$ is defined, in which case

$$(3.10) \quad [z]_l = (-1)^l (-z)_{-l}.$$

We construct the dual basis for $\mathrm{Hom}_{\mathbb{C}}(\mathcal{W}_0, \mathbb{C})$. Fix $\mathbf{b} \in \mathcal{B}$. Then

$$(3.11) \quad \mathbf{b} = \sum_{j=1}^m v_j \mathbf{a}_j$$

for unique $v_1, \dots, v_m \in [0, 1)$. For $j = 1, \dots, m$ we define series

$$(3.12) \quad g_j(t) = \sum_{s \in \mathbb{Z}_{\leq 0}} [-v_j]_s t^{-v_j+s}.$$

Set

$$(3.13) \quad g_{\mathbf{b}} = \prod_{j=1}^m \ell_j^{v_j} g_j(\ell_j x^{\mathbf{a}_j}).$$

A straightforward calculation establishes the following result.

Lemma 3.14. *We have $g_{\mathbf{b}} = \sum_{u \in M} g_u^{(\mathbf{b})} x^{-u}$, where $g_u^{(\mathbf{b})} = \prod_{j=1}^m ([-v_j]_{s_j} \ell_j^{s_j})$ if there exist $\{s_j\}_{j=1}^m \subseteq \mathbb{Z}_{\leq 0}$, necessarily unique, such that $-u = \sum_{j=1}^m (-v_j + s_j) \mathbf{a}_j$ and $g_u^{(\mathbf{b})} = 0$ otherwise.*

Proposition 3.15. *The set $\{g_{\mathbf{b}}\}_{\mathbf{b} \in \mathcal{B}}$ is the basis of $\text{Hom}_{\mathbb{C}}(\mathcal{W}_0, \mathbb{C})$ dual to the basis $X^{\mathcal{B}}$ of \mathcal{W}_0 .*

Proof. We first check that the $g_{\mathbf{b}}$ lie in $\text{Hom}_{\mathbb{C}}(\mathcal{W}_0, \mathbb{C})$. By Proposition 2.9 we need to check that $D_{i,0}^*(g_{\mathbf{b}}) = 0$ for $i = 1, \dots, n$. First note that

$$\begin{aligned} x_i \frac{\partial g_{\mathbf{b}}}{\partial x_i} &= \sum_{k=1}^m \left(\prod_{\substack{j=1 \\ j \neq k}}^m \ell_j^{v_j} g_j(\ell_j x^{\mathbf{a}_j}) \right) x_i \frac{\partial}{\partial x_i} (\ell_k^{v_k} g_k(\ell_k x^{\mathbf{a}_k})) \\ &= \sum_{k=1}^m \left(\prod_{\substack{j=1 \\ j \neq k}}^m \ell_j^{v_j} g_j(\ell_j x^{\mathbf{a}_j}) \right) (\ell_k^{v_k+1} a_{ik} x^{\mathbf{a}_k}) g'_k(\ell_k x^{\mathbf{a}_k}). \end{aligned}$$

Since $g'_k(t) = g_k(t) - t^{-v_j}$, this reduces to

$$(3.16) \quad x_i \frac{\partial g_{\mathbf{b}}}{\partial x_i} = \sum_{k=1}^m \left(\prod_{\substack{j=1 \\ j \neq k}}^m \ell_j^{v_j} g_j(\ell_j x^{\mathbf{a}_j}) \right) (\ell_k^{v_k+1} a_{ik} x^{\mathbf{a}_k} g_k(\ell_k x^{\mathbf{a}_k}) - a_{ik} \ell_k x^{(-v_k+1)\mathbf{a}_k}).$$

Because $(-v_k + 1) > 0$, no term in the product $x^{(-v_k+1)\mathbf{a}_k} \prod_{\substack{j=1 \\ j \neq k}}^m g_j(\ell_j x^{\mathbf{a}_j})$ contains a term x^{-u} with $u \in M$. It follows that

$$\begin{aligned} \gamma_- \left(x_i \frac{\partial g_{\mathbf{b}}}{\partial x_i} \right) &= \sum_{k=1}^m \left(\prod_{\substack{j=1 \\ j \neq k}}^m \ell_j^{v_j} g_j(\ell_j x^{\mathbf{a}_j}) \right) (\ell_k^{v_k+1} a_{ik} x^{\mathbf{a}_k} g_k(\ell_k x^{\mathbf{a}_k})) \\ &= \gamma_- \left(\sum_{k=1}^m \ell_k a_{ik} x^{\mathbf{a}_k} g_{\mathbf{b}} \right) = \gamma_- \left(x_i \frac{\partial f_0}{\partial x_i} g_{\mathbf{b}} \right). \end{aligned}$$

This equation implies that $D_{i,0}^*(g_{\mathbf{b}}) = 0$ for all i .

We also need to check that if $\mathbf{b}, \mathbf{b}' \in \mathcal{B}$, then

$$\langle g_{\mathbf{b}}, x^{\mathbf{b}'} \rangle = \begin{cases} 1 & \text{if } \mathbf{b} = \mathbf{b}', \\ 0 & \text{if } \mathbf{b} \neq \mathbf{b}'. \end{cases}$$

It follows from Lemma 3.14 that

$$\langle g_{\mathbf{b}}, x^{\mathbf{b}'} \rangle = g_{\mathbf{b}'}^{(\mathbf{b})} = \prod_{j=1}^m ([-v_j]_{s_j} \ell_j^{s_j}),$$

where the v_j are determined by the condition $\sum_{j=1}^m v_j \mathbf{a}_j = \mathbf{b}$ and the s_j are determined by the condition $-\mathbf{b}' = \sum_{j=1}^m (-v_j + s_j) \mathbf{a}_j$. If the second condition has a solution $\{s_j\}_{j=1}^m \subseteq \mathbb{Z}_{\leq 0}$, then $\mathbf{b} \equiv \mathbf{b}' \pmod{\mathbb{Z}A}$. But this can happen only if $\mathbf{b} = \mathbf{b}'$, so $g_{\mathbf{b}'}^{(\mathbf{b})} = 0$ if $\mathbf{b} \neq \mathbf{b}'$. And if $\mathbf{b} = \mathbf{b}'$, then $s_j = 0$ for all j so $g_{\mathbf{b}'}^{(\mathbf{b})} = 1$. \square

We can now apply the isomorphism of Theorem 2.1. Define

$$(3.17) \quad \xi_{\mathbf{b}} = \phi(g_{\mathbf{b}}) = \gamma_- (\exp(-\ell_0 \lambda x^{\mathbf{a}_0}) g_{\mathbf{b}}).$$

By Theorem 2.1, we have the following result.

Corollary 3.18. *The set $\{\xi_{\mathbf{b}}\}_{\mathbf{b} \in \mathcal{B}}$ is a basis for $\text{Hom}_{\mathcal{D}}(\mathcal{W}, \mathbb{C}[[\lambda]])$.*

Define $G_u^{(\mathbf{b})}(\lambda)$ by the formula

$$(3.19) \quad \xi_{\mathbf{b}} = \sum_{u \in M} G_u^{(\mathbf{b})}(\lambda) x^{-u}.$$

From Lemma 3.14, we have the following explicit formula, which will be simplified later.

$$(3.20) \quad \begin{aligned} G_u^{(\mathbf{b})}(\lambda) &= \sum_{\substack{w \in M, s \in \mathbb{Z}_{\geq 0} \\ -w + s\mathbf{a}_0 = -u}} g_w^{(\mathbf{b})} \frac{(-\ell_0 \lambda)^s}{s!} \\ &= \sum_{\substack{s_0 \in \mathbb{Z}_{\geq 0}, s_1, \dots, s_m \in \mathbb{Z}_{\leq 0} \\ s_0 \mathbf{a}_0 + \sum_{j=1}^m (-v_j + s_j) \mathbf{a}_j = -u}} \left(\prod_{j=1}^m [-v_j]_{s_j} \prod_{j=1}^m \ell_j^{s_j} \right) \frac{(-\ell_0)^{s_0}}{s_0!} \lambda^{s_0}. \end{aligned}$$

4. STRUCTURE OF SOLUTIONS

In this section we analyze Equation (3.20) to describe the $\xi_{\mathbf{b}}$ in more detail. We first make a general observation.

There is a direct sum decomposition of \mathcal{W} as \mathcal{D} -module. Put $N = [V_{\mathbb{Z}} : \mathbb{Z}A_+]$. We have

$$N = \frac{[V_{\mathbb{Z}} : \mathbb{Z}A]}{[\mathbb{Z}A_+ : \mathbb{Z}A]} = \frac{d}{\ell_0}.$$

Let \mathcal{C}_k , $k = 1, \dots, N$ be the cosets of $\mathbb{Z}A_+$ in $V_{\mathbb{Z}}$ and put $M_k = M \cap \mathcal{C}_k$. Let S_k be the $\mathbb{C}[[\lambda]]$ -module generated by the x^u with $u \in M_k$. The S_k are disjoint, stable under the action of the D_i and D_{λ} , and their union is S . Put $\mathcal{W}_k = S_k / \sum_{i=1}^n D_i S_k$. We have the following obvious result and corollaries.

Proposition 4.1. *There is a direct sum decomposition $\mathcal{W} \cong \bigoplus_{k=1}^N \mathcal{W}_k$.*

Let \mathcal{F} be a left \mathcal{D} -module.

Corollary 4.2. *There is a direct sum decomposition*

$$\text{Hom}_{\mathcal{D}}(\mathcal{W}, \mathcal{F}) \cong \bigoplus_{k=1}^N \text{Hom}_{\mathcal{D}}(\mathcal{W}_k, \mathcal{F}).$$

Put

$$\mathcal{K}_k(\mathcal{F}) = \left\{ \xi = \sum_{u \in M_k} A_u x^{-u} \mid A_u \in \mathcal{F} \text{ for } u \in M_k, D_i^*(\xi) = 0 \text{ for } i = 1, \dots, n, \right. \\ \left. \text{and } \partial_{\lambda}(A_u) = -\ell_0 A_{u+\mathbf{a}_0} \text{ for all } u \in M_k \right\}.$$

Corollary 4.3. *We have $\text{Hom}_{\mathcal{D}}(\mathcal{W}_k, \mathcal{F}) \cong \mathcal{K}_k(\mathcal{F})$.*

Let $\mathcal{B}_k = \mathcal{B} \cap \mathcal{C}_k$. Note that $|\mathcal{B}_k| = d/N = \ell_0$ for $k = 1, \dots, N$.

Corollary 4.4. *The set $\{\xi_{\mathbf{b}}\}_{\mathbf{b} \in \mathcal{B}_k}$ is a basis for $\text{Hom}_{\mathcal{D}}(\mathcal{W}_k, \mathbb{C}[[\lambda]])$.*

The condition on the second sum in Equation (3.20) shows that if $\mathbf{b} \in \mathcal{B}_k$, then

$$(4.5) \quad G_u^{(\mathbf{b})}(\lambda) = 0 \quad \text{if } u \notin M_k.$$

Equation (3.19) thus becomes, for $\mathbf{b} \in \mathcal{B}_k$,

$$(4.6) \quad \xi_{\mathbf{b}} = \sum_{u \in M_k} G_u^{(\mathbf{b})}(\lambda) x^{-u}.$$

The following theorem connects the $\xi_{\mathbf{b}}$ with classical hypergeometric series.

Theorem 4.7. *Let $\mathbf{b} \in \mathcal{B}_k$ and $u \in M_k$. Write $\mathbf{b} = \sum_{j=1}^m v_j \mathbf{a}_j$. Let $s_0 \in \{0, 1, \dots, \ell_0 - 1\}$ be the unique element satisfying $u + s_0 \mathbf{a}_0 \equiv \mathbf{b} \pmod{\mathbb{Z}A}$ and let $s_1, \dots, s_m \in \mathbb{Z}$ be the unique solution to the equation*

$$(4.8) \quad u + s_0 \mathbf{a}_0 = \mathbf{b} - \sum_{j=1}^m s_j \mathbf{a}_j.$$

Then

$$(4.9) \quad G_u^{(\mathbf{b})}(\lambda) = \left(\prod_{j=1}^m [-v_j]_{s_j} \prod_{j=1}^m \ell_j^{s_j} \right) \frac{(-\ell_0)^{s_0}}{s_0!} F_u^{(\mathbf{b})}(\lambda),$$

where

$$(4.10) \quad F_u^{(\mathbf{b})}(\lambda) = \lambda^{s_0} \sum_{s=0}^{\infty} \frac{\prod_{j=1}^m \left(\frac{v_j - s_j}{\ell_j} \right)_s \left(\frac{v_j - s_j + 1}{\ell_j} \right)_s \dots \left(\frac{v_j - s_j + \ell_j - 1}{\ell_j} \right)_s}{\left(\frac{s_0 + 1}{\ell_0} \right)_s \left(\frac{s_0 + 2}{\ell_0} \right)_s \dots \left(\frac{s_0 + \ell_0}{\ell_0} \right)_s} \lambda^{s \ell_0}.$$

Proof. The condition on the second summation in (3.20) is equivalent to (4.8). The solution $\{s_j\}_{j=1}^m$ of (4.8) must satisfy $s_j \leq 0$ for $j = 1, \dots, m$ because the left-hand side of (4.8) lies in $C(A)$ while the right-hand side of (4.8) equals $\sum_{j=1}^m (v_j - s_j) \mathbf{a}_j$ and $v_j \in [0, 1)$ for all j .

It now follows from (1.1) that all solutions of the condition on the summation in (3.20) are of the form

$$(4.11) \quad (s_0 + s \ell_0) \mathbf{a}_0 - \mathbf{b} + \sum_{j=1}^m (s_j - s \ell_j) \mathbf{a}_j = -u \quad \text{with } s \in \mathbb{Z}_{\geq 0}.$$

We can thus write (3.20) as

$$(4.12) \quad G_u^{(\mathbf{b})}(\lambda) = \lambda^{s_0} \sum_{s=0}^{\infty} \left(\prod_{j=1}^m [-v_j]_{s_j - s \ell_j} \prod_{j=1}^m \ell_j^{s_j - s \ell_j} \right) \frac{(-\ell_0)^{s_0 + s \ell_0}}{(s_0 + s \ell_0)!} \lambda^{s \ell_0}.$$

We simplify (4.12) by factoring out terms that do not depend on s . We have

$$[-v_j]_{s_j - s \ell_j} = [-v_j]_{s_j} [-v_j + s_j]_{-s \ell_j}$$

and

$$(s_0 + s \ell_0)! = s_0! (s_0 + 1)_{s \ell_0}.$$

We thus have

$$(4.13) \quad G_u^{(\mathbf{b})}(\lambda) = \left(\prod_{j=1}^m [-v_j]_{s_j} \prod_{j=1}^m \ell_j^{s_j} \right) \frac{(-\ell_0)^{s_0}}{s_0!} F_u^{(\mathbf{b})}(\lambda),$$

where

$$(4.14) \quad F_u^{(\mathbf{b})}(\lambda) = \lambda^{s_0} \sum_{s=0}^{\infty} \left(\prod_{j=1}^m [-v_j + s_j]_{-s\ell_j} \prod_{j=1}^m \ell_j^{-s\ell_j} \right) \frac{(-\ell_0)^{s\ell_0}}{(s_0 + 1)_{s\ell_0}} \lambda^{s\ell_0}.$$

Using (3.10) we can rewrite this in terms of Pochhammer symbols:

$$(4.15) \quad F_u^{(\mathbf{b})}(\lambda) = \lambda^{s_0} \sum_{s=0}^{\infty} \left(\prod_{j=1}^m \frac{(v_j - s_j)_{s\ell_j}}{\ell_j^{s\ell_j}} \right) \frac{\ell_0^{s\ell_0}}{(s_0 + 1)_{s\ell_0}} \lambda^{s\ell_0},$$

where we have used (1.2) to eliminate minus signs. Rearranging the right-hand side of (4.15) gives (4.10). \square

If $\mathbf{b} \in \mathcal{B}_k$ is an interior point of $P(A)$, then $v_j > 0$ for all j , so (4.9) and (4.10) imply that $G_u^{(\mathbf{b})}(\lambda) \neq 0$ for all $u \in M_k$. But if \mathbf{b} is a boundary point of $P(A)$ many of the $G_u^{(\mathbf{b})}(\lambda)$ vanish.

Let $\mathbf{b} \in \mathcal{B}_k$ and let $\sigma_{\mathbf{b}}$ be the smallest closed face of $C(A)$ containing \mathbf{b} . Denote by $\sigma_{\mathbf{b}}^\circ$ the relative interior of $\sigma_{\mathbf{b}}$, i. e., $\sigma_{\mathbf{b}}^\circ$ equals $\sigma_{\mathbf{b}}$ minus all its proper closed subfaces. Note that $\mathbf{b} \in \sigma_{\mathbf{b}}^\circ$. Since $\mathbf{b} = \sum_{j=1}^m v_j \mathbf{a}_j$, we have $v_j \neq 0$ if and only if $\mathbf{a}_j \in \sigma_{\mathbf{b}}$.

Corollary 4.16. *Let $\mathbf{b} \in \mathcal{B}_k$ and $u \in M_k$. Suppose that $\sigma_{\mathbf{b}} \neq C(A)$. Then $G_u^{(\mathbf{b})}(\lambda)$ is a nonzero constant if and only if $u \in \sigma_{\mathbf{b}}^\circ \cap (\mathbf{b} + \mathbb{Z}_{\geq 0}A)$, i. e.,*

$$(4.17) \quad u = \mathbf{b} + \sum_{\{j: \mathbf{a}_j \in \sigma_{\mathbf{b}}\}} t_j \mathbf{a}_j \quad \text{for (unique) } t_j \in \mathbb{Z}_{\geq 0}.$$

When (4.17) holds, one has

$$(4.18) \quad G_u^{(\mathbf{b})}(\lambda) = \left(\prod_{\{j: \mathbf{a}_j \in \sigma_{\mathbf{b}}\}} \ell_j^{-t_j} \right) \left(\prod_{\{j: \mathbf{a}_j \in \sigma_{\mathbf{b}}\}} [-v_j]_{-t_j} \right).$$

If u is not of the form (4.17), then $G_u^{(\mathbf{b})}(\lambda) = 0$.

Proof. Since $\sigma_{\mathbf{b}} \neq C(A)$, we must have $v_j = 0$ for some j . To fix ideas, suppose that $v_j \neq 0$ for $j = 1, \dots, J$ and $v_j = 0$ for $j = J + 1, \dots, m$, where $J < m$. Equation (4.8) gives

$$(4.19) \quad u + s_0 \mathbf{a}_0 = \sum_{j=1}^m (v_j - s_j) \mathbf{a}_j.$$

If $s_j < 0$ for some $j \in \{J + 1, \dots, m\}$, then the factor $[-v_j]_{s_j} = [0]_{s_j} = 0$ appears on the right-hand side of (4.13), so $G_u^{(\mathbf{b})}(\lambda) = 0$. So we assume $s_j = 0$ for $j = J + 1, \dots, m$ and (4.19) becomes

$$(4.20) \quad u + s_0 \mathbf{a}_0 = \sum_{j=1}^J (v_j - s_j) \mathbf{a}_j.$$

If $s_0 > 0$, then the left-hand side of (4.20) is an interior point of $C(A)$ while the right-hand side of (4.20) lies in $\sigma_{\mathbf{b}}^\circ$, so we must have $s_0 = 0$ and (4.20) becomes

$$(4.21) \quad u = \mathbf{b} - \sum_{j=1}^J s_j \mathbf{a}_j.$$

This proves that $G_u^{(\mathbf{b})}(\lambda) = 0$ if u is not of the form (4.17).

Equation (4.21) gives $v_j = s_j = 0$ for $j = J + 1, \dots, m$, so (4.10) implies that $F_u^{(\mathbf{b})}(\lambda) = 1$. Equation (4.18) then follows from (4.13). \square

Corollary 4.22. *Suppose in addition to the hypotheses of Corollary 4.16 that $u \in \mathcal{B}_k$. Then*

$$(4.23) \quad G_u^{(\mathbf{b})}(\lambda) = \begin{cases} 1 & \text{if } u = \mathbf{b}, \\ 0 & \text{if } u \neq \mathbf{b}. \end{cases}$$

Proof. Since $u, \mathbf{b} \in \mathcal{B}_k$, Equation (4.17) has a solution if and only if $u = \mathbf{b}$, in which case all t_j equal 0. The assertion then follows from Corollary 4.16. \square

Since $s_0 \in \{0, 1, \dots, \ell_0 - 1\}$, there are only finitely many possibilities for the denominator on the right-hand side of (4.10). It follows that for all but finitely many u there is no cancellation in the quotient on the right-hand side of (4.10). But for $u \in \mathcal{B}_k$ there is often cancellation, which we describe precisely in what follows.

5. THE DIFFERENTIAL MODULE \mathcal{M}'

The main purpose of this section is to show that the set $X^{\mathcal{B}}$, which is a \mathbb{C} -basis for \mathcal{W}_0 by Proposition 3.9, is also a $\mathbb{C}(\lambda)$ -basis for \mathcal{M}' .

Let $\Delta(A) \subseteq V$ be the convex hull of $A \cup \{\mathbf{0}\}$. For each face σ of $\Delta(A)$ not containing \mathbf{a}_0 , let $f_\lambda^\sigma = \sum_{\mathbf{a}_j \in \sigma} \ell_j x^{\mathbf{a}_j}$, a Laurent polynomial not involving λ . If σ contains \mathbf{a}_0 , we define $f_\lambda^\sigma = f_\lambda$. Recall that f_λ is *nondegenerate relative to $\Delta(A)$* if for each face σ of $\Delta(A)$ that does not contain the origin, the polynomials $\{x_i \partial f_\lambda^\sigma / \partial x_i\}_{i=1}^n$ have no common zero in $(\overline{\mathbb{C}(\lambda)}^\times)^n$, where $\overline{\mathbb{C}(\lambda)}$ denotes an algebraic closure of $\mathbb{C}(\lambda)$.

Proposition 5.1. *The Laurent polynomial f_λ is nondegenerate relative to $\Delta(A)$.*

Proof. Suppose first that σ is the (unique) face of $\Delta(A)$ not containing the origin but containing \mathbf{a}_0 . Then

$$x_i \frac{\partial f_\lambda^\sigma}{\partial x_i} = \sum_{j=1}^m \ell_j a_{ij} x^{\mathbf{a}_j} - \ell_0 a_{i0} \lambda x^{\mathbf{a}_0}$$

for $i = 1, \dots, n$. By (1.1), this can be rewritten as

$$x_i \frac{\partial f_\lambda^\sigma}{\partial x_i} = \sum_{j=1}^m \ell_j a_{ij} (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}),$$

or, in matrix form,

$$(5.2) \quad \begin{pmatrix} x_1 \frac{\partial f_\lambda^\sigma}{\partial x_1} \\ \vdots \\ x_n \frac{\partial f_\lambda^\sigma}{\partial x_n} \end{pmatrix} = \alpha \begin{pmatrix} \ell_1 (x^{\mathbf{a}_1} - \lambda x^{\mathbf{a}_0}) \\ \vdots \\ \ell_m (x^{\mathbf{a}_m} - \lambda x^{\mathbf{a}_0}) \end{pmatrix},$$

where α is the matrix defined in Section 3. Let β also be as defined in Section 3. Multiplying both sides of (5.2) on the left by β gives

$$(5.3) \quad \beta \begin{pmatrix} x_1 \frac{\partial f_\lambda^\sigma}{\partial x_1} \\ \vdots \\ x_n \frac{\partial f_\lambda^\sigma}{\partial x_n} \end{pmatrix} = \begin{pmatrix} \ell_1(x^{\mathbf{a}_1} - \lambda x^{\mathbf{a}_0}) \\ \vdots \\ \ell_m(x^{\mathbf{a}_m} - \lambda x^{\mathbf{a}_0}) \\ 0 \\ \vdots \\ 0 \end{pmatrix}.$$

Since β is invertible, this implies that the $x_i \partial f_\lambda^\sigma / \partial x_i$ have a common zero in $(\overline{\mathbb{C}(\lambda)})^\times$ if and only if the $x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}$ have a common zero in $(\overline{\mathbb{C}(\lambda)})^\times$. But if $x^{\mathbf{a}_j} = \lambda x^{\mathbf{a}_0}$ for $j = 1, \dots, m$, then

$$\prod_{j=1}^m x^{\ell_j \mathbf{a}_j} = \prod_{j=1}^m \lambda^{\ell_j} x^{\ell_j \mathbf{a}_0}.$$

Using (1.1) and (1.2), this reduces to the equation $(1 - \lambda^{\ell_0})x^{\ell_0 \mathbf{a}_0} = 0$. Since the x_i lie in $(\overline{\mathbb{C}(\lambda)})^\times$, it is not possible to satisfy this equation.

Now suppose σ is a face of $\Delta(A)$ containing neither the origin nor \mathbf{a}_0 . Arguing as above, one obtains an equation analogous to (5.3) but with the nonzero terms in the column on the right-hand side being $\{\ell_j x^{\mathbf{a}_j}\}_{\mathbf{a}_j \in \sigma}$. This shows that the $x_i \partial f_\lambda^\sigma / \partial x_i$ have a common zero in $(\overline{\mathbb{C}(\lambda)})^\times$ if and only if the $x^{\mathbf{a}_j}$ for $\mathbf{a}_j \in \sigma$ have a common zero in $(\overline{\mathbb{C}(\lambda)})^\times$. It is clear that no such zero exists, thus f_λ is nondegenerate. \square

Let $\text{vol}(\Delta(A))$ be the volume of $\Delta(A)$ relative to Lebesgue measure on V normalized by requiring the volume of a fundamental domain for $V_{\mathbb{Z}}$ to be 1. Thus

$$m! \text{vol}(\Delta(A)) = \text{vol}(P(A)) = [V_{\mathbb{Z}} : \mathbb{Z}A] = d.$$

From a result of Kouchnirenko [13] (see also [2, Theorem 2.17]) we get the following corollary.

Corollary 5.4. *One has $\dim_{\mathbb{C}(\lambda)} S' / \sum_{i=1}^n (x_i \frac{\partial f_\lambda}{\partial x_i}) S' = d$.*

Proposition 5.5. *The set $X^{\mathcal{B}}$ is a basis for $S' / \sum_{i=1}^n (x_i \frac{\partial f_\lambda}{\partial x_i}) S'$.*

Proof. By Corollary 5.4 the set $X^{\mathcal{B}}$ has the correct cardinality, so it suffices to prove that it spans the quotient. Since β is invertible, Equation (5.3) shows that

$$(5.6) \quad S' / \sum_{i=1}^n (x_i \frac{\partial f_\lambda}{\partial x_i}) S' = S' / \sum_{j=1}^m (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}) S'.$$

Let $u_0 \in M$ and consider $x^{u_0} \in S'$. If $u_0 \in \mathcal{B}$ then $x^{u_0} \in X^{\mathcal{B}}$, so there is nothing to prove. Suppose $u_0 \notin \mathcal{B}$. Then there exist $j_1 \in \{1, \dots, m\}$ and $u_1 \in M$ such that $u_0 = \mathbf{a}_{j_1} + u_1$, hence

$$x^{u_0} = (x^{\mathbf{a}_{j_1}} - \lambda x^{\mathbf{a}_0}) x^{u_1} + \lambda x^{u_1 + \mathbf{a}_0}.$$

If $u_1 + \mathbf{a}_0 \in \mathcal{B}$, then $x^{u_1 + \mathbf{a}_0} \in X^{\mathcal{B}}$ and we are done. Otherwise there exist $j_2 \in \{1, \dots, m\}$ and $u_2 \in M$ such that $u_1 + \mathbf{a}_0 = \mathbf{a}_{j_2} + u_2$, hence

$$\lambda x^{u_1 + \mathbf{a}_0} = (x^{\mathbf{a}_{j_2}} - \lambda x^{\mathbf{a}_0}) \lambda x^{u_2} + \lambda^2 x^{u_2 + \mathbf{a}_0}.$$

If $u_2 + \mathbf{a}_0 \in \mathcal{B}$, we are done, otherwise this procedure can be continued.

If this procedure continues for K steps with $u_k + \mathbf{a}_0 \notin \mathcal{B}$ for $k = 0, 1, \dots, K-1$, we get a sequence $u_0, u_1, \dots, u_K \in M$ with

$$(5.7) \quad \lambda^{k-1} x^{u_{k-1} + \mathbf{a}_0} = (x^{\mathbf{a}_{j_k}} - \lambda x^{\mathbf{a}_0}) \lambda^{k-1} x^{u_k} + \lambda^k x^{u_k + \mathbf{a}_0}.$$

Note that in this procedure we have inductively

$$(5.8) \quad x^{u_0} \equiv \lambda^k x^{u_k + \mathbf{a}_0} \pmod{\sum_{j=1}^m (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}) S'}$$

and

$$(5.9) \quad u_k + \mathbf{a}_0 = u_0 + k\mathbf{a}_0 - \mathbf{a}_{j_1} - \dots - \mathbf{a}_{j_k}.$$

If this process terminates for $K < \ell_0$, we are done. If we can continue to $K = \ell_0$, then Equations (5.8) and (5.9) imply that

$$(5.10) \quad x^{u_0} \equiv \lambda^{\ell_0} x^{u_{\ell_0} + \mathbf{a}_0} \pmod{\sum_{j=1}^m (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}) S'}$$

and

$$(5.11) \quad u_{\ell_0} + \mathbf{a}_0 = u_0 + \ell_0 \mathbf{a}_0 - \mathbf{a}_{j_1} - \dots - \mathbf{a}_{j_{\ell_0}}.$$

Note that for any pair $j, j' \in \{1, \dots, m\}$ we have

$$x^{\mathbf{a}_j} - x^{\mathbf{a}_{j'}} = (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}) - (x^{\mathbf{a}_{j'}} - \lambda x^{\mathbf{a}_0}) \in \sum_{j=1}^m (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}) S'.$$

So if $u - \mathbf{a}_j \in M$, we have

$$x^{u - \mathbf{a}_j} (x^{\mathbf{a}_j} - x^{\mathbf{a}_{j'}}) = x^u - x^{u + \mathbf{a}_{j'} - \mathbf{a}_j},$$

hence

$$(5.12) \quad x^u \equiv x^{u + \mathbf{a}_{j'} - \mathbf{a}_j} \pmod{\sum_{j=1}^m (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}) S'}.$$

Applying this repeatedly to (5.11) we get

$$(5.13) \quad x^{u_{\ell_0} + \mathbf{a}_0} \equiv x^{u_0 + \ell_0 \mathbf{a}_0 - \mathbf{a}_{j'_1} - \dots - \mathbf{a}_{j'_{\ell_0}}} \pmod{\sum_{j=1}^m (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}) S'}$$

for any sequence $\mathbf{a}_{j'_1}, \dots, \mathbf{a}_{j'_{\ell_0}}$. We now choose this sequence to be \mathbf{a}_1 taken ℓ_1 times, \mathbf{a}_2 taken ℓ_2 times, \dots , \mathbf{a}_m taken ℓ_m times. Equations (1.1) and (5.13) now imply

$$(5.14) \quad x^{u_{\ell_0} + \mathbf{a}_0} \equiv x^{u_0} \pmod{\sum_{j=1}^m (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}) S'}.$$

From Equation (5.10) we get

$$(5.15) \quad (\lambda^{\ell_0} - 1) x^{u_0} \equiv 0 \pmod{\sum_{j=1}^m (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}) S'},$$

which completes the proof. \square

We record for possible future use a corollary of the proof, which can be obtained by arguing with S in place of S' .

Corollary 5.16. *We have*

$$(1 - \lambda^{\ell_0})S \subseteq \sum_{\mathbf{b} \in \mathcal{B}} \mathbb{C}[\lambda]x^{\mathbf{b}} + \sum_{j=1}^m (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0})S.$$

Relative to the grading defined on S' in Section 3, the elements $x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}$ are homogeneous of degree 1. To show that these elements form a regular sequence on S' it suffices to show that if $\{g_j(\lambda, x)\}_{j=1}^m$ are elements of S' such that

$$(5.17) \quad \sum_{j=1}^m (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0})g_j(\lambda, x) = 0,$$

then there exists a skew-symmetric set $\{h_{ij}(\lambda, x)\}_{i,j=1}^m$ of elements of S' (i. e., $h_{ij} = -h_{ji}$) such that

$$(5.18) \quad g_j(\lambda, x) = \sum_{i=1}^m (x^{\mathbf{a}_i} - \lambda x^{\mathbf{a}_0})h_{ij}(\lambda, x) \quad \text{for } j = 1, \dots, m.$$

Proposition 5.19. *The elements $\{x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}\}_{j=1}^m$ form a regular sequence on S' .*

Proof. Since S' is a graded ring and the $x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}$ are homogeneous, it suffices to prove that (5.18) follows from (5.17) when the $g_j(\lambda, x)$ are homogeneous of the same degree. After multiplying (5.17) by an element of $\mathbb{C}[\lambda]$ we may assume that the $g_j(\lambda, x)$ lie in S . As such they may be written in the form

$$(5.20) \quad g_j(\lambda, x) = \sum_{k=0}^N \lambda^k g_j^{(k)}(x)$$

where the $g_j^{(k)}(x)$ are homogeneous elements of the same degree in S_0 . Equation (5.17) then implies that

$$\sum_{j=1}^m x^{\mathbf{a}_j} g_j^{(0)}(x) = 0.$$

Since the $x^{\mathbf{a}_j}$ form a regular sequence in S_0 , there exist $h_{ij}^{(0)}(x)$ in S_0 , skew-symmetric, such that

$$(5.21) \quad g_j^{(0)}(x) = \sum_{i=1}^m x^{\mathbf{a}_i} h_{ij}^{(0)}(x).$$

Define

$$\tilde{g}_j(\lambda, x) = g_j(\lambda, x) - \sum_{i=1}^m (x^{\mathbf{a}_i} - \lambda x^{\mathbf{a}_0})h_{ij}^{(0)}(x).$$

By (5.17) and skew-symmetry, we have

$$\sum_{j=1}^m (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0})\tilde{g}_j(\lambda, x) = 0.$$

Furthermore, from (5.20) and (5.21) we have

$$\lambda^{-1}\tilde{g}_j(\lambda, x) = x^{\mathbf{a}_0} \sum_{i=1}^m h_{ij}^{(0)}(x) + \sum_{k=0}^{N-1} \lambda^k g_j^{(k+1)}(x).$$

It follows that

$$\sum_{j=1}^m (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}) (\lambda^{-1} \tilde{g}_j(\lambda, x)) = 0,$$

hence it suffices to prove that (5.17) implies (5.18) when the $g_j(\lambda, x)$ are replaced by the $\lambda^{-1} \tilde{g}_j(\lambda, x)$. Since the $\lambda^{-1} \tilde{g}_j(\lambda, x)$ are of degree $N - 1$ in λ , we are reduced by induction to the case where $N = 0$, i. e., the $g_j(\lambda, x)$ do not involve λ .

So we now suppose that we have elements $g_j(x)$ of S_0 , homogeneous of the same degree, such that

$$(5.22) \quad \sum_{j=1}^m (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}) g_j(x) = 0$$

in S' . This implies that

$$(5.23) \quad \sum_{j=1}^m x^{\mathbf{a}_j} g_j(x) = 0$$

and

$$(5.24) \quad \sum_{j=1}^m g_j(x) = 0.$$

Since the $x^{\mathbf{a}_j}$ form a regular sequence in S_0 there exists a skew-symmetric set $\{h_{ij}(x)\}_{i,j=1}^m$ of elements of S_0 with $\deg h_{ij} = \deg g_j - 1$ such that

$$(5.25) \quad g_j(x) = \sum_{i=1}^m x^{\mathbf{a}_i} h_{ij}(x).$$

Define

$$(5.26) \quad \tilde{g}_j(x) = g_j(x) - \sum_{i=1}^m (x^{\mathbf{a}_i} - \lambda x^{\mathbf{a}_0}) h_{ij}(x).$$

By (5.22) and skew-symmetry we have

$$(5.27) \quad \sum_{j=1}^m (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}) \tilde{g}_j(x) = 0.$$

Furthermore, from (5.25), we have

$$\tilde{g}_j(x) = \lambda x^{\mathbf{a}_0} \sum_{i=1}^m h_{ij}(x),$$

so (5.27) implies

$$\sum_{j=1}^m (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}) \left(\sum_{i=1}^m h_{ij}(x) \right) = 0.$$

Since the $h_{ij}(x)$ have degree one less than the $g_j(x)$, we are reduced by induction to the case where the $g_j(x)$ have degree less than 1. In this case Equation (5.23) implies that the $g_j(x)$ equal 0 because the $x^{\mathbf{a}_j}$ form a regular sequence on S_0 . \square

We now follow the argument of Section 3. In analogy with (3.2) we define differential operators \tilde{D}_i , $i = 1, \dots, n$ by the formula

$$(5.28) \quad \begin{pmatrix} \tilde{D}_1 \\ \vdots \\ \tilde{D}_n \end{pmatrix} = \beta \begin{pmatrix} D_1 \\ \vdots \\ D_n \end{pmatrix} = \begin{pmatrix} \sum_{j=1}^n \beta_{1j} D_j \\ \vdots \\ \sum_{j=1}^n \beta_{nj} D_j \end{pmatrix}.$$

Lemma 5.29. *Let $u \in M$ and write $u = (u_1, \dots, u_n) = \sum_{j=1}^m u'_j \mathbf{a}_j$. We have*

$$\tilde{D}_j(x^u) = \begin{cases} u'_j x^u + \ell_j x^u (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}) & \text{for } j = 1, \dots, m, \\ 0 & \text{for } j = m+1, \dots, n. \end{cases}$$

Proof. We note that

$$(5.30) \quad \begin{pmatrix} x_1 \frac{\partial f_\lambda}{\partial x_1} \\ \vdots \\ x_n \frac{\partial f_\lambda}{\partial x_n} \end{pmatrix} = \alpha \begin{pmatrix} \ell_1(x^{\mathbf{a}_1} - \lambda x^{\mathbf{a}_0}) \\ \vdots \\ \ell_m(x^{\mathbf{a}_m} - \lambda x^{\mathbf{a}_0}) \end{pmatrix}$$

and

$$(5.31) \quad \alpha \begin{pmatrix} u'_1 \\ \vdots \\ u'_m \end{pmatrix} = \begin{pmatrix} u_1 \\ \vdots \\ u_n \end{pmatrix}.$$

From the definition we have

$$(5.32) \quad \begin{aligned} \begin{pmatrix} \tilde{D}_1 \\ \vdots \\ \tilde{D}_n \end{pmatrix} (x^u) &= \beta \begin{pmatrix} x_1 \frac{\partial}{\partial x_1} + x_1 \frac{\partial f_\lambda}{\partial x_1} \\ \vdots \\ x_n \frac{\partial}{\partial x_n} + x_n \frac{\partial f_\lambda}{\partial x_n} \end{pmatrix} (x^u) \\ &= \beta \begin{pmatrix} u_1 \\ \vdots \\ u_n \end{pmatrix} x^u + \beta \begin{pmatrix} x_1 \frac{\partial f_\lambda}{\partial x_1} \\ \vdots \\ x_n \frac{\partial f_\lambda}{\partial x_n} \end{pmatrix} x^u \\ &= \beta \alpha \begin{pmatrix} u'_1 \\ \vdots \\ u'_m \end{pmatrix} x^u + \beta \alpha \begin{pmatrix} \ell_1(x^{\mathbf{a}_1} - \lambda x^{\mathbf{a}_0}) \\ \vdots \\ \ell_m(x^{\mathbf{a}_m} - \lambda x^{\mathbf{a}_0}) \end{pmatrix} x^u \\ &= \begin{pmatrix} u'_1 x^u + \ell_1 x^u (x^{\mathbf{a}_1} - \lambda x^{\mathbf{a}_0}) \\ \vdots \\ u'_m x^u + \ell_m x^u (x^{\mathbf{a}_m} - \lambda x^{\mathbf{a}_0}) \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \end{aligned}$$

where the third equality follows from (5.30) and (5.31) and the fourth equality follows from (3.1). \square

Since β is invertible, we get the following result.

Corollary 5.33. *We have $\mathcal{W}' = S' / \sum_{j=1}^m \tilde{D}_j S'$.*

Proposition 5.34. *The set $X^{\mathcal{B}}$ is a basis for \mathcal{W}' .*

Proof. Consider the Koszul complex $\text{Kos}^\bullet(S', \{\tilde{D}_j\}_{j=1}^m)$ defined by the \tilde{D}_j . The grading defined in Section 3 makes it into a filtered complex for which (by Corollary 5.33)

$$H^m(\text{Kos}^\bullet(S', \{\tilde{D}_j\}_{j=1}^m)) = \mathcal{W}'.$$

The associated graded complex is $\text{Kos}^\bullet(S', \{\ell_j(x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0})\}_{j=1}^m)$. By Proposition 5.19 the set $\{x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0}\}_{j=1}^m$ is a regular sequence on S' , so

$$H^i(\text{Kos}^\bullet(S', \{\ell_j(x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0})\}_{j=1}^m)) = 0 \quad \text{for } 0 \leq i < m.$$

It follows that

$$H^i(\text{Kos}^\bullet(S', \{\tilde{D}_j\}_{j=1}^m)) = H^i(\text{Kos}^\bullet(S', \{\ell_j(x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0})\}_{j=1}^m))$$

for all i . In particular we have

$$\begin{aligned} \mathcal{W}' &= H^m(\text{Kos}^\bullet(S', \{\tilde{D}_j\}_{j=1}^m)) = H^m(\text{Kos}^\bullet(S', \{\ell_j(x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0})\}_{j=1}^m)) \\ &= S' / \sum_{j=1}^m (x^{\mathbf{a}_j} - \lambda x^{\mathbf{a}_0})S'. \end{aligned}$$

Proposition 5.5 and Equation (5.6) now imply the proposition. \square

6. FUNDAMENTAL SOLUTION MATRIX

Associated to the basis $X^{\mathcal{B}}$ of \mathcal{W}' is a Picard-Fuchs equation described as follows. Write $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_d\}$. For $i, j = 1, \dots, d$, define $C_{ij}(\lambda) \in \mathbb{C}(\lambda)$ by

$$(6.1) \quad D_\lambda(x^{\mathbf{b}_i}) = \sum_{j=1}^d C_{ij}(\lambda)x^{\mathbf{b}_j}$$

in \mathcal{W}' . Let Y be the column vector with entries y_1, \dots, y_d and let $C(\lambda)$ be the $(d \times d)$ -matrix $C(\lambda) = (C_{ij}(\lambda))_{i,j=1}^d$. The associated Picard-Fuchs equation is

$$(6.2) \quad \partial_\lambda Y = C(\lambda)Y.$$

Let $\xi \in \text{Hom}_{\mathbb{C}(\lambda)}(\mathcal{W}', \mathbb{C}((\lambda)))$. Then $\xi \in \text{Hom}_{\mathcal{D}'}(\mathcal{W}', \mathbb{C}((\lambda)))$ if and only if the column vector with entries $\xi(x^{\mathbf{b}_1}), \dots, \xi(x^{\mathbf{b}_d})$ is a solution of the system (6.2). Since $X^{\mathcal{B}}$ is a basis for \mathcal{W}' , the map that sends ξ to the column vector with entries $\xi(x^{\mathbf{b}_1}), \dots, \xi(x^{\mathbf{b}_d})$ identifies the \mathbb{C} -vector space $\text{Hom}_{\mathcal{D}'}(\mathcal{W}', \mathbb{C}((\lambda)))$ with the space of solutions of (6.2) that lie in $\mathbb{C}((\lambda))^d$. This latter space has dimension less than or equal to d , so

$$\dim_{\mathbb{C}} \text{Hom}_{\mathcal{D}'}(\mathcal{W}', \mathbb{C}((\lambda))) \leq d.$$

But there is a natural inclusion

$$\text{Hom}_{\mathcal{D}}(\mathcal{W}, \mathbb{C}[[\lambda]]) \hookrightarrow \text{Hom}_{\mathcal{D}'}(\mathcal{W}', \mathbb{C}((\lambda))),$$

and $\dim_{\mathbb{C}} \text{Hom}_{\mathcal{D}}(\mathcal{W}, \mathbb{C}[[\lambda]]) = d$ by Corollary 3.18. It follows that (6.2) has a full set of solutions in $\mathbb{C}[[\lambda]]^d$. Corollary 3.18 implies that the set $\{\xi_{\mathbf{b}_j}\}_{j=1}^d$ is a basis for $\text{Hom}_{\mathcal{D}'}(\mathcal{W}', \mathbb{C}((\lambda)))$. The associated fundamental solution matrix of (6.2) is the $(d \times d)$ -matrix whose (i, j) -entry is, by (3.19),

$$(6.3) \quad \xi_{\mathbf{b}_j}(x^{\mathbf{b}_i}) = G_{\mathbf{b}_i}^{(\mathbf{b}_j)}(\lambda).$$

As in Section 4, we have a direct sum decomposition of \mathcal{W}' . For $k = 1, \dots, N$, let S'_k be the $\mathbb{C}(\lambda)$ -vector space generated by the x^u with $u \in M_k$ and put $\mathcal{W}'_k = S'_k / \sum_{i=1}^n D_i S'_k$. The set $X^{\mathcal{B}_k}$ is a basis for \mathcal{W}'_k and the set $\{\xi_{\mathbf{b}}\}_{\mathbf{b} \in \mathcal{B}_k}$, with $\xi_{\mathbf{b}}$ given

by (4.6), is a basis for $\text{Hom}_{\mathcal{D}'}(\mathcal{W}'_k, \mathbb{C}((\lambda)))$. Fix k and write $\mathcal{B}_k = \{\mathbf{b}_1, \dots, \mathbf{b}_{\ell_0}\}$. The associated fundamental solution matrix is the $(\ell_0 \times \ell_0)$ -matrix whose (i, j) -entry is $G_{\mathbf{b}_i}^{(\mathbf{b}_j)}(\lambda)$.

As we observed in Section 4, if \mathbf{b}_j is an interior point of $P(A)$, then $G_{\mathbf{b}_i}^{(\mathbf{b}_j)}(\lambda) \neq 0$ for all $\mathbf{b}_i \in \mathcal{B}_k$ i. e., all entries in column j are nonzero. On the other hand, by Corollary 4.22, if \mathbf{b}_j is a boundary point of $P(A)$, then

$$(6.4) \quad G_{\mathbf{b}_i}^{(\mathbf{b}_j)}(\lambda) = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases}$$

Thus all entries in column j are zero except for the entry in row j , which equals 1.

We now describe the entries in each row. We fix i and let j vary. By the above, if \mathbf{b}_i is an interior point of $P(A)$, then $G_{\mathbf{b}_i}^{(\mathbf{b}_j)}(\lambda) \neq 0$ if and only if \mathbf{b}_j is an interior point of $P(A)$. If \mathbf{b}_i is a boundary point of $P(A)$, then $G_{\mathbf{b}_i}^{(\mathbf{b}_j)}(\lambda) \neq 0$ if and only if either \mathbf{b}_j is an interior point of $P(A)$ or $\mathbf{b}_j = \mathbf{b}_i$. Let R_k be the number of elements of \mathcal{B}_k that are interior to $P(A)$. Then row i has R_k nonzero entries if \mathbf{b}_i is an interior point of $P(A)$ and has $R_k + 1$ nonzero entries if \mathbf{b}_i is a boundary point of $P(A)$.

Since $F_{\mathbf{b}_i}^{(\mathbf{b}_j)}(\lambda)$ determines $G_{\mathbf{b}_i}^{(\mathbf{b}_j)}(\lambda)$ up to a constant factor, we examine the $F_{\mathbf{b}_i}^{(\mathbf{b}_j)}(\lambda)$ more closely. For each j , write

$$(6.5) \quad \mathbf{b}_j = \sum_{k=1}^m v_k^{(j)} \mathbf{a}_k.$$

Let $s_0^{(j)} \in \{0, 1, \dots, \ell_0 - 1\}$ be the unique element such that

$$\mathbf{b}_i + s_0^{(j)} \mathbf{a}_0 \equiv \mathbf{b}_j \pmod{\mathbb{Z}A}.$$

Then let $s_k^{(j)}$, $k = 1, \dots, m$, be the unique nonpositive integers such that

$$(6.6) \quad \mathbf{b}_i + s_0^{(j)} \mathbf{a}_0 = \mathbf{b}_j - \sum_{k=1}^m s_k^{(j)} \mathbf{a}_k.$$

The $s_k^{(j)}$ for $j = 0, 1, \dots, m$ depend on i , but since i is fixed this should not cause any confusion. Note that for fixed i , the correspondence $\mathbf{b}_j \leftrightarrow s_0^{(j)}$ is one-to-one between \mathcal{B}_k and the set $\{0, 1, \dots, \ell_0 - 1\}$. From (4.10) we have

$$(6.7) \quad F_{\mathbf{b}_i}^{(\mathbf{b}_j)}(\lambda) = \frac{\prod_{k=1}^m \left(\frac{v_k^{(j)} - s_k^{(j)}}{\ell_k} \right)_s \left(\frac{v_k^{(j)} - s_k^{(j)} + 1}{\ell_k} \right)_s \cdots \left(\frac{v_k^{(j)} - s_k^{(j)} + \ell_k - 1}{\ell_k} \right)_s}{\left(\frac{s_0^{(j)} + 1}{\ell_0} \right)_s \left(\frac{s_0^{(j)} + 2}{\ell_0} \right)_s \cdots \left(\frac{s_0^{(j)} + \ell_0}{\ell_0} \right)_s} \lambda^{s\ell_0}.$$

We first observe that there is a simple algebraic relation between the formula (6.7) and its special case $j = i$: since $s_k^{(i)} = 0$ for $k = 0, 1, \dots, m$ we have

$$(6.8) \quad F_{\mathbf{b}_i}^{(\mathbf{b}_i)}(\lambda) = \sum_{s=0}^{\infty} \frac{\prod_{k=1}^m \left(\frac{v_k^{(i)}}{\ell_k} \right)_s \left(\frac{v_k^{(i)} + 1}{\ell_k} \right)_s \cdots \left(\frac{v_k^{(i)} + \ell_k - 1}{\ell_k} \right)_s}{\left(\frac{1}{\ell_0} \right)_s \left(\frac{2}{\ell_0} \right)_s \cdots \left(\frac{\ell_0}{\ell_0} \right)_s} \lambda^{s\ell_0}.$$

Proposition 6.9. *The formula (6.7) for $F_{\mathbf{b}_i}^{(\mathbf{b}_j)}(\lambda)$ is obtained from the formula (6.8) for $F_{\mathbf{b}_i}^{(\mathbf{b}_i)}(\lambda)$ by multiplying the series (6.8) by $\lambda^{s_0^{(j)}}$ and adding $s_0^{(j)}/\ell_0$ to each term inside parenthesis.*

Proof. The only nonobvious part of this proposition is the assertion that

$$(6.10) \quad \frac{v_k^{(i)}}{\ell_k} + \frac{s_0^{(j)}}{\ell_0} = \frac{v_k^{(j)} - s_k^{(j)}}{\ell_k}$$

for $k = 1, \dots, m$. From (6.6) we have

$$(6.11) \quad \sum_{k=1}^m (v_k^{(j)} - s_k^{(j)}) \mathbf{a}_k = \mathbf{b}_i + s_0^{(j)} \mathbf{a}_0.$$

Using (1.1), this equation can be rewritten as

$$(6.12) \quad \sum_{k=1}^m \left(v_k^{(j)} - s_k^{(j)} - \frac{s_0^{(j)} \ell_k}{\ell_0} \right) \mathbf{a}_k = \mathbf{b}_i.$$

But $\mathbf{b}_i = \sum_{k=1}^m v_k^{(i)} \mathbf{a}_k$ and $\{\mathbf{a}_k\}_{k=1}^m$ is a linearly independent set, so

$$v_k^{(i)} = v_k^{(j)} - s_k^{(j)} - \frac{s_0^{(j)} \ell_k}{\ell_0}$$

for $k = 1, \dots, m$. This equation is equivalent to (6.10). \square

We now determine which terms in the ratio on the right-hand side of (6.8) cancel.

Lemma 6.13. (a) *Suppose that either \mathbf{b}_i is an interior point of $P(A)$ or $\mathbf{b}_i \neq \mathbf{b}_j$. Then $0 < v_k^{(j)} - s_k^{(j)} < \ell_k + 1$ for $k = 1, \dots, m$.*

(b) *If \mathbf{b}_i is a boundary point of $P(A)$ and $\mathbf{b}_i = \mathbf{b}_j$, then $0 \leq v_k^{(j)} - s_k^{(j)} < 1$ for $k = 1, \dots, m$.*

Proof. If \mathbf{b}_i is an interior point of $C(A)$, then, since $s_0^{(j)} \geq 0$, the right-hand side of (6.11) is an interior point of $C(A)$, so $v_k^{(j)} - s_k^{(j)} > 0$ for all k . If $\mathbf{b}_i \neq \mathbf{b}_j$, then $s_0^{(j)} > 0$ so again the right-hand side of (6.11) is an interior point of $C(A)$. Since $\mathbf{b}_i \in P(A)$, Equation (6.12) implies that

$$v_k^{(j)} - s_k^{(j)} - \frac{s_0^{(j)} \ell_k}{\ell_0} < 1$$

for all k , i. e.,

$$\frac{v_k^{(j)} - s_k^{(j)}}{\ell_k} - \frac{s_0^{(j)}}{\ell_0} < \frac{1}{\ell_k}.$$

Since $s_0^{(j)}/\ell_0 < 1$, this implies that $v_k^{(j)} - s_k^{(j)} < \ell_k + 1$. This proves part (a).

If $\mathbf{b}_i = \mathbf{b}_j$, then $s_k^{(j)} = 0$ for $k = 0, 1, \dots, m$, so $v_k^{(j)} - s_k^{(j)} = v_k^{(j)}$. Part (b) then follows from (6.5). \square

Proposition 6.14. (a) *Suppose that \mathbf{b}_i is an interior point of $P(A)$. There are exactly $\ell_0 - R_k$ factors that appear in both numerator and denominator on the right-hand side of (6.8), namely,*

$$\{(1 - s_0^{(j)}/\ell_0)_s \mid \mathbf{b}_j \text{ is a boundary point of } P(A)\}.$$

(b) *Suppose that \mathbf{b}_i is a boundary point of $P(A)$. There are exactly $\ell_0 - R_k - 1$ factors that appear in both numerator and denominator on the right-hand side of (6.8), namely,*

$$\{(1 - s_0^{(j)}/\ell_0)_s \mid \mathbf{b}_j \text{ is a boundary point of } P(A) \text{ and } \mathbf{b}_j \neq \mathbf{b}_i\}.$$

Proof. Since $s_0^{(j)}$ runs through the set $\{0, 1, \dots, \ell_0 - 1\}$ as \mathbf{b}_j runs through the set \mathcal{B}_k , the ratios $1 - s_0^{(j)}/\ell_0$ run through the set $\{1/\ell_0, 2/\ell_0, \dots, \ell_0/\ell_0\}$. We need to determine when

$$(6.15) \quad 1 - \frac{s_0^{(j)}}{\ell_0} = \frac{v_k^{(i)} + \sigma_k}{\ell_k}$$

for some k and some $\sigma_k \in \{0, 1, \dots, \ell_k - 1\}$. Solving (6.10) for $v_k^{(i)}/\ell_k$, substituting into (6.15), and simplifying gives the equation

$$(6.16) \quad \ell_k - (v_k^{(j)} - s_k^{(j)}) = \sigma_k.$$

Since ℓ_k and $s_k^{(j)}$ are integers, this equation has a solution $\sigma_k \in \mathbb{Z}$ if and only if $v_k^{(j)}$ is an integer. But $0 \leq v_k^{(j)} < 1$, so $v_k^{(j)}$ is integral if and only if $v_k^{(j)} = 0$, i. e., if and only if \mathbf{b}_j is a boundary point of $P(A)$. In this case we have $\sigma_k = \ell_k + s_k^{(j)}$, so we get cancellation if and only if $s_k^{(j)} \in \{-1, -2, \dots, -\ell_k\}$.

If \mathbf{b}_i is an interior point of $P(A)$, it follows from Lemma 6.13(a) that if $v_k^{(j)} = 0$ then $s_k^{(j)} \in \{-1, -2, \dots, -\ell_k\}$. Thus every boundary point \mathbf{b}_j produces a cancellation. This proves part (a) of the proposition.

If \mathbf{b}_i is a boundary point of $P(A)$ and \mathbf{b}_j is a boundary point different from \mathbf{b}_i , then Lemma 6.13(a) again implies that $s_k^{(j)} \in \{-1, -2, \dots, -\ell_k\}$ so we get a cancellation. But if $\mathbf{b}_j = \mathbf{b}_i$, then by Lemma 6.13(b) if $v_k^{(j)} = 0$, then $s_k^{(j)} = 0$ so there is no cancellation in that case. This proves part (b) of the proposition. \square

Suppose that \mathbf{b}_i is an interior point of $P(A)$. Applying Proposition 6.14(a) we let $\beta_1, \dots, \beta_{R_k}$ be the elements on the list $1/\ell_0, \dots, \ell_0/\ell_0$ that are not of the form $1 - s_0^{(j)}/\ell_0$ for \mathbf{b}_j a boundary point of $P(A)$. In particular, since $\ell_0/\ell_0 = 1$ is not of this form we may take $\beta_1 = 1$. We let $\alpha_1, \dots, \alpha_{R_k}$ be the elements remaining on the list

$$\frac{v_1^{(i)}}{\ell_1}, \frac{v_1^{(i)} + 1}{\ell_1}, \dots, \frac{v_1^{(i)} + \ell_1 - 1}{\ell_1}, \dots, \frac{v_m^{(i)}}{\ell_m}, \frac{v_m^{(i)} + 1}{\ell_m}, \dots, \frac{v_m^{(i)} + \ell_m - 1}{\ell_m}$$

after one copy of each element $1 - s_0^{(j)}/\ell_0$ with \mathbf{b}_j a boundary point of $P(A)$ has been removed.

Theorem 6.17. *Suppose that \mathbf{b}_i is an interior point of $P(A)$. The R_k series (6.7) with \mathbf{b}_j an interior point of $P(A)$ come from a full set of solutions at $x = 0$ to the hypergeometric operator*

$$(\delta_x + \beta_1 - 1) \cdots (\delta_x + \beta_{R_k} - 1) - x(\delta_x + \alpha_1) \cdots (\delta_x + \alpha_{R_k})$$

by replacing x by λ^{ℓ_0} .

Proof. Propositions 6.9 and 6.14(a) imply that the series (6.7) are obtained from the series (6.8) in exactly the way that the series (1.6) are obtained from the series (1.5). The conclusion of the theorem then follows from the discussion of the solutions of (1.4) in Section 1. \square

Suppose that \mathbf{b}_i is a boundary point of $P(A)$. Applying Proposition 6.14(b) we let $\beta_1, \dots, \beta_{R_k+1}$ be the elements on the list $1/\ell_0, \dots, \ell_0/\ell_0$ that are not of the form $1 - s_0^{(j)}/\ell_0$ for \mathbf{b}_j a boundary point of $P(A)$ with $\mathbf{b}_j \neq \mathbf{b}_i$. In particular, since $\ell_0/\ell_0 = 1$ is not of this form we may take $\beta_1 = 1$. We let $\alpha_1, \dots, \alpha_{R_k+1}$ be the elements remaining on the list

$$\frac{v_1^{(i)}}{\ell_1}, \frac{v_1^{(i)} + 1}{\ell_1}, \dots, \frac{v_1^{(i)} + \ell_1 - 1}{\ell_1}, \dots, \frac{v_m^{(i)}}{\ell_m}, \frac{v_m^{(i)} + 1}{\ell_m}, \dots, \frac{v_m^{(i)} + \ell_m - 1}{\ell_m}$$

after one copy of each element $1 - s_0^{(j)}/\ell_0$ for \mathbf{b}_j a boundary point of $P(A)$ with $\mathbf{b}_j \neq \mathbf{b}_i$ has been removed. The proof of the following result is analogous to the proof of Theorem 6.17.

Theorem 6.18. *Suppose that \mathbf{b}_i is a boundary point of $P(A)$. The $R_k + 1$ series (6.7) with \mathbf{b}_j an interior point of $P(A)$ or $\mathbf{b}_j = \mathbf{b}_i$ come from a full set of solutions at $x = 0$ to the hypergeometric equation*

$$(\delta_x + \beta_1 - 1) \cdots (\delta_x + \beta_{R_k+1} - 1) - x(\delta_x + \alpha_1) \cdots (\delta_x + \alpha_{R_k+1})$$

by replacing x by λ^{ℓ_0} .

7. EXAMPLE

We treat an example related to that of Gährs [9, Section 2.3]. Take $\mathbf{a}_1 = (5, 1, 0, 0)$, $\mathbf{a}_2 = (0, 4, 1, 0)$, $\mathbf{a}_3 = (0, 0, 8, 0)$, $\mathbf{a}_4 = (0, 0, 0, 2)$, and $\mathbf{a}_0 = (1, 1, 1, 1)$. Equation (1.1) becomes

$$(7.1) \quad 10\mathbf{a}_0 = 2\mathbf{a}_1 + 2\mathbf{a}_2 + \mathbf{a}_3 + 5\mathbf{a}_4,$$

so $\ell_1 = \ell_2 = 2$, $\ell_3 = 1$, $\ell_4 = 5$, $\ell_0 = 10$, and

$$f\lambda = 2x_1^5x_2 + 2x_2^4x_3 + x_3^8 + 5x_4^2 - 10\lambda x_1x_2x_3x_4.$$

We take $\mathbf{b}_1 = (1, 1, 1, 1) = \frac{1}{5}\mathbf{a}_1 + \frac{1}{5}\mathbf{a}_2 + \frac{1}{10}\mathbf{a}_3 + \frac{1}{2}\mathbf{a}_4$, an interior point of $P(A)$, so Equation (6.8) becomes

$$(7.2) \quad F_{\mathbf{b}_1}^{\mathbf{b}_1}(\lambda) = \sum_{s=0}^{\infty} \frac{\binom{1}{10}_s \binom{6}{10}_s \binom{1}{10}_s \binom{6}{10}_s \binom{1}{10}_s \binom{1}{10}_s \binom{3}{10}_s \binom{5}{10}_s \binom{7}{10}_s \binom{9}{10}_s}{\binom{1}{10}_s \binom{2}{10}_s \binom{3}{10}_s \binom{4}{10}_s \binom{5}{10}_s \binom{6}{10}_s \binom{7}{10}_s \binom{8}{10}_s \binom{9}{10}_s \binom{10}{10}_s} \lambda^{10s} \\ = {}_4F_3(1/10, 1/10, 1/10, 3/5; 1/5, 2/5, 4/5; \lambda^{10}).$$

By Theorem 6.17, the four series corresponding to the four interior points of $P(A)$ that are congruent to \mathbf{b}_1 modulo $\mathbb{Z}A_+$ and given by Equation (6.7) are obtained from a full set of solutions to

$$(7.3) \quad \delta_x(\delta_x - \frac{1}{5})(\delta_x - \frac{3}{5})(\delta_x - \frac{4}{5}) - x(\delta_x + \frac{1}{10})^3(\delta_x + \frac{3}{5})$$

by replacing x by λ^{10} . To find the equation satisfied by (7.2), make the change of variable $x \rightarrow \lambda^{10}$ in (7.3):

$$(7.4) \quad \delta_\lambda(\delta_\lambda - 2)(\delta_\lambda - 6)(\delta_\lambda - 8) - \lambda^{10}(\delta_\lambda + 1)^3(\delta_\lambda + 6),$$

where $\delta_\lambda = \lambda \frac{\partial}{\partial \lambda}$. We show in the next section that (7.4) is the differential equation satisfied by $x^{\mathbf{b}_1} \in \mathcal{W}'$.

For any two complex numbers c_1 and c_2 we have an equality of operators $(\delta_\lambda + c_1) \circ \lambda^{c_2} = \lambda^{c_2} \circ (\delta_\lambda + c_1 + c_2)$. Taking $c_2 = -1$, it follows that the series $\lambda {}_4F_3(\frac{1}{10}, \frac{1}{10}, \frac{1}{10}, \frac{3}{5}; \frac{1}{5}, \frac{2}{5}, \frac{4}{5}; \lambda^{10})$ satisfies the equation

$$(7.5) \quad (\delta_\lambda - 1)(\delta_\lambda - 3)(\delta_\lambda - 7)(\delta_\lambda - 9) - \lambda^{10}\delta_\lambda^3(\delta_\lambda + 5).$$

Equation (7.5) differs from the equation computed by Gährs ([9, unnumbered equation following Equation (2.4)]) by some constant factors, due to the fact that our normalization of f_λ differs from that of Gährs. The shift from our equation (7.4) to the Gährs equation (7.5) is owing to the fact that Gährs is computing, in the notation of [9], the differential equation satisfied by $s\Omega_0/f$ rather than the equation satisfied by Ω_0/f .

To further illustrate our results, we compute the elements of \mathcal{B} that are congruent to \mathbf{b}_1 modulo $\mathbb{Z}A_+$:

$$\begin{aligned} \mathbf{b}_2 &:= (2, 2, 2, 0) \equiv \mathbf{b}_1 + \mathbf{a}_0 & \mathbf{b}_3 &:= (3, 3, 3, 1) \equiv \mathbf{b}_1 + 2\mathbf{a}_0 & \mathbf{b}_4 &:= (4, 4, 4, 0) \equiv \mathbf{b}_1 + 3\mathbf{a}_0 \\ \mathbf{b}_5 &:= (0, 0, 4, 1) \equiv \mathbf{b}_1 + 4\mathbf{a}_0 & \mathbf{b}_6 &:= (1, 1, 5, 0) \equiv \mathbf{b}_1 + 5\mathbf{a}_0 & \mathbf{b}_7 &:= (2, 2, 6, 1) \equiv \mathbf{b}_1 + 6\mathbf{a}_0 \\ \mathbf{b}_8 &:= (3, 3, 7, 0) \equiv \mathbf{b}_1 + 7\mathbf{a}_0 & \mathbf{b}_9 &:= (4, 4, 8, 1) \equiv \mathbf{b}_1 + 8\mathbf{a}_0 & \mathbf{b}_{10} &:= (0, 0, 0, 0) \equiv \mathbf{b}_1 + 9\mathbf{a}_0. \end{aligned}$$

Of these, \mathbf{b}_j is an interior point of $P(A)$ for $j = 1, 3, 7, 9$ (so $R = 4$) and is a boundary point for $j = 2, 4, 5, 6, 8, 10$. By Proposition 6.14(a), the factors in (7.2) that cancel are

$$\left\{ \left(1 - \frac{s_0^{(j)}}{\ell_0}\right) \mid j = 2, 4, 5, 6, 8, 10 \right\} = \left\{ \frac{1}{10}, \frac{3}{10}, \frac{5}{10}, \frac{6}{10}, \frac{7}{10}, \frac{9}{10} \right\}.$$

8. DIFFERENTIAL EQUATION SATISFIED BY $x^{\mathbf{b}}$

Let $u \in M_k$ (for some k) and let R_k be the number of elements of \mathcal{B}_k that are interior to $P(A)$ and congruent to u modulo $\mathbb{Z}A_+$. In the next section, we prove the following result.

Proposition 8.1. (a) *If $u \in M_k$ is an interior point of $C(A)$, then $x^u \in \mathcal{W}'_k$ satisfies a differential equation of order R_k .*

(b) *If $u \in M_k$ is a boundary point of $C(A)$, then $x^u \in \mathcal{W}'_k$ satisfies a differential equation of order $R_k + 1$.*

Let $\mathbf{b} \in \mathcal{B}_k$ for some k and suppose that \mathbf{b} is an interior point of $P(A)$. By Proposition 8.1(a), there is a differential operator

$$\partial_\lambda^{R_k} + c_{R_k-1}(\lambda)\partial_\lambda^{R_k-1} + \cdots + c_0(\lambda),$$

where the $c_i(\lambda)$ lie in $\mathbb{C}(\lambda)$, that annihilates $x^{\mathbf{b}} \in \mathcal{W}'$. (Recall that ∂_λ acts on \mathcal{W}'_k as D_λ .) If $\xi \in \text{Hom}_{\mathcal{D}}(\mathcal{W}_k, \mathbb{C}[[\lambda]])$, then this differential operator will also annihilate the series $\xi(x^{\mathbf{b}})$. By Section 6, as \mathbf{b}' runs through the elements of \mathcal{B}_k , the power series $\xi_{\mathbf{b}'}(x^{\mathbf{b}'})$ that are nonzero (which occurs exactly when \mathbf{b}' is an interior point of $P(A)$) run through the nonzero elements of row \mathbf{b} of the fundamental solution matrix computed in Section 6. A similar argument applies when \mathbf{b} is a boundary point of $P(A)$ by using Proposition 8.1(b). This gives the following result.

Theorem 8.2. *For $\mathbf{b} \in \mathcal{B}_k$, the element $x^{\mathbf{b}} \in \mathcal{W}'_k$ satisfies the differential equation whose complete set of solutions is given by the nonzero elements of row \mathbf{b} of the fundamental solution matrix computed in Section 6.*

9. PROOF OF PROPOSITION 8.1

Proposition 8.1 is a consequence of the following result. Let $\widehat{\mathcal{W}}_k \subset \mathcal{W}'_k$ be the $\mathbb{C}(\lambda)$ -subspace generated by those x^u , $u \in M_k$, for which u is an interior point of $C(A)$.

Proposition 9.1. $\dim_{\mathbb{C}(\lambda)} \widehat{\mathcal{W}}_k = R_k$.

Proof of Proposition 8.1. If u is an interior point of $C(A)$, then the $R_k + 1$ elements

$$x^u, D_\lambda(x^u) = -\ell_0 x^{\mathbf{a}_0}, \dots, D_\lambda^{R_k}(x^u) = (-\ell_0)^{R_k} x^{u+R_k \mathbf{a}_0}$$

are interior points of $C(A)$, so by Proposition 9.1 there is a $\mathbb{C}(\lambda)$ -linear relation between them. This proves part (a).

If u is a boundary point of $C(A)$, then $D_\lambda(x^u), D_\lambda^2(x^u), \dots, D_\lambda^{R_k+1}(x^u)$ are interior points of $C(A)$, so by Proposition 9.1 there is a $\mathbb{C}(\lambda)$ -linear relation between them. This proves part (b). \square

Proposition 9.1 is a consequence of the appendix to [2], which contains a more general result [2, Theorem A.9]. We give here a simpler version of that result.

For any subset $B \subseteq A$, let $V_B \subseteq V$ be the subspace generated by B , so $\dim_{\mathbb{R}} V_B = \text{card}(B)$. Let $V_{B, \mathbb{Z}} = V_B \cap \mathbb{Z}$ and let $C(B)$ be the real cone generated by B . The cone $C(B)$ is a face of $C(A)$ of dimension $\text{card}(B)$. Put $M_B = V_{B, \mathbb{Z}} \cap C(B)$ and let S'_B be the $\mathbb{C}(\lambda)$ -algebra generated by $\{x^u \mid u \in M_B\}$. Let $\theta_B : S' \rightarrow S'_B$ be the $\mathbb{C}(\lambda)$ -homomorphism defined by

$$\theta_B(x^u) = \begin{cases} x^u & \text{if } u \in M_B, \\ 0 & \text{if } u \notin M_B. \end{cases}$$

For $j = 1, \dots, m$, we define $\tilde{D}_j^B = \theta_B \circ \tilde{D}_j$. It follows from Lemma 5.29 that for $u \in M_B$, say $u = \sum_{j: \mathbf{a}_j \in B} u'_j \mathbf{a}_j$, we have

$$(9.2) \quad \tilde{D}_j^B(x^u) = \begin{cases} u'_j x^u + \ell_j x^{u+\mathbf{a}_j} & \text{if } \mathbf{a}_j \in B, \\ 0 & \text{if } \mathbf{a}_j \notin B. \end{cases}$$

Consider $\text{Kos}^\bullet(S'_B, \{\tilde{D}_j^B\}_{\mathbf{a}_j \in B})$, the Koszul complex on S'_B defined by the operators $\{\tilde{D}_j^B\}_{\mathbf{a}_j \in B}$. It follows from (9.2) that the associated graded complex is the Koszul complex on S'_B defined by the $\{\ell_j \mathbf{a}_j\}_{\mathbf{a}_j \in B}$. Since these monomials form a regular sequence on S'_B , both these Koszul complexes are acyclic except in top dimension, where we have

$$(9.3) \quad \dim_{\mathbb{C}(\lambda)} H^{|B|}(\text{Kos}^\bullet(S'_B, \{\tilde{D}_j^B\}_{\mathbf{a}_j \in B})) = |\mathcal{B} \cap C(B)|.$$

We define subcomplexes of these Koszul complexes. Let B_1 be a subset of B and let S'_{B, B_1} be the $C(\lambda)$ -subalgebra of S'_B generated by those x^u , $u \in M_B$, for which $u = \sum_{j: \mathbf{a}_j \in B} u'_j \mathbf{a}_j$ with $u'_j > 0$ when $\mathbf{a}_j \in B_1$.

It follows from (9.2) that \tilde{D}_j for $\mathbf{a}_j \in B$ maps S_{B, B_1} to $S_{B, B_1 \cup \{\mathbf{a}_j\}}$. This allows us to define a subcomplex of $\text{Kos}^\bullet(S'_B, \{\tilde{D}_j^B\}_{\mathbf{a}_j \in B})$. Define

$$\mathcal{K}^k(B, B_1) = \bigoplus_{\substack{\mathbf{a}_{j_1}, \dots, \mathbf{a}_{j_k} \in B \\ j_1 < \dots < j_k}} S'_{B, B_1 \cap \{\mathbf{a}_{j_1}, \dots, \mathbf{a}_{j_k}\}} e_{j_1} \wedge \dots \wedge e_{j_k}.$$

Then $\mathcal{K}^\bullet(B, B_1)$ is a subcomplex of $\text{Kos}^\bullet(S'_B, \{\tilde{D}_j^B\}_{\mathbf{a}_j \in B})$. Note that, in particular,

$$(9.4) \quad \mathcal{K}^\bullet(B, \emptyset) = \text{Kos}^\bullet(S'_B, \{\tilde{D}_j^B\}_{\mathbf{a}_j \in B}).$$

Let $\mathbf{a}_j \in B_1$. The map $\theta_{B \setminus \{\mathbf{a}_j\}}$ maps S'_B onto $S'_{B \setminus \{\mathbf{a}_j\}}$ and induces a surjective map

$$\mathcal{K}^\bullet(B, B_1 \setminus \{\mathbf{a}_j\}) \rightarrow \mathcal{K}^\bullet(B \setminus \{\mathbf{a}_j\}, B_1 \setminus \{\mathbf{a}_j\})[1]$$

whose kernel is $\mathcal{K}^\bullet(B, B_1)$. There is thus a short exact sequence

$$(9.5) \quad 0 \rightarrow \mathcal{K}^\bullet(B, B_1) \rightarrow \mathcal{K}^\bullet(B, B_1 \setminus \{\mathbf{a}_j\}) \rightarrow \mathcal{K}^\bullet(B \setminus \{\mathbf{a}_j\}, B_1 \setminus \{\mathbf{a}_j\})[1] \rightarrow 0.$$

From (9.3) and (9.4) we have

$$(9.6) \quad H^i(\mathcal{K}^\bullet(B, \emptyset)) = 0 \quad \text{for } i = 0, 1, \dots, |B| - 1$$

and

$$(9.7) \quad \dim_{\mathbb{C}(\lambda)} H^{|B|}(\mathcal{K}^\bullet(B, \emptyset)) = |\mathcal{B} \cap C(B)|.$$

Using the long exact sequence of cohomology associated to (9.5) and induction on $|B_1|$ (the case $|B_1| = 0$ being (9.6) and (9.7)) gives the following result.

Theorem 9.8. *We have*

$$(9.9) \quad H^i(\mathcal{K}^\bullet(A, A)) = 0 \quad \text{for } i = 0, 1, \dots, m - 1$$

and

$$(9.10) \quad \dim_{\mathbb{C}(\lambda)} H^m(\mathcal{K}^\bullet(A, A)) = \sum_{B \subseteq A} (-1)^{|A \setminus B|} |\mathcal{B} \cap C(B)|.$$

Note that the right-hand side equals $|\mathcal{B} \cap C(A)^\circ|$, the number of points of \mathcal{B} that lie in the interior of the cone $C(A)$.

Corollary 9.11. *We have*

$$\dim_{\mathbb{C}(\lambda)} H^m(\mathcal{K}^\bullet(A, A)) = |\mathcal{B} \cap C(A)^\circ|.$$

Proof of Proposition 9.1. Since the elements of $\mathcal{B} \cap C(A)^\circ$ are linearly independent in $\widehat{\mathcal{W}}$, we have

$$\dim_{\mathbb{C}(\lambda)} \widehat{\mathcal{W}} \geq |\mathcal{B} \cap C(A)^\circ|.$$

On the other hand, there is a natural surjection

$$H^m(\mathcal{K}^\bullet(A, A)) = S'_{A, A} / \sum_{j=1}^m \tilde{D}_j S'_{A, A \setminus \{\mathbf{a}_j\}} \rightarrow \widehat{\mathcal{W}} = S'_{A, A} / (S'_{A, A} \cap \sum_{j=1}^m \tilde{D}_j S_{A, \emptyset}),$$

which implies that

$$\dim_{\mathbb{C}(\lambda)} \widehat{\mathcal{W}} \leq |\mathcal{B} \cap C(A)^\circ|.$$

Proposition 9.1 now follows from the direct sum decomposition discussed in Section 4. \square

10. SOLUTIONS AT INFINITY

We begin with some general remarks. Consider a differential operator

$$(10.1) \quad (\delta_x + \beta_1 - 1) \cdots (\delta_x + \beta_n - 1) - x(\delta_x + \alpha_1) \cdots (\delta_x + \alpha_n).$$

Let $F(x)$ be a solution of this operator and let ℓ be a positive integer.

Lemma 10.2. *The function $F(\lambda^\ell)$ satisfies the differential operator*

$$(10.3) \quad (\delta_\lambda + \ell\beta_1 - \ell) \cdots (\delta_\lambda + \ell\beta_n - \ell) - \lambda^\ell(\delta_\lambda + \ell\alpha_1) \cdots (\delta_\lambda + \ell\alpha_n).$$

Proof. Make the change of variable $x = \lambda^\ell$ in (10.1). □

We want to describe the solutions of (10.3) at infinity. Consider the differential operator

$$(10.4) \quad (\delta_x + \tau_1 - 1) \cdots (\delta_x + \tau_n - 1) - x(\delta_x + \sigma_1) \cdots (\delta_x + \sigma_n).$$

Let $G(x)$ be a solution of this differential operator.

Lemma 10.5. *The function $G(\lambda^{-\ell})$ satisfies the differential operator*

$$(10.6) \quad (\delta_\lambda - \ell\sigma_1) \cdots (\delta_\lambda - \ell\sigma_n) - \lambda^\ell(\delta_\lambda - \ell\tau_1 + \ell) \cdots (\delta_\lambda - \ell\tau_n + \ell).$$

Proof. Make the change of variable $x = \lambda^{-\ell}$ in (10.4). □

Corollary 10.7. *Let $G(x)$ be a solution of the differential operator*

$$(10.8) \quad (\delta_x - \alpha_1) \cdots (\delta_x - \alpha_n) - x(\delta_x + 1 - \beta_1) \cdots (\delta_x + 1 - \beta_n).$$

Then $G(\lambda^{-\ell})$ satisfies (10.3).

Proof. Taking $\sigma_i = 1 - \beta_i$ and $\tau_i = 1 - \alpha_i$ in (10.4) transforms (10.6) into (10.3). □

To make the solutions of (10.8) slightly simpler to express, we make one further change. Assume that the α_i are real numbers and that, say, $\alpha_1 \leq \alpha_i$ for all i . Consider the differential operator obtained from (10.8) by adding α_1 to each term:

$$(10.9) \quad \delta_x(\delta_x - \alpha_2 + \alpha_1) \cdots (\delta_x - \alpha_n + \alpha_1) - x(\delta_x + 1 - \beta_1 + \alpha_1) \cdots (\delta_x + 1 - \beta_n + \alpha_1).$$

One checks that (10.9) is the operator obtained from (10.8) by multiplying (10.8) on the right by x^{α_1} and on the left by $x^{-\alpha_1}$. This implies that if $H(x)$ is a solution of the operator (10.9), then $x^{\alpha_1}H(x)$ is a solution of the operator (10.8). Corollary 10.7 then implies the following result.

Corollary 10.10. *Let $H(x)$ be a solution of the operator (10.9). Then $\lambda^{-\ell\alpha_1}H(\lambda^{-\ell})$ satisfies (10.3).*

Example. We illustrate these observations with the example from Section 7. If we take $n = 4$ and $\beta_1 = 1$, $\beta_2 = 4/5$, $\beta_3 = 2/5$, $\beta_4 = 1/5$, $\alpha_1 = \alpha_2 = \alpha_3 = 1/10$, and $\alpha_4 = 3/5$ in (10.1), then (10.1) becomes (7.3). If we then take $\ell = 10$ in (10.3), then (10.3) becomes (7.4). Equation (10.9) then becomes

$$(10.11) \quad \delta_x^3(\delta_x - \frac{1}{2}) - x(\delta_x + \frac{1}{10})(\delta_x + \frac{3}{10})(\delta_x + \frac{7}{10})(\delta_x + \frac{9}{10}).$$

Corollary 10.10 implies that we can get a full set of solutions to (7.4) at infinity by taking a full set of solutions to (10.11) at the origin, replacing the variable by λ^{-10} , and multiplying by λ^{-1} . For example, one solution to (10.11) is ${}_4F_3(\frac{1}{10}, \frac{3}{10}, \frac{7}{10}, \frac{9}{10}; \frac{1}{2}, 1, 1; x)$, so one solution to (7.4) at infinity is

$$\lambda^{-1} {}_4F_3(\frac{1}{10}, \frac{3}{10}, \frac{7}{10}, \frac{9}{10}; \frac{1}{2}, 1, 1; \lambda^{-10}).$$

11. CONNECTION WITH DE RHAM COHOMOLOGY

We describe here an example where we apply the ideas of this paper to compute a fundamental solution matrix for the Picard-Fuchs equation of a projective variety. Consider the projective hypersurface $X \subseteq \mathbb{P}_{\mathbb{C}(\lambda)}^{n-1}$ defined by the equation

$$f_\lambda = \sum_{j=1}^n w_j x_j^d - d\lambda x_1^{w_1} \cdots x_n^{w_n} = 0,$$

where $w_j \geq 1$ for $j = 1, \dots, n$ and $\sum_{j=1}^n w_j = d$ (see Katz [11] for an ℓ -adic treatment of this hypersurface).

Let $\widetilde{\mathcal{W}} \subseteq \mathcal{W}'$ be the subspace generated by those x^u , $u = (u_1, \dots, u_n) \in M$, for which $u_j \geq 1$ for $j = 1, \dots, n$ and $\sum_{j=1}^n u_j$ is a multiple of d . Note that M in this case is the set of all lattice points in the first orthant. By Katz [10, Corollary 1.15], there is an isomorphism of \mathcal{D}' -modules $\widetilde{\mathcal{W}} \cong H_{\text{dR}}^{n-1}(\mathbb{P}_{\mathbb{C}(\lambda)}^{n-1} \setminus X)$. Specifically, the element $x^u \in \widetilde{\mathcal{W}}$ with $\sum_{j=1}^n u_j = kd$ corresponds (up to a scalar multiple) to the $(n-1)$ -form

$$\frac{x^u}{f_\lambda^k} \sum_{j=1}^n (-1)^j \frac{dx_1}{x_1} \cdots \frac{dx_j}{x_j} \cdots \frac{dx_n}{x_n}.$$

One can thus obtain a fundamental solution matrix for the Picard-Fuchs equation of $H_{\text{dR}}^{n-1}(\mathbb{P}_{\mathbb{C}(\lambda)}^{n-1} \setminus X)$, or, via the residue map, of the primitive part of $H_{\text{dR}}^{n-2}(X)$, using the results of this paper. By Section 9, a basis for $\widetilde{\mathcal{W}}$ is

$$\widetilde{\mathcal{B}} = \left\{ x^u \mid u = (u_1, \dots, u_n) \text{ with } 0 < u_j < d \text{ for } j = 1, \dots, n \text{ and } d \text{ divides } \sum_{j=1}^n u_j \right\}.$$

The set $\widetilde{\mathcal{B}}$ decomposes following the decomposition of \mathcal{W}' in Section 4. One can then write down the solution matrix of the Picard-Fuchs equation using Equations (6.7) and (4.9).

Example. Let $X \subseteq \mathbb{P}_{\mathbb{C}(\lambda)}^2$ be the plane curve of genus 10 defined by the equation

$$f_\lambda = x_1^6 + 2x_2^6 + 3x_3^6 - 6\lambda x_1 x_2^2 x_3^3 = 0.$$

Thus $\mathbf{a}_1 = (6, 0, 0)$, $\mathbf{a}_2 = (0, 6, 0)$, $\mathbf{a}_3 = (0, 0, 6)$, and $\mathbf{a}_0 = (1, 2, 3)$. The relation

$$6\mathbf{a}_0 = \mathbf{a}_1 + 2\mathbf{a}_2 + 3\mathbf{a}_3$$

gives $\ell_1 = 1$, $\ell_2 = 2$, $\ell_3 = 3$, and $\ell_0 = 6$. The set $\widetilde{\mathcal{B}}$ has 20 elements. Under the decomposition of Section 4, it decomposes into 6 subsets:

$$\begin{aligned} \widetilde{\mathcal{B}}_1 &= \{(1, 2, 3), (5, 4, 3)\}, & \widetilde{\mathcal{B}}_2 &= \{(2, 1, 3), (4, 5, 3)\}, \\ \widetilde{\mathcal{B}}_3 &= \{(2, 2, 2), (3, 4, 5), (5, 2, 5)\}, & \widetilde{\mathcal{B}}_4 &= \{(3, 2, 1), (4, 4, 4), (1, 4, 1)\}, \\ \widetilde{\mathcal{B}}_5 &= \{(1, 3, 2), (2, 5, 5), (3, 1, 2), (4, 3, 5), (5, 5, 2)\}, \\ \widetilde{\mathcal{B}}_6 &= \{(2, 3, 1), (3, 5, 4), (4, 1, 1), (5, 3, 4), (1, 1, 4)\}. \end{aligned}$$

The Picard-Fuchs equation thus has a block diagonal form, with two (2×2) -blocks, two (3×3) -blocks, and two (5×5) -blocks. We compute the solution matrix for one

of these blocks, the block associated to $\tilde{\mathcal{B}}_1$. From Equations (6.7) and (4.9) we get

$$\begin{pmatrix} {}_2F_1\left(\frac{1}{6}, \frac{1}{6}; \frac{1}{3}; \lambda^6\right) & -27\lambda^4 {}_2F_1\left(\frac{5}{6}, \frac{5}{6}; \frac{5}{3}; \lambda^6\right) \\ -\frac{1}{2}\lambda^2 {}_2F_1\left(\frac{7}{6}, \frac{7}{6}; \frac{4}{3}; \lambda^6\right) & {}_2F_1\left(\frac{5}{6}, \frac{5}{6}; \frac{2}{3}; \lambda^6\right) \end{pmatrix}.$$

The series in the first row are solutions of the operator

$$\delta_x(\delta_x - \frac{2}{3}) - x(\delta_x + \frac{1}{6})^2$$

with the variable x replaced by λ^6 . The series in the second row are solutions of the operator

$$\delta_x(\delta_x - \frac{1}{3}) - x(\delta_x + \frac{5}{6})^2$$

with the variable x replaced by λ^6 .

REFERENCES

- [1] Alan Adolphson. Hypergeometric functions and rings generated by monomials. *Duke Math. J.* **73** (1994), no. 2, 269–290.
- [2] Alan Adolphson and Steven Sperber. Exponential sums and Newton polyhedra: cohomology and estimates. *Ann. of Math. (2)* **130** (1989), no. 2, 367–406.
- [3] Alan Adolphson and Steven Sperber. Dwork cohomology, de Rham cohomology, and hypergeometric functions. *Amer. J. Math.* **122** (2000), no. 2, 319–348.
- [4] Alan Adolphson and Steven Sperber. Newton polytopes and algebraic hypergeometric series. With an appendix by Nicholas M. Katz. *Trans. Amer. Math. Soc.* **373** (2020), no. 12, 8365–8389.
- [5] Alan Adolphson and Steven Sperber. On solutions of codimension-one A -hypergeometric systems. Preprint (arXiv:2202.08389).
- [6] Victor Batyrev. Variations of the mixed Hodge structure of affine hypersurfaces in algebraic tori. *Duke Math. J.* **69** (1993), no. 2, 349–409.
- [7] Charles Doran, Tyler Kelly, Adriana Salerno, Steven Sperber, John Voight, and Ursula Whitcher. Hypergeometric decomposition of symmetric K3 quartic pencils. *Res. Math. Sci.* **7** (2020), no. 2, Paper No. 7, 81 pp.
- [8] Bernard Dwork. On the zeta function of a hypersurface. II. *Ann. of Math. (2)* **80** (1964), 227–299.
- [9] Swantje Gähns. Picard-Fuchs equations of special one-parameter families of invertible polynomials. Arithmetic and geometry of K3 surfaces and Calabi-Yau threefolds, 285–310, *Fields Inst. Commun.*, 67, Springer, New York, 2013.
- [10] Nicholas Katz. On the differential equations satisfied by period matrices. *Inst. Hautes Études Sci. Publ. Math.* No. 35 (1968), 223–258.
- [11] Nicholas Katz. Another look at the Dwork family. *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II*, 89–126, *Progr. Math.*, 270, Birkhäuser Boston, Boston, MA, 2009.
- [12] Remke Kloosterman. The zeta function of monomial deformations of Fermat hypersurfaces. *Algebra Number Theory* **1** (2007), no. 4, 421–450.
- [13] A. G. Kouchnirenko. Polyèdres de Newton et nombres de Milnor. *Invent. Math.* **32** (1976), no. 1, 1–31.
- [14] Kazuaki Miyatani. Monomial deformations of certain hypersurfaces and two hypergeometric functions. *Int. J. Number Theory* **11** (2015), no. 8, 2405–2430.
- [15] Ryo Negishi. Picard-Fuchs equations of the Dwork family. Preprint (arXiv:2208.01531).

DEPARTMENT OF MATHEMATICS, OKLAHOMA STATE UNIVERSITY, STILLWATER, OKLAHOMA 74078
 Email address: `adolphs@math.okstate.edu`

SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MINNESOTA 55455
 Email address: `sperber@math.umn.edu`