BIRATIONAL INDUCTION OF NILPOTENT ORBIT COVERS IN EXCEPTIONAL TYPES

MATTHEW WESTAWAY

ABSTRACT. Let G be a semisimple simply connected algebraic group over \mathbb{C} of exceptional type. For each G-equivariant nilpotent cover of a nilpotent coadjoint G-orbit \mathbb{O} , we determine the unique birationally rigid induction datum from which it is birationally induced.

1. INTRODUCTION

Lusztig-Spaltenstein induction was introduced in 1979 by Lusztig and Spaltenstein [LS]. While it was originally defined in the context of unipotent classes, it can be easily translated into the language of nilpotent orbits in reductive Lie algebras [CM], and in this context has had significant applications in Lie-theoretic representation theory. It works in this setting as follows. Let G be a reductive algebraic group over \mathbb{C} ; this acts on its Lie algebra $\mathfrak{g} = \text{Lie}(G)$ via the adjoint action¹ and thus partitions G into orbits. Of these, we pay particular attention to the **nilpotent** orbits, i.e. those orbits consisting of nilpotent elements of \mathfrak{g} . Given a parabolic subgroup P of G with Levi decomposition P = LU, where L is itself a reductive algebraic group over \mathbb{C} , we may also consider the nilpotent L-orbits in $\mathfrak{l} = \text{Lie}(L)$. Lusztig-Spaltenstein induction is then a process which takes as input a nilpotent L-orbit \mathbb{O}_L and produces from it a nilpotent G-orbit $\mathbb{O} = \text{Ind}_L^G(\mathbb{O}_L)$ (we call (L, \mathbb{O}_L) an induction datum for \mathbb{O}). This procedure satisfies a number of nice properties, including that it is transitive, independent of the parabolic subgroup P, and behaves predictably on the dimensions of orbits.

One reason why this procedure has turned out to be so useful is that it often allows us to reduce questions about nilpotent orbits to the case of **rigid** nilpotent orbits, i.e. those orbits that cannot be obtained (non-trivially) via Lusztig-Spaltenstein induction. For example, this was a significant tool in resolving the longstanding problem of showing that all finite W-algebras have a one-dimensional representation [L1, P2, P3] and the related problem in modular representation theory of finding the minimal dimension for representations of reduced enveloping algebras of Lie algebras over fields of positive characteristic [P1, PT2]. Lusztig-Spaltenstein induction also plays an essential role in the classification of sheets [Bo], i.e. the irreducible components of $\mathfrak{g}_{(m)} := \{x \in \mathfrak{g} \mid \dim G \cdot x = m\}$ for $m \in \mathbb{N}$, and is closely related to parabolic induction of primitive ideals [BJ].

One aspect of Lusztig-Spaltenstein induction which can be both a feature and a bug is that a given nilpotent orbit \mathbb{O} can be induced from multiple different rigid orbits. In order to avoid some of the problems this causes, one option is to look instead at **birational induction**. As initially introduced (see, for example, [L2]), being birational was a property that a given induction from one nilpotent orbit to another could have or not have, but this was refined in [LMM] to a procedure similar to Lusztig-Spaltenstein induction except applying to different objects: nilpotent orbit covers.

Date: December 5, 2023.

²⁰²⁰ Mathematics Subject Classification. Primary 17B08; Secondary 17B20; 22E60.

 $Key\ words\ and\ phrases.$ Lie algebras, nilpotent orbits, nilpotent covers, Lusztig-Spaltenstein induction, birational induction.

¹In the substance of this paper we prefer to work with the coadjoint action of G on \mathfrak{g}^* , but these two actions can be identified by means of a non-degenerate G-invariant symmetric bilinear form.

A (*G*-equivariant) nilpotent orbit cover (for a reductive algebraic group *G*) is a homogeneous space $\widetilde{\mathbb{O}}$ equipped with a *G*-equivariant finite morphism to a nilpotent *G*-orbit \mathbb{O} ; in particular, nilpotent orbits are (trivial) covers of themselves. Birational induction then sends a nilpotent orbit cover $\widetilde{\mathbb{O}}_L \to \mathbb{O}_L$ for a Levi subgroup *L* in *G* to a nilpotent orbit cover $\operatorname{Bind}_L^G(\widetilde{\mathbb{O}}_L) \to \operatorname{Ind}_L^G(\mathbb{O}_L)$, and we call $(L, \widetilde{\mathbb{O}}_L)$ a birational induction datum for $\operatorname{Bind}_L^G(\widetilde{\mathbb{O}}_L)$. This procedure has similar properties to Lusztig-Spaltenstein induction: it is transitive, it is independent of the parabolic subgroup *P* of which *L* is a Levi factor, and it behaves predictably on degrees of nilpotent covers. It also has an additional property which we don't see for Lusztig-Spaltenstein induction, namely that each nilpotent orbit cover has a **unique** birational induction datum (up to conjugacy) which is birationally rigid, i.e. which cannot be non-trivially obtained through birational induction.

This refinement of Lusztig-Spaltenstein induction is used significantly in [LMM, MM]. Losev, Mason-Brown and Matvieievskyi introduce in [LMM] a definition of unipotent ideals in $U(\mathfrak{g})$, in order to define the notion of unipotent representations of a complex reductive Lie algebra. Birational induction is one of the key tools which allows for the computation of the central characters of these unipotent ideals; this also has connections with the central characters for one-dimensional representations of finite W-algebras. Parabolic induction of unipotent ideals and of representations of finite W-algebras as in [L1] also appear to be related to birational induction. It has also proved key in constructing and understanding Losev's orbit method map [L2].

When G is a semisimple algebraic group of exceptional type, the rigid induction data for each nilpotent G-orbit \mathbb{O} are known and can be found in tables in [El, DE]. In this paper, for G a semisimple simply connected algebraic group of exceptional type, we determine the birationally rigid birational induction datum for each nilpotent orbit cover of an induced nilpotent orbit, which we compile in Tables 1 through 5. In other words, we have the following theorem.

Theorem 1.1. Let G be a semisimple simply connected algebraic group over \mathbb{C} of exceptional type and let $\widetilde{\mathbb{O}} \to \mathbb{O}$ be a G-equivariant nilpotent orbit cover, where \mathbb{O} is not a rigid orbit. Then the (unique) birationally rigid birational induction datum for $\widetilde{\mathbb{O}}$ is as given in Tables 1 through 5.

We note that Tables 1 through 5 only include the nilpotent orbit covers for induced nilpotent orbits, since nilpotent orbit covers of rigid nilpotent orbits are always birationally rigid. In order to describe the nilpotent covers, we appeal to the fact that such covers of an orbit \mathbb{O} (up to isomorphism) are in bijection with subgroups of the component group of the centralizer of an element $e \in \mathbb{O}$ (up to conjugacy).

Before ending the introduction, let us say a word about what happens when G is of classical type. When $G = \operatorname{SL}_n(\mathbb{C})$, each orbit is birationally induced from the zero orbit for a suitable Levi subgroup; to determine the birationally rigid birational induction datum for a non-trivial cover, one would want to determine the possible birationally rigid induction data which are not orbits and then use [MMY, Proposition 2.6]. It is likely that this could be done explicitly using [LMM, Proposition 7.6.4]. When $G = \operatorname{SO}_n(\mathbb{C})$ or $\operatorname{Sp}_{2n}(\mathbb{C})$, the component group of the centralizer of a nilpotent element is always a product of copies of $\mathbb{Z}/2\mathbb{Z}$. Since we only describe this group up to isomorphism, we cannot distinguish between subgroups of the same order. Perhaps the most we can hope for is to determine, for each birationally rigid induction datum $(L, \widetilde{\mathbb{O}}_L)$, the degree of the cover $\operatorname{Bind}_L^G(\widetilde{\mathbb{O}}_L)$. Each Levi subgroup of $\operatorname{Sp}_{2n}(\mathbb{C})$ is of the form

$$\operatorname{GL}_{i_1}(\mathbb{C}) \times \cdots \times \operatorname{GL}_{i_r}(\mathbb{C}) \times \operatorname{Sp}_{2m}(\mathbb{C})$$

for some i_1, \ldots, i_r and m such that $n = i_1 + \cdots + i_r + m$, and each birationally rigid cover for such Levi subgroup is of the form

$$\{0\} imes \cdots imes \{0\} imes \widetilde{\mathbb{O}}_L$$

where \mathbb{O}_L is a birationally rigid cover of a nilpotent $\operatorname{Sp}_{2m}(\mathbb{C})$ -orbit $\mathbb{O}_L \subseteq \mathfrak{sp}_{2m}(\mathbb{C})$. The orbits \mathbb{O}_L can be obtained from $\mathbb{O} = \operatorname{Ind}_L^G(\mathbb{O}_L)$ by means of (steps of) the Kempken-Spaltenstein algorithm introduced in [PT1]. This algorithm consists of making two types of reductions to a partition, called Type 1 and Type 2 reductions in [GTW]. Let k be the number of Type 2 reductions needed to get from \mathbb{O} to \mathbb{O}_L through the Kempken-Spaltenstein algorithm; it should follow from [N2, Claim 2.3.1, 2.3.2] and [MMY, Proposition 7.6.4] that

$$\deg(\operatorname{Bind}_{L}^{G}(\{0\}\times\cdots\times\{0\}\times\widetilde{\mathbb{O}}_{L})) = \deg(\widetilde{\mathbb{O}}_{L}) + 2k$$

For differences when $G = SO_n(\mathbb{C})$, see [N2, §3].

This paper is structured as follows. In Section 2, we discuss various preliminary matters, including the precise definition of birational induction and certain (mostly known) results which can be used to understand the effect of birational induction on a nilpotent orbit cover. In Section 3, we proceed case-by-case through the nilpotent orbits in exceptional Lie algebras, determining the birationally rigid covers which induce to each of their covers. Some classes of orbits we are able to deal with all at once, but we need to consider some orbits individually. Finally, in Section 4 we compile the results of Section 3 into tables.

Acknowledgements. The author would like to thank Simon Goodwin, Lucas Mason-Brown and Lewis Topley for discussions which were helpful for this paper. This author was supported during this research by a research fellowship from the Royal Commission for the Exhibition of 1851.

2. Preliminaries

2.1. Algebraic groups and Levi subgroups. In this subsection, we establish some notation and conventions for this paper and recall some well-known facts regarding algebraic groups and their Levi subgroups.

Throughout this paper, G denotes a reductive algebraic group over \mathbb{C} and \mathfrak{g} denotes its Lie algebra. We fix a maximal torus T of G and a positive Borel subgroup B of G containing T, and let \mathfrak{h} and \mathfrak{b} be the Lie algebras thereof. We denote by Φ the root system of G corresponding to T and by Φ^+ the subset of positive roots corresponding to B. Furthermore, we denote by Π the subset of simple roots of Φ^+ . Set W to be the Weyl group of G, and denote by $\mathfrak{g}(\mathfrak{g})$ the centre of \mathfrak{g} .

The **parabolic subgroups** of G are the closed subgroups of G which contain a conjugate of the Borel subgroup B; we call those parabolic subgroups containing B itself the **standard parabolic subgroups** of G. Such subgroups are in bijection with subsets of Π and each parabolic subgroup is conjugate to a standard parabolic subgroup. Each parabolic subgroup P has a Levi decomposition P = LU, where U is the unipotent radical of P and L is reductive. We call L the **Levi factor** of P; by a **Levi subgroup** of G we will mean a Levi factor of some parabolic subgroup of G. We call L a **standard Levi subgroup** of G when it is the Levi factor of a standard parabolic subgroup of G; clearly each Levi subgroup of G is conjugate to a standard Levi subgroup of G. To each subset $\Delta \subseteq \Pi$ we may define a standard Levi subgroup L_{Δ} as the Levi factor of the standard parabolic subgroup of G corresponding to this subset of simple roots; each Levi subgroup of G is conjugate to some L_{Δ} . The Weyl group W acts on the set of roots Φ ; given two subsets Δ , Γ of Π , the standard Levi subgroups L_{Δ} and L_{Γ} are G-conjugate if and only if there exists $w \in W$ such that $w(\Gamma) = \Delta$.

We may therefore associate to a Levi subgroup L of G the (unique) Dynkin type of the root system $\mathbb{Z}\Delta \cap \Phi$, where $\Delta \subseteq \Pi$ is such that L is conjugate to L_{Δ} . When G is of exceptional type, this determines L up to conjugacy in almost all cases; however, in type G_2 there are two non-conjugate Levi subgroups of type A_1 , in type F_4 there are two non-conjugate Levi subgroups for each type A_1 , A_2 and $A_2 + A_1$, and in type E_7 there are two non-conjugate Levi subgroups for each type $3A_1$, $A_3 + A_1$ and A_5 . In types G_2 and F_4 these Levi subgroups can be distinguished based on root lengths, and we use a tilde to denote the short roots. In type E_7 , we label the two conjugacy classes of Levi subgroups by $(3A_1)'$, $(A_3 + A_1)'$ and $(A_5)'$ or by $(3A_1)''$, $(A_3 + A_1)''$ and $(A_5)''$, where the latter notation is used when the subset of simple roots is W-conjugate to a subset of the black vertices in the below Dynkin diagram.



We use this notation throughout the paper, including in the tables in Section 4.

The **rank** of a reductive algebraic group G is the dimension of a maximal torus $T \subseteq G$; the semisimple rank of G is the rank of the derived subgroup [G, G]. Given a Levi subgroup L of G, the **semisimple corank** of L is defined to be the difference between the semisimple rank of G and the semisimple rank of L. This is equal to the difference between $|\Pi|$ and the number of simple roots in any subset of Π corresponding to L as above.

2.2. Nilpotent orbits and covers. In this subsection we recall the basics of the theory of nilpotent orbits and nilpotent covers for complex simple Lie algebras. The material on nilpotent orbits is standard and can be found, for example, in [CM]. For the discussion of orbit covers, we follow [LMM].

The algebraic group G acts on \mathfrak{g} via the adjoint action and on \mathfrak{g}^* via the coadjoint action. Since \mathfrak{g} and \mathfrak{g}^* are G-equivariantly isomorphic (via a G-invariant non-degenerate symmetric bilinear form) there is a correspondence between adjoint and coadjoint G-orbits (in \mathfrak{g} and \mathfrak{g}^* respectively); in this paper, we prefer to work with coadjoint G-orbits. A coadjoint G-orbit, usually denoted \mathbb{O} in this paper, is called **nilpotent** if $0 \in \overline{\mathbb{O}}$ (see [CM, §1.3] for various equivalent characterisations). There are only finitely many nilpotent G-orbits in \mathfrak{g}^* ; we denote by Nilp(G) the set of nilpotent G-orbits in \mathfrak{g}^* .

The following proposition is standard and describes what $\operatorname{Nilp}(G)$ looks like for each simple algebraic group G – note that $\operatorname{Nilp}(G)$ depends only on the root system Φ , since all nilpotent G-orbits lie inside $[\mathfrak{g},\mathfrak{g}]$ and the G-action factors through $\operatorname{Ad}(G)$.

Proposition 2.1. The nilpotent G-orbits, when Φ is of classical type, can be indexed as follows:

- (1) When $\Phi = A_n$, $n \ge 1$, there is a bijection between nilpotent G-orbits and partitions of n+1.
- (2) When $\Phi = B_n$, $n \ge 2$, there is a bijection between nilpotent G-orbits and partitions \underline{p} of 2n + 1 such that each even part of p occurs an even number of times.
- (3) When $\Phi = C_n$, $n \ge 3$, there is a bijection between nilpotent G-orbits and partitions \underline{p} of 2n such that each odd part of p occurs an even number of times.
- (4) When $\Phi = D_n$, $n \ge 4$, there is a bijection between nilpotent G-orbits and partitions \underline{p} of 2n such that each even part of \underline{p} occurs an even number of times, except that we count twice all partitions in which all parts of \underline{p} are even (we call such partitions **very even**). We use the labels I and II to differentiate such partitions.

When Φ is of exceptional type, the nilpotent G-orbits are listed explicitly (in Bala-Carter notation) in [CM, §8.4]. There are 20 non-zero nilpotent G-orbits in type E_6 , 44 in type E_7 , 69 in type E_8 , 15 in type F_4 and 4 in type G_2 .

We follow [CM] in labelling the nilpotent G-orbits in exceptional cases via the Bala-Carter labelling – details of this can be found in [CM, §8]. Nilpotent orbits in all cases can also be labelled by weighted Dynkin diagrams and we shall often use this notation as well (see [CM, §3.5] for an explanation of how this description works). This is a labelling of the nodes of the Dynkin diagram of \mathfrak{g} by numbers from the set $\{0, 1, 2\}$. We call \mathbb{O} **even** if all labels on the nodes on its weighted Dynkin diagram are either 0 or 2 (see [CM, §3.8] for equivalent definitions).

The G-equivariant fundamental group $\pi_1^G(\mathbb{O})$ of \mathbb{O} is defined by

$$G_1^G(\mathbb{O}) \coloneqq G_e/G_e^G$$

where e is an element in \mathbb{O} and G_e denotes the stabiliser of e in G. Different choices of $e \in \mathbb{O}$ give isomorphic G-equivariant fundamental groups, thus we only define $\pi_1^G(\mathbb{O})$ up to isomorphism.

Given a nilpotent G-orbit $\mathbb{O} \subseteq \mathfrak{g}^*$, a G-equivariant nilpotent cover of \mathbb{O} is defined to be a homogeneous space $\widetilde{\mathbb{O}}$ equipped with a finite G-equivariant map $\widetilde{\mathbb{O}} \to \mathbb{O}$ (see [LMM, §2.2] for more details). We often shorten this just to **nilpotent cover** when G is clear from context. Two such nilpotent covers $\widetilde{\mathbb{O}}$ and $\widetilde{\mathbb{O}}'$ are said to be isomorphic if there exists a G-equivariant isomorphism $\widetilde{\mathbb{O}} \to \widetilde{\mathbb{O}}'$ such that the following diagram commutes:



We denote by $\operatorname{Cov}(G, \mathbb{O})$ the set of G-equivariant nilpotent covers of \mathbb{O} up to isomorphism and

$$\operatorname{Cov}(G) := \bigcup_{\mathbb{O} \in \operatorname{Nilp}(G)} \operatorname{Cov}(G, \mathbb{O}).$$

Fix $e \in \mathbb{O}$. Isomorphism classes of *G*-equivariant nilpotent covers of \mathbb{O} are in bijection with conjugacy classes of subgroups of $\pi_1^G(\mathbb{O})$ via the following map:

(2.1)
$$\operatorname{Cov}(G, \mathbb{O}) \xrightarrow{\sim} \{\operatorname{Conjugacy classes of subgroups of } \pi_1^G(\mathbb{O})\}$$

$$\mathbb{O} \longmapsto \pi_1^G(\mathbb{O}) \coloneqq G_x/G_e^\circ \subseteq G_e/G_e^\circ = \pi_1^G(\mathbb{O}).$$

Here, $x \in \widetilde{\mathbb{O}}$ lies over $e \in \mathbb{O}$; different choices of x give rise different representatives of the conjugacy class of subgroups of $\pi_1^G(\mathbb{O})$. The inverse map is given by

$$G/H \longleftrightarrow H$$

where \widetilde{H} is the preimage of H under the map $G_e \to G_e/G_e^{\circ}$.

2.3. Lusztig-Spaltenstein induction. In this subsection, we recall the basics of Lusztig-Spaltenstein induction, following [CM, §7].

Let P be a parabolic subgroup of G with Levi decomposition P = LU, where L is the Levi factor of P and U is the unipotent radical of P. Denote the corresponding decomposition of Lie algebras by $\mathfrak{p} = \mathfrak{l} \oplus \mathfrak{u}$. Furthermore, set P^- to be the opposite parabolic of P, with unipotent radical $U^$ and Levi decompositions $P^- = U^-L$ and $\mathfrak{p} = \mathfrak{u}^- \oplus \mathfrak{l}$. Given a subspace V of \mathfrak{g} , denote by V^{\perp} the subspace of \mathfrak{g}^* consisting of those χ such that $\chi(V) = 0$; note that we may identify \mathfrak{p}^{\perp} with $(\mathfrak{u}^-)^*$ and \mathfrak{u}^{\perp} with $(\mathfrak{p}^-)^*$. Using a non-degenerate symmetric bilinear form, these subspaces of \mathfrak{g}^* may be identified with the subspaces \mathfrak{u} and \mathfrak{p} of \mathfrak{g} .

Let $\mathbb{O}_L \subseteq \mathfrak{l}^*$ be a nilpotent *L*-orbit. We may then form the variety $G \times^P (\overline{\mathbb{O}}_L \times \mathfrak{p}^{\perp})$, where *P* acts on *G* via right multiplication and on $\overline{\mathbb{O}}_L \times \mathfrak{p}^{\perp}$ via the coadjoint action. There exists a *G*-equivariant morphism

(2.2)
$$\mu: G \times^P (\overline{\mathbb{O}}_L \times \mathfrak{p}^\perp) \to \mathfrak{g}^*, \qquad (g, \chi) \mapsto g \cdot \chi.$$

The image of this map coincides with $\overline{\mathbb{O}}$ for some nilpotent *G*-orbit $\mathbb{O} \subseteq \mathfrak{g}^*$, by Joseph's irreducibility theorem [Jo]. We then say that \mathbb{O} is **induced** from \mathbb{O}_L and that (L, \mathbb{O}_L) is an **induction datum** for \mathbb{O} . This defines a map (called Lusztig-Spaltenstein induction, or induction)

$$\operatorname{Ind}_{L}^{G}: \operatorname{Nilp}(L) \to \operatorname{Nilp}(G), \quad \mathbb{O}_{L} \mapsto \operatorname{Ind}_{L}^{G}(\mathbb{O}_{L}) = \mathbb{O}.$$

This map has three properties of particular note. First, as implied by the notation, Lusztig-Spaltenstein induction is independent of the parabolic subgroup P of G in which L is a Levi factor.

Second, it is transitive, in the sense that $\operatorname{Ind}_L^G = \operatorname{Ind}_M^G \circ \operatorname{Ind}_L^M$ whenever $L \subseteq M \subseteq G$ is a tower of Levi subgroups in G. Finally, we have the equality

$$\operatorname{Codim}_{\mathfrak{l}^*}(\mathbb{O}_L) = \operatorname{Codim}_{\mathfrak{q}^*}(\operatorname{Ind}_L^G(\mathbb{O}_L)),$$

which can be rephrased as $\dim(\operatorname{Ind}_{L}^{G}(\mathbb{O}_{L})) = \dim \mathbb{O}_{L} + 2\dim \mathfrak{u}$.

A nilpotent *G*-orbit \mathbb{O} is called **rigid** if it cannot be induced from a nilpotent *L*-orbit for a proper Levi subgroup *L* of *G*, i.e. if the only induction datum for \mathbb{O} is (G, \mathbb{O}) . We call an induction datum (L, \mathbb{O}_L) a **rigid induction datum** if \mathbb{O}_L is a rigid nilpotent *L*-orbit in \mathfrak{l}^* . Let us denote the set of rigid induction data for \mathbb{O} by

$$\operatorname{rig}(\mathbb{O}) := \{ (L, \mathbb{O}_L) \mid \operatorname{Ind}_L^G(\mathbb{O}_L) = \mathbb{O} \}.$$

The adjoint (resp. coadjoint) action of G on G (resp. \mathfrak{g}^*) induces an action of G on rig(\mathbb{O}). We denote

$$\operatorname{Rig}(\mathbb{O}) := \operatorname{rig}(\mathbb{O})/G,$$

and we will abuse notation slightly to also refer to elements of $\operatorname{Rig}(\mathbb{O})$ as rigid induction data for \mathbb{O} . By the semisimple corank of an element of $\operatorname{Rig}(\mathbb{O})$, we will mean the semisimple corank of the Levi subgroup in any representative of such equivalence class.

The classification of rigid nilpotent G-orbits is given in [CM, Theorem 7.2.3], [CM, Corollary 7.3.5] and [DE], as follows. As with the classification of nilpotent orbits in general, it depends only on the root system Φ .

Proposition 2.2. The rigid nilpotent G-orbits, when Φ is of classical type, can be indexed as follows:

- (1) When $\Phi = A_n$, $n \ge 1$, the only rigid nilpotent G-orbit is the zero orbit.
- (2) When $\Phi = B_n$, $n \ge 2$, the rigid nilpotent G-orbits correspond to those partitions $\underline{p} = (p_1 \ge p_2 \ge \cdots \ge p_r)$ of 2n + 1 described in Proposition 2.1(2) which have the property that $p_i \le p_{i+1} + 1$ for all $i = 1, \ldots, r$ (setting $p_{r+1} = 0$) and which have no odd part of \underline{p} occurring exactly twice.
- (3) When $\Phi = C_n$, $n \ge 3$, the rigid nilpotent G-orbits correspond to those partitions $\underline{p} = (p_1 \ge p_2 \ge \cdots \ge p_r)$ of 2n described in Proposition 2.1(3) which have the property that $p_i \le p_{i+1} + 1$ for all $i = 1, \ldots, r$ (setting $p_{r+1} = 0$) and which have no even part of \underline{p} occurring exactly twice.
- (4) When $\Phi = D_n$, $n \ge 4$, the rigid nilpotent G-orbits correspond to those partitions $\underline{p} = (p_1 \ge p_2 \ge \cdots \ge p_r)$ of 2n described in Proposition 2.1(4) which have the property that $p_i \le p_{i+1}+1$ for all $i = 1, \ldots, r$ (setting $p_{r+1} = 0$) and which have no odd part of \underline{p} occurring exactly twice.

When Φ is of exceptional type, the rigid G-orbits are listed explicitly in [DE]. There are 3 non-zero rigid nilpotent G-orbits in type E_6 , 7 in type E_7 , 17 in type E_8 , 5 in type F_4 and 2 in type G_2 .

2.4. **Birational induction.** In this subsection we recall the notion of birational induction from [L2] and [LMM], which extends the idea of Lusztig-Spaltenstein induction to the setting of nilpotent covers.

Maintaining the notation from Subsection 2.3, let $\widetilde{\mathbb{O}}_L$ be an *L*-equivariant cover of \mathbb{O}_L . The covering map $\widetilde{\mathbb{O}}_L \to \mathbb{O}_L$ induces a map $\zeta : \operatorname{Spec}(\mathbb{C}[\widetilde{\mathbb{O}}_L]) \to \overline{\mathbb{O}}_L$. We hence get an action of *P* on $\operatorname{Spec}(\mathbb{C}[\widetilde{\mathbb{O}}_L]) \times \mathfrak{p}^{\perp}$ by letting *L* act diagonally and letting *U* act via

$$u \cdot (x, \chi) = (x, u \cdot \zeta(x) - \zeta(x) + u \cdot \chi)$$

for $u \in U$, $x \in \operatorname{Spec}(\mathbb{C}[\widetilde{\mathbb{O}}_L])$ and $\chi \in \mathfrak{p}^{\perp}$. This induces a map

$$\widetilde{\mu}: G \times^P (\operatorname{Spec}(\mathbb{C}[\widetilde{\mathbb{O}}_L]) \times \mathfrak{p}^{\perp}) \to G \times^P (\overline{\mathbb{O}}_L \times \mathfrak{p}^{\perp}) \xrightarrow{\mu} \mathfrak{g}^*.$$

Noting that $\mathbb{O} = \operatorname{Ind}_{L}^{G}(\mathbb{O}_{L})$ lies in the image of this map, we define

$$\widetilde{\mathbb{O}} = \widetilde{\mu}^{-1}(\mathbb{O}).$$

This is a G-equivariant nilpotent cover of \mathbb{O} . We have hence defined a map

$$\operatorname{Bind}_{L}^{G}:\operatorname{Cov}(L,\mathbb{O}_{L})\to\operatorname{Cov}(G,\operatorname{Ind}_{L}^{G}(\mathbb{O}_{L})),\quad \widetilde{\mathbb{O}}_{L}\mapsto\widetilde{\mu}^{-1}(\operatorname{Ind}_{L}^{G}(\mathbb{O}_{L}))=\widetilde{\mathbb{O}},$$

which we call **birational induction**. As with Lusztig-Spaltenstein induction, birational induction is independent of the parabolic subgroup P of G in which L is a Levi factor and is transitive by [LMM, Proposition 2.4.1]. We also have the following result from [LMM, Proposition 2.4.1] and [MMY, Proposition 2.6], which will be very useful in Section 3.

Proposition 2.3. Maintain the notation from above and suppose that $\widetilde{\mathbb{O}}_L \to \mathbb{O}_L$ is a degree m nilpotent cover of \mathbb{O}_L and that $\check{\mathbb{O}}_L \to \mathbb{O}_L$ is a degree mn nilpotent cover of \mathbb{O}_L which factors through $\check{\mathbb{O}}_L \to \widetilde{\mathbb{O}}_L$, for $m, n \in \mathbb{N}$. Then the degree of the covering map $\operatorname{Bind}_L^G(\widetilde{\mathbb{O}}_L) \to \operatorname{Ind}_L^G(\mathbb{O}_L)$ is divisible by m, and the degree of the covering map $\operatorname{Bind}_L^G(\check{\mathbb{O}}_L) \to \operatorname{Bind}_L^G(\check{\mathbb{O}}_L)$ is precisely n.

When $\widetilde{\mathbb{O}}$ is a *G*-equivariant nilpotent cover of a nilpotent *G*-orbit \mathbb{O} , we call $(L, \widetilde{\mathbb{O}}_L)$ a **birational** induction datum for $\widetilde{\mathbb{O}}$ if $\widetilde{\mathbb{O}} = \text{Bind}_L^G(\widetilde{\mathbb{O}}_L)$. We say that $\widetilde{\mathbb{O}}$ is **birationally rigid** if the only birational induction datum for $\widetilde{\mathbb{O}}$ is $(G, \widetilde{\mathbb{O}})$, i.e. if it cannot be birationally induced from a proper Levi subgroup. If $\widetilde{\mathbb{O}}_L$ is birationally rigid and $(L, \widetilde{\mathbb{O}}_L)$ is a birational induction datum for $\widetilde{\mathbb{O}}$, we call $(L, \widetilde{\mathbb{O}}_L)$ a **birationally rigid induction datum** for $\widetilde{\mathbb{O}}$.²

Analogous to the notation for rigid induction data, let us denote the set of birationally rigid induction data for $\widetilde{\mathbb{O}}$ by

$$\operatorname{brig}(\widetilde{\mathbb{O}}) := \{ (L, \widetilde{\mathbb{O}}_L) \mid \operatorname{Bind}_L^G(\widetilde{\mathbb{O}}_L) = \widetilde{\mathbb{O}} \}.$$

The adjoint action of G on G induces an action of G on $brig(\mathbb{O})$. We denote

$$\operatorname{Brig}(\mathbb{O}) := \operatorname{brig}(\mathbb{O})/G$$

and we will abuse notation slightly to also refer to elements of $\operatorname{Brig}(\widetilde{\mathbb{O}})$ as birationally rigid induction data for $\widetilde{\mathbb{O}}$.

The following result about birational induction is due to [L2, Theorem 4.4] and [LMM, Proposition 2.4.1], and forms the basis for the question presented in this paper.

Proposition 2.4. Let $\widetilde{\mathbb{O}}$ be a *G*-equivariant nilpotent cover of a nilpotent *G*-orbit \mathbb{O} . Then $\operatorname{Brig}(\widetilde{\mathbb{O}})$ consists of a unique birationally rigid induction datum for $\widetilde{\mathbb{O}}$.

A nilpotent *G*-orbit \mathbb{O} is a *G*-equivariant nilpotent cover of itself; this corresponds to the subgroup $\pi_1^G(\mathbb{O})$ under the bijection (2.1). Note that $\operatorname{Bind}_L^G(\mathbb{O}_L) = \mathbb{O}$ if and only if the map μ from (2.2) is birational – indeed, this was the notion of birational induction initially studied in [L2]. With this interpretation of birational induction, we can easily describe the birationally rigid induction datum for even \mathbb{O} . When \mathbb{O} is even we may form the Jacobson-Morozov Levi subgroup $L_{\mathbb{O}}$ of *G*; this is the standard Levi subgroup corresponding to the simple roots labelled by 0 in the weighted Dynkin diagram of \mathbb{O} . The following result is then standard and can be found in [LMM, §2.4] and references therein.

Proposition 2.5. Let \mathbb{O} be an even nilpotent *G*-orbit in \mathfrak{g}^* . Then \mathbb{O} is birationally induced from the zero orbit for the Jacobson-Morozov Levi subgroup $L_{\mathbb{O}}$ corresponding to \mathbb{O} .

²We should technically call this a birationally rigid birational induction datum, but for ease of reference we miss out the second "birational". In [L2] these are referred to as birationally minimal induction data, presumably to avoid confusion with an induction datum (L, \mathbb{O}_L) for \mathbb{O} in which \mathbb{O}_L is birationally rigid but $\mathbb{O} = \text{Ind}_L^G(\mathbb{O}_L) \neq \text{Bind}_L^G(\mathbb{O}_L)$. Since we shall never use the latter concept, we think this abuse of notation is forgivable.

Remark 1. Although each nilpotent G-orbit can also be viewed as a G-equivariant nilpotent cover of itself, it is worth highlighting here a distinction between the theory of nilpotent orbits and the theory of nilpotent orbit covers. When studying nilpotent orbits of an algebraic group G, we generally only care about the underlying root system Φ of G – in particular, for most questions we care about it doesn't usually matter whether G is reductive or semisimple, or what the isogeny type of G is. For example, this is true for most questions about Lusztig-Spaltenstein induction and so we don't normally need to concern ourselves with the precise form of Levi subgroups.

On the other hand, in the theory of nilpotent orbit covers the precise form of the algebraic group does matter. This is because G-equivariant nilpotent orbit covers of \mathbb{O} (up to isomorphism) are indexed by conjugacy classes of subgroups of $\pi_1^G(\mathbb{O}) = G_e/G_e^\circ$, and this latter group cannot be described purely in terms of the underlying root system of G. For example, when $G = \operatorname{PGL}_n$ for some $n \in \mathbb{N}$ we get that $\pi_1^G(\mathbb{O}) = 1$ for all nilpotent G-orbits \mathbb{O} , while for $G = \operatorname{SL}_n$ we get that $\pi_1^G(\mathbb{O}) = \mathbb{Z}/h\mathbb{Z}$ where h is the greatest common divisor of all the parts of the partition of n corresponding to \mathbb{O} .

When G is semisimple and the root system of G is indecomposable of exceptional type, we note that G is uniquely defined for types E_8 , F_4 and G_2 . On the other hand, for types E_6 and E_7 there are two isogeny classes: simply connected and adjoint. Given a nilpotent G-orbit in such cases, the G-equivariant fundamental group may include an extra $\mathbb{Z}/3\mathbb{Z}$ factor (in the E_6 case) or an extra $\mathbb{Z}/2\mathbb{Z}$ -factor (in the E_7 case) when G is simply connected versus when G is adjoint. In this paper, we focus exclusively on the simply connected exceptional groups; in such cases, the G-equivariant fundamental groups can be found in [CM, §8.4].³

Let L be a Levi subgroup of G and \mathbb{O}_L a nilpotent L-orbit. In Section 3, we often need to compute $\pi_1^L(\mathbb{O}_L)$ in order to determine the nilpotent covers of \mathbb{O}_L . We cannot do this by appealing to any list, instead we rely on the atlas software [At]. More details are provided at the beginning of Section 3.

2.5. Namikawa space and the Namikawa Weyl group. In this subsection we recall the basics on the Namikawa space and Namikawa Weyl group that are relevant for this paper, as can be found in [L2, LMM, MM, MMY, N1].

When \mathbb{O} is a *G*-equivariant nilpotent orbit cover of a nilpotent *G*-orbit \mathbb{O} , the affine variety $X := \operatorname{Spec}(\mathbb{C}[\widetilde{\mathbb{O}}])$ is a conical symplectic singularity by [L3, Lemma 2.5]. Let $X_1 := X \setminus X^{\operatorname{reg}}$, where X^{reg} is the regular locus of X, and let $\mathcal{L}_1, \ldots, \mathcal{L}_k$ be the irreducible components of X_1 ; we call these the **symplectic leaves of codimension 2** in X. Following [L2] and [N1] (see also [LMM, §4.5]), we may assign to each \mathcal{L}_i a simple complex Lie algebra \mathfrak{g}_k of type A, D or E. For such Lie algebra \mathfrak{g}_i , we fix a Cartan subalgebra \mathfrak{h}_i and denote by Φ_i the corresponding root system and by W_i the corresponding Weyl group. The fundamental group $\pi_1(\mathcal{L}_i)$ acts by monodromy on Φ_i and thus on \mathfrak{h}_i^* and on W_i . We define the **partial Namikawa space** for \mathcal{L}_i by

$$\mathfrak{P}_i := (\mathfrak{h}_i^*)^{\pi_1(\mathcal{L}_i)}$$

and define $\mathfrak{P}_0 := H^2(X^{\mathrm{reg}}, \mathbb{C})$. By [L2, Lemma 2.8] the Namikawa space of X is then

$$\mathfrak{P}(\widetilde{\mathbb{O}})\coloneqq\mathfrak{P}_0\oplusigoplus_{i=1}^k\mathfrak{P}_i.$$

Furthermore, the **Namikawa Weyl group** of X is defined by

$$W(\widetilde{\mathbb{O}}) := W_1^{\pi_1(\mathcal{L}_1)} \times \cdots \times W_k^{\pi_1(\mathcal{L}_k)},$$

which acts component-wise on $\mathfrak{P}(\widetilde{\mathbb{O}})$ (with trivial action on \mathfrak{P}_0).

³For the *G*-equivariant fundamental groups for the classical cases when *G* is simply connected or adjoint, the reader can consult [CM, Corollary 6.1.6].

The following result, allowing us to describe the Namikawa space of X using birational induction, is [LMM, Proposition 7.2.2(i)].

Proposition 2.6. Let $\widetilde{\mathbb{O}} \in \text{Cov}(G, \mathbb{O})$ and let $(L, \widetilde{\mathbb{O}}_L)$ be the birationally rigid induction datum for $\widetilde{\mathbb{O}}$. Then

$$\mathfrak{P}(\mathbb{O}) \cong \mathfrak{z}(\mathfrak{l})^*.$$

We may similarly describe the Namikawa Weyl group via birational induction. Since we will only need this in the case when $\widetilde{\mathbb{O}}$ is birationally induced from a nilpotent orbit \mathbb{O}_L , we only give the result in this case; for the more general case, see [L4], [LMM, §7], [MMY, §3.10].

We define

 $N_G(L, \mathbb{O}_L) = \{ g \in G \mid g \cdot L = L \text{ and } g \cdot \mathbb{O}_L = \mathbb{O}_L \}$

and define the **extended Namikawa Weyl group** by

$$\widetilde{W}(\widetilde{\mathbb{O}}) = N_G(L, \mathbb{O}_L)/L.$$

The following result is a special case of [MMY, Theorem 3.34] (see also [L4]), but is all we need in this paper.

Proposition 2.7. Let $\widetilde{\mathbb{O}} \to \mathbb{O}$ be a *G*-equivariant nilpotent cover of a nilpotent *G*-orbit \mathbb{O} , and suppose that $\operatorname{Brig}(\widetilde{\mathbb{O}}) = \{(L, \mathbb{O}_L)\}$ for some Levi subgroup *L* of *G* and rigid nilpotent *L*-orbit \mathbb{O}_L . Then $W(\widetilde{\mathbb{O}})$ is a normal subgroup of $N_G(L, \mathbb{O}_L)/L$. Furthermore, if $\widetilde{\mathbb{O}} = \mathbb{O}$ then $W(\widetilde{\mathbb{O}}) = W(\mathbb{O}) = N_G(L, \mathbb{O}_L)/L$.

2.6. Birationally rigid nilpotent covers. In this subsection, we give some results on birational induction and birationally rigid orbit covers. The vast majority of this material can be found in [LMM, MM, MMY].

In describing the birationally rigid nilpotent G-orbits (that is to say, nilpotent G-orbits which are birationally rigid when trivially viewed as nilpotent covers), the following result allows us to reduce to semisimple algebraic groups of a given isogeny type.

Theorem 2.8. Let G_1 and G_2 be reductive algebraic groups with the same root system Φ (associated to fixed choices of maximal tori). Let \mathfrak{g}_1 and \mathfrak{g}_2 be the Lie algebras of G_1 and G_2 respectively. Let \mathbb{O}_1 and \mathbb{O}_2 be corresponding nilpotent orbits in \mathfrak{g}_1^* and \mathfrak{g}_2^* . Then \mathbb{O}_1 is birationally rigid if and only if \mathbb{O}_2 is birationally rigid.

Proof. For i = 1, 2, let (L_i, \mathbb{O}_{L_i}) be the birationally rigid induction datum for \mathbb{O}_i and write $\mathfrak{l}_i = \text{Lie}(L_i)$. By Proposition 2.6, the Namikawa space $\mathfrak{P}(\mathbb{O}_i)$ for $\text{Spec}(\mathbb{C}[\mathbb{O}_i])$ is isomorphic to $\mathfrak{z}(\mathfrak{l}_i)^*$. Thus, \mathbb{O}_i is birationally rigid if and only if $\mathfrak{P}(\mathbb{O}_i) = 0$. On the other hand,

$$\mathfrak{P}(\mathbb{O}_i) = H^2(\operatorname{Spec}(\mathbb{C}[\mathbb{O}_i])^{\operatorname{reg}}, \mathbb{C}) \oplus (\mathfrak{h}_1^{i,*})^{\pi_1(\mathcal{L}_1^i)} \oplus \dots \oplus (\mathfrak{h}_{k_i}^{i,*})^{\pi_1(\mathcal{L}_{k_i}^i)}$$

where $\mathcal{L}_1^i, \ldots, \mathcal{L}_{k_i}^i$ are the symplectic leaves of codimension 2 in $\operatorname{Spec}(\mathbb{C}[\mathbb{O}_i])$ and the $\mathfrak{h}_j^{i,*}$ corresponding to these leaves are as defined at the beginning of Subsection 2.5. Hence $\mathfrak{P}(\mathbb{O}_i) = 0$ if and only if $H^2(\operatorname{Spec}(\mathbb{C}[\mathbb{O}_i])^{\operatorname{reg}}, \mathbb{C}) = 0$ and $k_i = 0$. These two properties are geometric in nature; in particular, since \mathbb{O}_1 and \mathbb{O}_2 are isomorphic as varieties, they hold for \mathbb{O}_1 if and only if they hold for \mathbb{O}_2 . Hence, \mathbb{O}_1 is birationally rigid if and only if \mathbb{O}_2 is.

As discussed in the prior proof, there is a geometric criterion for a nilpotent orbit cover to be birationally rigid due to [LMM, Corollary 7.6.1].

Proposition 2.9. Let G be a semisimple simply connected algebraic group, and let \mathbb{O} be a Gequivariant nilpotent cover of a nilpotent G-orbit \mathbb{O} in $\mathfrak{g} = \operatorname{Lie}(G)$. Then \mathbb{O} is birationally rigid if and only if $H^2(\mathbb{O}, \mathbb{C}) = 0$ and $\operatorname{Spec}(\mathbb{C}[\mathbb{O}])$ has no symplectic leaves of codimension 2. Using this result, [LMM, Proposition 7.6.3] and [MM, Proposition 3.8.3] give the following classification of birationally rigid G-orbits.⁴ By Theorem 2.8, this classification in exceptional types can in fact already be found in [Fu]. Note that rigid orbits are automatically birationally rigid.

Proposition 2.10. The birationally rigid nilpotent G-orbits can be indexed as follows:

- (1) When G = SL(n), $n \ge 1$, the only birationally rigid nilpotent G-orbit is the zero orbit.
- (2) When G = SO(2n+1), $n \ge 2$, the birationally rigid nilpotent G-orbits correspond to those partitions $\underline{p} = (p_1 \ge p_2 \ge \cdots \ge p_r)$ of 2n + 1 described in Proposition 2.1(2) with the property that $p_i \le p_{i+1} + 1$ for all $i = 1, \ldots, r$ (setting $p_{r+1} = 0$).
- (3) When G = Sp(2n), $n \ge 2$, the birationally rigid nilpotent G-orbits correspond to those partitions $\underline{p} = (p_1 \ge p_2 \ge \cdots \ge p_r)$ of 2n described in Proposition 2.1(3) with the property that $p_i \le p_{i+1} + 1$ for all $i = 1, \ldots, r$ (setting $p_{r+1} = 0$).
- (4) When G = SO(2n), $n \ge 4$, the birationally rigid nilpotent G-orbits correspond to those partitions $\underline{p} = (p_1 \ge p_2 \ge \cdots \ge p_r)$ of 2n described in Proposition 2.1(4) with the property that $p_i \le p_{i+1} + 1$ for all $i = 1, \ldots, r$, excluding those partitions of the form $(2^m, 1^2)$ for $m \in \mathbb{N}$.
- (5) When G is semisimple simply connected of type E_6 , F_4 or G_2 , the birationally rigid nilpotent G-orbits are precisely the rigid nilpotent G-orbits.
- (6) When G is semisimple simply connected of type E_7 , the birationally rigid nilpotents G-orbits are the rigid nilpotent G-orbits together with the nilpotent G-orbits with Bala-Carter labels $A_2 + A_1$ and $A_4 + A_1$.
- (7) When G is semisimple simply connected of type E_8 , the birationally rigid nilpotent G-orbits are the rigid nilpotent G-orbits together with the nilpotent G-orbits with Bala-Carter label $A_4 + A_1$ and $A_4 + 2A_1$.

Note that a birationally rigid nilpotent orbit may admit a birationally induced nilpotent cover (although rigid nilpotent orbits can only admit birationally rigid nilpotent covers) and, conversely, a birationally induced nilpotent orbit may admit a birationally rigid nilpotent cover. For the exceptional groups, the classification of birationally induced nilpotent orbits which admit birationally rigid nilpotent covers is given in [MM, Proposition 3.9.5], as follows:

Proposition 2.11. When G is semisimple simply connected of exceptional type, the birationally induced nilpotent G-orbits with birationally rigid nilpotent covers can be classified as follows:

- (1) When G is of type E_6 , such nilpotent G-orbits are the ones with Bala-Carter labels A_2 , $D_4(a_1)$, $2A_2$, A_5 and $E_6(a_3)$.
- (2) When G is of type E_7 , such nilpotent G-orbits are the ones with Bala-Carter labels $(3A_1)''$, A_2 , $A_2 + 3A_1$, $(A_3 + A_1)''$, $D_4(a_1)$, $A_3 + 2A_1$, $D_4(a_1) + A_1$, $A_3 + A_2 + A_1$, $A_5 + A_1$, $D_5(a_1) + A_1$, $E_7(a_5)$ and $E_7(a_4)$.
- (3) When G is of type E_8 , such nilpotent G-orbits are the ones with Bala-Carter labels A_2 , $2A_2$, $D_4(a_1)$, $D_4(a_1) + A_2$, $D_4 + A_2$, $D_6(a_2)$, $E_6(a_3) + A_1$, $E_7(a_5)$, $E_8(a_7)$ and $E_8(b_6)$.
- (4) When G is of type F_4 , such nilpotent G-orbits are the ones with Bala-Carter labels A_2 , B_2 , $C_3(a_1)$ and $F_4(a_3)$.
- (5) When G is of type G_2 , the only such nilpotent G-orbit has Bala-Carter label $G_2(a_1)$.

The proofs of [LMM, Proposition 7.6.16] and [MM, Proposition 3.9.5] show that in all these cases – except possibly $D_4(a_1) \subseteq E_6$ – the universal cover (i.e. the nilpotent cover corresponding to

⁴The result in [MM] actually misses out $A_4 + 2A_1 \subseteq E_8$, but the argument in that paper shows that this orbit is birationally rigid. Indeed, this can also be seen in [Fu].

 $1 \subseteq \pi_1^G(\mathbb{O})$) is birationally rigid.⁵ We see in Subsection 3.7.1 that the universal cover of $D_4(a_1) \subseteq E_6$ turns out not to be birationally rigid; in fact, the nilpotent cover of $D_4(a_1)$ corresponding to the subgroup $A_3 \subseteq S_3$ is the only birationally rigid one.

That the universal cover of an nilpotent orbit is birationally rigid generally neither implies nor is implied by the birational rigidity of any other nilpotent cover. However, there will be situations in which the birational rigidity of one nilpotent cover does tell us something about the birational rigidity of another. We say that a *G*-equivariant nilpotent cover $\widetilde{\mathbb{O}}$ is **2-leafless** if $\text{Spec}(\mathbb{C}[\widetilde{\mathbb{O}}])$ has no codimension 2 symplectic leaves. The following proposition shows how 2-leafless nilpotent covers interact with each other.

Proposition 2.12. Let \mathbb{O} be a nilpotent *G*-orbit in \mathfrak{g}^* . Then any *G*-equivariant nilpotent cover of \mathbb{O} which lies over a 2-leafless *G*-equivariant nilpotent cover of \mathbb{O} is also 2-leafless.

Proof. This Lemma 7.6.6 in [LMM].

The following straightforward result shows us how we can extend this property, in certain circumstances, to birational rigidity.

Corollary 2.13. Suppose G is semisimple and simply connected. Let \mathbb{O} be a nilpotent G-orbit in \mathfrak{g}^* and let \mathfrak{r} be the Lie algebra of the reductive part of the centraliser of any $\chi \in \mathbb{O}$. Suppose that \mathfrak{r} is semisimple. Then any G-equivariant nilpotent cover of \mathbb{O} which lies over a birationally rigid G-equivariant nilpotent cover of \mathbb{O} is also birationally rigid.

Proof. By [LMM, Corollary 7.6.1], a nilpotent cover $\widetilde{\mathbb{O}}$ is birationally rigid if and only if it is 2leafless and $H^2(\widetilde{\mathbb{O}}, \mathbb{C}) = 0$. Let $x \in \widetilde{\mathbb{O}}$, let R_x be the reductive part of the centraliser G_x of x in G, and let $\mathfrak{r}_x = \operatorname{Lie}(R_x)$. By [LMM, Lemma 7.2.7], $H^2(\widetilde{\mathbb{O}}, \mathbb{C}) = \mathfrak{X}(\mathfrak{r}_x)^{\pi_1^G(\widetilde{\mathbb{O}})}$. Note that $\mathfrak{r} = \mathfrak{r}_x$; therefore, \mathfrak{r} being semisimple implies $H^2(\widetilde{\mathbb{O}}, \mathbb{C}) = 0$. Since this applies for all nilpotent covers of \mathbb{O} , we conclude that being birationally rigid is equivalent to being 2-leafless for G-equivariant nilpotent covers of such \mathbb{O} . The result then follows from Proposition 2.12.

Since we use it a couple of times later, we emphasise the point made in this proof that

(2.3)
$$\mathfrak{r}$$
 semisimple $\Longrightarrow H^2(\mathbb{O}, \mathbb{C}) = 0$ for all nilpotent covers \mathbb{O} of \mathbb{O} .

Another important property of birational induction is discussed in [LMM, Lemma 2.5.1]. Suppose that $\widetilde{\mathbb{O}} \to \mathbb{O}$ is a *G*-equivariant nilpotent cover which is birationally induced from an *L*-equivariant nilpotent cover $\widetilde{\mathbb{O}}_L \to \mathbb{O}_L$ for some Levi subgroup *L* of *G*. Pick $x \in \widetilde{\mathbb{O}}$. Then the inclusion $\widetilde{\mathbb{O}} \hookrightarrow G \times^P (\widetilde{\mathbb{O}}_L \times \mathfrak{p}^{\perp})$ induces a surjective homomorphism

$$\pi_1(\widetilde{\mathbb{O}}) \twoheadrightarrow \pi_1(G \times^P (\widetilde{\mathbb{O}}_L \times \mathfrak{p}^{\perp})),$$

where we take x as the basepoint for both fundamental groups. Choose $x \in \widetilde{\mathbb{O}}$ and $y \in \widetilde{\mathbb{O}}_L$ such that x maps to the point (1, y) under the map $\widetilde{\mathbb{O}} \hookrightarrow G \times^P (\widetilde{\mathbb{O}}_L \times \mathfrak{p}^{\perp}) \twoheadrightarrow G \times^P \widetilde{\mathbb{O}}_L$. As in [LMM, §2.5], when we use $1 \in G$, $x \in \widetilde{\mathbb{O}}$ and $y \in \widetilde{\mathbb{O}}_L$ as the appropriate basepoints, the group homomorphism fits into the diagram

$$(2.4) \qquad \begin{array}{c} \pi_1(G) & \longrightarrow & \pi_1(\widetilde{\mathbb{O}}) & \longrightarrow & \pi_1^G(\widetilde{\mathbb{O}}) & \longrightarrow & 1 \\ \\ \parallel & & \downarrow & & \downarrow \\ & & & & \\ & & & \\ &$$

⁵The proof of [MM, Proposition 3.9.5] doesn't explicitly contain this statement for $A_3 + A_2 + A_1 \subseteq E_8$, but it follows immediately from the fact that the orbit itself is not birationally rigid and the universal cover is its only nontrivial cover.

where the rows are exact sequences of homotopy groups obtained from appropriate fibrations. The following result is [LMM, Lemma 2.5.1].

Proposition 2.14. Let \mathbb{O} be a nilpotent *G*-orbit in \mathfrak{g}^* and let $\widetilde{\mathbb{O}}$ be a *G*-equivariant nilpotent cover of \mathbb{O} . Suppose $(L, \widetilde{\mathbb{O}}_L)$ is a birational induction datum for $\widetilde{\mathbb{O}}$. Then there exists a surjective group homomorphism $\pi_1^G(\widetilde{\mathbb{O}}) \twoheadrightarrow \pi_1^L(\widetilde{\mathbb{O}}_L)$ such that Diagram (2.4) commutes.

3. CASE-BY-CASE ARGUMENTS

In this section G is a semisimple simply connected algebraic group with indecomposable root system of exceptional type. For each nilpotent G-orbit \mathbb{O} , we determine the birationally rigid induction datum for each nilpotent cover of \mathbb{O} . We proceed based on the isomorphism type of $\pi_1^G(\mathbb{O})$, with the conventions of Remark 1; with such conventions, there are 10 groups which can appear. These are 1, $\mathbb{Z}/2\mathbb{Z}$, S_2 , $\mathbb{Z}/3\mathbb{Z}$, $S_2 \times \mathbb{Z}/2\mathbb{Z}$, S_3 , $S_2 \times \mathbb{Z}/3\mathbb{Z}$, $S_3 \times \mathbb{Z}/2\mathbb{Z}$, S_4 and S_5 – as in Remark 1, we treat $\mathbb{Z}/2\mathbb{Z}$ and S_2 separately for consistency with [CM, §8.4] and to reflect whether or not such component exists when G is of adjoint type. We maintain this notation when determining $\pi_1^L(\mathbb{O}_L)$ for an induction datum (L, \mathbb{O}_L) , in that we write $\mathbb{Z}/2\mathbb{Z}$ for components which arise in $\pi_1^L(\mathbb{O}_L)$ only when G is simply connected and write S_2 for components which arise in $\pi_1^L(\mathbb{O}_L)$ in both the simply connected and adjoint cases (since G is exceptional, these are the only possible isogeny types).

In order to determine the *L*-equivariant fundamental group $\pi_1^L(\mathbb{O}_L)$ of an induction datum (L, \mathbb{O}_L) for \mathbb{O} , there are a couple of techniques we can use. One powerful tool is the atlas software [At]; we use the command Levi_datum to isolate the Levi subgroup, and the command component_datum(orbit).orders to get the orders of elements of $\pi_1^L(\mathbb{O}_L)$. More theoretically, we note that there are surjections

(3.1)
$$\pi_1^{[L,L]_{\rm sc}}(\mathbb{O}_L) \twoheadrightarrow \pi_1^L(\mathbb{O}_L) \twoheadrightarrow \pi_1^{[L,L]_{\rm ad}}(\mathbb{O}_L),$$

where $[L, L]_{sc}$ (resp. $([L, L]_{ad})$ denotes the simply connected (resp. adjoint) form of the derived subgroup [L, L] of L (see [CM, §1.2, §6.1] for more details). When $\pi_1^{[L,L]_{sc}}(\mathbb{O}_L) = \pi_1^{[L,L]_{ad}}(\mathbb{O}_L)$ this clearly allows us to determine the L-equivariant fundamental group $\pi_1^L(\mathbb{O}_L)$.

We make some further notational conventions. By conjugation, we may assume that all the Levi subgroups we consider are standard Levi subgroups. As discussed in Subsection 2.1, these may be described by Dynkin type (uniquely in most cases, although in some cases we will need additional decoration as discussed in that subsection); we therefore often refer to Levi subgroups via their Dynkin types. For example, we may write $\operatorname{Ind}_{D_4}^{E_8}$ to mean induction from the (unique to to conjugacy) standard Levi subgroup of G with Dynkin type D_4 to the standard Levi subgroup of G with Dynkin type E_8 . Based on these conventions, we often use the phrase "equivariant fundamental group" in this section rather than "L-equivariant fundamental group" when L is clear, in order to avoid undue clutter.

This section is structured as follows. We have one subsection for each group which can appear as $\pi_1^G(\mathbb{O})$. At the beginning of each of these sections we discuss how many nilpotent *G*-orbits in each exceptional type have such *G*-equivariant fundamental group, and we resolve as many cases as we can using class-wide arguments. For those nilpotent *G*-orbits which can't easily be dealt with using general arguments, we instead make orbit-by-orbit arguments in following (sub-)subsections. The results of this section are compiled into tables in Section 4.

We conclude the introduction to this section by noting that throughout the following subsections we prefer to cite to results as labelled in Section 2 rather than to their original sources, for ease of reference for the reader. Nonetheless, we emphasise that the vast majority of these results are due to other authors, and refer the reader back to Section 2 for the proper citations. 3.1. $\pi_1^G(\mathbb{O}) = 1$. There are 45 induced nilpotent orbits in exceptional cases with $\pi_1^G(\mathbb{O}) = 1$; that is to say, there is 1 such orbit for G_2 , 4 such orbits for F_4 , 10 such orbits for E_6 , 8 such orbits for E_7 and 22 such orbits for E_8 . That $\pi_1^G(\mathbb{O}) = 1$ means that \mathbb{O} has no non-trivial nilpotent covers. Since any rigid induction datum for \mathbb{O} must birationally induce to a nilpotent cover of \mathbb{O} , we must get that each \mathbb{O} with $\pi_1^G(\mathbb{O}) = 1$ has a unique rigid induction datum (we can also check this directly from [DE]) and that \mathbb{O} must be birationally induced from this rigid (and thus birationally rigid) induction datum.

3.2. $\pi_1^G(\mathbb{O}) = \mathbb{Z}/2\mathbb{Z}$. This case can only arise in type E_7 ; in this case, there are 16 induced nilpotent orbits with $\pi_1^G(\mathbb{O}) = \mathbb{Z}/2\mathbb{Z}$. That $\pi_1^G(\mathbb{O}) = \mathbb{Z}/2\mathbb{Z}$ implies that the only non-trivial nilpotent cover of \mathbb{O} is the universal cover $\widehat{\mathbb{O}}$. If the universal cover is birationally rigid, then arguing as in Subsection 3.1 shows that \mathbb{O} has a unique rigid induction datum and that \mathbb{O} is birationally induced from this rigid induction datum. This handles 7 nilpotent orbits (using Proposition 2.11 to see which induced nilpotent orbits for simply connected E_7 have birationally rigid nilpotent covers).

We tackle the remaining 9 nilpotent orbits individually.

3.2.1. $D_4 + A_1 \subseteq E_7$. The nilpotent orbit in E_7 with Bala-Carter label $D_4 + A_1$ has weighted Dynkin diagram

$$\begin{array}{c}1\\2&1&0&0&0&1\end{array}$$

and unique rigid induction datum

Note that this rigid induction datum has equivariant fundamental group $\mathbb{Z}/2\mathbb{Z}$; the universal cover of this nilpotent orbit must therefore also be birationally rigid. We thus have

$$\mathbb{O} = \operatorname{Bind}_{D_6}^{E_7} \left(\begin{array}{ccc} 1 & & \\ 1 & 0 & 0 & 0 & 1 \end{array} \right)$$

and

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{D_6}^{E_7} \left(\operatorname{univ. cover of} \begin{array}{ccc} 1 & & \\ 1 & 0 & 0 & 0 \end{array} \right).$$

3.2.2. $(A_5)'' \subseteq E_7$. The nilpotent orbit in E_7 with Bala-Carter label $(A_5)''$ has weighted Dynkin diagram

$$\begin{smallmatrix}&&0\\2&0&0&0&2&2\end{smallmatrix}$$

and unique rigid induction datum

$$\begin{array}{ccc} 0 \\ 0 & 0 \end{array} \subseteq D_4.$$

Since \mathbb{O} is even, we know that

$$\mathbb{O} = \operatorname{Bind}_{D_4}^{E_7} \left(\begin{array}{cc} 0 \\ 0 & 0 \end{array} \right).$$

To determine the birationally rigid induction datum for $\widehat{\mathbb{O}}$, it is enough to find the induction datum for \mathbb{O} with maximal semisimple corank such that the equivariant fundamental group is nontrivial. Arguing in this way, we get

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{D_5+A_1}^{E_7} \left(\operatorname{univ. cover of} \begin{array}{c} 0 \\ 2 \end{array} \right)$$
13

3.2.3. $D_6(a_2) \subseteq E_7$. The nilpotent orbit in E_7 with Bala-Carter label $D_6(a_2)$ has weighted Dynkin diagram

$$\begin{smallmatrix}&&1\\0&1&0&1&0&2\end{smallmatrix}$$

and unique rigid induction datum

$$\begin{array}{ccc} & 0 \\ 1 & 0 & 1 & 0 \end{array} \subseteq D_5.$$

Induction from this rigid induction datum passes through the induction datum

which is the induction datum of maximal semisimple corank such that its equivariant fundamental group is non-trivial. By Theorem 2.8 and Proposition 2.10, this nilpotent orbit is birationally induced; its universal cover must therefore be birationally rigid. We hence have

$$\mathbb{O} = \operatorname{Bind}_{D_5}^{E_7} \left(\begin{array}{cc} 0 \\ 1 & 0 & 1 & 0 \end{array} \right)$$

and

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{D_5+A_1}^{E_7} \left(\operatorname{univ. cover of} \begin{array}{ccc} 0 & & \\ 1 & 0 & 1 & 0 & 2 \end{array} \right).$$

3.2.4. $D_5 + A_1 \subseteq E_7$. The nilpotent orbit in E_7 with Bala-Carter label $D_5 + A_1$ has weighted Dynkin diagram

and unique rigid induction datum

$$\begin{array}{ccc} 0 \\ 0 \\ 0 \end{array} \quad \begin{array}{c} 0 \\ 0 \end{array} \subseteq 2A_2 \end{array}$$

Induction from this rigid induction datum passes through the induction datum

$$\begin{array}{cccc} 0 \\ 0 \\ 2 \\ 0 \\ 0 \end{array} \subseteq (A_5)'',$$

which is the induction datum of maximal semisimple corank such that its equivariant fundamental group is non-trivial. By Theorem 2.8 and Proposition 2.10, this nilpotent orbit is birationally induced; its universal cover must therefore be birationally rigid. We hence have

$$\mathbb{O} = \operatorname{Bind}_{2A_2}^{E_7} \left(\begin{array}{cc} 0 & & \\ 0 & & 0 \end{array} \right)$$

and

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{(A_5)''}^{E_7} \left(\operatorname{univ. cover of} \begin{array}{ccc} 0 & & \\ 0 & 2 & 0 & 0 \end{array} \right)$$

3.2.5. $D_6(a_1) \subseteq E_7$. The nilpotent orbit in E_7 with Bala-Carter label $D_6(a_1)$ has weighted Dynkin diagram

$$\begin{smallmatrix}&&1\\2&1&0&1&0&2\end{smallmatrix}$$

and unique rigid induction datum

$$0 \quad 0 \quad 0 \subseteq A_3$$

Induction from this rigid induction datum passes through the induction datum

$$\begin{array}{cccc} 0 \\ 2 & 0 & 0 \\ & & 2 \end{array} \subseteq D_4 + A_1,$$

which is the induction datum of maximal semisimple corank such that its equivariant fundamental group is non-trivial. By Theorem 2.8 and Proposition 2.10, this nilpotent orbit is birationally induced; its universal cover must therefore be birationally rigid. We hence have

$$\mathbb{O} = \operatorname{Bind}_{A_3}^{E_7} \left(\begin{array}{ccc} 0 & 0 \end{array} \right)$$

and

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{D_4+A_1}^{E_7} \left(\operatorname{univ. cover of} \begin{array}{cc} 0 \\ 2 \end{array} \right)$$

3.2.6. $D_6 \subseteq E_7$. The nilpotent orbit in E_7 with Bala-Carter label D_6 has weighted Dynkin diagram

and unique rigid induction datum

$$\begin{array}{ccc} 1 \\ 1 & 0 & 1 \end{array} \subseteq D_4.$$

Induction from this rigid induction datum passes through the induction datum

which is the induction datum of maximal semisimple corank such that its equivariant fundamental group is non-trivial. By Theorem 2.8 and Proposition 2.10, this nilpotent orbit is birationally induced; its universal cover must therefore be birationally rigid. We hence have

$$\mathbb{O} = \operatorname{Bind}_{D_4}^{E_7} \left(\begin{array}{cc} 1 \\ 1 & 0 & 1 \end{array} \right)$$

and

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{D_4+A_1}^{E_7} \left(\operatorname{univ. cover of} \begin{array}{cc} 1 \\ 1 \end{array} \right).$$

3.2.7. $E_7(a_2) \subseteq E_7$. The nilpotent orbit in E_7 with Bala-Carter label $E_7(a_2)$ has weighted Dynkin diagram

and unique rigid induction datum

$$0 \qquad 0 \subseteq 2A_1.$$

Since \mathbb{O} is even, we know that

$$\mathbb{O} = \operatorname{Bind}_{2A_1}^{E_7} \left(\begin{array}{cc} 0 & 0 \end{array} \right).$$

To determine the birationally rigid induction datum for $\widehat{\mathbb{O}}$, we can find the induction datum for \mathbb{O} with maximal semisimple corank such that the equivariant fundamental group is nontrivial. From this, we get

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{(A_3+A_1)''}^{E_7} \left(\operatorname{univ. cover of} \begin{array}{cc} 2 \\ & 0 \end{array} \right)$$
15

3.2.8. $E_7(a_1) \subseteq E_7$. The nilpotent orbit in E_7 with Bala-Carter label $E_7(a_1)$ has weighted Dynkin diagram

and unique rigid induction datum

 $0 \subseteq A_1.$

Since \mathbb{O} is even, we know that

$$\mathbb{O} = \operatorname{Bind}_{A_1}^{E_7} \left(\begin{array}{c} 0 \end{array} \right).$$

To determine the birationally rigid induction datum for \mathbb{O} , we can find the induction datum for \mathbb{O} with maximal semisimple corank such that the equivariant fundamental group is nontrivial. From this, we get

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{4A_1}^{E_7} \left(\operatorname{univ. cover of} \begin{array}{cc} 2 & & \\ 0 & 2 & 2 \end{array} \right).$$

3.2.9. $E_7 \subseteq E_7$. The nilpotent orbit in E_7 with Bala-Carter label E_7 has weighted Dynkin diagram

$$\begin{smallmatrix}&&2\\2&2&2&2&2&2\\\end{smallmatrix}$$

and unique rigid induction datum consisting of the zero orbit in the Lie algebra of the torus T. Since \mathbb{O} is even, we know that

$$\mathbb{O} = \operatorname{Bind}_{T}^{E_{7}}(\{0\}).$$

To determine the birationally rigid induction datum for $\widehat{\mathbb{O}}$, we can find the induction datum for \mathbb{O} with maximal semisimple corank such that the equivariant fundamental group is nontrivial. From this, we get

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{(3A_1)''}^{E_7} \left(\operatorname{univ. cover of} \begin{array}{c} 2 \\ 2 \end{array} \right).$$

3.3. $\pi_1^G(\mathbb{O}) = S_2$. There are 38 induced nilpotent orbits in exceptional cases with $\pi_1^G(\mathbb{O}) = S_2$; that is, there are no such nilpotent orbits in type G_2 , 5 such in type F_4 , 1 such in type E_6 , 8 such in type E_7 and 24 such in type E_8 . As in Subsection 3.2, the only non-trivial nilpotent cover of such \mathbb{O} is the universal cover $\widehat{\mathbb{O}}$ and if the universal cover is birationally rigid then \mathbb{O} is birationally induced from its unique rigid induction datum. Using Proposition 2.11, this handles 3 orbits for F_4 , the orbit for E_6 , 1 orbit for E_7 , and 6 orbits for E_8 .

By a similar argument, if \mathbb{O} is birationally rigid then \mathbb{O} must have a unique rigid induction datum and the universal cover of \mathbb{O} must be birationally induced from such rigid induction datum. This resolves another 2 orbits in type E_7 and 2 orbits in type E_8 . We thus have 2 remaining induced nilpotent orbits in type F_4 , 5 in type E_7 and 16 in type E_8 .

If \mathbb{O} is even then we know by Proposition 2.5 that \mathbb{O} is birationally induced from the zero orbit in the Jacobson-Morozov Levi. Thus, if \mathbb{O} is even and has exactly two rigid induction data, then we know that \mathbb{O} is birationally induced from the zero orbit in the Jacobson-Morozov Levi and the universal cover of \mathbb{O} must be birationally induced from the other rigid induction datum. This resolves the remaining 2 orbits in type F_4 , 3 more orbits in type E_7 and 8 more orbits in type E_8 .

We deal with the remaining 10 nilpotent orbits individually.

3.3.1. $A_3 + A_2 \subseteq E_7$. The nilpotent orbit \mathbb{O} in E_7 with Bala-Carter label $A_3 + A_2$ has weighted Dynkin diagram

and has two rigid induction data

By the proof of $[Fu, Proposition 3.1]^6$ we have

$$\mathbb{O} = \operatorname{Bind}_{D_5+A_1}^{E_7} \left(\begin{array}{ccc} 0 & & \\ 0 & 1 & 0 & 0 \end{array} \right)$$

and therefore

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{D_6}^{E_7} \left(\begin{array}{ccc} 0 & & \\ 0 & 0 & 1 & 0 & 1 \end{array} \right).$$

3.3.2. $D_5(a_1) \subseteq E_7$. The nilpotent orbit in E_7 with Bala-Carter label $D_5(a_1)$ has weighted Dynkin diagram

$$\begin{smallmatrix}&&0\\2&0&1&0&1&0\end{smallmatrix}$$

and has unique rigid induction datum

$$\begin{array}{ccc} & 0 \\ 0 & 0 & 0 \end{array} \subseteq A_4.$$

Induction from this rigid induction datum to \mathbb{O} passes through the nilpotent orbit

which has equivariant fundamental group S_2 . By Proposition 2.10, the nilpotent orbit in $\mathfrak{so}(12)$ with this weighted Dynkin diagram is birationally rigid; by Theorem 2.8, this nilpotent orbit is also birationally rigid in the standard Levi subalgebra of E_7 of type D_6 . Birational induction therefore sends the unique rigid induction datum to the universal cover of this induction datum, and thus we get

and

$$\mathbb{O} = \operatorname{Bind}_{D_6}^{E_7} \left(\begin{array}{ccc} 0 & & \\ 0 & 1 & 0 & 1 & 0 \end{array} \right)$$
$$\widehat{\mathbb{O}} = \operatorname{Bind}_{A_4}^{E_7} \left(\begin{array}{ccc} & 0 \\ 0 & 0 & 0 \end{array} \right).$$

3.3.3. $A_3 + A_2 \subseteq E_8$. The nilpotent orbit in E_8 with Bala-Carter label $A_3 + A_2$ has weighted Dynkin diagram

$$\begin{smallmatrix} 0\\1&0&0&1&0&0 \end{smallmatrix}$$

and has rigid induction data

By [Fu, Proposition 3.1], we have

$$\mathbb{O} = \operatorname{Bind}_{D_7}^{E_8} \left(\begin{array}{cccc} 0 & & \\ 0 & 0 & 0 & 1 & 0 \end{array} \right)$$

⁶The result in [Fu] is for E_7 of adjoint type, but the argument works equally well when G is simply connected.

and therefore

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{E_7}^{E_8} \left(egin{array}{ccccc} 0 & & \ 0 & 1 & 0 & 0 & 0 \end{array}
ight).$$

3.3.4. $A_4 \subseteq E_8$. The nilpotent orbit in E_8 with Bala-Carter label A_4 has weighted Dynkin diagram

$$\begin{smallmatrix}&&0\\2&0&0&0&0&0&2\end{smallmatrix}$$

and has unique rigid induction datum

Since \mathbb{O} is even, we know that

$$\mathbb{O} = \operatorname{Bind}_{D_6}^{E_8} \left(\begin{array}{ccc} 0 & & \\ 0 & 0 & 0 & 0 \end{array} \right).$$

To determine the birationally rigid induction datum for the universal cover of \mathbb{O} , we just need to find the induction datum for \mathbb{O} with maximal semisimple corank such that the equivariant fundamental group is non-trivial. This leads us to

$$\widehat{\mathbb{O}} = \text{Bind}_{E_7}^{E_8} \left(\text{univ. cover of} \quad \begin{array}{ccc} 0 & & \\ 2 & 0 & 0 & 0 & 0 \end{array} \right).$$

3.3.5. $D_5(a_1) \subseteq E_8$. The nilpotent orbit in E_8 with Bala-Carter label $D_5(a_1)$ has weighted Dynkin diagram

$$\begin{smallmatrix}&&0\\1&0&0&0&1&0&2\end{smallmatrix}$$

and has unique rigid induction datum

Induction from this rigid induction datum to \mathbb{O} passes through the nilpotent orbit

which has equivariant fundamental group S_2 . By Proposition 2.10, the nilpotent orbit in the Lie algebra of the semisimple simply connected algebraic group of type E_7 with the same weighted Dynkin diagram is birationally rigid; by Theorem 2.8, this is also true in this Levi subalgebra. Birational induction therefore sends the unique rigid induction datum to the universal cover of this induction datum, and thus we get

$$\mathbb{O} = \operatorname{Bind}_{E_7}^{E_8} \left(\begin{array}{cccc} 0 & & \\ 1 & 0 & 0 & 1 & 0 \end{array} \right)$$

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{E_6}^{E_7} \left(\begin{array}{ccc} 1 & & \\ 0 & 0 & 0 & 0 \end{array} \right).$$

3.3.6. $D_6(a_1) \subseteq E_8$. The nilpotent orbit in E_8 with Bala-Carter label $D_6(a_1)$ has weighted Dynkin diagram

$$\begin{smallmatrix}&&1\\0&1&0&0&0&1&2\end{smallmatrix}$$

and has unique rigid induction datum

$$0 \ 0 \ 0 \ 0 \ 0 \ \subseteq A_5.$$

We note that induction from this rigid induction datum to \mathbb{O} passes through the nilpotent orbit

which has equivariant fundamental group S_2 . By Proposition 2.5, this induction datum is birationally induced from the zero orbit in A_5 . We thus have

$$\mathbb{O} = \operatorname{Bind}_{A_5}^{E_8} \left(\begin{array}{ccccc} 0 & 0 & 0 & 0 \end{array} \right)$$

and

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{E_6}^{E_8} \left(\operatorname{univ. cover of} \begin{array}{c} 2 \\ 0 & 0 & 0 \end{array} \right).$$

By examining the equivariant fundamental groups for the induction data for this nilpotent orbit, we see that this birational induction datum for $\widehat{\mathbb{O}}$ is birationally rigid.

3.3.7. $E_7(a_4) \subseteq E_8$. The nilpotent orbit in E_8 with Bala-Carter label $E_7(a_4)$ has weighted Dynkin diagram

and has two rigid induction data

By the proof of [Fu, Proposition 3.1], we have

$$\mathbb{O} = \operatorname{Bind}_{E_7}^{E_8} \left(\begin{array}{cccc} 0 & & \\ 0 & 0 & 1 & 0 & 1 & 0 \end{array} \right).$$

We may therefore argue as in Subsection 3.3.1 to get

$$\mathbb{O} = \operatorname{Bind}_{D_5 + A_1}^{E_8} \left(\begin{array}{ccc} 0 & & \\ 0 & 1 & 0 & 0 \end{array} \right)$$

and therefore

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{D_6}^{E_8} \left(\begin{array}{ccc} 0 & & \\ 0 & 0 & 1 & 0 & 1 \end{array} \right).$$

3.3.8. $D_7(a_2) \subseteq E_8$. The nilpotent orbit in E_8 with Bala-Carter label $D_7(a_2)$ has weighted Dynkin diagram

and has two rigid induction data

$$\mathbb{O}_1 := \begin{array}{ccc} 0 \\ 0 & 0 \end{array} \begin{array}{ccc} 0 \\ 0 \end{array} = \begin{array}{ccc} 0 \\ 0 \end{array} = \begin{array}{ccc} 0 \\ 0 \end{array} = \begin{array}{cccc} A_4 + 2A_1 =: L_1 \end{array}$$

$$\mathbb{O}_2 \coloneqq 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ \subseteq 2A_3 =: L_2.$$

By [Ho], we get that

$$N_G(L_1)/L_1 \simeq \mathsf{S}_2 \times \mathsf{S}_2$$
 and $N_G(L_2)/L_2 \simeq \mathsf{D}_8$

Note that by $[FJLS] \overline{\mathbb{O}}$ contains a single codimension 2 nilpotent orbit \mathbb{O}' , which is the nilpotent orbit with Bala-Carter label $D_5 + A_1$. Let \mathcal{L}' be the corresponding symplectic leaf of codimension 2 in Spec($\mathbb{C}[\mathbb{O}]$), as in [MM, Lemma 3.2.2]. This symplectic leaf has singularity of type C_2 (i.e. of type D_3 with the unique non-trivial graph automorphism of the Dynkin diagram of D_3 acting non-trivially).

Let \mathfrak{g}' be a simple complex Lie algebra of type D_3 , let \mathfrak{h}' be a Cartan subalgebra thereof, and let W' be the corresponding Weyl group. Then we have $W' \cong S_4$ and $\pi_1(\mathcal{L}') = \pi_1(\mathbb{O}') = S_2$ acts on W' such that the non-trival element acts as conjugation by (1, 4)(2, 3). By definition, we then get

$$W(\mathbb{O}) = (W')^{\pi_1(\mathcal{L}')} = \langle (1,4), (1,2,4,3) \rangle \simeq \mathsf{D}_8.$$

Let (L, \mathbb{O}_L) be the birationally rigid induction datum for \mathbb{O} ; by Proposition 2.7, we get that $W(\mathbb{O}) = N_G(L, \mathbb{O}_L)/L$. Since both \mathbb{O}_1 and \mathbb{O}_2 are the zero orbit for their respective Levi subgroups, we in fact have $W(\mathbb{O}) = N_G(L)/L$. By above, this therefore means that $L = L_2$ and $\mathbb{O}_L = \mathbb{O}_2$. This shows that

$$\mathbb{O} = \text{Bind}_{2A_3}^{E_8} (\begin{array}{ccccccc} 0 & 0 & 0 & 0 \end{array})$$

and hence that

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{A_4+2A_1}^{E_8} \left(\begin{array}{ccc} 0 & & \\ 0 & 0 & 0 & 0 \end{array} \right).$$

3.3.9. $E_6(a_1) + A_1 \subseteq E_8$. The nilpotent orbit in E_8 with Bala-Carter label $E_6(a_1) + A_1$ has weighted Dynkin diagram

$$\begin{smallmatrix}&&0\\1&0&1&0&1&0&2\end{smallmatrix}$$

and has unique rigid induction datum

$$\begin{array}{ccc} & 0 \\ 0 & 0 & 0 \end{array} \quad \begin{array}{c} 0 \\ G \end{array} \subseteq A_4 + A_1.$$

Induction from this rigid induction datum to \mathbb{O} passes through the nilpotent orbit

which has equivariant fundamental group S_2 . By Proposition 2.10, the nilpotent orbit in the Lie algebra of the semisimple simply connected algebraic group of type E_7 with the same weighted Dynkin diagram is birationally rigid; by Theorem 2.8, this is also true in this Levi subalgebra. Birational induction therefore sends the rigid induction datum to the universal cover of this induction datum, and thus we get

$$\mathbb{O} = \operatorname{Bind}_{E_7}^{E_8} \left(\begin{array}{cccc} 0 & & \\ 1 & 0 & 1 & 0 & 1 & 0 \end{array} \right)$$

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{A_4+A_1}^{E_7} \begin{pmatrix} 0 & \\ 0 & 0 & 0 \end{pmatrix}.$$
₂₀

3.3.10. $E_7(a_3) \subseteq E_8$. The nilpotent orbit in E_8 with Bala-Carter label $E_7(a_3)$ has weighted Dynkin diagram

$$\begin{smallmatrix}&&0\\2&0&1&0&1&0&2\end{smallmatrix}$$

and has unique rigid induction datum

$$\begin{array}{ccc} & 0 \\ 0 & 0 & 0 \end{array} \subseteq A_4.$$

Induction from this rigid induction datum to \mathbb{O} passes through the nilpotent orbit

$$\mathbb{O}_1 := \begin{array}{cccc} 0 \\ 0 & 1 & 0 & 1 & 0 \end{array} \subseteq D_6$$

which has equivariant fundamental group S_2 . By Proposition 2.10, the nilpotent orbit in $\mathfrak{so}(12)$ with this weighted Dynkin diagram is birationally rigid; by Theorem 2.8, this nilpotent orbit is also birationally rigid in the standard Levi subalgebra of E_7 of type D_6 . Birational induction therefore maps the rigid induction datum to the universal cover of this induction datum, and thus we get

$$\mathbb{O} = \operatorname{Bind}_{D_6}^{E_8} \left(\begin{array}{ccc} 0 & & \\ 0 & 1 & 0 & 1 & 0 \end{array} \right)$$

and

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{A_4}^{E_7} \left(\begin{array}{cc} 0 \\ 0 & 0 \end{array} \right).$$

3.4. $\pi_1^G(\mathbb{O}) = \mathbb{Z}/3\mathbb{Z}$. This case can only arise in type E_6 ; in this case, there are 4 nilpotent orbits with $\pi_1^G(\mathbb{O}) = \mathbb{Z}/3\mathbb{Z}$. As in Subsections 3.2 and 3.3, the only non-trivial nilpotent cover of such \mathbb{O} is the universal cover and if the universal cover is birationally rigid then \mathbb{O} is birationally induced from its unique rigid induction datum. Using Proposition 2.11, this handles 2 of the orbits.

The remaining two orbits have Bala-Carter labels E_6 and $E_6(a_1)$. These are both even and thus the orbits are birationally induced from the zero orbit in the appropriate Jacobson-Morozov Levi subalgebras. To determine the birationally rigid induction data for the universal covers, it is sufficient to determine which induction data (M, \mathbb{O}_M) for \mathbb{O} have non-trivial $\pi_1^M(\mathbb{O}_M)$. The unique such (M, \mathbb{O}_M) with the largest semisimple corank then must have the property that the universal cover of \mathbb{O}_M is birationally rigid and birationally induces to the universal cover of \mathbb{O} . With the help of atlas [At], we compute that for the nilpotent orbit E_6 we have

$$\mathbb{O} = \operatorname{Bind}_{T}^{E_{6}}(\{0\})$$

and

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{2A_2}^{E_6} \left(\text{univ. cover of } 2 \quad 2 \quad 2 \quad 2 \quad \right),$$

while for the nilpotent orbit $E_6(a_1)$ we have

$$\mathbb{O} = \operatorname{Bind}_{A_1}^{E_6} \left(\begin{array}{c} 0 \end{array} \right)$$

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{2A_2+A_1}^{E_6} \left(\operatorname{univ. cover of} \begin{array}{ccc} 0 \\ 2 & 2 \end{array} \right).$$

3.5. $\pi_1^G(\mathbb{O}) = S_2 \times \mathbb{Z}/2\mathbb{Z}$. This case can only arise in type E_7 ; in this case, there are 3 nilpotent orbits with $\pi_1^G(\mathbb{O}) = S_2 \times \mathbb{Z}/2\mathbb{Z}$. We tackle these case-by-case.

Before doing so, however, let us establish some notation and conventions. We denote the generator of S_2 by a and the generator of $\mathbb{Z}/2\mathbb{Z}$ by b. The subgroups of $\pi_1^G(\mathbb{O})$ are then 1, $S_2 = \langle a \rangle$, tw(S_2) = $\langle ab \rangle$, $\mathbb{Z}/2\mathbb{Z} = \langle b \rangle$ and $S_2 \times \mathbb{Z}/2\mathbb{Z}$. Let us denote the corresponding nilpotent orbit covers by $\widehat{\mathbb{O}}_a$, $\widetilde{\mathbb{O}}_{ab}$, $\widetilde{\mathbb{O}}_b$ and \mathbb{O} .

Since in this paper we only describe the *G*-equivariant fundamental group $\pi_1^G(\mathbb{O})$ up to isomorphism, we cannot canonically distinguish between subgroups which are interchanged by an automorphism of $\pi_1^G(\mathbb{O})$. Since there are automorphisms of $S_2 \times \mathbb{Z}/2\mathbb{Z}$ which permute the subgroups S_2 , tw(S_2), and $\mathbb{Z}/2\mathbb{Z}$, we cannot distinguish amongst these subgroups. When (the image under birational induction of) a birationally rigid induction datum corresponds to such a subgroup, we thus assign it to one of S_2 , tw(S_2) and $\mathbb{Z}/2\mathbb{Z}$ in a largely arbitrary way. When the birationally rigid induction datum would also exist for E_7 of adjoint type, we prefer to say that the image under birational induction corresponds to $\mathbb{Z}/2\mathbb{Z}$; when it doesn't, we prefer S_2 or tw(S_2). We maintain our conventions from this section in Table 4.

3.5.1. $D_4(a_1) + A_1 \subseteq E_7$. The nilpotent orbit in E_7 with Bala-Carter label $D_4(a_1) + A_1$ has weighted Dynkin diagram

$$\begin{smallmatrix}&1\\0&1&0&0&0&1\end{smallmatrix}$$

This has unique rigid induction datum

$$0 \ 0 \ 0 \ 0 \ 0 \ \subseteq (A_5)'.$$

The only induction data for \mathbb{O} with semisimple corank 1 are therefore

1 0 0 0 0 1
$$\subseteq A_6$$
 and $\begin{array}{c} 2\\ 0 & 0 & 0 \end{array} \subseteq E_6;$

these have equivariant fundamental groups 1 and S_2 , respectively. Note that both these nilpotent orbits are birationally induced by Theorem 2.8 and Proposition 2.10, which means in particular that the universal cover of the latter orbit is birationally rigid.

By Proposition 2.11 and the discussion following it, the universal cover $\widehat{\mathbb{O}}$ of \mathbb{O} is birationally rigid. Therefore,

$$\mathbb{O} = \operatorname{Bind}_{(A_5)'}^{E_7} \left(\begin{array}{cccccc} 0 & 0 & 0 & 0 \end{array} \right)$$

while the universal cover of $\begin{pmatrix} 2 \\ 0 & 0 & 0 \end{pmatrix}$ birationally induces to a 2-fold nilpotent cover of \mathbb{O} .

We therefore conclude that one of the order 2 subgroups corresponds to this birationally induced 2-fold nilpotent cover and the other two correspond to birationally rigid nilpotent covers. Following the discussion at the beginning of this subsection, we make the convention that

$$\widetilde{\mathbb{O}}_b = \operatorname{Bind}_{E_6}^{E_7} \left(\operatorname{univ. cover of} \begin{array}{ccc} 2 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

and that $\widetilde{\mathbb{O}}_a$ and $\widetilde{\mathbb{O}}_{ab}$ are birationally rigid.

3.5.2. $E_7(a_4) \subseteq E_7$. The nilpotent orbit in E_7 with Bala-Carter label $E_7(a_4)$ has weighted Dynkin diagram

This nilpotent orbit has two rigid induction data:

The only induction data of semisimple corank 1 with non-trivial equivariant fundamental group are

which have equivariant fundamental groups S_2 and $\mathbb{Z}/2\mathbb{Z}$, respectively. The universal cover of the latter of these is birationally rigid (as the underlying nilpotent orbit is birationally induced by Theorem 2.8 and Proposition 2.10 and no other induction data have any nontrivial nilpotent covers). By Proposition 2.11 and the discussion following it, the universal cover of \mathbb{O} is birationally rigid, and since \mathbb{O} is even we have

$$\mathbb{O} = \operatorname{Bind}_{A_2+2A_1}^{E_7} \left(\begin{array}{cc} 0 \\ 0 & 0 \end{array} \right).$$

Therefore, $\begin{array}{cccc} 1 & & \\ 1 & 0 & 1 & 0 \end{array} \subseteq D_4 + A_1$ and the universal cover of $\begin{array}{ccccc} 0 & & \\ 0 & 0 & 2 & 0 \end{array} = \begin{array}{cccccc} D_5 + A_1 \\ \text{induce to distinct 2-fold nilpotent covers of } \mathbb{O}. \end{array}$

Following the conventions discussed at the beginning of this section, we say that

$$\pi_1^G \left(\operatorname{Bind}_{D_4+A_1}^{E_7} \begin{pmatrix} 1 & & \\ 1 & 0 & 1 & 0 \end{pmatrix} \right) = \langle b \rangle = \mathbb{Z}/2\mathbb{Z},$$

$$\pi_1^G \left(\operatorname{Bind}_{D_5+A_1}^{E_7} \left(\operatorname{univ. cover of} \begin{array}{c} & 0 & \\ 0 & 0 & 2 & 0 & 2 \end{array} \right) \right) = \langle a \rangle = \mathsf{S}_2$$

and that the nilpotent cover of \mathbb{O} corresponding to the subgroup tw(S₂) is birationally rigid.

3.5.3. $E_7(a_3) \subseteq E_7$. The nilpotent orbit in E_7 with Bala-Carter label $E_7(a_3)$ has weighted Dynkin diagram

There are two rigid induction data for this nilpotent orbit:

$$\begin{array}{cccc} 0 & & \\ 0 & & 0 \end{array} \subseteq (3A_1)' \quad \text{and} \quad 0 \quad 0 \ \subseteq A_2. \end{array}$$

Note that \mathbb{O} is even, and thus $\mathbb{O} = \operatorname{Bind}_{(3A_1)'}^{E_8} \begin{pmatrix} 0 \\ 0 & 0 \end{pmatrix}$. The induction detune

The induction datum

$$\mathbb{O}_0 := \begin{array}{cccc} 0 \\ 2 & 0 & 2 & 0 \end{array} \begin{array}{cccc} 0 \\ 2 & 0 & 2 \end{array} \subseteq D_5 + A_1 =: L$$

for \mathbb{O} has equivariant fundamental group equal to $S_2 \times \mathbb{Z}/2\mathbb{Z}$; determining the birationally rigid induction data for the nilpotent covers of \mathbb{O} therefore reduces to determining the same for the nilpotent covers of \mathbb{O}_0 .

The induction data for (L, \mathbb{O}_0) with semisimple corank 1 and non-trivial equivariant fundamental group are

The first and fourth of these have equivariant fundamental group S_2 , while the second and third have equivariant fundamental group $\mathbb{Z}/2\mathbb{Z}$. The induction data \mathbb{O}_1 and \mathbb{O}_4 are induced from both of the rigid induction data for \mathbb{O} ; on the other hand, \mathbb{O}_2 is only induced from the zero orbit in A_2 and \mathbb{O}_3 is only induced from the zero orbit in $(3A_1)'$. Since \mathbb{O}_2 and \mathbb{O}_3 are birationally induced by Theorem 2.8 and Proposition 2.10 and there are no other possible birational induction data, the universal covers of \mathbb{O}_2 and \mathbb{O}_3 are birationally rigid. We shall denote them by $\widehat{\mathbb{O}}_2$ and $\widehat{\mathbb{O}}_3$, respectively. The same argument also shows that \mathbb{O}_2 is birationally induced from the zero orbit in A_2 and \mathbb{O}_3 is birationally induced from the zero orbit in $(3A_1)'$.

Since the zero orbit in $(3A_1)'$ birationally induces to \mathbb{O} , the zero orbit in A_2 birationally induces to a non-trivial nilpotent cover of \mathbb{O} . Similarly $\widehat{\mathbb{O}}_2$ birationally induces to a nilpotent cover of $\operatorname{Bind}_{A_2+3A_1}^{E_8}(\mathbb{O}_2) = \operatorname{Bind}_{A_2}^{E_8}(0\ 0\)$. Therefore, we must have

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{A_2+3A_1}^{E_8}(\widehat{\mathbb{O}}_2).$$

The remaining birationally induced nilpotent covers of \mathbb{O} therefore correspond to subgroups of $\pi_1^G(\mathbb{O})$ of order 2.

Following the discussion in the introduction of Section 3.5, we make the conventions that

$$\widetilde{\mathbb{O}}_b = \operatorname{Bind}_{A_2}^{E_7} (\begin{array}{cc} 0 & 0 \end{array})$$

and

$$\widetilde{\mathbb{O}}_a = \operatorname{Bind}_{A_3+2A_1}^{E_7}(\widehat{\mathbb{O}}_3).$$

The only remaining nilpotent cover of \mathbb{O} is the nilpotent cover corresponding to tw(S₂); from what we have already seen, this must be induced from the nilpotent cover of \mathbb{O}_0 corresponding to tw(S₂),⁷ which itself must be birationally rigid. This resolves the final nilpotent cover of \mathbb{O} .

3.6. $\pi_1^G(\mathbb{O}) = S_2 \times \mathbb{Z}/3\mathbb{Z}$. This case can only arise when G is of type E_6 , in which case \mathbb{O} is the nilpotent orbit with Bala-Carter label $E_6(a_3)$ and weighted Dynkin diagram

There are two rigid induction data for this nilpotent orbit:

$$\begin{array}{ccc} 0 \\ 0 \\ 0 \end{array} \subseteq 3A_1 \quad \text{and} \quad 0 \quad 0 \subseteq A_2.$$

Induction from each of these induction data passes through the nilpotent orbit

$$\begin{array}{ccc} 0 \\ 0 & 2 & 0 \end{array} \subseteq D_4,$$

which has equivariant fundamental group S_2 . By Lemma 2.3, the (2-fold) universal cover of this nilpotent orbit must therefore birationally induce to a 2*n*-fold universal cover of \mathbb{O} for some $n \in \mathbb{N}$.

Note that there are four subgroups of $S_2 \times \mathbb{Z}/3\mathbb{Z}$: 1, S_2 , $\mathbb{Z}/3\mathbb{Z}$ and $S_2 \times \mathbb{Z}/3\mathbb{Z}$. Since \mathbb{O} is even, we must have

$$\pi_1^G \left(\text{Bind}_{3A_1}^{E_6} \begin{pmatrix} 0 \\ 0 & 0 \end{pmatrix} \right) = \mathsf{S}_2 \times \mathbb{Z}/3\mathbb{Z}$$

and, by Proposition 2.11 and the discussion following it, the universal cover of \mathbb{O} is birationally rigid. Putting this all together, we must therefore have that

$$\pi_1^G \left(\text{Bind}_{D_4}^{E_6} \left(\text{univ. cover of } \begin{array}{c} 0 \\ 0 \end{array} \right) \right) = \mathbb{Z}/3\mathbb{Z}$$

univ. cover of
$$\begin{array}{cc} 0\\ 0 & 2 & 0 \end{array} = \operatorname{Bind}_{A_2}^{D_4} \left(\begin{array}{cc} 0 & 0 \end{array} \right).$$

⁷We assume here that we have denoted the subgroups of $\pi_1^L(\mathbb{O})$ and $\pi_1^G(\mathbb{O})$ in such a way that birational induction corresponds to the trivial map on conjugacy classes of subgroups.

This thus implies that

$$\pi_1^G \left(\operatorname{Bind}_{A_2}^{E_6} \left(\begin{array}{cc} 0 & 0 \end{array} \right) \right) = \mathbb{Z}/3\mathbb{Z}$$

Finally, we note that the induction datum

1

$$0 \ 2 \ 0 \ 2 \ 0 \ \subseteq A_5$$

has equivariant fundamental group $\mathbb{Z}/3\mathbb{Z}$. By examining the induction data for this nilpotent orbit and using Theorem 2.8 and Proposition 2.10, we see that the universal cover of this nilpotent orbit must be birationally rigid. Since the universal cover of \mathbb{O} is birationally rigid, by process of elimination we must have

$$\tau_1^G \left(\operatorname{Bind}_{A_5}^{E_6} \left(3 \text{-fold cover of } 0 \ 2 \ 0 \ 2 \ 0 \ \right) \right) = \mathsf{S}_2.$$

3.7. $\pi_1^G(\mathbb{O}) = S_3$. There are 8 induced nilpotent orbits in exceptional types with $\pi_1^G(\mathbb{O}) = S_3$; more specifically, there is one in type G_2 , none in type F_4 , one in type E_6 , one in type E_7 and 5 in type E_8 . Note that S_3 (which we identify as permutations of the set $\{1, 2, 3\}$) has four subgroups up to conjugacy: the trivial group 1, the subgroup $S_2 := \langle (1, 2) \rangle$ (which is conjugate to $\langle (1, 3) \rangle$ and $\langle (2, 3) \rangle$), the alternating group A_3 and the whole group S_3 . These correspond to the four *G*-equivariant nilpotent covers of \mathbb{O} (up to isomorphism), which we denote respectively by $\widehat{\mathbb{O}}, \widetilde{\mathbb{O}}, \widetilde{\mathbb{O}}$ and \mathbb{O} .

For the nilpotent orbit with Bala-Carter label $G_2(a_1)$ in G_2 , the birationally rigid induction data are given in [LMM, Example 8.4.1]. The remaining nilpotent orbits we tackle individually.

3.7.1. $D_4(a_1) \subseteq E_6$. The nilpotent orbit in E_6 with Bala-Carter label $D_4(a_1)$ has weighted Dynkin diagram

$$\begin{smallmatrix}&&0\\0&0&2&0&0\end{smallmatrix}.$$

The rigid induction data for this nilpotent orbit are

Since \mathbb{O} is even, we have

$$\mathbb{O} = \operatorname{Bind}_{2A_2 + A_1}^{E_6} \left(\begin{array}{ccc} 0 & \\ 0 & 0 & 0 \end{array} \right).$$

Note furthermore that

$$\operatorname{Ind}_{A_3+A_1}^{D_5} \left(\begin{array}{cc} 0 \\ 0 & 0 \end{array}\right) = \operatorname{Ind}_{D_4}^{D_5} \left(\begin{array}{cc} 0 \\ 0 & 1 \end{array}\right) = \begin{array}{cc} 0 \\ 0 & 0 \end{array} \subseteq D_5$$

which has equivariant fundamental group S_2 . Since this nilpotent orbit is even, we have

$$\begin{array}{ccc} 0 \\ 0 & 0 & 2 & 0 \end{array} = \operatorname{Bind}_{A_3+A_1}^{E_6} \left(\begin{array}{ccc} 0 \\ 0 & 0 & 0 \end{array} \right)$$

and therefore the other rigid induction datum birationally induces to the (2-fold) universal cover. This therefore implies by Proposition 2.3 that $\operatorname{Bind}_{D_4}^{E_6} \begin{pmatrix} 0\\ 0 & 1 & 0 \end{pmatrix}$ is a 2-fold nilpotent cover of $\operatorname{Bind}_{A_3+A_1}^{E_6} \begin{pmatrix} 0\\ 0 & 0 & 0 \end{pmatrix}$. Amongst $\widehat{\mathbb{O}}, \widecheck{\mathbb{O}}$ and $\widetilde{\mathbb{O}}$, the only such possibility is therefore that

$$\widehat{\mathbb{O}} = \operatorname{Bind}_{D_4}^{E_6} \left(\begin{array}{cc} 0 \\ 0 & 1 & 0 \end{array} \right)_{25}$$

and

$$\check{\mathbb{O}} = \operatorname{Bind}_{A_3+A_1}^{E_6} \left(\begin{array}{cc} 0 & \\ 0 & 0 & 0 \end{array} \right).$$

By Proposition 2.11, \mathbb{O} has a birationally rigid nilpotent cover; this must therefore be \mathbb{O} .

3.7.2. $D_4(a_1) \subseteq E_7$. The nilpotent orbit in E_7 with Bala-Carter label $D_4(a_1)$ has weighted Dynkin diagram

The rigid induction data for this nilpotent orbit are

Since \mathbb{O} is even, we have

$$\mathbb{O} = \operatorname{Bind}_{A_5+A_1}^{E_7} \left(\begin{array}{ccc} 0 & & \\ 0 & 0 & 0 & 0 \end{array} \right).$$

Furthermore, Proposition 2.11 and the discussion following it show that $\widehat{\mathbb{O}}$ is birationally rigid. By examining the equivariant fundamental groups for the rigid induction data for \mathbb{O} , we see that one of $\widetilde{\mathbb{O}}$ and $\widetilde{\mathbb{O}}$ must be birationally rigid and the other birationally induced from $\begin{array}{c} 0\\ 0 & 1 & 0 & 0 \end{array} \subseteq D_6$. Note that by [FJLS] and [MM, Lemma 3.2.2] Spec($\mathbb{C}[\mathbb{O}]$) has a symplectic leaf of codimension 2 whose corresponding singularity has type A_1 . Since $\widetilde{\mathbb{O}}$, as a 3-fold nilpotent cover, cannot smooth any A_1 singularity, we must have that Spec($\mathbb{C}[\widetilde{\mathbb{O}}]$) has a codimension 2 leaf and thus by [LMM, Corollary 7.6.1] that $\widetilde{\mathbb{O}}$ is birationally induced. We conclude that $\widetilde{\mathbb{O}}$ is birationally rigid and that

$$\tilde{\mathbb{O}} = \operatorname{Bind}_{D_6}^{E_7} \left(\begin{array}{ccc} 0 & & \\ 0 & 1 & 0 & 0 \end{array} \right).$$

3.7.3. $D_4(a_1) \subseteq E_8$. The nilpotent orbit in E_8 with Bala-Carter label $D_4(a_1)$ has weighted Dynkin diagram

The rigid induction data for this nilpotent orbit are

This is even, and therefore we get

$$\mathbb{O} = \operatorname{Bind}_{E_6+A_1}^{E_8} \left(\begin{array}{cccc} 0 & & \\ 0 & 0 & 0 & 0 & 0 \end{array} \right)$$

Furthermore, Proposition 2.11 and the discussion following it show that $\widehat{\mathbb{O}}$ is birationally rigid. Looking at the equivariant fundamental groups for the rigid induction data for \mathbb{O} , we see that one of $\widetilde{\mathbb{O}}$ and $\widetilde{\mathbb{O}}$ must be birationally rigid and the other birationally induced from $\begin{array}{c} 0 \\ 0 \\ 0 \end{array} \subset \begin{array}{c} 0 \\ 0 \\ 0 \end{array} \subset \begin{array}{c} 0 \\ 0 \\ 0 \end{array} \subset \begin{array}{c} E_7. \end{array}$

Note that by [FJLS] and [MM, Lemma 3.2.2] Spec($\mathbb{C}[\mathbb{O}]$) has a symplectic leaf of codimension 2 whose corresponding singularity has type A_1 . Since \mathbb{O} , as a 3-fold nilpotent cover, cannot smooth

any A_1 singularity, we must have that $\text{Spec}(\mathbb{C}[\check{\mathbb{O}}])$ has a codimension 2 leaf and thus by [LMM, Corollary 7.6.1] that $\check{\mathbb{O}}$ is birationally induced. We conclude that $\widetilde{\mathbb{O}}$ is birationally rigid and that

$$\tilde{\mathbb{O}} = \operatorname{Bind}_{E_7}^{E_8} \left(\begin{array}{cccc} 0 & & \\ 0 & 0 & 0 & 0 & 1 & 0 \end{array} \right).$$

3.7.4. $E_7(a_5) \subseteq E_8$. The nilpotent orbit in E_8 with Bala-Carter label $E_7(a_5)$ has weighted Dynkin diagram

$$\begin{smallmatrix}&&0\\0&0&1&0&1&0&0\end{smallmatrix}.$$

There are two rigid induction data for this nilpotent orbit:

both of which have trivial equivariant fundamental group. Since the Levi subgroups have semisimple corank 1, only two nilpotent covers of \mathbb{O} can be birationally induced and therefore two nilpotent covers must be birationally rigid.

By Proposition 2.11 and the discussion following it, the universal cover $\widehat{\mathbb{O}}$ is birationally rigid, and by [Fu, Proposition 3.1] we have

$$\mathbb{O} = \operatorname{Bind}_{E_6+A_1}^{E_8} \left(\begin{array}{cccc} 0 & & \\ 0 & 0 & 1 & 0 & 0 \end{array} \right).$$

What remains is therefore to determine which of \mathbb{O} and \mathbb{O} are birationally rigid.

Note that by [FJLS] and [MM, Lemma 3.2.2] Spec($\mathbb{C}[\mathbb{O}]$) has a symplectic leaf of codimension 2 whose corresponding singularity has type $2A_1$. Since $\check{\mathbb{O}}$, as a 3-fold nilpotent cover, cannot smooth any $2A_1$ singularity, we must have that Spec($\mathbb{C}[\check{\mathbb{O}}]$) has a codimension 2 leaf and thus by [LMM, Corollary 7.6.1] that $\check{\mathbb{O}}$ is birationally induced. We conclude that $\widetilde{\mathbb{O}}$ is birationally rigid and that

$$\breve{\mathbb{O}} = \operatorname{Bind}_{E_7}^{E_8} \left(\begin{array}{cccc} 0 & & \\ 1 & 0 & 1 & 0 & 0 \end{array} \right).$$

3.7.5. $E_8(b_6) \subseteq E_8$. The nilpotent orbit in E_8 with Bala-Carter label $E_8(b_6)$ has weighted Dynkin diagram

There are three rigid induction data for this nilpotent orbit, which are

Since \mathbb{O} is even we have

$$\mathbb{O} = \operatorname{Bind}_{A_3 + A_2 + A_1}^{E_8} \left(\begin{array}{ccc} 0 & & \\ 0 & 0 & & 0 & 0 \end{array} \right),$$

and we know from Proposition 2.11 and the discussion following it that $\widehat{\mathbb{O}}$ is birationally rigid. It thus just remains to determine the birationally rigid induction data for $\check{\mathbb{O}}$ and $\widetilde{\mathbb{O}}$.

Note that

$$\mathbb{O} = \operatorname{Ind}_{D_5+A_2}^{E_8} \left(\begin{array}{ccc} 0 & & \\ 0 & 2 & 0 & 0 & 0 \end{array} \right)$$
27

and that

$$\begin{array}{cccc} 0 & & \\ 0 & 2 & 0 & 0 & 0 \end{array} = \operatorname{Ind}_{A_3+A_2+A_1}^{D_5+A_2} \left(\begin{array}{cccc} 0 & & \\ 0 & 0 & 0 & 0 \end{array} \right) = \operatorname{Ind}_{D_4+A_2}^{D_5+A_2} \left(\begin{array}{cccc} 0 & & \\ 0 & 1 & 0 & 0 \end{array} \right)$$

has equivariant fundamental group S_2 . By Proposition 2.5,

$$\begin{array}{cccc} 0 & & & \\ 0 & 2 & 0 & 0 & 0 \end{array} = \operatorname{Bind}_{A_3 + A_2 + A_1}^{D_5 + A_2} \left(\begin{array}{cccc} 0 & & & \\ 0 & 0 & 0 & 0 \end{array} \right),$$

$$\operatorname{Bind}_{D_4+A_2}^{E_8} \left(\begin{array}{ccc} 0 & & \\ 0 & 1 & 0 & 0 \end{array} \right)$$

is a 2-fold nilpotent cover of \mathbb{O} by Proposition 2.3. Since we know that $\widehat{\mathbb{O}}$ is birationally rigid, we must therefore have

$$\widetilde{\mathbb{O}} = \operatorname{Bind}_{D_4+A_2}^{E_8} \left(\begin{array}{ccc} 0 & & \\ 0 & 1 & 0 & 0 \end{array} \right)$$

and, by process of elimination,

$$\check{\mathbb{O}} = \operatorname{Bind}_{E_6+A_1}^{E_8} \left(\begin{array}{cccc} 0 & & \\ 1 & 0 & 1 & 0 & 1 & 0 \end{array} \right).$$

3.7.6. $E_8(a_6) \subseteq E_8$. The nilpotent orbit in E_8 with Bala-Carter label $E_8(a_6)$ has weighted Dynkin diagram

Note that

$$\begin{array}{cccccccc} 0 & & & \\ 0 & 0 & 2 & 0 & 0 & 2 & 0 \end{array} = \operatorname{Ind}_{E_6+A_1}^{E_8} \left(\begin{array}{ccccccccccccc} 0 & & & & \\ 0 & 0 & 2 & 0 & 0 & & 0 \end{array} \right)$$

and that $\pi_1^{E_6+A_1} \begin{pmatrix} 0 \\ 0 & 0 & 2 & 0 & 0 \end{pmatrix} = \mathsf{S}_3$ by (3.1). Using Proposition 2.3, it is easy to see the birationally rigid induction data for $E_8(a_6)$ in E_8 are the same as the birationally rigid induction data for $D_4(a_1) \times \{0\}$ in $E_6 + A_1$. Arguing as in Subsection 3.7.1, we therefore get

3.7.7. $E_8(b_5) \subseteq E_8$. The nilpotent orbit in E_8 with Bala-Carter label $E_8(b_5)$ has weighted Dynkin diagram

Note that

and that $\pi_1^{E_6} \begin{pmatrix} 0 \\ 0 & 0 & 2 & 0 & 0 \end{pmatrix} = S_3$. Using Proposition 2.3, it is easy to see the birationally rigid induction data for $E_8(b_5)$ in E_8 are the same as the birationally rigid induction data for $D_4(a_1)$ in E_6 . Arguing as in Subsection 3.7.1, we therefore get

3.8. $\pi_1^G(\mathbb{O}) = \mathsf{S}_3 \times \mathbb{Z}/2\mathbb{Z}$. This case can only arise in type E_7 ; in this case, there is one nilpotent orbit with $\pi_1^G(\mathbb{O}) = \mathsf{S}_3 \times \mathbb{Z}/2\mathbb{Z}$. This is the nilpotent orbit with Bala-Carter label $E_7(a_5)$, which has weighted Dynkin diagram

$$\begin{smallmatrix} & 0 \\ 0 & 0 & 2 & 0 & 0 & 2 \end{smallmatrix}$$

Since G-equivariant nilpotent covers of \mathbb{O} correspond to (conjugacy classes of) subgroups of $S_3 \times \mathbb{Z}/2\mathbb{Z}$, we record for reference the diagram of such subgroups and their inclusions (by which we mean that we draw a line between two subgroups when such an inclusion exists for some pair of subgroups in the conjugacy class). We view elements of S_3 as permutations of $\{1, 2, 3\}$ and denote the generator of $\mathbb{Z}/2\mathbb{Z}$ by b.



Here we use the notation S and A for the symmetric, and alternating. In most cases it should be clear which subgroup we are referring to (up to conjugacy), but we note

$$S_3 = \langle (1,2,3), (1,2) \rangle, \quad tw(S_3) = \langle (1,2,3), (1,2)b \rangle,$$

$$\mathfrak{S}_2 = \langle (1,2) \rangle, \quad \mathrm{tw}(\mathfrak{S}_2) = \langle (1,2)b \rangle, \quad \mathrm{and} \quad \mathbb{Z}/2\mathbb{Z} = \langle b \rangle.$$

There are three rigid induction data for \mathbb{O} , which are

Induction from each of these rigid nilpotent orbits passes through the nilpotent orbit

$$\mathbb{O}_0 := \begin{array}{cccc} 0 & 0 \\ 0 & 0 & 2 & 0 & 0 \end{array} \subseteq E_6 =: L_0,$$

which has equivariant fundamental group S_3 by (3.1). Labelling the nilpotent covers of \mathbb{O}_0 by $\widehat{\mathbb{O}}_0$, $\widetilde{\mathbb{O}}_0$ and \mathbb{O}_0 in the conventions of Subsection 3.7, we get from Subsection 3.7.1 that $\widetilde{\mathbb{O}}_0$ is birationally rigid and that

$$\mathbb{D}_{0} = \operatorname{Bind}_{2A_{2}+A_{1}}^{E_{7}} \begin{pmatrix} 0 \\ 0 & 0 & 0 \end{pmatrix},$$

$$\breve{\mathbb{D}}_{0} = \operatorname{Bind}_{(A_{3}+A_{1})'}^{E_{7}} \begin{pmatrix} 0 \\ 0 & 0 & 0 \end{pmatrix},$$

$$\widehat{\mathbb{D}}_{0} = \operatorname{Bind}_{D_{4}}^{E_{7}} \begin{pmatrix} 0 \\ 0 & 1 & 0 \end{pmatrix}.$$

Since \mathbb{O} and \mathbb{O}_0 are even, we have $\mathbb{O} = \text{Bind}_{E_6}^{E_7}(\mathbb{O}_0)$. We also know from Proposition 2.11 and the discussion following it that the universal cover $\widehat{\mathbb{O}}$ is birationally rigid.

As $\check{\mathbb{O}}_0$ is a 3-fold cover of \mathbb{O}_0 , Proposition 2.3 implies that $\operatorname{Bind}_{E_6}^{E_7}(\check{\mathbb{O}}_0)$ is a 3-fold cover of $\mathbb{O} = \operatorname{Bind}_{E_6}^{E_7}(\mathbb{O}_0)$. Thus, $\left|\pi_1^G(\mathbb{O}):\pi_1^G(\operatorname{Bind}_{E_6}^{E_7}(\check{\mathbb{O}}_0))\right| = 3$; the only option is therefore $\pi_1^G(\operatorname{Bind}_{E_6}^{E_7}(\check{\mathbb{O}}_0)) = S_2 \times \mathbb{Z}/2\mathbb{Z}$.

Similarly, as $\widetilde{\mathbb{O}}_0$ is a 2-fold cover of \mathbb{O}_0 we get $\left|\pi_1^G(\mathbb{O}):\pi_1^G(\operatorname{Bind}_{E_6}^{E_7}(\widetilde{\mathbb{O}}_0))\right| = 2$. Furthermore, since $\pi_1^{L_0}(\widetilde{\mathbb{O}}_0) \cong \mathsf{A}_3$, Proposition 2.14 implies that $\pi_1^G(\operatorname{Bind}_{E_6}^{E_7}(\widetilde{\mathbb{O}}))$ surjects onto A_3 . The only option is therefore $\pi_1^G(\operatorname{Bind}_{E_6}^{E_7}(\widetilde{\mathbb{O}})) = \mathsf{A}_3 \times \mathbb{Z}/2\mathbb{Z}$.

We also have that $\widehat{\mathbb{O}}_0$ is a 3-fold cover of $\widetilde{\mathbb{O}}_0$ and a 2-fold cover of $\widetilde{\mathbb{O}}_0$. This therefore implies that $\pi_1^G(\operatorname{Bind}_{E_6}^{E_7}(\widehat{\mathbb{O}}_0))$ is an index 2 subgroup of $S_2 \times \mathbb{Z}/2\mathbb{Z}$ and an index 3 subgroup of $A_3 \times \mathbb{Z}/2\mathbb{Z}$. We must therefore have $\pi_1^G(\operatorname{Bind}_{E_6}^{E_7}(\widehat{\mathbb{O}}_0)) = \mathbb{Z}/2\mathbb{Z}$.

Note furthermore that \mathbb{O} has induction datum

which has equivariant fundamental group $\mathbb{Z}/2\mathbb{Z}$. By Theorem 2.8 and Proposition 2.10, this nilpotent orbit must be birationally induced; since none of the induction data for this nilpotent orbit must therefore be birationally rigid. It must therefore birationally induce to a 2-fold cover of $\operatorname{Bind}_{(A_3+A_1)'}^{E_7}\begin{pmatrix} 0\\ 0 & 0 \end{pmatrix} = \operatorname{Bind}_{E_6}^{E_7}(\check{\mathbb{O}}_0)$. The corresponding subgroup of $S_3 \times \mathbb{Z}/2\mathbb{Z}$ must be an index 2 subgroup of $S_2 \times \mathbb{Z}/2\mathbb{Z}$ other than $\mathbb{Z}/2\mathbb{Z}$, hence either S_2 or tw(S_2). Note that there is an automorphism of $S_3 \times \mathbb{Z}/2\mathbb{Z}$ which interchanges these two subgroups; since $\pi_1^G(\mathbb{O})$ is only described

up to isomorphism we cannot distinguish between the two. We therefore make the convention (here and in Table 4) that

$$\pi_1^G \left(\operatorname{Bind}_{D_6}^{E_7} \left(\operatorname{univ. cover of} \begin{array}{cc} 2 \\ 0 & 0 \end{array} \right) \right) = \mathsf{S}_2.$$

Note that this nilpotent orbit in D_6 corresponds to the partition $(4^2, 2^2)$, which is very even. For the purposes of Table 4 we thus need to label it with either a I or II; we make the convention (consistent with [CM, Lemma 5.3.5]) that this nilpotent orbit has label I.

In the notation of Corollary 2.13, we have $\mathfrak{r} = \{0\}$ by [Ca, §13.1] and thus that covers of birationally rigid covers are birationally rigid. As a consequence of this, if a subgroup of $\pi_1^G(\mathbb{O})$ is an overgroup of a subgroup corresponding to a birationally induced cover that the overgroup must itself correspond to a birationally induced cover of \mathbb{O} . In our current setting, this means that the subgroup S_3 must correspond to a birationally induced cover of \mathbb{O} . By examining the induction data for \mathbb{O} of semisimple corank 1, we conclude that we must have

$$\mathsf{S}_{3} = \pi_{1}^{G} \left(\text{Bind}_{A_{5}+A_{1}}^{E_{7}} \left(\text{univ. cover of } \begin{array}{ccc} 0 & & \\ 0 & 0 & 2 & 0 & 0 \end{array} \right) \right).$$

We check that this induction datum is birationally rigid by examining the equivariant fundamental groups of the other induction data, noting that the underlying nilpotent orbit is not birationally rigid by Theorem 2.8 and Proposition 2.10. The same argument shows that none of the other nilpotent covers are birationally induced; they are thus all birationally rigid.

3.9. $\pi_1^G(\mathbb{O}) = \mathsf{S}_4$. This case only arises when G is of type F_4 , where there is a unique nilpotent orbit with $\pi_1^G(\mathbb{O}) = \mathsf{S}_4$. This is the nilpotent orbit with Bala-Carter label $F_4(a_3)$, which has weighted Dynkin diagram

$$0 \ 2 \ 0 \ 0$$

(where the first two nodes in the diagram correspond to long roots and the last two correspond to short roots). Since *G*-equivariant nilpotent covers of \mathbb{O} correspond to (conjugacy classes of) subgroups of S_4 , we record for reference the diagram of such subgroups and their inclusions (by which we mean that we draw a line between two subgroups when such an inclusion exists for some pair of subgroups in the conjugacy class).



Here we use the notation S, A, D and C for the symmetric, alternating, dihedral and cyclic groups, respectively (except that we prefer the notation S_2 to C_2 and A_3 to C_3). In most cases it should be clear to which subgroup we are referring (up to conjugacy), but we note

$$\begin{split} \mathsf{S}_2 \times \mathsf{S}_2 &= \langle (1,2), (3,4) \rangle, \quad \mathrm{tw}(\mathsf{S}_2 \times \mathsf{S}_2) &= \langle (1,2)(3,4), (1,3)(2,4) \rangle, \\ \mathsf{S}_2 &= \langle (1,2) \rangle, \quad \mathrm{and} \quad \mathrm{tw}(\mathsf{S}_2) &= \langle (1,2)(3,4) \rangle. \end{split}$$

Since \mathbb{O} is even it is birationally induced from the zero orbit in $A_2 + A_1$ (as described in Section 2, we use the notation \widetilde{A} for Levi subgroups in types G_2 and F_4 when we want to indicate that the corresponding simple roots are short). At the other extreme, we know from Proposition 2.11 and the discussion following it that the universal cover of \mathbb{O} is birationally rigid.

Note that by Corollary 2.13 any nilpotent cover of a birationally rigid nilpotent cover of \mathbb{O} is birationally rigid (since $\mathfrak{r} = \{0\}$ by [Ca, §13.1]). Therefore, any nilpotent cover of \mathbb{O} that is covered by a birationally induced nilpotent cover is itself birationally induced.

Other then the zero orbit in $A_2 + A_1$, the rigid induction data for \mathbb{O} are the nilpotent orbits

$$0 \ 0 \ 0 \ \subseteq A_2 + A_1$$
 and $0 \ \subseteq B_2 = C_2$.

We note also that

$$\operatorname{Ind}_{B_2}^{B_3}(0\ 0\) = 2\ 0\ 0$$
 and $\operatorname{Ind}_{C_2}^{C_3}(0\ 0\) = 0\ 1\ 0$

and that these two induced nilpotent orbits both have equivariant fundamental group S_2 . Finally, we note that by Theorem 2.8 and Proposition 2.10 the nilpotent orbit 2 0 0 in B_3 (corresponding to the partition $(3, 1^4)$) is birationally induced and the nilpotent orbit 0 1 0 in C_3 (corresponding to the partition $(2^2, 1^2)$) is birationally rigid. This therefore implies that

$$\operatorname{Bind}_{B_2}^{B_3} \left(\begin{array}{ccc} 0 & 0 \end{array} \right) = \begin{array}{ccc} 2 & 0 & 0 \end{array}$$

and

$$\operatorname{Bind}_{C_2}^{C_3} \left(\begin{array}{ccc} 0 & 0 \end{array} \right) = \operatorname{univ. \ cover \ of} \quad 0 \quad 1 \quad 0 \ .$$

Putting this all together, we get that that $\operatorname{Bind}_{B_3}^{F_4}$ (univ. cover of $2 \ 0 \ 0$) is a nilpotent cover of $\operatorname{Bind}_{B_2}^{F_4}$ ($0 \ 0$), which itself is a nilpotent cover of $\operatorname{Bind}_{C_3}^{F_4}$ ($0 \ 1 \ 0$). Passing to *G*-equivariant fundamental groups, this means that we get three nested subgroups of S_4 , and we know that neither 1 not S_4 can be in such list (as 1 corresponds to a birationally rigid nilpotent cover and S_4 corresponds to a nilpotent cover whose birationally rigid induction data we already know).

There are a total of 5 birationally induced nilpotent covers of \mathbb{O} (three coming from the rigid induction data, one coming from the universal cover of $0 \ 1 \ 0$ in C_3 and one coming from the birationally rigid nilpotent orbit $2 \ 0 \ 0$ in B_3). If the nilpotent cover corresponding to the subgroup tw(S₂) is birationally induced then all nilpotent covers corresponding to overgroups of tw(S₂) in S₄ are birationally induced. There are 6 such (conjugacy classes of) overgroups; since this exceeds the number of birationally induced nilpotent covers, we get a contradiction. Therefore, the nilpotent cover corresponding to tw(S₂) is birationally rigid.

With this in mind, the only chain of three nested (proper, non-trivial subgroups) in S_4 not including tw(S_2) is the chain

$$\mathsf{S}_2\subseteq\mathsf{S}_2\times\mathsf{S}_2\subseteq\mathsf{D}_8$$

This therefore implies

$$\pi_1^G \left(\operatorname{Bind}_{B_3}^{F_4} \left(\operatorname{univ. cover of} \ 2 \ 0 \ 0 \right) \right) = \mathsf{S}_2,$$
$$\pi_1^G \left(\operatorname{Bind}_{B_2}^{F_4} \left(\ 0 \ 0 \right) \right) = \mathsf{S}_2 \times \mathsf{S}_2,$$

and

$$\pi_1^G \left(\operatorname{Bind}_{C_3}^{F_4} \left(\begin{array}{ccc} 0 & 1 & 0 \end{array} \right) \right) = \mathsf{D}_8.$$

Since $S_2 \subseteq S_3$, the nilpotent cover corresponding to S_3 must also be birationally induced. There is only one remaining birationally rigid induction datum, and thus we must have

$$\mathsf{S}_3 = \pi_1^G \left(\operatorname{Bind}_{A_2 + \widetilde{A}_1}^{F_4} \left(\begin{array}{ccc} 0 & 0 \end{array} \right) \right).$$

3.10. $\pi_1^G(\mathbb{O}) = \mathsf{S}_5$. This case only arises in type E_8 , where there is a unique nilpotent orbit with $\pi_1^G(\mathbb{O}) = \mathsf{S}_5$. This is the nilpotent orbit with Bala-Carter label $E_8(a_7)$, which has weighted Dynkin diagram

Since G-equivariant nilpotent covers of \mathbb{O} correspond to (conjugacy classes of) subgroups of S_5 , we record for reference the diagram of such subgroups and their inclusions (by which we mean that we draw a line between two subgroups when such an inclusion exists for some pair of subgroups in the conjugacy class).



Here we use the notation S, A, D and C for the symmetric, alternating, dihedral and cyclic groups, respectively (except that we prefer the notation S_2 to C_2 and A_3 to C_3). In most cases it should be clear to which subgroup we are referring (up to conjugacy), but we note that

$$\begin{split} \mathsf{C}_5 \rtimes \mathsf{C}_4 &= \langle (1,2,3,4,5), (2,3,5,4) \rangle, \quad \mathsf{S}_3 = \langle (1,2,3), (1,2) \rangle, \quad \mathrm{tw}(\mathsf{S}_3) = \langle (1,2,3), (1,2)(4,5) \rangle, \\ &\mathsf{S}_2 \times \mathsf{S}_2 = \langle (1,2), (3,4) \rangle, \quad \mathrm{tw}(\mathsf{S}_2 \times \mathsf{S}_2) = \langle (1,2)(3,4), (1,3)(2,4) \rangle, \\ &\mathsf{S}_2 = \langle (1,2) \rangle \quad \mathrm{and} \quad \mathrm{tw}(\mathsf{S}_2) = \langle (1,2)(3,4) \rangle. \end{split}$$

The nilpotent orbit \mathbb{O} has four rigid induction data:

These nilpotent orbits all have trivial equivariant fundamental groups.

By induction from these rigid induction data, we see that the induction data for \mathbb{O} of semisimple corank 1 are the following:

Let us denote these Levi subgroups, in order, by L_1 , L_2 , L_3 , L_4 , L_5 , L_6 and L_7 . The equivariant fundamental groups of these nilpotent orbits are, respectively, 1, 1, S₂, 1, S₃, 1, and S₂. We deduce from this that \mathbb{O} admits eight birationally induced *G*-equivariant nilpotent covers: four birationally induced from the rigid nilpotent orbits, one birationally induced from a nilpotent cover of \mathbb{O}_3 , two birationally induced from nilpotent covers of \mathbb{O}_5 and one birationally induced from a nilpotent cover of \mathbb{O}_7 .

Let us establish some notation for the nilpotent covers of these induction data. For \mathbb{O}_3 and \mathbb{O}_7 (whose equivariant fundamental groups have 2 elements) we denote by $\widehat{\mathbb{O}}_3$ and $\widehat{\mathbb{O}}_7$ the (2-fold) universal covers. For \mathbb{O}_5 , which has equivariant fundamental group S_3 , we denote the nilpotent covers by $\widehat{\mathbb{O}}_5$, $\widetilde{\mathbb{O}}_5$, $\widetilde{\mathbb{O}}_5$, and \mathbb{O}_5 , following the conventions of Subsection 3.7.

By Proposition 2.5, the nilpotent orbits \mathbb{O}_3 and \mathbb{O}_5 are both birationally induced from the rigid nilpotent orbit $\begin{array}{ccc} 0 \\ 0 \end{array} \subseteq A_5 + A_1$. They therefore both birationally induce to the same nilpotent cover of \mathbb{O} . Furthermore, by Theorem 2.8 and Proposition 2.10, the nilpotent orbit \mathbb{O}_7 is birationally rigid and thus the rigid induction datum $\begin{array}{ccc} 0 \\ 0 \end{array} = \begin{array}{ccc} 0 \\ 1 \end{array} = \begin{array}{ccc} 0 \\ 0 \end{array} = \begin{array}{cccc} D_6 \end{array}$ birationally induces to $\widehat{\mathbb{O}}_7$. Arguing as in Subsection 3.7.2, we also get that $\check{\mathbb{O}}_5$ is birationally induced from this rigid induction datum; therefore both $\check{\mathbb{O}}_5$ and $\widehat{\mathbb{O}}_7$ birationally induce to the same nilpotent cover.

Since \mathbb{O} is even, we know that

$$\mathbb{O} = \operatorname{Bind}_{A_4+A_3}^{E_8}(\mathbb{O}_1).$$

We also know, by Proposition 2.11 and the discussion following it, that the universal cover of \mathbb{O} is birationally rigid. Furthermore, in the notation of Corollary 2.13 we have $\mathfrak{r} = \{0\}$ (by [Ca, §13.1]) and thus covers of birationally rigid covers are birationally rigid.

By Proposition 2.14, $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\mathbb{O}_5))$ surjects onto S_3 . Furthermore, since the universal cover of \mathbb{O} is birationally rigid the order of $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\mathbb{O}_5))$ must be at least 12 (were it 6, the universal cover of \mathbb{O}_5 would have to birationally induce to the universal cover of \mathbb{O}). Looking at the possible subgroups of S_5 , the only possibilities are that $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\mathbb{O}_5)) = S_2 \times S_3$ or $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\mathbb{O}_5)) = S_4$.

Let us first consider the case where $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\mathbb{O}_5)) = \mathsf{S}_4$. Then $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\widetilde{\mathbb{O}}_5))$ is an index 2 subgroup of S_4 (by Proposition 2.3); it must therefore be the alternating group $\mathsf{A}_4 \subseteq \mathsf{S}_4$.

We then get that $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\widehat{\mathbb{O}}_5) \leq \mathsf{A}_4$ is a subgroup of order 4 by Proposition 2.3. The only option is $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\widehat{\mathbb{O}}_5) = \operatorname{tw}(\mathsf{S}_2 \times \mathsf{S}_2)$. Then

$$\operatorname{tw}(\mathsf{S}_2 \times \mathsf{S}_2) \le \pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\breve{\mathbb{O}}_5)) \le \mathsf{S}_4,$$

which implies that we must have $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\breve{\mathbb{O}}_5)) = \mathsf{D}_8$. We note now that $\operatorname{Bind}_{L_5}^{E_8}(\breve{\mathbb{O}}_5) = \operatorname{Bind}_{L_7}^{E_8}(\widehat{\mathbb{O}}_7)$ is a nilpotent cover of $\operatorname{Bind}_{L_7}^{E_8}(\mathbb{O}_7)$ and so we must have $\mathsf{D}_8 \leq \pi_1^G(\operatorname{Bind}_{L_7}^{E_8}(\mathbb{O}_7))$. The only options are therefore that $\pi_1^G(\operatorname{Bind}_{L_7}^{E_8}(\mathbb{O}_7)) = \mathsf{S}_4$ or S_5 . Neither of these are possible, as S_4 and S_5 are already the equivariant fundamental groups of other birationally induced nilpotent orbits. It therefore cannot be the case that $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\mathbb{O}_5)) = \mathsf{S}_4$; we must have $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\mathbb{O}_5)) = \mathsf{S}_2 \times \mathsf{S}_3$.

Since $\widehat{\mathbb{O}}_5$ is a 6-fold nilpotent cover of \mathbb{O}_5 , $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\widehat{\mathbb{O}}_5))$ is a subgroup of $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\mathbb{O}_5)) = S_2 \times S_3$ of index 6 (by Proposition 2.3). Since $S_2 \times S_3$ has order 12, we must have that $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\widehat{\mathbb{O}}_5))$ has order 2. It must therefore be either (a) S_2 or (b) tw(S_2).

Suppose we are in case (b). Then we have

$$\operatorname{tw}(\mathsf{S}_2) \le \pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\widetilde{\mathbb{O}}_5)) \le \mathsf{S}_2 \times \mathsf{S}_3$$

and $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\widetilde{\mathbb{O}}_5))$ surjects onto $\pi_1^{L_5}(\widetilde{\mathbb{O}}_5) = A_3$. No such subgroups exist; we must therefore be in case (a), so

$$\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\widehat{\mathbb{O}}_5)) = \mathsf{S}_2$$

Once again, we must have

$$\mathsf{S}_2 \leq \pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\widetilde{\mathbb{O}}_5)) \leq \mathsf{S}_2 \times \mathsf{S}_3$$

and $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\widetilde{\mathbb{O}}_5))$ surjecting onto $\pi_1^{L_5}(\widetilde{\mathbb{O}}_5) = \mathsf{A}_3$. These two conditions force

$$\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\widetilde{\mathbb{O}}_5)) = \mathsf{C}_6$$

Similarly, we have

$$\mathsf{S}_2 \leq \pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\breve{\mathbb{O}}_5)) \leq \mathsf{S}_2 \times \mathsf{S}_3$$

and $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\breve{\mathbb{O}}_5))$ surjects onto $\pi_1^{L_5}(\breve{\mathbb{O}}_5) = \mathsf{S}_2$. Furthermore, as $\breve{\mathbb{O}}_5$ is a 3-fold nilpotent cover of \mathbb{O}_5 , $\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\breve{\mathbb{O}}_5))$ must be a subgroup of $\mathsf{S}_2 \times \mathsf{S}_3$ of index 3. This forces

$$\pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\breve{\mathbb{O}}_5)) = \mathsf{S}_2 \times \mathsf{S}_2.$$

Recall that $\tilde{\mathbb{O}}_5$ is not birationally rigid, but is instead birationally induced from the rigid induction datum whose Levi subgroup has type D_6 .

Recall that nilpotent covers of birationally rigid nilpotent covers are birationally rigid in this case. This implies that, since S_2 corresponds to a birationally induced nilpotent cover of \mathbb{O} , all overgroups of S_2 correspond to birationally induced nilpotent covers of \mathbb{O} . The only remaining overgroups of S_2 whose corresponding nilpotent cover we have not yet determined are S_3 , D_8 , and S_4 . Furthermore, there are three more birationally rigid induction data for covers of \mathbb{O} for which we have yet to determine the equivariant fundamental groups of their images under birational induction: $(L_2, \mathbb{O}_2), (L_3, \widehat{\mathbb{O}}_3)$, and (L_7, \mathbb{O}_7) . We therefore have

$$\{\pi_1^G(\operatorname{Bind}_{L_2}^{E_8}(\mathbb{O}_2)), \pi_1^G(\operatorname{Bind}_{L_3}^{E_8}(\widehat{\mathbb{O}}_3)), \pi_1^G(\operatorname{Bind}_{L_7}^{E_8}(\mathbb{O}_7))\} = \{\mathsf{S}_3, \mathsf{D}_8, \mathsf{S}_4\},\$$

and what remains is to determine which of these birationally induced nilpotent covers corresponds to which of these subgroups.

Since

$$\pi_1^G(\operatorname{Bind}_{L_3}^{E_8}(\widehat{\mathbb{O}}_3)) \le \pi_1^G(\operatorname{Bind}_{L_3}^{E_8}(\mathbb{O}_3)) = \pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\mathbb{O}_5)) = \mathsf{S}_2 \times \mathsf{S}_3,$$

the remaining options for $\pi_1^G(\operatorname{Bind}_{L_3}^{E_8}(\widehat{\mathbb{O}}_3))$ whose birationally rigid induction data aren't known are S_3 , tw(S_3), A_3 or tw(S_2). Since we precisely know which subgroups can arise from the such data, the only possibility is

$$\pi_1^G(\operatorname{Bind}_{L_2}^{E_8}(\widehat{\mathbb{O}}_3)) = \mathsf{S}_3$$

We therefore have

 $\{\pi_1^G(\operatorname{Bind}_{L_2}^{E_8}(\mathbb{O}_2)), \pi_1^G(\operatorname{Bind}_{L_7}^{E_8}(\mathbb{O}_7))\} = \{\mathsf{D}_8,\mathsf{S}_4\}.$

Since $S_2 \times S_2 = \pi_1^G(\operatorname{Bind}_{L_5}^{E_8}(\breve{\mathbb{O}}_5)) = \pi_1^G(\operatorname{Bind}_{L_7}^{E_8}(\widehat{\mathbb{O}}_7))$ is an index 2 subgroup of $\pi_1^G(\operatorname{Bind}_{L_7}^{E_8}(\mathbb{O}_7))$ by Proposition 2.3, we must have

$$\pi_1^G(\operatorname{Bind}_{L_7}^{E_8}(\mathbb{O}_7)) = \mathsf{D}_8.$$

Finally, by process of elimination we get

$$\pi_1^G(\operatorname{Bind}_{L_2}^G(\mathbb{O}_2)) = \mathsf{S}_4$$

The remaining covers are all birationally rigid.

4. TABLES

In this section, we record the birationally rigid induction data for the nilpotent orbit covers in exceptional Lie algebras. The tables are laid out as follows. There is one table for each type: G_2 , F_4, E_6, E_7 and E_8 . Each row corresponds to a G-equivariant nilpotent cover \mathbb{O} of an induced nilpotent G-orbit \mathbb{O} . For a given \mathbb{O} , the nilpotent covers are ordered starting with the trivial cover \mathbb{O} and ending with the universal cover \mathbb{O} . The first column contains the Bala-Carter label of \mathbb{O} – we only include this for the row corresponding to the nilpotent orbit. In the second column we give $\pi_1^G(\mathbb{O})$ as a subgroup of $\pi_1^G(\mathbb{O})$ – this represents the conjugacy class of subgroups which corresponds to the isomorphism class of $\widetilde{\mathbb{Q}}$. In the third column we indicate whether or not $\widetilde{\mathbb{Q}}$ is birationally rigid. If it is, we write 'Y' and leave all the remaining columns blank; if not, we write 'N' and fill in the remaining columns as follows. For a birationally induced nilpotent cover \mathbb{O} , let (L, \mathbb{O}_L) be the birationally rigid induction datum for $\widetilde{\mathbb{O}}$, where $\widetilde{\mathbb{O}}_L$ is a birationally rigid *L*-equivariant nilpotent cover of a nilpotent L-orbit \mathbb{O}_L . If $\widetilde{\mathbb{O}}$ is birationally induced, then in the fourth column we give the Dynkin type of L as described in Section 2; we write T when L is the maximal torus of G. In the fifth column we give either the partition or the Bala-Carter label corresponding to the nilpotent L-orbit \mathbb{O}_L (depending on whether L is of classical or exceptional type), except that we always write 0 for the zero orbit instead of giving the associated partition. In the sixth column we give the nilpotent cover $\widetilde{\mathbb{O}}_L$ by means of its L-equivariant fundamental group $\pi_1^L(\widetilde{\mathbb{O}}_L) \subseteq \pi_1^L(\mathbb{O}_L)$ (up to conjugation).

In describing the subgroups of *G*-equivariant fundamental groups, we use the following notation as in Section 3. When $\pi_1^G(\mathbb{O}) = S_2 \times \mathbb{Z}/2\mathbb{Z}$ we let $a \in S_2$ and $b \in \mathbb{Z}/2\mathbb{Z}$ generate $\pi_1^G(\mathbb{O})$. When $\pi_1^G(\mathbb{O}) = S_3 \times \mathbb{Z}/2\mathbb{Z}$, we denote the elements of S_3 as permutations of $\{1, 2, 3\}$ in cycle notation and let *b* be a generator of $\mathbb{Z}/2\mathbb{Z}$. We maintain the conventions of the relevant subsections of Section 3 for each of the orbits in the below tables, including the notation for subgroups of $S_2 \times \mathbb{Z}/2\mathbb{Z}$, $S_2 \times \mathbb{Z}/3\mathbb{Z}$, S_3 , $S_3 \times \mathbb{Z}/2\mathbb{Z}$, S_4 and S_5 .

Type	$\pi_1^G(\widetilde{\mathbb{O}})$	Birationally rigid?	L	\mathbb{O}_L	$\pi_1^L(\widetilde{\mathbb{O}}_L) \subseteq \pi_1^L(\mathbb{O}_L)$
$G_2(a_1)$	S_3	Ν	\widetilde{A}_1	0	$1 \subseteq 1$
	A_3	Y	-	-	-

	S_2	Ν	A_1	0	$1 \subseteq 1$
	1	Y	-	-	-
G_2	1	Ν	Т	0	$1 \subseteq 1$

Table 1: Birationally rigid induction data for G_2

Туре	$\pi_1^G(\widetilde{\mathbb{O}})$	Birationally rigid?	L	\mathbb{O}_L	$\pi_1^L(\widetilde{\mathbb{O}}_L)$
\widetilde{A}_2	1	Ν	B_3	0	$1 \subseteq 1$
A_2	S_2	Ν	C_3	0	$1 \subseteq 1$
	1	Y	-	-	-
B_2	S_2	Ν	C_3	$(2, 1^4)$	$1 \subseteq 1$
	1	Y	-	-	-
$C_3(a_1)$	S_2	Ν	B_3	$(2^2, 1^3)$	$1 \subseteq 1$
	1	Y	-	-	-
$F_4(a_3)$	S_4	Ν	$\widetilde{A}_2 + A_1$	0	$1 \subseteq 1$
	A_4	Υ	-	-	-
	D_8	Ν	C_3	$(2^2, 1^2)$	$S_2\subseteqS_2$
	S_3	Ν	$A_2 + \widetilde{A}_1$	0	$1 \subseteq 1$
	C_4	Y	-	-	-
	$S_2\timesS_2$	Ν	B_2	0	$1 \subseteq 1$
	$\operatorname{tw}(S_2\timesS_2)$	Y	-	-	-
	A_3	Y	-	-	-
	S_2	Ν	B_3	$(3, 1^4)$	$1 \subseteq S_2$
	$\operatorname{tw}(S_2)$	Y	-	-	-

	1	Y	_	_	_
C_3	1	Ν	A_2	0	$1 \subseteq 1$
B_3	1	Ν	\widetilde{A}_2	0	$1 \subseteq 1$
$F_4(a_2)$	S_2	Ν	$A_1 + \widetilde{A}_1$	0	$1 \subseteq 1$
	1	Ν	B_2	$(2^2, 1)$	$1 \subseteq 1$
$F_4(a_1)$	S_2	Ν	\widetilde{A}_1	0	$1 \subseteq 1$
	1	Ν	A_1	0	$1 \subseteq 1$
F_4	1	N	T	0	1 ⊂ 1

Table 2: Birationally rigid induction data for ${\cal F}_4$

Type	$\pi_1^G(\widetilde{\mathbb{O}})$	Birationally rigid?	L	\mathbb{O}_L	$\pi_1^L(\widetilde{\mathbb{O}}_L)$
$2A_1$	1	Ν	D_5	0	$1 \subseteq 1$
A_2	S_2	Ν	A_5	0	$1 \subseteq 1$
	1	Y	-	-	-
$A_2 + A_1$	1	Ν	D_5	$(2^2, 1^6)$	$1 \subseteq 1$
$2A_2$	$\mathbb{Z}/3\mathbb{Z}$	Ν	D_4	0	$1 \subseteq 1$
	1	Y	-	-	-
$A_2 + 2A_1$	1	Ν	$A_4 + A_1$	0	$1 \subseteq 1$
A_3	1	Ν	A_4	0	$1 \subseteq 1$
$A_3 + A_1$	1	Ν	D_5	$(3, 2^2, 1^3)$	$1 \subseteq 1$
$D_4(a_1)$	S_3	Ν	$2A_2 + A_1$	0	$1 \subseteq 1$
	A_3	Y	_	-	-
	S_2	Ν	$A_3 + A_1$	0	$1 \subseteq 1$

	1	Ν	D_4	$(2^2, 1^4)$	$1 \subseteq 1$
A_4	1	Ν	A_3	0	$1 \subseteq 1$
D_4	1	Ν	$2A_2$	0	$1 \subseteq 1$
$A_4 + A_1$	1	Ν	$A_2 + 2A_1$	0	$1 \subseteq 1$
$D_5(a_1)$	1	Ν	$A_2 + A_1$	0	$1 \subseteq 1$
A_5	$\mathbb{Z}/3\mathbb{Z}$	Ν	D_4	$(3, 2^2, 1)$	$1 \subseteq 1$
	1	Y	-	-	-
$E_6(a_3)$	$S_2 \times \mathbb{Z}/3\mathbb{Z}$	Ν	$3A_1$	0	$1 \subseteq 1$
	$\mathbb{Z}/3\mathbb{Z}$	Ν	A_2	0	$1 \subseteq 1$
	S_2	Ν	A_5	(3^2)	$1 \subseteq \mathbb{Z}/3\mathbb{Z}$
	1	Y	-	-	-
D_5	1	Ν	$2A_1$	0	$1 \subseteq 1$
$E_6(a_1)$	$\mathbb{Z}/3\mathbb{Z}$	Ν	A_1	0	$1 \subseteq 1$
	1	Ν	$2A_2 + A_1$	$(3) \times (3) \times 0$	$1 \subseteq \mathbb{Z}/3\mathbb{Z}$
E_6	$\mathbb{Z}/3\mathbb{Z}$	N	Т	0	$1 \subseteq 1$
	1	Ν	$2A_2$	$(3) \times (3)$	$1 \subseteq \mathbb{Z}/3\mathbb{Z}$

Table 3: Birationally rigid induction data for (simply connected) ${\cal E}_6$

Туре	$\pi_1^G(\widetilde{\mathbb{O}})$	BR?	L	\mathbb{O}_L	$\pi_1^L(\widetilde{\mathbb{O}}_L)$
$(3A_1)''$	$\mathbb{Z}/2\mathbb{Z}$	N	E_{6}^{8}	0	$1 \subseteq 1$
	1	Y	-	-	_
A_2	S_2	N	D_6	0	$1 \subseteq 1$

⁸In [DE] this rigid induction datum is erroneously listed as being the zero orbit in $2A_2 + A_1$ rather than in E_6 .

	1	Y	-	-	-
$A_2 + A_1$	S_2	Y	-	-	-
	1	N	E_6	A_1	$1 \subseteq 1$
$A_2 + 3A_1$	$\mathbb{Z}/2\mathbb{Z}$	N	A_6	0	$1 \subseteq 1$
	1	Y	-	-	_
2 <i>A</i> ₂	1	N	$D_5 + A_1$	0	$1 \subseteq 1$
A ₃	1	N	D_6	$(2^2, 1^8)$	$1 \subseteq 1$
$(A_3 + A_1)''$	$\mathbb{Z}/2\mathbb{Z}$	Ν	D_5	0	$1 \subseteq 1$
	1	Y	-	-	-
$D_4(a_1)$	S_3	Ν	$A_{5} + A_{1}$	0	$1 \subseteq 1$
	A ₃	Y	-	-	-
	S_2	N	D_6	$(2^4, 1^4)$	$1 \subseteq 1$
	1	Y	-	-	_
$A_3 + 2A_1$	$\mathbb{Z}/2\mathbb{Z}$	N	E_6	$3A_1$	$1 \subseteq 1$
	1	Y	-	-	_
D_4	1	N	$(A_5)''$	0	$1 \subseteq 1$
$D_4(a_1) + A_1$	$S_2 \times \mathbb{Z}/2\mathbb{Z}$	N	$(A_5)'$	0	$1 \subseteq 1$
	S_2	Y	-	-	-
	$\operatorname{tw}(S_2)$	Y	-	-	-
	$\mathbb{Z}/2\mathbb{Z}$	N	E_6	A_2	$1\subseteq S_2$
	1	Y	-	-	-
$A_3 + A_2$	S_2	N	$D_5 + A_1$	$(2^2, 1^6) \times 0$	$1 \subseteq 1$
	1	N	D_6	$(3, 2^2, 1^5)$	$1 \subseteq 1$
$A_3 + A_2 + A_1$	$\mathbb{Z}/2\mathbb{Z}$	N	$A_4 + A_2$	0	$1 \subseteq 1$

	1	Y	-	-	_
A_4	S_2	Ν	$D_4 + A_1$	0	$1 \subseteq 1$
	1	N	D_5	$(2^2, 1^6)$	$1 \subseteq 1$
$D_4 + A_1$	$\mathbb{Z}/2\mathbb{Z}$	N	D_6	$(3, 2^4, 1)$	$\mathbb{Z}/2\mathbb{Z}\subseteq\mathbb{Z}/2\mathbb{Z}$
	1	N	D_6	$(3, 2^4, 1)$	$1 \subseteq \mathbb{Z}/2\mathbb{Z}$
$(A_5)''$	$\mathbb{Z}/2\mathbb{Z}$	N	D_4	0	$1 \subseteq 1$
	1	Ν	$D_5 + A_1$	$(3,1^7)\times(2)$	$1\subseteq \mathbb{Z}/2\mathbb{Z}$
$A_4 + A_1$	S_2	Y	-	-	-
	1	N	$A_4 + A_1$	0	$1 \subseteq 1$
$A_4 + A_2$	1	Ν	$A_3 + A_2 + A_1$	0	$1 \subseteq 1$
$D_5(a_1)$	S_2	N	D_6	$(3^2, 2^2, 1^2)$	$S_2\subseteqS_2$
	1	Ν	A_4	0	$1 \subseteq 1$
$D_5(a_1) + A_1$	$\mathbb{Z}/2\mathbb{Z}$	N	$A_3 + A_2$	0	$1 \subseteq 1$
	1	Y	-	-	-
$(A_5)'$	1	N	$D_5 + A_1$	$(3, 2^2, 1^3) \times 0$	$1 \subseteq 1$
$A_5 + A_1$	$\mathbb{Z}/2\mathbb{Z}$	N	E_6	$2A_2 + A_1$	$1 \subseteq 1$
	1	Y	-	-	-
$E_{6}(a_{3})$	S_2	Ν	$A_3 + 2A_1$	0	$1 \subseteq 1$
	1	N	$D_4 + A_1$	$(2^2, 1^4) \times 0$	$1 \subseteq 1$
$D_6(a_2)$	$\mathbb{Z}/2\mathbb{Z}$	Ν	D_5	$(3, 2^2, 1^3)$	$1 \subseteq 1$
	1	N	$D_5 + A_1$	$(3, 2^2, 1^3) \times (2)$	$1\subseteq \mathbb{Z}/2\mathbb{Z}$
$E_7(a_5)$	$S_3 \times \mathbb{Z}/2\mathbb{Z}$	N	$2A_2 + A_1$	0	$1 \subseteq 1$
	$A_3 \times \mathbb{Z}/2\mathbb{Z}$	N	E_6	$D_4(a_1)$	$A_3 \subseteq S_3$
	S ₃	N	$A_5 + A_1$	$(2^3) \times 0$	$1 \subseteq \mathbb{Z}/2\mathbb{Z}$

	$\operatorname{tw}(S_3)$	Y	-	-	-
	$S_2 imes \mathbb{Z}/2\mathbb{Z}$	N	$(A_3 + A_1)'$	0	$1 \subseteq 1$
	A ₃	Y	-	-	-
	S_2	N	D_6	$(4^2, 2^2)_I^9$	$1 \subseteq \mathbb{Z}/2\mathbb{Z}$
	$\operatorname{tw}(S_2)$	Y	-	-	-
	$\mathbb{Z}/2\mathbb{Z}$	N	D_4	$(2^2, 1^4)$	$1 \subseteq 1$
	1	Y	-	-	-
D_5	1	N	$(A_3 + A_1)''$	0	$1 \subseteq 1$
A_6	1	N	$A_2 + 3A_1$	0	$1 \subseteq 1$
$D_5 + A_1$	$\mathbb{Z}/2\mathbb{Z}$	N	$2A_2$	0	$1 \subseteq 1$
	1	N	$(A_5)''$	(2^3)	$1\subseteq \mathbb{Z}/2\mathbb{Z}$
$D_6(a_1)$	$\mathbb{Z}/2\mathbb{Z}$	N	A_3	0	$1 \subseteq 1$
	1	N	$D_4 + A_1$	$(3,1^5)\times(2)$	$1\subseteq \mathbb{Z}/2\mathbb{Z}$
$E_7(a_4)$	$S_2 \times \mathbb{Z}/2\mathbb{Z}$	N	$A_2 + 2A_1$	0	$1 \subseteq 1$
	S_2	N	$D_5 + A_1$	$(3^3,1)\times(2)$	$1\subseteq \mathbb{Z}/2\mathbb{Z}$
	$\operatorname{tw}(S_2)$	Y	-	-	-
	$\mathbb{Z}/2\mathbb{Z}$	N	$D_4 + A_1$	$(3,2^2,1)\times 0$	$1 \subseteq 1$
	1	Y	-	-	-
$E_6(a_1)$	S_2	N	$4A_1$	0	$1 \subseteq 1$
	1	N	$A_2 + A_1$	0	$1 \subseteq 1$
D_6	$\mathbb{Z}/2\mathbb{Z}$	N	D_4	$(3, 2^2, 1)$	$1 \subseteq 1$
	1	N	$D_4 + A_1$	$(3, 2^2, 1) \times (2)$	$1\subseteq \mathbb{Z}/2\mathbb{Z}$
E_6	1	Ν	$(3A_1)''$	0	$1 \subseteq 1$

 $^{^9\}mathrm{The}$ meaning of the numeral I here is given in Subsection 3.8.

	/				
$E_7(a_3)$	$S_2 \times \mathbb{Z}/2\mathbb{Z}$	N	$(3A_1)'$	0	$1 \subseteq 1$
	S_2	N	$A_3 + 2A_1$	$(2^2) \times 0 \times (2)$	$1 \subseteq \mathbb{Z}/2\mathbb{Z}$
	$\operatorname{tw}(S_2)$	N	$D_5 + A_1$	$(5,3,1^2) \times (2)$	$\operatorname{tw}(S_2)\subseteqS_2\times\mathbb{Z}/2\mathbb{Z}$
	$\mathbb{Z}/2\mathbb{Z}$	N	A_2	0	$1 \subseteq 1$
	1	N	$A_2 + 3A_1$	$0 \times (2) \times (2) \times (2)$	$1\subseteq \mathbb{Z}/2\mathbb{Z}$
$E_7(a_2)$	$\mathbb{Z}/2\mathbb{Z}$	N	$2A_1$	0	$1 \subseteq 1$
	1	N	$(A_3 + A_1)''$	$(2^2) \times (2)$	$1 \subseteq \mathbb{Z}/2\mathbb{Z}$
$E_7(a_1)$	$\mathbb{Z}/2\mathbb{Z}$	N	A_1	0	$1 \subseteq 1$
	1	N	$4A_1$	$0 \times (2) \times (2) \times (2)$	$1\subseteq \mathbb{Z}/2\mathbb{Z}$
E7	$\mathbb{Z}/2\mathbb{Z}$	N	Т	0	$1 \subseteq 1$
	1	N	$(3A_1)''$	$(2) \times (2) \times (2)$	$1 \subseteq \mathbb{Z}/2\mathbb{Z}$

Table 4: Birationally rigid induction data for (simply connected) ${\cal E}_7$

Туре	$\pi_1^G(\widetilde{\mathbb{O}})$	BR?	L	\mathbb{O}_L	$\pi_1^L(\widetilde{\mathbb{O}}_L)$
A_2	S_2	N	E_7	0	$1 \subseteq 1$
	1	Y	-	-	-
A_3	1	Ν	E_7	A_1	$1 \subseteq 1$
$2A_2$	S_2	Ν	D_7	0	$1 \subseteq 1$
	1	Y	-	-	-
$D_4(a_1)$	S_3	Ν	$E_{6} + A_{1}$	0	$1 \subseteq 1$
	A_3	Y	-	-	-
	S_2	Ν	E_7	$2A_1$	$1 \subseteq 1$

	1	Y	-	-	-
D_4	1	Ν	E_6	0	$1 \subseteq 1$
$A_3 + A_2$	S_2	Ν	D_7	$(2^2, 1^{10})$	$1 \subseteq 1$
	1	Ν	E_7	$(3A_1)'$	$1 \subseteq 1$
A_4	S_2	Ν	D_6	0	$1 \subseteq 1$
	1	Ν	E_7	A_2	$1 \subseteq S_2$
$D_4(a_1) + A_2$	S_2	Ν	A_7	0	$1 \subseteq 1$
	1	Υ	-	-	-
$D_4 + A_1$	1	Ν	E_7	$4A_1$	0
$A_4 + A_1$	S_2	Y	-	-	-
	1	Ν	$E_6 + A_1$	$A_1 \times 0$	$1 \subseteq 1$
$D_5(a_1)$	S_2	Ν	E_7	$A_2 + A_1$	$S_2 \subseteq S_2$
	1	Ν	E_6	A_1	$1 \subseteq 1$
$A_4 + 2A_1$	S_2	Y	-	-	-
	1	Ν	D_7	$(2^4, 1^6)$	$1 \subseteq 1$
$A_4 + A_2$	1	Ν	$D_5 + A_2$	0	$1 \subseteq 1$
$A_4 + A_2 + A_1$	1	Ν	$A_{6} + A_{1}$	0	$1 \subseteq 1$
$D_5(a_1) + A_1$	1	Ν	E_7	$A_2 + 2A_1$	$1 \subseteq 1$
A_5	1	Ν	D_7	$(3, 2^2, 1^7)$	$1 \subseteq 1$
$D_4 + A_2$	S_2	Ν	A_6	0	$1 \subseteq 1$
	1	Υ	-	-	-
$E_{6}(a_{3})$	S_2	Ν	$D_5 + A_1$	0	$1 \subseteq 1$
	1	Ν	D_6	$(2^2, 1^8)$	$1 \subseteq 1$
D_5	1	Ν	D_5	0	$1 \subseteq 1$

$E_6(a_3) + A_1$	S_2	Ν	E_7	$A_1 + 2A_2$	$1 \subseteq 1$
	1	Y	-	-	-
$D_{6}(a_{2})$	S_2	Ν	D_7	$(3, 2^4, 1^3)$	$1 \subseteq 1$
	1	Y	-	-	-
$E_{7}(a_{5})$	S_3	Ν	$E_{6} + A_{1}$	$3A_1 \times 0$	$1 \subseteq 1$
	A_3	Υ	-	-	-
	S_2	Ν	E_7	$(A_1 + A_3)'$	$1 \subseteq 1$
	1	Y	-	-	-
$D_5 + A_1$	1	Ν	E_6	$3A_1$	$1 \subseteq 1$
$E_{8}(a_{7})$	${\sf S}_5$	Ν	$A_4 + A_3$	0	$1 \subseteq 1$
	A_5	Y	-	-	-
	S_4	Ν	$D_5 + A_2$	$(2^2, 1^6) \times 0$	$1 \subseteq 1$
	$C_5\rtimesC_4$	Y	-	-	-
	A_4	Y	-	-	-
	$S_2\timesS_3$	Ν	$A_{5} + A_{1}$	0	$1 \subseteq 1$
	D_{10}	Y	-	-	-
	D_8	Ν	D_7	$(3^2, 2^2, 1^4)$	$S_2\subseteqS_2$
	S_3	Ν	$E_{6} + A_{1}$	$A_2 \times 0$	$1 \subseteq S_2$
	$\mathrm{tw}S_3$	Υ	-	-	-
	C_6	Ν	E_7	$D_4(a_1)$	$A_3 \subseteq S_3$
	C_5	Υ	-	-	-
	C_4	Υ	-	-	-
	$S_2\timesS_2$	Ν	D_6	$(2^4, 1^4)$	$1 \subseteq 1$
	$\operatorname{tw}(S_2\timesS_2)$	Y	-	-	-

	A_3	Υ	-	-	-
	S_2	Ν	E_7	$D_4(a_1)$	$1\subseteq S_3$
	$\operatorname{tw}(S_2)$	Υ	-	-	-
	1	Y	_	_	-
A_6	1	Ν	$D_4 + A_2$	0	$1 \subseteq 1$
$D_6(a_1)$	S_2	Ν	A_5	0	$1 \subseteq 1$
	1	Ν	E_6	A_2	$1 \subseteq S_2$
$A_{6} + A_{1}$	1	Ν	$A_4 + A_2 + A_1$	0	$1 \subseteq 1$
$E_{7}(a_{4})$	S_2	Ν	$D_5 + A_1$	$(2^2, 1^6) \times 0$	$1 \subseteq 1$
	1	Ν	D_6	$(3, 2^2, 1^5)$	$1 \subseteq 1$
$D_5 + A_2$	S_2	Ν	$A_4 + A_2$	0	$1 \subseteq 1$
	1	Ν	D_7	$(3^3, 2^2, 1)$	$1 \subseteq 1$
$E_{6}(a_{1})$	S_2	Ν	$D_4 + A_1$	0	$1 \subseteq 1$
	1	Ν	D_5	$(2^2, 1^6)$	$1 \subseteq 1$
D_6	1	Ν	D_6	$(3, 2^4, 1)$	$1 \subseteq 1$
$D_7(a_2)$	S_2	Ν	$2A_3$	0	$1 \subseteq 1$
	1	Ν	$A_4 + 2A_1$	0	$1 \subseteq 1$
E_6	1	Ν	D_4	0	$1 \subseteq 1$
A_7	1	Ν	$D_5 + A_2$	$(3, 2^2, 1^3) \times 0$	$1 \subseteq 1$
$E_6(a_1) + A_1$	S_2	Ν	E_7	$A_4 + A_1$	$S_2\subseteqS_2$
	1	Ν	$A_4 + A_1$	0	$1 \subseteq 1$
$E_{8}(b_{6})$	S_3	Ν	$A_3 + A_2 + A_1$	0	$1 \subseteq 1$
	A_3	Ν	$D_4 + A_2$	$(2^2, 1^4) \times 0$	$1 \subseteq 1$
	S_2	Ν	$E_6 + A_1$	$\left (2A_2 + A_1) \times 0 \right $	$1 \subseteq 1$

	1	Y	-	-	-
$E_{7}(a_{3})$	S_2	Ν	D_6	$(3^2, 2^2, 1^2)$	$S_2 \subseteq S_2$
	1	Ν	A_4	0	$1 \subseteq 1$
$E_6 + A_1$	1	Ν	E_6	$2A_2 + A_1$	$1 \subseteq 1$
$D_7(a_1)$	S_2	Ν	$A_3 + A_2$	0	$1 \subseteq 1$
	1	Ν	$D_5 + A_1$	$(3, 2^2, 1^3) \times 0$	$1 \subseteq 1$
$E_{8}(a_{6})$	S_3	Ν	$2A_2 + 2A_1$	0	$1 \subseteq 1$
	A_3	Ν	$E_6 + A_1$	$D_4(a_1) \times 0$	$A_3 \subseteq S_3$
	S_2	Ν	$A_3 + 2A_1$	0	$1 \subseteq 1$
	1	Ν	$D_4 + A_1$	$(2^2, 1^4) \times 0$	$1 \subseteq 1$
$E_7(a_2)$	1	Ν	D_5	$(3, 2^2, 1^3)$	$1 \subseteq 1$
D_7	1	Ν	$D_4 + A_2$	$(3,2^2,1)\times 0$	$1 \subseteq 1$
$E_8(b_5)$	S_3	Ν	$2A_2 + A_1$	0	$1 \subseteq 1$
	A_3	Ν	E_6	$D_4(a_1)$	$A_3 \subseteq S_3$
	S_2	Ν	$A_3 + A_1$	0	$1 \subseteq 1$
	1	Ν	D_4	$(2^2, 1^4)$	$1 \subseteq 1$
$E_{8}(a_{5})$	S_2	Ν	$A_2 + 3A_1$	0	$1 \subseteq 1$
	1	Ν	$2A_2$	0	$1 \subseteq 1$
$E_{7}(a_{1})$	1	Ν	A_3	0	$1 \subseteq 1$
$E_{8}(b_{4})$	S_2	Ν	$A_2 + 2A_1$	0	$1 \subseteq 1$
	1	Ν	$D_4 + A_1$	$(3,2^2,1)\times 0$	$1 \subseteq 1$
$E_{8}(a_{4})$	S_2	Ν	$4A_1$	0	$1 \subseteq 1$
	1	Ν	$A_2 + A_1$	0	$1 \subseteq 1$
$\overline{E_7}$	1	Ν	D_4	$(3, 2^2, 1)$	$1 \subseteq 1$

$E_8(a_3)$	S_2	N	$3A_1$	0	$1 \subseteq 1$
	1	Ν	A_2	0	$1 \subseteq 1$
$E_8(a_2)$	1	Ν	$2A_1$	0	$1 \subseteq 1$
$E_8(a_1)$	1	Ν	A_1	0	$1 \subseteq 1$
E_8	1	Ν	Т	0	$1 \subseteq 1$

Table 5: Birationally rigid induction data for E_8

References

- [At] Atlas of Lie Groups and Representations, software, 2023, http://www.liegroups.org/. 8, 12, 21
- [Bo] W. Borho, Uber Schichten halbeinfacher Lie-Algebren, Invent. Math. 65, (1981), 283–317. 1
- [BJ] W. Borho and A. Joseph, Sheets and topology of primitive spectra for semisimple Lie algebras, J. Algebra 244 (2001), 76–167. 1
- [Ca] R. Carter, Finite Groups of Lie Type. Conjugacy Classes and Complex Characters, Reprint of the 1985 original, Wiley Classics Lib., John Wiley & Sons, Ltd., Chichester, 1993. 31, 32, 34
- [CM] D.H. Collingwood and W. McGovern, Nilpotent orbits in semisimple Lie algebras, Van Nostrand Reinhold, New York, 1993. 1, 4, 5, 6, 8, 12, 31
- [DE] W. A. de Graaf and A. Elashvili, Induced nilpotent orbits of the simple Lie algebras of exceptional type, Georgian Math. J. 2 (2009), 257–278. 2, 6, 13, 39
- [EI] A. Elashvili, Sheets of the exceptional Lie algebras. (Russian). Studies in Algebra, 171—194, Tbilisi University Press, Tbilisi, 1984. 2
- [FJLS] B. Fu, D. Juteau, P. Levy, and E. Sommers, Generic singularities of nilpotent orbit covers, Adv. Math. 305 (2017), 1–77. 20, 26, 27
- [Fu] B. Fu, On Q-factorial terminalizations of nilpotent orbits, J. Math. Pures App. 93 (2010), 623–635. 10, 17, 19, 27
- [GTW] S. M. Goodwin, L. Topley, M. Westaway, On induced completely prime primitive ideals in enveloping algebras of classical Lie algebras, arXiv:2311.10603, 2023. 3
- [Ho] R. Howlett, Normalizers of parabolic subgroups of reflection groups, J. London Math. Soc. 21 (1980), 62–80. 20
- [Jo] A. Joseph, On the associated variety of a primitive ideal, J. Algebra 93 (1985), 509–523. 5
- [L1] I. Losev, 1-dimensional representations and parabolic induction for W-algebras, Adv. Math. 226 (2011), 4841– 4883. 1, 2
- [L2] I. Losev, Deformations of symplectic singularities and orbit method for semisimple Lie algebras, Selecta Math. 28 (2022), Art. no. 30, 52pp. 1, 2, 6, 7, 8
- [L3] I. Losev, Harish-Chandra bimodules over quantized symplectic singularities, Transform. Groups 26 (2021), 565–600. 8
- [L4] I. Losev, Automorphisms and isomorphisms, continued. https://gauss.math.yale.edu/~il282/Math720_ Lec24.pdf, Nov. 2022. 9
- [LMM] I. Losev, D. Matvieievskyi and L. Mason-Brown, Unipotent ideals and Harish-Chandra bimodules, arXiv:2108.03453, 2021. 1, 2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 25, 26, 27
- [LS] G. Lusztig, N. Spaltenstein, Induced nilpotent classes, J. London Math. Soc. 19 (1979), 41–52. 1
- [MM] L. Mason-Brown, D. Matvieievskyi Unipotent ideals for spin and exceptional groups, J. Alg. 615 (2023), 358–454. 2, 8, 9, 10, 11, 20, 26, 27
- [MMY] L. Mason-Brown, D. Matvieievskyi, S. Yu, Unipotent representations of complex groups and extended Sommers duality, arXiv:2309.13853, 2023. 2, 3, 7, 8, 9
- [N1] Y. Namikawa, Poisson deformations of affine symplectic varieties, II, Kyoto J. Math. 50 (2010), 727–752. 8

- [N2] Y. Namikawa, Birational geometry for the covering of a nilpotent orbit closure, Selecta Math. (N.S.) 28 (2022), Paper No. 75, 59pp. 3
- [P1] A. Premet, Irreducible representations of Lie algebras of reductive groups and the Kac-Weisfeiler conjecture, Invent. Math. 121 (1995), 79–117. 1
- [P2] A. Premet, Commutative quotients of finite W-algebras, Adv. Math. 225 (2010), 269–306. 1
- [P3] A. Premet, Multiplicity-free primitive ideals associated with rigid nilpotent orbits, Transform. Groups 19 (2014), 569–641. 1
- [PT1] A. Premet and L. Topley, Derived subalgebras of centralisers and finite W-algebras, Compos. Math. 150 (2014), 1485–1548. 3
- [PT2] A. Premet, L. Topley, Modular representations of Lie algebras of reductive groups and Humphreys' conjecture, Adv. Math. 392 (2021), Paper No.108024, 40 pp. 1

Email address: M.P.Westaway@bham.ac.uk

School of Mathematics, University of Birmingham, Birmingham, B15 2TT, UK