Strong Convexity of Sets in Riemannian Manifolds

Damien Scieur^{*,‡}, David Martínez-Rubio^{*,§}, Thomas Kerdreux ^{*,§}, Alexandre d'Aspremont^{\$}, Sebastian Pokutta[§].

Abstract

Curvature properties of convex objects, such as strong convexity, are important in designing and analyzing convex optimization algorithms in the Hilbertian or Riemannian settings. In the case of the Hilbertian setting, strongly convex sets are well studied. Herein, we propose various definitions of strong convexity for uniquely geodesic sets in a Riemannian manifold. We study their relationship, propose tools to determine the geodesic strongly convex nature of sets, and analyze the convergence of optimization algorithms over those sets. In particular, we demonstrate that the Riemannian Frank-Wolfe algorithm enjoys a global linear convergence rate when the Riemannian scaling inequalities hold.

Contents

	1	Introduction	2
	2	Preliminaries and Notations	3
	3	Strong Convexity of Sets in Riemannian Manifolds	5
	4	Relations between the Definitions of Strong Convexity	7
	5	Sublevel Sets of Geodesically Strongly Convex Functions	10
	6	Double Geodesic Strong Convexity and Approximate Riemannian Scaling Inequality	12
	7	Frank-Wolfe on Geodesically Strongly Convex Sets	14
	8	Examples of Riemannian Strongly Convex Sets: Balls with Restricted Radius	19
	9	Example of a Simple Linear Minimization Oracle	21
	10	Conclusion	22
	A	Technical Proposition and Lemmas	28
	в	MATLAB code for the experiments	31
*Ec	qual	Contribution (randomized order).	

[‡]Samsung SAIL Montreal, Canada.

[°]CNRS & D.I. École Normale Supérieure, Paris, France.

[§]Zuse Institute Berlin & Technische Universität, Berlin, Germany.

1 Introduction

Strong convexity is a fundamental property characterizing convex objects such as normed spaces, functions, or sets. Algorithms can leverage strong (or uniform) convexity structures in optimization and learning problems to accelerate convergence, enhance generalization bounds, and strengthen concentration properties.

For instance, uniform convexity (that interpolates between plain and strong convexity) has a direct effect on the concentration inequalities of vector-valued martingales when measured via this space norm [52, 53]. Also, assuming strong convexity for objective functions has become standard for studying optimization algorithm performance in Hilbert spaces and proving generalization bounds. Similarly, the strong convexity of sets plays a role in optimization algorithms [23, 26, 47, 38, 39] or learning algorithms, e.g., to characterize action sets in online learning [34, 47, 37, 22, 9, 10] or bandit algorithms [16, 40]. These notions have a well-understood and well-exploited generalization in normed space settings [28].

However, many machine-learning problems are being formulated and solved beyond normed spaces, e.g., metric spaces and Riemannian manifolds. For instance, Sinkhorn divergences [21, 25] are regularized formulations of optimal transport distances [60]. Those methods approximate the optimal transport distance, breaking some computational bottlenecks. This has resulted in modeling many problems using these geometries over probability spaces, and hence, metric spaces. Similarly, some progress has recently been made in optimization algorithms on Riemannian manifolds [57, 3, 32, 33]. The definition of these optimization problems on these structures produces local-global properties when identifying geodesically convex problems that may be non-convex.

Unfortunately, the notion of strong convexity in these more complex frameworks is more difficult to formalize. Several definitions of strong convexity for metric spaces have been developed, such as those for the most notable *non-positively curved*, also known as CAT(0) spaces [2]. Those spaces require the square of the distance function to be geodesically strongly convex. Similar results, such as those in the Hilbertian case, have been obtained with concentration results [29, 1, 45] or in online learning [49, 50]. Nevertheless, the situation is not as well-behaved as in normed spaces, as several non-equivalent notions emerge when describing the problem structures.

Motivation. There are multiple consequences of analyzing strongly convex sets defined in manifolds. In the Euclidean case, such an analysis made the identification of the strong convexity of sets easier: sublevel sets of smooth, strong convex functions are strongly convex sets. In addition, the Frank–Wolfe (FW) algorithm [14] converges faster over strongly convex sets. Generalizing the notion of strong convexity for functions in metric spaces is relatively straightforward (Definition 2.7). However, to our knowledge, this notion has not been studied in the case of sets. Therefore, we propose to fill this gap in the context of a Riemannian manifold.

Contributions. We introduce novel structural properties of sets defined in Riemannian manifolds, provide examples, and demonstrate improved algorithmic performance when minimizing functions over those sets.

- 1. In Section 3, we introduce different definitions of strong convexity of a set in Riemannian manifolds. Each definition relies on two perspectives of uniquely geodesic subsets, either seen as a geodesic metric space or a smooth manifold.
- 2. We establish some relationships between the aforementioned various definitions of strongly convex sets in Section 4 and introduce the notion of approximate scaling inequality and its link with what we call *double geodesic strong convexity* property (Definition 3.4) in Section 6.
- 3. In Section 5, we prove that sublevel sets of geodesically smooth, strongly convex functions are strongly convex in Riemannian manifolds under mild conditions. Similar results have

been obtained in the case of Hilbert spaces [36, 26]. This perspective is valuable for identifying strongly convex sets in Riemannian manifolds.

- 4. We also provide examples for the strongest of our notions of Riemannian strong geodesic convexity. Hence, those examples also satisfy all other other notions, cf. Section 8.
- 5. In Section 7, we derive a global linear convergence bound for the Riemannian FW algorithm when the constraint set satisfies the *Riemannian scaling inequality* (Definition 3.5).

2 Preliminaries and Notations

Our primary goal is defining several notions of strong convexity for sets in metric spaces that are well-suited for analyzing and designing optimization algorithms. In this study, we first focus on the case of complete and connected *n*-dimensional Riemannian manifolds. This setting combines a simple manifold structure, e.g., *n*-dimensional tangent spaces with scalar products, with a *geodesically metric space* structure. We now recall some useful concepts.

2.1 Geodesic Metric Spaces

Metric space A metric space is a pair $(\mathcal{M}, d_{\mathcal{M}})$, where \mathcal{M} is a set, and $d_{\mathcal{M}} : \mathcal{M} \times \mathcal{M} \to \mathbb{R}^+$ is a distance function (the metric), i.e., it satisfies positivity, symmetry, and the triangle inequality.

Geodesics A geodesic $\gamma(t) : [0,1] \to \mathcal{M}$ between two points $x, y \in \mathcal{M}$ is a smooth curve such that $\gamma(0) = x, \gamma(1) = y$, and $\gamma''(t) = 0$. A metric set is a *geodesic metric space* if, for all $x, y \in \mathcal{M}$, there exists a geodesic that connects x and y. When this geodesic is unique, the set is considered uniquely geodesic.

Cartan–Hadamard spaces In particular, *Cartan–Hadamard spaces* are geodesic metric spaces generalizing Hilbert spaces which are well-suited for convex optimization purposes [35, 15, 5, 6]. By definition, a Cartan–Hadamard space \mathcal{M} (a.k.a. CAT(0) space or space of non-positive curvature) is a geodesically metric space that is non-positively curved. Non-positively curved spaces have the particularity that, for all geodesics γ and for all elements $x \in \mathcal{M}$, the metric $d_{\mathcal{M}}$ satisfies (e.g., [50, Corollary 2.5.])

$$d_{\mathcal{M}}(x,\gamma(t))^{2} \leq (1-t)d_{\mathcal{M}}(x,\gamma(0))^{2} + td_{\mathcal{M}}(x,\gamma(1))^{2} - t(1-t)d_{\mathcal{M}}(\gamma(0),\gamma(1))^{2}.$$
(non-positive curvature)

This non-positive curvature indicates that the function $f(\cdot) = d_{\mathcal{M}}(\cdot, x)^2$ is geodesically strongly convex (see Definition 2.7) for all reference point $x \in \mathcal{M}$.

2.2 Riemannian Manifolds

In the case of Banach spaces, wherein the strong convexity has been crucial to algorithm analysis, significantly fewer tools have been developed for directly characterizing sets that appear in geodesic metric spaces and, therefore, in Riemannian manifolds. Those tools are needed to analyze online learning algorithms or some constrained optimization algorithms. We recall some key concepts below.

Throughout this paper, we consider an *n*-dimensional Riemannian manifold \mathcal{M} , such as the Stiefel manifold $\operatorname{St}(n,p)$ for p < n; the group of rotations $\operatorname{SO}(n)$; hyperbolic spaces; spheres; or the manifold of symmetric positive definite matrices.

Connected manifolds A manifold \mathcal{M} is *connected* if, for all $x, y \in \mathcal{M}$, there exists a continuous path joining those points. An *n*-dimensional manifold is a topological space that is locally homeomorphic to the *n*-dimensional Euclidean space.

Tangent space For a point $x \in \mathcal{M}$, a tangent vector is the tangent of a parameterized curve passing through x, and all the tangent vectors at x form the tangent space $T_x \mathcal{M}$.

Metric tensor and inner product A Riemannian manifold (\mathcal{M}, g_x) is such that the Riemannian metric g is a C^{∞} metric tensor, where for all point $x \in \mathcal{M}$, g_x defines a positive definite inner product in the tangent space $T_x\mathcal{M}$, and $x \mapsto g_x$ is C^{∞} . For $u, v \in \mathcal{M}$, we write $g_x(u, v) := \langle u; v \rangle_x$, and $||u||_x := \langle u; u \rangle_x$.

Complete Riemannian manifold and Riemannian metric A connected Riemannian manifold is also a metric space. Furthermore, the length of a continuous piecewise smooth path $\gamma : [0,1] \to \mathcal{M}$ is defined as $L(\gamma) := \int_0^1 ||\gamma'(t)||_{\gamma(t)} dt$. We denote $\mathcal{P}_{x,y}$ as the set of continuous piecewise smooth paths joining x and y. We have the *Riemannian metric* $d_{\mathcal{M}}(x,y) := \inf_{\gamma \in \mathcal{P}_{x,y}} L(\gamma)$ as a distance [12, Theorem 10.2] and $(\mathcal{M}, d_{\mathcal{M}})$ is a metric space. We refer to *complete manifold* as a manifold that is complete as a metric space. Moreover, [12, Theorem 10.8] states that any connected and complete Riemannian metric is attained at a geodesic (w.r.t. $d_{\mathcal{M}}$) called the *minimizing geodesic*.

2.3 Sets in Riemannian Manifolds

Importance of sets when optimizing on Manifolds. When \mathcal{M} is a connected compact Riemannian manifold, and f is a geodesically convex function on \mathcal{M} (Definition 2.7), then f is constant [12, Corollary 11.10]. This result motivates the optimization of geodesically convex functions over subsets \mathcal{C} of these manifolds.

Assumptions for Sets in Riemannian Manifolds In this paper, we will make the assumption that we are working on a subset $C \subset M$ that is *uniquely geodesically convex*, defined as the set C being both convex and uniquely geodesic.

Definition 2.1 (Geodesic Convex Closed Subset of a Manifold). We say that a closed subset C of a manifold M is convex if for any two points $x, y \in C$, there exists a geodesic from x to y that is distance minimizing and is contained in C.

Definition 2.2 (Uniquely Geodesic Subset of a Manifold). We say that a subset C of a manifold \mathcal{M} is uniquely geodesic if for every two points $x, y \in C$, there is only one geodesic between x and y that is contained in C.

Assumption 2.3. The set $C \subseteq M$ is compact, convex, and uniquely geodesic.

For instance, open hemispheres and Cartan–Hadamard manifolds, that is, complete simply connected manifolds of non-positive sectional curvature everywhere, are uniquely geodesic. Hence, their compact counterparts are geodesically convex subsets. However, a closed hemisphere is not uniquely geodesic.

The exponential map parameterizes the manifold by mapping vectors in the tangent spaces to \mathcal{M} . For $x \in \mathcal{M}$, it is in general a local diffeomorphism; however, when \mathcal{M} is complete [30], it is defined on the whole tangent space as follows.

Definition 2.4 (Exponential Map). Let \mathcal{M} be a Riemannian manifold. For all $x \in \mathcal{M}$ and $v \in T_x \mathcal{M}$, let the geodesic $\gamma : [0,1] \to \mathcal{M}$ satisfying $\gamma(0) = x$ with $\gamma'(0) = v$. The exponential map at x, $\operatorname{Exp}_x : T_x \mathcal{M} \to \mathcal{M}$, is defined as $\operatorname{Exp}_x(v) := \gamma(1)$.

Remark 1 (Bijective Exponential Map and Logarithmic Map in the Right Domain). For uniquely geodesically convex sets C and $x \in C$, define the set

 $\mathcal{C}_x \stackrel{\text{\tiny def}}{=} \{ v \in T_x \mathcal{M} : t \mapsto \operatorname{Exp}_x(tv), \text{ for } t \in [0,1], \text{ is minimizing and } \operatorname{Exp}_x(v) \in \mathcal{C} \},\$

then, the exponential map from x restricted to this set is well defined and bijective in C, that is, $\operatorname{Exp}_x(y): \mathcal{C}_x \to \mathcal{C}$ is a bijective function. In this work we will always refer to the inverse of the exponential map, or logarithmic map, as the inverse of this restriction: $\operatorname{Exp}_x^{-1}: \mathcal{C} \to \mathcal{C}_x$. Note that generally in the literature the logarithmic map is not defined using this restriction.

The exponential map links the manifold and its tangent spaces, but vectors belonging to different tangent spaces are not directly comparable. Using the Levi-Civita connection, we can define a parallel transport operator between two points $x, y \in \mathcal{M}$ in the manifold [4].

Definition 2.5 (Parallel Transport Operator). For all $x, y \in \mathcal{M}$, the parallel transport map $\Gamma_x^y: T_x\mathcal{M} \to T_y\mathcal{M}$ combines vectors from different tangent spaces by transporting them along geodesics and such that $\langle \Gamma_x^y u; \Gamma_x^y v \rangle_y = \langle u; v \rangle_x$ for all $(u, v) \in T_x\mathcal{M}$.

2.4 Functions over Manifolds

We first define the Riemannian gradient of f at x as the unique element $\nabla f(x) \in T_x \mathcal{M}$, such that the directional derivative $Df(x)[v] = \langle v; \nabla f(x) \rangle_x$ for all $v \in T_x \mathcal{M}$.

We now recall the notion of relative geodesic (strong) convexity and smoothness for a function $f : \mathcal{M} \to \mathbb{R}$ w.r.t. a distance function $d : \mathcal{M} \times \mathcal{M} \to \mathbb{R}^+$ (see, for instance, [63]).

Definition 2.6 (Distance function). A function $d : \mathcal{M} \times \mathcal{M} \to \mathbb{R}^+$ is called a distance function if it satisfies positivity, symmetry, and the triangle inequality.

Remark 2. The distance d could differ from the distance function $d_{\mathcal{M}}$ associated with the manifold \mathcal{M} .

Definition 2.7 (Geodesic (Strong) Convexity). Let C be a uniquely geodesic set. A function $f : C \subseteq \mathcal{M} \to \mathbb{R}$ is geodesically μ -strongly convex in C (resp. convex if $\mu = 0$) w.r.t. d if, for all $x, y \in C$, we have

$$\forall t \in [0,1], \ f(\gamma(t)) \le (1-t)f(x) + tf(y) - \frac{\mu}{2}t(1-t)d^2(x,y), \ \mu \ge 0.$$
(1)

Definition 2.8 (Geodesic Smoothness). Let C be a uniquely geodesic set. A function $f : C \subseteq \mathcal{M} \to \mathbb{R}$ is geodesically L-smooth in C w.r.t. d if, for all $x, y \in C$, we have

$$|f(y) - f(x) - \langle \nabla f(x); \operatorname{Exp}_{x}^{-1}(y) \rangle_{x}| \le \frac{L}{2}d^{2}(x, y), \ L \ge 0.$$

Note that when the function is also geodesic (strongly) convex, the smoothness condition can be simplified by removing the absolute value. Indeed, (strong) convexity implies that the argument of the absolute value is nonnegative.

Finally, we state the following assumption on the relation between the distance function used to define smoothness and strong convexity and the distance function associated with the manifold \mathcal{M} . This assumption allows us to use distances whose value is within some constant factors from the Riemannian distance $\|\operatorname{Exp}_x^{-1}(y)\|$, which corresponds to the definition below with $\ell_{\mathcal{M}} = L_{\mathcal{M}} = 1$.

Assumption 2.9 (Distances Equivalence). Let \mathcal{M} be a Riemannian manifold and let $d : \mathcal{M} \times \mathcal{M} \to \mathbb{R}$ be a distance function, possibly different from the Riemannian distance. There exists $0 < l_{\mathcal{M}} \leq L_{\mathcal{M}}$ such that for all pair $(x, y) \in \mathcal{M}^2$,

$$\ell_{\mathcal{M}} \| \operatorname{Exp}_{x}^{-1}(y) \|_{x} \le d(x, y) \le L_{\mathcal{M}} \| \operatorname{Exp}_{x}^{-1}(y) \|_{x}.$$
(2)

3 Strong Convexity of Sets in Riemannian Manifolds

This section provides several variants of the definition of a strongly convex set in Riemannian manifolds. Some variants are instrumental in analyzing algorithms or proving a set's strongly convex nature.

3.1 Strong Convexity in Hilbert Spaces

Many equivalent characterizations of a set's strong convexity exist in Hilbert spaces [28]. We recall two characterizations in Proposition 3.1. Intuitively, a set is called strongly convex if, for all straight lines in the set and all points p in the line, there exists a ball around p of a certain radius that is contained in the set.

Proposition 3.1 (Equivalent Notions of Strong Convexity of Sets in Hilbert Spaces [28]). Consider a compact convex set $\mathcal{C} \subset \mathbb{R}^n$, $\alpha > 0$, and a norm $||x|| = \sqrt{\langle x, x \rangle}$.

We say that the set C is α -strongly convex w.r.t. $\|\cdot\|$ if and only if it satisfies the following equivalent assertions.

(a) For all $(x, y) \in \mathcal{C}$, $t \in [0, 1]$, and $z \in \mathbb{R}^n$ s.t. ||z|| = 1, we have

$$(1-t)x + ty + \alpha(1-t)t||x-y||^2 z \in \mathcal{C}.$$
(3)

(b) For all $(x, v) \in \mathcal{C} \times \partial \mathcal{C}$ and $w \in N_{\mathcal{C}}(v)$ (the normal cone of \mathcal{C} at x), we have

$$\langle w; v - x \rangle \ge \alpha \|w\| \|v - x\|^2.$$
 (scaling inequality)

Going back to the original definitions in the literature, one can find these properties where α takes the value 1/2R, where R > 0 is the radius of some balls that are used for an alternative equivalent definition of strong convexity of sets [59, 58, 28]. We use α in this work since it is simpler for our purposes. We also note that (scaling inequality) is equivalent to the following condition [38, Lemma 2.1]:

$$\langle w; v - x \rangle \ge \alpha \|w\| \|v - x\|^2 \text{ where } v \in \operatorname*{argmax}_{z \in \mathcal{C}} \langle w; z \rangle.$$
 (4)

3.2 Strong Convexity of Sets in Riemannian Manifolds

In this section, we propose several definitions that extend (3) for Riemannian manifolds, which all collapse to the known strong convexity notion when \mathcal{M} is the Euclidean space. We study their relationships and provide examples in Section 8 for the strongest conditions; therefore, the examples apply to all of our definitions.

3.2.1 Geodesic Strong Convexity

Definition 3.2 adapts (3) but relies on the geodesic metric structure of \mathcal{M} only, without considering the Riemannian metric structure (i.e., the definition does not even use the tangent space). We refer to it as the *geodesic strong convexity*.

Definition 3.2 (Geodesic Strong Convexity). Let \mathcal{M} be a Riemannian manifold, and let $\mathcal{C} \subseteq \mathcal{M}$ be a uniquely geodesic set. The set \mathcal{C} is geodesically α -strongly convex w.r.t. the distance function $d: \mathcal{M} \times \mathcal{M} \to \mathbb{R}^+$ if, for every geodesic γ joining $x, y \in \mathcal{C}$ and every $t \in [0, 1]$, we have that the following ball is in \mathcal{C} :

$$\{z \in \mathcal{M} \mid d(\gamma(t), z) \le \alpha t(1-t)d^2(x, y)\} \subseteq \mathcal{C}.$$
(5)

3.2.2 Riemannian Strong Convexity

The following definition leverages the Riemannian structure of \mathcal{M} via an assumption on the exponential map. The definition states that the inverse image of the set $\mathcal{C} \subset \mathcal{M}$ by the inverse exponential map at each $x \in \mathcal{M}$ must be strongly convex in $T_x \mathcal{M}$ for all x in the Euclidean sense. Recall that for a uniquely geodesic set \mathcal{C} , the inverse exponential map is always well defined for any two points in \mathcal{C} , cf. Remark 1.

Definition 3.3 (Riemannian Strong Convexity). Let \mathcal{M} be a Riemannian manifold, and $\mathcal{C} \subseteq \mathcal{M}$ be a uniquely geodesic set. Then, \mathcal{C} is a Riemannian α -strongly convex set if, for all $x \in \mathcal{C}$, the set

$$\operatorname{Exp}_x^{-1}(\mathcal{C}) := \{ y \in T_x \mathcal{M} : y = \operatorname{Exp}_x^{-1}(z), \ z \in \mathcal{C} \}$$

is α -strongly convex w.r.t. $\|\cdot\|_x$ in the Euclidean sense (3).

3.2.3 Double Geodesic Strong Convexity

In Definition 3.4, we now leverage the Riemannian structure through the exponential map to provide another notion of strong convexity of a set, analogous to the Euclidean formulation in (3). For $t \in [0, 1]$, we make the parallel between the term tx + (1 - t)y in (3) and $\gamma(t)$, the geodesic γ joining y, and x in \mathcal{M} . Then, (3) in \mathcal{M} ensures that, for all $z \in T_{\gamma(t)}\mathcal{M}$ with unit norm ||z|| = 1, we have $\operatorname{Exp}_{\gamma(t)} (\alpha t(1 - t)d^2(x, y)z) \in \mathcal{C}$.

Definition 3.4 (Double Geodesic Strong Convexity). Let \mathcal{M} be a Riemannian manifold equipped with a distance function $d(\cdot, \cdot)$, and let $\mathcal{C} \subseteq \mathcal{M}$ be a set that is uniquely geodesic. The set \mathcal{C} is a double geodesically α -strongly convex set w.r.t. $d(\cdot, \cdot)$ if, for every geodesic γ joining $x, y \in \mathcal{C}$,

$$\forall t \in [0,1], \ \forall z \in T_{\gamma(t)}\mathcal{M} : \|z\|_{\gamma(t)} \le \alpha t(1-t)d^2(x,y) \quad \Rightarrow \quad \operatorname{Exp}_{\gamma(t)}(z) \ exists \ and \ is \ in \ \mathcal{C}. \ (6)$$

The double geodesic strong convexity can also be rewritten in terms of the exponential map if $\operatorname{Exp}_{\gamma(t)}(\alpha t(1-t)d^2(x,y)z) \in \mathcal{C}$ for all z of unit norm in $T_{\gamma(t)}\mathcal{M}$. In this manner, it mirrors the algebraic expression of the Euclidean definition we provided in (3) but within the Riemannian setup. We denote it as the *double geodesic strong convexity* also because the point $\operatorname{Exp}_{\gamma(t)}(\alpha t(1-t)d^2(x,y)z)$ is built via two geodesics, one between x and y, and another starting at $\gamma(t)$. In Section 6, we define a characterization of the double geodesic convexity via the classical *double exponential map* (Definition 6.1).

3.2.4 Riemannian Scaling Inequality

In Euclidean space, (scaling inequality) is an equivalent definition of the strong convexity of the sets via Proposition 3.1, which helps in establishing convergence proofs of various algorithms. In Definition 3.5, we propose the notion of the Riemannian scaling inequality, which is the the Riemannian counterpart of (4).

Definition 3.5 (Riemannian Scaling Inequality). Let \mathcal{M} be a Riemannian manifold, and let $\mathcal{C} \subset \mathcal{M}$ be compact and uniquely geodesically convex. The elements in the set \mathcal{C} then satisfy the Riemannian scaling inequality if, for some $\alpha > 0$, for all $x \in \mathcal{C}$, $w \in T_x \mathcal{M}$, and $v \in \underset{z \in \mathcal{C}}{\operatorname{argmax}} \langle w; \operatorname{Exp}_x^{-1}(z) \rangle_x$,

 $\langle w; \operatorname{Exp}_{x}^{-1}(v) \rangle_{x} \ge \alpha \|w\|_{x} \|\operatorname{Exp}_{x}^{-1}(v)\|_{x}^{2}.$ (Riemannian scaling inequality)

4 Relations between the Definitions of Strong Convexity

This section establishes some implications and equivalences between strong convexity notions and scaling inequality for manifolds. We summarize the links between these notions here.

 $\begin{array}{cccc} \text{Riemannian} & \stackrel{\text{Prop. 4.2,}}{\underset{\text{Strong Convexity}}{\text{Convexity}}} & \stackrel{\text{Prop. 4.3}}{\underset{\text{Convexity}}{\overset{\text{Convex}}{\overset{\text{Convexity}}{\overset{\text{Convexity}$

The scaling inequality and global strong convexity of a set are equivalent notions (Proposition 3.1) in Hilbert spaces, but this won't be the case when working with Riemannian manifolds. Instead, Proposition 4.1 states that the Riemannian strong convexity implies a Riemannian scaling inequality, the latter being valuable for analyzing algorithms (Section 7).

Proposition 4.1 (Riemannian Strong Convexity implies Riemannian Scaling Inequality). Let \mathcal{M} be a Riemannian manifold, and $\mathcal{C} \subset \mathcal{M}$ be compact and uniquely geodesically convex. Let us assume that \mathcal{C} is a Riemannian α -strongly convex set (Definition 3.3). The elements in the set \mathcal{C} then also satisfy the Riemannian scaling inequality (Definition 3.5).

Proof. As the set is strongly convex in the Euclidean sense, we obtain from (scaling inequality) that, for all $x \in C$ and for all $w \in T_x \mathcal{M}$, we have

$$\langle w; u \rangle_x \ge \alpha \|w\|_x \|u - x\|_x^2, \quad u \in \operatorname*{argmax}_{z \in \operatorname{Exp}_x^{-1}(\mathcal{C})} \langle w; z \rangle_x.$$

As the exponential map is bijective over the set \mathcal{C} ,

$$\underset{z \in \operatorname{Exp}_x^{-1}(\mathcal{C})}{\operatorname{argmax}} \langle w; z \rangle_x = \operatorname{Exp}_x^{-1} \left(\underset{z \in \mathcal{C}}{\operatorname{argmax}} \langle w; \operatorname{Exp}_x^{-1}(z) \rangle_x \right).$$

Therefore, using $v = \text{Exp}_{x}(u)$ gives us

$$\langle w; \operatorname{Exp}_x^{-1}(v) \rangle_x \ge \alpha \|w\|_x \|\operatorname{Exp}_x^{-1}(v)\|_x^2, \quad v \in \operatorname{argmax}_{z \in \mathcal{C}} \langle w; \operatorname{Exp}_x^{-1}(z) \rangle_x.$$

Proposition 4.2 (Riemannian Strong Convexity implies Geodesic Strong Convexity). Let \mathcal{M} be a Riemannian manifold, and let $\mathcal{C} \subset \mathcal{M}$ be a set that is uniquely geodesically convex. Consider the distance $d(\cdot, \cdot)$ satisfying Assumption 2.9. Let us assume that the set \mathcal{C} is a Riemannian α -strongly convex set (Definition 3.3). The set \mathcal{C} is then a geodesic $(\alpha \ell_{\mathcal{M}}/L_{\mathcal{M}}^2)$ -strongly convex set (Definition 3.2).

Before proving the theorem, we introduce the important notion of geodesic triangle. For the three points $p, q, r \in \mathcal{M}$, the set Δpqr of the three minimizing geodesics joining these three points is a geodesic triangle. A comparison triangle $\Delta \bar{p}\bar{q}\bar{r}$ is then a triangle with the same side length as Δpqr in a metric space with a constant sectional curvature. Comparison theorems are then used to compare the angle between these triangles according to a lower (Toponogov's theorem) or an upper bound (Rauch's theorem) on the sectional curvature of the geodesic metric space \mathcal{M} . We refer to [18, 46, 17] for a detailed treatment of such comparison theorems and to [63, 64] for their use in optimization contexts. It should be noted that comparison theorems do not only compare triangles, but also hinges, and result in angle or length comparisons [46, Theorem 2.2. B].

Proof. As the set C is a Riemannian strongly convex set (Definition 3.3), by using the definition of the strong convexity of sets in Hilbert spaces (3), we obtain

$$\forall x \in \mathcal{M}, \ \forall p, q \in \operatorname{Exp}_{x}^{-1}(\mathcal{C}), \ \forall t \in [0,1],$$

if $z \in T_{x}\mathcal{M} : ||z - (tp + (1-t)q)|| \le \alpha t(1-t)||p - q||^{2}, \text{ then } z \in \operatorname{Exp}_{x}^{-1}(\mathcal{C}),$ (8)

for some parameter $\alpha > 0$. We now consider arbitrary points $x, y \in \mathcal{M}$ and $\tilde{z} \in \mathcal{M}$ s.t. $d(\gamma(t), \tilde{z}) \leq \tilde{\alpha}t(1-t)d^2(x, y)$, where $\gamma(t) : [0, 1] \to \mathcal{M}$ is the geodesic between x and y and $\tilde{\alpha} \stackrel{\text{def}}{=} \alpha \ell_{\mathcal{M}} L_{\mathcal{M}}^{-2} > 0$. Due to Assumption 2.9 we have

$$\ell_{\mathcal{M}} \| \operatorname{Exp}_{\gamma(t)}^{-1}(\tilde{z}) \| \le L_{\mathcal{M}}^2 \tilde{\alpha} t (1-t) \| \operatorname{Exp}_x^{-1}(y) \|^2.$$
(9)

Now, [44, Corollary 24] using the Riemannian cosine law inequality for our Cartan-Hadamard manifold in the geodesic triangle with vertices x, \tilde{z} and $\gamma(t)$, and the corresponding triangle in $T_x \mathcal{M}$ via $\operatorname{Exp}_x^{-1}(\cdot)$, we have

$$\begin{split} & 2\langle \operatorname{Exp}_x^{-1}(\tilde{z}), \operatorname{Exp}_x^{-1}(\gamma(t))\rangle \geq \|\operatorname{Exp}_x^{-1}(\tilde{z})\|^2 + \|\operatorname{Exp}_x^{-1}(\gamma(t))\|^2 - \|\operatorname{Exp}_{\gamma(t)}^{-1}(\tilde{z})\|^2 \\ & 2\langle \operatorname{Exp}_x^{-1}(\tilde{z}), \operatorname{Exp}_x^{-1}(\gamma(t))\rangle = \|\operatorname{Exp}_x^{-1}(\tilde{z})\|^2 + \|\operatorname{Exp}_x^{-1}(\gamma(t))\|^2 - \|\operatorname{Exp}_x^{-1}(\tilde{z}) - \operatorname{Exp}_x^{-1}(\gamma(t))\|^2, \end{split}$$

which implies

$$\|\operatorname{Exp}_{x}^{-1}(\tilde{z}) - \operatorname{Exp}_{x}^{-1}(\gamma(t))\|_{x} \le \|\operatorname{Exp}_{\gamma(t)}^{-1}(\tilde{z})\|_{\gamma(t)}.$$
(10)

Now, let $p = \exp_x^{-1}(y)$, $q = \exp_x^{-1}(x) = 0$, and $z = \exp_x^{-1}(\tilde{z})$, and note that $\exp_x^{-1}(\gamma(t)) = t \exp_x^{-1}(y)$. Hence, combining (9) with (10) and using our new notation, we obtain

$$\ell_{\mathcal{M}} \| z - (tp + (1-t)q) \|_{x} \le L_{\mathcal{M}}^{2} \tilde{\alpha} t (1-t) \| p - q \|_{x}^{2},$$

and after using the value of $\tilde{\alpha} = \alpha \ell_{\mathcal{M}} L_{\mathcal{M}}^{-2}$ and (8), we conclude that $z = \operatorname{Exp}_{x}^{-1}(\tilde{z}) \in \operatorname{Exp}_{x}^{-1}(\mathcal{C})$, and hence $\tilde{z} \in \mathcal{C}$.

By the same arguments presented in Proposition 4.2, one can establish that Riemannian strong convexity implies the geodesic strong convexity of sets in Cartan–Hadamard manifolds. This situation bears resemblance to the various notions of geodesically convex sets for Cartan–Hadamard manifolds [12]. It should be noted that geodesic and double geodesic strong convexity (Definitions 3.2 and 3.4) become equivalent under mild assumptions, which is noteworthy as Definition 3.2 relies on the geodesic metric space structure of \mathcal{M} , while Definition 3.4 leverages the manifold structure of \mathcal{M} .

Proposition 4.3 (Equivalence between Geodesic and Double Geodesic Strong Convexity). Let \mathcal{M} be a complete, connected Riemannian manifold, and let us assume that Assumption 2.9 holds. If the subset $\mathcal{C} \subset \mathcal{M}$ is a geodesic α -strongly convex set, then it is also a double geodesic $\frac{\alpha}{L_{\mathcal{M}}}$ -strongly convex set. If the set \mathcal{C} is a double geodesic α -strongly convex set, then it is also a geodesic $\ell_{\mathcal{M}} \alpha$ -strongly convex set.

Proof. (\Rightarrow) We start with $C \subset M$ being an α -geodesically strongly convex set. Now, let $z \in T_{\gamma(t)}M$ such that

$$||z|| \le \frac{\alpha}{L_{\mathcal{M}}} t(1-t)d^2(x,y).$$

As the distance function d can be bounded as $d(x, y) \leq L_{\mathcal{M}} \| \operatorname{Exp}_{x}^{-1}(y) \|$, we have

$$||z|| \le \frac{\alpha}{L_{\mathcal{M}}} t(1-t)d^2(x,y) \quad \Rightarrow \quad d\left(\gamma(t), \operatorname{Exp}_{\gamma(t)}(z)\right) \le \alpha t(1-t)d^2(x,y)$$

As the set is geodesically strongly convex, we have $\operatorname{Exp}_{\gamma(t)}(z) \in \mathcal{C}$.

 (\Leftarrow) Now, we assume C to be a doubly exponentially strongly convex set with the parameter α . We construct the point $z \in \mathcal{M}$ such that

$$d(\gamma(t), z) \le \ell_{\mathcal{M}} \alpha t (1-t) d^2(x, y).$$

As the distance function d can be bounded as $\ell_{\mathcal{M}} \| \operatorname{Exp}_{\gamma(t)}^{-1}(z) \| \leq d(\gamma(t), z)$, we have

$$d(\gamma(t), z) \le \ell_{\mathcal{M}} \alpha t(1-t) d^2(x, y) \quad \Rightarrow \quad \|\operatorname{Exp}_{\gamma(t)}^{-1}(z)\| \le \alpha t(1-t) d^2(x, y)$$

As the set is α -double exponentially strongly convex, we have $z \in \mathcal{C}$.

5 Sublevel Sets of Geodesically Strongly Convex Functions

Definitions 3.2 to **3.4** formalize the concept of strong convexity for subsets of a Riemannian manifold. Additionally, **Definition 3.5** serves as a key tool in Section 7 to establish linear convergence. However, proving that a set is strongly convex might be difficult in practice. Hence, this section aims to develop the necessary framework for demonstrating the strong convexity of sets within Riemannian manifolds.

5.1 Euclidean Case

In the context of Hilbert spaces, the uniform convexity of ℓ_p or p-Schatten balls has been studied in the theory of Banach spaces [19, 11, 7]. However, for less standard cases, the strong convexity of sets can often be most efficiently demonstrated by showing that they correspond to the sublevel sets of strongly convex functions [36, 26]. We present the Riemannian counterpart of [36, Theorem 12] for sublevel sets of geodesically strongly convex functions (Definition 2.7). These findings extend to the notion of geodesically convex functions, as observed in [54] and [12, Proposition 11.8].

In the Euclidean setting, [36, Theorem 12] demonstrates that the sublevel sets of *L*-smooth, μ -strongly convex functions are strongly convex sets. In particular, the following set is $\mu/2\sqrt{2Ls}$ strongly convex,

$$Q_s := \{x \mid f(x) - f^* \le s\}, \text{ where } f^* = \min_x f(x).$$

5.2 Non-Euclidean Case

We now demonstrate that the sublevel sets of a geodesically smooth, strongly convex function are geodesic strongly convex sets. This result relies heavily on the following lemma.

Lemma 5.1 (Smoothness property). Let us consider f as a geodesically L-smooth function on the geodesically closed convex subset $C \subset M$, where M is an Cartan–Hadamard manifold. We denote $x^* \in \operatorname{argmin}_{x \in C} f(x)$. Then,

$$\|\nabla f(x)\|_{x} \le \sqrt{2L(f(x) - f(x^{*}))}.$$
(11)

This result is based on the concept of functional duality in a Riemannian manifold, which has been comprehensively studied in [8]. The corresponding proof can be referred to in Appendix A.2.

Theorem 5.2 (Geodesic Strong Convexity of sublevel sets). Let \mathcal{M} be a Riemannian manifold. Suppose that $\mathcal{C} \subseteq \mathcal{M}$ is uniquely geodesically convex and $f : \mathcal{C} \subseteq \mathcal{M} \to \mathbb{R}$ is a proper, geodesically L-smooth, and μ -strongly convex function on \mathcal{C} w.r.t. the distance function d satisfying Assumption 2.9. Let $x^* \in \mathcal{C}$ satisfying $\nabla f(x^*) = 0$. Let $Q_s := \{x \mid f(x) - f^* \leq s\} \subseteq \mathcal{C}$ be geodesically strictly convex for some s > 0, that is, every geodesic segment in Q_s is in the interior of Q_s except possibly for its endpoints. Then, Q_s is a geodesic strongly convex set with $\alpha = \mu/2\sqrt{2sL \max\{\ell_{\mathcal{M}}^{-2}; 1\}}$ (Definition 3.2).

Proof. Let s > 0, and let us consider $(x, y) \in Q_s^2$ and write γ as the geodesic between x and y. On successively using the geodesic smoothness of f, Cauchy-Schwartz, and Lemma 5.1, for all

smooth curve $c_t(\tilde{t}): c_t(0) = \gamma(t)$ and $t, \tilde{t} \in [0, 1]$, we obtain

$$f(c_{t}(\tilde{t})) - f^{*} \leq f(\gamma(t)) - f^{*} + \langle \nabla f(\gamma(t)); \operatorname{Exp}_{\gamma(t)}^{-1}(c_{t}(\tilde{t})) \rangle_{\gamma(t)} + \frac{L}{2} d^{2}(\gamma(t), c_{t}(\tilde{t})) \\ \leq f(\gamma(t)) - f^{*} + \|\nabla f(\gamma(t))\|_{\gamma(t)}\| \operatorname{Exp}_{\gamma(t)}^{-1}(c_{t}(\tilde{t}))\|_{\gamma(t)} + \frac{L}{2} d^{2}(\gamma(t), c_{t}(\tilde{t})) \\ \leq f(\gamma(t)) - f^{*} + \sqrt{2L(f(\gamma(t)) - f^{*})} \ell_{\mathcal{M}}^{-1} d(\gamma(t), c_{t}(\tilde{t})) + \frac{L}{2} d^{2}(\gamma(t), c_{t}(\tilde{t})) \\ \leq \left(\sqrt{f(\gamma(t)) - f^{*}} + \sqrt{\frac{L \max\{\ell_{\mathcal{M}}^{-2}; 1\}}{2}} d(\gamma(t), c_{t}(\tilde{t}))\right)^{2}.$$
(12)

Therefore, to ensure that $c_t(\tilde{t}) \in Q_s$, we can identify a sufficient condition on $d(\gamma(t), c_t(\tilde{t}))$, such that

$$\left(\sqrt{f(\gamma(t)) - f^*} + \sqrt{\frac{L \max\{\ell_{\mathcal{M}}^{-2}; 1\}}{2}} d(\gamma(t), c_t(\tilde{t}))\right)^2 \le s,$$

$$\Leftarrow d(\gamma(t), c_t(\tilde{t})) \le \text{sufficient condition} \le \sqrt{\frac{2}{L \max\{\ell_{\mathcal{M}}^{-2}; 1\}}} \left(\sqrt{s} - \sqrt{f(\gamma(t)) - f^*}\right)$$
(13)

As f is strongly convex, on using Definition 2.7 and because $(x, y) \in Q_s$, we obtain

$$f(\gamma(t)) - f^* \le s - (1-t)t\frac{\mu}{2}d^2(x,y),$$

Since $\sqrt{\cdot}$ is a concave function, we have $\sqrt{x-y} \leq \sqrt{x} - \frac{y}{2\sqrt{x}}$. Therefore,

$$\sqrt{f(\gamma(t)) - f^*} \le \sqrt{s - (1 - t)t\frac{\mu d^2(x, y)}{2}} \le \sqrt{s} - \frac{(1 - t)t\mu d^2(x, y)}{4\sqrt{s}}$$

Hence, we have

$$\sqrt{\frac{2}{L \max\{\ell_{\mathcal{M}}^{-2}; 1\}}} \left(\sqrt{s} - \sqrt{f(\gamma(t)) - f^*}\right) \ge \sqrt{\frac{2}{L \max\{\ell_{\mathcal{M}}^{-2}; 1\}}} \left(\frac{(1-t)t\mu d^2(x, y)}{4\sqrt{s}}\right)$$
(14)

$$= (1-t)t \frac{\mu}{2\sqrt{2sL\max\{\ell_{\mathcal{M}}^{-2}; 1\}}} d^2(x,y).$$
(15)

Therefore,

$$d(\gamma(t), c_t(\tilde{t})) \leq (1-t)t \frac{\mu}{2\sqrt{2sL\max\{\ell_{\mathcal{M}}^{-2}; 1\}}} d^2(x, y)$$

$$\stackrel{(15)}{\Longrightarrow} d(\gamma(t), c_t(\tilde{t})) \leq \sqrt{\frac{2}{L\max\{\ell_{\mathcal{M}}^{-2}; 1\}}} \left(\sqrt{s} - \sqrt{f(\gamma(t)) - f^*}\right)$$

$$\stackrel{(13)}{\Longrightarrow} \left(\sqrt{f(\gamma(t)) - f^*} + \sqrt{\frac{L\max\{\ell_{\mathcal{M}}^{-2}; 1\}}{2}} d(\gamma(t), c_t(\tilde{t}))\right)^2 \leq s$$

$$\stackrel{(12)}{\Longrightarrow} f(c_t(\tilde{t})) - f^* \leq s.$$

Hence, $c_t(\tilde{t})$ is in the set Q_s , which is the definition of geodesic strong convexity.

Example: Unit sphere. Let us consider \mathbb{S}^{n-1} , the unit sphere manifold embedded in \mathbb{R}^n , with the distance function $d(x, y) = \arccos(\langle x; y \rangle)$. Let us fix $x_0 \in \mathcal{M}$, and let $f(x) = d^2(x_0, x)$. Let the set $Q_s := \{x : f(x) \leq s\}$. When $s < (\frac{\pi}{2})^2$, the squared distance function is a geodesically smooth and strongly convex function (the constants of which depend on s) [41, Lem. 12.15], [55, pp153–154]. As Q_s is also a strictly convex set for $s < (\pi/2)^2$, the set Q_s is a geodesically strongly convex set, as shown in Theorem 5.2.

Example: Symmetric Positive Definite Matrices. Let \mathcal{M} be the set of symmetric positive definite matrices with the affine-invariance metric, which yields the distance function $d_{\mathcal{M}}(X,Y) = \sqrt{\sum_{i} \log^2 \lambda_i(X^{-1}Y)}$ and is a Cartan–Hadamard manifold, cf. [31]. Let us fix $X_0 \in \mathcal{M}$, and let $f(X) = d_{\mathcal{M}}^2(X_0, X)$. As \mathcal{M} is a Cartan–Hadamard manifold, its distance function is strongly convex, and d is also smooth in bounded sets. Therefore, the sets Q_s , s < f(0), are strongly convex, as shown in Theorem 5.2.

6 Double Geodesic Strong Convexity and Approximate Riemannian Scaling Inequality

Under mild assumptions, Section 7 shows that the Riemannian FW algorithm (Algorithm 1) admits a global linear convergence rate (Theorem 7.1) when the feasible set in (OPT) satisfies a Riemannian scaling inequality. However, there is no apparent link between (double) geodesic strong convexity (Definition 3.3 or Definition 3.4) and the Riemannian scaling inequality. Therefore, in this section, we explore the link between double geodesic strong convexity and the Riemannian scaling inequality. We introduce the notion of approximate Riemannian scaling inequality (Definition 6.2) and demonstrate that the quality of the approximation depends on the exponential map operator (Definition 6.1).

6.1 Double Geodesic Strong Convexity and Double Exponential Map

In this section, we link the double geodesic strong convexity (Definition 3.4) with an approximate version of the scaling inequality (introduced subsequently in Definition 6.2). We first introduce the *double exponential map* [27, 24, 48] and rewrite the definition of the double geodesic strong convexity using two geodesic paths.

Definition 6.1 (Double Exponential Map). Let \mathcal{M} be a complete, connected Riemannian manifold. Let $T_x\mathcal{M}$ be the tangent space to \mathcal{M} at $x \in \mathcal{M}$, $\operatorname{Exp}_x(\cdot) : T_x\mathcal{M} \to \mathcal{M}$ be the exponential map at x, and $\Gamma_x^y : T_x\mathcal{M} \to T_y\mathcal{M}$ be the transportation map between the tangent spaces $T_x\mathcal{M}$ and $T_y\mathcal{M}$. We define the double exponential map at $x \in \mathcal{M}$ as the function $\operatorname{Exp}_x(u, v) : T_x\mathcal{M} \times T_x\mathcal{M} \to \mathcal{M}$, such that

$$\operatorname{Exp}_{x}(u, v) := \operatorname{Exp}_{\operatorname{Exp}_{x}(u)} \left(\Gamma_{x}^{\operatorname{Exp}_{x}(u)} v \right).$$
 (double exponential map)

We also define $h_x(\cdot, \cdot) : T_x \mathcal{M} \times T_x \mathcal{M} \to T_x \mathcal{M}$ as the (unique) exponential map operator, such that

$$\operatorname{Exp}_{x}(h_{x}(u, v)) := \operatorname{Exp}_{x}(u, v), \quad \forall u, v \in T_{x}\mathcal{M}.$$
(Exponential Map Operator)

In particular, we can rewrite the double geodesic strong convexity of a set with the double exponential map. Informally, Definition 3.4 takes into consideration the geodesic γ between x and y and other geodesics departing from a $\gamma(t)$ that moves in every z direction, thus describing a closed ball in the tangent space $T_{\gamma(t)}\mathcal{M}$. Therefore, (6) in Definition 3.4 becomes

$$\operatorname{Exp}_{x}\left(\operatorname{Exp}_{x}^{-1}(\gamma(t)), \alpha t(1-t)d^{2}(x,y)z\right) \in \mathcal{C}, \text{ for all } z \in \overline{B}(0,1) \subset T_{x}\mathcal{M}.$$
(16)

This expression motivates the use of the term *double* to describe the notion of strong convexity.

6.2 Link with Approximate Riemannian Scaling Inequality

This section presents a connection between double geodesic strong convexity and a weaker version of the Riemannian scaling inequality using the exponential map operator. When the exponential map operator satisfies h(u, v) = u + v (basically, when the set is Euclidean), the double geodesic set strong convexity implies the Riemannian scaling inequality (Proposition A.1). Instead, in [24, 48], explicit approximations of the *exponential map operator* were proposed when the Riemannian manifold is symmetric or has a constant curvature, e.g., the Euclidean sphere or Lobachevsky spaces [24]. These approximations provide expansions of $h_x(u, v)$ of the form

$$h_x(u,v) = u + v + R_x(u,v),$$
(17)

where the term $R_x(u, v)$ can be an order of magnitude smaller than $\sqrt{\|u\|_x^2 + \|v\|_x^2}$. In these cases, we can no longer prove that the double geodesic strong convexity would imply the Riemannian scaling inequality. Instead, we introduce an *approximate Riemannian scaling inequality* in Definition 6.2 and demonstrate in Proposition 6.3 that the double geodesic strong convexity implies an approximate Riemannian scaling inequality. The approximation quality is controlled by $R_x(u, v)$.

Definition 6.2 (Approximate Riemannian Scaling Inequality). Let \mathcal{M} be a Riemannian manifold and $\mathcal{C} \subset \mathcal{M}$. Let us consider a geodesically α -strongly convex set $\mathcal{C} \subset \mathcal{M}$. We then say that \mathcal{C} satisfies the approximate Riemannian scaling inequality w.r.t. the distance $d(\cdot, \cdot)$ and the residual $r(\cdot) : \mathcal{C} \to T\mathcal{M}$ if, for all $x \in \mathcal{C}$, $w \in T_x\mathcal{M}$, and $v \in \operatorname{argmax}_{z \in \mathcal{C}} \langle w; \operatorname{Exp}_x^{-1}(z) \rangle_x$, we have

$$\langle w; \operatorname{Exp}_x^{-1}(v) \rangle_x \ge \alpha \|w\|_x d(v, x)^2 + \langle w; r(x) \rangle_x$$
, (Approximate Riemannian Scaling Inequality)

In Proposition 6.3, we now show that the geodesic strong convexity of a set in the Riemannian manifold \mathcal{M} implies an approximate Riemannian scaling inequality that depends on this difference $R_x(u, v)$ of the exponential operator map.

Proposition 6.3 (Double Geodesic Str. Cvx. implies Approximate Riemannian Scaling Inequality). Let us consider $C \subset M$ as double geodesically α -strongly convex (Definition 3.4) in a Riemannian manifold M. We define $R_x : T_x \mathcal{M} \times T_x \mathcal{M} \to T_x \mathcal{M}$, such that, for all $x \in C$, the exponential map operator (Definition 6.1) is decomposed as

$$h_x(u,v) = u + v + R_x(u,v), \quad \forall (u,v) \in T_x \mathcal{M}, \ x \in \mathcal{C}.$$
(18)

The approximate scaling inequality (Definition 6.2) is then satisfied w.r.t. $d(\cdot, \cdot)$ with the residual $r(\cdot)$ s.t. for all $x \in C$ and $w \in T_x \mathcal{M}$

$$r(x) = R_x \left[\frac{1}{2} \gamma'_{x,v}(0), \Gamma^x_{\gamma_{x,v}(1/2)} \left(\frac{\alpha}{4} d^2(x,v) z^* \right) \right], \qquad (\text{Residual})$$

where $v \in \operatorname{argmax}_{z \in \mathcal{C}} \langle w; \operatorname{Exp}_x^{-1}(z) \rangle_x$ and $z^* \in \operatorname{argmax}_{\|z\|_{\gamma_{x,v}(1/2)}=1} \langle \Gamma_x^{\gamma_{x,v}(1/2)}w; z \rangle_{\gamma_{x,v}(1/2)}$.

Proof. This proof is similar to Proposition A.1 until (34). We then write

$$u = \frac{1}{2}\gamma'_{x,v}(0) \text{ and } \omega = \left(\Gamma_x^{\gamma_{x,v}(1/2)}\right)^{-1} \left(\frac{\alpha}{4}d^2(x,v)z\right)^{-1}$$

Hence, we have

$$\left\langle w; \operatorname{Exp}_{x}^{-1}(v) \right\rangle_{x} \geq \left\langle w; \operatorname{Exp}_{x}^{-1}\left(\operatorname{Exp}_{x}\left(u + \omega + R(u, v) \right) \right) \right\rangle_{x}$$

Using the same arguments as those in the proof of Proposition A.1, we obtain

$$\left\langle w; \operatorname{Exp}_{x}^{-1}(v) \right\rangle_{x} \geq \frac{\alpha}{2} d^{2}(x, v) \|w\|_{x} + \left\langle w; R(\frac{1}{2}\gamma'_{x, v}(0), \Gamma^{x}_{\gamma_{x, v}(1/2)}(\frac{\alpha}{4}d^{2}(x, v)z^{*}) \right\rangle_{x}.$$

The approximate scaling inequality is meaningful in the regime wherein x and v are close. In this situation, the scale of the residual $r(\cdot)$ is determined by $\gamma'_{x,v}(0)$ and $d^2(x,v)$. In some setting, the residual is such that the approximate term $\langle w; r(x) \rangle_x$ in Definition 6.2 becomes negligible w.r.t. the original term $\alpha ||w||_x d(x,v)^2$. It should be noted that, in the analysis of the FW algorithm in Section 7, we select $w := -\nabla f(x_t)$, where x_t are the FW iterates.

The general expression of $R_x(u, v) = h_x(u, v) - u - v$ as a series is given in [27]. Dzhepko and Nikonorov [24] provide the explicit Taylor series for the Euclidean sphere [24, (6)] and Lobachevskii spaces [24, Section 3].

7 Frank-Wolfe on Geodesically Strongly Convex Sets

The FW algorithm is a first-order method that is used to solve constrained optimization problems in Banach spaces. Each iteration relies on a linear minimization step over the constraint region. It has recently been extended for constrained optimization over Riemannian manifolds. The Riemannian Frank-Wolfe (RFW) algorithm [61, 62] solves the following smooth convex constrained problem.

minimize
$$f(x)$$
, for $x \in \mathcal{C}$ (OPT)

where $\mathcal{C} \subseteq \mathcal{M}$ is a compact geodesically convex set, and f is geodesically smooth and convex.

Algorithm 1 Riemannian Frank-Wolfe (RFW) algorithm Require: $x_0 \in C \subset \mathcal{M}$; assume access to the geodesic map $\gamma : [0, 1] \to \mathcal{M}$ for $t = 0, 1, \cdots$ do 1. $v_t \leftarrow \operatorname{argmax}_{v \in C} \langle \nabla f(x_t); \operatorname{Exp}_{x_t}^{-1}(v) \rangle$ 2. Let $s_t = \operatorname{argmin}_{s \in [0,1]} s \langle \nabla f(x_t); \operatorname{Exp}_{x_t}^{-1}(v_t) \rangle_x + s^2 \frac{L}{2} d^2(x_t, v_t).$ 3. $x_{t+1} \leftarrow \gamma(s_t)$, where $\gamma(0) = x_t$ and $\gamma(1) = v_t$. end for

Remarkably, [61, 62] proves similar convergence rates of RFW as the FW algorithm with comparable structural assumptions on the optimization problem as in the Hilbertian setting. For instance, when the function f is geodesically convex and smooth and the set is compact convex, the RFW algorithm converges in $\mathcal{O}(1/T)$ [61, Theorem 3.4.]. Similarly, Weber and Sra [61, Theorem 3.5.] show linear convergence of Algorithm 1 when using short-step sizes, and the objective function is geodesically strongly convex and the optimum in the interior of the set.

In the Hilbertian setting, the FW algorithm admits various accelerated convergence regimes when the set is strongly convex. When the unconstrained optimum of f is outside the constraint set, the FW algorithm converges linearly [23], or when the function is strongly convex, the convergence is in $\mathcal{O}(1/T^2)$ without an assumption on the unconstrained optimum location [26]. The previous sections establish possible notions of strong convexity for sets in Riemannian manifolds. We now demonstrate that analog convergence regimes for the RFW algorithm on geodesically convex sets hold as in the case of the Hilbertian setting. These results complete the work of [61, 62].

7.1 Linear convergence of RFW under the Riemannian scaling inequality

As outlined in [38], the *scaling inequality* (Proposition 3.1.(b)) is a convenient characterization of the strong convexity of the set (in Hilbertian setting) for establishing convergence rates. We follow the same path and establish the convergence rate of the RFW algorithm when the

constraint set satisfies (approximate) Riemannian scaling inequalities. In Theorem 7.1, we thus prove the linear convergence of the RFW algorithm when the set C satisfies a Riemannian scaling inequality and the unconstrained optimum of f is outside C. This provides a generalization of [23] in a Riemannian setting.

Theorem 7.1 (Linearly Convergent RFW with Riemannian Scaling Inequality). Consider a geodesically convex set C and a geodesically convex function f that is L-smooth in C. Assume that any unconstrained optimum of f lies outside the constraint set C, and in particular, there exists c > 0 s.t. $\operatorname{argmin}_{x \in C} \|\nabla f(x)\|_x > c$. Assume that, for every $x \in C$, the (Riemannian scaling inequality) holds. Then Algorithm 1 converges linearly:

$$f(x_{t+1}) - f(x^*) \le (f(x_t) - f(x^*)) \max\{1/2, 1 - \alpha c/(2L)\}$$

Proof. This proof is based on [23, 26, 38], but in in a Riemannian setting. Using the geodesic smoothness (Definition 2.8) of f at $x_{t+1} = \gamma(s_t)$, we obtain

$$f(x_{t+1}) \le f(x_t) - \left\langle -\nabla f(x_t); \operatorname{Exp}_{x_t}^{-1}(\gamma(s_t)) \right\rangle_x + \frac{L}{2} d^2(x_t, \gamma(s_t)).$$

As γ is a geodesic between x_t and v_t , we have $d(x_t, \gamma(s_t)) = s_t d(x_t, v_t)$ and $\operatorname{Exp}_{x_t}^{-1}(\gamma(s_t)) = s_t \operatorname{Exp}_{x_t}^{-1}(v_t)$. Hence, we now have

$$f(x_{t+1}) \le f(x_t) - s_t \langle -\nabla f(x_t); \operatorname{Exp}_{x_t}^{-1}(v_t) \rangle_x + \frac{L}{2} s_t^2 d^2(x_t, v_t)$$

According to the short-step rule for s_t (Algorithm 1), for all $s \in [0, 1]$, we now have

$$f(x_{t+1}) \le f(x_t) - s \langle -\nabla f(x_t); \operatorname{Exp}_{x_t}^{-1}(v_t) \rangle_x + \frac{L}{2} s^2 d^2(x_t, v_t)$$

After using the optimality of v_t , we have $\langle -\nabla f(x_t); \operatorname{Exp}_{x_t}^{-1}(v_t) \rangle_x \leq \langle -\nabla f(x_t); \operatorname{Exp}_{x_t}^{-1}(x^*) \rangle_x$, where $x^* \in \mathcal{C}$ is a solution to (OPT). Then, owing to the geodesic convexity of f, we have

$$f(x^*) - f(x) \ge \langle \nabla f(x); \operatorname{Exp}_x^{-1}(x^*) \rangle_x$$

Hence, as it is the case in the Hilbertian setting, the FW gap $\langle -\nabla f(x_t); \operatorname{Exp}_{x_t}^{-1}(v_t) \rangle_x$ upper bounds the primal gap at x_t , i.e.,

$$f(x_t) - f(x^*) \le \langle -\nabla f(x_t); \operatorname{Exp}_{x_t}^{-1}(v_t) \rangle_x.$$

We write $h_t = f(x_t) - f(x^*)$, and we hence have

$$h_{t+1} \le h_t (1 - s/2) - s/2 \langle -\nabla f(x_t); \operatorname{Exp}_{x_t}^{-1}(v_t) \rangle_x + \frac{L}{2} s^2 d^2(x_t, v_t).$$
(19)

Now, with $c = \operatorname{argmin}_{x \in \mathcal{C}} \|\nabla f(x)\|_x > 0$, after using the (Riemannian scaling inequality) at x_t and since $-\nabla f(x_t) \in T_{x_t} \mathcal{M}$, we obtain

$$\langle -\nabla f(x_t); \operatorname{Exp}_{x_t}^{-1}(v_t) \rangle_x \ge \alpha c d(x_t, v_t)^2,$$

such that, for all $s \in [0, 1]$, we have

$$h_{t+1} \le h_t(1-s/2) + \frac{s}{2} \Big(Ls - \alpha c \Big) d^2(x_t, v_t).$$
 (20)

Then, if $\alpha c/L < 1$, by choosing $s = \alpha c/L$ in (20), we have $h_{t+1} \leq h_t(1 - \alpha c/(2L))$; else, we have $L - \alpha c < 0$, and on selecting s = 1, we simply have $h_{t+1} \leq h_t/2$. Hence,

$$h_{t+1} \le h_t \max\{1/2, 1 - \alpha c/(2L)\}.$$

Examples of sets satisfying the Riemannian scaling inequality are, for instance, sets of restricted diameter in Riemannian manifolds with bounded sectional curvature (see Section 8). However, this condition might be restrictive.

7.2 Local Linear convergence of RFW under the approximate Riemannian scaling inequality

Since the "exact" Riemannian scaling inequality is quite restrictive, this section provides a similar convergence result when the feasible sets satisfy only approximate Riemannian scaling inequalities (Definition 6.2).

Theorem 7.2 (Linearly Convergent RFW on Double Geodesic Strongly Convex Sets). Consider a complete connected Riemannian manifold and a distance $d(\cdot, \cdot)$ satisfying Assumption 2.9 with parameters $0 < \ell_{\mathcal{M}} \leq L_{\mathcal{M}}$. Assume that $\mathcal{C} \subset \mathcal{M}$ is an α -double geodesically strong convex set w.r.t. the distance $d(\cdot, \cdot)$ (Definition 3.4). Assume that the function f is a geodesically convex L-smooth function, and there exists c > 0 s.t.

$$\min_{x \in \mathcal{C}} \|\nabla f(x)\|_x > c.$$

Let us assume that there exists C > 0 s.t. that the residual (18) of the exponential map operator $R_x(\cdot, \cdot)$ for all $x \in C$ and $u, w \in T_x \mathcal{M}$ is such that

$$||R_x(u,w)||_x \le C \cdot \max\left\{ ||u||_x^2 ||w||_x; ||w||_x^2 ||u||_x \right\}.$$
(21)

Let us assume that, for some diameter $\delta > 0$, $d(x_t, v_t)^2 \leq (\alpha c)/(2\delta L\tilde{C})$, where $\tilde{C} := C \max\left\{\frac{\alpha^2}{16\delta\ell_{\mathcal{M}}}; \frac{\alpha}{4\ell_{\mathcal{M}}^2}\right\}$. Then,

$$f(x_{t+1}) - f(x^*) \le (f(x_t) - f(x^*)) \max\left\{\frac{1}{2}, 1 - \frac{\alpha c}{2L}\right\}$$

Proof. As in the proof of Theorem (7.1), (19) is satisfied, i.e.,

$$h_{t+1} \le h_t (1 - s_t/2) - s_t/2 \langle -\nabla f(x_t); \operatorname{Exp}_{x_t}^{-1}(v_t) \rangle_{x_t} + \frac{L}{2} s_t^2 d^2(x_t, v_t).$$
(22)

Now, from Proposition 6.3, as C is double geodesically α -strongly convex, an approximate Riemannian scaling inequality is satisfied (Definition 6.2) at x_t with $-\nabla f(x_t) \in T_{x_t}\mathcal{M}$ with the residual $r(x_t)$ as in (Residual), i.e.,

$$r(x_t) = R_{x_t} \left[\frac{1}{2} \gamma'_{x_t, v_t}(0), \Gamma^{x_t}_{\gamma_{x_t, v_t}(1/2)} \left(\frac{\alpha}{4} d^2(x_t, v_t) z^* \right) \right].$$
(23)

Hence, on combining with $\operatorname{argmin}_{x \in \mathcal{C}} \|\nabla f(x)\|_x > c$, we can lower-bound $\langle -\nabla f(x_t); \operatorname{Exp}_{x_t}^{-1}(v_t) \rangle_{x_t}$ as follows:

$$\langle -\nabla f(x_t); \operatorname{Exp}_{x_t}^{-1}(v_t) \rangle_{x_t} \ge \alpha c d(x_t, v_t)^2 + \langle -\nabla f(x); r(x_t) \rangle_{x_t}$$

On substituting this inequality in (22), for all $s \in [0, 1]$, we have

$$h_{t+1} \le h_t (1 - s/2) + \frac{s}{2} \Big(Ls - \alpha c \Big) d^2(x_t, v_t) + \frac{s}{2} \langle \nabla f(x_t); r(x_t) \rangle_{x_t}.$$
 (24)

We use (21) to upper-bound the term $||r(x_t)||_{x_t}$. Hence, we are first required to obtain an upper bound on $||\gamma'_{x_t,v_t}(0)||_{x_t}$ and $||\frac{\alpha}{4}d^2(x_t,v_t)z^*||_{\gamma_{x_t,v_t}(1/2)}$. We first note that

$$\left\|\frac{\alpha}{4}d^{2}(x_{t},v_{t})z^{*}\right\|_{\gamma_{x_{t},v_{t}}(1/2)} = \frac{\alpha}{4}d^{2}(x_{t},v_{t}).$$

Furthermore, by definition of the exponential mapping, we have $\|\gamma'_{x_t,v_t}(0)\|_{x_t} = \|\operatorname{Exp}_{x_t}^{-1}(v_t)\|_{x_t}$, and according to Assumption 2.9, we have

$$\|\gamma'_{x_t,v_t}(0)\|_{x_t} \le \frac{1}{\ell_{\mathcal{M}}} d(x_t,v_t).$$

When plugging these two bounds in the residual $r(x_t)$ (23), with the growth condition on the residual (21), we have

$$\|r(x_t)\|_{x_t} \leq C \max\left\{\frac{\alpha^2}{16\ell_{\mathcal{M}}}d^5(x_t, v_t); \frac{\alpha}{4\ell_{\mathcal{M}}^2}d^4(x_t, v_t)\right\}$$
(25)

$$||r(x_t)||_{x_t} \leq Cd^4(x_t, v_t) \max\{\frac{\alpha^2}{16\ell_{\mathcal{M}}}d(x_t, v_t); \frac{\alpha}{4\ell_{\mathcal{M}}^2}\}.$$
 (26)

On using $d(x_t, v_t) \leq \delta$, and with $\tilde{C} := C \max \{ \alpha^2 / (16\ell_{\mathcal{M}})\delta; \alpha / (4\ell_{\mathcal{M}}^2) \}$, we have $||r(x_t)||_{x_t} \leq \tilde{C}d^4(x_t, v_t)$. Hence, using the Cauchy-Schwartz inequality and $||\nabla f(x_t)||_x \leq \delta L$, (24) becomes

$$h_{t+1} \leq h_t (1 - s/2) + \frac{s}{2} \left(Ls - \alpha c \right) d^2(x_t, v_t) + \frac{s}{2} \delta L \tilde{C} d^4(x_t, v_t)$$
(27)

$$\leq h_t(1 - s/2) + \frac{s}{2}d^2(x_t, v_t) \Big(Ls - \alpha c + \delta L \tilde{C} d^2(x_t, v_t) \Big).$$
(28)

As we assumed $d(x_t, v_t)^2 \leq (\alpha c)/(2\delta L\tilde{C})$, we have $-\alpha c + \delta L\tilde{C}d^2(x_t, v_t) \leq -(\alpha c)/2 < 0$. Let us consider $s^* := (\alpha c - \delta L\tilde{C}d^2(x_t, v_t))/L > 0$. If $s^* > 1$, then the choice of s = 1 results in $h_{t+1} \leq h_t/2$. Else, $s^* \in [0, 1]$, and we select $s = s^*$. We hence obtain $h_{t+1} \leq h_t(1 - s^*/2) \leq 1 - (\alpha c)/(2L)$. Overall, we obtain

$$h_{t+1} \le h_t \max\{1/2, 1 - \alpha c/(2L)\}.$$

The previous theorem thus states that, provided a burn-in phase (that follows from the general $\mathcal{O}(1/T)$ of the RFW algorithm) to ensure $d(x_t, v_t)^2 \leq (\alpha c)/(2\delta L\tilde{C})$, the iterates of the RFW algorithm converge linearly when the set satisfies the approximate Riemannian scaling inequality.

Example 7.3 (Riemannian trust-region subproblem.). When minimizing a function f on a Riemannian manifold, a common approach is to use the trust-region method. After t iterations, it approximates $\tilde{f}_t \approx f$, where \tilde{f}_t is usually a second-order approximation of f around x_t . Common trust region approaches minimize $\tilde{f}_t(x_t + \Delta)$, where Δ belongs to the tangent space of x_t under the constraint that $\|\Delta\| \leq \delta_t$; we then use a retraction on $x_t + \Delta$ to obtain the iterate x_{t+1} . Alternatively, it is also possible to solve the subproblem directly on the manifold as follows:

$$x_{t+1} = \operatorname*{argmin}_{x \in \mathcal{M}, \ d(x_t, x) \le \delta_t} f_t(x).$$

On using Theorem 5.2, we determine that the set is geodesically strongly convex. Therefore, if the set is sufficiently small, the RFW algorithm converges linearly on this subproblem.

Example 7.4 (Global Riemannian optimization through local subproblem solving). For manifolds of curvature bounded in $[\kappa_{\min}, \kappa_{\max}]$, and defining $K \stackrel{\text{def}}{=} \max\{|\kappa_{\min}|, \kappa_{\max}\}$, Martínez-Rubio [43] presented a reduction from global Riemannian g-convex optimization to optimization in Riemannian balls of radius $O(\frac{1}{\sqrt{K}})$. It is required to solve $O(\zeta_R)$ of such ball optimization problems to an accuracy proportional to the final global accuracy times low polynomial factors on other parameters. Here, R is the initial distance to an optimizer, and ζ_R is a natural geometric constant, cf. Proposition 8.2. If we can solve the subproblems using linear rates, the reduction only adds a $\widetilde{O}(\zeta_R)$ factor to these linear rates.



Figure 1: Numerical convergence of the RFW algorithm's iterates (Algo. 1) for minimizing a quadratic function over a double geodesic strongly convex set in a unit sphere. The dimension of the problem is n = 500, x_c is a vector of ones, and the function f is a random quadratic function parametrized as $f(x) \triangleq ||A(x - x^*)||^2$, where A is a random 250 × 500 matrix, and x^* is generated at random such that $dist(x^*, x_c) \leq \pi/2$. The parameter R is set such that $R = 0.9 \operatorname{dist}(x^*, x_c)$, which ensures that the solution lies on the boundary, and therefore, there exists c > 0 s.t. $\operatorname{argmin}_{x \in \mathcal{C}} ||\nabla f(x)||_x > c$. As predicted by Theorem 7.2, the rate is locally linear.

7.3 Numerical Experiment: Minimization Over a Sphere

3

This section presents a numerical experiment illustrating the rates stated in Theorem 7.2. Let the manifold \mathcal{M} be \mathbb{S}^{n-1} , the unit sphere embedded in \mathbb{R}^n . Let us consider the following problem.

$$\min_{c \in \mathbb{S}^{n-1}} f(x), \qquad \text{subject to } \operatorname{dist}(x, x_c) \le R < \frac{\pi}{2}.$$
(29)

Owing to the symmetries of the sphere, the linear minimization oracle for (29) can be formulated as a simple one-dimensional problem (Proposition 9.2). This problem appears, for instance, when training a neural network with spherical constraints over a hierarchical dataset.

Example 7.5 (Hierarchical neural network with sphere constraints). Scieur and Kim [56] trained a neural network on a hierarchical dataset: "We force the classifier (hyperplane) of each class to belong to a sphere manifold, whose center is the classifier of its super-class". Hence, in the case wherein one wants to fine-tune the last layer of such an architecture, the problem can be formulated as in (29), where x_c is the separating hyperplane of the super-class, R is a user-defined parameter, and f(x) is the loss of the neural network parametrized by x over the dataset.

The solution of Equation (29) is computed using the function fminbnd from Matlab (this function uses the secant method). The experimental results have been reported in Figure 1, wherein the two regimes (global rate of $O(t^{-1})$ and locally linearly convergent) are clearly distinguishable.

8 Examples of Riemannian Strongly Convex Sets: Balls with Restricted Radius

In this section, we always make use of the Riemannian distance $d_{\mathcal{M}}(x,y) = \|\operatorname{Exp}_{x}^{-1}(y)\|$. The Riemannian strongly convex set, in Equation (7), is the most restrictive of our definitions. It implies that, for all points x in the set \mathcal{C} , the logarithmic image of \mathcal{C} around x, i.e., $\operatorname{Exp}_{x}^{-1}(\mathcal{C})$, is strongly convex in the Euclidean sense. Intuitively, we can expect that such a situation arises when the logarithmic map $\operatorname{Exp}_{x}^{-1}$ does not affect the shape of \mathcal{C} too much, which is the case when the sectional curvature of \mathcal{M} is bounded and the set \mathcal{C} is not too large.

In this section, we show that some sublevel sets of the function $x \mapsto \frac{1}{2} d_{\mathcal{M}}(x, x_0)^2$ are Riemannian strongly convex. Before proving the main theorem, we first introduce some concepts and two known results: **1**) distance functions are locally geodesically smooth and strongly convex [44], and **2**) locally, Euclidean pulled-back functions of geodesically smooth, strongly convex functions are smooth and strongly convex in the Euclidean sense [20].

8.1 Bounded curvature

In the following, we make an assumption regarding the curvature of our manifolds. Let us recall that, given a *two*-dimensional subspace $V \subseteq T_x \mathcal{M}$ of the tangent space of a point x, the sectional curvature at x with respect to V is defined as the Gauss curvature for the surface $\operatorname{Exp}_x(V)$ at x. The Gauss curvature at a point x can be defined as the product of the maximum and minimum curvatures of the curves resulting from intersecting the surface with planes that are normal to the surface at x. See more details on the curvature tensor \mathfrak{R} in [51]. Our assumption is as follows.

Assumption 8.1. The sectional curvatures of \mathcal{M} are contained in the interval $[\kappa_{\min}, \kappa_{\max}]$ and the covariant derivative of the curvature tensor is bounded as $\|\nabla \mathfrak{R}\| \leq F$.

This assumption is not overly restrictive. The majority of the applications of Riemannian optimization are in locally symmetric spaces, which satisfy $\nabla \mathfrak{R} = 0$, for instance, constant curvature spaces, the SPD matrix manifold with the usual metric, SO(n), and the Grasmannian manifold [42].

8.2 Strong Convexity and Smoothness of Distance Functions

We now state a fact regarding the smoothness and strong convexity of the distance squared to a point, that is central to many Riemannian optimization algorithms. In the sequel, we use the notation $K \stackrel{\text{def}}{=} \max\{|\kappa_{\min}|; \kappa_{\max}\}$.

Proposition 8.2. (See [44]) Let us consider a uniquely geodesic Riemannian manifold \mathcal{M} of sectional curvature bounded in $[\kappa_{\min}, \kappa_{\max}]$ and a ball $B_{x_0}(r)$ in \mathcal{M} of radius r centered at x_0 . The function $x \mapsto \frac{1}{2} d_{\mathcal{M}}(x, x_0)^2$ is then δ_r -strongly convex and ζ_r -smooth in $B_{x_0}(r)$, where δ_r and ζ_r are the geometric constants defined by

$$\zeta_r \stackrel{\text{def}}{=} \begin{cases} r\sqrt{|\kappa_{\min}|} \coth(r\sqrt{|\kappa_{\min}|}) & \text{if } \kappa_{\min} \le 0\\ 1 & \text{if } \kappa_{\min} > 0 \end{cases}; \quad \delta_r \stackrel{\text{def}}{=} \begin{cases} 1 & \text{if } \kappa_{\max} \le 0\\ r\sqrt{\kappa_{\max}} \cot(r\sqrt{\kappa_{\max}}) & \text{if } \kappa_{\max} > 0 \end{cases}$$
(30)

In the case of Cartan–Hadamard manifolds, $\delta_r = 1$ and $\zeta_r \in [r\sqrt{|\kappa_{\min}|}, r\sqrt{|\kappa_{\min}|} + 1]$.

8.3 Smoothness and Strong Convexity of Euclidean Pulled-Back Function

Under Assumption 8.1, [20, Proposition 6.1] showed that the Euclidean pulled-back function $x \mapsto f(\operatorname{Exp}_{x_0}(x))$ of a smooth, strongly convex function f defined in a ball of restricted radius in \mathcal{M} is also smooth, strongly convex (in the Euclidean sense) in a ball of the same radius.

Proposition 8.3. (Informal, see [20, Proposition 6.1] for details). Let \mathcal{M} be a uniquely geodesic Riemannian manifold that satisfies Assumption 8.1 and that contains $\mathcal{B}(x_{ref}, r)$ defined as

 $\mathcal{B}(x_{ref}, r): \{x \in \mathcal{M} : d_{\mathcal{M}}(x, x_{ref}) \le r\}, \quad and \ let \quad \mathcal{B}_{x_{ref}}(0, r): \{v \in \mathcal{T}_{x_{ref}}\mathcal{M} : \|v\| \le r\}.$

As shown above, we also defined the pulled-back ball to $\mathcal{T}_{x_{ref}}\mathcal{M}$. Let us assume the function $f : \mathcal{M} \to \mathbb{R}$ is L-smooth, μ -strongly geodesically convex in the ball $\mathcal{B}(x_{ref}, r)$, and has its minimizer $x^* \in \mathcal{B}(x_{ref}, r)$. Then, if

$$r \le \frac{\mu}{L} \min\left\{\frac{1}{4K}, \frac{K}{4F}\right\}$$

the pulled-back Euclidean function $x \mapsto f(\operatorname{Exp}_{x_0}(x))$ is $\frac{3}{2}L$ -smooth, and $\frac{1}{2}\mu$ -strongly convex over the ball $\mathcal{B}_{x_{ref}}(0,r)$.

One can relax the assumption of the minimizer being in the ball as this is only used to show that the Lipschitz constant of the function in the ball is at most 2Lr. For instance, if the distance between the minimizer and x_{ref} is R, the Lipschitz constant can be bounded by O(LR), and we could use this bound to conclude a similar statement.

8.4 Main result

Using Propositions 8.2 and 8.3, we can prove that balls and other sets obtained from sublevel sets of smooth and strongly g-convex functions in small regions are Riemannian strongly convex.

Theorem 8.4. Let \mathcal{M} be a uniquely geodesic Riemannian manifold that satisfies Assumption 8.1, and let the function $f: x \mapsto \frac{1}{2} d_{\mathcal{M}}(x, x_0)^2$, $x_0 \in \mathcal{M}$. Then, the sublevel sets

$$\mathcal{C}_r = \{ x \in \mathcal{M} : f(x) \le \frac{1}{2}r^2 \}$$

are Riemannian strongly convex if $r \leq \frac{1}{2} \frac{\delta_r}{\zeta_r} \min\{\frac{1}{4K}, \frac{K}{4F}\}$, where ζ_r and δ_r are the smoothness and strong convexity parameters of the function $\frac{1}{2} d_{\mathcal{M}}(\cdot, x_0)^2$ in \mathcal{C}_r (see Equation (30)).

Proof. In this setting, pulling the function back to the tangent space of any point in C_r results in a Euclidean function that is strongly convex and smooth with condition number $O(\zeta_r/\delta_r)$ in C_r . Furthermore, for all point $x \in C_r$, the function $\hat{f}_x : \operatorname{Exp}_x^{-1}(C_r) \to \mathbb{R}$ defined as $y \mapsto f(\operatorname{Exp}_x(y))$ for all $y \in \operatorname{Exp}_x^{-1}(C_r)$ is strongly convex with the parameter $\delta_r/2$, and in particular, $\operatorname{Exp}_x^{-1}(C_r)$ is a strongly convex set. It should be noted that we considered that the distance from x to any point in C_r is at most 2r.

This implies that Riemannian balls in these generic manifolds are geodesic strongly convex sets. We note that the greatest r that satisfies the condition in Theorem 8.4 is roughly a constant, considering constant curvature bounds. Being able to optimize in these sets is important as Martínez-Rubio [43] proved that one can reduce global geodesically convex optimization to the optimization over these sets.

9 Example of a Simple Linear Minimization Oracle

When the curvature is constant (spheres or hyperbolic spaces) and the domain is a ball, the linear minimization oracle can be simplified into a one-dimensional problem.

Theorem 9.1 (Linear minimization oracle in a ball in a constant curvature manifold). Let $\bar{B} \stackrel{\text{\tiny def}}{=} \mathcal{B}(x_0, r) \subset \mathcal{M}$ be a closed Riemannian ball in a manifold \mathcal{M} of a constant sectional curvature. If the curvature K is positive, let us assume $r < \frac{\pi}{2\sqrt{K}}$, so that the ball is uniquely geodesically convex. Given a point x and a direction $v \in T_x \mathcal{M}$, we define

$$\Gamma \stackrel{\text{\tiny def}}{=} \operatorname{Exp}_x \left(\operatorname{span} \{ \operatorname{Exp}_x^{-1}(x_0), v \} \right) \cap \{ x \mid d_{\mathcal{M}}(x, x_0) = r \}.$$

The solution of $\operatorname{argmin}_{z \in \overline{B}} \langle v, \operatorname{Exp}_x^{-1}(z) \rangle$ can then be found in Γ .

Proof. A solution is in the first set of the definition of Γ owing to the symmetries of the manifold. Indeed, let us assume there is a solution z, such that

$$z \notin \operatorname{Exp}_x\left(\operatorname{span}\left\{\operatorname{Exp}_x^{-1}(x_0), v\right\}\right)$$

Now, let us consider the points $\operatorname{Exp}_x^{-1}(z) \in T_x \mathcal{M}$, and $\operatorname{Exp}_x^{-1}(z') \in T_x \mathcal{M}$, such that z' is the point resulting from the application of $\operatorname{Exp}_x(\cdot)$ to the symmetric of $\operatorname{Exp}_x^{-1}(z)$ with respect to the plane span{ $\operatorname{Exp}_x^{-1}(x_0), v$ }.

Then, z' is also a solution; therefore, $\operatorname{Exp}_x(\frac{1}{2}\operatorname{Exp}_x^{-1}(z) + \frac{1}{2}\operatorname{Exp}_x^{-1}(z'))$ would also be a solution, and it would be in $\operatorname{Exp}_x(\operatorname{span}\{\operatorname{Exp}_x^{-1}(x_0), v\})$.

Moreover, any solution z satisfies $d_{\mathcal{M}}(x,z) = r$ because, otherwise, there exists a neighborhood around z contained in \mathcal{B} , and we could further decrease the function value of $\langle v, \operatorname{Exp}_{x}^{-1}(\cdot) \rangle$ for some point in this neighborhood. Finally, we can parametrize Γ as $\theta \mapsto$ the point of intersection of \mathcal{B} with the geodesic segment starting at x from the direction $\cos(\theta) \operatorname{Exp}_{x}^{-1}(x_{0}) + \sin(\theta)v$, where $\theta \in [0, 2\pi)$. Except for the degenerate case wherein $\operatorname{Exp}_{x}^{-1}(x_{0})$ is parallel to v, where Γ is a single point and the solution we are looking for.

A direct consequence of Theorem 9.1 is that the problem of approximating a solution of the linear max oracle can be solved by solving an alternative one-dimensional problem.

Proposition 9.2. Let u_1, u_2 be an orthonormal basis for the linear subspace $\operatorname{span}\{\operatorname{Exp}_x^{-1}(x_0), v\}$. The solution of $\operatorname{argmin}_{z\in\bar{B}}\langle v, \operatorname{Exp}_x^{-1}(z)\rangle$ can then be obtained by solving the following onedimensional problem,

$$\min_{\phi \in [-\pi,\pi]} \alpha(\phi) \langle v; p(\phi) \rangle$$

where $p(\phi)$ and $\alpha(\phi)$ are defined as

$$p(\phi) = \cos(\phi)u_1 + \sin(\phi)u_2, \qquad \phi \in [-\pi, \pi], \tag{31}$$

$$\alpha(\phi) = \underset{\alpha>0}{\operatorname{argmin}} : \operatorname{Dist}\left(\operatorname{Exp}_{x}(\alpha p(\phi)), x_{0}\right) = r.$$
(32)

Proof. The proof is immediate. The subspace span $\{ Exp_x^{-1}(x_0), v \}$ is parametrized with polar coordinates: p is a direction of the unitary norm, and α is the length. The non-linear equation (32) represents the smallest α such that $\alpha \cdot p$ intersects $\mathcal{B}(x_0, r)$. П

The solution ϕ can be computed using a one-dimensional solver, e.g., bisection or the Newton-Raphson method. It should be noted that finding $\alpha(\phi)$ is also a one-dimensional problem, and it can also be solved using similar techniques.

In some cases, there are simple formulas for $\alpha(\phi)$, e.g., when the manifold is a sphere (Proposition 9.3).

Proposition 9.3. Let $\mathcal{M} = \mathbb{S}^{n-1}$ be the unit sphere. The formula of Equation (32) then reads as

$$\alpha(\phi) = 2tan^{-1} \left(\frac{b(\phi) + \sqrt{a^2 + b(\phi)^2 - c^2}}{a + c} \right), \quad a = x_0^T x, \quad b(\phi) = x_0^T p(\phi), \quad c = 1 - 2\sin\left(\frac{r}{2}\right).$$

Proof. The distance function and exponential map in the sphere read as

$$\operatorname{Dist}(x,y) = 2\operatorname{asin}\left(\frac{\|y-x\|}{2}\right), \qquad \operatorname{Exp}_x(tv) = x\cos(t\|v\|) + \frac{tv}{\|tv\|}\sin(t\|v\|), \qquad v \in T_x\mathcal{M}, \ t \in \mathbb{R}.$$

Hence, for all $x, y \in \mathcal{M}$,

$$Dist(x, y) = 2 \operatorname{asin}\left(\frac{\|y - x\|}{2}\right) = r,$$

$$\Rightarrow \|y - x\| = 2 \operatorname{sin}\left(\frac{r}{2}\right),$$

$$\Rightarrow \|y\|^2 + \|x\|^2 - 2x^T y = 4 \operatorname{sin}^2\left(\frac{r}{2}\right),$$

$$\Rightarrow x^T y = 1 - 2 \operatorname{sin}^2\left(\frac{r}{2}\right).$$

By replacing x, y by $\operatorname{Exp}_x(\alpha p), x_0$ and using the expression of the exponential map, we obtain the equation

$$x_0^T x \cos(\alpha) + x_0^T p(\phi) \sin(\alpha) = 4 \sin^2\left(\frac{r}{2}\right).$$

The desired result follows after identifying the terms a, b, c in

$$a\cos(\alpha) + b\sin(\alpha) = c,$$

the solution of which is

$$\alpha = 2\tan^{-1}\left(\frac{b \pm \sqrt{a^2 + b^2 - c^2}}{a + c}\right) + 2k\pi, \qquad k \in \mathbb{Z}.$$

-	
-	

10 Conclusion

We presented the first definitions for strong convexity of sets in Riemannian manifolds, studied their relationships, and have provided examples of these sets. Our definitions seek to be wellsuited for optimization and for establishing the strongly convex nature of the set. The global linear convergence of the RFW algorithm serves as a tangible demonstration of the impact of developing a theory around the strong convexity structure of these sets.

However, most importantly, we expect the development of a strongly convex structure to be helpful when developing Riemannian algorithms in the contexts wherein the Euclidean algorithm counterpart was leveraging such a structure, e.g., in the generalized power method [36], online learning [34, 22, 9, 10], and even more broadly, in the case of the use of *strongification* techniques, as in [47].

Acknowledgments

David Martínez-Rubio was partially funded by the DFG Cluster of Excellence MATH+ (EXC-2046/1, project id 390685689) funded by the Deutsche Forschungsgemeinschaft (DFG). AdA would like to acknowledge support from a Google focused award, as well as funding by the French government under management of Agence Nationale de la Recherche as part of the "Investissements d'avenir" program, reference ANR-19-P3IA-0001 (PRAIRIE 3IA Institute). We would like to thank Editage (www.editage.co.kr) for English language editing.

References

- A. Ahidar-Coutrix, T. Le Gouic, and Q. Paris. Convergence rates for empirical barycenters in metric spaces: curvature, convexity and extendable geodesics. *Probability theory and related fields*, 177(1):323–368, 2020.
- [2] A. D. Aleksandrov. Ruled surfaces in metric spaces. Vestnik Leningrad. Univ., 1957.
- [3] Z. Allen-Zhu, A. Garg, Y. Li, R. Oliveira, and A. Wigderson. Operator scaling via geodesically convex optimization, invariant theory and polynomial identity testing. In *Proceedings* of the 50th Annual ACM SIGACT Symposium on Theory of Computing, pages 172–181, 2018.
- [4] W. Ambrose. Parallel translation of riemannian curvature. Annals of Mathematics, 1956.
- [5] M. Bacák. Convex analysis and optimization in Hadamard spaces. de Gruyter, 2014.
- [6] M. Bacak. Old and new challenges in hadamard spaces. arXiv preprint arXiv:1807.01355, 2018.
- [7] K. Ball, E. A. Carlen, and E. H. Lieb. Sharp uniform convexity and smoothness inequalities for trace norms. *Inventiones mathematicae*, 115(1):463–482, 1994.
- [8] R. Bergmann, R. Herzog, M. S. Louzeiro, D. Tenbrinck, and J. Vidal-Núñez. Fenchel duality theory and a primal-dual algorithm on riemannian manifolds. *Foundations of Computational Mathematics*, pages 1–40, 2021.
- [9] A. Bhaskara, A. Cutkosky, R. Kumar, and M. Purohit. Online learning with imperfect hints. In *International Conference on Machine Learning*, pages 822–831. PMLR, 2020.
- [10] A. Bhaskara, A. Cutkosky, R. Kumar, and M. Purohit. Online linear optimization with many hints. arXiv:2010.03082, 2020.
- [11] R. P. Boas Jr. Some uniformly convex spaces. Bulletin of the American Mathematical Society, 46(4):304–311, 1940.
- [12] N. Boumal. An introduction to optimization on smooth manifolds. Available online, Aug, 2020.
- [13] N. Boumal, B. Mishra, P.-A. Absil, and R. Sepulchre. Manopt, a matlab toolbox for optimization on manifolds. *The Journal of Machine Learning Research*, 15(1), 2014.
- [14] G. Braun, A. Carderera, C. W. Combettes, H. Hassani, A. Karbasi, A. Mokhtari, and S. Pokutta. Conditional gradient methods. arXiv preprint arXiv:2211.14103, 2022.
- [15] M. R. Bridson and A. Haefliger. Metric spaces of non-positive curvature, volume 319. Springer Science & Business Media, 2013.
- [16] S. Bubeck, M. Cohen, and Y. Li. Sparsity, variance and curvature in multi-armed bandits. In *Proceedings of Algorithmic Learning Theory*, pages 111–127. PMLR, 2018.
- [17] D. Burago, I. D. Burago, Y. Burago, S. Ivanov, S. V. Ivanov, and S. A. Ivanov. A course in metric geometry, volume 33. American Mathematical Soc., 2001.
- [18] Y. Burago, M. Gromov, and G. Perel'man. Ad alexandrov spaces with curvature bounded below. *Russian mathematical surveys*, 47(2):1–58, 1992.
- [19] J. Clarkson. Uniformly convex spaces. Transactions of the American Mathematical Society, 40(3):396–414, 1936.
- [20] C. Criscitiello and N. Boumal. Negative curvature obstructs acceleration for geodesically convex optimization, even with exact first-order oracles. *CoRR*, abs/2111.13263v1, 2021. URL https://arxiv.org/abs/2111.13263v1.

- [21] M. Cuturi. Sinkhorn distances: Lightspeed computation of optimal transport. Advances in neural information processing systems, 26:2292–2300, 2013.
- [22] O. Dekel, A. Flajolet, N. Haghtalab, and P. Jaillet. Online learning with a hint. In Advances in Neural Information Processing Systems, pages 5299–5308, 2017.
- [23] V. F. Demyanov and A. M. Rubinov. Approximate methods in optimization problems. Modern Analytic and Computational Methods in Science and Mathematics, 1970.
- [24] V. Dzhepko and Y. G. Nikonorov. The double exponential map on spaces of constant curvature. Siberian Advances in Mathematics, 18(1):21–29, 2008.
- [25] J. Feydy, T. Séjourné, F.-X. Vialard, S.-i. Amari, A. Trouve, and G. Peyré. Interpolating between optimal transport and mmd using Sinkhorn divergences. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 2681–2690, 2019.
- [26] D. Garber and E. Hazan. Faster rates for the frank-wolfe method over strongly-convex sets. In International Conference on Machine Learning, pages 541–549. PMLR, 2015.
- [27] A. V. Gavrilov. The double exponential map and covariant derivation. Siberian Mathematical Journal, 48(1):56–61, 2007.
- [28] V. Goncharov and G. Ivanov. Strong and weak convexity of closed sets in a Hilbert space. Springer, 2017.
- [29] T. L. Gouic, Q. Paris, P. Rigollet, and A. J. Stromme. Fast convergence of empirical barycenters in alexandrov spaces and the wasserstein space. arXiv preprint arXiv:1908.00828, 2019.
- [30] H. Hopf and W. Rinow. Über den begriff der vollständigen differentialgeometrischen fläche. Commentarii Mathematici Helvetici, 3(1):209–225, 1931.
- [31] R. Hosseini and S. Sra. Matrix manifold optimization for Gaussian mixtures. In Advances inNeural Information Processing Systems 28: An-Conference Neural Information Processing nualonSystems 2015.Decem-7-12. 2015.Montreal, Quebec, Canada, pages 910–918, 2015.URL berhttp://papers.nips.cc/paper/5812-matrix-manifold-optimization-for-gaussian-mixtures.
- [32] R. Hosseini and S. Sra. An alternative to EM for gaussian mixture models: Batch and stochastic Riemannian optimization. *CoRR*, abs/1706.03267, 2017. URL http://arxiv.org/abs/1706.03267.
- [33] R. Hosseini and S. Sra. Recent advances in stochastic Riemannian optimization. Handbook of Variational Methods for Nonlinear Geometric Data, pages 527–554, 2020.
- [34] R. Huang, T. Lattimore, A. György, and C. Szepesvári. Following the leader and fast rates in online linear prediction: Curved constraint sets and other regularities. *The Journal of Machine Learning Research*, 18(1):5325–5355, 2017.
- [35] J. Jost. Nonpositive curvature: geometric and analytic aspects. Birkhäuser, 2012.
- [36] M. Journée, Y. Nesterov, P. Richtárik, and R. Sepulchre. Generalized power method for sparse principal component analysis. *Journal of Machine Learning Research*, 11(2), 2010.
- [37] T. Kerdreux, A. d'Aspremont, and S. Pokutta. Local and global uniform convexity conditions. arXiv preprint arXiv:2102.05134, 2021.
- [38] T. Kerdreux, A. d'Aspremont, and S. Pokutta. Projection-free optimization on uniformly convex sets. In *International Conference on Artificial Intelligence and Statistics*, pages 19–27. PMLR, 2021.
- [39] T. Kerdreux, L. Liu, S. Lacoste-Julien, and D. Scieur. Affine invariant analysis of Frank-Wolfe on strongly convex sets. In *International Conference on Machine Learning*, pages 5398–5408. PMLR, 2021.

- [40] T. Kerdreux, C. Roux, A. d'Aspremont, and S. Pokutta. Linear bandits on uniformly convex sets. *Journal of Machine Learning Research*, 22(284):1–23, 2021.
- [41] J. M. Lee. Introduction to Riemannian manifolds. Springer, 2018.
- [42] M. Lezcano-Casado. Curvature-dependant global convergence rates for optimization on manifolds of bounded geometry. arXiv preprint arXiv:2008.02517, 2020.
- [43] D. Martínez-Rubio. Global Riemannian acceleration in hyperbolic and spherical spaces. arXiv preprint arXiv:2012.03618, 2020. URL https://arxiv.org/abs/2012.03618.
- [44] D. Martínez-Rubio and S. Pokutta. Accelerated Riemannian optimization: Handling constraints with a prox to bound geometric penalties. arXiv preprint arXiv:2211.14645, 2022. URL https://arxiv.org/abs/2211.14645.
- [45] Q. Mérigot, A. Delalande, and F. Chazal. Quantitative stability of optimal transport maps and linearization of the 2-wasserstein space. In *International Conference on Artificial Intelligence and Statistics*, pages 3186–3196. PMLR, 2020.
- [46] W. Meyer. Toponogov's theorem and applications. Lecture Notes, Trieste, 1989.
- [47] M. Molinaro. Curvature of feasible sets in offline and online optimization. arXiv:2002.03213, 2020.
- [48] Y. G. Nikonorov. Double exponential map on symmetric spaces. Siberian Advances in Mathematics, 23(3):210–218, 2013.
- [49] Q. Paris. The exponentially weighted average forecaster in geodesic spaces of non-positive curvature. *arXiv:2002.00852*, 2020.
- [50] Q. Paris. Online learning with exponential weights in metric spaces. arXiv:2103.14389, 2021.
- [51] P. Petersen. *Riemannian geometry*, volume 171. Springer, 2006.
- [52] G. Pisier. Martingales with values in uniformly convex spaces. Israel Journal of Mathematics, 20(3-4):326–350, 1975.
- [53] G. Pisier. Martingales in Banach spaces (in connection with type and cotype). course IHP, Feb. 2–8, 2011.
- [54] T. Rapcsák. Smooth nonlinear optimization in Rn. Springer Science & Business Media, 2013.
- [55] T. Sakai. On riemannian manifolds admitting a function whose gradient is of constant norm. Kodai Mathematical Journal, 19(1):39–51, 1996.
- [56] D. Scieur and Y. Kim. Connecting sphere manifolds hierarchically for regularization. In International Conference on Machine Learning, pages 9399–9409. PMLR, 2021.
- [57] J. Sun, Q. Qu, and J. Wright. Complete dictionary recovery over the sphere II: recovery by Riemannian trust-region method. *IEEE Trans. Inf. Theory*, 63(2):885–914, 2017. doi: 10.1109/TIT.2016.2632149. URL https://doi.org/10.1109/TIT.2016.2632149.
- [58] J. Vial. Strong and weak convexity of sets and functions. Math. Oper. Res., 8(2):231-259, 1983. doi: 10.1287/MOOR.8.2.231. URL https://doi.org/10.1287/moor.8.2.231.
- [59] J.-P. Vial. Strong convexity of sets and functions. Journal of Mathematical Economics, 9 (1-2):187–205, 1982. doi: https://doi.org/10.1016/0304-4068(82)90026-X.
- [60] C. Villani. Optimal transport: old and new, volume 338. Springer, 2009.
- [61] M. Weber and S. Sra. Riemannian optimization via frank-wolfe methods. arXiv preprint arXiv:1710.10770, 2017.

- [62] M. Weber and S. Sra. Projection-free nonconvex stochastic optimization on riemannian manifolds. arXiv preprint arXiv:1910.04194, 2019.
- [63] H. Zhang and S. Sra. First-order methods for geodesically convex optimization. In Conference on Learning Theory, pages 1617–1638. PMLR, 2016.
- [64] H. Zhang and S. Sra. An estimate sequence for geodesically convex optimization. In Conference On Learning Theory, pages 1703–1723. PMLR, 2018.

A Technical Proposition and Lemmas

A.1 Double Geodesic Strong Convexity implies Riemannian Scaling Inequality

Proposition A.1 (Double Geodesic Str. Cvx. implies Riemannian Scaling Inequality). Let $C \subset \mathcal{M}$ be a double geodesic α -strongly convex set (Definition 3.4) in a complete connected Riemannian manifold \mathcal{M} . Let us assume that the double exponential map operator (Definition 6.1) is such that

$$h_x(u,v) = u + v, \quad \forall (u,v) \in T_x \mathcal{M}.$$
(33)

The Riemannian scaling inequality in Definition 3.5 is then satisfied.

Proof. Let us consider $x \in \mathcal{C}$, $w \in T_x \mathcal{M}$, and $v \in \mathcal{C}$ s.t.

$$v \in \operatorname*{argmax}_{z \in \mathcal{C}} \langle w; \operatorname{Exp}_x^{-1}(z) \rangle_x$$

Then, by (16), which is equivalent to Definition 3.4 with t = 1/2, we have $\operatorname{Exp}_{\gamma_{x,v}(1/2)}\left(\frac{\alpha}{4}d^2(x,v)z\right) \in \mathcal{C}$, where $\gamma_{x,v}(\cdot)$ is the geodesic joining x, and v and z are a unit norm vector in $T_{\gamma_{x,v}(1/2)}\mathcal{M}$. Then, by optimality of v, for all $z \in T_{\gamma_{x,v}(1/2)}\mathcal{M}$ with $\|z\|_{\gamma_{x,v}(1/2)} = 1$, we have

$$\left\langle w; \operatorname{Exp}_{x}^{-1}(v) \right\rangle_{x} \geq \left\langle w; \operatorname{Exp}_{x}^{-1}\left(\operatorname{Exp}_{\gamma_{x,v}(1/2)}\left(\frac{\alpha}{4} d^{2}(x,v)z \right) \right) \right\rangle_{x}$$

Let us first recall that, by definition of the exponential map, for the geodesic $\gamma_{x,v}$ and for all $t \in [0,1]$, we have $\operatorname{Exp}_x(t\gamma'_{x,v}(0)) = \gamma_{x,v}(t)$ such that we can write

$$\left\langle w; \operatorname{Exp}_{x}^{-1}(v) \right\rangle_{x} \geq \left\langle w; \operatorname{Exp}_{x}^{-1}\left(\operatorname{Exp}_{\operatorname{Exp}_{x}(\frac{1}{2}\gamma_{x,v}'(0))}\left(\frac{\alpha}{4}d^{2}(x,v)z\right) \right) \right\rangle_{x}.$$
 (34)

We can hence write this in terms of the double exponential map and subsequently in terms of the exponential operator map. We use

$$\operatorname{Exp}_{\operatorname{Exp}_{x}(\frac{1}{2}\gamma_{x,v}'(0))}\left(\frac{\alpha}{4}d^{2}(x,v)z\right) = \operatorname{Exp}_{x}\left(\frac{1}{2}\gamma_{x,v}'(0);\left(\Gamma_{x}^{\gamma_{x,v}(1/2)}\right)^{-1}\left(\frac{\alpha}{4}d^{2}(x,v)z\right)\right).$$

Hence, on using the assumption (33) on the exponential operator, we have

$$\operatorname{Exp}_{\operatorname{Exp}_{x}(\frac{1}{2}\gamma'_{x,v}(0))}\left(\frac{\alpha}{4}d^{2}(x,v)z\right) = \operatorname{Exp}_{x}\left(\frac{1}{2}\gamma'_{x,v}(0) + \left(\Gamma_{x}^{\gamma_{x,v}(1/2)}\right)^{-1}\left(\frac{\alpha}{4}d^{2}(x,v)z\right)\right).$$

When plugging the last equality in (34), we obtain

$$\left\langle w; \operatorname{Exp}_{x}^{-1}(v) \right\rangle_{x} \geq \frac{1}{2} \left\langle w; \gamma_{x,v}'(0) \right\rangle_{x} + \left\langle w; \left(\Gamma_{x}^{\gamma_{x,v}(1/2)} \right)^{-1} \left(\frac{\alpha}{4} d^{2}(x,v)z \right) \right\rangle_{x}.$$

It should be noted that $\gamma'_{x,v}(0) = \operatorname{Exp}_x^{-1}(v)$, such that

$$\left\langle w; \operatorname{Exp}_{x}^{-1}(v) \right\rangle_{x} \geq 2 \left\langle w; \left(\Gamma_{x}^{\gamma_{x,v}(1/2)} \right)^{-1} \left(\frac{\alpha}{4} d^{2}(x,v)z \right) \right\rangle_{x}.$$

Wth $(\Gamma_x^y)^{-1} = \Gamma_y^x$ and the isometry property of $\Gamma_x^{\gamma_{x,v}(1/2)}$, we have

$$\left\langle w; \Gamma_{\gamma_{x,v}(1/2)}^{x} \left(\frac{\alpha}{4} d^{2}(x,v)z\right) \right\rangle_{x} = \left\langle \Gamma_{\gamma_{x,v}(1/2)}^{x} \Gamma_{x}^{\gamma_{x,v}(1/2)}w; \Gamma_{\gamma_{x,v}(1/2)}^{x} \left(\frac{\alpha}{4} d^{2}(x,v)z\right) \right\rangle_{x}$$
(35)
$$= \left\langle \Gamma_{x}^{\gamma_{x,v}(1/2)}w; \frac{\alpha}{4} d^{2}(x,v)z \right\rangle_{\gamma_{x,v}(1/2)}.$$
(36)

Hence, for all z of a unit norm in $T_{\gamma_{x,v}(1/2)}\mathcal{M}$, we obtain

$$\left\langle w; \operatorname{Exp}_{x}^{-1}(v) \right\rangle_{x} \geq 2 \left\langle \Gamma_{x}^{\gamma_{x,v}(1/2)}w; \frac{\alpha}{4}d^{2}(x,v)z \right\rangle_{\gamma_{x,v(1/2)}}$$
(37)

$$= \frac{\alpha}{2} d^2(x,v) \left\langle \Gamma_x^{\gamma_{x,v}(1/2)} w; z \right\rangle_{\gamma_{x,v(1/2)}}.$$
(38)

Furthermore, by maximizing over z, for the best z^* , we obtain

$$\left\langle \Gamma_x^{\gamma_{x,v}(1/2)} w; z^* \right\rangle_{\gamma_{x,v}(1/2)} = \|\Gamma_x^{\gamma_{x,v}(1/2)} w\|_{\gamma_{x,v}(1/2)}.$$

Then, because the parallel transport $\Gamma_x^{\gamma_{x,v}(1/2)}$ is an isometry, we finally have

$$\left\langle w; \operatorname{Exp}_{x}^{-1}(v) \right\rangle_{x} \ge \frac{\alpha}{2} d^{2}(x, v) \| \Gamma_{x}^{\gamma_{x, v}(1/2)} \nabla f(x) \|_{\gamma_{x, v}(1/2)} = \frac{\alpha}{2} d^{2}(x, v) \| w \|_{x}.$$

A.2 Proof of Smoothness Property Lemma

First, we introduce the Fenchel conjugate of a function defined on a manifold.

Definition A.2. [8] Let us Suppose that $f : \mathcal{C} \to \mathbb{R}$, where $\mathcal{C} \subset \mathcal{M}$ is a strictly convex set, where \mathcal{M} is a Cartan-Hadamard manifold. For $m \in \mathcal{M}$, the m-Fenchel conjugate of f is defined as the function $f_m^* : T_m^* \mathcal{M} \to \mathbb{R}$ such that

$$f_m^*(\xi_m) := \sup_{x \in T_m \mathcal{M}} \left\{ \langle \xi_m, x \rangle - f\left(\operatorname{Exp}_m x \right) \right\},$$
(39)

where $T_m^*\mathcal{M}$ is the cotangent bundle of $T_m\mathcal{M}$

In particular, we need the following property.

Lemma A.3. [8, lem. 3.7] Let us suppose that $f, \tilde{f} : \mathcal{C} \to \mathbb{R}$ are proper functions, where $\mathcal{C} \subset \mathcal{M}$ is a strictly convex set, and \mathcal{M} is a Cartan–Hadamard manifold, and let $m \in \mathcal{C}$. Then,

if
$$f(p) \leq \tilde{f}(p) \ \forall \ p \in \mathcal{C}, \ then \ f_m^*(\xi_m) \geq \tilde{f}_m^*(\xi_m) \ \forall \ \xi_m \in T_m^*\mathcal{M}.$$
 (40)

We are now ready to prove Lemma 5.1.

Lemma 5.1 (Smoothness property). Let us consider f as a geodesically L-smooth function on the geodesically closed convex subset $C \subset M$, where M is an Cartan-Hadamard manifold. We denote $x^* \in \operatorname{argmin}_{x \in C} f(x)$. Then,

$$\|\nabla f(x)\|_{x} \le \sqrt{2L(f(x) - f(x^{*}))}.$$
(11)

Proof. As f is a geodesically smooth function, for all $p \in \mathcal{C}$, we have $f(p) \leq \tilde{f}(p)$, where

$$\tilde{f}(p) = f(x) + \langle \nabla f(x), \operatorname{Exp}_{x}^{-1}(p) \rangle + \frac{L}{2} d^{2}(x, p), \quad \forall p \in \mathcal{C}.$$

Therefore, according to Lemma A.3, we have $f_m^*(\xi_m) \geq \tilde{f}_m^*(\xi_m)$, for all $m \in \mathcal{M}$. Using the definition (39),

$$\tilde{f}_m^*(\xi_m) = \sup_{Z \in T_m \mathcal{M}} \langle \xi_m, Z \rangle_x - \tilde{f}(\operatorname{Exp}_m(Z))$$

In the particular case wherein m = x, we obtain

$$\tilde{f}_x^*(\xi_x) = \sup_{Z \in T_m \mathcal{M}} \langle \xi_x, Z \rangle - f(x) - \langle \nabla f(x), Z \rangle_x - \frac{L}{2} \|Z\|_x^2$$
$$= -f(x) + \frac{1}{2L} \|\xi_x - \nabla f(x)\|_x^2$$

Therefore, the inequality $f_x^*(\xi_x) \ge \tilde{f}_x^*(\xi_x)$ can be written as

$$\frac{1}{2L} \|\xi_x - \nabla f(x)\|_x^2 \le f_x^*(\xi_x) + f(x).$$

In particular, at $\xi_x = 0$, we have the desired result, as

$$f_x^*(0) = \sup_{Z \in T_m \mathcal{M}} \langle 0, Z \rangle_x - f(\operatorname{Exp}_x(Z)) = \sup_{Z \in T_m \mathcal{M}} -f(\operatorname{Exp}_x(Z)) = -\inf_{z \in \mathcal{C}} f(z) = -f^*.$$

14		

B MATLAB code for the experiments

B.1 Subroutine: Linear Max Oracle For a Ball In a Sphere

```
function v = linear_max_oracle(w, x, R, x0, manifold)
1
2
  % Solve the problem
3
  % max w<sup>Tz</sup> : z \in exp_x<sup>{</sup>{-1}(C)},
4
  % where C := \{x: dist(x, x_0) \setminus leq R\}
  % Assume that the sphere has radius = 1!
6
7
  \% Define the two basis vector of the subspace u_1 and u_2
8
9
   u1 = w;
   u1 = u1/norm(u1);
10
   u2 = manifold \cdot log(x, x0);
11
   u2 = u2 - u1 * (u1' * u2);
12
13
   if norm(u2) > 0
14
        u^{2} = u^{2} / norm(u^{2});
15
        p = @(phi) \cos(phi) * u1 + \sin(phi) * u2;
16
17
        alpha = @(p) solve_sincoseq(x0'*x, x0'*p,(1-2*sin(R/2)^2));
18
19
        fhandle = @(phi) - (alpha(p(phi)) * cos(phi)); %same minimum since w
20
           *u2 = 0.
        options = optimset('TolX', eps);
21
        [best_phi] = fminbnd(fhandle, -pi, pi, options);
22
        v = manifold.exp(x, p(best_phi), alpha(p(best_phi)));
^{23}
   else
24
        alpha = solve\_sincoseq(x0'*x, x0'*u1, (1-2*sin(R/2)^2)); \% best
25
           direction: u1
        v = manifold \cdot exp(x, u1, alpha);
26
27 end
```

B.2 Sample Script: Random Quadratic Over a Sphere With Ball Constraint

```
1 % Generate the problem data.
  d = 50;
2
  n = 25; % Slower convergence when n < d \rightarrow non-strongly convex case
3
  nIter = 500;
4
   dualgap = zeros(1, nIter);
5
6
  % Manifold: this code uses manop, see https://www.manopt.org
7
   manifold = spherefactory(d);
8
9
  % Define the problem cost function and its derivatives.
10
  A = randn(n,d);
11
^{12} A = A'*A;
  A = A/norm(A);
13
  xstar = manifold.rand();
14
  f = @(x) \quad 0.5 * (x - xstar) ' * A * (x - xstar);
15
  mgrad = @(x) manifold.egrad2rgrad(x, A*(x-xstar));
16
  L = norm(A); % Worst case
17
18
  % Create the problem set
19
  x_center = manifold.rand();
20
   radius ratio = 0.9; % <1: xstar is oustide
21
   radius max = manifold.dist(x center, xstar)*radius ratio;
22
   setFunction = @(x) manifold.dist(x center, x) <= radius max;
23
24
  x = x center;
25
  % Main loop RFW
26
   for i=1:(nIter-1)
27
       gradx = mgrad(x);
28
       v = linear_max_oracle(-gradx, x, radius_max, x_center, manifold);
29
       dualgap(i) = -manifold.inner(x, gradx, manifold.log(x, v));
30
31
       step_size = -\text{manifold.inner}(x, \text{ grad}x, \text{ manifold.log}(x, v)) / (L*
32
           manifold.dist(x, v)^2;
       step size = \min(\text{step size}, 1);
33
       x = manifold.exp(x, manifold.log(x, v), step_size);
34
   end
35
   v = linear_max_oracle(-gradx, x, radius_max, x_center, manifold);
36
   dualgap(end) = -manifold.inner(x, gradx, manifold.log(x, v));
37
38
  % Max at eps, otherwise the result is numerically meaningless
39
  semilogy(1:length(dualgap), max(dualgap, eps))
40
  legend({ 'FW Dual Gap'})
41
```