

Can the “basis vectors”, describing the internal space of point fermion and boson fields with the Clifford odd (for fermions) and Clifford even (for bosons) objects, be meaningfully extended to strings”?

N.S. Mankoč Borštnik¹, H.B. Nielsen²

¹Department of Physics, University of Ljubljana
SI-1000 Ljubljana, Slovenia

²Niels Bohr Institute, University of Copenhagen
Blegdamsvej 17, Copenhagen, Denmark

December 14, 2023

Abstract

Abstract: The *string theory* seems to be a mathematically consistent way for explaining so far observed fermion and boson second quantized fields, with gravity included, by offering the renormalizability of the theory by extending the point fermions and bosons into strings and by offering the supersymmetry among fermions and bosons. In a long series of works [1, 2, 3, 4, 5, 6, 7, 8] one of the authors in collaboration with another author and other collaborators, has found the phenomenological success with the model named the *spin-charge-family* theory with the properties: The creation and annihilation operators for fermions and bosons fields are described as tensor products of the Clifford odd (for fermions) and the Clifford even (for bosons) “basis vectors” and basis in ordinary space, explaining the second quantization postulates. The theory offers the explanation for the observed properties of fermion and bosons and for several cosmological observations. Since the number of creation and annihilation operators for fermions and bosons is in this theory the same, manifesting correspondingly a kind of supersymmetry, the authors start to study in this contribution the properties of the creation and annihilation operators if extending the point fermions and bosons into strings [14, 13], expecting that this theory offers the low energy limit for the *string theory*.

1 Introduction

In this contribution we start to study how far can the description of the internal space of fermion and boson fields with the Clifford odd and even “basis vectors” be able to reproduce the predictions of the *string theories*, provided that the creation and annihilation operators are the tensor products of the “basis vectors” and the basis in ordinary space, and if extending the points in the ordinary space to strings.

We are making the first steps in this study: We try to reproduce the internal wave function for the boson fields, represented in the “string theories” with the tensor products of the left and right movers, with the algebraic products of the Clifford odd “basis vectors” and their Hermitian conjugated partners.

Having in our case two kinds of the Clifford even “basis vectors”, ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ and ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$, we reproduce both:

- i. ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ as the algebraic product of the Clifford odd “basis vectors” $\hat{b}_{f^i}^{m'\dagger} *_A (\hat{b}_{f^i}^{m'\dagger})^\dagger$ for each of $2^{\frac{d}{2}-1}$ families of $\hat{b}_{f^i}^{m'\dagger}$ and $(\hat{b}_{f^i}^{m'\dagger})^\dagger$, f is kept unchanged;
 - ii. ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$ as the algebraic product of the Clifford odd “basis vectors” $(\hat{b}_{f^i}^{m'\dagger})^\dagger *_A \hat{b}_{f^i}^{m'\dagger}$, for each of $2^{\frac{d}{2}-1}$ members, with m kept unchanged.
- In i. each of $2^{\frac{d}{2}-1}$ families reproduces the same ${}^I\hat{\mathcal{A}}_f^{m\dagger}$, in ii. each of $2^{\frac{d}{2}-1}$ family members reproduces the same ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$.

Since in the *string theory* only bosons with the transverse momentum are considered, we study ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ and ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$ carrying only the transfer momenta.

To manifest properties of both kinds of boson field for $d = (9 + 1)$, we first discuss $d = (5 + 1)$ and then comment the case $d = (9 + 1)$.

We make first a short overview about the Clifford odd and even “basis vectors”, in any even dimensional space, describing the internal space of fermions by the Clifford odd “basis vectors” appearing in $2^{\frac{d}{2}-1}$ families, each family having $2^{\frac{d}{2}-1}$ members, and the same number $2^{\frac{d}{2}-1} \times 2^{\frac{d}{2}-1}$ of their Hermitian conjugated partners appearing in a separate group, and the internal space of bosons by the Clifford even “basis vectors” appearing in two groups each one with $2^{\frac{d}{2}-1} \times 2^{\frac{d}{2}-1}$ members), having their Hermitian conjugated partners within each group.

If the internal spaces of fermions and bosons, described by the Clifford odd and even “basis vectors”, respectively, are what the “nature” is using, then the internal space of *string theories* can be correlated to the one suggested by the Clifford algebra.

All the sections and subsections up to Subsect. 2.4 are shortly overviewing Refs. ([7, 6, 8] and the references therein) in the way needed in Subsects. (2.4, 2.5) and Sect. 3, where the authors start to present internal spaces of fermions and bosons, used in the *spin-charge-family* theory, looking for the parallelism with the internal spaces of boson strings. Presenting the boson internal space as the algebraic products of Clifford odd “basis vectors” and their Hermitian conjugated partners, they simulate left and right movers of IIA and IIB super *string theory*.

2 Creation and annihilation operators for fermions and bosons

The Clifford odd and even “basis vectors” and the creation and annihilation operators for fermions and bosons are shortly presented, overviewing Ref. [7], Sect. 2. up to 2.2.

2.1 Grassmann and Clifford algebras

The internal spaces of anti-commuting or commuting second quantized fields can be described by using either the Grassmann or the Clifford algebras [1, 6]

In Grassmann d -dimensional space there are d anti-commuting (operators) θ^a , and d anti-commuting operators which are derivatives with respect to θ^a , $\frac{\partial}{\partial\theta_a}$,

$$\begin{aligned} \{\theta^a, \theta^b\}_+ &= 0, & \left\{ \frac{\partial}{\partial\theta_a}, \frac{\partial}{\partial\theta_b} \right\}_+ &= 0, \\ \left\{ \theta_a, \frac{\partial}{\partial\theta_b} \right\}_+ &= \delta_{ab}, & (a, b) &= (0, 1, 2, 3, 5, \dots, d). \end{aligned} \tag{1}$$

Making a choice

$$(\theta^a)^\dagger = \eta^{aa} \frac{\partial}{\partial \theta_a}, \quad \text{leads to} \quad \left(\frac{\partial}{\partial \theta_a}\right)^\dagger = \eta^{aa} \theta^a, \quad (2)$$

with $\eta^{ab} = \text{diag}\{1, -1, -1, \dots, -1\}$.

θ^a and $\frac{\partial}{\partial \theta_a}$ are, up to the sign, Hermitian conjugated to each other. The identity is a self adjoint member of the algebra. The choice for the following complex properties of θ^a

$$\{\theta^a\}^* = (\theta^0, \theta^1, -\theta^2, \theta^3, -\theta^5, \theta^6, \dots, -\theta^{d-1}, \theta^d), \quad (3)$$

correspondingly requires $\left\{\frac{\partial}{\partial \theta_a}\right\}^* = \left(\frac{\partial}{\partial \theta_0}, \frac{\partial}{\partial \theta_1}, -\frac{\partial}{\partial \theta_2}, \frac{\partial}{\partial \theta_3}, -\frac{\partial}{\partial \theta_5}, \frac{\partial}{\partial \theta_6}, \dots, -\frac{\partial}{\partial \theta_{d-1}}, \frac{\partial}{\partial \theta_d}\right)$.

There are 2^d superposition of products of θ^a , the Hermitian conjugated partners of which are the corresponding superposition of products of $\frac{\partial}{\partial \theta_a}$ [10].

There exist two kinds of the Clifford algebra elements (operators), γ^a and $\tilde{\gamma}^a$, expressible with θ^a 's and their conjugate momenta $p^{\theta^a} = i \frac{\partial}{\partial \theta_a}$ [1], Eqs. (1, 2),

$$\begin{aligned} \gamma^a &= \left(\theta^a + \frac{\partial}{\partial \theta_a}\right), & \tilde{\gamma}^a &= i \left(\theta^a - \frac{\partial}{\partial \theta_a}\right), \\ \theta^a &= \frac{1}{2} (\gamma^a - i \tilde{\gamma}^a), & \frac{\partial}{\partial \theta_a} &= \frac{1}{2} (\gamma^a + i \tilde{\gamma}^a), \end{aligned} \quad (4)$$

offering together $2 \cdot 2^d$ operators: 2^d are superposition of products of γ^a and 2^d of $\tilde{\gamma}^a$. It is easy to prove if taking into account Eqs. (2, 4), that they form two anti-commuting Clifford subalgebras, $\{\gamma^a, \tilde{\gamma}^b\}_+ = 0$, Refs. ([6] and references therein)

$$\begin{aligned} \{\gamma^a, \gamma^b\}_+ &= 2\eta^{ab} = \{\tilde{\gamma}^a, \tilde{\gamma}^b\}_+, \\ \{\gamma^a, \tilde{\gamma}^b\}_+ &= 0, \quad (a, b) = (0, 1, 2, 3, 5, \dots, d), \\ (\gamma^a)^\dagger &= \eta^{aa} \gamma^a, \quad (\tilde{\gamma}^a)^\dagger = \eta^{aa} \tilde{\gamma}^a. \end{aligned} \quad (5)$$

While the Grassmann algebra offers the description of the ‘‘anti-commuting integer spin second quantized fields’’ and of the ‘‘commuting integer spin second quantized fields’’ [5, 6], the Clifford algebras which are superposition of odd products of either γ^a 's or $\tilde{\gamma}^a$'s offer the description of the second quantized half integer spin fermion fields, which from the point of the subgroups of the $SO(d-1, 1)$ group manifest spins and charges of fermions and antifermions in the fundamental representations of the group and subgroups [6].

The superposition of even products of either γ^a 's or $\tilde{\gamma}^a$'s offer the description of the commuting second quantized boson fields with integer spins (as we can see in [11] and shall see in this contribution) which from the point of the subgroups of the $SO(d-1, 1)$ group manifest spins and charges in the adjoint representations of the group and subgroups.

The following *postulate*, which determines how does $\tilde{\gamma}^a$ operate on γ^a , reduces the two Clifford subalgebras, γ^a and $\tilde{\gamma}^a$, to one, to the one described by γ^a [3, 1, 9]

$$\{\tilde{\gamma}^a B = (-)^B i B \gamma^a\} |\psi_{oc}\rangle, \quad (6)$$

with $(-)^B = -1$, if B is (a function of) odd products of γ^a 's, otherwise $(-)^B = 1$ [3], the vacuum state $|\psi_{oc}\rangle$ is defined in Eq. (14) of Subsect. 2.2.

After the postulate of Eq. (6) no vector space of $\tilde{\gamma}^a$'s needs to be taken into account for the description of the internal space of either fermions or bosons, in agreement with the observed properties of fermions and bosons. Also the Grassmann algebra is reduced to only one of the Clifford subalgebras.

The operator $\tilde{\gamma}^a$ will from now on be used to describe the properties of fermion “basis vectors”, determining by $\tilde{S}^{ab} = \frac{i}{4}(\tilde{\gamma}^a\tilde{\gamma}^b - \tilde{\gamma}^b\tilde{\gamma}^a)$ the “family” quantum numbers of the irreducible representations of the Lorentz group in internal space of fermions, S^{ab} , and the properties of bosons “basis vectors” determined by $\mathcal{S}^{ab} = S^{ab} + \tilde{S}^{ab}$. We shall see that while the fermion “basis vectors” appear in “families”, the boson “basis vectors” have no “families” and manifest properties of the gauge fields of the corresponding fermion fields.

Each irreducible representation of the Clifford odd “basis vectors” described by γ^a ’s are chosen to be equipped by the quantum numbers of the Cartan subalgebra members of \tilde{S}^{ab} , chosen in Eq. (7), as follows

$$\begin{aligned} S^{03}, S^{12}, S^{56}, \dots, S^{d-1 d}, \\ \tilde{S}^{03}, \tilde{S}^{12}, \tilde{S}^{56}, \dots, \tilde{S}^{d-1 d}, \\ \mathbf{S}^{ab} = S^{ab} + \tilde{S}^{ab} = i \left(\theta^a \frac{\partial}{\partial \theta_b} - \theta^b \frac{\partial}{\partial \theta_a} \right). \end{aligned} \quad (7)$$

2.2 “Basis vectors” and relations among Clifford even and Clifford odd “basis vectors”

This subsection is a short overview of similar sections of several articles of the author, like [12, 11, 8, 5].

After the reduction of the two Clifford subalgebras to only one, Eq. (6), we only need to define “basis vectors” for the case that the internal space of second quantized fields is described by superposition of odd or even products γ^a ’s ¹.

Let us use the technique which makes “basis vectors” products of nilpotents and projectors [1, 3] which are eigenvectors of the (chosen) Cartan subalgebra members, Eq. (7), of the Lorentz algebra in the space of γ^a ’s, either in the case of the Clifford odd or in the case of the Clifford even products of γ^a ’s.

There are in even-dimensional spaces $\frac{d}{2}$ members of the Cartan subalgebra, Eq. (7). In odd-dimensional spaces there are $\frac{d-1}{2}$ members of the Cartan subalgebra.

One finds in even dimensional spaces for any of the $\frac{d}{2}$ Cartan subalgebra member, S^{ab} , applying on a nilpotent $\binom{ab}{k}$ or on projector $[k]$

$$\begin{aligned} \binom{ab}{k}: &= \frac{1}{2}(\gamma^a + \frac{\eta^{aa}}{ik}\gamma^b), \quad (\binom{ab}{k})^2 = 0, \\ [k]: &= \frac{1}{2}(1 + \frac{i}{k}\gamma^a\gamma^b), \quad ([k])^2 = [k], \end{aligned} \quad (8)$$

the relations

$$\begin{aligned} S^{ab} \binom{ab}{k} &= \frac{k}{2} \binom{ab}{k}, & \tilde{S}^{ab} \binom{ab}{k} &= \frac{k}{2} \binom{ab}{k}, \\ S^{ab} [k] &= \frac{k}{2} [k], & \tilde{S}^{ab} [k] &= -\frac{k}{2} [k], \end{aligned} \quad (9)$$

with $k^2 = \eta^{aa}\eta^{bb}$ ², demonstrating that the eigenvalues of S^{ab} on nilpotents and projectors expressed with γ^a differ from the eigenvalues of \tilde{S}^{ab} on nilpotents and projectors expressed with γ^a , so that \tilde{S}^{ab} can be used to equip each irreducible representation of S^{ab} with the ”family” quantum number. ³

¹In Ref. [6], the reader can find in Subsects. (3.2.1 and 3.2.2) definitions for the “basis vectors” for the Grassmann and the two Clifford subalgebras, which are products of nilpotents and projectors chosen to be the eigenvectors of the corresponding Cartan subalgebra members of the Lorentz algebras presented in Eq. (7).

²Let us prove one of the relations in Eq. (9): $S^{ab} \binom{ab}{k} = \frac{i}{2}\gamma^a\gamma^b\frac{1}{2}(\gamma^a + \frac{\eta^{aa}}{ik}\gamma^b) = \frac{1}{2^2}\{-i(\gamma^a)^2\gamma^b + i(\gamma^b)^2\gamma^a\frac{\eta^{aa}}{ik}\} = \frac{1}{2}\frac{\eta^{aa}\eta^{bb}}{k}\frac{1}{2}\{\gamma^a + \frac{k^2}{\eta^{bb}ik}\gamma^b\}$. For $k^2 = \eta^{aa}\eta^{bb}$ the first relation follows.

³The reader can find the proof of Eq. (9) also in Ref. [6], App. (I).

Taking into account Eq. (5) one finds

$$\begin{aligned}
\gamma^a \begin{smallmatrix} ab \\ (k) \end{smallmatrix} &= \eta^{aa} \begin{smallmatrix} ab \\ [-k] \end{smallmatrix}, & \gamma^b \begin{smallmatrix} ab \\ (k) \end{smallmatrix} &= -ik \begin{smallmatrix} ab \\ [-k] \end{smallmatrix}, & \gamma^a \begin{smallmatrix} ab \\ [k] \end{smallmatrix} &= (-k), & \gamma^b \begin{smallmatrix} ab \\ [k] \end{smallmatrix} &= -ik\eta^{aa} \begin{smallmatrix} ab \\ (-k) \end{smallmatrix}, \\
\tilde{\gamma}^a \begin{smallmatrix} ab \\ (k) \end{smallmatrix} &= -i\eta^{aa} \begin{smallmatrix} ab \\ [k] \end{smallmatrix}, & \tilde{\gamma}^b \begin{smallmatrix} ab \\ (k) \end{smallmatrix} &= -k \begin{smallmatrix} ab \\ [k] \end{smallmatrix}, & \tilde{\gamma}^a \begin{smallmatrix} ab \\ [k] \end{smallmatrix} &= i \begin{smallmatrix} ab \\ (k) \end{smallmatrix}, & \tilde{\gamma}^b \begin{smallmatrix} ab \\ [k] \end{smallmatrix} &= -k\eta^{aa} \begin{smallmatrix} ab \\ (k) \end{smallmatrix}, \\
\begin{smallmatrix} ab \\ (k) \end{smallmatrix} \begin{smallmatrix} ab \\ (-k) \end{smallmatrix} &= \eta^{aa} \begin{smallmatrix} ab \\ [k] \end{smallmatrix}, & \begin{smallmatrix} ab \\ (-k) \end{smallmatrix} \begin{smallmatrix} ab \\ (k) \end{smallmatrix} &= \eta^{aa} \begin{smallmatrix} ab \\ [-k] \end{smallmatrix}, & \begin{smallmatrix} ab \\ (k) \end{smallmatrix} \begin{smallmatrix} ab \\ [k] \end{smallmatrix} &= 0, & \begin{smallmatrix} ab \\ (k) \end{smallmatrix} \begin{smallmatrix} ab \\ [-k] \end{smallmatrix} &= \begin{smallmatrix} ab \\ (k) \end{smallmatrix}, \\
\begin{smallmatrix} ab \\ (-k) \end{smallmatrix} \begin{smallmatrix} ab \\ [k] \end{smallmatrix} &= \begin{smallmatrix} ab \\ (-k) \end{smallmatrix}, & \begin{smallmatrix} ab \\ [k] \end{smallmatrix} \begin{smallmatrix} ab \\ (k) \end{smallmatrix} &= \begin{smallmatrix} ab \\ (k) \end{smallmatrix}, & \begin{smallmatrix} ab \\ [k] \end{smallmatrix} \begin{smallmatrix} ab \\ (-k) \end{smallmatrix} &= 0, & \begin{smallmatrix} ab \\ [k] \end{smallmatrix} \begin{smallmatrix} ab \\ [-k] \end{smallmatrix} &= 0, \\
\begin{smallmatrix} ab \\ (k) \end{smallmatrix}^\dagger &= \eta^{aa} \begin{smallmatrix} ab \\ (-k) \end{smallmatrix}, & \begin{smallmatrix} ab \\ ((k))^2 \end{smallmatrix} &= 0, & \begin{smallmatrix} ab \\ (k) \end{smallmatrix} \begin{smallmatrix} ab \\ (-k) \end{smallmatrix} &= \eta^{aa} \begin{smallmatrix} ab \\ [k] \end{smallmatrix}, \\
\begin{smallmatrix} ab \\ [k] \end{smallmatrix}^\dagger &= \begin{smallmatrix} ab \\ [k] \end{smallmatrix}, & \begin{smallmatrix} ab \\ ([k])^2 \end{smallmatrix} &= \begin{smallmatrix} ab \\ [k] \end{smallmatrix}, & \begin{smallmatrix} ab \\ [k] \end{smallmatrix} \begin{smallmatrix} ab \\ [-k] \end{smallmatrix} &= 0.
\end{aligned} \tag{10}$$

More relations are presented in App. A.

The relations in Eq. (10) demonstrate that the properties of “basis vectors” which include an odd number of nilpotents, differ essentially from the “basis vectors”, which include an even number of nilpotents.

2.3 Clifford odd and even “basis vectors”

Let us define Clifford odd and even “basis vectors” as products of nilpotents and projectors in even-dimensional spaces.

a. Clifford odd “basis vectors”

This part overviews several papers with the same topic ([6, 8] and references therein).

The Clifford odd “basis vectors” are chosen to be products of an odd number of nilpotents, and the rest, up to $\frac{d}{2}$, of projectors, each nilpotent and each projector are chosen to be the “eigenstate” of one of the members of the Cartan subalgebra, Eq. (7), correspondingly are the “basis vectors” eigenstates of all the members of the Lorentz algebra: S^{ab} 's determine $2^{\frac{d}{2}-1}$ members of one family, \tilde{S}^{ab} 's transform each member of one family to the same member of the rest of $2^{\frac{d}{2}-1}$ families.

Let us call the Clifford odd “basis vectors” $\hat{b}_f^{m\dagger}$, if it is the m^{th} membership of the family f . The Hermitian conjugated partner of $\hat{b}_f^{m\dagger}$ is called $\hat{b}_f^m (= (\hat{b}_f^{m\dagger})^\dagger)$.

Let us start in $d = 2(2n + 1)$ with the “basis vector” $\hat{b}_1^{1\dagger}$ which is the product of only nilpotents, all the rest members belonging to the $f = 1$ family follow by the application of $S^{01}, S^{03}, \dots, S^{0d}, S^{15}, \dots, S^{1d}, S^{5d}, \dots, S^{d-2d}$. They are presented on the left-hand side. Their Hermitian conjugated partners are presented on the right-hand side. The algebraic product mark $*_A$ among nilpotents and projectors is skipped.

$$\begin{aligned}
d &= 2(2n + 1), \\
\hat{b}_1^{1\dagger} &= \begin{smallmatrix} 03 & 12 & 56 & \dots & d-1d \\ (+i)(+)(+) \dots (+) \end{smallmatrix}, & \hat{b}_1^1 &= \begin{smallmatrix} 03 & 12 & \dots & d-1d \\ (-i)(-) \dots (-) \end{smallmatrix}, \\
\hat{b}_1^{2\dagger} &= \begin{smallmatrix} 03 & 12 & 56 & \dots & d-1d \\ [-i][-](+) \dots (+) \end{smallmatrix}, & \hat{b}_1^2 &= \begin{smallmatrix} 03 & 12 & 56 & \dots & d-1d \\ [-i]- \dots (-) \end{smallmatrix}, \\
\dots & & \dots & \\
\hat{b}_1^{2^{\frac{d}{2}-1}\dagger} &= \begin{smallmatrix} 03 & 12 & 56 & \dots & d-3d-2 & d-1d \\ [-i][-](+) \dots [-] [-] \end{smallmatrix}, & \hat{b}_1^{2^{\frac{d}{2}-1}} &= \begin{smallmatrix} 03 & 12 & 56 & 78 & \dots & d-3d-2 & d-1d \\ [-i]-[-] \dots [-] [-] \end{smallmatrix}, \\
\dots & & \dots &
\end{aligned} \tag{11}$$

In $d = 4n$ the choice of the starting “basis vector” with maximal number of nilpotents must have

one projector

$$\begin{aligned}
d &= 4n, \\
\hat{b}_1^{1\dagger} &= \begin{matrix} 03 & 12 & & d-1 & d \\ (+i)(+) & \cdots & & [+] \end{matrix}, & \hat{b}_1^1 &= \begin{matrix} 03 & 12 & & d-1 & d \\ (-i)(-) & \cdots & & [+] \end{matrix}, \\
\hat{b}_1^{2\dagger} &= \begin{matrix} 03 & 12 & 56 & & d-1 & d \\ [-i][-](+) & \cdots & & [+] \end{matrix}, & \hat{b}_1^2 &= \begin{matrix} 03 & 12 & 56 & & d-1 & d \\ [-i]- & \cdots & & [+] \end{matrix}, \\
\cdots, & & \cdots, & & & \\
\hat{b}_1^{2^{\frac{d}{2}-1}\dagger} &= \begin{matrix} 03 & 12 & 56 & & d-3 & d-2 & d-1 & d \\ [-i]- & \cdots & & [-] & & [+] \end{matrix}, & \hat{b}_1^{2^{\frac{d}{2}-1}} &= \begin{matrix} 03 & 12 & 56 & & d-3 & d-2 & d-1 & d \\ [-i]- & \cdots & & [-] & & [+] \end{matrix}, \\
\cdots, & & \cdots. & & & & & & (12)
\end{aligned}$$

The Hermitian conjugated partners of the Clifford odd “basis vectors” $\hat{b}_1^{m\dagger}$, presented in Eq. (12) on the right-hand side, follow if all nilpotents (k) of $\hat{b}_1^{m\dagger}$ are transformed into η^{aa} $(-k)$.

For either $d = 2(2n + 1)$ or for $d = 4n$ all the $2^{\frac{d}{2}-1}$ families follow by applying \tilde{S}^{ab} 's on all the members of the starting family. (Or one can find the starting $\hat{b}_f^{1\dagger}$ for all families f and then generate all the members \hat{b}_f^m from $\hat{b}_f^{1\dagger}$ by the application of S^{ab} on the starting member.)

It is not difficult to see that all the “basis vectors” within any family, as well as the “basis vectors” among families, are orthogonal; that is, their algebraic product is zero. The same is true within their Hermitian conjugated partners. Both can be proved by the algebraic multiplication using Eqs. (10, 23).

$$\hat{b}_f^{m\dagger} *_A \hat{b}_{f'}^{m'\dagger} = 0, \quad \hat{b}_f^m *_A \hat{b}_{f'}^{m'} = 0, \quad \forall m, m', f, f'. \quad (13)$$

When we choose the vacuum state equal to

$$|\psi_{oc}\rangle = \sum_{f=1}^{2^{\frac{d}{2}-1}} \hat{b}_f^m *_A \hat{b}_f^{m\dagger} |1\rangle, \quad (14)$$

for one of members m , which can be anyone of the odd irreducible representations f it follows that the Clifford odd “basis vectors” obey the relations

$$\begin{aligned}
\hat{b}_f^m *_A |\psi_{oc}\rangle &= 0 \cdot |\psi_{oc}\rangle, \\
\hat{b}_f^{m\dagger} *_A |\psi_{oc}\rangle &= |\psi_f^m\rangle, \\
\{\hat{b}_f^m, \hat{b}_{f'}^{m'}\} *_A |\psi_{oc}\rangle &= 0 \cdot |\psi_{oc}\rangle, \\
\{\hat{b}_f^{m\dagger}, \hat{b}_{f'}^{m'\dagger}\} *_A |\psi_{oc}\rangle &= 0 \cdot |\psi_{oc}\rangle, \\
\{\hat{b}_f^m, \hat{b}_{f'}^{m'\dagger}\} *_A |\psi_{oc}\rangle &= \delta^{mm'} \delta_{ff'} |\psi_{oc}\rangle, \quad (15)
\end{aligned}$$

b. Clifford even “basis vectors”

This part proves that the Clifford even “basis vectors” are in even-dimensional spaces offering the description of the internal spaces of boson fields — the gauge fields of the corresponding Clifford odd “basis vectors”: It is a new recognition, offering a new understanding of the second quantized fermion and **boson** fields [11].

The Clifford even “basis vectors” must be products of an even number of nilpotents and the rest, up to $\frac{d}{2}$, of projectors; each nilpotent and each projector is chosen to be the “eigenstate” of one of the members of the Cartan subalgebra of the Lorentz algebra, $\mathcal{S}^{ab} = S^{ab} + \tilde{S}^{ab}$, Eq. (7). Correspondingly the “basis vectors” are the eigenstates of all the members of the Cartan subalgebra of the Lorentz algebra.

The Clifford even “basis vectors” appear in two groups, each group has $2^{\frac{d}{2}-1} \times 2^{\frac{d}{2}-1}$ members. The members of one group can not be reached from the members of another group by either S^{ab} ’s or \tilde{S}^{ab} ’s or both.

S^{ab} and \tilde{S}^{ab} generate from the starting “basis vector” of each group all the $2^{\frac{d}{2}-1} \times 2^{\frac{d}{2}-1}$ members. Each group contains the Hermitian conjugated partner of any member; $2^{\frac{d}{2}-1}$ members of each group are products of only (self adjoint) projectors.

Let us call the Clifford even “basis vectors” ${}^i \hat{\mathcal{A}}_f^{m\dagger}$, where $i = (I, II)$ denotes the two groups of Clifford even “basis vectors”, while m and f determine membership of “basis vectors” in any of the two groups, I or II .

$$\begin{aligned}
d &= 2(2n+1) \\
{}^I \hat{\mathcal{A}}_1^{1\dagger} &= \begin{matrix} 03 & 12 & & d-1 & d \\ (+i)(+) & \cdots & & [+] & \end{matrix}, & {}^{II} \hat{\mathcal{A}}_1^{1\dagger} &= \begin{matrix} 03 & 12 & & d-1 & d \\ (-i)(+) & \cdots & & [+] & \end{matrix}, \\
{}^I \hat{\mathcal{A}}_1^{2\dagger} &= \begin{matrix} 03 & 12 & 56 & & d-1 & d \\ [-i][-](+) & \cdots & & [+] & \end{matrix}, & {}^{II} \hat{\mathcal{A}}_1^{2\dagger} &= \begin{matrix} 03 & 12 & 56 & & d-1 & d \\ [+i][-](+) & \cdots & & [+] & \end{matrix}, \\
{}^I \hat{\mathcal{A}}_1^{3\dagger} &= \begin{matrix} 03 & 12 & 56 & & d-3 & d-2 & d-1 & d \\ (+i)(+)(+) & \cdots & & [-] & (-) & \end{matrix}, & {}^{II} \hat{\mathcal{A}}_1^{3\dagger} &= \begin{matrix} 03 & 12 & 56 & & d-3 & d-2 & d-1 & d \\ (-i)(+)(+) & \cdots & & [-] & (-) & \end{matrix}, \\
&\dots & & \dots & & & & \\
d &= 4n \\
{}^I \hat{\mathcal{A}}_1^{1\dagger} &= \begin{matrix} 03 & 12 & & d-1 & d \\ (+i)(+) & \cdots & & (+) & \end{matrix}, & {}^{II} \hat{\mathcal{A}}_1^{1\dagger} &= \begin{matrix} 03 & 12 & & d-1 & d \\ (-i)(+) & \cdots & & (+) & \end{matrix}, \\
{}^I \hat{\mathcal{A}}_1^{2\dagger} &= \begin{matrix} 03 & 12 & 56 & & d-1 & d \\ [-i][-i](+) & \cdots & & (+) & \end{matrix}, & {}^{II} \hat{\mathcal{A}}_1^{2\dagger} &= \begin{matrix} 03 & 12 & 56 & & d-1 & d \\ [+i][-i](+) & \cdots & & (+) & \end{matrix}, \\
{}^I \hat{\mathcal{A}}_1^{3\dagger} &= \begin{matrix} 03 & 12 & 56 & & d-3 & d-2 & d-1 & d \\ (+i)(+)(+) & \cdots & & [-] & [-] & \end{matrix}, & {}^{II} \hat{\mathcal{A}}_1^{3\dagger} &= \begin{matrix} 03 & 12 & 56 & & d-3 & d-2 & d-1 & d \\ (-i)(+)(+) & \cdots & & [-] & [-] & \end{matrix} \\
&\dots & & \dots & & & &
\end{aligned} \tag{16}$$

There are $2^{\frac{d}{2}-1} \times 2^{\frac{d}{2}-1}$ Clifford even “basis vectors” of the kind ${}^I \hat{\mathcal{A}}_f^{m\dagger}$ and there are $2^{\frac{d}{2}-1} \times 2^{\frac{d}{2}-1}$ Clifford even “basis vectors” of the kind ${}^{II} \hat{\mathcal{A}}_f^{m\dagger}$.

Table 5, presented in App. B illustrates properties of the Clifford odd and Clifford even “basis vectors” on the case of $d = (5 + 1)$. Looking at this case it is easy to evaluate properties of either even or odd “basis vectors”. We shall discuss in this subsection the general case by carefully inspecting properties of both kinds of “basis vectors”.

Two “basis vectors”, ${}^i \hat{\mathcal{A}}_f^{m\dagger}$ and ${}^i \hat{\mathcal{A}}_{f'}^{m'\dagger}$, the algebraic product, $*_A$, of which gives non zero contribution, “scatter” into the third one ${}^i \hat{\mathcal{A}}_{f''}^{m''\dagger}$, for $i = (I, II)$.

It remains to evaluate the algebraic application, $*_A$, of the Clifford even “basis vectors” ${}^{I,II} \hat{\mathcal{A}}_f^{m\dagger}$ on the Clifford odd “basis vectors” $\hat{b}_{f'}^{m'\dagger}$. One finds, taking into account Eq. (23), for ${}^I \hat{\mathcal{A}}_f^{m\dagger}$

$${}^I \hat{\mathcal{A}}_f^{m\dagger} *_A \hat{b}_{f'}^{m'\dagger} \rightarrow \begin{cases} \hat{b}_{f''}^{m''\dagger}, \\ \text{or zero,} \end{cases} \tag{17}$$

For each ${}^I \hat{\mathcal{A}}_f^{m\dagger}$ there are among $2^{\frac{d}{2}-1} \times 2^{\frac{d}{2}-1}$ members of the Clifford odd “basis vectors” (describing the internal space of fermion fields) $2^{\frac{d}{2}-1}$ members, $\hat{b}_{f'}^{m'\dagger}$, fulfilling the relation of Eq. (17). All the rest $(2^{\frac{d}{2}-1} \times (2^{\frac{d}{2}-1} - 1))$ Clifford odd “basis vectors” give zero contributions. Or equivalently, there are $2^{\frac{d}{2}-1}$ pairs of quantum numbers (f, m') for which $\hat{b}_{f'}^{m'\dagger} \neq 0$.

Taking into account Eq. (23) one finds

$$\hat{b}_{f'}^{m'\dagger} *_A {}^I \hat{\mathcal{A}}_{f''}^{m''\dagger} = 0, \quad \forall (m, m', f, f'). \tag{18}$$

For “scattering” the Clifford even “basis vectors” ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$ on the Clifford odd “basis vectors” $\hat{b}_{f'}^{m'\dagger}$ it follows

$${}^{II}\hat{\mathcal{A}}_f^{m\dagger} *_A \hat{b}_{f'}^{m'\dagger} = 0, \quad \forall(m, m', f, f'), \quad (19)$$

while we get

$$\hat{b}_f^{m\dagger} *_A {}^{II}\hat{\mathcal{A}}_{f'}^{m'\dagger} \rightarrow \begin{cases} \hat{b}_{f''}^{m\dagger}, \\ \text{or zero,} \end{cases} \quad (20)$$

For each $\hat{b}_f^{m\dagger}$ there are among $2^{\frac{d}{2}-1} \times 2^{\frac{d}{2}-1}$ members of the Clifford even “basis vectors” (describing the internal space of boson fields), ${}^{II}\hat{\mathcal{A}}_{f'}^{m'\dagger}$, $2^{\frac{d}{2}-1}$ members (with appropriate f' and m') fulfilling the relation of Eq. (20).

All the rest ($2^{\frac{d}{2}-1} \times (2^{\frac{d}{2}-1} - 1)$) Clifford even “basis vectors” give zero contributions.

2.4 Symmetry relations between Clifford even and Clifford odd “basis vectors”

This part is, together with Subsect. 2.5 and Sect. 3, the only new part of this contribution. All the rest is the repetition of Refs. [7, 6, 8] and the references therein, needed for Subsects. (2.4, 2.5) and Sect. 3.

In Subsect. 2.2 the relations among the Clifford even “basis vectors”, ${}^i\hat{\mathcal{A}}_f^{m\dagger}$, $i = (I, II)$, as well as among the Clifford even ${}^i\hat{\mathcal{A}}_f^{m\dagger}$, $i = (I, II)$ and odd “basis vectors”, $\hat{b}_f^{m\dagger}$ and \hat{b}_f^m , are discussed.

This, the only new part up to now, looks for relations among the properties of *boson strings* (IIA and IIB) description of the internal space and the description of the internal space with the Clifford even “basis vectors”. We mostly treat the particular case of $d = (5 + 1)$. The case with $d = (9 + 1)$ is comment in the next subsection 2.5.

Let us look for the Clifford even “basis vectors”, ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ and ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$, which have in the internal space spin in transverse dimensions only, in order to relate “basis vectors” to the closed IIA and IIB *boson strings*. One finds ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ as the algebraic products of the Clifford odd “basis vectors” and their Hermitian conjugated partners, $\hat{b}_f^{m\dagger} *_A (\hat{b}_f^{m'\dagger})^\dagger$, for each of $2^{\frac{d}{2}-1}$ families, obtaining the same group of ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ for any f , while ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$ can be written as the algebraic products of the Clifford odd “basis vectors” and their hermitian conjugated partners in the opposite order, $(\hat{b}_{f'}^{m'\dagger})^\dagger *_A \hat{b}_{f'}^{m'\dagger}$, for each of $2^{\frac{d}{2}-1}$ family members, while f' and f “run over all the families. Again any m gives the same group of ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$.

We look for the relations among the internal spaces defined in *strings theory* and the internal spaces defined with the Clifford odd and even “basis vectors”, used in the *spin-charge-family* theory for only those fields ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ and ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$ with $\mathcal{S}^{12} = \pm 1$.

In this Subsect. we pay attention on:

A. How one can generate Clifford even “basis vectors” from the algebraic products, $*_A$, of the Clifford odd $\hat{b}_f^{m\dagger}$ and their Hermitian conjugated partners $\hat{b}_{f'}^{m'\dagger} = (\hat{b}_{f'}^{m'\dagger})^\dagger$.

B. How γ^a and $\tilde{\gamma}^a$ apply on the Clifford even and odd basis vectors”.

2.4.1 Generating Clifford even “basis vectors” from the algebraic products, $*_A$, of the Clifford odd “basis vectors”, $\hat{b}_f^{m\dagger}$, and their Hermitian conjugated partners, $\hat{b}_{f'}^{m'}$.

Let us start with the generation of the Clifford even “basis vectors” for ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ from the algebraic products of the Clifford odd “basis vectors” and their Hermitian conjugated partners, $\hat{b}_f^{m\dagger} *_A (\hat{b}_f^{m'\dagger})^\dagger$, for a particular case when $d = (5 + 1)$ to learn what we can expect after comparing these relations with the similar relations in the *string theory*.

And let us treat ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$ as the algebraic product, $(\hat{b}_f^{m\dagger})^\dagger *_A \hat{b}_f^{m\dagger}$, again for the case when $d = (5 + 1)$. The generalization to any even d is not difficult.

We shall comment our way of describing internal space and the *string theory* way for $d = (9 + 1)$ in Subject. 2.5.

We need in this $d = (5 + 1)$ case Tables 1, 2, 3 from Ref. [7] and Table 2 from the contribution of N.S.M.B. in this proceedings. Table 1 is repeated in App. B, Tables 2, 3 are repeated in this subsection.

In Table 1, and Table 2 (presented in the contribution of N.S.M.B. in this proceeding), the Clifford even “basis vectors” are presented as products of an even number of nilpotents $\binom{ab}{(k)}$, two or zero, and the rest of projectors $\binom{ab}{[k]}$, one or three, Eqs. (8, 10).

The “basis vectors” have in $d = (5+1)$ 16 members $(2^{\frac{d}{2}-1} \times 2^{\frac{d}{2}-1})$, which are chosen to be eigenvectors of all the Cartan subalgebra members, Eq. (7). These Clifford even “basis vectors” are either self adjoint (when they are products of only projectors, Eq. (10)), or they have their Hermitian conjugated partners within the same group.

We see in both above mentioned tables (1 and 2, presented in the contribution of the N.S.M.B. in this proceeding) that we can split the Clifford even “basis vectors”, either ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ or ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$, into two groups:

One group has the eigenvalues of $\mathcal{S}^{12} = \pm 1$. This represents bosons with internal momentum in transverse dimensions, $\mathcal{S}^{12} = \pm 1$. One notices that the members with $\mathcal{S}^{12} = 1$ have their Hermitian conjugated partners among those with $\mathcal{S}^{12} = -1$, and opposite. These “basis vectors” can transfer to fermions, the internal space of which is described by the Clifford odd “basis vectors”, their $\mathcal{S}^{12} = 1$ or -1 , respectively.

The second group has the eigenvalues of $\mathcal{S}^{12} = 0$. They have their Hermitian conjugated partners among those with $\mathcal{S}^{12} = 0$ (and \mathcal{S}^{03} and \mathcal{S}^{56} equal to ± 1) or are self adjoint (they are products of projectors only).

Table 2 represents the Clifford odd “basis vectors”, used in the *spin-charge-family* theory to describe internal space of fermions. The Clifford odd “basis vectors” have in $d = (5+1)$ 4 members $(2^{\frac{d}{2}-1})$ in each of 4 $(2^{\frac{d}{2}-1})$ families, presented in Table (2, 5) which are chosen to be the eigenvectors of all the Cartan subalgebra members, Eq. (7). Their Hermitian conjugated partners are presented (together with the Clifford odd and both Clifford even “basis vectors”) in Table 5. The Clifford odd “basis vectors”, and their Hermitian conjugated partners — they appear in a separate group — are presented as products of an odd number of nilpotents $\binom{ab}{(k)}$, three or one, and the rest of projectors $\binom{ab}{[k]}$, zero or two, Eqs. (8, 10).

We need to know the structure of the Clifford odd “basis vectors” and their Hermitian conjugated partners to be able to understand the description of the boson “basis vectors”, ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ and ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$, as the algebraic products of the Clifford odd “basis vectors” $\hat{b}_f^{m\dagger}$ and their Hermitian conjugated partners $\hat{b}_{f'}^{m'} = (\hat{b}_f^{m'\dagger})^\dagger$.

One can obtain the Clifford even “basis vectors”, ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ and ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$, as algebraic products of the

Table 1: This table is taken from Ref. [7] and the references therein. The Clifford even “basis vectors” ${}^I\hat{\mathcal{A}}_f^{m\dagger}$, each of them is the product of projectors and an even number of nilpotents, and each is the eigenvector of all the Cartan subalgebra members, \mathcal{S}^{03} , \mathcal{S}^{12} , \mathcal{S}^{56} , Eq. (7), are presented for $d = (5 + 1)$ -dimensional case. Indexes m and f determine $2^{\frac{d}{2}-1} \times 2^{\frac{d}{2}-1}$ different members ${}^I\hat{\mathcal{A}}_f^{m\dagger}$. In the third column the “basis vectors” ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ which are Hermitian conjugated partners to each other (and can therefore annihilate each other) are pointed out with the same symbol. For example, with $\star\star$ are equipped the first member with $m = 1$ and $f = 1$ and the last member of $f = 3$ with $m = 4$. The sign \bigcirc denotes the Clifford even “basis vectors” which are self-adjoint ($({}^I\hat{\mathcal{A}}_f^{m\dagger})^\dagger = {}^I\hat{\mathcal{A}}_f^{m\dagger}$). (It is obvious that \dagger has no meaning, since ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ are self adjoint or are Hermitian conjugated partner to another ${}^I\hat{\mathcal{A}}_{f'}^{m'\dagger}$.) This table also presents the eigenvalues of the three commuting operators τ^3, τ^8 and τ' of the subgroups $SU(3) \times U(1)$.

f	m	*	${}^I\hat{\mathcal{A}}_f^{m\dagger}$	\mathcal{S}^{03}	\mathcal{S}^{12}	\mathcal{S}^{56}	τ^3	τ^8	τ'
I	1	$\star\star$	$\begin{matrix} 03 & 12 & 56 \\ [+i] & (+) & (+) \end{matrix}$	0	1	1	$-\frac{1}{2}$	$-\frac{1}{2\sqrt{3}}$	$-\frac{2}{3}$
	2	\triangle	$\begin{matrix} 03 & 12 & 56 \\ (-i) & [-] & (+) \end{matrix}$	$-i$	0	1	$-\frac{1}{2}$	$-\frac{3}{2\sqrt{3}}$	0
	3	\ddagger	$\begin{matrix} 03 & 12 & 56 \\ (-i) & (+) & [-] \end{matrix}$	$-i$	1	0	-1	0	0
	4	\bigcirc	$\begin{matrix} 03 & 12 & 56 \\ [+i] & [-] & [-] \end{matrix}$	0	0	0	0	0	0
II	1	\bullet	$\begin{matrix} 03 & 12 & 56 \\ (+i) & [+] & (+) \end{matrix}$	i	0	1	$\frac{1}{2}$	$-\frac{1}{2\sqrt{3}}$	$-\frac{2}{3}$
	2	\otimes	$\begin{matrix} 03 & 12 & 56 \\ [-i] & (-) & (+) \end{matrix}$	0	-1	1	$\frac{1}{2}$	$-\frac{3}{2\sqrt{3}}$	0
	3	\bigcirc	$\begin{matrix} 03 & 12 & 56 \\ [-i] & [+] & [-] \end{matrix}$	0	0	0	0	0	0
	4	\ddagger	$\begin{matrix} 03 & 12 & 56 \\ (+i) & (-) & [-] \end{matrix}$	i	-1	0	1	0	0
III	1	\bigcirc	$\begin{matrix} 03 & 12 & 56 \\ [+i] & [+] & [+] \end{matrix}$	0	0	0	0	0	0
	2	$\bigcirc\bigcirc$	$\begin{matrix} 03 & 12 & 56 \\ (-i) & (-) & [+] \end{matrix}$	$-i$	-1	0	0	$-\frac{1}{\sqrt{3}}$	$\frac{2}{3}$
	3	\bullet	$\begin{matrix} 03 & 12 & 56 \\ (-i) & [+] & (-) \end{matrix}$	$-i$	0	-1	$-\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{2}{3}$
	4	$\star\star$	$\begin{matrix} 03 & 12 & 56 \\ [+i] & (-) & (-) \end{matrix}$	0	-1	-1	$\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{2}{3}$
IV	1	$\bigcirc\bigcirc$	$\begin{matrix} 03 & 12 & 56 \\ (+i) & (+) & [+] \end{matrix}$	i	1	0	0	$\frac{1}{\sqrt{3}}$	$-\frac{2}{3}$
	2	\bigcirc	$\begin{matrix} 03 & 12 & 56 \\ [-i] & [-] & [+] \end{matrix}$	0	0	0	0	0	0
	3	\otimes	$\begin{matrix} 03 & 12 & 56 \\ [-i] & (+) & (-) \end{matrix}$	0	1	-1	$-\frac{1}{2}$	$\frac{3}{2\sqrt{3}}$	0
	4	\triangle	$\begin{matrix} 03 & 12 & 56 \\ (+i) & [-] & (-) \end{matrix}$	i	0	-1	$\frac{1}{2}$	$\frac{3}{2\sqrt{3}}$	0

Table 2: This table is taken from Ref. [7]. The "basis vectors" $\hat{b}_f^{m\dagger}$ are presented for $d = (5 + 1)$ -dimensional case. Each $\hat{b}_f^{m\dagger}$ is a product of projectors and of an odd number of nilpotents and is the "eigenvector" of all the Cartan subalgebra members, (S^{03}, S^{12}, S^{56}) and $(\tilde{S}^{03}, \tilde{S}^{12}, \tilde{S}^{56})$, Eq. (7), m marks the members of each family, while f determines the family quantum numbers (the eigenvalues of $(\tilde{S}^{03}, \tilde{S}^{12}, \tilde{S}^{56})$). This table also presents in the columns ($8^{th}, 9^{th}, 10^{th}$) the eigenvalues of the three commuting operators (τ^3, τ^8 and τ') of the subgroups $SU(3) \times U(1)$, ($\tau^3 := \frac{1}{2}(-S^{12} - iS^{03}), \tau^8 = \frac{1}{2\sqrt{3}}(-iS^{03} + S^{12} - 2S^{56}), \tau' = -\frac{1}{3}(-iS^{03} + S^{12} + S^{56})$), as well as (in the last three columns) the corresponding ($\tilde{\tau}^3, \tilde{\tau}^8, \tilde{\tau}'$). $\Gamma^{(3+1)} = i\gamma^0\gamma^1\gamma^2\gamma^3$ is written in the 7^{th} column. $\Gamma^{(5+1)} = -1 (= -\gamma^0\gamma^1\gamma^2\gamma^3\gamma^5\gamma^6)$. Operators $\hat{b}_f^{m\dagger}$ and \hat{b}_f^m fulfil the anti-commutation relations for the second quantized fermion fields.

f	m	$\hat{b}_f^{m\dagger}$	S^{03}	S^{12}	S^{56}	Γ^{3+1}	τ^3	τ^8	τ'	\tilde{S}^{03}	\tilde{S}^{12}	\tilde{S}^{56}	$\tilde{\tau}^3$	$\tilde{\tau}^8$	$\tilde{\tau}'$
<i>I</i>	1	$\begin{matrix} 03 & 12 & 56 \\ (+i) & (+) & & (+) \end{matrix}$	$\frac{i}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	0	0	$-\frac{1}{2}$	$\frac{i}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{1}{6}$
	2	$\begin{matrix} 03 & 12 & 56 \\ [-i] & (-) & & (+) \end{matrix}$	$-\frac{i}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	1	0	$-\frac{1}{\sqrt{3}}$	$\frac{1}{6}$	$\frac{i}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{1}{6}$
	3	$\begin{matrix} 03 & 12 & 56 \\ [-i] & (+) & & (-) \end{matrix}$	$-\frac{i}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{1}{6}$	$\frac{i}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{1}{6}$
	4	$\begin{matrix} 03 & 12 & 56 \\ (+i) & (-) & & (-) \end{matrix}$	$\frac{i}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	-1	$\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{1}{6}$	$\frac{i}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{1}{6}$
<i>II</i>	1	$\begin{matrix} 03 & 12 & 56 \\ [+i] & (+) & & (+) \end{matrix}$	$\frac{i}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	0	0	$-\frac{1}{2}$	$-\frac{i}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	0	$-\frac{1}{\sqrt{3}}$	$\frac{1}{6}$
	2	$\begin{matrix} 03 & 12 & 56 \\ (-i) & (-) & & (+) \end{matrix}$	$-\frac{i}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	1	0	$-\frac{1}{\sqrt{3}}$	$\frac{1}{6}$	$-\frac{i}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	0	$-\frac{1}{\sqrt{3}}$	$\frac{1}{6}$
	3	$\begin{matrix} 03 & 12 & 56 \\ (-i) & (+) & & (-) \end{matrix}$	$-\frac{i}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{1}{6}$	$-\frac{i}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	0	$-\frac{1}{\sqrt{3}}$	$\frac{1}{6}$
	4	$\begin{matrix} 03 & 12 & 56 \\ [+i] & (-) & & (-) \end{matrix}$	$\frac{i}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	-1	$\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{1}{6}$	$-\frac{i}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	0	$-\frac{1}{\sqrt{3}}$	$\frac{1}{6}$
<i>III</i>	1	$\begin{matrix} 03 & 12 & 56 \\ [+i] & (+) & & (+) \end{matrix}$	$\frac{i}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	0	0	$-\frac{1}{2}$	$-\frac{i}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{1}{6}$
	2	$\begin{matrix} 03 & 12 & 56 \\ (-i) & (-) & & (+) \end{matrix}$	$-\frac{i}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	1	0	$-\frac{1}{\sqrt{3}}$	$\frac{1}{6}$	$-\frac{i}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{1}{6}$
	3	$\begin{matrix} 03 & 12 & 56 \\ (-i) & (+) & & (-) \end{matrix}$	$-\frac{i}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{1}{6}$	$-\frac{i}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{1}{6}$
	4	$\begin{matrix} 03 & 12 & 56 \\ [+i] & (-) & & (-) \end{matrix}$	$\frac{i}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	-1	$\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{1}{6}$	$-\frac{i}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{1}{6}$
<i>IV</i>	1	$\begin{matrix} 03 & 12 & 56 \\ (+i) & (+) & & (+) \end{matrix}$	$\frac{i}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	0	0	$-\frac{1}{2}$	$\frac{i}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	0	0	$-\frac{1}{2}$
	2	$\begin{matrix} 03 & 12 & 56 \\ [-i] & (-) & & (+) \end{matrix}$	$-\frac{i}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	1	0	$-\frac{1}{\sqrt{3}}$	$\frac{1}{6}$	$\frac{i}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	0	0	$-\frac{1}{2}$
	3	$\begin{matrix} 03 & 12 & 56 \\ [-i] & (+) & & (-) \end{matrix}$	$-\frac{i}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{1}{6}$	$\frac{i}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	0	0	$-\frac{1}{2}$
	4	$\begin{matrix} 03 & 12 & 56 \\ (+i) & (-) & & (-) \end{matrix}$	$\frac{i}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	-1	$\frac{1}{2}$	$\frac{1}{2\sqrt{3}}$	$\frac{1}{6}$	$\frac{i}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	0	0	$-\frac{1}{2}$

Clifford odd “basis vectors” and their Hermitian conjugated partners, ${}^I\hat{\mathcal{A}}_f^{m\dagger} = \hat{b}_{f^i}^{m'\dagger} *_A (\hat{b}_{f^i}^{m''\dagger})^\dagger$, while ${}^{II}\hat{\mathcal{A}}_f^{m\dagger} = (\hat{b}_{f^i}^{m'\dagger})^\dagger *_A \hat{b}_{f^i}^{m''\dagger}$. One can check that $\hat{b}_{f^i}^{m'\dagger} *_A (\hat{b}_{f^i}^{m''\dagger})^\dagger$ applying on $\hat{b}_{f^i}^{m''\dagger}$ obey Eq. (17) and that $\hat{b}_{f^i}^{m''\dagger}$ applying on $(\hat{b}_{f^i}^{m'\dagger})^\dagger *_A \hat{b}_{f^i}^{m''\dagger}$ obey Eq. (20).

We see in Table 3 and in Table 4 the same number, $8 = \frac{1}{2}2^{\frac{d-6}{2}-1} \times 2^{\frac{d-6}{2}-1}$, of the Clifford even “basis vectors” ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ and ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$, each with $\mathcal{S}^{12} = \pm 1$: Four of each of them have $\mathcal{S}^{12} = +1$ and four have $\mathcal{S}^{12} = -1$.

If the handedness of the Clifford odd “basis vectors” is chosen to be the right handedness, $\Gamma^{(d)} = \prod_a (\sqrt{\eta^{aa}} \gamma^a) \cdot \begin{cases} (i)^{\frac{d}{2}}, & \text{for } d \text{ even,} \\ (i)^{\frac{d-1}{2}}, & \text{for } d \text{ odd,} \end{cases}$ then their Hermitian conjugated partners have left handedness for either $\mathcal{S}^{12} = +1$ and $\mathcal{S}^{12} = -1$, resembling left and right movers contributing to boson strings in *string theories* AII and BII.

Let us point out that to both groups of the Clifford even “basis vectors” appearing in Table 3 and in Table 4 all family members m and all families f contribute:

- a. For ${}^I\hat{\mathcal{A}}_f^{m\dagger}$, all the family members m for a particular family f and their Hermitian conjugated partners contribute in $\hat{b}_{f^i}^{m'\dagger} *_A (\hat{b}_{f^i}^{m''\dagger})^\dagger$, using only half of possibilities, the other half contribute to $\mathcal{S}^{12} = 0$. Each family f' of $\hat{b}_{f^i}^{m'\dagger} *_A (\hat{b}_{f^i}^{m''\dagger})^\dagger$ generates the same eight Clifford even ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ as are the ones presented in Table 3 for $f^i = 1$.
- b. For ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$, all the families f of a particular member m and their Hermitian conjugated partners contribute in $(\hat{b}_{f^i}^{m'\dagger})^\dagger *_A \hat{b}_{f^i}^{m''\dagger}$, using only half of possibilities, the other half contribute to $\mathcal{S}^{12} = 0$. Each family member m' generates in $(\hat{b}_{f^i}^{m'\dagger})^\dagger *_A \hat{b}_{f^i}^{m''\dagger}$ the same eight Clifford even ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$ as are the ones presented in Table 4 for $m' = 1$.

2.4.2 How γ^a and $\tilde{\gamma}^a$ apply on Clifford odd and even “basis vectors”

Table 5, taken from Ref. [6], represents Clifford odd and Clifford even “basis vectors”. One easily sees that the algebraic multiplication of any Clifford odd “basis vector” by γ^a leads to the corresponding Clifford even “basis vector” ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$, in agreement with Eq. (10). The algebraic multiplication of any Clifford odd “basis vector” by $\tilde{\gamma}^a$ leads to the corresponding Clifford even “basis vector” ${}^I\hat{\mathcal{A}}_f^{m\dagger}$.

Multiplying the first column of *odd* ${}^I\hat{\mathcal{A}}_1^{m\dagger}$ in Table 5 by γ^0 : $\gamma^0 *_A \hat{b}_1^{m\dagger}$ leads to the third column of *even* ${}^{II}\hat{\mathcal{A}}_3^{m\dagger}$, according to Eq. (10).

Multiplying the first column of *odd* ${}^I\hat{\mathcal{A}}_1^{m\dagger}$ in Table 5 by $\tilde{\gamma}^0$: $\tilde{\gamma}^0 *_A \hat{b}_1^{m\dagger}$ leads to the third column of *even* ${}^I\hat{\mathcal{A}}_3^{m\dagger}$, according to Eq. (10).

The opposite way, the algebraic multiplication of any Clifford even “basis vector” ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$ by γ^a leads to the corresponding Clifford odd “basis vector” $\hat{b}_{f^i}^{m\dagger}$.

Multiplying the first column of *even* ${}^{II}\hat{\mathcal{A}}_1^{m\dagger}$ in Table 5 by γ^0 , for example, $\gamma^0 *_A {}^{II}\hat{\mathcal{A}}_1^{m\dagger}$, leads to the fourth column of *odd* ${}^I\hat{\mathcal{A}}_1^{m\dagger}$, according to Eq. (10).

And similarly if multiplying any Clifford even “basis vector” ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ by $\tilde{\gamma}^a$ leads to the corresponding Clifford odd “basis vector” $\hat{b}_{f^i}^{m\dagger}$.

These resemble a kind of supersymmetry: The same number of the Clifford odd and the Clifford even “basis vectors”, and the simple relations between fermions and bosons.

Table 3: The Clifford even “basis vectors” $I\hat{\mathcal{A}}_f^{m\dagger}$, belonging to transverse momentum in internal space, \mathcal{S}^{12} equal to 1, the first half $I\hat{\mathcal{A}}_f^{m\dagger}$, and -1 , the second half $I\hat{\mathcal{A}}_f^{m\dagger}$, for $d = (5 + 1)$, are presented as algebraic products of the $f = 1$ family “basis vectors” $\hat{b}_1^{m'\dagger}$ and their Hermitian conjugated partners $(\hat{b}_1^{m''\dagger})^\dagger$: $\hat{b}_1^{m'\dagger} *_A (\hat{b}_1^{m''\dagger})^\dagger$. The Hermitian conjugated partners of two $I\hat{\mathcal{A}}_f^{m\dagger}$ are marked with the same symbol. The Clifford even “basis vectors” $I\hat{\mathcal{A}}_f^{m\dagger}$ are the product of one projector and two nilpotents, the Clifford odd “basis vectors” and their Hermitian conjugated partners are products of one nilpotent and two projectors or of three nilpotents. The Clifford even and Clifford odd objects are eigenvectors of all the corresponding Cartan subalgebra members, Eq. (7). There are $2^{\frac{6}{2}-1} \times 2^{\frac{6}{2}-1}$ algebraic products of $\hat{b}_1^{m'\dagger} *_A (\hat{b}_1^{m''\dagger})^\dagger$. The rest 8 of 16 members present $I\hat{\mathcal{A}}_f^{m\dagger}$ with $\mathcal{S}^{12} = 0$. The members $\hat{b}_f^{m'\dagger}$ together with their Hermitian conjugated partners of each of the four families, $f = (1, 2, 3, 4)$, offers the same $I\hat{\mathcal{A}}_f^{m\dagger}$ with $\mathcal{S}^{12} = \pm 1$ as the ones presented in this table. (And equivalently for $\mathcal{S}^{12} = 0$.)

\mathcal{S}^{12}	symbol	$I\hat{\mathcal{A}}_f^{m\dagger} =$	$\hat{b}_f^{m'\dagger} *_A (\hat{b}_f^{m''\dagger})^\dagger$
1	**	$I\hat{\mathcal{A}}_1^{1\dagger} =$	$\hat{b}_1^{1\dagger} *_A (\hat{b}_1^{4\dagger})^\dagger$ $\begin{matrix} 03 & 12 & 56 \\ [+i] & (+) & (+) \end{matrix}$ $\begin{matrix} 03 & 12 & 56 \\ (+i) & [+][+] & *_A \end{matrix}$ $\begin{matrix} 03 & 12 & 56 \\ (-i) & (+) & (+) \end{matrix}$
1	‡	$I\hat{\mathcal{A}}_1^{3\dagger} =$	$\hat{b}_1^{3\dagger} *_A (\hat{b}_1^{4\dagger})^\dagger$ $\begin{matrix} 03 & 12 & 56 \\ (-i) & (+) & [-] \end{matrix}$ $\begin{matrix} 03 & 12 & 56 \\ [-i] & [+][(-) & *_A \end{matrix}$ $\begin{matrix} 03 & 12 & 56 \\ (-i) & (+) & (+) \end{matrix}$
1	⊙⊙	$I\hat{\mathcal{A}}_4^{1\dagger} =$	$\hat{b}_1^{1\dagger} *_A (\hat{b}_1^{2\dagger})^\dagger$ $\begin{matrix} 03 & 12 & 56 \\ (+i) & (+) & [+] \begin{matrix} 03 & 12 & 56 \\ (+i) & [+][+] & *_A \end{matrix} \begin{matrix} 03 & 12 & 56 \\ [-i] & (+) & [+]$
1	⊗	$I\hat{\mathcal{A}}_4^{3\dagger} =$	$\hat{b}_1^{3\dagger} *_A (\hat{b}_1^{2\dagger})^\dagger$ $\begin{matrix} 03 & 12 & 56 \\ [-i] & (+) & (-) \end{matrix}$ $\begin{matrix} 03 & 12 & 56 \\ [-i] & [+][(-) & *_A \end{matrix}$ $\begin{matrix} 03 & 12 & 56 \\ [-i] & (+) & [+]$
-1	⊗	$I\hat{\mathcal{A}}_2^{2\dagger} =$	$\hat{b}_1^{2\dagger} *_A (\hat{b}_1^{3\dagger})^\dagger$ $\begin{matrix} 03 & 12 & 56 \\ [-i] & (-) & (+) \end{matrix}$ $\begin{matrix} 03 & 12 & 56 \\ [-i] & (-) & [+] \begin{matrix} 03 & 12 & 56 \\ [-i] & [+][+] \end{matrix}$
-1	‡	$I\hat{\mathcal{A}}_2^{4\dagger} =$	$\hat{b}_1^{4\dagger} *_A (\hat{b}_1^{3\dagger})^\dagger$ $\begin{matrix} 03 & 12 & 56 \\ (+i) & (-) & [-] \end{matrix}$ $\begin{matrix} 03 & 12 & 56 \\ (+i) & (-) & (-) \end{matrix}$ $\begin{matrix} 03 & 12 & 56 \\ [-i] & [+][+] \end{matrix}$
-1	⊙⊙	$I\hat{\mathcal{A}}_3^{2\dagger} =$	$\hat{b}_1^{2\dagger} *_A (\hat{b}_1^{1\dagger})^\dagger$ $\begin{matrix} 03 & 12 & 56 \\ (-i) & (-) & [+] \begin{matrix} 03 & 12 & 56 \\ [-i] & (-) & [+] \begin{matrix} 03 & 12 & 56 \\ (-i) & [+][+] \end{matrix}$
-1	**	$I\hat{\mathcal{A}}_3^{4\dagger} =$	$\hat{b}_1^{4\dagger} *_A (\hat{b}_1^{1\dagger})^\dagger$ $\begin{matrix} 03 & 12 & 56 \\ [+i] & (-) & (-) \end{matrix}$ $\begin{matrix} 03 & 12 & 56 \\ (+i) & (-) & (-) \end{matrix}$ $\begin{matrix} 03 & 12 & 56 \\ (-i) & [+][+] \end{matrix}$

Table 4: The Clifford even “basis vectors” ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$, belonging to transverse momentum in internal space, \mathcal{S}^{12} equal to 1, the first half ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$, and -1 , the second half ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$, for $d = (5 + 1)$, are presented as algebraic products of the first, $m = 1$, member of “basis vectors” $\hat{b}_{f'}^{m'=1\dagger}$ and the Hermitian conjugated partners $(\hat{b}_{f''}^{m'=1\dagger})^\dagger$. The Hermitian conjugated partners of two ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$ are marked with the same symbol. The Clifford even “basis vectors” ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$ are the product of one projector and two nilpotents, the Clifford odd “basis vectors” and the Hermitian conjugated partners are products of one nilpotent and two projectors or of three nilpotents. Clifford even and Clifford odd objects are eigenvectors all the corresponding Cartan subalgebra members, Eq. (7). There are $2^{\frac{6}{2}-1} \times 2^{\frac{6}{2}-1}$ algebraic products of $\hat{b}_{f'}^{m'\dagger} *_A (\hat{b}_{f''}^{m''\dagger})^\dagger$, f' and f'' run over all four families. The rest of 16 members presents ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$ with $\mathcal{S}^{12} = 0$. The members $\hat{b}_{f'}^{m'\dagger}$ together with $(\hat{b}_{f''}^{m''\dagger})^\dagger$ $m' = (1, 2, 3, 4)$, offers the same ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ with $\mathcal{S}^{12} = \pm 1$ as the ones presented in this table. (And equivalently for $\mathcal{S}^{12} = 0$.)

\mathcal{S}^{12}	symbol	${}^{II}\hat{\mathcal{A}}_f^{m\dagger} =$	$(\hat{b}_{f'}^{1\dagger})^\dagger *_A \hat{b}_{f''}^{1\dagger}$
1	**	${}^{II}\hat{\mathcal{A}}_1^{1\dagger} =$	$(\hat{b}_1^{1\dagger})^\dagger *_A \hat{b}_4^{1\dagger}$ $\begin{matrix} 03 & 12 & 56 \\ [-i] & (+) & (+) \end{matrix} \quad \begin{matrix} 03 & 12 & 56 \\ (-i) & [+][+] & *_A \end{matrix} \quad \begin{matrix} 03 & 12 & 56 \\ (+i) & (+) & (+) \end{matrix}$
1	⊙⊙	${}^{II}\hat{\mathcal{A}}_1^{3\dagger} =$	$(\hat{b}_2^{1\dagger})^\dagger *_A \hat{b}_4^{1\dagger}$ $\begin{matrix} 03 & 12 & 56 \\ (+i) & (+) & [-] \end{matrix} \quad \begin{matrix} 03 & 12 & 56 \\ [+i] & [+][(-) & *_A \end{matrix} \quad \begin{matrix} 03 & 12 & 56 \\ (+i) & (+) & (+) \end{matrix}$
1	‡	${}^{II}\hat{\mathcal{A}}_4^{1\dagger} =$	$(\hat{b}_1^{1\dagger})^\dagger *_A \hat{b}_3^{1\dagger}$ $\begin{matrix} 03 & 12 & 56 \\ (-i) & (+) & [+]$
1	⊗	${}^{II}\hat{\mathcal{A}}_4^{3\dagger} =$	$(\hat{b}_2^{1\dagger})^\dagger *_A \hat{b}_3^{1\dagger}$ $\begin{matrix} 03 & 12 & 56 \\ [+i] & (+) & (-) \end{matrix} \quad \begin{matrix} 03 & 12 & 56 \\ [+i] & [+][(-) & *_A \end{matrix} \quad \begin{matrix} 03 & 12 & 56 \\ [+i] & (+) & [+]$
-1	⊗	${}^{II}\hat{\mathcal{A}}_2^{2\dagger} =$	$(\hat{b}_3^{1\dagger})^\dagger *_A \hat{b}_2^{1\dagger}$ $\begin{matrix} 03 & 12 & 56 \\ [+i] & (-) & (+) \end{matrix} \quad \begin{matrix} 03 & 12 & 56 \\ [+i] & (-) & [+]$
-1	⊗⊗	${}^{II}\hat{\mathcal{A}}_2^{4\dagger} =$	$(\hat{b}_4^{1\dagger})^\dagger *_A \hat{b}_2^{1\dagger}$ $\begin{matrix} 03 & 12 & 56 \\ (-i) & (-) & [-] \end{matrix} \quad \begin{matrix} 03 & 12 & 56 \\ (-i) & (-) & (-) \end{matrix} \quad \begin{matrix} 03 & 12 & 56 \\ [+i] & [+][+] \end{matrix}$
-1	‡	${}^{II}\hat{\mathcal{A}}_3^{2\dagger} =$	$(\hat{b}_3^{1\dagger})^\dagger *_A \hat{b}_1^{1\dagger}$ $\begin{matrix} 03 & 12 & 56 \\ (+i) & (-) & [+]$
-1	**	${}^{II}\hat{\mathcal{A}}_3^{4\dagger} =$	$(\hat{b}_4^{1\dagger})^\dagger *_A \hat{b}_1^{1\dagger}$ $\begin{matrix} 03 & 12 & 56 \\ [-i] & (-) & (-) \end{matrix} \quad \begin{matrix} 03 & 12 & 56 \\ (-i) & (-) & (-) \end{matrix} \quad \begin{matrix} 03 & 12 & 56 \\ (+i) & [+][+] \end{matrix}$




Figure 1: This figure is taken from Ref. [7]. The representations of the subgroups $SU(3)$ and $U(1)$ of the group $SO(5, 1)$, the properties of which appear in Tables (5, 2) for the Clifford odd “basis vectors”, are presented. (τ^3, τ^8, τ') can be calculated if using the relations $\tau^3 = \frac{1}{2}(-S^{12} - iS^{03})$, $\tau^8 = \frac{1}{2\sqrt{3}}(S^{12} - iS^{03} - 2S^{56})$, $\tau' = -\frac{1}{3}(S^{12} - iS^{03} + S^{56})$. On the abscissa axis, on the ordinate axis and on the third axis, the eigenvalues of the superposition of the three Cartan subalgebra members, (τ^3, τ^8, τ') , are presented. One notices one triplet, denoted by \bigcirc with the values $\tau' = \frac{1}{6}$, $(\tau^3 = -\frac{1}{2}, \tau^8 = \frac{1}{2\sqrt{3}}, \tau' = \frac{1}{6})$, $(\tau^3 = \frac{1}{2}, \tau^8 = \frac{1}{2\sqrt{3}}, \tau' = \frac{1}{6})$, $(\tau^3 = 0, \tau^8 = -\frac{1}{\sqrt{3}}, \tau' = \frac{1}{6})$, respectively, and one singlet denoted by the square. $(\tau^3 = 0, \tau^8 = 0, \tau' = -\frac{1}{2})$. The triplet and the singlet appear in four families, with the family quantum numbers presented in the last three columns of Table 2.

It is useful to point out again, that the Clifford odd and the Clifford even “basis vectors” differ essentially in their properties:

- i. While the Clifford odd “basis vectors” in even dimensional spaces appear in $2^{\frac{d}{2}-1}$ families, each family having $2^{\frac{d}{2}-1}$ members, and have their Hermitian conjugated partners in a separate group, again with $2^{\frac{d}{2}-1} \times 2^{\frac{d}{2}-1}$ contributions, appear the Clifford even “basis vectors” in even dimensional spaces in two groups, each with $2^{\frac{d}{2}-1} \times 2^{\frac{d}{2}-1}$ members, having the Hermitian conjugated partners within the same group. They have no families.*
- ii. The Clifford odd “basis vectors” in even dimensional spaces carry the eigenvalues of the Cartan subalgebra members, Eq. (7), $\pm\frac{i}{2}$ or $\pm\frac{1}{2}$. The Clifford even “basis vectors” in even dimensional spaces carry the eigenvalues of the Cartan subalgebra members, Eq. (7), $(\pm i, 0)$ or $(\pm 1, 0)$.*

The two figures, Fig. 1 and Fig. 2. manifest the differences among the Clifford odd and the Clifford even “basis vectors”.

In the case that the group $SO(5, 1)$ — manifesting as $SU(3) \times U(1)$ and representing the colour group with quantum numbers (τ^3, τ^8) and the “fermion” group with the quantum number τ' — is embedded into $SO(13, 1)$ the triplet would represent quarks (and antiquarks), and the singlet leptons (and antileptons).

The corresponding gauge fields, presented in Table 1 and Fig. 2, if belonging to the sextet, would transform the triplet of quarks among themselves, changing the colour and leaving the “fermion” quantum number equal to $\frac{1}{6}$.

We can see that $I\hat{\mathcal{A}}_3^{m\dagger}$ with $(m = 2, 3, 4)$, if applied on the $SU(3)$ singlet $\hat{b}_4^{1\dagger}$ with $(\tau' = -\frac{1}{2}, \tau^3 = 0, \tau^8 = 0)$, transforms it to $\hat{b}_4^{m(=2,3,4)\dagger}$, respectively, which are members of the $SU(3)$ triplet. All these Clifford even “basis vectors” have τ' equal to $\frac{2}{3}$, changing correspondingly $\tau' = -\frac{1}{2}$ into $\tau' = \frac{1}{6}$ and bringing the needed values of τ^3 and τ^8 .




Figure 2: This figure is taken from Ref. [7]. The Clifford even "basis vectors" ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ in the case that $d = (5+1)$ are presented by the eigenvalues of the commuting operators of the subgroups $SU(3)$ and $U(1)$ of the group $SO(5, 1)$: $(\tau^3 = \frac{1}{2}(-\mathcal{S}^{12} - i\mathcal{S}^{03}), \tau^8 = \frac{1}{2\sqrt{3}}(\mathcal{S}^{12} - i\mathcal{S}^{03} - 2\mathcal{S}^{56}), \tau' = -\frac{1}{3}(\mathcal{S}^{12} - i\mathcal{S}^{03} + \mathcal{S}^{56}))$. Their properties appear in Table 1. The abscissa axis carries the eigenvalues of τ^3 , the ordinate axis carries the eigenvalues of τ^8 and the third axis carries the eigenvalues of τ' . One notices four singlets with $(\tau^3 = 0, \tau^8 = 0, \tau' = 0)$, denoted by \bigcirc , representing four self adjoint Clifford even "basis vectors" ${}^I\hat{\mathcal{A}}_f^{m\dagger}$, with $(f = 1, m = 4), (f = 2, m = 3), (f = 3, m = 1), (f = 4, m = 2)$, one sextet of three pairs, Hermitian conjugated to each other, with $\tau' = 0$, denoted by \triangle (${}^I\hat{\mathcal{A}}_1^{2\dagger}$ with $(\tau' = 0, \tau^3 = -\frac{1}{2}, \tau^8 = -\frac{3}{2\sqrt{3}})$ and ${}^I\hat{\mathcal{A}}_4^{4\dagger}$ with $(\tau' = 0, \tau^3 = \frac{1}{2}, \tau^8 = \frac{3}{2\sqrt{3}})$), by \dagger (${}^I\hat{\mathcal{A}}_1^{3\dagger}$ with $(\tau' = 0, \tau^3 = -1, \tau^8 = 0)$ and ${}^I\hat{\mathcal{A}}_2^{4\dagger}$ with $\tau' = 0, \tau^3 = 1, \tau^8 = 0$), and by \otimes (${}^I\hat{\mathcal{A}}_2^{2\dagger}$ with $(\tau' = 0, \tau^3 = \frac{1}{2}, \tau^8 = -\frac{3}{2\sqrt{3}})$ and ${}^I\hat{\mathcal{A}}_4^{3\dagger}$ with $(\tau' = 0, \tau^3 = -\frac{1}{2}, \tau^8 = \frac{3}{2\sqrt{3}})$), and one triplet, denoted by $\star\star$ (${}^I\hat{\mathcal{A}}_3^{4\dagger}$ with $(\tau' = \frac{2}{3}, \tau^3 = \frac{1}{2}, \tau^8 = \frac{1}{2\sqrt{3}})$), by \bullet (${}^I\hat{\mathcal{A}}_3^{3\dagger}$ with $(\tau' = \frac{2}{3}, \tau^3 = -\frac{1}{2}, \tau^8 = \frac{1}{2\sqrt{3}})$), and by $\odot\odot$ (${}^I\hat{\mathcal{A}}_3^{2\dagger}$ with $(\tau' = \frac{2}{3}, \tau^3 = 0, \tau^8 = -\frac{1}{\sqrt{3}})$), as well as one antitriplet, Hermitian conjugated to triplet, denoted by $\star\star$ (${}^I\hat{\mathcal{A}}_1^{1\dagger}$ with $(\tau' = -\frac{2}{3}, \tau^3 = -\frac{1}{2}, \tau^8 = -\frac{1}{2\sqrt{3}})$), by \bullet (${}^I\hat{\mathcal{A}}_2^{1\dagger}$ with $(\tau' = -\frac{2}{3}, \tau^3 = \frac{1}{2}, \tau^8 = -\frac{1}{2\sqrt{3}})$), and by $\odot\odot$ (${}^I\hat{\mathcal{A}}_1^{4\dagger}$ with $(\tau' = -\frac{2}{3}, \tau^3 = 0, \tau^8 = \frac{1}{\sqrt{3}})$).

In Table 1 we find $(6 + 4)$ Clifford even “basis vectors” ${}^I \hat{\mathcal{A}}_f^{m\dagger}$ with $\tau' = 0$. Six of them are Hermitian conjugated to each other — the Hermitian conjugated partners are denoted by the same geometric figure on the third column. Four of them are self-adjoint and correspondingly with $(\tau' = 0, \tau^3 = 0, \tau^8 = 0)$, denoted in the third column of Table 1 by \bigcirc . The rest 6 Clifford even “basis vectors” belong to one triplet with $\tau' = \frac{2}{3}$ and (τ^3, τ^8) equal to $[(0, -\frac{1}{\sqrt{3}}), (-\frac{1}{2}, \frac{1}{2\sqrt{3}}), (\frac{1}{2}, \frac{1}{2\sqrt{3}})]$ and one antitriplet with $\tau' = -\frac{2}{3}$ and (τ^3, τ^8) equal to $[(-\frac{1}{2}, -\frac{1}{2\sqrt{3}}), (\frac{1}{2}, -\frac{1}{2\sqrt{3}}), (0, \frac{1}{\sqrt{3}})]$.

Each triplet has Hermitian conjugated partners in anti-triplet and opposite. In Table 1 the Hermitian conjugated partners of the triplet and antitriplet are denoted by the same signum: $({}^I \hat{\mathcal{A}}_1^{1\dagger}, {}^I \hat{\mathcal{A}}_3^{4\dagger})$ by $\star\star$, $({}^I \hat{\mathcal{A}}_2^{1\dagger}, {}^I \hat{\mathcal{A}}_3^{3\dagger})$ by \bullet , and $({}^I \hat{\mathcal{A}}_3^{2\dagger}, {}^I \hat{\mathcal{A}}_4^{1\dagger})$ by $\odot\odot$.

The octet, two triplets and four singlets are presented in Fig. 2.

The same Fig. 2 represents also the Clifford even ${}^{II} \hat{\mathcal{A}}_f^{m\dagger}$, although their expressions with the products of two nilpotents and one projector or three projectors, presented on Table 5, differ from ${}^I \hat{\mathcal{A}}_f^{m\dagger}$.

The way of application of the Clifford even “basis vectors” ${}^{II} \hat{\mathcal{A}}_f^{m\dagger}$ on $\hat{b}_f^{m'\dagger}$ follow Eqs. (19, 20), while the application of ${}^I \hat{\mathcal{A}}_f^{m\dagger}$ on $\hat{b}_f^{m'\dagger}$ follow Eqs. (17, 18).

iii. In odd dimensional spaces, $d = 2n + 1$, only half of “basis vectors” demonstrate the properties which they demonstrate in even dimensional spaces, the properties which empower the Clifford odd “basis vectors” to describe the internal space of fermion fields and the Clifford even “basis vectors” to describe the internal space of the corresponding gauge fields: This half belongs to $d' = 2n$ and does demonstrate these properties. The other half, obtained from the first half by the application of S^{02n+1} , this second half of the Clifford odd “basis vectors”, although anticommuting, demonstrate properties of the Clifford even “basis vectors”, and the second half of the Clifford even “basis vectors”, although commuting, demonstrate properties of the Clifford odd “basis vectors” in even dimensional spaces: The still anticommuting Clifford odd “basis vectors” (the Clifford even operators S^{02n+1} do not change either oddness or evenness of the “basis vectors”) appear in two separate groups with $2^{\frac{2n}{2}-1} \times 2^{\frac{2n}{2}-1}$, each with their Hermitian conjugated partners within the same group having no families; and still commuting Clifford even “basis vectors” appear in $2^{\frac{2n}{2}-1}$ families, each with $2^{\frac{2n}{2}-1}$ members, having their Hermitian conjugated partners $2^{\frac{2n}{2}-1} \times 2^{\frac{2n}{2}-1}$ in a separate group.

These second half of “basis vectors” obviously resemble properties of the internal spaces of the ghost scalar fields, used in the quantum field theory to make contributions of the Feynman diagrams finite, Ref. [8].

2.5 Relations between $SO(9, 1)$ description of the internal spaces in *string theories* and with the Clifford even and odd “basis vectors”

In this subsection the first step is done to compare the properties of the Clifford odd and the Clifford even “basis vectors” with the properties of the *string theories* in the case of $d = (9 + 1)$.

Let us first repeat what we learned about the Clifford even and the Clifford odd “basis vectors” in even dimensional spaces.

There are in $d = 2(2n + 1)$ and $d = 4n$ dimensional spaces $2^{\frac{d}{2}-1}$ Clifford odd families, each family having $2^{\frac{d}{2}-1}$ members. The Clifford odd “basis vectors” have their Hermitian conjugated partners in a separate group of $2^{\frac{d}{2}-1}$ families with $2^{\frac{d}{2}-1}$ members. In a tensor product with the basis in ordinary space the Clifford odd “basis vectors” together with their Hermitian conjugated partners form creations and annihilation operators, which fulfil on the vacuum state the postulates of the second quantized fermion fields, manifesting all the properties of the fermion fields [7], Eq. (34) and Subsect. 2.4.

There are in even dimensional spaces two times $2^{\frac{d}{2}-1} \times 2^{\frac{d}{2}-1}$ Clifford even basis vectors, with their Hermitian conjugated partners within the same group. In a tensor product with the basis in ordinary space the Clifford even “basis vectors” form creations and annihilation operators, which fulfil the postulates of the second quantized boson fields, manifesting all the properties of the boson fields to the corresponding fermion fields [7], Subsect. 2.4.

Both are represented by the points in the ordinary space.

Let us generalise to any even d what we learned about the description of the Clifford even ‘basis vectors’ with the algebraic products of the Clifford odd “basis vectors” and their Hermitian conjugated partners in Subsect. 2.4 when we treated the case $d = (5 + 1)$.

Looking at the properties of both kinds of the Clifford even “basis vectors”, ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ and ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$, manifesting momentum in only transverse dimensions (with \mathcal{S}^{ab} not equal \mathcal{S}^{03}), we found in Subsect. 2.4, in Table 3 and in Table 4 that to both groups of the Clifford even “basis vectors” all family members m and all families f contribute:

- a. To ${}^I\hat{\mathcal{A}}_f^{m\dagger}$, all the family members m for a particular family f and their Hermitian conjugated partners contribute in $\hat{b}_{f'}^{m'\dagger} *_{A} (\hat{b}_{f'}^{m''\dagger})^\dagger$, using only half of possibilities ($\frac{1}{2} \times 2^{\frac{d}{2}-1} \times 2^{\frac{d}{2}-1}$), the other half possibilities contribute to $\mathcal{S}^{12} = 0$. Each family f' of $\hat{b}_{f'}^{m'\dagger} *_{A} (\hat{b}_{f'}^{m''\dagger})^\dagger$ generates the same eight Clifford even ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ as are the ones presented in Table 3 for $f' = 1$.
- b. To ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$, all the families f' of a particular member m' and their Hermitian conjugated partners contribute in $(\hat{b}_{f'}^{m'\dagger})^\dagger *_{A} \hat{b}_{f'}^{m'\dagger}$, using only half of possibilities, the other half contribute to $\mathcal{S}^{12} = 0$. Each family member m' generates in $(\hat{b}_{f'}^{m'\dagger})^\dagger *_{A} \hat{b}_{f'}^{m'\dagger}$ the same eight Clifford even ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$ as are the ones presented in Table 4 for $m' = 1$.

Let us now try to relate the description of the internal spaces of bosons described by the Clifford even “basis vectors”, as explained in Subsect. 2.4 and the way how *string theories* consider the IIA or IIB superstring model for closed RNS strings as presented in Ref. [15] in Subsect. 11.6.3, where the spectrum is described.

We find in $d = (9 + 1)$, according to what it is discussed in this subsection, in the case that we are interested only on those internal degrees of freedom of the Clifford even basis vectors of each of the two kinds, ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ and ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$, which manifest momentum in only transverse dimensions (with \mathcal{S}^{ab} not equal \mathcal{S}^{03}), $\frac{1}{2} \times 2^{\frac{d=10}{2}-1} \times 2^{\frac{d=10}{2}-1} = 8 \times 16 = 128$ of ${}^I\hat{\mathcal{A}}_f^{m\dagger}$ and 128 of ${}^{II}\hat{\mathcal{A}}_f^{m\dagger}$, together 256 of both kinds of the Clifford even “basis vectors”, representing the boson fields. These are also possibilities suggested in Ref. [15] for closed strings in $d = (9 + 1)$; for the left-right movers or right-left movers forming the closed boson strings of AII and BII kind, manifesting the momentum in only transverse dimensions they found 256 possibilities.

3 Conclusions

Let be repeated: In a long series of works [1, 2, 3, 4, 5, 6, 7, 8] one of the authors in collaboration with another author and other collaborators, has found the phenomenological success with the model named the *spin-charge-family* theory with the properties: The creation and annihilation operators for fermion and boson fields are described as tensor products of the Clifford odd (for fermions) and the Clifford even (for bosons) “basis vectors” and basis in ordinary space, explaining the second quantization postulates. The theory, assuming in $d \geq (13 + 1)$ a simple starting action with fermions which interact with gravity only, offers the explanation for all the assumptions of the *standard model* before the electroweak break: For the properties of quarks and leptons and antiquarks and antileptons, for the corresponding gauge fields, for the appearance of the scalar higgs and Yukawa couplings, as well as for several cosmological

observations, making several predictions.

Since the *strings theories* are offering the way for explaining so far observed fermion and boson second quantized fields, with gravity included, by offering the renormalizability of the theory by extending the point fermions and bosons into strings and by offering the supersymmetry among fermions and bosons, we expect that in the low energy limit the *string theories* coincide with our predictions provided that we extend points in the ordinary space to strings, hoping that this would help to solve the problem of renormalisability of the *spin-charge-family* theory.

What we present in this contribution is the first step towards relating the spin-charge-family theory with the *string theories*. This first step is promising: We represent the two groups of the Clifford even “basis vectors”, $I\hat{\mathcal{A}}_f^{m\dagger}$ and $II\hat{\mathcal{A}}_f^{m\dagger}$, as the algebraic products of the Clifford odd “basis vectors” and their Hermitian conjugated partners.

The products $\hat{b}_{f'}^{m'\dagger} *_A (\hat{b}_{f_i}^{m''\dagger})^\dagger$ represent $I\hat{\mathcal{A}}_f^{m\dagger}$ and $(\hat{b}_{f'}^{m'\dagger})^\dagger *_A \hat{b}_{f_i}^{m''\dagger}$ represent $II\hat{\mathcal{A}}_f^{m\dagger}$. The discussions in Subsect. 2.4 and Subsect.2.5 demonstrate that in both cases *all the families and family members participate* when describing the Clifford even “basis vectors”, either $I\hat{\mathcal{A}}_f^{m\dagger}$ or $II\hat{\mathcal{A}}_f^{m\dagger}$, in the case that both boson fields demonstrate spins in the transverse dimensions only ($\mathcal{S}^{03} = 0$, the rest of Cartan subalgebra members \mathcal{S}^{ab} , Eq.(7), are in general non zero and can represent $\mathcal{S}^{12} = \pm 1$ as well as charges in adjoint representations). But only half products of $\hat{b}_{f_i}^{m'\dagger}$ and $(\hat{b}_{f_i}^{m''\dagger})^\dagger$ are involved, the other half determine the cases when \mathcal{S}^{03} is nonzero.

In Subsect. 2.5, we find in the case $d = (9 + 1)$, and that we are interested only on those internal degrees of freedom of the Clifford even basis vectors of each of the two kinds, $I\hat{\mathcal{A}}_f^{m\dagger}$ and $II\hat{\mathcal{A}}_f^{m\dagger}$, which manifest momentum in only transverse dimensions (with \mathcal{S}^{ab} not all equal to $\mathcal{S}^{03} = 0$), $\frac{1}{2} \times 2^{\frac{d-10}{2}-1} \times 2^{\frac{d-10}{2}-1} = 8 \times 16 = 128$ possibilities of $I\hat{\mathcal{A}}_f^{m\dagger}$ and 128 possibilities of $II\hat{\mathcal{A}}_f^{m\dagger}$, together 256 members of both kinds of the Clifford even “basis vectors”, representing the boson fields. These are also possibilities suggested in Ref. [15] for closed strings in $d = (9 + 1)$.

*The algebraic products $\hat{b}_{f_i}^{m'\dagger} *_A (\hat{b}_{f_i}^{m''\dagger})^\dagger$ can be interpreted as left and right movers and $(\hat{b}_{f_i}^{m'\dagger})^\dagger *_A \hat{b}_{f_i}^{m''\dagger}$ as right and left movers of the closed boson strings.*

Let us add that the relation among $SO(n)$ and all the rest groups is known for a long time [1, 17, 16], but not in this way as presented in this paper and in [7, 8]. This way offers, to our understanding, the new way towards understanding the second quantized fermion and boson fields, with the gravity included.

Still a hard work is needed to make the next step towards relating the *string theories* and the *spin-charge-family* theory. But since the description of the internal spaces of fermion and boson fields with the Clifford odd and Clifford even “basis vectors”, respectively, is simple and well defined, it might bring a new understanding of the theory of our world.

A Useful relations

This appendix, taken from Ref. [7], contains some helpful relations needed for the reader of this paper. For more detailed explanations and for proofs, the reader is kindly asked to read [6] and the references therein.

For fermions, the operator of handedness Γ^d is determined as follows:

$$\Gamma^{(d)} = \prod_a (\sqrt{\eta^{aa}} \gamma^a) \cdot \begin{cases} (i)^{\frac{d}{2}}, & \text{for } d \text{ even,} \\ (i)^{\frac{d-1}{2}}, & \text{for } d \text{ odd,} \end{cases} \quad (21)$$

The vacuum state for the Clifford odd "basis vectors", $|\psi_{oc}\rangle$, is defined as

$$|\psi_{oc}\rangle = \sum_{f=1}^{2^{\frac{d}{2}-1}} \hat{b}_f^m *_A \hat{b}_f^{m\dagger} |1\rangle. \quad (22)$$

Taking into account that the Clifford objects γ^a and $\tilde{\gamma}^a$ fulfil relations of Eq. 5, one obtains beside the relations presented in Eq. (10) some new ones.

$$\begin{aligned} \overset{ab}{(k)}\overset{ab}{(-k)} &= \eta^{aa} \overset{ab}{[k]}, & \overset{ab}{(-k)}\overset{ab}{(k)} &= \eta^{aa} \overset{ab}{[-k]}, & \overset{ab}{(k)}\overset{ab}{[k]} &= 0, & \overset{ab}{(k)}\overset{ab}{[-k]} &= \overset{ab}{(k)}, \\ \overset{ab}{(-k)}\overset{ab}{[k]} &= \overset{ab}{(-k)}, & \overset{ab}{[k]}\overset{ab}{(k)} &= \overset{ab}{(k)}, & \overset{ab}{[k]}\overset{ab}{(-k)} &= 0, & \overset{ab}{[k]}\overset{ab}{[-k]} &= 0, \\ \overset{ab}{(\tilde{k})}\overset{ab}{(k)} &= 0, & \overset{ab}{(\tilde{k})}\overset{ab}{(-k)} &= -i\eta^{aa} \overset{ab}{[-k]}, & \overset{ab}{(-k)}\overset{ab}{(k)} &= -i\eta^{aa} \overset{ab}{[k]}, & \overset{ab}{(\tilde{k})}\overset{ab}{[k]} &= i \overset{ab}{(k)}, \\ \overset{ab}{(\tilde{k})}\overset{ab}{[-k]} &= 0, & \overset{ab}{(-k)}\overset{ab}{[k]} &= 0, & \overset{ab}{(-k)}\overset{ab}{[-k]} &= i \overset{ab}{(-k)}, & \overset{ab}{[\tilde{k}]}\overset{ab}{(k)} &= \overset{ab}{(\tilde{k})}, \\ \overset{ab}{[\tilde{k}]}\overset{ab}{(-k)} &= 0, & \overset{ab}{[\tilde{k}]}\overset{ab}{[k]} &= 0, & \overset{ab}{[-k]}\overset{ab}{[k]} &= \overset{ab}{[k]}, & \overset{ab}{[\tilde{k}]}\overset{ab}{[-k]} &= \overset{ab}{[-k]}, \end{aligned} \quad (23)$$

The algebraic multiplication among $\overset{ab}{(\tilde{k})}$ and $\overset{ab}{[\tilde{k}]}$ goes as in the case of $\overset{ab}{(k)}$ and $\overset{ab}{[k]}$

$$\begin{aligned} \overset{ab}{(\tilde{k})}\overset{ab}{[\tilde{k}]} &= 0, & \overset{ab}{[\tilde{k}]}\overset{ab}{(\tilde{k})} &= \overset{ab}{(\tilde{k})}, & \overset{ab}{(\tilde{k})}\overset{ab}{[-k]} &= \overset{ab}{(k)}, & \overset{ab}{[\tilde{k}]}\overset{ab}{(-k)} &= 0, \\ \overset{ab}{(-k)}\overset{ab}{(\tilde{k})} &= \eta^{aa} \overset{ab}{[-k]}, & \overset{ab}{(-k)}\overset{ab}{[-k]} &= 0, \end{aligned} \quad (24)$$

One can further find that

$$\begin{aligned} S^{ac} \overset{ab}{(k)}\overset{cd}{(k)} &= -\frac{i}{2}\eta^{aa}\eta^{cc} \overset{ab}{[-k]}\overset{cd}{[-k]}, & S^{ac} \overset{ab}{[k]}\overset{cd}{[k]} &= \frac{i}{2} \overset{ab}{(-k)}\overset{cd}{(-k)}, \\ S^{ac} \overset{ab}{(k)}\overset{cd}{[k]} &= -\frac{i}{2}\eta^{aa} \overset{ab}{[-k]}\overset{cd}{(-k)}, & S^{ac} \overset{ab}{[k]}\overset{cd}{(k)} &= \frac{i}{2}\eta^{cc} \overset{ab}{(-k)}\overset{cd}{[-k]}. \end{aligned} \quad (25)$$

B Some properties of the Clifford odd and even "basis vectors" are presented

In this App. the properties of the Clifford odd and Clifford even "basis vectors" for the case that $d = (5 + 1)$ are presented in the table 5, taken from Ref. [7].

Table 5 presents the 64 ($= 2^{d=6}$) "eigenvectors" of the Cartan subalgebra members of the Lorentz algebra, S^{ab} and \mathcal{S}^{ab} , Eq. (7).

The Clifford odd "basis vectors" — they appear in 4 ($= 2^{\frac{d=6}{2}-1}$) families, each family has 4 members — are products of an odd number of nilpotents, either of three or one. They appear in the group named *odd I* $\hat{b}_f^{m\dagger}$. Their Hermitian conjugated partners appear in the second group named *odd II* \hat{b}_f^m . Within each of these two groups the members are mutually orthogonal (which can be checked by using Eq. (23)); $\hat{b}_f^{m\dagger} *_A \hat{b}_{f'}^{m'\dagger} = 0$ for all (m, m', f, f') . Equivalently, $\hat{b}_f^m *_A \hat{b}_{f'}^{m'} = 0$ for all (m, m', f, f') .

The Clifford even "basis vectors" are products of an even number of nilpotents — of either two or none in this case. They are presented in Table 5 in two groups, each with 16 ($= 2^{\frac{d=6}{2}-1} \times 2^{\frac{d=6}{2}-1}$) members, as *even I* $\mathcal{A}_f^{m\dagger}$ and *even II* $\mathcal{A}_f^{m\dagger}$. One can easily check, using Eq. (23), that the algebraic product $I \mathcal{A}_f^{m\dagger} *_A II \mathcal{A}_{f'}^{m'\dagger} = 0 = II \mathcal{A}_f^{m\dagger} *_A I \mathcal{A}_{f'}^{m'\dagger}, \forall (m, m', f, f')$, Eq. (??). An overview of the Clifford

Table 5: $2^d = 64$ “eigenvectors” of the Cartan subalgebra of the Clifford odd and even algebras — the superposition of odd and even products of γ^a 's — in $d = (5 + 1)$ -dimensional space are presented, divided into four groups. The first group, *odd I*, is chosen to represent “basis vectors”, named $\hat{b}_f^{m\dagger}$, appearing in $2^{\frac{d}{2}-1} = 4$ “families” ($f = 1, 2, 3, 4$), each ”family” with $2^{\frac{d}{2}-1} = 4$ “family” members ($m = 1, 2, 3, 4$). The second group, *odd II*, contains Hermitian conjugated partners of the first group for each family separately, $\hat{b}_f^m = (\hat{b}_f^{m\dagger})^\dagger$. Either *odd I* or *odd II* are products of an odd number of nilpotents (one or three) and projectors (two or none). The “family” quantum numbers of $\hat{b}_f^{m\dagger}$, that is the eigenvalues of $(\tilde{S}^{03}, \tilde{S}^{12}, \tilde{S}^{56})$, are for the first *odd I* group appearing above each “family”, the quantum numbers of the family members (S^{03}, S^{12}, S^{56}) are written in the last three columns. For the Hermitian conjugated partners of *odd I*, presented in the group *odd II*, the quantum numbers (S^{03}, S^{12}, S^{56}) are presented above each group of the Hermitian conjugated partners, the last three columns tell eigenvalues of $(\tilde{S}^{03}, \tilde{S}^{12}, \tilde{S}^{56})$. The two groups with the even number of γ^a 's, *even I* and *even II*, each group has their Hermitian conjugated partners within its group, have the quantum numbers f , that is the eigenvalues of $(\tilde{S}^{03}, \tilde{S}^{12}, \tilde{S}^{56})$, written above column of four members, the quantum numbers of the members, (S^{03}, S^{12}, S^{56}) , are written in the last three columns. To find the quantum numbers of $(\mathbf{S}^{03}, \mathbf{S}^{12}, \mathbf{S}^{56})$ one has to take into account that $\mathbf{S}^{ab} = S^{ab} + \tilde{S}^{ab}$.

“basis vectors” $(\tilde{S}^{03}, \tilde{S}^{12}, \tilde{S}^{56})$	m →	$f = 1$ $(\frac{i}{2}, -\frac{1}{2}, -\frac{1}{2})$	$f = 2$ $(-\frac{i}{2}, -\frac{1}{2}, \frac{1}{2})$	$f = 3$ $(-\frac{i}{2}, \frac{1}{2}, -\frac{1}{2})$	$f = 4$ $(\frac{i}{2}, \frac{1}{2}, \frac{1}{2})$	S^{03}	S^{12}	S^{56}
<i>odd I</i> $\hat{b}_f^{m\dagger}$	1	03 12 56 (+i)[+][+]	03 12 56 [+i][+][+]	03 12 56 [+i][+][+]	03 12 56 (+i)(+)(+)	$\frac{i}{2}$	$\frac{1}{2}$	$\frac{1}{2}$
	2	[-i](-)[+]	(-i)(-)(+)	(-i)[-][+]	[-i](-)(+)	$-\frac{i}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$
	3	[-i][+][-]	(-i)[+][-]	(-i)(+)(-)	[-i][+][-]	$-\frac{i}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$
	4	(+i)(-)(-)	[+i](-)[-]	[+i](-)[-]	(+i)[-][-]	$\frac{i}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$
(S^{03}, S^{12}, S^{56})	→	$(-\frac{i}{2}, \frac{1}{2}, \frac{1}{2})$ 03 12 56	$(\frac{i}{2}, \frac{1}{2}, -\frac{1}{2})$ 03 12 56	$(\frac{i}{2}, -\frac{1}{2}, \frac{1}{2})$ 03 12 56	$(-\frac{i}{2}, -\frac{1}{2}, -\frac{1}{2})$ 03 12 56	S^{03}	S^{12}	S^{56}
<i>odd II</i> \hat{b}_f^m	1	(-i)[+][+]	[+i][+][-]	[+i](-)[+]	(-i)(-)(-)	$-\frac{i}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$
	2	[-i](+)[+]	(+i)(+)(-)	(+i)[-][+]	[-i](-)(-)	$\frac{i}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$
	3	[-i][+][+]	(+i)[+][-]	(+i)(-)(+)	[-i](-)[-]	$-\frac{i}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$
	4	(-i)(+)(+)	[+i](+)[-]	[+i](-)[+]	(-i)[-][-]	$-\frac{i}{2}$	$\frac{1}{2}$	$\frac{1}{2}$
$(\tilde{S}^{03}, \tilde{S}^{12}, \tilde{S}^{56})$	→	$(-\frac{i}{2}, \frac{1}{2}, \frac{1}{2})$ 03 12 56	$(\frac{i}{2}, -\frac{1}{2}, \frac{1}{2})$ 03 12 56	$(-\frac{i}{2}, -\frac{1}{2}, -\frac{1}{2})$ 03 12 56	$(\frac{i}{2}, \frac{1}{2}, -\frac{1}{2})$ 03 12 56	S^{03}	S^{12}	S^{56}
<i>even I</i> \mathcal{A}_f^m	1	[+i](+)(+)	(+i)+	[+i][+][+]	(+i)(+)[+]	$\frac{i}{2}$	$\frac{1}{2}$	$\frac{1}{2}$
	2	(-i)[-](+)	[-i](-)(+)	(-i)(-)[+]	[-i](-)[+]	$-\frac{i}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$
	3	(-i)(+)[-]	[-i][+][-]	(-i)[+][-]	[-i](+)(-)	$-\frac{i}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$
	4	[+i](-)[-]	(+i)(-)[-]	[+i](-)(-)	(+i)[-][-]	$\frac{i}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$
$(\tilde{S}^{03}, \tilde{S}^{12}, \tilde{S}^{56})$	→	$(\frac{i}{2}, \frac{1}{2}, \frac{1}{2})$ 03 12 56	$(-\frac{i}{2}, -\frac{1}{2}, \frac{1}{2})$ 03 12 56	$(\frac{i}{2}, -\frac{1}{2}, -\frac{1}{2})$ 03 12 56	$(-\frac{i}{2}, \frac{1}{2}, -\frac{1}{2})$ 03 12 56	S^{03}	S^{12}	S^{56}
<i>even II</i> \mathcal{A}_f^m	1	[-i](+)(+)	(-i)+	[-i][+][+]	(-i)(+)[+]	$-\frac{i}{2}$	$\frac{1}{2}$	$\frac{1}{2}$
	2	(+i)[-](+)	[+i](-)(+)	(+i)(-)[+]	[+i](-)[+]	$\frac{i}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$
	3	(+i)(+)[-]	[+i][+][-]	(+i)[+][-]	[+i](+)(-)	$-\frac{i}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$
	4	[-i](-)[-]	(-i)(-)[-]	[-i](-)(-)	(-i)[-][-]	$-\frac{i}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$

even “basis vectors” and their Hermitian conjugated partners for the case $d = (5 + 1)$ can be found in Ref. [11].

While the Clifford odd “basis vectors” are (chosen to be) left handed, $\Gamma^{(5+1)} = -1$, their Hermitian conjugated partners have opposite handedness, Eq. 21 in App. A ⁴.

Acknowledgments

The authors thank Department of Physics, FMF, University of Ljubljana, Society of Mathematicians, Physicists and Astronomers of Slovenia, for supporting the research on the *spin-charge-family* theory by offering the room and computer facilities and Matjaž Breskvar of Beyond Semiconductor for donations, in particular for the annual workshops entitled “What comes beyond the standard models”.

References

- [1] N. Mankoč Borštnik, “Spinor and vector representations in four dimensional Grassmann space”, *J. of Math. Phys.* **34** (1993) 3731-3745, “Unification of spin and charges in Grassmann space?”, hep-th 9408002, IJS.TP.94/22, *Mod. Phys. Lett.***A** (10) No.7 (1995) 587-595;
- [2] A. Borštnik Bračić, N. S. Mankoč Borštnik, “On the origin of families of fermions and their mass matrices”, hep-ph/0512062, *Phys. Rev.* **D 74** 073013-28 (2006).
- [3] N.S. Mankoč Borštnik, H.B.F. Nielsen, *J. of Math. Phys.* **43**, 5782 (2002) [arXiv:hep-th/0111257]. “How to generate families of spinors”, *J. of Math. Phys.* **44** 4817 (2003) [arXiv:hep-th/0303224].
- [4] N.S. Mankoč Borštnik, D. Lukman, “Vector and scalar gauge fields with respect to $d = (3 + 1)$ in Kaluza-Klein theories and in the *spin-charge-family theory*”, *Eur. Phys. J. C* **77** (2017) 231.
- [5] N.S. Mankoč Borštnik, H.B.F. Nielsen, “Understanding the second quantization of fermions in Clifford and in Grassmann space”, *New way of second quantization of fermions — Part I and Part II*, [arXiv:2007.03517, arXiv:2007.03516].
- [6] N. S. Mankoč Borštnik, H. B. Nielsen, “How does Clifford algebra show the way to the second quantized fermions with unified spins, charges and families, and with vector and scalar gauge fields beyond the *standard model*”, *Progress in Particle and Nuclear Physics*, <http://doi.org/10.1016.j.pnp.2021.103890> .
- [7] N. S. Mankoč Borštnik, “How Clifford algebra helps understand second quantized quarks and leptons and corresponding vector and scalar boson fields, opening a new step beyond the standard model”, Reference: NUPHB 994 (2023) 116326 , [arXiv: 2210.06256, physics.gen-ph V2].
- [8] N. S. Mankoč Borštnik, “Clifford odd and even objects in even and odd dimensional spaces”, *Symmetry* 2023,15,818-12-V2 94818, <https://doi.org/10.3390/sym15040818>, [arxiv.org/abs/2301.04466] , <https://www.mdpi.com/2073-8994/15/4/818> Manuscript ID: symmetry-2179313.
- [9] N.S. Mankoč Borštnik N S, “The spin-charge-family theory is explaining the origin of families, of the Higgs and the Yukawa couplings”, *J. of Modern Phys.* **4** (2013) 823 [arXiv:1312.1542].

⁴Let us check the handedness of the chosen representation: $\Gamma^{5+1}\hat{b}_1^{1\dagger}(\equiv(+i)[+][+]) = \sqrt{(-1)^5}i^3(\frac{2}{i})^3S^{03}S^{12}S^{56}((+i)[+][+]) = \frac{i^4 2^3}{i^3} \frac{i}{2} \frac{1}{2} \frac{1}{2} ((+i)[+][+]) = -1((+i)[+][+])$.

- [10] N. S. Mankoč Borštnik, "New way of second quantized theory of fermions with either Clifford or Grassmann coordinates and *spin-charge-family* theory " [arXiv:1802.05554v4, arXiv:1902.10628], N. S. Mankoč Borštnik, "How Clifford algebra can help understand second quantization of fermion and boson fields", [arXiv: 2210.06256. physics.gen-ph V1] .
- [11] N. S. Mankoč Borštnik, "How Clifford algebra can help understand second quantization of fermion and boson fields", [arXiv: 2210.06256. physics.gen-ph V1] ,
- [12] N. S. Mankoč Borštnik, "Clifford odd and even objects offer description of internal space of fermions and bosons, respectively, opening new insight into the second quantization of fields", The 13th Biental Conference on Classical and Quantum Relativistic Dynamics of Particles and Fields IARD 2022, Prague, 6 – 9 June, [<http://arxiv.org/abs/2210.07004>] .
- [13] M. Blagojević, *Gravitation and gauge symmetries*, IoP Publishing, Bristol 2002.
- [14] H.B. Nielsen, M. Ninomiya, "Novel String Field Theory and Bound State, Projective Line, and sharply 3-transitive group", [arXiv:2111.05106v1, physics.gen-ph].
- [15] K. Wray, "An Introduction to String Theory".
- [16] E.H.El Kinani, "Between Quantum Virasoro Algebra \mathcal{L}_1 and Generalized Clifford Algebras", [arXiv:math-ph/0310044].
- [17] Pavšič, M. *The Landscape of Theoretical Physics: Global View*; van der Merwe, A., Ed.; Kluwer Academic Publishers: New York, NY, USA, 2001.