

(Re-)reading Sklar (1959) – A personal view on Sklar’s theorem

GERY GEENENS*

School of Mathematics and Statistics,

UNSW Sydney, Australia

In probability and statistics, copula methods have become ubiquitous when it comes to analyse, model and quantify the dependence between variables. Systematically, any written (research paper) or verbal (conference talk) communication about copulas starts with a statement of so-called ‘*Sklar’s theorem*’, establishing existence of a copula for any multivariate probability distribution. After defining a d -dimensional copula ($d \in \mathbb{N}$) as a continuous cumulative distribution function supported on the unit hypercube $[0, 1]^d$ with uniform marginals, the theorem is typically stated under a form equivalent to the following:

Theorem 0 (‘**Sklar’s theorem**’). *a) Let $F_{1\dots d}$ be a d -dimensional ($d \in \mathbb{N}$) distribution function with marginals F_1, \dots, F_d . Then, (i) there exists a d -dimensional copula C such that*

$$F_{1\dots d}(x_1, \dots, x_d) = C(F_1(x_1), \dots, F_d(x_d)) \quad (\star)$$

for all $(x_1, \dots, x_d) \in \overline{\mathbb{R}}^d$; (ii) if each F_k ($k = 1, \dots, d$) is continuous, then C is unique; otherwise, C is uniquely determined on $\times_{k=1}^d \text{Ran } F_k$, where $\text{Ran } F_k = \{t \in [0, 1] : \exists x \in \overline{\mathbb{R}} \text{ s.t. } F_k(x) = t\}$.

b) Conversely, if C is a d -dimensional copula and F_1, \dots, F_d are univariate distribution functions, then the function $F_{1\dots d}$ defined via (\star) is a d -dimensional distribution function with marginals F_1, \dots, F_d .

(Here $\overline{\mathbb{R}}$ denotes the extended real line $[-\infty, \infty]$.) This is the theorem as it is stated in McNeil et al. (2005, Theorem 5.3) and (for $d = 2$) in Nelsen (2006, Theorem 2.3.3). Statements in other main references on copulas, such as Joe (2015, Theorem 1.1), Durante and Sempi (2015, Theorem 2.2.1) or Hofert et al. (2018, Theorem 2.3.1), differ only slightly. The reference provided is invariably Sklar (1959).

Now, not long ago, in the discussion which followed a seminar on copulas which I attended, the speaker argued that Sklar (1959) was certainly the most cited *unread* statistical paper. The argument holds water if we put in perspective the facts that (i) Sklar (1959) is referenced each time copulas are introduced, leading to a huge number of citations (close to 11,000 at the time of writing this note, according to *Google*

*ggeenens@unsw.edu.au

Scholar); and (ii) it is an ‘old’ paper in French, which was difficult to access for a long time, even after it was republished (Bosq, 2010). Thus, as Theorem 0 above is found (in English) in a multitude of other sources easily accessible, it may be reasonably conjectured that only a minor fraction of the authors citing Sklar (1959) put in the effort to access and read the original text.

Admittedly, I was not part of that minor fraction until recently; and Theorem 0 was reported as-is in Geenens et al. (2017) and Geenens (2020) with reference to Sklar (1959) (which is very bad practice, for that matter). Yet, the previous discussion prompted me to read the original paper in French, only to find out that Sklar (1959) does *not* contain any such ‘Sklar’s theorem’ under the above form – making a wider community aware of this fact may be the only purpose of this short note. In fact, the paper comprises five theorems, among which three (*Théorème 1*, *Théorème 2* and *Théorème 3*), when combined, allow one to reconstruct and/or deduce Theorem 0. For convenience, we translate¹ here in English Sklar (1959)’s *Théorème 1*, *Théorème 2* and *Théorème 3*, as well as the definition of a copula appearing in the sequence (*Définition 1*). (The notations and footnotes are original from Sklar (1959). The three *Théorèmes* and the *Définition* appear in this order. Nothing is omitted between the statements, given without proofs.)

Théorème 1. *Let G_n be an n -dimensional cumulative distribution function with margins F_1, F_2, \dots, F_n . Let R_k be the set of values of F_k , for $k = 1, \dots, n$. Then there exists a unique function H_n defined on the Cartesian product $R_1 \times R_2 \times \dots \times R_n$ and such that*

$$G_n(x_1, \dots, x_n) = H_n(F_1(x_1), F_2(x_2), \dots, F_n(x_n)).$$

Définition 1. *We call (n -dimensional) copula any function C_n , continuous, non-decreasing², defined on the Cartesian product of n closed intervals $[0, 1]$ and satisfying the conditions:*

$$C_n(0, 0, \dots, 0) = 0, \quad C_n(1, \dots, 1, \alpha, 1, \dots, 1) = \alpha.^3$$

Théorème 2. *The function H_n of Theorem 1 can be extended (in general, in more than one way) into a copula C_n . An extension of H_n , the copula C_n satisfies the condition:*

$$G_n(x_1, \dots, x_n) = C_n(F_1(x_1), F_2(x_2), \dots, F_n(x_n)).$$

¹Bearing in mind that ‘*traduire c’est trahir*’ – ‘translating is betraying’ – as my high-school English teacher used to say. Ironically, the statements in Sklar (1959) were themselves, presumably, French translations of Sklar’s initial thoughts, making all this an interesting instance of the ‘broken telephone game’.

²In the sense of an n -dimensional cumulative distribution function.

³Special cases of such functions were considered in Féron (1956).

Théorème 3. *Let be given univariate cumulative distribution functions F_1, F_2, \dots, F_n . Let C_n be an arbitrary n -dimensional copula. Then the function G_n defined as*

$$G_n(x_1, \dots, x_n) = C_n(F_1(x_1), F_2(x_2), \dots, F_n(x_n))$$

is an n -dimensional cumulative distribution function with margins F_1, F_2, \dots, F_n .

Théorème 1 establishes the existence and uniqueness of the function that would later be called *subcopula* – Sklar (1959) did not use that word, not introduced before Schweizer and Sklar (1974, Definition 3). This subcopula, denoted H below to stay consistent with *Théorème 1*, is defined only on the ‘Cartesian product of the sets of values of F_k ’, that is, $\times_{k=1}^d \text{Ran } F_k$ in the notation of Theorem 0, and satisfies

$$F_{1\dots d}(x_1, \dots, x_d) = H(F_1(x_1), \dots, F_d(x_d)) \quad \forall (x_1, \dots, x_d) \in \overline{\mathbb{R}}^d. \quad (**)$$

Although no details are given, *Théorème 2* states that the unique subcopula satisfying $(**)$ may be extended ‘*in general, in more than one way*’ beyond $\times_{k=1}^d \text{Ran } F_k$ into a function defined on the whole of the unit hypercube $[0, 1]^d$ and satisfying *Définition 1*, such a function being called *copula*. In other words, there exists at least one copula C coinciding exactly with the subcopula on $\times_{k=1}^d \text{Ran } F_k$:

$$C(u_1, \dots, u_d) = H(u_1, \dots, u_d) \quad \forall (u_1, \dots, u_d) \in \times_{k=1}^d \text{Ran } F_k. \quad (***)$$

Then $(*)$ follows immediately from $(**)$ and $(***)$. Evidently, the values taken by any such copula C outside $\times_{k=1}^d \text{Ran } F_k$ are totally irrelevant, as they do not even appear in $(*)$. All in all, *Théorème 2* is akin to part $a)(i)$ of Theorem 0, while clearly *Théoreme 3* is its part $b)$.

What about part $a)(ii)$? Sklar (1959) does not make any specific mention of the uniqueness of the copula in the continuous case. Rather the contrary, *Théorème 2* is stated in general with an explicit note about the non-uniqueness of the copula. Naturally, for a continuous univariate distribution F_k , $\text{Ran } F_k \equiv [0, 1]$, thus if each F_k ($k = 1, \dots, d$) is continuous, then $\times_{k=1}^d \text{Ran } F_k \equiv [0, 1]^d$. In that case, $(***)$ implies that the subcopula is a copula, and since there is no room for arbitrary extension, any copula C satisfying $(*)$ *must be* the subcopula, making such C unique. Hence part $a)(ii)$ follows from *Théorèmes 1-2* and is not an add-on *stricto sensu*, but was apparently not an essential point to make for Sklar (1959).

This illustrates that Theorem 0 should not be regarded as just a concise re-statement of the sequence *Théorème 1*, *Théorème 2* and *Théorème 3*. The substance may be equivalent, but the form is not exactly the same, and this may lead to subtly different reading and interpretation. What is notable is that, although Sklar (1959) gives a prominent place to the subcopula – with *Théorème 1* explicitly devoted to

it – it has totally disappeared from the ‘modern’ statement Theorem 0, largely consigning it to oblivion. Indeed, in the above classical references, either the subcopula is introduced only in the technical lemmas leading to Theorem 0 (Nelsen, 2006, Lemma 2.3.4; Durante and Sempi, 2015, Lemma 2.3.3), or it is not mentioned at all (McNeil et al., 2005, Joe, 2015, Hofert et al., 2018). Though, it is clear that the only informative part of the copula is the underlying subcopula; and therefore, understanding completely the whys and wherefores of (\star) seems conditional on a proper recognition of the role played by H . It is my opinion that short-circuiting the subcopula step as in Theorem 0 induces overemphasis on the copula(s) C and ultimately unwarranted exploitation of (\star) when $C \neq H$. This is, especially, the case when it comes to analysing or modelling dependence, which is the main if not only application of Sklar’s theorem.

Remarkably, Theorem 0 does not make any reference to dependence: (\star) is merely an analytical result providing an alternative representation of $F_{1\dots d}$ which may or may not be of any relevance. It is really the *interpretation* which we are willing to make of it which brings in the concept of dependence and relates it to copulas. Effectively, it appears from (\star) that C is to capture how the marginals F_1, \dots, F_d interlock inside $F_{1\dots d}$ – which seems fair to call the ‘dependence structure’. This explains why, early on, copulas were called ‘*dependence functions*’; e.g., in Galambos (1978, Definition 5.2.1) and Deheuvels (1979, 1980).

Though, for playing with the dependence structure of $F_{1\dots d}$, the subcopula H is the only function worth examining: it always exists, it is always unique, and it always describes unequivocally through $(\star\star)$ how to reconstruct $F_{1\dots d}$ from the marginals F_1, \dots, F_d . Thus, with *Théorème 1* in hand, it is not clear what is the added value of the copula extension promised by *Théorème 2* – that same extension (\star) implicitly but exclusively put forward by Theorem 0. Having said this, the fact that the subcopula H contains all necessary information for describing the dependence in $F_{1\dots d}$ does not imply that it is, in itself, a valid representation of that dependence. Indeed, defined on $\times_{k=1}^d \text{Ran } F_k$, the subcopula is in general not a stand-alone element which can be handled and analysed without reference to marginal distributions, and therefore, cannot isolate a dependence structure as such. In fact, H must adjust to F_1, \dots, F_d by definition – again, *in general*.

Now, it so happens that, when all the marginal distributions are continuous, the subcopula takes a very specific form which is invariably a d -variate distribution function with continuous uniform margins on $[0, 1]$ – this follows straightforwardly from standard results on functions of random variables applied to $(\star\star)$, in particular the Probability Integral Transform (PIT).⁴ In this case, the subcopula is a copula as per *Définition 1*, so $H \equiv C$ (and $(\star) \equiv (\star\star)$) as observed above, but *even more importantly* this (sub)copula is

⁴If X_k is a continuous variable with cumulative distribution F_k , then $F_k(X_k) \sim \mathcal{U}_{[0,1]}$ always.

‘marginal-distribution-free’,⁵ a.k.a. ‘margin-free’. Unbound from any marginal interference, the (sub)copula can now be genuinely understood as capturing the heart of $F_{1\dots d}$, that is, its dependence structure as such. The representation (\star) is then particularly appealing, as it provides an explicit breakdown of a joint distribution into the individual behaviour of the variables of interest (captured by F_1, \dots, F_d) on one hand, and their interdependence structure (captured by C) on the other – with no overlap/redundancy between the two. The entire copula methodology for dependence modelling developed around this neat decomposition and the desirable consequences thereof.

It cannot be stressed enough, though, that this pleasant situation only follows as a corollary of two favourable events which occur concurrently *when and only when* all the variables involved are continuous: first, the copula appearing in (\star) is the subcopula, and second, that subcopula is margin-free. In all other non-continuous situations, any copula C satisfying (\star) is nothing more than an arbitrary extension of the subcopula H in $(\star\star)$, which itself is *not* a satisfactory representation of the dependence of $F_{1\dots d}$ as it is not margin-free. The suitability of (\star) for analysing and/or modelling dependence becomes then highly questionable. In effect, the validity of any attempt at dependence modelling based on the typical interpretation of (\star) as a clear-cut decomposition ‘marginals vs. dependence’, is *critically* contingent on the continuity of all the variables.

Without any reference to the subcopula, the usual statement of ‘Sklar’s theorem’ as in Theorem 0 does not provide the clues to appreciate that. I questioned above the real benefit of the extension promised by *Théorème 2* when we have *Théorème 1*. The question may be rephrased as: why did (\star) become the universal baseline, in lieu of $(\star\star)$? The only reason I see is that, since a copula is always a distribution supported on $[0, 1]^d$ with standard uniform margins by definition, the function C in (\star) appears as a standard object in a invariant form (in particular: margin-free); as opposed to the function H in $(\star\star)$, whose exact nature is undefined and its specification requiring knowledge of $\times_{k=1}^d \text{Ran } F_k$. Yet, such invariance of C may only be granted in continuous cases – but then H enjoys the same desirable property, anyway – otherwise it is mostly a lure: in fact, the definition of C makes it into a blanket which concealed the fact that, ‘underneath’, its anchor points are fixed by H via $(\star\star\star)$. That the gaps between the nodes of $\times_{k=1}^d \text{Ran } F_k$ may be filled in such a way that C maintains uniform margins is actually little more than an analytical artefact of no obvious relevance when it comes to dependence.

What adds to the blur is part *a)(ii)* explicitly contrasting the continuous and non-continuous cases in terms of the (non-)uniqueness of the copula C in (\star) . This may give the feeling that this is the only

⁵Where ‘distribution-free’ is taken in the sense of Kendall and Sundrum (1953): *free of the parent distribution*. Thus, more specifically here, ‘marginal-distribution-free’, or ‘margin-free’, means *free of the marginal distributions of the parent distribution $F_{1\dots d}$* .

notable difference between the two situations, and may consequently divert attention from other questions. Indeed the lack of uniqueness of C and the ensuing problems of *model unidentifiability* have often been presented as the main hurdle for practical use of copula methods outside the continuous framework, and have consequently been abundantly commented (Genest and Nešlehová, 2007, Trivedi and Zimmer, 2017, Faugeras, 2017, Geenens, 2020, Nasri and Rémillard, 2023). In my current view, though, the only consequential difference between continuous and non-continuous cases is that the (sub)copula is margin-free in the former, and not in the latter – and this seems to have been much less frequently pinpointed as such.

What has been discussed is all the ‘little annoyances’ which follow directly from this; e.g., the fact that copula-based dependence measures, such as Kendall’s τ or Spearman’s ρ , depend on the margins in non-continuous settings (Marshall, 1996; Genest and Nešlehová, 2007, Section 4). Yet, these are only consequences of the lack of margin-free-ness of C which, *in itself*, appears to me as the real predicament: in effect we are losing the very reason-of-being of the copula approach, which is precisely its power to dissociate marginal behaviour and dependence structure via (\star) . For example, Joe (2015, Section 1.6) motivates resorting to copulas over alternative multivariate models as: “(...) *the copula approach has an advantage of having univariate margins of different types and the dependence structure can be modeled separately from univariate margins*”. Yet, outside the continuous framework, this alleged separation between dependence structure and margins is clearly violated. All in all, it seems that copula methods applied to non-continuous distributions miss entirely their own point.

It is, therefore, my opinion that copula-like methods for analysing, modelling and quantifying dependence in non-continuous multivariate distributions should *not* be based on (\star) . I elaborated on this in Geenens (2020), and proposed an alternative approach for discrete distributions. In a nutshell, the idea is to extract the information about dependence from the subcopula, and to reshape it under the form of a distribution with (discrete) uniform margins – hence ‘margin-free’ – this defining a *discrete copula*.

References

- Bosq, D., 2010. L’article fondateur des copules. *Annales de l’ISUP* 54 (1-2), 3–6.
- Deheuvels, P., 1979. La fonction de dépendance empirique et ses propriétés. un test non paramétrique d’indépendance. *Bulletins de l’Académie Royale de Belgique* 65 (1), 274–292.
- Deheuvels, P., 1980. Non parametric tests of independence. In: *Statistique non paramétrique asymptotique*. Springer, pp. 95–107.
- Durante, F., Sempi, C., 2015. *Principles of Copula Theory*. CRC Press, Boca Raton.
- Faugeras, O., 2017. Inference for copula modeling of discrete data: a cautionary tale and some facts. *Depend Model*.

- Féron, R., 1956. Sur les tableaux de corrélation dont les marges sont données. Cas de l'espace a trois dimensions. *Annales de l'Institut de Statistiques de l'Université de Paris V* (1), 3–12.
- Galambos, J., 1978. The Asymptotic Theory of Extreme Order Statistics. Wiley Series in Probability and Statistics - Applied Probability and Statistics Section.
- Geenens, G., 2020. Copula modeling for discrete random vectors. *Dependence Modeling* 8 (1), 417–440.
- Geenens, G., Charpentier, A., Paindaveine, D., 2017. Probit transformation for nonparametric kernel estimation of the copula density. *Bernoulli*.
- Genest, C., Nešlehová, J., 2007. A primer on copulas for count data. *ASTIN Bull.*
- Hofert, M., Kojadinovic, I., Mächler, M., Yan, J., 2018. Elements of Copula Modeling with R. Use R! Series, Springer Nature.
- Joe, H., 2015. Dependence Modeling with Copulas. Chapman and Hall/CRC.
- Kendall, M. G., Sundrum, R., 1953. Distribution-free methods and order properties. *Revue de l'Institut International de Statistique*, 124–134.
- Marshall, A., 1996. Copulas, marginals and joint distributions. In: Rüschendorf, L., Schweizer, B., Taylor, M. (Eds.), Distributions with Fixed Marginals and Related Topics. Institute of Mathematical Statistics, pp. 213–222.
- McNeil, A., Frey, R., Embrechts, P., 2005. Quantitative Risk Management: Concepts, Techniques and Tools. Princeton University Press.
- Nasri, B., Rémillard, B., 2023. Identifiability and inference for copula-based semiparametric models for random vectors with arbitrary marginal distributions. *arXiv e-prints*, arXiv:2301.13408.
- Nelsen, R., 2006. An Introduction to Copulas, 2nd Edition. Springer, New York.
- Schweizer, B., Sklar, A., 1974. Operations on distribution functions not derivable from operations on random variables. *Studia Mathematica* 52 (1), 43–52.
- Sklar, M., 1959. Fonctions de répartition à n dimensions et leurs marges. *Publ. inst. statist. univ. Paris* 8, 229–231.
- Trivedi, P., Zimmer, D., 2017. A note on identification of bivariate copulas for discrete count data. *Econometrics* 5 (1), 10.