Bayesian Fused Lasso Modeling for Binary Data

Yuko Kakikawa^a and Shuichi Kawano^b

^a Department of Statistical Science, Graduate University of Advanced Studies (SOKENDAI), 10-3 Midori-cho, Tachikawa, Tokyo 190-8562, Japan.

^b Faculty of Mathematics, Kyushu University, 744 Motooka, Nishi-ku, Fukuoka 819-0395, Japan.

ARTICLE HISTORY

Compiled December 15, 2023

ABSTRACT

 L_1 -norm regularized logistic regression models are widely used for analyzing data with binary response. In those analyses, fusing regression coefficients is useful for detecting groups of variables. This paper proposes a binomial logistic regression model with Bayesian fused lasso. Assuming a Laplace prior on regression coefficients and differences between adjacent regression coefficients enables us to perform variable selection and variable fusion simultaneously in the Bayesian framework. We also propose assuming a horseshoe prior on the differences to improve the flexibility of variable fusion. The Gibbs sampler is derived to estimate the parameters by a hierarchical expression of priors and a data-augmentation method. Using simulation studies and real data analysis, we compare the proposed methods with the existing method.

KEYWORDS

Horseshoe prior; Markov chain Monte Carlo; Variable fusion; Variable selection; Regularized logistic regression

1. Introduction

Recently, L_1 -norm regularized logistic regression models for binary response data have received considerable attention in various fields of research (Wu et al. 2009; Pereira et al. 2016; Xin et al. 2014; Yu et al. 2015). The lasso (Tibshirani 1996), which is the most well-known L_1 -norm regularized method, imposes the sum of absolute values of regression coefficients as a regularization term, and then estimates some regression coefficients as zeros. This induces variable selection. Tibshirani (1996) focused on the linear regression models, and subsequently many researchers have extended the lasso to logistic regression models (e.g., see Lokhorst (1999); Shevade and Keerthi (2003); Pereira et al. (2016)). Meanwhile, Tibshirani et al. (2005) proposed the fused lasso, which not only performs variable selection but also estimates the differences between adjacent regression coefficients as exactly zero, that is, variable fusion. Through variable fusion, the group of variables which have similar roles in the prediction can be identified. Yu et al. (2015) introduced a logistic regression model estimated by the fused lasso and applied it to spectral data.

CONTACT Yuko Kakikawa. Email: kakikawa.yuko@ism.ac.jp

From the perspective of Bayesian methods, a regularization term in the lasso corresponds to assuming a Laplace distribution on the coefficients. Park and Casella (2008) proposed the Bayesian lasso and enabled Gibbs sampling with the hierarchical expression of a Laplace prior by a scale mixture of normals (Andrews and Mallows 1974). Kyung et al. (2010) extended the Bayesian lasso to the Bayesian fused lasso, which assumes a Laplace prior not only on regression coefficients but also on the differences of adjacent regression coefficients. Genkin et al. (2007) proposed the Bayesian lasso for logistic regression models for analyzing text data. Betancourt et al. (2017) extended the Bayesian fused lasso to a logistic regression model, and then applied it to analyzing financial trading network data. However, these existing methods do not consider variable fusion and variable selection simultaneously.

It should be noted that a Laplace prior induces over-shrinkage on the coefficients when the true coefficients are non-zero, while the prior induces under-shrinkage on the coefficients when the true coefficients are zero. This is a drawback to using a Laplace prior. As a counterpart of a Laplace prior, a horseshoe prior (Carvalho et al. 2010) has been considered. A horseshoe prior has both a taller spike at zero and a heavier tail than a Laplace prior. These properties of this distribution overcome the above drawback of a Laplace prior. Makalic and Schmidt (2015) and Bhattacharyya et al. (2022) proposed Bayesian logistic regression models by assuming a horseshoe prior on regression coefficients. However, the existing Bayesian logistic regression models with a horseshoe prior ignore variable fusion.

In the present paper, we first propose a Bayesian logistic regression model by assuming a Laplace prior on both regression coefficients and differences between adjacent regression coefficients. To prevent under-shrinkage and over-shrinkage of the differences, we next propose assuming a horseshoe prior on the differences between adjacent regression coefficients. Subsequently, we derive the Gibbs sampler by using the hierarchical expression of a half-Cauchy distribution (Wand et al. 2011) and a data-augmentation method with a Pólya-Gamma distribution (Polson et al. 2013).

The remainder of this paper is organized as follows. Section 2 introduces L_1 -norm regularized logistic regression models. Section 3 describes a Bayesian approach for the L_1 -norm regularized methods. In Section 4, we present a Bayesian logistic regression model with both variable selection by a Laplace prior and variable fusion by a Laplace prior or a horseshoe prior. Section 5 conducts Monte Carlo simulations and makes an application to real data to investigate the performance of our proposed methods, and compares it to the existing method, the logistic regression model with the fused lasso. We conclude the paper in Section 6.

2. L_1 -norm regularized logistic regression

Suppose that we have a dataset $\{(y_i, x_i); i = 1, ..., n\}$, where $y_i \in \{0, 1\}$ is a binary response variable and x_i is a *p*-dimensional vector of explanatory variables. A logistic regression model is formulated as

$$\Pr(y_i = 1 \mid \boldsymbol{x}_i) = \frac{1}{1 + e^{-(\beta_0 + \boldsymbol{x}_i^T \boldsymbol{\beta})}}, \quad i = 1, \dots, n,$$
(1)

where β_0 is an intercept and $\boldsymbol{\beta} = (\beta_1, \beta_2, \dots, \beta_p)^T$ is a *p*-dimensional regression coefficient vector. Then, the likelihood function is given by

$$\prod_{i=1}^{n} f(y_i \mid \boldsymbol{x}_i; \beta_0, \boldsymbol{\beta}),$$
(2)

where

$$f(y_i \mid \boldsymbol{x}_i; \beta_0, \boldsymbol{\beta}) = \frac{(e^{\beta_0 + \boldsymbol{x}_i^T \boldsymbol{\beta}})^{y_i}}{1 + e^{\beta_0 + \boldsymbol{x}_i^T \boldsymbol{\beta}}}.$$
(3)

Then, the log-likelihood function is given by

$$\ell_{LR}(\beta_0, \boldsymbol{\beta}) = \sum_{i=1}^{n} \left[y_i(\beta_0 + \boldsymbol{x}_i^T \boldsymbol{\beta}) - \log \left\{ 1 + \exp(\beta_0 + \boldsymbol{x}_i^T \boldsymbol{\beta}) \right\} \right].$$
(4)

In a maximum likelihood estimation, the regression coefficient β is estimated by the maximization of the log-likelihood (4). However, the over-fitting problem often occurs when the sample size n is smaller than the number of variables p. L_1 -norm regularization is a useful tool for solving this problem. The lasso (Tibshirani 1996; Lokhorst 1999) is one of the most widely used L_1 -norm regularization methods. The lasso is formulated as the following optimization problem:

$$\hat{\boldsymbol{\beta}}_{L} = \operatorname*{arg\,min}_{\boldsymbol{\beta}_{0},\boldsymbol{\beta}} \left\{ -\ell_{LR}(\boldsymbol{\beta}_{0},\boldsymbol{\beta}) + \gamma \sum_{j=1}^{p} |\boldsymbol{\beta}_{j}| \right\},\tag{5}$$

where γ (> 0) is a regularization parameter that adjusts the strength of the shrinkage of regression coefficients. In the problem in (5), the sum of absolute values of regression coefficients is imposed as the regularization term, and thus the lasso estimates regression coefficients as exactly zero. Consequently, the lasso performs variable selection.

However, the lasso cannot consider the relationships among explanatory variables. When adjacent variables are strongly correlated, like in the case of spectral data (Yu et al. 2015), adjacent variables have similar roles for the response. For such data, regression coefficients which correspond to variables having similar roles should be estimated as the same values. To address this issue, the fused lasso (Tibshirani et al. 2005; Yu et al. 2015) has been introduced. The fused lasso is formulated as the following minimization problem:

$$\hat{\boldsymbol{\beta}}_{FL} = \operatorname*{arg min}_{\boldsymbol{\beta}_0,\boldsymbol{\beta}} \left\{ -\ell_{LR}(\boldsymbol{\beta}_0,\boldsymbol{\beta}) + \lambda_1 \sum_{j=1}^p |\boldsymbol{\beta}_j| + \lambda_2 \sum_{j=2}^p |\boldsymbol{\beta}_j - \boldsymbol{\beta}_{j-1}| \right\},\tag{6}$$

where $\lambda_1 (> 0)$ and $\lambda_2 (> 0)$ are regularization parameters. The regularization parameter λ_1 adjusts the amount of the shrinkage of variables, while the regularization parameter λ_2 adjusts the smoothness of the differences between adjacent variables. The fused lasso (6) can estimate some of the regression coefficients and differences of

adjacent regression coefficients as exactly zero. This means that the fused lasso can perform variable selection and variable fusion simultaneously.

3. Bayesian approach for regularized logistic regression

This section reviews a Bayesian logistic regression model and a data-augmentation method using a Pólya-Gamma distribution. We describe a Bayesian approach for L_1 -norm regularized methods. Specifically, we explain the Bayesian lasso and the Bayesian fused lasso. We also describe a horseshoe prior and a Bayesian regularized logistic regression with a horseshoe prior.

3.1. Data-augmentation method with Pólya-Gamma distribution

In the Bayesian framework, a logistic regression model is formulated as

$$y_i \mid \boldsymbol{x}_i, \boldsymbol{\beta}, \beta_0 \sim \operatorname{Binom}\left(1, \frac{1}{1 + e^{-(\beta_0 + \boldsymbol{x}_i^T \boldsymbol{\beta})}}\right),$$
(7)

where $Binom(\cdot, \cdot)$ represents a binomial distribution.

To obtain samples from the posterior distribution for (7) by Gibbs sampling, Polson et al. (2013) proposed a data-augmentation method with a Pólya-Gamma distribution. The probability density function of a Pólya-Gamma distribution is expressed by

$$PG(x \mid a, b) = \frac{1}{2\pi^2} \sum_{k=1}^{\infty} \frac{g_k}{\left(k - \frac{1}{2}\right)^2 + \frac{b^2}{(4\pi^2)}},$$
(8)

where $a \ (> 0)$ and b are hyper-parameters and g_k is an independent random variable with a Gamma distribution Ga(a, 1). In addition, the Pólya-Gamma distribution (8) can be expressed hierarchically with $PG(x \mid a, 0)$ as follows:

$$\operatorname{PG}(x \mid a, b) \propto \exp\left(-\frac{b^2 x}{2}\right) \operatorname{PG}(x \mid a, 0).$$
 (9)

By using the hierarchical expression (9), the function (3) can be expressed hierarchically with the latent variables w_i :

$$f(y_i \mid \boldsymbol{x}_i, \boldsymbol{\beta}, \beta_0) = \frac{(e^{\beta_0 + \boldsymbol{x}_i^T \boldsymbol{\beta}})^{y_i}}{1 + e^{\beta_0 + \boldsymbol{x}_i^T \boldsymbol{\beta}}}$$
$$= \frac{1}{2} \exp\{\kappa_i(\beta_0 + \boldsymbol{x}_i^T \boldsymbol{\beta})\} \int_0^\infty \exp\left\{-\frac{w_i(\beta_0 + \boldsymbol{x}_i^T \boldsymbol{\beta})^2}{2}\right\} \operatorname{PG}(w_i \mid 1, 0) dw_i,$$
(10)

where $\kappa_i = y_i - 0.5$. When the regression coefficient vector β has a Gaussian prior

 $N_p(\mathbf{0}_p, \mathbf{B})$, the full conditional distributions can be obtained as follows:

$$\boldsymbol{\beta} \mid \boldsymbol{y}, \boldsymbol{X}, \beta_0, \lambda_1, w_1, \dots, w_n \sim \mathcal{N}_p(\boldsymbol{A}^{-1}\boldsymbol{X}^T\boldsymbol{W}(\boldsymbol{z} - \beta_0 \boldsymbol{1}), \boldsymbol{A}^{-1}), \\ w_i \mid \boldsymbol{x}_i, \boldsymbol{\beta}, \beta_0 \sim \mathcal{PG}(1, \beta_0 + \boldsymbol{x}_i^T \boldsymbol{\beta}),$$
(11)

where $\mathbf{X} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)^T$, $\mathbf{A} = \mathbf{X}^T \mathbf{W} \mathbf{X} + \mathbf{B}^{-1}$, $\mathbf{W} = \text{diag}(w_1, \dots, w_n)$, $\mathbf{z} = (\kappa_1/w_1, \kappa_2/w_2, \dots, \kappa_n/w_n)^T$, and **1** is an *n*-dimensional vector of which all components are one. From the full conditional distributions (11), the Gibbs sampling can be performed.

3.2. Bayesian approach for L_1 -norm regularized methods

The lasso can be interpreted as a MAP estimation when a Laplace prior is assumed on regression coefficients independently. Park and Casella (2008) proposed the Bayesian lasso by assuming a Laplace prior Laplace $(0, 1/\lambda)$, where the location parameter is 0 and the scale parameter is $1/\lambda$, on regression coefficients in terms of linear regression models. Park and Casella (2008) also developed the Gibbs sampler by using a scale mixture of normals (Andrews and Mallows 1974). With a scale mixture of normals, a Laplace prior can be expressed by a normal distribution and an exponential distribution hierarchically:

$$\boldsymbol{\beta} \mid \tau_1^2, \dots, \tau_p^2 \sim \mathcal{N}_p(\boldsymbol{0}_p, \boldsymbol{C}),$$

$$\tau_j^2 \sim \mathrm{EXP}\left(\frac{\lambda^2}{2}\right),$$
(12)

where $C = \text{diag}(\tau_1^2, \tau_2^2, \dots, \tau_p^2)$ and $\text{EXP}(\cdot)$ represents an exponential distribution. There are several ways to extend the Bayesian lasso in the framework of logistic

There are several ways to extend the Bayesian lasso in the framework of logistic regression models. Makalic and Schmidt (2016) proposed a logistic regression model with the priors (12) in the case of $\lambda^2 = 2$ and $\mathbf{C} = \text{diag} \left(\psi^2 \tau_1^2, \psi^2 \tau_2^2, \ldots, \psi^2 \tau_p^2\right)$, where ψ is a hyper-parameter which determines the overall shrinkage of the regression coefficients. Tian et al. (2019) also proposed a logistic regression model with the Bayesian lasso. In Tian et al. (2019), the priors (12) were assumed and the hyper-parameter λ was selected to make a predictive probability distribution be an approximately uniform distribution.

Among the models, we note the logistic regression model with the Bayesian lasso having priors (12) on regression coefficients. First, as in Makalic and Schmidt (2016), a uniform distribution $U(\alpha, \alpha)$ with a hyper-parameter α is assumed on the intercept β_0 . In addition, a Gamma distribution $Ga(r_1, \delta_1)$ ($r_1 > 0$, $\delta_1 > 0$) is assumed on λ^2 , following Park and Casella (2008). By combining the data-augmentation method in Sec. 3.1 and the expression for the prior on regression coefficients as a Gaussian scale-mixture prior (12), the full conditional distributions can be obtained as follows:

$$\boldsymbol{\beta} \mid \boldsymbol{y}, \boldsymbol{X}, \beta_0, \tau_1^2, \dots, \tau_p^2, w_1, \dots, w_n \sim \mathrm{N}_p(\boldsymbol{A}^{-1}\boldsymbol{X}^T\boldsymbol{W}(\boldsymbol{z} - \beta_0 \mathbf{1}), \boldsymbol{A}^{-1})$$

$$w_i \mid \boldsymbol{x}_i, \boldsymbol{\beta}, \beta_0 \sim \mathrm{PG}(1, \beta_0 + \boldsymbol{x}_i^T \boldsymbol{\beta}),$$

$$\frac{1}{\tau_j^2} \mid \beta_j, \lambda^2 \sim \mathrm{IGauss}\left(\sqrt{\frac{\lambda^2}{\beta_j^2}}, \lambda^2\right),$$

$$\lambda^2 \mid \tau_1^2, \dots, \tau_p^2 \sim \mathrm{Ga}\left(p + r_1, \frac{1}{2}\sum_{j=1}^p \tau_j^2 + \delta_1\right),$$

$$\beta_0 \mid \boldsymbol{y}, \boldsymbol{X}, \tau_1^2, \dots, \tau_p^2, w_1, \dots, w_n \sim \mathrm{N}\left(\frac{1}{S}\sum_{i=1}^n v_i, \frac{1}{S}\right),$$

where $\mathbf{A} = \mathbf{X}^T \mathbf{W} \mathbf{X} + \mathbf{C}^{-1}$, $\mathbf{v} = (\mathbf{z} - \mathbf{X}\boldsymbol{\beta}) \odot \mathbf{W}$, $S = \sum_{i=1}^n w_i$, and IGauss (\cdot, \cdot) represents an inverse-Gaussian distribution and the symbol \odot means the Hadamard product.

As an extension of the Bayesian lasso, Kyung et al. (2010) proposed a linear regression model which assumes a Laplace prior on regression coefficients and differences of adjacent regression coefficients. This extension is called the Bayesian fused lasso. The posterior mode of regression coefficients in the Bayesian fused lasso is equivalent to the fused lasso solution. Betancourt et al. (2017) extended the Bayesian fused lasso to multinomial logistic regression by assuming a Laplace prior on the differences between parameters corresponding to adjacent points in a time series expressed by a directed binary network.

3.3. Bayesian logistic regression model with horseshoe prior

A Laplace prior tends to shrink regression coefficients too little when the true coefficients are zero and shrink the coefficients too much when the true coefficients are non-zero. This is why a Laplace prior has insufficient concentration at zero and an exponential tail. To avoid this problem, Carvalho et al. (2010) proposed a horseshoe prior given by

$$\beta_j |\lambda_j^2, \tau^2 \sim \mathcal{N}(0, \lambda_j^2 \tau^2), \ \lambda_j \sim \mathcal{C}^+(0, 1), \ \tau \sim \mathcal{C}^+(0, 1),$$
(13)

where $C^+(\cdot, \cdot)$ represents a half-Cauchy distribution, λ_j adjusts the degree of the shrinkage of each regression coefficient β_j , and τ adjusts the amount of the shrinkage of the overall regression coefficients β . By having a hyper-parameter λ_j following a half-Cauchy distribution with a pole at zero and a polynomial tail, a horseshoe prior realizes an infinite spike at zero and a heavier tail than a Laplace prior. Therefore, a horseshoe prior can strongly shrink the small regression coefficients towards zero and prevent over-shrinkage of large regression coefficients.

Makalic and Schmidt (2015) proposed a logistic regression model whose regression coefficients follow a horseshoe prior. They utilized the hierarchical expression of a half-Cauchy distribution (Wand et al. 2011). When x follows $C^+(0, 1)$, the relation

$$x^2 \mid y \sim \operatorname{IG}\left(\frac{1}{2}, \frac{1}{y}\right), \quad y \sim \operatorname{IG}\left(\frac{1}{2}, 1\right)$$

holds, where $IG(\cdot, \cdot)$ represents an inverse-Gamma prior. Therefore, the priors (13) can be rewritten as follows:

$$\beta \mid \lambda_1^2, \dots, \lambda_p^2, \tau^2 \sim \mathcal{N}_p(\mathbf{0}_p, \mathbf{D}),$$

$$\lambda_j^2 \mid \nu_j \sim \mathrm{IG}\left(\frac{1}{2}, \frac{1}{\nu_j}\right),$$

$$\tau^2 \mid \xi \sim \mathrm{IG}\left(\frac{1}{2}, \frac{1}{\xi}\right),$$

$$\nu_1, \dots, \nu_p, \xi \sim \mathrm{IG}\left(\frac{1}{2}, 1\right),$$
(14)

where $\mathbf{D} = \text{diag} (\tau^2 \lambda_1^2, \tau^2 \lambda_2^2 \dots, \tau^2 \lambda_p^2)$. By using the priors (14) and the dataaugmentation method with a Pólya-Gamma distribution, the full conditional distributions can be obtained. The details are found in Makalic and Schmidt (2015).

4. Proposed methods

In this section, we propose a Bayesian logistic regression model whose regression coefficients and differences between adjacent coefficients follow a Laplace prior. We also propose an alternative Bayesian logistic regression model with a horseshoe prior on the differences.

4.1. Logistic regression model with the Bayesian fused lasso

We propose a logistic regression model which assumes a Laplace prior on regression coefficients and differences between adjacent regression coefficients.

First, we assume the following priors:

$$\beta_{j} \mid \lambda_{1} \sim \text{Laplace}\left(0, \frac{1}{\lambda_{1}}\right),$$

$$\beta_{j} - \beta_{j-1} \mid \lambda_{2} \sim \text{Laplace}\left(0, \frac{1}{\lambda_{2}}\right).$$
(15)

The priors (15) induce shrinkage of both regression coefficients and differences of adjacent regression coefficients towards zero. By using a scale mixture of normals, the priors (15) can be expressed as follows:

$$\boldsymbol{\beta} \mid \tau_1^2, \dots, \tau_p^2, \tilde{\tau}_2^2, \dots, \tilde{\tau}_p^2 \sim \mathcal{N}_p(\boldsymbol{0}_p, \boldsymbol{E}),$$

$$\tau_j^2 \sim \mathrm{EXP}\left(\frac{\lambda_1^2}{2}\right),$$

$$\tilde{\tau}_j^2 \sim \mathrm{EXP}\left(\frac{\lambda_2^2}{2}\right),$$

(16)

where the inverse matrix of E is expressed as

$$\boldsymbol{E}^{-1} = \begin{pmatrix} \frac{1}{\tau_1^2} + \frac{1}{\tilde{\tau}_2^2} & -\frac{1}{\tilde{\tau}_2^2} & 0 & \dots & 0 & 0\\ -\frac{1}{\tilde{\tau}_2^2} & \frac{1}{\tau_2^2} + \frac{1}{\tilde{\tau}_2^2} + \frac{1}{\tilde{\tau}_3^2} & -\frac{1}{\tilde{\tau}_3^2} & \dots & 0 & 0\\ 0 & -\frac{1}{\tilde{\tau}_3^2} & \frac{1}{\tau_3^2} + \frac{1}{\tilde{\tau}_3^2} + \frac{1}{\tilde{\tau}_4^2} & \dots & 0 & 0\\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots\\ 0 & 0 & 0 & \dots & \frac{1}{\tau_{p-1}^2} + \frac{1}{\tilde{\tau}_{p-1}^2} + \frac{1}{\tilde{\tau}_p^2} & -\frac{1}{\tilde{\tau}_p^2} \\ 0 & 0 & 0 & \dots & -\frac{1}{\tilde{\tau}_p^2} + \frac{1}{\tilde{\tau}_p^2} + \frac{1}{\tilde{\tau}_p^2} + \frac{1}{\tilde{\tau}_p^2} \end{pmatrix}$$

We assume priors on the intercept and hyper-parameters as follows:

$$\beta_0 \sim U(\alpha, \alpha),$$

$$\lambda_1^2 \sim Ga(r_1, \delta_1),$$

$$\lambda_2^2 \sim Ga(r_2, \delta_2),$$

(17)

where r_1, r_2, δ_1 , and δ_2 are positive parameters. The priors (17) enable fully Bayesian estimation. This formulation can be regarded as an extension of the Bayesian fused lasso to a logistic regression model.

By combining a data-augmentation method with a Pólya-Gamma distribution and the hierarchical expression of the priors (16), the full conditional distributions can be obtained as follows:

$$\boldsymbol{\beta} \mid \boldsymbol{y}, \boldsymbol{X}, \beta_{0}, \tau_{1}^{2}, \dots, \tau_{p}^{2}, \tilde{\tau}_{2}^{2}, \dots, \tilde{\tau}_{p}^{2}, w_{1}, \dots, w_{n} \sim \mathrm{N}_{p}(\boldsymbol{A}^{-1}\boldsymbol{X}^{T}\boldsymbol{W}(\boldsymbol{z}-\beta_{0}\boldsymbol{1}), \boldsymbol{A}^{-1}),$$

$$w_{i} \mid \boldsymbol{x}_{i}, \boldsymbol{\beta}, \beta_{0} \sim \mathrm{PG}(1, \beta_{0} + \boldsymbol{x}_{i}^{T}\boldsymbol{\beta}),$$

$$\frac{1}{\tau_{j}^{2}} \mid \beta_{j}, \lambda_{1}^{2} \sim \mathrm{IGauss}\left(\sqrt{\frac{\lambda_{1}^{2}}{\beta_{j}^{2}}}, \lambda_{1}^{2}\right),$$

$$\lambda_{1}^{2} \mid \tau_{1}^{2}, \dots, \tau_{p}^{2} \sim \mathrm{Ga}\left(p + r_{1}, \frac{1}{2}\sum_{j=1}^{p}\tau_{j}^{2} + \delta_{1}\right),$$

$$\frac{1}{\tilde{\tau}_{j}^{2}} \mid \beta_{j}, \beta_{j-1}, \lambda_{2}^{2} \sim \mathrm{IGauss}\left(\sqrt{\frac{\lambda_{2}^{2}}{(\beta_{j} - \beta_{j-1})^{2}}}, \lambda_{2}^{2}\right),$$

$$\lambda_{2}^{2} \mid \tilde{\tau}_{2}^{2}, \dots, \tilde{\tau}_{p}^{2} \sim \mathrm{Ga}\left(p - 1 + r_{2}, \frac{1}{2}\sum_{j=2}^{p}\tilde{\tau}_{j}^{2} + \delta_{2}\right),$$

$$\beta_{0} \mid \boldsymbol{y}, \boldsymbol{X}, \boldsymbol{\beta}, \tau_{1}^{2}, \dots, \tau_{p}^{2}, \tilde{\tau}_{2}^{2}, \dots, \tilde{\tau}_{p}^{2}, w_{1}, \dots, w_{n} \sim \mathrm{N}\left(\frac{1}{S}\sum_{i=1}^{n} v_{i}, \frac{1}{S}\right),$$
(18)

where $\boldsymbol{A} = \boldsymbol{X}^T \boldsymbol{W} \boldsymbol{X} + \boldsymbol{E}^{-1}$, $\boldsymbol{v} = (\boldsymbol{z} - \boldsymbol{X}\boldsymbol{\beta}) \odot \boldsymbol{W}$, and $S = \sum_{i=1}^n w_i$. From these full conditional distributions, Gibbs sampling can be performed.

4.2. Logistic regression model with Bayesian fused lasso with horseshoe prior

We also propose an alternative Bayesian model in Sec. 4.1 by assuming a horseshoe prior on differences between adjacent regression coefficients.

First, we introduce the priors given by

$$\beta_{j} \mid \tilde{\lambda}_{1} \sim \text{Laplace}\left(0, \frac{1}{\tilde{\lambda}_{1}}\right),$$

$$\beta_{j} - \beta_{j-1} \mid \lambda_{j}^{2}, \tilde{\tau}^{2} \sim \mathcal{N}\left(0, \lambda_{j}^{2} \tilde{\tau}^{2}\right),$$

$$\lambda_{j} \sim \mathcal{C}^{+}(0, 1),$$

$$\tilde{\tau} \sim \mathcal{C}^{+}(0, 1).$$
(19)

With the hierarchical expressions of a Laplace distribution and a half-Cauchy distribution, the priors (19) are expressed as

$$\beta \mid \tau_1^2, \dots, \tau_p^2, \lambda_2^2, \dots, \lambda_p^2, \tilde{\tau}^2 \sim N_p(\mathbf{0}_p, \mathbf{F}),$$

$$\tau_j^2 \sim EXP\left(\frac{\tilde{\lambda}_1^2}{2}\right),$$

$$\tilde{\tau}^2 \mid \xi \sim IG\left(\frac{1}{2}, \frac{1}{\xi}\right),$$

$$\lambda_j^2 \mid \nu_j \sim IG\left(\frac{1}{2}, \frac{1}{\nu_j}\right),$$

$$\xi, \nu_j \sim IG\left(\frac{1}{2}, 1\right),$$
(20)

where the inverse matrix of F is given by

$$\boldsymbol{F}^{-1} = \begin{pmatrix} \frac{1}{\tau_1^2} + \frac{1}{\lambda_2^2 \tilde{\tau}^2} & -\frac{1}{\lambda_2^2 \tilde{\tau}^2} & 0 & \dots & 0 & 0\\ -\frac{1}{\lambda_2^2 \tilde{\tau}^2} & \frac{1}{\tau_2^2} + \frac{1}{\lambda_2^2 \tilde{\tau}^2} + \frac{1}{\lambda_3^2 \tilde{\tau}^2} & -\frac{1}{\lambda_3^2 \tilde{\tau}^2} & \dots & 0 & 0\\ 0 & -\frac{1}{\lambda_3^2 \tilde{\tau}^2} & \frac{1}{\tau_3^2} + \frac{1}{\lambda_3^2 \tilde{\tau}^2} + \frac{1}{\lambda_4^2 \tilde{\tau}^2} & \dots & 0 & 0\\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots\\ 0 & 0 & 0 & \dots & \frac{1}{\tau_{p-1}^2} + \frac{1}{\lambda_{p-1}^2 \tilde{\tau}^2} + \frac{1}{\lambda_p^2 \tilde{\tau}^2} & -\frac{1}{\lambda_p^2 \tilde{\tau}^2} \\ 0 & 0 & 0 & \dots & -\frac{1}{\lambda_p^2 \tilde{\tau}^2} & \frac{1}{\tau_p^2} + \frac{1}{\lambda_p^2 \tilde{\tau}^2} \end{pmatrix}$$

By assuming a horseshoe prior on differences between adjacent regression coefficients, small differences can be shrunk more, while large differences are shrunk less, compared to a Laplace prior.

From the priors (20) and the data-augmentation method, we can get the full con-

ditional distributions as follows:

$$\beta \mid \boldsymbol{y}, \boldsymbol{X}, \beta_0, \tau_1^2, \dots, \tau_p^2, \lambda_2^2, \dots, \lambda_p^2, \tilde{\tau}^2, w_1, \dots, w_n \sim \mathcal{N}_p(\boldsymbol{A}^{-1}\boldsymbol{X}^T \boldsymbol{W}(\boldsymbol{z} - \beta_0 \mathbf{1}), \boldsymbol{A}^{-1}),$$
$$w_i \mid \boldsymbol{x}_i, \boldsymbol{\beta}, \beta_0 \sim \mathcal{PG}(1, \beta_0 + \boldsymbol{x}_i^T \boldsymbol{\beta}),$$
$$\frac{1}{\tau_j^2} \mid \beta_j, \tilde{\lambda}_1^2 \sim \mathcal{IGauss}\left(\sqrt{\frac{\tilde{\lambda}_1^2}{\beta_j^2}}, \tilde{\lambda}_1^2\right),$$
$$\tilde{\lambda}_1^2 \mid \tau_1^2, \dots, \tau_p^2 \sim \mathcal{Ga}\left(p + r_1, \frac{1}{2}\sum_{j=1}^p \tau_j^2 + \delta_1\right),$$

$$\begin{split} \tilde{\tau}^2 \mid \beta_1, \dots, \beta_p, \lambda_2^2, \dots, \lambda_p^2, \xi \sim \mathrm{IG} \left(\frac{p}{2}, \frac{1}{2} \sum_{j=2}^p \frac{(\beta_j - \beta_{j-1})^2}{\lambda_j^2} + \frac{1}{\xi} \right), \\ \lambda_j^2 \mid \beta_j, \beta_{j-1}, \tilde{\tau}^2, \nu_j \sim \mathrm{IG} \left(1, \frac{(\beta_j - \beta_{j-1})^2}{2\tilde{\tau}^2} + \frac{1}{\nu_j} \right), \\ \nu_j \mid \lambda_j^2 \sim \mathrm{IG} \left(1, \frac{1}{\lambda_j^2} + 1 \right), \\ \xi \mid \tilde{\tau}^2 \sim \mathrm{IG} \left(1, \frac{1}{\lambda_j^2} + 1 \right), \\ \beta_0 \mid \boldsymbol{y}, \boldsymbol{X}, \tau_1^2, \dots, \tau_p^2, \lambda_2^2, \dots, \lambda_p^2, \tilde{\tau}^2, w_1, \dots, w_n \sim \mathrm{N} \left(\frac{1}{S} \sum_{i=1}^n v_i, \frac{1}{S} \right), \end{split}$$

where $\mathbf{A} = \mathbf{X}^T \mathbf{W} \mathbf{X} + \mathbf{F}^{-1}$, $\mathbf{v} = (\mathbf{z} - \mathbf{X}\boldsymbol{\beta}) \odot \mathbf{W}$, and $S = \sum_{i=1}^n w_i$. Note that we did not assume a horseshoe prior on regression coefficients, because the MCMC chain did not converge.

5. Numerical studies

In this section, we describe the settings and results of Monte Carlo simulations and compare the performance of our proposed methods with the existing method. We also apply our proposed methods to anomaly detection in the field of semiconductor microelectronics manufacturing (Olszewski 2001; Deng et al. 2014).

5.1. Monte Carlo simulations

We generated y_i (i = 1, 2, ..., n) according to the true model:

$$\Pr(y_i = 1 \mid \boldsymbol{x}_i) = \frac{1}{1 + e^{-\boldsymbol{x}_i^T \boldsymbol{\beta}^*}},$$
(21)

where $\boldsymbol{\beta}^* = (\beta_1^*, \beta_2^*, \dots, \beta_p^*)^T$ is a *p*-dimensional regression coefficient vector. The explanatory variable $\boldsymbol{x}_i \ (i = 1, 2, \dots, n)$ followed the multivariate normal distribution $N_p(\mathbf{0}_p, \Sigma)$. For $\boldsymbol{\beta}^*$ and Σ , we considered the following cases:

Case 1: $\beta^* = \beta_1^* \text{ or } \beta_2^*$, $\Sigma_{ii} = 1$, $\Sigma_{ij} = \rho$, $(i \neq j)$, Case 2: $\beta^* = \beta_1^* \text{ or } \beta_2^*$, $\Sigma_{ii} = 1$, $\Sigma_{ij} = \begin{cases} 0.5 & (\beta_i^* = \beta_j^* \text{ and } 1 \leq |i - j| \leq 4) \\ 0 & \text{otherwise} \end{cases}$ $(i \neq j)$, Case 3: $\beta^* = \beta_1^* \text{ or } \beta_2^*$, $\Sigma_{ii} = 1$, $\Sigma_{ij} = \begin{cases} 0.5^{|i-j|} & (\beta_i^* = \beta_j^* \text{ and } 1 \leq |i - j| \leq 4) \\ 0 & \text{otherwise} \end{cases}$ $(i \neq j)$, Case 4: $\beta^* = (\mathbf{1.0}_{20}^T, -\mathbf{1.0}_{20}^T, \mathbf{0.0}_{170}^T, \mathbf{1.5}_{20}^T, \mathbf{0.0}_{170}^T)^T$, $\Sigma_{ii} = 1$, $\Sigma_{ij} = 0$, $(i \neq j)$,

where $\beta_1^* = (\mathbf{1.0}_5^T, \mathbf{0.0}_5^T, \mathbf{1.0}_5^T, \mathbf{0.0}_5^T)^T$, $\beta_2^* = (-\mathbf{1.0}_5^T, \mathbf{2.0}_5^T, \mathbf{1.0}_5^T, \mathbf{0.0}_5^T)^T$, Σ_{ij} is the (i, j)-th element of Σ , and $\rho = 0.0, 0.5$. We considered n = 500 for Cases 1, 2, and 3, which correspond to n > p cases, while n = 300 for Case 4, which corresponds to an n < p case. We simulated 100 datasets for each case. Note that Case 4 is based on the simulation in Bhattacharyya et al. (2022).

We compared our proposed methods, which are the logistic regression model with the Bayesian fused lasso (LBFL) and that with the Bayesian fused lasso with horseshoe prior (LBFH), to the logistic regression model with the fused lasso (LFL). For LFL, we used the package penalized of the software \mathbf{R} , which is available from https://cran.r-project.org/web/packages/penalized/index.html. The values of the hyper-parameters λ_1 and λ_2 for LFL were selected by the Bayesian information criterion (BIC). As the term of the degrees of freedom in BIC, we used the number of groups which consist of non-zero fused estimated regression coefficients (Tibshirani et al. 2005; Tibshirani and Taylor 2011). For LBFL and LBFH, the Gibbs sampling was run with 10,000 iterations, and then the first 6,000 iterations were discarded as burn-in.

To measure the accuracy of the estimation of regression coefficients, we computed the mean squared error (MSE):

$$MSE = \frac{1}{100} \sum_{k=1}^{100} \left[(\hat{\beta}_0^{(k)})^2 + \left\{ \hat{\beta}^{(k)} - \beta^* \right\}^T \left\{ \hat{\beta}^{(k)} - \beta^* \right\} \right],$$
(22)

where $\hat{\beta}_0^{(k)}$ and $\hat{\beta}^{(k)} = (\hat{\beta}_1^{(k)}, \dots, \hat{\beta}_p^{(k)})^T$ are an intercept and a vector of regression coefficients estimated from the k-th dataset, respectively. To evaluate the prediction accuracy, we used the negative expected log-likelihood. We generated 1,000 test data, and then computed the mean of the empirical negative expected log-likelihood:

$$\mathrm{EL} = \frac{1}{100} \sum_{k=1}^{100} \left\{ -\frac{1}{1000} \sum_{j=1}^{1000} \left[y_j^{\dagger(k)} (\boldsymbol{x}_j^{\dagger(k)T} \hat{\boldsymbol{\beta}}^{(k)} + \hat{\boldsymbol{\beta}}_0^{(k)}) - \log\{1 + \exp(\hat{\boldsymbol{\beta}}^{(k)} + \hat{\boldsymbol{\beta}}_0^{(k)})\} \right] \right\},$$
(23)

where $y_j^{\dagger(k)}$ and $x_j^{\dagger(k)}$ are the *j*-th data in the test data for the *k*-th dataset. To assess the performance of variable selection and variable fusion, we computed

the following measures:

$$\begin{aligned} \mathrm{PV} &= \frac{1}{100} \sum_{k=1}^{100} \frac{|\{j \mid (\beta_j^* \neq 0) \land (0 \notin \mathrm{CIV}_j(k))\}|}{|\{j \mid \beta_j^* \neq 0\}|} \quad (1 \leq j \leq p), \\ \mathrm{PZV} &= \frac{1}{100} \sum_{k=1}^{100} \frac{|\{j \mid (\beta_j^* = 0) \land (0 \in \mathrm{CIV}_j(k))\}|}{|\{j \mid \beta_j^* = 0\}|} \quad (1 \leq j \leq p), \\ \mathrm{AV} &= \frac{1}{100} \sum_{k=1}^{100} \frac{|\{j \mid (\beta_j^* - \beta_{j-1}^* \neq 0) \land (0 \notin \mathrm{CIV}_j(k))\}| + |\{j \mid (\beta_j^* = 0) \land (0 \in \mathrm{CIV}_j(k))\}|}{p} \quad (1 \leq j \leq p), \\ \mathrm{PF} &= \frac{1}{100} \sum_{k=1}^{100} \frac{|\{j \mid (\beta_j^* - \beta_{j-1}^* \neq 0) \land (0 \notin \mathrm{CIF}_j(k))\}|}{|\{j \mid \beta_j^* - \beta_{j-1}^* \neq 0\}|} \quad (2 \leq j \leq p), \\ \mathrm{PNF} &= \frac{1}{100} \sum_{k=1}^{100} \frac{|\{j \mid (\beta_j^* - \beta_{j-1}^* = 0) \land (0 \in \mathrm{CIF}_j(k))\}|}{|\{j \mid \beta_j^* - \beta_{j-1}^* = 0\}|} \quad (2 \leq j \leq p), \\ \mathrm{AF} &= \frac{1}{100} \sum_{k=1}^{100} \frac{|\{j \mid (\beta_j^* - \beta_{j-1}^* \neq 0) \land (0 \notin \mathrm{CIF}_j(k))\}|}{|\{j \mid \beta_j^* - \beta_{j-1}^* = 0\}|} \quad (2 \leq j < p). \end{aligned}$$

where $\operatorname{CIV}_j(k)$ denotes the 95% credible interval for $\hat{\beta}_j^{(k)}$, while $\operatorname{CIF}_j(k)$ denotes the 50% credible interval for the difference between $\hat{\beta}_j^{(k)}$ and $\hat{\beta}_{j-1}^{(k)}$. PV and PZV measure the accuracy of estimating regression coefficients when the corresponding true regression coefficients are non-zero and zero, respectively. PF and PNF measure the accuracy of the differences between estimated adjacent regression coefficients when the differences between the true adjacent regression coefficients are non-zero and zero, respectively. AV and AF measure the accuracy of variable selection and variable fusion, respectively.

Note that we utilized credible intervals of MCMC samples in evaluating the performance of variable selection and variable fusion. The reason why LBFL and LBFH do not estimate regression coefficients and their differences as exactly zero is that the posteriors of them are continuous. For variable selection, we computed 95% credible intervals as in Bhattacharyya et al. (2022), and then judged that the regression coefficient is estimated as zero when the credible interval of its MCMC samples includes zero. For variable fusion, we computed 50% credible intervals of the differences between adjacent regression coefficients as in Banerjee (2022), and then judged that the difference is estimated as zero, similar to variable selection.

Table 1.: MSE (standard deviation), EL, PV, PZV, AV, PF, PNF, and AF for Case 1 and $\rho = 0$. Bold font indicates the smallest value of MSE and EL and the largest value of PV, PZV, AV, PF, PNF, and AF before rounding among LFL, LBFL, and LBFH.

		MSE	EL	PV	PZV	AV	\mathbf{PF}	PNF	AF
		(sd)	(sd)	(sd)	(sd)	(sd)	(sd)	(sd)	(sd)
	IFI	0.619	180.551	1.000	0.501	0.751	1.000	0.766	0.803
		(0.287)	(3.032)	(0.000)	(0.288)	(0.144)	(0.000)	(0.113)	(0.095)
A *	I DEI	0.490	183.965	1.000	0.911	0.956	1.000	0.541	0.613
ρ_1	LDFL	(0.303)	(4.997)	(0.000)	(0.094)	(0.047)	(0.000)	(0.143)	(0.120)
	IDDU	0.252	178.379	1.000	0.961	0.981	1.000	0.850	0.874
	LDFП	(0.167)	(3.135)	(0.000)	(0.076)	(0.038)	(0.000)	(0.105)	(0.089)
	ITI	3.722	123.790	1.000	0.480	0.870	1.000	0.769	0.805
		(0.898)	(3.398)	(0.000)	(0.417)	(0.104)	(0.000)	(0.101)	(0.085)
Q *	IDFI	0.905	122.723	1.000	0.944	0.986	1.000	0.541	0.613
$ ho_2$	LBFL	(0.494)	(3.846)	(0.000)	(0.107)	(0.027)	(0.000)	(0.135)	(0.114)
	IDFU	0.809	117.895	1.000	0.972	0.993	1.000	0.868	0.888
	LDFT	(0.522)	(3.084)	(0.000)	(0.103)	(0.026)	(0.000)	(0.083)	(0.069)

Table 2.: MSE (standard deviation), EL, PV, PZV, AV, PF, PNF, and AF for Case 1 and $\rho = 0.5$. Bold font indicates the smallest value of MSE and EL and the largest value of PV, PZV, AV, PF, PNF, and AF before rounding among LFL, LBFL, and LBFH.

		MSE	EL	PV	PZV	AV	\mathbf{PF}	PNF	AF
		(sd)	(sd)	(sd)	(sd)	(sd)	(sd)	(sd)	(sd)
	IFI	1.735	94.449	1.000	0.236	0.618	0.940	0.901	0.907
		(0.634)	(4.134)	(0.000)	(0.261)	(0.130)	(0.137)	(0.066)	(0.065)
A *	IBEI	1.716	96.289	0.979	0.920	0.950	0.967	0.581	0.642
$ ho_1$	LDLL	(0.870)	(5.362)	(0.048)	(0.090)	(0.051)	(0.101)	(0.145)	(0.126)
	IDEII	1.062	92.329	0.995	0.905	0.950	0.970	0.871	0.887
	LDFII	(0.480)	(3.813)	(0.022)	(0.134)	(0.067)	(0.096)	(0.089)	(0.078)
	IFI	6.180	91.536	1.000	0.482	0.871	0.980	0.847	0.868
		(1.205)	(3.603)	(0.000)	(0.447)	(0.112)	(0.080)	(0.097)	(0.083)
Q *	IDFI	1.742	88.395	0.977	0.930	0.966	0.977	0.598	0.657
ρ_2	LDLL	(0.791)	(3.967)	(0.037)	(0.104)	(0.039)	(0.085)	(0.125)	(0.107)
	IDEII	1.520	84.530	0.997	0.966	0.989	0.970	0.881	0.895
	LDFH	(0.914)	(3.086)	(0.015)	(0.110)	(0.029)	(0.107)	(0.091)	(0.079)

Table 3.: MSE (standard deviation), EL, PV, PZV, AV, PF, PNF, and AF for Case 2. Bold font indicates the smallest value of MSE and EL and the largest value of PV, PZV, AV, PF, PNF, and AF before rounding among LFL, LBFL, and LBFH.

		MSE	EL	PV	PZV	AV	PF	PNF	AF
		(sd)							
	IFI	0.718	118.113	1.000	0.481	0.741	1.000	0.898	0.914
		(0.297)	(2.373)	(0.000)	(0.306)	(0.153)	(0.000)	(0.083)	(0.070)
Q *	IDEI	0.949	121.116	0.999	0.937	0.968	0.993	0.581	0.646
$ ho_1$	LDLL	(0.435)	(3.396)	(0.010)	(0.087)	(0.045)	(0.047)	(0.148)	(0.125)
	IDEII	0.381	116.649	1.000	0.985	0.993	1.000	0.888	0.905
	LDFII	(0.194)	(2.220)	(0.000)	(0.046)	(0.023)	(0.000)	(0.094)	(0.079)
	IFI	5.350	80.432	1.000	0.364	0.841	1.000	0.846	0.871
		(0.790)	(2.902)	(0.000)	(0.424)	(0.106)	(0.000)	(0.078)	(0.066)
Q *	IDEI	1.860	79.103	0.982	0.942	0.972	0.980	0.561	0.627
ρ_2	LDL L	(0.894)	(4.442)	(0.031)	(0.118)	(0.040)	(0.080)	(0.138)	(0.117)
	IDFU	1.375	74.886	0.994	0.990	0.993	0.987	0.871	0.889
	LDFT	(0.730)	(3.396)	(0.019)	(0.044)	(0.019)	(0.066)	(0.094)	(0.082)

Table 4.: MSE (standard deviation), EL, PV, PZV, AV, PF, PNF, and AF for Case 3. Bold font indicates the smallest value of MSE and EL and the largest value of PV, PZV, AV, PF, PNF, and AF before rounding among LFL, LBFL, and LBFH.

		MSE	EL	PV	PZV	AV	\mathbf{PF}	PNF	AF
		(sd)	(sd)	(sd)	(sd)	(sd)	(sd)	(sd)	(sd)
	IFI	0.690	134.037	1.000	0.465	0.733	1.000	0.846	0.870
		(0.243)	(2.457)	(0.000)	(0.254)	(0.127)	(0.000)	(0.089)	(0.075)
A *	IDEI	0.815	137.327	0.999	0.925	0.962	1.000	0.564	0.633
$ ho_1$	LDLL	(0.458)	(4.360)	(0.010)	(0.093)	(0.046)	(0.000)	(0.147)	(0.123)
	IDEII	0.297	132.313	1.000	0.982	0.991	1.000	0.883	0.902
	LDFII	(0.167)	(2.760)	(0.000)	(0.046)	(0.023)	(0.000)	(0.086)	(0.072)
	IFI	4.681	89.716	1.000	0.440	0.860	1.000	0.807	0.837
		(0.973)	(3.554)	(0.000)	(0.405)	(0.101)	(0.000)	(0.094)	(0.080)
Q *	IDFI	1.531	88.550	0.996	0.940	0.982	0.993	0.626	0.684
$ ho_2$	LBL	(0.923)	(4.763)	(0.019)	(0.104)	(0.029)	(0.047)	(0.149)	(0.127)
	IDEII	1.030	84.256	0.999	0.980	0.994	1.000	0.904	0.919
	LDFП	(0.657)	(3.792)	(0.009)	(0.060)	(0.016)	(0.000)	(0.077)	(0.065)

Table 5.: MSE (standard deviation), EL, PV, PZV, AV, PF, PNF, and AF for Case 4. Bold font indicates the smallest value of MSE and EL and the largest value of PV, PZV, AV, PF, PNF, and AF before rounding among LFL, LBFL, and LBFH.

	MSE	EL	PV	PZV	AV	PF	PNF	AF
	(sd)	(sd)	(sd)	(sd)	(sd)	(sd)	(sd)	(sd)
IFI	66.735	124.512	0.998	0.947	0.954	0.890	0.989	0.988
ΓL	(0.984)	(3.965)	(0.008)	(0.061)	(0.051)	(0.144)	(0.005)	(0.005)
IDFI	42.765	115.630	0.578	0.999	0.936	0.840	0.924	0.923
$\Gamma DL \Gamma$	(1.639)	(11.928)	(0.068)	(0.001)	(0.010)	(0.157)	(0.013)	(0.013)
IDEII	60.976	109.058	0.857	1.000	0.978	0.563	0.999	0.994
LDFΠ	(1.855)	(10.577)	(0.151)	(0.0004)	(0.023)	(0.231)	(0.002)	(0.004)

The results are summarized in Tables 1, 2, 3, 4, and 5. LBFH gives the smallest ELs in all cases. LBFH also gives the smallest MSEs in almost all cases except for Case 4. These results show that LBFH outperformed LFL and LBFL in terms of the estimation and prediction accuracy. In addition, LBFH achieves the largest AVs and AFs in most cases. This shows that LBFH provides superior performance of variable selection and variable fusion compared to the other methods. Comparing LBFL to LFL, LBFL gives the smaller MSEs and ELs in more than half of the cases. LFL often gives larger PVs than LBFL, but LBFL gives larger PZVs and AVs in almost all cases. These results show that LBFL performs better than LFL in terms of the accuracy of estimation, prediction, and variable selection. For the performance of variable fusion, PFs, PNFs, and AFs of LFL are competitive or larger than those of LBFL.

5.2. Application

We applied our proposed methods LBFL and LBFH to the Wafer dataset, which was formatted in Olszewski (2001) and analyzed in Deng et al. (2014). The dataset can be obtained from the UCR Time Series Classification Archive (https://www.cs.ucr.edu/%7Eeamonn/time_series_data_2018). Utilizing the dataset, the normal and abnormal etching processes of a wafer in semiconductor microelectronics manufacturing were classified based on time series data from six sensors (which monitor radio frequency forward power, radio frequency reflected power, chamber pressure, 405 nanometer emission, 520 nanometer emission, and direct current bias, respectively). Each time series data contains the value from one of the six sensors for one wafer and has length p = 152. We labeled abnormal data as one and normal data as zero. The dataset contains n = 1,000 training data and 6,164 test data. The abnormal data constitute 10.7% of the training data and 12.1% of the test data, meaning that the dataset has a large class imbalance.

We compared LBFL and LBFH to LFL. As in Deng et al. (2014), we selected the hyper-parameters λ_1 and λ_2 for LFL from four candidates, 0.05, 0.1, 0.3, and 0.5, by BIC. For LBFL and LBFH, the Gibbs sampler was run with 10,000 iterations, and then the first 6,000 samples were discarded as burn-in.

We evaluated the performance of LBFL, LBFH, and LFL by Area Under the ROC Curve (AUC). AUCs tend to be large in situations where the true positive rate is large when the false positive rate is small. The values of AUC are summarized in Table 6. From Table 6, LBFL gives the largest AUC. Meanwhile, AUC is not enough to evaluate the performance of the model when there is a large class imbalance in the dataset. In the case of such a large class imbalance, Area Under the Precision-Recall Curve (PR-AUC) is a more suitable indicator. PR-AUCs tend to be large in situations where the true positive rate is large when the precision is large. For the details of PR-AUC, we refer the reader to Davis and Goadrich (2006). The values of PR-AUC are also summarized in Table 6, which shows that LBFH gives the largest PR-AUC.

As with the Monte Carlo simulations in Sec. 5.1, we judged that the regression coefficient is estimated as zero if the corresponding 95% credible interval includes zero for LBFL and LBFH. We also judged that the difference between adjacent regression coefficients is regarded as non-zero when the 50% credible interval for the difference does not include zero. The estimated regression coefficients are shown in Figure 1. From Figure 1, we observe that LFL estimated 60 regression coefficients as zero, whereas LBFL estimated the most as zero, at 133, and LBFH estimated the second most, at 127. Thus, the number of points which are considered to be unnecessary for the model

is larger for LBFL and LBFH than for LFL, meaning that the former two make clearer which points are necessary for the prediction.

From Figure 1(c), we see that LBFH split the regression coefficients into four groups. The first group contains the 1st to 27th coefficients, the second contains the 28th to 35th coefficients, the third one the 36th to 44th, and the fourth one the 45th to 152nd. On the other hand, from Figure 1(b), we see that LBFL split the regression coefficients into 21 groups, whereas from Figure 1(a), we see that LFL split the coefficients into 57 groups. Thus, LBFH gave the smallest number of groups of variables, whereas LBFL gave the second smallest. Based on the number of detected groups, LBFH seems to provide smoother and more simplified estimation than the other methods. Many of the groups detected by LFL contain only one variable, meaning that LFL has weak performance regarding grouping multiple variables.

Table 6.: AUC and PR-AUC for the Wafer dataset. Bold font indicates the largest value among LFL, LBFL, and LBFH.

	AUC	PR-AUC
LFL	0.880	0.615
LBFL	0.886	0.592
LBFH	0.864	0.626



Figure 1.: Estimated regression coefficients for LFL, LBFL, and LBFH. Dots indicate the values of the estimated regression coefficients. For LFL, a red dot indicates a zero element of the estimated regression coefficients and a black vertical line indicates where a non-zero difference between adjacent estimated regression coefficients exists. For LBFL and LBFH, a red dot indicates an estimated regression coefficient whose MCMC samples give a 95% credible interval including zero and a black vertical line indicates where a difference between adjacent regression coefficients where a difference between adjacent regression coefficients where a 50% credible interval not including zero exists.

6. Conclusions

We proposed a logistic regression model which enables variable selection by a Laplace prior and variable fusion by a Laplace prior or a horseshoe prior. Using a dataaugmentation method with a Pólya-Gamma distribution and the hierarchical structure of the priors, we developed the Gibbs sampler. Through numerical studies, we showed that our proposed methods gave more accurate estimation and prediction than the existing method. Our proposed methods also outperformed the existing method in terms of variable selection and variable fusion without the selection of the hyper-parameters.

We used credible intervals for the evaluation of the performance of variable selection and variable fusion for Bayesian methods. It would be interesting to investigate the theoretical aspects of credible intervals for determining whether the target can be considered as zero or not. We leave that as future work.

Acknowledgements

Y. K. was supported by JST, Establishment of University Fellowships towards the Creation of Science Technology Innovation, Grant Number JP-MJFS2136. S. K. was supported by JSPS KAKENHI Grant Numbers JP23K11008, JP23H03352, and JP23H00809. Computational resources were provided by the Super Computer System, Human Genome Center, Institute of Medical Science, The University of Tokyo. The authors thank FORTE Science Communications (https://www.forte-science.co.jp/) for English language editing.

References

- Andrews DF, Mallows CL (1974) Scale mixtures of normal distributions. Journal of the Royal Statistical Society: Series B (Methodological) 36(1):99–102
- Banerjee S (2022) Horseshoe shrinkage methods for Bayesian fusion estimation. Computational Statistics and Data Analysis 174:107450
- Betancourt B, Rodríguez A, Boyd N (2017) Bayesian fused lasso regression for dynamic binary networks. Journal of Computational and Graphical Statistics 26(4):840–850
- Bhattacharyya A, Pal S, Mitra R, Rai S (2022) Applications of Bayesian shrinkage prior models in clinical research with categorical responses. BMC Medical Research Methodology 22(1):1–19
- Carvalho CM, Polson NG, Scott JG (2010) The horseshoe estimator for sparse signals. Biometrika 97(2):465–480
- Davis J, Goadrich M (2006) The relationship between Precision-Recall and ROC curves. In: Proceedings of the 23rd international conference on Machine learning, pp 233–240
- Deng H, Baydogan MG, Runger G (2014) SMT: Sparse multivariate tree. Statistical Analysis and Data Mining: The ASA Data Science Journal 7(1):53–69
- Genkin A, Lewis DD, Madigan D (2007) Large-scale Bayesian logistic regression for text categorization. Technometrics 49(3):291–304
- Kyung M, Gill J, Ghosh M, Casella G (2010) Penalized regression, standard errors, and Bayesian lassos. Bayesian Analysis 5(2):369–411
- Lokhorst J (1999) The lasso and generalised linear models. Honors Project, The University of Adelaide, Australia
- Makalic E, Schmidt DF (2015) A simple sampler for the horseshoe estimator. IEEE Signal Processing Letters 23(1):179–182

- Makalic E, Schmidt DF (2016) High-dimensional Bayesian regularised regression with the BayesReg package. arXiv preprint arXiv:161106649
- Olszewski RT (2001) Generalized feature extraction for structural pattern recognition in timeseries data. PhD thesis, Carnegie Mellon University
- Park T, Casella G (2008) The Bayesian lasso. Journal of the American Statistical Association 103(482):681–686
- Pereira JM, Basto M, da Silva AF (2016) The logistic lasso and ridge regression in predicting corporate failure. Procedia Economics and Finance 39:634–641
- Polson NG, Scott JG, Windle J (2013) Bayesian inference for logistic models using Pólya– Gamma latent variables. Journal of the American Statistical Association 108(504):1339– 1349
- Shevade SK, Keerthi SS (2003) A simple and efficient algorithm for gene selection using sparse logistic regression. Bioinformatics 19(17):2246–2253
- Tian Y, Bondell HD, Wilson A (2019) Bayesian variable selection for logistic regression. Statistical Analysis and Data Mining: The ASA Data Science Journal 12(5):378–393
- Tibshirani R (1996) Regression shrinkage and selection via the lasso. Journal of the Royal Statistical Society: Series B (Methodological) 58(1):267–288
- Tibshirani R, Saunders M, Rosset S, Zhu J, Knight K (2005) Sparsity and smoothness via the fused lasso. Journal of the Royal Statistical Society: Series B (Statistical Methodology) 67(1):91–108
- Tibshirani RJ, Taylor J (2011) The solution path of the generalized lasso. The Annals of Statistics 39(3):1335 1371,
- Wand MP, Ormerod JT, Padoan SA, Frühwirth R (2011) Mean field variational Bayes for elaborate distributions. Bayesian Analysis 6(4):847–900
- Wu TT, Chen YF, Hastie T, Sobel E, Lange K (2009) Genome-wide association analysis by lasso penalized logistic regression. Bioinformatics 25(6):714–721
- Xin B, Kawahara Y, Wang Y, Gao W (2014) Efficient generalized fused lasso and its application to the diagnosis of Alzheimer's disease. In: Proceedings of the AAAI Conference on Artificial Intelligence, pp 28, 2163–2169
- Yu D, Lee SJ, Lee WJ, Kim SC, Lim J, Kwon SW (2015) Classification of spectral data using fused lasso logistic regression. Chemometrics and Intelligent Laboratory Systems 142:70–77