Natural Gradient Variational Bayes without Fisher Matrix Analytic Calculation and Its Inversion

A. Godichon-Baggioni

Laboratoire de Probabilités Statistique et Modélisation Sorbonne-Université, 75005, Paris, France Email: antoine.godichon_baggioni@sorbonne-universite.fr D. Nguyen Department of Mathematics Marist College, 3399 North Road Poughkeepsie NY 12601, United States Email: nducduy@gmail.com M-N. Tran Business Analytics discipline The University of Sydney Business School Australia Email: minh-ngoc.tran@sydney.edu.au

April 29, 2024

Abstract

This paper introduces a method for efficiently approximating the inverse of the Fisher information matrix, a crucial step in achieving effective variational Bayes inference. A notable aspect of our approach is the avoidance of analytically computing the Fisher information matrix and its explicit inversion. Instead, we introduce an iterative procedure for generating a sequence of matrices that converge to the inverse of Fisher information. The natural gradient variational Bayes algorithm without analytic expression of the Fisher matrix and its inversion is provably convergent and achieves a convergence rate of order $\mathcal{O}(\log s/s)$, with s the number of iterations. We also obtain a central limit theorem for the iterates. Implementation of our method does not require storage of large matrices, and achieves a linear complexity in the number of variational parameters. Our algorithm exhibits versatility, making it applicable across a diverse array of variational Bayes domains, including

Gaussian approximation and normalizing flow Variational Bayes. We offer a range of numerical examples to demonstrate the efficiency and reliability of the proposed variational Bayes method.

Keywords: Bayesian computation, Stochastic gradient descent, Bayesian neural network, Normalizing flow

1 Introduction

The growing complexity of models used in modern statistics and machine learning has spurred the demand for more efficient Bayesian estimation techniques. Among the array of Bayesian tools available, Variational Bayes [Waterhouse et al., 1995, Jordan et al., 1999] has gained prominence as a remarkably versatile alternative to traditional Monte Carlo methods for tackling statistical inference in intricate models. Variational Bayes (VB) operates by approximating the posterior probability distribution using a member selected from a family of tractable distributions, characterized by variational parameters. The optimal member is determined through minimization of the Kullback-Leibler divergence, which quantifies the disparity between the chosen candidate and the posterior distribution. VB is a fast alternative to Markov chain Monte Carlo (MCMC) methods, and has found diverse applications, encompassing variational autoencoders [Kingma and Welling, 2013], text analysis [Hoffman et al., 2013], Bayesian synthetic likelihood [Ong et al., 2018], deep neural networks [Graves, 2011, Tran et al., 2020], to name a few. For recent advances in the field of VB and Bayesian approximation in general, please refer to the excellent survey papers of Blei et al. [2017] and Martin et al. [2023].

VB turns the Baysesian inference problem into an optimization problem, and a large class of VB methods use stochastic gradient descent (SGD) as their backbone. In the recent decades, a great deal of effort has been devoted to developing and improving optimization algorithms for big and high dimensional data. As a result, various first order stochastic optimization algorithms have been developed in response to these new demands; notable examples include AdaGrad of Duchi et al. [2011], Adam of Kingma and Ba [2014], Adadelta of Zeiler [2012], and their variance reduction variations [Defazio et al., 2014, Johnson and Zhang, 2013, Nguyen et al., 2017]. For a detailed discussion on stochastic optimization, please refer to the excellent books of Kushner and Yin [2003], Goodfellow et al. [2016] and Murphy [2012].

Gradient descent methods in VB rely on the gradient of the objective lower bound function, whose definition depends upon the metric on the variational parameter space. Optimization in conventional VB methods uses the Euclidean gradient defined using the usual Euclidean metric. It turns out that the natural gradient, the term coined by Amari [1998], represents a more adequate direction of ascent in the VB context as it takes into account the information geometry of the variational family [Martens, 2020, Khan and Lin, 2017]. The natural gradient is defined using the Fisher-Rao metric, which resembles the Kullback-Leibler divergence between probability distributions parameterized by the variational parameters. More precisely, the natural gradient is the steepest ascent direction of the objective function on the variational parameter space equipped with the Fisher-Rao metric. Martens [2020] sheds light on the concept that natural gradient descent can be viewed as a second-order optimization method where the Fisher information assumes the role of the Hessian matrix. Because of this, natural gradients take into account the curvature information (through the Fisher-Rao metric) of the variational parameter space; therefore the number of iteration steps required to find a local optimum is often found significantly reduced [Tran et al., 2017]. According to Tan [2021], stochastic optimization guided by natural gradients has proven more resilient, capable of circumventing or escaping plateaus, ultimately resulting in faster convergence; see also[Rattray et al., 1998,

Hoffman et al., 2013, Khan and Lin, 2017, Wilkinson et al., 2023].

The natural gradient is calculated by pre-multiplying the Euclidean gradient of the lower bound function with the inverse Fisher information matrix, a process that is notably intricate. Computing the Fisher matrix, not to mention its inverse, is challenging. In the realm of Gaussian variational approximation, where the posterior is approximated by a Gaussian distribution, the natural gradient can be calculated efficiently. Tran et al. [2020] consider a factor structure for the covariance matrix, and derive a closed-form approximation for the natural gradient. Tan [2021] employs a Cholesky factor structure for the covariance matrix and the precision matrix, and derives an analytic natural gradient; see also Khan and Lin [2017] and Magris et al. [2022]. On the other hand, for broader cases where the variational distribution is based on neural networks, Martens and Grosse [2015] approximate the Fisher matrix with a block diagonal matrix. It is important to highlight that existing techniques for computing the natural gradient are primarily restricted to certain contexts (like the Gaussian variational approximations mentioned above) or are heavily dependent on simplified approximations (such as employing a blockdiagonal matrix). These constraints restrict the broader application of natural gradients. For a large class of VB methods, e.g., when the variational distribution is a mixture [Giordani et al., 2013], a copula [Gunawan et al., 2023] or a normalizing flow [Rezende and Mohamed, 2015, it is challenging to use the natural gradient as the Fisher matrix is not available.

This paper makes several important contributions that significantly improve the natural gradient VB method. First, we present an approach for efficiently approximating the *inverse* of Fisher information matrix. We emphasize that there are two main difficulties in calculating the natural gradient: (i) analytical calculation of the Fisher matrix that often involves intractable expectations, and (ii) computing its matrix inversion. A notable aspect of our approach is the avoidance of these two difficulties altogether. Instead, we introduce an iterative procedure for generating a sequence of positive definite matrices that converge to the inverse of Fisher information. Pre-multiplying the Euclidean gradient with these matrices provides estimates of the natural gradient. Our method of approximating the natural gradient is general, easy to implement, asymptotically exact and applies to any variational distribution including Gaussian distributions, mixtures and normalizing flow based distributions. It is important to note that, for high-dimensional applications, implementation of our method does not require storage of large matrices because the estimate of inverse Fisher matrix can be written using outer products. Second, we propose a VB method that streamlines the natural gradient estimation without matrix inversion within the VB training iteration. This leads to an efficient natural gradient VB algorithm, referred to as inversion-free variational Bayes (IFVB). We also present a weighted averaged estimate version of IFVB, called AIFVB, that converges faster than IFVB. Both IFVB and AIFVB are provably convergent, with AIFVB being shown asymptotically efficient and achieving a central limit theorem. Third, to substantiate the effectiveness and robustness of our proposed method, we offer a range of numerical examples to demonstrate its efficiency and reliability.

The rest of the paper is organized as follows: Section 2 provides a brief overview of the variational Bayesian inference problem. Section 3 presents natural gradient and discusses its advantages as well as its computational difficulty. We introduce inversion free natural variational Bayes in Section 4. Section 5 is concerned with convergence analysis. Numerical examples are provided in Section 6. Section 7 concludes the paper. Section 8 provides more numerical examples, and Section 9 contains the proofs of the main theorems; further technical details are in the Appendix.

Notation. We denote by $||x|| = (x_1^2 + \ldots + x_d^2)^{1/2}$ the ℓ^2 -norm of the vector $x = (x_1, \ldots, x_d)^\top \in \mathbb{R}^d$. For a function f on \mathbb{R}^d , $\nabla_x f = (\frac{\partial f}{\partial x_1}, \ldots, \frac{\partial f}{\partial x_d})^\top$ denotes the gradient vector, and $\nabla_x^2 f = (\frac{\partial^2 f}{\partial x_i \partial x_j})_{i,j=1,\ldots,d}$ is the Hessian. $||A||_{op} = \max_{||x|| \leq 1} ||Ax||$ denotes the operator norm of a matrix A; $\lambda_{min}(A), \lambda_{max}(A)$ denote the minimum eigenvalue and maximum eigenvalue of matrix A, respectively. \mathbb{I}_d denotes a $d \times d$ identity matrix. $\mathbb{E}_f(g(X)) = \int g(x)f(x)dx$ with $X \sim f$. We write $a = \mathcal{O}(b)$ to denote $a \leq Cb$ for some constant C > 0, and $f(x) = o(g(x) \text{ means } |f(x)| \leq \epsilon |g(x)|$ for all $\epsilon > 0$. We use $\mathcal{N}(\mu, \Sigma)$ to denote a Gaussian random variable, or a Gaussian distribution, with mean μ and covariance Σ .

2 Variational Bayes

This section gives a brief overview of the VB method. Let y be the data and $p(y|\theta)$ the likelihood function, with θ the set of model parameters. Let $p(\theta)$ be the prior. Bayesian inference requires computing expectations with respect to the posterior distribution with density

$$p(\theta|y) = \frac{p(\theta)p(y|\theta)}{p(y)},$$

where $p(y) = \int p(\theta)p(y|\theta)d\theta$ is often called the marginal likelihood. It is often difficult to compute such expectations, partly because the density $p(\theta|y)$ itself is intractable as the normalizing constant p(y) is unknown. For simple models, Bayesian inference can be performed using Markov Chain Monte Carlo (MCMC), which estimates expectations with respect to $p(\theta|y)$ by sampling from it. For models where θ is high dimensional or has a complicated structure, MCMC methods in their current development are either not applicable or very time consuming. In the latter case, VB is an attractive alternative to MCMC. VB approximates the posterior $p(\theta|y)$ by a probability distribution with density $q_{\lambda}(\theta), \lambda \in \mathcal{M}$ - the variational parameter space, belonging to some tractable family of distributions such as Gaussian. The best λ is found by minimizing the Kullback-Leibler (KL) divergence of $p(\theta|y)$ from $q_{\lambda}(\theta)$

$$\lambda^* = \arg\min_{\lambda \in \mathcal{M}} \left\{ \operatorname{KL}(q_\lambda \| p(\cdot | y)) = \int q_\lambda(\theta) \log \frac{q_\lambda(\theta)}{p(\theta|y)} d\theta \right\}.$$
(2.1)

One can easily check that

$$\mathrm{KL}(q_{\lambda} \| p(\cdot | y)) = -\int q_{\lambda}(\theta) \log \frac{p(\theta)p(y|\theta)}{q_{\lambda}(\theta)} d\theta + \log p(y).$$

Thus minimizing KL is equivalent to maximizing the lower bound which is also called ELBO on $\log p(y)$

$$LB(\lambda) = \int q_{\lambda}(\theta) \log \frac{p(\theta)p(y|\theta)}{q_{\lambda}(\theta)} d\theta = \mathbb{E}_{q_{\lambda}} \left[\log \frac{p(\theta)p(y|\theta)}{q_{\lambda}(\theta)} \right] = \mathbb{E}_{q_{\lambda}} \left[h_{\lambda}(\theta) \right], \quad (2.2)$$

where $h_{\lambda}(\theta) := \log p(\theta) + \log p(y|\theta) - \log q_{\lambda}(\theta)$. Using the fact that $\mathbb{E}_{q_{\lambda}}[\nabla_{\lambda} \log q_{\lambda}(\theta)] = 0$, it can be seen that

$$\nabla_{\lambda} \text{LB}(\lambda) = \mathbb{E}_{q_{\lambda}} \left[\nabla_{\lambda} \log q_{\lambda}(\theta) \times h_{\lambda}(\theta) \right].$$
(2.3)

One then can obtain an unbiased estimate of $\nabla_{\lambda} \text{LB}(\lambda)$ by sampling from q_{λ} ,

$$\widehat{\nabla_{\lambda} \text{LB}}(\lambda) = \frac{1}{B} \sum_{s=1}^{B} \nabla_{\lambda} \log q_{\lambda}(\theta_s) \times h_{\lambda}(\theta_s), \quad \theta_s \sim q_{\lambda}(\theta), \quad s = 1, ..., B.$$
(2.4)

Alternative to (2.3), the Euclidean gradient $\nabla_{\lambda} \text{LB}(\lambda)$ can be computed using the so-called reparameterization-trick method [Kingma and Welling, 2013, Titsias and Lázaro-Gredilla, 2014]. The method for estimating the natural gradient proposed in this paper is applicable in both cases.

Stochastic gradient ascent (SGA) techniques are often employed to solve the maximization problem in (2.1). More specifically, one can iteratively update λ as follows

$$\lambda^{(k+1)} = \lambda^{(k)} + \tau_{k+1} \widehat{\nabla_{\lambda} \text{LB}}(\lambda^{(k)}), \qquad (2.5)$$

with the stepsize τ_k satisfying $\sum_{k=1}^{\infty} \tau_k = \infty$ and $\sum_{k=1}^{\infty} \tau_k^2 < \infty$. The convergence of the update in (2.5) has been studied in the literature [see, e.g. Robbins and Monro, 1951, Spall, 2005].

SGA approximates the exact gradient at each iteration by an estimate using a minibatch of the full sample (in big data settings) or by sampling from q_{λ} (as in (2.4)). This reduces computational cost, and facilitates on-the-fly (online) learning as new samples arrive. Note that, in practice, the data-dependent term $h_{\lambda}(\theta)$ is often estimated by using a mini-batch of the data. It is documented extensively in the literature [see, e.g. Bercu et al., 2020, Kirkby et al., 2022, Chau et al., 2024] that plain SGA as in (2.5) can lead to unsatisfactory estimates, as it is highly sensitive to the choice of hyper-parameters such as the step size or mini-batch size. In addition, SGA is known to have slow convergence when the Hessian of the cost function is ill-conditioned [Bottou et al., 2018], and even in the best case SGA converges no faster than sublinearly [Agarwal et al., 2009, Saad, 2009, Pelletier, 1998]. Significant effort in enhancing the plain SGA focuses on deriving adaptive learning step sizes; notable methods include Adam [Kingma and Ba, 2014], AdaGrad [Duchi et al., 2011] and Adadelta [Zeiler, 2012]. An alternative approach involves employing the natural gradient, which we will discuss in the following section.

3 Natural Gradient

Let $\mathcal{Q} = \{q_{\lambda}(\theta) : \lambda \in \mathcal{M} \subset \mathbb{R}^D\}$ be the set of VB approximating probability distributions parameterized by λ . We denote by d the dimension of the model parameter θ , and by D the dimension of the variational parameter λ . Gradient-based search for the optimal λ relies on the concept of gradient whose definition depends upon the metric on \mathcal{M} . It turns out that the regular Euclidean metric may not be appropriate for measuring the distance between two densities indexed by different variational parameters. For instance, by adapting examples given in Salimbeni et al. [2018], the pair of two Gaussians $\mathcal{N}(0, 0.1)$ and $\mathcal{N}(0, 1.1)$ look significantly different from each other, compared to the pair $\mathcal{N}(0, 1000)$ and $\mathcal{N}(0, 1001)$. Both pairs have the same Euclidean distance, while their KL divergences highlight a significant difference: the first pair exhibits a KL divergence of 1.6, whereas the second pair has a KL divergence of 2.5×10^{-7} .

Now consider two variational parameters λ , $\lambda + \delta \lambda$ and the KL divergence KL $(q_{\lambda}||q_{\lambda+\delta\lambda})$. From Tran et al. [2021], Tan [2021], it can be seen that

$$\operatorname{KL}(q_{\lambda}||q_{\lambda+\delta\lambda}) \approx \frac{1}{2} (\delta\lambda)^{\top} I_F(\lambda) \delta\lambda,$$

where,

$$I_F(\lambda) := -\mathbb{E}_{q_\lambda} \Big[\nabla_\lambda^2 \log q_\lambda(\theta) \Big] = \mathbb{E}_{q_\lambda} \Big[\nabla_\lambda \log q_\lambda(\theta) (\nabla_\lambda \log q_\lambda(\theta))^\top \Big], \tag{3.1}$$

is the Fisher information matrix of q_{λ} . This shows that the local KL divergence around the point $q_{\lambda} \in \mathcal{Q}$ is characterized by the Fisher matrix $I_F(\lambda)$. Therefore, a suitable metric between λ and $\lambda + \delta \lambda$ is the Fisher-Rao metric $(\delta \lambda)^{\top} I_F(\lambda) \delta \lambda$. As a result, assuming the objective function LB is smooth enough and for l > 0, if one considers the following optimization problem,

$$\arg \max_{\delta\lambda:(\delta\lambda)^{\top}I_{F}(\lambda)\delta\lambda=l} \left\{ \nabla_{\lambda} \mathrm{LB}(\lambda)^{\top}\delta\lambda \right\},\tag{3.2}$$

then through the method of Lagrangian multipliers, the steepest ascent is

$$\delta \lambda = \nabla_{\lambda}^{\text{nat}} \text{LB}(\lambda) := I_F^{-1}(\lambda) \nabla_{\lambda} \text{LB}(\lambda).$$
(3.3)

Amari [1998] termed this the natural gradient and popularized it in machine learning.

Using the natural gradient, the update in (2.5) becomes

$$\lambda^{(k+1)} = \lambda^{(k)} + \tau_{k+1} I_F^{-1}(\lambda^{(k)}) \nabla_\lambda \text{LB}(\lambda^{(k)}).$$
(3.4)

In the statistics literature, the steepest ascent in the form (3.4) has been used for a long time and is often known as Fisher's scoring in the context of maximum likelihood estimation [see, e.g. Longford, 1987]. The efficiency of the natural gradient over the Euclidean gradient has been well documented [Sato, 2001, Hoffman et al., 2013, Tran et al., 2017, Martens, 2020, Tan, 2021]. The natural gradient is invariant under parameterization [Martens, 2020], meaning it remains unchanged across different coordinate systems and is an intrinsic geometric object. This property makes it particularly suitable for use when the variational parameter space \mathcal{M} is a Riemannian manifold [Tran et al., 2021] as it is coordinate-free and leverages the underlying geometry of the space.

In the special case of Gaussian approximations with a full covariance matrix or a Cholesky-factor covariance matrix, it is possible to obtain the inverse Fisher matrix in closed form [Tan, 2021, Magris et al., 2022]. Beyond these limited cases, however, it is challenging to accurately compute the natural gradient. The natural gradient method

requires the analytic computation of the Fisher information matrix and its inversion. Even if an analytical expression of the Fisher matrix is obtained, computing its inverse has a complexity of $O(D^{\kappa})$, with $2 < \kappa \leq 3$ depending on various algorithms. It is therefore either analytically infeasible or prohibitively computationally expensive to use natural gradient in many modern statistical applications; current practice resorts to heuristic workarounds that can affect the results of Bayesian inference [Martens, 2020, Lopatnikova and Tran, 2023].

4 Inversion Free Natural Gradient Variational Bayes

This section first presents the approach for approximating the inverse of Fisher matrix. We then present the inversion free natural gradient Variational Bayes method, referred to as IFVB, and its weighted averaged version AIFVB. The IFVB and AIFVB methods are stochastic natural gradient descent algorithms that avoid computing the Fisher matrix and its inversion altogether. As explained later, these methods also enable us to deal with situations where the estimate of Fisher matrix has eigenvalues with significantly different orders of magnitude (i.e., poor conditioning). In this section and Section 5, we will denote $\mathcal{L}(\lambda) = -\text{LB}(\lambda)$, which can be viewed as the loss function, and the problem of maximizing the lower bound becomes the minimization of \mathcal{L} .

4.1 Recursive Estimation of $I_F^{-1}(\lambda)$

For each $s = 1, 2 \dots$, let

$$H_s = H_0 + \sum_{j=1}^s \nabla_\lambda \log q_\lambda(\theta_j) (\nabla_\lambda \log q_\lambda(\theta_j))^\top, \quad \theta_j \sim q_\lambda, \tag{4.1}$$

where H_0 is some positive definite matrix, e.g, $H_0 = \epsilon \mathbb{I}_D$ with $\epsilon > 0$ and \mathbb{I}_D the identity matrix of size D. Theorem 4.1 below says that $\mathbf{H}_s := H_s/s$ is a consistent estimate of I_F defined in (3.1) as $s \to \infty$, and provides a recursive procedure for obtaining H_s^{-1} . This recursive procedure computes H_{s+1}^{-1} from H_s^{-1} without resorting to the usual (expensive and error prone) matrix inversion. Additionally, the symmetry and positivity of H_s , and hence of H_s^{-1} , is preserved, which is an important property.

Theorem 4.1. Let $\phi_s = \nabla_\lambda \log q_\lambda(\theta_s)$. We have that

$$H_{s+1}^{-1} = H_s^{-1} - \left(1 + \phi_{s+1}^{\top} H_s^{-1} \phi_{s+1}\right)^{-1} H_s^{-1} \phi_{s+1} \phi_{s+1}^{\top} H_s^{-1}, \quad s = 0, 1, \dots$$
(4.2)

In particular, the positivity and symmetry of H_s is preserved for all s. Furthermore,

$$\mathbf{H}_{s} = \frac{1}{s} H_{s} \xrightarrow[s \to +\infty]{a.s} I_{F}(\lambda) \quad and \quad \mathbf{H}_{s}^{-1} \xrightarrow[s \to +\infty]{a.s} I_{F}^{-1}(\lambda)$$

The proof of Theorem 4.1 can be found in Appendix A.1. The expression (4.2) suggests that H_s^{-1} is a sum of outer products - a property that can be exploited to avoid storage of large matrices; see Remark 4.2.

4.2 Inversion Free Natural Gradient Variational Bayes

Theorem 4.1 suggests that one can approximate the inverse Fisher matrix I_F^{-1} by $\mathbf{H}_s^{-1} = sH_s^{-1}$ for some large s, where H_s^{-1} is calculated recursively as in (4.2). This method does not require an analytic calculation of I_F and its inversion; also, the estimate \mathbf{H}_s^{-1} is guaranteed to be symmetric and positive definite. However, a direct application of Theorem 4.1 for computing the natural gradient can be inefficient for two reasons.

First, in order to ensure the consistency of estimates $\lambda^{(k)}$ from (3.4), where the inverse Fisher matrix is replaced by its estimate \mathbf{H}_s^{-1} , one must control the eigenvalues of the estimate \mathbf{H}_s^{-1} . See Bercu et al. [2020] and Boyer and Godichon-Baggioni [2023] for related discussion in the context of stochastic Newton's method. With this aim, we follow Boyer and Godichon-Baggioni [2023] and modify (4.1) as follows

$$A_s(\lambda) = H_0 + \sum_{j=1}^s \nabla_\lambda \log q_\lambda(\theta_j) (\nabla_\lambda \log q_\lambda(\theta_j))^\top + c_\beta \sum_{j=1}^s j^{-\beta} Z_j Z_j^\top, \quad s = 1, 2, \dots$$
(4.3)

where $\theta_j \sim q_{\lambda}(\cdot), Z_1, \ldots, Z_s \sim \mathcal{N}(0, \mathbb{I}_D)$ are independent standard Gaussian vectors of dimension $D, c_{\beta} \geq 0$ and $\beta \in (0, \alpha - 1/2)$ for some $\alpha \in (1/2, 1)$. Theorem A.1 in the Appendix shows that A_s/s converges almost surely to $I_F(\lambda)$, and that A_{s+1}^{-1} can be computed recursively as follows

$$\begin{cases} A_{s+\frac{1}{2}}^{-1} = A_{s}^{-1} - \left(1 + \phi_{s+1}^{\top} A_{s}^{-1} \phi_{s+1}\right)^{-1} A_{s}^{-1} \phi_{s+1} \phi_{s+1}^{\top} A_{s}^{-1} \\ A_{s+1}^{-1} = A_{s+\frac{1}{2}}^{-1} - c_{\beta} (s+1)^{-\beta} \left(1 + c_{\beta} (s+1)^{-\beta} Z_{s+1}^{\top} A_{s+\frac{1}{2}}^{-1} Z_{s+1}\right)^{-1} A_{s+\frac{1}{2}}^{-1} Z_{s+1} Z_{s+1}^{\top} A_{s+\frac{1}{2}}^{-1} \\ (4.4)$$

with $\phi_s = \nabla_\lambda \log q_\lambda(\theta_s)$. As being shown later in the proof of Theorem 5.1, taking $c_\beta > 0$ ensures the smallest eigenvalue of the Fisher matrix estimate not going to zero faster than $O(s^{-\beta})$ almost surely, hence enabling strongly consistent estimates.

Second, a direct use of (4.3) would require a separate recursive procedure (4.4) to calculate the inverse Fisher estimate $sA_s^{-1}(\lambda^{(k)})$ in each update $\lambda^{(k)}$, k = 1, 2, ..., in (3.4). This might make the VB training procedure in (3.4) computationally expensive. Instead, we propose a "streamlined" version that updates the Fisher matrix estimate along with the iterates $\lambda^{(k)}$. To this end, we define

$$\tilde{H}_{s} = H_{0} + \sum_{k=0}^{s-1} \nabla_{\lambda} \log q_{\lambda^{(k)}}(\theta_{k+1}) \left(\nabla_{\lambda} \log q_{\lambda^{(k)}}(\theta_{k+1}) \right)^{\top} + c_{\beta} \sum_{k=1}^{s} k^{-\beta} Z_{k} Z_{k}^{\top}, \quad s = 0, 1, ...,$$
(4.5)

where $\theta_{k+1} \sim q_{\lambda^{(k)}}(\cdot)$. Note that, unlike (4.3), \tilde{H}_s in (4.5) depends on the iterates $\lambda^{(k)}$, k < s. Similar to (4.4), \tilde{H}_{s+1}^{-1} can be computed recursively as follows

$$\begin{cases} \tilde{H}_{s+\frac{1}{2}}^{-1} = \tilde{H}_{s}^{-1} - \left(1 + \phi_{s+1}^{\top} \tilde{H}_{s}^{-1} \phi_{s+1}\right)^{-1} \tilde{H}_{s}^{-1} \phi_{s+1} \phi_{s+1}^{\top} \tilde{H}_{s}^{-1} \\ \tilde{H}_{s+\frac{1}{2}}^{-1} = \tilde{H}_{s+\frac{1}{2}}^{-1} - c_{\beta} (s+1)^{-\beta} \left(1 + c_{\beta} (s+1)^{-\beta} Z_{s+1}^{\top} \tilde{H}_{s+\frac{1}{2}}^{-1} Z_{s+1}\right)^{-1} \tilde{H}_{s+\frac{1}{2}}^{-1} Z_{s+1} Z_{s+1}^{\top} \tilde{H}_{s+\frac{1}{2}}^{-1},$$

$$(4.6)$$

where $\phi_{s+1} = \nabla_{\lambda} \log q_{\lambda^{(s)}}(\theta_{s+1}).$

The Euclidean gradient of the loss function can be estimated as

$$\widehat{\nabla_{\lambda}\mathcal{L}}(\lambda) = -\frac{1}{B}\sum_{i=1}^{B} \nabla_{\lambda}\log q_{\lambda}\left(\theta_{i}\right) \times h_{\lambda}\left(\theta_{i}\right), \qquad (4.7)$$

where the θ_i 's are *B* i.i.d samples from q_{λ} .

Putting these together, we propose the Inverse Free Natural Gradient VB algorithm, referred to as IFVB and outlined in Algorithm 1. The performance of this algorithm is found not sensitive to ϵ and c_{β} ; both are set to 1 in our examples below unless stated otherwise. We found that the algorithm works well for β in the range (0.1,0.5).

Algorithm 1: Inversion-Free Natural Gradient Variational Baye	s (I)	FVF	3)
---	-------	-----	----

 $\begin{array}{ll} \textbf{Require:} \text{ Choose an initial value } \lambda^{(0)}, \ \epsilon > 0 \ \text{and } c_{\beta} \ge 0. \\ 1: \ H_0 = \epsilon \mathbb{I}_D \\ 2: \ \textbf{for } s = 0, 1, \dots, \ \textbf{do} \\ 3: \quad \text{Compute } \widehat{\nabla_{\lambda} \mathcal{L}}(\lambda^{(s)}). \\ 4: \quad \text{Sample } \theta_{s+1} \sim q_{\lambda^{(s)}}(\theta), \ Z_{s+1} \sim \mathcal{N}(0, \mathbb{I}_D) \ \text{and let } \phi_{s+1} = \nabla_{\lambda} \log q_{\lambda^{(s)}}(\theta_{s+1}). \\ \quad \text{Calculate } \widehat{H}_{s+1}^{-1} \ \text{as in } (4.6), \ \text{and } \widetilde{\mathbf{H}}_{s+1}^{-1} = (s+1) \widetilde{H}_{s+1}^{-1}. \\ 5: \quad \text{Update } \lambda^{(s+1)} = \lambda^{(s)} - \tau_{s+1} \widetilde{\mathbf{H}}_{s+1}^{-1} \widehat{\nabla_{\lambda} \mathcal{L}}(\lambda^{(s)}). \\ 6: \ \textbf{end for} \end{array}$

Remark 4.1. Some remarks are in order. Line #4 in Algorithm 1 updates \tilde{H}_{s+1}^{-1} from \tilde{H}_{s}^{-1} using only one sample from $q_{\lambda(s)}$. Depending on applications, however, it might be beneficial to use S samples (S > 1) to update \tilde{H}_{s+1}^{-1} . One then needs to run (4.6) for S times, and compute $\tilde{\mathbf{H}}_{s+1}^{-1} = S(s+1)\tilde{H}_{s+1}^{-1}$. The factor S is not practically important as gradient clipping, that keeps the norm of the natural gradient $\tilde{\mathbf{H}}_{s+1}^{-1} \widehat{\nabla}_{\lambda} \mathcal{L}(\lambda^{(s)})$ below some certain value, is often used in the SGD literature [see, e.g., Goodfellow et al., 2016].

Remark 4.2. For applications such as deep learning where the size D of variational parameter λ is large, it is important to note that implementation of Algorithm 1 does not require storage of the matrices \tilde{H}_{s+1}^{-1} . Let us take $c_{\beta} = 0$ to simplify the exposition; the extension to the case $c_{\beta} > 0$ is straightforward. From (4.6), \tilde{H}_{s+1}^{-1} takes the form

$$\tilde{H}_{s+1}^{-1} = \frac{1}{\epsilon} \mathbb{I}_D - \sum_{k=1}^{s+1} \psi_k \psi_k^{\top}, \quad \psi_k = \left(1 + \phi_{k+1}^{\top} \tilde{H}_k^{-1} \phi_{k+1}\right)^{-1/2} \tilde{H}_k^{-1} \phi_{k+1}, \quad (4.8)$$

which is presented by outer products. As the result, all the required matrix-vector multiplications can be performed efficiently. That is,

$$\widetilde{H}_{s+1}^{-1}\widehat{\nabla_{\lambda}\mathcal{L}}\left(\lambda^{(s)}\right) = \frac{1}{\epsilon}\widehat{\nabla_{\lambda}\mathcal{L}}\left(\lambda^{(s)}\right)^{\top}\widehat{\nabla_{\lambda}\mathcal{L}}\left(\lambda^{(s)}\right) - \sum_{k=1}^{s+1}\left(\psi_{k}^{\top}\widehat{\nabla_{\lambda}\mathcal{L}}\left(\lambda^{(s)}\right)\right)\psi_{k}$$
$$\widetilde{H}_{s}^{-1}\phi_{s+1} = \frac{1}{\epsilon}\phi_{s+1}^{\top}\phi_{s+1} - \sum_{k=1}^{s}\left(\psi_{k}^{\top}\phi_{s+1}\right)\psi_{k}$$

which does not necessitate storage of the matrices \tilde{H}_s^{-1} . It is also natural to only keep the last K outer products, for some $K \ge 1$ (K = 100 in our examples below), in (4.8) to further reduce the computation

$$\tilde{H}_{s+1}^{-1} \approx \frac{1}{\epsilon} \mathbb{I}_D - \sum_{k=s-K+2}^{s+1} \psi_k \psi_k^\top.$$
(4.9)

The computational complexity of Algorithm 1 is therefore $O(s \times D)$, with s the number of iterations. This complexity is the same as that of the popular adaptive learning methods such as Adam and AdaGrad.

4.3 Weighted Averaged IFVB Algorithm

It is a common practice in the SGD literature to use an average of the iterates $\lambda^{(k)}$ to form the final estimate of the optimal λ^* [Goodfellow et al., 2016, Boyer and Godichon-Baggioni, 2023]. The weighted averaged estimate is of the form

$$\overline{\lambda}^{(s+1)} = \frac{1}{\sum_{k=1}^{s+1} w_k} \sum_{k=1}^{s+1} w_k \lambda^{(k)} = \overline{\lambda}^{(s)} + \frac{w_{s+1}}{\sum_{k=1}^{s+1} w_k} \left(\lambda^{(s+1)} - \overline{\lambda}^{(s)} \right),$$

where the weights $w_k > 0$. Inspired by Boyer and Godichon-Baggioni [2023], we select $w_k = (\log(k))^w$ with $w \ge 0$. This averaging scheme puts more weight on recent estimates $\lambda^{(k)}$ if w > 0. If one chooses w = 0, this leads to the uniform averaging technique, i.e.,

$$\overline{\lambda}^{(s)} = \frac{1}{s} \sum_{k=1}^{s} \lambda^{(k)}.$$

We use w = 2 in all the numerical examples in Section 6. We arrive at the Weighted Averaged IFVB Algorithm (AIFVB) defined recursively for all $s \ge 0$ by

$$\lambda^{(s+1)} = \lambda^{(s)} - \tau_{s+1} \widetilde{\mathbf{H}}_{s+1}^{-1} \widehat{\nabla_{\lambda} \mathcal{L}} \left(\lambda^{(s)} \right)$$
$$\overline{\lambda}^{(s+1)} = \overline{\lambda}^{(s)} + \frac{\left(\log(s+1) \right)^w}{\sum_{k=0}^s \left(\log(k+1) \right)^w} \left(\lambda^{(s+1)} - \overline{\lambda}^{(s)} \right).$$

For the AIFVB algorithm, we consider the following estimate of the Fisher matrix

$$\tilde{\mathbf{H}}_{s} = \frac{1}{s} \left(H_{0} + \sum_{k=0}^{s-1} \nabla_{\lambda} \log q_{\overline{\lambda}^{(k)}}(\overline{\theta}_{k+1}) (\nabla_{\lambda} \log q_{\overline{\lambda}^{(k)}}(\overline{\theta}_{k+1}))^{\top} + c_{\beta} \sum_{k=1}^{s} k^{-\beta} Z_{k} Z_{k}^{T} \right), \quad (4.10)$$

for s = 1, 2, ..., where $\overline{\theta}_{k+1} \sim q_{\overline{\lambda}^{(k)}}(\theta)$, the Z_k 's are defined as before. Similar to (4.6), $\widetilde{\mathbf{H}}_{s+1}^{-1}$ can be updated recursively.

Remark 4.3. In the estimate (4.10), we compute $\tilde{\mathbf{H}}_s$ using the averaged iterates $\overline{\lambda}^{(k)}$, as this can lead to a faster convergence. Nonetheless, one can also use $\lambda^{(k)}$ instead of $\overline{\lambda}^{(k)}$ in (4.10).

Putting these together, we have the Weighted Averaged IFVB Algorithm, outlined in Algorithm 2.

Algorithm 2: Weighted Averaged IFVB Algorithm (AIFVB)

 $\begin{aligned} & \text{Require: Choose an initial value } \lambda^{(0)} = \overline{\lambda}^{(0)}, \ \epsilon > 0 \text{ and } c_{\beta} \ge 0. \\ & 1: \ H_0 = \epsilon \mathbb{I}_D \\ & 2: \ \text{for } s = 0, 1, \dots, \ \text{do} \\ & 3: \quad \text{Compute } \widehat{\nabla_{\lambda} \mathcal{L}}(\lambda^{(s)}). \\ & 4: \quad \text{Sample } \overline{\theta}_{s+1} \sim q_{\overline{\lambda}^{(s)}}(\theta), \ Z_{s+1} \sim \mathcal{N}(0, \mathbb{I}_D) \text{ and let } \phi_{s+1} = \nabla_{\lambda} \log q_{\overline{\lambda}^{(s)}}(\overline{\theta}_{s+1}). \\ & \text{Calculate } \widetilde{H}_{s+1}^{-1} \text{ as in } (4.6), \text{ and } \widetilde{\mathbf{H}}_{s+1}^{-1} = (s+1)\widetilde{H}_{s+1}^{-1}. \\ & 5: \quad \text{Update } \lambda^{(s+1)} = \lambda^{(s)} - \tau_{s+1}\widetilde{\mathbf{H}}_{s+1}^{-1}\widehat{\nabla_{\lambda} \mathcal{L}}(\lambda^{(s)}). \\ & 6: \quad \text{Update } \overline{\lambda}^{(s+1)} = \overline{\lambda}^{(s)} + \frac{\left(\log(s+1)\right)^w}{\sum_{k=0}^s \left(\log(k+1)\right)^w} \left(\lambda^{(s+1)} - \overline{\lambda}^{(s)}\right). \\ & 7: \ \text{end for} \end{aligned}$

5 Convergence analysis

For the convergence analysis in this section, we consider a step size of the form $\tau_k = \frac{c_{\alpha}}{(c'_{\alpha}+k)^{\alpha}}$ with $c_{\alpha} > 0$, $c'_{\alpha} \ge 0$ and $\alpha \in (1/2, 1)$. Write

$$\nabla_{\lambda} \mathcal{L}(\lambda) = \mathbb{E}_{q_{\lambda}}[-\nabla_{\lambda} \log q_{\lambda}(\theta) \times h_{\lambda}(\theta)] =: \mathbb{E}_{q_{\lambda}}[\ell(\theta, \lambda)]$$

It can be seen that¹

$$abla_{\lambda}^{2}\mathcal{L}(\lambda) = I_{F}(\lambda) + \int
abla_{\lambda}^{2}q_{\lambda}(\theta) \big(\log q_{\lambda}(\theta) - \log p(\theta|y)\big)d\theta.$$

At the optimal $\lambda = \lambda^*$, $q_{\lambda^*}(\theta) \approx p(\theta|y)$, the second term above expects to be close to zero. We can therefore expect that, in a neighbourhood of λ^* , the Fisher $I_F(\lambda)$ behaves like the Hessian $\nabla^2_{\lambda} \mathcal{L}(\lambda)$. This motivates us to adapt the results from stochastic Newton algorithms [see, e.g., Bercu et al., 2020, Boyer and Godichon-Baggioni, 2023] for convergence analysis of IFVB and AIFVB proposed in this paper.

We now provide the convergence analysis for the AIFVB algorithm; a minor modification provides convergence results for IFVB.

Theorem 5.1. Assume that \mathcal{L} is twice differentiable and that there is L_0 such that for all λ , $\|\nabla_{\lambda}^2 \mathcal{L}(\lambda)\|_{op} \leq L_0$. Suppose that $\nabla_{\lambda} \mathcal{L}(\lambda^*) = 0$ and $c_{\beta} > 0$. Assume also that there are non negative constants C_0, C_1 such that for all λ

$$\mathbb{E}\left[\left\|\ell(\theta,\lambda)\right\|^{2}\right] \leq C_{0} + C_{1}(\mathcal{L}(\lambda) - \mathcal{L}(\lambda^{*}))$$

and that there are positive constants C'_0, C'_1 such that for all λ ,

$$\mathbb{E}\left[\left\|\nabla_{\lambda}\log q_{\lambda}(\theta)\right\|^{4}\right] \leq C_{0}' + C_{1}'\left(\mathcal{L}(\lambda) - \mathcal{L}(\lambda^{*})\right)^{2}.$$
(5.1)

Suppose that $C'_1 = 0$ or that \mathcal{L} is convex. Then the estimates $\lambda^{(k)}$, $\overline{\lambda}^{(k)}$ k = 0, 1, ..., from Algorithm 2 satisfy:

¹See Appendix A.3 for detailed calculations; see also Tang and Ranganath [2019], Tan [2021].

(i) $\mathcal{L}(\lambda^{(k)}) - \mathcal{L}(\lambda^*)$ converges almost surely to a finite random variable.

(*ii*)
$$\min_{k=0}^{s} \left\| \nabla \mathcal{L} \left(\lambda^{(k)} \right) \right\|^2 = o \left(s^{-(1-\alpha)} \right) a.s.$$

(*iii*)
$$\min_{k=0}^{s} \left\| \nabla \mathcal{L} \left(\overline{\lambda}^{(k)} \right) \right\|^{2} = o\left(s^{-(1-\alpha)} \right) a.s.$$

If the convexity of \mathcal{L} is satisfied, conclusions (ii) and (iii) in Theorem 5.1 imply the almost sure convergence of $\lambda^{(k)}$ and $\overline{\lambda}^{(k)}$ to λ^* . The assumption in (5.1) might appear to look irrelevant, as the right-hand side term is model-dependent, i.e. depending on the prior and likelihood, while the left-hand side is not. One can simply replace (5.1) with an assumption on the uniform bound of the fourth order moment of $\nabla_{\lambda} \log q_{\lambda}$, which obviously implies (5.1). We use (5.1) to keep the result as general as possible.

We now give the consistency of the estimates of the Fisher information given in (4.10).

Corollary 5.1. Suppose that $\lambda^{(k)}$ converges almost surely to λ^* , and that the map $\lambda \mapsto I_F(\lambda)$ is continuous at λ^* . Suppose also that (5.1) is satisfied. Then

$$\tilde{\mathbf{H}}_{s} \xrightarrow[s \to +\infty]{a.s} I_{F}\left(\lambda^{*}\right).$$

Note that we can obtain the convergence rate of $\tilde{\mathbf{H}}_s$ to $I_F(\lambda^*)$, or even the rate of convergence of $\tilde{\mathbf{H}}_s$ to the Hessian $\nabla^2_{\lambda} \mathcal{L}(\lambda^*)$. Please see Appendix A.4 for more details.

Without the convexity assumption, the proof of Theorem 5.1 implies that the estimates $\lambda^{(k)}$ converge to a stationary point λ^* of the objective \mathcal{L} , i.e. $\nabla_{\lambda} \mathcal{L}(\lambda^*) = 0$. The result below gives a convergence rate of the estimates $\lambda^{(k)}$ to such λ^* .

Theorem 5.2. Suppose that $\lambda^{(k)}$ converges almost surely to λ^* . Assume that the functional \mathcal{L} is differentiable with $\nabla_{\lambda}\mathcal{L}(\lambda^*) = 0$ and twice continuously differentiable in a neighborhood of λ^* . Suppose also that there are $\eta > \frac{1}{\alpha} - 1$ and positive constants $C_{\eta,0}, C_{\eta,1}$ such that for all λ ,

$$\mathbb{E}\left[\left\|\ell(\theta,\lambda)\right\|^{2+2\eta}\right] \leq C_{\eta,0} + C_{\eta,1} \left(\mathcal{L}(\lambda) - \mathcal{L}(\lambda^*)\right)^{1+\eta}$$

and that inequality (5.1) holds. Suppose also that $\nabla^2_{\lambda} \mathcal{L}(\lambda^*)$ and $I_F(\lambda^*)$ are positive. Then

$$\left\|\lambda^{(s)} - \lambda^*\right\|^2 = \mathcal{O}\left(\frac{\log s}{s^{\alpha}}\right) \quad a.s.$$
(5.2)

Observe that (5.2) is the usual rate of convergence for (adaptive) stochastic gradient type algorithms [Bercu et al., 2020, Nguyen et al., 2021]. The following theorem shows that the weighted averaged estimates achieve a better convergence rate and are asymptotically efficient.

Theorem 5.3. Suppose that the assumptions in Theorem 5.2 hold. Assume that the functional

$$\lambda \longmapsto \Sigma(\lambda) := \mathbb{E}_{q_{\lambda}} \left[\ell(\theta, \lambda) \ell(\theta, \lambda)^{\top} \right]$$

is continuous at λ^* , and there are a neighborhood V^* of λ^* and a constant L_{δ} such that for all $\lambda \in V^*$,

$$\left\|\nabla_{\lambda}\mathcal{L}(\lambda) - \nabla^{2}\mathcal{L}(\lambda^{*})(\lambda - \lambda^{*})\right\| \leq L_{\delta} \left\|\lambda - \lambda^{*}\right\|^{2}.$$
(5.3)

Then

$$\left\|\overline{\lambda}^{(s)} - \lambda^*\right\|^2 = \mathcal{O}\left(\frac{\log s}{s}\right) \quad a.s.$$

In addition,

$$\sqrt{Bs}\left(\overline{\lambda}^{(s)}-\lambda^*\right)\xrightarrow[n\to+\infty]{law}\mathcal{N}\left(0,\nabla_{\lambda}^2\mathcal{L}\left(\lambda^*\right)^{-1}\Sigma\left(\lambda^*\right)\nabla_{\lambda}^2\mathcal{L}\left(\lambda^*\right)^{-1}\right).$$

Remark 5.1. Comparing Theorem 5.2 and Theorem 5.3, it is intriguing to observe that the averaging technique clearly contributes to an improvement in the rate of convergence. Moreover, observe that Bs represent the total number of samples generated for the AIFVB algorithm (without taking into account the ones for estimating the inverse of the Fisher information). In addition, following Godichon-Baggioni and Werge [2023], i.e taking B = D and generating only one sample to estimate the inverse of the Fisher information, it leads to an algorithm with O(sBD) operations, which is the same computational complexity as stochastic gradient type algorithms. To be more precise, taking B = D, and denoting by N = sB the total number of simulated data used for estimating the gradients at each step of the algorithm, one has

$$\sqrt{N}\left(\overline{\lambda}^{N/D} - \lambda^*\right) \xrightarrow[N \to +\infty]{law} \mathcal{N}\left(0, \nabla_{\lambda}^2 \mathcal{L}\left(\lambda^*\right)^{-1} \Sigma\left(\lambda^*\right) \nabla_{\lambda}^2 \mathcal{L}\left(\lambda^*\right)^{-1}\right),$$

with O(ND) operations, which is exactly the same rate and complexity as averaged stochastic gradient algorithms.

Remark 5.2. To the best of our knowledge, this paper is the first to give a central limit theorem for the estimate of variational parameters in VB. This result can have some interesting implications. Given a variational family Q and data y, one can define the "final" variational approximation of the posterior distribution $p(\theta|y)$ as

$$q(\theta|y, Q) = \int q_{\lambda}(\theta) p(\lambda|y) d\lambda$$
(5.4)

with $p(\lambda|y)$ approximated by $\mathcal{N}\left(\lambda^*, \frac{1}{Bs} \nabla_{\lambda}^2 \mathcal{L}\left(\lambda^*\right)^{-1} \Sigma\left(\lambda^*\right) \nabla_{\lambda}^2 \mathcal{L}\left(\lambda^*\right)^{-1}\right)$, who in turn can be further approximated by $\mathcal{N}\left(\lambda^*, \frac{1}{Bs}I_F\left(\lambda^*\right)^{-1}\Sigma\left(\lambda^*\right)I_F\left(\lambda^*\right)^{-1}\right)$. The posterior approximation in (5.4) not only takes into account the uncertainty in the SGD training procedure of λ , but also enlarges the variance in $q_{\lambda^*}(\theta)$. We recall that underestimating the posterior variance is a well perceived problem in the VB literature [Blei et al., 2017].

6 Numerical Examples

This section provides a range of examples to demonstrate the applicability of the inversion free variational Bayes methods. In Sections 4 and 5, to be consistent with the optimization literature, we presented the VB problem in terms of optimizing the negative lower bound $\mathcal{L}(\lambda)$. In this section, to be consistent with the VB literature, we present the numerical examples in terms of maximizing the lower bound $\text{LB}(\lambda)$. The five examples encompass a diverse array of domains ranging from Gaussian approximation to normalizing flow Variational Bayes. We provide four examples in this section; another example is provided in the Supplementary Material. The implementation code is available at https://github.com/VBayesLab/Inversion-free-VB.

Example 1 (Beta variational approximation). We consider an example in which the posterior distribution and the natural gradient are given in closed-form. This helps facilitate the comparison among the considered approximation approaches. Following Tran et al. [2017], we generated n = 200 observations $y_1, y_2, \ldots, y_{200} \sim \text{Bernoulli}(1, \theta = 0.3)$, and let $\kappa = \sum_{i=1}^{n} y_i = 57$. The prior distribution of θ is chosen to be the uniform distribution on (0, 1). The posterior distribution is $\text{Beta}(\kappa + 1, n - \kappa + 1)$. Let $\lambda^* = (\kappa + 1, n - \kappa + 1)^{\top}$. The variational distribution $q_{\lambda}(\theta)$ is chosen to be $\text{Beta}(\alpha, \beta)$, which belongs to the exponential family with the natural parameter $\lambda = (\alpha, \beta)^{\top}$. We have

$$\log q_{\lambda}(\theta) = \log \Gamma(\alpha + \beta) - \log \Gamma(\alpha) - \log \Gamma(\beta) + (\alpha - 1) \log \theta + (\beta - 1) \log(1 - \theta).$$

Hence

$$\nabla_{\lambda} \log q_{\lambda}(\theta) = (\psi(\alpha + \beta) - \psi(\alpha) + \log(\theta), \psi(\alpha + \beta) - \psi(\beta) + \log(1 - \theta))^{\top}$$

where ψ is the digamma function. In this case, the Fisher information matrix is available in a closed-form

$$I_F(\lambda) = \begin{pmatrix} \varphi_1(\alpha) - \varphi_1(\alpha + \beta) & -\varphi_1(\alpha + \beta) \\ -\varphi_1(\alpha + \beta) & \varphi_1(\beta) - \varphi_1(\alpha + \beta) \end{pmatrix},$$

where $\varphi_1(x)$ is the trigamma function. In addition, we have that

$$\nabla_{\lambda} \text{LB}(\lambda) = \begin{pmatrix} (\kappa + 1 - \alpha) (\varphi_1(\alpha) - \varphi_1(\alpha + \beta)) - (n - \kappa + 1 - \beta) \varphi_1(\alpha + \beta) \\ (n - \kappa + 1 - \beta) (\varphi_1(\beta) - \varphi_1(\alpha + \beta)) - (\kappa + 1 - \alpha) \varphi_1(\alpha + \beta) \end{pmatrix}.$$

The Euclidean gradient ascent update is given by

$$\lambda^{(k+1)} = \lambda^{(k)} + \tau_k \nabla_\lambda \text{LB}(\lambda^{(k)}),$$

and the natural gradient ascent update is given by

$$\lambda^{(k+1)} = \lambda^{(k)} + \tau_k I_F^{-1}(\lambda^{(k)}) \nabla_\lambda \text{LB}(\lambda^{(k)}).$$

Figure 1 compares the true posterior density with those obtained by the Euclidean gradient ascent, the (exact) natural gradient VB algorithm (NGVB) and IFVB, using two initializations for λ : $\lambda^{(0)} = (5; 45)^{\top}$ (Figure 1, left) and $\lambda^{(0)} = (25; 25)^{\top}$ (Figure 1, right). We set $c_{\beta} = 0$ in this example. The step size for IFVB is set as $\tau_k = 10/(1+k)^{\alpha}$ with $\alpha = 0.6$ (see Section 5), while the step size for other methods is set to 1/(1+k). All the algorithms used the same stopping rule, which terminates the training if the difference in l_2 -norm of two successive updates is less than a tolerance of 10^{-5} . The figure shows that



Figure 1: Comparing the true posterior density versus those obtained by Euclidean gradient ascent, NGVB and IFVB.

the posterior densities obtained by NGVB and IFVB are superior to using the traditional Euclidean gradient. Furthermore, these algorithms are insensitive to the initial $\lambda^{(0)}$.

As we know the exact posterior distribution in this case, we now further compare the performance of NGVB, IFVB and AIFVB using the initialization $\lambda^{(0)} = (5; 45)^{\top}$. Figure 2 plots the error $\|\lambda^{(s)} - \lambda^*\|$ ($\|\overline{\lambda}^{(s)} - \lambda^*\|$ for AIFVB) as the function of the number of iterations. It is clear from Figure 2 that both the IFVB and AIFVB algorithms require more steps to achieve a (relatively) equivalent error level compared to the exact natural gradient ascent algorithm. This observation aligns with the fact that both algorithms rely on an approximation of the inverse Fisher information matrix. As time advances, the quality of this approximation improves. Furthermore, it is evident that the incorporation of an averaging technique significantly aids in minimizing the error associated with the approximation.

Example 2 (Gaussian approximation). We consider a Gaussian approximation example which was considered in Tan [2021]. Specifically, it is assumed that $q_{\lambda}(\theta) = \mathcal{N}(\theta|\mu, \Sigma)$ which belongs to the exponential family and can be written as

$$q_{\lambda}(\theta) = \exp\left(s(\theta)^{\top}\lambda - a(\lambda)\right),$$

where $s(\theta) = (\theta^{\top}, \operatorname{vech}(\theta\theta^{\top})^{\top})^{\top}$ is the sufficient statistics, $\lambda = (\mu^{\top}\Sigma^{-1}, -\frac{1}{2}\operatorname{vec}(\Sigma^{-1})^{\top}\Gamma)^{\top}$ is the natural parameter, with Γ the duplication matrix. It can be seen that $a(\lambda) = \frac{1}{2}\mu^{\top}\Sigma^{-1}\mu + \frac{1}{2}\log|\Sigma| + \frac{d}{2}\log(2\pi)$ is the log-partition function. This implies the natural



Figure 2: Error versus iterations

gradient

$$\nabla_{\lambda}^{nat} LB = I_F^{-1}(\lambda) \nabla_{\lambda} LB(\lambda) = I_F^{-1}(\lambda) I_F(\lambda) \nabla_m LB(m) = \nabla_m LB = \begin{pmatrix} \nabla_{\mu} LB - 2(\nabla_{\Sigma} LB)\mu \\ \Gamma^{\top} vec(\nabla_{\Sigma} LB) \end{pmatrix},$$

where $\operatorname{vec}(\nabla_{\Sigma} LB) = \nabla_{\operatorname{vec}(\Sigma)} LB$. From here Tan [2021] derives the updates for λ as follows,

- 1. $\Sigma^{-1} \leftarrow \Sigma^{-1} 2\tau_k \nabla_{\Sigma} \text{LB}$
- 2. $\mu \leftarrow \mu + \tau_k \Sigma \nabla_\mu \text{LB}.$

Obviously, the traditional (Euclidean) gradient ascent is given by

- 1. $\mu \leftarrow \mu + \tau_k \nabla_\mu \text{LB}$
- 2. $\Sigma \leftarrow \Sigma + \tau_k \nabla_{\Sigma} \text{LB}.$

Let us consider a concrete example: Consider the loglinear model for counts, $y_i \sim \text{Poisson}(\delta_i), \delta_i = \exp(x_i^\top \theta)$ where $x_i, \theta \in \mathbb{R}^d$ are the vector of covariates and regression coefficients, respectively. Assume the prior of θ is $p(\theta) \sim \mathcal{N}(0, \sigma_0^2 \mathbb{I}_d)$. We use a Gaussian $q_\lambda(\theta) = \mathcal{N}(\theta|\mu, \Sigma)$ to approximate the true posterior distribution of θ . Let $y = y_{1:n}$ and $X = (x_1, x_2, \ldots, x_n)^\top$. For each $i = 1, \ldots, n$ let $w_i = \exp(x_i^\top \mu + \frac{1}{2}x_i^\top \Sigma x_i)$ and $W = \text{diag}(w_1, \ldots, w_n)$. By computing directly Tan [2021] showed that the closed form for the ELBO is,

$$LB(\lambda) = y^{\top} X \mu - \sum_{i=1}^{n} (w_i + \log(y_i!)) - \frac{\mu^{\top} \mu + Tr(\Sigma)}{2\sigma_0^2} + \frac{1}{2} \log|\Sigma| + \frac{d}{2} (1 - \log(\sigma_0^2)).$$

From here it is immediate to see that $\nabla_{\mu} LB$, $\nabla_{\Sigma} LB$, and $\nabla_{\text{vec}(\Sigma)} LB$ are given by

$$\begin{split} \nabla_{\mu} \mathbf{L} \mathbf{B} &= X^{\top} (y - w) - \frac{\mu}{\sigma_0^2}, \\ \nabla_{\Sigma} \mathbf{L} \mathbf{B} &= -\frac{1}{2} X^{\top} W X - \frac{1}{2\sigma_0^2} \mathbb{I} + \frac{1}{2} \Sigma^{-1} \\ \nabla_{\mathrm{vec}(\Sigma)} \mathbf{L} \mathbf{B} &= \frac{1}{2} \mathrm{vec} (\Sigma^{-1} - \frac{1}{\sigma_0^2} \mathbb{I} - X^{\top} W X) \end{split}$$

Also we have

$$\log q_{\lambda}(\theta) = -\frac{d}{2}\log(2\pi) - \frac{1}{2}\log|\Sigma| - \frac{1}{2}(\theta - \mu)^{\top}\Sigma^{-1}(\theta - \mu).$$

Hence

$$\nabla_{\lambda} q_{\lambda}(\theta) = (\nabla_{\mu} q_{\lambda}(\theta), \nabla_{\operatorname{vec}(\Sigma)} q_{\lambda}(\theta))$$
$$= \left((\theta - \mu)^{\top} \Sigma^{-1}, \operatorname{vec} \left(-\frac{1}{2} \Sigma^{-1} + \frac{1}{2} \Sigma^{-1} (\theta - \mu) (\theta - \mu)^{\top} \Sigma^{-1} \right)^{\top} \right)^{\top} \in \mathbb{R}^{d+d^{2}}.$$

We choose $n = 200, d = 3, \sigma_0^2 = 100, \theta = (1, \ldots, 1), x_{ij} \sim \mathcal{N}(0, 1)$, and simulate y_1, \ldots, y_{200} from those parameters. In this case $\lambda \in \mathbb{R}^{12}$ hence $I_F \in \mathbb{R}^{12 \times 12}$. The initial guesses are $\mu^{(0)} = (0, \ldots, 0)^{\top}$ and $\Sigma^{(0)} = 10^{-2}\mathbb{I}_d$. Moreover, we choose the step size $1/(1000+k)^{0.75}$ and $c_{\beta} = 1$. As we know the analytical form of the ELBO in this example, in Figure 3 we plot the graphs of $\mathrm{LB}(\lambda)$ (as a function of the number of iterations) obtained from the exact natural gradient ascent algorithm (NGVB), inversion free gradient algorithm (IFVB) and weighted inversion free algorithm (AIFVB). It can be seen that the three algorithms have almost identical performance in this case. One striking note is that AIFVB even outperforms (obtaining a larger lower bound value) the exact natural gradient algorithm in this case. It again helps to confirm that the averaging technique is useful. We note that we are tempted to include the ELBO obtained from the Euclidean gradient ascent but its performance is too unstable to include. This phenomenon has been observed in Tan [2021] as the matrix Σ obtained from the traditional gradient ascent is usually not positive definite, which results in an unstable performance.

Example 3 (Normalizing flow Variational Bayes). The choice of flexible variational distributions q_{λ} is important in VB. Normalizing flows [Rezende and Mohamed, 2015] are a class of techniques for designing flexible and expressive q_{λ} . We consider a flexible VB framework where the VB approximation distribution $q_{\lambda}(\theta)$ is constructed based on a normalizing flow as follows

$$\epsilon \sim \mathcal{N}_d(0, I), \quad Z = \psi(W_1 \epsilon + b_1), \quad \theta = W_2 Z + b_2$$

$$(6.1)$$

where $W_1, W_2 \in \text{Mat}(d, d)$ and b_1, b_2 are *d*-vectors, and $\psi(\cdot)$ is an activation function such as sigmoid. The transformation from ϵ to θ in (6.1) can be viewed as a neural network



Figure 3: Gaussian Approximation: Comparing NGVB, IFVB and AIFVB

with one hidden layer Z; it might be desirable to consider deeper neural nets with more than one hidden layers, but we do not consider it here. To be able to compute the density $q_{\lambda}(\theta)$ resulting from the transformations in (6.1), these transformations should be invertibe and the determinants of the Jacobian matrices should be easy to compute. To this end, we impose the orthogonality constraint on W_1 and W_2 : $W_1^{\top}W_1 = W_2^{\top}W_2 = \mathbb{I}_d$, i.e. the columns are orthonormal. Details on the derivation of the lower bound gradient and score function $\nabla \log q_{\lambda}(\theta)$ can be found in Appendix A.5.

Numerical results

We apply the manifold normalizing flow VB (NLVB) (6.1) to approximate the posterior distribution in a neural network classification problem, using the German Credit dataset. This dataset, available on the UCI Machine Learning Repository:

https://archive.ics.uci.edu/ml/index.php,

consists of observations on 1000 customers, each was already rated as being "good credit" (700 cases) or "bad credit" (300 cases). We create 10 predictors out of the available covariate variables including credit history, education, employment status, etc. The classification problem is based on a neural network with one hidden layer of 5 units. As W_1 and W_2 belong to the Stiefel manifold, for a comparison we use the VB on manifold algorithm of Tran et al. [2021] for updating these parameters. Figure 4 plots the lower bounds of the IFVB algorithm (solid red) together with the conventional VB algorithm (dash blue) using the Euclidean gradient (A.4) in the Appendix. As shown, the IFVB algorithm converges quicker and achieves a higher lower bound. Note that we do not

consider the AIFVB algorithm in this example, as the variational parameters belong to the Stiefel manifold, making the averaging technique challenging to use. It is interesting to extend the work to handle cases where the parameter space is a Riemannian manifold; however, we do not consider this in the present paper.



Figure 4: Normalizing flow VB: Lower bound values of Euclidean gradient VB (dash blue) versus IFVB (solid red) over the iterations.

Example 4 (Bayesian neural network). This example considers a Bayesian neural network for regression

$$y = \eta(x,\omega) + \epsilon, \ \epsilon \sim \mathcal{N}(0,\sigma^2),$$
(6.2)

where y is a real-valued response variable, $\eta(x, \omega)$ denotes the output of a neural network with the input vector $x = (x_1, ..., x_p)^\top \in \mathbb{R}^p$ and the vector of weights w. As neural networks are prone to overfitting, we follow Tran et al. [2020] and place a Bayesian adaptive group Lasso prior on the first-layer weights

$$w_{x_j}|\kappa_j \sim \mathcal{N}(0,\kappa_j \mathbb{I}_m), \quad \kappa_j|\gamma_j \sim \text{Gamma}\left(\frac{m+1}{2},\frac{\gamma_j^2}{2}\right), \quad j = 1,...,p,$$
 (6.3)

with the $\gamma_j > 0$ the shrinkage parameters; no regularization prior is put on the rest of the network weights. Here w_{x_j} denotes the vector of weights that connect the input x_j to the *m* units in the first hidden layer. An inverse-Gamma prior is used for σ^2 . We use empirical Bayes for selecting the shrinkage parameters γ_j , and the posterior of κ_j and σ^2 is approximated by a fixed-form within mean-filed VB procedure. See Tran et al. [2020] for the details.

The main task is to approximate the posterior of the network weights ω . Let d be the dimension of ω . We choose to approximate this posterior by a Gaussian variational distribution of the form $q_{\lambda}(\omega) = \mathcal{N}(\mu, \Sigma)$ with covariance matrix Σ having a factor form $\Sigma = bb^{\top} + \text{diag}(c)^2$, where b and c are vectors in \mathbb{R}^d . The vector of the variational parameters is $\lambda = (\mu^{\top}, b^{\top}, c^{\top})^{\top}$. This factor structure of the covariance matrix significantly reduces the size of variational parameters, making the Gaussian variational approximation method computationally efficient for Bayesian inference in large models such as Bayesian neural networks.

Tran et al. [2020] exploit the factor structure of Σ , and by setting certain sub-blocks of the Fisher information matrix $I_F(\lambda)$ to zero, to be able to derive a closed-form approximation of the inverse $I_F^{-1}(\lambda)$. Their VB method, termed the NAtural gradient Gaussian Variational Approximation with factor Covariance method (NAGVAC), is highly computationally efficient; however, the approximation of $I_F^{-1}(\lambda)$ might offset the VB approximation accuracy.

We now fit the Bayesian neural network model (6.2)-(6.3) to the Direct Marketing dataset [Jank, 2011] that consists of 1000 observations, of which 800 were used for training, and the rest for validation. The response y is the amount (in \$1000) a customer spends on the company's products per year, and 11 covariates include gender, income, married status, etc. We use a neural network with two hidden layers, each with ten units. Figure 5 plots the mean squared error (MSE) values, computed on the validation set, of the VB training using ADAM [Kingma and Ba, 2014], AIFVB, IFVB and NAGVAC. With the same stopping rule, the four methods, ADAM, AIFVB, IFVB and NAGVAC, stop after 425, 221, 480 and 700 iterations, respectively. This confirms the theoretical result in Theorem 5.3 that the averaging technique speeds up the convergence of AIFVB conpared to IFVB. On the validation set, the smallest MSE values produced by ADAM, AIFVB, IFVB and NAGVAC are 0.2273, 0.1750, 0.1749 and 0.1992, respectively. Both IFVB and AIFVB perform better than ADAM and NAGVAC, probably because the natural gradient approximation in NAGVAC, although being highly computationally efficient, might offset the approximation accuracy.



Figure 5: Bayesian neural network: Validation MSE values of ADAM (dotted black), AIFVB (dot greenish), IFVB (red solid) and NAGVAC (dash blue) over the iterations.

7 Conclusion

The paper introduced an efficient approach for approximating the inverse of Fisher information, a crucial component in variational Bayes used for approximating posterior distributions. An outstanding feature of our algorithm is its avoidance of calculating the Fisher matrix and its inversion. Instead, our approach generates a sequence of matrices converging to the inverse of Fisher information. Implementation of our method for natural gradient estimate does not require storage of large matrices. Our inversion free VB framework showcases versatility, enabling its application in a wide range of domains, including Gaussian approximation and normalizing flow Variational Bayes, and makes the natural gradient VB method applicable in cases that were impossible before. To demonstrate the efficiency and reliability of the method, we provided numerical examples as evidence of its effectiveness. We find it intriguing to consider expanding the scope of our approach to scenarios where the variational parameter space is a Riemannian manifold and to develop a rigorous theoretical framework for such cases. We plan to explore this avenue in our future research studies.

References

- Alekh Agarwal, Martin J Wainwright, Peter Bartlett, and Pradeep Ravikumar. Information-theoretic lower bounds on the oracle complexity of convex optimization. Advances in Neural Information Processing Systems, 22:1–9, 2009.
- Shun-Ichi Amari. Natural gradient works efficiently in learning. *Neural computation*, 10 (2):251–276, 1998.
- Kimon Antonakopoulos, Panayotis Mertikopoulos, Georgios Piliouras, and Xiao Wang. Adagrad avoids saddle points. In *International Conference on Machine Learning*, pages 731–771. PMLR, 2022.
- Bernard Bercu, Antoine Godichon, and Bruno Portier. An efficient stochastic Newton algorithm for parameter estimation in logistic regressions. *SIAM Journal on Control and Optimization*, 58(1):348–367, 2020.
- David M Blei, Alp Kucukelbir, and Jon D McAuliffe. Variational inference: A review for statisticians. *Journal of the American statistical Association*, 112(518):859–877, 2017.
- Léon Bottou, Frank E Curtis, and Jorge Nocedal. Optimization methods for large-scale machine learning. *Siam Review*, 60(2):223–311, 2018.
- Claire Boyer and Antoine Godichon-Baggioni. On the asymptotic rate of convergence of stochastic Newton algorithms and their weighted averaged versions. *Computational Optimization and Applications*, 84(3):921–972, 2023.
- Peggy Cénac, Antoine Godichon-Baggioni, and Bruno Portier. An efficient averaged stochastic Gauss-Newtwon algorithm for estimating parameters of non linear regressions models. arXiv preprint arXiv:2006.12920, 2020.
- Huy N Chau, J Lars Kirkby, Dang H Nguyen, Duy Nguyen, Nhu N Nguyen, and Thai Nguyen. On the inversion-free Newton's method and its applications. *International Statistical Review*, 2024.

- Aaron Defazio, Francis Bach, and Simon Lacoste-Julien. Saga: A fast incremental gradient method with support for non-strongly convex composite objectives. In Advances in neural information processing systems, pages 1646–1654, 2014.
- John Duchi, Elad Hazan, and Yoram Singer. Adaptive subgradient methods for online learning and stochastic optimization. *Journal of machine learning research*, 12(7), 2011.
- Marie Duflo. *Random iterative models*, volume 34. Springer Science & Business Media, 2013.
- Paolo Giordani, Xiuyan Mun, Minh-Ngoc Tran, and Robert Kohn. Flexible multivariate density estimation with marginal adaptation. *Journal of Computational and Graphical Statistics*, 22(4):814–829, 2013.
- Antoine Godichon-Baggioni and Nicklas Werge. On adaptive stochastic optimization for streaming data: A newton's method with o (dn) operations. *arXiv preprint arXiv:2311.17753*, 2023.
- Ian Goodfellow, Yoshua Bengio, and Aaron Courville. Deep learning. MIT press, 2016.
- Alex Graves. Practical variational inference for neural networks. Advances in neural information processing systems, 24, 2011.
- David Gunawan, Robert Kohn, and David Nott. Flexible variational Bayes based on a copula of a mixture. *Journal of Computational and Graphical Statistics*, pages 1–16, 2023.
- Matthew D Hoffman, David M Blei, Chong Wang, and John Paisley. Stochastic variational inference. *Journal of Machine Learning Research*, 2013.
- Wolfgang Jank. Business Analytics for Managers. Springer-Verlag New York, 2011.
- Rie Johnson and Tong Zhang. Accelerating stochastic gradient descent using predictive variance reduction. Advances in neural information processing systems, 26:315–323, 2013.
- Michael I Jordan, Zoubin Ghahramani, Tommi S Jaakkola, and Lawrence K Saul. An introduction to variational methods for graphical models. *Machine learning*, 37:183–233, 1999.
- Mohammad Khan and Wu Lin. Conjugate-computation variational inference: Converting variational inference in non-conjugate models to inferences in conjugate models. In *Artificial Intelligence and Statistics*, pages 878–887. PMLR, 2017.
- Diederik P Kingma and Jimmy Ba. Adam: A method for stochastic optimization. arXiv preprint arXiv:1412.6980, 2014.
- Diederik P Kingma and Max Welling. Auto-encoding variational Bayes. arXiv preprint arXiv:1312.6114, 2013.

- J Lars Kirkby, Dang H Nguyen, Duy Nguyen, and Nhu N Nguyen. Inversion-free subsampling Newton's method for large sample logistic regression. *Statistical Papers*, 63 (3):943–963, 2022.
- Harold Kushner and G George Yin. *Stochastic approximation and recursive algorithms and applications*, volume 35. Springer Science & Business Media, 2003.
- Jun Liu and Ye Yuan. On almost sure convergence rates of stochastic gradient methods. In *Conference on Learning Theory*, pages 2963–2983. PMLR, 2022.
- Nicholas T Longford. A fast scoring algorithm for maximum likelihood estimation in unbalanced mixed models with nested random effects. *Biometrika*, 74(4):817–827, 1987.
- Anna Lopatnikova and Minh-Ngoc Tran. Quantum variational Bayes on manifolds. In 2023 IEEE International Conference on Acoustics, Speech, and Signal Processing (ICASSP 2023), pages 1–5, 2023.
- Martin Magris, Mostafa Shabani, and Alexandros Iosifidis. Exact manifold Gaussian variational Bayes. arXiv preprint arXiv:2210.14598, 2022.
- James Martens. New insights and perspectives on the natural gradient method. The Journal of Machine Learning Research, 21(1):5776–5851, 2020.
- James Martens and Roger Grosse. Optimizing neural networks with Kronecker-factored approximate curvature. In *International conference on machine learning*, pages 2408–2417. PMLR, 2015.
- Gael M Martin, David T Frazier, and Christian P Robert. Approximating Bayes in the 21st century. *Statistical Science*, 1(1):1–26, 2023.
- Kevin P Murphy. Machine learning: a probabilistic perspective. MIT press, 2012.
- Lam M Nguyen, Jie Liu, Katya Scheinberg, and Martin Takáč. Sarah: A novel method for machine learning problems using stochastic recursive gradient. In *International Conference on Machine Learning*, pages 2613–2621. PMLR, 2017.
- Lam M Nguyen, Quoc Tran-Dinh, Dzung T Phan, Phuong Ha Nguyen, and Marten Van Dijk. A unified convergence analysis for shuffling-type gradient methods. *The Journal of Machine Learning Research*, 22(1):9397–9440, 2021.
- Victor MH Ong, David J Nott, Minh-Ngoc Tran, Scott A Sisson, and Christopher C Drovandi. Variational Bayes with synthetic likelihood. *Statistics and Computing*, 28: 971–988, 2018.
- Mariane Pelletier. On the almost sure asymptotic behaviour of stochastic algorithms. Stochastic processes and their applications, 78(2):217–244, 1998.
- Magnus Rattray, David Saad, and Shun-ichi Amari. Natural gradient descent for on-line learning. *Physical review letters*, 81(24):5461, 1998.

- Danilo Rezende and Shakir Mohamed. Variational inference with normalizing flows. In *International conference on machine learning*, pages 1530–1538. PMLR, 2015.
- Herbert Robbins and Sutton Monro. A stochastic approximation method. The annals of mathematical statistics, pages 400–407, 1951.
- Herbert Robbins and David Siegmund. A convergence theorem for non negative almost supermartingales and some applications. In *Optimizing methods in statistics*, pages 233–257. Elsevier, 1971.
- David Saad. On-line learning in neural networks. Cambridge University Press, 2009.
- Hugh Salimbeni, Stefanos Eleftheriadis, and James Hensman. Natural gradients in practice: Non-conjugate variational inference in Gaussian process models. In *International Conference on Artificial Intelligence and Statistics*, pages 689–697. PMLR, 2018.
- Masa-Aki Sato. Online model selection based on the variational Bayes. *Neural computation*, 13(7):1649–1681, 2001.
- Othmane Sebbouh, Robert M Gower, and Aaron Defazio. Almost sure convergence rates for stochastic gradient descent and stochastic heavy ball. In *Conference on Learning Theory*, pages 3935–3971. PMLR, 2021.
- James C Spall. Introduction to stochastic search and optimization: estimation, simulation, and control. John Wiley & Sons, 2005.
- Linda SL Tan. Analytic natural gradient updates for Cholesky factor in Gaussian variational approximation. arXiv preprint arXiv:2109.00375, 2021.
- Da Tang and Rajesh Ranganath. The variational predictive natural gradient. In International Conference on Machine Learning, pages 6145–6154. PMLR, 2019.
- Michalis Titsias and Miguel Lázaro-Gredilla. Doubly stochastic variational Bayes for nonconjugate inference. In *International conference on machine learning*, pages 1971–1979. PMLR, 2014.
- M.-N. Tran, N. Nguyen, D. Nott, and R. Kohn. Bayesian deep net GLM and GLMM. Journal of Computational and Graphical Statistics, 29(1):97–113, 2020. doi: 10.1080/ 10618600.2019.1637747.
- Minh-Ngoc Tran, David J Nott, and Robert Kohn. Variational Bayes with intractable likelihood. *Journal of Computational and Graphical Statistics*, 26(4):873–882, 2017.
- Minh-Ngoc Tran, Dang H Nguyen, and Duy Nguyen. Variational Bayes on manifolds. *Statistics and Computing*, 31:1–17, 2021. doi: doi.org/10.1007/s11222-021-10047-1.
- Steve Waterhouse, David MacKay, and Anthony Robinson. Bayesian methods for mixtures of experts. Advances in neural information processing systems, 8, 1995.

- William J Wilkinson, Simo Särkkä, and Arno Solin. Bayes–Newton methods for approximate Bayesian inference with psd guarantees. *Journal of Machine Learning Research*, 24(83):1–50, 2023.
- Matthew D Zeiler. Adadelta: an adaptive learning rate method. arXiv preprint arXiv:1212.5701, 2012.
- Fengshuo Zhang and Chao Gao. Convergence rates of variational posterior distributions. Annals of Statistics, 48(4):2180–2207, 2020.

SUPPLEMENTARY MATERIAL

8 Example 5 (Gaussian and Inverse Gamma variational approximation)

Let y = (11; 12; 8; 10; 9; 8; 9; 10; 13; 7) be observations from $\mathcal{N}(\mu, \sigma^2)$, the normal distribution with mean μ and variance σ^2 . We use the prior $\mathcal{N}(\mu_0, \sigma_0^2)$ for μ and Inverse-Gamma (α_0, β_0) for σ^2 with the hyperparameters $\mu_0 = 0, \sigma_0 = 10, \alpha_0 = 1, \beta_0 = 1$. The posterior distribution is written as

$$p(\mu, \sigma^2 | y) \propto p(\mu) p(\sigma^2) p(y | \mu, \sigma^2).$$

Assume that the VB approximation is $q_{\lambda}(\theta) = q(\mu)q(\sigma^2)$ with $q(\mu) = \mathcal{N}(\mu_{\mu}, \sigma_{\mu}^2)$, $q(\sigma^2) =$ Inverse-Gamma $(\alpha_{\sigma^2}, \beta_{\sigma^2})$, model parameter $\theta = (\mu, \sigma^2)^{\top}$ and the variational parameters $\lambda = (\mu_{\mu}, \sigma_{\mu}^2, \alpha_{\sigma^2}, \beta_{\sigma^2})$. Note that we have $h_{\lambda}(\theta) = h(\theta) - \log q_{\lambda}(\theta)$ with

$$h(\theta) = \log(p(\mu)p(\sigma^2)p(y|\mu, \sigma^2))$$

= $-\frac{n+1}{2}\log(2\pi) - \frac{1}{2}\log(\sigma_0^2) - \frac{(\mu-\mu_0)^2}{2\sigma_0^2} + \alpha_0\log(\beta_0) - \log\Gamma(\alpha_0)$
 $- (n/2 + \alpha_0 + 1)\log(\sigma^2) - \frac{\beta_0}{\sigma^2} - \frac{1}{2\sigma^2}\sum_{i=1}^n (y_i - \mu)^2,$

and

$$\log q_{\lambda}(\theta) = \alpha_{\sigma^2} \log \beta_{\sigma^2} - \log \Gamma(\alpha_{\sigma^2}) - (\alpha_{\sigma^2} + 1) \log \sigma^2 - \frac{\beta_{\sigma^2}}{\sigma^2} - \frac{1}{2} \log(2\pi) - \frac{1}{2} \log(\sigma_{\mu}^2) - \frac{(\mu - \mu_{\mu})^2}{\sigma_{\mu}^2}.$$

From here it can be seen that

$$\nabla_{\lambda} \log q_{\lambda}(\theta) = \left(\frac{\mu - \mu_{\mu}}{\sigma_{\mu}^2}, -\frac{1}{2\sigma_{\mu}^2} + \frac{(\mu - \mu_{\mu})^2}{2\sigma_{\mu}^4}, \log \beta_{\sigma^2} - \frac{\Gamma'(\alpha_{\sigma^2})}{\Gamma(\alpha_{\sigma^2})} - \log \sigma^2, \frac{\alpha_{\sigma^2}}{\beta_{\sigma^2}} - \frac{1}{\sigma^2}\right)^{\top}.$$

By direct calculation, it can be seen that the Fisher information matrix I_F is a diagonal block matrix with two main blocks

$$\begin{pmatrix} \frac{1}{\sigma_{\mu}^2} & 0\\ 0 & \frac{1}{2\sigma_{\mu}^4} \end{pmatrix}, \quad \text{and} \quad \begin{pmatrix} \frac{\partial^2 \log \Gamma(\alpha_{\sigma^2})}{\partial(\alpha_{\sigma^2})^2} & -\frac{1}{\beta_{\sigma^2}}\\ -\frac{1}{\beta_{\sigma^2}} & \frac{\alpha_{\sigma^2}^2}{\beta_{\sigma^2}^2} \end{pmatrix}.$$

In this example, the natural gradient can be computed in closed-form, which facilitates the testing of our IFVB and AIFVB algorithms. We use $(\bar{y}, 0.5, 1, 1)^{\top}$ as the initial guess for λ . The first and second panel of Figure 6 plot the posterior densities of μ and σ^2 using all four different approaches: MCMC, exact natural gradient VB (NGVB), inversion free VB (IFVB) and averaged inversion free VB (AIFVB). It can be seen that the posterior estimates obtained by IFVB and AIFVB are close to that of NGVB and MCMC. The last panel of Figure 6 plots the lower bound obtained from the VB methods. As shown, both IFVB and AIFVB converge almost as fast as NGVB even though NGVB uses the exact natural gradient in its computation.



Figure 6: The first two panels show that the IFVB and AIFVB estimates are close to that of the exact natural gradient ascent VB (NGVB) and MCMC. The last panel plots the lower bound estimates obtained from NGVB, IFVB and AIFVB.

9 Main proofs

Let us recall that $\tau_k = \frac{c_{\alpha}}{(c'_{\alpha}+k)^{\alpha}}$ with $c_{\alpha} > 0$, $c'_{\alpha} \ge 0$ and $\alpha \in (1/2, 1)$. In addition, $c_{\beta} \ge 0$ and $\beta \in (0, \alpha - 1/2)$. Recall that

$$\nabla_{\lambda} \mathcal{L}(\lambda) = \mathbb{E}_{q_{\lambda}}[-\nabla_{\lambda} \log q_{\lambda}(\theta)h_{\lambda}(\theta)] =: \mathbb{E}_{q_{\lambda}}[\ell(\theta, \lambda)].$$

We then have the recursive scheme

$$\lambda^{(k+1)} - \lambda^{(k)} = -\tau_{k+1} \tilde{\mathbf{H}}_{k+1}^{-1} \frac{1}{B} \sum_{i=1}^{B} \ell(\theta_{k+1,i}, \lambda^{(k)}), \quad \theta_{k+1,i} \sim q_{\lambda^{(k)}}(\theta)$$

9.1 Proof of Theorem 5.1

Proof. From Taylor's expansion of $\mathcal{L}(\lambda)$, since its Hessian is uniformly bounded, we have

$$\begin{split} \mathcal{L}(\lambda^{(k+1)}) &= \mathcal{L}(\lambda^{(k)}) + \nabla_{\lambda} \mathcal{L}^{\top}(\lambda^{(k)})(\lambda^{(k+1)} - \lambda^{(k)}) \\ &+ (\lambda^{(k+1)} - \lambda^{(k)})^{\top} \Big(\int_{0}^{1} (1-t) \nabla_{\lambda}^{2} \mathcal{L}(\lambda^{(k+1)} + t(\lambda^{(k)} - \lambda^{(k+1)})) dt \Big) (\lambda^{(k+1)} - \lambda^{(k)}) \\ &\leq \mathcal{L}(\lambda^{(k)}) + \nabla_{\lambda} \mathcal{L}^{\top}(\lambda^{(k)})(\lambda^{(k+1)} - \lambda^{(k)}) + \frac{1}{2} L_{0} \left\| (\lambda^{(k+1)} - \lambda^{(k)}) \right\|^{2} \\ &\leq \mathcal{L}(\lambda^{(k)}) - \alpha_{k+1} \nabla_{\lambda} \mathcal{L}^{\top}(\lambda^{(k)}) \tilde{\mathbf{H}}_{k+1}^{-1} \frac{1}{B} \sum_{i=1}^{B} \ell(\theta_{k+1,i}, \lambda^{(k)}) \\ &+ \frac{1}{2B} \alpha_{k+1}^{2} L_{0} \left\| \tilde{\mathbf{H}}_{k+1}^{-1} \right\|_{op}^{2} \sum_{i=1}^{B} \left\| \ell(\theta_{k+1,i}, \lambda^{(k)}) \right\|^{2} \end{split}$$

Hence

$$\mathcal{L}(\lambda^{(k+1)}) - \mathcal{L}(\lambda^{*}) \leq \mathcal{L}(\lambda^{(k)}) - \mathcal{L}(\lambda^{*}) - \alpha_{k+1} \nabla_{\lambda} \mathcal{L}^{\top}(\lambda^{(k)}) \tilde{\mathbf{H}}_{k+1}^{-1} \frac{1}{B} \sum_{i=1}^{B} \ell(\theta_{k+1,i}, \lambda^{(k)}) \\ + \frac{1}{2B} \alpha_{k+1}^{2} L_{0} \left\| \tilde{\mathbf{H}}_{k+1}^{-1} \right\|_{op}^{2} \sum_{i=1}^{B} \left\| \ell(\theta_{k+1,i}, \lambda^{(k)}) \right\|^{2}.$$

Let us denote $W_k = \mathcal{L}(\lambda^{(k)}) - \mathcal{L}(\lambda^*)$. From the above inequality we have

$$W_{k+1} \leq W_k - \alpha_{k+1} \nabla_{\lambda} \mathcal{L}^{\top}(\lambda^{(k)}) \tilde{\mathbf{H}}_{k+1}^{-1} \frac{1}{B} \sum_{i=1}^{B} \ell(\theta_{k+1,i}, \lambda^{(k)}) \\ + \frac{1}{2B} \alpha_{k+1}^2 L_0 \left\| \tilde{\mathbf{H}}_{k+1}^{-1} \right\|_{op}^2 \sum_{i=1}^{B} \left\| \ell(\theta_{k+1,i}, \lambda^{(k)}) \right\|^2.$$

Let us consider the filtration $\mathcal{F}_k = \sigma(\theta_{s,i}, \overline{\theta}_{s'}, Z_s : s \leq k : s' \leq k+1)$. As $\theta_{k+1,i}$ are independent of \mathcal{F}_k , taking the conditional expectation from the both sides of the above inequality, we have

$$\mathbb{E}[W_{k+1}|\mathcal{F}_{k}] \leq \mathbb{E}[W_{k}|\mathcal{F}_{k}] - \alpha_{k+1}\frac{1}{B}\sum_{i=1}^{B}\mathbb{E}[\nabla_{\lambda}\mathcal{L}^{\top}(\lambda^{(k)})\tilde{\mathbf{H}}_{k+1}^{-1}\ell(\theta_{k+1,i},\lambda^{(k)})|\mathcal{F}_{k}] \\ + \frac{1}{2B}\alpha_{k+1}^{2}L_{0}\sum_{i=1}^{B}\left\|\tilde{\mathbf{H}}_{k+1}^{-1}\right\|_{op}^{2}\mathbb{E}[\left\|\ell(\theta_{k+1,i},\lambda^{(k)})\right\|^{2}|\mathcal{F}_{k}] \\ \leq W_{k} - \alpha_{k+1}\nabla_{\lambda}\mathcal{L}^{\top}(\lambda^{(k)})\tilde{\mathbf{H}}_{k+1}^{-1}\nabla_{\lambda}\mathcal{L}(\lambda^{(k)}) \\ + \frac{1}{2B}\alpha_{k+1}^{2}L_{0}\left\|\tilde{\mathbf{H}}_{k+1}^{-1}\right\|_{op}^{2}\sum_{i=1}^{B}\mathbb{E}\left[\left\|\ell(\theta_{k+1,i},\lambda^{(k)})\right\|^{2}|\mathcal{F}_{k}\right].$$

Since $\mathbb{E}[\|\ell(\theta,\lambda)\|^2] \leq C_0 + C_1(\mathcal{L}(\lambda) - \mathcal{L}(\lambda^*))$, we have

$$\mathbb{E}[W_{k+1}|\mathcal{F}_{k}] \leq \left(1 + \frac{C_{1}L_{0}}{2}\alpha_{k+1}^{2} \left\|\tilde{\mathbf{H}}_{k+1}^{-1}\right\|_{op}^{2}\right)W_{k} - \alpha_{k+1}\lambda_{\min}(\tilde{\mathbf{H}}_{k+1}^{-1})\left\|\nabla_{\lambda}\mathcal{L}(\lambda^{(k)})\right\|^{2} + \frac{1}{2}\alpha_{k+1}^{2}C_{0}L_{0}\left\|\tilde{\mathbf{H}}_{k+1}^{-1}\right\|_{op}^{2}.$$

In order to apply the Robbins-Siegmund theorem [Robbins and Siegmund, 1971], we now focus on the behavior of the eigenvalues of $\tilde{\mathbf{H}}_s$. Observe that with the help of Toeplitz lemma (e.g., see the proof of Theorem A.1 in Appendix A.2).

$$(1-\beta)s^{1-\beta}\sum_{k=1}^{s}k^{-\beta}Z_kZ_k^\top \xrightarrow[s \to +\infty]{} I.$$

Then, as soon as $\lambda_{\min}\left(\tilde{\mathbf{H}}_{s}\right) \geq \lambda_{\min}\left(\frac{1}{s}H_{0} + \frac{1}{s}\sum_{k=1}^{s}c_{\beta}k^{-\beta}Z_{k}Z_{k}^{\top}\right)$, it comes

$$\lambda_{\max}\left(\tilde{\mathbf{H}}_{s}^{-1}\right) = O\left(s^{\beta}\right) \quad a.s$$

Then, since $\beta < \alpha - 1/2$, i.e $2\alpha - 2\beta > 1$, it comes

$$\sum_{k} \alpha_{k+1}^2 \left\| \tilde{\mathbf{H}}_{k+1}^{-1} \right\|^2 \le C \sum_{k} k^{2\beta - 2\alpha} < +\infty \quad a.s.$$

Using the Robbins-Siegmund theorem, one has that W_k converges almost surely to a finite random variable W_{∞} , which proves (i). We also conclude that

$$\sum_{k=0}^{\infty} \alpha_{k+1} \lambda_{\min}(\tilde{\mathbf{H}}_{k+1}^{-1}) \left\| \nabla \mathcal{L} \left(\lambda^{(k)} \right) \right\|^2 < \infty, \quad \text{a.s.}$$
(9.1)

We now have to control the smallest eigenvalue of $\tilde{\mathbf{H}}_{s+1}^{-1}$. In this aim, let us remark that the estimates of the Fisher Information can be written as

$$\tilde{\mathbf{H}}_{s} = \frac{1}{s} \left(H_{0} + \sum_{k=1}^{s} \beta_{k} Z_{k} Z_{k}^{T} \right) + \underbrace{\frac{1}{s} \sum_{k=0}^{s-1} \mathbb{E} \left[\nabla_{\lambda} \log q_{\overline{\lambda}^{(k)}}(\overline{\theta}_{k+1}) \nabla_{\lambda} \log q_{\overline{\lambda}^{(k)}}(\overline{\theta}_{k+1})^{\top} | \tilde{\mathcal{F}}_{k} \right]}_{=:R_{s}} + \underbrace{\frac{1}{s} \sum_{k=0}^{s-1} \Xi_{k+1}}_{=:M_{s}},$$

where $\tilde{\mathcal{F}}_{k} = \sigma\left(\theta_{s,i}, \overline{\theta}_{s}, Z_{s} : s \leq k, i \leq B\right)$ and $\Xi_{k+1} := -\mathbb{E}\left[\nabla_{\lambda} \log q_{\overline{\lambda}^{(k)}}(\overline{\theta}_{k+1}) \nabla_{\lambda} \log q_{\overline{\lambda}^{(k)}}(\overline{\theta}_{k+1})^{\top} | \tilde{\mathcal{F}}_{k}\right] + \nabla_{\lambda} \log q_{\overline{\lambda}^{(k)}}(\overline{\theta}_{k+1}) \nabla_{\lambda} \log q_{\overline{\lambda}^{(k)}}(\overline{\theta}_{k+1})^{\top}$ is a sequence of martingale differences. In addition

$$\mathbb{E}\left[\left\|M_{s+1}\right\|_{F}^{2}\left|\mathcal{F}_{s}\right] \leq \left(\frac{s}{s+1}\right)^{2}\left\|M_{s}\right\|_{F}^{2} + \frac{1}{(s+1)^{2}}\mathbb{E}\left[\left\|\nabla_{\lambda}\log q_{\overline{\lambda}^{(s)}}(\overline{\theta}_{s+1})\right\|^{4}\left|\tilde{\mathcal{F}}_{s}\right]\right].$$
 (9.2)

Since there are positive constants C'_0, C'_1 such that for all λ ,

$$\mathbb{E}\left[\left\|\nabla_{\lambda}\log q_{\lambda}(\theta)\right\|^{4}\right] \leq C_{0}' + C_{1}'\left(\mathcal{L}(\lambda) - \mathcal{L}(\lambda^{*})\right)^{2},$$

we have that $\mathbb{E}\left[\left\|\nabla_{\lambda}\log q_{\overline{\lambda}^{(k)}}(\overline{\theta}_{k+1})\right\|^{4} |\tilde{\mathcal{F}}_{k}\right] \leq C'_{0} + C'_{1}\overline{W}_{k}^{2}$ where $\overline{W}_{k} = \mathcal{L}\left(\overline{\lambda}^{(k)}\right) - \mathcal{L}(\lambda^{*})$. Then, if \mathcal{L} is not convex, one has $C'_{1} = 0$, otherwise, by the convexity of \mathcal{L} and with the help of the Toeplitz lemma,

$$\overline{W}_s \le \frac{1}{\sum_{k=0}^s \log(k+1)^w} \sum_{k=0}^s \log(k+1)^w \left(\mathcal{L}\left(\lambda^{(k)}\right) - \mathcal{L}\left(\lambda^*\right) \right) \xrightarrow[n \to +\infty]{a.s} W_{\infty}$$

Then, from (9.2), applying the Robbins-Siegmund theorem, $||M_s||_F^2 = \mathcal{O}(1)$ a.s. In a same way, one can check that $||R_s||_F^2 = \mathcal{O}(1)$ a.s. This means that at least,

$$\liminf_{k} \lambda_{\min}\left(\tilde{\mathbf{H}}_{k}^{-1}\right) > 0$$

so (9.1) implies that

$$\sum_{k=0}^{+\infty} \alpha_{k+1} \left\| \nabla \mathcal{L} \left(\lambda^{(k)} \right) \right\|^2 < +\infty \quad a.s.$$

Then, thanks to Lemma 2 in Liu and Yuan [2022] (or following the proof of Theorem 4.3 in Sebbouh et al. [2021]), it comes that

$$\min_{k=0,\dots,s} \left\| \nabla \mathcal{L} \left(\lambda^{(k)} \right) \right\|^2 = o\left(\frac{1}{\sum_{k=0}^s \alpha_{k+1}} \right) \quad a.s.$$

i.e. one has

$$\min_{k=0,\dots,s} \left\| \nabla \mathcal{L} \left(\lambda^{(k)} \right) \right\|^2 = o\left(\frac{1}{s^{1-\alpha}} \right) \quad a.s.$$

which proves (ii). One can apply Lemma 2 in Liu and Yuan [2022] again to obtain the result on $\overline{\lambda}^{(k)}$. This completes the proof of the theorem.

9.2 Proof of Corollary 5.1

Proof. Let us recall from the proof of Theorem 5.1 that $\tilde{\mathbf{H}}_s = \frac{1}{s} \left(H_0 + c_\beta \sum_{k=1}^s k^{-\beta} Z_k Z_k^T \right) + R_s + M_s$. Similar to the proof of Theorem A.1 in Appendix A.2, the norm of the first term can be estimated as

$$\left\|\frac{1}{s}\left(H_0 + c_\beta \sum_{k=1}^s k^{-\beta} Z_k Z_k^T\right)\right\| = \mathcal{O}\left(\max\left\{c_\beta s^{-\beta}, s^{-1}\right\}\right) \quad a.s.$$

which is is negligible. By the continuity of $I_F(\lambda)$ and since the convergence of $\lambda^{(k)}$ implies that $\overline{\lambda}^{(k)}$ converges almost surely to λ^* , it comes

$$R_s = \frac{1}{s} \sum_{k=0}^{s} \mathbb{E} \left[\nabla_{\lambda} \log q_{\overline{\lambda}^{(k)}}(\overline{\theta}_{k+1}) \nabla_{\lambda} \log q_{\overline{\lambda}^{(k)}}(\overline{\theta}_{k+1})^{\top} | \tilde{\mathcal{F}}_k \right] \xrightarrow[n \to +\infty]{a.s} I_F(\lambda^*).$$

Finally, applying Theorem 6.2 in Cénac et al. [2020], it comes that for any $\delta > 0$,

$$\|M_s\|_F^2 = o\left(\frac{\log s^{1+\delta}}{s}\right) \quad a.s.$$

which is negligible.

9.3 Proof of Theorem 5.2

Proof. The proof is adapted from Boyer and Godichon-Baggioni [2023] and Bercu et al. [2020]. Denoting $I_F = I_F(\lambda^*)$,

$$\lambda^{(k+1)} - \lambda^{*} = \lambda^{(k)} - \lambda^{*} - \tau_{k+1} \tilde{\mathbf{H}}_{k+1}^{-1} \frac{1}{B} \sum_{i=1}^{B} \ell(\theta_{k+1,i}, \lambda^{(k)})$$

$$= \lambda^{(k)} - \lambda^{*} - \tau_{k+1} \tilde{\mathbf{H}}_{k+1}^{-1} \nabla_{\lambda} \mathcal{L}(\lambda^{(k)}) + \tau_{k+1} \tilde{\mathbf{H}}_{k+1}^{-1} \underbrace{\left(\nabla_{\lambda} \mathcal{L}(\lambda^{(k)}) - \frac{1}{B} \sum_{i=1}^{B} \ell(\theta_{k+1,i}, \lambda^{(k)})\right)}_{\xi_{k+1}} \right)$$

$$= \lambda^{(k)} - \lambda^{*} - \tau_{k+1} I_{F}^{-1} \nabla_{\lambda} \mathcal{L}(\lambda^{(k)}) - \tau_{k+1} \underbrace{\left(\tilde{\mathbf{H}}_{k+1}^{-1} - I_{F}^{-1}\right) \nabla_{\lambda} \mathcal{L}(\lambda^{(k)})}_{=:\tau_{k}} + \tau_{k+1} \tilde{\mathbf{H}}_{k+1}^{-1} \xi_{k+1}} \right)$$

$$= \left(\mathbb{I} - \tau_{k+1} I_{F}^{-1} \nabla^{2} \mathcal{L}(\lambda^{*})\right) \left(\lambda^{(k)} - \lambda^{*}\right) - \tau_{k+1} r_{k} + \tau_{k+1} \tilde{\mathbf{H}}_{k+1}^{-1} \xi_{k+1} - \tau_{k+1} I_{F}^{-1} \underbrace{\left(\nabla \mathcal{L}\left(\lambda^{(k)}\right) - \nabla^{2} \mathcal{L}\left(\lambda^{*}\right) \left(\lambda^{(k)} - \lambda^{*}\right)\right)}_{=:\delta_{k}}$$

$$(9.3)$$

As explained in Antonakopoulos et al. [2022], since I_F and $\nabla^2 \mathcal{L}(\lambda^*)$ are symmetric and positive, $I_F^{-1} \nabla^2 \mathcal{L}(\lambda^*)$ and $I_F^{-1/2} \nabla^2 \mathcal{L}(\lambda^*) I_F^{-1/2}$ have the same eigenvalues, i.e there is a positive diagonal matrix Λ (of the eigenvalues of $I_F^{-1/2} \nabla^2 \mathcal{L}(\lambda^*) I_F^{-1/2}$) and a matrix Qsuch that $I_F^{-1} \nabla^2 \mathcal{L}(\lambda^*) = Q^{-1} \Lambda Q$. Then one can rewrite the previous decomposition as $Q(\lambda^{(k+1)} - \lambda^*) = (I - \tau_{k+1}\Lambda) Q(\lambda^{(k)} - \lambda^*) - \tau_{k+1}Qr_k + \tau_{k+1}Q\tilde{\mathbf{H}}_{k+1}^{-1}\xi_{k+1} - \tau_{k+1}QI_F^{-1}\delta_k.$ (9.4)

Then, with the help of an induction, it comes that

$$Q\left(\lambda^{(s)} - \lambda^{*}\right) = \beta_{s,0}Q\left(\lambda^{(0)} - \lambda^{*}\right) + \underbrace{\sum_{k=0}^{s-1} \beta_{s,k+1}\tau_{k+1}Q\tilde{\mathbf{H}}_{k+1}^{-1}\xi_{k+1}}_{=:M'_{s}} - \underbrace{\left(\sum_{k=0}^{s-1} \beta_{s,k+1}\tau_{k+1}Qr_{k} + \sum_{k=0}^{s-1} \beta_{s,k+1}\tau_{k+1}QI_{F}^{-1}\delta_{k}\right)}_{\Delta_{s}}.$$
(9.5)

_

with $\beta_{s,k} = \prod_{j=k+1}^{s} (\mathbb{I} - \alpha_j \Lambda)$ and $\beta_{s,s} = \mathbb{I}$. We now give the rate of convergence for each term in the decomposition (9.5).

Rate of convergence of $\beta_{s,0}Q(\lambda^{(0)} - \lambda^*)$. Since $\lambda_{\min}(\Lambda) > 0$ (because I_F and $\nabla^2 \mathcal{L}(\lambda^*)$ are positive), one can easily check that

$$\left\|\beta_{s,0}Q\left(\lambda^{(0)}-\lambda^{*}\right)\right\| = \mathcal{O}\left(\exp\left(-\lambda_{\min}(\Lambda)\frac{c_{\alpha}}{1-\alpha}s^{1-\alpha}\right)\right) \quad a.s$$

Rate of convergence of M'_s . Recall there are $\eta > \frac{1}{\alpha} - 1$ and positive constants $C_{\eta,0}, C_{\eta,1}$ such that for all λ ,

$$\mathbb{E}\left[\left\|l(\theta,\lambda)\right\|^{2+2\eta}\right] \leq C_{\eta,0} + C_{\eta,1} \left(\mathcal{L}(\lambda) - \mathcal{L}(\lambda^*)\right)^{1+\eta}$$

Then, one has with the help of Holder's inequality

$$\mathbb{E}\left[\left\|\xi_{k+1}\right\|^{2+2\eta}|\mathcal{F}_{k}\right] \leq 2^{2\eta+1}\frac{1}{B}\sum_{i=1}^{B}\mathbb{E}\left[\left\|\ell\left(\theta_{k+1,i},\lambda^{(k)}\right)\right\|^{2+2\eta}|\mathcal{F}_{k}\right] + 2^{2\eta+1}\left\|\nabla_{\lambda}\mathcal{L}\left(\lambda^{(k)}\right)\right\|^{2+2\eta}\right] \\ \leq 2^{2+2\eta}C_{\eta,0} + 2^{2+2\eta}C_{\eta,1}\left(\mathcal{L}\left(\overline{\lambda}^{(k)}\right) - \mathcal{L}\left(\lambda^{*}\right)\right)^{1+\eta}.$$

Since $\lambda^{(k)}$ is strongly consistent, the second term on the right-hand side of previous inequality converges almost surely to 0. In addition, thanks to Corollary 5.1, $\tilde{\mathbf{H}}_{k}^{-1}$ converges almost surely to I_{F}^{-1} . Then, with the help of Theorem 6.1 in Cénac et al. [2020],

$$\left\|M'_{s}\right\|^{2} = \mathcal{O}\left(\frac{\log s}{s^{\alpha}}\right) \quad a.s.$$

Rate of convergence of Δ_s . For *s* large enough, one has

$$\|\Delta_{s+1}\| \le (1 - \lambda_{\min}(\Lambda)\tau_{s+1}) \|\Delta_s\| + \tau_{s+1} \|Q\| \left(\|r_s\| + \|I_F^{-1}\| \|\delta_s\| \right).$$

Observe that since \mathcal{L} is twice continuously differentiable on a neighborhood of λ^* and since $\lambda^{(k)}$ is strongly consistent, it comes that $\|\delta_k\| = o\left(\|\lambda^{(k)} - \lambda^*\|\right)$ a.s. In addition, since $\tilde{\mathbf{H}}_k^{-1}$ converges almost surely to I_F^{-1} and since the gradient of \mathcal{L} is locally Lipschitz on a neighborhood of λ^* (since the Hessian is locally bounded by continuity), one has that $\|r_k\| = o\left(\|\lambda^{(s)} - \lambda^*\|\right)$ a.s. Then, $\|r_s\| + \|I_F^{-1}\| \|\delta_s\| = o\left(\|\lambda^{(s)} - \lambda^*\|\right)$ a.s, and with the help of decomposition (9.5), it comes that

$$\|r_s\| + \|I_F^{-1}\| \|\delta_s\| = o\left(\|\beta_{s,0}Q\left(\lambda^{(0)} - \lambda^*\right)\| + \|M_s'\| + \|\Delta_s\|\right) \quad a.s.$$

Thanks to previous convergence results, there exists a sequence of random variables r_s^\prime converging almost surely to 0 such that

$$\|\Delta_{s+1}\| \le (1 - \lambda_{\min}(\Lambda)\tau_{s+1}) \|\Delta_s\| + \tau_{s+1}r_{s+1} \left(\sqrt{\frac{\log s}{s^{\alpha}}} + \|\Delta_s\|\right).$$

Then, thanks to a stabilization Lemma (see Duflo [2013]), it comes

$$\|\Delta_s\| = \mathcal{O}\left(\sqrt{\frac{\log s}{s^{\alpha}}}\right) \quad a.s,$$

which concludes the proof.

9.4 Proof of Theorem 5.3

Proof. Observe that one can rewrite decomposition (9.3) as

$$\lambda^{(k+1)} - \lambda^{*} = \lambda^{(k)} - \lambda^{*} - \tau_{k+1} \tilde{\mathbf{H}}_{k+1}^{-1} \nabla_{\lambda} \mathcal{L} \left(\lambda^{(k)} \right) + \tau_{k+1} \tilde{\mathbf{H}}_{k+1}^{-1} \xi_{k+1} \\ = \lambda^{(k)} - \lambda^{*} - \tau_{k+1} \tilde{\mathbf{H}}_{k+1}^{-1} \nabla_{\lambda}^{2} \mathcal{L} (\lambda^{*}) \left(\lambda^{(k)} - \lambda^{*} \right) - \tau_{k+1} \tilde{\mathbf{H}}_{k+1}^{-1} \delta_{k} + \tau_{k+1} \tilde{\mathbf{H}}_{k+1}^{-1} \xi_{k+1},$$

where $\delta_k = \nabla \mathcal{L}(\lambda^{(k)}) - \nabla^2 \mathcal{L}(\lambda^*) (\lambda^{(k)} - \lambda^*)$ and $\xi_{k+1} = \nabla_\lambda \mathcal{L}(\lambda^{(k)}) - \frac{1}{B} \sum_{i=1}^B \ell(\theta_{k+1,i}, \lambda^{(k)})$. Denoting $u_k := \lambda^{(k)} - \lambda^*$ and $L^{-1} := \nabla_\lambda^2 \mathcal{L}(\lambda^*)^{-1}$, one can rewrite

$$\lambda^{(k)} - \lambda^* = L^{-1} \tilde{\mathbf{H}}_{k+1}^{-1} \frac{u_k - u_{k+1}}{\tau_{k+1}} + L^{-1} \xi_{k+1} - L^{-1} \delta_k.$$

Multiplying by $\log(k+1)^w$, then summing these equalities and dividing by $\sum_{k=0}^{s} \log(k+1)^w$, it comes

$$\overline{\lambda}^{(s)} - \lambda^* = L^{-1} \underbrace{\frac{1}{\sum_{k=0}^s \log(k+1)^w} \sum_{k=0}^s \log(k+1)^w \widetilde{\mathbf{H}}_{k+1}^{-1} \frac{u_k - u_{k+1}}{\tau_{k+1}}}_{=:A_{1,s}}}_{=:A_{1,s}} - L^{-1} \underbrace{\frac{1}{\sum_{k=0}^s \log(k+1)^w} \sum_{k=0}^s \log(k+1)^w \delta_k}_{A_{2,s}} + \underbrace{\frac{L^{-1}}{\sum_{k=0}^s \log(k+1)^w} \sum_{k=0}^s \log(k+1)^w \xi_{k+1}}_{M_{2,s}}}_{M_{2,s}}$$

The aim is to give the rate of convergence for each term on the right-hand side of the decomposition above. Let us first denote $t_s = \sum_{k=0}^s \log(k+1)^w$ and observe that

$$t_s \sim s \log(s+1)^w. \tag{9.6}$$

Rate of convergence of $A_{1,s}$. First, note that

$$A_{1,s} = \frac{1}{t_s} \sum_{k=0}^s \log(k+1)^w \frac{\tilde{\mathbf{H}}_k^{-1} u_k - \tilde{\mathbf{H}}_{k+1}^{-1} u_{k+1}}{\tau_{k+1}} + \frac{1}{t_s} \sum_{k=0}^s \log(k+1)^w \left(\tilde{\mathbf{H}}_{k+1}^{-1} - \tilde{\mathbf{H}}_k^{-1}\right) \frac{u_k}{\tau_{k+1}}$$

Concerning the first term on the right hand-side of previous equality, with the help of Abel's transform,

$$\frac{1}{t_s} \sum_{k=0}^s \log(k+1)^w \frac{\tilde{\mathbf{H}}_k^{-1} u_k - \tilde{\mathbf{H}}_{k+1}^{-1} u_{k+1}}{\tau_{k+1}} = -\frac{\log(s+1)^w \tilde{\mathbf{H}}_{s+1}^{-1} u_{s+1}}{t_s \tau_{s+1}} + \frac{\mathbf{1}_{w=0} \tilde{\mathbf{H}}_0^{-1} u_0}{t_s \alpha_1} + \sum_{k=1}^s \tilde{\mathbf{H}}_k^{-1} u_k \left(\frac{\log(k+1)^w}{\tau_{k+1}} - \frac{\log(k)^w}{\tau_k}\right).$$

With the help of an Abel's transform, one has

$$\frac{1}{t_s} \sum_{k=0}^s \log(k+1)^w \frac{\tilde{\mathbf{H}}_k^{-1} u_k - \tilde{\mathbf{H}}_{k+1}^{-1} u_{k+1}}{\tau_{k+1}} = \frac{-\log(s+1)^w \tilde{\mathbf{H}}_{s+1}^{-1} u_{s+1}}{t_s \tau_{s+1}} + \frac{\tilde{\mathbf{H}}_0^{-1} u_0 \mathbf{1}_{w=0}}{t_s \alpha_1} + \frac{1}{t_s} \sum_{k=1}^s \log(k+1)^w \tilde{\mathbf{H}}_k u_k \left(\frac{1}{\tau_{k+1}} - \frac{1}{\tau_k}\right)$$
(9.7)

Since $\tilde{\mathbf{H}}_s^{-1}$ converges almost surely to I_F^{-1} which is positive, and $|\tau_{k+1}^{-1} - \tau_k^{-1}| \leq \alpha c_{\alpha}^{-1} k^{1-\alpha}$ and with the help of Theorem 5.2 and equation (9.6), one can check that

$$\left\|\frac{1}{t_s}\sum_{k=0}^s \log(k+1)^w \frac{\tilde{\mathbf{H}}_k^{-1} u_k - \tilde{\mathbf{H}}_{k+1}^{-1} u_{k+1}}{\tau_{k+1}}\right\|^2 = \mathcal{O}\left(\frac{\log s}{s^{2-\alpha}}\right) \quad a.s.$$

which is negligible since $\alpha < 1$. For any $\delta > 0$, consider the event

$$E_k = \left\{ \left\| \lambda^{(k)} - \lambda^* \right\|^2 \le \frac{\log k^{1+\delta}}{k^{\alpha}}, \left\| \overline{\lambda}^{(k)} - \lambda^* \right\|^2 \le \frac{\log k^{1+\delta}}{k^{\alpha}} \right\}.$$

Thanks to Theorem 5.2, $\mathbf{1}_{E_k^C}$ converges almost surely to 0, and consequently

$$\left\|\frac{1}{t_s}\sum_{k=0}^s \log(k+1)^w \left(\tilde{\mathbf{H}}_{k+1}^{-1} - \tilde{\mathbf{H}}_k^{-1}\right) \frac{u_{k+1}}{\gamma_{k+1}} \mathbf{1}_{E_{k+1}^C}\right\|^2 = \mathcal{O}\left(\frac{1}{s^2 \log s^{2w}}\right) \quad a.s.$$

In addition,

$$\begin{split} \left\| \tilde{\mathbf{H}}_{k+1}^{-1} - \tilde{\mathbf{H}}_{k}^{-1} \right\|_{op} &\leq \left\| \tilde{\mathbf{H}}_{k+1}^{-1} \right\|_{op} \left\| \tilde{\mathbf{H}}_{k}^{-1} \right\|_{op} \left\| \tilde{\mathbf{H}}_{k}^{-1} \right\|_{op} \left\| \tilde{\mathbf{H}}_{k}^{-1} \right\|_{op} \left\| \tilde{\mathbf{H}}_{k}^{-1} \right\|_{op} \left(\frac{1}{k+1} \left\| \tilde{\mathbf{H}}_{k} \right\|_{op} + \underbrace{\frac{1}{k+1} \left\| \nabla_{\lambda} \log q_{\overline{\lambda}^{(k)}} \left(\overline{\theta}_{k+1} \right) \right\|^{2} + c_{\beta}(k+1)^{-\beta} \left\| Z_{k+1} \right\|^{2}}_{=\tilde{r}_{k+1}} \right\| \\ \end{split}$$

Then, since $\tilde{\mathbf{H}}_k^{-1}$ converges almost surely to a positive matrix, one can easily check that

$$\left\|\frac{1}{t_s}\sum_{k=0}^s \log(k+1)^w \frac{1}{k+1} \left\|\tilde{\mathbf{H}}_{k+1}^{-1}\right\|_{op} \left\|\tilde{\mathbf{H}}_{k}^{-1}\right\|_{op} \left\|\tilde{\mathbf{H}}_{k+1}\right\|_{op} \frac{\|u_k\|}{\tau_{k+1}} \mathbf{1}_{E_{k+1}} \mathbf{1}_{E_k}\right\|^2 = \mathcal{O}\left(\frac{\log s^{1+\delta}}{s^{2-\alpha}}\right) \quad a.s.$$

In addition, considering the filtration $\tilde{\mathcal{F}}_k$, and by hypothesis, one has

$$\frac{1}{t_s} \sum_{k=0}^{s} \frac{\log(k+1)^w}{k+1} \frac{\|\mathbf{u}_{k+1}\|}{\tau_{k+1}} \left\| \tilde{\mathbf{H}}_{k+1}^{-1} \right\|_{op} \left\| \tilde{\mathbf{H}}_{k}^{-1} \right\|_{op} \mathbf{1}_{E_{k+1}} \tilde{r}_{k+1} \\
\leq \underbrace{\frac{1}{t_s} \sum_{k=0}^{s} \frac{\log(k+1)^w}{k+1} \frac{\|\mathbf{u}_{k+1}\|}{\tau_{k+1}} \left\| \tilde{\mathbf{H}}_{k+1}^{-1} \right\|_{op} \left\| \tilde{\mathbf{H}}_{k}^{-1} \right\|_{op} \mathbf{1}_{E_{k+1}} \left(\underbrace{\frac{C_0'^{\frac{1}{2}} + C_1'^{\frac{1}{2}} (\mathcal{L}(\bar{\lambda}^{(k+1)}) - \mathcal{L}(\lambda^*))}{k+1} + dc_\beta(k+1)^{-\beta} \right)}_{=:(*)}}_{=:(*)}$$

where $\tilde{\xi}_{k+1} := \tilde{r}_{k+1} - \mathbb{E}\left[\tilde{r}_{k+1}|\tilde{\mathcal{F}}_k\right]$ is a martingale difference. Then, since $\overline{\lambda}^{(k)}$ converges almost surely to λ^* , it comes (at least)

$$\|(*)\|^2 = \mathcal{O}\left(\frac{\log s^{1+\delta}}{s^{2-\alpha}}\right) \quad a.s.$$

In a same way, since $\tilde{\xi}_k$ is a martingale difference satisfying

$$\mathbb{E}\left[\left\|\tilde{\xi}_{k+1}\right\|^{2}|\tilde{\mathcal{F}}_{k}\right] \leq \frac{2}{(k+1)^{2}}\left(C_{0}'+C_{1}\left(\mathcal{L}\left(\overline{\lambda}^{(k+1)}\right)-\mathcal{L}\left(\lambda^{*}\right)\right)^{2}\right)+6d^{2}c_{\beta}^{2}(k+1)^{-2\beta}$$

applying Theorem 6.2 in Cénac et al. [2020], it comes that at least

$$\|(**)\|^2 = \mathcal{O}\left(\frac{\log s^{1+\delta}}{s^{2-\alpha}}\right) \quad a.s$$

Rate of convergence of $A_{2,s}$. Thanks to inequality (5.3) and with the help of Theorem 5.2, it comes

$$\|\delta_k\| = \mathcal{O}\left(\frac{\log k}{k^{\alpha}}\right) \quad a.s.$$

Then, one can check that

$$||A_{2,s}||^2 = \mathcal{O}\left(\frac{\log s^2}{s^{2\alpha}}\right) \quad a.s$$

which is negligible as soon as $\alpha > 1/2$.

Rate of convergence of $M_{2,s}$. Let us denote $t'_s = \sum_{k=0}^s \log(k+1)^{2w}$, then considering the filtration \mathcal{F}_k ,

$$\frac{1}{t'_s} \sum_{k=0}^s \log(k+1)^{2w} \mathbb{E}\left[\xi_{k+1}\xi_{k+1}^T | \mathcal{F}_k\right] = \frac{1}{t'_s} \sum_{k=0}^s \log(k+1)^{2w} \nabla_\lambda \mathcal{L}\left(\lambda^{(k)}\right) \nabla_\lambda \mathcal{L}\left(\lambda^{(k)}\right)^T \\ + \underbrace{\frac{1}{t'_s} \sum_{k=0}^s \log(k+1)^{2w} \mathbb{E}\left[\frac{1}{B^2} \sum_{i=1}^B \ell\left(\theta_{k+1,i}, \lambda^{(k)}\right) \sum_{i=1}^B \ell\left(\theta_{k+1,i}, \lambda^{(k)}\right)^T | \mathcal{F}_k\right]}_{=:\langle M \rangle_s}.$$

Since the gradient of \mathcal{L} is continuous at λ^* and since $\lambda^{(k)}$ converges almost surely to λ^* , it comes that

$$\frac{1}{t'_s} \sum_{k=0}^s \log(k+1)^{2w} \nabla_{\lambda} \mathcal{L}\left(\lambda^{(k)}\right) \nabla_{\lambda} \mathcal{L}\left(\lambda^{(k)}\right)^T \xrightarrow[s \to +\infty]{a.s.} 0.$$

In addition, since $\theta_{k+1,1}, \ldots, \theta_{k+1,B}$ are i.i.d, it comes

$$\langle M \rangle_s = \frac{1}{Bt'_s} \sum_{k=0}^s \log(k+1)^{2w} \underbrace{\mathbb{E}_{q_{\lambda(k)}} \left[\ell\left(\theta, \lambda^{(k)}\right) \ell\left(\theta, \lambda^{(k)}\right)^T \right]}_{=\Sigma\left(\lambda^{(k)}\right)}$$

$$+ \underbrace{\frac{B-1}{Bt'_s} \sum_{k=0}^s \log(k+1)^{2w} \nabla_\lambda \mathcal{L}\left(\lambda^{(k)}\right) \nabla_\lambda \mathcal{L}\left(\lambda^{(k)}\right)^T}_{\xrightarrow{as}{s \to +\infty} 0} .$$

and since $\lambda^{(k)}$ converges almost surely to λ^* and since Σ is continuous, it comes

$$\langle M \rangle_s \xrightarrow[s \to +\infty]{a.s} \frac{1}{B} \Sigma \left(\lambda^* \right).$$

Then, thanks to the law of large numbers, one has

$$\left\|\sum_{k=0}^{s} \log(k+1)^{w} \xi_{k+1}\right\|^{2} = \mathcal{O}\left(\sum_{k=0}^{s} \log(k+1)^{2w} \log\left(\sum_{k=0}^{s} \log(k+1)^{2w}\right)\right) \quad a.s$$

ans since

$$\sum_{k=0}^{s} \log(k+1)^{2w} \sim \frac{1}{s+1} \left(\sum_{k=0}^{s} \log(k+1)^{w} \right)^{2} \sim (s+1) \log(s+1)^{2w},$$

it comes that

$$M_{2,s} = \mathcal{O}\left(\frac{\log s}{s}\right) \quad a.s.$$

In addition, with the help of a Central Limit Theorem for Martingales [Duflo, 2013], it comes that

$$\sqrt{Bs}M_{2,s} \xrightarrow[s \to +\infty]{\mathcal{L}} \mathcal{N}\left(0, L^{-1}\Sigma(\lambda^*)L^{-1}\right),$$

which concludes the proof.

A Appendix

A.1 Proof of Theorem 4.1

Proof. First, Riccati's equation (also known as the Sherman-Morrison formula) for matrix inversion in [Duflo, 2013, p. 96] states that for any $d \times d$ invertible matrix S, $d \times d$ invertible matrix T, $p \times d$ matrix U, and $p \times d$ matrix V, one has that the matrix S+UTV is invertible if $VS^{-1}U + T^{-1}$ is invertible and that, in this case,

$$(S + UTV)^{-1} = S^{-1} - S^{-1}U(VS^{-1}U + T^{-1})^{-1}VS^{-1}.$$

We will prove the claim by induction. Obviously it is true when s = 0. Now assume that it is true for s > 0. We have

$$\begin{aligned} H_{s+1}^{-1} &= H_s^{-1} - \left(1 + \phi_{i+1}^{\top} H_s^{-1} \phi_{s+1}\right)^{-1} H_s^{-1} \phi_{s+1} \phi_{s+1}^{\top} H_s^{-1} \\ &= \left(H_s + \phi_{s+1} \phi_{s+1}^{\top}\right)^{-1} \\ &= \left(H_0 + \sum_{j=1}^s \phi_j \phi_j^{\top} + \phi_{s+1} \phi_{s+1}^{\top}\right)^{-1} \\ &= \left(H_0 + \sum_{j=1}^{s+1} \phi_j \phi_j^{\top}\right)^{-1}. \end{aligned}$$

As a result, $H_{s+1} = H_0 + \sum_{j=1}^{s+1} \phi_j \phi_j^{\top}$. From this, it is clear that $H_k^{\top} = H_k$ for $k \in \mathbb{N}$. Lastly, we have for a vector v, $v^{\top}(\phi_j \phi_j^{\top})v = |v^{\top}\phi_j|^2 \ge 0$, the sum of positive definite matrices is positive definite, and the inverse of a positive definite matrix is positive definite. Hence the positivity and symmetry of H_s are preserved. Second, we have

$$\frac{1}{s}H_s = \frac{1}{s}H_0 + \frac{1}{s}\sum_{j=1}^s \nabla_\lambda \log q_\lambda(\theta_j) (\nabla_\lambda \log q_\lambda(\theta_j))^\top \longrightarrow I_F(\lambda), \quad \text{a.s.}$$

A.2 Proof of (4.4)

Recall that, for a fixed λ ,

$$A_s = A_s(\lambda) = H_0 + \sum_{j=1}^s \nabla_\lambda \log q_\lambda(\theta_j) (\nabla_\lambda \log q_\lambda(\theta_j))^\top + c_\beta \sum_{j=1}^s j^{-\beta} Z_j Z_j^\top$$

where Z_1, \ldots, Z_s are independent standard Gaussian vectors, $c_{\beta} \ge 0$ and $\beta \in (0, \alpha - 1/2)$, $\alpha \in (1/2, 1)$. One can update A_{s+1}^{-1} using the following scheme:

Theorem A.1. Let $\phi_s = \nabla_\lambda \log q_\lambda(\theta_s)$ and

$$A_{s+\frac{1}{2}}^{-1} = A_s^{-1} - \left(1 + \phi_{s+1}^{\top} A_s^{-1} \phi_{s+1}\right)^{-1} A_s^{-1} \phi_{s+1} \phi_{s+1}^{\top} A_s^{-1}$$

then

$$A_{s+1}^{-1} = A_{s+\frac{1}{2}}^{-1} - c_{\beta}(s+1)^{-\beta} \left(1 + c_{\beta}(s+1)^{-\beta} Z_{s+1}^{\top} A_{s+\frac{1}{2}}^{-1} Z_{s+1} \right)^{-1} A_{s+\frac{1}{2}}^{-1} Z_{s+1} Z_{s+1}^{\top} A_{s+\frac{1}{2}}^{-1}.$$

In particular, the positivity and symmetry of A_s 's is preserved. Moreover,

$$\frac{1}{s}A_s \longrightarrow I_F(\lambda) \quad a.s.$$

Proof. We will prove the claim by induction as in the proof of Theorem 4.1. The claim is obviously true when s = 0. Assume that it is true for some s > 0. By Riccati's formula,

$$A_{s+\frac{1}{2}}^{-1} = A_s^{-1} - \left(1 + \phi_{s+1}^{\top} A_s^{-1} \phi_{s+1}\right)^{-1} A_s^{-1} \phi_{s+1} \phi_{s+1}^{\top} A_s^{-1} = \left(A_s + \phi_{s+1} \phi_{s+1}^{\top}\right)^{-1}.$$
 (A.1)

Similarly,

$$A_{s+1}^{-1} = A_{s+\frac{1}{2}}^{-1} - c_{\beta}(s+1)^{-\beta} \left(1 + c_{\beta}(s+1)^{-\beta} Z_{s+1}^{\top} A_{s+\frac{1}{2}}^{-1} Z_{s+1} \right)^{-1} A_{s+\frac{1}{2}}^{-1} Z_{s+1} Z_{s+1}^{\top} A_{s+\frac{1}{2}}^{-1}$$

$$= \left(A_{s+\frac{1}{2}} + c_{\beta}(s+1)^{-\beta} Z_{s+1} Z_{s+1}^{\top} \right)^{-1}$$
(A.2)

Plugging (A.1) into (A.2), we have

$$A_{s+1}^{-1} = \left(A_s + \phi_{s+1}\phi_{s+1}^{\top} + c_{\beta}(s+1)^{-\beta}Z_{s+1}Z_{s+1}^{\top}\right)^{-1} \\ = \left(H_0 + \sum_{k=1}^{s+1}\phi_k\phi_k^{\top} + c_{\beta}\sum_{k=1}^{s+1}k^{-\beta}Z_kZ_k^{\top}\right)^{-1}.$$

Hence,

$$A_{s+1} = H_0 + \sum_{k=1}^{s+1} \phi_k \phi_k^\top + c_\beta \sum_{k=1}^{s+1} k^{-\beta} Z_k Z_k^\top.$$

From this, it is clear that $A_s^{\top} = A_s$ and $A_s > 0$ for $s \in \mathbb{N}$. Hence the positivity and symmetry of A_s are preserved.

Second, we have

$$\frac{1}{s}A_s = \frac{1}{s}H_0 + \frac{1}{s}\sum_{k=1}^s \phi_k \phi_k^{\top} + \frac{c_\beta}{s}\sum_{k=1}^s k^{-\beta} Z_k Z_k^{\top}.$$

It can be seen that

$$\frac{1}{s}H_0 + \frac{1}{s}\sum_{k=1}^s \phi_k \phi_k^\top \to 0 + \mathbb{E}[\phi\phi^\top] = \mathbb{E}_{q_\lambda} \left(\nabla_\lambda \log q_\lambda(\theta) \nabla_\lambda \log q_\lambda(\theta)^\top \right) = I_F(\lambda) \quad \text{a.s.}$$

Consider the term $\frac{c_{\beta}}{s} \sum_{k=1}^{s} k^{-\beta} Z_k Z_k^{\top}$, we have

$$\frac{c_{\beta}}{s} \sum_{k=1}^{s} k^{-\beta} Z_k Z_k^{\top} = \frac{c_{\beta}}{s} \left(\sum_{k=1}^{s} k^{-\beta} \right) \frac{1}{\sum_{k=1}^{s} k^{-\beta}} \sum_{k=1}^{s} \frac{1}{k^{\beta}} Z_k Z_k^{\top}.$$

Similar to Lemma 6.1 in Bercu et al. [2020] and recall that Z_k 's are independent copies of $Z \sim \mathcal{N}(0, \mathbb{I}_D)$, we have

$$\frac{1}{\sum_{k=1}^{s} k^{-\beta}} \sum_{k=1}^{s} k^{-\beta} Z_k Z_k^{\top} \to \mathbb{E}[ZZ^{\top}] = \mathbb{I}_D, \quad \text{a.s. as} \quad s \to \infty.$$

Also note that

$$\frac{1}{s}\sum_{k=1}^{s}k^{-\beta} = \frac{1}{s^{\beta}}\frac{1}{s^{1-\beta}}\sum_{k=1}^{s}k^{-\beta} \to 0 \quad \text{as} \quad s \to \infty,$$

due to the fact that $\frac{1}{s^{1-\beta}}\sum_{k=1}^{s}k^{-\beta} \to \frac{1}{1-\beta}$. As a result, we have

$$\frac{1}{s}\sum_{k=1}^{s}k^{-\beta}Z_kZ_k^{\top} \to 0 \quad \text{a.s. as} \quad s \to \infty.$$

Therefore

$$\frac{1}{s}A_s \longrightarrow I_F(\lambda)$$
 a.s as $s \to \infty$.

This completes the proof.

A.3 Calculation of the Hessian

Using the log derivative trick, $\nabla q_{\lambda}(\theta) = q_{\lambda}(\theta) \nabla \log q_{\lambda}(\theta)$, it can be seen that

$$\nabla_{\lambda}^{2} q_{\lambda}(\theta) = \nabla_{\lambda}^{2} \log q_{\lambda}(\theta) q_{\lambda}(\theta) + (\nabla_{\lambda} \log q_{\lambda}(\theta))^{2} q_{\lambda}(\theta)$$

Note that $h_{\lambda}(\theta)$ is a scalar. Hence

$$\begin{split} \nabla_{\lambda}^{2} \text{LB}(\lambda) &= \nabla_{\lambda} \left(\mathbb{E}_{q_{\lambda}} \left[\nabla_{\lambda} \log q_{\lambda}(\theta) h_{\lambda}(\theta) \right] \right) \\ &= \nabla_{\lambda} \int \nabla_{\lambda} \log q_{\lambda}(\theta) h_{\lambda}(\theta) q_{\lambda}(\theta) d\theta \\ &= \int \nabla_{\lambda}^{2} \log q_{\lambda}(\theta) h_{\lambda}(\theta) q_{\lambda}(\theta) d\theta + \int \nabla_{\lambda} h_{\lambda}(\theta) \nabla_{\lambda} \log q_{\lambda}(\theta)^{\top} q_{\lambda}(\theta) d\theta \\ &+ \int \nabla_{\lambda} \log q_{\lambda}(\theta) h_{\lambda}(\theta) \nabla_{\lambda} q_{\lambda}(\theta) d\theta \\ &= -\int \nabla_{\lambda} \log q_{\lambda}(\theta) \nabla_{\lambda} \log q_{\lambda}(\theta)^{\top} q_{\lambda}(\theta) d\theta \\ &+ \int \left(\nabla_{\lambda}^{2} \log q_{\lambda}(\theta) q_{\lambda}(\theta) + (\nabla_{\lambda} \log q_{\lambda}(\theta))^{2} q_{\lambda}(\theta) \right) h_{\lambda}(\theta) d\theta \\ &= -\mathbb{E}[\nabla_{\lambda} q_{\lambda}(\theta) \nabla_{\lambda} q_{\lambda}(\theta)^{\top}] - \int \nabla_{\lambda}^{2} q_{\lambda}(\theta) \cdot (\log q_{\lambda}(\theta) - \log p(y, \theta)) d\theta \\ &= -I_{F}(\lambda) - \int \nabla_{\lambda}^{2} q_{\lambda}(\theta) \cdot (\log q_{\lambda}(\theta) - \log p(\theta|y)) d\theta, \end{split}$$

where we have used the fact that $\nabla q_{\lambda}(\theta) = q_{\lambda}(\theta) \nabla \log q_{\lambda}(\theta)$ in the third line. Recall that $\mathcal{L}(\lambda) = -\text{LB}(\lambda)$, therefore

$$\nabla_{\lambda}^{2} \mathcal{L}(\lambda) = I_{F}(\lambda) + \int \nabla_{\lambda}^{2} q_{\lambda}(\theta) \cdot \big(\log q_{\lambda}(\theta) - \log p(\theta|y)\big) d\theta.$$

A.4 Convergence rate to Hessian

In this section we consider the convergence rate of $\tilde{\mathbf{H}}_s$ to $I_F := I_F(\lambda^*)$ and $\nabla^2_{\lambda} \mathcal{L}(\lambda^*)$.

Theorem A.2. Suppose that $\lambda^{(k)}$ converges almost surely to λ^* , and $\frac{\|\nabla_{\lambda}^2 q_{\lambda^*}(\theta)\|}{q_{\lambda^*}(\theta)}$ is bounded above by M > 0. Then

$$\left\|\tilde{\mathbf{H}}_{s}-I_{F}\right\|=\mathcal{O}\left(\max\left\{c_{\beta}s^{-\beta},s^{-1}\right\}\right)\quad a.s.$$

Furthermore, under the assumptions in Theorem 2.1 of Zhang and Gao [2020], the following holds true

$$\left\|\tilde{\mathbf{H}}_{s}-\nabla_{\lambda}^{2}\mathcal{L}(\lambda^{*})\right\|=\mathcal{O}\left(\max\left\{c_{\beta}s^{-\beta},s^{-1},n^{-1}\right\}\right)\quad a.s$$

with n the size of the data.

Proof. First recall that

$$\nabla_{\lambda}^{2} \mathcal{L}(\lambda) = I_{F}(\lambda) + \int \nabla_{\lambda}^{2} q_{\lambda}(\theta) \cdot \left(\log q_{\lambda}(\theta) - \log p(\theta|y)\right) d\theta.$$

and that

$$\tilde{\mathbf{H}}_{s} = \frac{1}{s} \left(H_{0} + \sum_{k=0}^{s-1} \nabla_{\lambda} \log q_{\overline{\lambda}^{(k)}}(\theta_{k+1}) (\nabla_{\lambda} \log q_{\overline{\lambda}^{(k)}}(\theta_{k+1}))^{\top} + c_{\beta} \sum_{k=1}^{s} k^{-\beta} Z_{k} Z_{k}^{T} \right)$$

Hence

$$\left\|\tilde{\mathbf{H}}_{s}-\nabla_{\lambda}^{2}\mathcal{L}(\lambda^{*})\right\| \leq \left\|\frac{1}{s}\left(H_{0}+\sum_{k=0}^{s-1}\nabla_{\lambda}\log q_{\overline{\lambda}^{(k)}}(\theta_{k+1})(\nabla_{\lambda}\log q_{\overline{\lambda}^{(k)}}(\theta_{k+1}))^{\top}+c_{\beta}\sum_{k=1}^{s}k^{-\beta}Z_{k}Z_{k}^{T}\right)-I_{F}\right\|$$
$$+\left\|\int\nabla_{\lambda}^{2}q_{\lambda^{*}}(\theta)\cdot\left(\log q_{\lambda^{*}}(\theta)-\log p(\theta|y)\right)d\theta\right\|$$
(A.3)

First, similar to the proof of Theorems 5.3 and A.1, we have

$$\begin{split} & \left\| \frac{1}{s} \left(H_0 + \sum_{k=0}^{s-1} \nabla_\lambda \log q_{\overline{\lambda}^{(k)}}(\theta_{k+1}) (\nabla_\lambda \log q_{\overline{\lambda}^{(k)}}(\theta_{k+1}))^\top + c_\beta \sum_{k=1}^s k^{-\beta} Z_k Z_k^T \right) - I_F \right\| \\ & \leq \left\| \frac{1}{s} \sum_{k=0}^{s-1} \nabla_\lambda \log q_{\overline{\lambda}^{(k)}}(\theta_{k+1}) (\nabla_\lambda \log q_{\overline{\lambda}^{(k)}}(\theta_{k+1}))^\top - \mathbb{E}[\nabla_\lambda \log q_\lambda(\theta) (\nabla_\lambda \log q_\lambda(\theta))^\top] \right\| \\ & + \left\| \frac{1}{s} \left(H_0 + c_\beta \sum_{k=1}^s k^{-\beta} Z_k Z_k^T \right) \right\| \\ & = O\left(\max\left\{ c_\beta s^{-\beta}, s^{-1} \right\} \right) \quad a.s. \end{split}$$

For the second term of(A.3),

$$\begin{split} \left\| \int \nabla_{\lambda}^{2} q_{\lambda^{*}}(\theta) \cdot \left(\log q_{\lambda^{*}}(\theta) - \log p(\theta|y)\right) d\theta \right\| \\ &\leq \int \frac{\left\| \nabla_{\lambda}^{2} q_{\lambda^{*}}(\theta) \right\|}{q_{\lambda^{*}}(\theta)} q_{\lambda^{*}}(\theta) \left(\log q_{\lambda^{*}}(\theta) - \log p(\theta|y)\right) | d\theta \\ &\leq M \int q_{\lambda^{*}}(\theta) \left|\log q_{\lambda^{*}}(\theta) - \log p(\theta|y)\right| d\theta \\ &= M \mathrm{KL}(q_{\lambda^{*}} \| p(\cdot|y)) \\ &\leq M C/n \end{split}$$

where the last inequality is from Theorem 2.1 of Zhang and Gao [2020]. By combining the estimates together, the desired result is followed immediately. \Box

A.5 Technical details for Example 4

Denote by $p_{\epsilon}(\epsilon)$, $p_Z(z)$ and $q_{\lambda}(\theta)$ the density function of random vectors ϵ , Z and θ , respectively, $\lambda = (\operatorname{vec}(W_1)^{\top}, b_1^{\top}, \operatorname{vec}(W_2)^{\top}, b_2^{\top})^{\top}$. Write D for the length of λ . We have that

$$p_{Z}(z) = p_{\epsilon}(\epsilon) \left| \frac{\partial \epsilon}{\partial z} \right| = p_{\epsilon}(\epsilon) \prod_{i=1}^{d} \frac{1}{h'(h^{-1}(z_{i}))}, \quad \epsilon = W_{1}^{\top}(h^{-1}(z) - b_{1})$$
$$= p_{\epsilon} \left(W_{1}^{\top}(h^{-1}(z) - b_{1}) \right) \prod_{i=1}^{d} \frac{1}{h'(h^{-1}(z_{i}))},$$

and

$$q_{\lambda}(\theta) = p_Z (z = W_2^{\top}(\theta - b_2)).$$

If we use $q_{\lambda}(\theta)$ to approximate a posterior distribution with prior $p(\theta)$ and log-likelihood $\ell(\theta)$, the lower bound is

$$LB(\lambda) = \mathbb{E}_{p_{\epsilon}}\left[\log p(\theta) + \ell(\theta) - \log p_{\epsilon}(\epsilon) + \sum_{1}^{d} \log h'(h^{-1}(z_{i}))\right]$$

where $z = h(W_1\epsilon + b_1)$, $\theta = W_2z + b_2$. It's straightforward to estimate $\nabla_{\lambda} \text{LB}(\lambda)$, if both $\log p(\theta)$ and $\ell(\theta)$ are differentiable in θ . After some algebra

$$\frac{\partial \theta}{\partial \operatorname{vec}(W_1)} = \epsilon^\top \otimes \left(W_2 \operatorname{diag}(h'(W_1 \epsilon + b_1)) \right), \quad \frac{\partial \theta}{\partial b_1} = W_2 \operatorname{diag}(h'(W_1 \epsilon + b_1))$$
$$\frac{\partial \theta}{\partial \operatorname{vec}(W_2)} = z^\top \otimes I_d, \quad \frac{\partial \theta}{\partial b_2} = I_d,$$

with \otimes the Kronecker product. Also,

$$\frac{\partial z}{\partial \operatorname{vec}(W_1)} = \epsilon^\top \otimes \operatorname{diag}(h'(W_1\epsilon + b_1)), \quad \frac{\partial z}{\partial b_1} = I_d, \frac{\partial z}{\partial \operatorname{vec}(W_2)} = 0, \quad \frac{\partial z}{\partial b_2} = 0.$$

Then $\frac{\partial \theta}{\partial \lambda}$ and $\frac{\partial z}{\partial \lambda}$ are the $d \times D$ matrices formed by

$$\frac{\partial \theta}{\partial \lambda} = \left[\frac{\partial \theta}{\partial \operatorname{vec}(W_1)}, \frac{\partial \theta}{\partial b_1}, \frac{\partial \theta}{\partial \operatorname{vec}(W_2)}, \frac{\partial \theta}{\partial b_2}, \right], \quad \frac{\partial z}{\partial \lambda} = \left[\frac{\partial z}{\partial \operatorname{vec}(W_1)}, \frac{\partial z}{\partial b_1}, \frac{\partial z}{\partial \operatorname{vec}(W_2)}, \frac{\partial z}{\partial b_2}\right]$$

It's now readily to compute the gradient of the lower bound

$$\nabla_{\lambda} \text{LB}(\lambda) = \mathbb{E}_{p_{\epsilon}} \left[\left(\frac{\partial \theta}{\partial \lambda} \right)^{\top} \nabla_{\theta} \left(\log p(\theta) + \ell(\theta) \right) + \left(\frac{\partial z}{\partial \lambda} \right)^{\top} h''(h^{-1}(z)) \right], \quad (A.4)$$

which can be estimated by sampling from p_{ϵ} .

In order to use the IFVB algorithm, we now derive the gradient $\nabla_{\lambda} \log q_{\lambda}(\theta)$. First, note that

$$\log q_{\lambda}(\theta) = -\frac{d}{2}\log(2\pi) - \frac{1}{2}(h^{-1}(z) - b_1)^{\top} W_1 W_1^{\top}(h^{-1}(z) - b_1) - \sum_{i=1}^d \log h'(h^{-1}(z_i)),$$

where $z = W_2^{\top}(\theta - b_2)$. It is easy to see that

$$\begin{aligned} \frac{\partial \log q_{\lambda}(\theta)}{\partial W_{1}} &= -(h^{-1}(z) - b_{1})(h^{-1}(z) - b_{1})^{\top}W_{1} = -\epsilon\epsilon^{\top}W_{1},\\ \frac{\partial \log q_{\lambda}(\theta)}{\partial b_{1}} &= W_{1}W_{1}^{\top}(h^{-1}(z) - b_{1}) = W_{1}\epsilon,\\ \frac{\partial \log q_{\lambda}(\theta)}{\partial W_{2}} &= (\theta - b_{2})\delta_{z}^{\top},\\ \frac{\partial \log q_{\lambda}(\theta)}{\partial b_{2}} &= -W_{2}\delta_{z}, \end{aligned}$$

where

$$\delta_{z} = -\operatorname{diag}\left(1/h'(h^{-1}(z))\right)W_{1}W_{1}^{\top}(h^{-1}(z) - b_{1}) - h''(h^{-1}(z))$$
$$= -\operatorname{diag}\left(1/h'(h^{-1}(z))\right)W_{1}\epsilon - h''(h^{-1}(z)).$$

Vectorizing these four terms and stacking them together gives $\nabla_{\lambda} \log q_{\lambda}(\theta)$.