Phylogeny of Twenty-One Mammals

Ray Han

Introduction

Phylogeny can be inferred using two sources of data from an organism: morphological data and molecular data. Historically, phylogenies were usually inferred using morphological characters (Lee and Palci, 2015), but some morphological features may not necessarily indicate shared heritage (Zou and Zhang, 2016). With the introduction of molecular phylogenies, the base sequence of genes, or amino acid sequence of proteins can be compared to find the number of similarities or differences to ascertain levels of relatedness between species. These two types of phylogenies are to be taken as a data-driven hypothesis about the evolutionary history of the studied organisms, and a consensus is drawn from the comparison between the different phylogenies built from the two sources of data, utilizing different methods.

It has been suggested that molecular phylogenies are more accurate than morphological phylogenies because it is believed that molecular characters are less susceptible to convergent evolution (Zou and Zhang, 2016). However, this does not mean that molecular phylogenies are to be considered over morphological phylogenies without indication to do so (Pisani et al., 2007). In some cases, morphological phylogenies can be the only phylogenies created, such as when the organism is extinct, and molecular data is unable to be gathered (Zou and Zhang, 2016). Therefore, using both morphological phylogenies and molecular phylogenies can provide independent evidence to support a hypothesis (Bucklin and Frost, 2009).

Two studies that related whales to taxa in the order Artiodactyla are good examples of using morphological phylogenies and molecular phylogenies to provide a unique view to support a hypothesis. The first study used a molecular approach, by characterizing two families of non-coding DNA sequences. Their findings resolved the phylogenetic relationships between whales, ruminants, hippopotamuses, and pigs, and provided evidence to support the hypothesis that whales, ruminants, and hippopotamuses formed a monophyletic group. Their findings challenged the traditional view that the order Artiodactyla, which included even-toed ungulates, is monophyletic (Shimamura et al., 1997). Their findings were consistent with an earlier 1 study, that used a morphological approach, comparing the morphological traits of whales and Basilosaurus, and found that the feet were paraxonic, i.e., bearing weight in 2 parallel axes within the foot in the order Cetacea which includes whales, which was consistent with serological evidence of a relationship to the order Artiodactyla (Gingrich et al., 1990).

This paper's goal is to infer the evolutionary history of twenty-one mammals by comparing a morphological phylogeny to a molecular phylogeny made by using the cytochrome c oxidase subunit 1 (COX-1) protein. From the literature, it is expected that the molecular and morphological phylogenies will have many clades overlapping, with minor differences (Pisani et al., 2007).

Materials and methods

Proteomic phylogeny - Cytochrome c oxidase subunit 1 (COX-1) protein is a component of cytochrome c oxidase, the last enzyme in the mitochondrial electron transport chain, that helps facilitate oxidative phosphorylation inside the mitochondria. The gene that encodes this protein is among the most conserved mitochondrial protein-coding genes in animals (Folmer et al., 1994), meaning its sequence remains relatively unchanged through evolution. This makes it a good protein to compare for evolutionary studies (Miya and Nishida, 2000) and is commonly used in research about animals (Girard et al., 2022).

To compare the twenty-one mammal taxa (Table 1), the amino acid sequences were obtained (The UniProt Consortium, 2023). These sequences were aligned by Clustal Omega (Sievers et al., 2011; McWilliam et al., 2013), which uses a method to align sequences using mathematical models and identifies areas within the sequences that are similar (Sievers et al., 2011; McWilliam et al., 2013). The percent identity matrix generated by Clustal Omega was saved (Table 2), and the alignment results were also saved in FASTA format.

The computer software Molecular Evolutionary Genetic Analysis (Tamura et al., 2021; Stecher et al., 2020) was used to construct the proteomic phylogenetic tree, using the maximum likelihood statistical method. Using this, an initial tree is built using a fast but suboptimal method. The branch lengths of this initial tree are then adjusted for maximum likelihood for the data set using a model of evolution, and variants of the branching pattern of the tree are then created using different methods to search for branching patterns that fit the data 2

better. Branch lengths are then computed for these variant trees and the greatest likelihood is retained as the best choice so far, and this search continues until no greater likelihoods are found. Some settings were set within the tree window for the initial tree method, model of evolution, and branching pattern variant methods. These settings were all on default settings, except for the branching pattern variant method, which was set as extensive subtree-pruning-regrafting. Subtree-pruning-regrafting prunes parts of the initial tree and then rearranges them, and then calculates the likelihood. This is done until the likelihoods of all possible tree orientations are calculated, and the branching pattern with the highest likelihood value is deemed the best tree (Wu, 2009). This was chosen because it provided a more expansive search for tree branching patterns than the other available option (Zhou et al., 2017).

Morphological Phylogeny - A morphological phylogeny was produced and obtained from the morphological traits of the mammals. It was made to be most parsimonious.

Comparison - The taxa were compared by clades, and which taxa were placed differently. Similar and identical clades between the proteomic phylogenies and the morphological phylogenies were noted and highlighted. Major differences in taxa placement between the phylogenies were also identified. The clades in the phylogenies were summarized in a table (Table 3), with taxa higher on the column being more closely related within the clade (e.g. the top two taxa are sister taxa), and differences highlighted in color.

Results

After comparing the morphological phylogeny (Figure 1), and the proteomic phylogeny (Figure 2), it was noted that the phylogenies mostly had similar clades, with some minor differences, as summarized in Table 3. Out of twenty-one taxa, fifteen taxa were grouped into the labeled clades I, II, III, IV, and V, which were present in and marked in both of the phylogenies, with the sister taxa arrangement between the taxa in clades II, III, IV, and V being identical. Six taxa were not grouped into clades, with five being identical, being *Colobus polykomos* (Colobus Monkey), *Cavia porcellus* (Guinea Pig), *Rattus norvegicus* (Rat), *Oryctolagus cuniculus* (Rabbit), and *Ornithorhynchus anatinus* (Platypus). The non-identical taxa were *Tamandua tetradactyla* 3

(Anteater) in the morphological phylogeny and the *Equus caballus* (Horse) in the proteomic phylogeny. It was also noted that clade I had two versions, clade I_1 in the morphological phylogeny, and clade I_2 in the proteomic phylogeny.

As shown in Table 3, clade I had two versions conflicting between both of them. Clade I₁ included *E*. *caballus*, *Dendrohyrax dorsalis* (Tree hyrax), *Loxodonta africana* (Elephant), *Trichechus manatus* (Manatee), and *E. caballus*, but in clade I₂, *E. caballus* was not present and was replaced with *T. tetradactyla*. In the morphological phylogeny, *T. tetradactyla* was more closely related to the taxa in clade IV, and in the proteomic phylogeny, *E. caballus* was more closely related to clade II. Also, in clade I₁, *D. dorsalis*, and *L. africana* were sister taxa, while in clade I₂, *D. dorsalis*, and *T. manatus* were the sister taxa. Through comparison of the proteomic phylogeny (Figure 2) and morphological phylogeny (Figure 1), it was observed that although not forming a clade in the morphological phylogeny, as they do in the proteomic phylogeny, *C. porcellus* (Guinea pig), and *O. cuniculus* were closely related in both phylogenies.

A major difference between the proteomic and morphological phylogenies was that in the proteomic phylogeny, *C. polykomos* was the most basal lineage, and *O. anatinus* was nested in other taxa, but the opposite was seen in the morphological phylogeny, with *O. anatinus* being the most basal lineage, while *C. polykomos* was nested in other taxa.

It was also observed that an even amount of toes, carnassial teeth, and the presence of a marsupium during gestation were associated with a clade that was also present in the proteomic phylogeny. Also observed was that aquatic taxa were mostly more closely related to terrestrial taxa than other aquatic taxa.




Figure 1: Morphological phylogeny of the twenty-one mammals, with common clades marked.





Figure 2: Proteomic phylogeny of twenty-one mammals, with common clades marked.

Common Name	Scientific Name	CO1 (UniProt)	CO1 (NCBI ID)		
Нірро	Hippopotamus amphibius	Q9ZZY9	9833		
Horse	Equus caballus	P48659	9796		
Dolphin	Tursiops truncatus	B9UD78	9739		
House Cat	Felis silvestris	P48888	9685		
Red Deer	Cervus elaphus	Q2V097	9860		
Guinea Pig	Cavia porcellus	Q9TEG9	10141		
Rat	Rattus norvegicus	P05503	10116		
Rabbit	Oryctolagus cuniculus	079429	9986		
Colobus Monkey	Colobus polykomos	099041	9572		
Racoon	Procyon lotor	Q3L6R3	9654		
Manatee	Trichechus manatus	B0JDY8	9778		
Anteater	Tamandua tetradactyla	Q8LWP0	48850		
Opossum	Didelphis virginiana	P41310	9267		
Sugar Glider	Petaurus breviceps	Q1MWH1	34899		
Tree Hyrax	Dendrohyrax dorsalis	B0JDW6	42325		
Elephant	Loxodonta africana	Q9TA27	9785		
Shrew	Sorex unguiculatus	Q94YE2	62275		
Red Panda	Ailurus fulgens	A7DXW2	9649		
Platypus	Ornithorhynchus anatinus	Q36452	9258		
Marsupial Mole	Notoryctes typhlops	Q5QRZ9	37699		
Mole	Urotrichus talpoides	Q7Y8E0	106106		

Mammal Phylogeny Taxa List



	Colobus	Marsupial Mole	Sugar Glider	Oppossum	Mole	Shrew	Platypus	Elephant	Rat	Tree Hyrax	Dolphin	Guinea Pig	Manatee	Ant eater	Rabbit	Hippo	Red Panda	Raccoon	Cat	Red deer	Horse
Colobus																					
Marsupial Mole	88.5	5																			
Sugar Glider	88.88	96.3																			
Opossum	90.06	96.49	96.69																		
Vole	88.13	93.96	93.76	93.57																	
Shrew	87.74	93.37	92.79	92.7	96.1	1															
Platypus	87.72	92.98	92.4	92.1	91.8	1 92.	2														
Elephant	87.94	91.42	90.84	91.03	91.2	5 91.6	3 90.0	6													
Rat	89.69	91.81	91.42	91.2	92.	8 93.3	9 91.8	1 90.6	6												
Tree Hyrax	88.72	93.18	92.59	92.79	92.	8 91.6	3 90.6	4 91.8	3 92.2	2											
Dolphin	87.16	91.42	91.42	91.6	92.0	2 90.8	6 90.2	5 90.2	9 91.8	3 92.2	2										
Guinea Pig	90.45	5 92.79	92.4	93.5	93.5	7 92.7	9 92.7	9 92.7	9 94.3	5 93.5	92.	1									
Manatee	89.67	92.79	93.18	93.76	93.7	6 93.1	8 92.	2 93.7	6 93.1	8 95.3	2 93.9	3 94.5	4								
Ant eater	89.67	93.18	92.79	93.3	93.5	7 93.1	8 92.5	9 92.9	8 93.5	7 94.7	92.9	94.9	3 96	5.1							
Rabbit	90.84	93.18	93.18	93.57	93.9	6 93.3	7 93.3	7 93.3	7 94.1	5 94.5	92.7	96.	3 95.	52 96	.3						
Hippo	89.1	92.98	92.59	93.18	93.5	8 92.	8 91.6	2 92.8	2 92.	8 94.7	5 94.9	5 94.1	5 94.	93 94.	74 94.	74					
Red Panda	90.08	93.76	93.96	93.7	95.1	4 93.3	9 92.5	9 92.	8 94.5	5 93.7	93.5	95.5	2 95.	32 95.	52 96	5.3 94.	94				
Raccoon	89.3	93.37	93.18	93.18	94.1	6 9	3 91.8	1 92.4	1 93.9	7 93.3	9 93.1	9 94.9	3 94.	54 94.	54 95.	52 94.	55 98.4	4			
Cat	90.08	93.57	93.76	93.3	95.3	3 93.7	7 92.7	9 9	3 93.9	7 94.5	5 94.1	3 95.1	3 95.	91 95.	32 96.4	49 96.	11 97.0	8 97.	47		
Red deer	89.49	94.15	94.15	93.76	95.3	3 93.5	8 92.	4 93.3	9 93.9	7 94.9	96.1	94.9	3 96	5.3 96	.1 95.	91 97.	08 96.8	9 96	.3 97.4	7	
Horse	89.88	93.37	93.96	93.3	95.3	3 93.3	9 92.5	9 92.6	1 94.1	6 94.3	95.1	95.1	3 96	5.1 95.	32 96	5.1 95. ³	53 97.2	8 96.	69 97.6	7 98.	05

Table 2: Percent identity matrix generated from Clustal Omega alignment.

I	П	Ш	IV	V
		Morphologic Phylogen	у	
Dendrohyrax dorsalis	Tursiops truncates	Ailurus fulgens	Nororyctes typhlops	
(Tree Hyrax)	(Dolphin)	(Red Panda)	(Shrew)	(Marsupial Mole)
Loxodonta Africana	Hippopotamus	Procyon lotor	Urotrichus talpoides	Petaurus breviceps
(Elephant)	amphibius(Hippo)	(Racoon)	(Mole)	(Sugar Glider)
Trichechus manatus	Cervus elaphus	Felis silvestris		Didelphis virginiana
(Manatee)	(Red Deer)	(House Cat)		(Opossum)
Equus caballus				
(Horse)				
		Molecular Phylogeny		
Dendrohyrax dorsalis	Tursiops	Ailurus fulgens	Urotrichus talpoides	Nororyctes typhlops
(Tree Hyrax)	truncatus(Dolphin)	(Red Panda)	(Mole)	(Marsupial Mole)
Trichechus manatus	Hippopotamus	Procyon lotor	Sorex unguiculatus	Petaurus breviceps
(Manatee)	amphibius(Hippo)	(Racoon)	(Shrew)	(Sugar Glider)
Loxodonta africana	Cervus elaphus	Felis silvestris		Didelphis virginiana
(Elephant)	(Red Deer)	(House Cat)		(Opossum)
Tamandua tetradactyla				
(Anteater)				

 Table 3: Table with the taxa in the clades, decreasing relatedness down the column, and differences highlighted.

Discussion

The goal of the paper was to form a consensus about the evolutionary histories of twenty-one mammals by comparing a morphological phylogeny created using morphological traits, and a molecular phylogeny created using the COX-1 protein. A comparison of these phylogenies showed large similarities between the proteomic and morphological phylogenies, with smaller differences. These differences could be the result of convergent evolution in the morphological traits, which is believed to happen more often than in molecular traits (Zou and Zhang, 2016), as well as different methods of processing data.

The presence of an even amount of toes, carnassial teeth, and the presence of a marsupium during gestation was associated with a clade that was also present in the proteomic phylogeny. Since each one of these defines a single clade, it can be hypothesized that these traits are affected by changes in the gene that encodes for the COX-1 protein. However, there is no such research currently to support this.

As previously mentioned, not all of the aquatic taxa were not closely related, and instead of being in a clade together, they were usually more closely related to other terrestrial taxa. These aquatic taxa are *T. manatus* in clade I₁, as well as *T. truncatus*, and *Hippopotamus amphibius* (Hippo) in clade II. It can be hypothesized that the aquatic taxa and the closely related terrestrial taxa had a common ancestor which was terrestrial, since there are many more terrestrial taxa, and eventually evolved to become aquatic. In the case of clade I₁, there is evidence that *L. africana*, *D. dorsalis*, and *T. manatus* were descended from a common ancestor (Prothero, 2009). In the case of clade II, this hypothesis is also supported by the findings that taxa of the order Cetacea, which includes *Tursiops truncatus* (Dolphin), are related to taxa of the order Artiodactyla, which includes *H. amphibius* and *Cervus elaphus* (Red deer) (Gingrich et al., 1990). It has also been well-documented that Cetaceans and *H. amphibius* had land ancestors (Thewissin et al., 2009; Orliac et al., 2023). This is also supported by the fact that it would have been more parsimonious to evolve the character of being aquatic than to lose the character, therefore it can be suggested that these aquatic mammals evolved from terrestrial ancestors.

It has been suggested that morphological phylogenies are less accurate than molecular ones. However, this is not to say that morphological phylogenies are to be completely disregarded. Instead, they are to be taken into account as much as molecular phylogenies when their results conflict and molecular phylogenies should

not immediately be considered to be more reliable without evidence (Pisani et al., 2007). One major difference between the proteomic and morphological phylogenies was the most basal taxon, with the most basal taxon in the morphological phylogeny being *O. anatinus*, and the most basal taxon in the proteomic phylogeny being *C. polykomos*. In the case of the different basal taxa, Research showed that the subfamily Colibinae, which included *C. polykomos* diverged about 5-9 million years ago (van der Kuyl et al., 1994), and families of platypus, which included *O. anatinus*, diverged about 17-90 million years ago (Bino et al., 2019). From this evidence, it is proposed that *O. anatinus* is the most basal taxon within the phylogeny of the twenty-one mammals.

One difference observed between the morphological and molecular phylogenies was the conflict between the 2 versions of clade I, namely the morphological version, clade I₁, and the proteomic version, clade I₂. There has been little evidence to suggest that one version of the clade is more accurate than the other, however, it has been well-documented that *D. dorsalis, L. africana, and T. manatus* are distantly related to each other (Kellogg et al., 2007). To resolve this conflict between the clades, it is proposed that *E. caballus* be excluded from the morphological clade I₁, *T. tetradactyla* be excluded from the proteomic clade I₂, and clade I simplified only to include *D. dorsalis, L. africana, and T. manatus*. Future research should be done to fully resolve the sister taxa arrangement of *D. dorsalis, L. africana, and T. manatus*, as it is still unclear as to whether the resolved I₁ or the resolved I₂ is more accurate in terms of sister taxa placement.

In this paper, the comparison and combination of two phylogenies, molecular and morphological, has been shown to be an appropriate method to come to a consensus about the evolutionary histories of twenty-one mammals, as the morphological and molecular phylogenies independently demonstrated the presence of four identical clades and one similar clade. Future research into this field could be done to determine the relatedness between the morphological traits, and changes in the COX-1 gene, as well as to fully resolve the differences between the proteomic clade I₂, and the morphological clade I₁. Furthermore, more research could be done to apply the phylogeny of the twenty-one mammal taxa to the universal tree of life, by comparing them with more taxa.

10

Bibliography

Bucklin, A; Frost, BW. 2009. Morphological and Molecular Phylogenetic Analysis of Evolutionary Lineages within Clausocalanus (Copepoda: Calanoida). Journal of Crustacean Biology. 29:1. 111-120.

Folmer, O; Black, M; Hoeh, W; Lutz, R; Vrijenhoek, R. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. Molecular Marine Biology and Biotechnology. 3:5. 294-299.

Gingrich, PD; Smith, HB; Simons, EL. 1990. Hind Limbs of Eocene Basilosaurus: Evidence of Feet in Whales. Science. 240:4965. 154-157.

Girard, EB; Langerak, A; Jompa, J; Wangensteen, OS; Macher; JN, Renema, W. 2022. Mitochondrial Cytochrome Oxidase Subunit 1: A Promising Molecular Marker for Species Identification in Foraminifera. Frontiers in Marine Science. 9:809659.

Kellogg, ME; Burkett, S; Dennis, TR; Stone, G; Gray, BA; McGuire, PM; Zori, RT; Stanyon, R. 2007. Chromosome painting in the manatee supports Afrotheria and Paenungulata. BMC Evolutionary Biology. 7:6.

Lee, MSY; Palci, A. 2015. Morphological Phylogenetics in the Genomic Age. Current Biology. 25:19. R922-R929.

McWilliam, H; Li, W; Uludag, M; Squizzato, S; Park, YM; Buso, N; Lopez, R. 2013. Analysis Tool Web Services from the EMBL-EBI. Nucleic Acids Research, 41:W1, W597–W600.

Miya, M; Nishida, M. 2000. Use of Mitogenomic Information in Teleostean Molecular Phylogenetics: A Tree-Based Exploration under the Maximum-Parsimony Optimality Criterion. Molecular Phylogenetics and Evolution. 17:3. 437-455.

Orliac, JM; Mourlam, JM; Boisserie, JR; Costeur, L; Lihoreau, F. 2023. Evolution of semiaquatic habits in hippos and their extinct relatives: insights from the ear region. Zoological Journal of the Linnean Society. X:X. 1-14.

Pisani, D; Benton, MJ; Wilkinson, M. 2007. Congruence of Morphological and Molecular Phylogenies. Acta Biotheoretica. 55:3. 269-281.

Prothero, DR. 2009. Evolutionary Transitions in the Fossil Record of Terrestrial Hoofed Mammals. Evolution: Education and Outreach. 2:2. 289-302

Shimamura, M; Yasue, H; Ohshima, K; Abe, H; Kato, H; Kishiro, T; Goto, M; Munechika, I; Okada, N. 1997. Molecular evidence from retroposons that whales form a clade within even-toed ungulates. Nature. 388:6643. 666-670.

Sievers, F; Wilm, A; Dineen, D; Gibson, TJ; Karplus, K; Li, W; Higgins, DG. 2014 Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. Molecular Systems Biology, 7:1, 539–539.

Stecher, G; Tamura, K; Kumar, S. 2020. Molecular Evolutionary Genetics Analysis (MEGA) for macOS. 37:4. 1237–1239.

Tamura, K; Stecher, G; Kumar, S. (2021). MEGA11: Molecular Evolutionary Genetics Analysis Version 11. Molecular Biology and Evolution. 38:7. 3022–3027.

Thewissen, JGM; Cooper, LN; George, JC; Bajpai, S. 2009. From Land to Water: the Origin of Whales, Dolphins, and Porpoises. Evolution: Education and Outreach, 2:2, 272–288.

UniProt Consortium, T. 2023. UniProt: the Universal Protein Knowledgebase in 2023. Nucleic Acids Research. 51:D1. D523–D531.

van der Kuyl, AC; Kuiken, C; Dekker, JT; Gouldsmit, J. 1994. Phylogeny of African Monkeys Based upon Mitochondrial 12S rRNA Sequences. Journal of Molecular Evolution. 40:2. 173-180

Wu, Y. 2009. A practical method for exact computation of subtree prune and regraft distance. Bioinformatics. 25:2. 190-196.

Zhou, X; Shen, X; Hittinger, CT; Rokas, A. 2017. Evaluating Fast Maximum Likelihood-Based Phylogenetic Programs Using Empirical Phylogenomic Data Sets. Molecular Biology and Evolution. 35:2. 486-503

Zou, Z; Zhang, J. 2016. Morphological and molecular convergences in mammalian phylogenetics. Nature Communications. 7:12758.