Focusing and defocusing mKdV equation with fully asymmetric nonzero boundary conditions: the inverse scattering transform

Yi ZHAO¹*, Dinghao ZHU¹

Abstract

In this paper, we investigate the inverse scattering transform(IST) for the focusing and defocusing mKdV equation with fully asymmetric nonzero boundary conditions. Our analysis focuses on the properties of the Jost function, allowing us to establish the associated Riemann-Hilbert (RH) problem for both the focusing and defocusing cases. Furthermore, we recover the potential from the RH problem, which plays a crucial role in calculating the long-time asymptotic behavior of the solution. In the symmetric case, the eigenvalues are given by $\lambda^2 = k^2 \pm q_0^2$. However, in the asymmetric case, the eigenvalues are represented as $\lambda_{\pm}^2 = k^2 \pm q_{\pm}^2$. As a result, we directly handle the branch cut instead of utilizing the four-sheeted Riemann surface. Naturally, this approach leads to the discontinuity of the functions across their respective branch cuts, which significantly impacts the entire development of the IST.

Keywords: Modified KdV equation; Inverse scattering tansform; Riemann-Hilbert problem; Integrable systems; Asymmetric nonzero boundary conditions.

¹ School of Mathematical Sciences, Laboratory of Mathematics and Complex Systems, MOE, Beijing Normal University, 100875 Beijing, China

[∗] Corresponding author and email address: zhaoy23@bnu.edu.cn

Contents

1 Introduction

In this paper, we study the Cauchy problem to the modified Korteweg-de Viries(mKdV) equation

$$
q_t - 6\sigma q^2 q_x + q_{xxx} = 0,
$$

\n
$$
q \to q_{\pm}, \quad x \to \pm \infty,
$$
\n(1.1)

where $q(x, t) \in \mathbb{R}$, $\sigma = \pm 1$ and q_{\pm} are real numbers. Without loss of generality, in the subsequent content, we assume that $q_{-} > q_{+} > 0$.

The Korteweg-de Vries(KdV) equation

$$
u_t + 6uu_x + u_{xxx} = 0 \t\t(1.2)
$$

was proposed by Korteweg and de Vries to describe wave propagation on the surface of shallow water[\[1](#page-20-0)].

It is well-known that the mKdV equation (1.1) admits the Lax pair $[2]$ $[2]$

$$
\phi_x = X\phi,\tag{1.3}
$$

$$
\phi_t = T\phi,\tag{1.4}
$$

where

$$
X = ik\sigma_3 + Q,
$$

\n
$$
T = 4k^2X - 2ik\sigma_3(Q_x - Q^2) + 2Q^3 - Q_{xx},
$$

with $Q =$ $\left(\begin{array}{cc} 0 & q(x,t) \\ \sigma q(x,t) & 0 \end{array}\right).$

In 1967, Miura presented an explicit nonlinear transformation(the well-known Miura transformation) $\begin{bmatrix} 3 \end{bmatrix}$ which related solutions of the KdV equation (1.2) and the mKdV equation (1.1) .

The mKdV equation [\(1.1\)](#page-1-1) arises in various physical fields, such as acoustic waves, phonons in a anharmonic lattices $(4, 5)$, Alfvén wave in cold collision-free plasma $(6, 7)$, dynamics of traffic flow $[8, 9]$ $[8, 9]$ and meandering ocean currents [\[10](#page-21-6)].

There has been significant research on the mKdV equation (1.1) in the field of integrable systems[\[11](#page-21-7)[–15](#page-22-0)]. The inverse scattering transform(IST) has been widely used to obtain the N-soliton solution for the mKdV equation [\(1.1\)](#page-1-1). Under zero boundary conditions(ZBCs), the N-soliton solution has been explicitly derived[\[16](#page-22-1), [17\]](#page-22-2). Under nonzero boundary conditions(NZBCs), Yeung et al obtained N-soliton solution for mKdV equation [\(1.1\)](#page-1-1) with nonzero vaccum parameter[\[18](#page-22-3), [19](#page-22-4)], while Alejo obtained the explicit breather solution for the focusing mKdV equation $[20]$ $[20]$. For both focusing and defocusing mKdV equation with symmetric NZBCs, Zhang and Yan developed the direct and inverse scattering problem using a suitable uniformization variable [\[21\]](#page-22-6). He et al reexamined the Hamiltonian formalism with NZBC[\[22\]](#page-22-7). Additionally, several studies have explored the asymptotic analysis of the mKdV equation (1.1) . The long-time asymptotic behavior with ZBCs was derived by the steepest descent method[\[23](#page-22-8)]. Baldwin discovered the long-time-asymptotic dispersive shock wave solution for the mKdV equation in 2013[\[24](#page-22-9)]. In 2016, Germain proved a full asymptotic stability result for solitary wave solutions of the mKdV equation[\[25](#page-22-10)].

To the best of our knowledge, no results are available for the mKdV equation with fully asymmetric boundary conditions.

The paper is organized as follows. In section [2,](#page-3-0) we focus on the focusing mKdV equation. Instead of using the Riemann surface approach, we directly handle the branch cut to establish the direct and inverse scattering transformation. We solve the Riemann-Hilbert (RH) problem and recover the potential from it. In section [3,](#page-13-0) we consider the defocusing mKdV equation. We present a different branch cut configuration, which leads to a key difference compared to the focusing case: the analyticity of the eigenfunction. Additionally, the self-adjoint operator property of the defocusing case results in a distinct discrete spectrum. We obtain the expression for the potential in this case as well.

2 The focusing mKdV equation

In this section, we consider the focusing case, i.e. $\sigma = -1$ First, let us establish and define some notations that will be used throughout this section.

$$
\begin{split} &\Sigma_{\pm}=[-iq_{\pm},iq_{\pm}],\\ &\Sigma_{0}=[-iq_{-},-iq_{+}]\cup[iq_{+},iq_{-}],\\ &\overset{\circ}{\Sigma}_{\pm}=(-iq_{\pm},iq_{\pm}),\\ &\overset{\circ}{\Sigma}_{0}=\left(-iq_{-},-iq_{+}\right)\cup(iq_{+},iq_{-}). \end{split}
$$

And three Pauli matrices are defined as

$$
\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.
$$

2.1 Direct scattering problem

Next, we will develop the spectral analysis and introduce the scattering matrix to prepare for the set-up of the RH problem.

2.1.1 Spectral analysis

To obtain the Jost functions of the spectral problem [\(1.3\)](#page-2-0), we need to diagonalize the asymptotic matrix $X_{\pm} = ik\sigma_3 + Q_{\pm}$. By calculation, the eigenvalues of the matrix X_{\pm} are $\pm i\lambda_{\pm}$, where

$$
\lambda_{\pm}^2 = k^2 + q_{\pm}^2.
$$

To avoid the multivalues of the function λ_{\pm} , we take the branch cut for λ_{\pm} on Σ_{\pm} . For all $k \in \mathbb{C}\setminus\Sigma_{\pm}$, we define λ_{\pm} as the single-valued, analytic and continuous as k approaches Σ_{\pm} from right. For $k \in \Sigma_{\pm}$, we take the positive value of the real square root for λ_{\pm} .

Fig. 1. The branch cuts Σ ₋ and Σ ₊ to the focusing case

The eigenvector matrix is

$$
E_{\pm}=I+\frac{i}{k-\lambda_{\pm}}\sigma_3Q_{\pm}
$$

corresponds to the eigenvalues $\mp i\lambda_{\pm}$, and the determinant is $\det E_{\pm} = \frac{2\lambda_{\pm}}{\lambda_{\pm}-k}$ \triangleq $\gamma_{\pm}(k)$.

After diagonalizition, we have

$$
\phi_{\pm} \sim E_{\pm} e^{-i\lambda_{\pm} x \sigma_3}, \quad x \to \pm \infty. \tag{2.1}
$$

Define the normalized Jost functions μ_\pm as

$$
\mu_{\pm} = \phi_{\pm} e^{i\lambda_{\pm} x \sigma_3},\tag{2.2}
$$

then we have

$$
\mu_{\pm} \to E_{\pm}, \quad x \to \pm \infty.
$$

And they satisfy the Volterra integral equation

$$
\mu_{\pm} = E_{\pm}(k) + \int_{\pm\infty}^{x} E_{\pm}(k) e^{-i\lambda_{\pm}(x-y)\hat{\sigma}_{3}} \left[E_{\pm}^{-1}(k) \Delta Q_{\pm}(y) \mu_{\pm}(y,k) \right] dy \tag{2.3}
$$

with $\Delta Q_{\pm} = Q - Q_{\pm}$.

Lemma 2.1. If $(1+|x|)(q-q_{\pm}) \in L^1(\mathbb{R}_{\pm})$, then

- Analyticity: the function $\mu_{-,1}(x,t,k)$ and $\mu_{+,2}(x,t,k)$ are analytic in $\mathbb{C}^+ \setminus [0, iq_-]$, whereas the function $\mu_{+,1}(x,t,k)$ and $\mu_{-,2}(x,t,k)$ are analytic in $\mathbb{C}^-\setminus[-iq_-,0]$.
- Symmetry: Let $\widetilde{\phi}_{\pm} = \phi_{\pm}(x, t, -\lambda_{\pm}(k))$, then we have

$$
\phi_{\pm} = \tilde{\phi}_{\pm} \frac{i}{k - \lambda_{\pm}} \sigma_3 Q_{\pm}, \quad k \in \mathring{\Sigma}_{\pm}, \tag{2.4}
$$

Column-wise reads

$$
\phi_{\pm,1} = \frac{i q_{\pm}}{k - \lambda_{\pm}} \widetilde{\phi}_{\pm,2}, \quad k \in \mathring{\Sigma}_{\pm}, \tag{2.5}
$$

$$
\phi_{\pm,2} = \frac{i q_{\pm}}{k - \lambda_{\pm}} \widetilde{\phi}_{\pm,1}, \quad k \in \mathring{\Sigma}_{\pm}.
$$
 (2.6)

Proof. We only prove the symmetry. Note that in defining λ_{\pm} , we could have taken the opposite sign of the comlex square roots. With some abuse of notation, we temporaily express the dependence of the Jost functions on the choice of sign explicitly.

From the above discussion, if ϕ_{\pm} solves the spectral problem [\(1.3\)](#page-2-0), so does $\widetilde{\phi}_{\pm}$. Recall the asymptotic $\phi_{\pm} \sim \left(I + \frac{i}{k - \lambda_{\pm}} \sigma_3 Q_{\pm} \right) e^{-i\lambda_{\pm} x \sigma_3}$ as $x \to \pm \infty$, thus we have $\widetilde{\phi}_{\pm} \sim \left(I + \frac{i}{k + 1} \right)$ $\frac{i}{k+\lambda_{\pm}}\sigma_3Q_{\pm}\bigg)\,e^{i\lambda_{\pm}x\sigma_3}$. Direct check leads to

$$
\phi_{\pm} = \tilde{\phi}_{\pm} \frac{i}{k - \lambda_{\pm}} \sigma_3 Q_{\pm}, \quad k \in \mathring{\Sigma}_{\pm}, \tag{2.7}
$$

which can be rewittten column-wise as

$$
\phi_{\pm,1} = \frac{i q_{\pm}}{k - \lambda_{\pm}} \widetilde{\phi}_{\pm,2}, \quad k \in \mathring{\Sigma}_{\pm}, \tag{2.8}
$$

$$
\phi_{\pm,2} = \frac{i q_{\pm}}{k - \lambda_{\pm}} \widetilde{\phi}_{\pm,1}, \quad k \in \mathring{\Sigma}_{\pm}.
$$
 (2.9)

 \Box

2.1.2 Scattering matrix

There exists a scattering matrix $S(k)$ such that

$$
\phi_{-}(x,t,k) = \phi_{+}(x,t,k)S(k), \quad k \in \Sigma_{+}.
$$
\n(2.10)

Thus, we have det $S = \frac{\gamma}{\gamma}$ $\frac{\gamma}{\gamma_+}$. And the scattering coefficients can be written as the Wronskians expressions

$$
s_{11} = \frac{\text{Wr}(\phi_{-,1}, \phi_{+,2})}{\gamma_+}, \quad s_{12} = \frac{\text{Wr}(\phi_{-,2}, \phi_{+,2})}{\gamma_+}, s_{21} = \frac{\text{Wr}(\phi_{+,1}, \phi_{-,1})}{\gamma_+}, \quad s_{22} = \frac{\text{Wr}(\phi_{+,1}, \phi_{-,2})}{\gamma_+}.
$$
\n(2.11)

Define

$$
\rho = \frac{s_{21}}{s_{11}}, \quad \tilde{\rho} = \frac{s_{12}}{s_{22}}.
$$
\n(2.12)

Lemma 2.2. Let $s_{11}^-(k) = \lim_{\epsilon \uparrow 0} s_{11}(\epsilon + k)$ for $k \in \Sigma_-$, then we have

$$
s_{11}^{-}(k) = s_{22}(k)\frac{(k + \lambda_{+})(k - \lambda_{-})}{-q_{+}q_{-}}, \quad k \in \Sigma_{+},
$$

\n
$$
s_{11}^{-}(k) = s_{12}(k)\frac{k - \lambda_{-}}{iq_{-}}, \quad k \in \Sigma_{0}.
$$
\n(2.13)

Let $s_{22}^-(k) = \lim_{\epsilon \uparrow 0} s_{22}(\epsilon + k)$ for $k \in \Sigma_-$, then we have

$$
s_{22}^{-}(k) = s_{11}(k)\frac{(k + \lambda_{+})(k - \lambda_{-})}{-q_{+}q_{-}}, \quad k \in \Sigma_{+},
$$

\n
$$
s_{22}^{-}(k) = s_{21}(k)\frac{k - \lambda_{-}}{iq_{-}}, \quad k \in \Sigma_{0}.
$$
\n(2.14)

Proof. For $k \in \Sigma_+$, we know $\gamma_+^{\perp} \triangleq \lim_{\epsilon \uparrow 0} \gamma_+(k+\epsilon) = \frac{2\lambda_+}{k+\lambda_+}$. It follows from the Wronskians expression of s_{11} [\(2.11\)](#page-6-1), [\(2.5\)](#page-5-0) and [\(2.6\)](#page-5-1) that

$$
s_{11}^{-} = \frac{\text{Wr}(\phi_{-,1}^{-}, \phi_{+,2}^{-})}{\gamma_{+}^{-}} = \frac{k + \lambda_{+}}{2\lambda_{+}} \text{Wr}\left(\frac{k - \lambda_{-}}{iq_{-}}\phi_{-,2}, \frac{k - \lambda_{+}}{iq_{+}}\phi_{+,1}\right) = s_{22} \frac{(k + \lambda_{+})(k - \lambda_{-})}{-q_{+}q_{-}}.
$$
\n(2.15)

For $k \in \Sigma_0$, the function $\phi_{+,2}$ has no jump, i.e, $\phi_{+,2}^- = \phi_{+,2}$, thus we have

$$
s_{11}^{-} = \frac{\lambda_{+} - k}{2\lambda_{+}} \text{Wr}(\phi_{-,1}^{-}, \phi_{+,2})
$$

=
$$
\frac{\lambda_{+} - k}{2\lambda_{+}} \text{Wr}\left(\frac{k - \lambda_{-}}{iq_{-}} \phi_{-,2}, \phi_{+,2}\right)
$$

=
$$
s_{12} \frac{k - \lambda_{-}}{iq_{-}}.
$$
 (2.16)

Performing a similar procedure leads to

$$
s_{22}^{-} = \frac{\text{Wr}(\phi_{+,1}^{-}, \phi_{-,2}^{-})}{\gamma_{+}^{-}} = \frac{k + \lambda_{+}}{2\lambda_{+}} \text{Wr}\left(\frac{k - \lambda_{+}}{iq_{+}}\phi_{+,2}, \frac{k - \lambda_{-}}{iq_{-}}\phi_{-,1}\right) = s_{11} \frac{(k + \lambda_{+})(k - \lambda_{-})}{-q_{+}q_{-}}, \quad k \in \Sigma_{+}
$$
\n(2.17)

and

$$
s_{22}^{-} = \frac{\lambda_{+} - k}{2\lambda_{+}} \text{Wr}\left(\phi_{+,1}, \frac{k - \lambda_{-}}{iq_{-}} \phi_{-,1}\right)
$$

= $\frac{\lambda_{+} - k}{2\lambda_{+}} \text{Wr}\left((\phi_{+,1}, \frac{k - \lambda_{-}}{iq_{-}} \phi_{-,1}\right)$
= $s_{21} \frac{k - \lambda_{-}}{iq_{-}}, \quad k \in \Sigma_{0}.$ (2.18)

2.1.3 Discrete spectrum and residue condition

Suppose $s_{11}(k, t)$ has J simple zeros in $\mathbb{C}^+\setminus[0, iq_-]$ denoted by k_j , $j = 1, 2, ..., J$. From the Wronskians expression of $s_{11}(k, t)$, there exists a constant only depending on t such that

$$
\phi_{-,1}(x,t,k_j) = b_j(t)\phi_{+,2}(x,t,k_j). \tag{2.19}
$$

On account of (3.2) , we have

$$
\mu_{-,1}(x,t,k_j) = b_j(t)\mu_{+,2(x,t,k_j)}e^{i(\lambda_{+,j}+\lambda_{-,j})x},
$$

thus for $j = 1, 2, \ldots, J$, we get

$$
\operatorname{Res}_{k=k_j} \left[\frac{\mu_{-,1}(x,t,k)}{s_{11}(k,t)} \right] = \frac{\mu_{-,1}(k_j)}{s'_{11}(k_j)} = C_j(t) e^{i(\lambda_{+,j} + \lambda_{-,j})x} \mu_{+,2}(x,t,k_j), \tag{2.20}
$$

with $C_j = \frac{b_j}{e^{\prime}}$ $\frac{b_j}{s'_{11}(k_j)}$ and $\lambda_{\pm,j} = \lambda_{\pm}(k_j)$.

From the symmetry of the scattering matrix(see (35) in [\[21](#page-22-6)]), we know that k_j^* are also zeros of $s_{22}(k)$. Similarly, we get

$$
\operatorname{Res}_{k=k_j^*} \left[\frac{\mu_{+,2}(x,t,k)}{s_{22}(k,t)} \right] = \frac{\mu_{+,2}(k_j^*)}{s_{22}'(k_j^*)} = -C_j^* e^{i(\lambda_{+,j} + \lambda_{-,j})x} \mu_{-,1}(x,t,k_j^*),\tag{2.21}
$$

2.2 Time evolution

Define the solution of the both parts of the Lax pair $(1.3)(1.4)$ $(1.3)(1.4)$ as

$$
\psi(x,t,k) = \phi_{\pm}(x,t,k)C_{\pm}(k,t),\tag{2.22}
$$

where the function C is independent of x . Taking the derivative with respect to t on both sides gives

$$
(C_{\pm})_t = R_{\pm}C_{\pm}
$$

where $R_{\pm} = \phi_{\pm}^{-1}(T\phi_{\pm} - (\phi_{\pm})_t)$. Since the function C is independent of x, the function R_{\pm} is also independent of x. Therefore we have

$$
R_{\pm} = \lim_{x \to \pm \infty} \phi_{\pm}^{-1} (T \phi_{\pm} - (\phi_{\pm})_t) = i f_{\pm}(k) \sigma_3,
$$
\n(2.23)

where

$$
f_{\pm}(k) = 2(q_{\pm}^2 - 2k^2)\lambda_{\pm}(k).
$$

Thus we can express $(\phi_{\pm})_t$ as

$$
(\phi_{\pm})_t = T\phi_{\pm} - \phi_{\pm}R_{\pm}.
$$
\n(2.24)

Taking the derivative with respect to t of Eq. (3.7) gives

$$
S_t = R_+ S - S R_-,
$$

element-wise reads

$$
s_{11}(k,t) = s_{11}(k,0) \exp\left[i(t+(k) - f_{-}(k))\right],
$$

\n
$$
s_{21}(k,t) = s_{21}(k,0) \exp\left[-it(f_{+}(k) + f_{-}(k))\right],
$$
\n(2.25)

which result in

$$
\rho(k,t) = \rho(k,0) \exp(-2\mathrm{itf}_{+}(k))
$$

and

$$
s'_{11}(k_j, t) = s'_{11}(k_j, 0) \exp[{\rm it(f}_{+}(k_j) - f_{-}(k_j))].
$$

Taking the derivative with respect to t of Eq. (2.19) , we have

$$
b_j(t) = b_j(0) \exp[-it(f_{+}(k_j) + f_{-}(k_j))],
$$

therefore we obtain

$$
C_j(t) = C_j(0) \exp(-2\mathrm{i} f_+(k_j)t).
$$

2.3 Inverse scattering problem

Next, we will establish the RH problem associated to the focusing modified KdV equation [\(1.1\)](#page-1-1) and recover the potential.

2.3.1 Riemann-Hilbert problem

Define a meromorphic function as

$$
M(x;k) = \begin{cases} \left[\frac{\mu_{-,1}}{s_{11}}, \ \mu_{+,2}\right], & k \in \mathbb{C}^+\backslash [0, iq_-],\\ \left[\mu_{+,1}, \ \frac{\mu_{-,2}}{s_{22}}\right], & k \in \mathbb{C}^-\backslash [-iq_-,0]. \end{cases}
$$
(2.26)

Jump on R For $k \in \mathbb{R}$, we write the RH problem as $M_+(x, k) = M_-(x, k)V_0(x, k)$, i.e. $\sqrt{a^2 + b^2}$

$$
\left(\frac{\mu_{-,1}}{s_{11}}, \mu_{+,2}\right) = \left(\mu_{+,1}, \frac{\mu_{-,2}}{s_{22}}\right) V_0, \quad k \in \mathbb{R},\tag{2.27}
$$

where the subscripts \pm denote limiting values from the upper/lower complex plane and

$$
V_0(x,k) = \begin{pmatrix} (1 - \rho \tilde{\rho})e^{i(\lambda - \lambda_+)x} & -\tilde{\rho}e^{-2i\lambda_+ x} \\ \rho e^{2i\lambda_- x} & e^{i(\lambda_- - \lambda_+)x} \end{pmatrix}.
$$
 (2.28)

Jump on the Σ^+ For $k \in \Sigma^+$:= $[0, iq_+]$, the RH problem can be written as $M_+(x, k) = M_-(x, k)V_1(x, k)$, where the subscripts \pm denote limiting values from the right/left complex plane, i.e.

$$
\left(\frac{\mu_{-,1}}{s_{11}}, \mu_{+,2}\right) = \left(\frac{\mu_{-,1}^-}{s_{11}^-}, \mu_{+,2}^-\right) V_1, \quad k \in \Sigma_+^+.
$$
\n(2.29)

Due to the scattering relation (3.7) and the symmetris, we have

$$
\frac{\phi_{-,1}}{s_{11}} = (1 - \rho \tilde{\rho})\phi_{+,1} + \rho \frac{\phi_{-,2}}{s_{22}}
$$
\n
$$
= (1 - \rho \tilde{\rho})\frac{iq_+}{k - \lambda_+}\phi_{+,2} - \rho \frac{(k + \lambda_+)}{iq_+} \frac{\phi_{-,1}^-}{s_{11}^-}
$$
\n(2.30)

and

$$
\phi_{+,2} = \frac{\phi_{-,2}}{s_{22}} - \tilde{\rho}\phi_{+,1}
$$
\n
$$
= \frac{k + \lambda_+}{iq_+} \frac{\phi_{-,1}^-}{s_{11}^-} - \tilde{\rho}\frac{iq_+}{k - \lambda_+}\phi_{+,2}^- \tag{2.31}
$$

which results in

$$
V_1 = \begin{pmatrix} \frac{k+\lambda_+}{iq_+} \rho e^{2i\lambda_- x} & \frac{k+\lambda_+}{iq_+} e^{i(\lambda_- - \lambda_+)x} \\ (1 - \rho \tilde{\rho}) \frac{iq_+}{k-\lambda_+} e^{i(\lambda_- - \lambda_+)x} & -\tilde{\rho} \frac{iq_+}{k-\lambda_+} e^{-2i\lambda_+ x} \end{pmatrix}.
$$
 (2.32)

Jump on the Σ_{+}^{-} For $k \in \Sigma_{+}^{-} := [-iq_{+}, 0]$, the jump can be determined by $M_{+}(x, k) = M_{-}(x, k)V_{2}(x, k)$, where the subscripts \pm denote limiting values from the right/left complex plane, i.e.

$$
\left(\mu_{+,1}, \frac{\mu_{-,2}}{s_{22}}\right) = \left(\mu_{+,1}^-, \frac{\mu_{-,2}^-}{s_{22}^-}\right) V_2, \quad k \in \Sigma_+^-. \tag{2.33}
$$

Due to (3.7) and the symmetris, we have

$$
\phi_{+,1} = \frac{\phi_{-,1}}{s_{11}} - \rho \phi_{+,2}
$$
\n
$$
= \frac{k + \lambda_+}{iq_+} \frac{\phi_{-,2}^2}{s_{22}^2} - \rho \frac{iq_+}{k - \lambda_+} \phi_{+,1}^-
$$
\n(2.34)

and

$$
\frac{\phi_{-,2}}{s_{22}} = \tilde{\rho}\phi_{+,1} + \phi_{+,2}
$$
\n
$$
= \tilde{\rho}\frac{\phi_{-,1}}{s_{11}} + (1 - \rho\tilde{\rho})\phi_{+,2}
$$
\n
$$
= \tilde{\rho}\frac{k + \lambda_+}{iq_+}\frac{\phi_{-,2}^-}{s_{22}^-} + (1 - \rho\tilde{\rho})\frac{iq_+}{k - \lambda_+}\phi_{+,1}^-,\tag{2.35}
$$

then we have

$$
V_2 = \begin{pmatrix} -\rho \frac{iq_+}{k-\lambda_+} e^{2i\lambda_+ x} & (1-\rho \tilde{\rho}) \frac{iq_+}{k-\lambda_+} e^{i(\lambda_+ - \lambda_-)x} \\ \frac{k_+ \lambda_+}{iq_+} e^{i(\lambda_+ - \lambda_-)x} & \tilde{\rho} \frac{k_+ \lambda_+}{iq_+} e^{-2i\lambda_- x} \end{pmatrix}.
$$
 (2.36)

Jump on the Σ_0^+ For $k \in \Sigma_0^+ := [iq_+, iq_-]$, the jump can be determined by $M_+(x, k) = M_-(x, k)V_2(x, k)$, where the subscripts \pm denote limiting values from the right/left complex plane, i.e.

$$
\left(\frac{\mu_{-,1}}{s_{11}}, \mu_{+,2}\right) = \left(\frac{\mu_{-,1}^-}{s_{11}^-}, \mu_{+,2}\right) V_3, \quad k \in \Sigma_0^+, \tag{2.37}
$$

Due to [\(3.7\)](#page-15-2) and the symmetris, we have

$$
\frac{\phi_{-,1}}{s_{11}} = \phi_{+,1} + \rho \phi_{+,2}
$$
\n
$$
= \frac{\phi_{-,2}}{s_{12}} + (\rho - \frac{1}{\tilde{\rho}})\phi_{+,2}
$$
\n
$$
= \frac{\phi_{-,1}}{s_{11}^{-}} + (\rho - \frac{1}{\tilde{\rho}})\phi_{+,2},
$$
\n(2.38)

that is

$$
V_3 = \begin{pmatrix} e^{2i\lambda - x} & 0\\ (\rho - \frac{1}{\tilde{\rho}})e^{i(\lambda - \lambda +)x} & 1 \end{pmatrix}
$$
 (2.39)

Jump on the Σ_0^- For $k \in \Sigma_0^ \overline{0}$:= $[-iq_-, -iq_+]$, the jump can be determined by $M_{+}(x, k) = M_{-}(x, k)V_{4}(x, k)$, where the subscripts \pm denote limiting values from the right/left complex plane, i.e.

$$
\left(\mu_{+,1}, \frac{\mu_{-,2}}{s_{22}}\right) = \left(\mu_{+,1}, \frac{\mu_{-,2}^-}{s_{22}^-}\right) V_4, \quad k \in \Sigma_0^-.
$$
\n(2.40)

Due to [\(3.7\)](#page-15-2) and the symmetris, we have

$$
\frac{\phi_{-,2}}{s_{22}} = \tilde{\rho}\phi_{+,1} + \phi_{+,2}
$$
\n
$$
= (\tilde{\rho} - \frac{1}{\rho})\phi_{+,1} + \frac{\phi_{-,1}}{s_{21}}
$$
\n
$$
= (\tilde{\rho} - \frac{1}{\rho})\phi_{+,1} - \frac{\phi_{-,2}^{-}}{s_{22}^{-}},
$$
\n(2.41)

that is

$$
V_4 = \begin{pmatrix} 1 & (\tilde{\rho} - \frac{1}{\rho})e^{-i(\lambda_- + \lambda_+)x} \\ 0 & -e^{-2i\lambda_- x} \end{pmatrix}.
$$
 (2.42)

Summarize the results, we have

RH problem 1. Find a 2×2 matrix-valued function $M(x; k)$ with the following properties:

- $M(x; \cdot)$: $\mathbb{C}\backslash\Sigma \to \mathbb{C}^{2\times 2}$ is analytic, where $\Sigma = \mathbb{R} \cup \Sigma_-$.
- The limits of $M(x; k)$ as k approaches Σ from the positive and negative side exist and are continuous on Σ , moreover, they satisfy

$$
M_{+}(x;k) = M_{-}(x;k)V(x;k), \quad k \in \Sigma,
$$
\n(2.43)

where

$$
V = \begin{cases} V_0, & k \in \mathbb{R}, \\ V_1, & k \in \Sigma_+^+, \\ V_2, & k \in \Sigma_+^-, \\ V_3, & k \in \Sigma_0^+, \\ V_4, & k \in \Sigma_0^-, \end{cases}
$$

• $M(x; k)$ satisfies the asymptotic(see (53) in [\[21](#page-22-6)])

$$
M(x; k) \to I, \quad |k| \to \infty, k \in \mathbb{C}\backslash\Sigma.
$$

$$
M(x; k) \to \frac{i}{k}\sigma_3 Q_- + O(1), \quad k \to 0.
$$

2.3.2 Reconstruction formula

Using the Plemelj formula, we can express the solution M as follows:

$$
M(x,t,k) = I + \frac{i}{k}\sigma_3 Q_- + \sum_{j=1}^J \frac{1}{k - k_j} \operatorname{Res}_{k=k_j} M^+ + \sum_{j=1}^J \frac{1}{k - k_j^*} \operatorname{Res}_{k=k_j^*} M^- + \frac{1}{2\pi i} \int_{\Sigma} \frac{[M^-(V-I)](x,t,\zeta)}{\zeta - k} d\zeta, \quad k \in \mathbb{C}\setminus\Sigma,
$$
\n(2.44)

By a standard argument, we can recover the potential $q(x, t)$

$$
q(x,t) = -i \lim_{k \to \infty} k M_{12}(x,t,k).
$$
 (2.45)

Substituting [\(2.44\)](#page-12-1) into it, we obtain

$$
q(x,t) = q_{-}(t) - i \sum_{j=1}^{J} C_j^*(t) e^{i(\lambda_{+,j}^* + \lambda_{-,j}^*)x} M_{11}(x,t,k_j^*) + \frac{1}{2\pi} \int_{\Sigma} \left[M^-(V-I) \right]_{12} dk
$$
\n(2.46)

Remark 2.1 When $q_ - = q_ + = q_0$, the reconstruction formula [\(2.46\)](#page-13-2) matches Eq. (65) in [\[21](#page-22-6)].

3 The defocusing mKdV equation

In this section, we consider the defocusing case, i.e. $\sigma = 1$. First, let us establish and define some notations that will be used throughout this section.

$$
\Sigma_{\pm} = (-\infty, -q_{\pm}] \cup [q_{\pm}, \infty),
$$

\n
$$
\Sigma_{0} = [-q_{-}, -q_{+}] \cup [q_{+}, q_{-}],
$$

\n
$$
\dot{\Sigma}_{\pm} = (-\infty, -q_{\pm}) \cup (q_{\pm}, \infty),
$$

\n
$$
\dot{\Sigma}_{0} = (-q_{-}, -q_{+}) \cup (q_{+}, q_{-}).
$$

Fig. 2. The branch cuts $\Sigma_-\$ and Σ_+ to the defocusing case

3.1 Direct scattering problem

Next, we will develop the spectral analysis and introduce the scattering matrix to prepare for the set-up of the RH problem.

3.1.1 Spectral analysis

To obtain the Jost functions of the spectral problem [\(1.3\)](#page-2-0), we need to diagonalize the asymptotic matrix $X_{\pm} = ik\sigma_3 + Q_{\pm}$. By calculation, the eigenvalues of the matrix X_{\pm} are $\pm i\lambda_{\pm}$, where

$$
\lambda_{\pm}^2 = k^2 - q_{\pm}^2.
$$

To avoid the multivalues of the function λ_{\pm} , we take the branch cut for λ_{\pm} on Σ_{\pm} . For all $k \in \mathbb{C}\setminus\Sigma_{\pm}$, we define λ_{\pm} as the single-valued, analytic and continuous as k approaches Σ_{\pm} from above. For $k \in \Sigma_{\pm}$, we take the positive value of the real square root for λ_{\pm} .

The eigenvector matrix is

$$
E_{\pm} = I + \frac{i}{k - \lambda_{\pm}} \sigma_3 Q_{\pm}
$$

corresponding to $\mp i\lambda_{\pm}$ and det $E_{\pm} = -\frac{2\lambda_{\pm}}{k-\lambda_{\pm}}$ $\frac{2\lambda_{\pm}}{k-\lambda_{\pm}} \triangleq \gamma_{\pm}(k).$

After diagonalizition, we have

$$
\phi_{\pm} \sim E_{\pm} e^{-i\lambda_{\pm} x \sigma_3}, \quad x \to \pm \infty. \tag{3.1}
$$

Define the normalized Jost functions μ_{\pm} as

$$
\mu_{\pm} = \phi_{\pm} e^{i\lambda_{\pm} x \sigma_3},\tag{3.2}
$$

then we have

$$
\mu_{\pm} \to E_{\pm}, \quad x \to \pm \infty.
$$

And they satisfy the Volterra integral equation

$$
\mu_{\pm} = E_{\pm}(k) + \int_{\pm\infty}^{x} E_{\pm}(k) e^{-i\lambda_{\pm}(x-y)\hat{\sigma}_{3}} \left[E_{\pm}^{-1}(k) \Delta Q_{\pm}(y) \mu_{\pm}(y,k) \right] dy \tag{3.3}
$$

with $\Delta Q_{\pm} = Q - Q_{\pm}$.

Lemma 3.1. If $(1+|x|)(q-q_{\pm}) \in L^1(\mathbb{R}_{\pm})$, then

• Analyticity: the function $\mu_{-1}(x,t,k)$ is well-defined in Σ_- and analytic in $\mathbb{C}\setminus\Sigma_+$, whereas the function $\mu_{+,2}(x,t,k)$ is well-defined in Σ_+ and analytic in $\mathbb{C}\backslash\Sigma_{+}.$

• Symmetry: Let $\widetilde{\phi}_{\pm} = \phi_{\pm}(x, t, -\lambda_{\pm}(k))$, we have

$$
\phi_{\pm} = \tilde{\phi}_{\pm} \frac{i}{k - \lambda_{\pm}} \sigma_3 Q_{\pm}, \quad k \in \mathring{\Sigma}_{\pm}, \tag{3.4}
$$

Column-wise reads

$$
\phi_{\pm,1} = \frac{-iq_{\pm}}{k - \lambda_{\pm}} \widetilde{\phi}_{\pm,2}, \quad k \in \mathring{\Sigma}_{\pm}, \tag{3.5}
$$

$$
\phi_{\pm,2} = \frac{i q_{\pm}}{k - \lambda_{\pm}} \widetilde{\phi}_{\pm,1}, \quad k \in \mathring{\Sigma}_{\pm}.
$$
 (3.6)

3.1.2 Scattering matrix

There exists a scattering matrix $S(k, t)$ such that

$$
\phi_{-}(x,t,k) = \phi_{+}(x,t,k)S(k,t), \quad k \in \Sigma_{-}.
$$
\n(3.7)

Thus, we have det $S = \frac{\gamma}{\gamma}$ $\frac{\gamma_-}{\gamma_+}$. And the scattering coefficients can be written as the Wronskians expressions

$$
s_{11} = \frac{\text{Wr}(\phi_{-,1}, \phi_{+,2})}{\gamma_+}, \quad s_{12} = \frac{\text{Wr}(\phi_{-,2}, \phi_{+,2})}{\gamma_+}, s_{21} = \frac{\text{Wr}(\phi_{+,1}, \phi_{-,1})}{\gamma_+}, \quad s_{12} = \frac{\text{Wr}(\phi_{+,1}, \phi_{-,2})}{\gamma_+}.
$$
(3.8)

Define

$$
\rho = \frac{s_{21}}{s_{11}}, \quad \tilde{\rho} = \frac{s_{12}}{s_{22}}.
$$
\n(3.9)

Lemma 3.2. Let $s_{11}^{-}(k) = \lim_{\epsilon \uparrow 0} s_{11}(k + i\epsilon, t)$, then we have

$$
s_{11}^{-}(k) = s_{22} \frac{(k + \lambda_{+})(k - \lambda_{-})}{q_{+}q_{-}}, \quad k \in \Sigma_{-},
$$

\n
$$
s_{11}^{-}(k) = s_{21}(k) \frac{k + \lambda_{+}}{-iq_{+}}, \quad k \in \Sigma_{0}.
$$
\n
$$
(3.10)
$$

3.1.3 Discrete spectrum and residue condition

Assume that there exists no spectral singularities. We can prove that the discrete eigenvalues don't exist in the interval Σ_0 , also can not be the branch points $\pm q_{\pm}$ (see more details in [\[26\]](#page-23-0)). And we assume $s_{11}(k, t)$ has J simple zeros in $(-q_+, q_+)$ denoted by k_j , $j = 1, 2, ..., J$. From the Wronskians expression of $s_{11}(k, t)$, there exists a constant only depending on t such that

$$
\phi_{-,1}(x,t,k_j) = b_j(t)\phi_{+,2}(x,t,k_j).
$$

On account of (3.2) , we have

$$
\mu_{-,1}(x,t,k_j) = b_j(t)\mu_{+,2(x,t,k_j)}e^{i(\lambda_{+,j}+\lambda_{-,j})x},
$$

where $\lambda_{\pm,j} = \lambda_{\pm}(k_j)$. Thus for $j = 1, 2, ..., J$, we get

$$
\operatorname{Res}_{k=k_j} \left[\frac{\mu_{-,1}(x,t,k)}{s_{11}(k,t)} \right] = \frac{\mu_{-,1}(k_j)}{s'_{11}(k_j)} = C_j(t) e^{i(\lambda_{+,j} + \lambda_{-,j})x} \mu_{+,2}(x,t,k_j), \tag{3.11}
$$

with $C_j = \frac{b_j}{e^{\prime}}$ $\frac{b_j}{s'_{11}(k_j)}$. Therefore, we have

$$
\underset{k=k_j}{\text{Res}} M(x,t,k) = \left(C_j e^{i(\lambda_{+,j} + \lambda_{-,j})x} M_2(k_j) , 0 \right). \tag{3.12}
$$

3.2 Time evolution

Performing a similar maniputation of the time evolution [2.2](#page-8-0) in the focusing case, we obtain

$$
f_{\pm}(k) = -2(q_{\pm}^2 + 2k^2)\lambda_{\pm}(k).
$$

and

$$
s_{11}(k,t) = s_{11}(k,0) \exp\left[\text{it}(f_{+}(k) - f_{-}(k))\right],
$$

\n
$$
s_{21}(k,t) = s_{21}(k,0) \exp\left[-\text{it}(f_{+}(k) + f_{-}(k))\right],
$$

\n
$$
\rho(k,t) = \rho(k,0) \exp(-2\text{it}f_{+}(k)),
$$

\n
$$
s'_{11}(k_{j},t) = s'_{11}(k_{j},0) \exp\left[\text{it}(f_{+}(k_{j}) - f_{-}(k_{j}))\right],
$$

\n
$$
b_{j}(t) = b_{j}(0) \exp\left[-\text{it}(f_{+}(k_{j}) + f_{-}(k_{j}))\right].
$$

\n(3.13)

Therefore we obtain

$$
C_j(t) = C_j(0) \exp(-2\mathrm{i} f_+(k_j)t).
$$

3.3 Inverse scattering problem

Next, we will establish the RH problem associated to the defocusing modified KdV equation [\(1.1\)](#page-1-1) and recover the potential.

3.3.1 Riemann-Hilbert problem

Define a meromorphic function as

$$
M(x,t,k) = \left(\frac{\mu_{-,1}}{s_{11}}, \ \mu_{+,2}\right), \quad k \in \Sigma_+.
$$
 (3.14)

Jump on the Σ _− Using the scattering relation [\(3.7\)](#page-15-2), the second symmetry [\(3.5\)](#page-15-3) (3.6) of the Jost function and (3.10) of the scattering coefficients, we have

$$
\frac{\phi_{-,1}}{s_{11}} = (1 - \rho \tilde{\rho})\phi_{+,1} + \rho \frac{\phi_{-,2}}{s_{22}} \n= (1 - \rho \tilde{\rho})\frac{-iq_+}{k - \lambda_+}\tilde{\phi}_{+,2} + \rho \frac{(k + \lambda_+)(k - \lambda_-)}{q_+q_-} \frac{iq_-}{k - \lambda_-} \frac{\tilde{\phi}_{-,1}}{s_{11}^-} \n= (1 - \rho \tilde{\rho})\frac{-iq_+}{k - \lambda_+}\tilde{\phi}_{+,2} + \rho \frac{i(k + \lambda_+)}{q_+} \frac{\tilde{\phi}_{-,1}}{s_{11}^-}
$$
\n(3.15)

and

$$
\phi_{+,2} = \frac{\phi_{-,2}}{s_{22}} - \tilde{\rho}\phi_{+,1}
$$
\n
$$
= \frac{(k+\lambda_{+})(k-\lambda_{-})}{q_{+}q_{-}} \frac{iq_{-}}{k-\lambda_{-}} \frac{\tilde{\phi}_{-,1}}{s_{11}^{-}} - \tilde{\rho}\frac{-iq_{+}}{k-\lambda_{+}} \tilde{\phi}_{+,2}
$$
\n
$$
= \frac{i(k+\lambda_{-})}{q_{+}} \frac{\tilde{\phi}_{-,1}}{s_{11}^{-}} + \tilde{\rho}\frac{iq_{+}}{k-\lambda_{+}} \tilde{\phi}_{+,2}.
$$
\n(3.16)

Combining (3.15) and (3.16) leads to

$$
\left(\frac{\phi_{-,1}}{s_{11}}, \phi_{+,2}\right) = \left(\frac{\phi_{-,1}^-}{s_{11}^-}, \phi_{+,2}^-\right) \left(\begin{array}{cc} \rho \frac{i(k+\lambda_+)}{q_+} & \frac{i(k+\lambda_+)}{q_+} \\ (1-\rho\tilde{\rho})\frac{-iq_+}{k-\lambda_+} & \tilde{\rho}\frac{iq_+}{k-\lambda_+} \end{array}\right), \quad k \in \Sigma_-. \quad (3.17)
$$

Jump on the Σ_0 According to [\(3.10\)](#page-15-5), we get

$$
\frac{\phi_{-,1}}{s_{11}} = \frac{\phi_{-,1}}{s_{11}^{-}} \frac{s_{11}^{-}}{s_{11}} = \rho \frac{k + \lambda_{+}}{-i q_{+}} \frac{\phi_{-,1}}{s_{11}^{-}}.
$$
\n(3.18)

Again, using the scattering relation (3.7) and (3.10) , we have

$$
\phi_{+,2} = \frac{1}{s_{21}} \phi_{-,1} - \frac{1}{\rho} \phi_{+,1}
$$
\n
$$
= \frac{k + \lambda_+}{-iq_+} \frac{\phi_{-,1}}{s_{11}^-} - \frac{1}{\rho} \frac{-iq_+}{k - \lambda_+} \widetilde{\phi}_{+,2}.
$$
\n(3.19)

Combining (3.18) and (3.19) leads to

$$
\left(\frac{\phi_{-,1}}{s_{11}}, \phi_{+,2}\right) = \left(\frac{\phi_{-,1}}{s_{11}^-}, \phi_{+,2}^-\right) \left(\begin{array}{cc} \rho \frac{(k+\lambda_+)}{-iq_+} & \frac{(k+\lambda_+)}{-iq_+} \\ 0 & \frac{1}{\rho} \frac{iq_+}{k-\lambda_+} \end{array}\right), \quad k \in \Sigma_0.
$$
 (3.20)

Note the relation for μ_\pm and ϕ_\pm and calculate direct, we arrive at

$$
\left(\frac{\mu_{-,1}}{s_{11}}, \mu_{+,2}\right) = \begin{cases} \left(\frac{\mu_{-,1}^-}{s_{11}^-}, \mu_{+,2}^-\right) V_{\Sigma_-}, & k \in \Sigma_-\\ \left(\frac{\mu_{-,1}}{s_{11}^-}, \mu_{+,2}^-\right) V_{\Sigma_0}, & k \in \Sigma_0 \end{cases}
$$
(3.21)

where

$$
V_{\Sigma-} = (E_{+} - I) \begin{pmatrix} e^{-i\lambda_{+}x} & e^{i\lambda_{-}x} \end{pmatrix} \begin{pmatrix} 1 - \rho\tilde{\rho} & -\tilde{\rho} \\ \rho & 1 \end{pmatrix} \begin{pmatrix} e^{i\lambda_{-}x} & e^{-i\lambda_{+}x} \end{pmatrix},
$$
\n
$$
V_{\Sigma_{0}} = (E_{+} - I) \begin{pmatrix} e^{-i\lambda_{+}x} & e^{-i\lambda_{-}x} \end{pmatrix} \begin{pmatrix} 0 & \frac{1}{\rho} \\ \rho & -1 \end{pmatrix} \begin{pmatrix} e^{i\lambda_{-}x} & e^{-i\lambda_{+}x} \end{pmatrix}.
$$

Therefore, we obtain a jump condition

$$
M^{+} = M^{-}(E_{+} - I)(I - V_{0}), \quad k \in \Sigma_{+}, \tag{3.22}
$$

with

$$
V_0 = \begin{cases} \begin{pmatrix} 1 - (1 - \rho \tilde{\rho})e^{i(\lambda_- - \lambda_+)x} & \tilde{\rho} e^{-2i\lambda_+} \\ \rho e^{2i\lambda_-} & 1 - e^{-i(\lambda_- - \lambda_+)x} \end{pmatrix}, & k \in \Sigma_- \\ \begin{pmatrix} 1 & \frac{1}{\rho} e^{-2i\lambda_+} \\ -\rho & 1 - e^{-i(\lambda_+ + \lambda_-)x} \end{pmatrix}, & k \in \Sigma_0 \end{cases}
$$

Due to the different asymptotics as $k \to \infty$ between $k \in \mathbb{C}^+$ and $k \in \mathbb{C}^-$

$$
\lambda_{\pm} = \begin{cases}\nk - \frac{q_{\pm}^2}{2k} + o\left(\frac{1}{k}\right), & k \to \infty, \quad \text{Imk} > 0, \\
-k + \frac{q_{\pm}^2}{2k} + o\left(\frac{1}{k}\right), & k \to \infty, \quad \text{Imk} < 0,\n\end{cases}
$$
\n(3.23)

we obtain the asymptotic of ${\cal M}(x,t,k)$

$$
M(x,t,k) = \begin{cases} I + O(\frac{1}{k}), & k \to \infty, \quad \text{Imk} > 0, \\ \frac{i}{k - \lambda_+} \sigma_3 Q_+ + O(1), & k \to \infty, \quad \text{Imk} < 0. \end{cases}
$$
 (3.24)

which can be rewritten as

$$
M(x, t, k) = E_+ + O(1), \quad k \to \infty.
$$

To deal with the different limit, we introduce a new function ${\cal N}(x, t, k)$

$$
M(x, t, k) = N(x, t, k)E_{+}(k, t),
$$
\n(3.25)

thus we have

$$
N(x, t, k) = I + O(\frac{1}{k}).
$$

On account of the jump [\(3.22\)](#page-18-0), we have

$$
N^{+} = N^{-} \tilde{V}, \quad k \in \Sigma_{+}, \tag{3.26}
$$

where $\tilde{V} = E_+^- V E_+^{-1}$.

3.3.2 Reconstruction formula

Using the Plemelj formula, we can express the solution M as follows:

$$
N(x,t,k) = I + \sum_{j=1}^{J} \frac{1}{k - k_j} \operatorname{Res}_{k=k_j} N(x,t,k)
$$

$$
- \frac{1}{2\pi i} \int_{\Sigma_+} \frac{[N^-(I - \tilde{V})](x,t,\zeta)}{\zeta - k} d\zeta, \quad k \in \mathbb{C} \setminus \Sigma_+,
$$
 (3.27)

from (3.25) , we obtain

$$
M(x,t,k) = E_{+}(k,t) + \sum_{j=1}^{J} \frac{1}{k - k_j} \operatorname{Res}_{k=k_j} M(x,t,k) E_{+}^{-1}(k_j) E_{+}(k,t)
$$

$$
- \frac{E_{+}(k,t)}{2\pi i} \int_{\Sigma_{+}} \frac{[M^{-(}E_{+} - I)V_0 E_{+}^{-1}](x,t,\zeta)}{\zeta - k} d\zeta, \quad k \in \mathbb{C}\setminus\Sigma_{+},
$$
\n(3.28)

By substituting the residue conditions, we have

$$
M(x,t,k) = E_{+}(k,t) + \sum_{j=1}^{J} C_{j}(t)e^{i(\lambda_{+,j} + \lambda_{-,j})x} (M_{2}(k_{j}), 0) E_{+}^{-1}(k_{j}, t)E_{+}(k, t) + \frac{1}{2\pi i k} \int_{\Sigma_{+}} [M^{-(}E_{+} - I)V_{0}E_{+}^{-1}](x, t, \zeta) d\zeta + O\left(\frac{1}{k^{2}}\right), k \to \infty, \quad \text{Im}k > 0.
$$
\n(3.29)

By a standard argument, we can recover the potential $q(x, t)$

$$
q(x,t) = -i \lim_{\substack{k \to \infty \\ \text{Im}k > 0}} k M_{12}(x,t,k),
$$
\n(3.30)

by substituting (3.29) into it, we obtain

$$
q(x,t) = q_{+}(t) \left(1 - \sum_{j=1}^{J} C_{j}(t) e^{i(\lambda_{+,j} + \lambda_{-,j})x} M_{12}(x,t,k_{j}) \right) + \frac{1}{2\pi} \int_{\Sigma_{+}} \frac{1}{\lambda_{+}(k)} \left[\left(\frac{iq_{+}}{k + \lambda_{+}} V_{0,11} - V_{0,12} \right) q_{+}(t) M_{12}^{-}(x,t,k) \right) - \left(\frac{iq_{+}}{k + \lambda_{+}} V_{0,21} + V_{0,22} \right) q_{+}(t) M_{11}^{-}(x,t,k) \right] dk
$$
\n(3.31)

Remark 3.1 When $q_ - = q_ + = q_0$, the reconstruction formula [\(2.46\)](#page-13-2) matches Eq. (128) in [\[21\]](#page-22-6).

Acknowledgements

This work is supported by the National Science Foundation of China (Grant No.12371247).

Data Availability Statements

The data that supports the findings of this study are available within the article.

Conflict of Interest

The authors have no conflicts to disclose.

References

- [1] M. J. Ablowitz, H. Segur. Solitons and the inverse scattering transform. Society for Industrial and Applied Mathematics, 1981.
- [2] M. J. Ablowitz, D. J. Kaup, A. C. Newell, H. Segur. Nonlinear-evolution equations of physical significance. Physical Review Letters 31.2 (1973): 125.
- [3] R. M. Miura. Korteweg-de Vries equation and generalizations. I. A remarkable explicit nonlinear transformation. Journal of Mathematical Physics 9.8 (1968): 1202-1204.
- [4] N. J, Zabusky. A synergetic approach to problems of nonlinear dispersive wave propagation and interaction. Nonlinear partial differential equations. Academic Press, 1967. 223-258.
- [5] H. Ono. Soliton fission in anharmonic lattices with reflectionless inhomogeneity. Journal of the Physical Society of Japan 61.12 (1992): 4336-4343.
- [6] T. Kakutani, H. Ono. Weak non-linear hydromagnetic waves in a cold collisionfree plasma. Journal of the physical society of Japan 26.5 (1969): 1305-1318.
- [7] A. H. Khater, O. H. El-Kalaawy, D. K. Callebaut. Bäcklund transformations and exact solutions for Alfvén solitons in a relativistic electron?positron plasma. Physica Scripta 58.6 (1998): 545.
- [8] T. S. Komatsu, S. Sasa. Kink soliton characterizing traffic congestion. Physical Review E 52.5 (1995): 5574.
- [9] H. X. Ge, S. Q. Dai, Y. Xue, L. Y. Dong. Stabilization analysis and modified Korteweg?de Vries equation in a cooperative driving system. Physical Review E 71.6 (2005): 066119.
- [10] E. A. Ralph, L. Pratt. Predicting eddy detachment for an equivalent barotropic thin jet. Journal of Nonlinear Science 4 (1994): 355-374.
- [11] A. A. Himonas, F. Yan. A higher dispersion KdV equation on the half-line. Journal of Differential Equations 333 (2022): 55-102.
- [12] N. Liu, B. L. Guo. Painlevé-type asymptotics of an extended modified KdV equation in transition regions. Journal of Differential Equations 280 (2021): 203-235.
- [13] T. Y. Xu, Z. C. Zhang, E. G. Fan. On the Cauchy problem of defocusing mKdV equation with finite density initial data: Long time asymptotics in soliton-less regions. Journal of Differential Equations 372 (2023): 55-122.
- [14] T. Y Xu, E. G. Fan. Large-time asymptotics to the focusing nonlocal modified Kortweg-de Vries equation with step-like boundary conditions. Studies in Applied Mathematics 150.4 (2023): 1217-1273.
- [15] L. M. Ling, X. Sun. Stability of elliptic function solutions for the focusing modified KdV equation. Advances in Mathematics 435 (2023): 109356.
- [16] F. Demontis. Exact solutions of the modified Korteweg-de Vries equation. Theoretical and Mathematical Physics 168.1 (2011): 886-897.
- [17] M. Wadati. The modified Korteweg-de Vries equation. Journal of the Physical Society of Japan 34.5 (1973): 1289-1296.
- [18] T. Au-Yeung, P. Fung, C. Au. Modified KdV solitons with non-zero vacuum parameter obtainable from the ZS-AKNS inverse method. Journal of Physics A: Mathematical and General 17.7 (1984): 1425.
- [19] T. Au-Yeung, P. Fung. Hamiltonian formulation of the inverse scattering method of the modified KdV equation under the non-vanishing boundary condition u (x, t) to b as x to $+$ or -infinity. Journal of Physics A: Mathematical and General 21.18 (1988): 3575.
- [20] M. A. Alejo. Focusing mKdV breather solutions with nonvanishing boundary condition by the inverse scattering method. Journal of Nonlinear Mathematical Physics 19.1 (2012): 119-135.
- [21] G. Q. Zhang, Z. Y. Yan. Focusing and defocusing mKdV equations with nonzero boundary conditions: Inverse scattering transforms and soliton interactions. Physica D: Nonlinear Phenomena 410 (2020): 132521.
- [22] J. S. He, S. R. Chen. Hamiltonian formalism of mKdV equation with nonvanishing boundary values. Communications in Theoretical Physics 44.2 (2005): 321.
- [23] P. Deift, X. Zhou. A steepest descent method for oscillatory Riemann–Hilbert problems. Asymptotics for the MKdV equation. Annals of Mathematics 137.2 (1993): 295-368.
- [24] D. E. Baldwin. Dispersive shock wave interactions and two-dimensional oceanwave soliton interactions. Diss. Ph. D. thesis, University of Colorado, 2013.
- [25] P. Germain, F. Pusateri, F. Rousset. Asymptotic stability of solitons for mKdV. Advances in Mathematics 299 (2016): 272-330.

[26] C. Biondini, E. Fagerstrom, B. Prinari. Inverse scattering transform for the defocusing nonlinear Schrödinger equation with fully asymmetric non-zero boundary conditions. Physica D: Nonlinear Phenomena 333 (2016): 117-136.