

# Jupiter Mass Binaries and Cosmic Ray Viscosity

J. I. Katz

Department of Physics and McDonnell Center for the Space Sciences, Washington  
University, St. Louis, Mo. 63130

[katz@wuphys.wustl.edu](mailto:katz@wuphys.wustl.edu)

Received \_\_\_\_\_; accepted \_\_\_\_\_

arXiv:2312.10834v2 [astro-ph.GA] 15 Feb 2024

## ABSTRACT

The fraction of planetary mass objects in the Trapezium cluster that are in wide binaries is much greater than implied by extrapolation to lower masses of the fraction of stars that are wide binaries. Wide binaries may be produced by gravitational collapse of a medium with fluid vorticity. In a uniform medium with uniform vorticity the collapse criterion is independent of the size and mass of the collapsing region, which would imply a wide binary fraction independent of mass, in contradiction to observation. Angular momentum, rather than thermal pressure, may be the chief obstacle to star formation. The excess of Jupiter Mass Binary Objects in the Trapezium cluster may be attributed to cosmic ray viscosity that transports angular momentum to surrounding material. Viscosity is more effective in smaller and less massive collapsing regions, preferentially producing planetary mass wide binaries. Energetic particles trapped on magnetic field lines in a corona contribute to viscosity, dissipation and angular momentum flow in accretion discs.

*Subject headings:* Planet formation, binaries, cosmic rays

## 1. Introduction

Pearson & McCaughrean (2023) discovered that an unexpectedly large fraction of Jupiter mass objects in the Trapezium are in wide (projected separation  $\gtrsim 100$  AU) binaries. At the lowest masses ( $M \lesssim 0.002M_{\odot}$ ) this fraction is  $9 \pm 3\%$ . Below about  $0.1M_{\odot}$  the fraction is an increasing function of decreasing mass, while above  $0.1M_{\odot}$  it is an decreasing function of decreasing mass; at the minimum around  $0.1M_{\odot}$  the wide binary fraction is  $\lesssim 2\%$ . The change in sign of the dependence on mass around  $0.1M_{\odot}$  indicates a characteristic mass scale of wide binary formation, at least in this star- and planet-forming region.

Star and planet formation in the Trapezium cluster is comparatively recent. These dynamically fragile planetary mass wide binaries with escape velocities  $\mathcal{O}(100 \text{ m/s})$  have been little affected by gravitational interaction with other stellar or planetary objects. Their present-day statistics of orbital parameters provides information about their processes of formation, information that has been lost from the statistics of wide field binaries.

Formation from three unbound slow-moving single objects is unlikely because it is rare that three unbound objects would have such low relative velocities and be close enough (within approximately the final binary separation) for two to be captured with expulsion of the third. That process would also preferentially produce more massive binaries, in contradiction to observation, because such three-body interactions preferentially expel the lowest mass object. Because of their dynamical fragility, wide binary planetary mass objects cannot be explained by expulsion from more tightly bound multiple systems.

The theory of the formation of wide binaries remains controversial and uncertain, and presents different issues than the formation of close binaries from the late-stage collapse of a rotating cloud (Tohline 2002; Offner *et al.* 2023). Their formation is attributed to the collapse of a subregion of the interstellar cloud to a density comparable to the density

of the binary’s mass spread over its orbital volume, at which point angular momentum arrests further collapse on a dynamical time scale. Unlike field objects (Offner *et al.* 2023; Fontanive *et al.* 2023), the Trapezium wide binaries are a homogeneous population with recent coeval formation, perhaps from a nearly uniform medium, and likely have been unaffected by subsequent interactions. Their binding energies and masses are orders of magnitude below those discussed by Jumper & Fisher (2013) as the result of turbulent fragmentation. Because of the complexity of the problem and the heterogeneity of the interstellar medium, I consider scaling relations, with the objective of explaining the observed (Pearson & McCaughrean 2023) increase of the wide binary fraction toward planetary masses.

## 2. Angular Momentum

Angular momentum is a critical obstacle to collapse of a gas cloud (Katz 1987). Thermal energy can be radiated away, rapidly at high density, and magnetic energy turned into thermal energy by resistivity, but angular momentum cannot be. The angular momentum of a collapsing cloud almost entirely appears as the orbital angular momentum of the binary. Rotation of the final hydrostatic objects can only take up a tiny fraction ( $\mathcal{O}(10^{-3})$  in the case of binary Jupiters at 100 AU separation) of the orbital angular momentum. Here we are concerned with the conditions under which a subregion of an extended cloud can begin to collapse and the viscous transport of some of its angular momentum to its surroundings that enables that collapse.

If a gas cloud collapses to form a single star or planetary mass object, it is not possible to infer the pre-collapse conditions in the cloud from the properties of the condensed object; the mass is known, but the pre-collapse length scale, density and angular momentum are not. Because the moment of inertia of a single star or planet is orders of magnitude less

than that of its much larger pre-collapse gas cloud, nearly all of the pre-collapse angular momentum must have been lost.

The pre-collapse angular momentum of a newly formed binary appears as orbital angular momentum. The mean density of the binary’s orbital volume  $\langle \rho \rangle \equiv M/(4\pi R^3/3)$ , where  $M$  is its total mass and  $R$  the binary separation, provides an estimate of the pre-collapse density. This follows from the assumption that the cloud’s angular momentum does not stop collapse until it reaches this density, at which point the cloud fragments into proto-stars or proto-planets whose relative motion takes up the angular momentum. If there are more than two fragments, those of the lowest mass are generally expelled gravitationally, leaving a surviving binary (Pearson & McCaughrean (2023) report 40 binaries but only two triples).

These wide binaries are dynamically fragile, readily disrupted by stellar encounters, and the observed binary fraction likely underestimates the fraction at birth. For a binary of mass  $0.002M_\odot$  ( $2M_{Jupiter}$ ) and separation 100 AU, like those discovered by Pearson & McCaughrean (2023), the pre-collapse density of molecular hydrogen  $n \sim 10^9 \text{ cm}^{-3}$ .

For a cloud with uniform vorticity  $\vec{\omega} \equiv \vec{\nabla} \times \vec{v}$ , where  $\vec{v}$  is the fluid velocity, the velocity difference  $\Delta v_\omega \equiv |\vec{\omega}|R$  over a length  $R$  may be compared to the velocity of gravitational collapse  $\Delta v_G \equiv \sqrt{GM/R} = \sqrt{G4\pi\langle\rho\rangle R^2/3}$ . The necessary condition for collapse of a subregion within a larger homogeneous cloud  $\Delta v_\omega < \Delta v_G$  becomes

$$|\vec{\omega}| < \sqrt{\frac{4\pi G\langle\rho\rangle}{3}}. \quad (1)$$

This could be estimated on dimensional grounds, because the express on the right hand side is the only form containing  $G$ ,  $M$ ,  $R$  and  $\rho$  with the dimensions ( $\text{s}^{-1}$ ) of vorticity.

Collapse, and formation of a wide binary whose orbit contains the angular momentum

of the collapsing cloud, depends only on the comparison of  $|\vec{\omega}|$  to  $\langle\rho\rangle$  in (1). These parameters, averaged over a length scale containing a planetary or stellar mass of gas, may be approximately uniform through a much larger and more massive cloud comprising the entire star- and planet-forming region. The temperature, set by cosmic ray heating and thermal emission by grains, is confined to a narrow range, and on these length scales pressure equilibration is rapid and implies near-uniform density. This is supported by the observations of Pearson & McCaughrean (2023) of binaries with similar mass densities over the extended region they observed.

Conservation of specific angular momentum  $\propto VR \propto |\vec{\omega}|R^2$  implies that as a cloud collapses  $|\vec{\omega}| \propto R^{-2}$ . The characteristic gravitational frequency  $\sqrt{4\pi G\langle\rho\rangle/3} \propto R^{-3/2}$ . At the beginning of collapse the inequality (1) must be satisfied, but as collapse proceeds  $R$  becomes small enough that (1) is no longer satisfied, and angular momentum stops collapse. The likely result is formation of a flattened, disc-like, structure (Offner *et al.* 2023). Because it is self-gravitating, it separates into two or more denser structures that each collapse into proto-planets or proto-stars. If there are more than two such objects all but two are expelled, and the remaining two, usually the most massive two, become the components of a wide binary. Tohline (2002) and Offner *et al.* (2023) discuss complications of this picture, but these complications do not affect the control of collapse by angular momentum and the fact that collapse must stop when the inequality (1) is no longer satisfied.

For the general interstellar medium the rotation of the Galaxy contributes  $|\vec{\omega}| \sim 10^{-15} \text{ s}^{-1}$ . For a hydrogen density  $n \sim 1 \text{ cm}^{-3}$  the collapse condition is satisfied by a factor of 5–10. Comparing the increase  $\propto R^{-2}$  on the left hand side of Eq. 1 to that  $\propto R^{-3/2}$  on the right hand side shows that after a decrease in length scale by the square of this factor, and an increase in density by its sixth power, angular momentum becomes sufficient to stop collapse. At least qualitatively, the high power of  $R$  may explain the high

inferred density ( $n \sim 10^9 \text{ cm}^{-3}$ ) of the subregions forming planetary mass binary objects in the Trapezium cluster. In fact, most of the interstellar medium likely has  $|\vec{\omega}|$  greater than that of the general Galactic rotation because the medium is stirred by supernova shocks, the gravitational fields of spiral arms and massive clouds, expansion of regions warmed by photoionization and contraction of those cooled by radiation, and other complex hydrodynamics.

### 3. Collapse

The collapse inequality (1), unlike the Jeans criterion for collapse of clouds without angular momentum (Binney & Tremaine 2008), does not depend on the mass or size of the cloud, and predicts no dependence of the wide binary fraction on the mass of the binary formed. It also applies to clouds of arbitrary size; if the condition is satisfied in a homogeneous cloud, a large number of wide binaries will be formed, while if not satisfied no collapse can occur. Collapse and planetary or star formation occurs when both criteria are satisfied; the Jeans criterion is satisfied for Jupiter-mass regions when  $n \sim 10^9 \text{ cm}^{-3}$ , consistent with the pre-collapse density inferred from the present mean density of these binaries.

In a cloud of uniform density and vorticity the collapse condition (1) is satisfied either everywhere or nowhere. But no real astronomical gas cloud is uniform. The vorticity varies in this complex turbulent flow. A general cloud may comprise regions in which this condition is satisfied and those in which it is not. The density increases as a result of compressive external pressure, self-gravity, and the radiative loss of internal energy, and even subregions not initially satisfying the inequality (1) may come to do so. The distribution of masses of these subregions determines the masses of the objects formed. If there is no natural (*a priori*) scale on which  $|\vec{\omega}|$  or  $\rho$  vary, then the fraction of wide binaries

must be a power law function of the stellar or planetary mass, the only function without a characteristic scale. This is not consistent with the change in slope of the observed wide binary fraction (Pearson & McCaughrean 2023) around  $0.1M_{\odot}$ . That change in slope implies the existence of a characteristic spatial scale in the cloud.

Collapse begins when (1) is first satisfied; the total mass of the objects formed is nearly the mass of the gas in the collapsing region. Because collapse is limited by angular momentum, the resulting condensed objects form a binary with separation comparable to the size of the collapsing subregion when it first satisfies Eq. 1. Subsequent gravitational interactions with other objects disrupt most of these wide binaries, and lower mass binaries are particularly susceptible to disruption because of their weak binding. Disruption is a random process and in a very young cloud with active star formation like the Trapezium cluster many wide binaries are still present, despite the high density of potentially disrupting stellar or planetary mass objects.

#### 4. Vorticity

The observations of Pearson & McCaughrean (2023) contradict the prediction, based on the absence of a characteristic scale of angular-momentum limited collapse (condition 1), that the wide binary fraction should be independent of  $R$  (and hence of  $M$ ) in an initially homogeneous cloud. This prediction does not apply if  $|\langle\vec{\omega}\rangle|$  depends on the length scale  $R$  because, if so, whether a region satisfies the collapse criterion would depend on its size. Viscosity diffuses vorticity, with a characteristic time scale  $\propto R^2$ . The evolution of vorticity is described by

$$\frac{d\vec{\omega}}{dt} = (\vec{\omega} \cdot \vec{\nabla})\vec{v} - \vec{\omega}(\vec{\nabla} \cdot \vec{v}) + \frac{\eta}{\rho}\nabla^2\vec{\omega}, \quad (2)$$

where we have dropped the baroclinic source term  $(\vec{\nabla}\rho \times \vec{\nabla}p)/\rho^2$  because the pressure field  $p$  is unknown and likely small compared to  $\rho v^2$  (as a result of radiation of thermal energy),

and also ignore body forces because self-gravity conserves angular momentum.

In a simple (two-dimensional) geometry in which  $\vec{v}$  lies in the  $x$ - $y$  plane and is independent of  $z$ ,  $\vec{\omega}$  is in the  $z$  direction and the first term on the right hand side is zero. The most important part of Eq. 2 is the second term on the right hand side, which is large and positive ( $\vec{\nabla} \cdot \vec{v} < 0$ ) in a contracting subregion. Scaling or dimensional analysis shows that, if viscosity is unimportant, as the subregion contracts the vorticity grows in proportion to the density:

$$\begin{aligned} \frac{d\vec{\omega}}{dt} &\approx \vec{\omega} \frac{1}{\rho} \frac{\partial \rho}{\partial t}, \\ |\vec{\omega}| &\propto \rho. \end{aligned} \tag{3}$$

As a result, the left hand side of (1) grows faster than the right hand side until angular momentum prevents further contraction.

If self-gravity is negligible, as is likely for the length scales and masses considered here ( $\sqrt{4\pi G\rho/3} \sim 2 \times 10^{-11} \text{ s}^{-1}$ ), viscosity homogenizes the vorticity of an isolated subregion, leaving it in a state of uniform (rigid-body) rotation. Contraction may resume if viscosity diffuses vorticity away from the contracting medium and transports angular momentum to its less vortical surroundings. Viscous transport of vorticity (and angular momentum) from the collapsing subregion competes with the amplification of vorticity by collapse.

Viscosity is more effective on smaller spatial scales because in Eq. 2 it is a diffusive term, reducing the vorticity and angular momentum of smaller, less massive, subregions more effectively than those of larger, more massive, regions. As a result, less massive subregions more readily contract, increasing the wide binary fraction of less massive objects. This scaling may account for the larger wide binary fraction of the least massive objects discovered by Pearson & McCaughrean (2023).

When self-gravity becomes significant, angular momentum flows from the inner parts

of the cloud to its outer parts, and it tends towards an accretion disc in which Kepler’s laws are inconsistent with uniform vorticity. In a Keplerian disc  $|\vec{\omega}| \propto r^{-3/2}$ . This occurs later in the collapse, after most of the initial angular momentum has been transferred to the orbital angular momentum of the binary proto-planets.

## 5. Cosmic Ray Viscosity

It remains to find a significant source of viscosity. The atomic viscosity of gases is negligible on astronomical scales, but interstellar gas contains cosmic rays that contribute a large effective viscosity because of their long free paths and relativistic velocities (Earl, Jokipii & Morfill 1988). The magnetic fields permeating the gas prevent the free escape of the cosmic rays, although a quantitative description would require knowledge of the magnetic field topology—the particles are free to travel along field lines, so transport in that direction depends on whether they form closed loops and the dimensions of such loops, and on whether there are magnetic mirrors along their lengths.

As a rough approximation, the cosmic ray dynamic viscosity

$$\eta \sim n_{CR}(\gamma - 1)m_{CR}v_{CR}\ell_{CR}, \quad (4)$$

where  $n_{CR}$ ,  $m_{CR}$ ,  $\gamma$ ,  $v_{CR} \approx c$  and  $\ell_{CR}$  are respectively the cosmic ray number density, mass, Lorentz factor, speed and scattering length, integrated over the cosmic ray spectrum. For the general interstellar medium  $n_{CR}(\gamma - 1)m_{CR}c^2 \sim 1 \text{ eV/cm}^3$  (Webber 1998).

The scattering length  $\ell_{CR}$  is strongly anisotropic. In the directions perpendicular to the magnetic field it may be estimated (in the spirit of Bohm diffusion (Chen 1984))

$$\ell_{\perp} \sim r_g = \frac{m_{CR}c^2}{eB}, \quad (5)$$

where the gyroradius  $r_g$  of a cosmic ray proton with 1 GeV kinetic energy in a representative

interstellar  $3 \mu\text{G}$  magnetic field  $r_g \sim 2 \times 10^{12} \text{ cm}$ . If  $\ell_{CR}$  is taken as  $\ell_{\perp}$  then  $\eta \sim 10^{-10} \text{ g/cm-s}$ , which is negligible.

But in the direction parallel to the magnetic field a cosmic ray may travel a length comparable to the radius of curvature of the field. We know from comparing the rotation measures of pulsars to the magnitude of the interstellar field that the interstellar field is ordered on long scales. Hence we estimate

$$\ell_{\parallel} \sim R, \tag{6}$$

where  $R$  is the radius of a collapsing subregion; once that subregion has reached a density a few times higher than that of its surroundings the field must bend by an angle  $\mathcal{O}(1 \text{ rad})$  between the subregion and its less dense surroundings. That bend will not scatter the cosmic rays (its radius of curvature is  $\sim R \gg r_g$ ), but in being bent the cosmic rays couple their momentum into the surrounding gas that anchors the magnetic field and provide an effective viscosity;  $R$  is the length on which the cosmic rays couple momentum.

The separations and masses of binary planetary mass objects in the Trapezium cluster indicate that they formed at a density  $\sim 10^9$  times that of the general interstellar medium. The column density of the collapsing cloud  $nR \sim 10^{24} \text{ cm}^{-2}$  is less than the collisional interaction length  $\sim 3 \times 10^{25} \text{ cm}^{-2}$  of cosmic rays with hydrogen, so cosmic rays may flow through the collapsing cloud, viscously coupling it to its surroundings. Because cosmic rays are inextricably bound (by the magnetic field) to the conducting interstellar gas, their density  $n_{CR} \propto n$  as the cloud collapses. The energy of individual cosmic rays is increased by compression of the cosmic ray-gas fluid; because cosmic rays are relativistic  $\gamma \propto n^{1/3}$ . Viscosity scales  $\eta \propto n\gamma\ell_{\parallel} \propto n\gamma R \propto n \propto R^{-3}$ , so that in a proto-JuMBO subregion with  $n \sim 10^9 \text{ cm}^{-3}$ ,  $\eta \sim 0.5 \times 10^5 \text{ g/cm-s}$ .

The effect of viscosity is quantified by the Reynolds Number

$$\text{Re} \sim \frac{\rho R V}{\eta} \sim \frac{\rho R^2 |\vec{\omega}|}{\eta}, \quad (7)$$

where  $V \sim R|\vec{\omega}|$  is a characteristic velocity and  $\eta$  is the (dynamic) viscosity. In a region forming a wide Jupiter-mass binary with separation  $R \sim 100$  AU and mass  $M \sim 10^{-3}M_{\odot}$  the density  $\rho \sim 10^{-15}$  g/cm<sup>3</sup>. For a subregion on the threshold of collapse to a binary planet, estimating  $|\vec{\omega}| \sim 2 \times 10^{-11}$  s<sup>-1</sup> from (1), leads to a Reynolds number  $\text{Re} \sim 1$ , implying effective removal of angular momentum from the subregion to its surroundings.

This  $\text{Re}$  is small enough to imply that, in clouds with little enough vorticity to permit collapse to begin, viscosity can transfer enough of their angular momentum to surrounding matter to permit further collapse to a wide binary (the binary orbit taking up the remaining angular momentum). The  $R^2$  factor in Eq. 7 explains why this process is more effective for smaller (lower mass) clouds, and leads to formation of a larger fraction of wide binaries at low masses. This scaling facilitates the collapse of smaller clouds more than that of larger clouds, and hence prefers the formation of less massive wide binaries over more massive wide binaries, explaining the observations of Pearson & McCaughrean (2023).

In the subsequent collapse of any particular cloud,  $\text{Re}$  increases  $\propto 1/R$  (Eq. 7, noting that  $|\vec{\omega}| \propto \rho \propto R^{-3}$  (Eq. 3) and  $\gamma \propto R^{-1}$ ). The cloud will rotate nearly rigidly, until the increasing importance of gravity forces its matter into a proto-Keplerian accretion disc. Cosmic ray viscosity will continue to transfer angular momentum from the cloud to its surroundings, facilitating its collapse.

## 6. Discussion

The maximum  $|\vec{\omega}|$  consistent with collapse (1) to form the wide planetary binaries discovered by Pearson & McCaughrean (2023) is several orders of magnitude less than

values implied by the scaling relation  $|\vec{\omega}| \propto \rho$  derived from Eq. 2. Star or planet (even wide binary) formation only occurs in subregions whose complex hydrodynamics has fortuitously led to vorticity much less than its mean in the extended cloud. At any moment only a tiny fraction of the mass of a cloud like that in the Trapezium cluster has sufficiently low vorticity to form stars or planets.

Pearson & McCaughrean (2023) found that the  $A_V$  of the primary and secondary objects typically differ by  $\sim 5$  magnitudes. Combining this with a nominal interstellar extinction per unit hydrogen column density of  $2 \text{ mag/kpc-cm}^{-3}$  implies a gas density over a 100 AU difference in distance  $n \sim 5 \times 10^6 \text{ cm}^{-3}$ , about 200 times less than the density required to form the binary. The intra-binary space must have been largely cleared of gas since the binary was formed, either by accretion onto the planetary objects or by expulsion by their changing gravitational potential. An alternative explanation is that binaries still immersed in the very dense pre-collapse gas are difficult to detect, and even more difficult to recognize as binaries (requiring detection of both objects), because of their excessive extinction, even in the infrared, so that observational selection favors detection of binaries in less dense regions.

An implication of the arguments presented here is that nearly all stars (and free-floating planets) form as wide binaries because of the difficulty and necessity of removing angular momentum. Some survive to be observed as common proper motion binaries, but most are likely disrupted by gravitational interaction with other objects in their star-forming regions of birth or subsequently in the interstellar medium. This predicts an anticorrelation of wide binary fraction in star-forming regions with these regions' age and also with their stellar density.

Another implication is that energetic particle viscosity may contribute to angular momentum transport in other astronomical environments. It may occur in the dilute

coronæ of accretion discs where relativistic particles are accelerated, and may contribute significantly to their viscosity and energy dissipation. Particles may be trapped on field lines with feet in the dense disc plasma, whose differential rotation accelerates the particles in analogy to second-order Fermi acceleration. This leads to a mutually amplifying feedback among particle acceleration, radial flow of mass and angular momentum, and viscous dissipation. Escape of particles along open field lines that result from magnetic reconnection powers the extended radio sources of quasars and microquasars (Katz 2024).

### **Compliance with Ethical Standards**

The author has no potential conflicts of interest.

### **Data Availability**

This theoretical study generated no new data.

## REFERENCES

- Binney, J. & Tremaine, S. 2008 *Galactic Dynamics* 2nd ed. (Princeton U. Press, Princeton).
- Chen, F. F. 1984 *Plasma Physics and Controlled Fusion* 2nd ed. (Plenum, New York).
- Earl, J. A., Jokipii, J. R. & Morfill, G. 1988 ApJ 331, L91.
- Fontanive, C., Bedin, L. R., De Furio, M., Biller, B., Anderson, J., Bonavita, M., Allers, K. & Pantoja, B. 2023 MNRAS 526, 1783.
- Han, J. L., Manchester, R. N. & Qiao, G. J. 1999 MNRAS 306, 371.
- Jumper, P. H. & Fisher, R. T. 2013 ApJ 769, 3.
- Katz, J. I. 1987 *High Energy Astrophysics* §3.2 (Addison-Wesley, Menlo-Park).
- Katz, J. I. 2024 Ap. Sp. Sci. submitted arXiv:2312.10834.
- Offner, S. S. R., Moe, M., Kratter, K. M., Sadavoy, S. T., Jensen, E. L. N. & Tobin, J. J. 2023 *Protostars and Planets VII* eds. Inutsuka, I., Aikawa, Y., Muto, T., Tomida, K. & Tamura, M. Ast. Soc. Pac. Conf. Ser. 534, 275 (arXiv:2203.10066).
- Pearson, S. G. & McCaughrean, M. J. 2023 arXiv:2310.01231.
- Tohline, J. E. 2002 ARA&A 40, 349.
- Webber, W. R. 1998 ApJ 506, 329.