

# Laguerre inequalities and determinantal inequalities for the finite difference of the partition functions

Eve Y.Y. Yang

Department of Mathematics

Tianjin University

Tianjin 300072, P. R. China

Email: yangyaoyao@tju.edu.cn

**Abstract.** The paper aims to establish the Turán inequalities, the Laguerre inequalities (order 2), and the determinantal inequalities (order 3) for  $\Delta p(n)$  and  $\Delta \bar{p}(n)$ , where  $\Delta f(n)$  is the first-order forward difference of a sequence  $f(n)$ . The functions  $p(n)$  and  $\bar{p}(n)$  denote the partition function and overpartition function, respectively. Conjectures for thresholds of Laguerre inequalities (order  $m$ ) and positivity of  $m$ -order determinants are proposed, extending to  $\Delta^k p(n)$  and  $\Delta^k \bar{p}(n)$ , with  $1 \leq m \leq 11$  and  $1 \leq k \leq 5$ .

**Keywords:** the finite difference, partition function, overpartition function, Laguerre inequality, determinant inequality

**AMS Classification:** 05A20, 11P82

## 1 Introduction

A positive integer partition of  $n$  is a non-increasing sequence of positive integers  $\lambda = (\lambda_1, \dots, \lambda_\ell)$  with  $\sum_{i=1}^{\ell} \lambda_i = n$ . An overpartition of  $n$  allows overlining the first occurrence of each distinct part in the partition. The number of partitions and overpartitions of  $n$  are denoted by  $p(n)$  and  $\bar{p}(n)$ , respectively.

Various inequalities concerning  $p(n)$  and  $\bar{p}(n)$  have been explored using analytical tools, incorporating the Hardy-Ramanujan-Rademacher formulae for  $p(n)$  and  $\bar{p}(n)$  and associated error bounds detailed in [16, 37, 38, 40, 56].

The first-order forward difference of a sequence  $f(n)$  is defined as

$$\Delta f(n) = f(n+1) - f(n).$$

Iterating this operation yields the  $k$ -th order forward difference,

$$\Delta^k f(n) = \Delta(\Delta^{k-1} f(n)).$$

In 1977, Good [25] conjectured  $\Delta^r p(n) \geq 0$  for a positive integer  $n(r)$ . Employing the Hardy-Ramanujan-Rademacher formula for  $p(n)$ , Gupta [27] proved  $\Delta^r p(n) > 0$  for sufficiently large  $n$ . Subsequently, Odlyzko [43] confirmed Good's conjecture and derived the asymptotic formula for  $n(r)$ :

$$n(r) \sim (6/\pi^2)r^2 \log^2 r \quad \text{as } r \rightarrow \infty.$$

Knessl and Keller [33, 34] refined this asymptotic approximation  $n'(r)$  of  $n(r)$  by deriving a recursion equation for  $\Delta^r p(n)$  with respect to  $n$  and solving it asymptotically. They showed that

$$|n'(r) - n(r)| \leq 2, \quad \text{for all } r \leq 75.$$

Almkvist [1, 2] discovered a more explicit formula for  $\Delta^r p(n) > 0$  and established specific equations satisfied by  $n(r)$ . Wang, Xie and Zhang [50] extended this to  $\Delta^r \bar{p}(n) > 0$  and provided an upper bound for  $\Delta^r \bar{p}(n)$  for any  $r \geq 1$ . Yang [54] extended this approach effectively to  $\Delta^r q(n)$  and  $\{\Delta^r \Delta_k(n)\}_{k=1,2}$ , where  $q(n)$  is the distinct partition function and  $\Delta_k(n)$  is the broken  $k$ -diamond partition function.

Consider the  $j$ -shifted differences of a sequence  $f(n)$ , denoted as  $\Delta_j f(n) = f(n) - f(n-j)$ , where  $1 \leq j < n$ . Notably, for  $j = 1$ , this corresponds to the first-order backward difference, distinct from the forward difference defined earlier. Gomez, Males and Rolén [24] proved that

$$\Delta_j^2 p(n) = \Delta_j (\Delta_j p(n)) = p(n) - 2p(n-j) + p(n-2j) > 0$$

by using the Hardy-Ramanujan-Rademacher formula for  $p(n)$  and Lehmer's error bound. They conjectured  $\Delta_j^r p(n) \geq 0$  for a fixed  $j$ , any given  $r$ , and sufficiently large  $n$ . The case for  $j = 1$  had been previously proved by Gupta [27]. Banerjee [3] proved  $\Delta_j^2 p(n) \geq 0$  and  $\Delta_j^2 \bar{p}(n) > 0$  using an elementary combinatorial approach. Numerous related works exist [4, 8, 32, 41].

The Turán-type inequalities arise in the study of the Maclaurin coefficients of real entire functions in the Laguerre-Pólya class. For a more detailed study, we refer to [10, 18, 39, 46, 47].

A sequence  $\{a_n\}_{n \geq 0}$  of real numbers is log-concave if it satisfies the (second order) Turán inequality  $a_n^2 \geq a_{n-1}a_{n+1}$  for  $n \geq 1$ . It satisfies the third-order Turán inequality if, for  $n \geq 1$ ,

$$4(a_n^2 - a_{n-1}a_{n+1})(a_{n+1}^2 - a_n a_{n+2}) \geq (a_n a_{n+1} - a_{n-1} a_{n+2})^2.$$

Note that the log-concavity of a sequence  $f(n)$  implies that  $-\Delta^2 \log f(n) \geq 0$ .

Define the operator  $\mathcal{L}$  as

$$\mathcal{L}a_n = a_n^2 - a_{n-1}a_{n+1} \quad \text{and} \quad \mathcal{L}^r a_n = \mathcal{L}(\mathcal{L}^{r-1}a_n).$$

A sequence  $\{a_n\}_{n \geq 0}$  is said to be  $r$ -log-concave if, for  $1 \leq k \leq r$  and  $n \geq k$ ,  $\{\mathcal{L}^k a_n\}_{n \geq 1}$  are all nonnegative sequences. It should be noted that 2-log-concavity is consistent with the double Turán inequality.

The Turán-type inequalities are closely related to the Jensen polynomials. The Jensen polynomials of degree  $d$  and shift  $n$  associated with an arbitrary real sequence  $\{a_n\}_{n \geq 0}$  are defined by

$$J_a^{d,n}(X) := \sum_{j=0}^d \binom{d}{j} a_{n+j} X^j.$$

In general, a sequence  $\{a_n\}_{n \geq 0}$  satisfies the Turán inequality of order  $d$  at  $n$  if and only if  $J_a^{d,n-1}(X)$  is hyperbolic. A real polynomial is said to be hyperbolic if all of its zeros are real.

Pólya [46] proved that the Riemann hypothesis (RH) is equivalent to the hyperbolicity of the polynomials  $J_{\hat{\gamma}_k}^{d,n}$  for all  $d, n \geq 0$ , where  $\{\hat{\gamma}_k\}_{k \geq 0}$  is the sequence of Maclaurin coefficients of the Riemann Xi-function. Due to the difficulty of proving RH, research has concentrated on proving hyperbolicity for all  $n \geq 0$  when  $d$  is small. The cases  $d = 2$  and  $d = 3$  imply that  $\{\hat{\gamma}_k\}_{k \geq 0}$  satisfies both the Turán inequality and the third-order Turán inequality, as proved by Csordas, Norfolk and Varga [13], and Dimitrov and Lucas [19], respectively.

Further properties of the Turán-type inequalities, the Laguerre-Pólya class, the Jensen polynomials, and the Riemann Xi-function are detailed in [10, 13, 14, 18, 19, 29, 46].

The Laguerre inequality, also relevant to the Laguerre-Pólya class, initially formulated for a polynomial function  $f(x)$ , is expressed as

$$L_1(f(x)) := f'(x)^2 - f(x)f''(x) \geq 0, \quad x \in \mathbb{R}. \quad (1.1)$$

Laguerre [35] stated that if  $f(x)$  is hyperbolic, then it satisfies (1.1). Skovgaard [48] asserted that if  $f(x)$  belongs to the Laguerre-Pólya class, then for any  $p \in \mathbb{Z}_{\geq 0}$ ,  $f^p(x)$  satisfies (1.1), where  $f^p(x)$  represents the  $p$ -th derivative of  $f(x) = f^0(x)$ .

In 1913, Jensen [29] defined a higher order generalization of the Laguerre inequality, namely, for  $f(x)$  in the Laguerre-Pólya class,

$$L_n(f(x)) := \frac{1}{2} \sum_{k=0}^{2n} (-1)^{n+k} \binom{2n}{k} f^k(x) f^{2n-k}(x) \geq 0.$$

These inequalities, often referred to as Laguerre-type inequalities, yield the classical Laguerre inequality (1.1) for  $n = 1$ . Csordas and Varga [14] showed

that if a function  $f(x)$  satisfies  $L_n(f(x)) \geq 0$  for all  $n$  and all  $x \in \mathbb{R}$ , then  $f(x)$  belongs to the Laguerre-Pólya class.

Recently, Wang and Yang [51] considered whether the discrete sequence  $\{a_n\}_{n \geq 0}$  has the similar results with higher order Laguerre inequalities. They defined that a sequence  $\{a_n\}_{n \geq 0}$  satisfies the Laguerre inequality of order  $m$  if

$$L_m(a_n) := \frac{1}{2} \sum_{k=0}^{2m} (-1)^{k+m} \binom{2m}{k} a_{n+k} a_{2m-k+n} \geq 0.$$

For  $m = 1$ , this reduces to

$$a_{n+1}^2 - a_n a_{n+2} \geq 0,$$

which corresponds to the log-concavity of  $\{a_n\}_{n \geq 0}$ .

Wang and Yang [51, 53] established the Laguerre inequality of order 2 for some discrete sequences, including  $p(n)$ ,  $\bar{p}(n)$ ,  $q(n)$ , and others. Wagner [49] proved that  $p(n)$  satisfies the Laguerre inequalities of any order as  $n \rightarrow \infty$  and conjectured the thresholds of the  $m$ -rd Laguerre inequalities of  $p(n)$  for  $m \leq 10$ . Dou and Wang [22] proved Wagner's conjecture for  $3 \leq m \leq 9$ . For more work on the relation between the Laguerre inequality, the Turán inequality and the Laguerre-Pólya class, refer to [5, 10, 11, 12, 17, 23, 44, 45].

Chen [6] conducted a comprehensive study on inequalities related to invariants of a binary form. He examined the following three invariants associated with the hyperbolicity of the Jensen polynomial of degree 4.

$$A(a_0, a_1, a_2, a_3, a_4) = a_0 a_4 - 4a_1 a_3 + 3a_2^2,$$

$$B(a_0, a_1, a_2, a_3, a_4) = -a_0 a_2 a_4 + a_2^3 + a_0 a_3^2 + a_1^2 a_4 - 2a_1 a_2 a_3,$$

$$I(a_0, a_1, a_2, a_3, a_4) = A(a_0, a_1, a_2, a_3, a_4)^3 - 27B(a_0, a_1, a_2, a_3, a_4)^2.$$

Throughout the rest of this paper, we shall use the notations  $A$ ,  $B$ , and  $I$  to denote  $A(a_0, a_1, a_2, a_3, a_4)$ ,  $B(a_0, a_1, a_2, a_3, a_4)$ , and  $I(a_0, a_1, a_2, a_3, a_4)$  respectively, associated with a sequence  $\{a_n\}_{n \geq 0}$ .

Indeed,  $A > 0$  coincides with the Laguerre inequality of order 2.  $B > 0$  can be expressed as

$$\begin{vmatrix} a_2 & a_3 & a_4 \\ a_1 & a_2 & a_3 \\ a_0 & a_1 & a_2 \end{vmatrix} > 0,$$

which has been proven to be equivalent to 2-log-concavity in combinatorics. For  $p(n)$ ,  $B > 0$  has been showed by Hou and Zhang [28], and Jia and Wang [31] independently. Recently, Wang and Yang [52] proved  $\det(a_{n-i+j})_{1 \leq i, j \leq 4} >$

0 for the sequence  $\{a_n\}_{n \geq 0}$  involving  $p(n)$  and  $\bar{p}(n)$  and provided an iterated approach to compute  $\det(a_{n-i+j})_{1 \leq i, j \leq m}$  for any positive integer  $m$ .

Recent work by various mathematicians has established Turán inequalities of order  $d$  for  $p(n)$ . Nicolas [42] and DeSalvo and Pak [15] independently confirmed the log-concavity of  $\{p(n)\}_{n \geq 25}$ . Chen [6] conjectured and Chen, Jia and Wang [7] proved the third-order Turán inequality for  $\{p(n)\}_{n \geq 95}$ . Chen, Jia and Wang [7] extended this to propose a conjecture for Turán inequalities of order  $d$  for  $p(n)$  with  $d \geq 4$  and sufficiently large  $n$ , which was proven by Griffin, Ono, Rolin and Zagier [26]. Larson and Wagner [36] provided the thresholds for Turán inequalities of order 4 and 5.

Turán inequalities for other partition functions have been extensively investigated, refer to [9, 20, 21, 30, 53, 55] for further details.

Let

$$\Delta p(n) = p(n+1) - p(n) \quad \text{and} \quad \Delta \bar{p}(n) = \bar{p}(n+1) - \bar{p}(n).$$

This paper establishes the Turán inequalities, the Laguerre inequalities of order 2, and the determinantal inequalities of order 3 for  $\Delta p(n)$  and  $\Delta \bar{p}(n)$ . Our main tool, due to Dou and Wang [22], relies on Lemma 1.1 for  $p(n)$  and Lemma 1.2 for  $\bar{p}(n)$ , both provided by Wang and Yang [52].

**Lemma 1.1** ([52, Lemma 2.1]). *For any given integer  $l$ , there exists  $N(l)$  such that for all  $n \geq N(l)$ ,*

$$\frac{\sqrt{12}\pi^2 e^\mu}{36\mu^2} \left(1 - \frac{1}{\mu} - \frac{1}{\mu^l}\right) < p(n) < \frac{\sqrt{12}\pi^2 e^\mu}{36\mu^2} \left(1 - \frac{1}{\mu} + \frac{1}{\mu^l}\right),$$

where  $\mu$  is the abbreviation for  $\mu(n)$ , and

$$\mu(n) = \frac{\pi\sqrt{24n-1}}{6}. \tag{1.2}$$

**Lemma 1.2** ([52, Lemma 5.1]). *For any given integer  $t$ , there exists  $N(t)$  such that for all  $n \geq N(t)$ ,*

$$\frac{\pi^2 e^{\bar{\mu}}}{8\bar{\mu}^2} \left(1 - \frac{1}{\bar{\mu}} - \frac{1}{\bar{\mu}^t}\right) < \bar{p}(n) < \frac{\pi^2 e^{\bar{\mu}}}{8\bar{\mu}^2} \left(1 - \frac{1}{\bar{\mu}} + \frac{1}{\bar{\mu}^t}\right),$$

where  $\bar{\mu}$  is the abbreviation for  $\bar{\mu}(n)$ , and

$$\bar{\mu}(n) = \pi\sqrt{n}.$$

Wang and Yang [52] inferred that  $n \geq N(k)$  ( $k = l, t$ ) in the above Lemmas are solutions to  $6e^{-\frac{u}{2}} < \frac{1}{u^k}$  ( $u = \mu, \bar{\mu}$ ).

The remaining of this paper is organized as follows. Section 2 establishes the Turán inequalities for  $\Delta p(n)$  and  $\Delta \bar{p}(n)$ . Section 3, following the spirit of Section 2, establishes the Laguerre inequalities for  $\Delta p(n)$  and  $\Delta \bar{p}(n)$ . Similarly, Section 4 establishes the determinantal inequalities of order 3 for  $\Delta p(n)$  and  $\Delta \bar{p}(n)$ . We conclude in Section 5 with open problems for further exploration.

## 2 The Turán inequalities for $\Delta p(n)$ and $\Delta \bar{p}(n)$

In this section, we will establish the Turán inequalities for  $\Delta p(n)$  and  $\Delta \bar{p}(n)$ , detailed in Theorem 2.1 and Theorem 2.2, respectively.

**Theorem 2.1.** *For  $n \geq 71$ ,  $\Delta p(n)$  satisfies the Turán inequality,*

$$(\Delta p(n))^2 > (\Delta p(n-1)) (\Delta p(n+1)). \quad (2.1)$$

**Theorem 2.2.** *For  $n \geq 8$ ,  $\Delta \bar{p}(n)$  satisfies the Turán inequality,*

$$(\Delta \bar{p}(n))^2 > (\Delta \bar{p}(n-1)) (\Delta \bar{p}(n+1)). \quad (2.2)$$

In fact, the entire approach used to prove Theorem 2.1 is also applicable to Theorem 2.2. Thus, we will provide a detailed proof only for the former.

*Proof of Theorem 2.1.* Note that (2.1) can be rewritten as

$$\begin{aligned} p(n)^2 - p(n-1)p(n+1) - p(n)p(n+1) + p(n+1)^2 \\ + p(n-1)p(n+2) - p(n)p(n+2) > 0. \end{aligned} \quad (2.3)$$

To prove it, setting  $l = 6$  in Lemma 1.1, obtaining the bounds for  $n \geq 391$ ,

$$e^\mu \frac{\sqrt{12}\beta(\mu)\pi^2}{36\mu^8} < p(n) < e^\mu \frac{\sqrt{12}\alpha(\mu)\pi^2}{36\mu^8}, \quad (2.4)$$

where

$$\alpha(t) = t^6 - t^5 + 1, \quad \beta(t) = t^6 - t^5 - 1. \quad (2.5)$$

Define

$$f(n) := e^\mu \frac{\sqrt{12}\beta(\mu)\pi^2}{36\mu^8}, \quad g(n) := e^\mu \frac{\sqrt{12}\alpha(\mu)\pi^2}{36\mu^8}. \quad (2.6)$$

Then, to establish (2.1), we aim to show that

$$\begin{aligned} f(n)^2 - g(n-1)g(n+1) - g(n)g(n+1) + f(n+1)^2 \\ + f(n-1)f(n+2) - g(n)g(n+2) > 0, \end{aligned}$$

which is equivalent to

$$\begin{aligned} & \frac{f(n)^2}{f(n+1)^2} - \frac{g(n-1)g(n+1)}{f(n+1)^2} - \frac{g(n)g(n+1)}{f(n+1)^2} + 1 \\ & + \frac{f(n-1)f(n+2)}{f(n+1)^2} - \frac{g(n)g(n+2)}{f(n+1)^2} > 0. \end{aligned} \quad (2.7)$$

For convenience, let

$$w = \mu(n-1), \quad x = \mu(n), \quad y = \mu(n+1), \quad z = \mu(n+2).$$

Simplify the left-hand side of the inequality (2.7) as

$$\frac{-h_1 e^{w-y} - h_2 e^{x-y} - h_3 e^{x-2y+z} + h_4 e^{2x-2y} + h_5 e^{w-2y+z} + h_6}{h_6}, \quad (2.8)$$

where

$$\begin{aligned} h_1 &= x^{16} y^8 z^8 \alpha(w) \alpha(y), & h_2 &= w^8 x^8 y^8 z^8 \alpha(x) \alpha(y), \\ h_3 &= w^8 x^8 y^{16} \alpha(x) \alpha(z), & h_4 &= w^8 y^{16} z^8 \beta(x)^2, \\ h_5 &= x^{16} y^{16} \beta(w) \beta(z), & h_6 &= w^8 x^{16} z^8 \beta(y)^2. \end{aligned} \quad (2.9)$$

Clearly,  $h_6 > 0$  for  $n \geq 1$ . Thus, our focus is only on proving the positivity of the numerator in (2.8), i.e.,

$$-h_1 e^{w-y} - h_2 e^{x-y} - h_3 e^{x-2y+z} + h_4 e^{2x-2y} + h_5 e^{w-2y+z} + h_6 > 0. \quad (2.10)$$

To achieve this, we need to estimate  $h_1, h_2, h_3, h_4, h_5, h_6, e^{w-y}, e^{x-y}, e^{x-2y+z}, e^{2x-2y}$  and  $e^{w-2y+z}$ . We prefer to use the following equations to estimate  $w, x,$  and  $z$ . For  $n \geq 2$ ,

$$w = \sqrt{y^2 - \frac{4\pi^2}{3}}, \quad x = \sqrt{y^2 - \frac{2\pi^2}{3}}, \quad z = \sqrt{y^2 + \frac{2\pi^2}{3}}. \quad (2.11)$$

Their Taylor expansions are

$$\begin{aligned} w &= y - \frac{2\pi^2}{3y} - \frac{2\pi^4}{9y^3} - \frac{4\pi^6}{27y^5} + O\left(\frac{1}{y^7}\right), \\ x &= y - \frac{\pi^2}{3y} - \frac{\pi^4}{18y^3} - \frac{\pi^6}{54y^5} + O\left(\frac{1}{y^7}\right), \\ z &= y + \frac{\pi^2}{3y} - \frac{\pi^4}{18y^3} + \frac{\pi^6}{54y^5} + O\left(\frac{1}{y^7}\right). \end{aligned}$$

Then, it can be checked that for  $n \geq 2$ ,

$$w_1 < w < w_2, \quad x_1 < x < x_2, \quad z_1 < z < z_2, \quad (2.12)$$

where

$$\begin{aligned}
w_1 &= y - \frac{2\pi^2}{3y} - \frac{2\pi^4}{9y^3} - \frac{4\pi^6}{27y^5}, & w_2 &= y - \frac{2\pi^2}{3y} - \frac{2\pi^4}{9y^3} - \frac{3\pi^6}{27y^5}, \\
x_1 &= y - \frac{\pi^2}{3y} - \frac{\pi^4}{18y^3} - \frac{\pi^6}{54y^5}, & x_2 &= y - \frac{\pi^2}{3y} - \frac{\pi^4}{18y^3} - \frac{\pi^6}{108y^5}, \\
z_1 &= y + \frac{\pi^2}{3y} - \frac{\pi^4}{18y^3}, & z_2 &= y + \frac{\pi^2}{3y} - \frac{\pi^4}{18y^3} + \frac{\pi^6}{54y^5}.
\end{aligned}$$

Next, we estimate  $h_1, h_2, h_3, h_4$  and  $h_5$ . Applying (2.12) to the definitions (2.5) of  $\alpha(t)$  and  $\beta(t)$ , we obtain that for  $n \geq 2$ ,

$$\begin{aligned}
w^6 - w_2w^4 + 1 &< \alpha(w) < w^6 - w_1w^4 + 1, \\
x^6 - x_2x^4 + 1 &< \alpha(x) < x^6 - x_1x^4 + 1, \\
z^6 - z_2z^4 + 1 &< \alpha(z) < z^6 - z_1z^4 + 1, \\
w^6 - w_2w^4 - 1 &< \beta(w) < w^6 - w_1w^4 - 1, \\
x^6 - x_2x^4 - 1 &< \beta(x) < x^6 - x_1x^4 - 1, \\
z^6 - z_2z^4 - 1 &< \beta(z) < z^6 - z_1z^4 - 1.
\end{aligned}$$

Substituting these  $\alpha(t)$  and  $\beta(t)$  into  $h_1, h_2, h_3, h_4, h_5$ , we get

$$\begin{aligned}
h_1 &< x^{16}y^8z^8 (w^6 - w_1w^4 + 1) (y^6 - y^5 + 1), \\
h_2 &< w^8x^8y^8z^8 (x^6 - x_1x^4 + 1) (y^6 - y^5 + 1), \\
h_3 &< w^8x^8y^{16} (x^6 - x_1x^4 + 1) (z^6 - z_1z^4 + 1), \\
h_4 &> w^8y^{16}z^8 (x^6 - x_2x^4 - 1)^2, \\
h_5 &> x^{16}y^{16} (w^6 - w_2w^4 - 1) (z^6 - z_2z^4 - 1).
\end{aligned} \tag{2.13}$$

Next we turn to estimate  $e^{w-y}, e^{x-y}, e^{x-2y+z}, e^{2x-2y}$  and  $e^{w-2y+z}$ . By (2.12), it can be seen that for  $n \geq 2$ ,

$$\begin{aligned}
w_1 - y &< w - y < w_2 - y, \\
x_1 - y &< x - y < x_2 - y, \\
x_1 - 2y + z_1 &< x - 2y + z < x_2 - 2y + z_2,
\end{aligned}$$



$$w_1 - 2y + z_1 < w - 2y + z < w_2 - 2y + z_2.$$

This implies that

$$\begin{aligned} e^{w_1-y} &< e^{w-y} < e^{w_2-y}, \\ e^{x_1-y} &< e^{x-y} < e^{x_2-y}, \\ e^{2x_1-2y} &< e^{2x-2y} < e^{2x_2-2y}, \\ e^{x_1-2y+z_1} &< e^{x-2y+z} < e^{x_2-2y+z_2}, \\ e^{w_1-2y+z_1} &< e^{w-2y+z} < e^{w_2-2y+z_2}. \end{aligned} \tag{2.14}$$

To provide a feasible bound for  $e^{x-2y+z}$  and  $e^{w-2y+z}$ , we define

$$\begin{aligned} \Phi(t) &= 1 + t + \frac{t^2}{2} + \frac{t^3}{6} + \frac{t^4}{24} + \frac{t^5}{120}, \\ \phi(t) &= 1 + t + \frac{t^2}{2} + \frac{t^3}{6} + \frac{t^4}{24} + \frac{t^5}{120} + \frac{t^6}{720}. \end{aligned}$$

For  $t < 0$ , it is evident that

$$\phi(t) < e^t < \Phi(t). \tag{2.15}$$

To apply this to (2.14), it suffices to show the negativity of  $w_2 - y$ ,  $x_2 - y$ ,  $2x_2 - 2y$ ,  $x_2 - 2y + z_2$  and  $w_2 - 2y + z_2$ . A direct calculation reveals that

$$\begin{aligned} w_2 - y &= -\frac{\pi^6 + 2\pi^4 y^2 + 6\pi^2 y^4}{9y^5}, \\ x_2 - y &= -\frac{\pi^6 + 6\pi^4 y^2 + 36\pi^2 y^4}{108y^5}, \\ x_2 - 2y + z_2 &= \frac{\pi^6 - 12\pi^4 y^2}{108y^5}, \\ w_2 - 2y + z_2 &= -\frac{5\pi^6 + 15\pi^4 y^2 + 18\pi^2 y^4}{54y^5}. \end{aligned} \tag{2.16}$$

Clearly,  $w_2 - y$ ,  $x_2 - y$ ,  $2x_2 - 2y$ ,  $x_2 - 2y + z_2$  and  $w_2 - 2y + z_2$  are negative for all  $n$ . Notably, for their negativity, a judicious choice of the denominator in (2.7) is essential. Thus, applying (2.15) to (2.14), we deduce

that for  $n \geq 2$ ,

$$\begin{aligned}
\phi(w_1 - y) &< e^{w-y} < \Phi(w_2 - y), \\
\phi(x_1 - y) &< e^{x-y} < \Phi(x_2 - y), \\
\phi(2x_1 - 2y) &< e^{2x-2y} < \Phi(2x_2 - 2y), \\
\phi(x_1 - 2y + z_1) &< e^{x-2y+z} < \Phi(x_2 - 2y + z_2), \\
\phi(w_1 - 2y + z_1) &< e^{w-2y+z} < \Phi(w_2 - 2y + z_2).
\end{aligned} \tag{2.17}$$

Now, we are ready for proving (2.10). Combing expressions (2.9) and (2.11), we can get that  $h_1, h_2, h_3, h_4, h_5, h_6, e^{w-y}, e^{x-y}, e^{x-2y+z}, e^{2x-2y}$  and  $e^{w-2y+z}$  are all functions of  $y$ . For convenience, let

$$A(y) = -h_1 e^{w-y} - h_2 e^{x-y} - h_3 e^{x-2y+z} + h_4 e^{2x-2y} + h_5 e^{w-2y+z} + h_6,$$

we need to show  $A(y) > 0$ . Using (2.13) and (2.17), it holds that for  $n \geq 2$ ,

$$\begin{aligned}
A(y) &> -x^{16}y^8z^8(w^6 - w_1w^4 + 1)(y^6 - y^5 + 1)\Phi(w_2 - y) \\
&\quad - w^8x^8y^8z^8(x^6 - x_1x^4 + 1)(y^6 - y^5 + 1)\Phi(x_2 - y) \\
&\quad - w^8x^8y^{16}(x^6 - x_1x^4 + 1)(z^6 - z_1z^4 + 1)\Phi(x_2 - 2y + z_2) \\
&\quad + w^8y^{16}z^8(x^6 - x_2x^4 - 1)^2\phi(2x_1 - 2y) \\
&\quad + x^{16}y^{16}(w^6 - w_2w^4 - 1)(z^6 - z_2z^4 - 1)\phi(w_1 - 2y + z_1) \\
&\quad + w^8x^{16}z^8\beta(y)^2.
\end{aligned}$$

Represent the right-hand side of the above inequality as  $A_1(y)$ . With Mathematica, we can quickly simplify  $A_1(y)$  as

$$A_1(y) = \frac{\sum_{k=0}^{66} a_k y^k}{2^{18}3^{37}5y^{27}}.$$

Note that all the coefficients of  $A_1(y)$  can be known, and the values of  $a_{64}, a_{65}, a_{66}$  are given below,

$$a_{64} = 2^{15}3^{29}5(629856 + 209952\pi^2 - 7776\pi^6 + 62856\pi^8 - 2133\pi^{10} + 55\pi^{12}),$$

$$a_{65} = -2^{14}3^{29}5(1259712 - 15552\pi^6 + 11988\pi^8 + 432\pi^{10} + 13\pi^{12}),$$

$$a_{66} = 2^{18}3^{33}5\pi^6(-6 + \pi^2).$$

Thus, for  $n \geq 2$ , we have

$$A(y) > \frac{\sum_{k=0}^{66} a_k y^k}{2^{18} 3^{37} 5^{27}}. \quad (2.18)$$

Since  $y$  is positive for  $n \geq 1$ , it follows that

$$\sum_{k=0}^{66} a_k y^k > \sum_{k=0}^{64} -|a_k| y^k + a_{65} y^{65} + a_{66} y^{66}.$$

For  $0 \leq k \leq 64$ , it can be verified that

$$-|a_k| y^k > -a_{64} y^{64}$$

holds when  $y > 7.68$ . This leads to

$$\sum_{k=0}^{66} a_k y^k > \sum_{k=0}^{64} -|a_k| y^k + a_{65} y^{65} + a_{66} y^{66} > (-65a_{64} + a_{65}y + a_{66}y^2)y^{64}. \quad (2.19)$$

Combing (2.18) and (2.19),  $A(y)$  is positive provided

$$-65a_{64} + a_{65}y + a_{66}y^2 > 0,$$

which holds for  $y > 126.22$ . From the relation between  $y$  and  $n$  in (1.2) and (2.11), we deduce that  $y > 126.22$  is equivalent to  $n \geq 2421$ . Thus, (2.10) is true for  $n \geq 2421$ , which implies (2.1). Numerical verification further confirms the truth of (2.1) for  $71 \leq n \leq 2420$ . Consequently, we conclude the proof of Theorem 2.1.  $\blacksquare$

To prove Theorem 2.2, we employ Lemma 1.2 and adopt notations consistent with the proof of Theorem 2.1, except adjusting  $p(n)$  and  $\mu(n)$  to  $\bar{p}(n)$  and  $\bar{\mu}(n)$ , respectively. The proof follows the framework established for Theorem 2.1. Hence, in the remaining of this section, we will omit some tedious formulae and statements and focus on presenting key modifications resulting from substituting  $p(n)$  and  $\mu(n)$  with  $\bar{p}(n)$  and  $\bar{\mu}(n)$ , respectively.

Considering the substitution of  $\bar{p}(n)$  for  $p(n)$ , we first adjust (2.3) to

$$\begin{aligned} & \bar{p}(n)^2 - \bar{p}(n-1)\bar{p}(n+1) - \bar{p}(n)\bar{p}(n+1) + \bar{p}(n+1)^2 \\ & + \bar{p}(n-1)\bar{p}(n+2) - \bar{p}(n)\bar{p}(n+2) > 0. \end{aligned}$$

Furthermore, with the substitution of  $\mu(n)$  for  $\bar{\mu}(n)$ , (2.4) transforms to

$$e^{\bar{\mu}} \frac{\beta(\bar{\mu})\pi^2}{8\bar{\mu}^8} < \bar{p}(n) < e^{\bar{\mu}} \frac{\alpha(\bar{\mu})\pi^2}{8\bar{\mu}^8},$$

where

$$\alpha(t) = t^6 - t^5 + 1, \quad \beta(t) = t^6 - t^5 - 1.$$

Then (2.6) transforms to

$$f(n) := e^{\bar{\mu}} \frac{\beta(\bar{\mu})\pi^2}{8\bar{\mu}^8}, \quad g(n) := e^{\bar{\mu}} \frac{\alpha(\bar{\mu})\pi^2}{8\bar{\mu}^8}.$$

As  $\mu(n)$  changes to  $\bar{\mu}(n)$ , (2.11) transforms to, for  $n \geq 2$ ,

$$w = \sqrt{y^2 - 2\pi^2}, \quad x = \sqrt{y^2 - \pi^2}, \quad z = \sqrt{y^2 + \pi^2}.$$

Their Taylor expansions are given by

$$\begin{aligned} w &= y - \frac{\pi^2}{y} - \frac{\pi^4}{2y^3} - \frac{\pi^6}{2y^5} + O\left(\frac{1}{y^7}\right), \\ x &= y - \frac{\pi^2}{2y} - \frac{\pi^4}{8y^3} - \frac{\pi^6}{16y^5} + O\left(\frac{1}{y^7}\right), \\ z &= y + \frac{\pi^2}{2y} - \frac{\pi^4}{8y^3} + \frac{\pi^6}{16y^5} + O\left(\frac{1}{y^7}\right). \end{aligned}$$

For  $n \geq 2$ , it can be checked that

$$w_1 < w < w_2, \quad x_1 < x < x_2, \quad z_1 < z < z_2,$$

where

$$\begin{aligned} w_1 &= y - \frac{\pi^2}{y} - \frac{\pi^4}{2y^3} - \frac{\pi^6}{2y^5}, & w_2 &= y - \frac{\pi^2}{y} - \frac{\pi^4}{2y^3} - \frac{\pi^6}{4y^5}, \\ x_1 &= y - \frac{\pi^2}{2y} - \frac{\pi^4}{8y^3} - \frac{\pi^6}{16y^5}, & x_2 &= y - \frac{\pi^2}{2y} - \frac{\pi^4}{8y^3} - \frac{\pi^6}{32y^5}, \\ z_1 &= y + \frac{\pi^2}{2y} - \frac{\pi^4}{8y^3}, & z_2 &= y + \frac{\pi^2}{2y} - \frac{\pi^4}{8y^3} + \frac{\pi^6}{16y^5}. \end{aligned}$$

Furthermore, (2.16) should be adjusted to

$$\begin{aligned} w_2 - y &= -\frac{\pi^6 + 2\pi^4 y^2 + 4\pi^2 y^4}{4y^5}, \\ x_2 - y &= -\frac{\pi^6 + 4\pi^4 y^2 + 16\pi^2 y^4}{32y^5}, \\ x_2 - 2y + z_2 &= \frac{\pi^6 - 8\pi^4 y^2}{32y^5}, \end{aligned}$$

$$w_2 - 2y + z_2 = -\frac{3\pi^6 + 10\pi^4 y^2 + 8\pi^2 y^4}{16y^5}.$$

Finally, applying similar arguments as in the proof of Theorem 2.1, we observe that  $A_1(y)$  is a certain polynomial of degree 39 with a positive leading coefficient. Verification confirms the positivity of  $A(y)$  for  $n \geq 1641$ , establishing the Turán inequality for  $\Delta\bar{p}(n)$  in this range. For  $8 \leq n \leq 1640$ , numerical verification additionally supports the Turán inequality for  $\Delta\bar{p}(n)$ . Thus, the proof of Theorem 2.2 is completed.  $\blacksquare$

### 3 The Laguerre inequalities for $\Delta p(n)$ and $\Delta\bar{p}(n)$

In this section, we will consider the Laguerre inequalities of order 2 for  $\Delta p(n)$  and  $\Delta\bar{p}(n)$  as showed in Theorem 3.1 and Theorem 3.2, respectively. In fact, the approach given in Section 2 still works here. The key point in the proof is to determine suitable values for  $l$  (resp.  $t$ ) in Lemma 1.1 (resp. Lemma 1.2), choose an appropriate denominator as in (2.7), and adjust the terms of the Taylor expansion for the exponential function, as well as the upper and lower bounds for  $\sqrt{w^2 - \frac{2\delta\pi^2}{3}}$  and  $\sqrt{w^2 + \frac{2\eta\pi^2}{3}}$ , where  $\delta = 1, 2, 3$  and  $\eta = 1, 2$ .

Since the procedure resembles that of Section 2, we will provide the proof only for Theorem 3.1, omitting details and presenting the value of  $l$ , the chosen denominator as in (2.7), the terms of the Taylor expansion of  $e^t$ , and the terms of the Taylor expansion of the upper and lower bounds of  $\sqrt{w^2 - \frac{2\delta\pi^2}{3}}$  and  $\sqrt{w^2 + \frac{2\eta\pi^2}{3}}$ , where  $\delta = 1, 2, 3$  and  $\eta = 1, 2$ .

**Theorem 3.1.** *For  $n \geq 301$ ,  $\Delta p(n)$  satisfies the Laguerre inequality of order 2,*

$$3(\Delta p(n+2))^2 - 4(\Delta p(n+1))(\Delta p(n+3)) + (\Delta p(n))(\Delta p(n+4)) > 0. \quad (3.1)$$

**Theorem 3.2.** *For  $n \geq 50$ ,  $\Delta\bar{p}(n)$  satisfies the Laguerre inequality of order 2,*

$$3(\Delta\bar{p}(n+2))^2 - 4(\Delta\bar{p}(n+1))(\Delta\bar{p}(n+3)) + (\Delta\bar{p}(n))(\Delta\bar{p}(n+4)) > 0. \quad (3.2)$$

*Proof of Theorem 3.1.* Setting  $l = 8$  in Lemma 1.1 yields, for  $n \geq 789$ ,

$$e^\mu \frac{\sqrt{12}\beta(\mu)\pi^2}{36\mu^{10}} < p(n) < e^\mu \frac{\sqrt{12}\alpha(\mu)\pi^2}{36\mu^{10}},$$

where

$$\alpha(t) = t^8 - t^7 + 1, \quad \beta(t) = t^8 - t^7 - 1.$$

Let

$$f(n) := e^\mu \frac{\sqrt{12}\beta(\mu)\pi^2}{36\mu^{10}}, \quad g(n) := e^\mu \frac{\sqrt{12}\alpha(\mu)\pi^2}{36\mu^{10}}.$$

Then, when we reach the step (2.7), we select  $f(n+3)^2$  as the denominator.

Using the first 7 and 8 terms of the Taylor expansion of the exponential function, we approximate the upper and lower bounds for  $t < 0$ ,

$$\phi(t) < e^t < \Phi(t),$$

where

$$\begin{aligned} \Phi(t) &= 1 + t + \frac{t^2}{2} + \frac{t^3}{6} + \frac{t^4}{24} + \frac{t^5}{120} + \frac{t^6}{720} + \frac{t^7}{5040}, \\ \phi(t) &= 1 + t + \frac{t^2}{2} + \frac{t^3}{6} + \frac{t^4}{24} + \frac{t^5}{120} + \frac{t^6}{720} + \frac{t^7}{5040} + \frac{t^8}{40320}. \end{aligned}$$

More precise estimates for  $x$ ,  $y$ ,  $z$ ,  $r$ , and  $j$  are required, as in the proof of Theorem 2.1, where

$$\begin{aligned} x &= \sqrt{w^2 - 2\pi^2}, & y &= \sqrt{w^2 - \frac{4\pi^2}{3}}, & z &= \sqrt{w^2 - \frac{2\pi^2}{3}}, \\ r &= \sqrt{w^2 + \frac{2\pi^2}{3}}, & j &= \sqrt{w^2 + \frac{4\pi^2}{3}}. \end{aligned}$$

We use the first 5 and 6 terms of the Taylor expansion of  $x$ ,  $y$ ,  $z$ ,  $r$  and  $j$  to approximate their upper and lower bounds. It can be checked that for  $n \geq 1$ ,

$$x_1 < x < x_2, \quad y_1 < y < y_2, \quad z_1 < z < z_2, \quad r_1 < r < r_2, \quad j_1 < j < j_2,$$

where

$$\begin{aligned} x_1 &= w - \frac{\pi^2}{w} - \frac{\pi^4}{2w^3} - \frac{\pi^6}{2w^5} - \frac{5\pi^8}{8w^7} - \frac{7\pi^{10}}{8w^9}, \\ x_2 &= w - \frac{\pi^2}{w} - \frac{\pi^4}{2w^3} - \frac{\pi^6}{2w^5} - \frac{5\pi^8}{8w^7} - \frac{6\pi^{10}}{8w^9}, \\ y_1 &= w - \frac{2\pi^2}{3w} - \frac{2\pi^4}{9w^3} - \frac{4\pi^6}{27w^5} - \frac{10\pi^8}{81w^7} - \frac{28\pi^{10}}{243w^9}, \\ y_2 &= w - \frac{2\pi^2}{3w} - \frac{2\pi^4}{9w^3} - \frac{4\pi^6}{27w^5} - \frac{10\pi^8}{81w^7} - \frac{27\pi^{10}}{243w^9}, \\ z_1 &= w - \frac{\pi^2}{3w} - \frac{\pi^4}{18w^3} - \frac{\pi^6}{54w^5} - \frac{5\pi^8}{648w^7} - \frac{7\pi^{16}}{1944w^9}, \end{aligned}$$

$$\begin{aligned}
z_2 &= w - \frac{\pi^2}{3w} - \frac{\pi^4}{18w^3} - \frac{\pi^6}{54w^5} - \frac{5\pi^8}{648w^7} - \frac{6\pi^{16}}{1944w^9}, \\
r_1 &= w + \frac{\pi^2}{3w} - \frac{\pi^4}{18w^3} + \frac{\pi^6}{54w^5} - \frac{5\pi^8}{648w^7}, \\
r_2 &= w + \frac{\pi^2}{3w} - \frac{\pi^4}{18w^3} + \frac{\pi^6}{54w^5} - \frac{5\pi^8}{648w^7} + \frac{7\pi^{19}}{1944w^9}, \\
j_1 &= w + \frac{2\pi^2}{3w} - \frac{2\pi^4}{9w^3} + \frac{4\pi^6}{27w^5} - \frac{10\pi^8}{81w^7}, \\
j_2 &= w + \frac{2\pi^2}{3w} - \frac{2\pi^4}{9w^3} + \frac{4\pi^6}{27w^5} - \frac{10\pi^8}{81w^7} + \frac{28\pi^{10}}{243w^9}.
\end{aligned}$$

Similar to Section 2, we observe  $A_1(w)$  is a certain polynomial of degree 68 with positive coefficient for the first term. For  $n \geq 4277$ ,  $A(w)$  is positive. Thus, for  $n \geq 4277$ , (3.1) holds. Direct calculation reveals that for  $301 \leq n \leq 4276$ ,  $\Delta p(n)$  also satisfies (3.1). The proof of Theorem 3.1 is thus completed. ■

The proof of Theorem 3.2 closely follows the structure of Theorem 3.1. Utilizing Lemma 1.2 and maintaining the notations introduced in the proof of Theorem 3.1, we adapt  $p(n)$  and  $\mu(n)$  to  $\bar{p}(n)$  and  $\bar{\mu}(n)$ , respectively, in line with Section 2. Details are omitted for brevity, and we present the final result.

Applying arguments similar to those in the proof of Theorem 3.1, we confirm that  $A_1(w)$ , a polynomial of degree 68 with a positive leading coefficient, remains positive for  $n \geq 2868$ . This establishes (3.2) for  $\Delta \bar{p}(n)$  within this range. Further numerical evidence reveals that for  $50 \leq n \leq 2867$ ,  $\Delta \bar{p}(n)$  also satisfies (3.2). Thus, the proof of Theorem 3.2 is concluded. ■

## 4 The determinantal inequalities for $\Delta p(n)$ and $\Delta \bar{p}(n)$

In this section, we investigate the determinantal inequalities of order 3 for  $\Delta p(n)$  and  $\Delta \bar{p}(n)$ , as stated in Theorem 4.1 and Theorem 4.2, respectively. The proof methodology for Theorem 4.2 follows the structure established in proving Theorem 4.1. Thus, we concentrate on a detailed proof for the former, with details analogous to those in Section 2. Here, we present only the value of  $l$ , the denominator as in (2.7), the terms of the Taylor expansion of  $e^t$ , and the terms of the Taylor expansion of the upper and lower bounds of  $\sqrt{w^2 - \frac{2\delta\pi^2}{3}}$  and  $\sqrt{w^2 + \frac{2\eta\pi^2}{3}}$ , where  $\delta = 1, 2, 3$  and  $\eta = 1, 2$ .

**Theorem 4.1.** For  $n \geq 345$ ,  $\Delta p(n)$  satisfies the determinant inequality,

$$\begin{vmatrix} \Delta p(n+2) & \Delta p(n+3) & \Delta p(n+4) \\ \Delta p(n+1) & \Delta p(n+2) & \Delta p(n+3) \\ \Delta p(n) & \Delta p(n+1) & \Delta p(n+2) \end{vmatrix} > 0. \quad (4.1)$$

**Theorem 4.2.** For  $n \geq 62$ ,  $\Delta \bar{p}(n)$  satisfies the determinant inequality,

$$\begin{vmatrix} \Delta \bar{p}(n+2) & \Delta \bar{p}(n+3) & \Delta \bar{p}(n+4) \\ \Delta \bar{p}(n+1) & \Delta \bar{p}(n+2) & \Delta \bar{p}(n+3) \\ \Delta \bar{p}(n) & \Delta \bar{p}(n+1) & \Delta \bar{p}(n+2) \end{vmatrix} > 0. \quad (4.2)$$

*Proof of Theorem 4.1.* Setting  $l = 12$  in Lemma 1.1 yields, for  $n \geq 2120$ ,

$$e^\mu \frac{\sqrt{12}\beta(\mu)\pi^2}{36\mu^{14}} < p(n) < e^\mu \frac{\sqrt{12}\alpha(\mu)\pi^2}{36\mu^{14}},$$

where

$$\alpha(t) = t^{12} - t^{11} + 1, \quad \beta(t) = t^{12} - t^{11} - 1.$$

Let

$$f(n) := e^\mu \frac{\sqrt{12}\beta(\mu)\pi^2}{36\mu^{14}}, \quad g(n) := e^\mu \frac{\sqrt{12}\alpha(\mu)\pi^2}{36\mu^{14}},$$

and choose  $f(n+1)^3$  as the denominator when we reach the step (2.7).

Denote  $\Phi(t)$  and  $\phi(t)$  as the first 11 and 12 terms of the Taylor expansion of  $e^t$ . Then, for  $t < 0$ , we have  $\phi(t) < e^t < \Phi(t)$ . We employ the first 7 and 8 terms of the Taylor expansion of  $\sqrt{w^2 - \frac{2\delta\pi^2}{3}}$  and  $\sqrt{w^2 + \frac{2\eta\pi^2}{3}}$  to approximate their upper and lower bounds, where  $\delta = 1, 2, 3$  and  $\eta = 1, 2$ , as in the proof of Theorem 3.1.

With the analogous arguments from Section 3, we determine that  $A_1(w)$  is a certain polynomial of degree 150 with a positive leading coefficient. For  $n \geq 45284$ ,  $A(w)$  is positive, leading to (4.1) for  $\Delta p(n)$ . The case for  $345 \leq n \leq 45283$  can be verified with Mathematica. Hence, the proof of Theorem 4.1 is completed.  $\blacksquare$

The proof of Theorem 4.2 closely follows the structure of Theorem 4.1. Utilizing Lemma 1.2 and maintaining notations from the proof of Theorem 4.1, we adapt  $p(n)$  and  $\mu(n)$  to  $\bar{p}(n)$  and  $\bar{\mu}(n)$ , respectively, in line with Section 2. Details are omitted for brevity, and we present the final result.

Applying arguments akin to those in the proof of Theorem 4.1, we observe that  $A_1(w)$  is a polynomial of degree 150 with a positive leading coefficient. Verification confirms  $A(w) > 0$  for  $n \geq 22275$ , establishing (4.2) for  $\Delta \bar{p}(n)$  within this range. Further numerical evidence reveals that for  $62 \leq n \leq 22274$ ,  $\Delta \bar{p}(n)$  also satisfies (4.2). Thus, the proof of Theorem 4.2 is concluded.  $\blacksquare$



## 5 Open problems

In this section, we conjecture thresholds for Laguerre inequalities of order  $m$  and the positivity of  $m$ -th order determinants. Moreover, we conjecture thresholds for the third-order Turán inequalities and the positivity of invariant  $I$ . These conjectures apply to the  $k$ -th order differences of  $p(n)$  and  $\bar{p}(n)$ , where  $1 \leq m \leq 11$  and  $1 \leq k \leq 5$ .

**Conjecture 5.1.** For  $1 \leq m \leq 11$  and  $1 \leq k \leq 5$ :

1.  $\Delta^k p(n)$  (resp.  $\Delta^k \bar{p}(n)$ ) satisfies the Laguerre inequality of order  $m$  when  $n \geq L_p(k, m)$  (resp.  $n \geq L_{\bar{p}}(k, m)$ ).
2. The  $m$ -th order determinants of (resp.  $\Delta^k \bar{p}(n)$ ) are positive for  $n \geq D_p(k, m)$  (resp.  $n \geq D_{\bar{p}}(k, m)$ ), except when both  $k \geq 3$  and  $m$  are odd.

The values of  $L_p(k, m)$ ,  $L_{\bar{p}}(k, m)$ ,  $D_p(k, m)$  and  $D_{\bar{p}}(k, m)$  are specified in Tables 1, 2, 3 and 4, respectively. The symbol  $\times$  indicates that there does not exist a value satisfying corresponding conjecture.

**Conjecture 5.2.** For  $1 \leq k \leq 5$ :

1.  $\Delta^k p(n)$  (resp.  $\Delta^k \bar{p}(n)$ ) satisfies the third-order Turán inequality when  $n \geq T_p(k)$  (resp.  $n \geq T_{\bar{p}}(k)$ ).
2.  $\Delta^k p(n)$  (resp.  $\Delta^k \bar{p}(n)$ ) satisfies  $I > 0$  when  $n \geq I_p(k)$  (resp.  $n \geq I_{\bar{p}}(k)$ ).

The values of  $T_p(k)$ ,  $T_{\bar{p}}(k)$ ,  $I_p(k)$  and  $I_{\bar{p}}(k)$  are specified in Table 5.

$m \backslash k$ $L_p(k, m)$	1	2	3	4	5	6	7	8	9	10	11
1	70	301	738	1413	2346	3557	5062	6873	9006	11467	14270
2	138	451	986	1767	2816	4151	5786	7733	10006	12613	15566
3	234	637	1272	1767	3334	4795	6562	8649	11064	13819	16922
4	362	859	1602	2609	2346	5491	7394	9619	12180	15083	18340
5	522	1121	1974	3101	4518	6241	8280	10649	13356	16411	19822

Table 1: The values of  $L_p(k, m)$ .

**Conclusion.** The proof strategy for all inequalities in this paper relies on the key Lemma 1.1 and Lemma 1.2. The process involved determining an appropriate  $l$  (resp.  $t$ ) in Lemma 1.1 (resp. Lemma 1.2), reasonably selecting the denominator as in (2.7) to ensure the negativity of the power part of  $e^t$ , truncating the terms of the Taylor expansion for  $e^t$ ,  $\sqrt{w^2 - \frac{2\delta\pi^2}{3}}$  and  $\sqrt{w^2 + \frac{2\eta\pi^2}{3}}$  (resp.  $\sqrt{w^2 - \delta\pi^2}$  and  $\sqrt{w^2 + \eta\pi^2}$ ), where  $\delta = 1, 2, \dots, m$  and  $\eta = 1, 2, \dots, m-1$ , at a suitable point depending on the Laguerre inequality of order  $m-1$  or the positivity of  $m$ -th order determinants under consideration.

$k \backslash m$ $L_{\bar{p}}(k, m)$	1	2	3	4	5	6	7	8	9	10	11
1	7	50	142	294	509	799	1167	1616	2146	2778	3497
2	21	89	208	390	641	967	1371	1859	2440	3111	3785
3	45	131	277	489	773	1126	1575	2105	2722	3435	4241
4	69	179	352	588	908	1300	1779	2345	3001	3753	4601
5	98	232	429	698	1045	1473	1985	2587	3282	4073	4963

Table 2: The values of  $L_{\bar{p}}(k, m)$ .

$k \backslash m$ $D_p(k, m)$	1	2	3	4	5	6	7	8	9	10	11
1	1	69	345	879	1709	2857	4347	6197	8419	11029	14039
2	7	137	503	1145	2091	3369	4995	6987	9359	12123	15293
3	×	233	×	1451	×	3929	×	7831	×	13275	×
4	67	361	929	1797	2997	4537	6443	8729	11405	14485	17979
5	×	521	×	2189	×	5197	×	9683	×	15755	×

Table 3: The values of  $D_p(k, m)$ .

$k \backslash m$ $D_{\bar{p}}(k, m)$	1	2	3	4	5	6	7	8	9	10	11
1	1	6	62	185	389	674	1055	1535	2120	2813	3620
2	1	20	104	257	494	821	1241	1766	2396	3134	3992
3	×	44	×	335	×	965	×	1991	×	3449	×
4	10	68	200	416	716	1115	1613	2216	2930	3758	4706
5	×	97	×	496	×	1264	×	2440	×	4066	×

Table 4: The values of  $D_{\bar{p}}(k, m)$ .

$k$	1	2	3	4	5
$T_p(k)$	174	284	424	600	810
$T_{\bar{p}}(k)$	33	57	87	118	173
$I_p(k)$	329	483	675	903	1171
$I_{\bar{p}}(k)$	64	98	143	194	255

Table 5: The values of Conjecture 5.2.

## Acknowledgments

I am grateful to Prof. Qing-Hu Hou for his dedicated guidance and to the referees for their valuable feedback. This research was financially supported by the National Natural Science Foundation of China (Grant 11921001).

## References

- [1] G. Almkvist, Exact Asymptotic Formulas for the Coefficients of Non-modular Functions, *J. Number Theory*. 38 (1991), 145–160.
- [2] G. Almkvist, On the differences of the partition function, *Acta Arith.* 61 (1992), 173–181.
- [3] K. Banerjee, Inequalities for the modified bessel function of first kind of non-negative order, *J. Math. Anal. Appl.* 524 (1) (2023), 127082.
- [4] R. Canfield, S. Corteel and P. Hitczenko, Random partitions with non-negative  $r$ -th differences, *Adv. Appl. Math.* 27 (2001), 298–317.
- [5] D.A. Cardon and A. Rich, Turán inequalities and subtraction-free expressions, *JIPAM. J. Inequal. Pure Appl. Math.* 9 (4) (2008), Artical 91. 11 pp.
- [6] W.Y.C. Chen, The  $spt$ -Function of Andrews, *Sueveys in combinatorics*, London Math. Soc. Lecture Note Ser., 440, Cambridge Univ. Press, Cambridge, 2017, 141–203.
- [7] W.Y.C. Chen, D.X.Q. Jia and L.X.W. Wang, Higher order Turán inequalities for the partition function, *Trans. Amer. Math. Soc.* 372 (3) (2019), 2143–2165.
- [8] W.Y.C. Chen, L.X.W. Wang and G.Y.B. Xie, Finite differences of the logarithm of the partition function, *Math. Comp.* 85 (2016), 825–847.
- [9] W. Craig and A. Pun, A note on the higher order Turán inequalities for  $k$ -regular partitions, *Res. Number Theory*. 7 (5) (2021), 1–7.
- [10] T. Craven and G. Csordas, Jensen polynomials and the Turán and Laguerre inequalities, *Pacific J. Math.* 136 (2) (1989), 241–260.
- [11] T. Craven and G. Csordas, Iterated Laguerre and Turán inequalities, *JIPAM. J. Inequal. Pure Appl. Math.* 3 (3) (2002), Artical 39. 14 pp.
- [12] G. Csordas, Complex zero decreasing sequences and the Riemann Hypothesis II, *Analysis and applications-ISAAC 2001 (Berlin)*, Int. Soc.

- Anal. Appl. Comput., vol. 10, Kluwer Acad. Publ., Dordrecht, 2003, 121–134.
- [13] G. Csordas, T.S. Norfolk and R.S. Varga, The Riemann hypothesis and the Turán inequalities, *Trans. Amer. Math. Soc.* 296 (2) (1986), 521–541.
  - [14] G. Csordas and R.S. Varga, Necessary and sufficient conditions and the Riemann hypothesis, *Adv. Appl. Math.* 11 (3) (1990), 328–357.
  - [15] S. DeSalvo and I. Pak, Log-concavity of the partition function, *Ramanujan J.* 38 (1) (2015), 61–73.
  - [16] B. Engel, Log-concavity of the overpartition function [J], *Ramanujan J.* 43 (2) (2017), 229–241.
  - [17] K. Dilcher and K.B. Stolarsky, On a class of nonlinear differential operators acting on polynomials, *J. Math. Anal. Appl.* 170 (2) (1992), 382–400.
  - [18] D.K. Dimitrov, Higher order Turán inequalities, *Proc. Amer. Math. Soc.* 126 (7) (1998), 2033–2037.
  - [19] D.K. Dimitrov and F.R. Lucas, Higher order Turán inequalities for the Riemann  $\xi$ -function, *Proc. Amer. Math. Soc.* 139 (3) (2011), 1013–1022.
  - [20] J.J.W. Dong and K.Q. Ji, Higher order Turán inequalities for the distinct partition function, 2023, arXiv:2303.05243.
  - [21] J.J.W. Dong, K.Q. Ji and D.X.Q. Jia, Turán inequalities for the broken  $k$ -diamond partition function, *Ramanujan J.* 62 (2) (2023), 593–615.
  - [22] L.M. Dou and L.X.W. Wang, Higher order Laguerre inequalities for the partition function, *Discret. Math.* 346 (6) (2023), 113366.
  - [23] W.H. Foster and I. Krasikov, Inequalities for real-root polynomials and entire functions, *Adv. Appl. Math.* 29 (1) (2002), 102–114.
  - [24] K. Gomez, J. Males and L. Rolén, The second shifted difference of partitions and its applications, *Bull. Aust. Math. Soc.* 107 (2023), 66–78.
  - [25] I.J. Good, The difference of the partition function, Problem 6137, *Am. Math. Mon.* 84 (1997), 141.
  - [26] M. Griffin, K. Ono, L. Rolén and D. Zagier, Jensen polynomials for the Riemann zeta function and other sequences, *Proc. Natl. Acad. Sci.* 116 (23) (2019), 11103–11110.

- [27] H. Gupta, Finite differences of the partition function, *Math. Comp.* 32 (1978), 1241–1243.
- [28] Q. Hou and Z. Zhang,  $r$ -log-concavity of partition functions, *Ramanujan J.* 48 (1) (2019), 117–129.
- [29] J.L.W.V. Jensen, Recherches sur la théorie des équations, *Acta Math.* 36 (1913), 181–195.
- [30] D.X.Q. Jia, Inequalities for the broken  $k$ -diamond partition function, *J. Number Theory.* 249 (2023), 314–347.
- [31] D.X.Q. Jia and L.X.W. Wang, Determinantal inequalities of partition function, *Proc. Royal Soc. Edinb. A.* 150 (3) (2020), 1451–1466.
- [32] C. Knessl, Asymptotic behavior of high-order differences of the plane partition function, *Discret. Math.* 126 (1994), 179–193.
- [33] C. Knessl and J.B. Keller, Partition asymptotics from recursion equations, *SIAM J. Appl. Math.* 50 (1990), 323–338.
- [34] C. Knessl and J.B. Keller, Asymptotic behavior of high-order differences of the partition function, *Commun. Pure Appl. Math.* 44 (1991), 1033–1045.
- [35] E. Laguerre, *Oeuvres*, vol.1, (Paris: Gauthier-Villars, 1898).
- [36] H. Larson and I. Wagner, Hyperbolicity of the partition Jensen polynomials, *Res. Number Theory.* 5 (19) (2019), 1–12.
- [37] D.H. Lehmer, On the series for the partition function, *Trans. Amer. Math. Soc.*, 43 (2) (1938), 271–295.
- [38] D.H. Lehmer, On the Remainders and Convergence of the Series for the Partition Function [J], *Trans. Amer. Math. Soc.* 46 (1939), 362–373.
- [39] B.J. Levin, *Distribution of zeros of entire functions*, Gosudarstv. Izdat. Tehn.-Teor. Lit., Moscow, 1956, 632 pp.
- [40] E.Y.S. Liu and H.W.J. Zhang, Inequalities for the overpartition function, *Ramanujan J.* 54 (3) (2021), 485–509.
- [41] M. Merca and J. Katriel, A general method for proving the non-trivial linear homogeneous partition inequalities, *Ramanujan J.* 51 (2020), 245–266.
- [42] J.L. Nicolas, Sur les entiers  $N$  pour lesquels il y a beaucoup de groupes abéliens d'ordre  $N$ , *Ann. Inst. Fourier.* 28 (4) (1978), 1–16.

- [43] A.M. Odlyzko, Differences of the partition function, *Acta Arith.* 49 (1988), 237–254.
- [44] M.L. Patrick, Some inequalities concerning Jacobi polynomials, *SIAM J. Math. Anal.* 2 (2) (1971), 213–220.
- [45] M.L. Patrick, Extensions of inequalities of the Laguerre and Turán type, *Pacific J. Math.* 44 (2) (1973), 675–682.
- [46] G. Pólya, Über die algebraisch-funktionentheoretischen Untersuchungen von J.L.W.V. Jensen, *Kgl. Danske Vid. Sel. Math.-Fys. Medd.* 7 (1927), 3–33.
- [47] Q.I. Rahman and G. Schmeisser, *Analytic theory of polynomials*, Oxford University Press, Oxford, 2002. xiv+742 pp.
- [48] H. Skovgaard, On inequalities of the Turán type, *Math. Scand.* 2 (1954), 65–73.
- [49] I. Wagner, On a new class of Laguerre-Pólya type functions with applications in number theory, *Pacific J. Math.* 320 (1) (2022), 177–192.
- [50] L.X.W. Wang, G.Y.B. Xie and A.Q. Zhang, Finite difference of the overpartition function, *Adv. Appl. Math.* 92 (2018), 51–72.
- [51] L.X.W. Wang and E.Y.Y. Yang, Laguerre inequalities for discrete sequences, *Adv. Appl. Math.* 139 (2022), 102357.
- [52] L.X.W. Wang and N.N.Y. Yang, Positivity of the determinants of the partition function and the overpartition function, *Math. Comput.* 341 (92) (2023), 1383–1402.
- [53] L.X.W. Wang and E.Y.Y. Yang, Laguerre inequality and determinantal inequality for the distinct partition function, submitted.
- [54] E.Y.Y. Yang, Finite differences of the distinct partition function and the broken  $k$ -diamond partition function, submitted.
- [55] E.Y.Y. Yang, Laguerre inequality and determinantal inequality for the broken  $k$ -diamond partition function, submitted.
- [56] H.S. Zuckerman, On the coefficients of certain modular forms belonging to subgroups of the modular group. *Trans. Am. Math. Soc.* 45 (2) (1939), 298–321.