# A SEMIDEFINITE PROGRAMMING HIERARCHY FOR COVERING PROBLEMS IN DISCRETE GEOMETRY

CORDIAN RIENER, JAN ROLFES, AND FRANK VALLENTIN

ABSTRACT. In this paper we present a new semidefinite programming hierarchy for covering problems in compact metric spaces.

Over the last years, these kind of hierarchies were developed primarily for geometric packing and for energy minimization problems; they frequently provide the best known bounds.

Starting from a semidefinite programming hierarchy for the dominating set problem in graph theory, we derive the new hierarchy for covering and show some of its basic properties: The hierarchy converges in finitely many steps, but the first level collapses to the volume bound when the compact metric space is homogeneous.

### 1. INTRODUCTION

Let (X, d) be a compact metric space. Frequently one asks for specific configurations of finitely many points in X. Various performance measures for such finite point distributions are investigated. Homogeneous measures, like maximizing minimal distance between pairs of distinct points (*packing*), or minimizing energy with respect to a potential function (*energy minimization*), are most commonly used. These homogeneous measures have inhomogeneous counterparts: Finding a point configuration which minimizes the maximal distance of an arbitrary point in X to the finite point distribution (*covering*), or maximizing the minimal potential energy of an arbitrary point in X in the force field given by the charges of the point configuration (*max-min polarization*). For a more detailed overview, we refer to the book of Borodachov, Hardin, Saff [4].

Over the last years, semidefinite programming hierarchies, in the spirit of the moment-SOS hierarchy in polynomial optimization, due to Lasserre, were developed primarily for providing bounds for geometric packing and for energy minimization problems; see de Laat, Vallentin [9], and de Laat [11]. Computing these semidefinite programming bounds frequently yield the best known bounds.

In this paper we want to demonstrate that this approach is also applicable to geometric covering problems.

We start by defining the covering problem formally. Let X be a compact metric space with distance function d. We denote closed metric balls of radius r and center x by  $\overline{B}(x,r) = \{y \in X : d(x,y) \leq r\}$ . A set of balls of radius r is uniquely determined by the collection of its centers Y. A set Y of centers with the property  $\bigcup_{y \in Y} \overline{B}(y,r) = X$  is called a *covering* of X. Since X is compact, there always exists a finite covering. The *covering number*  $\mathcal{N}(X,r)$  of the space X and a positive

Date: December 18, 2023.

number r is the smallest number of such balls with radius r one needs to cover X. So,

$$\mathcal{N}(X,r) = \min\left\{ |Y| : Y \subseteq X, \bigcup_{y \in Y} \overline{B}(y,r) = X \right\}.$$

Determining the covering number is a fundamental problem in metric geometry (see for example the classical book by Rogers [22]). Among others, applications arise in the field of compressive sensing [13], approximation theory and machine learning [7], in probability theory [16] and theoretical quantum computing [20].

So far, upper bounds for the covering number of several specific metric spaces are known. For a survey on this, we refer to Naszódi [19]. Using a greedy approach, in the spirit Chvátal's greedy approximation algorithm for the set-covering problem, Rolfes and Vallentin [23] provide upper bounds for a wider class of compact metric spaces. They showed that for every  $\varepsilon$  with  $r/2 > \varepsilon > 0$  the inequalities

(1) 
$$\frac{1}{\omega_r} \le \mathcal{N}(X, r) \le \frac{1}{\omega_{r-\varepsilon}} \left( \ln \left( \frac{\omega_{r-\varepsilon}}{\omega_{\varepsilon}} \right) + 1 \right),$$

holds, where X is equipped with a probability measure  $\omega$  satisfying the following two conditions:

- (a)  $\omega(\overline{B}(x,s)) = \omega(\overline{B}(y,s))$  for all  $x, y \in X$ , and for all  $s \ge 0$ ,
- (b)  $\omega(\overline{B}(x,\varepsilon)) > 0$  for all  $x \in X$ , and for all  $\varepsilon > 0$ .

By (a) the measure of a ball does only depend on the radius s and not on the center x, so we simply denote  $\omega(\overline{B}(x,s))$  by  $\omega_s$ . These conditions are fulfilled when X is a *homogeneous* space, that is, when the automorphism group acts transitively. Then the Haar measure of the automorphism group induces an invariant probability measure on X satisfying (a) and (b).

The trivial lower bound above is known as the *volume bound*. Stronger and more sophisticated lower bounds have not yet been established. The central aim of this paper is to develop a semidefinite programming hierarchy for geometric covering problems giving such lower bounds.

We start by motivating our definition of the covering hierarchy by recalling the Lasserre hierarchy for general 0/1 integer linear programs in Section 2. Then we also recall the packing hierarchy of de Laat, Vallentin [9] in Section 3. The covering hierarchy is based on the packing hierarchy. We will define our covering hierarchy in Section 4 and show its basic properties in Theorem 4.2, which is the principal result of the paper. In Section 5 we derive the dual covering hierarchy and use the dual to show that for homogeneous compact metric spaces the first step of the hierarchy collapses to the volume bound.

## 2. The Lasserre hierarchy for 0/1 integer linear programs

Consider the following integer linear program with 0/1-variables

$$ILP = \min\{c^{\mathsf{T}}x : x \in \{0, 1\}^n, \ Ax \ge b\},\$$

where the matrix  $A \in \mathbb{R}^{m \times n}$  and the vectors  $b \in \mathbb{R}^m$  and  $c \in \mathbb{R}^n$  are given as input.

In the by now classical paper [14] Lasserre showed how to systematically compute a sequence of stronger and stronger lower bounds, converging in n + 1 steps to ILP by solving semidefinite programs of growing size. Lasserre's original convergence proof relies on the Positivstellensatz of Putinar [21] and on the flat extension theorem of Curto, Fialkow [8]. Laurent [15] provided an elementary, combinatorial proof of the convergence. We shall reproduce her arguments below because they gave the inspiration for the semidefinite hierarchies for geometric packing and covering problems. For good behavior we note that the results of Lasserre and Laurent are much more general, as they apply to nonlinear (polynomial) optimization problems with 0/1-variables.

2.1. Combinatorial moment and localizing matrices. To define the Lasserre hierarchy for solving ILP we need a bit of notation.

Set  $X = [n] = \{1, ..., n\}$ . We write  $\mathcal{P}(X)$  for the power set of X. By  $\mathcal{P}_t(X)$  we denote the subsets of  $\mathcal{P}(X)$  containing all  $J \in \mathcal{P}(X)$  with  $|J| \leq t$ . For t = 0, ..., n and  $J \subseteq X$  we define the *(truncated) characteristic vector*  $\chi_t^J \in \{0, 1\}^{\mathcal{P}_t(X)}$  by

$$\chi_t^J(J') = \begin{cases} 1 & \text{if } J' \subseteq J, \\ 0 & \text{otherwise} \end{cases}$$

A vector  $y \in \mathbb{R}^{\mathcal{P}_{2t}(X)}$  defines the *(truncated) combinatorial moment matrix*  $M_t(y) \in \mathcal{S}^{\mathcal{P}_t(X)}$ , which is a symmetric matrix with  $\sum_{i=0}^t \binom{n}{i}$  many rows and columns, by

$$[M_t(y)]_{J,J'} = y(J \cup J'), \quad \text{with } J, J' \in \mathcal{P}_t(X).$$

The following lemma dealing with t = n, the final step of Lasserre's hierarchy, is crucial. It is due to Lindström [17] and Wilf [27].

**Lemma 2.1.** A vector  $y \in \mathbb{R}^{\mathcal{P}(X)}$  defines a positive semidefinite combinatorial moment matrix  $M_n(y)$  if and only if y lies in the (simplicial) polyhedral cone generated by the characteristic vectors  $\chi_n^J$ , with  $J \in \mathcal{P}(X)$ . In short:

$$M_n(y) \succeq 0 \iff y \in \operatorname{cone}\{\chi_n^J : J \in \mathcal{P}(X)\}.$$

*Proof.* Sufficiency follows easily by the homomorphism property of the characteristic vectors, which is

$$\chi_n^J(J'\cup J'') = \chi_n^J(J')\cdot \chi_n^J(J'') \text{ for all } J, J', J'' \in \mathcal{P}(X).$$

Now any y with

$$y = \sum_{J \in \mathcal{P}(X)} \alpha_J \chi_n^J \text{ and } \alpha_J \ge 0$$

defines a positive semidefinite combinatorial moment matrix

$$M_n(y) = \sum_{J \in \mathcal{P}(X)} \alpha_J \chi_n^J (\chi_n^J)^{\mathsf{T}}.$$

For necessity, we first note that the characteristic vectors  $\chi_n^J$  form a basis of  $\mathbb{R}^{\mathcal{P}(X)}$ . Let  $\psi_n^J$  be the dual basis with respect to the standard inner product, so that

$$(\chi_n^J)^{\mathsf{T}}\psi_n^{J'} = \delta_{J,J'}, \text{ where } \delta_{J,J'} = \begin{cases} 1 & \text{if } J = J' \\ 0 & \text{otherwise.} \end{cases}$$

Now consider a vector  $y \in \mathbb{R}^{\mathcal{P}(X)}$  which defines a positive semidefinite combinatorial moment matrix  $M_n(y)$ . Expand

$$y = \sum_{J \in \mathcal{P}(X)} \alpha_J \chi_n^J,$$

and so

$$0 \le (\psi_n^J)^{\mathsf{T}} M_n(y) \psi_n^J = \alpha_J.$$

Consider the *j*-th row of the system  $Ax \geq b$  which is the constraint  $a_j^{\mathsf{T}}x \geq b_j$ , where the *j*-th row  $a_j$  of matrix A is viewed as a column vector. Using this information we can define the *j*-th *(truncated) localizing matrix*  $M_t^j(y)$  of  $y \in \mathbb{R}^{\mathcal{P}_{2t+2}(X)}$  by

$$[M_t^j(y)]_{J,J'} = \sum_{i \in X} a_{ji} y(J \cup J' \cup \{i\}) - b_j y(J \cup J'),$$

where  $J, J' \in \mathcal{P}_t(X)^1$ .

2.2. The Lasserre hierarchy for ILP. For t = 1, ..., n, the *t*-th step of the Lasserre hierarchy for ILP is the (value of the) following semidefinite program

$$las_t = \min\left\{\sum_{i \in X} c_i y(\{i\}) : y \in \mathbb{R}^{\mathcal{P}_{2t}(X)}, \ y(\emptyset) = 1, \\ M_t(y) \succeq 0, \ M_{t-1}^j(y) \succeq 0 \ (j \in [m])\right\}.$$

The Lasserre hierarchy provides a monotonically increasing sequence of lower bounds for ILP:

$$ILP \ge las_{n+1} \ge las_n \ge \ldots \ge las_1$$
.

The sequence is monotonically increasing because every feasible solution  $y \in \mathbb{R}^{\mathcal{P}_{2t}(X)}$  of  $\operatorname{las}_t$  can be truncated to a feasible solution  $y' \in \mathbb{R}^{\mathcal{P}_{2t-2}(X)}$  of  $\operatorname{las}_{t-1}$  because  $M_{t-1}(y')$  and  $M_{t-2}^j(y')$  are principal submatrices of  $M_t(y)$  and  $M_{t-1}^j(y)$ .

We have ILP  $\geq \log_{n+1}$  because if  $x^* \in \{0,1\}^n$  is an optimal solution of ILP, then  $y = \chi_n^J$  with  $J = \{i \in X : x_i^* = 1\}$  is feasible for  $\log_{n+1}$ ; Positive semidefiniteness of  $M_n(y)$  follows from Lemma 2.1 and positive semidefiniteness of  $M_n^j(y)$  is again due to the homomorphism property of the characteristic vector

$$\begin{split} [M_n^j(y)]_{J',J''} &= \sum_{i \in X} a_{ji} \chi_n^J (J' \cup J'' \cup \{i\}) - b_j \chi_n^J (J' \cup J'') \\ &= \chi_n^J (J' \cup J'') \left( \sum_{i \in X} a_{ji} \chi_n^J (\{i\}) - b_j \right) \\ &= \chi_n^J (J' \cup J'') \left( \sum_{i \in X} a_{ji} x_i^* - b_j \right), \end{split}$$

and hence

$$M_n^j(y) = \left(a_j^{\mathsf{T}} x^* - b_j\right) \chi_n^J(\chi_n^J)^{\mathsf{T}} \succeq 0,$$

because  $a_j^\mathsf{T} x^* \ge b_j$ .

Moreover, the Lasserre hierarchy converges to ILP in (at most) n + 1 steps because the reverse inequality ILP  $\leq |as_{n+1}|$  holds. This can be seen by again applying Lemma 2.1, which says that we can expand any feasible solution y of  $|as_{n+1}|$  as  $y = \sum_{J \in \mathcal{P}(X)} \alpha_J \chi_n^J$  with nonnegative coefficients  $\alpha_J$ . If  $\alpha_J > 0$ , then  $x^J \in \{0,1\}^n$  with  $x_i^J = \chi_n^J(\{i\})$  is a feasible solution of ILP. This follows from

4

<sup>&</sup>lt;sup>1</sup>In fact, the entries of y which are indexed by 2t + 2-element subsets of X are not used here. We keep them because the notation for the Lasserre hierarchy becomes simpler.

the positive semidefiniteness of the localizing matrix. In particular, using the dual basis  $\psi_n^J$ ,

$$0 \le (\psi_n^J)^\mathsf{T} M_n^j(y) \psi_n^J = \alpha_J \left( \sum_{i \in X} a_{ji} \chi_n^J(\{i\}) - b_j \right)$$

holds, which implies  $a_j^{\mathsf{T}} x^J \ge b_j$ . The normalization  $y(\{\emptyset\}) = 1$  translates into  $\sum_{J \in \mathcal{P}(X)} \alpha_J = 1$  and therefore the objective value of  $\operatorname{las}_{n+1}$  satisfies

$$\sum_{i \in X} c_i y(\{i\}) = \sum_{i \in X} c_i \sum_{J \in \mathcal{P}(X)} \alpha_J \chi_n^J(\{i\}) = \sum_{J \in \mathcal{P}(X)} \alpha_J c^\mathsf{T} x^J \ge \mathrm{ILP} \,.$$

### 3. SDP HIERARCHY FOR GEOMETRIC PACKING PROBLEMS

Let X be a compact metric space with distance function d. By  $B(x,r) = \{y \in X : d(x,y) < r\}$  we denote open metric balls with radius r and center x.

One important geometric parameter of X and a positive r is the *packing number* 

$$\alpha(X,r) = \max\{|Y| : Y \subseteq X : B(x,r) \cap B(y,r) = \emptyset \text{ for all } x, y \in Y, x \neq y\}.$$

The idea behind the paper [9] of de Laat and Vallentin is to consider  $\alpha(X, r)$  as an infinite integer linear program where every point in X corresponds to a 0/1-variable. Then an appropriate semidefinite programming hierarchy is constructed which can be used to determine upper bounds for  $\alpha(X, r)$  and which eventually converges to  $\alpha(X, r)$ . In this section we recall the main steps of [9] because our semidefinite hierarchy for covering builds on this packing hierarchy. Whereas the packing hierarchy only uses moment constraints, the covering hierarchy needs to incorporate localizing constraints.

The packing number can also be seen as the independence number of the, potentially infinite, packing graph P(X,r) with vertex set X in which two distinct vertices x, y are adjacent whenever  $B(x,r) \cap B(y,r) \neq \emptyset$ . Generally, the *independence number* of a graph is the maximum cardinality of an independent set, where a subset of the vertex set is called an *independent set* if it does not contain a pair of adjacent vertices.

For a finite graph G = (V, E), with vertex set  $V = \{1, ..., n\}$  and edge set E, one can formulate the independence number of G as an integer linear program with 0/1-variables, which is

$$\alpha(G) = \max\left\{\sum_{i=1}^{n} x_i : x \in \{0,1\}^n, x_i + x_j \le 1 \text{ for all } \{i,j\} \in E\right\},\$$

and apply the Lasserre hierarchy.

Laurent [15, Lemma 20] observed that one can write the Lasserre hierarchy for  $\alpha(G)$  more compactly, only by considering combinatorial moment matrices, without using localizing matrices. To state this compact version of the Lasserre hierarchy, we slightly change the definitions introduced in Section 2. We hope that this will not cause confusion. The first change is that instead of working with the complete power set of the vertex set we consider only independent subsets. By  $I_t$  we denote the set of all independent sets with at most t elements, then the t-th step of the Lasserre hierarchy is

(2) 
$$las_t^{\alpha}(G) = max\left\{\sum_{i=1}^n y(\{i\}) : y \in \mathbb{R}^{I_{2t}}_{\geq 0}, \ y(\emptyset) = 1, \ M_t(y) \succeq 0\right\},$$

where we define  $M_t(y) \in \mathbb{R}^{I_t \times I_t}$  by

$$[M_t(y)]_{J,J'} = \begin{cases} y_{J\cup J'} & \text{if } J \cup J' \in I_{2t}, \\ 0 & \text{otherwise,} \end{cases}$$

which is the second change. The third and last change, requiring y to be a nonnegative vector, enables us to say that the first step  $las_1^{\alpha}(G)$  coincides with  $\vartheta'(G)$ , the strengthened version of Lovász  $\vartheta$ -number [18] due to Schrijver [25]. We obtain

$$\vartheta'(G) = \operatorname{las}_1^{\alpha}(G) \ge \operatorname{las}_2^{\alpha}(G) \ge \ldots \ge \operatorname{las}_{\alpha(G)}^{\alpha}(G) = \alpha(G).$$

In [9] this semidefinite programming hierarchy (2) is generalized to give a hierarchy for the packing number  $\alpha(X, r)$ . In particular, independence numbers of topological packing graphs are introduced, which include the packings graphs P(X, r) on compact metric spaces.

Since X can have infinitely many points, some topology is needed to generalize the objects in (2). The natural distance on nonempty subsets of X is the Hausdorff distance

$$d_H(J, J') = \inf\{\varepsilon : J \subseteq J'_{\varepsilon} \text{ and } J' \subseteq J_{\varepsilon}\},\$$

where  $J_{\varepsilon} = \bigcup_{x \in J} B(x, \varepsilon)$  is the  $\varepsilon$ -thickening of the set J.

Using the Hausdorff distance on  $I_t$  instead of  $\mathcal{P}_t(X)$  has an important advantage. By considering  $\alpha(X, r)$  we are interested in cardinalities and generally elements in  $\mathcal{P}_t(X)$  with non-equal cardinality can be topologically close. However, because we work with  $I_t$  and not with  $\mathcal{P}_t(X)$ , independent sets having different cardinality, lie in separate connected components; see the discussion in [9, Section 2].

In the generalization, the vector  $y \in \mathbb{R}_{\geq 0}^{I_{2t}}$  is replaced by a measure  $\lambda \in \mathcal{M}(I_{2t})_{\geq 0}$ in the cone of non-negative Radon measure. The idea is that an independent set S determines a feasible solution

$$\lambda = \sum_{R \in I_{2t}: R \subseteq S} \delta_R, \quad \text{where } \delta_R \text{ is the delta measure at } R,$$

of the *t*-th step of the packing hierarchy.

The map which assigns y the truncated combinatorial moment matrix  $y \mapsto M_t(y)$  is replaced by a dual construction using the adjoint map.

By the Riesz representation theorem the dual space of the space of real-valued continuous functions  $C(I_{2t})$ , equipped with the supremum norm, is the space of signed Radon measures  $\mathcal{M}(I_{2t})$ . The dual cone of nonnegative continuous functions we denote by  $\mathcal{M}(I_{2t})_{\geq 0}$ . Similarly, the dual space of symmetric, real-valued continuous kernels  $C(I_t \times I_t)_{\text{sym}}$  is the space of symmetric Radon measures  $\mathcal{M}(I_t \times I_t)_{\text{sym}}$ . The dual cone of positive definite kernels, that is kernels  $K \in C(I_t \times I_t)_{\text{sym}}$  which satisfy

$$(K(J_i, J_j))_{i,j=1}^m \succeq 0 \quad \text{for all } J_1, \dots, J_m \in I_{2t}, \ m \in \mathbb{N},$$

we denote by  $\mathcal{M}(I_t \times I_t)_{\succ 0}$ .

We define the map

$$A_t: \mathcal{C}(I_t \times I_t)_{\text{sym}} \to \mathcal{C}(I_{2t}) \quad \text{by} \quad A_t K(S) = \sum_{J, J' \in I_t: J \cup J' = S} K(J, J'),$$

where  $K \in \mathcal{C}(I_t \times I_t)_{\text{sym}}$  and  $S \in I_{2t}$  Note that the above sum has at most  $2^{2t}$  summands. Therefore, the adjoint map

$$A_t^* : \mathcal{M}(I_{2t}) \to \mathcal{M}(I_t \times I_t)_{\text{sym}},$$

(well-)defined by

$$\langle K, A^*\lambda \rangle = \langle A_t K, \lambda \rangle,$$

is the natural replacement for  $y \mapsto M_t(y)$ .

After these preparations one can define the *t*-th step of the *packing hierarchy* by

$$\operatorname{las}_{t}^{\alpha}(P(X,r)) = \max\{\lambda(I_{=1}) : \lambda \in \mathcal{M}(I_{2t})_{\geq 0}, \ \lambda(\{\emptyset\}) = 1, \ A_{t}^{*}(\lambda) \in \mathcal{M}(I_{t} \times I_{t})_{\geq 0}, \}$$

where  $I_{=1}$  denotes the set of independent subsets with cardinality exactly 1. Again, and this is the main result of [9],

$$\vartheta'(P(X,r)) = \operatorname{las}_1^{\alpha}(P(X,r)) \ge \operatorname{las}_2^{\alpha}(P(X,r)) \ge \ldots \ge \operatorname{las}_{\alpha(X,r)}^{\alpha}(P(X,r)) = \alpha(X,r)$$

holds.

The first step coincides with the generalization of the  $\vartheta'$ -number for the graph P(X, r), defined in [2]. If for instance the compact metric space is the unit sphere,  $X = S^{n-1}$ , then the first step coincides with the linear programming bound of Delsarte, Goethals, Seidel [10].

The convergence follows from a Choquet-type variant (cf. Simon [26]) of the crucial Lemma 2.1, where finite convex combinations are replaced by integrals with respect to probability measures, which reads:

**Lemma 3.1.** (de Laat, Vallentin [9, Proposition 1]) Let  $I = I_{\alpha(X,r)}$  be the set of all independent sets of P(X,r). Let  $\lambda \in \mathcal{M}(I)$  be so that  $\lambda(\{\emptyset\}) = 1$  and  $A^*_{\alpha(X,r)}\lambda \in \mathcal{M}(I \times I)_{\geq 0}$ . Then there exists a unique probability measure  $\sigma \in \mathcal{M}(I)_{\geq 0}$ so that  $\lambda$  can be represented as

$$\lambda = \int \chi^R \, d\sigma(R), \quad where \quad \chi^R = \sum_{Q \subseteq R} \delta_Q.$$

Next to putting the linear programming bound into perspective and giving theoretical convergence, the packing hierarchy has turned out to be useful in various applications, especially for the unit sphere or for Euclidean space where the presence of symmetries makes it possible to simplify the computations. Bachoc, Vallentin [1] used semidefinite constraints from  $las_{2}^{\alpha}$  to find new upper bounds for the kissing number. de Laat [11] used the second step of the hierarchy in the context of energy minimization on the sphere  $S^{2}$ . de Laat, Machado, de Munick Keizer [12] computed  $las_{2}^{\alpha}$  and  $las_{3}^{\alpha}$  for the problem of equiangular lines; they were even able to turn these computations in an asymptotic analysis. Cohn, Salmon [6] showed that the packing hierarchy converges to the packing density for Euclidean space. Cohn, de Laat, Salmon [5] used constraints from  $las_{2}^{\alpha}$  to find new upper bounds for sphere packing densities in dimensions 4 through 7 and 9 through 16.

## 4. SDP HIERARCHY FOR GEOMETRIC COVERING PROBLEMS

The finite graph analog of the covering number is the domination number. For a finite graph G = (V, E), with vertex set  $V = \{1, \ldots, n\}$  and edge set E a subset of the vertex set  $D \subseteq V$  is called a *dominating set* if for every vertex  $i \in V$  either the vertex lies itself in the dominating set or there is vertex  $j \in D$ , in the dominating set, which is adjacent to i. The *domination number*  $\gamma(G)$  of G is the cardinality of a smallest dominating set. One can formulate  $\gamma(G)$  as an integer linear program

with 0/1-variables:

$$\gamma(G) = \min\left\{\sum_{i=1}^{n} x_i : x \in \{0,1\}^n, \sum_{i \in V : \{i,j\} \in E} x_i \ge 1 \ (j \in V)\right\},\$$

and apply the Lasserre hierarchy. Here the t-th step of the Lasserre hierarchy is

$$las_t^{\gamma}(G) = \min\left\{ \sum_{i=1}^n y(\{i\}) : y \in \mathbb{R}^{\mathcal{P}_{2t}(V)}, y(\emptyset) = 1, \\ M_t(y) \succeq 0, \ M_{t-1}^j(y) \succeq 0 \ (j \in V) \right\}$$

where the linear constraint that every vertex  $j \in V$  either should lie in the dominating set or should be adjacent to a vertex in the dominating set determines a truncated localizing matrix

$$[M_{t-1}^{j}(y)]_{J,J'} = \sum_{i \in V: \{i,j\} \in E} y(J \cup J' \cup \{i\}) - y(J \cup J'),$$

where  $J, J' \in \mathcal{P}_{t-1}(V)$ .

In the geometric setting we consider the covering graph C(X, r) with vertex set X in which two vertices  $x, y \in X$  are adjacent whenever  $d(x, y) \leq r$ . Then, as dominating sets in C(X, r) determine the centers of balls with radius r of a covering of X, and vice versa, we have  $\gamma(C(X, r)) = \mathcal{N}(X, r)$ .

We want to define a semidefinite hierarchy which converges in finitely many steps to the covering number  $\mathcal{N}(X, r)$ . For the proof we will make use of Lemma 3.1. Therefore, we have to assume that we know a lower bound  $\varepsilon > 0$  so that an optimal covering X determines a packing of balls with radius  $\varepsilon$ . Clearly such a positive lower bound always exists as optimal coverings consist of finitely many pairwise distinct balls. We define  $I_t$  to be the set of subsets of X with at most t elements which are independent in the packing graph  $P(X, \varepsilon)$ .

For  $y \in X$  we define the map

$$B_t^y : \mathcal{C}(I_t \times I_t)_{\text{sym}} \to \mathcal{C}(I_{2t+2})$$

by

$$B^y_tK(S) = \sum_{x\in \overline{B}(y,r),J,J':J\cup J'\cup \{x\}=S} K(J,J') - \sum_{J,J'\in I_t:J\cup J'=S} K(J,J'),$$

where  $K \in \mathcal{C}(I_t \times I_t)_{\text{sym}}$  and  $S \in I_{2t+2}$ . The adjoint map  $(B_t^y)^* : \mathcal{M}(I_{2t+2}) \to \mathcal{M}(I_t \times I_t)_{\text{sym}}$  gives the analog of the map which assigns y to the truncated combinatorial moment matrix.

**Definition 4.1.** For t = 1, 2, ..., we define the t-th step of the covering hierarchy by

$$\begin{aligned} \operatorname{las}_{t}^{\gamma}(C(X,r)) &= \inf \left\{ \lambda(I_{=1}) : \lambda \in \mathcal{M}(I_{2t})_{\geq 0}, \ \lambda(\{\emptyset\}) = 1, \\ A_{t}^{*}(\lambda) \in \mathcal{M}(I_{t} \times I_{t})_{\succeq 0}, \\ (B_{t-1}^{y})^{*}(\lambda) \in \mathcal{M}(I_{t-1} \times I_{t-1})_{\succeq 0} \ (y \in X) \right\} \end{aligned}$$

**Theorem 4.2.** The covering hierarchy gives a monotonically increasing sequence of lower bounds for the covering number and converges to the covering number in at most  $\alpha(P(X, \varepsilon))$  steps:

$$\operatorname{las}_{1}^{\gamma}(C(X,r)) \leq \operatorname{las}_{2}^{\gamma}(C(X,r)) \leq \ldots \leq \operatorname{las}_{\alpha(P(X,\varepsilon))}^{\gamma}(C(X,r)) = \mathcal{N}(X,r).$$

We split the proof of this theorem in three lemmas.

# 4.1. Every step of the covering hierarchy gives a lower bound for the covering number.

**Lemma 4.3.** For every t = 1, 2, ... the inequality  $\operatorname{las}_t^{\gamma}(C(X, r)) \leq \mathcal{N}(X, r)$  holds.

*Proof.* We verify this inequality by showing that every finite covering Y of X gives a feasible solution

$$\lambda = \sum_{Z \in I_{2t}: \ Z \subseteq Y} \delta_Z$$

of the *t*-step of the covering hierarchy with objective  $\lambda(I_{=1}) = |Y|$ . Indeed, we have

$$\lambda\left(\{\emptyset\}\right) = \sum_{Z \in I_{2t}: Z \subseteq Y} \delta_Z\left(\{\emptyset\}\right) = \delta_{\emptyset}\left(\{\emptyset\}\right) = 1.$$

Then, for every  $K \in \mathcal{C}(I_t \times I_t)_{\succ 0}$ :

$$\begin{split} \langle A_t^* \lambda, K \rangle &= \langle \lambda, A_t K \rangle \\ &= \sum_{Z \in I_{2t}: Z \subseteq Y} \sum_{J, J' \in I_t: Z = J \cup J'} K(J, J') \\ &= \sum_{J, J' \in I_t: J, J' \subseteq Y} K(J, J') \\ &\geq 0, \end{split}$$

as well as, for every  $y \in X$  and every  $K' \in \mathcal{C}(I_{t-1} \times I_{t-1})_{\succeq 0}$ 

$$\begin{split} &\langle (B_{t-1}^{y})^{*}\lambda, K' \rangle \\ &= \langle \lambda, B_{t-1}^{y}K' \rangle \\ &= \sum_{Z \in I_{2t}: Z \subseteq Y} \left( \sum_{x \in \overline{B}(y,r), J, J': Z = J \cup J' \cup \{x\}} K'(J,J') - \sum_{J,J': Z = J \cup J'} K'(J,J') \right) \\ &= \sum_{x \in Y \cap \overline{B}(y,r)} \sum_{J,J': J, J' \subseteq Y} K'(J,J') - \sum_{J,J': J, J' \subseteq Y} K'(J,J') \\ &= \left( |Y \cap \overline{B}(y,r)| - 1 \right) \sum_{J,J' \in I_{t-1}: J, J' \subseteq Y} K'(J,J') \\ &> 0. \end{split}$$

where the last inequality is implied by the fact that  $Y \cap \overline{B}(y,r) \neq \emptyset$  since Y is a covering.

### 4.2. The covering hierarchy is monotonically increasing with t.

**Lemma 4.4.** For every t = 1, 2, ... the inequality  $\operatorname{las}_t^{\gamma}(C(X, r)) \leq \operatorname{las}_{t+1}^{\gamma}(C(X, r))$  holds.

Proof. Increasing t strengthens the bound by imposing more constraints to the set of feasible solutions. To be precise, if the measure  $\lambda \in \mathcal{M}(I_{2t+2})_{\geq 0}$  is feasible for  $\operatorname{las}_{t+1}^{\gamma}(C(X,r))$ , then its restriction to  $\mathcal{M}(I_{2t})$  is also feasible for  $\operatorname{las}_{t}^{\gamma}(C(X,r))$ . Furthermore,  $\lambda$  and its restriction have the same objective value, namely  $\lambda(I_{=1})$ .  $\Box$  4.3. The covering hierarchy converges to the covering number. Before we can prove this, we need a technical, topological observation.

We equip the non-empty sets in  $I_t$  with the Hausdorff distance

$$d_H(Y,Z) = \max\left\{\sup_{y\in Y}\inf_{z\in Z}d(y,z),\sup_{z\in Z}\inf_{y\in Y}d(y,z)\right\}$$

and extend this distance to a metric

$$d_H(Y,Z) = \begin{cases} d_H(Y,Z) & \text{if } Y, Z \neq \emptyset \\ 0 & \text{if } Y = Z = \emptyset \\ \sup_{x,y \in X} d(x,y) & \text{otherwise,} \end{cases}$$

which also takes care of  $\emptyset \in I_t$ .

**Lemma 4.5.** Let  $B = \overline{B}(x, r)$  be a ball in X. Consider the function  $k : I_t \to \mathbb{R}$  defined by

$$k(J) = \begin{cases} (-1)^{|J|} & \text{if } J \subseteq B\\ 0 & \text{otherwise} \end{cases}$$

Then there exists a continuous function  $g \in C(I_t)$  such that k and g agree  $\nu$ -almost surely for every  $\nu \in \mathcal{M}(I_t)$ .

*Proof.* First we observe that one can write k in terms of indicator functions

$$k(J) = \left(\sum_{r=0, r \text{ even}}^{|J|} \mathbf{1}_{I_{=r}}(J) - \sum_{r=0, r \text{ odd}}^{|J|} \mathbf{1}_{I_{=r}}(J)\right) \mathbf{1}_{B}(J).$$

Due to the continuity of  $\mathbf{1}_{I=r}$ , given by [9, Lemma 2], it suffices to show that there is a continuous function g such that  $g(J) = \mathbf{1}_B(J)$ ,  $\nu$ -almost surely.

We further consider  $\{J \in I_t : J \nsubseteq B\}$  and show that it is an open set in  $I_t$ . Because  $J \nsubseteq B$ , implying  $J \neq \emptyset$ , there is an element  $y_0 \in J \setminus B$ . Due to the Hausdorff property of X we have  $d(y_0, b) > \delta$  for every  $b \in B$ . Thus we can consider the ball  $B(J, \delta)$  defined by the Hausdorff metric and show that  $d_H(B, B(J, \delta)) \ge \sup_{y \in B(J,\delta)} \inf_{b \in B} d(b, y) \ge \inf_{b \in B} d(b, y_0) > 0$  holds. This implies that  $B(J, \delta) \subseteq \{J \in I_t : J \nsubseteq B\}$  and thus

 $C = \{J \in I_t : J \nsubseteq B\}$  is open and  $D = \{J \in I_t : J \subseteq B\}$  is closed.

As a metric space  $(I_t, d_H)$  is a normal space and thus we can apply Urysohn's lemma, which gives us that for any compact subset of C, say C' and D there exists a continuous function g with g(c) = 0 for every  $c \in C'$  and g(d) = 1 for every  $d \in D$ . In the case of our metric space  $(I_t, d_H)$  one concrete g that satisfies the constraints can be shown to be  $g(J) = \frac{d_H(J,C')}{d_H(J,C')+d_H(J,D)}$ . This g is bounded by 0 and 1. For an arbitrary Radon measure  $\nu \in \mathcal{M}(I_t)$  and every compact  $C' \subseteq C$  we consider

$$\int_{I_t} |\mathbf{1}_B - g| d\nu = \int_{C \setminus C'} |\mathbf{1}_B - g| d\nu \le \nu(C \setminus C').$$

Thus

$$\int_{I_t} |\mathbf{1}_B - g| d\nu \leq \inf_{C' \subseteq C, \ C' \text{ compact}} \nu(C \setminus C') \leq \nu(C) - \sup_{C' \subseteq C, \ C' \text{ compact}} \nu(C') = 0,$$

where the last step is due to the inner regularity of every Radon measure  $\nu$ .

**Lemma 4.6.** Equality  $\operatorname{las}_{\alpha(P(X,\varepsilon))}^{\gamma}(C(X,r)) = \mathcal{N}(X,r)$  holds.

*Proof.* Set  $\alpha = \alpha(P(X, \varepsilon))$  We consider a feasible measure  $\lambda$  for the program  $las_{\alpha}^{\gamma}(C(X, r))$  and its representation  $\lambda = \int \chi_R d\sigma(R)$ , which exists due to Lemma 3.1.

Now we have to verify that  $\sigma$ -supported subsets  $R \subseteq X$  determine a covering. For contradiction, suppose R does not determine a covering. Then there exists  $y \in X$  so that  $\overline{B}(y,r) \cap R = \emptyset$ . Consider the map  $k : I_{\alpha} \to \mathbb{R}$ , with  $k(J) = (-1)^{|J|}$  if  $J \subseteq \overline{B}(y,r)$  and k(J) = 0 otherwise, as defined in Lemma 4.5 and let g be its continuous version. Then

$$\begin{aligned} 0 &\leq \langle (B_{g}^{y})^{*}\chi_{R}, g \otimes g \rangle \\ &= \chi^{R}(B_{\alpha}^{y}(g \otimes g)) \\ &= \sum_{x \in \overline{B}(y,r), J, J': J \cup J' \cup \{x\} \subseteq R} g \otimes g(J, J') - \sum_{J, J' \in I_{t}: J \cup J' \subseteq R} g \otimes g(J, J'). \\ &= |\overline{B}(y,r) \cap R| \sum_{J, J': J \cup J' \cup \subseteq R} g \otimes g(J, J') - \sum_{J, J' \in I_{t}: J \cup J' \subseteq R} g \otimes g(J, J') \\ &= -\sum_{J, J' \in I_{t}: J \cup J' \subseteq R} g \otimes g(J, J') \\ &= -\left(\sum_{J: J \subseteq \overline{B}(y,r) \cap R} (-1)^{|J|}\right)^{2} \\ &= -1. \end{aligned}$$

We finish the proof of the lemma by looking at the objective value of  $\lambda$  which is

$$\lambda(I_{=1}) = \int \chi_R(I_{=1}) \, d\sigma(R) = \int |R| \, d\sigma(R) \ge \int \mathcal{N}(X, r) \, d\sigma(R) = \mathcal{N}(X, r),$$

because every R in the support of  $\sigma$  determines a covering and  $\sigma$  is a probability measure.

### 5. The dual covering hierarchy

In this section we assume that the automorphism group G of X acts transitively on X so that X is the homogenous space G/H where H is the stabilizer subgroup of a point in G. This is a natural assumption which is fulfilled in many examples; for example when X is the unit sphere  $S^{n-1}$  and G is the orthogonal group O(n)and H is isomorphic to O(n-1). Then the Haar measure of G induces a probability measure  $\omega$  on G/H which satisfies conditions (a) and (b) given in Section 1.

Looking at the covering hierarchy this has the advantage, due to convexity, that the constraints coming from the localizing matrix only have to required for one arbitrary point of X. Let  $e \in X$  be this point, then

$$\begin{aligned} \operatorname{las}_{t}^{\gamma}(C(X,r)) &= \inf \left\{ \lambda(I_{=1}) \ : \lambda \in \mathcal{M}(I_{2t})_{\geq 0}, \ \lambda(\{\emptyset\}) = 1, \\ A_{t}^{*}(\lambda) \in \mathcal{M}(I_{t} \times I_{t})_{\succeq 0}, \\ (B_{t-1}^{e})^{*}(\lambda) \in \mathcal{M}(I_{t-1} \times I_{t-1})_{\succeq 0} \right\} \end{aligned}$$

We define the *t*-th step of the dual covering hierarchy by

$$\begin{aligned} \operatorname{las}_{t}^{\gamma,*}(C(X,r)) &= \sup \left\{ y : y \in \mathbb{R}, \\ \mathbf{1}_{I_{l=1}}(S) - y\mathbf{1}_{\{\emptyset\}}(S) - A_{t}K(S) - B_{t-1}^{e}K'(S) \geq 0 \\ & \text{for all } S \in I_{2t}, \\ K \in \mathcal{C}(I_{t} \times I_{t})_{\geq 0}, \ K' \in \mathcal{C}(I_{t-1} \times I_{t-1})_{\geq 0} \right\}. \end{aligned}$$

One can easily check that weak duality  $las_t^{\gamma,*} \leq las_t^{\gamma}$  holds. Although we do not need this here, we note that also strong duality holds; this was shown by Rolfes in [24] using the framework of infinite dimensional conic optimization presented in Barvinok [3, Chapter IV].

With the help of the dual covering hierarchy  $las_t^{\gamma,*}$  it is further possible to show that the first step of the hierarchy coincides with the volume bound.

**Theorem 5.1.** If the automorphism group G acts transitively on X, then the first step of the covering hierarchy collapses to the volume bound,

$$\operatorname{las}_{1}^{\gamma,*}(C(X,r)) = \frac{1}{\omega_{r}}.$$

*Proof.* Consider a probability measure  $\omega$  satisfying properties (a) and (b). Then, if t = 1, we observe that the measure

$$\lambda(\mathcal{S}) = \begin{cases} 1 & \text{if } \mathcal{S} = \{\emptyset\}, \\ \frac{\omega(\bigcup_{\{x\}\in\mathcal{S}}\{x\})}{\omega_r} & \text{if } \mathcal{S} \subseteq I_{=1}, \\ 0 & \text{otherwise.} \end{cases}$$

is a feasible measure for  $las_1^{\gamma}(C(X, r))$ . Its objective value is  $\lambda(I_{=1}) = \frac{1}{\omega_r}$  and thus  $\begin{array}{l} \text{implies } \operatorname{las}_1^\gamma(C(X,r)) \leq \frac{1}{\omega_r}.\\ \text{For the reverse inequality we have} \end{array}$ 

$$B_0^e K'(\emptyset) = -K'(\emptyset, \emptyset),$$

and

$$\begin{split} B_0^e K'\left(\{z\}\right) &= \sum_{x \in \overline{B}(e,r): \ \{x\} = \{z\}} K'(\emptyset, \emptyset) \\ &= \begin{cases} K'(\emptyset, \emptyset) & \text{if } z \in \overline{B}(e, r) \\ 0 & \text{otherwise,} \end{cases} \end{split}$$

and

$$B_0^e K'(\{z_1, z_2\}) = \sum_{x \in \overline{B}(e,r): \{x\} = \{z_1, z_2\}} K'(\emptyset, \emptyset) = 0.$$

Thus,

$$K = 0, \ K'(\emptyset, \emptyset) = \frac{1}{\omega_r} \text{ and } y = K'(\emptyset, \emptyset)$$

is a feasible solution for  $las_1^{\gamma,*}(C(X,r))$ . So  $las_1^{\gamma,*}(C(X,r)) \geq \frac{1}{\omega_r}$  and by weak duality  $las_1^{\gamma}(C(X, r)) \ge \frac{1}{\omega_r}$ . 

For the packing number  $\alpha(S^{n-1}, r)$  for the unit sphere, the first step of the packing hierarchy  $las_1^{\alpha}(P(S^{n-1}, r))$  coincides with the linear programming bound

12

 $\vartheta'(P(S^{n-1},r))$  of Delsarte, Goethals, Seidel, which often provides strong and sometimes even tight upper bounds. Now Theorem 5.1 gives an explanation why the corresponding first step of the covering hierarchy  $las_1^{\gamma}(C(S^{n-1},r))$ , which also is, after symmetry reduction, a linear programming bound, is not a strong bound for the covering number. On the other hand, the first step of the dual covering hierarchy is implicitly used in the analysis of the greedy approach to construct efficient coverings, see [23].

### Acknowledgements

The authors are grateful to David de Laat for helpful discussions. The first author was supported by the Tromsø Research Foundation project "Symmetry in Real Algebraic Geometry" and the UiT Aurora project MASCOT. The second author was supported by a "Short-Term Grant" funded by DAAD. The third named author is partially supported by the SFB/TRR 191 "Symplectic Structures in Geometry, Algebra and Dynamics" funded by the DFG.

### References

- C. Bachoc, F. Vallentin, New upper bounds for kissing numbers from semidefinite programming, J. Amer. Math. Soc. 21 (2008), 909–924.
- [2] C. Bachoc, G. Nebe, F.M. de Oliveira Filho, F. Vallentin, Lower bounds for measurable chromatic numbers, Geom. Funct. Anal. 19 (2009), 645–661.
- [3] A. Barvinok, A course in convexity, AMS, 2002.
- [4] S.V. Borodachov, D.P. Hardin, E.B. Saff, Discrete energy on rectifiable sets, Springer, 2019.
- [5] H. Cohn, D. Laat, A. Salmon, Three-point bounds for sphere packing, arXiv:2206.15373 [math.MG]
- [6] H. Cohn, A. Salmon, Sphere packing bounds via rescaling, arXiv:2108.10936 [math.MG]
- [7] F. Cucker, S. Smale, On the mathematical foundations of learning, Bull. Amer. Math. Soc. 39 (2002), 1–49.
- [8] R.E. Curto, L.A. Fialkow, The truncated complex K-moment problem, Trans. Amer. Math. Soc. 352 (2000), 2825–2855.
- [9] D. de Laat, F. Vallentin, A semidefinite programming hierarchy for packing problems in discrete geometry, Math. Program., Ser. B 151 (2015), 529–553.
- [10] P. Delsarte, J.M. Goethals, J.J. Seidel, Spherical codes and designs, Geom. Dedicata 6 (1977), 363–388.
- [11] D. de Laat, Moment methods in energy minimization: New bounds for Riesz minimal energy problems, Trans. Amer. Math. Soc. 373 (2020), 1407–1453
- [12] D. de Laat, F Caluza Machado, W. de Muinck Keizer, The Lasserre hierarchy for equiangular lines with a fixed angle, arXiv:2211.16471 [math.MG]
- [13] S. Foucart, H. Rauhut, A mathematical introduction to compressive sensing, Birkhäuser/Springer, 2013.
- [14] J.B. Lasserre, An explicit equivalent positive semidefinite program for nonlinear 0-1 programs, SIAM J. Optim. 12 (2002), 756–769.
- [15] M. Laurent, A comparison of the Sherali-Adams, Lovász-Schrijver, and Lasserre relaxations for 0-1 programming, Math. Oper. Res. 28 (2003), 470–496.
- [16] M. Ledoux, M. Talagrand, Probability in Banach spaces, Springer, 1991.
- [17] B. Lindström, Determinants on semilattices, Proc. Amer. Math. Soc. 20 (1969) 207–208.
- [18] L. Lovász, On the Shannon capacity of a graph, IEEE Trans. Inf. Th. 25 (1979), 1-7.
- [19] M. Naszódi, Flavors of Translative Coverings, pp. 335–358 in: New Trends in Intuitive Geometry, Springer, 2018.
- [20] M.A. Nielsen, I.L. Chuang, Quantum computation and quantum information, Cambridge University Press, 2000.
- [21] M. Putinar, Positive polynomials on compact semi-algebraic sets, Ind. Univ. Math. J. 42 (1993), 969–984.
- [22] C.A. Rogers, Packing and covering, Cambridge University Press, 1964.

- [23] J.H. Rolfes, F. Vallentin, Covering compact metric spaces greedily, Acta Math. Hungar. 155 (2018), 130–140.
- [24] J.H. Rolfes, Convex optimization techniques for geometric covering problems, PhD thesis, University of Cologne, 2019.
- [25] A. Schrijver, A comparison of the Delsarte and Lovász bounds, IEEE Trans. Inf. Th. 25 (1979), 425–429.
- [26] B. Simon, Convexity: An analytic viewpoint, Cambridge University Press, 2011
- [27] H.S. Wilf, Hadamard determinants, Möbius functions, and the chromatic number of a graph, Bull. Amer. Math. Soc. 74 (1968) 960–964.

CORDIAN RIENER, DEPARTMENT OF MATHEMATICS AND STATISTICS, UIT THE ARCTIC UNIVER-SITY OF NORWAY, 9037 TROMSØ, NORWAY Email address: cordian.riener@uit.no

Jan Rolfes, Friedrich-Alexander-Universität Erlangen-Nürnberg, Cauerstr. 11, 91058 Erlangen, Germany

 $Email \ address: \ \texttt{jan.rolfes@fau.de}$ 

FRANK VALLENTIN, UNIVERSITÄT ZU KÖLN, DEPARTMENT MATHEMATIK/INFORMATIK, ABTEILUNG MATHEMATIK, WEYERTAL 86-90, 50931 KÖLN, GERMANY Email address: frank.vallentin@uni-koeln.de

14