Diagonalizing the Born-Oppenheimer Hamiltonian via Moyal Perturbation Theory, Nonadiabatic Corrections and Translational Degrees of Freedom

Robert Littlejohn,¹ Jonathan Rawlinson,² and Joseph Subotnik³ ¹⁾Department of Physics, University of California, Berkeley, California 94720 USA ²⁾School of Mathematics, University of Manchester, Manchester UK ³⁾Department of Chemistry, University of Pennsylvania, Philadelphia, PA, USA

(*Electronic mail: subotnik@sas.upenn.edu)

(*Electronic mail: jonathan.rawlinson@manchester.ac.uk)

(*Electronic mail: robert@wigner.berkeley.edu)

(Dated: 21 December 2023)

This article describes a method for calculating higher order or nonadiabatic corrections in Born-Oppenheimer theory and its interaction with the translational degrees of freedom. The method uses the Wigner-Weyl correspondence to map nuclear operators into functions on the classical phase space and the Moyal star product to represent operator multiplication on those functions. The result is a power series in κ^2 , where $\kappa = (m/M)^{1/4}$ is the usual Born-Oppenheimer parameter. The lowest order term is the usual Born-Oppenheimer approximation while higher order terms are nonadiabatic corrections. These are needed in calculations of electronic currents, momenta and densities. The method was applied to Born-Oppenheimer theory by Littlejohn and Weigert (1993), in a treatment that notably produced the correction K_{22} to the Born-Oppenheimer Hamiltonian (see *infra*). Recently Mátyus and Teufel (2019) have applied an improved and more elegant version of the method to Born-Oppenheimer theory, and have calculated the Born-Oppenheimer Hamiltonian for multiple potential energy surfaces to order κ^6 . One of the shortcomings of earlier methods is that the separation of nuclear and electronic degrees of freedom takes place in the context of the exact symmetries (for an isolated molecule) of translations and rotations, and these need to be a part of the discussion. This article presents an independent derivation of the Moyal expansion in molecular Born-Oppenheimer theory, with special attention to the translational degrees of freedom. We show how electronic currents and momenta can be calculated within the framework of Moyal perturbation theory; we derive the transformation laws of the electronic Hamiltonian, the electronic eigenstates, and the derivative couplings under translations; we discuss in detail the rectilinear motion of the molecular center of mass in the Born-Oppenheimer representation; and we show how the elimination of the translational components of the derivative couplings leads to a unitary transformation that has the effect of exactly separating the translational degrees of freedom. Our discussion is framed in terms of dressing transformations, that is, unitary transformations that change the physical meanings of the operators of the theory. We show, for example, that in the Born-Oppenheimer representation, the operator that looks like the momentum of the nuclei actually includes, from a physical standpoint, the momentum of the electrons. Moreover, this result is exact. In regard to hybrid classical-quantum models such as used in surface hopping calculations, we note that the theory presented here will be useful in developing new semiclassical approximations going beyond the Born-Oppenheimer approximation, as well as taking into account a glaring error of existing schemes, namely, the

failure to conserve linear and angular momentum.

I. INTRODUCTION

In the Born-Oppenheimer theory of molecules the Hamiltonian becomes an infinite-dimensional matrix of nuclear operators once an electronic basis is chosen. Moyal perturbation theory provides a method of block-diagonalizing this matrix to obtain effective dynamics on one or a small number of potential energy surfaces. This is an elegant way of deriving Born-Oppenheimer dynamics on a small number of surfaces in a manner that is more rigorous than the usual approaches, with other advantages enumerated below.

By "Moyal perturbation theory" we refer to the use of the Wigner-Weyl correspondence and the Moyal star product to block-diagonalize matrices of operators. It would be more descriptive to call it "Wigner-Weyl-Moyal-van Vleck" perturbation theory, but we will use the shorter designation. The Wigner-Weyl correspondence maps quantum operators into functions on a classical phase space, and the Moyal star product is used to represent operator multiplication on those functions. The Wigner-Weyl-Moyal formalism is fairly well known, and for convenience is summarized in Appendix A, along with literature citations. The application in this article of Moyal perturbation theory to Born-Oppenheimer problems is based on the treatment of Weigert and Littlejohn (1993), with many improvements.

Moyal perturbation theory is an efficient way of obtaining higher order or nonadiabatic corrections in Born-Oppenheimer theory. Such corrections are needed in cases where the lowest order contribution vanishes, notably in the calculation of electronic currents, densities, linear and angular momenta, and matrix elements needed for radiative transitions. There is an extensive literature on such problems, including Mead and Moscowitz (1967); Nafie (1983); Stephens (1985); Stephens and Lowe (1985); Freedman and Nafie (1986); Buckingham, Fowler, and Galwas (1987); Nafie (1992, 1997); Barth *et al.* (2009); Okuyama and Takatsuka (2009); Patchkovskii (2012); Diestler (2012a,b, 2013); Diestler *et al.* (2013); Bredtmann *et al.* (2015); Pohl and Tremblay (2016); Albert, Hader, and Engel (2017); Schaupp, Albert, and Engel (2018); Schaupp and Engel (2022a,b). In recent years the method of exact factorization has been developed and promoted (Abedi, Maitra, and Gross (2010, 2012); Cederbaum (2013, 2014, 2015); Parashar, Sajeev, and Ghosh (2015); Jecko, Sutcliffe, and Woolley (2015); Requist and Gross (2016); Requist, Tandetzky, and Gross (2016); Curchod and Agostini (2017); Requist, Proetto, and Gross (2017); Agostini and Curchod (2018); Lorin (2021); Martinazzo and Burghardt (2023)). A notable application of this method is the calculation of higher order or nonadiabatic corrections (Scherrer, Vuilleumier, and Sabastiani (2013); Scherrer *et al.* (2015); Schild, Agostini, and Gross (2016)). Other approaches to higher order corrections include Pachucki and Komasa (2008); Mátyus (2018); Mátyus and Teufel (2019); Requist (2023). Of these, Mátyus and Teufel (2019); Requist (2023) have used a version of what we are calling Moyal perturbation theory.

The advantages of Moyal perturbation theory, as we see them, for accessing nonadiabatic corrections and for related topics are the following. The method is clean and elegant, allowing one to visualize the structure of the theory as a systematic procedure that can be carried to any order in the Born-Oppenheimer parameter $\kappa = (m/M)^{1/4}$. The method automatically incorporates terms in the effective Hamiltonian, such as K_{22} that is discussed below, that should be there but are usually neglected. The method automatically accommodates large-amplitude motions, such as occur in scattering, isomerization and photoexcitation problems, and is not restricted to small-amplitude expansions about an equilibrium. It works in simple cases such as motion on a single surface in the electrostatic model, as well as in generalizations including multiple surfaces, fine-structure and spin. Many of the first order results presented in this article, for example, the first order corrections to the wave function and formulas for electronic momenta and currents, are known in the case of the electrostatic model on a single surface, but generalizations to multiple surfaces and spin are incomplete. These are of current interest, for example, in hybrid classical-quantum models used in surface hopping (Mannouch and Richardson (2023); Runeson and Manolopoulos (2023); Bian et al. (2021, 2023)). The method interacts well with the exact symmetries of the system, including overall translations and rotations. The case of rotational invariance was treated by Littlejohn, Rawlinson, and Subotnik (2023), and translational invariance is covered in this article. The method allows one to make simple statements about the conservation of linear and angular momentum that are valid to all orders in κ . The method leads to a perspective in which Born-Oppenheimer theory consists of a series of dressing transformations, that is, unitary transformations that change the physical meanings of the variables employed, for example, "dressing" nuclear dynamical variables to include the effects of the electrons. The notion of the Born-Oppenheimer transformation as a dressing transformation was introduced by Cederbaum (2004), and we have developed the idea considerably. We feel that it makes an important contribution to perspectives on Born-Oppenheimer theory.

For example, in a recent study (Littlejohn, Rawlinson, and Subotnik (2023)) we showed that in the usual Born-Oppenheimer representation the operator that looks like the orbital angular momentum of the nuclei actually includes the orbital angular momentum of the electrons. This is in the electrostatic model; in fine-structure models the electron spin is included as well. These statements are exact in the sense of being valid to all orders of the Born-Oppenheimer expansion parameter κ . For another example, in this article we display (in subsection V J) a unitary transformation that has the effect of separating the translational degrees of freedom, after which the the operator that looks like the nuclear center of mass is actually, from a physical standpoint, the molecular center of mass, including the electrons.

Finally, there are questions regarding three steps that may be distinguished in the Born-Oppenheimer theory of isolated molecules, namely, the separation of the translational degrees of freedom, the separation of the rotational degrees of freedom, and the expansion in the Born-Oppenheimer parameter κ . Note that the third step amounts to the separation of the nuclear and electronic degrees of freedom. For example, there is the question of the advantages and disadvantages of carrying out these steps in various orders. We feel that treating the third step by Moyal perturbation theory clarifies the analysis of this question.

The commutativity of the first and third steps (the separation of translations and the κ expansion) is explicitly addressed in this article. It is sometimes stated that the Born-Oppenheimer
approximation violates the separation of translational degrees of freedom (Cederbaum (2013)),
but this does not mean that translational invariance or the associated conservation laws are lost,
simply that they take on different forms in the different representations. Some of these are very
interesting.

Moyal perturbation theory was first developed by Blount (1962a), who applied it to various problems in condensed matter physics as well as to the Foldy-Wouthuysen transformation (Blount (1962b); Foldy and Wouthuysen (1950); Bjorken and Drell (1964)), the latter of which is the block-diagonalization of the Dirac Hamiltonian. The method was rediscovered by Littlejohn and Flynn (1991b,a), who applied it to the WKB theory of multi-component wave equations and to several other problems in adiabatic or semiclassical theory, including spin-orbit coupling and multi-dimensional Landau-Zener transitions (Littlejohn and Flynn (1992, 1993)). Later Littlejohn and Weigert (1993) applied the method to the adiabatic motion of a neutral spinning particle in an inhomogeneous magnetic field, otherwise a Stern-Gerlach apparatus. That problem is formally almost identical to the Born-Oppenheimer approximation in molecules, in which the spin of the neutral particle corresponds to the electronic degrees of freedom, while the spatial variables correspond to the nuclear degrees of freedom. A notable outcome of that work was the production of a second-order contribution to the Hamiltonian, what we call K_{22} in this article. This term had been missing in the analysis by other authors (Aharonov and Stern (1992)) of the Stern-Gerlach problem.

Later Weigert and Littlejohn (1993) applied Moyal theory to the Born-Oppenheimer approximation in molecules, producing among other things the contribution K_{22} to the Hamiltonian. This term is notable for the fact that for large-amplitude motions it is of the same order of magnitude as the well known diagonal Born-Oppenheimer correction, but it has received much less attention than the latter.

Later Panati, Spohn, and Teufel (2002) and Teufel (2003) developed the functional analysis of Moyal perturbation theory, making several applications and innovations. One of the latter is the breaking of the problem into two parts, the computation of the formally invariant subspace and the representation of the dynamics on that subspace. Another is the use of operator-valued Weyl symbols, which leads to a compact and largely coordinate- and basis-independent formalism. This formalism was applied by Mátyus and Teufel (2019) to Born-Oppenheimer theory, producing an expansion of the Born-Oppenheimer Hamiltonian on one or a few potential energy surfaces out to order κ^6 (described as "third order").

The term K_{22} is discussed more fully in Sec. III J below. Its history and its role in Born-Oppenheimer theory have been discussed by Mátyus (2018), whose citations place its discovery in the 1960's, although several of the earlier references are specific to small vibrations about an equilibrium in diatomic or triatomic molecules. To put this term into general language, applying to a molecule with any number of atoms, it is necessary to work in lab or Cartesian coordinates. In addition, the importance of this term grows with energy, and it is more important for large-amplitude motions than for small vibrations. Therefore to treat this term in generality it is necessary not to expand about an equilibrium position. As far as we can see the first authors to derive this term in such generality were Moody, Shapere, and Wilczek (1989), who used a path-integral formalism for treating the Born-Oppenheimer approximation and its higher order corrections and who noted that this term modifies the kinetic-energy metric, making it non-Euclidean at second order. In addition, there is the independent discovery by Goldhaber (2005), who derived K_{22} in a discussion that links Berry's phase, geometry, and averaging techniques in quantum field theory. In this article we will show that the contribution K_{22} to the Hamiltonian is necessary in the Born-Oppenheimer representation to account for the proper (rectilinear) evolution of the center of mass of the molecule.

The original treatments of Born-Oppenheimer theory (Born and Oppenheimer (1927); Born and Huang (1954)) and large parts of the vast literature on this subject envision an expansion in powers of $\kappa = (m/M)^{1/4}$, in which the nuclear displacements from an equilibrium position are small, of order κ , while nuclear momenta are of order κ^{-1} . We call this the "small-amplitude" ordering. It was pointed out by Littlejohn and Flynn (1992), however, that large-amplitude motions such as occur in scattering theory required a different ordering, in which nuclear momenta are of order κ^{-2} . Later Mead (2006) discussed the large-amplitude ordering in detail and its contrast to the small-amplitude ordering. Another analysis has been given recently by Mátyus and Teufel (2019), and we have returned to the subject in Sec. II D below. To summarize the situation, Moyal perturbation theory is an expansion in powers of κ^2 , it accommodates large-amplitude motions, and it is equivalent to a semiclassical theory in the nuclear degrees of freedom, that is, it is an expansion in the "nuclear \hbar " (but not the electronic).

The outline of this paper is as follows. Section II contains the setup of the problem, including notation, the relevant Hamiltonians, the electronic bases that will be used (adiabatic, diabatic, etc.), the coordinate transformations that bring about the exact elimination of the translational degrees of freedom, and the several representations we will use. In this article the word "representation" is used in a particular sense, implying a space of wave functions, a mapping between physical states (for example, the ground state of the molecule or an eigenstate of energy and angular momentum) and wave functions in that space, and a mapping between physical observables (energy, momentum, etc.) and specific linear operators that act on those wave functions. Different representations are connected by unitary operators.

In Sec. III we describe Moyal perturbation theory in a manner that is hopefully as painless as possible, based on the approach of Littlejohn and Flynn (1991b); Littlejohn and Weigert (1993); Weigert and Littlejohn (1993). Much the same territory has been covered by Mátyus and Teufel (2019), based on the formalism of Panati, Spohn, and Teufel (2002); Teufel (2003). In comparison, our formalism is more coordinate-based and less abstract. Our discussion of the relation between the generator of the dressing transformation and the usual Landau-Zener transmission probability seems to be new, as is our presentation in subsection III M of the second order, nonadiabatic corrections to the wave function. For simplicity in this section we treat only single-surface problems.

In Sec. IV we generalize parts of the single-surface calculation in Sec. III to multiple surfaces, and we review how the Moyal theory is applied to the calculation of electronic momenta and currents. We offer an approach in which the higher order corrections are obtained, not by finding corrections to the wave function, but by transforming ("dressing") the operators.

In Sec. V we begin by treating a number of issues regarding the translational degrees of freedom, in a manner similar to our treatment of rotations in Littlejohn, Rawlinson, and Subotnik (2023). In particular, we establish phase conventions for the electronic basis states, both adiabatic and diabatic, in effect showing how they transform under translations. We also give the transformation laws for the electronic Hamiltonian and the derivative couplings under translations. These results are placed in the geometrical context of the translational fiber bundle, which we explain as a certain decomposition of the nuclear configuration space. A notable consequence of this work is that in the Born-Oppenheimer representations, both the original and all the dressed versions of it, the operator that looks like the total momentum of the nuclei actually includes, from a physical standpoint, the momentum of the electrons. A similar statement regarding angular momentum was a principal point of Littlejohn, Rawlinson, and Subotnik (2023).

We then address several topics concerning the translational degrees of freedom and Moyal perturbation theory. First we study the molecular center of mass (including the electrons) and show how its uniform, rectilinear motion is expressed in the dressed Born-Oppenheimer representation. It turns out that the term K_{22} in the Hamiltonian is required for this to work out right. Next we study the translational components of the electronic current and the derivative couplings, in which the resolvent operator can be eliminated and results obtained in closed form. The existence of the translational components of the derivative couplings leads to some apparent paradoxes, which we resolve by engineering a unitary transformation that removes them. This transformation is revealed by the use of Moyal perturbation theory, but ultimately is expressed in closed form (not as a power series). This produces an operator that has the effect of exactly separating the translational degrees of freedom. As far as we can tell this is a new perspective on the separation of the translational degrees of freedom, in that the procedure can be seen as a dressing transformation.

Finally, in Sec. VI we present some conclusions and plans for future work.

A note on notation and terminology. Starting in subsection III D we use hats on nuclear operators, for example $\hat{\mathbf{P}}_{\alpha}$, to distinguish them from classical quantities or *c*-numbers. See the introduction to Sec. III. The Hamiltonian H_{BO} is the usual (scalar) Born-Oppenheimer Hamiltonian on a single surface; see (53). An overbar is used to indicate fully dressed operators, for example the Hamiltonian \bar{H} (see (58)) or $\bar{\mathbf{p}}_i$. The scalar Hamiltonian *K* is the diagonal element of \bar{H} , the effective Hamiltonian that replaces H_{BO} (see above (93)). In this article, "diabatic" means what is sometimes called "quasi-diabatic," that is, there is no implication that the diabatic basis eliminates the derivative couplings (which is usually impossible anyway).

II. PRELIMINARIES

In this section we describe the setup of the problem and establish notation. We adopt the electrostatic model for the electronic Hamiltonian (2). For simplicity, we treat the nuclei as distinguishable, spinless particles.

A. Hamiltonians and Notation

We consider a molecule with N nuclei. The molecular Hamiltonian in the lab frame is

$$H = \sum_{\alpha=1}^{N} \frac{\mathbf{P}_{\alpha}^{2}}{2M_{\alpha}} + H_{e}(x; \mathbf{r}, \mathbf{p}), \qquad (1)$$

where the electronic Hamiltonian is

$$H_e(x;\mathbf{r},\mathbf{p}) = \sum_{i=1}^{N_e} \frac{\mathbf{p}_i^2}{2m_e} + V_{\text{Coul}}(\mathbf{X},\mathbf{r}).$$
(2)

Here N_e is the number of electrons; \mathbf{X}_{α} , $\mathbf{P}_{\alpha} = -i\hbar\nabla_{\alpha}$ and M_{α} are respectively the nuclear positions, momenta and masses; and \mathbf{r}_i and $\mathbf{p}_i = -i\hbar\nabla_i$ are respectively the electron positions and momenta. Gradient operators are $\nabla_{\alpha} = \partial/\partial \mathbf{X}_{\alpha}$ and $\nabla_i = \partial/\partial \mathbf{r}_i$. All positions and momenta are taken with respect to a lab frame. Indices α , β label nuclei and range over $1, \ldots, N$, while indices i, j label electrons and range over $1, \ldots, N_e$. The electron mass is m_e . The bare symbols \mathbf{X} and \mathbf{r} stand for the collections ($\mathbf{X}_1, \ldots, \mathbf{X}_N$) and ($\mathbf{r}_1, \ldots, \mathbf{r}_{N_e}$), respectively, and similarly for the momenta \mathbf{P} and \mathbf{p} . The potential V_{Coul} contains all the Coulomb interactions among all the particles, nuclei and electrons. The nuclear configuration space is the space upon which the \mathbf{X}_{α} are coordinates; it is \mathbb{R}^{3N} . We denote a point of nuclear configuration space by the symbol x, whose coordinates are $\mathbf{X} = (\mathbf{X}_1, \ldots, \mathbf{X}_N)$. In our usage, x and \mathbf{X} mean almost the same thing (x is the point, \mathbf{X} , the coordinates of that point).

B. The Adiabatic Basis and the Working Basis

We define the "adiabatic basis" as the energy eigenbasis of the electronic Hamiltonian,

$$H_e(x) |ax;k\rangle = \varepsilon_k(x) |ax;k\rangle.$$
(3)

The adiabatic basis vectors in ket language are $|ax;k\rangle$, where *a* stands for "adiabatic," *x* indicates the parametric dependence on the nuclear configuration, and *k* is the quantum number (a sequencing number for energy eigenvalues). In (3) we write simply $H_e(x)$ for the electronic Hamiltonian, indicating the *x*-dependence of the operator, which elsewhere is written $H_e(x; \mathbf{r}, \mathbf{p})$. The energy eigenvalues themselves are $\varepsilon_k(x)$. The energy eigenfunctions in wave function language will be denoted by

$$\phi_{ak}(x;\mathbf{r}) = \langle \mathbf{r} | ax; k \rangle. \tag{4}$$

Although we shall treat the index k notationally as if it were discrete, in fact above a threshold the spectrum of $H_e(x)$ becomes continuous and formal sums over k that employ the adiabatic basis must be interpreted as discrete sums over the discrete spectrum plus integrals over the continuous spectrum. In addition, one should note that the continuum threshold is a function of x.

In the following we shall make use of a basis of electronic states that we call the "working basis," which we denote simply by $|x;k\rangle$ (without the *a*). It is defined to be either the adiabatic basis or else a substitute, which will be defined as the need arises. We define the working basis wave functions by

$$\phi_k(x;\mathbf{r}) = \langle \mathbf{r} | x; k \rangle \tag{5}$$

(again, without the *a*).

We choose the phase conventions of the adiabatic basis states so that these states are invariant under time reversal. We also assume that the coefficients of the linear transformation taking us from the adiabatic basis to the working basis are real, so that the working basis states are also invariant under time reversal. In the electrostatic model, this just means that the basis wave functions $\phi_{ak}(x; \mathbf{r})$ or $\phi_k(x; \mathbf{r})$ are real.

Here and below when we use bra-ket notation for scalar products and matrix elements, it is understood that we are integrating only over the electronic coordinates \mathbf{r} , not the nuclear coordinates, except as noted. For example, we have the orthonormality relations,

$$\langle x;k|x;l\rangle = \delta_{kl}.\tag{6}$$

We introduce the derivative couplings,

$$\mathbf{F}_{\alpha;kl}(x) = \langle x; k | \nabla_{\alpha} | x; l \rangle = -(\nabla_{\alpha} \langle x; k |) | x; l \rangle, \tag{7}$$

where the alternative form follows from the orthogonality relations (6). The derivative couplings

are defined to the extent that the basis states $|x;k\rangle$ are differentiable with respect to x. The orthonormality relations (6) imply that $\mathbf{F}_{\alpha;kl}(x)$, regarded as a matrix in kl, is anti-Hermitian,

$$\mathbf{F}_{\alpha;kl}(x) = -\mathbf{F}_{\alpha;lk}^*(x). \tag{8}$$

In addition, the invariance of the basis vectors under time reversal implies that $\mathbf{F}_{\alpha;kl}(x)$ is real, so that as a matrix in (kl), it is real and antisymmetric. Thus, the derivative couplings vanish on the diagonal,

$$\mathbf{F}_{\alpha;kk}(x) = 0. \tag{9}$$

C. Two Representations

We introduce two representations for the quantum state of the molecule, the "molecular" and the "Born-Oppenheimer." These are described in more detail in Sec. V of Littlejohn, Rawlinson, and Subotnik (2023). In the molecular representation the state of the molecule is represented by a wave function $\Psi(\mathbf{X}, \mathbf{r})$ that depends on the coordinates of both nuclei and electrons. In the Born-Oppenheimer representation the state of the molecule is represented by an infinite-dimensional vector of purely nuclear wave functions, $\psi_k(\mathbf{X})$, indexed by *k* and defined by

$$\Psi(\mathbf{X},\mathbf{r}) = \sum_{k} \psi_k(\mathbf{X}) \,\phi_k(\mathbf{X};\mathbf{r}). \tag{10}$$

The Born-Oppenheimer representation depends on the choice made for the working basis, and, in particular, on the phase conventions of that basis. We denote the association between the two representations of the state of the molecule by

$$\Psi(\mathbf{X}, \mathbf{r}) \longleftrightarrow \psi_k(\mathbf{X}),\tag{11}$$

where the notation \longleftrightarrow is a reminder that the relationship is one-to-one and that no information is lost by transforming to the Born-Oppenheimer representation. For time-dependent problems both Ψ and ψ_k depend on *t*, but the basis kets $|x;k\rangle$ or wave functions $\phi_k(\mathbf{X};\mathbf{r})$ do not.

Similarly, let *A* be an operator acting in the molecular representation, say, $\Psi'(\mathbf{X}, \mathbf{r}) = (A\Psi)(\mathbf{X}, \mathbf{r})$, and let $\Psi \leftrightarrow \psi_k$ and $\Psi' \leftrightarrow \psi'_k$. Then in the Born-Oppenheimer representation *A* is represented by an infinite-dimensional matrix of operators A_{kl} that act on nuclear wave functions, such that

$$\boldsymbol{\psi}_{k}'(\mathbf{X}) = \sum_{l} (A_{kl} \, \boldsymbol{\psi}_{l})(\mathbf{X}). \tag{12}$$

We denote the association between these two representations of the operator by $A \leftrightarrow A_{kl}$. The components A_{kl} of the operator in the Born-Oppenheimer representation are purely nuclear operators, that can be thought of as functions of **X** and **P**.

Several examples of this association are discussed in Littlejohn, Rawlinson, and Subotnik (2023), including the following. A purely multiplicative operator in the molecular representation depending only on the nuclear coordinates \mathbf{X}_{α} , call it f(x), corresponds, in the Born-Oppenheimer representation, to a multiple of the identity matrix, that is, $f(x) \leftrightarrow f(x) \delta_{kl}$. In particular, this applies to the nuclear coordinates, the components of \mathbf{X}_{α} . A purely electronic operator in the molecular representation, that is, a function of \mathbf{r} and \mathbf{p} , corresponds to a purely multiplicative operator in the Born-Oppenheimer representation, that is, a function of \mathbf{r} and \mathbf{p} , corresponds to a purely multiplicative operator in the Born-Oppenheimer representation, that is, a matrix whose matrix elements are functions of \mathbf{X} only. These are nothing but the matrix elements of the original operator in the working basis. For example, in the case of the momentum \mathbf{p}_i of one of the electrons, we have $\mathbf{p}_i \longleftrightarrow \langle x; k | \mathbf{p}_i | x; l \rangle$. This also applies to electronic operators that are parameterized by x, such as the electronic Hamiltonian. We define

$$\langle x; k | H_e(x) | x; l \rangle = W_{kl}(x), \tag{13}$$

so that $H_e(x) \longleftrightarrow W_{kl}(x)$. Note that if the working basis is the adiabatic basis (at least for some range of indices k, l), then

$$W_{kl}(x) = \varepsilon_k(x)\,\delta_{kl}.\tag{14}$$

The momentum $\mathbf{P}_{\alpha} = -i\hbar \nabla_{\alpha}$, which in the molecular representation represents the kinetic momentum of nucleus α , is transformed to the Born-Oppenheimer representation according to

$$\mathbf{P}_{\alpha} \longleftrightarrow \mathbf{P}_{\alpha} \, \delta_{kl} - i\hbar \, \mathbf{F}_{\alpha;kl}(x). \tag{15}$$

Equation (15) illustrates an important theme of this article. The operator \mathbf{P}_{α} is the differential operator $-i\hbar\nabla_{\alpha}$ in all representations, but it only represents physically the kinetic momentum of nucleus α in the molecular representation. In the Born-Oppenheimer representation, there is a correction term, as shown (which, it turns out, is related to the electronic momentum).

Now suppose that we have operators *A*, *B* and *C*, and that $A \leftrightarrow A_{kl}$, $B \leftrightarrow B_{kl}$ and $C \leftrightarrow C_{kl}$. Then we have some theorems. First, if $B = A^{\dagger}$, then $B_{kl} = (A_{lk})^{\dagger}$. Next, if C = AB, then

$$C_{kl} = \sum_{p} A_{kp} B_{pl}.$$
 (16)

Next, if $\Psi_1 \longleftrightarrow \psi_{1k}$ and $\Psi_2 \longleftrightarrow \psi_{2k}$, then

$$\langle \Psi_1 | \Psi_2 \rangle_{Xr} = \sum_k \langle \psi_{1k} | \psi_{2k} \rangle_X \tag{17}$$

where the subscripts X, Xr indicate the variables that are integrated over in taking the scalar product. (If no subscript is given, integration over **r** alone is implied, as in (6).) Finally, matrix elements of operators transform between the two representations according to

$$\langle \Psi_1 | A | \Psi_2 \rangle_{Xr} = \sum_{kl} \langle \psi_{1k} | A_{kl} | \psi_{2l} \rangle_X.$$
(18)

As for the molecular Hamiltonian (1), when transformed to the Born-Oppenheimer representation, $H \longleftrightarrow H_{kl}$, it becomes

$$H_{kl} = \sum_{\alpha=1}^{N} \frac{1}{2M_{\alpha}} \sum_{p} \left[\mathbf{P}_{\alpha} \, \delta_{kp} - i\hbar \, \mathbf{F}_{\alpha;kp}(x) \right] \cdot \left[\mathbf{P}_{\alpha} \, \delta_{pl} - i\hbar \, \mathbf{F}_{\alpha;pl}(x) \right] + W_{kl}(x), \tag{19}$$

where the sum on p indicates a multiplication of matrices of operators, as in (16). We expand this according to

$$H_{kl} = H_{0,kl} + \hbar H_{1,kl} + \hbar^2 H_{2,kl},$$
(20)

where

$$H_{0,kl} = \left(\sum_{\alpha=1}^{N} \frac{\mathbf{P}_{\alpha}^{2}}{2M_{\alpha}}\right) \delta_{kl} + W_{kl}(\mathbf{X}), \qquad (21a)$$

$$H_{1,kl} = -i \sum_{\alpha=1}^{N} \frac{1}{2M_{\alpha}} [\mathbf{P}_{\alpha} \cdot \mathbf{F}_{\alpha;kl}(\mathbf{X}) + \mathbf{F}_{\alpha;kl}(\mathbf{X}) \cdot \mathbf{P}_{\alpha}], \qquad (21b)$$

$$H_{2,kl} = -\sum_{\alpha=1}^{N} \sum_{p} \frac{1}{2M_{\alpha}} \mathbf{F}_{\alpha;kp}(\mathbf{X}) \cdot \mathbf{F}_{\alpha;pl}(\mathbf{X}).$$
(21c)

Note that, since $\mathbf{F}_{\alpha;kl}(x)$ is real, the two terms in the square brackets in (21b) are Hermitian conjugates of each other, so the sum is Hermitian. Overall, we have $(H_{1,kl})^{\dagger} = H_{1,lk}$, so that H_1 , as a matrix of nuclear operators, is Hermitian.

The molecular Schrödinger equation, which in the molecular representation is

$$H\Psi(\mathbf{X},\mathbf{r}) = E\Psi(\mathbf{X},\mathbf{r}),\tag{22}$$

or, in its time-dependent version,

$$H\Psi(\mathbf{X},\mathbf{r},t) = i\hbar \frac{\partial \Psi(\mathbf{X},\mathbf{r},t)}{\partial t},$$
(23)

becomes, in the Born-Oppenheimer representation,

$$\sum_{l} H_{kl} \, \psi_l(\mathbf{X}) = E \, \psi_k(\mathbf{X}), \tag{24}$$

or, in its time-dependent version,

$$\sum_{l} H_{kl} \, \psi_l(\mathbf{X}, t) = i\hbar \frac{\partial \psi_k(\mathbf{X}, t)}{\partial t}.$$
(25)

D. Born-Oppenheimer Ordering

The standard Born-Oppenheimer ordering parameter is $\kappa = (m_e/M)^{1/4}$, where *M* is an average or typical nuclear mass (Born and Oppenheimer (1927); Born and Huang (1954)). By this definition, κ has the numerical value of roughly 10^{-1} . A different approach is to scale the nuclear masses by κ^{-4} , that is, to replace the masses M_{α} in the Hamiltonian by M_{α}/κ^4 , where κ is a scaling parameter that ranges from $\kappa = 0$ (for complete adiabatic separation of the length and time scales) to $\kappa = 1$ (which gives the physical values). Actually, the limit $\kappa \to 0$ is singular since the nuclear de Broglie wavelength goes to zero and the nuclear wave function is no longer defined. But many physical quantities are well defined in the limit $\kappa \to 0$. By making κ a variable the Hamiltonian becomes parameterized by κ , and thereby represents a family of dynamical systems. We can think of a curve through the space of Hamiltonians, parameterized by κ . Then we have a solvable or at least simplified system as $\kappa \to 0$.

In the following we shall adopt the second approach, in which κ is a variable. Then we can speak of the order of magnitude of various quantities in terms of the power of κ by which they scale as $\kappa \to 0$. For example, since the electronic Hamiltonian does not depend on the nuclear masses, it is independent of κ and everything associated with it (the energy eigenvalues $\varepsilon_k(x)$, the derivative couplings $\mathbf{F}_{\alpha:kl}(x)$, etc.), are independent of κ and therefore of order $\kappa^0 = 1$.

To say a quantity is of order $\kappa^0 = 1$ in this sense does not necessarily imply that its numerical value is close to unity, since there may be other reasons (besides the values of the nuclear masses or momenta) why it is large or small. For example, in a scattering problem X_{α} may take on large values, independent of κ , and the derivative couplings in the adiabatic basis diverge as a degeneracy or conical intersection is approached. In terms of the κ scaling, however, these quantities are of order unity. On the other hand, if there is no other reason why a given quantity should be

large or small, and if it is of order κ^n in the scaling sense, then it should have a numerical value in atomic units of roughly 10^{-n} .

In addition to the scaling of the nuclear masses, we must also indicate how the dynamical variables scale with κ as we take $\kappa \to 0$. For small vibrations about an equilibrium, the displacement from equilibrium $\mathbf{X}_{\alpha} - \mathbf{X}_{0,\alpha}$ is of order $\kappa^1 = \kappa$, as discussed by Born and Huang (1954), while the nuclear momentum \mathbf{P}_{α} is order κ^{-1} and the nuclear velocity $\mathbf{V}_{\alpha} = \mathbf{P}_{\alpha}/M_{\alpha}$ is of order κ^3 . Thus, the time require to execute an oscillation of amplitude of order κ is κ^{-2} , and the vibrational frequency is of order κ^2 . In this analysis we are imagining that as we move through the space of dynamical systems by taking $\kappa \to 0$, we are also changing the space of wave functions or quantum states that we are considering. For the problem of small vibrations, we take this limit in such a way that the vibrational quantum number is constant, which causes the spatial extent of the wave function to shrink in proportion to κ . A nice discussion of this (small-amplitude) ordering is given by Mead and Moscowitz (1967).

Modern applications such as isomerization, scattering and photoexcitation problems involve motions in which the nuclei are not restricted to a small range around an equilibrium, but rather move over distances of the order of an atomic unit or larger. For these applications a different assumption about the scaling of the dynamical variables is required. The modified scaling assumptions required for such applications have been discussed in detail by Mead (2006). See also Mátyus and Teufel (2019). In the following we add to this discussion and show how it applies to our approach to Moyal perturbation theory.

We shall call the modified scaling law the "large-amplitude" scaling. In this scaling we imagine that the kinetic or potential energy of the nuclei remains constant as $\kappa \to 0$. For example, if a nucleus climbs half way up a potential well when $\kappa = 1$, we require that it continue to do so as we take $\kappa \to 0$. We do this because we want the qualitative features of the simplified problem as $\kappa \to 0$ to be the same as those of the physical problem at $\kappa = 1$. This means, for example, that bound state quantum numbers, to the extent those are meaningful for large-amplitude motions, go to infinity as $\kappa \to 0$.

If the kinetic energy $\mathbf{P}_{\alpha}^2/2M_{\alpha}$ is independent of κ as $\kappa \to 0$, as required by the large-amplitude ordering, then, given the scaling of M_{α} , we conclude that \mathbf{P}_{α} is order κ^{-2} (therefore larger by a factor of $1/\kappa$ than in the small-amplitude scaling). Nuclear velocities $\mathbf{V}_{\alpha} = \mathbf{P}_{\alpha}/M_{\alpha}$ are therefore of order κ^2 and the time required for a nucleus to travel a distance of the order of one atomic unit is of order κ^{-2} . Thus nuclear motions are executed ever more slowly as $\kappa \to 0$, as expected in a system that can be considered "adiabatic."

The spatial scale of an electronic energy eigenfunction $\phi_{ak}(\mathbf{X}; \mathbf{r})$ for fixed *k*, that is, the electronic de Broglie wave length, is of order $\kappa^0 = 1$, since the electronic Hamiltonian is independent of the nuclear masses, as explained. The same applies to the operator ∇_i . The same also applies to the **X**-dependence of the electronic energy eigenfunctions and the operator ∇_{α} when acting on them. We shall assume that the transformation from the adiabatic basis to the working basis is independent of κ , so the same scale lengths (in both **r** and **X**) apply to the working basis functions $\phi_k(\mathbf{X}; \mathbf{r})$. The nuclear wave functions $\psi_k(\mathbf{X})$ in the Born-Oppenheimer representation, however, have a de Broglie wave length of order κ^2 , since the operator $\mathbf{P}_{\alpha} = -i\hbar \nabla_{\alpha}$ is of order κ^{-2} . Thus, in the expansion (10) the scale lengths of the two factors on the right (ψ_k and ϕ_k) with respect to **X** are quite different.

In the large-amplitude scaling, the nuclear de Broglie wave length is of order κ^2 times smaller than the scale length of the environment (the potential, derivative couplings, etc.) in which the nuclei move. These are otherwise the conditions for WKB theory, which would be more useful for molecular dynamics were it not for the usual difficulties of WKB theory in multidimensional problems (chaotic orbits and the lack of global, invariant Lagrangian manifolds in phase space). Nevertheless, the basic method of this paper, the use of Moyal perturbation theory, is a part of semiclassical theory.

In this paper we use the large-amplitude scaling exclusively. For simplicity we will not insert explicit factors of κ^n to indicate the order of various terms. Instead, the ordering of any term can be obtained by applying the two rules, $M_{\alpha} = O(\kappa^{-4})$ and $\mathbf{P}_{\alpha} = O(\kappa^{-2})$.

In the large-amplitude ordering, the three terms in (21), $H_{0,kl}$, $H_{1,kl}$ and $H_{2,kl}$, are of order $\kappa^0 = 1$, κ^2 and κ^4 , respectively. This ordering will be the basis of the perturbation theory we will carry out in this paper. It is not, however, ordinary perturbation theory, which is based solely on the magnitudes of the unperturbed Hamiltonian and the perturbation, because the nuclear wave functions also have a dependence on κ , that is, the nuclear de Broglie wave length is of order κ^2 .

E. Translational Degrees of Freedom

The Hamiltonian (1) commutes with translations and thus conserves total linear momentum. The manner in which this invariance and conservation law are manifested is straightforward in the molecular representation but more obscure in the Born-Oppenheimer representation. Therefore one is motivated to carry out the exact elimination of the translational degrees of freedom before switching to the Born-Oppenheimer representation. This is a standard procedure (Sutcliffe (2000)) but it requires some algebra and attention to notation.

We begin with the nuclear configuration space, the 3*N*-dimensional space upon which the coordinates are \mathbf{X}_{α} , $\alpha = 1, ..., N$. This is the space upon which purely nuclear wave functions $\psi(\mathbf{X})$ are defined. A coordinate transformation on this space that is useful for separating the translational degrees of freedom is $\mathbf{X}_{\alpha} \rightarrow (\mathbf{X}_{CM}, \mathbf{Y}_{\kappa})$, where \mathbf{X}_{CM} is the center of mass of the nuclei,

$$\mathbf{X}_{\rm CM} = \frac{1}{M_t} \sum_{\alpha=1}^N M_\alpha \mathbf{X}_\alpha, \tag{26}$$

and where \mathbf{Y}_{κ} , $\kappa = 1, ..., N - 1$, are a set of Jacobi vectors (see Appendix B). Here M_t is the total nuclear mass,

$$M_t = \sum_{\alpha=1}^N M_\alpha.$$
 (27)

We let the momenta conjugate to $(\mathbf{X}_{CM}, \mathbf{Y}_{\kappa})$ be $(\mathbf{P}_t, \mathbf{Q}_{\kappa})$. Then it turns out that \mathbf{P}_t , the momentum conjugate to \mathbf{X}_{CM} , is the total momentum of the nuclei,

$$\mathbf{P}_t = \sum_{\alpha=1}^N \mathbf{P}_{\alpha}.$$
 (28)

The other details of this coordinate transformation are summarized in Appendix B.

When this transformation is applied to the nuclear kinetic energy, we find

$$\sum_{\alpha=1}^{N} \frac{\mathbf{P}_{\alpha}^2}{2M_{\alpha}} = \frac{\mathbf{P}_t^2}{2M_t} + \sum_{\kappa=1}^{N-1} \frac{\mathbf{Q}_{\kappa}^2}{2\mu_{\kappa}},\tag{29}$$

where μ_{κ} are reduced masses associated with Jacobi coordinates \mathbf{Y}_{κ} .

We require a different coordinate transformation on the molecular configuration space, the $3(N + N_e)$ -dimensional space with coordinates $(\mathbf{X}_{\alpha}, \mathbf{r}_i)$ upon which the molecular wave function $\Psi(\mathbf{X}, \mathbf{r})$ is defined. This can be seen as the composition of two transformations, $(\mathbf{X}_{\alpha}, \mathbf{r}_i) \rightarrow (\mathbf{X}_{\text{CM}}, \mathbf{Y}_{\kappa}, \mathbf{r}_i) \rightarrow (\mathbf{R}_{\text{CM}}, \mathbf{Y}_{\kappa}, \mathbf{s}_i)$, where in the first step we transform the nuclear coordinates as described above, leaving the electronic coordinates \mathbf{r}_i alone, while in the second step we leave the nuclear Jacobi vectors \mathbf{Y}_{κ} alone while introducing two new sets of coordinates defined by

$$\mathbf{R}_{\rm CM} = \frac{M_t \mathbf{X}_{\rm CM} + N_e m_e \mathbf{r}_{\rm CM}}{M_{\rm mol}}$$
(30)

and

$$\mathbf{s}_i = \mathbf{r}_i - \mathbf{X}_{\rm CM}.\tag{31}$$

Here $M_{\rm mol}$ is the total molecular mass,

$$M_{\rm mol} = M_t + N_e m_e, \tag{32}$$

and \mathbf{r}_{CM} is the center of mass of the electrons,

$$\mathbf{r}_{\rm CM} = \frac{1}{N_e} \sum_{i=1}^{N_e} \mathbf{r}_i.$$
(33)

Evidently \mathbf{R}_{CM} is the molecular center of mass (not to be confused with \mathbf{X}_{CM} , the nuclear center of mass) and \mathbf{s}_i is the position of electron *i* relative to the nuclear center of mass.

We denote the momenta conjugate to coordinates $(\mathbf{R}_{CM}, \mathbf{Y}_{\kappa}, \mathbf{s}_i)$ by $(\mathbf{P}_{CM}, \mathbf{Q}_{\kappa}, \mathbf{q}_i)$, where \mathbf{Q}_{κ} , the momentum conjugate to Jacobi vector \mathbf{Y}_{κ} , is the same as produced by the first (purely nuclear) transformation (see (B9)). As for \mathbf{P}_{CM} , the momentum conjugate to \mathbf{R}_{CM} , it turns out to be the total linear momentum of the molecule,

$$\mathbf{P}_{\rm CM} = \mathbf{P}_t + \mathbf{p}_t,\tag{34}$$

where \mathbf{P}_t , the total nuclear momentum, is given by (28), and \mathbf{p}_t , the total electronic momentum, is given by

$$\mathbf{p}_t = \sum_{i=1}^{N_e} \mathbf{p}_i. \tag{35}$$

The other details of this coordinate transformation are given in Appendix C.

When we transform the molecular Hamiltonian (1) to the new coordinates, we obtain

$$H = \frac{\mathbf{P}_{\rm CM}^2}{2M_{\rm mol}} + H_{\rm TR}(\mathbf{Y}, \mathbf{s}, \mathbf{Q}, \mathbf{q}), \tag{36}$$

where the "translation-reduced" molecular Hamiltonian is

$$H_{\mathrm{TR}}(\mathbf{Y}, \mathbf{s}, \mathbf{Q}, \mathbf{q}) = \sum_{\kappa=1}^{N-1} \frac{\mathbf{Q}_{\kappa}^2}{2\mu_{\kappa}} + H_{\mathrm{TR}e}(y; \mathbf{s}, \mathbf{q}),$$
(37)

and the translation-reduced electronic Hamiltonian is

$$H_{\mathrm{TR}e}(y;\mathbf{s},\mathbf{q}) = \sum_{i=1}^{N_e} \frac{\mathbf{q}_i^2}{2m_e} + \sum_{i,j=1}^{N_e} \frac{\mathbf{q}_i \cdot \mathbf{q}_j}{2M_t} + V_{\mathrm{Coul}}(\mathbf{Y},\mathbf{s}).$$
(38)

Here y means a point of the translation-reduced configuration space whose coordinates are \mathbf{Y}_{κ} , and V_{Coul} is independent of \mathbf{R}_{CM} because it is translationally invariant. The translation-reduced configuration space will be discussed more fully in subsection V B (see also Fig. 1). The middle term on the right in (38) is the mass-polarization term.

F. The Translation-Reduced Molecular and Born-Oppenheimer Representations

The Hamiltonian H_{TR} in (37) gives us a "translation-reduced" molecular representation, in which the wave function $\Psi_{\text{TR}}(\mathbf{Y}, \mathbf{s})$ is defined by

$$\Psi(\mathbf{X},\mathbf{r}) = \Psi(\mathbf{R}_{CM},\mathbf{Y},\mathbf{s}) = \exp(i\mathbf{K}\cdot\mathbf{R}_{CM})\Psi_{TR}(\mathbf{Y},\mathbf{s}),\tag{39}$$

which is an eigenfunction of \mathbf{P}_{CM} with eigenvalue $\hbar \mathbf{K}$. This is in the time-independent version; in the time-dependent version we define Ψ_{TR} by

$$\Psi(\mathbf{X},\mathbf{r},t) = \Psi(\mathbf{R}_{\mathrm{CM}},\mathbf{Y},\mathbf{s},t) = \exp[i(\mathbf{K}\cdot\mathbf{R}_{\mathrm{CM}}-\hbar\mathbf{K}^{2}t/2M_{\mathrm{mol}})]\Psi_{\mathrm{TR}}(\mathbf{Y},\mathbf{s},t).$$
(40)

The wave function Ψ_{TR} satisfies a translation-reduced version of the Schrödinger equation,

$$H_{\rm TR}\Psi_{\rm TR}(\mathbf{Y},\mathbf{s}) = E_{\rm TR}\Psi_{\rm TR}(\mathbf{Y},\mathbf{s}),\tag{41}$$

or its time-dependent version,

$$H_{\rm TR}\Psi_{\rm TR}(\mathbf{Y},\mathbf{s},t) = i\hbar \frac{\partial \Psi_{\rm TR}(\mathbf{Y},\mathbf{s},t)}{\partial t}.$$
(42)

Once we have achieved the translation-reduced molecular representation of the system we can expand in an electronic basis and create a translation-reduced Born-Oppenheimer representation, in which the state of the system is represented by an infinite-dimensional vector of nuclear wave functions $\psi_{\text{TR},k}(\mathbf{Y})$, defined on the translation-reduced configuration space. The potential energy surfaces defined by the eigenvalues $\varepsilon_{\text{TR},k}(y)$ are not the same as before, because the translationreduced electronic Hamiltonian (37) contains mass-polarization terms, in contrast to the original electronic Hamiltonian (2), whose eigenvalues are $\varepsilon_k(x)$.

Thus we now have several representations of the quantum mechanics of the molecule, including the original or "lab" molecular and Born-Oppenheimer representations, and now "translationreduced" versions of these. The situation is summarized in Table I. Once we have achieved the Born-Oppenheimer representation, either the lab or translation-reduced, the Hamiltonian is an infinite-dimensional matrix of nuclear operators, which we can block-diagonalize by means of Moyal perturbation theory. This carries us to the last column of Table I, the "dressed" or blockdiagonalized Born-Oppenheimer representation. We turn now to a detailed description of this last step.

	Molecular	Born-Opp.	Dressed Born-Opp.
Lab	$\Psi(\mathbf{X},\mathbf{r})$	$\pmb{\psi}_k(\mathbf{X})$	$ar{m{\psi}}_k({f X})$
	Н	H_{kl}	$ar{H}_{kl}$
	(1)	(10), (11)	(58)
Translation- Reduced	$\Psi_{\rm TR}({\bf Y},{\bf s})$	$\psi_{\mathrm{TR},k}(\mathbf{Y})$	$ar{\psi}_{\mathrm{TR},k}(\mathbf{Y})$
	H_{TR}	$H_{\mathrm{TR},kl}$	$ar{H}_{\mathrm{TR},kl}$
	(37), (39), (40)	_	_

TABLE I. Six representations of the quantum mechanics of a molecule. In each case, the relevant wave function and Hamiltonian are shown, as well as references to relevant equations. In the molecular representation (first column) the wave function Ψ or Ψ_{TR} is a function defined on the combined nuclear-electronic configuration space. In the Born-Oppenheimer representation (second column), the wave function ψ_k or $\psi_{TR,k}$ is an infinite-dimensional vector of wave functions defined on the nuclear configuration space. The same is true of the dressed representations (third column); in this representation, the Hamiltonian is blockdiagonal. Sometimes we will call the Born-Oppenheimer representation (second column) the "original" Born-Oppenheimer representation, to distinguish it from the dressed representations in the third column.

III. MOYAL PERTURBATION THEORY ON A SINGLE SURFACE

The purpose of this section is to describe in detail the Moyal method for the block diagonalization of the Born-Oppenheimer Hamiltonian. For a somewhat different formalism, see Panati, Spohn, and Teufel (2002); Mátyus and Teufel (2019).

For this purpose we can work either with the lab representations or the translation-reduced. In some ways it is preferable to carry out the exact separation of the translational degrees of freedom first, and then to block-diagonalize the Hamiltonian. But much of the current literature, for example, that devoted to surface hopping, uses the lab representations (so also the treatment of Mátyus and Teufel (2019)). Therefore in this section we shall illustrate Moyal diagonalization of the Born-Oppenheimer Hamiltonian in the lab representations, effectively working across the first row of Table I from left to right. For simplicity we will do this for a single surface problem, deferring multiple surfaces to subsection IV A. We select one surface k = n, so that *n* is fixed for the duration of this section (Sec. III).

In subsections III A–III C we use unaccented symbols for operators, for example, H for the Hamiltonian. Starting in subsection III D, however, and continuing to the end of the paper, it is necessary to distinguish between operators, which are represented with a hat, and their Weyl symbols, which are *c*-number functions on the classical phase space and which are denoted without the hat. For example, starting in subsection III D, the Hamiltonian operator is \hat{H} while its Weyl symbol is H. Weyl symbols are explained in Appendix A.

A. Basis Vectors and the Privileged Subspace

We let $|x;n\rangle = |ax;n\rangle$, so that in the one case k = n the working basis vector and the adiabatic basis vector are the same. We define the "privileged subspace" of the electronic Hilbert space as

$$\mathscr{S}(x) = \operatorname{span}\{|ax;n\rangle\} \tag{43}$$

and we define $\mathscr{S}^{\perp}(x)$ as the complementary, orthogonal subspace. The working basis vectors $|x;k\rangle$ for $k \neq n$ are either the adiabatic basis vectors $|ax;k\rangle$ for $k \neq n$ or another choice to be specified. Both the working basis vectors $|x;k\rangle$ and the adiabatic basis vectors $|ax;k\rangle$ for $k \neq n$ span $\mathscr{S}^{\perp}(x)$, so they are linear combinations of one another. The subspace $\mathscr{S}(x)$ is one-dimensional, while $\mathscr{S}^{\perp}(x)$ is infinite-dimensional. The projection operator onto the privileged subspace is

$$P(x) = |ax;n\rangle\langle ax;n|,\tag{44}$$

while the complementary projector is Q(x) = 1 - P(x). Equation (44) gives P(x) in the molecular representation; on transforming to the Born-Oppenheimer representation it becomes

$$P(x) \longleftrightarrow P_{kl} = \delta_{nk} \,\delta_{nl}. \tag{45}$$

B. The Born-Oppenheimer Approximation

Since $|x;n\rangle = |ax;n\rangle$, the diagonal matrix element W_{nn} of the electronic Hamiltonian satisfies

$$W_{nn}(x) = \mathcal{E}_n(x). \tag{46}$$

In addition, since $|x;n\rangle$ is orthogonal to $\mathscr{S}^{\perp}(x)$, we have $W_{nk}(x) = W_{kn}(x) = 0$ for $k \neq n$, and the matrix $W_{kl}(x)$ is block-diagonal. The blocks in question are the 1×1 block and the infinitedimensional block corresponding to $\mathscr{S}(x)$ and $\mathscr{S}^{\perp}(x)$, respectively. An operator that is blockdiagonal does not couple $\mathscr{S}(x)$ with $\mathscr{S}^{\perp}(x)$. The terms in the Hamiltonian (21) that do couple $\mathscr{S}(x)$ and $\mathscr{S}^{\perp}(x)$ are the off-block-diagonal elements of H_1 and H_2 , that is, elements with indices (nk) or (kn) for $k \neq n$. These matrix elements involve the derivative couplings and are small in the κ ordering, either of order κ^2 or κ^4 . In particular, they are small in comparison to the diagonal element at zeroth order,

$$H_{0,nn} = \sum_{\alpha=1}^{N} \frac{\mathbf{P}_{\alpha}^2}{2M_{\alpha}} + \varepsilon_n(x), \qquad (47)$$

which is of order κ^0 .

In the literature the "Born-Oppenheimer approximation" usually means approximating the molecular wave function as

$$\Psi(\mathbf{X},\mathbf{r}) = \psi(\mathbf{X})\,\phi_{an}(\mathbf{X};\mathbf{r}),\tag{48}$$

that is, a version of (10) with a single term k = n and employing the adiabatic, energy eigenfunction $\phi_{an}(\mathbf{X};\mathbf{r})$. But if we throw away the small, off-block-diagonal terms of the molecular Hamiltonian (21), then the molecular Schrödinger equation (22) has solutions of the form (48), that is, (24) has solutions in which ψ_k is nonzero only in the one slot k = n. The same is true of the time-dependent versions of the molecular Schrödinger equation, (23) and (25), which have solutions of the form (48) in which both $\Psi(\mathbf{X}, \mathbf{r}, t)$ and $\psi(\mathbf{X}, t)$ depend on time but the adiabatic basis state $\phi_{an}(\mathbf{X}; \mathbf{r})$ does not.

Therefore we shall regard the "Born-Oppenheimer approximation," in the context of a singlesurface problem, as one of throwing away the off-block-diagonal matrix elements of H_{kl} . In this approximation, the molecular Schrödinger equation (24) or (25) on the privileged subspace $\mathscr{S}(x)$ becomes

$$H_{\rm BO}(\mathbf{X}, \mathbf{P}) \,\psi(\mathbf{X}) = E \,\psi(\mathbf{X}),\tag{49}$$

or its time-dependent version,

$$H_{\rm BO}(\mathbf{X}, \mathbf{P}) \, \boldsymbol{\psi}(\mathbf{X}, t) = i\hbar \frac{\partial \boldsymbol{\psi}(\mathbf{X}, t)}{\partial t}, \tag{50}$$

where $\psi = \psi_n$ is the one, nonvanishing component of ψ_k , and where $H_{BO} = H_{nn} = H_{0,nn} + \hbar H_{1,nn} + \hbar^2 H_{2,nn}$ is the diagonal element of the Hamiltonian matrix, the effective Hamiltonian operator on this subspace.

To write out H_{BO} explicitly we note first that $H_{1,nn} = 0$ because of (9). As for $H_{2,nn}$, we take the divergence of $\mathbf{F}_{\alpha;nn}(x) = \langle x; n | \nabla_{\alpha} | x; n \rangle = 0$ to obtain the identity,

$$\nabla_{\alpha} \cdot \mathbf{F}_{\alpha;nn}(x) = (\nabla_{\alpha} \langle x; n |) \cdot (\nabla_{\alpha} | x; n \rangle) + \langle x; n | \nabla_{\alpha}^2 | x; n \rangle = 0.$$
(51)

Then using (7) we have

$$\sum_{p} \mathbf{F}_{\alpha;np}(x) \cdot \mathbf{F}_{\alpha;pn}(x) = -\sum_{p} (\nabla_{\alpha} \langle x; n |) | x; p \rangle \cdot \langle x; p | (\nabla_{\alpha} | x; n \rangle)$$
$$= -(\nabla_{\alpha} \langle x; n |) \cdot (\nabla_{\alpha} | x; n \rangle) = \langle x; n | \nabla_{\alpha}^{2} | x; n \rangle,$$
(52)

where in the first step we use (7) and in the last step, (51). Altogether, we obtain

$$H_{\rm BO} = H_{nn} = \sum_{\alpha=1}^{N} \frac{\mathbf{P}_{\alpha}^2}{2M_{\alpha}} + \varepsilon_n(x) - \sum_{\alpha=1}^{N} \frac{1}{2M_{\alpha}} \langle x; n | \nabla_{\alpha}^2 | x; n \rangle.$$
(53)

This is a standard Hamiltonian in Born-Oppenheimer theory. It is the same as the projection of the original molecular Hamiltonian (1) onto the subspace $\mathscr{S}(x)$, that is,

$$P(x)HP(x) \longleftrightarrow \delta_{kn} \,\delta_{nl} \,H_{\rm BO}. \tag{54}$$

The final term in (53) is the "diagonal Born-Oppenheimer correction" (Valeev and Sherrill (2003); Gauss *et al.* (2006); Meek and Levine (2016); Gherib *et al.* (2016); Culpitt *et al.* (2022)). It is otherwise $H_{2,nn}$. It appears to be a contribution to the potential energy, but physically it is a part of the kinetic energy. Away from conical intersections, it is small, being of order κ^4 , and many authors take liberties with it (throwing it away, or throwing part of it away, etc.). In the adiabatic basis it diverges at conical intersections, for the same reason as the derivative couplings (Matsunaga and Yarkony (1998); Yarkony (1997b)), but in our treatment of single-surface problems we shall work in regions far enough from conical intersections that this term is small. See Sec. III F for more details. Near conical intersections we must use a multi-surface approach (see subsection IV A).

The terms in the molecular Hamiltonian that we have thrown away to obtain the Born-Oppenheimer approximation are of order κ^2 or higher and thus are small compared to the diagonal element $H_{0,nn}$ (see (47)). On the other hand, they are large in comparison to $H_{2,nn}$, the diagonal Born-Oppenheimer correction. Therefore there is a question of whether $H_{2,nn}$ is even the correct, second order contribution to the Hamiltonian when the wave function has a single component ψ_n . In fact, a perturbative treatment of the off-block-diagonal terms, one that transforms them away rather than throwing them away, generates a contribution at second order, what we call K_{22} , that is of the same order in κ as $H_{2,nn}$. See Sec. III J.

C. Dressing Transformations

As an alternative to the Born-Oppenheimer approximation, that is, just throwing away terms in the molecular Hamiltonian (21), we will transform them away by means of a series of unitary transformations. Since the terms to be eliminated are small, each unitary transformation is close to the identity and can be conveniently represented as the exponential of another operator, the generator.

In the first step we transform the Hamiltonian by the unitary operator $U_1 = \exp(\hbar G_1)$, where for convenience we split off a factor of \hbar from the anti-Hermitian generator G_1 . The transformed Hamiltonian H' can be written as an exponential series of iterated commutators,

$$H' = U_1 H U_1^{\dagger} = e^{\hbar G_1} H e^{-\hbar G_1} = H + \hbar [G_1, H] + \frac{\hbar^2}{2!} [G_1, [G_1, H]] + \dots$$
(55)

Although we are writing H, G_1 and U_1 without matrix indices, we will work in the Born-Oppenheimer representation and all such symbols should be understood to be matrices of nuclear operators, for example, H_{kl} , $G_{1,kl}$, etc. Likewise, commutators of operators are interpreted as matrix commutators, for example,

$$[G_1, H]_{kl} = \sum_{p} (G_{1,kp} H_{pl} - H_{kp} G_{1,pl}).$$
(56)

The products that appear in (56) are operator products, as in (16), and the ordering must be respected.

The strategy will be to choose G_1 so as to eliminate the off-block-diagonal terms in H to order κ^2 . As it turns out, this means that G_1 itself is of order κ^2 . This leaves behind off-block-diagonal terms in H' that are of order κ^4 . To eliminate these, we perform a second unitary transformation, $H'' = U_2 H' U_2^{\dagger}$, where $U_2 = \exp(\hbar^2 G_2)$. Operator G_2 is a second generator that turns out to be of order κ^4 . Proceeding this way we envision an infinite sequence of unitary operators,

$$U = \dots U_3 U_2 U_1, \tag{57}$$

and a transformed Hamiltonian \bar{H} that is formally free of off-block-diagonal terms to all orders,

$$\bar{H} = UHU^{\dagger} = \dots U_3 U_2 U_1 H U_1^{\dagger} U_2^{\dagger} U_3^{\dagger} \dots$$
(58)

These unitary transformations create a sequence of new representations for the state of the system, what we will call "dressed" representations. Thus we can distinguish between the "original" (that is, undressed) Born-Oppenheimer representation, and successive dressed versions of it. See Table I. If the wave function is ψ in the original Born-Oppenheimer representation (where ψ is short for the infinite-dimensional vector ψ_k), then the wave function in the final, dressed representation is $\bar{\psi} = U\psi$ (that is, it is the vector $\bar{\psi}_k = \sum_l U_{kl}\psi_l$). Then the Schrödinger equation in the final, dressed representation is $\bar{H}\bar{\psi} = E\bar{\psi}$ or its time-dependent version, where \bar{H} has no offblock-diagonal terms. Therefore there are solutions $\bar{\psi}$ that are nonzero only in the one slot k = n, and the effective Hamiltonian for this component is the diagonal element \bar{H}_{nn} . This is the (scalar) Hamiltonian driving this one component, which takes the place of H_{BO} within dressing theory. We now explain the dressing transformations in detail.

D. First Order Transformation and Weyl Symbols

In this subsection we begin by assuming that the working basis is the adiabatic basis, that is, $|x;k\rangle = |ax;k\rangle$ for all *k* (not just k = n). This assumption causes (14) to hold for all *k*, *l*. It simplifies the derivation somewhat but leads to difficulties that we will explain when we encounter them.

In this subsection and henceforth we must change notation in order to distinguish between nuclear operators and their corresponding Weyl symbols. See Appendix A for the definition of the Weyl symbol of an operator and for the invertible mapping between operators and their corresponding Weyl symbols. As in Appendix A, we will place hats on nuclear operators such as $\hat{\mathbf{X}}_{\alpha}$ or $\hat{\mathbf{P}}_{\alpha}$, while omitting the hats on the corresponding classical quantities or Weyl symbols \mathbf{X}_{α} or \mathbf{P}_{α} . The notation \hat{x} or \hat{p} of Appendix A is to be identified with $\hat{\mathbf{X}}$ or $\hat{\mathbf{P}}$ here. We use the Weyl symbol correspondence only for nuclear operators, not electronic ones, so we omit the hats on purely electronic operators such as the electronic Hamiltonian H_e .

Thus, we should go back and put hats on operators in equations such as (21) to conform with the new notation. We will not do this, but we note that when the (hatted, operator) versions of (21) are converted into Weyl symbols, we obtain

$$H_{0,kl} = \left(\sum_{\alpha=1}^{N} \frac{\mathbf{P}_{\alpha}^{2}}{2M_{\alpha}}\right) \delta_{kl} + W_{kl}(\mathbf{X}),$$
(59a)

$$H_{1,kl} = -i \sum_{\alpha=1}^{N} \frac{1}{M_{\alpha}} \mathbf{P}_{\alpha} \cdot \mathbf{F}_{\alpha;kl}(\mathbf{X}),$$
(59b)

$$H_{2,kl} = -\sum_{\alpha=1}^{N} \sum_{p} \frac{1}{2M_{\alpha}} \mathbf{F}_{\alpha;kp}(\mathbf{X}) \cdot \mathbf{F}_{\alpha;pl}(\mathbf{X}), \qquad (59c)$$

where now $W_{kl}(x) = \delta_{kl} \varepsilon_k(x)$ on account of our assumptions about the working basis. These look almost the same as (21) but now they are symbol equations and note that the expression for $H_{1,kl}$ has simplified. In fact, this is an example of (A7), illustrating the convenient manner in which Weyl symbols handle operator ordering. Weyl symbols are *c*-numbers, and can be multiplied in any order. Equations (59) contain exactly the same information as (21); no information is lost on transforming to Weyl symbols.

Likewise, in the case of (55), to conform with the new notation we replace H, H' and G_1 by \hat{H} , \hat{H}' and \hat{G}_1 , respectively, indicating operators (actually, matrices of operators). Then converting these operator equations into symbols, we have the symbol equation,

$$H' = H + \hbar [G_1, H]_* + \frac{\hbar^2}{2} [G_1, [G_1, H]_*]_* + \dots,$$
(60)

where the notation $[G_1, H]_*$ is explained in Appendix A. Note in particular the definition (A8) of the *-product and the discussion in subsection A3 of commutators of matrices of operators, especially (A15).

Let us now write out the expansion of the symbol of the molecular Hamiltonian, $H = H_0 + \hbar H_1 + \hbar^2 H_2$, where the individual terms are given by (59), and substitute it into (60). We obtain

$$H' = H_0 + \hbar \left(H_1 + [G_1, H_0]_* \right) + \hbar^2 \left(H_2 + [G_1, H_1]_* + \frac{1}{2} [G_1, [G_1, H_0]_*]_* \right) + \dots$$
(61)

The commutators shown contain an implicit \hbar -ordering. To expand this out it is convenient to define

$$T_1 = [G_1, H_0]_* = T_{10} + \hbar T_{11} + \dots,$$
(62)

where the 1-subscript on T_1 is a reminder that this term is associated with generator G_1 , and where the expansion follows (A15). In particular, we have

$$T_{10,kl} = [G_1, H_0]_{kl} = \sum_p (G_{1,kp} H_{0,pl} - H_{0,kp} G_{1,pl}),$$
(63a)

$$T_{11,kl} = \frac{i}{2} \sum_{p} \left(\{ G_{1,kp}, H_{0,pl} \} - \{ H_{0,kp}, G_{1,pl} \} \right),$$
(63b)

where $T_{10} = [G_1, H_0]$ is the ordinary commutator of matrices of Weyl symbols (without the *) and where the brackets in (63b) are Poisson brackets. Then (61) becomes

$$H' = H_0 + \hbar (H_1 + T_{10}) + \hbar^2 \left(H_2 + [G_1, H_1] + T_{11} + \frac{1}{2} [G_1, T_{10}] \right) + \dots$$
(64)

where we have dropped the *-subscript in commutators at order \hbar^2 since the ordinary, matrix commutator is the leading term in \hbar (see subsection A 3) and the higher order corrections are beyond the order to which we are working.

The Moyal expansion of the *-product (A10) obviously generates a power series in \hbar . As pointed out at the end of subsection II D, however, each power of \hbar in the \hbar -ordering of the molecular Hamiltonian (21) is coupled to one power of κ^2 . That is, terms H_0 , H_1 and H_2 are of order κ^0 , κ^2 and κ^4 , respectively. Likewise, in the Moyal expansion (A12), which is nominally an expansion in \hbar , each successive bracket involves one higher derivative with respect to momentum (see (A11)). In this work the operators of interest are polynomials in the nuclear momentum $\hat{\mathbf{P}}_{\alpha}$, which is of order κ^{-2} , and their Weyl symbols are corresponding polynomials in \mathbf{P}_{α} . Therefore each momentum derivative kills one factor of \mathbf{P}_{α} and raises the order of κ by 2. In other words, the Moyal expansion couples every power of \hbar with a power of κ^2 . The result is that every term of order \hbar^n that we encounter in the Moyal expansion behaves in the large-amplitude scaling as of order κ^{2n} . In the following when we say, "first order," we shall mean, order \hbar or κ^2 , while "second order" means order \hbar^2 or κ^4 , etc.

The fact that the \hbar ordering is closely related to the κ ordering should not be surprising, in view of the close connection between Born-Oppenheimer theory, especially in the large-amplitude ordering, with semiclassical theory (for example, the fact that the nuclear de Broglie wavelength is of order κ^2). But it is a semiclassical theory only in the nuclear variables, not the electronic.

E. The Generator G_1

Returning now to (64) we wish to choose G_1 to kill the off-block-diagonal elements of H_1 , so that we will have $H'_{1,nk} = H'_{1,kn} = 0$ for $k \neq n$. Thus, we require

$$[G_1, H_0]_{nk} = -H_{1,nk}, \qquad (k \neq n).$$
(65)

Of the two terms in H_0 (see (59a)) the kinetic energy is a multiple of the identity matrix which commutes with the matrix G_1 , while for the potential energy we use (14) since we are working (for the time being) in the adiabatic basis. Then the commutator in (65) is easy to evaluate and we obtain,

$$G_{1,nk}(\varepsilon_k - \varepsilon_n) = -H_{1,nk}, \qquad (k \neq n), \tag{66}$$

or,

$$G_{1,nk} = -i\sum_{\alpha=1}^{N} \frac{1}{M_{\alpha}} \frac{\mathbf{P}_{\alpha} \cdot \mathbf{F}_{\alpha;nk}(\mathbf{X})}{\varepsilon_{n}(\mathbf{X}) - \varepsilon_{k}(\mathbf{X})}, \qquad (k \neq n).$$
(67)

If we carry out the same procedure for the (kn)-block, we obtain

$$G_{1,kn} = -i\sum_{\alpha=1}^{N} \frac{1}{M_{\alpha}} \frac{\mathbf{P}_{\alpha} \cdot \mathbf{F}_{\alpha;kn}(\mathbf{X})}{\varepsilon_{k}(\mathbf{X}) - \varepsilon_{n}(\mathbf{X})}, \qquad (k \neq n),$$
(68)

so that $G_{1,kn} = -G_{1,nk}^*$, as required if \hat{G}_1 is to be anti-Hermitian. These are symbol equations; if we want the operator \hat{G}_1 then we can convert back to operators. For example, we find

$$\hat{G}_{1,nk} = -i\sum_{\alpha=1}^{N} \frac{1}{2M_{\alpha}} \left[\hat{\mathbf{P}}_{\alpha} \cdot \frac{\mathbf{F}_{\alpha;nk}(\hat{\mathbf{X}})}{\varepsilon_{n}(\hat{\mathbf{X}}) - \varepsilon_{k}(\hat{\mathbf{X}})} + \frac{\mathbf{F}_{\alpha;nk}(\hat{\mathbf{X}})}{\varepsilon_{n}(\hat{\mathbf{X}}) - \varepsilon_{k}(\hat{\mathbf{X}})} \cdot \hat{\mathbf{P}}_{\alpha} \right], \qquad (k \neq n).$$
(69)

in which the ordering of products must be respected. It will be appreciated that Weyl symbols are easier to work with than operators.

Equation (65) is all we require of G_1 ; since it does not determine the on-block components of G_1 , we set these to zero, that is, $G_{1,nn} = 0$ and $G_{1,kl} = 0$ for $k \neq n$ and $l \neq n$. The result is that G_1 is purely off-block-diagonal.

F. Energy Denominators, the Excluded Region, and Landau-Zener

The solution (67) diverges where the energy denominator $\varepsilon_n(x) - \varepsilon_k(x)$ goes to zero. Since $k \neq n$ this can only happen if the energy denominator goes to zero for the nearest neighbor, $k = n \pm 1$. We define the "degeneracy manifold" as the subset of nuclear configuration space where $\varepsilon_n(x) = \varepsilon_k(x)$ for $k = n \pm 1$, that is, where level *n* is degenerate with one of its immediate neighbors. (If *n* is the ground state, then it has only one neighbor, the first excited state.) Degeneracy manifolds are usually called "seams." As *x* approaches the degeneracy manifold, *G*₁ diverges and the perturbation expansion (55) breaks down. Actually, not only does *G*₁ diverge, but so do the derivative couplings and therefore *H*₁ itself. This is shown by the Feynman-Hellman formula,

$$\mathbf{F}_{\alpha;nk}(x) = -\frac{\langle ax; n | \nabla_{\alpha} H_e(x) | ax; k \rangle}{\varepsilon_n(x) - \varepsilon_k(x)}, \qquad (k \neq n),$$
(70)

which applies in the adiabatic basis, as shown. Substituting this into (67), we see that G_1 diverges as $(\varepsilon_n - \varepsilon_k)^{-2}$ as we approach the degeneracy manifold.

If the nearest neighbor energy denominator $\varepsilon_n(x) - \varepsilon_k(x)$ in (67) or in (70) is of order $\kappa^0 = 1$, then the size of the generators $G_{1,nk}$ is determined by the factor $\mathbf{P}_{\alpha}/M_{\alpha}$, the nuclear velocity, which is of order κ^2 . In this case the generators $G_{1,nk}$ (all of them, for all k) are of order κ^2 , and therefore small. But the nearest neighbor $\varepsilon_n(x) - \varepsilon_k(x)$ is of order unity if the distance from the degeneracy manifold, call it Δx , is also of order unity. (This is true in the sense of the κ scaling.) At such distances, the generators $G_{1,nk}$ are small and the first few terms of the series (55) are rapidly decreasing. This is all we require of the series for practical purposes; we do not require convergence. Thus, Moyal perturbation theory, applied to a single-surface problem for the adiabatic separation of the privileged surface n from the others, is valid when Δx is of order $\kappa^0 = 1$.

It is also valid for smaller Δx , but not too much. As we approach the degeneracy manifold, the nearest-neighbor $G_{1,nk}$ grows and when it becomes of order unity the series (55) breaks down, that is, the terms are no longer rapidly decreasing. Since $G_{1,nk}$ goes as $(\varepsilon_n - \varepsilon_k)^{-2}$, the breakdown occurs when the energy difference is of order κ , which implies that Δx is also of order κ . This defines, at least in terms of its κ -scaling, the region that must be excluded for the validity of single-surface methods. Inside the excluded region it will be necessary to use multiple-surface methods, which are discussed in subsection IV A.

Since the nuclear de Broglie wave length is of order κ^2 , the excluded region, which is within a distance of order κ from the degeneracy manifold, contains on the order of κ^{-1} de Broglie wave lengths. The actual number in a practical situation depends on the magnitudes of various matrix elements, the actual value of the nuclear momentum and other quantities. But making a first stab based on the κ -scaling alone, we estimate that the excluded region extends roughly 0.1 atomic unit away from the degeneracy manifold, and that it contains roughly 10 nuclear de Broglie wavelengths in this distance. This is all in the large-amplitude scaling.

In the case of small molecules, it is possible to visualize the excluded region geometrically, but it is necessary to distinguish between the nuclear configuration space and "shape space." Nuclear configuration space is the space \mathbb{R}^{3N} upon which the nuclear positions in the lab frame, \mathbf{X}_{α} , are coordinates. Nuclear configuration space contains information about the center-of-mass position and the orientation of the molecule. Shape space, on the other hand, is the (3N - 6)-dimensional space in which the center-of-mass position and orientation of the molecule are ignored. Coordinates on shape space are translationally and rotationally invariant quantities such as bond lengths and angles. In the case of triatomics (N = 3), shape space is topologically one half of \mathbb{R}^3 (Littlejohn and Reinsch (1995)), while the codimension 2 degeneracy manifold is a curve in this space (Yarkony (1996, 2001, 2004b,a)). The excluded region is therefore a tube whose width is of order κ surrounding this curve. This curve may bifurcate (Gordon, Glezaku, and Yarkony

(1998); Yarkony (1997a)), and, where it does, the excluded region surrounds it like the sleeves of a shirt. The geometry of shape space and its relation to nuclear configuration space is explained in detail by Littlejohn and Reinsch (1997).

The size of the excluded region depends on the nuclear momentum, and grows in proportion to the square root of the latter. If we change the scaling of the nuclear momentum, we find other scalings for the size of the excluded region. In the small-amplitude scaling \mathbf{P}_{α} is of order κ^{-1} , as explained above. This means that the size of the excluded region is of order $\kappa^{3/2}$, or roughly 0.03 atomic units. Conversely, if we make \mathbf{P}_{α} larger than the large-amplitude value of κ^{-2} , we can make the excluded region larger. If we make \mathbf{P}_{α} large enough, we can make the size of the excluded region as large as order $\kappa^0 = 1$, which means that the excluded region is everywhere and the perturbation expansion fails everywhere. This occurs when \mathbf{P}_{α} is of order κ^{-4} , so that the nuclear velocity $\mathbf{P}_{\alpha}/M_{\alpha}$ is of order $\kappa^0 = 1$. This means that the nuclear velocity is of the same order as the electron velocity, at which point naturally there is no adiabatic separation of time scales. Such momenta are too large to be of much interest for the usual chemical processes, but it is nice to see that the theory breaks down completely where it should.

The generator G_1 has the same structure as the one-dimensional, Landau-Zener transition probability (Landau (1932); Zener (1932); Stückelberg (1932); Nikitin and Umanskii (1984)), which can be written as

$$\exp\left(-\frac{2\pi V_{12}^2}{\hbar v_0 |V_{11}' - V_{22}'|}\right).$$
(71)

Apart from the notation and numerical constants, the dimensionless quantity in the exponential has the same structure as the reciprocal of $\hbar G_{1,nk}$, which is also dimensionless. To see this we identify v_0 in (71), which refers to the velocity with which the one-dimensional transition point is passed, with $\mathbf{P}_{\alpha}/M_{\alpha}$; we identify V_{12} in (71), which is otherwise one half the minimum energy difference between two levels at the transition point, with $(\varepsilon_n - \varepsilon_k)/2$; and we identify $V'_{11} - V'_{22}$ in (71), which is the *x*-derivative of the difference in the diagonal elements of the Hamiltonian matrix, with $\langle x; n | \nabla_{\alpha} H_e | x; k \rangle$. Thus, the quantity $\exp[-\pi/(2\hbar |G_{1,nk}|)]$, for nearest neighbor *k*, behaves something like a local Landau-Zener transition probability, and the excluded region is where this quantity is of order unity.

The excluded region defines what we may think of as the "size of a conical intersection," that is, the distance over which it exerts its influence in preventing the adiabatic separation of potential energy surfaces.

G. Diabatic Basis on $\mathscr{S}^{\perp}(x)$

When nondegenerate, the adiabatic basis vector $|ax;k\rangle$ is only determined by (3) to within a phase, and when degenerate the set of degenerate vectors is only determined to within the choice of an orthonormal frame inside the degenerate eigenspace. Phase and frame conventions can be chosen at each and every point *x* of nuclear configuration space, and some such choice is implied in the notation $|ax;k\rangle$, including in the continuum. But Born-Oppenheimer theory generates derivatives of the basis vectors, for example, inside the derivative couplings, so we must ask whether the phase and frame conventions for the basis $\{|ax;k\rangle\}$ can be made as smooth functions of *x* over some region of interest of the nuclear configuration space.

The answer is no, for topological reasons it is impossible, in general, to make a smooth assignment of phase and frame conventions for the vectors of the adiabatic basis, and singularities (places where the basis vectors are not differentiable with respect to *x*) are inevitable. Such singularities occur on surfaces of two different types. First, when level *k* is degenerate with level $k \pm 1$, the derivatives $\nabla_{\alpha} | ax; k \rangle$ are not defined. The manifolds where this occurs are like the degeneracy manifolds introduced earlier, except they apply to any *k*, not just k = n. In the electrostatic model these manifolds have codimension 2. As we approach such a manifold along a curve, the frame has a limit, but the limit depends on the direction of approach so derivatives are not defined on the manifold itself. (The singularity is like that of the orthonormal frame in 3-dimensional space in spherical coordinates, $(\hat{\mathbf{r}}, \hat{\boldsymbol{\theta}}, \hat{\boldsymbol{\phi}})$, as we approach the *z*-axis.) The second type of surface upon which the adiabatic basis is singular consists of codimension-1 manifolds that emanate from the degeneracy manifolds. These manifolds can be moved about by gauge transformations like a branch cut in complex variable theory. The adiabatic basis vectors suffer a change in sign when crossing such a manifold and so are not differentiable there.

These manifolds (of both types) proliferate as k increases, and present serious conceptual complications when derivatives of the adiabatic basis arise. In addition there is the problem of the continuum, where the concept of a nearest neighbor breaks down. For these reasons we advocate replacing the adiabatic basis on $\mathscr{S}^{\perp}(x)$ with a diabatic basis, which is a choice of an orthonormal frame $\{|dx;k\rangle, k \neq n\}$, with d for "diabatic," that is a smooth function of x. Diabatic bases are usually used only in the subspace $\mathscr{S}(x)$, but here we are using them in $\mathscr{S}^{\perp}(x)$. The fact that a Hilbert space always possesses a discrete basis means that the diabatic basis in $\mathscr{S}^{\perp}(x)$ can be chosen to be discrete; there is no continuum. The fact that the diabatic basis on $\mathscr{S}^{\perp}(x)$ can be chosen to be a smooth function of *x*, at least in some region of nuclear configuration space, is discussed by Littlejohn, Rawlinson, and Subotnik (2022). (For example, one can use parallel transport to create such a basis.) In addition, one can choose a diabatic basis that is invariant under time reversal, so that the derivative couplings $\mathbf{F}_{\alpha;kl}(x)$ are real.

Therefore we now change the definition of the working basis, defining

$$|x;k\rangle = \begin{cases} |ax;n\rangle & k = n, \\ |dx;k\rangle & \text{otherwise.} \end{cases}$$
(72)

In the working basis, the electronic Hamiltonian has matrix elements,

$$\langle x; k | H_e(x) | x; l \rangle = \begin{cases} \varepsilon_n(x) & k = l = n, \\ W_{kl}(x) & k \neq n \text{ and } l \neq n, \\ 0 & \text{otherwise.} \end{cases}$$
(73)

The derivative couplings are still defined by (7) but are now taken with respect to the (new) working basis, and the expressions (21) for the Hamiltonian are still valid.

The adiabatic basis is widely used in the literature on Born-Oppenheimer theory, in which formal sums over an infinite number of basis states, including the continuum, are of frequent occurrance. Our scruples over the use of this basis have nothing to do with the practical impossibility of the numerical evaluation of such sums, but rather with the singularities of the adiabatic basis as x is varied. These present such a confusing picture that it is hard to know what one is really talking about when derivatives with respect to x are taken. Our purpose in switching to a discrete, diabatic basis is to obtain a formalism in which we can be confident that the answers we obtain are correct.

On the other hand, for a fixed value of x the adiabatic basis states do form a complete set, and there is no harm in expanding vectors or operators in terms of them. Therefore for $k \neq n$ we will consider it permissible to use the adiabatic basis vectors $|ax;k\rangle$ in cases where no derivatives of the basis vectors appear and the results are independent of the phase and frame conventions of those vectors, for example, in (76) below. In cases where derivatives of the basis vectors do appear we shall require the diabatic basis (or the working basis, as we are now defining it).

The fact that the adiabatic basis is complete follows from the self-adjointness of the electronic Hamiltonian $H_e(x)$. It has been claimed that the electronic Hamiltonian is not self-adjoint (Sutcliffe (2012)), but we believe the arguments for this are not well founded and that the matter is settled by Jecko (2014).

However, since we are using the one adiabatic basis vector $|ax;n\rangle$ on the privileged subspace, we should address its singularites. We will assume that we are working in the region where the Moyal perturbation theory is valid, so that Δx is larger than order κ and so that we avoid the degeneracy manifold and its singularities. We can also avoid the -1 jumps in the phase of the adiabatic basis if our region is not so large as to encircle a conical intersection; this can be achieved if our region is simply connected (Juanes-Marcos, Althorpe, and Wrede (2005); Althorpe (2006, 2012); Littlejohn, Rawlinson, and Subotnik (2023)). In the following we will assume that $|ax;n\rangle$ is free of singularities in the region under consideration, so that its derivatives are well defined.

H. The Resolvent $R(\varepsilon, x)$

Now that we have changed the working basis, we must return to the calculation of the generator G_1 , starting with (65), and revise it. The (nk)-component of the commutator for $k \neq n$ is now

$$[G_{1}, H_{0}]_{nk} = \sum_{p} G_{1,np} H_{0,pk} - \sum_{p} H_{0,np} G_{1,pk} = \sum_{p \neq n} G_{1,np} W_{pk}(x) - \varepsilon_{n}(x) G_{1,nk}$$

$$= \sum_{p \neq n} G_{1,np} [W_{pk}(x) - \varepsilon_{n}(x) \delta_{pk}] = -H_{1,nk},$$
(74)

where again the kinetic energy part of H_0 does not contribute and where we use (73) for the potential energy. In addition, in the first sum only terms $p \neq n$ contribute since W_{pk} is block-diagonal, and in the second sum only the term p = n contributes, for the same reason.

We now introduce the matrix of the resolvent operator in the working basis, $R_{kl}(\varepsilon, x)$ for $k, l \neq n$, defined by

$$\sum_{p \neq n} \left[\varepsilon \, \delta_{kp} - W_{kp}(x) \right] R_{pl}(\varepsilon, x) = \delta_{kl}, \qquad (k, l \neq n), \tag{75}$$

so that $R(\varepsilon, x)$ is basically the inverse of $\varepsilon - H_e(x)$ on the space $\mathscr{S}^{\perp}(x)$. We also define $R_{kl}(\varepsilon, x) = 0$ when k = n or l = n, so that $R(\varepsilon, x) = 0$ on $\mathscr{S}(x)$ and the matrix $R_{kl}(\varepsilon, x)$ is block-diagonal. The operator $R(\varepsilon, x)$ is given explicitly in the adiabatic basis by

$$R(\varepsilon, x) = \sum_{k \neq n} \frac{|ax; k\rangle \langle ax; k|}{\varepsilon - \varepsilon_k(x)};$$
(76)

by $R_{kl}(\varepsilon, x)$ we mean the matrix elements of this operator in the working basis. This operator is defined as long as ε is not equal to any of the eigenvalues of $H_e(x)$ on $\mathscr{S}^{\perp}(x)$. In particular, $R(\varepsilon_n, x)$ is defined over the region in which we are working, which avoids the degeneracy manifold.

Now we may solve (74) for $G_{1,nk}$,

$$G_{1,nk} = \sum_{l \neq n} H_{1,nl} R_{lk}(\varepsilon_n, x) = -i \sum_{l \neq n} \sum_{\alpha=1}^{N} \frac{1}{M_{\alpha}} \mathbf{P}_{\alpha} \cdot \langle x; n | \nabla_{\alpha} | x; l \rangle \langle x; l | R(\varepsilon_n, x) | x; k \rangle$$

$$= +i \sum_{\alpha=1}^{N} \frac{1}{M_{\alpha}} \mathbf{P}_{\alpha} \cdot (\nabla_{\alpha} \langle x; n |) R(\varepsilon_n, x) | x; k \rangle, \qquad (k \neq n).$$
(77)

In the last step we have used (7) to transfer the derivative to $\langle x; n \rangle$, we have extended the *l*-sum to include the term l = n which vanishes anyway, and then removed the resolution of the identity.

The Feynman-Hellman formula (70) applies only in the adiabatic basis but we can obtain an equivalent result in the diabatic basis. First we note that the resolvent satisfies

$$P(x)R(\varepsilon,x) = R(\varepsilon,x)P(x) = 0, \qquad (78a)$$

$$R(\varepsilon, x) = Q(x)R(\varepsilon, x) = R(\varepsilon, x)Q(x) = Q(x)R(\varepsilon, x)Q(x),$$
(78b)

as well as

$$R(\varepsilon, x) \left[\varepsilon - H_e(x) \right] = \left[\varepsilon - H_e(x) \right] R(\varepsilon, x) = Q(x).$$
(79)

Next we write $[H_e(x) - \varepsilon_n(x)]|x;n\rangle = 0$ and apply ∇_{α} to obtain

$$\left[\nabla_{\alpha}H_{e}(x) - \nabla_{\alpha}\varepsilon_{n}(x)\right]|x;n\rangle + \left[H_{e}(x) - \varepsilon_{n}(x)\right]\nabla_{\alpha}|x;n\rangle = 0.$$
(80)

We multiply this by $R(\varepsilon_n, x)$ to obtain

$$R(\varepsilon_n, x) \nabla_{\alpha} H_e(x) |x; n\rangle = Q(x) \nabla_{\alpha} |x; n\rangle,$$
(81)

where we use (78b) and where *R* moves past the *c*-number $\nabla_{\alpha} \varepsilon_n(x)$ and annihilates $|x;n\rangle$. This can be regarded as a result of first-order perturbation theory, in which $\delta H_e = \sum_{\alpha} \xi_{\alpha} \cdot \nabla_{\alpha} H_e(x)$, where ξ_{α} is a small displacement in nuclear configuration space. It is often needed in Hermitian conjugated form,

$$\left(\nabla_{\alpha}\langle x; n|\right)Q(x) = \langle x; n|\nabla_{\alpha}H_e(x)R(\varepsilon_n, x).$$
(82)

For example, the off-block-diagonal derivative couplings can now be written

$$\mathbf{F}_{\alpha;nk}(x) = -(\nabla_{\alpha}\langle x;n|)|x;k\rangle = -(\nabla_{\alpha}\langle x;n|)Q(x)|x;k\rangle$$
$$= -\langle x;n|\nabla_{\alpha}H_{e}(x)R(\varepsilon_{n},x)|x;k\rangle, \qquad (k \neq n).$$
(83)

This is the analog of the Feynman-Hellman formula (70) in the diabatic basis. The other offdiagonal block is also useful,

$$\mathbf{F}_{\alpha;kn}(x) = -\mathbf{F}_{\alpha;nk}(x)^* = \langle x; k | R(\varepsilon_n, x) \nabla_\alpha H_e(x) | x; n \rangle, \qquad (k \neq n).$$
(84)

Similarly, we can replace R by QR in (77) and then substitute (82) to obtain

$$G_{1,nk} = i \sum_{\alpha=1}^{N} \frac{1}{M_{\alpha}} \mathbf{P}_{\alpha} \cdot \langle x; n | \nabla_{\alpha} H_{e}(x) R(\varepsilon_{n}, x)^{2} | x; k \rangle, \qquad (k \neq n).$$
(85)

Equations (77) and (85) give us alternative versions of the (nk)-components, for $k \neq n$, of G_1 in the diabatic basis. The derivation has nowhere relied on dubious derivatives of the adiabatic basis vectors $|ax;k\rangle$ for $k \neq n$. The components $G_{1,kn} = -G_{1,nk}^*$ of the other off-diagonal block are obtained by the anti-Hermiticity of G_1 . The block-diagonal components of G_1 vanish, as in the calculation using the adiabatic basis, so that G_1 is purely off-block-diagonal.

The operator $R(\varepsilon_n, x)$, when acting on an arbitrary wave function, has an effect whose order of magnitude is its largest eigenvalue, which by (76) is the nearest neighbor value of $1/(\varepsilon_n - \varepsilon_k)$. Thus (85) shows that $G_{1,nk}$ in the diabatic basis behaves like the nearest neighbor value of $(\varepsilon_n - \varepsilon_k)^{-2}$, as in the adiabatic basis.

I. The Second Order Hamiltonian

We have found the generator \hat{G}_1 , whose Weyl symbol is given by (77) or (85), that kills the off-block-diagonal components of \hat{H}_1 . To describe this situation it is convenient to say that an operator or its symbol is "even" if it is block-diagonal, and "odd" if it is purely off-block-diagonal. This terminology is borrowed from the Foldy-Wouthuysen transformation (Foldy and Wouthuysen (1950); Bjorken and Drell (1964)), which closely resembles the procedure we are following here in block-diagonalizing the Born-Oppenheimer Hamiltonian. An arbitrary symbol or operator matrix is neither even nor odd but can be uniquely decomposed into its even and odd parts, which for example in the case of a symbol matrix we write as $A = A^e + A^o$. When we multiply matrices of operators or their Weyl symbols, the product of an even times an even or an odd times an odd matrix is even, while the product of an odd times an even matrix or vice versa is odd. That is, $(AB)^e = A^e B^e + A^o B^o$ and $(AB)^o = A^e B^o + A^o B^e$. The same rule applies to commutators of matrices.

Concerning the operators or symbol matrices discussed so far, H_0 is purely even, $H_0 = H_0^e$, and G_1 is purely odd, $G_1 = G_1^o$. As for H_1 (see (59b)), it vanishes on the \mathscr{S} -block due to the time-reversal invariance of the basis states, that is, $H_{1,nn} = 0$, but it has nonvanishing elements in the \mathscr{S}^{\perp} -block. Altogether, it has both even and odd parts, $H_1 = H_1^e + H_1^o$. Likewise we have $H_2 = H_2^e + H_2^o$. The symbol matrix $T_1 = [G_1, H_0]_* = [G_1^o, H_0^e]_*$ (see (62)) is purely odd, $T_1 = T_1^o$,
as are the terms T_{10} , T_{11} , etc., of its \hbar -expansion. The condition that \hat{G}_1 kill the off-block-diagonal terms in the Hamiltonian through order κ^2 can be written as

$$T_{10} = T_{10}^o = [G_1, H_0] = -H_1^o.$$
(86)

This makes H'_1 (see (64)) purely even,

$$H'_1 = H_1 + T_{10} = H_1 + [G_1, H_0] = H^e_1,$$
(87)

which was the point of the first transformation.

As for H' at second order, we can use (64) and (86) to write

$$H_2' = H_2 + [G_1, H_1^e] + \frac{1}{2}[G_1, H_1^o] + T_{11},$$
(88)

which decomposes into even and odd parts,

$$H_2^{\prime e} = H_2^e + \frac{1}{2} [G_1, H_1^o],$$
(89a)

$$H_2^{\prime o} = H_2^o + [G_1, H_1^e] + T_{11}.$$
(89b)

The unitary transformation $\hat{U}_1 = \exp(\hbar \hat{G}_1)$ has killed the off-block-diagonal elements of the Hamiltonian at first order, but not at second order, as we see from (89b). To kill these we apply a second unitary transformation, $\hat{U}_2 = \exp(\hbar^2 \hat{G}_2)$, defining a second transformed Hamiltonian,

$$\hat{H}'' = e^{\hbar^2 \hat{G}_2} \hat{H}' e^{-\hbar^2 \hat{G}_2} = \hat{H}' + \hbar^2 [\hat{G}_2, \hat{H}'] + \dots$$
(90)

Converting this operator equation to symbols and expanding in \hbar , we find $H_0'' = H_0'$, $H_1'' = H_1'$ and

$$H_2'' = H_2' + [G_2, H_0].$$
(91)

Thus, to kill the off-block-diagonal terms in the Hamiltonian to second order we require

$$[G_2, H_0] = -H_2^{\prime o}, (92)$$

a condition on G_2 that may be compared to condition (86) on G_1 . Combined with (91), (92) shows that $H_2'' = H_2'^e$, which is given by (89a).

Thus we do not need G_2 in order to get H_2'' , the transformed Hamiltonian at second order, which is the more interesting object. Note also that the higher order transformations (\hat{U}_3 , etc.) will not change the Hamiltonian through second order, so that \bar{H} and H'' agree through second order (see (58)). Therefore we will express the answers in terms of \overline{H} , not H''. Finally, we are only interested in the (nn)-component of \overline{H} , which is the effective (scalar) Hamiltonian driving the wave function on surface n.

We will call the Weyl symbol of this scalar Hamiltonian *K*, so that $K = \bar{H}_{nn}$. (The operator itself is \hat{K} .) Then we have $K_0 = H_{0,nn}$, which is given by (47), and $K_1 = H_{1,nn} = 0$. As for $K_2 = \bar{H}_{2,nn}$, it is the (nn)-component of (89a), which has two terms. We write $K_2 = K_{21} + K_{22}$ for these, of which the first term, $K_{21} = H_{2,nn}$, is the diagonal Born-Oppenheimer correction seen in (53). In fact, we have $K = H_{BO} + K_{22}$ to second order, so that K_{22} is another correction, in addition to the diagonal Born-Oppenheimer correction. The new correction K_{22} is

$$K_{22} = \frac{1}{2} [G_1, H_1^o]_{nn} = \frac{1}{2} \sum_{k \neq n} G_{1,nk} H_{1,kn} - \frac{1}{2} \sum_{k \neq n} H_{1,nk} G_{1,kn}.$$
(93)

The sums are only over $k \neq n$ because $G_{1,nn} = H_{1,nn} = 0$.

We work on the first sum. The quantity $G_{1,nk}$ is given by (85), while $H_{1,kn}$ is given by (59b) with $l \rightarrow n$, into which we substitute (84). This gives

$$H_{1,kn} = -i\sum_{\beta} \frac{1}{M_{\beta}} \mathbf{P}_{\beta} \cdot \langle x; k | R(\varepsilon_n, x) \nabla_{\beta} H_e(x) | x; n \rangle.$$
(94)

With this expression for $H_{1,kn}$ the first sum in (93) can be extended to all k since the term k = n vanishes anyway, after which we have a resolution of the identity. In this way we evaluate the first sum in (93). When we evaluate the second sum, including the minus sign, we find that it is equal to the first sum.

Overall, we obtain

$$K_{22} = \sum_{\alpha,\beta} \frac{1}{M_{\alpha}M_{\beta}} \mathbf{P}_{\alpha} \cdot \langle x; n | \nabla_{\alpha}H_{e}(x) R(\varepsilon_{n}, x)^{3} \nabla_{\beta}H_{e}(x) | x; n \rangle \cdot \mathbf{P}_{\beta}$$
$$= \sum_{\alpha,\beta} \frac{1}{M_{\alpha}M_{\beta}} \mathbf{P}_{\alpha} \cdot (\nabla_{\alpha} \langle x; n |) R(\varepsilon_{n}, x) (\nabla_{\beta} | x; n \rangle) \cdot \mathbf{P}_{\beta},$$
(95)

where in the second form we have used (78b) and (81). By inserting resolutions of the identity in the working basis this can be written

$$K_{22} = -\sum_{\alpha,\beta} \frac{1}{M_{\alpha}M_{\beta}} \sum_{kl} [\mathbf{P}_{\alpha} \cdot \mathbf{F}_{\alpha;nk}(x)] R_{kl}(\varepsilon_{n}, x) [\mathbf{F}_{\beta;ln}(x) \cdot \mathbf{P}_{\beta}], \qquad (96)$$

where the derivative couplings are evaluated in the working basis. Finally, transforming this to the adiabatic basis we obtain

$$K_{22} = -\sum_{\alpha,\beta} \frac{1}{M_{\alpha}M_{\beta}} \sum_{k \neq n} \frac{[\mathbf{P}_{\alpha} \cdot \mathbf{F}_{\alpha;nk}(x)][\mathbf{F}_{\beta;kn}(x) \cdot \mathbf{P}_{\beta}]}{\varepsilon_{n}(x) - \varepsilon_{k}(x)},$$
(97)

where now the derivative couplings are evaluated in the adiabatic basis. This is the result we would have obtained if we had worked strictly in the adiabatic basis, without worrying about its singularities.

J. Discussion of *K*₂₂

Equations (95)–(97) give the symbol of the operator \hat{K}_{22} . The operator itself can be obtained by replacing \mathbf{X}_{α} and \mathbf{P}_{α} by $\hat{\mathbf{X}}_{\alpha}$ and $\hat{\mathbf{P}}_{\alpha}$, in the order shown, plus commutator terms which are of higher order in \hbar and can be neglected. The resulting operator is Hermitian.

The term K_{22} should be included in Born-Oppenheimer descriptions of nuclear motion, but it has usually been neglected, in spite of the fact that it is of the same order of magnitude as the diagonal Born-Oppenheimer correction, which has received extensive attention. (At least, this is true in the large-amplitude ordering.) Admittedly both terms are small, and in many applications it may be reasonable to neglect one or both. Nevertheless, K_{22} grows with nuclear momentum while the diagonal Born-Oppenheimer correction does not, and at sufficiently large nuclear momenta it will become important.

This term has been discussed in depth by Mátyus (2018), who calls it the "nonadiabatic mass correction." The idea is that this term modifies the expression for the kinetic energy of the nuclei by adding a contribution due to the electrons that are carried along by them during nuclear motions. We refer to Mátyus (2018) for more details, mentioning also Kutzelnigg (2007). Here we add a few further comments about K_{22} .

One is that this term is more difficult to evaluate numerically than the other terms in the Born-Oppenheimer Hamiltonian. Mátyus (2018) has given ways of evaluating it, basically by inverting the operator $\varepsilon_n - H_e$ numerically. This requires careful attention to the choice of the subspace on which the inversion is carried out, and to numerical issues due to the fact that $\varepsilon_n - H_e$ has a nontrivial kernel.

Actually the resolvent operator is ubiquitous in higher order Born-Oppenheimer calculations, appearing not only in K_{22} but also in matrix elements of the electronic momentum and angular momentum (Nafie (1983); Freedman and Nafie (1986)). These are needed in studies of radiative processes such as vibrational circular dichroism (Stephens and Lowe (1985); Buckingham, Fowler, and Galwas (1987); Nafie (1992)), as well as in expressions for electronic currents (Barth *et al.* (2009); Okuyama and Takatsuka (2009)). A method for evaluating such matrix elements has been implemented by Stephens (1985) and similar ideas have been proposed in a more general context by Patchkovskii (2012). Basically, one calls on the fact that matrix elements involving the resolvent occur in perturbation theory. By concocting a perturbation of the right form, one can make the energy shifts in perturbation theory or their derivatives agree with the desired matrix elements.

Another remark is that including K_{22} , the entire $K = \bar{H}_{nn}$, that is, the Weyl symbol of the operator \hat{K} , is the eigenvalue of the symbol matrix H_{kl} , computed through second order. Since H_{kl} is already diagonal to lowest order in \hbar its eigenvalues to higher order can be computed by means of perturbation theory, taking into account the off-diagonal contributions from H_1 and H_2 . At second order this gives both K_{21} and K_{22} . This is interesting in terms of the work of Littlejohn and Flynn (1991b), who show that the Weyl symbol of the operator driving a given mode or polarization of a system of coupled wave equations is the eigenvalue of the symbol matrix of the wave operator, plus two corrections. The first is a correction due to Berry's phase, and the second, sometimes called the "no-name" term (Emmrich and Weinstein (1996)), involves the Poisson bracket of the polarization vectors. In the Born-Oppenheimer case, the Berry's phase term vanishes on a single surface, due to time-reversal invariance, while the Poisson bracket or no-name term is of third order in \hbar and so does not appear. This gives a complete accounting through second order of the Hamiltonian K relevant for a single surface in Born-Oppenheimer theory, including K_{22} .

K. Tensor Notation and the Generator G₂

We now introduce tensor notation for the kinetic energy metric, as explained in Appendix D, which is convenient especially for second-order calculations. Note that henceforth there will be an implied summation on repeated indices μ , ν , etc (one lower, one upper), where μ , $\nu = 1, ..., 3N$. Note also that P_{μ} (with a lower or covariant index) is the usual momentum, which is of order κ^{-2} , while P^{μ} (with an upper or contravariant index) is the same as the velocity, which is of order κ^2 . That is, $P^{\mu} = \dot{X}^{\mu}$. In the new notation Eqs. (59) become

$$H_{0,kl} = \left(\frac{1}{2}P^{\mu}P_{\mu}\right)\delta_{kl} + W_{kl}(x), \qquad (98a)$$

$$H_{1,kl} = -iP^{\mu}F_{\mu;kl}(x) = -iP^{\mu}\langle x;k|\partial_{\mu}|x;l\rangle,$$
(98b)

$$H_{2,kl} = -\frac{1}{2} \sum_{p} F^{\mu}{}_{;kp}(x) F_{\mu;pl}(x) = \frac{1}{2} (\partial^{\mu} \langle x;k|) (\partial_{\mu} |x;l\rangle),$$
(98c)

where we write the derivative couplings as

$$F_{\mu;kl}(x) = \langle x;k|\partial_{\mu}|x;l\rangle = -(\partial_{\mu}\langle x;k|)|x;l\rangle, \tag{99}$$

with

$$F^{\mu}{}_{;kl}(x) = G^{\mu\nu} F_{\nu;kl}(x). \tag{100}$$

Notice that $F_{\mu;kl} = O(1)$ while $F^{\mu}_{;kl} = O(\kappa^4)$. The metric $G^{\mu\nu}$ or $G_{\mu\nu}$ (see (D1 and (D2)) is not to be confused with the generators G_1 , G_2 , etc.

In the new notation the Feynman-Hellman formula (81) becomes

$$R(\varepsilon_n, x) \left[\partial_\mu H_e(x)\right] |x; n\rangle = Q(x) \partial_\mu |x; n\rangle, \tag{101}$$

and the generator G_1 becomes

$$G_{1,nk} = i P^{\mu}(\partial_{\mu}\langle x; n |) R(\varepsilon_n, x) | x; k\rangle, \qquad (k \neq n),$$
(102)

which is equivalent to (77), while the diagonal element $K = \bar{H}_{nn} = K_0 + \hbar K_1 + \hbar^2 K_2 + ...$ of the diagonalized Hamiltonian is given by

$$K_0 = \frac{1}{2} P^{\mu} P_{\mu} + \varepsilon_n(x), \qquad (103a)$$

$$K_1 = 0, \tag{103b}$$

$$K_2 = K_{21} + K_{22}. (103c)$$

Here $K_{21} = H_{2,nn}$ is the diagonal Born-Oppenheimer correction,

$$K_{21} = \frac{1}{2} (\partial^{\mu} \langle x; n |) (\partial_{\mu} | x; n \rangle) = -\frac{1}{2} \langle x; n | \partial^{\mu} \partial_{\mu} | x; n \rangle,$$
(104)

which is equivalent to the last term in (53), and K_{22} is the extra correction,

$$K_{22} = P^{\mu} P^{\nu}(\partial_{\mu} \langle x; n |) R(\varepsilon_n, x) (\partial_{\nu} | x; n \rangle), \qquad (105)$$

which is a version of (95).

The second generator G_2 is worked out in Appendix E. It is purely odd (off-block-diagonal) and anti-Hermitian. Its (nk)-block for $k \neq n$ is given by

$$G_{2,nk} = -P^{\mu}P^{\nu} \partial_{\nu} [(\partial_{\mu} \langle x; n |) R(\varepsilon_{n}, x)] R(\varepsilon_{n}, x) |x; k\rangle + [\partial^{\mu} \varepsilon_{n}(x)] (\partial_{\mu} \langle x; n |) R(\varepsilon_{n}, x)^{2} |x; k\rangle + \frac{1}{2} (\partial_{\mu} \langle x; n |) \partial^{\mu} [R(\varepsilon_{n}, x) |x; k\rangle], \quad (k \neq n).$$
(106)

Notice that the final term contains derivatives of the basis vectors $|x;k\rangle$ on \mathscr{S}^{\perp} . The other offdiagonal block is given by $G_{2,kn} = -G_{2,nk}^*$.

L. Changes of Diabatic Basis in \mathscr{S}^{\perp}

On \mathscr{S}^{\perp} , the working basis vectors are the same as the diabatic basis vectors, that is, $|x;k\rangle = |dx;k\rangle$ for $k \neq n$ (see (72)). But the diabatic basis on \mathscr{S}^{\perp} is highly arbitrary, that is, we can change it according to

$$|dx;k\rangle = \sum_{l\neq n} |d'x;l\rangle V_{lk}^{\perp}(x), \qquad (k\neq n),$$
(107)

where V_{lk}^{\perp} is a smooth function of *x*. Here $|dx;k\rangle$ is the old diabatic basis on \mathscr{S}^{\perp} and $|d'x;k\rangle$ is the new one. We restrict V_{lk}^{\perp} to be unitary so that the new basis is orthonormal, and restrict it further to be real and orthogonal so that the new basis is invariant under time reversal (we are assuming that the old basis is invariant under time reversal).

According to (10), under the change of basis (107) the wave function transforms according to

$$\psi'_k(x) = \sum_{l \neq n} V_{kl}^{\perp}(x) \, \psi_l(x), \qquad (k \neq n).$$
 (108)

The wave function ψ_k is not invariant under the change of diabatic basis on \mathscr{S}^{\perp} , but, as we shall say, it transforms as a vector.

Physical results cannot depend on the choice of diabatic basis vectors on \mathscr{S}^{\perp} . Note that the second order Hamiltonians K_{21} and K_{22} , given by (104) and (105), make no reference to the diabatic basis on \mathscr{S}^{\perp} and thus are invariant under a change of that basis. The wave function ψ_k is not invariant but transforms in a simple manner (as a vector) under a change of basis on \mathscr{S}^{\perp} . This makes it easy to construct invariants, for example, $\sum_{k\neq n} |\psi_k(x)|^2$. The components $G_{1,nk}$ of the generator G_1 are not invariant but do transform as a vector in the index k, as seen by (102). This means that physical results will never involve individual components $G_{1,nk}$, but rather sums over $k \neq n$ of $G_{1,nk}$ times something that transforms as a vector in k, in order to make an invariant.

On the other hand, it is clear from (107) that derivatives of the diabatic basis vectors in \mathscr{S}^{\perp} , things that look like $\partial_{\mu}|x;k\rangle$ for $k \neq n$, have a more complicated transformation law, one that involves derivatives of the transformation matrix $V_{kl}^{\perp}(x)$. Derivatives of the basis vectors on \mathscr{S}^{\perp} do not occur in the components $G_{1,nk}$ of the generator G_1 , but they do occur in G_2 , as shown in Appendix E. Such terms complicate the construction of quantities that are invariant under the change of basis (107).

This complication arises because we are using ordinary derivatives ∂_{μ} instead of a properly defined covariant derivative, which is the ordinary derivative plus a correction term. The correction term is, however, of higher order in κ^2 than the ordinary derivative, so the use of covariant

derivatives mixes up the expansion in powers of κ^2 . We will stick with ordinary derivatives, to keep the formalism as simple as possible.

M. Transformation of the Wave Function

Following the discussion of Sec. III C, we write $\hat{H}\bar{\Psi} = E\bar{\Psi}$ or $\hat{H}\bar{\Psi} = i\hbar\partial\bar{\Psi}/\partial t$ for the Schrödinger equation in the completely dressed representation (with hats on operators), where $\bar{\Psi}$ is an infinite-dimensional vector and \hat{H} an infinite-dimensional matrix. But \hat{H} is block-diagonal and the wave function $\bar{\Psi}$ is nonzero only in the slot k = n, so we write $\bar{\Psi}_k = \delta_{kn} f$ where f is the single wave function on surface n, satisfying $\hat{K}f = Ef$ or $\hat{K}f = i\hbar\partial f/\partial t$, where $\hat{K} = \hat{H}_{nn}$. The symbol K of operator \hat{K} is given by (103). If we solve this equation, we obtain f and hence $\bar{\Psi}$ in the fully dressed Born-Oppenheimer representation.

The wave function in the original Born-Oppenheimer representation is given by $\Psi = \hat{U}^{\dagger} \bar{\Psi}$, that is, $\Psi_k = \sum_l (\hat{U}^{\dagger})_{kl} \bar{\Psi}_l$. All the components Ψ_k are nonzero. The component k = n is the largest and approximately equal to f. Keeping this one and neglecting all the others is equivalent to the usual Born-Oppenheimer approximation. Thus, finding the other components, Ψ_k for $k \neq n$, as well as the corrections to Ψ_n , amounts to finding corrections to the Born-Oppenheimer approximation. It is feasible to calculate these corrections by hand to second order, as we shall now do.

Since $\bar{\psi}_k = \delta_{nk} f$, the transformation equations are

$$\Psi_n = (\hat{U}^{\dagger})_{nn} f = (\hat{U}_{nn})^{\dagger} f = (\hat{U}_{nn}^e)^{\dagger} f, \qquad (109a)$$

$$\Psi_k = (\hat{U}^{\dagger})_{kn} f = (\hat{U}_{nk})^{\dagger} f = (\hat{U}_{nk}^o)^{\dagger} f, \qquad (k \neq n),$$
 (109b)

where, as we see, only the even part of \hat{U} contributes to ψ_n while only the odd part contributes to ψ_k for $k \neq n$. On the other hand, we have

$$\hat{U} = \dots \hat{U}_2 \hat{U}_1 = \dots \exp(\hbar^2 \hat{G}_2) \exp(\hbar \hat{G}_1) = 1 + \hbar \hat{G}_1 + \hbar^2 \left(\hat{G}_2 + \frac{1}{2} \hat{G}_1^2 \right) + \dots,$$
(110)

so, taking into account the fact that \hat{G}_1 and \hat{G}_2 are purely odd, we can write (109) as

$$\Psi_n = f + \frac{\hbar^2}{2} [(\hat{G}_1^2)_{nn}]^{\dagger} f, \qquad (111a)$$

$$\Psi_k = \hbar (\hat{G}_{1,nk})^{\dagger} f + \hbar^2 (\hat{G}_{2,nk})^{\dagger} f, \qquad (k \neq n),$$
 (111b)

which are valid through second order. Note that the first correction to $\psi_n = f$ occurs at second order.

We work first on the term involving \hat{G}_1^2 . Since \hat{G}_1^2 is Hermitian (as a matrix of operators) its diagonal elements are Hermitian (as scalar operators), and the \dagger is unnecessary in (111a). Converting operators into symbols, we have

$$(\hat{G}_{1}^{2})_{nn} = \sum_{k} \hat{G}_{1,nk} \hat{G}_{1,kn} \longleftrightarrow \sum_{k} G_{1,nk} * G_{1,kn} = -\sum_{k} |G_{1,nk}|^{2} + \dots$$
$$= -P^{\mu} P^{\nu} \left(\partial_{\mu} \langle x; n | \right) R(\varepsilon_{n}, x)^{2} (\partial_{\nu} | x; n \rangle) + \dots,$$
(112)

where we use \leftrightarrow as in Appendix A, where we have used $G_{1,kn} = -G_{1,nk}^*$ and where the ellipsis indicates higher order terms in the expansion of the *-product, which we neglect since we are working at second order. In the final step we have used (102) and evaluated the sum. When we convert the final expression, a Weyl symbol, back into an operator, we are free to put the factors \hat{P}^{μ} , \hat{P}^{ν} in any order, since the differences among the choices are commutators that are of higher order in \hbar . With this understanding, we can write (111a) as

$$\psi_n = f - \frac{\hbar^2}{2} (\partial_\mu \langle x; n |) R(\varepsilon_n, x)^2 (\partial_\nu | x; n \rangle) \hat{P}^\mu \hat{P}^\nu f.$$
(113)

As for (111b), we will need

$$(\hat{G}_{1,nk})^{\dagger} \longleftrightarrow G_{1,nk}^{*} = -iP^{\mu}\langle x;k|R(\varepsilon_{n},x)(\partial_{\mu}|x;n\rangle) \longleftrightarrow -i\langle x;k|R(\varepsilon_{n},x)(\partial_{\mu}|x;n\rangle)\hat{P}^{\mu} - \frac{\hbar}{2}\partial^{\mu}[\langle x;k|R(\varepsilon_{n},x)(\partial_{\mu}|x;n\rangle)], \qquad (114)$$

where we use (102) and a version of (A5b). Since G_1 occurs at first order, we must keep the commutator term (the final term) in (114) to obtain results valid to second order.

A similar treatment of G_2 (see (106)) gives rise to three terms, one of which is

$$\frac{1}{2} \{ \partial^{\mu} [\langle x; k | R(\varepsilon_n, x)] \} (\partial_{\mu} | x; n \rangle),$$
(115)

the term that depends on the derivatives of $\langle x; k |$. But so does the commutator term in (114), and when the two are combined the derivatives of $\langle x; k |$ drop out. Thus (111b) becomes

$$\Psi_{k} = -i\hbar \langle x; k | R(\varepsilon_{n}, x) (\partial_{\mu} | x; n \rangle) \hat{P}^{\mu} f + \hbar^{2} \{ -\langle x; k | R(\varepsilon_{n}, x) \partial_{\nu} [R(\varepsilon_{n}, x) (\partial_{\mu} | x; n \rangle)] \hat{P}^{\mu} \hat{P}^{\nu} + [\partial^{\mu} \varepsilon_{n}(x)] \langle x; k | R(\varepsilon_{n}, x)^{2} (\partial_{\mu} | x; n \rangle) - \frac{1}{2} \langle x; k | R(\varepsilon_{n}, x) (\partial^{\mu} \partial_{\mu} | x; n \rangle) \} f, \quad (k \neq n).$$
(116)

We see that through second order, ψ_k transforms as a vector, as it must.

We can also write out the expansion of the wave function $\Psi(\mathbf{X}, \mathbf{r})$, the solution of the original molecular Schrödinger equation, (22) or (23). We write

$$\Psi(\mathbf{X},\mathbf{r}) = \Psi_0(\mathbf{X},\mathbf{r}) + \Psi_1(\mathbf{X},\mathbf{r}) + \dots, \qquad (117)$$

where the subscripts 0, 1, etc., indicate the power of κ^2 . Then $\Psi_0(\mathbf{X}, \mathbf{r}) = \phi_n(\mathbf{X}; \mathbf{r}) f(\mathbf{X}) = \langle \mathbf{r} | x; n \rangle f(\mathbf{X})$, which is the usual Born-Oppenheimer approximation, and

$$\Psi_{1}(\mathbf{X}, \mathbf{r}) = -i\hbar \sum_{k \neq n} \phi_{k}(\mathbf{X}; \mathbf{r}) \langle x; k | R(\varepsilon_{n}, x)(\partial_{\mu} | x; n \rangle) \hat{P}^{\mu} f$$

$$= -i\hbar \langle \mathbf{r} | R(\varepsilon_{n}, x)(\partial_{\mu} | x; n \rangle) \hat{P}^{\mu} f$$

$$= -i\hbar \sum_{\alpha=1}^{N} \frac{1}{M_{\alpha}} \langle \mathbf{r} | R(\varepsilon_{n}, x)(\nabla_{\alpha} | x; n \rangle) \cdot \hat{\mathbf{P}}_{\alpha} f \qquad (118)$$

is the first order correction, written in several different ways. For brevity we omit the second order correction, but it is easily written out with the help of (113) and (116).

The first order correction to the Born-Oppenheimer wave function was apparently first derived by Nafie (1983), in the context of small vibrations, and has been rederived many times since. Our results apply in the more general context of large-amplitude motions, and our second order results (113) and (116) seem to be new.

IV. GENERALIZATIONS AND APPLICATIONS

In this section we outline the generalization of Moyal perturbation theory to multisurface problems and we discuss some applications including the calculation of electronic currents.

A. Multiple Surfaces

For multi-surface problems we define an index set I of N_l adjacent electronic energy levels,

$$I = \{k_0, k_0 + 1, \dots, k_0 + N_l - 1\},$$
(119)

which are regarded as "strongly coupled," at least somewhere in the region of nuclear configuration space under consideration. These levels define a privileged subspace $\mathscr{S}(x)$,

$$\mathscr{S}(x) = \operatorname{span}\{|ax;k\rangle, k \in I\},\tag{120}$$

which replaces (43). As before, the complementary, orthogonal subspace is $\mathscr{S}^{\perp}(x)$. Most of the formulas of Sec. III, which was devoted to single surface problems in which the privileged subspace was the single level k = n, are generalized to the multi-surface case by replacing k = n and $k \neq n$ by $k \in I$ and $k \notin I$, respectively. For example, the resolvent is still given by (76), except

now summed over $k \notin I$, and the projector onto $\mathscr{S}(x)$ becomes

$$P(x) = \sum_{k \in I} |ax;k\rangle \langle ax;k|.$$
(121)

Now we can distinguish between "internal degeneracies," those that occur among levels $k \in I$, and "external degeneracies," those that occur between a level $k \in I$ and a level $k \notin I$. Now the region of nuclear configuration space must be restricted to exclude external degeneracies and a margin around them, while internal degeneracies are allowed.

Ever since (72) we have been using a working basis which is the diabatic basis on \mathscr{S}^{\perp} . Now we must introduce a diabatic basis also on \mathscr{S} , that is, a basis $|dx;k\rangle$ for $k \in I$, so that derivatives of the basis states are defined. Now we define the working basis by

$$|x;k\rangle = \begin{cases} |dx;k\rangle & k \in I, \\ |dx;k\rangle & k \notin I. \end{cases}$$
(122)

To be clear about the notation, there are two diabatic bases, one on \mathscr{S} and one on \mathscr{S}^{\perp} . The first is a linear combination of the adiabatic basis vectors $|ax;k\rangle$ for $k \in I$, and the second is a linear combination of those for $k \notin I$. For the transformation between the diabatic and adiabatic bases on \mathscr{S} we write

$$|dx;k\rangle = \sum_{l\in I} |ax;l\rangle V_{lk}, \qquad k\in I,$$
(123)

where V_{lk} is an $N_l \times N_l$ orthogonal matrix,

$$V_{lk} = \langle ax; l | dx; k \rangle. \tag{124}$$

Now "even" and "odd" are defined relative to the sets of indices, $k \in I$ and $k \notin I$.

The derivation proceeds much as in the case of single surface. The generator G_1 (that is, its Weyl symbol) is purely odd and satisfies $[G_1, H_0] = -H_1^o$. This can be solved for one off-diagonal block,

$$G_{1,kp} = \sum_{l,m\in I} \sum_{q\notin I} (V^{\dagger})_{kl} V_{lm} H_{1,mq} R_{qp}(\varepsilon_l, x), \qquad k \in I, \ p \notin I,$$
(125)

while the other is determined by the anti-Hermiticity of G_1 . The presence of the matrix V in this formula means that effectively we have carried out the linear algebra of the solution for G_1 in the adiabatic basis, then transformed the result to the working (that is, diabatic) basis. The calculation would be formally simpler if we had defined the working basis to be the adiabatic basis on S, but we wish to avoid any derivatives of these adiabatic basis vectors since they are not defined at internal degeneracies (that is, on the seams of the single surface problem). Our presentation of the multisurface problem is designed so that such derivatives never occur (but derivatives of the diabatic basis states on \mathscr{S} are allowed).

Equation (125) raises the question of whether the solution $G_{1,kp}$ is a smooth function of x, since the matrix V is not smooth. (Matrix V connects the diabatic basis, which is smooth, with the adiabatic basis, which is not.) To answer this question we use the logic explained more fully by Littlejohn, Rawlinson, and Subotnik (2022), that the singularities of the adiabatic basis are caused by the topological impossibility of making smooth phase conventions for energy eigenstates in the neighborhood of a degeneracy, and that correspondingly expressions that are independent of such phase conventions are smooth. It is easily seen that $G_{1,kp}$ given by (125) is independent of the phase conventions for the adiabatic basis, and so is a smooth function of x. A more sophisticated argument can be based on a contour integral (see Appendix A of Littlejohn, Rawlinson, and Subotnik (2022)). Now substituting (98b) into (125) and carrying out some manipulations we find

$$G_{1,kp} = -iP^{\mu} \sum_{l \in I} (V^{\dagger})_{kl} \langle ax; l | [\partial_{\mu} R(\varepsilon_l, x)] | x; p \rangle, \qquad k \in I, \, p \notin I.$$
(126)

Results equivalent to (125) have previously been reported by Mátyus and Teufel (2019), who work with an electronic operator that in our notation could be written as

$$\sum_{k \in I} \sum_{p \notin I} |x;k\rangle G_{1,kp} \langle x;p| + \text{h.c.}$$
(127)

This is part of their formalism of operator-valued Weyl symbols. The use of this formalism does not obviate questions of smoothness, for example, the projector onto individual energy eigenstates, which appears in their formulas, is not smooth at internal degeneracies.

The Hamiltonian \hat{K} on the privileged subspace is an $N_l \times N_l$ matrix, that is, with indices $k, l \in I$, whose symbol matrix can be written $K = K_0 + \hbar K_1 + \hbar^2 K_2$, to second order. As for K_0 , it is a version of (98a),

$$K_{0,kl} = \left(\frac{1}{2}P^{\mu}P_{\mu}\right)\delta_{kl} + W_{kl}, \qquad k,l \in I,$$
(128)

where W_{kl} is a full matrix because we are using the diabatic basis for $k \in I$. As for K_1 , it is the same as H_1 on the diagonal block corresponding to \mathscr{S} ,

$$K_{1,kl} = -iP^{\mu} \langle x; k | \partial_{\mu} | x; l \rangle, \qquad k, l \in I.$$
(129)

Note that in multi-surface problems, K_1 has non-vanishing, off-diagonal components with derivative couplings. Finally, $K_2 = K_{21} + K_{22}$ consists of two terms, of which K_{21} is a version of (98c),

$$K_{21,kl} = \frac{1}{2} (\partial^{\mu} \langle x; k |) (\partial_{\mu} | x; l \rangle), \qquad k, l \in I,$$
(130)

which is a multi-surface version of the diagonal Born-Oppenheimer correction. The second correction K_{22} can be written

$$K_{22,kl} = \frac{1}{2} P^{\mu} P^{\nu} \sum_{m,n \in I} (V^{\dagger})_{km} \langle ax; m | [\partial_{\mu} P(x)] [R(\varepsilon_m, x) + R(\varepsilon_n, x)] [\partial_{\nu} P(x)] | ax; n \rangle V_{nl}, \qquad k, l \in I.$$
(131)

In this expression we have taken derivatives of the projection operator P(x), which is smooth, but not of the adiabatic basis states, which are not. (The projection operator is not to be confused with the momenta P^{μ} , P^{ν} .) A result equivalent to (131) has been previously reported by Mátyus and Teufel (2019).

The second generator G_2 and the corrections to the wave function may also be computed.

B. Electronic Momentum

In single-surface problems the usual (lowest order) Born-Oppenheimer approximation gives a vanishing result for the matrix elements of electron momenta (linear and angular) and in the calculation of electronic currents. For such problems it is necessary to go to higher order. The same applies to the single-surface matrix elements of any Hermitian electronic operator that is odd under time-reversal. In this subsection we explain how the issue may be treated within Moyal perturbation theory.

We begin with the momentum \mathbf{p}_i of one of the electrons, which is odd under time reversal, $T\mathbf{p}_i T^{\dagger} = -\mathbf{p}_i$. This is a useful practice case, although it has a special feature not shared with other such operators, namely, the operator \mathbf{p}_i can be represented as an exact time derivative. That is, let $\Psi(\mathbf{X}, \mathbf{r}, t)$ be the time-dependent wave function of the molecule. Then

$$\langle \Psi | \mathbf{p}_i | \Psi \rangle_{Xr} = \frac{m_e}{i\hbar} \langle \Psi | [\mathbf{r}_i, H] | \Psi \rangle_{Xr} = m_e \frac{d}{dt} \langle \Psi | \mathbf{r}_i | \Psi \rangle_{Xr}.$$
(132)

This is exact in the electrostatic model, but if the Born-Oppenheimer approximation (48) is used in it (where Ψ and ψ are allowed to depend on *t*) then the left-hand side vanishes while the right-hand side does not. To reconcile the two it is necessary to go to higher order in the Born-Oppenheimer

expansion in the evaluation of the left-hand side. Note that \mathbf{r}_i is even under time reversal, so its matrix element does not vanish at lowest order.

To evaluate $\langle \Psi | \mathbf{p}_i | \Psi \rangle$ we use a version of (18),

$$\langle \Psi | A | \Psi \rangle_{Xr} = \sum_{kl} \langle \psi_k | \hat{A}_{kl} | \psi_l \rangle_X = \sum_{kl} \langle \bar{\psi}_k | \hat{\bar{A}}_{kl} | \bar{\psi}_l \rangle_X,$$
(133)

showing how the expectation value of an operator A changes as the representation moves from the molecular to the original Born-Oppenheimer to the dressed Born-Oppenheimer (that is, across the first row of Table I). Here we define $\hat{A} = \hat{U}\hat{A}\hat{U}^{\dagger}$, as in (58), using hats to indicate nuclear operators or matrices thereof. If the original A is a purely electronic operator then $\hat{A}_{kl} = \langle x; k | A | x; l \rangle$ (see subsection II C), which is a purely multiplicative nuclear operator (that is, a function of **X** only). In this case there is hardly a difference between the operator \hat{A}_{kl} and its Weyl symbol A_{kl} .

We identify the original operator A with \mathbf{p}_i , and write (\mathbf{p}_i) for the matrix representing it in the original Born-Oppenheimer representation, so that

$$(\mathbf{p}_i)_{kl} = \langle x; k | \mathbf{p}_i | x; l \rangle.$$
(134)

If we wish to emphasize that this is a nuclear operator, we will write this matrix as $(\widehat{\mathbf{p}_i})$ or its components as $(\widehat{\mathbf{p}_i})_{kl}$, but these are purely multiplicative operators (functions of **X** only). We write

$$\widehat{(\mathbf{\bar{p}}_i)} = \hat{U}(\mathbf{\bar{p}}_i)\hat{U}^{\dagger}$$
(135)

(with an overbar) for the matrix of nuclear operators representing \mathbf{p}_i in the dressed Born-Oppenheimer representation, as in (58). Now it is necessary to distinguish the nuclear operators $(\mathbf{\bar{p}}_i)_{kl}$ from their Weyl symbols, which are $(\mathbf{\bar{p}}_i)_{kl}$ (without the hat), because the conjugation by \hat{U} introduces a momentum dependence.

We can write the expectation value of the momentum in three different representations, moving across the first row of Table I,

$$\langle \Psi | \mathbf{p}_i | \Psi \rangle_{Xr} = \sum_{kl} \langle \psi_k | (\mathbf{p}_i)_{kl} | \psi_l \rangle_X = \langle f | \widehat{(\mathbf{\bar{p}}_i)}_{nn} | f \rangle_X,$$
(136)

where we have used $\bar{\psi}_k = \delta_{kn} f$ in the dressed Born-Oppenheimer representation, since we are working on a single surface. In this representation the double sum has reduced to a single term.

Now there is the question of which representation is best for the calculation of $\langle \mathbf{p}_i \rangle$. For the molecular representation we have the wave function (117) and (118) and for the original Born-Oppenheimer representation we have the wave function (113) and (116), where *f* is a solution

of $\hat{K}f = Ef$ or $\hat{K}f = i\hbar\partial f/\partial t$. But we suggest that the fully dressed Born-Oppenheimer representation is optimal for the calculation, since it does not require us to transform wave functions. Instead, it requires us to transform operators, to obtain the one (scalar) nuclear operator

$$\widehat{(\mathbf{\bar{p}}_i)}_{nn} = [\hat{U}(\widehat{\mathbf{p}_i})\hat{U}^{\dagger}]_{nn} = \widehat{(\mathbf{p}_i)}_{nn} + \hbar[\hat{G}_1, \widehat{(\mathbf{p}_i)}]_{nn} + \dots,$$
(137)

or, in terms of Weyl symbols,

$$(\mathbf{\bar{p}}_i)_{nn} = (\mathbf{p}_i)_{nn} + \hbar[G_1, (\mathbf{p}_i)]_{nn} + \dots$$
(138)

The first term of this expansion vanishes,

$$(\mathbf{p}_i)_{nn} = \langle x; n | \mathbf{p}_i | x; n \rangle = 0, \tag{139}$$

because of time-reversal (this is the vanishing of the expectation value to lowest order). With the help of (102) the commutator in the correction term is easily worked out, giving altogether

$$(\bar{\mathbf{p}}_i)_{nn} = i\hbar P^{\mu} [(\partial_{\mu} \langle x; n |) R(\boldsymbol{\varepsilon}_n, x) \mathbf{p}_i | x; n \rangle - \langle x; n | \mathbf{p}_i R(\boldsymbol{\varepsilon}_n, x) (\partial_{\mu} | x; n \rangle)],$$
(140)

valid to first order. Actually, due to time reversal, the two terms shown (including the minus sign in the second term) are equal to one another, but it is convenient to leave the answer as shown.

Now we may use the fact that $\mathbf{p}_i = m_e(d/dt)\mathbf{r}_i = (m_e/i\hbar)[\mathbf{r}_i, H_e]$ is an exact time derivative (this is like (132) but using H_e instead of the full molecular Hamiltonian H). With this substitution the first matrix element in (140) becomes

$$\frac{m_e}{i\hbar}(\partial_{\mu}\langle x;n|)R(\varepsilon_n,x)(\varepsilon_n-H_e)\mathbf{r}_i|x;n\rangle.$$
(141)

Treating the second matrix element similarly and using (79) and $(\partial_{\mu}\langle x; n|)Q(x) = \partial_{\mu}\langle x; n|$ we can rewrite (140) as

$$(\bar{\mathbf{p}}_i)_{nn} = m_e P^{\mu} \partial_{\mu} (\langle x; n | \mathbf{r}_i | x; n \rangle).$$
(142)

Now using (A7) to convert the Weyl symbol into an operator, we can use (136) to write

$$\langle \mathbf{p}_i \rangle = \frac{m_e}{2} \int dX f(\mathbf{X}, t)^* [\hat{P}^{\mu}(\partial_{\mu} \langle x; n | \mathbf{r}_i | x; n \rangle) + (\partial_{\mu} \langle x; n | \mathbf{r}_i | x; n \rangle) \hat{P}^{\mu}] f(\mathbf{X}, t)$$

$$= m_e \int dX J_n^{\mu}(\mathbf{X}, t) \partial_{\mu} \langle x; n | \mathbf{r}_i | x; n \rangle,$$
(143)

where $J_n^{\mu}(\mathbf{X}, t)$ is the nuclear current,

$$J_n^{\mu}(\mathbf{X},t) = f(\mathbf{X},t)^* \left(\frac{-i\hbar\partial^{\mu}}{2}\right) f(\mathbf{X},t) + \text{c.c.}$$
(144)

The current satisfies the continuity equation,

$$\partial_{\mu}J_{n}^{\mu}(\mathbf{X},t) + \frac{\partial\rho_{n}(\mathbf{X},t)}{\partial t} = 0, \qquad (145)$$

where the nuclear probability density is

$$\boldsymbol{\rho}_n(\mathbf{X},t) = |f(\mathbf{X},t)|^2. \tag{146}$$

Thus we can integrate (143) by parts and discard boundary terms (assuming $f(\mathbf{X},t)$ is normalizable) to obtain

$$\langle \mathbf{p}_i \rangle = -m_e \int dX \left[\partial_\mu J_n^\mu(\mathbf{X}, t) \right] \langle x; n | \mathbf{r}_i | x; n \rangle = m_e \frac{d}{dt} \int dX \, \rho_n(\mathbf{X}, t) \, \langle x; n | \mathbf{r}_i | x; n \rangle = m_e \frac{d}{dt} \langle \mathbf{r}_i \rangle, \quad (147)$$

thereby recovering (132).

We have introduced the usual definitions of the nuclear probability density and current and invoked the continuity equation that they satisfy, but we are working in the dressed Born-Oppenheimer representation and these operations should be justified. That is, one should take the usual definitions and properties in the molecular representation and track them through the changes of representation across the first row of Table I. It turns out that what we have done is correct through the order to which we are working but for brevity we omit the details.

The integral over X on the right-hand side of (136) is nominally taken over all of configuration space, but the transformation to the dressed representation on a single surface is only valid outside the excluded region, that is, away from degeneracies. Thus, the manipulations presented are valid if f effectively has support in the allowed region, but if the wave function extends to the neighborhood of a degeneracy then we must use a multi-surface approach. In multisurface problems the expectation value of the electronic momentum does not vanish at lowest order and the answers cannot be expressed so simply in terms of nuclear currents. Time reversal forces the diagonal matrix elements, $\langle x; k | \mathbf{p}_i | x; l \rangle$ for k = l, to vanish, but not the off-diagonal elements ($k \neq l$). If both f_k and f_l are nearly zero in the excluded region then the X integral in $\langle f_k | (\mathbf{p}_i)_{kl} | f_l \rangle_X$ for $k \neq l$ contains an integrand that is a rapidly oscillating function times a slowly varying one. In this case the integral is nearly zero, and the off-diagonal elements do not contribute. But if both f_k and f_l extend to the neighborhood of the degeneracy, then the off-diagonal elements will make a contribution, nominally at lowest order. On the other hand, speaking of a two-surface problem, the off-diagonal matrix elements of the momentum vanish at internal degeneracies and are proportional to the energy difference as we move away from those degeneracies. This should be true over the extent of the excluded region, which, as explained, is of order κ . Overall, the κ -ordering of the expectation value of the electronic momentum displays a somewhat complicated picture in multi-surface problems. Such problems are currently of considerable interest, and this is a topic that deserves further development.

C. Electronic Current

We turn now to the electronic probability current, which also vanishes at lowest order in the Born-Oppenheimer expansion. We show this by expressing it as the expectation value of an Hermitian operator that is odd under time reversal. We continue with the convention of using hats on nuclear operators, and omitting them on the corresponding Weyl symbols. We do not use hats on electronic operators, but in this subsection it is important to distinguish between electronic operators and corresponding quantities that are *c*-numbers.

In this subsection we use index $\sigma = 1, ..., 3N_e$ to label the coordinates in the $3N_e$ -dimensional electronic configuration space, writing r^{σ} for the components of the vectors \mathbf{r}_i , $i = 1, ..., N_e$ and $p_{\sigma} = -i\hbar\partial/\partial r^{\sigma}$ for the corresponding momenta. We employ an implied summation over repeated indices σ .

We define a purely electronic operator ρ_{r_0} , parameterized by a *c*-number r_0 representing a point of the electronic configuration space,

$$\rho_{r_0} = \delta(r - r_0). \tag{148}$$

Really r_0 is a $3N_e$ -vector of *c*-numbers, the components r_0^{σ} of r_0 , while *r* represents the $3N_e$ dimensional vector of electronic operators whose components are multiplication by r^{σ} . Then we define the electronic probability density $\rho(r_0, t)$ by

$$\boldsymbol{\rho}(r_0,t) = \langle \Psi | \boldsymbol{\rho}_{r_0} | \Psi \rangle_{Xr} = \int dX \, |\Psi(\mathbf{X},\mathbf{r}_0,t)|^2, \tag{149}$$

which is the usual definition. Note that ρ_{r_0} is an operator while $\rho(r_0, t)$ is a *c*-number.

Now we compute the time derivative,

$$\frac{\partial \rho(r_0, t)}{\partial t} = \frac{d}{dt} \langle \Psi | \rho_{r_0} | \Psi \rangle_{Xr} = \frac{1}{i\hbar} \langle \Psi | [\rho_{r_0}, H] | \Psi \rangle_{Xr}$$
$$= \frac{1}{2m_e} \langle \Psi | [p_\sigma \delta_\sigma (r - r_0) + \delta_\sigma (r - r_0) p_\sigma] | \Psi \rangle_{Xr}, \tag{150}$$

where *H* is the molecular Hamiltonian (1), where we have worked out the commutator, and where $\delta_{\sigma}(r-r_0)$ means $(\partial/\partial r^{\sigma})\delta(r-r_0)$. The commutator comes entirely from the electronic kinetic energy in the Hamiltonian, which can be written as $p_{\sigma}p_{\sigma}/2m_e$. But since

$$\frac{\partial}{\partial r^{\sigma}}\delta(r-r_0) = -\frac{\partial}{\partial r_0^{\sigma}}\delta(r-r_0), \qquad (151)$$

we can write

$$\frac{\partial \rho(r_0,t)}{\partial t} = -\frac{\partial J^{\sigma}(r_0,t)}{\partial r_0^{\sigma}}$$
(152)

where

$$J^{\sigma}(r_0,t) = \langle \Psi | J^{\sigma}_{r_0} | \Psi \rangle_{Xr}$$
(153)

and

$$J_{r_0}^{\sigma} = \frac{1}{2m_e} [p_{\sigma} \delta(r - r_0) + \delta(r - r_0) p_{\sigma}].$$
(154)

Note that $J_{r_0}^{\sigma}$ is an operator parameterized by r_0 while $J^{\sigma}(r_0,t)$ is the usual electronic current, which is a *c*-number. Diestler (2012a) has obtained results equivalent to (153) and (154) by performing an *ad hoc* quantization of the classical expression for the current.

Equation (153) expresses the electronic current as the expectation value of a purely electronic operator (154) that is odd under time reversal, so the result vanishes to lowest order of the Born-Oppenheimer expansion. We can treat this operator exactly as we did \mathbf{p}_i in subsection IV B, except that $J_{r_0}^{\sigma}$, unlike \mathbf{p}_i , cannot be expressed as an exact time derivative. That is, we write $(J_{r_0}^{\sigma})$ for the matrix of operators representing the electronic current in the Born-Oppenheimer representation, with components $(J_{r_0}^{\sigma})_{kl}$. These are functions of \mathbf{X} and if we wish to emphasize that they are nuclear operators we can place a hat over them, writing $(\widehat{J_{r_0}^{\sigma}})$ for the matrix or $(\widehat{J_{r_0}^{\sigma}})_{kl}$ for its components. These are then transformed to the dressed Born-Oppenheimer representation according to $(\widehat{J_{r_0}^{\sigma}}) = \widehat{U}(\widehat{J_{r_0}^{\sigma}})\widehat{U}^{\dagger}$, and we write $(\widehat{J_{r_0}^{\sigma}})$ for the corresponding matrix of Weyl symbols (without the hat). Finally, the *nn*-component of this is the effective (nuclear) operator representing the electronic current on the given surface *n*; we give this a special name, writing

$$\hat{I}_{r_0}^{\sigma} = \left(\overline{\bar{J}_{r_0}^{\sigma}}\right)_{nn},\tag{155}$$

or $I_{r_0}^{\sigma}$ for the corresponding Weyl symbol. Then we have

$$I_{r_0}^{\sigma} = -2i\hbar P^{\mu} \langle x; n | J_{r_0}^{\sigma} R(\varepsilon_n, x) (\partial_{\mu} | x; n \rangle), \qquad (156)$$

which is derived just like (140) except that the two equal terms in the result have been combined. Then we have

$$J^{\sigma}(r_{0},t) = \langle f | \hat{I}_{r_{0}}^{\sigma} | f \rangle_{X} = -2i\hbar \int dX J_{n}^{\mu}(\mathbf{X},t) \, \langle x; n | J_{r_{0}}^{\sigma} R(\varepsilon_{n},x)(\partial_{\mu} | x; n \rangle),$$
(157)

which is derived just like (143).

Results equivalent to (157) have been derived previously by Nafie (1983) and Patchkovskii (2012), although it takes some extrapolation to compare our results with Nafie's because he worked in the small-amplitude regime and expanded his results about an equilibrium. What Nafie calls "complete adiabatic" versions of operators such as the dipole and current are the same as our dressed operators. He emphasizes that these have a dependence on nuclear momentum. Nafie employs an *ad hoc* quantization and dequantization of nuclear operators, roughly equivalent to our use of Weyl symbols, and he has presented some differential equations satisfied by the current operators in their dequantized versions (Nafie (2020)). Patchkovskii (2012) allows large-amplitude motions, as we do. He evaluates the matrix element in (157) by inserting a resolution of the identity in the adiabatic basis just before the resolvent operator. This gives

$$J^{\sigma}(r_{0},t) = \frac{\hbar^{2}}{m_{e}} \int dX J^{\mu}_{n}(\mathbf{X},t) \sum_{k \neq n} \frac{\langle ax;k|\partial_{\mu}|ax;n\rangle}{\varepsilon_{n}(x) - \varepsilon_{k}(x)} \times \left[\frac{\partial \phi^{*}_{an}(x;r_{0})}{\partial r_{0\sigma}} \phi_{ak}(x;r_{0}) - \phi^{*}_{an}(x;r_{0})\frac{\partial \phi_{ak}(x;r_{0})}{\partial r_{0\sigma}}\right],$$
(158)

which is essentially the expression given by him.

The sum over all adiabatic states engages the continuum and is not useful for direct evaluation, but in any case it is unlikely that one will want to compute the $3N_e$ -dimensional electronic current. Instead, reduced versions obtained by integrating over all electron coordinates but one are more useful. See the remarks in subsection III J on the evaluation of matrix elements involving the resolvent. Diestler (2013) evaluates the sum in (158) by replacing the energy denominator $\varepsilon_n - \varepsilon_k$ by an appropriately chosen average energy ΔE .

If the current operator $J_{r_0}^{\sigma}$ were an exact time derivative we could use the trick employed with the electronic momentum \mathbf{p}_i to get rid of the resolvent; but it is not. On the other hand, the divergence $\partial_{\mu}J_{r_0}^{\mu}$ of the current operator is an exact time derivative. If we use this fact to eliminate the resolvent operator, we just recover the continuity equation. This is a trivial observation but it emphasizes a point made by Barth *et al.* (2009), that normally one is interested in the integral of the current over surfaces that extend to infinity. The evaluation of these does not require the use of the resolvent operator. In the special case that all the nuclei are displaced by the same displacement $\boldsymbol{\xi}$ (a 3-vector), it is clear that the electrons are just carried along by the nuclei and that the expression for the electronic current should simplify. This is another case in which we can get rid of the resolvent operator. To treat this problem we must make a digression into the behavior of the basis states $|x;k\rangle$ under translations, including their phase conventions.

V. TRANSLATIONAL DEGREES OF FREEDOM, IN GREATER DEPTH

We return to the subject of the translational degrees of freedom, showing that their elimination can be achieved by means of a unitary transformation and showing how the rectilinear motion of the center of mass is manifested in the Born-Oppenheimer and dressed Born-Oppenheimer representations. See also our treatment of rotations in Littlejohn, Rawlinson, and Subotnik (2023).

A. Transformation of $H_e(x)$ Under Translations

Let $\boldsymbol{\xi}$ be a displacement vector in 3-dimensional space, and let $\boldsymbol{\xi} = (\boldsymbol{\xi}, \boldsymbol{\xi}, \dots, \boldsymbol{\xi})$ be the corresponding displacement vector in the 3*N*-dimensional nuclear configuration space. Also, let $\mathbf{r} + \boldsymbol{\xi}$ stand for the collection $(\mathbf{r}_1 + \boldsymbol{\xi}, \dots, \mathbf{r}_{N_e} + \boldsymbol{\xi})$. Then the translational invariance of the electronic Hamiltonian is expressed by

$$H_e(x+\boldsymbol{\xi};\mathbf{r}+\boldsymbol{\xi},\mathbf{p}) = H_e(x;\mathbf{r},\mathbf{p}).$$
(159)

This is a statement about the functional form of $H_e(x; \mathbf{r}, \mathbf{p})$, which depends only on vector differences among \mathbf{X}_{α} and \mathbf{r}_i .

We define electronic, nuclear, and total translation operators by

$$T_e(\boldsymbol{\xi}) = e^{-i\boldsymbol{\xi}\cdot\mathbf{p}_t/\hbar}, \qquad T_n(\boldsymbol{\xi}) = e^{-i\boldsymbol{\xi}\cdot\mathbf{P}_t/\hbar}, \qquad T(\boldsymbol{\xi}) = T_e(\boldsymbol{\xi})T_n(\boldsymbol{\xi}) = e^{-i\boldsymbol{\xi}\cdot(\mathbf{P}_t+\mathbf{p}_t)/\hbar}.$$
 (160)

See (35) and (28) for definitions of \mathbf{p}_t and \mathbf{P}_t . These are in the molecular representation. The electronic translation operators satisfy

$$T_e(\boldsymbol{\xi}) \mathbf{r}_i T_e(\boldsymbol{\xi})^{\dagger} = \mathbf{r}_i - \boldsymbol{\xi}, \qquad T_e(\boldsymbol{\xi}) \mathbf{p}_i T_e(\boldsymbol{\xi})^{\dagger} = \mathbf{p}_i.$$
(161)

Thus we have

$$T_e(\boldsymbol{\xi}) H_e(x; \mathbf{r}, \mathbf{p}) T_e(\boldsymbol{\xi})^{\dagger} = H_e(x; \mathbf{r} - \boldsymbol{\xi}, \mathbf{p}) = H_e(x + \boldsymbol{\xi}; \mathbf{r}, \mathbf{p}),$$
(162)



FIG. 1. The translational fiber bundle.

where in the first equality the conjugation by $T_e(\boldsymbol{\xi})$ does nothing to *x*, which is just the set of parameters of the electronic Hamiltonian, and where the second equality follows from (159). We abbreviate this by writing

$$T_e(\boldsymbol{\xi})H_e(x)T_e(\boldsymbol{\xi})^{\dagger} = H_e(x+\boldsymbol{\xi}), \qquad (163)$$

which is the transformation law of the electronic Hamiltonian under electronic translations.

B. The Translational Fiber Bundle

The transformation law (163) has a geometrical interpretation. A translation acts on the nuclear configuration space according to $x \mapsto x + \xi$, or, in coordinates, $(\mathbf{X}_1, \dots, \mathbf{X}_N) \mapsto (\mathbf{X}_1 + \xi, \dots, \mathbf{X}_N + \xi)$. In the alternative coordinate system on the nuclear configuration space (see subsection II E) this becomes $(\mathbf{X}_{CM}, \mathbf{Y}_{\kappa}) \mapsto (\mathbf{X}_{CM} + \xi, \mathbf{Y}_{\kappa})$, that is, the center of mass is translated and the translationally invariant Jacobi vectors do not change. Nuclear configuration space in this coordinate system is illustrated in Fig. 1, in which the vertical \mathbf{X}_{CM} -axis is really a 3-dimensional subspace, while the complementary plane, upon which the Jacobi vectors \mathbf{Y}_{κ} are coordinates, has dimension 3N - 3. We will call this plane the "translation-reduced configuration space" (TRCS in the figure). These two subspaces are orthogonal to one another in the kinetic energy metric.

According to (163), conjugation by $T_e(\xi)$ maps the electronic Hamiltonian at nuclear configuration x into that at configuration $x + \xi$. This calls attention to the surface swept out when the entire translation group is allowed to act on x, that is, the set of all points $x + \xi$ for fixed x when ξ runs over \mathbb{R}^3 . It is the set of points that can be reached from x by applying translation operators. It is otherwise the *orbit* of the point x under the action of the translation group; it is a 3-dimensional subspace parallel to the X_{CM} -axis. As drawn it is not a vector subspace because it does not pass through the origin, but it does pass through a unique point of the translation-reduced configuration space, labeled x_0 in the figure, whose X_{CM} coordinate is 0. The action of the translation group foliates nuclear configuration space into a (3N - 3)-parameter family of 3-dimensional orbits, which are the fibers of the translational fiber bundle. In this context, "fibers" and "orbits" are the same thing; we will call them the "translational fibers." Any point in a given fiber can be reached from any other point of the same fiber by means of a translation operator. The 3N - 3 parameters labeling the translational fibers are the components of the Jacobi vectors Y_{κ} . The translation-reduced configuration space is an example of a *section* of the fiber bundle, that is, it is a surface of complementary dimensionality that cuts through the fibers, intersecting each at a unique point. The section plays the role of an initial value surface.

C. Transformation of Basis Vectors

The phase and frame conventions for the adiabatic basis vectors $|ax;k\rangle$, assumed to be invariant under time reversal, are discussed in subsection III G. Thus there is the question of the phase and frame conventions for these basis vectors as we vary *x*. We would like these vectors to be smooth functions of *x*.

Suppose $|ax_0;k\rangle$ is an eigenvector of $H_e(x_0)$ at point x_0 on the translation-reduced configuration space, so that $H_e(x_0)|ax_0;k\rangle = \varepsilon_k(x_0)|ax_0;k\rangle$. Let us define $|ax;k\rangle$ at other points of the same translational fiber by

$$|ax;k\rangle = T_e(\boldsymbol{\xi})|ax_0;k\rangle,\tag{164}$$

where $x = x_0 + \xi$. Then $|ax;k\rangle$ is automatically an eigenstate of $H_e(x)$,

$$H_e(x)|ax;k\rangle = \mathcal{E}_k(x_0)|ax;k\rangle,\tag{165}$$

with an eigenvalue that has not changed. That is, the eigenvalue is invariant under translations,

$$\boldsymbol{\varepsilon}_k(\boldsymbol{x}) = \boldsymbol{\varepsilon}_k(\boldsymbol{x} + \boldsymbol{\xi}). \tag{166}$$

The use of translation operators to define the phase of the electronic eigenstates, as in (164), makes those eigenstates smooth functions of x along the translational fibers. If we make a smooth choice of phase conventions for $|x_0;k\rangle$ for x_0 in some region of the translation-reduced configuration space of dimensionality 3N - 3, and then extend those definitions along translational fibers according to (164), then we will have smooth conventions over a region of dimensionality 3N of the full nuclear configuration space.

Now let $x_1 = x_0 + \xi_0$, let ξ_1 be another displacement, and consider

$$T(\boldsymbol{\xi}_{1})|ax_{1};k\rangle = T(\boldsymbol{\xi}_{1})T(\boldsymbol{\xi}_{0})|ax_{0};k\rangle = T(\boldsymbol{\xi}_{1} + \boldsymbol{\xi}_{0})|ax_{0};k\rangle$$
$$= |a,x_{0} + \boldsymbol{\xi}_{0} + \boldsymbol{\xi}_{1};k\rangle = |a,x_{1} + \boldsymbol{\xi}_{1};k\rangle.$$
(167)

With a change of notation this is

$$T(\boldsymbol{\xi})|ax;k\rangle = |a,x+\boldsymbol{\xi};k\rangle, \tag{168}$$

which is like (164) except that it applies to any point *x* on a translational fiber, not just the initial point x_0 .

If we assume that the matrix that transforms us from the adiabatic to the diabatic basis is independent of \mathbf{X}_{CM} , then the transformation law (168) also applies to the diabatic basis or any working basis we construct. This means that the transformation matrix is constant along translational fibers and that it depends only on the Jacobi vectors \mathbf{Y}_{κ} . Then we have

$$T(\boldsymbol{\xi})|\boldsymbol{x};\boldsymbol{k}\rangle = |\boldsymbol{x} + \boldsymbol{\xi};\boldsymbol{k}\rangle,\tag{169}$$

for all electronic bases that we consider. This is the transformation law for electronic basis states under translations.

D. Molecular Momentum in Various Representations

Equation (169) leads to an important result when we let the displacement vector $\boldsymbol{\xi}$ be infinitesimal. In that case we can expand the translation operator $T_e(\boldsymbol{\xi})$ (see (160)) and the state $|x + \boldsymbol{\xi}; k\rangle$, obtaining

$$\left[1 - \frac{i}{\hbar} \boldsymbol{\xi} \cdot \mathbf{p}_t\right] |x; k\rangle = |x; k\rangle + \sum_{\alpha=1}^N \boldsymbol{\xi} \cdot \nabla_\alpha |x; k\rangle,$$
(170)

or, since $\boldsymbol{\xi}$ is arbitrary,

$$\mathbf{p}_t |x;k\rangle = i\hbar \sum_{\alpha=1}^N \nabla_\alpha |x;k\rangle.$$
(171)

Now writing $\mathbf{P}_{\alpha} = -i\hbar \nabla_{\alpha}$ for the operator that acts on the parametric *x*-dependence of $|x;k\rangle$ and using (28), we can express the result as

$$(\mathbf{P}_t + \mathbf{p}_t)|x;k\rangle = 0. \tag{172}$$

Equation (172) leads to an important conclusion. In the molecular representation the operator $\mathbf{P}_{CM} = \mathbf{P}_t + \mathbf{p}_t$ is the total momentum of the system, which is conserved. Let us allow this operator to act on a molecular wave function $\Psi(\mathbf{X}, \mathbf{r})$, which we expand according to (10). This gives

$$\mathbf{P}_{\mathrm{CM}}\Psi(\mathbf{X},\mathbf{r}) = \sum_{k} [(\mathbf{P}_{t} + \mathbf{p}_{t})\psi_{k}(\mathbf{X})]\phi_{k}(x;\mathbf{r}) + \psi_{k}(\mathbf{X})[(\mathbf{P}_{t} + \mathbf{p}_{t})\phi_{k}(x;\mathbf{r})]$$

$$= \sum_{k} [\mathbf{P}_{t}\psi_{k}(\mathbf{X})]\phi_{k}(x;\mathbf{r}), \qquad (173)$$

where we distribute $\mathbf{P}_t + \mathbf{p}_t$, a first-order, linear differential operator, over the two factors according to the product rule, where the second major term vanishes according to (172), and where the operator \mathbf{p}_t in the first major term does not contribute since $\psi_k(\mathbf{X})$ is independent of \mathbf{r} .

We summarize this by writing out the transformation of P_{CM} from the molecular to the Born-Oppenheimer representation,

$$\mathbf{P}_{\rm CM} = \mathbf{P}_t + \mathbf{p}_t \longleftrightarrow \hat{\mathbf{P}}_t \,\delta_{kl},\tag{174}$$

where we put a hat on the nuclear operator that appears in the Born-Oppenheimer representation. In other words, in the Born-Oppenheimer representation, the operator that looks like the total nuclear momentum actually represents the total momentum of the molecule, including that of the electrons. Moreover, this is exact. This in turn implies that

$$T(\boldsymbol{\xi}) = e^{-i\boldsymbol{\xi}\cdot(\mathbf{P}_l + \mathbf{p}_l)/\hbar} \longleftrightarrow \widehat{T_n(\boldsymbol{\xi})} \,\delta_{kl} = e^{-i\boldsymbol{\xi}\cdot\hat{\mathbf{P}}_l/\hbar} \,\delta_{kl} \tag{175}$$

(see (160)). That is, in the Born-Oppenheimer representation what looks like a purely nuclear translation is actually a translation of the whole molecule, including the electrons.

E. Transformation of Derivative Couplings Under Translations

Finally, we consider how the derivative couplings transform under translations, that is, we ask how $\mathbf{F}_{\alpha;kl}(x+\xi)$ is related to $\mathbf{F}_{\alpha;kl}(x)$. As above we let $\xi = (\xi, \xi, ..., \xi)$. In addition we let $\eta = (\eta_1, \eta_2, ..., \eta_N)$ be an infinitesimal displacement in which the component vectors η_{α} are allowed to be different. Then

$$|x+\eta;l\rangle = |x;l\rangle + \sum_{\alpha=1}^{N} \eta_{\alpha} \cdot \nabla_{\alpha} |x;l\rangle,$$
 (176)

so that

$$\langle x; k | x + \eta; l \rangle = \delta_{kl} + \sum_{\alpha=1}^{N} \eta_{\alpha} \cdot \mathbf{F}_{\alpha;kl}(x).$$
 (177)

But x is a dummy variable in this equation, so we can replace it by $x + \xi$. If we do, the left-hand side does not change, since by (169) we have

$$\langle x + \boldsymbol{\xi}; k | x + \boldsymbol{\eta} + \boldsymbol{\xi}; l \rangle = \langle x; k | T_e(\boldsymbol{\xi})^{\dagger} T_e(\boldsymbol{\xi}) | x + \boldsymbol{\eta}; l \rangle = \langle x; k | x + \boldsymbol{\eta}; l \rangle.$$
(178)

Therefore the right-hand side of (177) does not change, either, and since the η_{α} are arbitrary we have

$$\mathbf{F}_{\alpha;kl}(x) = \mathbf{F}_{\alpha;kl}(x+\xi). \tag{179}$$

That is, the derivative couplings are invariant under nuclear translations, they are constant along translational fibers and they are independent of \mathbf{X}_{CM} , depending only on the Jacobi vectors \mathbf{Y}_{κ} . This means that, regarded as nuclear operators, that is, functions of \mathbf{X} or $\hat{\mathbf{X}}$, they commute with both $\hat{\mathbf{P}}_t$ and nuclear translations $\widehat{T_n(\boldsymbol{\xi})}$, where we are careful to put hats on nuclear operators that appear in the Born-Oppenheimer representation.

Nuclear translation operators satisfy the commutation relations,

$$\widehat{T_n(\boldsymbol{\xi})}\,\hat{\mathbf{X}}_{\alpha}\,\widehat{T_n(\boldsymbol{\xi})}^{\dagger} = \hat{\mathbf{X}}_{\alpha} - \boldsymbol{\xi}, \qquad \widehat{T_n(\boldsymbol{\xi})}\,\hat{\mathbf{P}}_{\alpha}\,\widehat{T_n(\boldsymbol{\xi})}^{\dagger} = \hat{\mathbf{P}}_{\alpha}, \tag{180}$$

so from (69) we see that the generator \hat{G}_1 commutes with nuclear translations and hence with $\hat{\mathbf{P}}_t$. More precisely, all operators are represented by matrices, but the matrices for $\hat{\mathbf{P}}_t$ and $\widehat{T_n(\boldsymbol{\xi})}$ are multiples of the identity, so each component $\hat{G}_{1,kl}$ commutes with both $\hat{\mathbf{P}}_t$ and $\widehat{T_n(\boldsymbol{\xi})}$. The same, it can be shown, applies to all the higher order generators \hat{G}_n for any n > 1.

This in turn means that all the commutators and iterated commutators in the dressing of $\hat{\mathbf{P}}_t$ vanish (see (55) for the series of commutators), so that the dressing transformation does nothing to $\hat{\mathbf{P}}_t$ or to the nuclear translation operators $\widehat{T_n}(\boldsymbol{\xi})$. Thus, the total linear momentum of the molecule, represented by $\mathbf{P}_{\text{CM}} = \mathbf{P}_t + \mathbf{p}_t$ in the molecular representation, is represented by $\hat{\mathbf{P}}_t \, \delta_{kl}$ in the Born-Oppenheimer and in all dressed Born-Oppenheimer representations, to all orders. Moreover, this is exact.

We can summarize this situation in a notation like that used above for the electronic momentum and current. We transform the molecular operator $\mathbf{P}_{\rm CM} = \mathbf{P}_t + \mathbf{p}_t$ to the Born-Oppenheimer representation, writing $(\mathbf{P}_{\rm CM})$ for the resulting matrix of nuclear operators with matrix elements $(\mathbf{P}_{\rm CM})_{kl} = \mathbf{\hat{P}}_t \,\delta_{kl}$. The corresponding matrix of Weyl symbols is $(\mathbf{P}_{\rm CM})$ (without the hat) with components $(\mathbf{P}_{\rm CM})_{kl} = \mathbf{P}_t \,\delta_{kl}$. We define the dressed versions of these by the matrix $(\mathbf{\bar{P}}_{\rm CM}) = \hat{U}(\mathbf{P}_{\rm CM})\hat{U}^{\dagger}$ with components $(\mathbf{\bar{P}}_{\rm CM})_{kl} = \mathbf{\hat{P}}_t \,\delta_{kl}$, or as Weyl symbols, $(\mathbf{\bar{P}}_{\rm CM})_{kl} = \mathbf{P}_t \,\delta_{kl}$.

F. The Molecular Center of Mass

We can treat the molecular center of mass \mathbf{R}_{CM} similarly. We write $\mathbf{R}_{CM} = \mathbf{R}_{1,CM} + \mathbf{R}_{2,CM}$ for the two terms seen in (30). When we transform these to the Born-Oppenheimer representation, we obtain

$$(\mathbf{R}_{1,\mathrm{CM}})_{kl} = \left(\frac{1}{M_{\mathrm{mol}}}\sum_{\alpha=1}^{N} M_{\alpha} \mathbf{X}_{\alpha}\right) \delta_{kl}, \qquad (181a)$$

$$(\mathbf{R}_{2,\mathrm{CM}})_{kl} = \frac{m_e}{M_{\mathrm{mol}}} \sum_{i=1}^{N_e} \langle x; k | \mathbf{r}_i | x; l \rangle.$$
(181b)

These are purely multiplicative nuclear operators, that is, functions of **X** only, and if we wish to emphasize their operator aspect we can put hats over them. Note that $\mathbf{R}_{2,CM}$ is of order κ^4 in comparison to $\mathbf{R}_{1,CM}$.

We transform the center of mass to the dressed Born-Oppenheimer representation according to $\widehat{(\mathbf{R}_{CM})} = \widehat{U}(\widehat{\mathbf{R}_{CM}})\widehat{U}^{\dagger}$, working with Weyl symbols and carrying out the expansion through order \hbar^2 . We start with $(\mathbf{R}_{1,CM})$. Since this is a multiple of the identity, the first correction term is

$$\hbar[G_1, (\mathbf{R}_{1, \text{CM}})]_{*,kl} = i\hbar^2 \{G_{1,kl}, \mathbf{R}_{1, \text{CM}}\} + O(\hbar^4),$$
(182)

which is of order \hbar^2 as shown, and all other corrections are of order \hbar^3 or higher. The bracket shown is the Poisson bracket and the series is a version of (A13). But G_1 is purely off-blockdiagonal, so through order \hbar^2 , the corrections are, too. The (nk)-components of these corrections, for $k \neq n$, are, according to (77) and (181a),

$$\frac{\hbar^2}{M_{\rm mol}} \sum_{\alpha=1}^N (\nabla_\alpha \langle x; n |) R(\varepsilon_n, x) | x; k \rangle.$$
(183)

But

$$\sum_{i=1}^{N} \nabla_{\alpha} |x;n\rangle = \frac{i}{\hbar} \mathbf{P}_{t} |x;n\rangle = -\frac{i}{\hbar} \mathbf{p}_{t} |x;n\rangle = -\frac{m_{e}}{\hbar^{2}} \sum_{i=1}^{N_{e}} (\varepsilon_{n} - H_{e}) \mathbf{r}_{i} |x;n\rangle,$$
(184)

so with this substitution the resolvent in (183) is canceled and the correction becomes

$$-\frac{m_e}{M_{\rm mol}} \sum_{i=1}^{N_e} \langle x; n | \mathbf{r}_i | x; k \rangle.$$
(185)

This exactly cancels the off-diagonal contribution of (181b), and we see that $(\bar{\mathbf{R}}_{CM})$ is purely even through order \hbar^2 . In other words, the dressing transformation that block-diagonalizes the Hamiltonian also block-diagonalizes the molecular center-of-mass position, at least through order \hbar^2 . We believe this must be true to all orders. When the equation of motion $(d/dt)\mathbf{R}_{CM} = (1/i\hbar)[\mathbf{R}_{CM}, H] = \mathbf{P}_{CM}/M_{mol}$, which is valid in the molecular representation, is transformed to the dressed Born-Oppenheimer representation it becomes

$$\frac{1}{i\hbar}[\widehat{(\mathbf{\bar{R}}_{CM})},\widehat{H}] = \frac{\widehat{\mathbf{P}}_t}{M_{mol}}I,$$
(186)

where *I* is the identity matrix. But since \hat{H} and the identity are purely even, the odd part of $(\mathbf{\bar{R}}_{CM})$ must commute with \hat{H} and therefore be a vector constant of motion. The only vector constants of motion are the linear and angular momentum, both of which are purely even in the dressed Born-Oppenheimer representation. Therefore the odd part of $(\mathbf{\bar{R}}_{CM})$ cannot be proportional to these, and the only choice left is zero.

Assuming this is true, the center of mass of the molecule \mathbf{R}_{CM} is represented on surface *n* by a scalar operator (that is, not a matrix), which is the one component $(\mathbf{\bar{R}}_{CM})_{nn}$. This is just as the Hamiltonian is represented by the operator \hat{K} and the total momentum of the molecule \mathbf{P}_{CM} is represented by $\mathbf{\hat{P}}_t$. Therefore we must have an exact equation of motion for the center-of-mass position in the dressed Born-Oppenheimer representation,

$$\frac{d(\bar{\mathbf{R}}_{\rm CM})_{nn}}{dt} = \frac{1}{i\hbar} [(\bar{\mathbf{R}}_{\rm CM})_{nn}, K]_* = \frac{\mathbf{P}_t}{M_{\rm mol}},\tag{187}$$

here expressed in terms of Weyl symbols. But, as we have shown, the diagonal elements $(\bar{\mathbf{R}}_{CM})_{nn}$ and $(\mathbf{R}_{CM})_{nn}$ are the same through order \hbar^2 , so we can write

$$(\bar{\mathbf{R}}_{\mathrm{CM}})_{nn} = (\bar{\mathbf{R}}_{1,\mathrm{CM}})_{nn} + (\bar{\mathbf{R}}_{2,\mathrm{CM}})_{nn} = \frac{1}{M_{\mathrm{mol}}} \sum_{\alpha=1}^{N} M_{\alpha} \mathbf{X}_{\alpha} + \frac{m_{e}}{M_{\mathrm{mol}}} \sum_{i=1}^{N_{e}} \langle x; n | \mathbf{r}_{i} | x; n \rangle.$$
(188)

This is the explicit form for the molecular center of mass on surface *n* in the dressed Born-Oppenheimer representation, valid through order κ^4 .

The molecular center of mass must exhibit uniform, rectilinear motion, which means that it must satisfy the equation of motion (187). We now check that it does. We write the symbol of the Hamiltonian as $K = K_0 + \hbar K_1 + \hbar^2 (K_{21} + K_{22})$, where $K_0 = H_{0,nn}$ is given by (47), where $K_1 = 0$, where $K_{21} = H_{2,nn}$ is the diagonal Born-Oppenheimer correction seen in (53), and where K_{22} is given by (95). The leading term is

$$\frac{1}{i\hbar}[(\bar{\mathbf{R}}_{1,\text{CM}})_{nn}, K_0]_* = \{(\bar{\mathbf{R}}_{1,\text{CM}})_{nn}, K_0\} = \frac{\mathbf{P}_t}{M_{\text{mol}}},$$
(189)

which is the right answer, so the correction terms must cancel. In all these calculations the Moyal *-bracket reduces to a Poisson bracket. One correction term is

$$\frac{1}{i\hbar}[(\bar{\mathbf{R}}_{2,\mathrm{CM}})_{nn},K_0]_* = \{(\bar{\mathbf{R}}_{2,\mathrm{CM}})_{nn},K_0\} = \frac{m_e}{M_{\mathrm{mol}}}\sum_{i=1}^{N_e}\sum_{\alpha=1}^{N_e}\frac{1}{M_\alpha}\mathbf{P}_\alpha\cdot\nabla_\alpha(\langle x;n|\mathbf{r}_i|x;n\rangle).$$
(190)

The contribution from K_{21} vanishes, and that from K_{22} is

$$\frac{i}{\hbar}[(\bar{\mathbf{R}}_{1,\mathrm{CM}})_{nn},\hbar^2 K_{22}]_* = \{(\bar{\mathbf{R}}_{1,\mathrm{CM}})_{nn},\hbar^2 K_{22}\} = \frac{2}{M_{\mathrm{mol}}} \sum_{\alpha,\beta} \frac{1}{M_{\beta}} (\nabla_{\alpha} \langle x;n|) R(\varepsilon_n,x) (\nabla_{\beta} |x;n\rangle) \cdot \mathbf{P}_{\beta}.$$
(191)

But with the aid of (184) this is shown to be equal to the first correction (190), with the opposite sign.

Thus the corrections do cancel, and the equation (187) of uniform, rectilinear motion is shown to be valid through order κ^4 . Notice that the correction K_{22} to the Hamiltonian is required for this result.

G. Translational Component of the Current

We return to the point raised at the end of subsection IV C, that it should be possible to simplify the translational component of the current. We return to the expression (156) for the Weyl symbol of the operator that represents the electronic current on a single surface n in the dressed Born-Oppenheimer representation. We note that

$$P^{\mu}\partial_{\mu} = \sum_{\alpha=1}^{N} \frac{1}{M_{\alpha}} \mathbf{P}_{\alpha} \cdot \nabla_{\alpha} = \frac{1}{M_{t}} \mathbf{P}_{t} \cdot \frac{\partial}{\partial \mathbf{X}_{\mathrm{CM}}} + \sum_{\kappa=1}^{N-1} \frac{1}{\mu_{\alpha}} \mathbf{Q}_{\kappa} \cdot \frac{\partial}{\partial \mathbf{Y}_{\kappa}},$$
(192)

where we use Appendix D and where in the second step we carry out the coordinate transformation $X_{\alpha} \rightarrow (X_{CM}, Y_{\kappa})$ on the nuclear configuration space (see subsection II E). The first term on the right gives the translational component of the current; we note that

$$\frac{\partial}{\partial \mathbf{X}_{\rm CM}} = \sum_{\alpha=1}^{N} \nabla_{\alpha},\tag{193}$$

and we define the translational component of the current operator $I_{r_0}^{\sigma}$ by

$$I_{\mathrm{CM},r_0}^{\sigma} = -\frac{2i\hbar}{M_t} \mathbf{P}_t \cdot \langle x; n | J_{r_0}^{\sigma} R(\varepsilon_n, x) \sum_{\alpha=1}^N \nabla_{\alpha} | x; n \rangle.$$
(194)

We substitute (184) into this, thereby eliminating the resolvent and obtaining

$$I_{\mathrm{CM},r_0}^{\sigma} = \frac{2im_e}{\hbar M_t} \mathbf{P}_t \cdot \langle x; n | J_{r_0}^{\sigma} Q(x) \sum_{i=1}^{N_e} \mathbf{r}_i | x; n \rangle.$$
(195)

Since $\langle x; n | J_{r_0}^{\sigma} | x; n \rangle = 0$ the projector Q(x) can be dropped. The current is a vector in the electronic configuration space with $3N_e$ components that we can organize as a collection of N_e 3-vectors, one

associated with each electron. For that purpose let us replace σ by the double index (jJ), as in Appendix D. We also substitute the definition (154) of $J_{r_0}^{\sigma}$. Then we have

$$I_{\text{CM},r_0}^{jJ} = \frac{i}{\hbar M_t} \sum_{i=1}^{N_e} \sum_{I=1}^{3} P_t^I \langle x; n | [p_{jJ} \delta(r - r_0) + \delta(r - r_0) p_{jJ}] r_{iI} | x; n \rangle.$$
(196)

The operator in the matrix element can be written

$$p_{jJ}\delta(r-r_0)r_{iI}+\delta(r-r_0)r_{iI}p_{jJ}-i\hbar\delta(r-r_0)\delta_{ij}\,\delta_{IJ},$$
(197)

whereupon the first two terms constitute an Hermitian operator that is odd under time-reversal, whose *nn*-matrix element vanishes. Only the last term in (197) contributes, and we obtain

$$I_{\text{CM},r_0}^{jJ} = \frac{P_t^J}{M_t} |\phi_n(x;r_0)|^2.$$
(198)

The current vector is independent of j, that is, it is the same for all the electrons, and it is given by the velocity of the nuclear center of mass \mathbf{P}_t/M_t times the electronic probability density at r_0 . This is just what we expect for the translational component of the current.

H. Translational Components of the Derivative Couplings

Let us return to the original Born-Oppenheimer representation (the middle entry of the first row of Table I), for which the Hamiltonian is given by (59) in Weyl symbol form. The first order Hamiltonian (59b) is essentially the nuclear velocity contracted against the derivative couplings, which are the components of a differential form on nuclear configuration space. The linear differential operator that appears in H_1 is the same as in (192), which allows us to pick out the translational component of the derivative couplings. We define

$$\mathbf{F}_{\mathrm{CM},kl}(x) = \langle x; k | \frac{\partial}{\partial \mathbf{X}_{\mathrm{CM}}} | x; l \rangle, \qquad \mathbf{F}_{\kappa;kl}(x) = \langle x; k | \frac{\partial}{\partial \mathbf{Y}_{\kappa}} | x; l \rangle, \tag{199}$$

and we note that the center-of-mass component can be written in several different ways,

$$\mathbf{F}_{\mathrm{CM},kl}(x) = \langle x;k | \sum_{\alpha=1}^{N} \nabla_{\alpha} | x;l \rangle = \sum_{\alpha=1}^{N} \mathbf{F}_{\alpha,kl}(x) = \frac{i}{\hbar} \langle x;k | \hat{\mathbf{P}}_{t} | x;l \rangle = -\frac{i}{\hbar} \langle x;k | \mathbf{p}_{t} | x;l \rangle,$$
(200)

where we use (172).

Thus the translational component of $H_{1,kl}$ is

$$-i\frac{\mathbf{P}_t}{M_t} \cdot \mathbf{F}_{\mathrm{CM},kl}(x). \tag{201}$$

This component sometimes causes confusion, especially when the basis vectors $|x;k\rangle$ (some of them, at least) are the adiabatic basis, since it implies transitions between potential energy surfaces when the nuclear motion is purely translational. This is most clear in models in which the nuclear coordinates are treated as classical variables, since in that case \mathbf{P}_t/M_t is the nuclear velocity and the term (201) becomes simply $-i\langle x;k|(d/dt)|x;l\rangle$. For example, it would appear that in the uniform, rectilinear motion of a hydrogen atom we have transitions between the 1*s* and 2*p* levels, which does not make sense physically. This leads to the suggestion, which we now pursue, of eliminating the translational component (201) of H_1 by means of a unitary transformation. We explain the hydrogen-atom paradox in subsection V K.

To this end, let the working basis $|x;k\rangle$ be an arbitrary discrete basis. We do not assume that it diagonalizes or block-diagonalizes the electronic Hamiltonian $H_e(x)$, and there is no privileged subspace or concept of even or odd operators. As above, we let \hat{G}_1 be a generator and we transform the Hamiltonian according to $\hat{H}' = \hat{U}_1 \hat{H} \hat{U}_1^{\dagger}$, where $\hat{U}_1 = \exp(\hbar \hat{G}_1)$. Upon translating to Weyl symbols and expanding to first order, this gives

$$H_1' = H_1 + [G_1, H_0], (202)$$

which is expressed in terms of matrices of Weyl symbols. We wish to choose G_1 so that

$$[G_1, H_0]_{kl} = H'_{1,kl} - H_{1,kl} = i \frac{\mathbf{P}_t}{M_t} \cdot \mathbf{F}_{\mathrm{CM},kl}(x) = -\frac{iN_e m_e}{\hbar^2 M_t} \mathbf{P}_t \cdot \langle x; k | [\mathbf{r}_{\mathrm{CM}}, H_e(x)] | x; l \rangle,$$
(203)

for all *k*, *l*, to remove the translational component of *H*₁. In the last step we have used (200) and $\mathbf{p}_t = N_e m_e (d/dt) \mathbf{r}_{\text{CM}}.$

The left-hand side of (203) can be written $[G_1, W]_{kl}$ where $W_{kl} = \langle x; k | H_e(x) | x; l \rangle$, since the kinetic energy part of H_0 , a multiple of the identity, does not contribute, while the right-hand side is proportional to $[(\mathbf{r}_{\text{CM}}), W]_{kl}$, where $(\mathbf{r}_{\text{CM}})_{kl} = \langle x; k | \mathbf{r}_{\text{CM}} | x; l \rangle$ is the matrix representing \mathbf{r}_{CM} in the Born-Oppenheimer representation. Therefore a solution of (203) is given by

$$G_{1,kl} = -\frac{iN_e m_e}{\hbar^2 M_t} \mathbf{P}_t \cdot \langle x; k | \mathbf{r}_{\rm CM} | x; l \rangle.$$
(204)

We can add to this any matrix that commutes with W, of which a multiple of the identity is an obvious choice.

The procedure being carried out here, the elimination of the translational components of the derivative couplings, bears some relation to the introduction of electron translation factors in electronic structure calculations (Delos (1981); Fatehi *et al.* (2011)), which we will explore in future

publications. In that context, the electron translation factors are also not unique, although a meaningful choice can be made in an atomic orbital basis.

One obvious shortcoming of (204) is that the right-hand side does not commute with translations. We wish G_1 to be translationally invariant, so that U_1 will map the old Hamiltonian H, which is translationally invariant, into a new one H' with the same property. Also, making G_1 translationally invariant means that $\mathbf{P}_t \delta_{kl}$, the matrix representing the total linear momentum of the molecule, is not changed by the transformation U_1 . To show that the right-hand side of (204) is not translationally invariant we conjugate it with a nuclear translation operator, whereupon \mathbf{P}_t goes into itself and the matrix element transforms according to

$$T_{n}(\boldsymbol{\xi})\langle x;k|\mathbf{r}_{\mathrm{CM}}|x;l\rangle T_{n}(\boldsymbol{\xi})^{\dagger} = \langle x-\boldsymbol{\xi};k|\mathbf{r}_{\mathrm{CM}}|x-\boldsymbol{\xi};l\rangle = \langle x;k|T_{e}(\boldsymbol{\xi})\mathbf{r}_{\mathrm{CM}}T_{e}(\boldsymbol{\xi})^{\dagger}|x;l\rangle$$
$$= \langle x;k|\mathbf{r}_{\mathrm{CM}}|x;l\rangle - \boldsymbol{\xi}\,\delta_{kl}, \qquad (205)$$

where we use (180), (169) and (161).

Now for a given x let x_0 be the point on the translation-reduced configuration space on the same translational fiber as x, as illustrated in Fig. 1. This means that x and x_0 are connected by a translation; let $\boldsymbol{\xi}$ be the vector such that $T(\boldsymbol{\xi})x_0 = x$, that is, $x = x_0 + \boldsymbol{\xi}$. Then $\boldsymbol{\xi}$ is the same as the **X**_{CM} coordinate of x, and we have

$$\langle x_0; k | \mathbf{r}_{\rm CM} | x_0; l \rangle = \langle x; k | \mathbf{r}_{\rm CM} | x; l \rangle - \mathbf{X}_{\rm CM} \,\delta_{kl}.$$
(206)

The final term is a multiple of the identity, so we obtain a solution of (203) that is also translationally invariant if we replace (204) by

$$G_{1,kl} = -\frac{iN_e m_e}{\hbar^2 M_t} \mathbf{P}_t \cdot (\langle x; k | \mathbf{r}_{\mathrm{CM}} | x; l \rangle - \mathbf{X}_{\mathrm{CM}} \, \delta_{kl}).$$
(207)

I. An Exact Unitary Operator for Translations

Although in the practice of perturbation theory we expand the unitary operator $\hat{U}_1 = \exp(\hbar \hat{G}_1)$ in a power series, in fact for the generator shown in (207) it is possible to carry out the unitary transformation \hat{U}_1 exactly. That is, we can give exact formulas for the action of \hat{U}_1 on wave functions and for its action by conjugation on operators. This is easiest to see if we transform (207) to the molecular representation, whereupon we find

$$G_1 = -\frac{iN_e m_e}{\hbar^2 M_t} \left(\mathbf{P}_t + \mathbf{p}_t \right) \cdot \left(\mathbf{r}_{\rm CM} - \mathbf{X}_{\rm CM} \right), \tag{208}$$

where we use (174). To be clear about the notation, (207) gives the Weyl symbols of the components of the matrix that represents the operator in the Born-Oppenheimer representation. When these are converted into operators, the operator $\hat{\mathbf{P}}_t$ can be placed before or after the factor $\mathbf{r}_{CM} - \mathbf{X}_{CM}$, which is translationally invariant. When these operators are converted to the molecular representation, using the rules explained in subsection II C, we obtain (208). We omit the hats on operators when working in the molecular representation, but G_1 in (208) is an operator. Recall that $\mathbf{P}_{CM} = \mathbf{P}_t + \mathbf{p}_t$ in this representation (see (34)). For the rest of this subsection we work in the molecular representation.

Let us write the unitary operator as $U_1(\lambda) = \exp(\hbar G_1) = \exp(-i\lambda g_1/\hbar)$, where

$$g_1 = \mathbf{P}_{\mathrm{CM}} \cdot (\mathbf{r}_{\mathrm{CM}} - \mathbf{X}_{\mathrm{CM}}) \tag{209}$$

and where

$$\lambda = \lambda_0 = \frac{N_e m_e}{M_t}.$$
(210)

Let A = A(0) be an arbitrary operator and let

$$A(\lambda) = U_1(\lambda)A(0)U_1(\lambda)^{\dagger}.$$
(211)

We can solve for $A(\lambda)$ by promoting λ into a variable and solving the differential equation,

$$\frac{dA(\lambda)}{d\lambda} = -\frac{1}{i\hbar}[A, g_1], \qquad (212)$$

a version of the Heisenberg equations of motion.

For example, evaluating the commutators $[\mathbf{r}_i, g_1]$ and $[\mathbf{X}_{\alpha}, g_1]$ we find

$$\frac{d\mathbf{r}_{i}(\lambda)}{d\lambda} = \frac{d\mathbf{X}_{\alpha}(\lambda)}{d\lambda} = -(\mathbf{r}_{\rm CM} - \mathbf{X}_{\rm CM}).$$
(213)

As derived, the vector $\mathbf{r}_{CM} - \mathbf{X}_{CM}$ on the right-hand side should be evaluated at λ , but, in fact, it is easy to show that this vector is independent of λ so it can be evaluated anywhere, such as $\lambda = 0$. Thus we have the solutions,

$$\mathbf{r}_i(\lambda) = \mathbf{r}_i - \lambda (\mathbf{r}_{\rm CM} - \mathbf{X}_{\rm CM}), \qquad (214a)$$

$$\mathbf{X}_{\alpha}(\lambda) = \mathbf{X}_{\alpha} - \lambda(\mathbf{r}_{\mathrm{CM}} - \mathbf{X}_{\mathrm{CM}}), \qquad (214b)$$

where on the right-hand side all vectors are understood to be evaluated at $\lambda = 0$.

Similarly, we find

$$\frac{d\mathbf{p}_{i}(\lambda)}{d\lambda} = \frac{1}{N_{e}}\mathbf{P}_{\mathrm{CM}}, \qquad \frac{d\mathbf{P}_{\alpha}(\lambda)}{d\lambda} = -\frac{M_{\alpha}}{M_{t}}\mathbf{P}_{\mathrm{CM}}.$$
(215)

As before, we find that \mathbf{P}_{CM} is independent of λ . The solutions are

$$\mathbf{p}_i(\lambda) = \mathbf{p}_i + \frac{\lambda}{N_e} \mathbf{P}_{\text{CM}}, \qquad (216a)$$

$$\mathbf{P}_{\alpha}(\lambda) = \mathbf{P}_{\alpha} - \frac{\lambda M_{\alpha}}{M_{t}} \mathbf{P}_{\mathrm{CM}}, \qquad (216b)$$

where all vectors on the right-hand side are evaluated at $\lambda = 0$.

J. Eliminating $\mathbf{F}_{CM,kl}$ to All Orders

In this subsection we continue working in the molecular representation, omitting hats on nuclear operators.

The transformation laws (214) and (216) allow us to evaluate $U_1(\lambda)HU_1(\lambda)^{\dagger}$, where *H* is the molecular Hamiltonian (1). For $\lambda = \lambda_0$ this transformation should eliminate the translational components of the derivative couplings to order κ^2 . We find that the potential energy does not change,

$$U_1(\lambda) V_{\text{Coul}}(\mathbf{X}, \mathbf{r}) U_1(\lambda)^{\dagger} = V_{\text{Coul}}(\mathbf{X}, \mathbf{r}), \qquad (217)$$

since it depends only on vector differences among X_{α} and r_i .

As for the kinetic energy, we find

$$U_{1}(\lambda) \left(\sum_{\alpha=1}^{N} \frac{\mathbf{P}_{\alpha}^{2}}{2M_{\alpha}} + \sum_{i=1}^{N_{e}} \frac{\mathbf{p}_{i}^{2}}{2m_{e}} \right) U_{1}(\lambda)^{\dagger} = \frac{1}{2M_{t}} (\mathbf{P}_{\rm CM} - \mathbf{p}_{t})^{2} + \sum_{\kappa=1}^{N-1} \frac{\mathbf{Q}_{\kappa}^{2}}{2\mu_{\kappa}} + \sum_{i=1}^{N_{e}} \frac{\mathbf{p}_{i}^{2}}{2m_{e}} + \lambda \left[\left(\frac{1}{N_{e}m_{e}} + \frac{1}{M_{t}} \right) \mathbf{p}_{t} \cdot \mathbf{P}_{\rm CM} - \frac{\mathbf{P}_{\rm CM}^{2}}{M_{t}} \right] + \lambda^{2} \left(\frac{1}{N_{e}m_{e}} + \frac{1}{M_{t}} \right) \frac{\mathbf{P}_{\rm CM}^{2}}{2},$$
(218)

where we have transformed the nuclear kinetic energy according to (29). In this result we have used $\mathbf{P}_t = \mathbf{P}_{CM} - \mathbf{p}_t$ to eliminate \mathbf{P}_t , and we note that $\mathbf{P}_{CM} \longleftrightarrow \mathbf{P}_t \delta_{kl}$ and $\mathbf{p}_t \longleftrightarrow i\hbar \mathbf{F}_{CM,kl}$, the latter of which follows from (200).

Thus in the molecular representation the terms that will produce the translational component (201) of $H_{1,kl}$ in the Born-Oppenheimer representation are those proportional to $\mathbf{p}_t \cdot \mathbf{P}_{CM}$. The coefficient of these terms in (218) is

$$-\frac{1}{M_t} + \lambda \left(\frac{1}{N_e m_e} + \frac{1}{M_t}\right). \tag{219}$$

If we set $\lambda = \lambda_0 = N_e m_e / M_t$, the coefficient of this term is reduced by a factor of order κ^4 , which is better than expected, since U_1 was only supposed to eliminate this term through order κ^2 . But

we see that if we choose the slightly different value,

$$\lambda = \lambda_1 = \frac{N_e m_e}{M_{\rm mol}} = \frac{N_e m_e}{M_t + N_e m_e} \tag{220}$$

then the translational component of the derivative couplings is eliminated exactly (to all orders in κ).

With this choice (220) of λ (and henceforth dropping the 1-subscript on λ and U), we find for the transformed molecular Hamiltonian,

$$U(\lambda)HU(\lambda)^{\dagger} = \frac{\mathbf{P}_{CM}^{2}}{2M_{mol}} + \sum_{\kappa=1}^{N-1} \frac{\mathbf{Q}_{\kappa}^{2}}{2\mu_{\kappa}} + \sum_{i=1}^{N_{e}} \frac{\mathbf{p}_{i}^{2}}{2m_{e}} + \frac{\mathbf{p}_{t}^{2}}{2M_{t}} + V_{Coul}(\mathbf{X}, \mathbf{r}).$$
(221)

Apart from the meanings of the symbols, the result is the same as the molecular Hamiltonian seen in (36), (37) and (38). In particular, the fourth term of (221) is the mass polarization term. Our unitary transformation $U(\lambda)$ reproduces the effects of the coordinate transformation discussed in subsection IIE, the purpose of which was to separate the translational degrees of freedom.

With the value (220) of λ we note that (30) can be written $\mathbf{R}_{CM} = (1 - \lambda)\mathbf{X}_{CM} + \lambda \mathbf{r}_{CM}$. Thus the transformation equations (214) and (216) become

$$U(\lambda)\mathbf{r}_{i}U(\lambda)^{\dagger} = \mathbf{r}_{i} - (\mathbf{R}_{\rm CM} - \mathbf{X}_{\rm CM}), \qquad U(\lambda)\mathbf{p}_{i}U(\lambda)^{\dagger} = \mathbf{p}_{i} + \frac{m_{e}}{M_{\rm mol}}\mathbf{P}_{\rm CM},$$
$$U(\lambda)\mathbf{X}_{\alpha}U(\lambda)^{\dagger} = \mathbf{X}_{\alpha} - (\mathbf{R}_{\rm CM} - \mathbf{X}_{\rm CM}), \qquad U(\lambda)\mathbf{P}_{\alpha}U(\lambda)^{\dagger} = \mathbf{P}_{\alpha} - \frac{N_{e}m_{e}M_{\alpha}}{M_{t}M_{\rm mol}}\mathbf{P}_{\rm CM}.$$
(222)

The unitary transformation itself becomes

$$U(\lambda) = \exp\left[-\frac{i}{\hbar}\mathbf{P}_{\rm CM} \cdot (\mathbf{R}_{\rm CM} - \mathbf{X}_{\rm CM})\right].$$
 (223)

An example of this transformation appeared in Cederbaum (2013), who used it in his study of translational invariance in the method of exact factorization.

K. Unitary Transformation as a Coordinate Transformation

Equations (222) look like a coordinate transformation taking us from variables \mathbf{r}_i , \mathbf{X}_{α} , etc., on the right to variables $U(\lambda)\mathbf{r}_i U(\lambda)^{\dagger}$, etc., on the left. It is convenient to call the variables on the right the "new" variables and those on the left the "old" ones. This seems counter-intuitive but it is the interpretation needed to obtain the transformation law (225) given below. We note that the transformation of the momenta follows from that of the coordinates, via the chain rule for the operators $\mathbf{p}_i = -i\hbar \partial/\partial \mathbf{r}_i$, etc. We will denote the new coordinates with a prime and the old without a prime. We obtain the coordinate transformation in this language if we replace \mathbf{r}_i , etc., on the right side of (222) with \mathbf{r}'_i , etc., and then replace $U(\lambda)\mathbf{r}_iU(\lambda)^{\dagger}$, etc., on the left with simply \mathbf{r}_i , etc. This gives us the old variables as functions of the new. Inverting these equations to solve for the new variables as functions of the old, we obtain

$$\mathbf{r}'_{i} = \mathbf{r}_{i} + (\mathbf{R}_{\rm CM} - \mathbf{X}_{\rm CM}), \qquad \mathbf{p}'_{i} = \mathbf{p}_{i} - \frac{m_{e}}{M_{\rm mol}} \mathbf{P}_{\rm CM},$$

$$\mathbf{X}'_{\alpha} = \mathbf{X}_{\alpha} + (\mathbf{R}_{\rm CM} - \mathbf{X}_{\rm CM}), \qquad \mathbf{P}'_{\alpha} = \mathbf{P}_{\alpha} + \frac{N_{e}m_{e}M_{\alpha}}{M_{t}M_{\rm mol}} \mathbf{P}_{\rm CM},$$
(224)

where we note that $\mathbf{R}'_{CM} - \mathbf{X}'_{CM} = \mathbf{R}_{CM} - \mathbf{X}_{CM}$ and $\mathbf{P}'_{CM} = \mathbf{P}_{CM}$.

Now if we call (1) the "old" Hamiltonian H, a function of the old variables \mathbf{r}_i , \mathbf{X}_{α} , etc., and (221) the "new" Hamiltonian H', regarded as a function of the new variables \mathbf{r}'_i , \mathbf{X}'_{α} , etc., then we have

$$H(\mathbf{X}, \mathbf{P}, \mathbf{r}, \mathbf{p}) = H'(\mathbf{r}', \mathbf{p}', \mathbf{X}', \mathbf{P}'), \qquad (225)$$

in which the primed variables are functions of the unprimed ones as shown in (224) and vice versa.

Alternatively, we may regard (222) as a dressing transformation, taking us from the lab molecular representation to the lab, translation-reduced representation (down the first column of Table I). The two points of view are complementary; in the dressing point of view the operators do not change but the wave functions do, while in the coordinate-transformation point of view it is the opposite. The dressing point of view has some interesting features, for example, in the translationreduced, molecular representation the operator that looks like the center of mass of the nuclei is physically the center of mass of the molecule, since

$$U(\lambda) \mathbf{R}_{\rm CM} U(\lambda)^{\dagger} = \mathbf{X}_{\rm CM}.$$
(226)

This reminds us of the fact that in the Born-Oppenheimer representation, the operator that looks like the nuclear angular momentum actually includes the electronic angular momentum, a main point of Littlejohn, Rawlinson, and Subotnik (2023). See also the end of subsection V D for the situation regarding the total momentum of the molecule.

Having achieved the translation-reduced molecular representation (the lower left corner of Table I) we can move to the right (across the lower row) in much the same manner as we did previously along the upper row, except that the the electronic Hamiltonian is now the translationreduced version (38) instead of the original version (2). In particular, the potential energy surfaces and adiabatic basis states are those of $H_{\text{TR}e}$ and not H_e , the former of which contains the masspolarization terms. In the case of hydrogen, mentioned in subsection V H, the mass-polarization term provides the correction between the true electron mass and the reduced mass; the latter is needed for the correct energy eigenvalues. Naturally, if we use the wrong electronic Hamiltonian then there will be transitions among the surfaces, even for uniform translational motion of the atom.

Once we have reached the dressed, translation-reduced representation (the lower right-hand corner of Table I) we will be in a position to separate the rotational degrees of freedom. This involves a coordinate transformation on the translation-reduced configuration space, taking us from the \mathbf{Y}_{κ} coordinates to orientational (Euler angle) and shape coordinates. This is often described as the construction of the kinetic energy operator on the internal space. There is an extensive literature on this procedure for single-surface problems in the electrostatic model, for example, Wang and Tucker Carrington (2000). Kendrick (2018) has given a partial analysis of the case of two coupled surfaces in the electrostatic model in his study of low-temperature scattering. The separation of rotational degrees of freedom in the case of fine structure effects has apparently not been worked out, but, as noted by Littlejohn, Rawlinson, and Subotnik (2023), in the odd-electron case the nuclear Born-Oppenheimer wave function is a 2-component pseudo-spinor. We will report on this problem in more detail in the future.

VI. CONCLUSIONS

This article is a part of a series in which we hope to introduce a geometrical flavor to Born-Oppenheimer theory and to reveal some new results, building on older work. The latter includes Littlejohn and Reinsch (1997), which dealt with rotations in the *n*-body problem but not specifically with Born-Oppenheimer theory. That article presented a rather complete explanation of the rotational fiber bundle structure of the nuclear configuration space. The first article of the new series was Littlejohn, Rawlinson, and Subotnik (2022), which was devoted to diabatic bases. This was necessary preparation since it is impossible to talk about the theory without diabatic bases, and because the existing literature on the subject has several shortcomings. The second, Littlejohn, Rawlinson, and Subotnik (2023), dealt with angular momentum and its conservation within Born-Oppenheimer theory. This article, the third, has explained Moyal theory for the separation of nuclear and electronic degrees of freedom, and translational invariance. Future articles

will treat the separation of rotational degrees of freedom, which practically speaking means the construction of kinetic energy operators on the internal space. We also plan an article on the interaction between molecular point groups and the geometry of the nuclear configuration space, especially the rotational fiber bundle. We feel that this is an important step in the development of a proper geometrical understanding of aspects of the Jahn-Teller effect and the bifurcation of degeneracy manifolds. Finally, we plan a parallel treatment of some of the same issues in the context of hybrid classical-quantum models, which are currently popular in surface hopping and related algorithms. For such models the unitary transformations implemented by the Moyal star product are replaced by canonical transformations on a phase space that incorporates both classical and quantum variables.

ACKNOWLEDGMENTS

JES was supported by U.S. Department of Energy (DoE), Office of Science, Office of Basic Energy Sciences, under Award No. DE-SC0019397.

Appendix A: Weyl-Moyal Formalism

Original articles and those that review the Wigner-Weyl-Moyal formalism include Weyl (1927); Wigner (1932); Groenewold (1946); Moyal (1949); Balazs and Jennings (1984); Littlejohn (1986); McDonald (1988); Osborn and Molzahn (1995). The notation in this appendix is independent of that of the rest of the paper.

1. The Weyl Symbol Correspondence

We work with wave functions $\psi(x)$, where $x = (x_1, ..., x_n) \in \mathbb{R}^n$. The conjugate momentum is $p = (p_1, ..., p_n)$. Operators are indicated by hats, for example, $\hat{x}_i =$ multiplication by x_i and $\hat{p}_i = -i\hbar \partial/\partial x_i$, when acting on wave functions $\psi(x)$. Classical quantities or *c*-numbers are indicated without the hat.

If \hat{A} is an operator then we define the *Weyl symbol* of \hat{A} as the function A(x, p) on the classical phase space given by

$$A(x,p) = \int ds \, e^{-ip \cdot s/\hbar} \, \langle x + s/2 | \hat{A} | x - s/2 \rangle, \tag{A1}$$
where $s \in \mathbb{R}^n$ and where the matrix element shown is the kernel of the integral operator \hat{A} in the *x*-representation, that is,

$$(\hat{A}\psi)(x) = \int dx' \langle x|\hat{A}|x'\rangle \,\psi(x'). \tag{A2}$$

Equation (A1) specifies the *Weyl transform* A(x, p) of the operator \hat{A} . The inverse Weyl transform is given by

$$\langle x|\hat{A}|x'\rangle = \int \frac{dp}{(2\pi\hbar)^n} e^{ip\cdot(x-x')/\hbar} A\left(\frac{x+x'}{2}, p\right). \tag{A3}$$

We denote the correspondence between operators and their Weyl symbols by $\hat{A} \longleftrightarrow A(x, p)$. This is not the same usage of the symbol \longleftrightarrow as in other parts of this paper, but what the notation has in common is that it represents a one-to-one correspondence between two sets of objects, in this case, operators and functions on phase space.

As special cases, we note that the operators $\hat{1}$ (the identity operator), \hat{x}_i and \hat{p}_i have Weyl symbols 1, x_i and p_i , respectively.

No information is lost by mapping operators to their Weyl symbols or vice versa. In particular, there is no "ordering ambiguity" on passing from symbols (objects that are nominally classical) to operators. For example, we have

$$\hat{x}_i \hat{p}_j \longleftrightarrow x_i p_j + \frac{i\hbar}{2} \delta_{ij},$$
 (A4a)

$$\hat{p}_j \hat{x}_i \longleftrightarrow x_i p_j - \frac{i\hbar}{2} \delta_{ij},$$
 (A4b)

which shows that the ordering of operators is indicated by terms in the symbol that are of higher order in \hbar . A useful generalization of (A4) is

$$f(\hat{x})\hat{p}_i \longleftrightarrow f(x)p_i + \frac{i\hbar}{2}\frac{\partial f(x)}{\partial x_i},$$
 (A5a)

$$\hat{p}_i f(\hat{x}) \longleftrightarrow f(x) p_i - \frac{i\hbar}{2} \frac{\partial f(x)}{\partial x_i}.$$
 (A5b)

If \hat{A} and \hat{B} are operators such that $\hat{A} = \hat{B}^{\dagger}$, then the corresponding symbols satisfy

$$A(x,p) = B(x,p)^*.$$
 (A6)

In particular, the Weyl symbol of a Hermitian operator is real. A special case of this follows from (A5),

$$\frac{1}{2}[\hat{p}_i f(\hat{x}) + f(\hat{x})\hat{p}_i] \longleftrightarrow f(x)p_i.$$
(A7)

2. The *-Product

If \hat{A} , \hat{B} and \hat{C} are operators such that $\hat{C} = \hat{A}\hat{B}$, then we write the corresponding relation between the symbols as

$$C(x,p) = A(x,p) * B(x,p), \tag{A8}$$

where * is the "Moyal star product." It has an expansion in powers of \hbar ,

$$C(x,p) = \sum_{n=0}^{\infty} \frac{1}{n!} \left(\frac{i\hbar}{2}\right)^n \{A,B\}_n,$$
 (A9)

where

$$\{A,B\}_n = A(x,p) \left(\frac{\overleftarrow{\partial}}{\partial x} \cdot \frac{\overrightarrow{\partial}}{\partial p} - \frac{\overleftarrow{\partial}}{\partial p} \cdot \frac{\overrightarrow{\partial}}{\partial x} \right)^n B(x,p).$$
(A10)

The arrows indicate the direction in which the partial derivatives act. The first few of these brackets are

$$\{A,B\}_0 = AB, \{A,B\}_1 = \{A,B\} = \frac{\partial A}{\partial x} \cdot \frac{\partial B}{\partial p} - \frac{\partial A}{\partial p} \cdot \frac{\partial B}{\partial x}, \{A,B\}_2 = \sum_{ij} \left(\frac{\partial^2 A}{\partial x_i \partial x_j} \frac{\partial^2 B}{\partial p_i \partial p_j} - 2 \frac{\partial^2 A}{\partial x_i \partial p_j} \frac{\partial^2 B}{\partial p_i \partial x_j} + \frac{\partial^2 A}{\partial p_i \partial p_j} \frac{\partial^2 B}{\partial x_i \partial x_j} \right),$$

etc. The zeroth order bracket is the ordinary product and the first order bracket is the ordinary Poisson bracket. If no subscript is given, the Poisson bracket is implied. The *n*-th order bracket is symmetric or antisymmetric in the exchange of the operands as n is even or odd, respectively. Writing out the first few terms of (A10), we have

$$A * B = AB + \frac{i\hbar}{2} \{A, B\} - \frac{\hbar^2}{8} \{A, B\}_2 + \dots$$
 (A12)

We also define the "Moyal bracket,"

$$[A,B]_* = A * B - B * A = i\hbar\{A,B\} - \frac{i\hbar^3}{24}\{A,B\}_3 + \dots,$$
(A13)

which is the Weyl symbol of the commutator $[\hat{A}, \hat{B}]$.

3. Matrices of Operators and Weyl Symbols

In the main body of the paper we make frequent use of commutators of matrices of operators. We now make some comments about what happens to such commutators when translated into symbols. Let \hat{A}_{kl} and \hat{B}_{kl} be two matrices of operators. By the commutator $\hat{C} = [\hat{A}, \hat{B}]$, we mean the matrix of operators,

$$\hat{C}_{kl} = [\hat{A}, \hat{B}]_{kl} = \sum_{p} \left(\hat{A}_{kp} \hat{B}_{pl} - \hat{B}_{kp} \hat{A}_{pl} \right).$$
(A14)

When this is translated into symbols, we obtain

$$C_{kl} = [A,B]_{*kl} = \sum_{p} (A_{kp} * B_{pl} - B_{kp} * A_{pl}) = \sum_{p} (A_{kp} B_{pl} - B_{kp} A_{pl}) + \frac{i\hbar}{2} \sum_{p} (\{A_{kp}, B_{pl}\} - \{B_{kp}, A_{pl}\}) - \frac{\hbar^2}{8} \sum_{p} (\{A_{kp}, B_{pl}\}_2 - \{B_{kp}, A_{pl}\}_2) + \dots$$
(A15)

This equation defines the Moyal bracket $C = [A, B]_*$ of matrices of symbols. Notice that the lowest, order \hbar^0 term in the expansion is otherwise just the commutator [A, B] of the symbol matrices, in which symbols are multiplied as *c*-numbers. That is, the first term of $[A, B]_*$ is [A, B] (without the star). Notice also that in the case of scalar operators, the expansion of the symbol of $[\hat{A}, \hat{B}]$ involves only odd powers of \hbar , as shown by (A13), whereas in the case of matrices of operators, all powers occur, including the power \hbar^0 .

Appendix B: Coordinates on Nuclear Configuration Space

The coordinate transformation $\mathbf{X}_{\alpha} \rightarrow (\mathbf{X}_{CM}, \mathbf{Y}_{\kappa})$ was introduced in subsection II E, where the definitions of \mathbf{X}_{CM} and \mathbf{P}_t were given (in (26) and (28)).

The new coordinates \mathbf{Y}_{κ} , $\kappa = 1, ..., N - 1$, are a set of N - 1 translationally invariant vectors defined by an $(N - 1) \times N$ matrix $B_{\kappa\alpha}$,

$$\mathbf{Y}_{\kappa} = \sum_{\alpha=1}^{N} B_{\kappa\alpha} \mathbf{X}_{\alpha}, \qquad \kappa = 1, \dots, N-1.$$
(B1)

The matrix $B_{\kappa\alpha}$ must satisfy

$$\sum_{\alpha=1}^{N} B_{\kappa\alpha} = 0, \tag{B2}$$

in order that the vectors \mathbf{Y}_{κ} be translationally invariant.

We require that the \mathbf{Y}_{κ} be Jacobi vectors (Delves (1960); Hirschfelder (1969); Aquilanti and Cavalli (1986); Gatti *et al.* (1998)). This means that the matrix $B_{\kappa\alpha}$ satisfies

$$\sum_{\alpha=1}^{N} \frac{B_{\kappa\alpha} B_{\lambda\alpha}}{M_{\alpha}} = \frac{\delta_{\kappa\lambda}}{\mu_{\kappa}},\tag{B3}$$

which is a kind of an orthogonality condition on $B_{\kappa\alpha}$ with respect to the kinetic energy metric. Here the μ_{κ} , $\kappa = 1, ..., N-1$, are reduced masses associated with the Jacobi vectors \mathbf{Y}_{κ} . The inverse coordinate transformation $(\mathbf{X}_{CM}, \mathbf{Y}_{\kappa}) \to \mathbf{X}_{\alpha}$ is given by

$$\mathbf{X}_{\alpha} = \mathbf{X}_{\mathrm{CM}} + \frac{1}{M_{\alpha}} \sum_{\kappa=1}^{N-1} \mu_{\kappa} B_{\kappa\alpha} \mathbf{Y}_{\kappa}.$$
 (B4)

The matrix $B_{\kappa\alpha}$ satisfies a kind of inverse orthogonality relation,

$$\frac{M_{\alpha}M_{\beta}}{M_{t}} + \sum_{\kappa=1}^{N-1} \mu_{\kappa} B_{\kappa\alpha} B_{\kappa\beta} = M_{\alpha} \,\delta_{\alpha\beta}. \tag{B5}$$

This may be proven by writing out the $N \times N$ matrix of the linear transformation $\mathbf{X}_{\alpha} \to (\mathbf{X}_{CM}, \mathbf{Y}_{\kappa})$ and the one for its inverse, $(\mathbf{X}_{CM}, \mathbf{Y}_{\kappa}) \to \mathbf{X}_{\alpha}$, and multiplying the two matrices in both orders.

For example, in the case of the water molecule, we let atoms 1 and 2 be hydrogens and atom 3 be the oxygen. Then we define Jacobi vectors $\mathbf{Y}_1 = \mathbf{X}_2 - \mathbf{X}_1$ and $\mathbf{Y}_2 = \mathbf{X}_3 - (\mathbf{X}_1 + \mathbf{X}_2)/2$, which implies

$$B_{\kappa\alpha} = \begin{pmatrix} -1 & 1 & 0\\ -1/2 & -1/2 & 1 \end{pmatrix}.$$
 (B6)

These imply reduced masses $\mu_1 = M_{\rm H}/2$, and

$$\frac{1}{\mu_2} = \frac{1}{2M_{\rm H}} + \frac{1}{M_{\rm O}}.$$
(B7)

The momenta conjugate to \mathbf{X}_{α} are \mathbf{P}_{α} , and those conjugate to $(\mathbf{X}_{CM}, \mathbf{Y}_{\kappa})$ are $(\mathbf{P}_t, \mathbf{Q}_{\kappa})$. The momenta acting on wave functions are defined by $\mathbf{P}_{\alpha} = -i\hbar\partial/\partial \mathbf{X}_{\alpha}$, $\mathbf{P}_t = -i\hbar\partial/\partial \mathbf{X}_{CM}$, and $\mathbf{Q}_{\alpha} = -i\hbar\partial/\partial \mathbf{Y}_{\kappa}$. One way to transform the momenta, which is consistent with the chain rule, is to demand the equality of the differential forms,

$$\sum_{\alpha=1}^{N} \mathbf{P}_{\alpha} \cdot d\mathbf{X}_{\alpha} = \mathbf{P}_{t} \cdot d\mathbf{X}_{CM} + \sum_{\kappa=1}^{N-1} \mathbf{Q}_{\kappa} \cdot d\mathbf{Y}_{\kappa}.$$
(B8)

Substituting the coordinate transformation (B4) into this gives (28), showing that \mathbf{P}_t is the total nuclear momentum, and it gives

$$\mathbf{Q}_{\kappa} = \sum_{\alpha=1}^{N} \frac{\mu_{\kappa}}{M_{\alpha}} B_{\kappa\alpha} \mathbf{P}_{\alpha}.$$
 (B9)

Substituting the the inverse transformation (26) and (B1) gives

$$\mathbf{P}_{\alpha} = \frac{M_{\alpha}}{M_t} \mathbf{P}_t + \sum_{\kappa=1}^{N-1} B_{\kappa\alpha} \mathbf{Q}_{\kappa}.$$
 (B10)

Appendix C: Coordinates on the Molecular Configuration Space

The transformation on the molecular configuration space proceeds in two steps, $(\mathbf{X}_{\alpha}, \mathbf{r}_i) \rightarrow (\mathbf{X}_{CM}, \mathbf{Y}_{\kappa}, \mathbf{r}_i) \rightarrow (\mathbf{R}_{CM}, \mathbf{Y}_{\kappa}, \mathbf{s}_i)$. The first step is explained in subsection II E and Appendix B. In the second step \mathbf{R}_{CM} , the molecular center of mass, is given by (30) and \mathbf{s}_i , the electron position relative to the nuclear center of mass, is given by (31), while \mathbf{Y}_{κ} , the nuclear Jacobi coordinates, do not change. The inverse transformation is given by

$$\mathbf{X}_{\rm CM} = \mathbf{R}_{\rm CM} - \frac{m_e}{M_{\rm mol}} \sum_{i=1}^{N_e} \mathbf{s}_i,\tag{C1}$$

and $\mathbf{r}_i = \mathbf{s}_i + \mathbf{X}_{CM}$.

The momenta conjugate to $(\mathbf{R}_{CM}, \mathbf{Y}_{\kappa}, \mathbf{s}_i)$ are $(\mathbf{P}_{CM}, \mathbf{Q}_{\kappa}, \mathbf{q}_i)$, which are functions of the momenta $(\mathbf{P}_t, \mathbf{Q}_{\kappa}, \mathbf{p}_i)$ after the first coordinate transformation. Of these \mathbf{P}_{CM} , the total molecular momentum, is given by (34) and \mathbf{Q}_{κ} does not change from the first transformation (it is given by (B9)). As for \mathbf{q}_i , it is given by

$$\mathbf{q}_i = \mathbf{p}_i - \frac{m_e}{M_{\text{mol}}} (\mathbf{P}_t + \mathbf{p}_t).$$
(C2)

The inverse transformations are

$$\mathbf{P}_t = \frac{M_t}{M_{\text{mol}}} \mathbf{P}_{\text{CM}} - \sum_{i=1}^{N_e} \mathbf{q}_i, \tag{C3}$$

and

$$\mathbf{p}_i = \frac{m_e}{M_{\rm mol}} \mathbf{P}_{\rm CM} + \mathbf{q}_i. \tag{C4}$$

Appendix D: Covariant Notation for the Kinetic Energy Metric

We write $X_{\alpha I}$ for the *I*-th component of Jacobi vector \mathbf{X}_{α} , for I = 1, 2, 3, and we write $x^{\mu} = X^{\mu} = X_{\alpha I}$, where $\mu = (\alpha I)$ and $\mu = 1, ..., 3N - 3$. The kinetic energy can be expressed in terms of the metric,

$$G_{\mu\nu} = G_{(\alpha I)(\beta J)} = M_{\alpha} \,\delta_{\alpha\beta} \,\delta_{IJ},\tag{D1}$$

that is, the kinetic energy itself is $(1/2)G_{\mu\nu}\dot{x}^{\mu}\dot{x}^{\nu}$, where here and throughout this article we use the summation convention on indices μ , ν , etc. The tensor $G_{\mu\nu}$ (with lower indices) is the covariant

metric tensor. The contravariant metric tensor (with upper indices) is the inverse of (D1),

$$G^{\mu\nu} = G^{(\alpha I)(\beta J)} = \frac{1}{M_{\alpha}} \,\delta_{\alpha\beta} \,\delta_{IJ},\tag{D2}$$

so that $G^{\mu\sigma}G_{\sigma\nu} = \delta^{\mu}_{\nu}$.

As for the momenta, we define $P_{\mu} = P_{(\alpha I)}$ as the *I*-th component of the momentum \mathbf{P}_{α} , and we define P^{μ} (with an upper index) as

$$P^{\mu} = G^{\mu\nu} P_{\nu} = \frac{P_{\alpha I}}{M_{\alpha}},\tag{D3}$$

where $\mu = (\alpha I)$, so that P^{μ} , with an upper or contravariant index, is the velocity $V^{\mu} = \dot{x}^{\mu}$. Similarly we define the derivative operators,

$$\partial_{\mu} = \frac{\partial}{\partial x^{\mu}}, \qquad \partial^{\mu} = G^{\mu\nu} \partial_{\nu},$$
 (D4)

so that as an operator, the momentum is $\hat{P}_{\mu} = -i\hbar \partial_{\mu}$. Equations (D3) and (D4) are examples of raising an index with the metric tensor.

The kinetic energy can be expressed in terms of the momenta by

$$\frac{1}{2}\sum_{\alpha=1}^{N}\frac{\mathbf{P}_{\alpha}^{2}}{M_{\alpha}} = \frac{1}{2}G^{\mu\nu}P_{\mu}P_{\nu} = \frac{1}{2}P^{\mu}P_{\mu}.$$
(D5)

Appendix E: Derivation of *G*₂

In this Appendix we use the working basis (72) and for brevity we write the resolvent $R(\varepsilon_n(x), x)$ as R(x).

The condition on G_2 , that it kill the second-order, off-block diagonal terms in H', is expressed by (92), where $H_2'^o$ is given by (89b). As in the solution for G_1 , we find that (92) is consistent with the anti-Hermiticity of G_2 and that it determines the odd (off-block-diagonal) part of G_2 , We set the even part to zero so that G_2 is purely odd. The solution for the (nk)-block, $k \neq n$, is

$$G_{2,nk} = \sum_{l \neq n} (H_2^{\prime o})_{nl} R_{lk}(\varepsilon_n, x), \qquad (k \neq n), \tag{E1}$$

which may be compared to (77). The other off-diagonal block is given by anti-Hermiticity, $G_{2,kn} = -G_{2,nk}^*$, $k \neq n$. To find G_2 , we will first find $(H_2'^o)_{nk}$ for $k \neq n$ and then apply (E1).

According to (89b) there are three terms in $(H_2'^o)_{nk}$ for $k \neq n$. The first is

$$H_{2,nk} = \frac{1}{2} (\partial^{\mu} \langle x; n |) (\partial_{\mu} | x; k \rangle), \qquad (k \neq n), \tag{E2}$$

a special case of (98c). This term represents a purely multiplicative operator, that is, it is independent of the nuclear momenta P_{μ} . Notice that it involves derivatives $\partial_{\mu}|x;k\rangle$ of the diabatic basis on \mathscr{S}^{\perp} , whose problematical aspects are discussed in Sec. III L. The derivatives $\partial^{\mu}|x;n\rangle$ of the basis vector on \mathscr{S} present no such difficulties. The latter can be expressed in terms of the basis vector $|x;n\rangle$ itself, via the Feynman-Hellman formula (101). Similar formulas can be worked out for higher derivatives of the basis vector $|x;n\rangle$ on \mathscr{S} .

For the second term of (89b) we require

$$[G_1, H_1^e]_{nk} = \sum_l G_{1,nl} H_{1,lk}^e - \sum_l H_{1,nl}^e G_{1,lk}, \qquad (k \neq n),$$
(E3)

where the first sum can be restricted to $l \neq n$ since $G_{1,nn} = 0$ and where the second sum can be restricted to l = n since H_1^e is even. In fact, the second sum vanishes, since $H_{1,nn} = 0$. As for the first sum, we use (102) and (98b), whereupon the sum can be extended to l = n and a resolution of the identity removed. The result is

$$[G_1, H_1^e]_{nk} = P^{\mu} P^{\nu}(\partial_{\mu} \langle x; n |) R(x) (\partial_{\nu} | x; k \rangle), \qquad (k \neq n),$$
(E4)

another expression that involves derivatives of $|x;k\rangle$ on \mathscr{S}^{\perp} but now the symbol of a second-order differential operator.

We see that $(H_2'^o)_{nk}$ contains terms that are independent of the momentum P^{μ} , that is, constant in the momentum, and terms that are quadratic in the momentum. In fact, it turns out that this is all it contains, so we will write $(H_2'^o)_{nk} = (H_2'^o)_{nk}^C + (H_2'^o)_{nk}^Q$ for the two parts. Then (E2) is a contribution to $(H_2'^o)_{nk}^C$ and (E4) is a contribution to $(H_2'^o)_{nk}^Q$.

For the third term of (89b) we require $T_{11,nk}$ for $k \neq n$, which by (63b) can be written

$$T_{11,nk} = \frac{i}{2} \sum_{l} \{G_{1,nl}, H_{0,lk}\} + \frac{i}{2} \sum_{l} \{G_{1,lk}, H_{0,nl}\},$$
(E5)

where we have used the antisymmetry of the Poisson bracket in the second sum. It is not hard to see that the kinetic energy part of H_0 (see (98a)) produces a contribution to $T_{11,nk}$ that is quadratic in P^{μ} while the potential energy produces one that is independent of P^{μ} , so we shall write $T_{11,nk} = T_{11,nk}^C + T_{11,nk}^Q$ for the two contributions.

The kinetic energy part of H_0 is a multiple of the identity which allows the sums in (E5) to be done, whereupon they turn out to be equal. Thus we find

$$T^{Q}_{11,nk} = i\{G_{1,nk}, (1/2)P^{\mu}P_{\mu}\} = -P^{\mu}P^{\nu}\partial_{\nu}[(\partial_{\mu}\langle x; n|)R(x)|x;k\rangle], \qquad (k \neq n),$$
(E6)

where we have used (102) and evaluated the Poisson bracket. When this is added to (E4) we obtain the quadratic part of $(H_2'^o)_{nk}$,

$$(H_2'^{o})_{nk}^Q = -P^{\mu}P^{\nu} \partial_{\nu} [(\partial_{\mu} \langle x; n |) R(x)] | x; k \rangle. \qquad (k \neq n),$$
(E7)

in which all derivatives of the basis vectors $|x;k\rangle$ on \mathscr{S}^{\perp} have disappeared. The tensor contracting against $P^{\mu}P^{\nu}$ is not symmetric in $(\mu\nu)$ but it may be replaced by its symmetric part.

As for the potential energy contribution to (E5), we find that the first sum in (E5) can be restricted to $l \neq n$ and the second to l = n, since H_0 is purely even. This gives

$$T_{11,nk}^{C} = -\frac{1}{2} \sum_{l \neq n} \{ P^{\mu}(\partial_{\mu}\langle x; n |) R(x) | x; l \rangle, W_{lk}(x) \} - \frac{1}{2} \{ P^{\mu}(\partial_{\mu}\langle x; n |) R(x) | x; k \rangle, \varepsilon_{n}(x) \}$$

$$= \frac{1}{2} \sum_{l \neq n} (\partial_{\mu}\langle x; n |) R(x) | x; l \rangle [\partial^{\mu} W_{lk}(x)] + \frac{1}{2} (\partial_{\mu}\langle x; n |) R(x) | x; k \rangle [\partial^{\mu} \varepsilon_{n}(x)]$$

$$= -\frac{1}{2} \sum_{l \neq n} (\partial_{\mu}\langle x; n |) R(x) | x; l \rangle \partial^{\mu} [\varepsilon_{n}(x) \delta_{lk} - W_{lk}(x)]$$

$$+ (\partial_{\mu}\langle x; n |) R(x) | x; k \rangle [\partial^{\mu} \varepsilon_{n}(x)], \quad (k \neq n), \qquad (E8)$$

where in the first step we have evaluated the Poisson brackets and in the second we have added and subtracted a term to make the derivative of the inverse of the resolvent appear.

Now $(H_2'^{o})_{nk}^C$ is the sum of (E2) and the two terms on the right of (E8). We expand the derivative appearing there,

$$\partial^{\mu}[\varepsilon_{n}(x) \,\delta_{lk} - W_{lk}(x)] = \partial^{\mu} \left\{ \langle x; l | [\varepsilon_{n}(x) - H_{e}(x)] | x; k \rangle \right\}$$

= $(\partial^{\mu} \langle x; l |) [\varepsilon_{n}(x) - H_{e}(x)] | x; k \rangle + \langle x; l | \left\{ \partial^{\mu} [\varepsilon_{n}(x) - H_{e}(x)] \right\} | x; k \rangle$
+ $\langle x; l | [\varepsilon_{n}(x) - H_{e}(x)] (\partial^{\mu} | x; k \rangle),$ (E9)

giving what we will call the "left," "middle" and "right" contributions to the first term on the right of (E8). The right contribution to that term is

$$-\frac{1}{2}\sum_{l\neq n}(\partial_{\mu}\langle x;n|)R(x)|x;l\rangle\langle x;l|[\varepsilon_{n}(x)-H_{e}(x)](\partial^{\mu}|x;k\rangle) = -\frac{1}{2}(\partial_{\mu}\langle x;n|)(\partial^{\mu}|x;k\rangle, \quad (E10)$$

where we have extended the *l* sum to include l = n (since $R(x)|x;n\rangle = 0$), removed the resolution of the identity, applied (79) and used $(\partial_{\mu}\langle x;n|)Q(x) = \partial_{\mu}\langle x;n|$. We see that the right contribution cancels (E2).

We simplify the middle contribution to the first term on the right in (E8) by first removing the same resolution of the identity and then applying ∂^{μ} to the identity (79). Thus the middle

contribution becomes

$$-\frac{1}{2}(\partial_{\mu}|x;n\rangle)R(x)\left\{\partial^{\mu}[\varepsilon_{n}(x)-H_{e}(x)]\right\}|x;k\rangle$$
$$=-\frac{1}{2}(\partial_{\mu}\langle x;n|)[\partial^{\mu}Q(x)]|x;k\rangle+\frac{1}{2}(\partial_{\mu}\langle x;n|)[\partial^{\mu}R(x)][\varepsilon_{n}(x)-H_{e}(x)]|x;k\rangle.$$
 (E11)

The term involving $\partial^{\mu}Q(x) = -\partial^{\mu}P(x) = -(\partial^{\mu}|x;n\rangle)\langle x;n|-|x;n\rangle(\partial^{\mu}\langle x;n|)$ vanishes since $\langle x;n|x;k\rangle = 0$ and $(\partial_{\mu}\langle x;n|)|x;n\rangle = -F_{\mu;nn} = 0$. Altogether, we now have

$$(H_{2}^{\prime o})_{nk}^{C} = \left[\partial^{\mu} \varepsilon_{n}(x)\right] \left(\partial_{\mu} \langle x; n| \right) R(x) |x; k\rangle -\frac{1}{2} \sum_{l \neq n} \left(\partial_{\mu} \langle x; n| \right) R(x) |x; l\rangle \left(\partial^{\mu} \langle x; l| \right) \left[\varepsilon_{n}(x) - H_{e}(x)\right] |x; k\rangle +\frac{1}{2} \left(\partial_{\mu} \langle x; n| \right) \left[\partial^{\mu} R(x)\right] \left[\varepsilon_{n}(x) - H_{e}(x)\right] |x; k\rangle, \qquad (k \neq n).$$
(E12)

Now we may return to (E1) and compute G_2 . It also has constant and quadratic contributions, referring to the dependence on P^{μ} . The execution of the sum in (E1) amounts to multiplying the operators seen in the matrix elements in $(H_2'^o)_{nk}$ on the right by R(x), for example, (E7) gives the quadratic contribution to G_2 as

$$(G_2^Q)_{nk} = -P^{\mu}P^{\nu} \partial_{\nu} [(\partial_{\mu} \langle x; n |) R(x)] R(x) | x; k \rangle, \qquad (k \neq n),$$
(E13)

while the first term on the right in (E12) gives the contribution $[\partial^{\mu} \varepsilon_n(x)] (\partial_{\mu} \langle x; n |) R(x)^2 | x; k \rangle$ to $(G_2^C)_{nk}$. As for the second and third terms on the right in (E12), applying the sum in (E1) effectively cancels the factor of $[\varepsilon_n(x) - H_e(x)]$, first replacing it by Q(x) which then disappears since $Q(x)|x;k\rangle = |x;k\rangle$. Thus the third term on the right in (E12) gives the contribution to $(G_2^C)_{nk}$,

$$\frac{1}{2}(\partial_{\mu}\langle x;n|)[\partial^{\mu}R(x)]|x;k\rangle.$$
(E14)

As for the second term on the right in (E12), it gives

$$-\frac{1}{2}\sum_{l\neq n}(\partial_{\mu}\langle x;n|)R(x)|x;l\rangle\,(\partial^{\mu}\langle x;l|)|x;k\rangle = \frac{1}{2}(\partial_{\mu}\langle x;n|)R(x)\,(\partial^{\mu}|x;k\rangle). \tag{E15}$$

This contribution to G_2 contains derivatives of the basis vectors $|x;k\rangle$ on \mathscr{S}^{\perp} that do not cancel. These pieces of G_2 are assembled and the final solution is displayed in (106).

REFERENCES

Abedi, A., Maitra, N. T., and Gross, E. K. U., "Exact factorization of the time-dependent electronnuclear wave function," Phys. Rev. Lett. **105**, 123002 (2010).

- Abedi, A., Maitra, N. T., and Gross, E. K. U., "Correlated electron-nuclear dynamics: Exact factorization of the molecular wave function," J. Chem. Phys. **137**, 22A530 (2012).
- Agostini, F. and Curchod, B. F. E., "When the exact factorization meets conical intersections ...," Eur. Phys. J. B **91**, 141 (2018).
- Aharonov, Y. and Stern, A., "Origin of the geometric forces accompanying Berry's geometric potentials," Phys. Rev. Lett. **69**, 3593–3597 (1992).
- Albert, J., Hader, K., and Engel, V., "Communication: On the calculation of time-dependent electron flux within the Born-Oppenheimer approximation: A flux-flux reflection principle," J. Chem. Phys. 147, 241101 (2017).
- Althorpe, S. C., "General explanation of geometric phase effects in reactive systems: Unwinding the nuclear wave function using simple topology," J. Chem. Phys. **124**, 084105 (2006).
- Althorpe, S. C., "Symmetry analysis of geometric-phase effects in quantum dynamics," in *Conical Intersections, Theory, Computation and Experiment*, edited by W. Domcke, D. R. Yarkony, and H. Köppel (World Scientific, Singapore, 2012) pp. 155–194.
- Aquilanti, V. and Cavalli, S., "Coordinates for molecular dynamics: Orthogonal local systems," J. Chem. Phys. **85**, 1355–1361 (1986).
- Balazs, N. L. and Jennings, B. K., "Wigner's function and other distribution functions in mock phase spaces," Phys. Rep. **104**, 347–391 (1984).
- Barth, I., Hege, H.-C., Ikeda, H., Kenfack, A., Koppitz, M., Manz, J., Marquardt, F., and Paramonov, G. K., "Concerted quantum effects of electronic and nuclear fluxes in molecules," Chem. Phys. Lett. 481, 118–123 (2009).
- Bian, X., Tao, Z., Wu, Y., Rawlinson, J., Littlejohn, R. G., and Subotnik, J. E., "Total angular momentum conservation in *ab initio* Born-Oppenheimer molecular dynamics," (2023), arXiv:2303.13787v1 [physics.chem-ph].
- Bian, X., Wu, Y., Teh, H.-H., Zhou, Z., Chen, H.-T., and Subotnik, J. E., "Modeling nonadiabatic dynamics with degenerate electronic states, intersystem crossing, and spin separation: A key goal for chemical physics," J. Chem. Phys. 154, 110901 (2021).
- Bjorken, J. D. and Drell, S. D., Relativistic Quantum Mechanics (McGraw-Hill, New York, 1964).
- Blount, E. I., "Bloch electrons in a magnetic field," Phys. Rev. 126, 1636–1653 (1962a).
- Blount, E. I., "Extension of the Foldy-Wouthuysen transformation," Phys. Rev. **128**, 2454–2458 (1962b).
- Born, M. and Huang, K., Dynamical Theory of Crystal Lattices (Oxford University Press, New

York, 1954).

- Born, M. and Oppenheimer, R., "Zur Quantentheorie der Molekeln," Ann. Phys. (Leipzig) **84**, 457 (1927).
- Bredtmann, T., Diestler, D. J., Li, S.-D., Manz, J., Pérez-Torres, J. F., Tian, W.-J., Wu, Y.-B., Yang,
 Y., and Zhai, H.-J., "Quantum theory of concerted electronic and nuclear fluxes associated with adiabatic intramolecular processes," Phys. Chem. Chem. Phys. 17, 29421–29464 (2015).
- Buckingham, A. D., Fowler, P. W., and Galwas, P. A., "Velocity-dependent property surfaces and the theory of vibrational circular dichroism," Chemical Physics **112**, 1–14 (1987).
- Cederbaum, L. S., "Born-Oppenheimer approximation and beyond," in *Conical Intersections: Electronic Structure, Dynamics and Spectroscopy*, edited by W. Domcke, D. R. Yarkony, and H. Köppel (World Scientific, Singapore, 2004) p. 3.
- Cederbaum, L. S., "The exact molecular wave function as a product of an electronic and a nuclear wavefunction," J. Chem. Phys. **138**, 224110 (2013).
- Cederbaum, L. S., "Erratum: "The exact molecular wavefunction as a product of an electronic and a nuclear wavefunction" [J. Chem. Phys. 138, 224110(2013)]," J. Chem. Phys. 141, 029902 (2014).
- Cederbaum, L. S., "The exact wavefunction of interacting *N* degrees of freedom as a product of *N* single-degree-of-freedom wavefunctions," Chem. Phys. **457**, 129–132 (2015).
- Culpitt, T., Peters, L. D. M., Tellgren, E. I., and Helgaker, T., "Analytic calculation of the Berry curvature and diagonal Born-Oppenheimer correction for molecular systems in uniforms magnetic fields," J. Chem. Phys. 156, 044121 (2022).
- Curchod, B. F. E. and Agostini, F., "On the dynamics through a conical intersection," J. Phys. Chem. Lett. **8**, 831–837 (2017).
- Delos, J. B., "Theory of electronic transitions in slow atomic collisions," Rev. Mod. Phys. **53**, 287 (1981).
- Delves, L. M., "Tertial and general-order collisions II," Nucl. Phys. 20, 275 (1960).
- Diestler, D. J., "Coupled-channels quantum theory of electronic flux density in electronically adiabatic processes: Fundamentals," J. Phys. Chem. A **116**, 2728–2735 (2012a).
- Diestler, D. J., "Quasi-classical theory of electronic flux density in electronically adiabatic molecular processes," J. Phys. Chem. A **116**, 11161–11166 (2012b).
- Diestler, D. J., "Beyond the Born-Oppenheimer approximation: A treatment of electronic flux density in electronically adiabatic molecular processes," J. Phys. Chem. A **117**, 4698–4708 (2013).

- Diestler, D. J., Kenfack, A., Manz, J., Paulus, B., Pérez-Torres, J. F., and Pohl, V., "Computation of the electronic flux density in the Born-Oppenheimer approximation," J. Phys. Chem. A **117**, 8519–8527 (2013).
- Emmrich, C. and Weinstein, A., "Geometry of the transport equation in multicomponent WKB approximations," Comm. Math. Phys. **176**, 701–711 (1996).
- Fatehi, S., Alguire, E., Shao, Y., and Subotnik, J. E., "Analytic derivative couplings between configuration-interaction-singles states with built-in electron-translation factors for translational invariance," J. Chem. Phys. 135, 234105 (2011).
- Foldy, L. L. and Wouthuysen, S. A., "On the Dirac theory of spin 1/2 particles and its nonrelativistic limit," Phys. Rev. **78**, 29–36 (1950).
- Freedman, T. B. and Nafie, L. A., "Molecular orbital calculations of infrared intensities beyond the Born-Oppenheimer approximation using the dipole moment operator," Chem. Phys. Lett. 126, 441–446 (1986).
- Gatti, F., Iung, C., Menou, M., Justum, Y., Nauts, A., and Chapuisat, X., "Vector parameterization of the *N*-atom problem in quantum mechanics, I. Jacobi vectors," J. Chem. Phys. **108**, 8804–8820 (1998).
- Gauss, J., Tajti, A., Kállay, M., Stanton, J. F., and Szalay, P. G., "Analytic calculation of the diagonal Born-Oppenheimer correction within configuration-interaction and coupled-cluster theory," J. Chem. Phys. 125, 144111 (2006).
- Gherib, R., Ye, L., Ryabinkin, I. G., and Izmaylov, A. F., "On the inclusion of the diagonal Born-Oppenheimer correction in surface hopping methods," J. Chem. Phys. **144**, 154103 (2016).
- Goldhaber, A. S., "Newtonian adiabatics unified," Phys. Rev. A 71, 062102 (2005).
- Gordon, M. S., Glezaku, V.-A., and Yarkony, D. R., "Systematic location of intersecting seams of conical intersection in triatomic molecules: The $1^2A'-2^2A'$ conical intersection in BH₂," J. Chem. Phys. **108**, 5657 (1998).
- Groenewold, H. J., "On the principles of elementary quantum mechanics," Physica **12**, 405–460 (1946).
- Hirschfelder, J. O., "Coordinates which diagonalize the kinetic energy of relative motion," Int. J. Quantum Chem. **3**, 17–31 (1969).
- Jecko, T., "On the mathematical treatment of the Born-Oppenheimer approximation," J. Math. Phys. **55**, 053504 (2014).
- Jecko, T., Sutcliffe, B. T., and Woolley, R. G., "On factorization of molecular wavefunctions," J.

Phys. A 48, 445201 (2015).

- Juanes-Marcos, J. C., Althorpe, S. C., and Wrede, E., "Theoretical study of geometric phase effects in the hydrogen-exchange reaction," Science **309**, 1227–1230 (2005).
- Kendrick, B. K., "Non-adiabatic quantum reactive scattering in hyperspherical coordinates," J. Chem. Phys. **148**, 044116 (2018).
- Kutzelnigg, W., "Which masses are vibrating or rotating in a molecule?" Mol. Phys. **105**, 2627–2647 (2007).
- Landau, L. D., "Zur Theorie der Energieübertragung bei Stössen," Phys. Z. Sowjetunion 1, 88 (1932).
- Littlejohn, R., Rawlinson, J., and Subotnik, J., "The parallel-transported (quasi)-diabatic basis," J. Chem. Phys. **157**, 184303 (2022).
- Littlejohn, R., Rawlinson, J., and Subotnik, J., "Representation and conservation of angular momentum in the Born-Oppenheimer theory of polyatomic molecules," J. Chem. Phys. 158, 104302 (2023).
- Littlejohn, R. G., "The semiclassical evolution of wave packets," Phys. Rep. 138, 193–291 (1986).
- Littlejohn, R. G. and Flynn, W. G., "Geometric phases and the Bohr-Sommerfeld quantization of multicomponent wave fields," Phys. Rev. Lett. **66**, 2839–2842 (1991a).
- Littlejohn, R. G. and Flynn, W. G., "Geometric phases in the asymptotic theory of coupled wave equations," Phys. Rev. A 44, 5239 (1991b).
- Littlejohn, R. G. and Flynn, W. G., "Phase integral theory, coupled wave equations, and mode conversion," Chaos **2**, 149 (1992).
- Littlejohn, R. G. and Flynn, W. G., "General linear mode conversion coefficient in one dimension," Phys. Rev. Lett. **70**, 1799–1802 (1993).
- Littlejohn, R. G. and Reinsch, M., "Internal or shape coordinates in the *n*-body problem," Phys. Rev. A **52**, 4035 (1995).
- Littlejohn, R. G. and Reinsch, M., "Gauge fields in the separation of rotations and internal motions in the *n*-body problem," Rev. Mod. Phys. **69**, 213 (1997).
- Littlejohn, R. G. and Weigert, S., "Adiabatic motion of a neutral spinning particle in an inhomogeneous magnetic field," Phys. Rev. A **48**, 924–940 (1993).
- Lorin, E., "Numerical analysis of the exact factorization of molecular time-dependent Schrödinger wave functions," Commun. Nonlinear Sci. Numer. Simulat. **95**, 105627 (2021).
- Mannouch, J. R. and Richardson, J. O., "A mapping approach to surface hopping," J. Chem. Phys.

158, 104111 (2023).

- Martinazzo, R. and Burghardt, I., "Quantum hydrodynamics of coupled electron-nuclear systems," (2023), arXiv:2310.08766v1 [physics.chem-ph].
- Matsunaga, N. and Yarkony, D. R., "Energies and derivative couplings in the vicinity of a conical intersection 3. The 'most' diabatic basis," Mol. Phys. **93**, 79–84 (1998).
- Mátyus, E., "Non-adiabatic mass correction to the rovibrational states of molecules: Numerical application for the H₂⁺ molecular ion," J. Chem. Phys. **149**, 194111 (2018).
- Mátyus, E. and Teufel, S., "Effective non-adiabatic Hamiltonians for the quantum nuclear motion over coupled electronic states," J. Chem. Phys. **151**, 014113 (2019).
- McDonald, S. W., "Phase-space representations of wave equations with applications to the eikonal approximation for short-wavelength waves," Phys. Rep. **158**, 337–416 (1988).
- Mead, C. A., "Born-Oppenheimer expansion at constant energy," J. Chem. Phys. **125**, 204109 (2006).
- Mead, C. A. and Moscowitz, A., "Dipole length versus dipole velocity in the calculation of infrared intensities with Born-Oppenheimer wave functions," Int. J. Quantum Chem. **1**, 243 (1967).
- Meek, G. A. and Levine, B. G., "Wave function continuity and the diagonal Born-Oppenheimer correction at conical intersections," J. Chem. Phys. **144**, 184109 (2016).
- Moody, J., Shapere, A., and Wilczek, F., "Adiabatic effective Lagrangians," in *Geometric phases in physics*, edited by A. Shapere and F. Wilczek (World Scientific, Singapore, 1989) p. 160.
- Moyal, J. E., "Quantum mechanics as a statistical theory," Proc. Cambridge Phil. Soc. **45**, 99–124 (1949).
- Nafie, L. A., "Adiabatic molecular properties beyond the Born-Oppenheimer approximation. Complete adiabatic wave functions and vibrationally induced electronic current density," J. Chem. Phys. 79, 4950–4957 (1983).
- Nafie, L. A., "Velocity-gauge formalism in the theory of vibrational circular dichroism and infrared absorption," J. Chem. Phys. 96, 5687–5702 (1992).
- Nafie, L. A., "Electron transition current density in molecules. I. Non-Born-Oppenheimer theory of vibronic and vibrational transitions," J. Phys. Chem. A **101**, 7826–7833 (1997).
- Nafie, L. A., "Vibrational optical activity: From discovery and development to future challenges," Chirality **32**, 667–692 (2020).
- Nikitin, E. E. and Umanskii, S. Y., *Theory of Slow Atomic Collisions* (Springer-Verlag, Berlin, 1984).

- Okuyama, M. and Takatsuka, K., "Electron flux in molecules induced by nuclear motion," Chem. Phys. Lett. **476**, 109–115 (2009).
- Osborn, T. A. and Molzahn, F. H., "Moyal quantum mechanics: The semiclassical Heisenberg dynamics," Ann. Phys. **141**, 79–127 (1995).
- Pachucki, K. and Komasa, J., "Nonadiabatic corrections to wave function and energy," J. Chem. Phys. **129**, 034102 (2008).
- Panati, G., Spohn, H., and Teufel, S., "Space-adiabatic perturbation theory in quantum dynamics," Phys. Rev. Lett. **88**, 250405 (2002).
- Parashar, S., Sajeev, Y., and Ghosh, S. K., "Towards an exact factorization of the molecular wave function," Mol. Phys. 113, 3067–3072 (2015).
- Patchkovskii, S., "Electronic currents and Born-Oppenheimer molecular dynamics," J. Chem. Phys. **137**, 084109 (2012).
- Pohl, V. and Tremblay, C., "Adiabatic electron flux density: A Born-Oppenheimer broken symmetry ansatz," Phys. Rev. A **93**, 012504 (2016).
- Requist, R., "Adiabatic perturbation theory for two-component systems with one heavy component," (2023), arXiv:2310.09189v1 [quant-ph].
- Requist, R. and Gross, E. K. U., "Exact factorization-based density functional theory of electrons and nuclei," Phys. Rev. Lett. **117**, 193001 (2016).
- Requist, R., Proetto, C. R., and Gross, E. K. U., "Asymptotic analysis of Berry curvature in the $E \otimes e$ Jahn-Teller model," Phys. Rev. A **96**, 062503 (2017).
- Requist, R., Tandetzky, F., and Gross, E. K. U., "Molecular geometric phase from the exact electron-nuclear factorization," Phys. Rev. A **93**, 042108 (2016).
- Runeson, J. E. and Manolopoulos, D. E., "A multi-state mapping approach to surface hopping," (2023), arXiv:2305.08835v2 [physics.chem-ph].
- Schaupp, T., Albert, J., and Engel, V., "Time-dependent elecron momenta from Born-Oppenheimer calculations," Eur. Phys. J. B **91**, 97 (2018).
- Schaupp, T. and Engel, V., "On the calculation of time-dependent electron momenta within the Born-Oppenheimer approximation," J. Chem. Phys. **150**, 164110 (2019).
- Schaupp, T. and Engel, V., "Born-Oppenheimer and non-Born-Oppenheimer contributions to timedependent electron momenta," J. Chem. Phys. **152**, 204310 (2020).
- Schaupp, T. and Engel, V., "Correlated three-dimensional electron-nuclear motion: Adiabatic dynamics versus passage of conical intersection," J. Chem. Phys. **156**, 074302 (2022a).

- Schaupp, T. and Engel, V., "Quantum flux densities for electronic-nuclear motion: Exact versus Born-Oppenheimer dynamics," Phil. Trans. R. Soc. A **380**, 20200385 (2022b).
- Schaupp, T., Renziehausen, K., Barth, I., and Engel, V., "Time-dependent momentum expectation values from different quantum probability and flux densities," J. Chem. Phys. **154**, 064307 (2021).
- Scherrer, A., Agostini, F., Sabastiani, D., Gross, E. K. U., and Vuilleumier, R., "Nuclear velocity perturbation theory for vibrational circular dichroism: An approach based on the exact factorization of the electron-nuclear wave function," J. Chem. Phys. **143**, 074106 (2015).
- Scherrer, A., Vuilleumier, R., and Sabastiani, D., "Nuclear velocity perturbation theory of vibrational circular dichroism," J. Chem. Theory Comput. **9**, 5305–5312 (2013).
- Schild, A., Agostini, F., and Gross, E. K. U., "Electronic flux density beyond the Born-Oppenheimer approximation," J. Phys. Chem. A **120**, 3316–3325 (2016).
- Stephens, P. J., "Theory of vibrational circular dichroism," J. Phys. Chem. 89, 748–752 (1985).
- Stephens, P. J. and Lowe, M. A., "Vibrational circular dichroism," Ann. Rev. Phys. Chem. 36, 213–241 (1985).
- Stückelberg, E. C. G., "Theorie der unelastischen Stösse zwischen Atomen," Helv. Phys. Acta 5, 369–422 (1932).
- Sutcliffe, B., "The decoupling of electronic and nuclear motions in the isolated molecule Schrödinger Hamiltonian," Adv. Chem. Phys. **114**, 1–121 (2000).
- Sutcliffe, B., "Some difficulties in considering rotation motion within the Born-Oppenheimer approximation for polyatomic molecules," Int. J. Quantum Chem. **112**, 2894–2903 (2012).
- Teufel, S., Adiabatic Perturbation Theory in Quantum Dynamics (Springer-Verlag, Berlin, 2003).
- Valeev, E. F. and Sherrill, C. D., "The diagonal Born-Oppenheimer correction beyond the Hartree-Fock approximation," J. Chem. Phys. **118**, 3921–3927 (2003).
- Wang, X.-G. and Tucker Carrington, J., "A simple method for deriving kinetic energy operators,"J. Chem. Phys. 113, 7097 (2000).
- Weigert, S. and Littlejohn, R. G., "Diagonalization of multicomponent wave equations with a Born-Oppenheimer example," Phys. Rev. A **47**, 3506 (1993).
- Weyl, H., "Quantenmechanik und Gruppentheorie," Z. Phys. 46, 1–46 (1927).
- Wigner, E. P., "On the quantum correction for thermodynamic equilibrium," Phys. Rev. **40**, 749–759 (1932).
- Yarkony, D. R., "Diabolical conical intersections," Rev. Mod. Phys. 68, 985 (1996).

- Yarkony, D. R., "A criterion for the confluence of two seams of conical intersection in triatomic molecules," Theor. Chem. Acc. 98, 197 (1997a).
- Yarkony, D. R., "Energies and derivative couplings in the vicinity of a conical intersection using degenerate perturbation theory and analytic gradient techniques. 1," J. Phys. Chem. A 101, 4263 (1997b).
- Yarkony, D. R., "Conical intersections: The new conventional wisdom," J. Phys. Chem. A **105**, 6277 (2001).
- Yarkony, D. R., "Conical intersections: Their description and consequences," in *Conical Intersections: Electronic Structure, Dynamics and Spectroscopy*, edited by W. Domcke, D. R. Yarkony, and H. Köppel (World Scientific, Singapore, 2004) p. 41.
- Yarkony, D. R., "Determination of potential energy surface intersections and derviative couplings in the adiabatic representation," in *Conical Intersections: Electronic Structure, Dynamics and Spectroscopy*, edited by W. Domcke, D. R. Yarkony, and H. Köppel (World Scientific, Singapore, 2004) p. 129.
- Zener, C., "Non-adiabatic crossing of energy levels," Proc. Royal Soc. A 137, 696–702 (1932).