

Masses and Magnetic Moments of Doubly Heavy Tetraquarks via Diffusion Monte Carlo Method

Halil Mutuk^{1,*}

¹*Department of Physics, Faculty of Science, Ondokuz Mayıs University, 55200 Samsun, Türkiye*

We present mass spectrum and magnetic moments of the $\bar{n}\bar{n}QQ$ states, where $n = u, d, s$ and $Q = c, b$. We solve four-body Schrödinger equation with a quark potential model by using diffusion Monte Carlo (DMC) method. The quark potential is based on the Coulomb, confinement and spin-spin interaction terms. We find mass and magnetic moment of the T_{cc}^+ state as $M_{T_{cc}^+} = 3892$ MeV and $\mu = 0.28\mu_N$, respectively. We also find the mass and magnetic moment of T_{bb}^- as $M_{T_{bb}^-} = 10338$ MeV and $\mu = -0.32\mu_N$, respectively. We find some bound state candidates of doubly heavy tetraquark systems with $I(J^P) = 0(1)^+ nn\bar{b}\bar{b}$, $I(J^P) = 0(0)^+ nn\bar{c}\bar{b}$, $I(J^P) = 0(1)^+ nn\bar{c}\bar{b}$, and $I(J^P) = 1/2(1)^+ ns\bar{b}\bar{b}$. We compare our results with other approaches in the literature.

I. PROLOGUE

After the breakthrough works of Gell-Mann [1] and Zweig [2] in 1964, hadrons can be classified into two groups in terms of their valence quarks and antiquarks: quark-antiquark ($q\bar{q}$) configuration as mesons and three-quark configuration (qqq) as baryons. This classification was called as *quark model* and reproduced reasonably well many hadron observables such as baryon and meson spectrum, strong decays, magnetic moments of baryons, etc. However, the Quantum Chromodynamics (QCD) allows exotic hadrons beyond the scope of quark model. The existence of multi-quark states is consistent with QCD. In fact, in Refs. [1, 2], multi-quark states beyond conventional ones (mesons and baryons) were conjectured as combinations of quarks. The era of conventional hadrons directed in a different route when the observation of $\chi_{c1}(3872)$ by the Belle Collaboration [3] announced in 2003. This state has a narrow width, decays into $J/\psi 2\pi$ and $J/\psi 3\pi$ and does not fall into the traditional charmonium spectra. Many unconventional states have been reported by Belle, BaBar, CLEO, BESIII, CDF, D0, LHCb, ATLAS, and CMS collaborations since 2003. These unconventional charmonium and bottomonium states are named as XYZ mesons. For a recent naming convention of these states, see Ref. [4]. These unconventional states mostly decay into final states which include ($c\bar{c}$) and ($b\bar{b}$) pairs. At this point, heavy quarkonium physics becomes important for elucidating these states. The current status and prospects of the heavy quarkonium physics can be found in Refs. [5–10].

The recent discovery of T_{cc}^+ tetraquark is of particular importance due to its quark content [11, 12]. The quark configuration is $c\bar{c}u\bar{d}$ and its mass is a few MeV below the lowest two-meson threshold of DD^* . Since the lowest comparable decay threshold consists of heavy-light meson pairings rather than a threshold consisting of a light meson and quarkonium, such $QQ\bar{q}\bar{q}$ multi-quark states are flavor unusual. Exotic hadrons with two heavy quarks, $QQ\bar{q}\bar{q}$, were in the to-do list starting from 1980s [13–18]. Especially during the last decade, doubly heavy tetraquarks were studied by means of quark level models [19–62], chromomagnetic interaction models [63–71], QCD sum rule [72–93], lattice QCD simulations [94–113], and so on. $QQ\bar{q}\bar{q}$ tetraquarks should be stable when the quark masses are sufficiently large. In other words, when the ratio of m_{QQ}/m_{qq} is large in the four-quark state, then it is natural to consider the stability of this multi-quark state [18, 114]. In this limit two heavy quarks (antiquarks) form an anticolor (color) triplet due to the Coulomb term of the potential at small QQ ($\bar{Q}\bar{Q}$) separations. Then the four-quark structure is quite similar to $\bar{Q}\bar{q}\bar{q}$ (Qqq) baryons. The same logic applies to the ($\bar{Q}\bar{Q}q$) baryons which are related to heavy-light mesons $Q\bar{q}$ [115–118]. Furthermore, heavy quark symmetry (HQS) also implies stable $QQ\bar{q}\bar{q}$ tetraquark states [36].

Hadron spectroscopy constitutes basis to test predictions of QCD. Hadron spectroscopy is related to masses and decays of the hadrons. The abundance and clear signals in the charmonium and bottomonium states prompt hadron spectroscopy. Charmonium and bottomonium states are heavy quarkonium states. Heavy quarkonium is a good laboratory for QCD: at high energies (small distances) where a perturbative calculations are possible and at low energies (large distances) nonperturbative effects take place [119]. Since charm and bottom quarks are heavy, nonrelativistic approach of QCD can be used for these mesons.

There are various methods to study physical properties of exotic states. QCD motivated phenomenological models can be used to classify and specify the components (coloured constituents) and forces between them in order to study these exotic states. In that line, many important problems related to these states can be addressed by solving Schrödinger equation. Since these exotic states include at least more than three quarks, solving many-body Schrödinger equation is quite challenging. In general four-quark states are often taken as diquark-antidiquark states or meson-meson molecules. Such quark clusterings yield a two-body Schrödinger equation in the context of potential models. In this present work we rely on a different method for solving four-quark Schrödinger equation. In our model, none of the valence quarks will be assumed to be clustered together.

* hmutuk@omu.edu.tr

The interaction in the context of potential model for this system is the widely accepted and used one: Coulomb + linear confining + spin-spin term. We solve four-quark Schrödinger equation by using Diffusion Monte Carlo (DMC) method which takes into account full correlations between the constituents of the system and is capable of producing exactly the lowest eigenenergy since the effect of trial wave function is diminished by the method.

The order of the paper is as follows: We describe the potential model, numerical approach in Sec. II. We present and explain the mass spectrum and magnetic moments of the doubly heavy tetraquark states in Sec. III. The conclusion is reserved for Sec. IV.

II. THEORETICAL FRAMEWORK

A. Potential Model

Phenomenological models with a simple relativistic kinetic energy term and a scalar potential term which incorporates the so called linear confinement, and a term related to short distance which incorporates color-Coulomb interaction stemmed from QCD, give good results and descriptions of the observed spectra of both heavy and light quark mesons and baryons.

The Hamiltonian which describes doubly heavy tetraquark states reads as follows [120]

$$H = \sum_i \left(m_i + \frac{\mathbf{p}_i^2}{2m_i} \right) - \frac{3}{16} \sum_{i < j} \tilde{\lambda}_i \tilde{\lambda}_j v_{ij}(r_{ij}) \quad (1)$$

with the potential

$$v_{ij}(r) = -\frac{\kappa(1 - e^{-\frac{r}{r_c}})}{r} + \lambda r^p + \Lambda \quad (2)$$

$$+ \frac{2\pi}{3m_i m_j} \kappa' (1 - e^{-\frac{r}{r_c}}) \frac{e^{-\frac{r^2}{r_0^2}}}{\pi^{3/2} r_0^3} \sigma_i \sigma_j, \quad (3)$$

where $r_0(m_i, m_j) = A \left(\frac{2m_i m_j}{m_i + m_j} \right)^{-B}$, A and B are constant parameters, κ and κ' are parameters, r_{ij} is the interquark distance $|\mathbf{r}_i - \mathbf{r}_j|$, σ_i are the Pauli matrices and $\tilde{\lambda}_i$ are Gell-Mann matrices. κ and κ' are related to the hyperfine interactions, just like the replacement of $\delta(\vec{r})$ with a smearing function in the Breit-Fermi approximation. There are four potentials referred to the p and r_c :

$$\begin{aligned} \text{AL1} &\rightarrow p = 1, r_c = 0, \\ \text{AP1} &\rightarrow p = 2/3, r_c = 0, \\ \text{AL2} &\rightarrow p = 1, r_c \neq 0, \\ \text{AP2} &\rightarrow p = 2/3, r_c \neq 0. \end{aligned}$$

This potential was developed under the nonrelativistic quark model (NRQM) and used for exploratory studies. It compose of ‘‘Coulomb + linear’’ or ‘‘Coulomb + 2/3-power’’ term and a strong but smooth hyperfine term. As mentioned in Ref. [120], potentials with $r_c \neq 0$ give results slightly better than those with $r_c = 0$, where $r_c = 0$ means $\kappa(r)$ and $\kappa(r)'$ are constant parameters. Therefore we choose AL2 potential with the following parameter listed in Table I:

TABLE I. Parameters of the AL2 potential

	AL2
$m_u = m_d$	0.320 GeV
m_s	0.587 GeV
m_c	1.851 GeV
m_b	5.231 GeV
κ	0.5871
κ'	1.8475
λ	0.1673 GeV ²
Λ	-0.8182 GeV
B	0.2132
A	1.6560 GeV ^{B-1}
r_c	0.1844 GeV ⁻¹

B. Diffusion Monte Carlo Method

Many important aspects of microscopic phenomena can be provided by the Schrödinger equation. Unfortunately, there are very few instances in which the Schrödinger equation can be solved exactly. For the rest which include more realistic cases, one needs to use approximations and numerical descriptions. Diffusion Monte Carlo (DMC) technique may help on this endeavour.

The imaginary-time Schrödinger equation (in $\hbar = c = 1$) can be written as

$$-\frac{\partial \Psi_{\alpha'}(\mathbf{R}, t)}{\partial t} = (H_{\alpha'\alpha} - E_R) \Psi_{\alpha}(\mathbf{R}, t), \quad (4)$$

with the Hamiltonian defined as

$$H_{\alpha'\alpha} = -\frac{\nabla_{\mathbf{R}}^2}{2m} \delta_{\alpha'\alpha} + V_{\alpha'\alpha}(\mathbf{R}). \quad (5)$$

In the above equations, $\mathbf{R} \equiv (\vec{r}_1, \dots, \vec{r}_n)$ represents the positions of n particles, E_R is the shifted energy, and α denotes each possible spin-color channel of the system. The wave function in Eq. (4) can be expanded as

$$\Psi_{\alpha}(\mathbf{R}, t) = \sum_i c_{i,\alpha} \Phi_{i,\alpha}(\mathbf{R}) e^{-[E_i - E_R]t}, \quad (6)$$

where E_i are the eigenvalues and $\Phi_{i,\alpha}(\mathbf{R})$ are the eigenfunctions. If E_R is taken close to the ground state energy E_0 , i.e. $E_R \simeq E_0$, the wave function $\Psi_{\alpha}(\mathbf{R}, t)$ will approach to $\Phi_{0,\alpha}(\mathbf{R})$ when sufficiently long time elapses, i.e. $t \rightarrow \infty$.

An important feature of the quantum monte carlo (QMC) method is to use sampling technique to reduce statistical fluctuation [121]. In this approach, we use the function

$$f_{\alpha}(\mathbf{R}, t) \equiv \psi(\mathbf{R}) \Psi_{\alpha}(\mathbf{R}, t), \quad (7)$$

where $\psi(\mathbf{R})$ is the time-independent trial function which in principle should be chosen as close to $\Phi_{0,\alpha}(\mathbf{R})$ as possible. Since $\psi(\mathbf{R})$ is a trial function, it should be chosen according to the sake of the calculations. In this present work, we formulate it as [122]

$$\psi(\mathbf{R}) = \prod_{i < j} \Phi(\vec{r}_{ij}) = \prod_{i < j} e^{-a_{ij} r_{ij}}, \quad (8)$$

where the values of the adjustable constants a_{ij} are set to minimize fluctuation.

With this new definition of the trial function (Eq. 7), the Schrödinger equation turns out to be

$$\begin{aligned} -\frac{\partial f_{\alpha'}(\mathbf{R}, t)}{\partial t} &= -\frac{1}{2m} \nabla_{\mathbf{R}}^2 f_{\alpha'}(\mathbf{R}, t) + \frac{1}{2m} \nabla_{\mathbf{R}} [F(\mathbf{R}) f_{\alpha'}(\mathbf{R}, t)] \\ &+ [E_L(\mathbf{R}) - E_s] f_{\alpha'}(\mathbf{R}, t) \\ &+ V_{\alpha'\alpha}(\mathbf{R}) f_{\alpha}(\mathbf{R}, t), \end{aligned} \quad (9)$$

where $E_L(\mathbf{R}) = \psi(\mathbf{R})^{-1} H_0 \psi(\mathbf{R})$ is the local energy and $F(\mathbf{R}) = 2\psi(\mathbf{R})^{-1} \nabla_{\mathbf{R}} \psi(\mathbf{R})$ is the drift force. The solution to Eq. (9) in path integral formalism reads as

$$\begin{aligned} f_{\alpha'}(\mathbf{R}', t + \Delta t) &= \int d\mathbf{R}_1 d\mathbf{R}_2 d\mathbf{R}_3 d\mathbf{R}_4 d\mathbf{R} \\ &\times \left[G^{(3)}\left(\mathbf{R}', \mathbf{R}_1, \frac{\Delta t}{2}\right) G^{(2)}\left(\mathbf{R}_1, \mathbf{R}_2, \frac{\Delta t}{2}\right) \right. \\ &\times G^{(1)}\left(\mathbf{R}_2, \mathbf{R}_3, \Delta t\right) \\ &\times \left. G^{(2)}\left(\mathbf{R}_3, \mathbf{R}_4, \frac{\Delta t}{2}\right) G^{(3)}\left(\mathbf{R}_4, \mathbf{R}, \frac{\Delta t}{2}\right) \right] \\ &\times \sum_{\alpha} e^{-V_{\alpha'\alpha}(\mathbf{R})\Delta t} f_{\alpha}(\mathbf{R}, t), \end{aligned} \quad (10)$$

with

$$G^{(1)}(\mathbf{R}', \mathbf{R}, t) = \left(\frac{2\pi t}{m}\right)^{-\frac{3n}{2}} e^{-\frac{m(\mathbf{R}' - \mathbf{R})^2}{2t}}, \quad (11)$$

$$G^{(2)}(\mathbf{R}', \mathbf{R}, t) = \delta(\mathbf{R}' - \mathbf{R}(t)), \text{ where } \begin{cases} \mathbf{R}(0) = \mathbf{R}, \\ \frac{d(t)}{dt} = \frac{F(\mathbf{R}(t))}{2m}, \end{cases} \quad (12)$$

$$G^{(3)}(\mathbf{R}', \mathbf{R}, t) = \exp[-(E_L(\mathbf{R}) - E_s)t] \delta(\mathbf{R}' - \mathbf{R}). \quad (13)$$

Note herein that in the case of having more than one α -channel, we propagate the quantity [123]

$$\mathcal{F}(\mathbf{R}, t) = \sum_{\alpha} f_{\alpha}(\mathbf{R}, t), \quad (14)$$

such as

$$\begin{aligned} \mathcal{F}(\mathbf{R}', t + \Delta t) &= \int d\mathbf{R} \langle \mathbf{R}' | e^{-A\Delta t} | \mathbf{R} \rangle \sum_{\alpha'\alpha} e^{-V_{\alpha'\alpha}(\mathbf{R})\Delta t} f_{\alpha}(\mathbf{R}, t) \\ &= \int d\mathbf{R} \langle \mathbf{R}' | e^{-A\Delta t} | \mathbf{R} \rangle \omega(\mathbf{R}, t) \mathcal{F}(\mathbf{R}, t), \end{aligned} \quad (15)$$

where the weight factor is introduced as

$$\omega(\mathbf{R}, t) = \frac{\sum_{\alpha'\alpha} e^{-V_{\alpha'\alpha}(\mathbf{R})\Delta t} f_{\alpha}(\mathbf{R}, t)}{\sum_{\alpha} f_{\alpha}(\mathbf{R}, t)}. \quad (16)$$

The application of DMC method to hadron physics has been scarce. For an introduction of the DMC method see Ref. [124] and for applications to hadron physics see Refs. [122, 125–130].

C. Wave function

The total wave function of a tetraquark state can written as

$$\Psi = |space\rangle \otimes |spin\rangle \otimes |color\rangle \otimes |flavor\rangle. \quad (17)$$

by taking into account possible color, spin and flavor configurations. The construction of the total wave function needs elaboration of Pauli principle. The possible spin wave functions for the $(q_1 q_2 \bar{q}_3 \bar{q}_4)$ tetraquark states with the $(qq\bar{Q}\bar{Q})$ configuration are

$$\begin{aligned}
\chi_1 &= |(q_1 q_2)_1 (\bar{Q}_3 \bar{Q}_4)_1 \rangle_2, \\
\chi_2 &= |(q_1 q_2)_1 (\bar{Q}_3 \bar{Q}_4)_1 \rangle_1, \\
\chi_3 &= |(q_1 q_2)_1 (\bar{Q}_3 \bar{Q}_4)_1 \rangle_0, \\
\chi_4 &= |(q_1 q_2)_1 (\bar{Q}_3 \bar{Q}_4)_0 \rangle_1, \\
\chi_5 &= |(q_1 q_2)_0 (\bar{Q}_3 \bar{Q}_4)_1 \rangle_1, \\
\chi_6 &= |(q_1 q_2)_0 (\bar{Q}_3 \bar{Q}_4)_0 \rangle_0,
\end{aligned} \tag{18}$$

with the spin alignments

$$\begin{aligned}
\chi_1 &= \uparrow\uparrow\uparrow\uparrow, \\
\chi_2 &= \frac{1}{2} (\uparrow\uparrow\uparrow\downarrow + \uparrow\uparrow\downarrow\uparrow - \uparrow\downarrow\uparrow\uparrow - \downarrow\uparrow\uparrow\uparrow), \\
\chi_3 &= \frac{1}{\sqrt{3}} (\uparrow\uparrow\downarrow\downarrow + \downarrow\downarrow\uparrow\uparrow) \\
&\quad - \frac{1}{2\sqrt{3}} (\uparrow\downarrow\uparrow\downarrow + \uparrow\downarrow\downarrow\uparrow + \downarrow\uparrow\uparrow\downarrow + \downarrow\uparrow\downarrow\uparrow) \\
\chi_4 &= \frac{1}{\sqrt{2}} (\uparrow\uparrow\uparrow\downarrow - \uparrow\uparrow\downarrow\uparrow), \\
\chi_5 &= \frac{1}{\sqrt{2}} (\uparrow\downarrow\uparrow\uparrow - \downarrow\uparrow\uparrow\uparrow), \\
\chi_6 &= \frac{1}{2} (\uparrow\downarrow\uparrow\downarrow - \uparrow\downarrow\downarrow\uparrow - \downarrow\uparrow\uparrow\downarrow + \downarrow\uparrow\downarrow\uparrow),
\end{aligned} \tag{19}$$

where subscript outside the paranthesis and bracket denotes the spin of the hadron.

The only color singlet configurations of $q_1 q_2 \bar{q}_3 \bar{q}_4$ tetraquark states are $\mathbf{6}_c \otimes \bar{\mathbf{6}}_c$ or $\bar{\mathbf{3}}_c \otimes \mathbf{3}_c$ color configurations with the respective color wave functions

$$\phi_1 = |(q_1 q_2)^6 (\bar{q}_3 \bar{q}_4)^{\bar{6}} \rangle, \quad \phi_2 = |(q_1 q_2)^{\bar{3}} (\bar{q}_3 \bar{q}_4)^3 \rangle. \tag{20}$$

For the $qq\bar{Q}\bar{Q}$ configuration, we can write Eq. (20) as

$$\begin{aligned}
\phi_1 &= |(q_1 q_2)^6 (\bar{Q}_3 \bar{Q}_4)^{\bar{6}} \rangle \\
&= \frac{1}{2\sqrt{6}} [2(rr\bar{r}\bar{r} + gg\bar{g}\bar{g} + bb\bar{b}\bar{b}) + rb\bar{b}\bar{r} + br\bar{r}\bar{b} \\
&\quad + gr\bar{g}\bar{r} + rg\bar{g}\bar{r} + gb\bar{b}\bar{g} + bg\bar{b}\bar{g} + gr\bar{r}\bar{g} + rg\bar{r}\bar{g} \\
&\quad + gb\bar{g}\bar{b} + bg\bar{g}\bar{b} + rb\bar{r}\bar{b} + br\bar{r}\bar{b}]
\end{aligned} \tag{21}$$

and

$$\begin{aligned}
\phi_2 &= |(q_1 q_2)^{\bar{3}} (\bar{Q}_3 \bar{Q}_4)^3 \rangle \\
&= \frac{1}{2\sqrt{3}} (rb\bar{b}\bar{r} - br\bar{r}\bar{b} - gr\bar{g}\bar{r} + rg\bar{g}\bar{r} + gb\bar{b}\bar{g} - bg\bar{b}\bar{g} \\
&\quad + gr\bar{r}\bar{g} - rg\bar{r}\bar{g} - gb\bar{g}\bar{b} + bg\bar{g}\bar{b} - rb\bar{r}\bar{b} + br\bar{r}\bar{b}).
\end{aligned} \tag{22}$$

Concerning the color degree of freedom, the heavy quark is treated as an $SU(3)$ singlet.

Armed with these considerations, we give the wave function bases in the notation of $|(q_1 q_2)_{\text{spin}}^{\text{color}} (\bar{Q}_3 \bar{Q}_4)_{\text{spin}}^{\text{color}} \rangle_{\text{total spin}}$:

$$\begin{aligned}
\phi_1\chi_1 &= |(q_1q_2)_1^6(\bar{Q}_3\bar{Q}_4)_1^{\bar{6}}\rangle_2\delta_{12}^S\delta_{34}, \\
\phi_2\chi_1 &= |(q_1q_2)_1^{\bar{3}}(\bar{Q}_3\bar{Q}_4)_1^3\rangle_2\delta_{12}^A, \\
\phi_1\chi_2 &= |(q_1q_2)_1^6(\bar{Q}_3\bar{Q}_4)_1^{\bar{6}}\rangle_1\delta_{12}^S\delta_{34}, \\
\phi_2\chi_2 &= |(q_1q_2)_1^{\bar{3}}(\bar{Q}_3\bar{Q}_4)_1^3\rangle_1\delta_{12}^A, \\
\phi_1\chi_3 &= |(q_1q_2)_1^6(\bar{Q}_3\bar{Q}_4)_1^{\bar{6}}\rangle_0\delta_{12}^S\delta_{34}, \\
\phi_2\chi_3 &= |(q_1q_2)_1^{\bar{3}}(\bar{Q}_3\bar{Q}_4)_1^3\rangle_0\delta_{12}^A, \\
\phi_1\chi_4 &= |(q_1q_2)_1^6(\bar{Q}_3\bar{Q}_4)_0^{\bar{6}}\rangle_1\delta_{12}^S, \\
\phi_2\chi_4 &= |(q_1q_2)_1^{\bar{3}}(\bar{Q}_3\bar{Q}_4)_0^3\rangle_1\delta_{12}^A\delta_{34}, \\
\phi_1\chi_5 &= |(q_1q_2)_0^6(\bar{Q}_3\bar{Q}_4)_1^{\bar{6}}\rangle_1\delta_{12}^A\delta_{34}, \\
\phi_2\chi_5 &= |(q_1q_2)_0^{\bar{3}}(\bar{Q}_3\bar{Q}_4)_1^3\rangle_1\delta_{12}^S, \\
\phi_1\chi_6 &= |(q_1q_2)_0^6(\bar{Q}_3\bar{Q}_4)_0^{\bar{6}}\rangle_0\delta_{12}^A, \\
\phi_2\chi_6 &= |(q_1q_2)_0^{\bar{3}}(\bar{Q}_3\bar{Q}_4)_0^3\rangle_0\delta_{12}^S\delta_{34}.
\end{aligned} \tag{23}$$

One should be aware of the wave functions which are not allowed for a given quantum numbers. It can be seen in Eq. (24) that, δ_{12}^S , δ_{12}^A , and δ_{34} factors are introduced to reflect flavor symmetry. If light quarks in (q_1q_2) are symmetric then $\delta_{12}^S = 0$. Accordingly if light quarks in (q_1q_2) are antisymmetric then $\delta_{12}^A = 0$. If the two quarks are heavy and identical, then we have $\delta_{34} = 0$. Apart from these considerations, when the factor cannot be equal to 0, one can use its value as 1. A similar logic was used in Ref. [131].

D. Magnetic Moment

The magnetic moment of a hadron arises from its charge distribution and angular momentum distribution. Not only it presents valuable information about the internal structure of the hadron but also the shape of the hadron. Furthermore, magnetic moment is an excellent probe for quark-gluon dynamics inside the hadron since it is the leading-order response of a bound state to a weak external magnetic moment [132].

The magnetic moment of a doubly-heavy tetraquark system can be written as

$$\mu = \langle \phi_{IJ} | \hat{\mu}_m | \phi_{IJ} \rangle, \tag{24}$$

where $\hat{\mu}_m = \sum_{i=1}^4 \frac{\hat{Q}_i}{2m_i} \hat{\sigma}_i^z$ is the magnetic moment operator, \hat{Q}_i is the electric charge operator of the i -th quark, $\hat{\sigma}_i^z$ is the third component (z -component) of Pauli matrix and ϕ_{IJ} is the eigenvector of those states.

III. NUMERICAL RESULTS AND DISCUSSION

The DMC algorithm can be carried out to obtain the desired solution to Eq. (15) equation with the formalism described in Ref. [122]. We follow the same algorithm.

The main idea can be described as follows: The kinetic and potential energy terms of the Schrödinger equation correspond to the diffusion term (source/sink) in the diffusion equation

$$\frac{\partial u(\mathbf{r}, t)}{\partial t} = \nabla \cdot [D(u(\mathbf{r}, t), \mathbf{r}) \nabla u(\mathbf{r}, t)] \tag{25}$$

where $u(\mathbf{r}, t)$ is the density of the diffusing matter and $D(u(\mathbf{r}, t), \mathbf{r})$ is the diffusion term or coefficient. The Schrödinger equation with imaginary time is a kind of diffusion equation with source/sink mechanism. It can be solved by simulating random walks which are subject to birth/death processes resulting from the source/sink term. In the DMC algorithm, the wave function is represented by random walks of many particles. Each walker will walk randomly in finite small time steps and experience death or birth. After sufficiently long time, the diffusion of the walkers will approach the ground state wave function.

There are two theoretical errors in this model which have different sources. The first one is statistical related to DMC method itself and the second one is related to quark model parameters. We utilize Monte Carlo simulations comprising 50000 steps. We use 1×10^4 walkers in the simulations to implement the movement of each quark. Any implementation of DMC method leads to a point where the distribution of the walkers should be uncorrelated. To avoid this situation, we consider in the averages the

values for every 500 steps and we drop the first 10^5 values to prevent any effect of chosen initial configuration. The resulting energy is averaged over the last 10000 steps with a standard deviation $\simeq 1$ MeV.

Before we present our results, we give possible corresponding thresholds in Table II regarding the states that are studied. We also present our results according to the strangeness quantum number.

TABLE II. Corresponding thresholds for the doubly heavy tetraquark systems of this work.

System	I	Thresholds		
		$J^P = 0^+$	$J^P = 1^+$	$J^P = 2^+$
$nn\bar{c}\bar{c}$	0		DD^*	
	1	DD	DD^*	D^*D^*
$ss\bar{c}\bar{c}$	0	D_sD_s	$D_sD_s^*$	$D_s^*D_s^*$
$nn\bar{b}\bar{b}$	0		$\bar{B}\bar{B}^*$	
	1	$\bar{B}\bar{B}$	$\bar{B}\bar{B}^*$	$\bar{B}^*\bar{B}^*$
$ns\bar{b}\bar{b}$	$\frac{1}{2}$	$\bar{B}\bar{B}_s$	$\bar{B}_s^*\bar{B}/\bar{B}^*\bar{B}_s$	$\bar{B}^*\bar{B}_s^*$
$ss\bar{b}\bar{b}$	0	$\bar{B}_s\bar{B}_s$	$\bar{B}_s\bar{B}_s^*$	$\bar{B}_s^*\bar{B}_s^*$
$nn\bar{c}\bar{b}$	0	$D\bar{B}$	$D^*\bar{B}/D\bar{B}^*$	$D^*\bar{B}^*$
	1	$D\bar{B}$	$D^*\bar{B}/D\bar{B}^*$	$D^*\bar{B}^*$
$ss\bar{c}\bar{b}$	0	\bar{B}_sD_s	$D_s^*\bar{B}_s/D_s\bar{B}_s^*$	$\bar{B}_s^*D_s^*$

A. Tetraquarks with Strangeness-Zero (S=0)

The quark content for these tetraquark states are $nn\bar{c}\bar{c}$, $nn\bar{b}\bar{b}$, and $nn\bar{c}\bar{b}$. In Table ??, our results for the mass and magnetic moment values are listed.

It can be seen from the table that, the mass splittings in the $nn\bar{c}\bar{c}$ system are 221 MeV for $I = 1$ and 238 MeV for $I = 0$. In the $nn\bar{b}\bar{b}$ system, for $I = 1$ the mass splitting is 124 MeV and for $I = 0$ it is 286 MeV. For the mixed heavy quark system, $nn\bar{c}\bar{b}$, the mass splittings are 224 MeV for $I = 1$ and 276 MeV. One can also compare mass spectra with the corresponding thresholds which is shown in Fig. 1.

We can see from the figure that obtained mass values for $I = 1$ case are above corresponding thresholds. They should be resonance states. A special attention should be made on the $I = 0$ states. In the case of $I = 0$, one of the $J^P = 1^+$ states is high above than the DD^* threshold. The other is just a few MeV above. The mass and magnetic moment of the T_{cc}^+ are predicted to be $M = 3892$ MeV and $\mu = 0.28\mu_N$, respectively. The obtained mass agree well with the observed T_{cc}^+ mass, $M = 3875$ MeV. Ref. [60] calculated $I(J^P) = 0(1^+) T_{cc}^+$ magnetic moment as $\mu = 0.18\mu_N$ in compact configuration and $\mu = 0.13\mu_N$ in deuteron-like configuration. In Ref. [88], the magnetic moments of T_{cc}^+ state are obtained in both diquark-antidiquark and molecular configurations via light-cone QCD sum rule formalism. They obtained magnetic moment as $\mu = 0.66_{-0.23}^{+0.34}\mu_N$ with diquark-antidiquark configuration and $\mu = 0.43_{-0.22}^{+0.23}\mu_N$ with molecular configuration. Ref. [131] obtained magnetic moment of T_{cc}^+ as $\mu = 0.88\mu_N$ by using MIT bag model. Ref. [133] calculated magnetic moment as $\mu = -0.09\mu_N$ by using constituent quark model. Ref. [134] obtained a magnetic moment of $\mu = 0.732\mu_N$ by using heavy antiquark-diquark symmetry in the constituent quark model. The calculated results for the magnetic moment of T_{cc}^+ in general are not consistent with each other.

In Figure 2, mass spectrum of $nn\bar{b}\bar{b}$ states can be seen. The results of $I = 1$ case are above corresponding thresholds. They should be resonance states. In the case of $I = 0$, the state which has a mass of 10624 MeV is 20 MeV above $\bar{B}\bar{B}^*$ threshold whereas the other is 266 MeV below. The obtained mass of this state in our model is $M = 10338$ MeV. This should be a deeply bound state. Considering magnetic moment of this state which is named as T_{bb}^- , we obtain $\mu = -0.32\mu_N$. Ref. [60] calculated $I(J^P) = 0(1^+) T_{bb}^-$ magnetic moment as $\mu = 0.64\mu_N$ in compact configuration and as $\mu = 0.49\mu_N$ in deuteron-like configuration. In Ref. [92], magnetic moments of T_{bb} states with $J^P = 1^+$ are obtained in light-cone QCD sum rule using molecular picture. The results follow as $\mu = (1.72 \pm 0.67)\mu_N$ for B^-B^{*-} , $\mu = (1.38 \pm 0.56)\mu_N$ for B^0B^{*-} , $\mu = (-0.44 \pm 0.17)\mu_N$ for B^-B^{*0} , and $\mu = (-0.77 \pm 0.28)\mu_N$ for B^0B^{*0} . Ref. [131] obtained magnetic moment of T_{bb}^- as




FIG. 1. Mass spectrum of $nn\bar{c}\bar{c}$ states with $I = 0$ (left), $I = 1$ (right) and corresponding J^P values. Red color refers to threshold values whereas blue one refers to obtained results.

$\mu = 0.18\mu_N$ whereas Ref. [134] obtained $\mu = -0.124\mu_N$ for T_{bb}^- . As in the T_{cc}^+ case, the calculated results for the magnetic moment of T_{bb}^- in general are not consistent with each other.




FIG. 2. Mass spectrum of $nn\bar{b}\bar{b}$ states with $I = 0$ (left), $I = 1$ (right) with corresponding J^P values. Red color refers to threshold values whereas blue one refers to obtained results.

In Figure 3, mass spectrum of $nn\bar{c}\bar{b}$ states can be seen. All the results corresponding to $I = 1$ are above corresponding thresholds. These states should be resonances. We have bound state candidates in the $I = 0$ channel. The $J^P = 0^+$ state which has a mass of 7108 MeV is 39 MeV below $D\bar{B}$ threshold. In $J^P = 1^+$ states, the state which has a mass of 7084 is 108 MeV below $D\bar{B}^*$ threshold and 204 MeV below $D^*\bar{B}$ threshold. The state having a mass of 7204 MeV is 84 MeV below $D^*\bar{B}$ threshold. We also obtain magnetic moments of $I = 0$ and $I = 1$ $nn\bar{c}\bar{b}$ states. Focusing on $I = 0$ states, we obtain $\mu = 0.74\mu_N$ and $\mu = 0.45\mu_N$ for two low-lying $J^P = 1^+$ states. Ref. [60] obtained $I(J^P) = 0(1^+) T_{bc}$ magnetic moment as $\mu = 0.66\mu_N$ in compact configuration and as $\mu = 0.59\mu_N$ in deuteron-like configuration. Our results agree with these results.




FIG. 3. Mass spectrum of $nn\bar{c}\bar{b}$ states with $I = 0$ (left), $I = 1$ (right) with corresponding J^P values. Red color refers to threshold values whereas blue one refers to obtained results.

B. Tetraquarks with Strangeness-One (S=-1)

The quark content for these tetraquark states are $ns\bar{c}\bar{c}$ and $ns\bar{b}\bar{b}$. We present mass spectrum and magnetic moment results in Table IV. As can be seen from table, in the $ns\bar{c}\bar{c}$ system the mass splitting is 285 MeV for $J^P = 0^+$ states and 145 MeV for $J^P = 1^+$ states. In the $ns\bar{b}\bar{b}$ system, the mass splitting is 131 MeV for $J^P = 0^+$ states and 94 MeV for $J^P = 1^+$ states.

TABLE IV. Mass spectrum and magnetic moments of doubly heavy $ns\bar{c}\bar{c}$ and $ns\bar{b}\bar{b}$ tetraquarks, where $n = u$ or d . Mass results are in units of MeV. Magnetic moments (in units of nuclear magneton $\mu_N, \mu_p = 2.79285\mu_N$) are organized in the order of $I_3 = 1/2, -1/2$ for $I = 1$.

State	J^P	Mass	μ
$ns\bar{c}\bar{c}$	0^+	4349	-
		4064	-
		4056	0.43, -0.67
	1^+	4201	1.24, -1.96
		4128	0.52, -0.75
2^+	4314	0.24, -2.76	
$ns\bar{b}\bar{b}$	0^+	10852	-
		10721	-
		10736	0.48, -0.66
	1^+	10778	1.11, -1.35
		10684	-0.09, -0.12
	2^+	10747	1.36, -1.98

In Figure 4, mass spectrum of $ns\bar{c}\bar{c}$ and $ns\bar{b}\bar{b}$ tetraquarks can be seen. In the $ns\bar{c}\bar{c}$ systems, all obtained mass values are above corresponding thresholds. They should be resonance states. In the $ns\bar{b}\bar{b}$ systems, the $J^P = 1^+$ state with a mass of $M = 10684$ is below corresponding thresholds $\bar{B}_s^*\bar{B}/\bar{B}^*\bar{B}_s$. This could be a bound state candidate. The $J^P = 2^+$ state is just 7 MeV above corresponding $\bar{B}^*\bar{B}_s^*$ state. This state seems to be a resonance state but also could be a shallow bound state candidate. Focusing on this state, in Ref. [93], $J^P = 1^+$ $T_{QQ\bar{q}\bar{s}}$ and $T_{QQ\bar{s}\bar{s}}$ states are studied as diquark-antidiquark tetraquarks and magnetic moments are calculated with using heavy axial vector diquark-light scalar antidiquark type current denoted as J^1 , heavy scalar diquark-light axial vector antidiquark type current denoted as J^2 , heavy tensor diquark-light axial vector antidiquark type current denoted as J^3 and heavy vector diquark-light pseudotensor antidiquark type current denoted as J^4 . According to their work, J^1 gives $\mu = (-0.46 \pm 0.07)\mu_N$, J^2 gives $\mu = (-0.34 \pm 0.12)\mu_N$, J^3 gives $\mu = (-2.37 \pm 0.50)\mu_N$ and J^4 gives $\mu = (-2.90 \pm 0.36)\mu_N$ for $T_{bb\bar{u}\bar{s}}$ state. For $T_{bb\bar{d}\bar{s}}$ state, J^1 gives $\mu = (-0.59 \pm 0.09)\mu_N$, J^2 gives $\mu = (-0.68 \pm 0.24)\mu_N$,

J^3 gives $\mu = (-1.37 \pm 0.17)\mu_N$ and J^4 gives $\mu = (-1.45 \pm 0.33)\mu_N$. Our results are compatible with this work. In the same study considering $T_{cc\bar{u}\bar{s}}$ state, J^1 gives $\mu = (0.22 \pm 0.07)\mu_N$, J^2 gives $\mu = (0.18 \pm 0.06)\mu_N$, J^3 gives $\mu = (1.09 \pm 0.36)\mu_N$ and J^4 gives $\mu = (1.04 \pm 0.35)\mu_N$. For $T_{cc\bar{d}\bar{s}}$ state, J^1 gives $\mu = (0.15 \pm 0.05)\mu_N$, J^2 gives $\mu = (0.36 \pm 0.11)\mu_N$, J^3 gives $\mu = (1.70 \pm 0.57)\mu_N$ and J^4 gives $\mu = (2.08 \pm 0.64)\mu_N$. Our results are compatible with this work.




FIG. 4. Mass spectrum of $ns\bar{c}\bar{c}$ (left) and $ns\bar{b}\bar{b}$ (right) tetraquarks with corresponding J^P values. Red color refers to threshold values whereas blue one refers to obtained results.

C. Tetraquarks with Strangeness-Two (S=-2)

The quark content for these tetraquark states are $ss\bar{c}\bar{c}$ and $ss\bar{b}\bar{b}$. We present mass spectrum and magnetic moment results in Table V. As can be seen from table, in the $ss\bar{c}\bar{c}$ system the mass splittings are 269 MeV for $J^P = 0^+$ states, 104 MeV for $ss\bar{b}\bar{b}$ $J^P = 0^+$ states. In the $ss\bar{c}\bar{b}$ systems, the mass splittings are 180 MeV for $J^P = 0^+$ states and 105 MeV for $J^P = 1^+$ states.

TABLE V. Mass spectrum and magnetic moments of doubly heavy $ns\bar{c}\bar{c}$ and $ns\bar{b}\bar{b}$ tetraquarks, where $n = u$ or d . Mass results are in units of MeV. Magnetic moments are presented in units of nuclear magneton μ_N , $\mu_p = 2.79285\mu_N$.

State	J^P	Mass	μ
$ss\bar{c}\bar{c}$	0^+	4474	-
		4205	-
	1^+	4323	-1.33
	2^+	4381	-1.76
$ss\bar{b}\bar{b}$	0^+	10972	-
		10868	-
	1^+	10908	0.73
	2^+	10926	1.42
$ss\bar{c}\bar{b}$	0^+	7833	-
		7653	-
		7764	-0.86
	1^+	7816	-0.35
		7711	-0.21
	2^+	7862	-2.09

In Figures 5, 6, and 7 mass values of $ss\bar{c}\bar{c}$, $ss\bar{b}\bar{b}$, and $ss\bar{c}\bar{b}$ states are located. As can be seen from figures all the states are above their corresponding thresholds, therefore they should be resonance states. Considering magnetic moments, Ref. [93] calculated

magnetic moments of $T_{cc\bar{s}\bar{s}}$ and $T_{bb\bar{s}\bar{s}}$ states in light cone QCD sum rule formalism. The results are $\mu = (1.52 \pm 0.46)\mu_N$ for J^3 type current and $\mu = (1.90 \pm 0.56)\mu_N$ for J^4 type current. Our magnetic moment results of $ss\bar{c}\bar{c}$ are compatible with these results. For the $T_{bb\bar{s}\bar{s}}$ state, they obtained $\mu = (-2.90 \pm 0.23)\mu_N$ for J^3 type current and $\mu = (-3.20 \pm 0.48)\mu_N$ for J^4 type current. Our results are not as compatible as in the previous case.




FIG. 5. Mass spectrum of $ss\bar{c}\bar{c}$ tetraquark states. Red color refers to threshold values whereas blue one refers to obtained results.




FIG. 6. Mass spectrum of $ss\bar{b}\bar{b}$ tetraquark states. Red color refers to threshold values whereas blue one refers to obtained results.

For the wave functions in this work, the general ground state can be the mixing of $\phi_2\chi_3$ and $\phi_1\chi_6$ for the isotriplet tetraquarks with $J^P = 0^+$. In the case of isosinglet tetraquarks with $J^P = 1^+$, the wave function can be mixing of $\phi_2\chi_5$ and $\phi_1\chi_4$ for $nn\bar{Q}\bar{Q}$ tetraquarks and mixing of $\phi_2\chi_2$, $\phi_2\chi_5$, $\phi_1\chi_4$ for $nn\bar{Q}\bar{Q}$ tetraquarks. In the $nn\bar{c}\bar{b}$ tetraquarks, for $I(J^P) = 1(0^+)$ the wave function can be mixing of $\phi_2\chi_3$ and $\phi_1\chi_6$, for $I(J^P) = 0(0^+)$ the wave function can be mixing of $\phi_1\chi_3$ and $\phi_2\chi_6$, for $I(J^P) = 1(1^+)$ the wave function can be mixing of $\phi_2\chi_2$, $\phi_2\chi_4$, $\phi_1\chi_5$, and for $I(J^P) = 0(1^+)$ the wave function can be mixing




FIG. 7. Mass spectrum of $ss\bar{c}\bar{b}$ tetraquark states. Red color refers to threshold values whereas blue one refers to obtained results.

of $\phi_2\chi_2$, $\phi_2\chi_5$, $\phi_1\chi_4$.

We compare our results with some of the works that studied same states. In Table VI we present our results with some other works conducted in different frameworks. As can be seen from table, our results are in agreement with the cited works. We also compare our mass results in Table VII with a recent work conducted in DMC [130]. The agreement of the results can be seen.

Finally, we compare our results of magnetic moments with the available results in Table VIII. Our results are in general compatible with the cited work.

IV. CONCLUDING REMARKS

We employ a diffusion Monte Carlo (DMC) method to investigate systematically masses and magnetic moments of doubly heavy tetraquark states in constituent quark model. This method has some advantages compared to other methods; (i) reduces the uncertainty of the numerical procedure, (ii) accounts for multiparticle correlations, and (iii) avoids quark clustering in the relevant states.

Most QCD states arise as bound states or as resonances of multihadron configurations. Studying mass spectrum is an important aspect for the general view point of the related states. To obtain possible mass locations around the corresponding thresholds is not only crucial for understanding the underlying dynamics of the exotic hadrons and the nature of strong interactions of QCD but also useful for experimental researches for their existence.

In the $nn\bar{c}\bar{c}$ systems, the $I(J^P) = 0(1^+)$ $nn\bar{c}\bar{c}$ state with mass 3892 MeV is the candidate of the experimentally observed T_{cc}^+ state. Its mass is only 16 MeV above DD^* threshold. T_{cc}^+ is argued to be a shallow bound state. Although the level of the uncertainty is less than 1 MeV, one cannot conclude this state is bound or not. Our results suggest $I(J^P) = 0(1^+)$ $nn\bar{c}\bar{c}$ could be a shallow bound state. All the other predicted states with $I(J^P) = 0(1^+)$ and $I(J^P) = 1(1^+)$ for $nn\bar{c}\bar{c}$ tetraquarks are well above corresponding thresholds. The magnetic moment presents information about the inner quark structure and can be used to distinguish the preferred configuration from predictions of different theoretical models. Our prediction of the magnetic moment for T_{cc}^+ state is $\mu = 0.28\mu_N$.

In the $nn\bar{b}\bar{b}$ systems, we find a deeply bound state candidate, namely $I(J^P) = 0(1^+)$ $nn\bar{b}\bar{b}$ state with mass 10338 MeV. It is just 266 MeV below $\bar{B}\bar{B}^*$ threshold. All the other $nn\bar{c}\bar{c}$ states are above their corresponding thresholds and could be resonances. The existence T_{bb}^- was conjectured long time ago and recent lattice QCD studies confirm the existence of $ud\bar{b}\bar{b}$ tetraquark with $I(J^P) = 0(1^+)$ [102, 106, 107, 109]. We obtain magnetic moment of T_{bb}^- as $\mu = -0.32\mu_N$.

In the $nn\bar{c}\bar{b}$ systems, we observe some bound state candidates. For $I = 0$, low-lying $J^P = 0^+$ $nn\bar{c}\bar{b}$ has a mass of 7108 MeV and is 39 MeV below $D\bar{B}$ threshold. One of the $J^P = 1^+$ $nn\bar{c}\bar{b}$ has a mass of 7084 MeV and is 108 below $D\bar{B}^*$ threshold and 204 MeV below $D^*\bar{B}^*$ threshold. The $J^P = 1^+$ $nn\bar{c}\bar{b}$ state with mass of 7204 MeV is 12 MeV above $D\bar{B}^*$ threshold and 84 MeV below $D^*\bar{B}^*$ threshold. The two low-lying $nn\bar{c}\bar{b}$ states with $J^P = 1^+$ could be a bound or loosely bound state with

TABLE VI. Comparison of calculated mass (in MeV) with different calculations conducted in different models for double heavy tetraquarks. The masses before and after slash stand for color states splitting. Refs. [131], [66], [48] employ the CMI model and Ref. [29] employ relativistic quark model.

State	J	This work	[66]	[131]	[48]	[29]
$(nn\bar{c}\bar{c})^{I=1}$	0	4062/4283	4078/4356	4032/4342	4195/4414	4056
	1	4104	4201	4117	4268	4079
	2	4207	4271	4179	4318	4118
$(nn\bar{c}\bar{c})^{I=0}$	1	3892/4130	4007/4204	3925/4205	4041/4313	3935
$(nn\bar{b}\bar{b})^{I=1}$	0	10624/10748	10841/10937	10834/11092	10765/11019	10648
	1	10680	10875	10854	10779	10657
	2	10702	10897	10878	10799	10673
$(nn\bar{b}\bar{b})^{I=0}$	1	10338/10624	10686/10821	10654/10982	10550/10951	10502
$(nn\bar{c}\bar{b})^{I=1}$	0	7396/7620	7457/7643	7438/7714	7519/7740	7383
	1	7510/7542	7473/7548	7465/7509	7537/7561	7396/7403
	2	7448	7609	7699	7729	7422
$(nn\bar{c}\bar{b})^{I=0}$	0	7108/7201	7256/7429	7260/7502	7297/7580	7239
	1	7084/7204	7321/7431	7288/7518	7.325/7.607	7246
	2	7324	7516	7605	7666	-
$ns\bar{c}\bar{c}$	0	4165/4349	4236/4514	4165/4429	4323/4512	4221
	1	4056/4128	4225/4363	4091/4247	4232/4394	4143/4239
	2	4201	4400	4314	4427	4271
$ns\bar{b}\bar{b}$	0	4314	4434	4305	4440	4271
	0	10721/10852	10999/11095	10955/11160	10883/11098	10802
	1	10684/10736	10911/11010	10811/10974	10734/10897	10706/10809
$ss\bar{c}\bar{c}$	0	10778	11037	11068	11046	10823
	0	4205/4474	4395/4672	4300/4521	4417/4587	4359
	1	4323	4526	4382	4493	4375
$ss\bar{b}\bar{b}$	2	4381	4597	4433	4536	4402
	0	10868/10972	11157/11254	11078/11232	10972/11155	10932
	1	10908	11199	11099	10986	10939
$ss\bar{c}\bar{b}$	2	10926	11224	11119	11004	10950
	0	7653/7833	7774/7960	7693/7875	7735/7894	7673
	1	7711/7764	7793/7872	7716/7757	7752/7775	7683/7684
	2	7816	7924	7858	7881	7701
	2	7862	7908	7779	7798	7701

TABLE VII. Comparison of calculated mass (in MeV) with calculations conducted in DMC. “NB” represents no bound solution. We only took the results which are bound state solutions from Ref. [130].

State	J^P	This work	AL1 [130]	AP1 [130]	SLM [130]
$nn\bar{c}\bar{b}^{I=0}$	0^+	7108/7201	7136	7164	6986
$nn\bar{c}\bar{c}^{I=0}$	1^+	3892/4130	3880	3916	3759
$nn\bar{b}\bar{b}^{I=0}$	1^+	10338/10624	10500	10510	10249
$ns\bar{b}\bar{b}^{I=1/2}$	1^+	10736/10778/10684	10660	10667	10653
$nn\bar{c}\bar{b}^{I=0}$	1^+	7084/7324/7204	7200	7224	7012
$nn\bar{c}\bar{b}^{I=0}$	2^+	7360	77367	7400	NB

regard to corresponding threshold. Some of the recent lattice QCD calculations [108, 113, 135] did not find any bound state indications of $I(J^P) = 0(0^+)$ and $I(J^P) = 0(1^+)$ $ud\bar{c}\bar{b}$ states but cannot rule out their existence. However, Refs. [105, 136] support $I(J^P) = 0(1^+)$ $ud\bar{c}\bar{b}$ existence. A very recent lattice QCD study also found evidence for $J = 0$ shallow bound state and found hints for $J = 1$ broad resonance of $ud\bar{c}\bar{b}$ [137].

In the $ns\bar{c}\bar{c}$ and $ns\bar{b}\bar{b}$ systems, we find all the $ns\bar{c}\bar{c}$ states above corresponding thresholds. They could be resonances. In the

TABLE VIII. Comparison of magnetic moments (in units of nuclear magneton μ_N , $\mu_p = 2.79285\mu_N$).

State	J^P	Present work	μ [131]	
$(nn\bar{c}\bar{c})^{I=1}$	0 ⁺	-	-	
	1 ⁺	1.14, -0.07, 1.26	1.36, -0.03, -1.43	
	2 ⁺	2.66, -0.06, 2.39	2.80, -0.05, -2.90	
$(nn\bar{c}\bar{c})^{I=0}$	1 ⁺	-0.28 0.35	-0.88 0.83	
	$(nn\bar{b}\bar{b})^{I=1}$	0 ⁺	-	
$(nn\bar{b}\bar{b})^{I=1}$	1 ⁺	-1.02, 0.28, -0.64	1.81, 0.52, -0.78	
	2 ⁺	-2.32, 0.47, -1.28	3.65, 1.04, -1.57	
	$(nn\bar{b}\bar{b})^{I=0}$	1 ⁺	0.32 0.69	0.18 0.86
$(nn\bar{c}\bar{b})^{I=1}$	0 ⁺	-	-	
	1 ⁺	1.84, 0.12, -1.36 0.09, 0.12, 0.21	2.33, 0.21, -1.90 -0.17, -0.32, -0.47	
	2 ⁺	1.94, 0.41, -0.59 2.47, 0.32, -1.68	2.57, 0.82, -0.93 3.24, 0.50, -2.24	
	$(nn\bar{c}\bar{b})^{I=0}$	0 ⁺	-	
$(nn\bar{c}\bar{b})^{I=0}$	1 ⁺	0.74 0.70 0.45	0.55 0.50 -0.33	
	2 ⁺	-0.54	0.50	
	$ns\bar{c}\bar{c}$	0 ⁺	-	-
		1 ⁺	0.43, -0.67 1.24, -1.96	0.32, -1.33 0.86, -1.58
2 ⁺		0.52, -0.75 0.24, -2.76	-0.95, -1.03 0.19, -2.68	
$ns\bar{b}\bar{b}$		0 ⁺	-	
$ns\bar{b}\bar{b}$	1 ⁺	0.48, -0.66 1.11, -1.35 -0.09, -0.12	0.65, -0.68 1.02, -1.50 0.14, 0.16	
	2 ⁺	1.36, -1.98	1.25, -1.39	
	$ss\bar{c}\bar{c}$	0 ⁺	-	-
		1 ⁺	-1.33	-1.22
2 ⁺		-1.76	-2.46	
$ss\bar{b}\bar{b}$	0 ⁺	-	-	
	1 ⁺	0.73	-0.60	
	2 ⁺	1.42	-1.20	
$ss\bar{c}\bar{b}$	0 ⁺	-	-	
	1 ⁺	-0.86 -0.35 -0.21	-1.54 -0.48 -0.66	
	2 ⁺	-2.09	-1.84	

case of $ns\bar{b}\bar{b}$ systems, the low-lying $J^P = 1^+$ state is 8 MeV below $\bar{B}^*\bar{B}_s$ threshold and 11 MeV below $\bar{B}_s^*\bar{B}$ threshold. This could be a bound state candidate. The $J^P = 2^+$ state is 7 MeV above $\bar{B}^*\bar{B}_s$ threshold. This could be a shallow bound state

candidate. In Ref. [113], a clear evidence for the existence of $us\bar{b}\bar{b}$ with $J^P = 1^+$ is found.

Lattice QCD studies present important evidences about the existence of the states. For the $I = 0 T_{bb}$, $I = 0, 1 T_{bc}$ and T_{bbs} tetraquark states, lattice QCD results support deeply bound state or very narrow resonances (see a recent review on lattice QCD results and references therein [138]). Our results are in agreement with those predictions.

In the $ss\bar{c}\bar{c}$, $ss\bar{b}\bar{b}$, and $ss\bar{c}\bar{b}$ systems, all the states are above their corresponding thresholds. They all could be resonances. We have no bound state candidates in these systems.

The experimental progress on the multiquark states open a new era for our understanding of QCD. We hope that our results will be useful for not only for experimental studies but also other theoretical studies.

- [1] M. Gell-Mann, A Schematic Model of Baryons and Mesons, *Phys. Lett.* **8**, 214 (1964).
- [2] G. Zweig, An SU(3) model for strong interaction symmetry and its breaking. Version 2, in *DEVELOPMENTS IN THE QUARK THEORY OF HADRONS. VOL. 1. 1964 - 1978*, edited by D. B. Lichtenberg and S. P. Rosen (1964) pp. 22–101.
- [3] S. K. Choi *et al.* (Belle), Observation of a narrow charmonium-like state in exclusive $B^\pm \rightarrow K^\pm \pi^+ \pi^- J/\psi$ decays, *Phys. Rev. Lett.* **91**, 262001 (2003), arXiv:hep-ex/0309032.
- [4] T. Gershon (LHCb), Exotic hadron naming convention, (2022), arXiv:2206.15233 [hep-ex].
- [5] R. F. Lebed, R. E. Mitchell, and E. S. Swanson, Heavy-Quark QCD Exotica, *Prog. Part. Nucl. Phys.* **93**, 143 (2017), arXiv:1610.04528 [hep-ph].
- [6] A. Ali, J. S. Lange, and S. Stone, Exotics: Heavy Pentaquarks and Tetraquarks, *Prog. Part. Nucl. Phys.* **97**, 123 (2017), arXiv:1706.00610 [hep-ph].
- [7] S. L. Olsen, T. Skwarnicki, and D. Zieminska, Nonstandard heavy mesons and baryons: Experimental evidence, *Rev. Mod. Phys.* **90**, 015003 (2018), arXiv:1708.04012 [hep-ph].
- [8] F.-K. Guo, C. Hanhart, U.-G. Meißner, Q. Wang, Q. Zhao, and B.-S. Zou, Hadronic molecules, *Rev. Mod. Phys.* **90**, 015004 (2018), [Erratum: *Rev. Mod. Phys.* **94**, 029901 (2022)], arXiv:1705.00141 [hep-ph].
- [9] Y.-R. Liu, H.-X. Chen, W. Chen, X. Liu, and S.-L. Zhu, Pentaquark and Tetraquark states, *Prog. Part. Nucl. Phys.* **107**, 237 (2019), arXiv:1903.11976 [hep-ph].
- [10] N. Brambilla, S. Eidelman, C. Hanhart, A. Nefediev, C.-P. Shen, C. E. Thomas, A. Vairo, and C.-Z. Yuan, The XYZ states: experimental and theoretical status and perspectives, *Phys. Rept.* **873**, 1 (2020), arXiv:1907.07583 [hep-ex].
- [11] R. Aaij *et al.* (LHCb), Observation of an exotic narrow doubly charmed tetraquark, *Nature Phys.* **18**, 751 (2022), arXiv:2109.01038 [hep-ex].
- [12] R. Aaij *et al.* (LHCb), Study of the doubly charmed tetraquark T_{cc}^+ , *Nature Commun.* **13**, 3351 (2022), arXiv:2109.01056 [hep-ex].
- [13] J. P. Ader, J. M. Richard, and P. Taxil, DO NARROW HEAVY MULTI - QUARK STATES EXIST?, *Phys. Rev. D* **25**, 2370 (1982).
- [14] J. I. Ballot and J. M. Richard, FOUR QUARK STATES IN ADDITIVE POTENTIALS, *Phys. Lett. B* **123**, 449 (1983).
- [15] H. J. Lipkin, A MODEL INDEPENDENT APPROACH TO MULTI - QUARK BOUND STATES, *Phys. Lett. B* **172**, 242 (1986).
- [16] S. Zouzou, B. Silvestre-Brac, C. Gignoux, and J. M. Richard, FOUR QUARK BOUND STATES, *Z. Phys. C* **30**, 457 (1986).
- [17] L. Heller and J. A. Tjon, On the Existence of Stable Dimesons, *Phys. Rev. D* **35**, 969 (1987).
- [18] J. Carlson, L. Heller, and J. A. Tjon, Stability of Dimesons, *Phys. Rev. D* **37**, 744 (1988).
- [19] B. Silvestre-Brac and C. Semay, Systematics of $L = 0$ q-2 anti-q-2 systems, *Z. Phys. C* **57**, 273 (1993).
- [20] C. Semay and B. Silvestre-Brac, Diquonia and potential models, *Z. Phys. C* **61**, 271 (1994).
- [21] C.-K. Chow, Semileptonic decays of heavy tetraquarks, *Phys. Rev. D* **51**, 3541 (1995), arXiv:hep-ph/9411221.
- [22] C.-K. Chow, From tetraquark to hexaquark: A Systematic study of heavy exotics in the large $N(c)$ limit, *Phys. Rev. D* **51**, 6327 (1995), arXiv:hep-ph/9412242.
- [23] S. Pepin, F. Stancu, M. Genovese, and J. M. Richard, Tetraquarks with color blind forces in chiral quark models, *Phys. Lett. B* **393**, 119 (1997), arXiv:hep-ph/9609348.
- [24] D. M. Brink and F. Stancu, Tetraquarks with heavy flavors, *Phys. Rev. D* **57**, 6778 (1998).
- [25] B. A. Gelman and S. Nussinov, Does a narrow tetraquark cc anti-u anti-d state exist?, *Phys. Lett. B* **551**, 296 (2003), arXiv:hep-ph/0209095.
- [26] J. Vijande, F. Fernandez, A. Valcarce, and B. Silvestre-Brac, Tetraquarks in a chiral constituent quark model, *Eur. Phys. J. A* **19**, 383 (2004), arXiv:hep-ph/0310007.
- [27] D. Janc and M. Rosina, The $T_{cc} = DD^*$ molecular state, *Few Body Syst.* **35**, 175 (2004), arXiv:hep-ph/0405208.
- [28] J. Vijande, A. Valcarce, and K. Tsushima, Dynamical study of bf QQ - anti-u anti-d mesons, *Phys. Rev. D* **74**, 054018 (2006), arXiv:hep-ph/0608316.
- [29] D. Ebert, R. N. Faustov, V. O. Galkin, and W. Lucha, Masses of tetraquarks with two heavy quarks in the relativistic quark model, *Phys. Rev. D* **76**, 114015 (2007), arXiv:0706.3853 [hep-ph].
- [30] M. Zhang, H. X. Zhang, and Z. Y. Zhang, QQ anti-q anti-q four-quark bound states in chiral SU(3) quark model, *Commun. Theor. Phys.* **50**, 437 (2008), arXiv:0711.1029 [nucl-th].
- [31] J. Vijande, A. Valcarce, and J. M. Richard, Stability of multiquarks in a simple string model, *Phys. Rev. D* **76**, 114013 (2007), arXiv:0707.3996 [hep-ph].
- [32] J. Vijande, A. Valcarce, and N. Barnea, Exotic meson-meson molecules and compact four-quark states, *Phys. Rev. D* **79**, 074010 (2009), arXiv:0903.2949 [hep-ph].

- [33] Y. Yang, C. Deng, J. Ping, and T. Goldman, S-wave Q Q anti-q anti-q state in the constituent quark model, *Phys. Rev. D* **80**, 114023 (2009).
- [34] G. Q. Feng, X. H. Guo, and B. S. Zou, $QQ'\bar{u}\bar{d}$ bound state in the Bethe-Salpeter equation approach, (2013), arXiv:1309.7813 [hep-ph].
- [35] M. Karliner and J. L. Rosner, Discovery of doubly-charmed Ξ_{cc} baryon implies a stable $(bb\bar{u}\bar{d})$ tetraquark, *Phys. Rev. Lett.* **119**, 202001 (2017), arXiv:1707.07666 [hep-ph].
- [36] E. J. Eichten and C. Quigg, Heavy-quark symmetry implies stable heavy tetraquark mesons $Q_i Q_j \bar{q}_k \bar{q}_l$, *Phys. Rev. Lett.* **119**, 202002 (2017), arXiv:1707.09575 [hep-ph].
- [37] X. Yan, B. Zhong, and R. Zhu, Doubly charmed tetraquarks in a diquark–antidiquark model, *Int. J. Mod. Phys. A* **33**, 1850096 (2018), arXiv:1804.06761 [hep-ph].
- [38] A. Ali, Q. Qin, and W. Wang, Discovery potential of stable and near-threshold doubly heavy tetraquarks at the LHC, *Phys. Lett. B* **785**, 605 (2018), arXiv:1806.09288 [hep-ph].
- [39] W. Park, S. Noh, and S. H. Lee, Masses of the doubly heavy tetraquarks in a constituent quark model, *Nucl. Phys. A* **983**, 1 (2019), arXiv:1809.05257 [nucl-th].
- [40] C. Deng, H. Chen, and J. Ping, Systematical investigation on the stability of doubly heavy tetraquark states, *Eur. Phys. J. A* **56**, 9 (2020), arXiv:1811.06462 [hep-ph].
- [41] T. F. Caramés, J. Vijande, and A. Valcarce, Exotic $bc\bar{q}\bar{q}$ four-quark states, *Phys. Rev. D* **99**, 014006 (2019), arXiv:1812.08991 [hep-ph].
- [42] E. Hernández, J. Vijande, A. Valcarce, and J.-M. Richard, Spectroscopy, lifetime and decay modes of the T_{bb}^- tetraquark, *Phys. Lett. B* **800**, 135073 (2020), arXiv:1910.13394 [hep-ph].
- [43] G. Yang, J. Ping, and J. Segovia, Doubly-heavy tetraquarks, *Phys. Rev. D* **101**, 014001 (2020), arXiv:1911.00215 [hep-ph].
- [44] M. A. Bedolla, J. Ferretti, C. D. Roberts, and E. Santopinto, Spectrum of fully-heavy tetraquarks from a diquark+antidiquark perspective, *Eur. Phys. J. C* **80**, 1004 (2020), arXiv:1911.00960 [hep-ph].
- [45] M.-T. Yu, Z.-Y. Zhou, D.-Y. Chen, and Z. Xiao, Possible molecular states in $B^{(*)}B^{(*)}$ scatterings, *Phys. Rev. D* **101**, 074027 (2020), arXiv:1912.07348 [hep-ph].
- [46] P. C. Wallbott, G. Eichmann, and C. S. Fischer, Disentangling different structures in heavy-light four-quark states, *Phys. Rev. D* **102**, 051501 (2020), arXiv:2003.12407 [hep-ph].
- [47] Y. Tan, W. Lu, and J. Ping, Systematics of $QQ\bar{q}\bar{q}$ in a chiral constituent quark model, *Eur. Phys. J. Plus* **135**, 716 (2020), arXiv:2004.02106 [hep-ph].
- [48] Q.-F. Lü, D.-Y. Chen, and Y.-B. Dong, Masses of doubly heavy tetraquarks $T_{QQ'}$ in a relativized quark model, *Phys. Rev. D* **102**, 034012 (2020), arXiv:2006.08087 [hep-ph].
- [49] E. Braaten, L.-P. He, and A. Mohapatra, Masses of doubly heavy tetraquarks with error bars, *Phys. Rev. D* **103**, 016001 (2021), arXiv:2006.08650 [hep-ph].
- [50] G. Yang, J. Ping, and J. Segovia, $QQ\bar{s}\bar{s}$ tetraquarks in the chiral quark model, *Phys. Rev. D* **102**, 054023 (2020), arXiv:2007.05190 [hep-ph].
- [51] J. F. Giron and R. F. Lebed, Spectrum of the hidden-bottom and the hidden-charm-strange exotics in the dynamical diquark model, *Phys. Rev. D* **102**, 014036 (2020), arXiv:2005.07100 [hep-ph].
- [52] Q. Qin, Y.-F. Shen, and F.-S. Yu, Discovery potentials of double-charm tetraquarks, *Chin. Phys. C* **45**, 103106 (2021), arXiv:2008.08026 [hep-ph].
- [53] Q. Meng, E. Hiyama, A. Hosaka, M. Oka, P. Gubler, K. U. Can, T. T. Takahashi, and H. S. Zong, Stable double-heavy tetraquarks: spectrum and structure, *Phys. Lett. B* **814**, 136095 (2021), arXiv:2009.14493 [nucl-th].
- [54] Q. Meng, M. Harada, E. Hiyama, A. Hosaka, and M. Oka, Doubly heavy tetraquark resonant states, *Phys. Lett. B* **824**, 136800 (2022), arXiv:2106.11868 [hep-ph].
- [55] X. Chen and Y. Yang, Doubly-heavy tetraquark states and \ast , *Chin. Phys. C* **46**, 054103 (2022), arXiv:2109.02828 [hep-ph].
- [56] Y. Jin, S.-Y. Li, Y.-R. Liu, Q. Qin, Z.-G. Si, and F.-S. Yu, Color and baryon number fluctuation of preconfinement system in production process and T_{cc} structure, *Phys. Rev. D* **104**, 114009 (2021), arXiv:2109.05678 [hep-ph].
- [57] J. F. Giron, R. F. Lebed, and S. R. Martinez, Spectrum of hidden-charm, open-strange exotics in the dynamical diquark model, *Phys. Rev. D* **104**, 054001 (2021), arXiv:2106.05883 [hep-ph].
- [58] K. Chen, R. Chen, L. Meng, B. Wang, and S.-L. Zhu, Systematics of the heavy flavor hadronic molecules, *Eur. Phys. J. C* **82**, 581 (2022), arXiv:2109.13057 [hep-ph].
- [59] O. Andreev, $QQ\bar{q}\bar{q}$ potential in string models, *Phys. Rev. D* **105**, 086025 (2022), arXiv:2111.14418 [hep-ph].
- [60] C. Deng and S.-L. Zhu, $T_{cc}+$ and its partners, *Phys. Rev. D* **105**, 054015 (2022), arXiv:2112.12472 [hep-ph].
- [61] P. G. Ortega, J. Segovia, D. R. Entem, and F. Fernandez, Nature of the doubly-charmed tetraquark $T_{cc}+$ in a constituent quark model, *Phys. Lett. B* **841**, 137918 (2023), [Erratum: *Phys.Lett.B* 847, 138308 (2023)], arXiv:2211.06118 [hep-ph].
- [62] Y. Kim, M. Oka, D. Suenaga, and K. Suzuki, Strong decays of singly heavy baryons from a chiral effective theory of diquarks, *Phys. Rev. D* **107**, 074015 (2023), arXiv:2212.08338 [hep-ph].
- [63] S. H. Lee, S. Yasui, W. Liu, and C. M. Ko, Charmed exotics in Heavy Ion Collisions, *Eur. Phys. J. C* **54**, 259 (2008), arXiv:0707.1747 [hep-ph].
- [64] S. H. Lee and S. Yasui, Stable multi-quark states with heavy quarks in a diquark model, *Eur. Phys. J. C* **64**, 283 (2009), arXiv:0901.2977 [hep-ph].
- [65] T. Hyodo, Y.-R. Liu, M. Oka, K. Sudoh, and S. Yasui, Production of doubly charmed tetraquarks with exotic color configurations in electron-positron collisions, *Phys. Lett. B* **721**, 56 (2013), arXiv:1209.6207 [hep-ph].
- [66] S.-Q. Luo, K. Chen, X. Liu, Y.-R. Liu, and S.-L. Zhu, Exotic tetraquark states with the $qq\bar{Q}\bar{Q}$ configuration, *Eur. Phys. J. C* **77**, 709 (2017), arXiv:1707.01180 [hep-ph].
- [67] T. Hyodo, Y.-R. Liu, M. Oka, and S. Yasui, Spectroscopy and production of doubly charmed tetraquarks, (2017), arXiv:1708.05169 [hep-ph].

- [68] J.-B. Cheng, S.-Y. Li, Y.-R. Liu, Y.-N. Liu, Z.-G. Si, and T. Yao, Spectrum and rearrangement decays of tetraquark states with four different flavors, *Phys. Rev. D* **101**, 114017 (2020), arXiv:2001.05287 [hep-ph].
- [69] J.-B. Cheng, S.-Y. Li, Y.-R. Liu, Z.-G. Si, and T. Yao, Double-heavy tetraquark states with heavy diquark-antiquark symmetry, *Chin. Phys. C* **45**, 043102 (2021), arXiv:2008.00737 [hep-ph].
- [70] X.-Z. Weng, W.-Z. Deng, and S.-L. Zhu, Doubly heavy tetraquarks in an extended chromomagnetic model *, *Chin. Phys. C* **46**, 013102 (2022), arXiv:2108.07242 [hep-ph].
- [71] T. Guo, J. Li, J. Zhao, and L. He, Mass spectra of doubly heavy tetraquarks in an improved chromomagnetic interaction model, *Phys. Rev. D* **105**, 014021 (2022), arXiv:2108.10462 [hep-ph].
- [72] F. S. Navarra, M. Nielsen, and S. H. Lee, QCD sum rules study of QQ - anti-u anti-d mesons, *Phys. Lett. B* **649**, 166 (2007), arXiv:hep-ph/0703071.
- [73] J. M. Dias, S. Narison, F. S. Navarra, M. Nielsen, and J. M. Richard, Relation between $T_{cc,bb}$ and $X_{c,b}$ from QCD, *Phys. Lett. B* **703**, 274 (2011), arXiv:1105.5630 [hep-ph].
- [74] M.-L. Du, W. Chen, X.-L. Chen, and S.-L. Zhu, Exotic $QQ\bar{q}\bar{q}$, $QQ\bar{q}\bar{s}$ and $QQ\bar{s}\bar{s}$ states, *Phys. Rev. D* **87**, 014003 (2013), arXiv:1209.5134 [hep-ph].
- [75] W. Chen, T. G. Steele, and S.-L. Zhu, Exotic open-flavor $bc\bar{q}\bar{q}$, $bc\bar{s}\bar{s}$ and $qc\bar{q}\bar{b}$, $sc\bar{s}\bar{b}$ tetraquark states, *Phys. Rev. D* **89**, 054037 (2014), arXiv:1310.8337 [hep-ph].
- [76] Z.-G. Wang, Analysis of the axialvector doubly heavy tetraquark states with QCD sum rules, *Acta Phys. Polon. B* **49**, 1781 (2018), arXiv:1708.04545 [hep-ph].
- [77] Z.-G. Wang and Z.-H. Yan, Analysis of the scalar, axialvector, vector, tensor doubly charmed tetraquark states with QCD sum rules, *Eur. Phys. J. C* **78**, 19 (2018), arXiv:1710.02810 [hep-ph].
- [78] S. S. Agaev, K. Azizi, B. Barsbay, and H. Sundu, Weak decays of the axial-vector tetraquark $T_{bb;\bar{u}\bar{d}}^-$, *Phys. Rev. D* **99**, 033002 (2019), arXiv:1809.07791 [hep-ph].
- [79] H. Sundu, S. S. Agaev, and K. Azizi, Semileptonic decays of the scalar tetraquark $Z_{bc;\bar{u}\bar{d}}^0$, *Eur. Phys. J. C* **79**, 753 (2019), arXiv:1903.05931 [hep-ph].
- [80] S. S. Agaev, K. Azizi, and H. Sundu, Double-heavy axial-vector tetraquark $T_{bc;\bar{u}\bar{d}}^0$, *Nucl. Phys. B* **951**, 114890 (2020), arXiv:1905.07591 [hep-ph].
- [81] L. Tang, B.-D. Wan, K. Maltman, and C.-F. Qiao, Doubly Heavy Tetraquarks in QCD Sum Rules, *Phys. Rev. D* **101**, 094032 (2020), arXiv:1911.10951 [hep-ph].
- [82] S. S. Agaev, K. Azizi, B. Barsbay, and H. Sundu, Heavy exotic scalar meson $T_{bb;\bar{u}\bar{s}}^-$, *Phys. Rev. D* **101**, 094026 (2020), arXiv:1912.07656 [hep-ph].
- [83] S. S. Agaev, K. Azizi, B. Barsbay, and H. Sundu, Stable scalar tetraquark $T_{bb;\bar{u}\bar{d}}^-$, *Eur. Phys. J. A* **56**, 177 (2020), arXiv:2001.01446 [hep-ph].
- [84] Q.-N. Wang and W. Chen, Fully open-flavor tetraquark states $bc\bar{q}\bar{s}$ and $sc\bar{q}\bar{b}$ with $J^P = 0^+, 1^+$, *Eur. Phys. J. C* **80**, 389 (2020), arXiv:2002.04243 [hep-ph].
- [85] S. S. Agaev, K. Azizi, B. Barsbay, and H. Sundu, A family of double-beauty tetraquarks: Axial-vector state $T_{bb;\bar{u}\bar{s}}^-$, *Chin. Phys. C* **45**, 013105 (2021), arXiv:2002.04553 [hep-ph].
- [86] S. S. Agaev, K. Azizi, B. Barsbay, and H. Sundu, Semileptonic and nonleptonic decays of the axial-vector tetraquark $T_{bb;\bar{u}\bar{d}}^-$, *Eur. Phys. J. A* **57**, 106 (2021), arXiv:2008.02049 [hep-ph].
- [87] S. S. Agaev, K. Azizi, and H. Sundu, Newly observed exotic doubly charmed meson Tcc+, *Nucl. Phys. B* **975**, 115650 (2022), arXiv:2108.00188 [hep-ph].
- [88] K. Azizi and U. Özdem, Magnetic dipole moments of the Tcc+ and ZV++ tetraquark states, *Phys. Rev. D* **104**, 114002 (2021), arXiv:2109.02390 [hep-ph].
- [89] T. M. Aliev, S. Bilmis, and M. Savci, Determination of the spectroscopic parameters of beauty-partners of Tcc from QCD, *Phys. Rev. D* **105**, 074038 (2022), arXiv:2111.01081 [hep-ph].
- [90] U. Özdem, Magnetic moments of the doubly charged axial-vector Tcc++ states, *Phys. Rev. D* **105**, 054019 (2022), arXiv:2112.10402 [hep-ph].
- [91] S. Bilmiş, Mixing angles between the tetraquark states with two heavy quarks within QCD sum rules, *Turk. J. Phys.* **45**, 390 (2021), arXiv:2201.06109 [hep-ph].
- [92] U. Özdem, Magnetic dipole moments of states, *Chin. Phys. C* **46**, 113106 (2022), arXiv:2203.07759 [hep-ph].
- [93] K. Azizi and U. Özdem, Exploring the magnetic dipole moments of $T_{QQ\bar{q}\bar{s}}$ and $T_{QQ\bar{s}\bar{s}}$ states in the framework of QCD light-cone sum rules, *JHEP* **03**, 166, arXiv:2301.07713 [hep-ph].
- [94] W. Detmold, K. Orginos, and M. J. Savage, BB Potentials in Quenched Lattice QCD, *Phys. Rev. D* **76**, 114503 (2007), arXiv:hep-lat/0703009.
- [95] M. Wagner (ETM, Y), Forces between static-light mesons, *PoS LATTICE2010*, 162 (2010), arXiv:1008.1538 [hep-lat].
- [96] G. Bali and M. Hetzenegger (QCDSF), Static-light meson-meson potentials, *PoS LATTICE2010*, 142 (2010), arXiv:1011.0571 [hep-lat].
- [97] P. Bicudo and M. Wagner (European Twisted Mass), Lattice QCD signal for a bottom-bottom tetraquark, *Phys. Rev. D* **87**, 114511 (2013), arXiv:1209.6274 [hep-ph].
- [98] Z. S. Brown and K. Orginos, Tetraquark bound states in the heavy-light heavy-light system, *Phys. Rev. D* **86**, 114506 (2012), arXiv:1210.1953 [hep-lat].
- [99] Y. Ikeda, B. Charron, S. Aoki, T. Doi, T. Hatsuda, T. Inoue, N. Ishii, K. Murano, H. Nemura, and K. Sasaki, Charmed tetraquarks T_{cc} and T_{cs} from dynamical lattice QCD simulations, *Phys. Lett. B* **729**, 85 (2014), arXiv:1311.6214 [hep-lat].
- [100] P. Bicudo, K. Cichy, A. Peters, B. Wagenbach, and M. Wagner, Evidence for the existence of $ud\bar{b}\bar{b}$ and the non-existence of $ss\bar{b}\bar{b}$ and

- $c\bar{c}b\bar{b}$ tetraquarks from lattice QCD, *Phys. Rev. D* **92**, 014507 (2015), arXiv:1505.00613 [hep-lat].
- [101] P. Bicudo, K. Cichy, A. Peters, and M. Wagner, BB interactions with static bottom quarks from Lattice QCD, *Phys. Rev. D* **93**, 034501 (2016), arXiv:1510.03441 [hep-lat].
- [102] A. Francis, R. J. Hudspith, R. Lewis, and K. Maltman, Lattice Prediction for Deeply Bound Doubly Heavy Tetraquarks, *Phys. Rev. Lett.* **118**, 142001 (2017), arXiv:1607.05214 [hep-lat].
- [103] P. Bicudo, J. Scheunert, and M. Wagner, Including heavy spin effects in the prediction of a $\bar{b}\bar{b}ud$ tetraquark with lattice QCD potentials, *Phys. Rev. D* **95**, 034502 (2017), arXiv:1612.02758 [hep-lat].
- [104] P. Bicudo, M. Cardoso, A. Peters, M. Pflaumer, and M. Wagner, $ud\bar{b}\bar{b}$ tetraquark resonances with lattice QCD potentials and the Born-Oppenheimer approximation, *Phys. Rev. D* **96**, 054510 (2017), arXiv:1704.02383 [hep-lat].
- [105] A. Francis, R. J. Hudspith, R. Lewis, and K. Maltman, Evidence for charm-bottom tetraquarks and the mass dependence of heavy-light tetraquark states from lattice QCD, *Phys. Rev. D* **99**, 054505 (2019), arXiv:1810.10550 [hep-lat].
- [106] P. Jannan, N. Mathur, and M. Padmanath, Study of doubly heavy tetraquarks in Lattice QCD, *Phys. Rev. D* **99**, 034507 (2019), arXiv:1810.12285 [hep-lat].
- [107] L. Leskovec, S. Meinel, M. Pflaumer, and M. Wagner, Lattice QCD investigation of a doubly-bottom $\bar{b}\bar{b}ud$ tetraquark with quantum numbers $I(J^P) = 0(1^+)$, *Phys. Rev. D* **100**, 014503 (2019), arXiv:1904.04197 [hep-lat].
- [108] R. J. Hudspith, B. Colquhoun, A. Francis, R. Lewis, and K. Maltman, A lattice investigation of exotic tetraquark channels, *Phys. Rev. D* **102**, 114506 (2020), arXiv:2006.14294 [hep-lat].
- [109] P. Mohanta and S. Basak, Construction of $bb\bar{u}\bar{d}$ tetraquark states on lattice with NRQCD bottom and HISQ up and down quarks, *Phys. Rev. D* **102**, 094516 (2020), arXiv:2008.11146 [hep-lat].
- [110] P. Bicudo, A. Peters, S. Velten, and M. Wagner, Importance of meson-meson and of diquark-antidiquark creation operators for a $\bar{b}\bar{b}ud$ tetraquark, *Phys. Rev. D* **103**, 114506 (2021), arXiv:2101.00723 [hep-lat].
- [111] N. Mathur and M. Padmanath, $\bar{b}\bar{c}q_1q_2$ four-quark states from Lattice QCD, *PoS LATTICE2021*, 443 (2022), arXiv:2111.01147 [hep-lat].
- [112] J. Hoffmann, A. Zimmermann-Santos, and M. Wagner, Inclusion of heavy spin effects in the $ud\bar{b}\bar{b} I(J^P)=0(1^-)$ four-quark channel in the Born-Oppenheimer approximation, *PoS LATTICE2022*, 262 (2023), arXiv:2211.15765 [hep-lat].
- [113] S. Meinel, M. Pflaumer, and M. Wagner, Search for $b\bar{b}u\bar{s}$ and $b\bar{b}c\bar{u}$ tetraquark bound states using lattice QCD, *Phys. Rev. D* **106**, 034507 (2022), arXiv:2205.13982 [hep-lat].
- [114] A. V. Manohar and M. B. Wise, Exotic Q Q anti-q anti-q states in QCD, *Nucl. Phys. B* **399**, 17 (1993), arXiv:hep-ph/9212236.
- [115] M. J. Savage and M. B. Wise, Spectrum of baryons with two heavy quarks, *Phys. Lett. B* **248**, 177 (1990).
- [116] N. Brambilla, A. Vairo, and T. Rosch, Effective field theory Lagrangians for baryons with two and three heavy quarks, *Phys. Rev. D* **72**, 034021 (2005), arXiv:hep-ph/0506065.
- [117] T. D. Cohen and P. M. Hohler, Doubly heavy hadrons and the domain of validity of doubly heavy diquark-anti-quark symmetry, *Phys. Rev. D* **74**, 094003 (2006), arXiv:hep-ph/0606084.
- [118] T. Mehen, Implications of Heavy Quark-Diquark Symmetry for Excited Doubly Heavy Baryons and Tetraquarks, *Phys. Rev. D* **96**, 094028 (2017), arXiv:1708.05020 [hep-ph].
- [119] N. Brambilla *et al.*, Heavy Quarkonium: Progress, Puzzles, and Opportunities, *Eur. Phys. J. C* **71**, 1534 (2011), arXiv:1010.5827 [hep-ph].
- [120] B. Silvestre-Brac, Spectrum and static properties of heavy baryons, *Few Body Syst.* **20**, 1 (1996).
- [121] M. Kalos, D. Levesque, and L. Verlet, Helium at zero temperature with hard-sphere and other forces, *Phys. Rev. A* **9**, 2178 (1974).
- [122] M. C. Gordillo, F. De Soto, and J. Segovia, Diffusion Monte Carlo calculations of fully-heavy multi-quark bound states, *Phys. Rev. D* **102**, 114007 (2020), arXiv:2009.11889 [hep-ph].
- [123] J. Sánchez-Baena, J. Boronat, and F. Mazzanti, Diffusion Monte Carlo methods for spin-orbit-coupled ultracold Bose gases, *Phys. Rev. A* **98**, 053632 (2018).
- [124] I. Kosztin, B. Faber, and K. Schulten, Introduction to the diffusion Monte Carlo method, *Am. J. Phys.* **64**, 633 (1996), arXiv:physics/9702023.
- [125] Y. Bai, S. Lu, and J. Osborne, Beauty-full Tetraquarks, *Phys. Lett. B* **798**, 134930 (2019), arXiv:1612.00012 [hep-ph].
- [126] M. C. Gordillo, F. De Soto, and J. Segovia, Structure of the X(3872) as explained by a diffusion Monte Carlo calculation, *Phys. Rev. D* **104**, 054036 (2021), arXiv:2105.11976 [hep-ph].
- [127] J. M. Alcaraz-Pelegri and M. C. Gordillo, Diffusion Monte Carlo calculations of fully heavy compact hexaquarks, *Phys. Rev. D* **106**, 114028 (2022), arXiv:2205.13886 [hep-ph].
- [128] Y. Ma, L. Meng, Y.-K. Chen, and S.-L. Zhu, Ground state baryons in the flux-tube three-body confinement model using diffusion Monte Carlo, *Phys. Rev. D* **107**, 054035 (2023), arXiv:2211.09021 [hep-ph].
- [129] M. C. Gordillo and J. M. Alcaraz-Pelegri, Asymptotic mass limit of large fully heavy compact multi-quarks, *Phys. Rev. D* **108**, 054027 (2023), arXiv:2307.08408 [hep-ph].
- [130] Y. Ma, L. Meng, Y.-K. Chen, and S.-L. Zhu, Doubly heavy tetraquark states in the constituent quark model using diffusion Monte Carlo method, (2023), arXiv:2309.17068 [hep-ph].
- [131] W.-X. Zhang, H. Xu, and D. Jia, Masses and magnetic moments of hadrons with one and two open heavy quarks: Heavy baryons and tetraquarks, *Phys. Rev. D* **104**, 114011 (2021), arXiv:2109.07040 [hep-ph].
- [132] U. Özdem, Analysis of the $Z_b(10650)$ state based on electromagnetic properties, *Eur. Phys. J. C* **84**, 45 (2024), arXiv:2311.11327 [hep-ph].
- [133] Y.-D. Lei and H.-S. Li, Electromagnetic properties of the T_{cc}^+ molecular states, (2023), arXiv:2312.01332 [hep-ph].
- [134] T.-W. Wu and Y.-L. Ma, Doubly heavy tetraquark multiplets as heavy antiquark-diquark symmetry partners of heavy baryons, *Phys. Rev. D* **107**, L071501 (2023), arXiv:2211.15094 [hep-ph].
- [135] M. Pflaumer, L. Leskovec, S. Meinel, and M. Wagner, Existence and Non-Existence of Doubly Heavy Tetraquark Bound States, *PoS*

- [LATTICE2021](#), 392 (2022), [arXiv:2108.10704 \[hep-lat\]](#).
- [136] N. Mathur and M. Padmanath, $\bar{b}\bar{c}q_1q_2$ four-quark states from Lattice QCD, *PoS LATTICE2021*, 443 (2022), [arXiv:2111.01147 \[hep-lat\]](#).
- [137] C. Alexandrou, J. Finkenrath, T. Leontiou, S. Meinel, M. Pflaumer, and M. Wagner, Evidence for shallow bound states and hints for broad resonances with quark content $\bar{b}\bar{c}ud$ in $B-\bar{D}$ and $B^*-\bar{D}$ scattering from lattice QCD, (2023), [arXiv:2312.02925 \[hep-lat\]](#).
- [138] P. Bicudo, Tetraquarks and pentaquarks in lattice QCD with light and heavy quarks, *Phys. Rept.* **1039**, 1 (2023), [arXiv:2212.07793 \[hep-lat\]](#).