

REAL FORMS OF MINIMAL SL_2 -THREEFOLDS

LUCAS MOULIN

ABSTRACT. We complete the classification of the real forms of almost homogeneous SL_2 -threefolds. More precisely, we use the Luna-Vust theory to determine the real forms of minimal smooth complete SL_2 -varieties containing an orbit isomorphic to SL_2/H , where H is a finite cyclic subgroup of SL_2 . Moreover, we study the rationality and the set of real points of those varieties.

CONTENTS

1. Introduction	1
2. Preliminaries	5
3. Equivariant automorphism groups	9
4. Determination of real forms	12
5. Proof of the Main Theorem	19
Appendix: Diagrams of minimal smooth completions of SL_2/A_k	24
References	28

1. INTRODUCTION

1.1. **Setting.** Let X be a complex algebraic variety. A classical problem in algebraic geometry is to determine the real forms of X , i.e. the real algebraic varieties W such that $W_{\mathbb{C}} := W \times_{\text{Spec}(\mathbb{R})} \text{Spec}(\mathbb{C}) \simeq X$. This problem is usually divided in two parts: whether X admits a real form or not, and if so, determine how many there are (up to isomorphism).

In the present article, we work on a slightly different problem. We are interested in algebraic varieties endowed with an algebraic group action, and thus we only consider real forms which are compatible with this algebraic group action. More precisely, let G be a complex reductive algebraic group and let F be a real algebraic group such that $G \simeq F_{\mathbb{C}}$ as complex algebraic groups. Now let X be a complex algebraic G -variety. We say that W is a *real F -form* of X if $W_{\mathbb{C}} \simeq X$ as complex algebraic G -varieties. When the real algebraic group F is clear from the context, we will just refer to real F -forms as real forms.

An important invariant of G -varieties is the complexity. For $B \subset G$ a Borel subgroup, the *complexity* of a complex algebraic G -variety is defined as the maximal codimension of a B -orbit. Note that this does not depend on the choice of B . Forms over any perfect base field of complexity-zero varieties (more commonly known as spherical varieties) have already been classified by Huruguen [Hur11], Wedhorn

The IMB receives support from the EIPHI Graduate School (contract ANR-17-EURE-0002).

[Wed18], and Borovoi-Gagliardi [BG21]. For T a torus, forms of complexity-one T -varieties have also been studied by Langlois in [Lan15] and by Gillard in [Gil22].

Another invariant of a homogeneous space G/H is the rank, which we denote $\text{rk}(G/H)$. Suppose that $\text{rk}(G/H) \leq 1$. Then, Panyushev proved in [Pan95] that either G/H is a spherical homogeneous space or G/H is obtained from a homogeneous SL_2 -threefold by parabolic induction. In this sense, almost homogeneous SL_2 -threefolds are therefore the easiest case of complexity-one almost homogeneous varieties. Thus, let us now suppose that $G = \text{SL}_2$ and that X is a complex almost homogeneous SL_2 -threefold, i.e. an SL_2 -variety with a dense open orbit $X_0 = \text{SL}_2/H$, where H is a finite subgroup of SL_2 . Note that, since a Borel subgroup of SL_2 is of codimension 1, these varieties are of complexity one. In this setting, the Luna-Vust theory provides a combinatorial description of all the G -equivariant embeddings $X_0 \hookrightarrow X$ for a given X_0 , up to G -equivariant isomorphisms (see [LV83, § 9] for details or §2.2 for an overview).

More specifically, we will study the real forms of minimal smooth completions of G/H . Here we call a *minimal smooth completion* of G/H a smooth complete almost homogeneous SL_2 -threefold, with open orbit G/H , such that any SL_2 -equivariant birational morphism $X \rightarrow X'$, with X' smooth, is an isomorphism. To simplify, we will also refer to a minimal smooth completion as a *model*. Those varieties are well-known, they were first studied by Umemura in [Ume88] as the connected component of their automorphism groups are maximal connected algebraic subgroups of the Cremona group $\text{Bir}(\mathbb{P}^3)$, and the projective ones were then classified by Nakano in [Nak89] as an application of Mori theory. They arise naturally in many ways in algebraic geometry, and the list includes Fano varieties such as \mathbb{P}^3 , the smooth quadric Q_3 and the degree five del Pezzo surface Y_5 , as well as Umemura \mathbb{P}^1 -bundles and the projectivization of Schwarzenberger bundles (see [Ume88] and [BFT23]). From the Luna-Vust theory point of view, the full classification and the description of the corresponding combinatorial data were obtained by Moser-Jauslin [MJ87] when $H \subset \{\pm Id\}$ and by Bousquet [Bou00] in the general case.

Using a generalization of the Luna-Vust theory over arbitrary perfect fields, Moser-Jauslin and Terpereau determined in [MJT] the real forms of the minimal smooth completions of SL_2/H when H is a non-cyclic finite subgroup. In their case, there is always a unique smooth completion which has one or two real F -forms for a fixed F . Here we will use a similar approach to determine the real forms of SL_2/H when H is a cyclic subgroup of order n . However, in this case, a few new difficulties arise compared to the non-cyclic case since the automorphism group $\text{Aut}^G(X)$ can be infinite and there are many more cases to consider. Also the underlying complete G -varieties are not all projective, so we have to investigate more carefully the effectiveness of the Galois descent.

For a cyclic subgroup $H \subset \text{SL}_2$ of cardinal k , H is conjugate to the group A_k generated by $\omega_k = \begin{bmatrix} \zeta_k & 0 \\ 0 & \zeta_k^{-1} \end{bmatrix}$, where ζ_k is a primitive k -th root of unity. A diagrammatic representation of the combinatorial data of all the models of SL_2/H , as well as the correspondence with the description of Nakano, can be found in the [Appendix](#). The list of the projective ones given by Nakano (cf. [Nak89]) is the following:

- 1) **The Umemura \mathbb{P}^1 -bundles** $X_k(n, m) \rightarrow \mathbb{F}_k$, $k \geq 1$, $n, m \geq 0$. We note \mathbb{F}_k the Hirzebruch surface defined by $\mathbb{F}_k := \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-k))$, $p_k : \mathbb{F}_k \rightarrow \mathbb{P}^1$

the projection, and C_∞ the negative section of p_k defined by the projection $\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-k) \rightarrow \mathcal{O}_{\mathbb{P}^1}(-k)$. There is a natural SL_2 -action on \mathbb{F}_k and C_∞ is a 1-dimensional closed orbit. There is a unique homogeneous non-trivial extension (cf. [Ume85])

$$0 \rightarrow \mathcal{O}_{\mathbb{F}_k} \rightarrow E_k(n, m) \rightarrow p_k^*(\mathcal{O}_{\mathbb{P}^1}(2 - nk)) \otimes \mathcal{O}_{\mathbb{F}_k}(-mC_\infty) \rightarrow 0.$$

We get a \mathbb{P}^1 -bundle $\pi_k(n, m) : X_k(n, m) \rightarrow \mathbb{F}_k$ by projectivizing the vector bundle $E_k(n, m) \rightarrow \mathbb{F}_k$. The variety $X_k(n, m)$ is a smooth completion of SL_2/A_k , and it is minimal if and only if $m \neq 1$ and $(n \neq m \text{ if } k = 1)$.

- 2) **The \mathbb{P}^1 -bundles** $W(k) \rightarrow \mathbb{P}^2$, $k \geq 0$. As we just saw, $X_1(k, k)$ is not minimal and we can equivariantly contract $\pi_1(k, k)^{-1}(C_\infty)$ to \mathbb{P}^1 to get a new smooth completion of SL_2/A_k , which we call $W(k)$. The variety $W(k)$ is minimal if and only if $k \neq 2$.
- 3) **The decomposable \mathbb{P}^1 -bundles** $Y(a, b) \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$, $a > b, a + b \leq 0$. We set $Y(a, b) := \mathbb{P}(\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(a, b))$ where $\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(a, b)$ is the line bundle over $\mathbb{P}^1 \times \mathbb{P}^1$ of bidegree (a, b) . We can define an SL_2 -action on $Y(a, b)$ such that the projection $\pi(a, b) : Y(a, b) \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ is equivariant (we consider the diagonal action on $\mathbb{P}^1 \times \mathbb{P}^1$). The variety $Y(a, b)$ is an almost homogeneous threefold of SL_2/A_{-a-b} . It is minimal if and only if $a \neq \pm 1$ and $b \neq \pm 1$.
- 4) **The pull-backs of Schwarzenberger \mathbb{P}^1 -bundles** $Y_0(k)$, $k = 2m > 0$. Given $k = 2m > 0$, define $a = 1 + m$ and $b = 1 - m$. Then we can show that $\mathrm{Ext}^1(\mathcal{O}(a, b), \mathcal{O})$ has a unique SL_2 -invariant 1-dimensional subspace V , which determines an extension

$$0 \rightarrow \mathcal{O} \rightarrow G_0(k) \rightarrow M \rightarrow 0,$$

where $G_0(k)$ is a homogeneous vector bundle of rank 2 and $M := \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(a, b)$. $G_0(k)$ has an SL_2 -linearization such that the above exact sequence is SL_2 -exact. We define $Y_0(k) := \mathbb{P}(G_0(k))$. There is an induced SL_2 -action on $Y_0(k)$ such that the \mathbb{P}^1 -bundle projection $\pi_0(k) : Y_0(k) \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ is equivariant. $Y_0(k)$ has an open orbit equivariantly isomorphic to SL_2/A_k . It is minimal if and only if $k \neq 4$.

- 5) **The decomposable \mathbb{P}^2 -bundles** $\pi(r) : Z(r) \rightarrow \mathbb{P}^1$, $r \neq 1$. Define $E(r) := \mathcal{O}_{\mathbb{P}^1}^{\oplus 2} \oplus \mathcal{O}_{\mathbb{P}^1}(r)$ and $Z(r) := \mathbb{P}(E(r))$. We can define an SL_2 -action on $Z(r)$ such that the projection $\pi(r)$ is equivariant. $Z(r)$ is minimal if and only if $r \neq 1$. For $r \neq 1$, it has an open orbit isomorphic to $\mathrm{SL}_2/A_{|r-1|}$.
- 6) $\mathbb{P}^2 \times \mathbb{P}^1 \simeq \mathbb{P}(R_2) \times \mathbb{P}(R_1)$. We note R_i the unique irreducible representation of SL_2 of dimension $i + 1$. Then $\mathbb{P}(R_2) \times \mathbb{P}(R_1)$ is naturally equipped with a diagonal action of SL_2 . It is a minimal smooth completion of SL_2/A_4 .
- 7) Finally we have two more isolated models, namely $\mathbb{P}^3 \simeq \mathbb{P}(R_1 \oplus R_1)$ and $Q_3 \subset \mathbb{P}^4 \simeq \mathbb{P}(R_1 \oplus R_1 \oplus R_0)$. The group PGL_2 identifies with an open subset of \mathbb{P}^3 . The natural SL_2 -action on PGL_2 extends to the one on $\mathbb{P}^3 \simeq \mathbb{P}(R_1 \oplus R_1)$ so we get a smooth completion of PGL_2 . In a similar way, SL_2 is defined by the equation $ad - bc = 1$ in \mathbb{A}^4 so we set $Q_3 := \{ad - bc = e^2\} \subset \mathbb{P}^4$. It is easy to show that the action of SL_2 on itself by translation extends to Q_3 . Furthermore, Q_3 and \mathbb{P}^3 are both minimal since they have Picard number equal to 1.

1.2. Our results. In this paper, we determine the equivariant real forms of all these models using the Luna-Vust theory. The group SL_2 has two real forms up to

isomorphism, namely $\mathrm{SL}_{2,\mathbb{R}}$ and $\mathrm{Spin}_{3,\mathbb{R}}$. Their set of real points are $\mathrm{SL}_2(\mathbb{R})$ and $\mathrm{SU}_2(\mathbb{C})$ respectively. Therefore, we classify the equivariant real forms for these two groups. Moreover we determine which real forms (in the classical sense) appear as an equivariant real form. Finally, we also determine which equivariant real forms are rational and whether they have real points. Recall that a rational variety always has real points. In this case we show that the converse is true. The main result of this article is the following (see Section 5 for the proof):

Main Theorem. *Let $G = \mathrm{SL}_{2,\mathbb{C}}$, $H \subset G$ a finite cyclic subgroup, and X a complex projective minimal smooth completion of G/H . Then the equivariant real forms of X are either rational, or they have no real points. Furthermore the number of rational real forms and real forms with an empty real locus, up to isomorphism, are listed in the table below for each model X . Each column indicates the following number :*

- (i) Number of real forms.
- (ii) Number of $\mathrm{SL}_{2,\mathbb{R}}$ -real forms.
- (iii) Number of $\mathrm{Spin}_{3,\mathbb{R}}$ -real forms.
- (iv) Number of real forms that appear in columns (ii) or (iii).

Models	Rational real forms				Real forms w/o real points			
	(i)	(ii)	(iii)	(iv)	(i)	(ii)	(iii)	(iv)
$Z(1-k), X_k(n, m), k$ even	1	1	0	1	1	0	1	1
$Z(1-k), X_k(n, m), k$ odd	1	1	0	1	0	0	0	0
$Y(a, -a)$	2	2	1	2	3	2	1	3
$Y(a, b), a + b \neq 0$ even	1	1	0	1	2	0	1	1
$Y(a, b), a + b$ odd	1	1	0	1	2	0	0	0
$Y_0(2)$	2	2	0	2	4	2	0	2
$Y_0(k),$	2	1	1	2	2	1	1	2
$W(1)$	2	2	1	2	1	0	1	1
$W(n+1), n \geq 2$	1	1	0	1	0	0	0	0
$Q_3 \subset \mathbb{P}(R_1 \oplus R_1 \oplus R_0)$	2	1	1	2	1	1	1	1
$\mathbb{P}^3 \simeq \mathbb{P}(R_1 \oplus R_1)$	1	1	1	1	1	1	1	1
$\mathbb{P}(R_2) \times \mathbb{P}(R_1)$	1	1	1	1	1	0	0	0

Remark. Note that for $\mathbb{P}(R_2) \times \mathbb{P}(R_1) \simeq \mathbb{P}^2 \times \mathbb{P}^1$ there are two real forms in total, but only the rational one is equivariant under an action of a real form of SL_2 . Actually, we see that the rational real form is equivariant under an action of both $\mathrm{SL}_{2,\mathbb{R}}$ and $\mathrm{Spin}_{3,\mathbb{R}}$.

Remark. We also determine the equivariant real forms of the nonprojective minimal smooth completions of G/H , however we do not study the rationality nor the real points of those varieties (see Section 4).

1.3. Structure of the article. Section 2 contains preliminary results necessary for the rest of the paper. In Section 2.1, we review real group structures on reductive algebraic groups as well as real structures on varieties endowed with an algebraic

group action. We also explain how they correspond to real forms. Then, in Section 2.2 we see how Luna-Vust theory specializes to classify the equivariant embeddings of homogeneous SL_2 -threefolds, and how a generalization of this theory can be used to classify real forms. In Section 3, we explain how to compute the equivariant automorphism groups of equivariant embeddings of homogeneous spaces using the combinatorial data of the Luna-Vust theory. We then compute those groups for all the minimal smooth completions of SL_2/H , where H is a cyclic subgroup. In Section 4, we determine all the real forms of the SL_2 -threefolds listed previously. Finally, Section 5, we study the rationality and the real locus of the real forms we obtained previously, and then prove the [Main Theorem](#).

1.4. Acknowledgements. I would like to thank my advisors Ronan Terpereau and Lucy Moser-Jauslin for their comments and suggestions, which helped in writing and improving this article.

2. PRELIMINARIES

2.1. Real structures and real forms. We start by recalling the notions of real forms and real structures on a given complex algebraic group G and then on a G -variety, and how the notions are related. More details can be found in [MJT22].

Definition 2.1. Let G be a complex algebraic group.

- (i) A *real form* of G is a pair (F, Θ) with F a real algebraic group and $\Theta : G \rightarrow F_{\mathbb{C}} = F \times_{\text{Spec}(\mathbb{R})} \text{Spec}(\mathbb{C})$ an isomorphism of algebraic groups. (To simplify notation we will drop the isomorphism Θ and simply write that F is a form of G).
- (ii) A *real group structure* σ on G is a scheme involution on G such that the diagram

$$\begin{array}{ccc} G & \xrightarrow{\sigma} & G \\ \downarrow & & \downarrow \\ \text{Spec}(\mathbb{C}) & \xrightarrow{\gamma^* = \text{Spec}(z \mapsto \bar{z})} & \text{Spec}(\mathbb{C}) \end{array}$$

commutes and

$$\iota_G \circ \sigma = \sigma \circ \iota_G \text{ and } m_G \circ (\sigma \times \sigma) = \sigma \circ m_G,$$

where $\iota_G : G \rightarrow G$ is the inverse morphism and $m_G : G \times G \rightarrow G$ is the multiplication morphism.

- (iii) Two real group structures σ and σ' are equivalent if there exists a group automorphism $\varphi \in \text{Aut}(G)$ such that $\sigma' = \varphi \circ \sigma \circ \varphi^{-1}$.
- (iv) If G is a complex algebraic group with a real structure σ , then the set of fixed point $G(\mathbb{C})^\sigma$ is called the *real locus* of (G, σ) .

There is a correspondence between real group structures on G and real forms of G given as follows (see [FSS98, § 1.4]).

- If σ is a real group structure on G , one associates the real algebraic group $G_0 := G/\langle \sigma \rangle$ and the isomorphism Θ given by (q, f) , where $q : G \rightarrow G_0$ is the quotient morphism and $f : G \rightarrow \text{Spec}(\mathbb{C})$ is the structure morphism.
- To a real form F of G , one associates the real group structure given by $\sigma : G \simeq F_{\mathbb{C}} \xrightarrow{\text{Id} \times \text{Spec}(z \mapsto \bar{z})} F_{\mathbb{C}} \simeq G$.

Moreover, two real forms of G are isomorphic (as real algebraic groups) if and only if the corresponding real group structures are equivalent.

Example 2.2. For $G = \mathrm{SL}_2$, let σ_s be the real group structure defined by $\sigma_s(g) = \bar{g}$ where \bar{g} is the complex conjugate of g and let σ_c be the real group structure defined by $\sigma_c(g) = {}^t\sigma_s(g)^{-1}$. Note that $\sigma_c(g) = e\sigma_s(g)e^{-1}$ for all $g \in G$, where $e = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$. The real loci of σ_s and σ_c are respectively $\mathrm{SL}_2(\mathbb{R})$ and $\mathrm{SU}(2)$, and the corresponding real forms are respectively $\mathrm{SL}_{2,\mathbb{R}}$ and $\mathrm{Spin}_{3,\mathbb{R}}$. Also, any real group structure on G is strongly equivalent to either σ_s or σ_c .

We now define real forms and equivariant real structures for a G -variety.

Definition 2.3. Let (G, σ) be a complex algebraic group with a real structure, and let X be a G -variety. We denote by $F = G/\langle\sigma\rangle$ the corresponding real form of G .

- An (\mathbb{R}, F) -form of X is an F -variety Z together with an isomorphism $Z_{\mathbb{C}} \simeq X$ of G -varieties, where G acts on $Z_{\mathbb{C}}$ through $F_{\mathbb{C}} \simeq G$.

- A (G, σ) -equivariant real structure on X is a scheme involution μ on X such that the diagram

$$\begin{array}{ccc} X & \xrightarrow{\mu} & X \\ \downarrow & & \downarrow \\ \mathrm{Spec}(\mathbb{C}) & \xrightarrow{\gamma^* = \mathrm{Spec}(z \mapsto \bar{z})} & \mathrm{Spec}(\mathbb{C}) \end{array}$$

commutes, and such that

$$\forall g \in G(\mathbb{C}), \forall x \in X(\mathbb{C}), \mu(g \cdot x) = \sigma(g) \cdot \mu(x).$$

- Two (G, σ) -equivariant real structure μ_1 and μ_2 are called equivalent if there exists a G -equivariant automorphism φ such that $\mu_2 = \varphi \circ \mu_1 \circ \varphi^{-1}$.
- A (G, σ) -equivariant real structure μ on X is *effective* if X is covered by $\langle\mu\rangle$ -stable affine open subsets. In this case, the categorical quotient $X/\langle\mu\rangle$, which always exists as a real algebraic space, is in fact a real algebraic variety.
- The *real locus* of (G, σ) -equivariant real structure μ on X is the set of fixed points $X(\mathbb{C})^{\mu}$; it identifies with the set of \mathbb{R} -points of the corresponding real form $X/\langle\mu\rangle$. Moreover, it is endowed with an action of $G(\mathbb{C})^{\sigma}$.

As for algebraic groups, there is a bijection between isomorphism classes of real F -forms of X (as real algebraic varieties) and equivalence classes of effective (G, σ) -equivariant real structures on X (see [Bor20, § 5])

Given the existence of a (G, σ) -equivariant real structure, one can use Galois cohomology to parametrize all of them. We note $\Gamma = \mathrm{Gal}(\mathbb{C}/\mathbb{R}) = \{1, \gamma\}$. We define a Γ -group structure on $\mathrm{Aut}^G(X)$ as follows:

$$\mathrm{inn}_{\mu} : \Gamma \times \mathrm{Aut}^G(X) \rightarrow \mathrm{Aut}^G(X), (\gamma, \varphi) \mapsto \mu_{\gamma} \circ \varphi \circ \mu_{\gamma}^{-1}.$$

The next result is then a straightforward consequence of the definition of 1-cocycles.

Proposition 2.4. ([MJT22, Lemma 2.11]) *Let (G, σ) be a complex algebraic group with a real group structure, and let (X, μ_0) be a homogeneous space with a (G, σ) -equivariant real structure. There is a bijection of pointed sets*

$$\begin{array}{ccc} Z^1(\Gamma, \mathrm{Aut}_{\mathbb{C}}^G(X)) & \xrightarrow{\sim} & \{(G, \sigma)\text{-equivariant real structure on } X\} \\ \varphi & \mapsto & \varphi \circ \mu_0 \end{array}$$

Remark 2.5. In fact the correspondence above induces a bijection between the equivalence classes of (G, σ) -equivariant real structure on X and $H^1(\Gamma, \mathrm{Aut}_{\mathbb{C}}^G(X))$. However, we will not use this result in this paper.

2.2. Luna-Vust Theory for equivariant embeddings of SL_2/H . We now recall how the Luna-Vust theory specializes to classify the equivariant embeddings of homogeneous SL_2 -threefolds. This section is inspired by [MJT, Appendix B]. For more information on the Luna-Vust theory in the general case, see [LV83] and [Tim11, § 3].

This description was given by Luna-Vust [LV83, § 9] for equivariant embeddings of SL_2 , and by Moser-Jauslin [MJ87] and Bousquet [Bou00] for equivariant embeddings of SL_2/H .

- *The colors.* We start by describing the set of colors of G/H where $H \subset G = \mathrm{SL}_2$ is a finite subgroup. We fix $B = \begin{bmatrix} * & 0 \\ * & * \end{bmatrix}$ and we denote the coordinate ring of G by $\mathbb{C}[G] = \mathbb{C}[w, x, y, z]/(xy - wz - 1)$ with $g = \begin{bmatrix} x & w \\ z & y \end{bmatrix} \in G$. In general, the set of colors is defined by

$$\mathcal{D}^B = \mathcal{D}^B(G/H) = \{B\text{-stable prime divisors of } G/H\}.$$

In our case, we have a right action of H on $\mathbb{P}^1 \simeq B \backslash G$ and this set identifies with \mathbb{P}^1/H via the map:

$$\begin{aligned} \mathbb{P}^1/H &\xrightarrow{\sim} \mathcal{D}^B \\ [\alpha : \beta]H &\mapsto Z(\alpha w + \beta x)H, \end{aligned}$$

where $Z(\alpha w + \beta x)$ denotes the zero locus of the regular function $\alpha w + \beta x$ in G .

- *The invariant valuations.* We now describe the set

$$\mathcal{V}^G = \mathcal{V}^G(G/H) = \{G\text{-invariant geometric valuations of } \mathbb{C}(G/H)\}.$$

We start with the case where H is trivial. A G -invariant geometric valuation on $\mathbb{C}(G)$ is determined by its restriction on the subset

$$\mathbb{P}^1 \simeq \{\alpha w + \beta x \mid [\alpha : \beta] \in \mathbb{P}^1\} \subset \mathbb{C}[G].$$

Furthermore, we can show that for any $\nu \in \mathcal{V}^G$, there exists $[\alpha_0 : \beta_0] \in \mathbb{P}^1$ and $-1 \leq r = \frac{p}{q} \leq 1$ (with p and q relatively prime and $q \geq 1$) such that $\nu(\alpha_0 w + \beta_0 x) = p$ and $\nu(\alpha w + \beta x) = -q$ for all $[\alpha : \beta] \neq [\alpha_0 : \beta_0]$. Therefore,

$$\mathcal{V}^G \simeq \{\mathbb{P}^1 \times ([-1, 1] \cap \mathbb{Q})\} / \sim$$

where $([\alpha : \beta], -1) \sim ([0 : 1], -1)$ for all $[\alpha : \beta] \in \mathbb{P}^1$. For $j \in \mathbb{P}^1$ and $r \in [-1 : 1] \cap \mathbb{Q}$, we denote by $\nu(j, r)$ the G -invariant geometric valuation corresponding to (j, r) . The valuation corresponding to $(j, -1)$ does not depend on j ; we denote this valuation by ν_0 .

Now for any finite subgroup $H \subset G$, any G -invariant geometric valuations on $\mathbb{C}(G)^H$ extends to a valuation on $\mathbb{C}(G)$. Thus we can also associate a couple (j, r) with $j \in \mathbb{P}^1/H$ and $r \in [-1, 1] \cap \mathbb{Q}$ to a G -invariant geometric valuation on $\mathbb{C}(G)^H$. We again denote by $\nu(j, r)$ the valuation of $\mathcal{V}^G(G/H)$ corresponding to $j \in \mathbb{P}^1/H$ and $r \in [-1, 1] \cap \mathbb{Q}$.

We let $b(j) \in \mathbb{Q}$ be the maximum rational number r such that $\mathcal{V}^G(G/H)$ has a valuation of the form $\nu(j, r)$. For example, for $H = A_k$ we have

$$b([0 : 1]) = b([1 : 0]) = 1$$

$$b(j) = \frac{4}{m} - 1 \text{ if } j \notin \{[0 : 1], [1 : 0]\}$$

where $m = n$ if n is even, and $m = 2n$ if n is odd.

- *The colored data.* Now let X be a G -variety in the G -birational class of G/H , and let $\phi : X_0 \dashrightarrow X$ be a G -equivariant birational map. The pair (X, ϕ) is called a G -model of X_0 . In the general theory, we associate to each G -orbit $Y \subset X$ a pair

- $\mathcal{V}_Y^G = \{\nu_D \in \mathcal{V}^G(X) \mid Y \subset D, \text{ where } D \text{ is a } G\text{-stable prime divisor of } X\}$
 $\subset \mathcal{V}^G(X) \simeq \mathcal{V}^G$; and
- $\mathcal{D}_Y^B = \{D \in \mathcal{D}^B(X) \mid Y \subset D\} \subset \mathcal{D}^B(X) \simeq \mathcal{D}^B$.

Such a pair is called the colored data of the G -orbit Y , and the sets of all those pairs is called the colored equipment of X .

We give a description of all possible colored data in our particular case, i.e. where X is G -equivariant embedding of G/H with $H \subset G = \mathrm{SL}_2$ a finite subgroup. We call the facet of an orbit the set of valuations in $\mathcal{V}_1(G/H) = \mathcal{V}^G(G/H) \setminus \{\nu_0\}$ which dominate the local ring of the closure of the given orbit.

The (non-open) G -orbits are divided into the six following types:

- **Type $\mathcal{C}(j, r)$** (2-dimensional orbit): $j \in \mathbb{P}^1/H$ and $r \in \mathbb{Q} \cap]-1, b(j)[$.
 For this case, $\mathcal{V}_Y^G = \{\nu(j, r)\}$ is a single element of \mathcal{V}_1 , $\mathcal{D}_Y^B = \emptyset$, and the facet of Y is $\{\nu(j, r)\}$.
- **Type $\mathcal{A}_N(j_1, \dots, j_N, r_1, \dots, r_N)$** (orbit isomorphic to \mathbb{P}^1): j_1, \dots, j_N are N distinct elements of \mathbb{P}^1/H , and $r_i \in \mathbb{Q} \cap]-1, b(j_i)[$ for each $1 \leq i \leq N$.
 Suppose also that $\sum_{i=1}^N \frac{1}{1+r_i} \geq 1$. Then $\mathcal{V}_Y^G = \{\nu(j_1, r_1), \dots, \nu(j_N, r_N)\}$ and $\mathcal{D}_Y^B = \mathbb{P}^1/H \setminus \{j_1, \dots, j_N\}$. The facet of Y is

$$\bigcup_{i=1}^N \{\nu(j_i, s_i) \mid -1 < s_i < r_i\} \cup \{\nu(j, s) \mid j \notin \{j_1, \dots, j_N\}, -1 < s \leq b(j)\}.$$
- **Type $\mathcal{AB}(j, r_1, r_2)$** (orbit isomorphic to \mathbb{P}^1): $j \in \mathbb{P}^1/H$, and r_1, r_2 are two rational numbers satisfying $-1 \leq r_1 < r_2 \leq b(j)$. Then $\mathcal{V}_Y^G = \{\nu(j, r_1), \nu(j, r_2)\}$ and $\mathcal{D}_Y^B = \emptyset$. The facet of Y is $\{\nu(j, s) \mid r_1 < s < r_2\}$.
- **Type $\mathcal{B}_+(j, r)$** (orbit isomorphic to \mathbb{P}^1): $j \in \mathbb{P}^1/H$, and $r \in \mathbb{Q} \cap [-1, b(j)[$.
 Then $\mathcal{V}_Y^G = \{\nu(j, r)\}$ and $\mathcal{D}_Y^B = \{j\}$. The facet of Y is $\{\nu(j, s) \mid r < s \leq b(j)\}$.
- **Type $\mathcal{B}_-(j, r)$** (orbit isomorphic to \mathbb{P}^1): $j \in \mathbb{P}^1/H$, and $r \in \mathbb{Q} \cap]0, b(j)[$.
 Then $\mathcal{V}_Y^G = \{\nu(j, r)\}$ and $\mathcal{D}_Y^B = \mathbb{P}^1/H \setminus \{j\}$. The facet of Y is $\{\nu(j, s) \mid r < s \leq b(j)\}$.
- **Type $\mathcal{B}_0(j, r)$** (fixed point): $j \in \mathbb{P}^1/H$, and $r \in \mathbb{Q} \cap]0, b(j)[$. Then $\mathcal{V}_Y^G = \{\nu(j, r)\}$ and $\mathcal{D}_Y^B = \mathbb{P}^1/H$. The facet of Y is $\{\nu(j, s) \mid r < s \leq b(j)\}$.

The facets of the colored data of a G -model do not intersect, and their unions is closed in \mathcal{V}^G . Therefore, we can now represent the colored equipment of equivariant embeddings of homogeneous SL_2 -threefolds. We represent the set $\mathcal{V}^G(G/H)$ by a skeleton diagram, and we represent each orbit by its facet. Since the facets

for orbits of types \mathcal{B}_- , \mathcal{B}_0 and \mathcal{B}_+ are the same, we differentiate them with a sign 0, + or -. See the [Appendix](#) for many examples of such diagrams.

- *Luna-Vust over the field of real numbers.* Finally, we present how the Luna-Vust theory allows us to find real forms of equivariant embeddings of SL_2/H . To do this, Moser-Jauslin and Terpereau defined the following action of the Galois group Γ on the colored equipment of G/H in [MJT, § 4.3], generalizing the action defined by Huruguen [Hur11] in the case of spherical varieties.

Let $\sigma \in \{\sigma_s, \sigma_c\}$ and let μ be a (G, σ) -equivariant real structure on G/H . By Proposition 2.4, we have $\mu(gH) = \sigma(g)tH$ for a certain $t \in G$. Then the action of the Galois group Γ on the set of colors $\mathcal{D}^B(G/H) \simeq \mathbb{P}^1/H$ is given by

$$\begin{cases} [\alpha : \beta]H \mapsto [\bar{\alpha} : \bar{\beta}]tH & \text{if } \sigma = \sigma_s; \text{ and} \\ [\alpha : \beta]H \mapsto [-\bar{\beta} : \bar{\alpha}]tH & \text{if } \sigma = \sigma_c \end{cases}$$

and the Γ -action on $\mathcal{V}^G(G/H)$ is defined by

$$\forall j \in \mathbb{P}^1/H, \forall \nu(j, r) \in \mathcal{V}^G(G/H), \gamma \cdot \nu(j, r) = \nu(\gamma \cdot j, r).$$

Note that this action preserves the type of the G -orbits. It can be visualised on the skeleton diagrams where it corresponds to a permutation of the spokes.

We can now give the following result that determines when an equivariant real structure on a homogeneous space X_0 extends to an effective equivariant real structure on an equivariant embedding of X_0 , using the previously defined Γ -action on the colored equipment of that embedding.

Theorem 2.6. ([MJT, Theorem 4.7]) *Let H be a finite subgroup of $G = \mathrm{SL}_2$, let σ be a real group structure on G corresponding to the real form F of G , and let μ be a (G, σ) -equivariant real structure on $X_0 = G/H$. Let $X_0 \hookrightarrow X$ be a G -equivariant embedding of X_0 . Then μ extends to an effective (G, σ) -equivariant real structure $\tilde{\mu}$ on X , which corresponds to an (\mathbb{R}, F) -form of X through the map $X \mapsto X/\langle \tilde{\mu} \rangle$, if and only if*

- (1) *the Γ -actions on $\mathcal{V}^G(G/H)$ and $\mathcal{D}^B(G/H)$ induced by μ preserves the collection of colored data of the G -orbits of X ; and*
- (2) *every G -orbit of X of type \mathcal{B}_0 or \mathcal{B}_- is stabilized by the Γ -action.*

3. EQUIVARIANT AUTOMORPHISM GROUPS

In this section, we first see how we can use the Luna-Vust theory to compute the equivariant automorphism groups of equivariant embeddings of homogeneous spaces. Then, we apply to the particular case of the minimal smooth completions of SL_2/A_k listed in Section 1.1.

Let k be an algebraic closed field, G a reductive group over k , $H \subset G$ an algebraic subgroup of G and $B \subset G$ a Borel subgroup of G . We denote by K the field $k(G)^H = k(G/H)$.

We have an isomorphism (see [Tim11, Proposition 1.8])

$$\begin{aligned} N_G(H)/H &\longrightarrow \mathrm{Aut}^G(G/H) \\ nH &\longmapsto (\varphi_n : kH \mapsto kn^{-1}H). \end{aligned}$$

We consider two group actions on G/H :

- The action of $A = \mathrm{Aut}^G(G/H) \cong N_G(H)/H$

- The action of G by left multiplication.

Those actions induce actions of A and G on K . Indeed, for $f \in K$, $n \in N_G(H)$ and $g \in G$, we define:

- $(g \cdot f)(x) = f(g^{-1}x)$ and,
- $(\varphi_n(f))(kH) = f(\varphi_n^{-1}(kH)) = f(knH)$.

We can now define an action of A on \mathcal{V}^G and on \mathcal{D}^B .

- Action of A on \mathcal{V}^G : If $\nu \in \mathcal{V}^G$, and if $n \in N_G(H)$, then $((\varphi_n\nu)(f)) = \nu(\varphi_n^{-1}(f))$. Since G acts on K on the left, this does indeed give a G -stable geometric valuation on G/H .
- Action of A on \mathcal{D}^B : Again, since B acts on the left and the action of A is induced by a right action of $N_G(H)$, the action of A on G/H induces an action on B -stable divisors.

Let $X_0 = G/H$. We can now determine when a G -equivariant automorphism of X_0 extends to an automorphism on a G -model of X_0 .

Theorem 3.1. *Let (X, ψ) be a G -model of X_0 , and $\varphi \in A$. Let $\mathbb{F}(X, \psi) = \{(\mathcal{V}^G, \mathcal{D}^B)\}$ be the colored data of (X, ψ) . Then, the automorphism φ extends to an automorphism of X if and only if $\mathbb{F}(X, \psi)$ is preserved by the action of φ on \mathcal{V}^G and \mathcal{D}^B , i.e.*

$$\forall (\mathcal{V}_Y^G, \mathcal{D}_Y^B) \in \mathbb{F}(X, \psi), (\varphi(\mathcal{V}_Y^G), \varphi(\mathcal{D}_Y^B)) \in \mathbb{F}(X, \psi).$$

For the proof, we will need the following result that gives a combinatorial description of the B -charts of X . We recall that a B -chart of a G -variety X is a B -stable affine open subset of X .

Proposition 3.2. [Tim11, Corollary 13.9] *The map*

$$\{B\text{-charts of } X\} \rightarrow \mathfrak{P}(\mathcal{V}_X^G) \times \mathfrak{P}(\mathcal{D}_X^B), \hat{X} \mapsto \bigcup_{\substack{G\text{-orbits } Y \subseteq X, \\ Y \cap \hat{X} \neq \emptyset}} (\mathcal{V}_Y^G, \mathcal{D}_Y^B)$$

is a bijection between the B -charts of X and the pairs $(\mathcal{W}, \mathcal{R}) \in \mathfrak{P}(\mathcal{V}_X^G) \times \mathfrak{P}(\mathcal{D}_X^B)$ satisfying conditions (C), (F), and (W) (See [Tim11, Chapter 13]).

The converse map sends the pair $(\mathcal{W}, \mathcal{R})$ to $\text{Spec}(A[\mathcal{W}, \mathcal{R}])$, where

$$A[\mathcal{W}, \mathcal{R}] = \left(\bigcap_{w \in \mathcal{W}} \mathcal{O}_w \right) \cap \left(\bigcap_{D \in \mathcal{R}} \mathcal{O}_{\nu_D} \right) \cap \left(\bigcap_{D \in \mathcal{D} \setminus \mathcal{D}^B} \mathcal{O}_{\nu_D} \right) \subseteq k(X)$$

and \mathcal{D} is the set of prime divisors on X that are not G -stable.

Proof of the theorem. Let's suppose that φ extends to an automorphism of X . Let Y be a G -orbit of X , then $\varphi(Y)$ is also a G -orbit of X . We have $\varphi(\mathcal{D}_Y^B) = \mathcal{D}_{\varphi(Y)}^B$, and $\varphi(\mathcal{V}_Y^G) = \mathcal{V}_{\varphi(Y)}^G$. Therefore, $(\varphi(\mathcal{V}_Y^G), \varphi(\mathcal{D}_Y^B)) \in \mathbb{F}(X, \psi)$ and the colored data is preserved.

Conversely, assume that the colored data $\mathbb{F}(X, \psi)$ is preserved by the action of φ . Let Y be a G -orbit of X , and let \hat{X} be a B -chart of X such that $\hat{X} \cap Y \neq \emptyset$. By Proposition 3.2, there exists a unique pair $(\mathcal{W}, \mathcal{R})$ such that $\text{Spec}(A[\mathcal{W}, \mathcal{R}]) = \hat{X}$. Now, the pair $(\varphi^{-1}(\mathcal{W}), \varphi^{-1}(\mathcal{R}))$ also satisfies conditions (C), (F) and (W). Since $\mathbb{F}(X, \psi)$ is stable under φ , this means that $\hat{X}' = \text{Spec}(A[\varphi^{-1}(\mathcal{W}), \varphi^{-1}(\mathcal{R})])$ is a B -chart of X .

Now, φ sends $A[\varphi^{-1}(\mathcal{W}), \varphi^{-1}(\mathcal{R})]$ to $A[\mathcal{W}, \mathcal{R}]$ so φ is a regular map $\mathring{X} \rightarrow \mathring{X}'$. Indeed, let $f \in A[\varphi^{-1}(\mathcal{W}), \varphi^{-1}(\mathcal{R})]$ we have :

$$\begin{aligned} \forall \omega \in \mathcal{W}, \omega(\varphi(f)) &= (\varphi^{-1}(\omega))(f) \geq 0 \text{ and} \\ \forall D \in \mathcal{R}, \nu_D(\varphi(f)) &= \nu_{\varphi^{-1}(D)}(f) \geq 0. \end{aligned}$$

Moreover, $\mathcal{D} \setminus \mathcal{D}^B$ is preserved by A so $\varphi(f) \in A[\mathcal{W}, \mathcal{R}]$ by Proposition 3.2.

Thus, the open locus $U_\varphi \subset X$ over which φ is defined contains \mathring{X} . Since, φ is G -equivariant, U_φ actually contains $G \cdot \mathring{X}$, in particular it contains Y . This is true for all G -orbit Y of X so φ is a regular morphism on X . Moreover, since $\varphi^{-1} \in A$ and $\varphi^{-1} \cdot \mathbb{F}(X, \psi) = \varphi^{-1} \cdot (\varphi \cdot \mathbb{F}(X, \psi)) = \mathbb{F}(X, \psi)$, φ^{-1} is also a regular morphism on X . Therefore, φ is an automorphism of X . \square

We now let $k = \mathbb{C}$, $G = \text{SL}_2$, H be a finite cyclic subgroup of G and X be a minimal smooth completion of G/H . The diagrams corresponding to those varieties can be found in the Appendix and the groups $\text{Aut}^G(X_0)$ have been computed in [MJT, Lemma 4.3] for all H . We do not give the equivariant automorphism group in cases (b1) and (b2), because the result depends heavily on the different parameters and is not required for the rest of this paper.

Corollary 3.3.

- If X corresponds to model (c1), (g3) or (i4) with $n = m$, we have

$$\text{Aut}^G(X) = \begin{bmatrix} * & 0 \\ 0 & * \end{bmatrix} / H.$$

- If X corresponds to model (c1), (g3) or (i4) with $n \neq m$, we have

$$\text{Aut}^G(X) = \left(\begin{bmatrix} * & 0 \\ 0 & * \end{bmatrix} \cup \begin{bmatrix} 0 & * \\ * & 0 \end{bmatrix} \right) / H.$$

- For the other varieties, the equivariant automorphism group are given in the following table.

Models	$\text{Aut}^G(X)$
(a1)	SL_2
(a2)	PGL_2
(d1),(f1),(c2)	$\begin{bmatrix} * & * \\ 0 & * \end{bmatrix} / H$
(e1),(h4)	$\left(\begin{bmatrix} * & 0 \\ 0 & * \end{bmatrix} \cup \begin{bmatrix} 0 & * \\ * & 0 \end{bmatrix} \right) / H$
(g1),..., (k1), (d2), (e2), (f2) (a3),..., (f3), (a4),..., (f4)	$\begin{bmatrix} * & 0 \\ 0 & * \end{bmatrix} / H$
(g2)	\mathfrak{S}_3
(g4)	$\mathbb{Z}/2\mathbb{Z}$
(g4')	$\{1\}$

Proof. We use Theorem 3.1. Let $n = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in N_G(H)$. For cases (a1) and (a2), the diagram is preserved by the action of φ_n for all n so we have $\text{Aut}^G(X_0) = \text{Aut}^G(X)$, hence the result.

We have $\varphi_n \cdot [0 : 1]H = [0 : 1]n^{-1}H = [-c : a]H$ and $\varphi_n \cdot [1 : 0]H = [d : -b]H$. Therefore, $[0 : 1]$ (resp. $[1 : 0]$) is fixed by the action of φ_n if and only if $c = 0$ (resp. $b = 0$), and $[0 : 1]$ and $[1 : 0]$ are permuted if and only if $a = d = 0$. This gives us the result for all models except (g2), (g4) and (g4').

For model (g2), the equivariant automorphism group has been computed in [MJT, Example 4.11] so we are left with models (g4) and (g4'). We can now suppose that $k \geq 4$ and is even and so we have $N_G(H)/H = \left(\begin{bmatrix} * & 0 \\ 0 & * \end{bmatrix} \cup \begin{bmatrix} 0 & * \\ * & 0 \end{bmatrix} \right) / H$. Let $\alpha \in \mathbb{C}^*$, $\varphi_n \in \text{Aut}^G(X)$ and suppose that $nH = \begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix} H$. For (g4'), $[0 : 1]$ and $[1 : 0]$ both have to be fixed so n must be of this form. Moreover, we have $\varphi_n \cdot [1 : \alpha]H = [1 : a^2\alpha]H$ so $[1 : \alpha]$ is fixed if and only if $a = \pm 1$, i.e. if and only if $nH = I_2H$.

Now suppose that $nH = \begin{bmatrix} 0 & b \\ -b^{-1} & 0 \end{bmatrix} H$. Then $\varphi_n \cdot [1 : \alpha] = [1 : -b^2\alpha^{-1}]$ so $[1 : \alpha]$ is fixed by the action of φ_n if and only if $b = \pm i\alpha$, i.e. $nH = \begin{bmatrix} 0 & i\alpha \\ i\alpha^{-1} & 0 \end{bmatrix} H$. \square

4. DETERMINATION OF REAL FORMS

Let $G = \text{SL}_2$, H a finite cyclic subgroup of G of order k , $X_0 = G/H$, $\Gamma = \text{Gal}(\mathbb{C}/\mathbb{R}) = \{1, \gamma\}$, and $\sigma \in \{\sigma_s, \sigma_c\}$. H is conjugate to the cyclic subgroup A_k of SL_2 generated by $\omega_k = \begin{bmatrix} \zeta_k & 0 \\ 0 & \zeta_k^{-1} \end{bmatrix}$ where ζ_k is a primitive k -th root of unity. For the rest of this section, we will suppose that $H = A_k$. We also note $e = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \in \text{SL}_2(\mathbb{C})$ et $f = \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix} \in \text{SL}_2(\mathbb{C})$.

In this section, we will classify the equivariant real structures of the minimal smooth completions of X_0 and their equivalence classes. We recall that the diagrams of those models are in the [Appendix](#). We start by the case where H is trivial.

Proposition 4.1. *Let X be a minimal smooth completion of $G = \text{SL}_2$, and let $\sigma \in \{\sigma_s, \sigma_c\}$. Then the following table gives, in each case, a representant μ for each equivalence class of effective (G, σ) -equivariant real structures on X .*

Models	μ for $\sigma = \sigma_s$	μ for $\sigma = \sigma_c$
(a1)	$g \mapsto \sigma_s(g)$	$g \mapsto \sigma_c(g)$ $g \mapsto -\sigma_c(g)$
(c1),(d1),(f1)(g1),(h1),(i1),(j1),(k1)	$g \mapsto \sigma_s(g)$	\emptyset
(e1)	$g \mapsto \sigma_s(g)$ $g \mapsto \sigma_s(g)f$	$g \mapsto \sigma_c(g)$ $g \mapsto -\sigma_c(g)$
(b1) and all the x_i are congruent modulo π	$g \mapsto \pm \sigma_s(g) \begin{bmatrix} e^{-i\pi x} & 0 \\ 0 & e^{i\pi x} \end{bmatrix}$	\emptyset
(b1) and not all the x_i are congruent modulo π	\emptyset	\emptyset

Proof. - Step 1: We start by looking for all the (G, σ) -equivariant real structures on X_0 . For this, we use Proposition 2.4 and so we just have to compute $Z^1(\Gamma, N_G(H)/H)$. In this case, $N_G(H)/H = SL_2$ ([MJT, Lemma 4.3]) and since $\Gamma \simeq \mathbb{Z}/2\mathbb{Z}$ we have

$$Z^1(\Gamma, G) = \{g \in G \mid g^{-1} = \gamma \cdot g\}.$$

Let $g = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL_2$. We start by the case $\sigma = \sigma_s$. Then, $\gamma \cdot g = \begin{bmatrix} \bar{a} & \bar{b} \\ \bar{c} & \bar{d} \end{bmatrix}$, where \bar{x} denotes the complex conjugation of x for $x \in \mathbb{C}$, and $g^{-1} = \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$ so we get

$$\begin{aligned} \gamma \cdot g = g^{-1} &\Leftrightarrow \bar{a} = d, \bar{d} = a, c = -\bar{c}, b = -\bar{b} \\ &\Leftrightarrow \bar{a} = d \text{ and } b, c \in i\mathbb{R}. \end{aligned}$$

Therefore,

$$Z^1(\Gamma, G) = \left\{ \begin{bmatrix} a & ix \\ iy & \bar{a} \end{bmatrix}; x, y \in \mathbb{R}, |a|^2 + xy = 1 \right\}.$$

For $\sigma = \sigma_c$, $\gamma \cdot g = {}^t\sigma_s(g)^{-1} = \begin{bmatrix} \bar{d} & -\bar{c} \\ -\bar{b} & \bar{a} \end{bmatrix}$ so

$$\begin{aligned} \gamma \cdot g = g^{-1} &\Leftrightarrow \bar{d} = d, \bar{a} = a, b = \bar{c}, c = \bar{b} \\ &\Leftrightarrow c = \bar{b} \text{ and } a, d \in \mathbb{R}. \end{aligned}$$

which gives

$$Z^1(\Gamma, G) = \left\{ \begin{bmatrix} a & b \\ \bar{b} & d \end{bmatrix}; a, d \in \mathbb{R}, ad - |b|^2 = 1 \right\}$$

- *Step 2:* We will now see in each case, using Theorem 2.6 which one of those equivariant real structures on X_0 extend to an equivariant real structure on X .

First, for model (a1), the result is immediate in both cases because the diagram is necessarily preserved by the Γ -action on it. Let us also notice that except for model (b1), we only need to look at whether the action of Γ fixes or permutes $[0 : 1]$ and $[1 : 0]$.

- Let us first look at the case where $\sigma = \sigma_s$. Let $t = \begin{bmatrix} a & ix \\ iy & \bar{a} \end{bmatrix} \in Z^1(\Gamma, G)$.

$$\gamma \cdot [0 : 1] = [0 : 1]t = [iy : \bar{a}] \text{ and } \gamma \cdot [1 : 0] = [1 : 0]t = [a : ix].$$

So,

$$\begin{aligned} \gamma \cdot [0 : 1] = [0 : 1] &\Leftrightarrow y = 0 \text{ and } a \neq 0 \\ \text{and } \gamma \cdot [1 : 0] = [1 : 0] &\Leftrightarrow x = 0 \text{ and } a \neq 0. \end{aligned}$$

Therefore, the action defined by μ_t fixes $[0 : 1]$ and $[1 : 0]$ if and only if $t = \begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix}$

with $|a| = 1$.

Furthermore,

$$\begin{aligned} \gamma \cdot [0 : 1] = [1 : 0] &\Leftrightarrow y \neq 0 \text{ and } a = 0 \\ \text{and } \gamma \cdot [1 : 0] = [0 : 1] &\Leftrightarrow x \neq 0 \text{ and } a = 0. \end{aligned}$$

So γ permutes $[1 : 0]$ and $[0 : 1]$ if and only if $t = \begin{bmatrix} 0 & ix \\ ix^{-1} & 0 \end{bmatrix}$ with $x \in \mathbb{R}^*$.

We can now conclude for all models other than model (b1). Indeed, for models (d1) and (f1), γ has to fix $[0 : 1]$ for μ_t to extend effectively to X . For models (c1), (g1), (h1), (i1), (j1) and (k1), γ has to fix both $[0 : 1]$ and $[1 : 0]$. And for model (e1), γ can either permute $[0 : 1]$ and $[1 : 0]$ or fix them.

Now let $m \geq 3$ and consider model (b1). We note $(\omega_1, \dots, \omega_n)$ the elements of \mathbb{P}^1 corresponding to the orbits of type \mathcal{B}_- , with $\omega_1 = [0 : 1]$, $\omega_n = [1 : 0]$ and $\omega_i = [1 : \alpha_i]$ for $i \in \{2, \dots, m-1\}$. We also note $x_i = \arg(\alpha_i)$. For an equivariant real structure on X_0 to extend to an effective equivariant real structure on X , all of the ω_i have to be fixed by the corresponding action. For ω_1 and ω_n to be fixed, we already know that t must be of the form $\begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix}$. For such a t , $\gamma \cdot \omega_i = [1 : a^{-2}\overline{\alpha_i}]$ so $\gamma \cdot \omega_i = \omega_i$ if and only if $a = \pm e^{-i\pi x_i}$. Therefore we have a (G, σ_s) -equivariant real structure on X if and only if all of the x_i are congruent modulo π , in which case we actually have two real structures corresponding to $t = \pm \begin{bmatrix} e^{-i\pi x} & 0 \\ 0 & e^{i\pi x} \end{bmatrix}$.

Let us now look at the case $\sigma = \sigma_c$. Let $t = \begin{bmatrix} a & b \\ \bar{b} & d \end{bmatrix} \in Z^1(\Gamma, G)$. The conditions for μ_t to extend effectively to X are the same as in the case $\sigma = \sigma_s$, but the action is now different :

$$\begin{aligned} \gamma \cdot [0 : 1] &= [-1 : 0]t = [-a : -b] = [a : b] \\ \text{and } \gamma \cdot [1 : 0] &= [0 : 1]t = [\bar{b} : d]. \end{aligned}$$

So $\gamma \cdot [0 : 1] = [0 : 1] \Rightarrow a = 0$, which is impossible since $t \in \text{SL}_2$. This shows that $[0 : 1]$ cannot be fixed by γ . Notice that this is enough to conclude for model (b1) too. Then, we also have that

$$\begin{aligned} \gamma \cdot [0 : 1] &= [1 : 0] \Leftrightarrow a \neq 0 \text{ and } b = 0 \\ \text{and } \gamma \cdot [1 : 0] &= [0 : 1] \Leftrightarrow d \neq 0 \text{ and } b = 0. \end{aligned}$$

So the action defined by μ_t permutes $[0 : 1]$ and $[1 : 0]$ if and only if $t = \begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix}$ with $a \in \mathbb{R}^*$.

To recap, the following table gives in each case for which $t \in Z^1(\Gamma, G)$ the corresponding equivariant real structure $\mu_t : g \mapsto \sigma(g)t$ on $X_0 = G$ extends to an effective (G, σ) -equivariant real structure on X :

Models	$\sigma = \sigma_s$	$\sigma = \sigma_c$
(a1)	$\left\{ \begin{bmatrix} a & ix \\ iy & \bar{a} \end{bmatrix}; x, y \in \mathbb{R}, a ^2 + xy = 1 \right\}$	$\left\{ \begin{bmatrix} a & b \\ \bar{b} & d \end{bmatrix}; a, d \in \mathbb{R}, ad - b ^2 = 1 \right\}$
(e1)	$\left\{ \begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix}; a = 1 \right\} \sqcup \left\{ \begin{bmatrix} 0 & ix \\ ix^{-1} & 0 \end{bmatrix}; x \in \mathbb{R} \right\}$	$\left\{ \begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix}; a = 1 \right\}$
(d1) and (f1)	$\left\{ \begin{bmatrix} a & ix \\ 0 & a^{-1} \end{bmatrix}; x \in \mathbb{R}, a = 1 \right\}$	\emptyset
(c1), (g1), (h1), (i1), (j1) and (k1)	$\left\{ \begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix}; a = 1 \right\}$	\emptyset
(b1) and all the x_i are congruent modulo π	$\pm \begin{bmatrix} e^{-i\pi x} & 0 \\ 0 & e^{i\pi x} \end{bmatrix}$	\emptyset
(b1) and all the x_i are not congruent modulo π	\emptyset	\emptyset

- *Step 3:* Finally, we determine the equivalence classes of the effective equivariant real structures for each model using the results of Corollary 3.3.

For model (a1), $\text{Aut}_{\mathbb{C}}^G(X) = \text{Aut}_{\mathbb{C}}^G(X_0)$, so we have the same results as in the homogeneous case.

Suppose that X corresponds to diagrams (c1),(g1),(h1),(i1),(j1) or (k1). Then we have no (G, σ_c) -equivariant real structures, so we suppose that $\sigma = \sigma_s$. Let $t = \begin{bmatrix} s & 0 \\ 0 & s^{-1} \end{bmatrix}$ with $|s| = 1$ and $n = \begin{bmatrix} x & 0 \\ 0 & x^{-1} \end{bmatrix}$ with x such that $x^{-2} = s$. By Corollary 3.3, $\varphi_n \in \text{Aut}_{\mathbb{C}}^G(X)$. For all $g \in G$, we have $\varphi_n \circ \mu_{Id} \circ \varphi_n^{-1}(g) = \sigma_s(g)t = \mu_t(g)$ so μ_{Id} is equivalent to μ_t . Therefore, all (G, σ_s) -equivariant real structures are therefore equivalent to μ_{Id} .

Suppose now that X corresponds to (d1) or (f1). We have no equivariant real structures for $\sigma = \sigma_c$. Let $\sigma = \sigma_s$, $t = \begin{bmatrix} s & ix \\ 0 & s^{-1} \end{bmatrix}$ with $|s| = 1$ and $x \in \mathbb{R}$. Let $\alpha, \beta \in \mathbb{C}$ such that $\alpha^{-2} = s$ and $\text{Im}(\alpha\bar{\beta}) = \frac{x}{2}$, and let $n = \begin{bmatrix} \alpha & \beta \\ 0 & \alpha^{-1} \end{bmatrix} \in \text{Aut}_{\mathbb{C}}^G(X)$. Then we have $\varphi_n \circ \mu_{Id} \circ \varphi_n^{-1} = \mu_t$ so all (G, σ_s) -equivariant real structures on X are equivalent to the one given by the identity matrix.

For model (e1), we recall that $\text{Aut}_{\mathbb{C}}^G(X) = \begin{bmatrix} * & 0 \\ 0 & * \end{bmatrix} \sqcup \begin{bmatrix} 0 & * \\ * & 0 \end{bmatrix}$. Let $g = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in G$ and let $\sigma = \sigma_c$. For $n = \begin{bmatrix} x & 0 \\ 0 & x^{-1} \end{bmatrix} \in \text{Aut}_{\mathbb{C}}^G(X)$ we have

$$\begin{aligned} \varphi_n \circ \mu_{Id} \circ \varphi_n^{-1}(g) &= \begin{bmatrix} \overline{ax}x^{-1} & \overline{bx^{-1}x} \\ \overline{cx}x^{-1} & \overline{dx^{-1}x} \end{bmatrix} \text{ and} \\ \varphi_n \circ \mu_f \circ \varphi_n^{-1}(g) &= \begin{bmatrix} i\bar{b}|x|^{-2} & i\bar{a}|x|^2 \\ i\bar{d}|x|^{-2} & i\bar{c}|x|^2 \end{bmatrix}. \end{aligned}$$

For $t = \begin{bmatrix} s & 0 \\ 0 & s^{-1} \end{bmatrix}$ with $|s| = 1$, $\mu_t(g) = \begin{bmatrix} \bar{a}s & \bar{b}s^{-1} \\ \bar{c}s & \bar{d}s^{-1} \end{bmatrix}$ so μ_t is equivalent to μ_{Id} (by taking $x^{-2} = s$), and for $t = \begin{bmatrix} 0 & iy \\ iy^{-1} & 0 \end{bmatrix}$ with $y \in \mathbb{R}$, $\mu_t(g) = \begin{bmatrix} i\bar{b}y^{-1} & i\bar{a}y \\ i\bar{d}y^{-1} & i\bar{c}y \end{bmatrix}$ so μ_t is equivalent to μ_f when $y < 0$. Now for $n = \begin{bmatrix} x & 0 \\ 0 & x^{-1} \end{bmatrix} \in \text{Aut}_{\mathbb{C}}^G(X)$ we get

$$\begin{aligned} \varphi_n \circ \mu_{Id} \circ \varphi_n^{-1}(g) &= \begin{bmatrix} \overline{ax}x^{-1} & \overline{bx^{-1}x} \\ \overline{cx}x^{-1} & \overline{dx^{-1}x} \end{bmatrix} \text{ and} \\ \varphi_n \circ \mu_f \circ \varphi_n^{-1}(g) &= - \begin{bmatrix} i\bar{b}|x|^{-2} & i\bar{a}|x|^2 \\ i\bar{d}|x|^{-2} & i\bar{c}|x|^2 \end{bmatrix}. \end{aligned}$$

The second equality shows that μ_t is also equivalent to μ_f for $t = \begin{bmatrix} 0 & iy \\ iy^{-1} & 0 \end{bmatrix}$ when $y > 0$. The previous computations show that μ_{Id} and μ_f are not equivalent.

Now let $\sigma = \sigma_c$, and fix $t = \begin{bmatrix} s & 0 \\ 0 & s^{-1} \end{bmatrix}$ with $|s| = 1$. We have $\mu_t(g) = \begin{bmatrix} s\bar{d} & -s^{-1}\bar{c} \\ -s\bar{b} & s^{-1}\bar{a} \end{bmatrix}$. For $n = \begin{bmatrix} x & 0 \\ 0 & x^{-1} \end{bmatrix}$, $\varphi_n \circ \mu_{Id} \circ \varphi_n^{-1} = \begin{bmatrix} |x|^{-2}\bar{d} & -|x|^2\bar{c} \\ -|x|^{-2}\bar{b} & |x|^2\bar{a} \end{bmatrix}$. Therefore, μ_t is equivalent to μ_{Id} if $s > 0$, and to $-\mu_{Id}$ if $s < 0$. Since those two real

structures are not equivalent in the homogeneous case [MJT], they are also not equivalent on X .

Finally for model (b1), the two classes we found when $\sigma = \sigma_s$ and all the x_i are congruent modulo π are equivalent if and only if $n = \begin{bmatrix} i & 0 \\ 0 & i \end{bmatrix} \in \text{Aut}_{\mathbb{C}}^G(X)$, which is not the case by 3.1. \square

Remark 4.2. Models (b1) and (c1) have other equivariant real structures, but those real structures are not effective, and therefore do not correspond to a real form of X .

Using the same strategy we can also determine the real forms of the minimal smooth completions of SL_2/A_k when $k \geq 2$.

Proposition 4.3. *Let $k = 2$, $\sigma \in \{\sigma_s, \sigma_c\}$ and X be a minimal smooth completions of PGL_2 . Then the following table gives, in each case, a representant μ for each equivalence class of effective (G, σ) -equivariant real structures on X .*

Models	μ for $\sigma = \sigma_s$	μ for $\sigma = \sigma_c$
(a2)	$gH \mapsto \sigma_s(g)H$ $gH \mapsto \sigma_s(g)eH$	$gH \mapsto \sigma_c(g)H$ $gH \mapsto \sigma_c(g)eH$
(c2),(d2),(e2),(f2)	$gH \mapsto \sigma_s(g)H$	$gH \mapsto \sigma_c(g)eH$
(g2)	$gH \mapsto \sigma_s(g)H$ $gH \mapsto \sigma_s(g)fH$	$gH \mapsto \sigma_c(g)eH$ $gH \mapsto \sigma_c(g)efH$
(b2 with $m \geq 3$) and all the x_i are congruent modulo π	$g \mapsto \sigma_s(g) \begin{bmatrix} e^{-i\pi x} & 0 \\ 0 & e^{i\pi x} \end{bmatrix} H$	$g \mapsto \sigma_c(g) \begin{bmatrix} 0 & e^{i\pi x} \\ -e^{-i\pi x} & 0 \end{bmatrix} H$
(b2 with $m \geq 3$) and not all the x_i are congruent modulo π	\emptyset	\emptyset

Proof. As in the trivial case, we start by computing $Z^1(\Gamma, N_G(H)/H)$. Here $N_G(H)/H \simeq \text{PGL}_2$ by [MJT, Lemma 4.3] and we obtain

$$\begin{aligned}
Z^1(\Gamma, G/H) &= \left\{ \begin{bmatrix} a & ix \\ iy & \bar{a} \end{bmatrix} H; x, y \in \mathbb{R}, |a|^2 + xy = 1 \right\} && \text{if } \sigma = \sigma_s \\
&\sqcup \left\{ \begin{bmatrix} a & b \\ c & -\bar{a} \end{bmatrix} H; b, c \in \mathbb{R}, |a|^2 + bc = -1 \right\} \\
Z^1(\Gamma, G/H) &= \left\{ \begin{bmatrix} a & b \\ \bar{b} & d \end{bmatrix} H; a, d \in \mathbb{R}, ad - |b|^2 = 1 \right\} && \text{if } \sigma = \sigma_c \\
&\sqcup \left\{ \begin{bmatrix} ix & b \\ -\bar{b} & iy \end{bmatrix} H; x, y \in \mathbb{R}, |b|^2 - xy = 1 \right\}
\end{aligned}$$

Then we use Theorem 2.6 to find which of those real structures extend to an effective equivariant real structure on X , and we get the following

Models	$\sigma = \sigma_s$	$\sigma = \sigma_c$
(a2)	$\sqcup \left\{ \begin{array}{l} \left[\begin{array}{cc} a & ix \\ iy & \bar{a} \end{array} \right] H; x, y \in \mathbb{R}, a ^2 + xy = 1 \\ \left[\begin{array}{cc} a & b \\ c & -\bar{a} \end{array} \right] H; b, c \in \mathbb{R}, a ^2 + bc = -1 \end{array} \right\}$	$\sqcup \left\{ \begin{array}{l} \left[\begin{array}{cc} a & b \\ \bar{b} & d \end{array} \right] H; a, d \in \mathbb{R}, ad - b ^2 = 1 \\ \left[\begin{array}{cc} ix & b \\ -\bar{b} & iy \end{array} \right] H; x, y \in \mathbb{R}, b ^2 - xy = 1 \end{array} \right\}$
(c2),(b2 with $m = 1$)	$\left\{ \left[\begin{array}{cc} a & ix \\ 0 & a^{-1} \end{array} \right] H; a = 1, x \in \mathbb{R} \right\}$	$\left\{ \left[\begin{array}{cc} 0 & b \\ -b^{-1} & iy \end{array} \right] H; b = 1, y \in \mathbb{R} \right\}$
(d2), (e2) (f2) and (b2 with $m = 2$)	$\left\{ \left[\begin{array}{cc} a & 0 \\ 0 & a^{-1} \end{array} \right] H; a = 1 \right\}$	$\left\{ \left[\begin{array}{cc} 0 & b \\ -b^{-1} & 0 \end{array} \right] H; b = 1 \right\}$
(g2)	$\left\{ I_2 H; \left[\begin{array}{cc} 0 & i \\ i & 0 \end{array} \right] H; \left[\begin{array}{cc} i & i \\ 0 & -i \end{array} \right] H; \left[\begin{array}{cc} i & 0 \\ -i & -i \end{array} \right] H \right\}$	$\left\{ \left[\begin{array}{cc} 0 & 1 \\ -1 & 0 \end{array} \right] H; \left[\begin{array}{cc} -i & 0 \\ 0 & i \end{array} \right] H; \left[\begin{array}{cc} -i & i \\ i & 0 \end{array} \right] H; \left[\begin{array}{cc} 0 & i \\ -i & i \end{array} \right] H \right\}$
(b2 with $m \geq 3$) and all the x_i are congruent modulo π	$\left[\begin{array}{cc} e^{-i\pi x} & 0 \\ 0 & e^{i\pi x} \end{array} \right] H$	$\left[\begin{array}{cc} 0 & e^{i\pi x} \\ -e^{-i\pi x} & 0 \end{array} \right] H$
(b2 with $m \geq 3$) and not all the x_i are congruent modulo π	\emptyset	\emptyset

Finally we use the automorphism groups computed in Corollary 3.3 to compute the equivalence classes of the previous effective equivariant real structures. \square

When H is cyclic of order $k \geq 3$, the diagrams of the minimal smooth completions of SL_2/H have been determined by Bousquet. They depend on whether k is odd or even, we will start by assuming that k is odd. In this case, the diagrams of the minimal smooth completions of SL_2/H , up to conjugacy, are given in Figure 3 of the [Appendix](#).

Proposition 4.4. *Let $k \geq 3$ be an odd integer and X be a minimal smooth completions of SL_2/H . Then there is no effective (G, σ_c) -equivariant real structures on X , and an unique effective (G, σ_s) -equivariant real structures on X up to equivalence: $\mu_{Id} : gH \mapsto \sigma_s(g)H$.*

Proof. First, we have in this case $N_G(H) = \begin{bmatrix} * & 0 \\ 0 & * \end{bmatrix}$ and

$$Z^1(\Gamma, N_G(H)/H) = \begin{cases} \left\{ \left[\begin{array}{cc} a & 0 \\ 0 & a^{-1} \end{array} \right] H; |a| = 1 \right\} & \text{if } \sigma = \sigma_s \\ \sqcup \left\{ \left[\begin{array}{cc} 0 & ix \\ ix^{-1} & 0 \end{array} \right] H; x \in \mathbb{R}^* \right\} \\ \left\{ \left[\begin{array}{cc} x & 0 \\ 0 & x^{-1} \end{array} \right] H; x \in \mathbb{R}^* \right\} & \text{if } \sigma = \sigma_c. \end{cases}$$

Then by Theorem 2.6, there are no effective equivariant real structures on X for σ_c , and the ones for σ_s are given by $tH \in \left\{ \left[\begin{array}{cc} s & 0 \\ 0 & s^{-1} \end{array} \right] H; |s| = 1 \right\}$. We have $\begin{bmatrix} * & 0 \\ 0 & * \end{bmatrix} \subset \text{Aut}_{\mathbb{C}}^G(X)$ by Corollary 3.3. Taking $nH = \begin{bmatrix} x & 0 \\ 0 & x^{-1} \end{bmatrix} \in \text{Aut}_{\mathbb{C}}^G(X)$ with $x^{-2} = s$ gives the equivalence we need. \square

Finally when $k \geq 4$ and is even, the diagrams of the minimal smooth completions of SL_2/H are given, up to conjugacy, in Figure 4 of the [Appendix](#).

Proposition 4.5. *Let $k \geq 4$ be an even integer, $\sigma \in \{\sigma_s, \sigma_c\}$ and X be a minimal smooth completions of SL_2/A_k , Then the following table gives, in each case, a representant μ for each class of effective (G, σ) -equivariant real structures on X .*

Models	μ for $\sigma = \sigma_s$	μ for $\sigma = \sigma_c$
(a4),..., (f4), (i4)	$gH \mapsto \sigma_s(g)H$	$gH \mapsto \sigma_c(g)eH$
(g4) ($x = \arg(\alpha)$)	$gH \mapsto \sigma_s(g) \begin{bmatrix} e^{-i\pi x} & 0 \\ 0 & e^{i\pi x} \end{bmatrix} H$ $gH \mapsto \sigma_s(g) \begin{bmatrix} 0 & i \alpha \\ i \alpha ^{-1} & 0 \end{bmatrix} H$	$gH \mapsto \sigma_c(g) \begin{bmatrix} 0 & e^{i\pi x} \\ -e^{-i\pi x} & 0 \end{bmatrix} H$ $gH \mapsto \sigma_c(g) \begin{bmatrix} i \alpha ^{-1} & 0 \\ 0 & -i \alpha \end{bmatrix} H$
(g4') ($x = \arg(\alpha)$)	$gH \mapsto \sigma_s(g) \begin{bmatrix} e^{-i\pi x} & 0 \\ 0 & e^{i\pi x} \end{bmatrix} H$	$gH \mapsto \sigma_c(g) \begin{bmatrix} 0 & e^{i\pi x} \\ -e^{-i\pi x} & 0 \end{bmatrix} H$
(h4)	$gH \mapsto \sigma_s(g)H$ $gH \mapsto \sigma_s(g)eH$ $gH \mapsto \sigma_s(g)e\omega_{2n}H$	$gH \mapsto \sigma_c(g)H$ $gH \mapsto \sigma_c(g)eH$ $gH \mapsto \sigma_c(g)\omega_{2n}H$

Proof. Here we have

$$\begin{aligned}
Z^1(\Gamma, N_G(H)/H) &= \left\{ \begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix} H; |a| = 1 \right\} && \text{if } \sigma = \sigma_s \\
&\sqcup \left\{ \begin{bmatrix} 0 & ix \\ ix^{-1} & 0 \end{bmatrix} H; x \in \mathbb{R}^* \right\} \\
&\sqcup \left\{ \begin{bmatrix} 0 & ix\zeta_{2k} \\ ix^{-1}\zeta_{2k}^{-1} & 0 \end{bmatrix} H; x \in \mathbb{R}^* \right\} \\
Z^1(\Gamma, N_G(H)/H) &= \left\{ \begin{bmatrix} x & 0 \\ 0 & x^{-1} \end{bmatrix} H; x \in \mathbb{R}^* \right\} && \text{if } \sigma = \sigma_c \\
&\sqcup \left\{ \begin{bmatrix} x\zeta_{2k} & 0 \\ 0 & x^{-1}\zeta_{2k}^{-1} \end{bmatrix} H; x \in \mathbb{R}^* \right\} \\
&\sqcup \left\{ \begin{bmatrix} 0 & b \\ -b^{-1} & 0 \end{bmatrix} H; |b| = 1 \right\}.
\end{aligned}$$

Then the following table gives, in each cases, for which $tH \in Z^1(\Gamma, G/H)$ the corresponding equivariant real structure $\mu_t : g \mapsto \sigma(g)tH$ on $X_0 = G/H$ extends to an effective (G, σ) -equivariant real structure on X :

Models	$\sigma = \sigma_s$	$\sigma = \sigma_c$
(a4),..., (f4), (i4)	$\left\{ \begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix} H; a = 1 \right\}$	$\left\{ \begin{bmatrix} 0 & b \\ -b^{-1} & 0 \end{bmatrix} H; b = 1 \right\}$
(h4)	$Z^1(\Gamma, G/A_k)$	$Z^1(\Gamma, G/A_k)$
(g4) ($x = \arg(\alpha)$)	$\left\{ \begin{bmatrix} e^{-i\pi x} & 0 \\ 0 & e^{i\pi x} \end{bmatrix}; \begin{bmatrix} 0 & i \alpha \\ i \alpha ^{-1} & 0 \end{bmatrix} \right\}$	$\left\{ \begin{bmatrix} 0 & e^{i\pi x} \\ -e^{-i\pi x} & 0 \end{bmatrix}; \begin{bmatrix} i \alpha ^{-1} & 0 \\ 0 & -i \alpha \end{bmatrix} \right\}$
(g4') ($x = \arg(\alpha)$)	$\begin{bmatrix} e^{-i\pi x} & 0 \\ 0 & e^{i\pi x} \end{bmatrix}$	$\begin{bmatrix} 0 & e^{i\pi x} \\ -e^{-i\pi x} & 0 \end{bmatrix}$

As in the previous case, it is relatively straightforward to find the equivalence classes of those real structures with Corollary 3.3. Note that for model (g4) the two real structures we obtained in both cases are not equivalent in the homogeneous case [MJT], so they are not equivalent for X . \square

5. PROOF OF THE MAIN THEOREM

We can now use the previous results to prove the main theorem. In this section, we always consider real structures up to equivalence and real forms up to isomorphism. We say that a real form (in the classical sense) Y of a complex variety is equivariant under an action of $\mathrm{SL}_{2,\mathbb{R}}$ (resp of $\mathrm{Spin}_{3,\mathbb{R}}$) when there is an action of $\mathrm{SL}_{2,\mathbb{R}}$ (resp of $\mathrm{Spin}_{3,\mathbb{R}}$) on Y such that $Y_{\mathbb{C}}$ is an almost homogeneous $\mathrm{SL}_{2,\mathbb{C}}$ -threefold.

We recall that the correspondence between the geometric description and the description from the Luna-Vust point of view is given in the Appendix.

To begin with, note that all the geometrical descriptions given in Section 1.1 are valid over any field of characteristic zero. In particular, all the minimal SL_2 -threefolds have a trivial real form given by the same description. Furthermore, it is easily checked, by a case by case study, that those real forms are all rational.

We first prove the following lemma, which helps us determine when a real form has no real points. Recall that a rational variety always has a real point (in fact a dense subset of real points).

Lemma 5.1. *Let X be a smooth almost homogeneous SL_2 -variety. If the intersection of $X(\mathbb{R})$ with the open orbit of X is empty, then $X(\mathbb{R}) = \emptyset$.*

Proof. By [Man20, Corollary 2.2.10], the real locus of a smooth real variety is either dense or empty. Hence, either $X(\mathbb{R})$ intersects the open orbit of X or $X(\mathbb{R}) = \emptyset$. \square

Remark 5.2. For H a finite subgroup of SL_2 and $\sigma \in \{\sigma_s, \sigma_c\}$, the list of all (G, σ) -equivariant real structures on SL_2/H , as well as the corresponding real loci, can be found in [MJT, Appendix C]. We can therefore use the previous lemma to determine if X has a real point for all minimal smooth completions X of SL_2/H .

We now split the proof of the Main Theorem into a case by case study, starting with the Umemura \mathbb{P}^1 -bundles $X_k(n, m) \rightarrow \mathbb{F}_k$.

Proposition 5.3. *The \mathbb{P}^1 -bundle $X_k(n, m) \rightarrow \mathbb{F}_k$ always has a unique real $\mathrm{SL}_{2,\mathbb{R}}$ -form, which is rational. Moreover, it has no real $\mathrm{Spin}_{3,\mathbb{R}}$ -form when k is odd, and a unique real $\mathrm{Spin}_{3,\mathbb{R}}$ -form, which has no real points, when k is even. Finally, all real forms of $X_k(n, m)$ are equivariant under an action of either $\mathrm{SL}_{2,\mathbb{R}}$ or $\mathrm{Spin}_{3,\mathbb{R}}$.*

Proof. We first suppose that k is odd. Then, by Propositions 4.1 and 4.4, there is only one (G, σ_s) -equivariant real structure on $X_k(n, m)$ and no (G, σ_c) -equivariant real structure. Hence, when k is odd, $X_k(n, m)$ has one real $\mathrm{SL}_{2, \mathbb{R}}$ -form which must be the trivial one and is therefore rational.

On the other hand when k is even, by Propositions 4.3 and 4.5, we have one (G, σ) -equivariant real structure for each $\sigma \in \{\sigma_s, \sigma_c\}$. For σ_c , the real locus of the open orbit is empty by [MJT, Appendix C]. Therefore by Lemma 5.1, when k is even, $X_k(n, m)$ has a rational real $\mathrm{SL}_{2, \mathbb{R}}$ -form and a real $\mathrm{Spin}_{3, \mathbb{R}}$ -form with no real points. Finally, there are no other real forms by [TZ24, Proposition 7.1] and its proof. \square

Proposition 5.4. *The decomposable \mathbb{P}^2 -bundle $Z(1 - k) \rightarrow \mathbb{P}^1$ has a unique real $\mathrm{SL}_{2, \mathbb{R}}$ -form, which is rational, for all k . Moreover, it has no real $\mathrm{Spin}_{3, \mathbb{R}}$ -form when k is odd, and a unique real $\mathrm{Spin}_{3, \mathbb{R}}$ -form, which has no real points, when k is even. Finally, all real forms of $Z(1 - k)$ are equivariant under an action of either $\mathrm{SL}_{2, \mathbb{R}}$ or $\mathrm{Spin}_{3, \mathbb{R}}$.*

Proof. The proof is identical to the case of the Umemura \mathbb{P}^1 -bundles. The non-equivariant real forms are in [TZ24, Proposition 5.3 and Remark 5.4]. \square

Proposition 5.5. *For $n \geq 2$, the \mathbb{P}^1 -bundle $W(n + 1) \rightarrow \mathbb{P}^2$ has a unique real form. This real form is rational and is equivariant under an action of $\mathrm{SL}_{2, \mathbb{R}}$, but not under an action of $\mathrm{Spin}_{3, \mathbb{R}}$. On the other hand, the variety $W(1)$ has three real forms :*

- *The trivial real form which is rational and is equivariant under an action of $\mathrm{SL}_{2, \mathbb{R}}$ but not of $\mathrm{Spin}_{3, \mathbb{R}}$.*
- *A rational real form which is equivariant under an action of both $\mathrm{Spin}_{3, \mathbb{R}}$ and $\mathrm{SL}_{2, \mathbb{R}}$.*
- *A real form with no real points which is equivariant under an action of $\mathrm{Spin}_{3, \mathbb{R}}$ but not of $\mathrm{SL}_{2, \mathbb{R}}$.*

Proof. For $n \geq 2$, there is, by Proposition 4.1, only one (G, σ_s) -equivariant real structure on $W(n + 1)$ and no (G, σ_c) -equivariant real structure. Hence, $W(n + 1)$ only has one real $\mathrm{SL}_{2, \mathbb{R}}$ -form, which is the trivial one and is in particular rational. By [TZ24, Corollary 7.2], there are no other real forms for $W(n + 1)$.

The variety $W(1)$ can also be defined as the complex flag variety $\mathrm{PGL}_{3, \mathbb{C}}/B$ where B is a Borel subgroup $\mathrm{PGL}_{3, \mathbb{C}}$. By [TZ24, Proposition 8.1], this variety has three real forms, two rational ones and one with no real points. We only obtained one real structure with an empty real locus which therefore corresponds to the unique real form with no real points. For the other real structures, the Γ -action given by the real structure defined on the open orbit by $g \mapsto \sigma_s(g)$ acts trivially on the combinatorial data, whereas the other two real structures with a non-empty real locus do not. Therefore the real structure given by $g \mapsto \sigma_s(g)$ corresponds to the trivial real form and the other two real structures correspond to the nontrivial real form. \square

Let us now look at the three isolated models, namely $\mathbb{P}^3 \simeq \mathbb{P}(R_1 \oplus R_1)$, $\mathbb{P}(R_2) \times \mathbb{P}(R_1)$ and $Q_3 \subset \mathbb{P}^4 \simeq \mathbb{P}(R_1 \oplus R_1 \oplus R_0)$.

Proposition 5.6. *The variety $\mathbb{P}^3 \simeq \mathbb{P}(R_1 \oplus R_1)$ has one rational real form and one real form with no real points. Those two varieties are both equivariant under an action of $\mathrm{SL}_{2, \mathbb{R}}$ and an action of $\mathrm{Spin}_{3, \mathbb{R}}$.*

Proof. Let $\sigma \in \{\sigma_s, \sigma_c\}$, by Proposition 4.1 there are two (G, σ) -equivariant real structures, with one of them having an empty real locus. However, it is known that \mathbb{P}^3 has two real forms, one is rational while the other has no real points. Therefore the real $\mathrm{SL}_{2, \mathbb{R}}$ -form with no real points corresponds to the same real form as the real $\mathrm{Spin}_{3, \mathbb{R}}$ -form with no real points. The same can be said for the rational real form of \mathbb{P}^3 . \square

Proposition 5.7. *The variety $\mathbb{P}(R_2) \times \mathbb{P}(R_1)$ has two real forms. One is rational and is equivariant under an action of both $\mathrm{SL}_{2, \mathbb{R}}$ and $\mathrm{Spin}_{3, \mathbb{R}}$, while the other one has no real points and is not equivariant under an action of a real form of $\mathrm{SL}_{2, \mathbb{C}}$.*

Proof. In this case there is one (G, σ_s) -equivariant real structure given by $gH \mapsto \sigma_s(g) \begin{bmatrix} e^{-i\pi x} & 0 \\ 0 & e^{i\pi x} \end{bmatrix} H$. On the open orbit, this real structure is equivalent to $gH \mapsto \sigma_s(g)H$ which has a non-empty real locus. Therefore this real structure corresponds to a real form with real points. Similarly, there is one (G, σ_c) -equivariant real structure which corresponds to a real form with real points. Since $\mathbb{P}^2 \times \mathbb{P}^1 \simeq \mathbb{P}(R_2) \times \mathbb{P}(R_1)$ has only one real form with real points, which is rational, the previous two equivariant real structures correspond to the same rational real form. Moreover, $\mathbb{P}^2 \times \mathbb{P}^1$ has a unique real form with no real points, which is therefore not equivariant under an action of a real form of $\mathrm{SL}_{2, \mathbb{C}}$. \square

Let $r, s \in \mathbb{N}$, we denote by $Q^{r,s}$ the smooth real quadric hypersurface of equation $\{x_1^2 + \dots + x_r^2 - x_{r+1}^2 - \dots - x_{r+s}^2 = 0\} \subset \mathbb{P}^{r+s}$.

Proposition 5.8. *The complex quadric $Q_3 \subset \mathbb{P}^4$ has three real forms :*

- $Q^{5,0}$, which has no real points, and is equivariant both under an action of $\mathrm{SL}_{2, \mathbb{R}}$ and of $\mathrm{Spin}_{3, \mathbb{R}}$.
- $Q^{4,1}$, which is rational, and is equivariant under an action of $\mathrm{Spin}_{3, \mathbb{R}}$ but not of $\mathrm{SL}_{2, \mathbb{R}}$.
- $Q^{3,2}$, which is rational, and is equivariant under an action of $\mathrm{SL}_{2, \mathbb{R}}$ but not of $\mathrm{Spin}_{3, \mathbb{R}}$.

Proof. It is classical that Q_3 has three real forms: $Q^{3,2}$ and $Q^{4,1}$, which are both rational, and $Q^{5,0}$, which has no real points (see e.g. [TZ24, Proposition 10.2]). The equivariant real structures on Q_3 are given in Proposition 4.3 (model (a2)). The real structures corresponding to $g \mapsto \sigma_c(g)eH$ and to $g \mapsto \sigma_s(g)eH$ have an empty real locus and therefore correspond to the real form $Q^{5,0}$. Moreover, by [Nak89, Lemma 1.19], the real form $Q^{4,1}$ is not equivariant under any action of $\mathrm{SL}_{2, \mathbb{R}}$. Hence, the real structure $g \mapsto \sigma_s(g)H$ corresponds to the real form $Q^{3,2}$. Furthermore, the group PSU_2 cannot be embedded into $\mathrm{SO}(3, 2)$, which is the automorphism group of $Q^{3,2}$. Therefore, the real structure given by $g \mapsto \sigma_c(g)H$ corresponds to the real form $Q^{4,1}$. \square

We now consider the decomposable \mathbb{P}^1 -bundles $Y(a, b) \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$.

Proposition 5.9. *Let a and b be two integers such that $a > b, a + b \leq 0$.*

- *If $a + b$ is odd, then $Y(a, b)$ has three real forms. One is rational (the trivial one) and equivariant under an action of $\mathrm{SL}_{2, \mathbb{R}}$ but not of $\mathrm{Spin}_{3, \mathbb{R}}$, while the other two have no real points and are not equivariant under an action of a real form of $\mathrm{SL}_{2, \mathbb{C}}$.*

- If $a + b \neq 0$ and $a + b$ is even, then $Y(a, b)$ has three real forms. The trivial one, which is equivariant under an action of $\mathrm{SL}_{2, \mathbb{R}}$ but not of $\mathrm{Spin}_{3, \mathbb{R}}$, and two without real points: one is equivariant under an action of $\mathrm{Spin}_{3, \mathbb{R}}$ but not of $\mathrm{SL}_{2, \mathbb{R}}$, while the other one is not equivariant under an action of a real form of $\mathrm{SL}_{2, \mathbb{C}}$.
- If $a + b = 0$, then $Y(a, b)$ has five real forms. Two that are equivariant under an action of $\mathrm{SL}_{2, \mathbb{R}}$ but not of $\mathrm{Spin}_{3, \mathbb{R}}$, a rational one and one without real points. Two that are equivariant under an action of $\mathrm{Spin}_{3, \mathbb{R}}$ but not of $\mathrm{SL}_{2, \mathbb{R}}$, a rational one and one without real points. And one without real points that is not equivariant under an action of a real form of $\mathrm{SL}_{2, \mathbb{C}}$.

Proof. We use [TZ24, Propositions 5.7 and 5.12, and Remark 5.8] and its proof to determine all the real forms of $Y(a, b)$ and which ones are rational or without real points. Then, similarly as in the previous cases, we deduce which ones are equivariant under an action of $\mathrm{SL}_{2, \mathbb{R}}$ (respectively of $\mathrm{Spin}_{3, \mathbb{R}}$) from the results of Section 4. \square

It remains to treat the case of the models $Y_0(k)$ with $k \geq 2$ even. The variety $Y_0(2)$ corresponds to $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \simeq \mathbb{P}(R_1) \times \mathbb{P}(R_1) \times \mathbb{P}(R_1)$. This case was studied in [MJT, Example 4.11] and [TZ24, Lemma 5.9]. Recall that $Y_0(k)$ is not minimal.

Proposition 5.10. *Let $k \geq 6$ be an even integer. Then $Y_0(k)$ has one rational $\mathrm{SL}_{2, \mathbb{R}}$ -real form, one rational $\mathrm{Spin}_{3, \mathbb{R}}$ -real form, one $\mathrm{SL}_{2, \mathbb{R}}$ -real form with no real points and one $\mathrm{Spin}_{3, \mathbb{R}}$ -real form with no real points. Moreover, $Y_0(k)$ has no other real form.*

Proof. Those varieties appear as \hat{S}_b in [BFT23] with $b = k/2$, and by Lemma 3.5.5 of loc.cit. we have that $\mathrm{Aut}^\circ(\hat{S}_b) \simeq \mathrm{PGL}_{2, \mathbb{C}}$. Hence, by [TZ24, Proposition 2.20], if X is a real form of \hat{S}_b , then $\mathrm{Aut}^\circ(X)$ is a real form of $\mathrm{PGL}_{2, \mathbb{C}}$, i.e. $\mathrm{Aut}^\circ(X) \simeq \mathrm{PGL}_{2, \mathbb{R}}$ or $\mathrm{Aut}^\circ(X) \simeq \mathrm{SO}_{3, \mathbb{R}}$. However, there is a nontrivial homomorphism from $\mathrm{SL}_{2, \mathbb{R}}$ to $\mathrm{PGL}_{2, \mathbb{R}}$, the natural $2 : 1$ cover, but none to $\mathrm{SO}_{3, \mathbb{R}}$. Similarly, there is a nontrivial homomorphism from $\mathrm{Spin}_{3, \mathbb{R}}$ to $\mathrm{SO}_{3, \mathbb{R}}$, the natural $2 : 1$ cover, but none to $\mathrm{PGL}_{2, \mathbb{R}}$. Therefore, X is equivariant under an action of either $\mathrm{SL}_{2, \mathbb{R}}$ or $\mathrm{Spin}_{3, \mathbb{R}}$.

Recall by Proposition 4.5 that there are four equivariant real structures on $Y_0(k)$:

$$\begin{aligned} \mu_1 : gH \mapsto \sigma_s(g) \begin{bmatrix} e^{-i\pi x} & 0 \\ 0 & e^{i\pi x} \end{bmatrix} H, \quad \mu_2 : gH \mapsto \sigma_s(g) \begin{bmatrix} 0 & i|\alpha| \\ i|\alpha|^{-1} & 0 \end{bmatrix} H, \quad \mu_3 : gH \mapsto \\ \sigma_c(g) \begin{bmatrix} 0 & e^{-i\pi x} \\ -e^{i\pi x} & 0 \end{bmatrix} H \quad \text{and} \quad \mu_4 : gH \mapsto \sigma_c(g) \begin{bmatrix} i|\alpha|^{-1} & 0 \\ 0 & i|\alpha| \end{bmatrix} H. \end{aligned}$$

By Lemma 5.1, μ_2 and μ_4 correspond to real forms with no real points.

Let us prove that the real form corresponding to the real structure μ_1 is rational. By [BFT23, Lemma 3.5.5], we have the following commutative diagram

$$\begin{array}{ccc} & X & \\ \epsilon \swarrow & & \searrow \eta \\ Y_0(k) & \overset{\psi}{\dashrightarrow} & Y\left(\frac{k}{2}, -\frac{k}{2}\right) \\ & \searrow & \swarrow \\ & \mathbb{P}^1 \times \mathbb{P}^1 & \end{array}$$

where all maps are PGL_2 -equivariant, and the morphisms ϵ and η are blow-ups of curves. The morphism $\psi^{-1} \circ \mu \circ \psi$ is a birational antiregular involution on $Y_0(k)$.

By [MJT22, Proposition 3.9], this map is actually an antiregular automorphism because it preserves the skeleton diagram associated to $Y_0(k)$. Therefore, since μ corresponds to a rational real form of $Y(\frac{k}{2}, -\frac{k}{2})$, μ corresponds to a rational real form of $Y_0(k)$; this real form is the trivial one. Similarly, we prove that the real form corresponding to μ_3 is rational. \square

APPENDIX: DIAGRAMS OF MINIMAL SMOOTH COMPLETIONS OF SL_2/A_k




FIGURE 1. List of diagrams of the minimal smooth completions of SL_2 .




FIGURE 2. List of diagrams of the minimal smooth completions of PGL_2 .




FIGURE 3. List of diagrams of the minimal smooth completions of SL_2/A_k , where $k \geq 3$ and k is odd.




FIGURE 4. List of diagrams of the minimal smooth completions of SL_2/A_k , where $k \geq 4$ and k is even.

REFERENCES

- [BFT23] J. Blanc, A. Fanelli, and R. Terpereau. Automorphisms of \mathbb{P}^1 -bundles over rational surfaces. *Épjournal de Géométrie Algébrique*, Volume 6, 2023.
- [Bor20] Mikhail V. Borovoi. Equivariant models of spherical varieties. *Transform. Groups*, 25(2):391–439, 2020.
- [BG21] Mikhail V. Borovoi and Giuliano Gagliardi. Existence of equivariant models of spherical varieties and other G-varieties. *International Mathematics Research Notices*, 07 2021.
- [Bou00] Gilles Bousquet. *Plongements homogènes de $SL(2)$ modulo un sous-groupe fini* (available on <https://tel.archives-ouvertes.fr/tel-02899185>). PhD thesis, 2000.
- [FSS98] Yuval Z. Flicker, Claus Scheiderer, and R. Sujatha. Grothendieck’s on non-abelian H^2 and local-global principles. *J. Am. Math. Soc.*, 11(3):731–750, 1998.
- [Gil22] Pierre-Alexandre Gillard. Torus actions on affine varieties over characteristic zero fields. *arXiv:2208.01495v1*, preprint.
- [Hur11] Mathieu Huruguen. Toric varieties and spherical embeddings over an arbitrary field. *J. Algebra*, 342:212–234, 2011.
- [Lan15] Kevin Langlois. Polyhedral divisors and torus actions of complexity one over arbitrary fields. *J. Pure Appl. Algebra*, 219(6):2015–2045, 2015.
- [LV83] Domingo Luna and Thierry Vust. Plongements d’espaces homogènes. *Comment. Math. Helv.*, 58(2):186–245, 1983.
- [Man20] Frédéric Mangolte. *Real Algebraic Varieties*, in *Springer Monographs in Mathematics*. Springer Nature Switzerland AG 2020
- [MJ87] Lucy Moser-Jauslin. *Normal embeddings of $SL(2)/\Gamma$* . PhD thesis, 1987. University of Geneva.
- [MJT] Lucy Moser-Jauslin and Ronan Terpereau. Forms of almost homogeneous varieties over perfect fields. *arXiv:2008.05197v4*, preprint.
- [MJT22] Lucy Moser-Jauslin and Ronan Terpereau. Real Structures on Horospherical Varieties. *Michigan Mathematical Journal*, 71(2): 283–320, May 2022.
- [Nak89] Tetsuo Nakano. On equivariant completions of 3-dimensional homogeneous spaces of $SL(2, \mathbb{C})$. *Japan. J. Math. (N.S.)*, 15(2):221–273, 1989.
- [Pan95] Dmitri Panyushev. On homogeneous spaces of rank one. *Indag. Math. (N.S.)*, 6(3):315–323, 1995.
- [TZ24] Ronan Terpereau and Suzanna Zimmermann. Real forms of Mori fiber spaces with many symmetries. *arXiv:2403.14493v1*, preprint.
- [Tim11] Dmitry A. Timashev. *Homogeneous spaces and equivariant embeddings*, volume 138 of *Encyclopaedia of Mathematical Sciences*. Springer, Heidelberg, 2011. Invariant Theory and Algebraic Transformation Groups, 8.
- [Ume85] Hiroshi Umemura. On the maximal connected algebraic subgroups of the Cremona group. II. *Algebraic groups and related topics (Kyoto/Nagoya, 1983)*, volume 6 of *Adv. Stud. Pure Math.*, pages 349–436, 1985.
- [Ume88] Hiroshi Umemura. Minimal rational threefolds. II. *Nagoya Math. J.*, 110:15–80, 1988.
- [Wed18] Torsten Wedhorn. Spherical spaces. *Ann. Inst. Fourier (Grenoble)*, 68(1):229–256, 2018.

INSTITUT DE MATHÉMATIQUES DE BOURGOGNE, UMR 5584 CNRS, UNIVERSITÉ DE BOURGOGNE, F-21000 DIJON, FRANCE

Email address: lucas.moulin@u-bourgogne.fr