

Strictly Positive Fragments of the Provability Logic of Heyting Arithmetic

Ana de Almeida Borges*  Joost J. Joosten† 

University of Barcelona

Abstract

We determine the strictly positive fragment $\text{QPL}^+(\text{HA})$ of the quantified provability logic $\text{QPL}(\text{HA})$ of Heyting Arithmetic. We show that $\text{QPL}^+(\text{HA})$ is decidable and that it coincides with $\text{QPL}^+(\text{PA})$, which is the strictly positive fragment of the quantified provability logic of Peano Arithmetic. This positively resolves a previous conjecture of the authors described in [3].

On our way to proving these results, we carve out the strictly positive fragment $\text{PL}^+(\text{HA})$ of the provability logic $\text{PL}(\text{HA})$ of Heyting Arithmetic, provide a simple axiomatization, and prove it to be sound and complete for two types of arithmetical interpretations.

The simple fragments presented in this paper should be contrasted with a recent result by Mojtabehi [42], where an axiomatization for $\text{PL}(\text{HA})$ is provided. This axiomatization, although decidable, is of considerable complexity.

1 Introduction

The provability logic of Heyting Arithmetic (**HA**) has recently been axiomatized by Mojtabehi [42], solving a fifty-year-old problem. This axiomatization is decidable but also quite technically complex.

One way of simplifying a logic L without losing all of its expressive power is to restrict its language to the strictly positive fragment L^+ . This is the fragment containing only \top , propositional or predicate symbols (depending on whether L is propositional or quantified, respectively), conjunctions, universal quantifiers (if L is quantified), and diamonds (if L is modal). Theorems of L^+ are implications between strictly positive formulas. For example, $\forall x \diamond P(x) \vdash_{L^+} \diamond \forall x P(x)$ is a grammatical statement that might or not hold depending on L^+ . There are several examples of pairs $\langle L, L^+ \rangle$ where

*ana.agvb@gmail.com

†jjoosten@ub.edu

L^+ has lower complexity than L (see Table 1 in Section 2.3 for a survey), and for at least some of them L^+ has been actively studied and used in its own right.

In [3], we studied the strictly positive fragment of the quantified provability logic of Peano Arithmetic (PA) and conjectured that it might coincide with the strictly positive fragment of the quantified provability logic of HA. In the present paper, we show that this is indeed the case, by providing a link between a PA-extension and an HA-extension of a given arithmetical realization.

We further study the strictly positive fragment of the (propositional) provability logic of HA, and show that it is quite simple and also coincides with the strictly positive fragment of the provability logic of PA.

In what follows, we introduce the main concepts related to provability logics, providing some more context and history, and end this section with a more detailed plan of the paper.

1.1 Provability logics

The arithmetical predicate $\Box_T(\cdot)$ has been often used ever since Gödel. This predicate is a natural and Σ_1 -complete formalization of the concept of provability in the theory T , in that $\mathbb{N} \models \Box_T(n)$ holds exactly when n is the Gödel number of a formula that is provable in T (as long as T is a reasonable enough theory, for example as long as T has a decidable set of axioms).

The provability logic $\text{PL}(T)$ of a theory T describes the structural behavior of $\Box_T(\cdot)$ as can be proven in T itself. This structural behavior is expressed in a propositional modal logic where, for example, we have $(\Box(p \wedge q) \rightarrow \Box p) \in \text{PL}(T)$, since, for every possible arithmetical reading p^* and q^* of p and q , respectively, we have that $T \vdash \Box_T(p^* \wedge q^*) \rightarrow \Box_T p^*$. In a similar spirit, one defines the set $\text{TPL}(T)$ to be the set of those propositional logical formulas that induce corresponding formulas that are always true on the standard model of arithmetic.

The definition of $\text{PL}(T)$ is of complexity Π_2 (“for all readings \cdot^* , it is provable that φ^* ”), and the definition of $\text{TPL}(T)$ is not even arithmetical (“for all readings \cdot^* , it is true that φ^* ”). However, it turns out that both $\text{PL}(T)$ and $\text{TPL}(T)$ can be generated by the decidable modal logics GL (thus named in reference to Gödel and Löb) and GLS (thus named in reference to GL and Solovay) respectively, for a large class of classical logical theories T , including Peano Arithmetic. See [13] for a more thorough overview.

1.2 Quantified provability logic and its fragments

The quantified provability logic $\text{QPL}(T)$ of a theory T is defined in an analogous way to $\text{PL}(T)$. We restrict ourselves to the languages of predicate logic without function symbols or identity.

As an example, the converse Barcan formula is valid:

$$(\Box_T \forall x \varphi(x) \rightarrow \forall x \Box_T \varphi(x)) \in \text{QPL}(T),$$

but the Barcan formula is not:

$$(\forall x \Box_T \varphi(x) \rightarrow \Box_T \forall x \varphi(x)) \notin \text{QPL}(T).$$

Recall that, in the classical case, $\text{PL}(T)$ allows for a reduction from a Π_2 -definition to a decidable one. Vardanyan proved in [51] that this is not the case for $\text{QPL}(\text{PA})$, which is actually Π_2 -complete. George Boolos and Vann McGee [14] proved that $\text{TQPL}(\text{PA})$ is Π_1 -complete in the set of true arithmetic. Vardanyan's result allows for various generalizations [52, 12, 55] and for a few exceptions of very limited range [6, 58, 27].

In [3], the strictly positive fragment $\text{QPL}^+(\text{PA})$ of the quantified provability logic of PA was presented and proven to be decidable, axiomatized by a logic called QRC_1 . The same paper proved soundness of QRC_1 for HA and conjectured completeness. The current paper settles that conjecture positively and proves completeness of QRC_1 for HA for two different arithmetical readings: the *finitary reading* and the *infinitary reading*.

The finitary reading translates predicates to arithmetical formulas with the same arity, as is usual in the case of quantified provability logic. The infinitary reading falls in the tradition so-called reflection calculi. Propositional reflection calculus translates propositional variables to elementary axiomatizations of theories. Generalizing this to predicate logic, the infinitary reading translates a predicate S to the axiomatization of a collection of theories parametrized by a number of parameters equal to the arity of S .

1.3 Constructive provability logics

For about fifty years, the provability logic of Heyting Arithmetic remained an open problem. Recently, Mojtabehi presented a solution, with a complete axiomatization of $\text{PL}(\text{HA})$ described in [42]. Though this axiomatization is proven to be decidable, it is of considerable complexity. To give an impression of the nature of $\text{PL}(\text{HA})$, we mention an incomplete selection of landmark results here and refer the reader to [4] for a more detailed overview.

De Jongh proved in [16] the non-trivial result that the propositional fragment of $\text{PL}(\text{HA})$ is exactly IPC , the set of intuitionistically valid propositional formulas.¹ It was easy to observe that $\text{iGL} \subseteq \text{PL}(\text{HA})$, where iGL is exactly as GL except that it is defined over IPC instead of over the classical propositional logic CPC .

Since HA has different meta-mathematical properties than PA , different provability logical principles were expected to be found. For example, Markov's rule is admissible for HA in that $\text{HA} \vdash \neg\neg\pi$ implies $\text{HA} \vdash \pi$ whenever $\pi \in \Pi_2$. Then the admissibility of Markov's rule translates to $(\Box\neg\neg\Box\varphi \rightarrow \Box\Box\varphi) \in \text{PL}(\text{HA})$, since $\Box_{\text{HA}}\varphi$ is always Σ_1 regardless of φ . More generally:

$$\left(\Box\neg\neg \left(\Box\varphi \rightarrow \bigvee_i \Box\varphi_i \right) \rightarrow \Box \left(\Box\varphi \rightarrow \bigvee_i \Box\varphi_i \right) \right) \in \text{PL}(\text{HA}).$$

¹The result is not as stable as in the classical logical case and, for example, the propositional fragment of the provability logic of HA together with Markov's principle and with Church's thesis is unknown.

There is actually a more general relationship between $\text{PL}(\text{HA})$ and admissible rules. Visser showed that the admissible rules of HA and the admissible rules of IPC are the same so that each admissible rule $\frac{\varphi}{\psi}$ of HA gives rise to $(\Box\varphi \rightarrow \Box\psi) \in \text{PL}(\text{HA})$. Although Rybakov showed in [46] that the collection of admissible rules of IPC is decidable, it was only later that concrete characterizations were found. In particular, Iemhoff showed in [29] how the admissible rules can be defined in terms of Visser's rule.

The exact definition of Visser's rule is not of real importance in the context of this paper. However, we wish to convey an impression of the complexity of $\text{PL}(\text{HA})$ and as such we find it instructive to formulate it. For an arbitrary natural number n , using the abbreviation $(\varphi)(\psi_1, \dots, \psi_n) := (\varphi \rightarrow \psi_1) \vee \dots \vee (\varphi \rightarrow \psi_n)$ and letting $\alpha := \bigwedge_{i=1}^n (\epsilon_i \rightarrow \zeta_i)$, Visser's rule (V_n) is defined as:

$$\frac{(\alpha \rightarrow (\beta \vee \gamma)) \vee \delta}{(\alpha)(\epsilon_1, \dots, \epsilon_n, \beta, \gamma) \vee \delta},$$

leading to:

$$(\Box((\alpha \rightarrow (\beta \vee \gamma)) \vee \delta) \rightarrow \Box((\alpha)(\epsilon_1, \dots, \epsilon_n, \beta, \gamma) \vee \delta)) \in \text{PL}(\text{HA}).$$

So far we have seen how various meta-mathematical properties led to new principles of the form $(\Box\varphi \rightarrow \Box\psi)$ in $\text{PL}(\text{HA})$. However, not every meta-mathematical property of HA directly translates into a principle in $\text{PL}(\text{HA})$. For example, Myhill [43] and Friedman [20] observed that the disjunction property of HA is not provable in HA itself, whence $(\Box(p \vee q) \rightarrow \Box p \vee \Box q) \notin \text{PL}(\text{HA})$. Notwithstanding this negative result, Leivant could prove in [39, 54] a weak version of the disjunction property to the extent that $(\Box(p \vee q) \rightarrow \Box(p \vee \Box q)) \in \text{PL}(\text{HA})$, whence also $(\Box(p \vee q) \rightarrow \Box(\Box p \vee \Box q)) \in \text{PL}(\text{HA})$.

Visser characterized the closed fragment of $\text{PL}(\text{HA})$ in [53] and the "logic" of Σ_1 -substitutions of HA in [54]. The latter turned out to be important for Mojtehed's result, in a sense, reflecting that the substitutions in Solovay's completeness proof are Boolean combinations of Σ_1 -formulas. In [42], Mojtaba Mojtehed could thus include and extend all the previous results, culminating in a complete and decidable axiomatization of $\text{PL}(\text{HA})$. This axiomatization is iGL together with axioms of the form $\Box\varphi \rightarrow \Box\psi$, where φ and ψ satisfy a complicated yet decidable condition.

The first result of this paper is a simple PTIME -decidable fragment of $\text{PL}(\text{HA})$: the strictly positive fragment $\text{PL}(\text{HA})^+$. The second result of this paper is that the first result can be extended to the quantified setting, leading to a decidable (although to our knowledge not in PTIME) fragment of $\text{QPL}(\text{HA})$. Since the propositional logic $\text{PL}(\text{HA})$ was already such a challenge, there has been hardly any work done on the quantified provability logic $\text{QPL}(\text{HA})$ of HA before now.

1.4 Plan of the paper

In Section 2 we introduce the most relevant modal logics for this paper, both propositional and quantified. For each modal logic, we define its corresponding strictly positive

fragment. As the main contribution of this section, we display some literature findings showing how in general a shift from the full logic to the strictly positive fragment comes with a drop in computational complexity of the problem of theoremhood.

In Section 3 we revisit relational and arithmetical semantics for modal provability logics for both the propositional and the predicate logical case. We state the main theorems relating the logics and their semantics. All of the results in this section are known, but they are needed for the sequel. The only novel detail is the exposition of QRC_1 in terms of Kripke sheaves as in [1].

In Section 4 we prepare for a move to the arithmetical semantics of provability logics for intuitionistic theories by first analyzing the behavior of fragments of the arithmetical hierarchy in the constructive setting.

In Section 5 we revisit the main ingredients of the arithmetical completeness proof for provability logic and observe that the needed substitutions are of low and constructively well-behaved complexity.

Sections 6 and 7 contain the main results of the paper, providing an axiomatization of the strictly positive fragment of the provability logic of Heyting Arithmetic, both in the propositional and in the quantified case.

Finally, Section 8 closes off with some observations on the interpolation and the fixpoint theorem for these fragments.

2 Modal logics and strictly positive fragments

In this section we introduce the most relevant modal logics for this paper. These are mostly related to the provability logic GL (see [13]). We dwell on both propositional logics and on their corresponding predicate logical versions. For each logic, we define its strictly positive fragment and display some findings from the literature showing how in general a shift from the full logic to its strictly positive fragment comes with a drop in the computational complexity of theoremhood.

2.1 Propositional modal logics

The language of propositional modal logic \mathcal{L}_\square is obtained from the language of propositional logic by extending it with a unary \square symbol:

$$\varphi ::= \perp \mid p \mid (\varphi \rightarrow \varphi) \mid (\varphi \wedge \varphi) \mid (\varphi \vee \varphi) \mid \square\varphi,$$

where p represents a propositional symbol. In classical logic, \perp and \rightarrow suffice to define \wedge and \vee , but not so in constructive logic. Negation is defined as usual, with $\neg\varphi := \varphi \rightarrow \perp$, and the diamond operator is defined as the dual of the box: $\diamond\varphi := \neg\square\neg\varphi$.

The only (non-strictly positive) modal logics that we consider in this paper are all normal modal logics. Thus, they are an extension of the basic normal modal logic K . The modal logic K has as axioms all propositional logical tautologies in \mathcal{L}_\square and all

distribution axioms of the form $\Box(\varphi \rightarrow \psi) \rightarrow (\Box\varphi \rightarrow \Box\psi)$. The rules are *modus ponens* and necessitation, to wit, $\mathbf{K} \vdash \varphi \implies \mathbf{K} \vdash \Box\varphi$.

The modal logic **K4** is defined from **K** by adding all the transitivity axioms (also called **4**), which are of the form $\Box\varphi \rightarrow \Box\Box\varphi$. The modal logic **GL** is defined by adding Löb's axiom scheme $\Box(\Box\varphi \rightarrow \varphi) \rightarrow \Box\varphi$ to **K4** (or, equivalently, to **K**).

For a given ordinal Λ , the modal logic \mathbf{GLP}_Λ [30, 7] with language $\mathcal{L}_{[\alpha], \alpha < \Lambda}$ has a modality $[\alpha]$ for each $\alpha < \Lambda$ (with corresponding dual $\langle \alpha \rangle$) and is defined such that each modality $[\alpha]$ behaves like the \Box of **GL** and furthermore the following two axiom schemes are available, both with $\beta < \alpha$: monotonicity $[\beta]\varphi \rightarrow [\alpha]\varphi$; and negative introspection $\langle \beta \rangle\varphi \rightarrow [\alpha]\langle \beta \rangle\varphi$.

2.2 Quantified modal logics

The language of modal logic can be extended to a quantified (first-order) setting in the natural way. We describe a simple version with no equality nor function symbols, $\mathcal{L}_{\Box, \forall}$. Fix a signature Σ , which in this case is simply a set of relation symbols \mathbf{REL}_Σ and their respective arities (we mostly omit specific mentions of Σ). Then $\mathcal{L}_{\Box, \forall}$ is defined as follows:

$$\varphi ::= \perp \mid S(\mathbf{x}) \mid (\varphi \rightarrow \varphi) \mid (\varphi \wedge \varphi) \mid (\varphi \vee \varphi) \mid \forall x \varphi \mid \exists x \varphi \mid \Box\varphi,$$

where $S \in \mathbf{REL}_\Sigma$ is a relation symbol with arity n , \mathbf{x} is an n -tuple of variables, and x is a variable. Again, $\neg\varphi := \varphi \rightarrow \perp$ and $\Diamond\varphi := \neg\Box\neg\varphi$. In classical settings, conjunction, disjunction and existential quantification can be defined from the other connectives.

The free variables of a formula φ are defined as usual, and denoted by $\text{fv}(\varphi)$. The expression $\varphi[x \leftarrow y]$ denotes the formula φ with all free occurrences of the variable x simultaneously replaced by the variable y . We say that y is free for x in φ if no occurrence of y becomes bound in $\varphi[x \leftarrow y]$.

Given a propositional modal logic L , one can define its quantified version QL as described in [28] by adding the following axiom scheme and rule to the first order formulation of L : $\forall x \varphi \rightarrow \varphi[x \leftarrow y]$ with y free for x in φ (**VE**); and $QL \vdash \varphi \rightarrow \psi \implies QL \vdash \varphi \rightarrow \forall x \psi$ with $x \notin \text{fv}(\varphi)$ (**VI**). This leads to the straightforward definition of logics such as **QK**, **QK4**, and **QGL**.

As is to be expected, quantified modal logics such as **QK** and **QK4** are undecidable (they typically contain plain predicate logic). Even small fragments of quantified modal logics are undecidable, such as the fragment with only two variables [32] and the one with only one monadic predicate symbol and two variables [45]. There are, however, some decidable fragments of these logics discussed in [57]. Another decidable fragment is given by the strictly positive quantified modal logic **QRC**₁ [2], further described below.

2.3 Strictly positive logics

Propositional strictly positive formulas (abbreviated p.s.p. formulas) are built up from \top , propositional variables, the binary \wedge and the unary \Diamond (or $\langle \alpha \rangle$ in the polymodal

setting). Thus, the strictly positive propositional language \mathcal{L}_{\square}^+ is defined as $\varphi ::= \top \mid p \mid (\varphi \wedge \varphi) \mid \Diamond \varphi$, and the polymodal version $\mathcal{L}_{[\alpha], \alpha < \Lambda}^+$ is defined as $\varphi ::= \top \mid p \mid (\varphi \wedge \varphi) \mid \langle \alpha \rangle \varphi$, with $\alpha < \Lambda$. We extend the definition to strictly positive quantified modal formulas over some given relational vocabulary Σ as $\varphi ::= \top \mid S(\mathbf{t}) \mid (\varphi \wedge \varphi) \mid \forall x \varphi \mid \Diamond \varphi$, where S is in REL_{Σ} , and denote this language by $\mathcal{L}_{\square, \forall}^+$. For practical reasons related with the study of QRC_1 , the signature Σ may allow for individual constants in the strictly positive case, and so terms \mathbf{t} are either variables or constants. We do not work with function symbols.

Note that the above are the only available connectives. The absence of implication and negation does not allow us to define any of the other ones. The statements of a strictly positive logic are implications between two strictly positive formulas, typically written $\varphi \vdash \psi$ (where φ, ψ are strictly positive) instead of $\varphi \rightarrow \psi$ to remind ourselves that the implication cannot be nested.

An important propositional strictly positive modal logic in the context of this paper is the Reflection Calculus RC_{Λ} , defined as follows.

Definition 2.1 (RC_{Λ} [15, 8, 19]). For φ, ψ and χ formulas in the language $\mathcal{L}_{[\alpha], \alpha < \Lambda}^+$ and $\alpha, \beta < \Lambda$, the axioms and rules of the Reflection Calculus (denoted by RC_{Λ}) are the following:

- (i) $\varphi \vdash_{\text{RC}_{\Lambda}} \varphi$ and $\varphi \vdash_{\text{RC}_{\Lambda}} \top$;
- (ii) $\varphi \wedge \psi \vdash_{\text{RC}_{\Lambda}} \varphi$ and $\varphi \wedge \psi \vdash_{\text{RC}_{\Lambda}} \psi$ (conjunction elimination);
- (iii) if $\varphi \vdash_{\text{RC}_{\Lambda}} \psi$ and $\varphi \vdash_{\text{RC}_{\Lambda}} \chi$, then $\varphi \vdash_{\text{RC}_{\Lambda}} \psi \wedge \chi$ (conjunction introduction);
- (iv) if $\varphi \vdash_{\text{RC}_{\Lambda}} \psi$ and $\psi \vdash_{\text{RC}_{\Lambda}} \chi$, then $\varphi \vdash_{\text{RC}_{\Lambda}} \chi$ (cut);
- (v) if $\varphi \vdash_{\text{RC}_{\Lambda}} \psi$, then $\langle \alpha \rangle \varphi \vdash_{\text{RC}_{\Lambda}} \langle \alpha \rangle \psi$ (necessitation);
- (vi) $\langle \alpha \rangle \langle \alpha \rangle \varphi \vdash_{\text{RC}_{\Lambda}} \langle \alpha \rangle \varphi$ (transitivity);
- (vii) $\langle \alpha \rangle \varphi \vdash_{\text{RC}_{\Lambda}} \langle \beta \rangle \varphi$, for $\alpha > \beta$ (monotonicity);
- (viii) $\langle \alpha \rangle \varphi \wedge \langle \beta \rangle \psi \vdash_{\text{RC}_{\Lambda}} \langle \alpha \rangle (\varphi \wedge \langle \beta \rangle \psi)$ for $\alpha > \beta$ (negative introspection).

The logic RC_{Λ} is also called the strictly positive fragment of GLP_{Λ} since one can prove that for any p.s.p. formulas φ and ψ :

$$\text{GLP}_{\Lambda} \vdash \varphi \rightarrow \psi \iff \varphi \vdash_{\text{RC}_{\Lambda}} \psi.$$

See [15, 8, 19, 11] for details.

The logic RC_1 has just one modality and is proven in [15] to be the strictly positive fragment of any modal logic between K4 and GL , inclusive.

In general, when considering a strictly positive logic P , one may ask whether there is some modal logic L whose strictly positive fragment L^+ coincides with P . In that case we would have:

$$\varphi \vdash_P \psi \iff L \vdash \varphi \rightarrow \psi, \tag{2.1}$$

where φ and ψ are strictly positive formulas. In the remainder of this paper, given a logic L we shall denote its strictly positive fragment by L^+ .

Definition 2.2. For a modal logic L the *strictly positive fragment* L^+ of L is defined as:

$$L^+ := \{\langle \varphi, \psi \rangle \mid L \vdash \varphi \rightarrow \psi \text{ and } \varphi \text{ and } \psi \text{ are strictly positive formulas}\}.$$

One can also consider strictly positive fragments for quantified modal logics. Of principal interest in this paper is the Quantified Reflection Calculus with one modality QRC_1 , which is an extension of RC_1 .

Definition 2.3 (QRC_1 [2]). Let φ , ψ , and χ be strictly positive formulas in $\mathcal{L}_{\square, \forall}^+$ with signature Σ . The axioms and rules of QRC_1 are the following:

- (i) $\varphi \vdash \top$ and $\varphi \vdash \varphi$;
- (ii) $\varphi \wedge \psi \vdash \varphi$ and $\varphi \wedge \psi \vdash \psi$ (conjunction elimination);
- (iii) if $\varphi \vdash \psi$ and $\varphi \vdash \chi$, then $\varphi \vdash \psi \wedge \chi$ (conjunction introduction);
- (iv) if $\varphi \vdash \psi$ and $\psi \vdash \chi$, then $\varphi \vdash \chi$ (cut);
- (v) if $\varphi \vdash \psi$, then $\diamond\varphi \vdash \diamond\psi$ (necessitation);
- (vi) $\diamond\diamond\varphi \vdash \diamond\varphi$ (transitivity);
- (vii) if $\varphi \vdash \psi$, then $\varphi \vdash \forall x\psi$, for $x \notin \text{fv}(\varphi)$ (quantifier introduction on the right);
- (viii) if $\varphi[x \leftarrow t] \vdash \psi$, then $\forall x\varphi \vdash \psi$, for t free for x in φ (quantifier introduction on the left);
- (ix) if $\varphi \vdash \psi$, then $\varphi[x \leftarrow t] \vdash \psi[x \leftarrow t]$, for t free for x in φ and ψ (term instantiation);
- (x) if $\varphi[x \leftarrow c] \vdash \psi[x \leftarrow c]$, then $\varphi \vdash \psi$, for c not in φ nor ψ (constant elimination).

Analogously to the propositional case, QRC_1 is the strictly positive fragment of any logic between QK4 and QGL , inclusive, and also the strictly positive fragment of the quantified provability logic of Peano Arithmetic [3]. In this paper, we show that QRC_1 is the strictly positive fragment of the quantified provability logic of Heyting Arithmetic as well.

Moving from a logic to its strictly positive fragment generally leads to a reduction in the computational complexity of the corresponding decision procedures. Many such reductions are listed in Table 1, although the strictness of most reductions depends on complexity theoretical assumptions such as $\text{PSPACE} \neq \text{PTIME}$. Notably, such assumptions are not required in the case of QRC_1 .

Table 1: Complexity of some modal logics and their strictly positive counterparts.

	Full logic		Strictly positive fragment		
K	PSPACE-complete	[38]	K^+	PTIME	[9, 10]
K4 and GL	PSPACE-complete	[23]	RC_1	PTIME	[15, 8]
K4.3 and GL.3	coNP-complete	[40]	$K4.3^+$	PTIME	[49]
GLP_Λ^0 ($\Lambda \geq \omega$)	PSPACE-complete	[44]	RC_Λ^0	PTIME	[15, 8]
GLP	PSPACE-complete	[47]	RC	PTIME	[15, 8]
T	PSPACE-complete	[38]	$\{e_{refl}\}$	PTIME	[31]
S4	PSPACE-complete	[38]	\mathcal{E}_{qo}	PTIME	[31]
S4.3	coNP-complete	[23]	\mathcal{E}_{lin}	PTIME	[31]
S5	coNP-complete	[18]	$S5^+$	PTIME	[49]
$S5_m$ ($m > 1$)	PSPACE-complete	[18]	$S5_m^+$	PTIME	[49]
QPL(PA)	Π_2 -complete	[51]	QRC_1	decidable	[3]

3 Semantics for modal logics

In this section, we revisit relational and arithmetical semantics for modal provability logics for the propositional and the quantified case and further state the main theorems relating the logics and their semantics.

3.1 Relational semantics

We briefly revisit relational semantics (also known as Kripke semantics) for classical modal logics and refer to [28] for details. We start with the propositional case.

Definition 3.1 (Kripke frame, Kripke model). A Kripke frame is a tuple $\langle W, R \rangle$ where W is a non-empty set (the worlds) and $R \subseteq W \times W$ is a binary relation (we write wRu instead of $\langle w, u \rangle \in R$). A Kripke model is a Kripke frame together with a valuation function $V : \text{PROP} \rightarrow \wp(W)$, determining which propositional symbols hold at which worlds.

The truth of a modal formula in a model $\mathcal{M} := \langle W, R, V \rangle$ at a world w is defined inductively as follows:

- (i) $\mathcal{M}, w \not\models \perp$;
- (ii) $\mathcal{M}, w \models p$ if and only if $w \in V(p)$, where $p \in \text{PROP}$;
- (iii) $\mathcal{M}, w \models \varphi \rightarrow \psi$ if and only if either $\mathcal{M}, w \not\models \varphi$ or $\mathcal{M}, w \models \psi$;
- (iv) $\mathcal{M}, w \models \Box\varphi$ if and only if for every u such that wRu we have $\mathcal{M}, u \models \varphi$.

If $\mathcal{M}, w \models \varphi$ for every world $w \in W$, we say that φ is valid in \mathcal{M} and write $\mathcal{M} \models \varphi$.

We say that a model $\mathcal{M} := \langle W, R, V \rangle$ is finite if W is finite, that it is transitive if R is transitive, that it is Noetherian if every non empty subset of W has an R -maximal

element, and that it is tree-like if R is acyclic and there is a root world that is connected to every other world and is not itself a successor of any world. Both RC_1 and GL are complete with respect to finite transitive Noetherian tree-like models.

Theorem 3.2 ([13, 15]). Let φ be an \mathcal{L}_\square formula and ψ, χ be \mathcal{L}_\square^+ formulas. Let quantifiers over \mathcal{M} range over finite transitive Noetherian tree-like models with root r . Then:

$$\begin{aligned} \text{GL} \vdash \varphi &\iff (\mathcal{M}, r \Vdash \varphi \text{ for any such } \mathcal{M}); \\ \psi \vdash_{\text{RC}_1} \chi &\iff ((\mathcal{M}, r \not\Vdash \psi \text{ or } \mathcal{M}, r \Vdash \chi) \text{ for any such } \mathcal{M}). \end{aligned}$$

There have been several proposals for relational semantics for quantified modal logics, from Kripke [34] to many others. Overviews can be found in [28, 24, 25]. The idea of the semantics we present is to glue together several first-order models. The precise way in which to do it can vary, due to different ways to interpret the relationship between quantifiers and modal symbols, as well as with the first-order domains. In particular, if different worlds are allowed different domains, there is more than one way of relating those domains with each other. Here we describe Kripke sheaves [24], which for any worlds w and u such that u is accessible from w includes a compatibility function $\eta_{w,u}$ translating the domain of w into the domain of u .

Definition 3.3 (Kripke sheaf, first-order Kripke model). A Kripke sheaf \mathcal{S} is a tuple $\langle W, R, \{M_w\}_{w \in W}, \{\eta_{w,u}\}_{wRu} \rangle$ where:

- (i) W is a non-empty set (the set of worlds, where individual worlds are referred to as w, u, v , etc);
- (ii) R is a binary relation on W (the accessibility relation);
- (iii) each M_w is a non-empty set (the domain of the world w , whose elements are referred to as d, d_0, d_1 , etc);
- (iv) if wRu , then $\eta_{w,u}$ is a function from M_w to M_u .

Furthermore, we require two conditions on the compatibility functions:

- (v) if wRu , uRv , and wRv , then for every $d \in M_w$ we have $\eta_{u,v}(\eta_{w,u}(d)) = \eta_{w,v}(d)$;
- (vi) if wRw then $\eta_{w,w}$ is the identity function on M_w .

Given a signature Σ , a first-order Kripke model \mathcal{M} is a tuple $\langle \mathcal{S}, \{I_w\}_{w \in W}, \{J_w\}_{w \in W} \rangle$ where $\mathcal{S} := \langle W, R, \{M_w\}_{w \in W}, \{\eta_{w,u}\}_{wRu} \rangle$ is a Kripke sheaf and:

- for each $w \in W$, the interpretation I_w assigns an element of the domain M_w to each constant c of Σ , written c^{I_w} ;
- I is concordant: if wRu , then $c^{I_u} = \eta_{w,u}(c^{I_w})$ for every constant c of Σ .

- for each $w \in W$, the interpretation J_w assigns a set of tuples $S^{J_w} \subseteq \wp((M_w)^n)$ to each n -ary relation symbol S of Σ .

In this case, we say that the model \mathcal{M} is based on the sheaf \mathcal{S} .

We say that a Kripke sheaf is finite if both W and the domains M_w for every $w \in W$ are finite, and that it has a constant domain if all the M_w coincide and all the $\eta_{w,u}$ are the identity function. We say that a model is finite (respectively has a constant domain) if it is based on a finite (respectively constant domain) sheaf.

In order to define the notion of satisfaction, we need one more ingredient, namely a way to interpret free variables. For this, we use assignments. Given a world w , a w -assignment g is a function from the set of variables to the domain M_w . Any w -assignment can be seen as a u -assignment as long as wRu , by composing it with $\eta_{w,u}$ on the left. We write g^u to shorten $\eta_{w,u} \circ g$ when w is clear from context. A w -assignment g is extended to terms by defining $g(c) := c^{I_w}$ for any constant c . Note that this meshes nicely with the concordance restriction of the model: for any term t , if wRu then $g^u(t) = \eta_{w,u}(g(t))$. Two w -assignments g and h are x -alternative, written $g \sim_x h$, if they coincide on all variables other than x .

We are now ready to define the notion of truth.

Definition 3.4 (Truth). Let $\mathcal{M} = \langle W, R, \{M_w\}_{w \in W}, \{\eta_{w,u}\}_{wRu}, \{I_w\}_{w \in W}, \{J_w\}_{w \in W} \rangle$ be a first-order Kripke model in some signature Σ , and let $w \in W$ be a world, g be a w -assignment, S be an n -ary relation symbol, and φ, ψ be $\mathcal{L}_{\Box, \forall}$ -formulas in the language of Σ .

We define $\mathcal{M}, w \Vdash^g \varphi$ (φ is true at w under g) by induction on φ as follows:

- $\mathcal{M}, w \not\Vdash^g \perp$;
- $\mathcal{M}, w \Vdash^g S(t_0, \dots, t_{n-1})$ if and only if $\langle g(t_0), \dots, g(t_{n-1}) \rangle \in S^{J_w}$;
- $\mathcal{M}, w \Vdash^g \varphi \rightarrow \psi$ if and only if either $\mathcal{M}, w \not\Vdash^g \varphi$ or $\mathcal{M}, w \Vdash^g \psi$;
- $\mathcal{M}, w \Vdash^g \forall x \varphi$ if and only if for all w -assignments h such that $h \sim_x g$ we have $\mathcal{M}, w \Vdash^h \varphi$;
- $\mathcal{M}, w \Vdash^g \Box \varphi$ if and only if for every $u \in W$ such that wRu we have $\mathcal{M}, u \Vdash^{g^u} \varphi$.

If $\mathcal{M}, w \Vdash^g \varphi$ for every w -assignment g we say that φ is satisfied at w and write $\mathcal{M}, w \Vdash \varphi$. If φ is satisfied at every world w , we say that φ is valid in \mathcal{M} and write $\mathcal{M} \models \varphi$. Finally, if φ is valid in every model \mathcal{M} based on a given sheaf \mathcal{S} , we say that φ is valid in \mathcal{S} and write $\mathcal{S} \models \varphi$.

The soundness of a propositional modal logic can be leveraged to obtain soundness with respect to Kripke sheaves for its quantified counterpart.

Theorem 3.5 (Soundness for QL). Let \mathcal{F} be a Kripke frame for \mathcal{L}_\square as described in Definition 3.1. If \mathcal{F} satisfies every theorem of a propositional modal logic L , then any sheaf based on \mathcal{F} (i.e., with the same set of worlds and accessibility relation) satisfies every theorem of QL.

Proof. By induction on a QL proof of a $\mathcal{L}_{\square, \forall}$ formula. The proof is analogous to the one for the non-sheaf quantified Kripke semantics described in [28]. See details in [1, Theorem 2.5.4]. \square

As always, completeness has to be independently established for each different logic. Modal completeness for QRC_1 is proved in [2], and this result is strengthened to constant-domain completeness in [3].

Theorem 3.6 ([3, Theorem 5.11]). If $\varphi \not\vdash_{\text{QRC}_1} \psi$, then there are a finite Kripke model \mathcal{M} with a constant finite domain where R is irreflexive and transitive, a world $w \in \mathcal{M}$, and a w -assignment g such that $\mathcal{M}, w \Vdash^g \varphi$ and $\mathcal{M}, w \not\Vdash^g \psi$.

3.2 Arithmetical semantics

We refer to [26, 50] for definitions of PA and HA, which are Robinson’s arithmetic with full induction over classical or constructive logic, respectively. As always, Δ_0 refers to formulas where all quantifiers are bounded by some term in our language and we set $\Pi_0 := \Sigma_0 := \Delta_0$. The class Σ_{n+1} arises from existentially quantifying Π_n formulas and the class Π_{n+1} arises from universally quantifying Σ_n formulas. Given a class Γ (for example Σ_1) and a theory T , we say that a formula φ is “ Γ over T ” if T proves that φ is equivalent to ψ , for some formula $\psi \in \Gamma$.

The theory EA is as PA with the induction axiom restricted to Δ_0 formulas. The intuitionistic version of EA is denoted by iEA and it only differs from EA in that it is defined over intuitionistic predicate logic instead of classical predicate logic. A theory is said to be Γ -sound if all formulas in Γ provable in the theory are valid in the standard model \mathbb{N} .

Modal logics are related to arithmetic via Gödel numbering and via interpreting the \square modality as formalized provability. For an arithmetical formula φ , we denote its Gödel number by $\ulcorner \varphi \urcorner$. We suggestively denote the arithmetical formalized provability predicate by \square_T so that we have $T \vdash \varphi$ if and only if $\mathbb{N} \models \square_T(\ulcorner \varphi \urcorner)$. Strictly speaking, an axiomatization τ of a theory T induces a provability predicate \square_τ , but we shall often write \square_T when the intended axiomatization τ for T is clear from context. We define $\diamond_\tau \varphi$ as $\neg \square_\tau \neg \varphi$.

Propositional variables in modal logics are taken to represent placeholders for arbitrary arithmetical formulas. To formalize this idea, we use so-called realizations.

Definition 3.7 (Arithmetical realization for $\mathcal{L}_\square, \cdot^*, \cdot^{*T}$). An arithmetical realization \cdot^* is a function from PROP to sentences in the language of arithmetic. Given a c.e. theory T in the language of arithmetic and an arithmetical realization \cdot^* , we define \cdot^{*T} on modal formulas as follows:

- $\perp^{*T} := (0 = 1)$;
- $p^{*T} := p^*$, where $p \in \mathbf{PROP}$;
- $(\varphi \rightarrow \psi)^{*T} := \varphi^{*T} \rightarrow \psi^{*T}$;
- $(\Box\varphi)^{*T} := \Box_T \varphi^{*T}$.

In the case of constructive theories such as **HA**, the definition is adapted in the usual way so that \cdot^{*T} commutes with the connectives \wedge and \vee .

We can now define the provability logic $\mathbf{PL}(T)$ and the strictly positive provability logic $\mathbf{PL}^+(T)$ of a theory T in the language as:

$$\mathbf{PL}(T) := \{\varphi \in \mathcal{L}_\Box \mid \text{for any } \cdot^*, \text{ we have } T \vdash \varphi^{*T}\},$$

$$\mathbf{PL}^+(T) := \{\langle \varphi, \psi \rangle \in \mathcal{L}_\Box^+ \times \mathcal{L}_\Box^+ \mid \text{for any } \cdot^*, \text{ we have } T \vdash \varphi^{*T} \rightarrow \psi^{*T}\}.$$

We observe that, in virtue of their definitions, $\mathbf{PL}^+(T) = (\mathbf{PL}(T))^+$.

We can now state that $\mathbf{PL}(T) = \mathbf{GL}$ for a large class of theories T . The version for **PA** is known as Solovay's completeness theorem.

Theorem 3.8 ([48, 17]). Let T be a c.e. Σ_1 -sound extension of **EA** and let $\varphi \in \mathcal{L}_\Box$. Then:

$$\mathbf{GL} \vdash \varphi \iff \text{for any } \cdot^{*T}, \text{ we have } T \vdash \varphi^{*T}.$$

Strictly positive propositional logics have an alternative arithmetical interpretation where propositional variables are not mapped to sentences but rather to formulas axiomatizing a theory.

Definition 3.9 (Infinitary arithmetical realization for $\mathcal{L}_\Box^+, \cdot^\circ, \cdot^{\circ\tau}$). Let \cdot° be a function from propositional symbols to axiomatizations of theories of arithmetic. Given a base axiomatization τ , we extend \cdot° to formulas of \mathcal{L}_\Box^+ as follows:

- $\top^{\circ\tau} := \tau(u)$;
- $p^{\circ\tau} := p^\circ \vee \tau(u)$;
- $(\varphi \wedge \psi)^{\circ\tau} := \varphi^{\circ\tau} \vee \psi^{\circ\tau}$;
- $(\Diamond\varphi)^{\circ\tau} := \tau(u) \vee (u = \ulcorner \Diamond_{\varphi^{\circ\tau}} \top \urcorner)$.

Given these infinitary realizations, we define the reflection logic of a theory:

$$\mathbf{RL}(T) := \{\langle \varphi, \psi \rangle \in \mathcal{L}_\Box^+ \times \mathcal{L}_\Box^+ \mid \text{for any } \cdot^\circ, \text{ we have } T \vdash \forall \theta (\Box_{\psi^{\circ\tau}} \theta \rightarrow \Box_{\varphi^{\circ\tau}} \theta)\}. \quad (3.1)$$

Clearly, infinitary realizations extend finitary ones, for given a finitary realization \cdot^* we can define $p^\circ := \ulcorner p^* \urcorner$. Then $p^{\circ\tau}$ axiomatizes the base theory extended by the single axiom p^* . Using this observation it is not hard to see that $\mathbf{RL}(\mathbf{PA}) = \mathbf{RC}_1$ and in Section 6 we show that $\mathbf{RL}(\mathbf{HA}) = \mathbf{RC}_1$ as well.

We extend the notion of (finitary) arithmetical realization to $\mathcal{L}_{\square, \forall}$ in the natural way. To simplify matters, in the sections dealing with both the quantified modal language and the language of arithmetic, we always use variables named x_i (for some i) in the modal setting and variables named y_i or z_i in the arithmetical setting.

Definition 3.10 (Arithmetical realization for $\mathcal{L}_{\square, \forall}$ and $\mathcal{L}_{\square, \forall}^+$, \cdot^* , \cdot^{*T}). An arithmetical realization \cdot^* is a function from relation symbols applied to a tuple of n variables (corresponding to the symbol's arity) to formulas in the language of arithmetic with n free variables. We require that a variable x_i in the modal language be translated to y_i in the arithmetical setting and that a constant c_i in the modal language be translated to z_i in the arithmetical setting.

Given a c.e. theory T extending iEA , and an arithmetical realization \cdot^* , we define \cdot^{*T} on quantified modal formulas as follows (extendable to the constructive setting as before):

- $\perp^{*T} := (0 = 1)$;
- $S(\mathbf{t})^{*T} := S(\mathbf{t})^*$;
- $(\varphi \rightarrow \psi)^{*T} := \varphi^{*T} \rightarrow \psi^{*T}$;
- $(\square\varphi)^{*T} := \square_T \varphi^{*T}$;
- $(\forall x_k \varphi)^{*T} := \forall y_k \varphi^{*T}$.

This definition can be restricted to $\mathcal{L}_{\square, \forall}^+$ in the obvious way.

After Solovay's completeness theorem (Theorem 3.8), it was natural to ask whether one could find a nice characterization for the counterpart of $\text{PL}(\text{PA})$ in the quantified modal language, defined as follows (see Boolos [13]).

Definition 3.11 (Quantified provability logic, QPL). If T is a c.e. extension of iEA , the quantified provability logic of T is defined as:²

$$\text{QPL}(T) := \{\varphi \in \mathcal{L}_{\square, \forall} \mid \text{for any } \cdot^*, \text{ we have } T \vdash \forall \mathbf{y} \varphi^{*T}\},$$

$$\text{QPL}^+(T) := \{\langle \varphi, \psi \rangle \in \mathcal{L}_{\square, \forall}^+ \times \mathcal{L}_{\square, \forall}^+ \mid \text{for any } \cdot^*, \text{ we have } T \vdash \forall \mathbf{y}, \mathbf{z} (\varphi^{*T} \rightarrow \psi^{*T})\}.$$

In general, $\text{QPL}(T)$ can be different depending on the chosen axiomatization τ for T (see [5, 35, 36]), but here we assume that an axiomatization is fixed for each relevant theory for simplicity. It is not exactly the case that $\text{QPL}^+(T) = (\text{QPL}(T))^+$ due to the difference in whether constants are allowed in the language. However, for a fixed signature Σ without constants, we indeed have $\text{QPL}^+(T) = (\text{QPL}(T))^+$.

²Recall that we assume all modal variables are of the form x_i and that these are translated to the arithmetical variables y_i . We also assume that all modal constants are translated to some z_i , but note that constants are not included in the language $\mathcal{L}_{\square, \forall}$, only in $\mathcal{L}_{\square, \forall}^+$. With $\forall \mathbf{y} \varphi^{*T}$ we mean to quantify over all the free variables of φ^{*T} , and similarly in the $\text{QPL}^+(T)$ case.

The natural candidate for a quantified provability logic for PA was QGL, but although $\text{QGL} \subseteq \text{QPL}(\text{PA})$, Vardanyan [51] and McGee [41] independently showed that $\text{QPL}(\text{PA})$ is as complex as it can possibly be: Π_2 -complete. However, in [3] it was proven that the strictly positive fragment of this Π_2 -complete logic is given by QRC_1 . In other words, for the corresponding signatures we have the following result.

Theorem 3.12 ([3, Theorem 9.1]). $\text{QPL}^+(\text{PA}) = \text{QRC}_1$.

In Section 7 we prove that likewise $\text{QPL}^+(\text{HA}) = \text{QRC}_1$.

The strictly positive fragment $\mathcal{L}_{\Box, \forall}^+$ again allows for an infinitary arithmetical interpretation. In that case, a predicate $S(\mathbf{x}, \mathbf{c})$ in the signature Σ is mapped to a formula $S^\circ(\mathbf{y}, \mathbf{z})$ that axiomatizes a collection of theories parametrized by \mathbf{y}, \mathbf{z} .

Definition 3.13 (Infinitary arithmetical realization for $\mathcal{L}_{\Box, \forall}^+, \cdot^\circ, \cdot^{\circ\tau}$). Let \cdot° be a realization such that, for a given n -ary predicate S and terms \mathbf{t} , $S(\mathbf{t})^\circ$ is an $(n+1)$ -ary Σ_1 arithmetical formula where modal variables x_k are interpreted as y_k and modal constants c_k are interpreted as z_k . Let τ be a Σ_1 axiomatization of some base theory T . We extend this realization to QRC_1 formulas as follows:

- $\top^{\circ\tau} := \tau(u)$;
- $S(\mathbf{x}, \mathbf{c})^{\circ\tau} := S(\mathbf{x}, \mathbf{c})^\circ \vee \tau(u)$;
- $(\psi \wedge \delta)^{\circ\tau} := \psi^{\circ\tau} \vee \delta^{\circ\tau}$;
- $(\Diamond\psi)^{\circ\tau} := \tau(u) \vee (u = \ulcorner \Diamond_{\psi^{\circ\tau}} \top \urcorner)$;
- $(\forall x_i \psi)^{\circ\tau} := \exists y_i \psi^{\circ\tau}$.

Using the infinitary semantics, we can define the quantified reflection logic $\text{QRL}(T)$ of a theory T in analogy with (3.1) as:

$$\text{QRL}(T) := \{ \langle \varphi, \psi \rangle \in \mathcal{L}_{\Box, \forall}^+ \times \mathcal{L}_{\Box, \forall}^+ \mid \text{for any } \cdot^\circ, T \vdash \forall \theta \forall \mathbf{y}, \mathbf{z} (\Box_{\psi^{\circ\tau}} \theta \rightarrow \Box_{\varphi^{\circ\tau}} \theta) \}.$$

In [3, Theorems 7.2 and 8.9] it is shown that $\text{QRL}(\text{PA}) = \text{QRC}_1$ and in Section 7 we prove that furthermore $\text{QRL}(\text{HA}) = \text{QRC}_1$.

4 The arithmetical hierarchy in HA

Heyting Arithmetic does not permit a simple treatment of the arithmetical hierarchy (see e.g. [22]). In this section we collect some basic observations on approximations to classical reasoning regarding formula complexity, which we use later in the paper.

4.1 Closure and semi-closure under double negations

Using standard coding techniques, it is possible to see that **HA** proves that Σ_1 sentences are closed under conjunction, and this fact is formalizable in **HA**. We furthermore recall that for any arithmetical formula φ we have:

$$\mathbf{HA} \vdash \forall x \neg \varphi \leftrightarrow \neg \exists x \varphi. \quad (4.1)$$

This equivalence can be contrasted with $\mathbf{HA} \not\vdash \exists x \neg \varphi \leftrightarrow \neg \forall x \varphi$.

Additionally, **HA** proves excluded middle for decidable formulas. Consequently, if δ is a decidable formula, then:

$$\mathbf{HA} \vdash \neg \neg \delta \leftrightarrow \delta. \quad (4.2)$$

In fact, this holds for arithmetical formulas δ of complexity Δ_1 . However, excluded middle fails to hold for genuine Π_1 sentences π , but double negation elimination is still possible:

$$\mathbf{HA} \vdash \neg \neg \pi \leftrightarrow \pi.$$

The same equivalence does not hold for genuine Σ_1 sentences σ :

$$\mathbf{HA} \not\vdash \neg \neg \sigma \leftrightarrow \sigma.$$

Even though double negation elimination fails in general for Σ_1 sentences, we do have a certain formalized version. In Lemma 4.2 we will see that $\mathbf{HA} \vdash \Box_{\mathbf{HA}} \sigma \leftrightarrow \Box_{\mathbf{HA}} \neg \neg \sigma$ for Σ_1 sentences σ . This is due to a certain amount of conservatism between **HA** and **PA** that shall be discussed in the next subsection.

The class of Σ_2 formulas is particularly important for our study. In pure constructive predicate logic, Σ_2 formulas are provably closed under conjunctions but not under disjunctions. However, in **HA** we do have the latter.

Lemma 4.1. For each $\varphi_0, \varphi_1 \in \Sigma_2$, there is $\varphi \in \Sigma_2$ such that:

$$\mathbf{HA} \vdash \varphi_0 \vee \varphi_1 \leftrightarrow \varphi.$$

Proof. Let $\varphi_0 = \exists s \forall t \tilde{\varphi}_0(s, t)$ and $\varphi_1 = \exists u \forall v \tilde{\varphi}_1(u, v)$, with $\tilde{\varphi}_1(u, v), \tilde{\varphi}_0(s, t) \in \Delta_0$. We can take:

$$\varphi := \exists x \forall y ((x)_0 = 0 \wedge \tilde{\varphi}_0((x)_1, y)) \vee ((x)_0 = 1 \wedge \tilde{\varphi}_1((x)_1, y)),$$

where x is being interpreted as a pair, of which $(x)_0$ and $(x)_1$ are the first and second projections, respectively. \square

4.2 Equi-consistency results in **HA**

When moving between **PA** and **HA**, one conservativity direction is trivial and formalizable in **HA**:

$$\mathbf{HA} \vdash \forall \varphi (\Box_{\mathbf{HA}} \varphi \rightarrow \Box_{\mathbf{PA}} \varphi). \quad (4.3)$$

The other direction is known to hold provably in **HA** for Π_2 sentences (see [21]), both in the non-formalized setting and in the formalized setting, so we have an equivalence:

$$\mathbf{HA} \vdash \forall \pi \in \Pi_2 \ (\Box_{\mathbf{PA}} \pi \leftrightarrow \Box_{\mathbf{HA}} \pi). \quad (4.4)$$

With this conservativity result, we obtain a certain amount of classical behavior for Π_2 sentences in a formalized setting. In particular, combining the above we have a **HA** provable semi-closure of a certain kind of double negation.

Lemma 4.2.

1. $\mathbf{HA} \vdash \forall \sigma \in \Sigma_1 \ (\Box_{\mathbf{HA}} \neg\neg\sigma \leftrightarrow \Box_{\mathbf{HA}} \sigma)$;
2. $\mathbf{HA} \vdash \forall \sigma \in \Sigma_1 \ (\Box_{\mathbf{HA}} \forall x \neg\neg\sigma \leftrightarrow \Box_{\mathbf{HA}} \forall x \sigma)$.

Proof. Clearly, the first statement follows from the second, and so we only focus on the latter. Reason in **HA** and fix an arbitrary $\sigma \in \Sigma_1$. Since clearly $\sigma \rightarrow \neg\neg\sigma$, we focus on the other implication. Consider:

$$\begin{aligned} \Box_{\mathbf{HA}} \forall x \neg\neg\sigma &\rightarrow \Box_{\mathbf{PA}} \forall x \neg\neg\sigma && \text{by (4.3)} \\ &\rightarrow \Box_{\mathbf{PA}} \forall x \sigma \\ &\rightarrow \Box_{\mathbf{HA}} \forall x \sigma && \text{by (4.4)}. \end{aligned}$$

□

Combining some of the above observations, it is easy to see that the negation of a Π_1 sentence is equivalent to the double negation of a Σ_1 sentence over **HA**.

Lemma 4.3. Given a Π_1 sentence π , there is a Σ_1 sentence σ such that:

$$\mathbf{HA} \vdash \neg\pi \leftrightarrow \neg\neg\sigma.$$

Proof. Let $\pi := \forall x \delta$ where $\delta \in \Delta_0$. Consider:

$$\begin{aligned} \mathbf{HA} \vdash \neg\forall x \delta &\leftrightarrow \neg\forall x \neg\neg\delta \\ &\leftrightarrow \neg\neg\exists x \neg\delta && \text{by (4.1)}. \end{aligned}$$

Then $\sigma := \exists x \neg\delta$ is the required Σ_1 sentence. □

We can now prove a dual version of (4.4).

Lemma 4.4.

1. $\mathbf{HA} \vdash \forall \varphi \in \Sigma_2 \ (\Box_{\mathbf{HA}} \neg\varphi \leftrightarrow \Box_{\mathbf{PA}} \neg\varphi)$;
2. $\mathbf{HA} \vdash \forall \varphi \in \Sigma_2 \ (\Diamond_{\mathbf{HA}} \varphi \leftrightarrow \Diamond_{\mathbf{PA}} \varphi)$.

Proof. The second statement easily follows from the first. Reason in HA and fix an arbitrary $\varphi \in \Sigma_2$. We can write φ as $\exists x \pi$ for some $\pi \in \Pi_1$. Let σ be the Σ_1 sentence given by Lemma (4.3) for our fixed π such that:

$$\Box_{\text{HA}}(\neg\pi \leftrightarrow \neg\neg\sigma). \quad (4.5)$$

Now consider:

$$\begin{aligned} \Box_{\text{HA}} \neg\varphi &\leftrightarrow \Box_{\text{HA}} \neg \exists x \pi && \\ &\leftrightarrow \Box_{\text{HA}} \forall x \neg\pi && \text{by (4.1)} \\ &\leftrightarrow \Box_{\text{HA}} \forall x \neg\neg\sigma && \text{by (4.5)} \\ &\leftrightarrow \Box_{\text{HA}} \forall x \sigma && \text{by Lemma 4.2} \\ &\leftrightarrow \Box_{\text{PA}} \forall x \sigma && \text{by (4.4)} \\ &\leftrightarrow \Box_{\text{PA}} \neg\neg \forall x \sigma && \\ &\leftrightarrow \Box_{\text{PA}} \neg \exists x \neg\sigma && \\ &\leftrightarrow \Box_{\text{PA}} \neg \exists x \pi && \text{by (4.5)} \\ &\leftrightarrow \Box_{\text{PA}} \neg\varphi. && \end{aligned}$$

□

The above lemma is reminiscent of the following result by Kreisel.

Theorem 4.5 (Kreisel [33]). For every formula φ in prenex normal form we have:

$$\text{HA} \vdash \Box_{\text{PA}} \neg\varphi \leftrightarrow \Box_{\text{HA}} \neg\varphi.$$

Proof. The right to left implication is clear, so we focus on the other direction. Reason in HA and assume:

$$\Box_{\text{PA}} \neg \forall x_1 \exists y_1 \forall x_2 \exists y_2 \cdots \delta,$$

taking φ as $\forall x_1 \exists y_1 \forall x_2 \exists y_2 \cdots \delta$ for some Δ_0 -formula δ , where the first universal quantification may be vacuous. By the Kuroda translation [37], we deduce:

$$\Box_{\text{HA}} \neg \forall x_1 \neg\neg \exists y_1 \forall x_2 \neg\neg \exists y_2 \cdots \delta.$$

Since $\text{HA} \vdash \forall x_1 \exists y_1 \forall x_2 \exists y_2 \cdots \delta \rightarrow \forall x_1 \neg\neg \exists y_1 \forall x_2 \neg\neg \exists y_2 \cdots \delta$, we obtain:

$$\Box_{\text{HA}} \neg \forall x_1 \exists y_1 \forall x_2 \exists y_2 \cdots \delta,$$

that is, $\Box_{\text{HA}} \neg\varphi$, as desired. □

From Kreisel's observation we directly obtain the following corollary.

Corollary 4.6. For every formula φ in prenex normal form we have:

$$\text{HA} \vdash \Diamond_{\text{PA}} \varphi \leftrightarrow \Diamond_{\text{HA}} \varphi.$$

5 Reflections on arithmetical completeness

In this section we briefly revisit the ingredients of Solovay's arithmetical completeness proof for **GL**. We have a particular interest in the complexity of the formulas occurring in that proof so that we can invoke conservativity of **PA** over **HA**. We first observe that the complexity of the arithmetical realizations of propositional strictly positive formulas can be kept low.

Lemma 5.1. For every realization $\cdot^* : \text{PROP} \rightarrow \Sigma_2$, every c.e. theory T and every p.s.p. formula φ we have that φ^{*T} is Σ_2 over **HA**, and this is provable in **HA**.

Proof. By an easy induction on the complexity of φ . The only interesting case is the complexity of $\diamond\psi$, which is taken care of by (4.1) since the negation of a Σ_1 sentence is Π_1 , whence also Σ_2 . \square

This simple observation can be used to see that extending a realization using **HA** provability or **PA** provability yields **HA**-equivalent formulas, provably in **HA**.

Lemma 5.2. For every realization $\cdot^* : \text{PROP} \rightarrow \Sigma_2$ and every p.s.p. formula φ we have:

$$\text{HA} \vdash \varphi^{*\text{HA}} \leftrightarrow \varphi^{*\text{PA}}.$$

Proof. By an easy induction on φ where the \diamond case is taken care of by Lemma 4.4. \square

Solovay proved the arithmetical completeness theorem for **GL** by embedding Kripke frames into arithmetic. Given a Kripke frame with worlds $1, \dots, N$, a new world 0 is added such that it can see all the other worlds. Then, each $i \in \{0, 1, \dots, N\}$ is assigned an arithmetical sentence λ_i such that each λ_i is independent from **PA** and that these λ_i sentences mimic the accessibility relations between their respective worlds as described by the following lemma. Then the realization \cdot^* of Solovay's proof is defined as $p^* := \bigvee_{i \Vdash p} \lambda_i$.

Lemma 5.3 (Embedding [13]).

1. $\text{PA} \vdash \bigvee_{i \leq N} \lambda_i$;
2. $\text{PA} \vdash \lambda_i \rightarrow \bigwedge_{j \leq N, j \neq i} \neg \lambda_j$, for $i \leq N$;
3. $\text{PA} \vdash \lambda_i \rightarrow \bigwedge_{j \leq N, iRj} \diamond_{\text{PA}} \lambda_j$, for $i \leq N$;
4. $\text{PA} \vdash \lambda_i \rightarrow \square_{\text{PA}} \bigvee_{j \leq N, iRj} \lambda_j$, for $0 < i \leq N$;
5. $\mathbb{N} \models \lambda_0$.

The sentence λ_i states that the limit of a so-called Solovay function $f : \mathbb{N} \rightarrow \{0, 1, \dots, N\}$ is i . Each sentence is thus *a priori* of complexity Σ_2 . However, by definition, the Solovay function can only move along R -transitions in the frame, so we can write λ_i as a conjunction of Π_1 and Σ_1 sentences as follows:

$$\lambda_i = (\exists x f(x) = i) \wedge \forall x, y (x \leq y \rightarrow (f(x) = i \rightarrow f(y) = i)). \quad (5.1)$$

We denote the class of formulas built of conjunctions and disjunctions of Π_1 and Σ_1 formulas by $\mathcal{DC}(\Pi_1, \Sigma_1)$. Since in Solovay's proof $p^* := \bigvee_{i \in \mathbb{N}} \lambda_i$, it is clear that the realization satisfies $\cdot^* : \text{PROP} \rightarrow \mathcal{DC}(\Pi_1, \Sigma_1)$. In fact, φ^{*T} still has complexity $\mathcal{DC}(\Pi_1, \Sigma_1)$, which can be proved by a simple induction on φ .

Remark 5.4. For any p.s.p. formula φ we have $\varphi^{*T} \in \mathcal{DC}(\Pi_1, \Sigma_1)$.

Note that $\mathcal{DC}(\Pi_1, \Sigma_1) \subseteq \Sigma_2$ over HA by Lemma 4.1. As a direct consequence of the previous observations we obtain the following result.

Corollary 5.5. Let \cdot^* be a realization obtained via Solovay's embedding of a finite transitive Noetherian tree-like model. Then for any p.s.p. formula φ :

$$\text{HA} \vdash \neg \varphi^{*\text{HA}} \iff \text{PA} \vdash \neg \varphi^{*\text{PA}}.$$

Moreover, this is formalizable in HA.

Proof. By Remark 5.4, Lemma 5.2, and Lemma 4.4. □

Note that we cannot prove a version of Corollary 5.5 without the negation: if $\text{PA} \vdash p^{*\text{PA}}$ implied $\text{HA} \vdash p^{*\text{HA}}$, then by the disjunction property we would have $\text{HA} \vdash \lambda_i$ for some i , which would contradict the independence of λ_i from PA.

6 The strictly positive fragment of PL(HA)

In this section we show that the strictly positive provability logic of HA and the reflection logic of HA both coincide with RC_1 .

Theorem 6.1. Let φ and ψ be p.s.p. formulas. We have $\varphi \vdash_{\text{RC}_1} \psi$ if and only if for all realizations \cdot^* we have $\text{HA} \vdash \varphi^{*\text{HA}} \rightarrow \psi^{*\text{HA}}$.

Proof. Soundness is easy given the derivability conditions of \Box_{HA} (see [56]) and it was already incidentally observed in [1, Theorem 5.4.3]. For completeness, reason as follows:

$$\begin{aligned} \varphi \not\vdash_{\text{RC}_1} \psi &\implies \text{there is } \cdot^* \text{ s.t. } \text{PA} \not\vdash \varphi^{*\text{PA}} \rightarrow \psi^{*\text{PA}} && \text{by Theorem 3.8} \\ &\implies \text{there is } \cdot^* \text{ s.t. } \text{PA} \not\vdash \varphi^{*\text{HA}} \rightarrow \psi^{*\text{HA}} && \text{by Lemma 5.2} \\ &\implies \text{there is } \cdot^* \text{ s.t. } \text{HA} \not\vdash \varphi^{*\text{HA}} \rightarrow \psi^{*\text{HA}} && \text{by (4.3).} \end{aligned}$$

□

We note that there is some room for improvement and generalization, arithmetically speaking. Can this be exploited to get a slightly larger fragment (disjunctions seem to come for free, and boxes with a bit of extra work)?

Recall that infinitary realizations extend finitary ones in the following sense. Given a finitary realization \cdot^* , we can define $p^\circ := (u = \ulcorner p^* \urcorner)$. Then $p^{\circ\tau}$ axiomatizes the base theory extended by the single axiom p^* . This infinitary realization behaves exactly like its finitary counterpart, as the following lemma illustrates.

Lemma 6.2. Let φ be a p.s.p. formula, \cdot^* be a finitary realization and \cdot° be its finitary counterpart as described above. Then:

$$\text{HA} \vdash \forall \theta (\Box_{\varphi^\circ \eta} \theta \leftrightarrow \Box_{\text{HA}}(\varphi^{*\text{HA}} \rightarrow \theta)).$$

Proof. By an easy external induction on φ . □

Using this the reflection logic of HA can be obtained as follows (recall (3.1)).

Theorem 6.3. $\text{RL}(\text{HA}) = \text{RC}_1$.

Proof. The soundness, that $\text{RC}_1 \subseteq \text{RL}(\text{HA})$, follows from a straightforward induction and was already incidentally observed in [3, Theorem 10.3]. For the other direction, assume that $\varphi \not\vdash_{\text{RC}_1} \psi$. With Theorem 6.1 we find \cdot^* such that $\text{HA} \not\vdash \varphi^{*\text{HA}} \rightarrow \psi^{*\text{HA}}$. We now set $p^\circ := (u = \ulcorner p^* \urcorner)$. We have $\text{HA} \vdash \Box_{\psi^\circ \eta} \psi^{*\text{HA}}$ by Lemma 6.2. We now check that $\text{HA} \not\vdash \Box_{\varphi^\circ \eta} \psi^{*\text{HA}}$ with the following reasoning:

$$\begin{aligned} \text{HA} \vdash \Box_{\varphi^\circ \eta} \psi^{*\text{HA}} &\implies \text{HA} \vdash \Box_{\text{HA}}(\varphi^{*\text{HA}} \rightarrow \psi^{*\text{HA}}) && \text{by Lemma 6.2} \\ &\implies \mathbb{N} \models \Box_{\text{HA}}(\varphi^{*\text{HA}} \rightarrow \psi^{*\text{HA}}) && \text{by } \Sigma_1\text{-soundness} \\ &\implies \text{HA} \vdash \varphi^{*\text{HA}} \rightarrow \psi^{*\text{HA}}. \end{aligned}$$

Since $\text{HA} \not\vdash \varphi^{*\text{HA}} \rightarrow \psi^{*\text{HA}}$ by the choice of \cdot^* , we conclude that $\text{HA} \not\vdash \Box_{\varphi^\circ \eta} \psi^{*\text{HA}}$. Consequently, $\text{HA} \not\vdash \forall \theta (\Box_{\psi^\circ \eta} \theta \rightarrow \Box_{\varphi^\circ \eta} \theta)$, as was to be shown. □

7 The strictly positive fragment of QPL(HA)

In this section we prove that $\text{QPL}^+(\text{HA}) = \text{QRL}(\text{HA}) = \text{QRC}_1$. The proofs are essentially the same as the ones in the propositional case as stated in Theorems 6.1 and 6.3, because we can reduce the quantified case to the propositional case. In [3] this reduction was done in PA but it can actually be obtained in HA.

The reduction boils down to the finite model property of QRC_1 (Theorem 3.6), which implies that a seemingly unrestricted universal quantification is equivalent to a finite conjunction. This is observed in Lemma 7.2 below. As such, the complexity of the arithmetical interpretations of our quantified modal formulas can be restricted, as observed in Lemma 7.3.

To obtain these results, we first recall the arithmetical interpretations used in the arithmetical completeness proof presented in [3], where finite first-order Kripke models are embedded in arithmetic.

For the remainder of this section, we fix a certain finite constant domain model \mathcal{M} . Let the domain of every world of \mathcal{M} be denoted by M . Let m be the size of M , and $\ulcorner \cdot \urcorner$ be a bijection between M and the set of numerals $\{\overline{0}, \dots, \overline{m-1}\}$.

For a given n -ary predicate symbol S and terms $\mathbf{t} := t_0, \dots, t_{n-1}$, the \cdot^{\otimes} interpretation of S is defined as follows:

$$S(\mathbf{t})^{\otimes} := \bigvee_{i \leq N} \left(\lambda_i \wedge \Phi_i^{S(\mathbf{t})} \right), \text{ where:}$$

$$\Phi_i^{S(\mathbf{t})} := \bigvee_{\langle a_0, \dots, a_{n-1} \rangle \in S^{J_i}} \bigwedge_{l < n} \left(\ulcorner a_l \urcorner = \begin{cases} y_k \bmod \bar{m} & \text{if } t_l = x_k \\ z_k \bmod \bar{m} & \text{if } t_l = c_k \end{cases} \right).$$

Here x_k is any QRC_1 variable, y_k is the corresponding arithmetical variable, c_k is any QRC_1 constant, z_k is the corresponding arithmetical variable, and \cdot^{J_i} is the denotation of a predicate symbol at world i . It is clear that \cdot^{\otimes} depends on the particular choice of \mathcal{M} , but this dependency is omitted in the notation for simplicity. We keep \mathcal{M} fixed throughout this section, so this should not lead to confusion.

Remark 7.1. For any given finite and constant domain model \mathcal{M} , we have that $S(\mathbf{t})^{\otimes} \in \mathcal{DC}(\Pi_1, \Sigma_1)$.

The \cdot^{\otimes} interpretation is extended to all strictly positive formulas as described in Definition 3.10. The interpretation $S(\mathbf{t})^{\otimes}$ depends only on the values of its free variables modulo m . This modular dependence propagates over all formulas, as shown by the following lemma.

Lemma 7.2. For any c.e. theory T , any QRC_1 formula φ , and any arithmetical variable u :

1. $\text{HA} \vdash \varphi^{\otimes T} \leftrightarrow \varphi^{\otimes T}[u \leftarrow (u \bmod \bar{m})]$;
2. $\text{HA} \vdash \forall u \varphi^{\otimes T} \leftrightarrow \bigwedge_{u \in \{\bar{0}, \dots, \bar{m}-1\}} \varphi^{\otimes T}$.

Proof. The second item is a straightforward consequence of the first, which is proved by an external induction on the complexity of φ , noting that **HA** proves:

$$(u \bmod \bar{m}) \bmod \bar{m} = u \bmod \bar{m}$$

for any u . □

Since quantification does not essentially increase complexity, we see that the arithmetical interpretations of $\mathcal{L}_{\square, \forall}^+$ are all of low complexity.

Lemma 7.3. For any c.e. theory T and any s.p. formula $\varphi \in \mathcal{L}_{\square, \forall}^+$, we have that $\varphi^{\otimes T} \in \mathcal{DC}(\Pi_1, \Sigma_1)$.

Proof. By an easy induction using Remark 7.1 for the base case and Lemma 7.2 for the only non-trivial inductive step. □

We can now obtain the analogous of Lemma 5.2 in the quantified case. Note however that this does not hold in general for all realizations; the claim here is only for the specific realization \cdot^{\otimes} .

Lemma 7.4. For any formula $\varphi \in \mathcal{L}_{\square, \forall}^+$, we have $\text{HA} \vdash \varphi^{\otimes \text{HA}} \leftrightarrow \varphi^{\otimes \text{PA}}$.

Proof. By an easy external induction using Lemma 4.4 for the diamond case and Lemma 7.2 for the quantifier case. \square

Finally, we can be more specific in the statement of the completeness theorem $\text{QPL}^+(\text{PA}) \subseteq \text{QRC}_1$.

Theorem 7.5 ([3, Theorem 8.7]). Let φ and ψ be QRC_1 formulas and let \mathbf{y}, \mathbf{z} represent all the free variables of $\varphi^{\otimes \text{PA}} \wedge \psi^{\otimes \text{PA}}$. If $\varphi \not\vdash_{\text{QRC}_1} \psi$, then there are standard numbers $\mathbf{y}_0, \mathbf{z}_0$ such that $\text{PA} \not\vdash (\varphi^{\otimes \text{PA}} \rightarrow \psi^{\otimes \text{PA}})[\mathbf{y} \leftarrow \mathbf{y}_0][\mathbf{z} \leftarrow \mathbf{z}_0]$.

With these preparations, we repeat our propositional completeness proof and obtain a quantified completeness proof.

Theorem 7.6. Let φ and ψ be formulas in $\mathcal{L}_{\square, \forall}^+$. We have $\varphi \vdash_{\text{QRC}_1} \psi$ if and only if for all realizations \cdot^* we have $\text{HA} \vdash \varphi^{*\text{HA}} \rightarrow \psi^{*\text{HA}}$.

Proof. Soundness is proven by a straightforward induction on $\varphi \vdash_{\text{QRC}_1} \psi$ and was already observed in [1, Theorem 5.4.3]. For completeness we reason as follows:

$$\begin{aligned} \varphi \not\vdash_{\text{QRC}_1} \psi &\implies \text{PA} \not\vdash (\varphi^{\otimes \text{PA}} \rightarrow \psi^{\otimes \text{PA}})[\mathbf{y} \leftarrow \mathbf{y}_0][\mathbf{z} \leftarrow \mathbf{z}_0] && \text{by Theorem 7.5} \\ &\implies \text{PA} \not\vdash (\varphi^{\otimes \text{HA}} \rightarrow \psi^{\otimes \text{HA}})[\mathbf{y} \leftarrow \mathbf{y}_0][\mathbf{z} \leftarrow \mathbf{z}_0] && \text{by Lemma 7.4} \\ &\implies \text{HA} \not\vdash (\varphi^{\otimes \text{HA}} \rightarrow \psi^{\otimes \text{HA}})[\mathbf{y} \leftarrow \mathbf{y}_0][\mathbf{z} \leftarrow \mathbf{z}_0] && \text{by (4.3),} \end{aligned}$$

where \mathbf{y}, \mathbf{z} are the free variables of $\varphi^{\otimes \text{PA}}$ and $\psi^{\otimes \text{PA}}$ (which are the same as the free variables of $\varphi^{\otimes \text{HA}}$ and $\psi^{\otimes \text{HA}}$) and $\mathbf{y}_0, \mathbf{z}_0$ are as given by Theorem 7.5. Thus there is a realization \cdot^{\otimes} such that $\text{HA} \not\vdash \forall \mathbf{y}, \mathbf{z} (\varphi^{\otimes \text{HA}} \rightarrow \psi^{\otimes \text{HA}})$, as we wanted to show. \square

We now determine $\text{QRL}(\text{HA})$, the quantified reflection logic of HA . Just as in the propositional case, we can interpret a finitary realization \cdot^* as an infinitary one by taking $S(\mathbf{t})^\circ := (u = \ulcorner S(\mathbf{t})^* \urcorner)$. However, unlike in the propositional case, the realizations $\varphi^{\circ\tau}$ and $\tau \vee (u = \ulcorner \varphi^* \urcorner)$ are not necessarily equivalent (contrast with Lemma 6.2). In general, one can only prove $T \vdash \forall \theta \forall \mathbf{y}, \mathbf{z} (\Box_{\varphi^{\circ\tau}} \theta \rightarrow \Box_\tau (\varphi^{*T} \rightarrow \theta))$ (see [1, Lemma 5.2.8] and the following discussion), but in the specific case of the realization \cdot^{\otimes} above we can indeed prove the equivalence.

Let \cdot^{\circledast} be the infinitary counterpart to \cdot^{\otimes} , i.e., $S(\mathbf{t})^{\circledast} := (u = \ulcorner S(\mathbf{t})^{\otimes} \urcorner)$. Note that $\varphi^{\circledast\tau}$ has same free variables as $\varphi^{\otimes T}$ for any φ .

Lemma 7.7. Let η denote a standard axiomatization of HA and let φ be a QRC_1 formula with free variables \mathbf{x} and constants \mathbf{c} . Then:

$$\text{HA} \vdash \forall \theta \forall \mathbf{y}, \mathbf{z} (\Box_{\varphi^{\circ\eta}} \theta \leftrightarrow \Box_{\text{HA}} (\varphi^{\otimes \text{HA}} \rightarrow \theta)),$$

where θ is (the Gödel number of) a closed formula.

Proof. By external induction on φ . The proof is presented for PA in [3, Lemma 8.8] and adapts without any problem to HA. \square

With this reduction from quantified to propositional we can now determine the quantified reflection logic of HA.

Theorem 7.8. $\text{QRL}(\text{HA}) = \text{QRC}_1$.

Proof. The inclusion $\text{QRC}_1 \subseteq \text{QRL}(\text{HA})$ is a straightforward induction and is proved in [3, Theorem 10.3]. For the other direction, we reason as in the proof of Theorem 6.3 and assume that $\varphi \not\vdash_{\text{QRC}_1} \psi$.

By Theorem 7.6 we find \cdot^{\otimes} and $\mathbf{y}_0, \mathbf{z}_0$ such that:

$$\text{HA} \not\vdash (\varphi^{\otimes_{\text{HA}}} \rightarrow \psi^{\otimes_{\text{HA}}})[\mathbf{y} \leftarrow \mathbf{y}_0][\mathbf{z} \leftarrow \mathbf{z}_0]. \quad (7.1)$$

By Lemma 7.7 we get for the corresponding \cdot^{\odot_η} that for arbitrary \mathbf{y}, \mathbf{z} we have $\text{HA} \vdash \square_{\psi^{\odot_\eta}} \psi^{\otimes_{\text{HA}}}$. Again by Lemma 7.7 and using (7.1), in complete analogy with the proof of Theorem 6.3, we see that $\text{HA} \not\vdash \square_{\varphi^{\odot_\eta}} \psi^{\otimes_{\text{HA}}}$, so $\text{HA} \not\vdash \forall \theta \forall \mathbf{y}, \mathbf{z} (\square_{\psi^{\odot_\eta}} \theta \rightarrow \square_{\varphi^{\odot_\eta}} \theta)$ as was to be shown. \square

8 On interpolation and the fixpoint theorem

The interpolation and fixpoint theorems are important properties of theories and logics. In this section we consider them in the setting of QRC_1 .

8.1 Interpolation

The predicate provability logic of Peano Arithmetic does not have the interpolation property (due to Vardanyan, see [13]). Since $\text{QPL}(\text{PA})$ fails interpolation, it is important to ask whether our fragment does satisfy interpolation. It turns out that indeed, both RC_1 and QRC_1 have interpolation, and the proof is rather trivial.

Theorem 8.1. Let T be either RC or QRC_1 . If $\varphi \vdash_T \psi$, then there is a formula χ in the common language of φ and ψ such that both $\varphi \vdash_T \chi$ and $\chi \vdash_T \psi$.

Proof. By an easy induction on a proof of $\varphi \vdash_T \psi$ we observe that all symbols occurring in ψ already occur in φ . Consequently, we can take $\chi := \psi$, that is, ψ is the required interpolant. \square

8.2 Fixpoint theorem

The fixpoint theorem for GL states that for each formula φ where all appearances of p occur under the scope of a modality, we can find a fixpoint χ of φ where p does not occur such that $\text{GL} \vdash \square(p \leftrightarrow \varphi) \rightarrow (p \leftrightarrow \chi)$ (where $\square\delta := \delta \wedge \square\delta$).

We observe that the fixpoint theorem cannot really be formulated in our framework. The best we could do is to demand both $p \vdash \varphi(p, \mathbf{q})$ and $\varphi(p, \mathbf{q}) \vdash p$ as antecedents (with p occurring modalized in φ) of our version of the fixpoint theorem. Note, however, that $p \not\vdash \varphi(p, \mathbf{q})$ for any φ where p occurs modalized. This is due to the following elementary observation.

Lemma 8.2 ([2, Lemma 2.4]). Let $d_\diamond(\varphi)$ be the modal depth (maximum number of nested diamonds) of φ . If $\varphi \vdash_{\text{RC}_1} \psi$ or $\varphi \vdash_{\text{QRC}_1} \psi$, then $d_\diamond(\varphi) \geq d_\diamond(\psi)$.

Proof. By an easy induction on the length of a RC_1 or QRC_1 proof. □

Thus, p cannot prove any formula φ that is modalized in p (since the only available modality is the diamond). Our framework is hence not expressive enough to formulate a sensible version of the fixpoint theorem. Alternatively, we could state that the fixpoint theorem holds vacuously true.

Acknowledgments

We are very grateful to Dick de Jongh, Mojtaba Mojtahedi, and Albert Visser, who helped with the intuitionistic mathematics involved in this work. Their reading, comments, suggestions and discussions substantially helped the presentation of this paper.

References

- [1] de Almeida Borges, A. (2023). *Suitable logics: provability, temporal laws, and formalization*. Ph.D. thesis, University of Barcelona.
URL https://mat.ub.edu/pop/doctorat/tesis/Thesis_Ana_Almeida.pdf
- [2] de Almeida Borges, A., & Joosten, J. J. (2020). Quantified Reflection Calculus with one modality. In N. Olivetti, R. Verbrugge, S. Negri, & G. Sandu (Eds.) *Advances in Modal Logic 13*, (pp. 13–32). College Publications.
URL <http://www.aiml.net/volumes/volume13/deAlmeidaBorges-Joosten.pdf>
- [3] de Almeida Borges, A., & Joosten, J. J. (2023). An escape from Vardanyan’s Theorem. *The Journal of Symbolic Logic*, 88(4), 1613–1638.
URL <https://www.cambridge.org/core/journals/journal-of-symbolic-logic/article/an-escape-from-wardanyans-theorem/03E13D4C282563F918AF77CF6E2830DA>
- [4] Artemov, S., & Beklemishev, L. (2004). Provability logic. In D. Gabbay, & F. Guenther (Eds.) *Handbook of Philosophical Logic, 2nd ed.*, vol. 13, (pp. 229–403). Dordrecht: Springer.

- [5] Artemov, S. N. (1986). Numerically correct logics of provability. *Doklady Akademii Nauk SSSR*, 290(6), 1289–1292. In Russian.
- [6] Artemov, S. N., & Japaridze, G. K. (1990). Finite Kripke models and predicate logics of provability. *Journal of Symbolic Logic*, 55(3), 1090–1098.
- [7] Beklemishev, L. D. (2005). Veblen hierarchy in the context of provability algebras. In P. Hájek, L. Valdés-Villanueva, & D. Westerståhl (Eds.) *Logic, Methodology and Philosophy of Science, Proceedings of the Twelfth International Congress*, (pp. 65–78). Kings College Publications.
- [8] Beklemishev, L. D. (2012). Calibrating provability logic: From modal logic to Reflection Calculus. In T. Bolander, T. Braüner, T. S. Ghilardi, & L. Moss (Eds.) *Advances in Modal Logic 9*, (pp. 89–94). London: College Publications.
- [9] Beklemishev, L. D. (2018). A note on strictly positive logics and word rewriting systems. In S. Odintsov (Ed.) *Larisa Maximova on Implication, Interpolation, and Definability*, vol. 15, (pp. 61–70). Berlin, Heidelberg: Springer.
- [10] Beklemishev, L. D. (2021). Strictly positive provability logics: Recent progress and open questions.
URL <http://yongcheng.whu.edu.cn/webPageContent/Goedel2021/Goedel-Lev.pdf>
- [11] Beklemishev, L. D., Fernández-Duque, D., & Joosten, J. J. (2014). On provability logics with linearly ordered modalities. *Studia Logica*, 102, 541–566.
- [12] Berarducci, A. (1989). Σ_n^0 -interpretations of modal logic. *Bollettino dell'Unione Matematica Italiana*, 7(3-A), 177–184.
- [13] Boolos, G. S. (1993). *The logic of provability*. Cambridge: Cambridge University Press.
- [14] Boolos, G. S., & McGee, V. R. (1987). The degree of the set of sentences of predicate provability logic that are true under every interpretation. *The Journal of Symbolic Logic*, 52, 165–171.
- [15] Dashkov, E. V. (2012). On the positive fragment of the polymodal provability logic GLP. *Mathematical Notes*, 91(3–4), 318–333.
- [16] de Jongh, D. (1970). The maximality of the intuitionistic predicate calculus with respect to Heyting's Arithmetic. *The Journal of Symbolic Logic*, 36, 606.
- [17] de Jongh, D., Jumelet, M., & Montagna, F. (1991). On the proof of Solovay's theorem. *Studia Logica*, 50, 51–69.
- [18] Fagin, R., Halpern, J. Y., Moses, Y., & Vardi, M. Y. (1995). *Reasoning About Knowledge*. Cambridge, MA: MIT Press.

- [19] Fernández-Duque, D., & Joosten, J. J. (2014). Well-orders in the transfinite Japaridze algebra. *Logic Journal of the IGPL*, 22(6), 933–963.
- [20] Friedman, H. (1975). The disjunction property implies the numerical existence property. *Proceedings of the National Academy of Sciences*, 72(8), 2877–2878.
- [21] Friedman, H. (1978). Classically and intuitionistically provably recursive functions. In D. S. Scott, & G. H. Muller (Eds.) *Higher Set Theory*, vol. 699, (pp. 21–28). Springer Verlag.
- [22] Fujiwara, M., & Kurahashi, T. (2023). Prenex normalization and the hierarchical classification of formulas. arXiv:2302.11808 [math.LO].
- [23] Gabbay, D. M., Kurucz, A., Wolter, F., & Zakharyashev, M. (2003). *Many-dimensional modal logics: Theory and applications*. Elsevier.
- [24] Gabbay, D. M., Shehtman, V. B., & Skvortsov, D. P. (2009). *Quantification in Nonclassical Logic, Volume 1*. Amsterdam: Elsevier.
- [25] Goldblatt, R. (2011). *Quantifiers, propositions and identity, admissible semantics for quantified modal and substructural logics*. Cambridge University Press.
- [26] Hájek, P., & Pudlák, P. (1993). *Metamathematics of First Order Arithmetic*. Berlin, Heidelberg, New York: Springer-Verlag.
- [27] Hao, Y., & Tournakis, G. (2021). An arithmetically complete predicate modal logic. *Bulletin of the Section of Logic*.
URL <https://czasopisma.uni.lodz.pl/bulletin/article/view/8441>
- [28] Hughes, G. E., & Cresswell, M. J. (1996). *A New Introduction to Modal Logic*. Routledge.
- [29] Iemhoff, R. (2001). On the admissible rules of intuitionistic propositional logic. *Journal of Symbolic Logic*, (pp. 281–294).
- [30] Japaridze, G. K. (1988). The polymodal provability logic. In *Intensional logics and logical structure of theories: material from the fourth Soviet-Finnish symposium on logic, Telavi, May 20–24, 1985*, (pp. 16–48). Metsniereba. In Russian.
- [31] Kikot, S., Kurucz, A., Tanaka, Y., Wolter, F., & Zakharyashev, M. (2019). Kripke completeness of strictly positive modal logics over meet-semilattices with operators. *The Journal of Symbolic Logic*, 84(2), 533–588.
- [32] Kontchakov, R., Kurucz, A., & Zakharyashev, M. (2005). Undecidability of first-order intuitionistic and modal logics with two variables. *Bulletin of Symbolic Logic*, 11(3), 428–438.

- [33] Kreisel, G. (1951). On the interpretation of non-finitist proofs — Part I. *The Journal of Symbolic Logic*, 16(4), 241–267.
- [34] Kripke, S. A. (1963). Semantical considerations on modal logic. *Acta Philosophica Fennica*, 16, 83–94.
- [35] Kurahashi, T. (2013). On predicate provability logics and binumerations of fragments of Peano Arithmetic. *Archive for Mathematical Logic*, 52, 871–880.
- [36] Kurahashi, T. (2022). On inclusions between quantified provability logics. *Studia Logica*, 110, 165–188.
- [37] Kuroda, S. (1951). Intuitionistische Untersuchungen der formalistischen Logik. *Nagoya Mathematical Journal*, 2, 35–47.
- [38] Ladner, R. E. (1977). The computational complexity of provability in systems of modal propositional logic. *SIAM journal on computing*, 6(3), 467–480.
- [39] Leivant, D. (1979). *Absoluteness in intuitionistic logic*, vol. 73. Amsterdam: Mathematical Centre Tract. The Thesis was originally published in 1975.
- [40] Litak, T., & Wolter, F. (2005). All finitely axiomatizable tense logics of linear time flows are coNP-complete. *Studia Logica*, 81, 153–165.
- [41] McGee, V. R. (1985). *Truth and Necessity in Partially Interpreted Languages*. Ph.D. thesis, University of California, Berkeley.
- [42] Mojtabedi, M. (2022). On provability logic of HA. arXiv:2206.00445 [math.LO].
- [43] Myhill, J. (1973). A note on indicator functions. *Proceedings of the AMS*, 39, 181–183.
- [44] Pakhomov, F. (2014). On the complexity of the closed fragment of Japaridze’s provability logic. *Archive for Mathematical Logic*, 53(7–8), 949–967.
- [45] Rybakov, M., & Shkatov, D. (2019). Undecidability of first-order modal and intuitionistic logics with two variables and one monadic predicate letter. *Studia Logica*, 107(4), 695–717.
- [46] Rybakov, V. V. (1997). *Admissibility of logical inference rules*. Studies in Logic. Amsterdam: Elsevier.
- [47] Shapirovsky, I. (2008). PSPACE-decidability of Japaridze’s polymodal logic. In C. Areces, & R. Goldblatt (Eds.) *Advances in Modal Logic* 7, (pp. 289–304). College Publications.
- [48] Solovay, R. M. (1976). Provability interpretations of modal logic. *Israel Journal of Mathematics*, 28, 33–71.

- [49] Svyatlovskiy, M. V. (2018). Axiomatization and polynomial solvability of strictly positive fragments of certain modal logics. *Mathematical Notes*, 103, 952–967.
- [50] Troelstra, A. S. (1973). *Metamathematical Investigation of Intuitionistic Arithmetic and Analysis*, vol. 344 of *Lecture Notes in Mathematics*. Berlin: Springer.
- [51] Vardanyan, V. A. (1986). Arithmetic complexity of predicate logics of provability and their fragments. *Doklady Akad. Nauk SSSR*, 288(1), 11–14. In Russian. English translation in *Soviet Mathematics Doklady* 33, 569–572 (1986).
- [52] Vardanyan, V. A. (1988). Bounds on the arithmetical complexity of predicate logics of provability. In S. N. Adyan (Ed.) *Questions of Cybernetics: Complexity of Computation and Applied Mathematical Logic*, vol. 134 of *Scientific Council of the USSR Academy of Sciences on the complex problem “Cybernetics”*, (pp. 46–72). Academy of Sciences of the USSR. In Russian.
- [53] Visser, A. (1985). Evaluation, provably deductive equivalence in Heyting’s Arithmetic of substitution instances of propositional formulas. Logic Group Preprint Series 4, Faculty of Humanities, Philosophy, Utrecht University, Janskerkhof 13, 3512 BL Utrecht.
- [54] Visser, A. (2002). Substitutions of Σ_1^0 -sentences: explorations between intuitionistic propositional logic and intuitionistic arithmetic. *Annals of Pure and Applied Logic*, 114, 227–271.
- [55] Visser, A., & de Jonge, M. (2006). No escape from Vardanyan’s theorem. *Archive for Mathematical Logic*, 45(5), 539–554.
- [56] Visser, A., & Zoethout, J. (2019). Provability logic and the completeness principle. *Annals of Pure and Applied Logic*, 170(6), 718–753.
- [57] Wolter, F., & Zakharyashev, M. (2001). Decidable fragments of first-order modal logics. *The Journal of Symbolic Logic*, 66(3), 1415–1438.
- [58] Yavorsky, R. E. (2002). On arithmetical completeness of first-order logics of provability. In F. Wolter, H. Wansing, M. de Rijke, & M. Zakharyashev (Eds.) *Advances in Modal Logic 3*, (pp. 1–16). World Scientific Publishing Co. Pte. Ltd.