

HOCHSCHILD COHOMOLOGY OF THE SECOND KIND: KOSZUL DUALITY AND MORITA INVARIANCE

J. HOLSTEIN, A. LAZAREV, AND A. GUAN

ABSTRACT. We define Hochschild cohomology of the second kind for differential graded (dg) or curved algebras as a derived functor in a compactly generated derived category of the second kind, and show that it is invariant under Morita equivalence of the second kind. A bimodule version of Koszul duality is constructed and used to show that Hochschild cohomology of the second kind is preserved under (nonconilpotent) Koszul duality. Hochschild cohomology of the second kind of an algebra often computes the ordinary Hochschild cohomology of geometrically meaningful dg categories. Examples include the category of infinity local systems on a topological space, the bounded derived category of a complex algebraic manifold and the category of matrix factorizations.

CONTENTS

1. Introduction	1
2. Notation and Conventions	3
2.1. Pseudocompact algebras	3
2.2. DG categories	4
3. Twisted modules	4
3.1. DG modules and twistings	4
3.2. DG Morita equivalence	5
4. Hochschild cohomology of the first and second kind	5
4.1. Hochschild cohomology of the first kind	6
4.2. Model structures of the second kind and Koszul duality	6
4.3. Hochschild cohomology of the second kind	10
5. Bimodule Koszul duality	12
5.1. Koszul duality for the cobar construction	12
5.2. Koszul duality for the bar constructions	14
6. Curved and non-augmented cases	16
7. Examples of Hochschild cohomology of the second kind	18
7.1. Preliminaries	18
7.2. Complex algebraic manifolds	19
7.3. Topological spaces	20
7.4. Matrix Factorizations	21
References	22

1. INTRODUCTION

Hochschild cohomology of an associative algebra (or, more generally, a dg algebra) is well-known to be invariant under Morita equivalence. A related fundamental result, or a

This work was partially supported by EPSRC grant EP/T029455/1. The second author acknowledges support by the Deutsche Forschungsgemeinschaft under Germany’s Excellence Strategy – EXC 2121 “Quantum Universe” – 390833306.

series of results, is the invariance of Hochschild cohomology under Koszul duality. In other words, Hochschild cohomology of a dg algebra and that of its Koszul dual dg coalgebra are isomorphic, together with various structures that they possess; we refer to [18] for the most structured version of this result and its history.

At the same time, there is another version of Hochschild cohomology (sometimes called Hochschild cohomology of the second kind or compactly supported Hochschild cohomology, [21]) which is used e.g. in the study of categories of matrix factorizations. There are also Koszul duality theorems involving coalgebras that are not necessarily conilpotent, cf. [14, 7]. In this paper we investigate the analogues of the above-mentioned results in the context of global (i.e. non-conilpotent) Koszul duality and Hochschild cohomology of the second kind.

In more detail, our results are as follows. After recalling in Section 3 the notion of a twisted complex over a dg category and, as a special case, that of a twisted finitely generated module over a dg algebra together with Morita equivalence for dg categories, we give a proof of the Morita invariance of Hochschild cohomology. Our proof is constructed in such a way that it is suitable for generalization to homological algebra of the second kind. We review coderived categories of coalgebras and two versions of nonconilpotent Koszul duality between comodules over a (not necessarily conilpotent) dg coalgebra and modules over its Koszul dual dg algebra, following [22] and [14]. In particular, we recall the compactly generated derived category of the second kind $D_c^{\text{II}}(A)$ for a dg algebra A , generated by its subcategory $\text{Perf}^{\text{II}}(A)$ of compact objects. $D_c^{\text{II}}(A)$ is equivalent to the coderived category $D^{\text{co}}(\check{B}A)$ of its Koszul dual.

Definition 4.18. *The Hochschild cohomology complex of the second kind of a dg algebra A is $\text{HH}_c^{\text{II}}(A) = \text{RHom}_{D_c^{\text{II}}(A \otimes A^{\text{op}})}(A, A)$.*

A map of dg or curved algebras $f : A \rightarrow B$ is a *Morita equivalence of the second kind* if it induces an equivalence $D_c^{\text{II}}(A) \simeq D_c^{\text{II}}(B)$ and we show Morita invariance for Hochschild cohomology:

Theorem 4.20. *A Morita equivalence of the second kind between dg algebras $F : A \rightarrow B$ induces a quasi-isomorphism $\text{HH}_c^{\text{II}}(A) \simeq \text{HH}_c^{\text{II}}(B)$ of dg algebras.*

This is the content of Section 4. At this point we should warn the reader that our version of Hochschild cohomology of the second kind is *not* the same as that of Positselski and Polishchuk [21] but rather lies between the ordinary Hochschild cohomology and that of Positselski-Polishchuk, cf. Remark 5.9. Our definition is less elementary than the one in [21] but has the advantage of being compatible with Koszul duality; in favourable cases the two definitions are equivalent.

Note that Hochschild cohomology, be it of the first or second kind, is constructed as a (co)derived functor of bi(co)modules; whereas Koszul duality is usually formulated as an equivalence between one-sided modules and comodules. It is natural, therefore, to establish Koszul duality as an equivalence between bimodules over an augmented algebra and bicomodules over a suitably Koszul dual coalgebra; note that this is the approach of [18]. This is done in Section 5, both in the context of conilpotent and non-conilpotent Koszul duality. In fact, our result is slightly more general and establishes compatibility of Koszul duality with tensor products of dg algebras and dg coalgebras. A consequence of this compatibility is a Quillen equivalence between the model categories of C -bicomodules for a dg coalgebra C and dg bimodules over its cobar-construction ΩC . In the case when C is conilpotent, the weak equivalences on the ΩC -bimodule side are the ordinary quasi-isomorphisms (and our result reduces to Keller's [18, Proposition 2.4]) but in the non-conilpotent case they are more subtle (closer to isomorphisms).

We deduce:

Corollary 5.7. *For any dg algebra A , there is a quasi-isomorphisms of dg algebras*

$$\mathrm{HH}_c^{\mathrm{II}}(A) \simeq \mathrm{HH}(\check{\mathrm{B}}A).$$

Here $\check{\mathrm{B}}A$ is the *extended* bar-construction of A , cf. [1, 14, 7] concerning this notion.

Similarly, for a dg algebra A , we establish two further types of Koszul duality. One is a Quillen equivalence between the category of dg A -bimodules with the ordinary model structure (i.e. with quasi-isomorphisms for weak equivalences) and bicomodules over $\mathrm{B}A$, the bar-construction of A ; this is also essentially Keller's result in op. cit. The other is a Quillen equivalence between the category of A -bimodules with the compactly generated model structure of the second kind and $\check{\mathrm{B}}A$ -bicomodules.

For simplicity we state and prove our results first in the differential graded and augmented case and formulate the general results for curved, not necessarily augmented algebras in Section 6.

In particular, Definition 4.18, Theorem 4.20 and Corollary 5.7 hold more generally for curved algebras. The proofs are mostly the same except that in the curved setting there is no Yoneda embedding available as a curved algebra is not a left or right module over itself. To circumvent this difficulty, we construct for every curved algebra A a dg algebra A' with $\mathrm{D}_c^{\mathrm{II}}(A) \cong \mathrm{D}_c^{\mathrm{II}}(A')$ and so, $\mathrm{HH}_c^{\mathrm{II}}(A) \cong \mathrm{HH}_c^{\mathrm{II}}(A')$. It is interesting to observe that the dg algebra A' is acyclic, in particular its ordinary derived category is trivial as well as the ordinary Hochschild cohomology.

Finally, in Section 7 we give some examples of Hochschild cohomology of the second kind of a dg or curved algebra A that can be reduced to Hochschild cohomology of the *first kind* of the dg category of perfect A -modules of the second kind, which often has geometric or topological meaning. Results of this sort were first obtained in [21]. Specifically, with this reduction we show the following:

- If A is the Dolbeault algebra of a smooth complex algebraic manifold X then $\mathrm{HH}_c^{\mathrm{II}}(A)$ is Hochschild cohomology of a dg model of the bounded derived category of coherent sheaves on X ,
- If $\mathcal{A}^*(M)$ is the de Rham algebra of a smooth compact manifold M then $\mathrm{HH}_c^{\mathrm{II}}(\mathcal{A}^*(M))$ is Hochschild cohomology of the category of infinity local systems on M ,
- If R_w is $\mathbb{Z}/2$ -graded regular commutative algebra concentrated in even degrees with curvature element w then, under a technical assumption, $\mathrm{HH}_c^{\mathrm{II}}(R_w)$ is Hochschild cohomology of the category of matrix factorizations $\mathrm{MF}(R, w)$ which encodes the structure of the hypersurface singularity defined by w .

2. NOTATION AND CONVENTIONS

We fix a ground field k throughout. The category of (cohomologically \mathbb{Z} -graded) differential graded (dg) vector spaces over k will be denoted by DGVect . The shift of a dg vector space A is denoted by $A[1]$ so that $A[1]^i = A^{i+1}$. The category DGVect is monoidal with respect to the tensor product; monoids and comonoids in it are dg algebras and dg coalgebras respectively. A *curved* algebra (A, d, h) is a graded algebra A equipped with an element $h \in A^2$ and a derivation d such that $d^2(a) = [h, a]$. If $h = 0$, a curved algebra can be viewed as a dg algebra. A curved module (M, d) over (A, d, h) is a graded A -module M with a derivation d such that $d^2(m) = hm$. The notions of a curved coalgebra and a comodule over it can be defined by duality. More details on curved (co)algebras and (co)modules can be found e.g. in [22].

2.1. Pseudocompact algebras. Instead of (dg or curved) coalgebras, we will consider (dg or curved) pseudocompact algebras, i.e. topological algebras arising as inverse limits of finite-dimensional discrete algebras.

Taking (continuous) duals gives a contravariant equivalence of categories between coalgebras and pseudocompact algebras, and similarly between right comodules over C and right pseudocompact modules over C^* . Local augmented pseudocompact algebras correspond to conilpotent coalgebras.

A tensor product of pseudocompact algebras or modules is always assumed completed, equivalently it is the linear dual of the tensor product of the dual coalgebras or comodules.

We denote the dual of a dg vector space V by V^* . If V is discrete this is $\text{Hom}(V, k)$ equipped with the natural inverse limit topology. If V is pseudocompact this is the discrete k -module of continuous maps $V \rightarrow k$. In particular this ensures $V^{**} \cong V$.

2.2. DG categories. A *dg category* \mathbf{A} is a category enriched over DGVect ; in particular, for each pair of objects $X, Y \in \mathbf{A}$, homomorphisms from X to Y form a dg k -module which we denote by $\underline{\text{Hom}}_{\mathbf{A}}(X, Y)$. A *dg functor* is a functor enriched over DGVect . Any dg category \mathbf{A} has an associated ordinary category $\text{Ho } \mathbf{A}$, called its *homotopy category*, whose objects are the same as \mathbf{A} but whose morphisms are defined by $\text{Ho } \mathbf{A}(X, Y) = H^0(\underline{\text{Hom}}_{\mathbf{A}}(X, Y))$.

A dg functor $F: \mathbf{A} \rightarrow \mathbf{B}$ is a *quasi-equivalence* if

- (1) F induces quasi-isomorphisms $\underline{\text{Hom}}_{\mathbf{A}}(X, Y) \rightarrow \underline{\text{Hom}}_{\mathbf{B}}(FX, FY)$ for any objects X, Y in \mathbf{A} .
- (2) $H^0(F)$ is essentially surjective (or, equivalently assuming (1), $H^0(F)$ is an equivalence of categories).

3. TWISTED MODULES

3.1. DG modules and twistings. Let \mathbf{A} be a dg category. A (*right*) *dg \mathbf{A} -module* is a dg functor $M: \mathbf{A}^{\text{op}} \rightarrow \text{DGVect}$.

We denote by $\text{DGMod-}\mathbf{A}$ the category of dg \mathbf{A} -modules. There is a natural map $h: \mathbf{A} \rightarrow \text{DGMod-}\mathbf{A}$, sending an object $Y \in \mathbf{A}$ to $\underline{\text{Hom}}(-, Y)$, called the *Yoneda embedding*; a dg version of the usual Yoneda lemma says that h is fully faithful. Similarly a *left* dg \mathbf{A} -module is a right dg \mathbf{A}^{op} -module, i.e. a dg functor $N: \mathbf{A} \rightarrow \text{DGVect}$. We will always specify if we consider a left dg module; by default a dg module will be a right module.

A dg \mathbf{A} -module M is *acyclic* if it is acyclic pointwise, i.e. $M(A)$ is an acyclic complex for all $A \in \mathbf{A}$. The *derived category* $\text{D}(\mathbf{A})$ of dg \mathbf{A} -modules is the Verdier quotient of $\text{Ho}(\text{DGMod-}\mathbf{A})$ by acyclic dg \mathbf{A} -modules. There is a model category structure on $\text{DGMod-}\mathbf{A}$, where weak equivalences are pointwise quasi-isomorphisms and fibrations are pointwise surjections. All objects are fibrant; we denote by $\text{Cof}(\mathbf{A})$ the cofibrant objects in $\text{DGMod-}\mathbf{A}$. By general results on model categories, $\text{Ho } \text{Cof}(\mathbf{A}) = \text{D}(\mathbf{A})$.

Twisted complexes were first defined in [6] and later redefined in [4].

Definition 3.1. A (*two-sided*) *twisted complex* over \mathbf{A} is a formal expression $(\bigoplus_{i=1}^n C_i[r_i], q)$, where $C_i \in \mathbf{A}$, $r_i \in \mathbb{Z}$, $n \geq 0$, $q = (q_{ij})$, $q_{ij} \in \text{Hom}(C_j[r_j], C_i[r_i])$ homogeneous of degree 1 such that $dq + q^2 = 0$. A twisted complex is *one-sided* if $q_{ij} = 0$ for all $i \geq j$.

For two twisted complexes C and C' , the space of *morphisms of twisted complexes* $\text{Hom}(C, C')$ is the \mathbb{Z} -graded k -module of matrices $f = (f_{ij})$, $f_{ij} \in \text{Hom}(C_j[r_j], C'_i[r'_i])$ with differential $df = (df_{ij}) + q'f - (-1)^{|f_{ij}|}fq$. Composition of morphisms is usual matrix multiplication.

We denote the dg category of twisted complexes over \mathbf{A} by $\text{Tw}(\mathbf{A})$ and the dg category of one-sided twisted complexes by $\text{tw}(\mathbf{A})$. In [6], these were respectively denoted $\text{Pre-Tr}(\mathbf{A})$ and $\text{Pre-Tr}^+(\mathbf{A})$; furthermore, $\text{tw}(\mathbf{A}) = \text{Pre-Tr}^+(\mathbf{A})$ can alternatively be defined as the pretriangulated hull of \mathbf{A} , that is, it is the closure of \mathbf{A} under shifts and cones.

There is a natural functor from twisted complexes over a dg category \mathbf{A} to right dg \mathbf{A} -modules: We send $(\bigoplus_{i=1}^n C_i[r_i], q)$ to $(\bigoplus_{i=1}^n \text{Hom}(A, C_i)[r_i], q_*)$.

As an example, consider the case where A is a dg algebra (A, d_A) , considered as a dg category with one object. Recall that a *Maurer–Cartan element* in A is an element $x \in A^1$ such that $dx + x^2 = 0$, and that the set of all Maurer–Cartan elements in A is denoted by $\text{MC}(A)$. Then a twisted complex over A is a pair (M, q) where $M \cong V \otimes A$ as an A -module for some finite-dimensional \mathbb{Z} -graded vector space V , and $q \in \text{MC}(\text{End } V \otimes A)$. The A -module M becomes a (right) dg A -module when equipped with the differential $1 \otimes d_A + q$, and in fact, every dg A -module structure on the A -module M arises this way, as noted in [10, Remark 3.2]. A twisted complex over A is therefore precisely a *finitely generated* twisted A -module in the following sense.

Definition 3.2. A *twisted A -module* over a dg algebra A is a dg A -module that is free as an A -module after forgetting the differential, that is, it is isomorphic as an A -module to $V \otimes A$ for some graded vector space V . A *finitely generated twisted A -module* is a twisted A -module $V \otimes A$ with V finite-dimensional.

Note that there is a slight clash of terminology here; nevertheless this is unavoidable as we wish to refer to non-finitely generated twisted modules later. Note also that $(M, 1 \otimes d_A)$ above is a dg $(\text{End } V \otimes A)$ - A -bimodule whose differential has been *twisted* by the element $q \in \text{MC}(\text{End } V \otimes A)$. More generally, we have the following.

Definition 3.3. Let (A, d_A) be a dg algebra and $x \in \text{MC}(A)$.

- (1) The *twisted algebra of A by x* , denoted $A^x = (A, d^x)$, is the dg algebra with the same underlying algebra as A and differential $d^x(a) = d_A(a) + [x, a]$.
- (2) Let (M, d_M) be a *left* dg A -module. The *twisted module of M by x* , denoted $M^{[x]} = (M, d^{[x]})$, is the left dg A^x -module with the same underlying module structure as M and differential $d^{[x]}(m) = d(m) + xm$.

The condition $x \in \text{MC}(A)$ ensures that $M^{[x]}$ is indeed a left dg A^x -module, and that furthermore, if M is a dg A - B -bimodule for some dg algebra B , then $M^{[x]}$ is a dg A^x - B -bimodule, that is, the right B -module action remains compatible with the new differential.

A *perfect* dg A -module for a dg category A is a cofibrant dg A -module that is homotopy equivalent to a direct summand of a module in $\text{tw}(A)$. We denote by $\text{Perf}(A)$ the full dg subcategory of $\text{Cof}(A)$ consisting of perfect modules. It can be shown that $\text{D}(A)$ is a *compactly generated* triangulated category, and that $\text{Ho Perf}(A)$ consists precisely the compact objects in $\text{D}(A)$, cf. [28].

3.2. DG Morita equivalence. A dg functor $F: A \rightarrow B$ is a (*dg*) *Morita equivalence* if the induced map $F_!: \text{D}(A) \rightarrow \text{D}(B)$ is an equivalence of triangulated categories. Equivalently, F is a Morita equivalence if it induces a quasi-equivalence $\text{tw}(A) \rightarrow \text{tw}(B)$, $\text{Perf}(A) \rightarrow \text{Perf}(B)$ or $\text{Cof}(A) \rightarrow \text{Cof}(B)$. (The first two statements follow from the fact that $\text{D } A$ is compactly generated by $\text{Perf}(A)$.) We will abuse notation and also denote these maps by $F_!$.

A main result of [6] is that the Yoneda embedding $h: A \rightarrow \text{tw}(A) \hookrightarrow \text{Tw}(A)$ induces dg equivalences $\text{tw}(A) \rightarrow \text{tw}(\text{tw}(A))$ and $\text{Tw}(A) \rightarrow \text{Tw}(\text{Tw}(A))$. In particular, this implies that $h_!: \text{Perf}(A) \rightarrow \text{Perf}(\text{Perf}(A))$ is a quasi-equivalence, so the Yoneda embedding is a Morita equivalence. In fact, $\text{Perf}(A)$ is a Morita fibrant replacement of A in the sense of [17, Section 4.6].

4. HOCHSCHILD COHOMOLOGY OF THE FIRST AND SECOND KIND

In this section we recall Hochschild cohomology for dg categories and define Hochschild cohomology of the second kind for dg algebras. It is well-known that Hochschild cohomology is Morita invariant; we will give an alternative proof of this fact and show how the proof can be adapted to give a Morita invariance result in the second kind case.

4.1. Hochschild cohomology of the first kind. Let \mathbf{A} be a dg category. The *Hochschild cohomology* of \mathbf{A} is $\mathrm{HH}(\mathbf{A}) = \mathrm{RHom}_{\mathbf{A} \otimes \mathbf{A}^{\mathrm{op}}}(\mathbf{A}, \mathbf{A})$. This is a dg algebra considered as the endomorphisms of one object.

We show that Hochschild cohomology is invariant under Morita equivalence. This is well-known, cf. [27, Corollary 8.2] but we give a proof that is different from the standard proofs and will be used as a model when proving the analogous result for Hochschild cohomology of the second kind.

Theorem 4.1. *A Morita equivalence $F: \mathbf{A} \rightarrow \mathbf{B}$ induces a quasi-isomorphism $\mathrm{HH}(\mathbf{A}) \simeq \mathrm{HH}(\mathbf{B})$ of dg algebras.*

Proof. Since $F: \mathbf{A} \rightarrow \mathbf{B}$ induces a map $(F \otimes F^{\mathrm{op}})_!: \mathrm{DGMod}(\mathbf{A} \otimes \mathbf{A}^{\mathrm{op}}) \rightarrow \mathrm{DGMod}(\mathbf{B} \otimes \mathbf{B}^{\mathrm{op}})$ sending \mathbf{A} to \mathbf{B} , and all bimodules admit cofibrant replacements, it suffices to show that $(F \otimes F^{\mathrm{op}})_!: \mathrm{Cof}(\mathbf{A} \otimes \mathbf{A}^{\mathrm{op}}) \rightarrow \mathrm{Cof}(\mathbf{B} \otimes \mathbf{B}^{\mathrm{op}})$ is a quasi-equivalence.

Indeed, the restriction of $(F \otimes F^{\mathrm{op}})_!$ to $\mathrm{Perf}(\mathbf{A} \otimes \mathbf{A}^{\mathrm{op}})$ fits into the commutative diagram

$$\begin{array}{ccc} \mathrm{Perf}(\mathbf{A} \otimes \mathbf{A}^{\mathrm{op}}) & \xrightarrow{(h \otimes h^{\mathrm{op}})_!} & \mathrm{Perf}(\mathrm{Perf}(\mathbf{A}) \otimes \mathrm{Perf}(\mathbf{A})^{\mathrm{op}}) \\ \downarrow (F \otimes F^{\mathrm{op}})_! & & \downarrow (F_! \otimes F_!^{\mathrm{op}})_! \\ \mathrm{Perf}(\mathbf{B} \otimes \mathbf{B}^{\mathrm{op}}) & \xrightarrow{(h \otimes h^{\mathrm{op}})_!} & \mathrm{Perf}(\mathrm{Perf}(\mathbf{B}) \otimes \mathrm{Perf}(\mathbf{B})^{\mathrm{op}}) \end{array}$$

where the rows are quasi-equivalences as the Yoneda embedding h induces a quasi-equivalence $h_!: \mathrm{Perf}(\mathbf{A}) \rightarrow \mathrm{Perf}(\mathrm{Perf}(\mathbf{A}))$, and $(F_! \otimes F_!^{\mathrm{op}})_!$ is a quasi-equivalence as F is a Morita equivalence. Hence $\mathrm{Ho}(F \otimes F^{\mathrm{op}})_!: \mathrm{D}(\mathbf{A} \otimes \mathbf{A}^{\mathrm{op}}) \rightarrow \mathrm{D}(\mathbf{B} \otimes \mathbf{B}^{\mathrm{op}})$ restricts to an equivalence of categories on the compact objects, and any such cocontinuous functor between compactly generated triangulated categories is an equivalence, by [25, Lemma 3.3]. Finally the endomorphisms of corresponding objects are multiplicatively quasi-isomorphic. \square

Corollary 4.2. *Let \mathbf{A} be a dg category. Then the Yoneda embedding $\mathbf{A} \rightarrow \mathrm{Perf}(\mathbf{A})$ induces a quasi-isomorphism*

$$\mathrm{HH}(\mathbf{A}) \simeq \mathrm{HH}(\mathrm{Perf}(\mathbf{A})).$$

Proof. Immediate as $\mathbf{A} \rightarrow \mathrm{Perf}(\mathbf{A})$ is a Morita equivalence. \square

4.2. Model structures of the second kind and Koszul duality. In this section we give the other model structures that will feature in the remainder of this paper. These model structures will all give rise to “derived categories of the second kind”, the collective name given to derived categories which arise from localizing at some collection of weak equivalences that are finer than quasi-isomorphism.

There are many such derived categories; we will recall the coderived category of a pseudo-compact algebra, following [22], and the compactly generated derived category of the second kind defined of a dg algebra, as in [14]. These have the feature of being compatible with Koszul duality, which we will also recall.

We first define, following [22], that in a category of dg or curved modules or comodules an object is *coacyclic* (resp. *contraacyclic*) if it lies in the smallest subcategory containing all totalizations of exact triples and closed under cones and direct sums (resp. direct products).

Let now A be dg algebra and C be a pseudocompact dg algebra over a field k . The categories $\mathrm{DGMod}\text{-}C$ and $\mathrm{DGMod}\text{-}A$, of pseudocompact dg C -modules and dg A -modules respectively, admit the following model category structures of the second kind.

Theorem 4.3 (see [22, Theorem 8.2]). *For any pseudocompact dg algebra C , the category $\mathrm{DGMod}\text{-}C$ is a model category where a morphism $f: M \rightarrow N$ is*

- (1) a weak equivalence if the cone of the dual map $f^*: N^* \rightarrow M^*$ of C^* -comodules is coacyclic;
- (2) a fibration if it is surjective;
- (3) a cofibration if it has the left lifting property with respect to acyclic fibrations.

Equivalently to the dual having a coacyclic cone, we can also characterize weak equivalences directly as having contraacyclic cone.

The homotopy category of $\text{DGMod-}C$ is the *coderived category* of C , denoted by $\text{D}^{\text{co}}(C)$.

Theorem 4.4 (see [14, Theorem 4.6]). *Let A be a dg algebra. There is a model category structure on $\text{DGMod-}A$, where a morphism $f: M \rightarrow N$ is*

- (1) a weak equivalence if it induces a quasi-isomorphism

$$\underline{\text{Hom}}_A(T, M) \rightarrow \underline{\text{Hom}}_A(T, N)$$

for any finitely generated twisted A -module T ;

- (2) a fibration if it is surjective;
- (3) a cofibration if it has the left lifting property with respect to acyclic fibrations.

Where it is necessary to distinguish between the usual model structure of the first kind on dg A -modules (where weak equivalences are quasi-isomorphisms) and the model structure of Theorem 4.4, we will denote these model categories respectively by $(\text{DGMod-}A)^{\text{I}}$ and $(\text{DGMod-}A)^{\text{II}}$. We say that a dg A -module is *cofibrant of the second kind* if it is cofibrant in $(\text{DGMod-}A)^{\text{II}}$, and denote these cofibrant objects by $\text{Cof}_c^{\text{II}}(A)$.

The (*compactly generated*) *derived category of the second kind* $\text{D}_c^{\text{II}}(A)$ of dg A -modules is the homotopy category of $(\text{DGMod-}A)^{\text{II}}$.

Recall that the cobar construction ΩC of C is the tensor graded algebra $T\bar{C}^*[-1]$ with differential defined using the differential and multiplication on C . The model structures above are then related by the following statement, referred to as Koszul duality for the cobar construction.

Theorem 4.5. *Let C be a pseudocompact algebra. There is a Quillen equivalence*

$$G: (\text{DGMod-}C)^{\text{op}} \rightleftarrows (\text{DGMod-}\Omega C)_c^{\text{II}} : F.$$

We will sometimes write $G_C(N)$ for GN when we want to make the dependence on C clear.

The functor G sends a pseudocompact dg C -module N to the dg ΩC -module

$$GN := (N^* \otimes \Omega C)^{[\xi]},$$

where ξ is the canonical Maurer–Cartan element in $C \otimes \Omega C$. The twist makes sense as we regard $N^* \otimes \Omega C$ as a $(C \otimes \Omega C)^{\text{op}} \otimes \Omega C$ -module, where the left action of C comes from the right action of C on N and the $\Omega C^{\text{op}} \otimes \Omega C$ -action arises from the left and right multiplication on ΩC . Hence the twisted module $(N^* \otimes \Omega C)^{[\xi]}$ is a left dg $(C \otimes \Omega C)^{\xi}$ module and a right dg ΩC -module, and GN is obtained by forgetting the left $(C \otimes \Omega C)^{\xi}$ -action. More explicitly, GN can be written as a complex as follows:

$$N^* \leftarrow N^* \otimes \bar{C} \leftarrow N^* \otimes \bar{C}^2 \leftarrow \dots$$

This is, in fact, a double complex where the vertical differential (not indicated explicitly) is induced by the internal differentials in C and N whereas the horizontal differential is induced by the multiplication in C and its action on N^* . Explicitly, the horizontal differential is given by

$$d(c_0 \otimes \dots \otimes c_k \otimes n) = \sum_{i=0}^{k-1} (-1)^i c_0 \otimes \dots \otimes c_i c_{i+1} \dots \otimes c_k \otimes n + (-1)^{k+1} c_0 \otimes \dots \otimes c_k n$$

where the last term in the above sum corresponds to the twisting by ξ . Similarly, the functor F sends a dg ΩC -module M to the pseudocompact dg C -module

$$FM := (M^* \otimes C)^{[\xi]}$$

where we use the left ΩC -module structure on M^* and the left C -module structure on C to define the twist.

Remark 4.6. A version of Theorem 4.5, formulated in the language of coalgebras and comodules, is due to [22, Theorem 6.7]. Combining Theorem 4.5 with Positselski's result we see that if $A = \Omega C$ then the homotopy category $D_c^{\text{II}}(A)$ is precisely the coderived category of A defined by localizing at coacyclic dg A -modules.

In the case that C is augmented local (equivalently, that C^* is conilpotent), the cobar construction ΩC is cofibrant and one recovers the more classical statement with $(\text{DGMod-}\Omega C)^{\text{I}}$ on the right hand side.

More on the relationship between different (co)derived categories can be found in [14, Section 3.3].

For the sake of convenience, we will now sketch the proof of Theorem 4.5. We will later note that the arguments also work for (pseudocompact) modules over a curved pseudocompact algebras C . First note that the functors F and G are adjoint in the sense that there are natural isomorphisms

$$\text{Hom}_{\text{DGMod-}\Omega C}(GN, M) \cong \text{Hom}_{\text{DGMod-}C}(FM, N),$$

for N and M as above, as both sides are identified with the dg vector spaces $(N \otimes M)^{[\xi]}$. Composing the functors gives the pseudocompact dg C -module

$$FGN = (N \otimes \Omega C^* \otimes C)^{[\xi \otimes 1 + 1 \otimes \xi]},$$

which as a (double) complex, can be written as follows:

$$N \otimes C \leftarrow N \otimes \bar{C}^* \otimes C \leftarrow N \otimes (\bar{C}^*)^2 \otimes C \leftarrow \dots$$

The horizontal differential is described by a similar formula as above. Note, however, that due to the tensor factor C , the horizontal differential is acyclic except at the first term; in fact it gives the standard cobar resolution of N , the map from the above complex to N being induced by the action map $N \otimes C \rightarrow N$. Of course, one has to be careful calling this a resolution since C is a pseudocompact algebra and N is a pseudocompact C -module. This can be made sense of as follows. Consider the homotopy cofiber of the map $FGN \rightarrow N$; this is the double complex that can be represented as follows:

$$N \leftarrow N \otimes C \leftarrow N \otimes \bar{C}^* \otimes C \leftarrow N \otimes (\bar{C}^*)^2 \otimes C \leftarrow \dots$$

Its horizontal differential is acyclic. By taking canonical truncations this complex can be represented as a homotopy inverse limit of acyclic complexes of finite horizontal length, which are, therefore, contra-acyclic as pseudocompact C -modules. Thus, we obtain the following result, from which Theorem 4.5 follows.

Proposition 4.7. *The adjunction unit $FGN \rightarrow N$ is a cofibrant resolution of the pseudocompact dg C -module N .*

Analogous statements can be obtained if we instead start with a dg algebra A ; however, depending on whether we consider model structures of the first or second kind over $\text{DGMod-}A$, we will have two different equivalences. Let us denote by $\widehat{B}A$ the bar construction of A and by $\check{B}A$ the extended bar construction of A , both viewed as pseudocompact dg algebras.

As graded algebras these are respectively defined as $\widehat{T}\bar{A}^*[-1]$ and $\check{T}\bar{A}^*[-1]$, where \widehat{T} denotes the free *local* pseudocompact functor while \check{T} denotes the free pseudocompact functor on a graded pseudocompact vector space. (When dualizing, this corresponds to the free conilpotent coalgebra and the free coalgebra, respectively. Note that we are using $\widehat{B}A$ rather than BA to distinguish this construction from the usual bar construction on A , which would be a coalgebra dual to our $\widehat{B}A$.) As before, bar differentials are defined using the differential and multiplication on A ; see [14, Definition 2.5] in the extended bar case. We then have the following Koszul duality statements for the bar constructions, see [14, Theorem 4.7]:

Theorem 4.8. *There are Quillen equivalences*

$$(\mathrm{DGMod}\text{-}\widehat{B}A)^{\mathrm{op}} \rightleftarrows (\mathrm{DGMod}\text{-}A)^{\mathrm{I}}$$

and

$$(\mathrm{DGMod}\text{-}\check{B}A)^{\mathrm{op}} \rightleftarrows (\mathrm{DGMod}\text{-}A)_{\mathrm{c}}^{\mathrm{II}}.$$

We collect some useful facts about the model structure of the second kind:

- Proposition 4.9.** (1) *Twisted modules are cofibrant of the second kind if they are a union of finitely generated twisted modules. All cofibrant modules are retracts of such.*
- (2) *Any map $f : M \rightarrow N$ in $\mathrm{DGMod}\text{-}A$ with coacyclic or contra-acyclic cone is also a weak equivalence in the sense of Theorem 4.3.*

Proof. (1) Any $\check{B}A$ -module is fibrant, and may be written as a filtered limit of finite-dimensional modules. Under Koszul duality the finite-dimensional $\check{B}A$ -modules correspond to finitely generated twisted A -modules. An arbitrary cofibrant object is a retract of its cofibrant replacement (which is obtained as a union of finitely generated twisted modules).

- (2) Let $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ be an exact triple and T a finitely generated twisted A -module. Then $0 \rightarrow \mathrm{Hom}_A(T, L) \rightarrow \mathrm{Hom}_A(T, M) \rightarrow \mathrm{Hom}_A(T, N) \rightarrow 0$ is exact. Therefore the dg space of homomorphisms from T into the totalization of $L \rightarrow M \rightarrow N$ is acyclic. Thus the weak equivalences are closed under totalization of exact triples. They are automatically closed under direct products, and as any finitely generated twisted module is compact they are also closed under direct sums. \square

We now define perfect modules of the second kind.

Definition 4.10. A dg A -module is *perfect of the second kind* if it is homotopy equivalent to a direct summand of a two-sided twisted module in $\mathrm{Tw}(A)$. We denote by $\mathrm{Perf}^{\mathrm{II}}(A)$ the full dg subcategory of $\mathrm{Cof}_{\mathrm{c}}^{\mathrm{II}}(A)$ consisting of perfect modules of the second kind.

We note that any homomorphism $f : A \rightarrow B$ induces a dg functor $f_! : \mathrm{Perf}^{\mathrm{II}}(A) \rightarrow \mathrm{Perf}^{\mathrm{II}}(B)$ and there is a quasi-fully faithful Yoneda embedding $h : A \rightarrow \mathrm{Perf}^{\mathrm{II}}(A)$.

The following statement for $\mathrm{Perf}^{\mathrm{II}}$ is analogous to the corresponding statement for Perf .

Lemma 4.11. *For any dg category A , the Yoneda embedding induces a quasi-equivalence $\mathrm{Perf}^{\mathrm{II}}(A) \rightarrow \mathrm{Perf}^{\mathrm{II}}(\mathrm{Perf}^{\mathrm{II}}(A))$.*

Proof. The natural map $i : \mathrm{Tw}(A) \rightarrow \mathrm{Perf}^{\mathrm{II}}(A)$ is a Morita fibrant replacement by [26]. It suffices to show that $i' : \mathrm{Tw}^2(A) \rightarrow (\mathrm{Perf}^{\mathrm{II}})^2(A)$ is also a Morita fibrant replacement: the map $\mathrm{Perf}^{\mathrm{II}}(A) \rightarrow (\mathrm{Perf}^{\mathrm{II}})^2(A)$ is then a Morita fibrant replacement of the the functor $\mathrm{Tw}(A) \rightarrow \mathrm{Tw}^2(A)$ induced by the Yoneda embedding, which is a quasi-equivalence by the Definition of Tw . So i' is a Morita equivalence between Morita fibrant categories, and hence a quasi-equivalence, as the Morita model structure is a (left) Bousfield localisation of the usual Dwyer-Kan model structure.

But the map i' is the composition $\mathrm{Tw}^2(A) \rightarrow \mathrm{Perf}^{\mathrm{II}}(\mathrm{Tw}(A)) \rightarrow (\mathrm{Perf}^{\mathrm{II}})^2(A)$ and each of the factors is a Morita equivalences, obtained by applying the map i to $\mathrm{Tw}(A)$ and by applying $\mathrm{Perf}^{\mathrm{II}}$ to i respectively. Hence i' is itself a Morita equivalence. \square

Theorem 4.12. *Let A be a dg algebra. Then $\mathrm{HoPerf}^{\mathrm{II}}(A)$ is the category of compact objects in $\mathrm{D}_c^{\mathrm{II}}(A)$, and $\mathrm{D}_c^{\mathrm{II}}(A)$ is compactly generated.*

Proof. By definition, $\mathrm{HoPerf}^{\mathrm{II}}(A)$ consists of dg A -modules that are direct summands of finitely generated twisted A -modules, and by Proposition 4.9 it is clear that these are compact as finite-dimensional twisted modules are.

To prove the that $\mathrm{Perf}^{\mathrm{II}}(A)$ generates we use that $\mathrm{D}_c^{\mathrm{II}}(A)$ is by Koszul duality (Theorem 4.8) anti-equivalent as a triangulated category to the coderived category of pseudocompact dg modules over a pseudocompact dg algebra, which is cocompactly cogenerated by finite dimensional dg modules by [22, Section 5.5] (which correspond to finitely generated twisted A -modules).

Finally, to show that $\mathrm{Perf}^{\mathrm{II}}(A)$ contains all compact objects we use [19, Lemma 2.2] in the case $\mathcal{S} = \mathcal{R} = \mathrm{D}_c^{\mathrm{II}}(A)$. \square

4.3. Hochschild cohomology of the second kind. We define Morita equivalence of the second kind. We will then define Hochschild cohomology of the second kind for dg algebras, and prove that it is invariant under Morita equivalence of the second kind.

Definition 4.13. Let A and B be dg algebras. A morphism $F: A \rightarrow B$ is a *Morita equivalence of the second kind* if it induces an equivalence of triangulated categories $\mathrm{D}_c^{\mathrm{II}}(A) \rightarrow \mathrm{D}_c^{\mathrm{II}}(B)$.

Example 4.14. For any dg algebra A the counit map $\Omega\check{B}A \rightarrow A$ is a Morita equivalence of the second kind. This follows directly by [7, Theorem 6.4 and Corollary 6.6].

Morita equivalence can equivalently be characterized as follows.

Proposition 4.15. *A morphism $F: A \rightarrow B$ is a Morita equivalence of the second kind if and only if it induces a quasi-equivalence $\mathrm{Cof}_c^{\mathrm{II}}(A) \rightarrow \mathrm{Cof}_c^{\mathrm{II}}(B)$, $\mathrm{Perf}^{\mathrm{II}}(A) \rightarrow \mathrm{Perf}^{\mathrm{II}}(B)$ or $\mathrm{Tw}(A) \rightarrow \mathrm{Tw}(B)$.*

Proof. It follows from the model category structure that F is a Morita equivalence of the second kind if and only if the induced left Quillen functor $F_!$ induces a quasi-equivalence $\mathrm{Cof}_c^{\mathrm{II}}(A) \rightarrow \mathrm{Cof}_c^{\mathrm{II}}(B)$. But this is a quasi-equivalence if and only if it is a quasi-equivalence when restricted to compact objects by Theorem 4.12, and $\mathrm{Perf}^{\mathrm{II}}(A)$ is the category of compact objects in $\mathrm{Cof}_c^{\mathrm{II}}(A)$ by Theorem 4.12 again. Finally, as $\mathrm{Perf}^{\mathrm{II}}(A)$ is the idempotent completion of $\mathrm{Tw}(A)$, $\mathrm{Perf}^{\mathrm{II}}(A) \simeq \mathrm{Perf}^{\mathrm{II}}(B)$ follows from the quasi-equivalence $\mathrm{Tw}(A) \rightarrow \mathrm{Tw}(B)$. \square

If A is a dg algebra, then by Lemma 4.11 the Yoneda embedding $A \rightarrow \mathrm{Perf}^{\mathrm{II}}(A)$ is a Morita equivalence of the second kind.

Lemma 4.16. *For curved algebras A, B there is a quasi-equivalence*

$$\mathrm{Perf}^{\mathrm{II}}(A \otimes B) \simeq \mathrm{Perf}^{\mathrm{II}}(\mathrm{Perf}^{\mathrm{II}}(A) \otimes \mathrm{Perf}^{\mathrm{II}}(B))$$

Proof. Consider the following diagram

$$\begin{array}{ccc} \mathrm{Perf}^{\mathrm{II}}(A \otimes B) & \xrightarrow{(h^A \otimes h^B)_!} & \mathrm{Perf}^{\mathrm{II}}(\mathrm{Perf}^{\mathrm{II}}(A) \otimes \mathrm{Perf}^{\mathrm{II}}(A)^{\mathrm{op}}) \\ & \searrow h_!^{A \otimes B} & \downarrow i_! \\ & & \mathrm{Perf}^{\mathrm{II}}(\mathrm{Perf}^{\mathrm{II}}(A \otimes B)) \end{array}$$

Here $h_!^{A \otimes B}$ is a quasi-equivalence by Proposition 4.15 as the Yoneda embedding is a Morita equivalence of the second kind by Lemma 4.11. The functor $i : \text{Perf}^{\text{II}}(A) \otimes \text{Perf}^{\text{II}}(B) \rightarrow \text{Perf}^{\text{II}}(A \otimes B)$ is quasi-fully faithful and thus so is $i_!$. It follows that $(h^A \otimes h^B)_!$ is quasi-fully faithful and thus, is a quasi-equivalence. The claimed statement is therefore proved. \square

We will now consider the category of A -bimodules with the model structure from Theorem 4.4, $A\text{-DGMod-}A := \text{DGMod-}(A^{\text{op}} \otimes A)$.

Lemma 4.17. *The Hom functor $\text{Hom}_{A \otimes A^{\text{op}}}(-, -) : (A\text{-DGMod-}A)_c^{\text{II}} \otimes (A\text{-DGMod-}A)_c^{\text{II}} \rightarrow \text{DGVect}$ is a Quillen bifunctor if we put the projective model structure on DGVect .*

Proof. By adjointness it suffices to show that

$$\otimes_k : (A\text{-DGMod-}A)_c^{\text{II}} \otimes \text{DGVect} \rightarrow (A\text{-DGMod-}A)_c^{\text{II}}$$

is a Quillen bifunctor. So let $g : P \rightarrow Q$ be a cofibration in DGVect and $f : L \rightarrow M$ a cofibration in $(A\text{-DGMod-}A)_c^{\text{II}}$ and consider

$$f \square g : M \otimes P \amalg_{L \otimes P} L \otimes Q \rightarrow M \otimes Q.$$

To check that $f \square g$ is a cofibration if f and g are, it suffices to check the case that g is a generating cofibration.

Then we can write $g : k[n] \rightarrow \text{cone}(\text{id}_{k[n]})$ and we obtain $f \square g : M \oplus L[1] \rightarrow M \oplus M[1]$ which is a cofibration if f is.

If g is a generating acyclic cofibration it is of the form $0 \rightarrow \text{cone}(\text{id}_{k[n]})$. Then $f \square g : L \otimes Q \rightarrow M \otimes Q$ has cokernel $\text{cone}(f) \otimes \text{cone}(\text{id}_{k[n]})$ which is a totalization of a short exact sequence and thus coacyclic, and thus $f \square g$ is a weak equivalence by Proposition 4.9.

Finally, it is clear from the definition in Theorem 4.4 that if f is a weak equivalence so is $f \otimes \text{id}_B$ for B bounded finitely generated in DGVect . Thus if $L \rightarrow M$ is a weak equivalence then $L \otimes P \rightarrow M \otimes P$ and its pushout along $L \otimes P \rightarrow L \otimes Q$ are weak equivalences, as is $L \otimes Q \rightarrow M \otimes Q$. By the 2-out-of-3 property, this gives a weak equivalence $f \square g$. \square

We now write $\text{RHom}_A^{\text{II}}(M, N)$ for the derived hom space between M and N in $(\text{DGMod-}A)_c^{\text{II}}$.

Definition 4.18. Let A be a dg algebra and M a dg bimodule. The *Hochschild cohomology of the second kind* of A with coefficients in M is

$$\text{HH}_c^{\text{II}}(A, M) = \text{RHom}_{A \otimes A^{\text{op}}}^{\text{II}}(A, M).$$

We write $\text{HH}_c^{\text{II}}(A)$ for $\text{HH}_c^{\text{II}}(A, A)$.

Remark 4.19. We are concerned with Hochschild cohomology in this paper, but could equally define *Hochschild homology of the second kind* to be $A \otimes_{A \otimes A^{\text{op}}}^{L, \text{II}} A$ where the tensor product is derived in the model category $(A\text{-DGMod-}A)_c^{\text{II}}$.

This makes sense as the tensor product over $A \otimes A^{\text{op}}$ is a left Quillen bifunctor into DGVect with the projective model structure. This can be shown using that generating cofibrations in $(A\text{-DGMod-}A)_c^{\text{II}}$ are the images of injection of finite-dimensional comodules under Koszul duality. The definition of weak equivalences ensures that tensoring with the image of a finite-dimensional comodule preserves weak equivalences.

Theorem 4.20. *A Morita equivalence of the second kind $F : A \rightarrow B$ between dg algebras induces an isomorphism $\text{HH}_c^{\text{II}}(A) \cong \text{HH}_c^{\text{II}}(B)$ of dg algebras.*

Proof. We proceed as in the proof of Theorem 4.1. As before, it suffices to show that $(F \otimes F^{\text{op}})_! : \text{Cof}_c^{\text{II}}(A \otimes A^{\text{op}}) \rightarrow \text{Cof}_c^{\text{II}}(B \otimes B^{\text{op}})$ is a quasi-equivalence. The restriction of

$(F \otimes F^{\text{op}})_!$ to $\text{Perf}^{\text{II}}(A \otimes A^{\text{op}})$ fits into the commutative diagram

$$\begin{array}{ccc} \text{Perf}^{\text{II}}(A \otimes A^{\text{op}}) & \xrightarrow{(h \otimes h^{\text{op}})_!} & \text{Perf}^{\text{II}}(\text{Perf}^{\text{II}}(A) \otimes \text{Perf}^{\text{II}}(A)^{\text{op}}) \\ (F \otimes F^{\text{op}})_! \downarrow & & \downarrow (F_! \otimes F_!^{\text{op}})_! \\ \text{Perf}^{\text{II}}(B \otimes B^{\text{op}}) & \xrightarrow{(h \otimes h^{\text{op}})_!} & \text{Perf}^{\text{II}}(\text{Perf}^{\text{II}}(B) \otimes \text{Perf}^{\text{II}}(B)^{\text{op}}) \end{array}$$

where the rows are quasi-equivalences by Lemma 4.16, and $(F_! \otimes F_!^{\text{op}})_!$ is a quasi-equivalence as F is a Morita equivalence of the second kind. Hence by Theorem 4.12, the functor $\text{Ho}(F \otimes F^{\text{op}})_!: \text{D}_c^{\text{II}}(A \otimes A^{\text{op}}) \rightarrow \text{D}_c^{\text{II}}(B \otimes B^{\text{op}})$ restricts to an equivalence of categories on the compact objects, so is an equivalence. \square

5. BIMODULE KOSZUL DUALITY

The aim of this section is to generalize the Koszul duality statements, Theorem 4.5 and Theorem 4.8, to bimodules. Throughout this section, A and E will denote two augmented dg algebras, and $A\text{-DGMod-}E := \text{DGMod-}(A^{\text{op}} \otimes E)$ will denote the category of dg A - E -bimodules. Similarly C and D will denote two augmented pseudocompact dg algebras, and the notation $C\text{-DGMod-}D$ will be understood to mean pseudocompact bimodules, i.e. pseudocompact left C -modules that are also pseudocompact right D -modules.

5.1. Koszul duality for the cobar construction. We now obtain an analogue of Proposition 4.7 for pseudocompact bimodules by defining the following adjunction:

$$G: (C\text{-DGMod-}D)^{\text{op}} \rightleftarrows \Omega C\text{-DGMod-}\Omega D : F.$$

Let $\xi_C \in \text{MC}(C^{\text{op}} \otimes \Omega C^{\text{op}})$ and $\xi_D \in \text{MC}(D \otimes \Omega D)$ be the canonical Maurer–Cartan elements corresponding to the counits $\widehat{\text{B}}\Omega C^{\text{op}} \rightarrow C^{\text{op}}$ and $\widehat{\text{B}}\Omega D \rightarrow D$ of the adjunction $\Omega \dashv \widehat{\text{B}}$. Define

$$\xi := \xi_C \otimes 1 + 1 \otimes \xi_D \in C^{\text{op}} \otimes \Omega C^{\text{op}} \otimes D \otimes \Omega D,$$

then $\xi \in \text{MC}(C^{\text{op}} \otimes \Omega C^{\text{op}} \otimes D \otimes \Omega D)$. The functor G sends a pseudocompact dg C - D -bimodule N to

$$GN := (\Omega C \otimes N^* \otimes \Omega D)^{[\xi]},$$

where as before, GN is a C - D -bimodule. It is the bimodule cobar-construction of the $C^{\text{op}} \otimes D$ -module N . It can be written as the direct sum totalization of a double complex as follows:

$$(5.1) \quad \begin{array}{ccccccc} & \downarrow & & \downarrow & & \downarrow & \\ N^* \otimes \bar{D}^2 & \longleftarrow & \bar{C} \otimes N^* \otimes \bar{D}^2 & \longleftarrow & \bar{C}^2 \otimes N^* \otimes \bar{D}^2 & \longleftarrow & \dots \\ & \downarrow & & \downarrow & & \downarrow & \\ N^* \otimes \bar{D} & \longleftarrow & \bar{C} \otimes N^* \otimes \bar{D} & \longleftarrow & \bar{C}^2 \otimes N^* \otimes \bar{D} & \longleftarrow & \dots \\ & \downarrow & & \downarrow & & \downarrow & \\ N^* & \longleftarrow & \bar{C} \otimes N^* & \longleftarrow & \bar{C}^2 \otimes N^* & \longleftarrow & \dots \end{array}$$

This is, in fact, a triple complex where the third differential (not indicated explicitly) is induced on each term by the internal differentials in C , D and N . The n th row of the above complex is $G_C(N \otimes \bar{D}^n)$ and the n th column is $G_D(\bar{C}^n \otimes N)$.

Similarly, the functor F sends a dg ΩC - ΩD -bimodule M to the pseudocompact dg C - D -bimodule

$$FM := (C \otimes M^* \otimes D)^{[\xi]},$$

and the functors F and G are adjoint in the sense that there is a natural isomorphism of dg vector spaces

$$\mathrm{Hom}_{\Omega C\text{-DGMod-}\Omega D}(GN, M) \cong \mathrm{Hom}_{C\text{-DGMod-}D}(FM, N).$$

Indeed, both sides above are identified with the dg vector spaces $(N \otimes M)^{[\xi]}$.

Composing the functors, we obtain the $\Omega C^{\mathrm{op}} \otimes \Omega D$ -module

$$FGN = (C \otimes \Omega C \otimes N \otimes \Omega D \otimes D)^{[\xi \otimes 1 + 1 \otimes \xi]}.$$

This is a double complex obtained from (5.1) by tensoring each entry with C from the left and D from the right.

Furthermore, this new double complex can be ‘augmented’ by adding to it as a (-1) -st row the cobar-resolution $F_C G_C(N)$ of the C -module N . We obtain

$$\begin{array}{ccccccc} & \downarrow & & \downarrow & & \downarrow & \\ C \otimes N \otimes \bar{D}^2 \otimes D & \longleftarrow & C \otimes \bar{C} \otimes N \otimes \bar{D}^2 \otimes D & \longleftarrow & C \otimes \bar{C}^2 \otimes N \otimes \bar{D}^2 \otimes D & \longleftarrow & \dots \\ & \downarrow & & \downarrow & & \downarrow & \\ C \otimes N \otimes \bar{D} \otimes D & \longleftarrow & C \otimes \bar{C} \otimes N \otimes \bar{D} \otimes D & \longleftarrow & C \otimes \bar{C}^2 \otimes N \otimes \bar{D} \otimes D & \longleftarrow & \dots \\ & \downarrow & & \downarrow & & \downarrow & \\ C \otimes N \otimes D & \longleftarrow & C \otimes \bar{C} \otimes N \otimes D & \longleftarrow & C \otimes \bar{C}^2 \otimes N \otimes D & \longleftarrow & \dots \\ & \downarrow & & \downarrow & & \downarrow & \\ C \otimes N & \longleftarrow & C \otimes \bar{C} \otimes N & \longleftarrow & C \otimes \bar{C}^2 \otimes N & \longleftarrow & \dots \end{array}$$

The resulting total complex can be viewed as the cofiber of a map $FGN \rightarrow F_C G_C(N)$. We canonically truncate in the vertical and horizontal direction and as in the one-sided case we obtain an inverse system of bounded acyclic complexes of finite length. This shows that this cofiber is contra-acyclic as a C - D -bimodule. Thus, we conclude that the following result holds:

Proposition 5.1. *The adjunction unit $FGN \rightarrow N$ is a cofibrant resolution of the pseudo-compact dg C - D -bimodule N .*

One proves similarly:

Proposition 5.2. *The adjunction counit $GFM \rightarrow M$ is a cofibrant resolution of the dg ΩC - ΩD -bimodule M .*

Proof. The argument proving Proposition 5.1 proves that the unit map has a contra-acyclic cone and Proposition 4.9.2 shows this is a weak equivalence. \square

We can now formulate the following bimodule version of Theorem 4.5.

Theorem 5.3. *The functor G is left adjoint to F and they form a Quillen equivalence*

$$G: (C\text{-DGMod-}D)^{\mathrm{op}} \rightleftarrows (\Omega C\text{-DGMod-}\Omega D)_c^{\mathrm{II}} : F.$$

Proof. We have an adjoint pair of functors (G, F) between $C^{\mathrm{op}} \otimes D$ -modules and $\Omega C^{\mathrm{op}} \otimes \Omega D$ -modules; it has already been argued above that this is indeed an adjoint pair. Moreover, the functor F clearly converts cofibrations of $\Omega C^{\mathrm{op}} \otimes \Omega D$ -modules into fibrations of $C^{\mathrm{op}} \otimes D$ -modules (since the latter are simply surjective maps) while G takes cofibrations of $C^{\mathrm{op}} \otimes D$ -modules to fibrations of $(\Omega C^{\mathrm{op}} \otimes \Omega D)$ -modules (since the latter are similarly surjective maps). This shows that (G, F) is a Quillen adjunction. By Propositions 5.1 and 5.2 F and G induce an isomorphism at the level of homotopy categories. \square

Corollary 5.4. *For any pseudocompact dg algebras C and D , there is an equivalence of triangulated categories between $D_c^{\text{II}}(\Omega C^{\text{op}} \otimes \Omega D)$ and $D_c^{\text{II}}(\Omega(C^{\text{op}} \otimes D))$.*

Proof. By Koszul duality there is a Quillen equivalence

$$(C\text{-DGMod-}D)^{\text{op}} \rightleftarrows (\text{DGMod-}\Omega(C^{\text{op}} \otimes D))_c^{\text{II}},$$

so by Theorem 5.3 there is an equivalence of homotopy categories $\text{Ho}(\Omega C\text{-DGMod-}\Omega D)$ and $\text{Ho}(\text{DGMod-}\Omega(C^{\text{op}} \otimes D))$. \square

Remark 5.5. Note that this subsection would have simplified drastically if the functor Ω was quasi-strong monoidal. This is known for local pseudocompact algebras and those dual to pointed coalgebras [15], but not in general.

5.2. Koszul duality for the bar constructions. Let A, E be two dg algebras. We define two functors

$$\widehat{G}: (\widehat{B}A\text{-DGMod-}\widehat{B}E)^{\text{op}} \rightleftarrows A\text{-DGMod-}E : \widehat{F}.$$

Let $\xi_A \in \text{MC}(A^{\text{op}} \otimes \widehat{B}A^{\text{op}})$ and $\xi_E \in \text{MC}(E \otimes \widehat{B}E)$ be the canonical Maurer–Cartan elements corresponding to the counits $\Omega \widehat{B}A^{\text{op}} \rightarrow A^{\text{op}}$ and $\Omega \widehat{B}E \rightarrow E$ of the adjunction $\Omega \dashv \widehat{B}$. Define

$$\xi := \xi_A \otimes 1 + 1 \otimes \xi_E \in A^{\text{op}} \otimes \widehat{B}A^{\text{op}} \otimes E \otimes \widehat{B}E,$$

then $\xi \in \text{MC}(A^{\text{op}} \otimes E \otimes \widehat{B}A^{\text{op}} \otimes \widehat{B}E)$. The functor \widehat{F} sends a dg A - E -bimodule M to

$$\widehat{F}M := (M^* \otimes \widehat{B}A^{\text{op}} \otimes \widehat{B}E)^{[\xi]}$$

and the functor \widehat{G} sends a pseudocompact dg $\widehat{B}A$ - $\widehat{B}E$ -bimodule N to

$$\widehat{G}N := (N^* \otimes A^{\text{op}} \otimes E)^{[\xi]}.$$

Analogously we can define two functors

$$\check{G}: (\check{B}A\text{-DGMod-}\check{B}E)^{\text{op}} \rightleftarrows A\text{-DGMod-}E : \check{F}$$

exactly as above, except replacing every occurrence of \widehat{B} by \check{B} .

The following statement is the bimodule version of Theorem 4.8.

Theorem 5.6. *The functors \widehat{G} and \check{G} are left adjoint to \widehat{F} and \check{F} respectively, and form Quillen equivalences*

$$\widehat{G}: (\widehat{B}A\text{-DGMod-}\widehat{B}E)^{\text{op}} \rightleftarrows (A\text{-DGMod-}E)^{\text{I}} : \widehat{F}$$

and

$$\check{G}: (\check{B}A\text{-DGMod-}\check{B}E)^{\text{op}} \rightleftarrows (A\text{-DGMod-}E)_c^{\text{II}} : \check{F}.$$

Furthermore for any dg algebras A and E , there are equivalences of triangulated categories

$$D_c^{\text{II}}(\widehat{B}A^{\text{op}} \otimes \widehat{B}E) \cong D_c^{\text{II}}(\widehat{B}(A^{\text{op}} \otimes E))$$

and

$$D_c^{\text{II}}(\check{B}A^{\text{op}} \otimes \check{B}E) \cong D_c^{\text{II}}(\check{B}(A^{\text{op}} \otimes E));$$

hence $(\widehat{G}, \widehat{F})$ and (\check{G}, \check{F}) are Quillen equivalences.

Proof. We prove the theorem for the pair of functors (\check{G}, \check{F}) ; the proof for $(\widehat{G}, \widehat{F})$ is the same. By Koszul duality there are Quillen equivalences

$$(\text{DGMod-}(\check{B}A^{\text{op}} \otimes \check{B}E))^{\text{op}} \rightleftarrows \text{DGMod-}\Omega(\check{B}A^{\text{op}} \otimes \check{B}E)$$

and

$$(\text{DGMod-}\check{B}(A^{\text{op}} \otimes E))^{\text{op}} \rightleftarrows \text{DGMod-}\Omega\check{B}(A^{\text{op}} \otimes E)$$

so it is equivalent to prove that $D_c^{\text{II}}(\Omega(\check{\mathcal{B}}A^{\text{op}} \otimes \check{\mathcal{B}}E)) \cong D_c^{\text{II}}(\Omega\check{\mathcal{B}}(A^{\text{op}} \otimes E))$. But by Corollary 5.4,

$$D_c^{\text{II}}(\Omega(\check{\mathcal{B}}A^{\text{op}} \otimes \check{\mathcal{B}}E)) \cong D_c^{\text{II}}(\Omega\check{\mathcal{B}}A^{\text{op}} \otimes \Omega\check{\mathcal{B}}E) \cong D_c^{\text{II}}(A^{\text{op}} \otimes E) \cong D_c^{\text{II}}(\Omega\check{\mathcal{B}}(A^{\text{op}} \otimes E)).$$

Here for the middle equivalence by Proposition 4.15 it suffices to check $\text{Perf}^{\text{II}}(\Omega\check{\mathcal{B}}A^{\text{op}} \otimes \Omega\check{\mathcal{B}}E) \simeq \text{Perf}^{\text{II}}(A^{\text{op}} \otimes E)$. This holds true as both sides are equivalent to $\text{Perf}^{\text{II}}(\text{Perf}^{\text{II}}(A^{\text{op}}) \otimes \text{Perf}^{\text{II}}(E))$ by Lemma 4.16 and since $\Omega\check{\mathcal{B}}A \rightarrow A$ is a Morita equivalence of the second kind (4.14).

Finally this implies that (\check{G}, \check{F}) is a Quillen equivalence as $D^{\text{co}}(\check{\mathcal{B}}(A^{\text{op}} \otimes E)) \cong D_c^{\text{II}}(A^{\text{op}} \otimes E)$. \square

As a corollary, we get the following result, the first part of which appears in [18, Proposition 2.4]. Here $\text{HH}(C)$ for a pseudocompact algebra C (sometimes called *coHochschild cohomology* of the dual coalgebra C^*) denotes the Hochschild cohomology $\text{RHom}_{C \otimes C^{\text{op}}}(C, C)$.

Corollary 5.7. *For any dg algebra A , there are isomorphisms of algebras*

$$\text{HH}(A) \cong \text{HH}(\widehat{\mathcal{B}}A)$$

and

$$\text{HH}_c^{\text{II}}(A) \cong \text{HH}(\check{\mathcal{B}}A).$$

Proof. We prove the second statement. By Theorem 5.6 $D^{\text{co}}(\check{\mathcal{B}}A \otimes \check{\mathcal{B}}A^{\text{op}}) \cong D_c^{\text{II}}(A \otimes A^{\text{op}})$. It suffices to check that $\check{G}(\check{\mathcal{B}}A) = (A \otimes \check{\mathcal{B}}A^* \otimes A)^{[\xi]}$ $\simeq A$. But this follows as $\check{G}_{A \otimes A^{\text{op}}}(\check{\mathcal{B}}A)$ is the same as $\check{G}_A \check{F}_A(A) = (A \otimes (\check{\mathcal{B}}A \otimes A^*)^*)^{[\xi_A \otimes 1 + 1 \otimes \xi_A]}$. The two expressions clearly have the same underlying graded object and the differential is induced in each case by two copies of the MC element $\xi_A \in \check{\mathcal{B}}A \otimes A$, acting by left multiplication on $\check{\mathcal{B}}A$ and A and by right multiplication on $\check{\mathcal{B}}A$ and the other copy of A respectively. \square

Corollary 5.8. $\text{HH}_c^{\text{II}}(A)$ is computed by the complex $(\check{\mathcal{B}}A \otimes A)^\xi$ where the superscript induces two-sided twisting by ξ_A .

Proof. The computation is the same as how one might compute Hochschild cohomology in terms of the usual tensor algebra.

We have $\text{HH}_c^{\text{II}}(A) \cong \text{RHom}_{\check{\mathcal{B}}A \otimes \check{\mathcal{B}}A^{\text{op}}}(\check{\mathcal{B}}A, \check{\mathcal{B}}A)$ and $\check{\mathcal{B}}A$ is freely resolved by $\check{\mathcal{B}}A \otimes \overline{A}^*[1] \otimes \check{\mathcal{B}}A \rightarrow \check{\mathcal{B}}A \otimes \check{\mathcal{B}}A$ via the multiplication map. Thus we obtain

$$\text{HH}_c^{\text{II}}(A) \simeq (\text{Hom}_k(A^*, \check{\mathcal{B}}A), d) \simeq (A \otimes \check{\mathcal{B}}A, d)$$

and the induced differential d is exactly the two-sided twisting by ξ_A , which is the usual Hochschild differential \square

Remark 5.9. There is another definition of Hochschild cohomology of the second kind, cf. [21] which is *not* equivalent to our notion. To define it, let $\mathcal{B}'(A) := \bigoplus_{n=0}^{\infty} (\overline{A}^{\otimes n})^*[-1]$ be the ‘semi-complete’ bar-construction of A ; it is a dg algebra that is neither discrete in general (because $(\overline{A}^{\otimes n})^*$ is a pseudocompact vector space) nor pseudocompact (because an infinite direct sum of pseudocompact vector spaces is not pseudocompact). Nevertheless, the complex $(\mathcal{B}'(A) \hat{\otimes} A)^\xi$ makes sense and can be taken as a definition of *Hochschild cohomology of the second kind* (in the sense of Polishchuk-Positselski), $\text{HH}_{\text{PP}}^{\text{II}}(A)$. This is also sometimes called *compactly supported Hochschild cohomology*, $\text{HH}_c^*(A)$, and the dual construction is called *Borel-Moore Hochschild homology* $\text{HH}_*^{\text{BM}}(A)$. (Note that the subscript c in $\text{HH}_c^*(A)$ stands for ‘compactly supported’ in contrast with its usage in the present paper which refers to ‘compactly generated’.)

The pseudocompact bar-construction $\check{\mathcal{B}}(A)$ is the pseudocompact completion of $\mathcal{B}'(A)$ and the ordinary bar-construction $\widehat{\mathcal{B}}(A)$ is a further completion at the maximal ideal. It

follows that there are maps of complexes $\mathrm{HH}_{\mathrm{pp}}^{\mathrm{II}}(A) \rightarrow \mathrm{HH}_c^{\mathrm{II}}(A) \rightarrow \mathrm{HH}(A)$. In other words, $\mathrm{HH}_c^{\mathrm{II}}(A)$ is a kind of a half-way house between $\mathrm{HH}_{\mathrm{pp}}^{\mathrm{II}}(A)$ and $\mathrm{HH}(A)$.

6. CURVED AND NON-AUGMENTED CASES

We note that the results in the previous two sections can be generalized to the curved and non-augmented settings, i.e. to curved, non-augmented algebras which are Koszul dual to non-local, curved pseudocompact algebras.

We restrict ourselves to the case of algebras and do not consider curved categories.

We now gather the results, and indicate where there is a difference in the proofs.

Definition 6.1. A *twisted module* over a curved algebra (A, d, w) is just a curved A -module whose underlying graded module has the form $V \otimes A$ for a graded k -module V . Explicitly a twisted module is of the form $(V \otimes A, 1 \otimes d + q)$ where $q \in (\mathrm{End} V \otimes A)^1$ satisfies $dq + q^2 = 1 \otimes w \in \mathrm{End} V \otimes A$.

With this definition we can define finitely-generated twisted modules, $\mathrm{Cof}_c^{\mathrm{II}}(A)$ and $\mathrm{Perf}^{\mathrm{II}}(A)$ for a curved algebra. Note that modules over A are somewhat more subtle if A is curved since A itself is no longer a (left or right) twisted module over itself. It is, however, always a bimodule over itself.

Koszul duality extends to the curved setting following [22, 14, 7]. As a first step one may extend the bar and cobar construction to the non-augmented case by choosing a section of the unit to define a decomposition $A \cong \check{A} \oplus k$ of an algebra as a k -module. This will introduce curvature. The Koszul dual of a non-augmented dg algebra is thus a curved pseudocompact algebra. Similarly the Koszul dual of a non-local pseudocompact algebra is a curved algebra.

Next, while the bar construction of a curved algebra is not a good notion in general, we always have the extended bar construction $\check{\check{B}}A$ for a curved algebra A , right adjoint to the cobar construction.

With this we still have the model structure on $(\mathrm{DGMod}\text{-}A)_c^{\mathrm{II}}$ by [14, Theorem 4.6] and the Quillen adjunction to $\mathrm{DGMod}\text{-}\check{\check{B}}A^{\mathrm{op}}$ by [14, Theorem 4.7].

There is also a Morita equivalence of the second kind $\Omega\check{\check{B}}A \simeq A$. This follows directly by [7, Theorem 6.4 and Corollary 6.6].

Theorem 4.12 still holds in the curved setting: sompact generation by twisted modules is again inherited via Koszul duality from the fact that comodules are compactly generated by finite-dimensional comodules. This also holds in the curved, nonconilpotent setting, see [22, Section 5.5].

Since a curved algebra is always a bimodule over itself, the following definition makes sense:

Definition 6.2. Let A be a curved algebra and M be an A -bimodule (i.e. a module over the curved algebra $A \otimes A^{\mathrm{op}}$). The *Hochschild cohomology of the second kind* of A with coefficients in M is

$$\mathrm{HH}_c^{\mathrm{II}}(A, M) = \mathrm{RHom}_{A \otimes A^{\mathrm{op}}}^{\mathrm{II}}(A, M).$$

We write $\mathrm{HH}_c^{\mathrm{II}}(A)$ for $\mathrm{HH}_c^{\mathrm{II}}(A, A)$.

The next goal is to transfer the remaining content of Section 4. Some results, namely Proposition 4.15 and Lemma 4.17 hold without any adjustments.

However, other results strongly rely on the fully faithful Yoneda embedding $A \rightarrow \mathrm{Perf}^{\mathrm{II}}(A)$. If A is curved, this is no longer available since A is not a (right) module over itself and so it has no Yoneda embedding. To get around this technical point, we need the following result, which is of some independent interest.

Lemma 6.3. *Given a curved algebra A , there is a dg algebra A' such that $\text{Perf}^{\text{II}}(A)$ and $\text{Perf}^{\text{II}}(A')$ are quasi-equivalent.*

Proof. Let $w \in A^2$ be the curvature element. Let $V := k \oplus k[1]$ and consider the A -module $M := A \oplus A[1] \cong A \otimes V$ with the differential d_M given on $1 \otimes V \subset M$ by the 2×2 matrix $x_w = -\begin{pmatrix} 0 & 1 \\ w & 0 \end{pmatrix}$. We have $d_M^2 = \begin{pmatrix} w & 0 \\ 0 & w \end{pmatrix}$ and so M is a curved (perfect) A -module. It follows that x_w is an MC element in the curved algebra $A \otimes \text{End}(V)$. Set $A' := \text{End}_A(M) \cong A \otimes \text{End}(V)^{x_w}$, the twist of $A \otimes \text{End}(V)$ by x_w . The curved algebra $A \otimes \text{End}(V)$ is clearly Morita equivalent of the second kind to A with the equivalence $\text{Perf}^{\text{II}}(A) \rightarrow \text{Perf}^{\text{II}}(A \otimes \text{End}(V))$ given by the usual prescription $? \mapsto \text{Hom}_A(?, M)$. Since A' is isomorphic to $A \otimes \text{End}(V)$ as a curved algebra, it follows that $\text{Perf}^{\text{II}}(A)$ and $\text{Perf}^{\text{II}}(A')$ are quasi-equivalent as desired (in fact, the constructed quasi-equivalence is even an equivalence as ordinary categories). \square

Remark 6.4. It is easy to see that the dg algebra A' constructed above, is acyclic (and so, the twisted module M is homotopically trivial). Indeed, the identity element $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ is the coboundary of the element $\begin{pmatrix} 0 & 0 \\ -1 & 0 \end{pmatrix}$ in A' . Note that the ordinary derived category of A' is, of course, trivial.

We now have the curved analogue of Lemma 4.16.

Lemma 6.5. *For curved algebras A, B there is a quasi-equivalence*

$$\text{Perf}^{\text{II}}(A \otimes B) \cong \text{Perf}^{\text{II}}(\text{Perf}^{\text{II}}(A) \otimes \text{Perf}^{\text{II}}(B))$$

Proof. We have a quasi-equivalence $\text{Perf}^{\text{II}}(A \otimes B) \simeq \text{Perf}(A' \otimes B')$ (where A' and B' are dg algebras II-Morita equivalent to A and B as constructed in Lemma 6.3) since $A' \otimes B'$ is isomorphic to the tensor product of $A \otimes B$ and a 4×4 matrix algebra. Similarly $\text{Perf}^{\text{II}}(\text{Perf}^{\text{II}}(A) \otimes \text{Perf}^{\text{II}}(B))$ is Morita equivalent of the second kind to $\text{Perf}^{\text{II}}(A \otimes B)$ which reduces the question to the uncurved case that has already been proved in Lemma 4.16. \square

Using Lemma 6.5 in place of Lemma 4.16, the curved analogue of Theorem 4.20 can now be shown with the same proof:

Theorem 6.6. *A Morita equivalence of the second kind $F: A \rightarrow B$ between curved algebras induces a quasi-isomorphism $\text{HH}_c^{\text{II}}(A) \simeq \text{HH}_c^{\text{II}}(B)$ of dg algebras.*

With the same adjustment of using Lemma 6.5 we also have the curved analogues of Theorem 5.6 and Corollary 5.7 by the same proof. We will just define the necessary functors and state the theorem:

As there is an adjunction $\Omega \dashv \check{B}$ also for curved pseudocompact algebras [7] we obtain a canonical MC element $\xi_A \in \text{MC}(A^{\text{op}} \otimes \check{B}A^{\text{op}})$ corresponding to the counit $\Omega \check{B}A^{\text{op}} \rightarrow A^{\text{op}}$. We may define

$$\xi := \xi_A \otimes 1 + 1 \otimes \xi_E \in A^{\text{op}} \otimes \check{B}A^{\text{op}} \otimes E \otimes \check{B}E,$$

then $\xi \in \text{MC}(A^{\text{op}} \otimes E \otimes \check{B}A^{\text{op}} \otimes \check{B}E)$. The functor \check{F} sends a dg A - E -bimodule M to

$$\check{F}M := (M^* \otimes \check{B}A^{\text{op}} \otimes \check{B}E)^{[\xi]}$$

and the functor \check{G} sends a pseudocompact dg $\check{B}A$ - $\check{B}E$ -bimodule N to

$$\check{G}N := (N^* \otimes A^{\text{op}} \otimes E)^{[\xi]}.$$

Theorem 6.7. *Let A, E be curved algebras. The functor \check{G} is left adjoint to \check{F} respectively, and forms a Quillen equivalence*

$$\check{G}: (\check{B}A\text{-DGMod-}\check{B}E)^{\text{op}} \rightleftarrows (A\text{-DGMod-}E)_c^{\text{II}} : \check{F}.$$

Furthermore for any dg algebras A and E , there is an equivalence of triangulated categories

$$D_c^{\text{II}}(\check{B}A^{\text{op}} \otimes \check{B}E) \cong D_c^{\text{II}}(\check{B}(A^{\text{op}} \otimes E)).$$

Corollary 6.8. *For any curved algebra A , there is a quasi-isomorphism of algebras*

$$\text{HH}_c^{\text{II}}(A) \simeq \text{HH}(\check{B}A)$$

and $\text{HH}_c^{\text{II}}(A)$ is computed by the complex $(\check{B}A \otimes A)^\xi$.

7. EXAMPLES OF HOCHSCHILD COHOMOLOGY OF THE SECOND KIND

7.1. Preliminaries. In this final section we compare Hochschild cohomology of the first and second kind in some situations. Let A and E be two curved algebras and M, N be two $A \otimes E$ -modules. Recall that $\text{RHom}_{A \otimes E}^{\text{II}}(M, N)$ is the derived complex of homomorphisms from M to N in the compactly generated model category of the second kind of $A \otimes E$ -modules. Thus, $\text{RHom}_{A \otimes E}^{\text{II}}(M, N)$ can be represented as $\text{Hom}_{A \otimes E}(\tilde{M}, N)$ where \tilde{M} is a cofibrant replacement of M as an $A \otimes E$ -module.

Recall that $\text{Perf}^{\text{II}}(?)$ stands for the dg category of perfect cofibrant $?$ -modules of the second kind (which can be represented as retracts of finitely generated twisted modules).

The modules M and N can be viewed as $\text{Perf}^{\text{II}}(A) \otimes \text{Perf}^{\text{II}}(E)$ -modules: the corresponding functor to dg vector spaces $\text{Perf}^{\text{II}}(A) \rightarrow \text{DGVect}$ is given by $L \mapsto \text{Hom}_{A \otimes E}(M, L)$ where L is a given $A \otimes E$ -module. We can, therefore, form $\text{RHom}_{\text{Perf}^{\text{II}}(A) \otimes \text{Perf}^{\text{II}}(E)}(M, N)$, the derived functor of homomorphisms in the model category (of the first kind) of $\text{Perf}^{\text{II}}(A) \otimes \text{Perf}^{\text{II}}(E)$ -modules.

We would like to compare $\text{RHom}_{A \otimes E}^{\text{II}}(M, N)$ and $\text{RHom}_{\text{Perf}^{\text{II}}(A) \otimes \text{Perf}^{\text{II}}(E)}(M, N)$. Note that it is well-known that $\text{RHom}_{A \otimes E}(M, N)$ and $\text{RHom}_{\text{Perf}(A) \otimes \text{Perf}(E)}(M, N)$ are naturally quasi-isomorphic, the reason being that the Yoneda embedding $A \otimes E \rightarrow \text{Perf}(A) \otimes \text{Perf}(E)$ is a Morita equivalence of categories. This argument, however, breaks down in our situation (e.g. because we wish to compare RHom of the first kind with RHom of the second kind). In fact, there is not even a natural map between $\text{RHom}_{A \otimes E}^{\text{II}}(M, N)$ and $\text{RHom}_{\text{Perf}^{\text{II}}(A) \otimes \text{Perf}^{\text{II}}(E)}(M, N)$.

Consider the natural functor

$$i: \text{Perf}^{\text{II}}(A) \otimes \text{Perf}^{\text{II}}(E) \rightarrow \text{Perf}^{\text{II}}(A \otimes E)$$

sending a pair M, N of perfect modules of the second kind to $M \otimes N$.

Then we have the following result, similar to [21, Theorem 3.5 C].

Lemma 7.1. *Suppose that i above is a Morita equivalence (of the first kind). Then for $A \otimes E$ -modules M and N we have that $\text{RHom}_{A \otimes E}^{\text{II}}(M, N)$ and $\text{RHom}_{\text{Perf}^{\text{II}}(A) \otimes \text{Perf}^{\text{II}}(E)}(M, N)$ are naturally quasi-isomorphic.*

Proof. We first note that there is an equivalence $D_c^{\text{II}}(A) \cong D(\text{Perf}^{\text{II}}(A))$ induced by the natural map $D_c^{\text{II}}(\text{Perf}^{\text{II}}(A)) \rightarrow D(\text{Perf}^{\text{II}}(A))$. This follows by comparing compact objects, which are $\text{Perf}^{\text{II}}(A)$ on both sides (using $\text{Perf}(\text{Perf}^{\text{II}}(A)) \cong \text{Perf}^{\text{II}}(A)$).

With this we can compute

$$\begin{aligned} \mathrm{RHom}_{A \otimes E}^{\mathrm{II}}(M, N) &\simeq \mathrm{RHom}_{\mathrm{D}_c^{\mathrm{II}}(A \otimes E)}(M, N) \\ &\simeq \mathrm{RHom}_D(\mathrm{Perf}^{\mathrm{II}}(A \otimes E))(M, N) \\ &\simeq \mathrm{RHom}_D(\mathrm{Perf}^{\mathrm{II}}(A) \otimes \mathrm{Perf}^{\mathrm{II}}(E))(M, N) \\ &\simeq \mathrm{RHom}_{\mathrm{Perf}^{\mathrm{II}}(A) \otimes \mathrm{Perf}^{\mathrm{II}}(E)}(M, N) \end{aligned}$$

where we used the assumption in the penultimate step. \square

Remark 7.2. There is no reason to believe that i is always a quasi-equivalence. In fact, the dg category $\mathrm{Perf}^{\mathrm{II}}(A \otimes E)$ should be viewed as a kind of a *completed* tensor product of the categories $\mathrm{Perf}^{\mathrm{II}}(A)$ and $\mathrm{Perf}^{\mathrm{II}}(E)$. There are, however, important situations when this completion is extraneous.

Let us take $E := A^{\mathrm{op}}$ and $M := A$. The above lemma specializes to the following statement.

Corollary 7.3. *Let A be a dg algebra such that $\mathrm{Perf}^{\mathrm{II}}(A) \otimes \mathrm{Perf}^{\mathrm{II}}(A^{\mathrm{op}})$ is Morita equivalent to $\mathrm{Perf}^{\mathrm{II}}(A \otimes A^{\mathrm{op}})$ and M be an A -bimodule. Then $\mathrm{HH}_c^{\mathrm{II}}(A, M)$ is naturally quasi-isomorphic to $\mathrm{HH}(\mathrm{Perf}^{\mathrm{II}}(A), M)$.*

In other words, under the assumptions of Lemma 7.1, Hochschild cohomology of the second kind of dg algebras reduce to Hochschild cohomology of the first kind of a suitable dg category. A version of this question was considered in [21] where some partial results were obtained (for a different notion of Hochschild cohomology of the second kind, see Remark 5.9).

7.2. Complex algebraic manifolds. Let X be a compact complex projective manifold and $A := (\mathcal{A}^{0,*}(X), \bar{\partial})$ its Dolbeault-algebra. It is well-known [3, 10] that the bounded derived category $\mathrm{DCoh}^b(X)$ of coherent sheaves on X is equivalent to the derived category of sheaves on X with bounded coherent cohomology and, since X is smooth, the latter coincides with the derived category of perfect complexes of sheaves on X . We will consider its dg model $\mathrm{Perf}(X)$ formed by taking Dolbeault resolutions of coherent sheaves. According to [5, Corollary 3.1.8], the latter is equivalent to the dg category of dg modules over some dg algebra (the endomorphism algebra of a generator of $\mathrm{Perf}(X)$).

The following result holds.

Proposition 7.4. *The categories $\mathrm{Perf}(X \times X)$ and $\mathrm{Perf}(X) \otimes \mathrm{Perf}(X)$ are Morita equivalent.*

Proof. The statement is well-known and follows from a very general result [2, Theorem 1.2(1)] valid for perfect derived stacks, not merely for complex projective manifolds. The argument goes back to [5]. We sketch a proof for the reader's convenience.

Let \mathcal{F} be a complex of sheaves representing a generator of $\mathrm{Perf}(X)$ and $B = R\mathrm{End}(\mathcal{F})$ be its endomorphism dg algebra. Then the external tensor product $\mathcal{F} \boxtimes \mathcal{F}$ is a generator of $\mathrm{Perf}(X \times X)$ and $B \otimes B \simeq R\mathrm{End}(\mathcal{F} \boxtimes \mathcal{F})$. So, $\mathrm{Perf}(X)$ is quasi-equivalent to $\mathrm{Perf}(B)$ and $\mathrm{Perf}(X \times X)$ is quasi-equivalent to $B \otimes B$. Thus we have reduced the question to that of a perfect derived category of a dg algebra. Since $\mathrm{Perf}(B \otimes B)$ is Morita equivalent to $\mathrm{Perf}(B) \otimes \mathrm{Perf}(B)$ (for any dg algebra B), the desired claim follows. \square

Theorem 7.5. *The dg algebra $A = \mathcal{A}^{0,*}$ satisfies the conditions of 7.3. Thus we have a natural quasi-isomorphism*

$$\mathrm{HH}_c^{\mathrm{II}}(A) \simeq \mathrm{HH}(\mathrm{Perf}(X)).$$

Proof. This follows at once from Proposition 7.4 together with the observation that A is graded commutative and so $A^{\mathrm{op}} \cong A$ and $\mathrm{Perf}(A) \simeq (\mathrm{Perf}^{\mathrm{II}}(A))^{\mathrm{op}}$. \square

7.3. Topological spaces. In the topological setting we consider higher local systems on a topological space X . To be precise we denote by $\mathrm{LC}(X)$ the dg category of fibrant cofibrant cohomologically locally constant (clc) sheaves of complexes over k whose cohomology sheaves have finite-dimensional fibers. We recall the following:

Theorem 7.6. *Let X be a connected locally contractible topological space and $C^*(X)$ its normalized singular cochain algebra with coefficients in k . Then $\mathrm{Perf}^{\mathrm{II}}(C^*(X)) \cong \mathrm{LC}(X)$.*

Let $k = \mathbb{R}$, X a connected manifold and $\mathcal{A}^(X)$ its de Rham algebra. Then $\mathrm{Perf}^{\mathrm{II}}(\mathcal{A}^*(X)) \cong \mathrm{LC}(X)$.*

Proof. For the first result it follows from [10, Theorem 8.4] that $\mathrm{LC}(X)$ is given by finitely generated twisted modules over $C^*(X)$ since finitely generated twisted modules correspond to clc sheaves whose fibers are bounded and finite dimensional in each degree. As $\mathrm{LC}(X)$ is idempotent complete it follows that it is quasi-equivalent to $\mathrm{Perf}^{\mathrm{II}}(C^*(X))$.

The second result is [10, Theorem 8.1]. \square

Remark 7.7. Note that as we restrict to finitely generated twisted complexes we did not need to consider $C^*(X)$ as a pseudocompact algebra in the theorem. If we consider $C^*(X)$ as a pseudocompact algebra we have $\mathrm{D}^{\mathrm{co}}(C^*(X)) \cong \mathrm{LC}^{\infty}(X)$, where the right hand side is the category of potentially infinite-dimensional local systems [10, Theorem 8.4].

But $\mathrm{D}^{\mathrm{co}}(C^*(X))$ is different from $\mathrm{D}_c^{\mathrm{II}}(C^*(X))$. E.g. the local system on S^1 associated to the regular representation of $\pi_1(S^1)$ exists in $\mathrm{D}^{\mathrm{co}}(C^*(X))$ but not in $\mathrm{D}_c^{\mathrm{II}}(C^*(X))$.

If $\pi_1(X)$ is finite then $\mathrm{LC}^{\infty}(X)$ is generated by $\mathrm{LC}(X)$ and their Hochschild cohomologies agree. It is known that $\mathrm{LC}^{\infty}(X) \simeq \mathrm{DGMod}\text{-}C_*\Omega X$ (e.g. [16]). Thus Hochschild cohomology of $\mathrm{LC}^{\infty}(X)$ is given by $\mathrm{HH}(C_*\Omega(X))$ which is known to be equal to the string topology of X [8, 12]. See also [24].

Proposition 7.8. *Let X be a CW complex with finitely many cells in each degree. Then $\mathrm{LC}(X \times X)$ and $\mathrm{LC}(X) \otimes \mathrm{LC}(X)$ are Morita equivalent.*

Proof. We let $\{G_i\}$ be a collection of generators for $\mathrm{LC}(X)$, e.g. the collection of all (classical) local systems on X . We first show that $G_i \boxtimes G_j$ generate $\mathrm{LC}(X \times X)$. To do this we follow again the argument by Bondal-Van den Bergh.

Let N be right orthogonal to $\{G_i \boxtimes G_j\}$, i.e. $\mathrm{RHom}_{X \times X}(G_i \boxtimes G_j, N) \simeq 0$ for all i, j . We need to show that N is trivial.

By adjunction we have

$$\begin{aligned} 0 &\simeq \mathrm{RHom}_{X \times X}(G_i \boxtimes G_j, N) \\ &\simeq \mathrm{RHom}_{X \times X}(\pi_1^* G_i, \mathrm{Hom}(\pi_2^* G_j, N)) \\ &\simeq \mathrm{RHom}_X(G_i, (\pi_1)_* \mathrm{Hom}(\pi_2^* G_j, N)). \end{aligned}$$

It follows that $(\pi_1)_* \mathrm{Hom}(\pi_2^* G_j, N) \simeq 0$ as the G_i generate $\mathrm{LC}(X)$ and $(\pi_1)_* \mathrm{Hom}(\pi_2^* G_j, N) \in \mathrm{LC}(X)$ by our finiteness assumption.

Thus the fibers of $(\pi_1)_* \mathrm{Hom}(\pi_2^* G_j, N)$ must be trivial and we have

$$\mathrm{Hom}_{\{x_1\} \times X}(\pi_2^* G_j|_{\{x_1\} \times X}, N|_{\{x_1\} \times X}) \simeq 0$$

which gives $\mathrm{Hom}_X(G_j, (\pi_2)_* N|_{\{x_1\} \times X}) \simeq 0$ and as the G_j generate we have that $N|_{\{x_1\} \times X}$ is trivial and it follows that N itself is trivial.

This shows that the comparison functor $\mathrm{LC}(X) \otimes \mathrm{LC}(X) \rightarrow \mathrm{LC}(X \times X)$ given by \boxtimes is essentially surjective.

It remains to be shown the functor is quasi-fully faithful. It suffices to check on generators, i.e. on local systems. We thus have to compare $\mathrm{RHom}(L, L') \otimes \mathrm{RHom}(M, M')$ with

$\mathrm{RHom}(L \boxtimes L, M \boxtimes M')$ for local systems L, L', M, M' on X . But this is equivalent to showing

$$C^*(X, L' \otimes L^*) \otimes C^*(X, M' \otimes M^*) \simeq C^*(X \times X, (L' \otimes L^*) \boxtimes (M' \otimes M^*)).$$

Thus the result follows from the Künneth theorem with local coefficients [13, Theorem 1.7]. Here we use the finiteness assumption. \square

Corollary 7.9. *Let X be a connected locally contractible topological space that has the homotopy type of a CW complex with finitely many cells in each degree. Then $\mathrm{HH}_c^{\mathrm{II}}(C^*(X)) \simeq \mathrm{HH}(\mathrm{LC}(X))$. If X is moreover a manifold $\mathrm{HH}_c^{\mathrm{II}}(\mathcal{A}^*(X)) \simeq \mathrm{HH}(\mathrm{LC}(X))$.*

7.4. Matrix Factorizations. We now turn to a curved example. Let R be a commutative algebra over a field k of characteristic 0, and assume R is regular, and let $w \in R$. Interpreting R as a $\mathbb{Z}/2$ graded algebra concentrated in even degrees we may consider w as curvature and define a curved ring $R_w = (R, 0, w)$.

Then the dg category of matrix factorizations $\mathrm{MF}(R, w)$ may be defined as the idempotent completion of the category of curved modules over R_w such that the underlying graded module is finitely generated and projective in each degree. Its homotopy category is known to be equivalent to the derived category of singularities $\mathrm{D}_{\mathrm{coh}}^b(Z)/\mathrm{Perf}(Z)$ where $Z = w^{-1}(0) \subset \mathrm{Spec}(R)$ is the zero locus with singular locus $\mathrm{crit}(w)$ [20].

For a $\mathbb{Z}/2$ -graded curved algebra we may define its $\mathbb{Z}/2$ -graded categories of twisted modules and perfect complexes of the second kind $\mathrm{Perf}_{\mathbb{Z}/2}^{\mathrm{II}}$ by just changing the grading in our definitions.

With these definitions we obtain the following lemma:

Lemma 7.10. *There is a quasi-equivalence $\mathrm{MF}(R, w) \cong \mathrm{Perf}_{\mathbb{Z}/2}^{\mathrm{II}}(R_w)$ of $\mathbb{Z}/2$ -graded dg categories.*

Proof. The definitions of $\mathrm{MF}(R, w)$ and $\mathrm{Perf}_{\mathbb{Z}/2}^{\mathrm{II}}(R_w)$ agree except that matrix factorizations are built out of projective modules for $R^\#$, the underlying graded of R , rather than free modules. But any curved R_w module whose underlying graded module P is finitely generated projective is a direct summand of a finitely generated twisted module, i.e. a curved module whose underlying graded is finitely generated free over $R^\#$.

To show this, pick a finitely generated projective $R^\#$ -module L such that $P \oplus L$ is free and consider $G(L)$ the free curved R -module on L which has elements formal sums $\ell + d\ell$ and differential $d(\ell + d\ell) = h\ell + d\ell$, cf. [22, Section 3.6]. Then the underlying graded of $G(L)$ is $L \oplus L[-1]$ and $P \oplus P[-1] \oplus G(L)$ is the desired R_w -module, proving the assertion. \square

We define Hochschild cohomology of the second kind of R_w as in the \mathbb{Z} -graded case, noting that $\mathrm{HH}_c^{\mathrm{II}}(R_w) = \mathrm{RHom}_{R_w \otimes R_w^{\mathrm{op}}}^{\mathrm{II}}(R_w, R_w)$ is now a $\mathbb{Z}/2$ -graded complex.

Proposition 7.11. *In the situation as above let furthermore w satisfy $\mathrm{crit}(w) \subset w^{-1}(0)$. Then $\mathrm{HH}_c^{\mathrm{II}}(R_w) \simeq \mathrm{HH}(\mathrm{MF}(R, w))$.*

Proof. The Thom-Sebastiani theorem [23, Theorem 4.1.3] says $\mathrm{MF}(R, w) \otimes \mathrm{MF}(R, w) \cong \mathrm{MF}(R \otimes R, w \otimes 1 + 1 \otimes w)$. Rewriting in terms of curved algebras this is exactly saying $\mathrm{MF}(R_w) \otimes \mathrm{MF}(R_w) = \mathrm{MF}(R_w \otimes R_w)$.

Using Lemma 7.10 we are in the setting of Corollary 7.3 and immediately obtain

$$\mathrm{HH}_c^{\mathrm{II}}(R_w) \simeq \mathrm{HH}(\mathrm{MF}(R, w)). \quad \square$$

Remark 7.12. Note that $\mathrm{HH}(\mathrm{MF}(R, w))$ has also been computed for compactly supported Hochschild cohomology (cf. Remark 5.9). For isolated singularities of $w^{-1}(0)$ this is [9, 29] and in general [21, Section 4.8]. They show $\mathrm{HH}(\mathrm{MF}(R, w)) = \mathrm{HH}_c(R_w)$. This is further used for example in [11].

It is notable that in this important case our definitions agree with the older definition that is different in many cases.

REFERENCES

- [1] M. ANEL AND A. JOYAL, *Sweedler theory for (co)algebras and the bar-cobar constructions*, arXiv:1309.695, (2013).
- [2] D. BEN-ZVI, J. FRANCIS, AND D. NADLER, *Integral transforms and Drinfeld centers in derived algebraic geometry*, *Journal of the American Mathematical Society*, 23 (2010), pp. 909–966.
- [3] J. BLOCK, *Duality and equivalence of module categories in noncommutative geometry*, in A celebration of the mathematical legacy of Raoul Bott, vol. 50 of CRM Proc. Lecture Notes, Amer. Math. Soc., Providence, RI, 2010, pp. 311–339.
- [4] A. BONDAL, M. LARSEN, AND V. LUNTS, *Grothendieck ring of pretriangulated categories*, *International Mathematics Research Notices*, (2004), pp. 1461–1495.
- [5] A. BONDAL AND M. VAN DEN BERGH, *Generators and representability of functors in commutative and noncommutative geometry*, *Moscow Mathematical Journal*, 3 (2003), pp. 1–36.
- [6] A. I. BONDAL AND M. M. KAPRANOV, *Enhanced triangulated categories*, *Mathematics of the USSR-Sbornik*, 70 (1991), pp. 93–107.
- [7] M. BOOTH AND A. LAZAREV, *Global Koszul duality*, arXiv:2304.08409, (2023).
- [8] D. BURGHELEA AND Z. FIEDOROWICZ, *Cyclic homology and algebraic K-theory of spaces. II*, *Topology*, 25 (1986), pp. 303–317.
- [9] A. CALDARARU AND J. TU, *Curved A-infinity algebras and Landau-Ginzburg models*, *New York J. Math*, 19 (2013), pp. 305–342.
- [10] J. CHUANG, J. HOLSTEIN, AND A. LAZAREV, *Maurer-Cartan moduli and theorems of Riemann-Hilbert type*, *Applied Categorical Structures*, (2021).
- [11] A. I. EFIMOV, *Cyclic homology of categories of matrix factorizations*, *Int. Math. Res. Not. IMRN*, (2018), pp. 3834–3869.
- [12] T. G. GOODWILLIE, *Cyclic homology, derivations, and the free loop space*, *Topology*, 24 (1985), pp. 187–215.
- [13] R. GREENBLATT, *Homology with local coefficients and characteristic classes*, *Homology, Homotopy and Applications*, 8 (2006), pp. 91 – 103.
- [14] A. GUAN AND A. LAZAREV, *Koszul duality for compactly generated derived categories of second kind*, *Journal of Noncommutative Geometry*, 15 (2021), pp. 1355–1371.
- [15] J. HOLSTEIN AND A. LAZAREV, *Enriched Koszul duality for dg categories*, arXiv preprint arXiv:2211.08118, (2022).
- [16] J. V. S. HOLSTEIN, *Morita cohomology*, *Math. Proc. Cambridge Philos. Soc.*, 158 (2015), pp. 1–26.
- [17] B. KELLER, *On differential graded categories*, in *Proceedings of the International Congress of Mathematicians Madrid, 2006*, pp. 151–190.
- [18] ———, *A remark on Hochschild cohomology and Koszul duality*, in *Advances in Representation Theory of Algebras, Contemporary Mathematics*, vol. 761, American Mathematical Society, 2021, pp. 131–136.
- [19] A. NEEMAN, *The connection between the k-theory localization theorem of thomason, trobaugh and yao and the smashing subcategories of bousfield and ravenel*, in *Annales scientifiques de l’Ecole normale supérieure*, vol. 25, 1992, pp. 547–566.
- [20] D. O. ORLOV, *Triangulated categories of singularities and D-branes in Landau-Ginzburg models*, *Tr. Mat. Inst. Steklova*, 246 (2004), pp. 240–262.
- [21] A. POLISHCHUK AND L. POSITSSELSKI, *Hochschild (co)homology of the second kind I*, *Transactions of the American Mathematical Society*, 364 (2012), pp. 5311–5368.
- [22] L. POSITSSELSKI, *Two kinds of derived categories, Koszul duality, and comodule-contramodule correspondence*, *Memoirs of the American Mathematical Society*, 212 (2011).
- [23] A. PREYGEL, *Thom-Sebastiani and Duality for Matrix Factorizations*, ArXiv e-prints, (2011), arXiv:1101.5834.
- [24] M. RIVERA, *An algebraic model for the free loop space*, arXiv preprint arXiv:2210.10096, (2022).
- [25] S. SCHWEDE, *The stable homotopy category has a unique model at the prime 2*, *Advances in Mathematics*, 164 (2001), pp. 24–40.
- [26] G. TABUADA, *Invariants additifs de dg-catégories*, *International Mathematics Research Notices*, 2005 (2005), pp. 3309–3339.
- [27] B. TOËN, *The homotopy theory of dg-categories and derived Morita theory*, *Inventiones mathematicae*, 167 (2006), pp. 615–667.

- [28] ———, *Lectures on dg-categories*, in Topics in algebraic and topological K -theory, vol. 2008 of Lecture Notes in Math., Springer, Berlin, 2011, pp. 243–302.
- [29] J. Tu, *Matrix factorizations via Koszul duality*, *Compositio Mathematica*, 150 (2014), pp. 1549–1578.

DEPARTMENT OF MATHEMATICS, UNIVERSITÄT HAMBURG, 20146 HAMBURG, GERMANY
Email address: `julian.holstein@uni-hamburg.de`

DEPARTMENT OF MATHEMATICS AND STATISTICS, LANCASTER UNIVERSITY, LANCASTER LA1 4YF, UNITED KINGDOM
Email address: `a.lazarev@lancaster.ac.uk`

SCHOOL OF MATHEMATICS, UNIVERSITY OF BIRMINGHAM, BIRMINGHAM B15 2TT, UNITED KINGDOM
Email address: `a.guan@bham.ac.uk`