DOUBLETHINK: SIMULTANEOUS BAYESIAN-FREQUENTIST MODEL-AVERAGED HYPOTHESIS TESTING

HELEN R. FRYER, NICOLAS ARNING, DANIEL J. WILSON

UNIVERSITY OF OXFORD BIG DATA INSTITUTE, OXFORD POPULATION HEALTH, AND DEPARTMENT FOR CONTINUING EDUCATION

26 July 2024

NB: This is a work-in-progress with some gaps left to be filled, notably the proof to Theorem 4.

Abstract

Establishing the frequentist properties of Bayesian approaches widens their appeal and offers new understanding. In hypothesis testing, Bayesian model averaging addresses the problem that conclusions are sensitive to variable selection. But Bayesian false discovery rate (FDR) guarantees are contingent on prior assumptions that may be disputed. Here we show that Bayesian model-averaged hypothesis testing is a closed testing procedure that controls the frequentist familywise error rate (FWER) in the strong sense. The rate converges pointwise as the sample size grows and, under some conditions, uniformly. The 'Doublethink' method computes simultaneous posterior odds and asymptotic *p*-values for model-averaged hypothesis testing. We explore its benefits, including post-hoc variable selection, and limitations, including finite-sample inflation, through a Mendelian randomization study and simulations comparing approaches like LASSO, stepwise regression, the Benjamini-Hochberg procedure and e-values.

Keywords: Bayesian model averaging, hypothesis tests, post-hoc variable selection, false discovery rate, familywise error rate, Doublethink

1. INTRODUCTION

Hypothesis testing is a fundamental scientific approach for identifying variables that affect an outcome of interest [Pearson, 1900, Fisher, 1925, Neyman and Pearson, 1933]. In observational studies, however, the statistical evidence that one variable directly affects an outcome typically depends on which other variables are included in the model, because of confounding and mediation [Vandenbroucke, 2002, Pearl et al., 2016]. Uncertainty in model selection can, therefore, strongly influence conclusions and should be accounted for, particularly in big data settings featuring thousands of variables. Bayesian model averaging provides a solution [Raftery, 1995], but in many settings Bayes has not become mainstream.

The advantages of Bayesian methods, like model-averaging and interpretability, are counterbalanced by various concerns [Gelman, 2008], chiefly the sensitivity to debatable prior assumptions. Whereas Bernstein-von Mises theorem [Borwanker et al., 1971] shows that Bayesian and frequentist parameter estimates and confidence regions can converge for large, informative samples, the influence of the prior on hypothesis tests remains uncomfortably direct. The posterior odds of one hypothesis versus another can be doubled by simply doubling the prior odds. Bayes factors are similarly manipulable through the prior on effect sizes [Gelman and Shalizi, 2013]. This is partly a problem of scale: in large samples, the prior contributes only a constant term to the log posterior odds and gets overwhelmed, and indeed, Bayesian hypothesis testing is consistent by Doob's theorem [Doob, 1949, Kass and Wasserman, 1995, O'Hagan, 1995]. Yet even in large samples, priors influence testing at the boundary of statistical significance, which can lead different researchers to different conclusions. Likewise, Bayesian false discovery rate (FDR) guarantees are contingent on the prior, which may be disputed.

Establishing the frequentist properties of Bayesian tests can broaden their appeal and afford new insights [Bayarri and Berger, 2004]. There is long-standing interest in bridging this Bayesian-frequentist divide [Held and Ott, 2018]. Formalizing earlier ideas expressed by Good [1992], Sellke, Bayarri, and Berger [2001] introduced methods that convert *p*-values to maximum Bayes factors and, by implication, Bayes factors to minimum *p*-values. Zhou and Guan [2018] derived the null distribution of Bayes factors in linear regression, enabling direct interconversion of Bayesian and frequentist measures of evidence. For models in which chi-squared statistics are sufficient, Johnson's [2005, 2008] approach (see also Wakefield [2009]) enabled one-to-one interconversion between Bayes factors and *p*-values via the maximized likelihood ratio (MLR), similarly to the Bayesian information criterion (BIC; Schwarz [1978]), to which it converges. Vovk and Wang [2021] developed conversion of Bayes factors involving a simple null and a possibly model-averaged alternative hypothesis (*e*-values) to maximum *p*-values, together with a closed testing procedure (CTP; Marcus et al. [1976]) to control the familywise error rate (FWER). FWER is considered standard control in frequentist multiple testing [Tukey, 1953]. Goeman and Solari [2011] developed general *p*-value-based CTPs for FWER control that allow multilevel testing, meaning that thresholds for arbitrary combinations of hypotheses are pre-determined and significant groups of variables can be freely identified post-hoc [Meinshausen, 2008, Goeman et al., 2019].

In this paper, we investigate the frequentist properties of Bayesian model-averaged hypothesis tests involving nested models, with uncertainty in both the null and alternative hypotheses. Surprisingly, we show that Bayesian tests are CTPs that control both the Bayesian FDR and the frequentist FWER in the strong sense at some rate. We quantify a large-sample asymptotic false positive rate and FWER, which enables interconversion of Bayesian model-averaged posterior odds and asymptotic *p*-values, on which we base a new method 'Doublethink'. We investigate the limitations of Doublethink, including the nature of asymptotic convergence, finite-sample inflation, and the requirement for minimum effect size and maximum correlation assumptions. We demonstrate the benefits of the approach, including multilevel testing and post-hoc variable selection, through a simulation study benchmarking alternative approaches and a Mendelian randomization study of age-related macular degeneration. Finally, we discuss the broader implications of this work.

2. Bayesian hypothesis testing is a closed testing procedure that controls the familywise error rate in the strong sense

Definition 1 (Frequentist familywise error rate control). Marcus, Peritz, and Gabriel [1976] introduced the closed testing procedure (CTP) to control the frequentist familywise error rate (FWER) in the strong sense. Suppose random variable \boldsymbol{y} has a probability mass or density function $p(\boldsymbol{y}; \boldsymbol{x}, \theta)$ that depends on auxiliary data \boldsymbol{x} and parameters $\theta \in \Theta$ that are partitioned into ν parameters-ofinterest, β , and ζ nuisance parameters, γ . The aim is to test a set of hypotheses about β defined by $W = \{\omega_s\}$, where $\omega_s \subset \Theta$ and \boldsymbol{s} is an index. W must be closed under intersection, meaning $\omega_s, \omega_{s'} \in W$ implies $\omega_s \cap \omega_{s'} \in W$. A 'local' test controls the false positive rate (FPR) at level α_s by rejecting $\theta \in \omega_s$ when $\psi_s(\boldsymbol{y}) = 1$ (rather than 0) such that:

$$FPR_{\boldsymbol{s}} := \sup_{\boldsymbol{\theta} \in \omega_{\boldsymbol{s}}} \Pr\left(\psi_{\boldsymbol{s}}(\boldsymbol{y}) = 1; \boldsymbol{x}, \boldsymbol{\theta}\right) = \alpha_{\boldsymbol{s}}.$$
 (1)

A CTP rejects $\theta \in \omega_s$ when $\phi_s(y) = 1$, which indicates rejection of all intersection hypotheses in W:

$$\phi_{\boldsymbol{s}}(\boldsymbol{y}) = \min_{\boldsymbol{\omega}_{\boldsymbol{r}} \subseteq \boldsymbol{\omega}_{\boldsymbol{s}}} \psi_{\boldsymbol{r}}(\boldsymbol{y}).$$
(2)

If $\omega_{\tilde{s}} \in W$ is the intersection of true null hypotheses, this CTP controls the FWER in the strong sense (i.e. $\tilde{s} \ge 0$) at level (see SI Section 2.1)

FWER :=
$$\sup_{\theta \in \omega_{\tilde{s}}} \Pr\left(\max_{\omega_{s} \subseteq \omega_{\tilde{s}}} \phi_{s}(\boldsymbol{y}) = 1; \boldsymbol{x}, \theta\right) \leq \alpha_{\tilde{s}}$$

 $\leq \max\left\{\alpha_{s} : \omega_{s} \in W\right\}.$ (3)

Definition 2 (Bayesian false discovery rate control). A Bayesian test rejects the null hypothesis that $\theta \in \omega_s$ when $\psi_s(\boldsymbol{y}) := I(\text{PO}_{\omega_s^c:\omega_s} \ge \tau)$, i.e. when the posterior odds

$$PO_{\omega_{\boldsymbol{s}}^{\mathsf{c}}:\omega_{\boldsymbol{s}}} = \frac{\int_{\omega_{\boldsymbol{s}}^{\mathsf{c}}} p(\boldsymbol{y};\boldsymbol{x},\theta) \, \mathrm{d}\Pi(\theta)}{\int_{\omega_{\boldsymbol{s}}} p(\boldsymbol{y};\boldsymbol{x},\theta) \, \mathrm{d}\Pi(\theta)},$$
(4)

exceed some threshold τ , where *I* is an indicator function, $\omega_s^{\mathsf{c}} = \Theta \setminus \omega_s$, and $\Pi(\theta)$ is the prior distribution on $\theta \in \Theta$. It controls the Bayesian false discovery rate (FDR) locally [Efron et al., 2001] at level

$$FDR_{s} := Pr(\theta \in \omega_{s} | \boldsymbol{y}, \boldsymbol{x}) \psi_{s}(\boldsymbol{y})$$

$$= \frac{1}{1 + PO_{\omega'_{s}:\omega_{s}}} I(PO_{\omega'_{s}:\omega_{s}} \ge \tau)$$

$$\leq \frac{1}{1 + \tau}$$
(5)

and globally at level

$$FDR := \frac{\sum_{\boldsymbol{s}} FDR_{\boldsymbol{s}}}{\max\left\{1, \sum_{\boldsymbol{s}} \psi_{\boldsymbol{s}}(\boldsymbol{y})\right\}} \leq \frac{1}{1+\tau}.$$
(6)

Theorem 1 (Bayesian hypothesis tests simultaneously control the Bayesian FDR and the frequentist FWER). Bayesian hypothesis tests are a type of CTP known as a shortcut CTP. That is, $\phi_s(\boldsymbol{y}) = \psi_s(\boldsymbol{y}) = 1$ (rejection of $\theta \in \omega_s$) automatically implies $\phi_r(\boldsymbol{y}) = \psi_r(\boldsymbol{y}) = 1$ for all intersection hypotheses $\omega_r \subset \omega_s$. Therefore by Definitions 1 and 2 they simultaneously control the frequentist FWER in the strong sense at level max{ $\alpha_s : \omega_s \in W$ } and the Bayesian FDR at level $1/(1 + \tau)$.

Proof. First, define the posterior odds versus the grand null hypothesis, s = 0:

$$PO_{\omega_{s}} = \frac{\int_{\omega_{s}} p(\boldsymbol{y}; \boldsymbol{x}, \theta) \, \mathrm{d}\Pi(\theta)}{\int_{\omega_{0}} p(\boldsymbol{y}; \boldsymbol{x}, \theta) \, \mathrm{d}\Pi(\theta)}.$$
(7)

By definition, $\omega_{\boldsymbol{r}} \subset \omega_{\boldsymbol{s}}$ and $\omega_{\boldsymbol{r}}^{\mathsf{c}} \supset \omega_{\boldsymbol{s}}^{\mathsf{c}}$, so $\mathrm{PO}_{\omega_{\boldsymbol{r}}} \leq \mathrm{PO}_{\omega_{\boldsymbol{s}}}$ and $\mathrm{PO}_{\omega_{\boldsymbol{r}}}^{\mathsf{c}} \geq \mathrm{PO}_{\omega_{\boldsymbol{s}}}^{\mathsf{c}}$. Since $\mathrm{PO}_{\omega_{\boldsymbol{s}}}^{\mathsf{c}} : \omega_{\boldsymbol{s}} = \mathrm{PO}_{\omega_{\boldsymbol{s}}}^{\mathsf{c}} / \mathrm{PO}_{\omega_{\boldsymbol{s}}}$ (by Equation 4) then $\mathrm{PO}_{\omega_{\boldsymbol{r}}}^{\mathsf{c}} : \omega_{\boldsymbol{r}} \geq \mathrm{PO}_{\omega_{\boldsymbol{s}}}^{\mathsf{c}} : \omega_{\boldsymbol{s}}$ so $\psi_{\boldsymbol{s}}(\boldsymbol{y}) = 1$ immediately implies $\psi_{\boldsymbol{r}}(\boldsymbol{y}) = 1$.

The rest of this paper focuses on quantifying the level at which a Bayesian test controls the FWER, asymptotically for large samples, in a general regression setting.

3. Frequentist false positive rate of a Bayesian model-averaged regression converges pointwise as the sample size grows

We apply Theorem 1 to construct a joint Bayesian-frequentist approach to model-averaged hypothesis testing in a regression setting.

Definition 3 (**Regression problem**). We consider a general regression with *n* observed outcomes, $\boldsymbol{y} = (y_1, \ldots, y_n)^T$ and ν regression coefficients, $\beta = (\beta_1, \ldots, \beta_\nu)^T$, for candidate explanatory variables $x_{ij}, i = 1 \ldots n, j = 1 \ldots \nu$. The aim is to identify which regression coefficients are non-zero. We index null hypotheses by a binary vector $\boldsymbol{s} \in \mathcal{S} = 2^{\nu}$ such that $\omega_{\boldsymbol{s}} = \{\theta : \beta_j = 0 \forall s_j = 0\} \subseteq \Theta = \mathbb{R}^{\nu+\zeta}$. We define models in terms of variable selection, so that model \boldsymbol{s} has parameter space $\Theta_{\boldsymbol{s}} = \{\theta : \beta_j = 0 \forall s_j = 0\}$ ($\theta : \beta_j = 0 \forall s_j = 0$); $\beta_j \neq 0 \forall s_j = 1\} \subseteq \Theta$. The models are disjoint with respect to β , unlike the null hypotheses. All models include ζ nuisance parameters, $\gamma = (\gamma_1, \ldots, \gamma_\zeta)^T$. Together, we write $\theta = {\beta \choose \gamma}$.

In general, the finite sample properties of frequentist methods, including FWER, are difficult to obtain. Instead it is standard to use approximations based on the asymptotic distribution of methods as the sample size, n, tends to infinity. For nested hypotheses, the workhorse is the likelihood ratio test.

Definition 4 (Likelihood ratio test; LRT). Wilks [1938] and Wald [1943] showed that the distribution of the maximized likelihood ratio (MLR) for null model s nested in alternative t (i.e. $t_j \ge s_j \forall j = 1 \dots \nu$ and $\sum_{j=1}^{\nu} t_j > \sum_{j=1}^{\nu} s_j$, or t > s for shorthand)

$$R_{\Theta_t:\Theta_s} := \frac{\sup_{\theta \in \Theta_t} p(\boldsymbol{y}; \boldsymbol{x}, \theta)}{\sup_{\theta \in \Theta_s} p(\boldsymbol{y}; \boldsymbol{x}, \theta)}$$
(8)

converges (not necessarily uniformly), such that the deviance,

$$2 \log R_{\Theta_t:\Theta_s} \xrightarrow{d} \chi^2_{|t|-|s|}, \qquad n \to \infty, \tag{9}$$

where χ_k^2 represents a chi-squared distribution with k degrees of freedom, assuming the n outcomes are independent realizations each with likelihood that satisfies standard regularity conditions (SI Section 1) under the 'local alternatives' assumption (Equation 47) [Davidson and Lever, 1970]. The FPR of this test converges,

$$\alpha_{t:s} := \sup_{\theta \in \Theta_s} \Pr\left(2 \log R_{\Theta_t:\Theta_s} \ge x_{\operatorname{crit}}; \theta\right) \sim \Pr\left(\chi_{t:s}^2 \ge x_{\operatorname{crit}}\right), \quad n \to \infty, \tag{10}$$

for some threshold $x_{\rm crit}$, in general pointwise meaning that

$$\sup_{\theta \in \Theta_{s}} \lim_{n \to \infty} \frac{\Pr\left(2 \log R_{\Theta_{t}:\Theta_{s}} \ge x_{\text{crit}}; \theta\right)}{\Pr\left(\chi_{t:s}^{2} \ge x_{\text{crit}}\right)} = 1.$$
(11)

As shorthand we will write $R_{t:s} := R_{\Theta_t:\Theta_s}$ and $R_s := R_{s:0}$.

One method that addresses the regression problem while controlling the FWER, approximately for large n, is given below.

Example 1 (FWER control of the regression problem). The LRT [Wilks, 1938, Wald, 1943] can be combined with Bonferroni correction [Bonferroni, 1936] to define a leave-one-out test that drops variables relative to the grand alternative:

$$\mathcal{P} = \{ p_{1:t} = \Pr\left(\chi_1^2 \ge 2 \log R_{1:t}\right) : t \in \mathcal{S}, |t| = \nu - 1 \}$$
(12)

$$\phi_{\boldsymbol{s}}(\boldsymbol{y}) = \phi_{\boldsymbol{s}}^{\text{Bon}}(\mathcal{P}) = I\left(\min_{\substack{\boldsymbol{t} \geq \boldsymbol{s} \\ |\boldsymbol{t}| = \nu - 1}} p_{\boldsymbol{1}:\boldsymbol{t}} \leq \frac{\alpha}{\nu}\right).$$
(13)

This controls the FWER at level α , to the order of the large *n* approximation. There is a large literature on combined tests for *p*-values that are more powerful than Bonferroni correction under certain conditions [Loughin, 2004], for example Simes [1986], Hommel [1988], Meijer et al. [2019], Wilson [2019a], Liu and Xie [2020].

Despite its familiarity, it can be hard to solve the regression problem while controlling the FWER and maintaining power, and Example 1 has various problems. When the candidate explanatory variables, \boldsymbol{x} , are correlated, parameter estimates can be noisy and *p*-values deflated (conservative), which reduces power. Under collinearity of \boldsymbol{x} , it may not be possible to fit the grand alternative model at all, in which case the method does not work.

Instead, it is common to select a model, s < 1, for example using expert opinion, univariable associations, machine learning, or statistical optimization [Miller, 2002, Porwal and Raftery, 2022], then perform |s| leave-one-out likelihood ratio tests and, perhaps, $(\nu - |s|)$ add-one-in tests. Sparse models $(|s| \ll \nu)$ are usually preferred on principle, and to mitigate variance and deflation due to correlated variables (see e.g. Tibshirani [1996]). Often, adjustment is made for multiple testing, as per Example 1, but uncertainty in model selection – which can affect conclusions – is rarely accounted for, partly because the effects of selecting s on the distribution of test statistics is not known, in general.

Bayesian model averaging presents a solution to these challenges and, by Theorem 1, controls the FWER. To be practical, we need to quantify the level at which the FWER is controlled. As a first step we pursue an asymptotic approximation to the FPR by extending work by Schwarz [1978] and Johnson [2005, 2008].

Definition 5 (Bayesian information criterion; BIC). Schwarz [1978] developed the BIC for model selection as an approximate Bayes factor in large samples. It implies the posterior odds of model t versus model s (not necessarily nested) are

$$\mathrm{PO}_{\Theta_t;\Theta_s} \approx \mu_{\Theta_t;\Theta_s} n^{-(|t|-|s|)/2} R_{t;s},$$
(14)

where $\mu_{\Theta_t:\Theta_s} = \int_{\Theta_t} d\Pi(\theta) / \int_{\Theta_s} d\Pi(\theta)$ are the prior odds. The BIC has error of order $O_p(1)$ when used to approximate the log posterior odds or log Bayes factor [Kass and Wasserman, 1995]. As shorthand we will write $PO_{t:s} := PO_{\Theta_t:\Theta_s}$ and $PO_s := PO_{s:0}$, and likewise for $\mu_{t:s}$.

Definition 6 (Joint Bayesian-frequentist test: Johnson model). Johnson [2005, 2008] developed simultaneous Bayesian-frequentist inference that defines the posterior odds of model t versus model s (not necessarily nested) via the MLR as

$$PO_{t:s} = \mu_{t:s} \xi_n^{(|t|-|s|)/2} R_{t:s}^{1-\xi_n}, \quad \xi_n = \frac{h}{n+h},$$
(15)

where h is the prior precision parameter. This closely resembles Equation 14, to which it converges as $n \to \infty$, but is proper in the sense that $\mathbb{E}_{\Pi}[\mathrm{PO}_{t:s} \mid \theta \in \Theta_s] = \mu_{t:s}$. In general, the model mirrors the

Classical q	uantity		Вау	vesian counterpart	
Quantity	Expression	SI Eqn.	Quantity	Expression	SI Eqn.
Maximum likelihood estimate	$\hat{ heta}^{s}$	22	Posterior mean	$rac{n}{n+h}\hat{ heta}^{m{s}}$	62
Estimator variance	$[J(\hat{\theta}^{\boldsymbol{s}})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}}]^{-1}$	23	Posterior variance	$\frac{n}{n+h}[J(\hat{\theta}^{\boldsymbol{s}})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}}]^{-1}$	63
Maximized likelihood ratio	<i>R</i> s	26		$\frac{n}{n+h} [J(\hat{\theta}^{s})_{\mathcal{F}_{s},\mathcal{F}_{s}}]^{-1}$ $\left(\frac{h}{n+h}\right)^{ s /2} R_{s}^{n/n+h}$	

TABLE 1. Connections between Bayesian and classical inference in Johnson's [2005, 2008] model. $J(\hat{\theta}^s)_{\mathcal{F}_s,\mathcal{F}_s}$ is the observed information matrix for model s at the MLE.

assumptions of the LRT (Definition 4), including large n and local alternatives. It is derived under the following conjugate prior:

$$\theta_{\mathcal{F}_{\boldsymbol{s}}} \mid \boldsymbol{\theta} \in \Theta_{\boldsymbol{s}} \stackrel{d}{=} \mathcal{N}_{|\boldsymbol{s}|+\zeta} \left(\boldsymbol{0}, h^{-1} \left[\mathcal{I}(\boldsymbol{0})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}} \right]^{-1} \right), \tag{16}$$

where \mathcal{F}_s is the index set of unconstrained parameters in model s, $\mathcal{N}_k(\boldsymbol{m}, \boldsymbol{\Sigma})$ represents a multivariate Normal distribution on k dimensions with mean vector \boldsymbol{m} and variance matrix $\boldsymbol{\Sigma}$, $\mathcal{I}(\theta)_{\mathcal{F}_s,\mathcal{F}_s}$ is the per-observation Fisher information matrix for model s evaluated at θ . The prior resembles Zellner's [1986] g-prior with g = n/h; h = 1 corresponds to the unit information prior [Kass and Wasserman, 1995, Liang et al., 2008]. The test PO_{t:s} $\geq \tau$ controls the local Bayesian FDR at level $1/(1 + \tau)$ (Definition 2) and the FPR converges in general pointwise

$$\sup_{\theta \in \Theta_{\boldsymbol{s}}} \Pr\left(\mathrm{PO}_{\boldsymbol{t}:\boldsymbol{s}} \ge \tau; \theta\right) \sim \Pr\left(\chi_{\boldsymbol{t}:\boldsymbol{s}}^2 \ge 2 \log \frac{\tau}{\mu_{\boldsymbol{t}:\boldsymbol{s}} \, \xi_n^{(|\boldsymbol{t}| - |\boldsymbol{s}|)/2}}\right), \quad n \to \infty,\tag{17}$$

meaning that

$$\sup_{\theta \in \Theta_{\boldsymbol{s}}} \lim_{n \to \infty} \frac{\Pr\left(\mathrm{PO}_{\boldsymbol{t}:\boldsymbol{s}} \ge \tau; \theta\right)}{\Pr\left(\chi_{\boldsymbol{t}:\boldsymbol{s}}^2 \ge 2 \log \frac{\tau}{\mu_{\boldsymbol{t}:\boldsymbol{s}} \,\xi_n^{\left(|\boldsymbol{t}| - |\boldsymbol{s}|\right)/2}}\right)} = 1.$$
(18)

Johnson's model formalizes the asymptotic relationships not only between classical and Bayesian hypothesis tests, but point estimates and variances too (Table 1).

Next we define a model-averaged extension of Johnson's [2005, 2008] model, including a prior on models. Notably, the posterior odds in Definition 6 would be pre-determined if the prior on θ were continuous since that would require $\int_{\Theta_s} d\Pi(\theta) = I(s = 1)$. Therefore pursuing Bayesian hypothesis testing with null hypotheses $\{\theta : \beta_j = 0 \forall s_j = 0\}$ implies a discontinuous prior.

Definition 7 (Bayesian model-averaged hypothesis testing: Doublethink model). We assume that the prior odds of model t versus model s take the form

$$\mu_{t:s} = \mu^{|t|-|s|} \tag{19}$$

meaning the prior probability that $\beta_j \neq 0$ is independent for each variable and equal to $\mu/(1 + \mu)$. We can write the model-averaged posterior odds of null hypothesis $\omega_{\boldsymbol{v}} = \{\theta : \beta_j = 0 \forall v_j = 0\}$, which constrains the variables indexed by set $\mathcal{V} = \{j : \boldsymbol{v}_j = 0\}$, via Equation 4 in terms of the Johnson model (Equation 15) as

$$PO_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} := PO_{\omega_{v}^{c}:\omega_{v}} = \frac{\int_{\omega_{v}^{c}} p(\boldsymbol{y};\boldsymbol{x},\theta) \, \mathrm{d}\Pi(\theta)}{\int_{\omega_{v}} p(\boldsymbol{y};\boldsymbol{x},\theta) \, \mathrm{d}\Pi(\theta)} \\ = \frac{\sum_{\boldsymbol{s}\in\mathcal{A}_{\mathcal{V}}} \int_{\Theta_{\boldsymbol{s}}} p(\boldsymbol{y};\boldsymbol{x},\theta) \, \mathrm{d}\Pi(\theta)}{\sum_{\boldsymbol{s}\in\mathcal{O}_{\mathcal{V}}} \int_{\Theta_{\boldsymbol{s}}} p(\boldsymbol{y};\boldsymbol{x},\theta) \, \mathrm{d}\Pi(\theta)} = \frac{\sum_{\boldsymbol{s}\in\mathcal{A}_{\mathcal{V}}} PO_{\boldsymbol{s}}}{\sum_{\boldsymbol{s}\in\mathcal{O}_{\mathcal{V}}} PO_{\boldsymbol{s}}}$$
(20)

where we define the set of models compatible with null hypothesis $\omega_{\boldsymbol{v}}$ as $\mathcal{O}_{\mathcal{V}} = \{\boldsymbol{s} \in \mathcal{S} : \Theta_{\boldsymbol{s}} \subseteq \omega_{\boldsymbol{v}}\}$, and the models consistent with the alternative hypothesis $\omega_{\boldsymbol{v}}^{\mathsf{c}}$ as $\mathcal{A}_{\mathcal{V}} = \mathcal{S} \setminus \mathcal{O}_{\mathcal{V}}$. The test of the realized model-averaged posterior odds, $PO_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}$,

$$\psi_{\boldsymbol{v}}(\boldsymbol{y}) = I\left(\mathrm{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau\right)$$
(21)

controls the Bayesian FDR locally and globally at level $1/(1+\tau)$ (by Definition 2) and (by Definition 1 and Theorem 1) the FWER at level

$$\max_{\boldsymbol{\omega}_{\boldsymbol{v}}\in W} \sup_{\boldsymbol{\theta}\in\boldsymbol{\omega}_{\boldsymbol{v}}} \Pr\left(\psi_{\boldsymbol{v}}(\boldsymbol{y})=1;\boldsymbol{x};\boldsymbol{\theta}\right).$$
(22)

It remains to quantify the rate at which the FPRs and FWER are controlled. We note that whereas the Johnson model is derived under a specific prior, its close connection to the BIC implies that the model-averaged posterior odds in Definition 7 may serve as a useful large sample approximation in a wide variety of settings.

Theorem 2 (Frequentist false positive rate of the Bayesian model-averaged hypothesis test in large samples). Asymptotically, for large sample size, the frequentist FPR of the Bayesian modelaveraged hypothesis test in Definition 7, which rejects the null hypothesis, $\omega_{v} = \{\theta : \beta_{j} = 0 \forall j \in \mathcal{V}\},$ (where $\mathcal{V} = \{j : v_{j} = 0\}$), when $\mathrm{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \geq \tau$, converges pointwise to

$$\alpha_{\boldsymbol{v}} = \sup_{\boldsymbol{\theta} \in \boldsymbol{\omega}_{\boldsymbol{v}}} \Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau ; \boldsymbol{\theta}\right) \sim \Pr\left(\chi_{1}^{2} \ge 2\log\frac{\tau}{|\mathcal{V}|\,\mu\sqrt{\xi_{n}}}\right), \quad n \to \infty$$
(23)

meaning

$$\sup_{\theta \in \omega_{\boldsymbol{v}}} \lim_{n \to \infty} \frac{\Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau; \; \theta\right)}{\Pr\left(\chi_{1}^{2} \ge 2\log \frac{\tau}{|\mathcal{V}| \, \mu \sqrt{\xi_{n}}}\right)} = 1.$$
(24)

Proof. See SI Section 3.1

Corollary 1 (The model-averaged deviance has a limiting chi-squared distribution in the tail). Define the model-averaged MLR as

$$R_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} := \frac{\mathrm{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}}{\left(\mu\sqrt{\xi}_{n}+1\right)^{|\mathcal{V}|}-1} \sim \frac{\mathrm{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}}{|\mathcal{V}|\,\mu\sqrt{\xi}_{n}}, \quad n \to \infty.$$
(25)

The model-averaged deviance converges pointwise in the tail to a chi-squared distribution with one degree of freedom:

$$2 \log R_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \xrightarrow{d} \chi_1^2, \qquad n \to \infty$$
(26)

in the sense that

$$\sup_{\theta \in \omega_{\boldsymbol{v}}} \lim_{n \to \infty} \frac{\Pr\left(2 \log R_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge x + \log n; \theta\right)}{\Pr\left(\chi_{1}^{2} \ge x + \log n\right)} = 1.$$
(27)

Corollary 2 (The model-averaged posterior odds can be transformed into an asymptotic *p*-value). The Bayesian model-averaged test in Definition 7, which rejects the null hypothesis $\omega_{\boldsymbol{v}} = \{\theta : \beta_j = 0 \forall j \in \mathcal{V}\}$, (where $\mathcal{V} = \{j : \boldsymbol{v}_j = 0\}$), when $\text{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \geq \tau$, is equivalent to a frequentist test that rejects the null when the asymptotic *p*-value $p_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}$ is less than or equal to the asymptotic false positive rate $\alpha_{\boldsymbol{v}}$ (Equation 23), where

$$p_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \sim \Pr\left(\chi_1^2 \ge 2\log\frac{\mathrm{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}}{|\mathcal{V}|\,\mu\sqrt{\xi_n}}\right), \qquad n \to \infty.$$
 (28)

This result extends Johnson's [2005, 2008] interconversion of MLRs and *p*-values for nested hypothesis tests to the model-averaged setting.

In the next section we derive the asymptotic FWER and adjusted *p*-values that account for multiple testing.

4. Strong-sense familywise error rate of a Bayesian model-averaged regression converges pointwise as the sample size grows

Theorem 3 (Familywise error rate of the Bayesian model-averaged hypothesis test in large samples). Asymptotically, for large sample size, the strong-sense frequentist FWER of the Bayesian model-averaged hypothesis test in Definition 7, which rejects all null hypotheses $\omega_{v} = \{\theta : \beta_{j} = 0 \forall v_{j} = 0\}, v \in 2^{\nu}$, for which $PO_{\omega_{s}^{c}:\omega_{v}} \geq \tau$, converges pointwise to

$$\alpha := \max_{\omega_{\boldsymbol{v}} \in W} \sup_{\theta \in \omega_{\boldsymbol{v}}} \Pr\left(\operatorname{PO}_{\omega_{\boldsymbol{v}}^{\mathsf{c}}:\omega_{\boldsymbol{v}}} \ge \tau; \theta\right) \sim \Pr\left(\chi_{1}^{2} \ge 2\log\frac{\tau}{\nu \mu \sqrt{\xi_{n}}}\right), \quad n \to \infty$$
(29)

meaning

$$\max_{\omega_{\boldsymbol{v}}\in W} \sup_{\theta\in\omega_{\boldsymbol{v}}} \lim_{n\to\infty} \frac{\Pr\left(\mathrm{PO}_{\omega_{\boldsymbol{v}}^{\mathsf{c}}:\omega_{\boldsymbol{v}}} \ge \tau; \theta\right)}{\Pr\left(\chi_{1}^{2} \ge 2\log\frac{\tau}{\nu \mu \sqrt{\xi_{n}}}\right)} \le 1,$$
(30)

 $\mathrm{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \equiv \mathrm{PO}_{\omega_{\boldsymbol{v}}^{\mathsf{c}}:\omega_{\boldsymbol{v}}} \text{ and } \mathcal{V} = \{j: \boldsymbol{v}_{j} = 0\} \text{ by Equation 20.}$

Proof. By Definition 1, Theorem 1, Definition 7 and Theorem 2. See SI Section 3.2. \Box

Corollary 3 (The model-averaged posterior odds can be transformed into an asymptotic adjusted *p*-value; Doublethink). The Bayesian model-averaged test in Definition 7, which rejects all null hypotheses $\{\theta : \beta_j = 0 \forall j \in \mathcal{V}\}, \mathcal{V} \subseteq \{1, \ldots, \nu\}$, for which $\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \geq \tau$, is equivalent to a frequentist test that rejects the same nulls when their asymptotic adjusted *p*-values $p^{\star}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}$ are less than or equal to the asymptotic FWER α (Equation 29), where

$$p_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}^{\star} \sim \Pr\left(\chi_{1}^{2} \ge 2\log\frac{\mathrm{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}}{\nu\,\mu\sqrt{\xi_{n}}}\right), \qquad n \to \infty.$$
 (31)

We call this approach, which (by Theorem 1) simultaneously controls the Bayesian FDR, locally and globally, at level $1/(1 + \tau)$, Doublethink.

Corollary 4 (Scaling of bounds on the Bayesian FDR and the asymptotic FWER). The asymptotic bound on the FWER, α , scales approximately linearly with the bound on the Bayesian FDR, $1/(1+\tau)$, the number of variables ν , the prior odds that $\beta_j \neq 0$, μ , the square root of the prior precision, \sqrt{h} , and the inverse square root of the sample size, $1/\sqrt{n}$:

$$\alpha \sim \frac{\nu \mu \sqrt{h}}{\tau} \sqrt{\frac{n_0}{n}} \Pr\left(\chi_1^2 \ge 2\log\sqrt{n_0}\right), \quad n \to \infty, \quad n_0 = O(n).$$
(32)

The bounds are not necessarily tight.

Proof. By Theorem 3, noting that the function $G(x) = \Pr(\chi_1^2 \ge 2\log x)$ is regularly varying in x with index $\lambda = -1$ which, by the definition of regular variation (Karamata [1933], Mikosch [1999] Definition 1.1.1), satisfies

$$\lim_{x \to \infty} \frac{G(c x)}{G(x)} = c^{\lambda}.$$
(33)

5. Uniform convergence of the false positive rate requires beta-min and rho-max assumptions in a two-variable model

Definition 8 (Two-variable model with simplifying assumptions). To study uniform convergence we consider a normal linear model with $\nu = 2$ regression coefficients (the variables-of-interest, β), an intercept (the nuisance parameter γ), a known variance σ^2 , and likelihood proportional to $p(\boldsymbol{y}; \boldsymbol{x}, \theta) = \prod_{i=1}^{n} p_{\mathcal{N}(0,1)} ((y_i - \gamma - \beta_1 x_{i1} - \beta_2 x_{i2})/\sigma)$. We assume the variables are standardized with means $\mathbb{E}[x_{\cdot 1}] = \mathbb{E}[x_{\cdot 2}] = 0$, variances $\mathbb{E}[x_{\cdot 1}^2] = \mathbb{E}[x_{\cdot 2}^2] = 1$, and correlation coefficient $\mathbb{E}[x_{\cdot 1} x_{\cdot 2}] = \rho$.




FIGURE 1. Inflation in the simplified two-variable model testing $\beta_1 = \beta_2 = 0$ (left) and $\beta_1 = 0$ (right). LEFT: FPR as a function of sample size: simulations (black line) and Theorem 2 (green line). Assuming $\mu = 1, h = 1, \tau = 9$. RIGHT: FPR as a function of β_2 for $\rho \in [0.0, 1.0]$ (shaded grey lines, labelled by ρ) and Theorem 2 (green dashed line). Assuming $n = 145, \mu = 0.1, h = 1, \sigma = 1, \tau = 9$. 10 million simulations in both.

There are three null hypotheses of interest, $\{\theta : \beta_j = 0 \forall j \in \mathcal{V}\}$, with $\mathcal{V} = \{1\}$, $\mathcal{V} = \{2\}$ or $\mathcal{V} = \{1, 2\}$. The first two scenarios are symmetric, so we only consider the first. As shorthand, we write the corresponding posterior odds as $PO_{\{1\}} = PO_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}$ with $\mathcal{V} = \{1\}$, and $PO_{\{12\}} = PO_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}$ with $\mathcal{V} = \{1, 2\}$.

Theorem 4 (Uniform convergence in a simplified two-variable model). In the simplified two-variable model (Definition 8), the FPR, and therefore the FWER, converge uniformly to the asymptotic expressions given by Theorems 2 and 3, subject to further assumptions: (i) $|\beta_j| \ge \beta_{\min} > 0$, and (ii) $|\rho| \le \rho_{\max} < 1$, meaning that

$$\lim_{n \to \infty} \sup_{\substack{\theta \in \omega_v^* \\ \rho \in P^*}} \frac{\Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau ; \theta\right)}{\Pr\left(\chi_1^2 \ge 2\log \frac{\tau}{|\mathcal{V}| \, \mu \sqrt{\xi_n}}\right)} = 1,$$
(34)

where $v_j = I(j \notin \mathcal{V}), \ \omega_{\boldsymbol{v}}^* = \{\theta \in \omega_{\boldsymbol{v}} : |\beta_j| \ge \beta_{\min} > 0 \ \forall j \notin \mathcal{V}\} \text{ and } P^* = \{\rho : |\rho| \le \rho_{\max} < 1\}.$

Proof. The proof is still a work-in-progress. See SI Section 3.3.

Corollary 5 (Inflation in the simplified two-variable model). In the simplified two-variable model (Definition 8), with fixed *n* and $\mathcal{V} = \{1\}$, the FPR can be inflated, with inflation factor ι , such that $\Pr(\operatorname{PO}_{\{1\}} \geq \tau; \theta) = \Pr\left(\chi_1^2 \geq 2\log\frac{\tau}{\mu\sqrt{\xi_n}\iota}\right) \sim \iota \Pr\left(\chi_1^2 \geq 2\log\frac{\tau}{\mu\sqrt{\xi_n}}\right)$. The inflation factor has range $\iota \in \left(\frac{\mu\sqrt{\xi_n}}{1+\mu\sqrt{\xi_n}}, \frac{1+\mu\sqrt{\xi_n}}{\mu\sqrt{\xi_n}}\right)$. At its worst, $\Pr(\operatorname{PO}_1 \geq \tau; \theta) \sim \Pr\left(\chi_1^2 \geq 2\log\tau\right)$.

There are two sources of inflation in the simplified two-variable model, illustrated by Figure 1. The first source (left panel) are higher-order tests involving more than one-degree-of-freedom (here, $PO_{11:00}$), whose contribution is assumed negligible by Theorem 2. As the sample size increases, this source of inflation recedes and the FPR tends to its asymptotic level.

The second source (right panel) is correlation between variables that have zero effect $(x_{.1} \text{ here})$ and those that have non-zero effect $(x_{.2} \text{ here})$. Inflation peaks at an effect size β_2 that depends on the value of ρ . For the smallest effect sizes, deflation is evident for higher $|\rho|$. For the largest effect sizes, inflation abates to its asymptotic level. Worst-case inflation increases with $|\rho|$, from none when $\rho = 0$, to the worst case as $|\rho| \to 1$. At the point $|\rho| = 1$, deflation is observed regardless of β_2 . Negative ρ and β_2 are not shown because the behaviour is symmetric. This second source of inflation does not abate as the sample size increases; instead, worst case inflation increases, and manifests at ever-smaller values of β_2 , inversely proportional to \sqrt{n} . At its worst, the FPR, instead of scaling inversely with \sqrt{n} , as per Corollary 4, remains on the order of the FDR (Corollary 5).

This explains the need to assume a minimum effect size $|\beta_2|$ and maximum correlation coefficient $|\rho|$ to ensure uniform convergence of the FPR to Theorem 2. Whereas it is impossible to control the minimum effect size in real data analysis it is possible to mitigate the inflating effect of highly correlated variables, as we will discuss later (Definition 9). Another countermeasure would be to increase either μ or ξ_n via h; however, this would diminish the advantages of Bayesian model-averaging in terms of sparsity, variance reduction and over-fitting.

6. Application to Mendelian randomization study of age-related macular degeneration

To trial Doublethink in a real data analysis, we applied it to a Mendelian randomization study of age-related macular degeneration (AMD).

Example 2 (Mendelian randomization study of age-related macular degeneration). Zuber et al. [2020] used two-sample multivariable Mendelian randomization [Sanderson et al., 2019] to study the causal effects of $\nu = 49$ biomarkers on AMD, and introduced a Bayesian model-averaging approach, MR-BMA. The aim was to identify which biomarkers had a non-zero effect on risk of AMD. In their model, the parameters-of-interest, β , were the direct causal effects of each biomarker on AMD. The outcome data **y** were z-scores for the direct effect of n = 145 genetic variants on AMD, estimated by a genome-wide association study (GWAS) [Fritsche et al., 2016]. For each genetic variant, the variables x_i were z-scores of its direct effects on the 49 biomarkers, estimated by another GWAS [Kettunen et al., 2016]. For each genetic variant, a common standard error was used in the denominator of all z-scores, applying the standard error from y_i to all the z-scores in x_i . There was one nuisance parameter, the error variance, γ , expected theoretically to equal one. This yielded the normal linear regression $p(\mathbf{y}; \mathbf{x}, \theta) = \prod_{i=1}^{n} \phi \left((y_i - \beta_1 x_{i1} - \cdots - \beta_{\nu} x_{i\nu}) / \sqrt{\gamma} \right)$.

In their Bayesian model-averaged analysis of AMD, the biomarkers for which Zuber et al. [2020] found the strongest evidence of a causal effect on AMD were XL.HDL.C and L.HDL.C (extra-large and large high-density lipoprotein cholesterol), with posterior probabilities 0.70 and 0.23, corresponding to Bayes factors of 21 and 2.7, respectively. Considering that neither would be considered 'strong' evidence according to Jeffreys' [1939] scale, and the prior influences Bayes factors anyway, this begs the question: (i) How would a frequentist evaluate this level of evidence? (ii) Would the evidence against the null hypothesis of no effect have been stronger by grouping XL.HDL.C and L.HDL.C together?

In partial answer to the first question, Julian et al. [2023] implemented a permutation procedure, which for this dataset yields unadjusted p-values below 0.005 for both XL.HDL.C and L.HDL.C (200 permutations, taking 80 hours). However, such procedures generally control the FWER only in the weak sense, i.e. when the grand null is true. The e-value approach of Vovk and Wang [2021] suggests maximum p-values (inverse Bayes factors) of 0.048 and 0.37 respectively, unadjusted for multiple testing, although its requirement for no nuisance parameters in the null hypothesis is not met here. The leave-one-out tests of Example 1 yielded unadjusted p-values of 0.90 and 0.16, probably a dilution of the signal due to many correlated variables.

Since the number of biomarkers was too large to fit all $2^{49} \approx 10^{15}$ models, and since many biomarkers were highly correlated, we selected a subset of 15 variables, for which we exhaustively analysed all $2^{15} = 32$ 768 models. We selected the 15 variables in a way that attempted to preserve most of the important variability, and the correlation structure, in the data: (1) We ranked the biomarkers by their two-sample univariable Mendelian randomization *p*-values for a causal association with AMD. (2) From the most to the least significant, we introduced 15 biomarkers iteratively. (3) We omitted biomarkers with an absolute correlation exceeding $|\rho| = 0.8$ with two or more biomarkers that we had

Biomarker	Posterior odds	Asymptotic p -value	Point estimate	Standard error
		(unadjusted)	(posterior mean)	(root posterior variance)
XL.HDL.C	1.75	0.001	0.33	0.26
L.HDL.C	0.58	0.004	0.14	0.20
Gln	0.08	0.033	-0.03	0.11
IDL.TG	0.06	0.048	-0.01	0.05
S.VLDL.TG	0.03	0.091	-0.01	0.04
TADLE 9 7	Two comple mult	iveriable Mondelian w	andomization in a l	Doublethink sub analysis

TABLE 2. Two-sample multivariable Mendelian randomization in a Doublethink sub-analysis of 15 biomarkers in Zuber et al. [2020] for direct effects on AMD risk. Top 5 variables shown.

already introduced. The 15 remaining biomarkers ranged from the first to the 24th most significantly associated with AMD. Three pairs were highly correlated: XL.HDL.C and L.HDL.C ($\rho = 0.82$); ApoB and IDL.TG ($\rho = 0.91$); S.HDL.TG and S.VLDL.TG ($\rho = 0.92$). The other nine biomarkers we included were LDL.D, Ace, XL.HDL.TG, ApoB, VLDL.D, M.HDL.C, His, Ala and Gln.

An important caveat is that the analysis of only a subset of the candidate variables affects the interpretation of those results. The next theorem clarifies the valid interpretation of such results.

Theorem 5 (Testing based on a sub-analysis). Suppose we conduct a sub-analysis in which a subset of variables, $\mathcal{V}' \subset \{1, \ldots, \nu\}$, are excluded from the analysis, in a Bayesian model-averaged setting (Definition 7). In the sub-analysis, the model-averaged posterior odds against the null hypothesis $\mathcal{V} \subset \{1, \ldots, \nu\}, \mathcal{V} \cap \mathcal{V}' = \emptyset$, could be written

$$\mathrm{PO}_{\mathcal{A}_{\mathcal{V}}^{-}:\mathcal{O}_{\mathcal{V}}^{-}} = \frac{\sum_{\boldsymbol{s}\in\mathcal{A}_{\mathcal{V}}^{-}}\mathrm{PO}_{\boldsymbol{s}}}{\sum_{\boldsymbol{s}\in\mathcal{O}_{\mathcal{V}}^{-}}\mathrm{PO}_{\boldsymbol{s}}},$$
(35)

where we define the reduced model space to be $S^- = \{s \in S, s_j = 0 : j \in \mathcal{V}'\}$, and the sets of models compatible with the null and alternative hypotheses in the sub-analysis to be $\mathcal{O}_{\mathcal{V}}^- \subset S^-$ and $\mathcal{A}_{\mathcal{V}}^- \subset S^-$. Then $\mathrm{PO}_{\mathcal{A}_{\mathcal{V}}^-:\mathcal{O}_{\mathcal{V}}^-}$ is a lower bound on the posterior odds against the intersection null hypothesis $\mathcal{V} \cup \mathcal{V}'$ because

$$\mathrm{PO}_{\mathcal{A}_{\mathcal{V}}^{-}:\mathcal{O}_{\mathcal{V}}^{-}} < \mathrm{PO}_{\mathcal{A}_{\mathcal{V}\cup\mathcal{V}}':\mathcal{O}_{\mathcal{V}\cup\mathcal{V}'}}.$$
(36)

The rejection of a null hypothesis \mathcal{V} in the sub-analysis therefore implies rejection of the null hypothesis $\mathcal{V} \cup \mathcal{V}'$ in the full analysis.

Proof.
$$\mathcal{A}_{\mathcal{V}}^{-} \subset \mathcal{A}_{\mathcal{V}\cup\mathcal{V}'}$$
 and $\mathcal{O}_{\mathcal{V}}^{-} = \mathcal{O}_{\mathcal{V}\cup\mathcal{V}'}$.

We analysed the 15 biomarkers with Doublethink, assuming a prior odds of variable inclusion of $\mu = 0.1$ and a prior precision parameter h = 1. Table 2 reports the five most significant associations between biomarkers and AMD risk. The top two variables were the same as Zuber et al. [2020] found, XL.HDL.C and L.HDL.C, with posterior odds of 1.75 and 0.58, corresponding to Bayes factors of 17 and 5.8, and unadjusted asymptotic *p*-values (by Corollary 2) of 0.001 and 0.004.

However, these tests should be properly interpreted as intersection null hypotheses with the 34 excluded variables, according to Theorem 5. This implies Bayes factors of only 0.045 and 0.015. After adjusting for $\nu = 49$ biomarkers (by Corollary 3 or Bonferroni correction), neither was significant, even at $\alpha = 0.05$. This begs the question, is there evidence that any of the biomarkers had non-zero causal effects on AMD risk?

In fact there was strong evidence that at least one of the biomarkers directly affected AMD risk. The posterior odds against the null hypothesis that none of the 15 biomarkers affected AMD risk were 3688, corresponding to a Bayes factor of 956, and an unadjusted asymptotic *p*-value (by Corollary 2) of 6.0×10^{-6} . After sub-analysis adjustment (Theorem 5), the Bayes factor was 21, and after adjusting for $\nu = 49$ variables, the asymptotic *p*-value (by Corollary 3) was 2.0×10^{-5} .

The smallest group of significant variables comprised XL.HDL.C and L.HDL.C. The posterior odds against the null hypothesis that neither affected AMD risk was 89, corresponding to a Bayes factor of 381 and an unadjusted asymptotic *p*-value of 3.4×10^{-5} . After sub-analysis adjustment (for 36 variables; Theorem 5), the Bayes factor was 2.1, and the asymptotic *p*-value, adjusted for $\nu = 49$ variables, was 1.0×10^{-3} . This shows that the combined test was able to establish significance (in a Bayesian sense at an FDR < 0.012 and in a frequentist sense at an asymptotic FWER < 0.0011), whereas the individual tests were not.

Not only is power increased by testing an intersection null hypothesis that no members of a group have an effect, but consideration of FWER control in the presence of correlation suggests that highly correlated variables should always be treated as a group, as the next section considers.

Note that the Doublethink combined test (Corollary 3) was significant when *p*-value-based combined tests were not. We applied four methods that control the FWER: Bonferroni [1936] correction, two multilevel procedures (Hommel [1988] implemented by Meijer et al. [2019], and Wilson [2019b]) based on Simes' [1986] test, and the harmonic mean *p*-value (HMP) procedure [Wilson, 2019a]. We also applied the Benjamini and Hochberg [1995] procedure, which controls a frequentist FDR. We furnished these procedures with unadjusted *p*-values from Corollary 2, which test the null hypothesis of no effect for each of the 15 variables individually (or rather, the intersection of each with the 34 excluded variables, as per Theorem 5). None of these tests rejected the null hypothesis that neither XL.HDL.C nor L.HDL (nor the 34 excluded variables) influence AMD risk, at $\alpha = 0.05$. None of them even rejected the grand null hypothesis that none of the biomarkers affected AMD risk. The much-reduced significance among these *p*-value-based combination tests compared to Doublethink likely reflects the information bottleneck when reducing the data to individual *p*-values before combining them.

7. INFLATION BETWEEN HIGHLY CORRELATED VARIABLES: SIMULATION APPROACH

In human GWAS, genetic variants that are physically linked tend to be correlated, so that signals of association often manifest as clusters of significant genetic variants. When several linked genetic variants are significant, the assumption is that most are not causally associated. So we reject the intersection hypothesis that none of the clustered variants are associated, but we do not reject all the elementary hypotheses, as a straightforward interpretation of statistical significance would suggest, because that would probably produce many false positives among the clustered variants. In this section we formalize the idea and show through simulation that the way statistical significance is interpreted has a direct bearing on FWER control.

Definition 9 (Frequentist control of FWER_{ρ}). In the regression problem (Definition 3), where the binary vector $s \in S = 2^{\nu}$ indexes the null hypotheses such that $\{\beta_j = 0 \forall s_j = 0\}$, where β_j is the coefficient for variable $x_{\cdot j}$, define the correlation coefficient between variables j and k to be $\rho_{j,k}$. Define the maximum absolute correlation between variables whose regression coefficients are constrained versus unconstrained under null hypothesis s to be

$$\rho_{\boldsymbol{s}}^{\max} = \max_{\boldsymbol{j},\boldsymbol{k}:s_{\boldsymbol{j}}\neq s_{\boldsymbol{k}}} |\rho_{\boldsymbol{j},\boldsymbol{k}}|,\tag{37}$$

with $\rho_{\mathbf{0}}^{\max} = \rho_{\mathbf{1}}^{\max} = 0$. The idea is group the most correlated variables indivisibly, in order to exclude null hypotheses with the highest ρ_{s}^{\max} from the set of all tested hypotheses, W, because they are most susceptible to inflation. We define $W_{\rho} = \{\omega_{s} : s \in S \setminus \mathbf{1}, \rho_{s}^{\max} \leq \rho\}$. By definition, W_{ρ} remains closed under intersection. We define the intersection of true null hypotheses in W_{ρ} to be $\omega_{\tilde{s}}$ and we define FWER_{ρ} to be

$$FWER_{\rho} := \sup_{\theta \in \omega_{\tilde{s}}} \Pr\left(\max_{\omega_{s} \subseteq \omega_{\tilde{s}}} \phi_{s}(\boldsymbol{y}) = 1; \boldsymbol{x}, \theta\right) \leq \alpha_{\tilde{s}}$$
$$\leq \max\left\{\alpha_{s} : \omega_{s} \in W_{\rho}\right\}.$$
(38)

This implies the regular FWER (Definition 1) equals FWER₁. Since FWER_{ρ} \leq FWER₁, it is a less stringent error rate to control, and control of FWER₁ implies control of FWER_{ρ}. In particular, notice that it limits the maximum absolute correlation between variables that are constrained versus

unconstrained in the intersection of true null hypotheses. It also reduces the size of the intersection of true null hypotheses, $|\tilde{s}|$.

Definition 10 (Asymptotically equivalent test statistic). The proof of Theorem 2 implies that the model-averaged posterior odds against the null hypothesis $\{\theta : \beta_j = 0 \ \forall j \in \mathcal{V}\}$, $\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}$, are asymptotically equivalent to the posterior odds of alternatives differing from the true model $\tilde{s}, \theta \in \Theta_{\tilde{s}}$ by one degree-of-freedom:

$$\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \to \sum_{j\in\mathcal{V}} \operatorname{PO}_{\tilde{\boldsymbol{s}}+\boldsymbol{e}_{j}:\tilde{\boldsymbol{s}}} =: \tilde{\operatorname{PO}}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}$$
(39)

as $n \to \infty$, assuming that $\tilde{s} \in \mathcal{O}_{\mathcal{V}}$, where $\{e_j\}_k = I(j = k)$. This allows us to partition the error of Doublethink into (i) the approximation to the distribution of $\tilde{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}$ and (ii) the rate of convergence to the asymptotically equivalent test statistic.

Remark 1 (The $\alpha \leq 0.025$ threshold in Doublethink). Simulations based on the asymptotically equivalent test statistic (Definition 10) indicate that the FWER should be set no larger than $\alpha = 0.025$ in Doublethink; equivalently, *p*-values that exceed 0.025 should be reported as not significant, or 1. This is because the chi-squared approximation (Theorem 2; Corollary 1) can be anti-conservative for $\alpha \geq 0.0259846$, for some values of $|\mathcal{V}|$, even under independence, even when applied to the asymptotically equivalent test statistic. See SI Section 4 and SI Figure 9.

Definition 11 (Bayes FWER). For the purpose of simulation it is convenient to define the FWER as an expectation over a (possibly degenerate) distribution of parameter values, rather than in a minimax sense (Definition 1). We define the Bayes FWER in a Bayes risk sense (see, e.g. Berger [1985]) as

BFWER :=
$$\int_{\Theta} \Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}_{0}(\theta)}:\mathcal{O}_{\mathcal{V}_{0}(\theta)}} \geq \tau; \boldsymbol{x}, \theta\right) \mathrm{d}\Pi(\theta), \tag{40}$$

where the notation $\mathcal{V}_0(\theta)$ emphasizes that the intersection of the true null hypotheses depends on the parameter θ . By analogy, we can define BFWER_{ρ} with reference to Definition 9, and the asymptotically equivalent Bayes FWER,

AFWER :=
$$\int_{\Theta} \Pr\left(\tilde{\mathrm{PO}}_{\mathcal{A}_{\mathcal{V}_{0}(\theta)}:\mathcal{O}_{\mathcal{V}_{0}(\theta)}} \geq \tau; \boldsymbol{x}, \theta\right) \mathrm{d}\Pi(\theta),$$
(41)

with the caveat that the asymptotically equivalent test (Definition 10), does not actually define a CTP, and therefore AFWER is not a familywise error rate, for finite n.

Definition 12 (Worst-case Bayes FWER in Doublethink). The *e*-value approach of Vovk and Wang [2021] yields a worst-case inflation for BFWER in Doublethink when Π is the 'true' distribution of parameter values, which provides a benchmark for simulation performance.

BFWER =
$$\Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}_{0}(\theta)}:\mathcal{O}_{\mathcal{V}_{0}(\theta)}} \geq \tau; \boldsymbol{x}, \Pi\right)$$

 $\leq \mathbb{E}\left[\operatorname{PO}_{\mathcal{A}_{\mathcal{V}_{0}(\theta)}:\mathcal{O}_{\mathcal{V}_{0}(\theta)}}; \boldsymbol{x}, \Pi\right] / \tau$
 $= \mathbb{E}\left[\mu^{|\mathcal{V}_{0}(\theta)|}; \Pi\right] / \tau$
 $= \left[\left(1 + \frac{\mu}{1 + \mu}\right)^{\nu} - 1\right] / \tau \sim \frac{\nu \mu}{\tau} \text{ when } \mu \ll 1.$ (42)

To investigate inflation in Doublethink in a real data setting, we performed simulations based on the AMD Mendelian randomization data [Zuber et al., 2020]. We simulated β for $\nu = 15$ variables from the Doublethink prior, assuming $\mu = 0.1$ and h = 1, fixing $\gamma = 1$, and selecting the variables as before. To assess performance, we calculated BFWER and AFWER (Definition 11).

Figure 2 shows that when n = 145 (left panel), BFWER_{ρ} (black points) exhibited circa five-fold inflation at $\rho = 1$, less than the 87-fold worst case inflation determined by the *e*-value approach (Definition 12). The inflation had subsided by $\rho = 0.5$, there was slight deflation around $\rho = 0.3$, and 1.6-fold inflation at $\rho = 0$. This pattern can be understood by comparing to the asymptotically equivalent error rate, AFWER_{ρ} (grey points). Near $\rho = 1$, BFWER_{ρ} far exceeded AFWER_{ρ}, indicating that the




FIGURE 2. Inflation in simulations based on the AMD Mendelian randomization example with $\nu = 15$ variables and n = 145 (left) or n = 14 500 (right). BFWER as a function of ρ (black points), versus AFWER (grey points) and Theorem 3 (green line). Assuming $\mu = 0.1$, h = 1, $\tau = 9$. Error bars based on 50 000 (left) and 500 000 (right) simulations.

model-averaged posterior odds were far from converging to the asymptotically equivalent test statistic. This reflects the difficulty in reliably attributing the causal effect between pairs of highly correlated variables. Near $\rho = 0.5$, BFWER_{ρ} had converged to AFWER_{ρ}. AFWER_{ρ} was slightly conservative for intermediate values of ρ , by the conservative nature of the CTP. Near $\rho = 0$ and $\rho = 1$, there was inflation in AFWER_{ρ} caused by small-sample error in the distribution of the asymptotically equivalent test statistic.

When n = 14500 (Figure 2, right panel), the pattern was similar but worst-case deflation was reduced (3.2-fold) and BFWER_{ρ} (black points) converged to AFWER_{ρ} (grey points) at larger values of ρ , reflecting the improved ability to correctly identify the true model with larger sample size. Theorem 3 (green line) was no longer an anti-conservative approximation to AFWER_{ρ} at any value of ρ , reflecting better performance of the asymptotic approximation at a larger sample size.

With either sample size, BFWER_{ρ} was less inflated for smaller ρ because the set of true null hypotheses was less likely to split groups of highly correlated variables. For $\rho = 1$, the set of true null hypotheses $(\mathcal{V}_0 = \{j : \beta_j = 0\})$ often included variables that were highly correlated with other variables outside the set. This caused inflation of the posterior odds. When $\rho = 0.5$, elementary hypotheses concerning highly correlated variables were removed from W, the set of null hypotheses under investigation. Although W still contained groups of the same hypotheses, these groups were indivisible. Therefore the set of true null hypotheses (which can be smaller than under $\rho = 1$, but not larger) no longer split highly correlated variables, so inflated variables were not left out of the set of true null hypotheses. When $\rho = 0$, there was a single group of all variables, the null hypothesis which corresponds to the grand null. Controlling FWER₀ is therefore equivalent to a single combined test with weak-sense control.

These results show that not only it is beneficial, in terms of power, to group correlated variables, but also advisable in terms of controlling the FWER. This formalizes existing practice in terms of interpreting statistically significant signals, as per the GWAS example. Interestingly, decisions concerning choice of ρ for FWER_{ρ} can be deferred until post-analysis, in the same way that decisions concerning choice of τ for the FDR and α for the FWER can be deferred. This means the analysis can be executed across all elementary hypotheses, and the level ρ in FWER_{ρ} control determined when the results come to be interpreted. Comparing these results to the sub-analysis of the AMD Mendelian randomization data in the previous section, it suggests that any attempt to attribute causal effects within the three pairs of highly correlated variables (XL.HDL.C and L.HDL.C, ApoB and IDL.TG, S.HDL.TG and S.VLDL.TG; $\rho > 0.8$) would be prone to inflated familywise error, although the Bayesian FDR would still be properly controlled, contingent on the prior. Therefore, not only do we conclude that XL.HDL.C and L.HDL.C form a significant group in the sub-analysis, but the frequentist would exercise great caution in attempting to attribute the signal to XL.HDL.C versus L.HDL.C.

8. Comparison to related approaches: simulations

We used simulations to compare the performance of Doublethink to related approaches: LASSO [Tibshirani, 1996], elastic net [Zou and Hastie, 2005], ridge regression [Hoerl and Kennard, 1970], backward elimination [Venables and Ripley, 2002] using the BIC [Schwarz, 1978], and MR-BMA [Zuber et al., 2020]. To these we added as benchmarks the grand null model, the grand alternative model, and the 'oracle' or true model [Fan and Li, 2001], as if it were known.

To compare methods for hypothesis testing, we deployed some approaches (LASSO, elastic net, MR-BMA and Doublethink) both natively, and as the first step in a model selection procedure. In the model selection version (taking the maximum *a posteriori* model for the Bayesian methods), we fitted the data without regularization and performed ν leave-one-out or add-one-in hypothesis tests. To investigate the effect of prior assumptions in Doublethink, we fitted the model at combinations of $\mu = 0.05$, 0.1 and 0.2, and h = 0.25, 1 and 4.

For testing intersection null hypotheses, which only Doublethink supported natively (Corollary 3), we employed p-value combination approaches: the Bonferroni [1936] procedure, a multilevel Simes' procedure [Wilson, 2019b], Hommel's [1988] procedure, which is also based on Simes' [1986] test, implemented by [Meijer et al., 2019], and the HMP procedure [Wilson, 2019a]. For comparison, we applied the same p-value combination procedures to Doublethink's unadjusted p-values (Corollary 2), substituting any p-values above 0.02 with 1 (as per Remark 1).

Simulations were based on the AMD Mendelian randomization data [Zuber et al., 2020]. We simulated β for $\nu = 15$ variables from the Doublethink prior, assuming $\mu = 0.1$ and h = 1, fixing $\gamma = 1$, and selecting 15 of the 49 biomarker variables as before. Again, we computed the Bayes FWER (Definition 11) as an average over the simulations.

We compared the following metrics: computation time, out-of-sample prediction error (measured as the root mean squared error, or L2 norm), estimator error (L2 norm), coverage of the standard error, Bayes FWER for type I (false positive) errors, and rates of strikeout (wrongly calling every test) for type II (false negative) errors. We calculated the FWER overall and for tests of individual variables only (marginal), pairs of variables only (pairwise), and the global test that all variables are null (headline). The target FWER was $\alpha = 0.01$. In total we performed 10 000 simulations.

Computation time. On average, most methods took under 0.01 seconds to run in R [R Core Team, 2024], the exceptions being LASSO (0.08s), elastic net (0.86s), MR-BMA (5.5s) and Doublethink (49s). The 5.5s run-time of MR-BMA, which is efficiently implemented, reflects its exhaustive search of all 32 768 models. The 49s run-time of Doublethink reflects its exhaustive search, native support for testing intersection null hypotheses, and lack of optimization.

Prediction error We computed out-of-sample predictive error (root mean squared error, i.e. L2 norm) by simulating pairs of datasets for each simulated parameter vector. The first was used for training the model, the second for testing prediction error. Most methods achieved an error below 1.05, the exceptions being ridge regression (1.16) and the grand null model (3.4) (Figure 3). The latter reflects the worst case because no coefficients were fit.

Estimator error. Methods differed substantially in their estimator error (Figure 3, LEFT). The oracle




FIGURE 3. Out-of-sample prediction error in 10 000 simulations from the Doublethink prior with $\mu = 0.01$ and h = 1. The key applies to all figures in this section.




FIGURE 4. Estimator error (LEFT) and standard error coverage (RIGHT) in 10 000 simulations from the Doublethink prior with $\mu = 0.01$ and h = 1. Expected coverage (black line) is shown, with allowance for Monte Carlo error (grey lines; 95% confidence interval).

model was close to optimal (root mean squared error 0.020). Doublethink's Bayesian model-averaged estimates and model selection-based estimates performed closely (below 0.033), for all assumed values of μ and h. Surprisingly, the model selection-based estimates out-performed the native estimates for LASSO and MR-BMA (error below 0.050). Backward elimination (0.067) and elastic net (0.100) followed. Poor estimates were obtained under the grand alternative model (0.257) and ridge regression (0.407), presumably due to over-fitting caused by lack of sparsity. The grand null model, in which all parameters were returned as zero, under-fitted, returning the worst case estimator error (0.654).

Standard error coverage. We defined coverage to be the proportion of simulations in which a 99% confidence interval, constructed from the standard error assuming a Normal sampling distribution, included the true parameter value (Figure 4, RIGHT). The oracle model was conservative (99.5% coverage), with Doublethink model averaging the best calibrated of the methods (98.7–99.3% coverage), followed by Doublethink model selection (97.8%–98.3%). LASSO, MR-BMA and elastic net model

selection were less well calibrated (97.3%, 96.8% and 96.3% respectively), while MR-BMA model averaging performed similarly (97.2%). Three methods produced anti-conservative standard errors: the grand alternative model (94.9%), backward elimination (93.2%) and the grand null model (90.9%) coverage).

Marginal Bayes FWER. To compare the false positive (type I error) rates across methods, we began by focusing only on marginal tests for the significance of individual variables. Figure 5, LEFT, circles, shows that again the oracle method was close to optimal in terms of marginal type I Bayes FWER, nearing the target of $\alpha = 0.01$ by achieving 0.0091. The grand alternative model was slightly anticonservative (0.015). Doublethink model averaging showed a range of errors, from the conservative, 0.002 (when assuming $\mu = 0.2$, h = 4), to the anti-conservative, 0.015 (assuming $\mu = 0.05$, h = 0.25). Doublethink, LASSO and elastic net model selection produced anti-conservative errors ranging from 0.021–0.0026, followed by MR-BMA model selection (0.038), backward elimination (0.081) and the grand null model (0.664), which was strongly anti-conservative, as expected.

For every method capable of producing *p*-values, we were able to apply *p*-value combination methods (Bonferroni, Simes, Hommel and HMP procedures). Results were usually similar across procedures, so Figure 5, LEFT shows these as vertical lines of differing heights that often overlap. For Doublethink model averaging, they combine the unadjusted *p*-values from Corollary 2, instead of the adjusted *p*-values from Corollary 3. They differed from the latter in the anti-conservative direction. For all other methods, the main result utilized Bonferroni correction, so these results coincided, and the other combination tests differed very little.

Marginal strikeout rates. To compare power, we defined the type II strikeout rate as the probability of failing to detect all true signals, which corresponds, for an individual dataset, to 100% false negative rate. We restrict attention to methods that did not exhibit appreciably anti-conservative BFWER. For marginal tests of the significance of individual variables, the oracle model showed a strikeout rate of 0.045, providing a reference point for other methods (Figure 5, RIGHT). The grand alternative model exhibited the worst strikeout rate of 0.129, demonstrating that it is slightly anti-conservative and strongly under-powered, an undesirable combination. Doublethink model-averaging exhibited strikeout rates of 0.052-0.070. This range of performance across different prior assumptions reflected the corresponding degree of conservatism of the tests (Figure 5, LEFT).

Pairwise Bayes FWER. Pairwise tests of the intersection null hypothesis that neither of a pair of variables has an effect are expected to be more powerful than marginal tests alone, but the additional multiple testing burden increases the risk of false positives. Figure 6, LEFT shows the BFWER for pairwise tests. Only Doublethink model averaging performed these tests natively (circles). For all other methods, including Doublethink model selection, combined tests were performed using the *p*-value combination methods (vertical bars). The pairwise BFWER was barely higher than the marginal BFWER for these combination tests, which is unsurprising since (i) no new information is introduced and (ii) the closed testing procedures on which they are based are designed to control the FWER. For Doublethink model averaging, the pairwise BFWERs were all higher than the marginal BFWERs (Figure 5, LEFT), ranging from the conservative, 0.005, to the anti-conservative, 0.021. This reflects the method's exploitation of the full data for pairwise tests, which is potentially more powerful, and the limitation of the asymptotic approximation for controlling the FWER.

Pairwise strikeout rates. The greater power of the Doublethink approach to pairwise testing can be seen in Figure 6, RIGHT. Pairwise strikeout rates were lower than marginal strikeout rates (Figure 5, RIGHT), ranging from 0.045–0.056, and rivalling the oracle strikeout rate of 0.045 (albeit for the anti-conservative prior $\mu = 0.05$, h = 0.25). The pairwise strikeout rate for the grand alternative model was nearly unchanged at 0.127.

Headline Bayes FWER. The headline FWER (Figure 7, LEFT) is calculated assuming the grand null hypothesis is true. As before, the oracle model was well calibrated (0.010–0.013) which coincided, in this case, with the grand null hypothesis. Methods that were not generally well calibrated (LASSO, elastic net, MR-BMA and Doublethink model selection) showed headline BFWER in the range 0.010–0.016.






FIGURE 5. Type I Bayes FWER (LEFT) and type II strikeout rate (RIGHT) for marginal tests of the significance of individual variables in 10 000 simulations from the Doublethink prior with $\mu = 0.01$ and h = 1. Expected type I BFWER (black line) is shown, with allowance for Monte Carlo error (grey lines; 95% confidence interval).








FIGURE 7. Type I Bayes FWER (LEFT) and type II strikeout rate (RIGHT) for headline tests.




FIGURE 8. Type I Bayes FWER for $\rho = 1$ (LEFT) and $\rho = 0.43$ (RIGHT) for all tests of the significance of any group of variables in 10 000 simulations from the Doublethink prior with $\mu = 0.01$ and h = 1. Expected type I BFWER (black line) is shown, with allowance for Monte Carlo error (grey lines; 95% confidence interval).

Backward elimination remained anti-conservative (0.067–0.084), as did the grand alternative model (0.018–0.019). Combination tests based on Doublethink Corollary 2 were conservative (0.002–0.009), but under-powered. Doublethink Corollary 3 was anti-conservative under the grand null hypothesis (0.011–0.029). While undesirable, earlier simulations suggested this source of inflation should abate for larger sample sizes (Figure 2, LEFT versus RIGHT).

Headline strikeout rates. In a closed testing procedure, when there are true signals to be found, the headline or 'global' test is the most powerful. The headline strikeout rates are calculated assuming the grand null hypothesis is false. The oracle model again provided a benchmark, 0.045. The grand alternative remained under-powered at 0.127. Headline strikeout error rates for Doublethink model averaging outperformed the oracle model, ranging from 0.037–0.043.

Overall Bayes FWER and FWER_{ρ}. We calculated the overall Bayes FWER and the corresponding Bayes FWER_{ρ}, in which variables with absolute correlation coefficients exceeding ρ are grouped together indivisibly, so that sub-groups cannot be tested (Definition 9). Figure 8 shows that BFWER_{ρ} was inflated for Doublethink model averaging at $\rho = 1$ (LEFT), but this was mitigated at $\rho = 0.43$ (0.010–0.017; RIGHT). BFWER_{0.43} \leq BFWER₁ for all methods, but some remained inflated, including backward elimination (0.061–0.071) and the grand null model (0.644–0.650). The oracle model was well-calibrated at both BFWER₁ and BFWER_{0.43} (0.008–0.011).

Importantly, the headline strikeout rate (Figure 7, RIGHT) prevails irrespective of choice of ρ , so power can be upheld while inflation in the FWER is mitigated by defining indivisible groups of null hypotheses. This allowed Doublethink performance to approach that of the oracle model.

9. DISCUSSION

Often, the goal of hypothesis testing is to identify a few influential variables among many candidates, reflecting the scientific principal of explaining complex observed patterns by simple unobserved processes. Recently, statistical inference has lost ground to artificial intelligence prediction engines. But for accurately estimating parameters, quantifying their uncertainty, and controlling false positives, model choice is vital, as the simulations showed. Bayesian model averaging is especially useful when the model is uncertain, but for decades frequentists and Bayesians have debated hypothesis testing,

seemingly irreconcilably (e.g. Edwards et al. [1963], Berger and Sellke [1987], Bayarri and Berger [2004], Gelman [2008]).

Perhaps it is surprising, therefore, that Bayesian hypothesis testing can be formulated as a closed testing procedure [Marcus et al., 1976] that simultaneously controls the Bayesian FDR and the strong-sense frequentist FWER (Theorem 1). Particularly because FWER, in contrast to FDR, is often characterized as "an extremely strict criterion which is not always appropriate" [Storey, 2003]. Theorem 1 is general, independent of prior or modelling assumptions, complementing known results like the frequentist admissibility of Bayesian procedures [Wald, 1950, Berger, 1985]. It addresses both the broad question of when Bayesian and frequentist philosophies converge, and the practical question of how to choose significance thresholds. Whereas it will usually be convenient to pre-determine one or the other, reporting both the FWER and FDR offers an opportunity to bridge Bayesian and frequentist approaches to hypothesis testing

In big data regression settings, Theorems 2 and 3 extend work by Johnson [2005, 2008] to interconvert FPR, FWER and FDR thresholds. Corollaries 2 and 3 interconvert the unadjusted and adjusted p-values and posterior odds. These results are less general than Theorem 1, comprising large-sample limits that require assumptions including an appropriate prior and independent observations. Convergence is pointwise, rather than uniform in general, subjecting their use to finite-sample inflation. Yet they afford insights into the relationship between the frequentist approach to multiple testing and Bayesian prior assumptions. Corollary 4 shows that the FWER threshold is asymptotically proportional to (i) the number of variables, ν ; (ii) the prior odds, μ ; and (iii) the square root of the prior precision h; and asymptotically inversely proportional to (iv) the Bayesian threshold, τ ; and (v) the square root of the sample size, n.

The FWER and FDR bounds – which are not necessarily tight – are approximately inversely proportional. For a Bayesian, ν , μ and h are components of the prior, whose effects must be counter-balanced if the FWER is to be held constant: in a Bayesian sense, Bonferroni correction modifies the prior odds μ to hold the expected number of variables $\nu \mu/(1 + \mu)$ constant while ν increases. Corollary 4 recapitulates the Jeffreys-Lindley paradox [Jeffreys, 1939, Lindley, 1957]: Bayesian inference is consistent for a fixed FDR threshold, τ , [Doob, 1949], whereas the practice of fixing the FWER threshold, α , irrespective of sample size, n (e.g. at 0.05 or 0.005 [Benjamin et al., 2018]), is not consistent because there exists a tangible probability of falsely rejecting a null hypothesis no matter how large the sample [O'Hagan, 1995]. This is solved by inversely scaling α with \sqrt{n} , originally advocated by Jeffreys [Wagenmakers and Ly, 2023], from which point one could report both the FWER and FDR bound.

Interconverting model-averaged posterior odds and adjusted p-values using Doublethink (Corollary 3) has several limitations that stem from the underlying assumptions, including (i) a large sample size, which implies that a Normal approximation to the likelihood is reasonable locally; (ii) data that comprise independent realizations of a process satisfying technical conditions which imply locally unique, consistent and asymptotically Normal maximum likelihood estimators [Cox and Hinkley, 1974]; (iii) a specific family of conjugate priors parameterized by μ and h, and (iv) local alternatives, which implies that the Normal approximation holds in the vicinity of null and alternative hypotheses. The local alternatives assumption is convenient for deriving an asymptotic distribution for the maximized likelihood ratio [Cox and Hinkley, 1974, Johnson, 2005, 2008], yet it implies that non-zero parameter values scale with \sqrt{n} , contrary to consistency and our prior. The conjugate prior is convenient for pursuing joint Bayesian-frequentist inference (e.g. Zellner [1986], Liang et al. [2008]), but its covariance structure, based on the Fisher information, is otherwise hard to justify. The independent data assumption restricts the generality of the approach. It is not recommended to apply Doublethink with $\alpha > 0.025$; p-values larger than 0.025 should be interpreted as 1 (Remark 1), but this is hardly limiting if α is to scale inversely with \sqrt{n} .

Of principal concern, the large sample assumption means the asymptotic results may be inaccurate for finite samples, as evidenced by the simulations. Two sources of inflation were found: (i) the contribution of higher order tests involving more than one-degree-of-freedom to the posterior odds, which vanishes asymptotically, and (ii) lack of convergence of the posterior odds to the asymptotically equivalent test statistic (Definition 10). The latter is caused by correlation between variables that have zero and non-zero effects. The simplified two-variable model (Definition 8) attributed this inflation to specific parameter values that scale inversely with \sqrt{n} , and therefore threaten inflation regardless of n. In simulations, all methods were affected by such inflation (Figure 8), which invalidates the fundamental assumption that test statistics are well calibrated under the null. To address it, we formalized the practice of making inferences about indivisible groups of correlated variables, rather than the individual variables concerned, through the notion of FWER_{ρ} (Definition 9). In simulations, this mitigated, but did not eradicate, inflation at $\rho = 0.43$.

Hypothesis testing, particularly multiple testing, represents a long difficult-to-reconcile conflict between Bayesian and frequentist approaches [Bayarri and Berger, 2004], unlike parameter estimation (e.g. Doob [1949], Borwanker et al. [1971]). Doublethink is connected to other approaches that bridge this gap, notably *e*-values [Vovk and Wang, 2021]. Application of the *e*-value approach to the Doublethink model (Definition 7) allows interconversion of the Bayesian FDR and the Bayes FWER, assuming the prior is correctly specified. The *e*-value bound on the Bayes FWER (Definition 12) is then approximately proportional to (i) the number of variables, ν ; and (ii) the prior odds, μ ; but not (iii) the square root of the prior precision h; and inversely proportional to (iv) the Bayesian threshold, τ ; but not (v) the square root of the sample size, n. Therefore the *e*-value bound on the Bayes FWER is conservative, in the large-sample limit, by a factor of approximately $\sqrt{h/n}$, but conversely, the Doublethink threshold on the Bayes FWER could be anti-conservative, in a finite sample, by the corresponding amount.

Another related approach, which extends FWER, is the false discovery proportion (FDP) approach of Goeman and Solari [2011], Goeman et al. [2019]. The FDP approach comprises a 'multilevel' closed testing procedure [Meinshausen, 2008] in which all possible combinations of hypotheses are simultaneously controlled via pre-determined thresholds. This means arbitrary combinations of hypotheses can be tested and significant groups of variables identified freely identified post-hoc. The approach is inherent to Bayesian control of the FDR and, by Theorem 1, the FWER. Multilevel testing challenges concepts like fishing for significance, data dredging, and *p*-hacking [Andrade, 2021].

The Bayesian FDR considered here motivated, but remains distinct from, frequentist FDR concepts. Doublethink controls both the 'local' and 'global' Bayesian FDR [Efron et al., 2001], rather than a frequentist FDR controlled by Benjamini and Hochberg's [1995] or Storey's [2003] procedures. One connection to Theorem 1 is that controlling the frequentist FDR controls the FWER at the same level, but only in the weak sense [Benjamini and Hochberg, 1995]. Whereas Corollary 4 indicates the FWER bound is $O(\sqrt{n})$ smaller than the Bayesian FDR bound.

Doublethink belongs to a class of combination tests that exploit heavy-tailed distributions, including the Cauchy combination test [Liu and Xie, 2020] and the HMP procedure (Wilson [2019a]), through its reliance on Karamata's theory of regular variation [Karamata, 1933]. The HMP offers model averaging, starting with *p*-values, whereas Doublethink pursues joint Bayesian-frequentist model-averaging, beginning with nested maximized likelihood ratios. A theoretical advance over the HMP is the ability of Doublethink to allow model uncertainty in the null hypothesis, as well as in the alternative hypothesis. Another is the clarification that the relevant limit is $n \to \infty$, rather than $\nu \to \infty$. By Corollary 1, the model-averaged deviance asymptotically follows a chi-squared distribution with one degree of freedom. This mirroring of the null distribution of the classical likelihood ratio test statistic [Wilks, 1938, Wald, 1943] emerges from the self-similarity or fractal property of sums of heavy-tailed random variables [Mandelbrot and Van Ness, 1968, Taqqu, 1978].

This paper focused on the frequentist properties of Bayesian model averaging. To the extent it defends p-values, it should be read only as a defence of LRT-based p-values. Equally, it could be read as an inquiry into the circumstances under which p-values are useful, despite their flaws. The ability to interconvert model-averaged p-values with sufficient statistics like the model-averaged deviance or posterior odds implies a reprieve from the central criticisms of p-values – their lack of evidentiary value and violation of the likelihood principal (see, e.g. Birnbaum [1962]). Simultaneous Bayesian-frequentist hypothesis testing addresses another major flaw of p-values – their indifference towards

power – because the prior determines the optimal performance envelope of the test. By characterizing the asymptotic properties of Bayesian tests under the null hypothesis, even when the prior is wrong, we have aimed to exploit the benefits of Bayesian model-averaged hypothesis testing while mitigating the subjective influence of the prior by controlling both the FWER and the FDR.

10. Acknowledgements

The authors wish to thank colleagues for their comments and advice on the paper, and the Mathematisches Forschungsinstitut Oberwolfach, organizers and participants of workshop 2308 Design and Analysis of Infectious Disease Studies. This work was funded by the Robertson Foundation, the Wellcome Trust, and the Royal Society (grant no. 101237/Z/13/B).

References

- C. Andrade. Harking, cherry-picking, p-hacking, fishing expeditions, and data dredging and mining as questionable research practices. *The Journal of clinical psychiatry*, 82(1):25941, 2021.
- M. J. Bayarri and J. O. Berger. The interplay of bayesian and frequentist analysis. *Statistical Science*, 19(1):58–80, 2004.
- D. J. Benjamin, J. O. Berger, M. Johannesson, B. A. Nosek, E.-J. Wagenmakers, R. Berk, K. A. Bollen, B. Brembs, L. Brown, C. Camerer, et al. Redefine statistical significance. *Nature Human Behaviour*, 2(1):6–10, 2018.
- Y. Benjamini and Y. Hochberg. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B*, 57(1):289–300, 1995.
- J. O. Berger. *Statistical Decision Theory and Bayesian Analysis*. Springer-Verlag, New York, second edition, 1985.
- J. O. Berger and T. Sellke. Testing a point null hypothesis: The irreconcilability of p values and evidence. *Journal of the American Statistical Association*, 82(397):112–122, 1987.
- A. Birnbaum. On the foundations of statistical inference. Journal of the American Statistical Association, 57(298):269–306, 1962.
- C. Bonferroni. Teoria statistica delle classi e calcolo delle probabilita. Pubblicazioni del R. istituto superiore di scienze economiche e commericiali di firenze, 8:3–62, 1936.
- J. Borwanker, G. Kallianpur, and B. P. Rao. The bernstein-von mises theorem for markov processes. *The Annals of Mathematical Statistics*, pages 1241–1253, 1971.
- D. Cox and D. Hinkley. Theoretical Statistics. Chapman and Hall, London, 1974.
- R. R. Davidson and W. E. Lever. The limiting distribution of the likelihood ratio statistic under a class of local alternatives. *Sankhyā: The Indian Journal of Statistics, Series A*, pages 209–224, 1970.
- R. A. Davis and S. I. Resnick. Limit theory for bilinear processes with heavy-tailed noise. The Annals of Applied Probability, 6(4):1191–1210, 1996.
- A. C. Davison. *Statistical models*, volume 11. Cambridge university press, 2003.
- J. L. Doob. Application of the theory of martingales. Le calcul des probabilites et ses applications, pages 23–27, 1949.
- W. Edwards, H. Lindman, and L. J. Savage. Bayesian statistical inference for psychological research. Psychological review, 70(3):193, 1963.
- B. Efron, R. Tibshirani, J. D. Storey, and V. Tusher. Empirical bayes analysis of a microarray experiment. *Journal of the American Statistical Association*, 96(456):1151–1160, 2001.
- J. Fan and R. Li. Variable selection via nonconcave penalized likelihood and its oracle properties. Journal of the American statistical Association, 96(456):1348–1360, 2001.
- R. A. Fisher. Statistical Methods for Research Workers. Oliver and Boyd, Edinburgh, 1925.
- L. G. Fritsche, W. Igl, J. N. C. Bailey, F. Grassmann, S. Sengupta, J. L. Bragg-Gresham, K. P. Burdon, S. J. Hebbring, C. Wen, M. Gorski, et al. A large genome-wide association study of age-related macular degeneration highlights contributions of rare and common variants. *Nature genetics*, 48(2): 134–143, 2016.
- A. Gelman. Objections to bayesian statistics. *Bayesian Analysis*, 3(3):445–449, 2008.

- A. Gelman and C. R. Shalizi. Philosophy and the practice of bayesian statistics. British Journal of Mathematical and Statistical Psychology, 66(1):8–38, 2013.
- J. J. Goeman and A. Solari. Multiple testing for exploratory research. *Statistical Science*, 26(4): 584–597, 2011.
- J. J. Goeman, R. J. Meijer, T. J. Krebs, and A. Solari. Simultaneous control of all false discovery proportions in large-scale multiple hypothesis testing. *Biometrika*, 106(4):841–856, 2019.
- I. J. Good. The bayes/non-bayes compromise: A brief review. Journal of the American Statistical Association, 87(419):597–606, 1992.
- L. Held and M. Ott. On p-values and Bayes factors. Annual Review of Statistics and Its Application, 5:393–419, 2018.
- A. E. Hoerl and R. W. Kennard. Ridge regression: Biased estimation for nonorthogonal problems. *Technometrics*, 12(1):55–67, 1970.
- G. Hommel. A stagewise rejective multiple test procedure based on a modified bonferroni test. *Biometrika*, 75(2):383–386, 1988.
- R. A. Horn and C. R. Johnson. Matrix analysis. Cambridge university press, 2012.
- J. Hu and V. E. Johnson. Bayesian model selection using test statistics. Journal of the Royal Statistical Society: Series B (Statistical Methodology), 71(1):143–158, 2009.
- H. Jeffreys. The Theory of Probability. Clarendon, Oxford, first edition, 1939.
- V. E. Johnson. Bayes factors based on test statistics. Journal of the Royal Statistical Society Series B: Statistical Methodology, 67(5):689–701, 2005.
- V. E. Johnson. Properties of bayes factors based on test statistics. Scandinavian Journal of statistics, 35(2):354–368, 2008.
- T. H. Julian, J. Cooper-Knock, S. MacGregor, H. Guo, T. Aslam, E. Sanderson, G. C. Black, and P. I. Sergouniotis. Phenome-wide mendelian randomisation analysis identifies causal factors for age-related macular degeneration. *Elife*, 12:e82546, 2023.
- J. Karamata. Sur un mode de croissance régulière. théorèmes fondamentaux. Bull. Soc. Math. France, 61:55–62, 1933.
- R. E. Kass and L. Wasserman. A reference Bayesian test for nested hypotheses and its relationship to the schwarz criterion. *Journal of the American Statistical Association*, 90(431):928–934, 1995.
- J. Kettunen, A. Demirkan, P. Würtz, H. H. Draisma, T. Haller, R. Rawal, A. Vaarhorst, A. J. Kangas, L.-P. Lyytikäinen, M. Pirinen, et al. Genome-wide study for circulating metabolites identifies 62 loci and reveals novel systemic effects of lpa. *Nature communications*, 7(1):11122, 2016.
- P. Kevei. Regularly varying functions. 2019.
- F. Liang, R. Paulo, G. Molina, M. A. Clyde, and J. O. Berger. Mixtures of g priors for Bayesian variable selection. *Journal of the American Statistical Association*, 103(481):410–423, 2008.
- D. V. Lindley. A statistical paradox. *Biometrika*, 44(1/2):187–192, 1957.
- Y. Liu and J. Xie. Cauchy combination test: a powerful test with analytic p-value calculation under arbitrary dependency structures. *Journal of the American Statistical Association*, 115(529):393–402, 2020.
- T. M. Loughin. A systematic comparison of methods for combining p-values from independent tests. Computational Statistics & Data Analysis, 47(3):467–485, 2004.
- B. B. Mandelbrot and J. W. Van Ness. Fractional brownian motions, fractional noises and applications. SIAM review, 10(4):422–437, 1968.
- R. Marcus, E. Peritz, and K. R. Gabriel. On closed testing procedures with special reference to ordered analysis of variance. *Biometrika*, 63(3):655–660, 1976.
- R. J. Meijer, T. J. Krebs, and J. J. Goeman. Hommel's procedure in linear time. *Biometrical Journal*, 61(1):73–82, 2019.
- N. Meinshausen. Hierarchical testing of variable importance. Biometrika, 95(2):265–278, 2008.
- T. Mikosch. *Regular variation, subexponentiality and their applications in probability theory*. Eindhoven University of Technology, Eindhoven, The Netherlands, 1999.
- A. Miller. Subset selection in regression. CRC Press, 2002.
- J. W. Miller. A detailed treatment of doob's theorem. arXiv preprint arXiv:1801.03122, 2018.
- J. Neyman and E. S. Pearson. Ix. on the problem of the most efficient tests of statistical hypotheses. Philosophical Transactions of the Royal Society of London. Series A, Containing Papers of a Mathematical or Physical Character, 231(694–706):289–337, 1933.

- 24 DOUBLETHINK: SIMULTANEOUS BAYESIAN-FREQUENTIST MODEL-AVERAGED HYPOTHESIS TESTING
- A. O'Hagan. Fractional Bayes factors for model comparison. Journal of the Royal Statistical Society: Series B (Methodological), 57(1):99–118, 1995.
- J. Pearl, M. Glymour, and N. P. Jewell. *Causal inference in statistics: A primer*. John Wiley & Sons, 2016.
- K. Pearson. X. on the criterion that a given system of deviations from the probable in the case of a correlated system of variables is such that it can be reasonably supposed to have arisen from random sampling. *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 50 (302):157–175, 1900.
- A. Porwal and A. E. Raftery. Comparing methods for statistical inference with model uncertainty. Proceedings of the National Academy of Sciences, 119(16):e2120737119, 2022.
- R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria, 2024. URL https://www.R-project.org/.
- A. E. Raftery. Bayesian model selection in social research. Sociological methodology, pages 111–163, 1995.
- E. Sanderson, G. Davey Smith, F. Windmeijer, and J. Bowden. An examination of multivariable mendelian randomization in the single-sample and two-sample summary data settings. *International journal of epidemiology*, 48(3):713–727, 2019.
- G. Schwarz. Estimating the dimension of a model. Annals of Statistics, 6(2):461–464, 1978.
- T. Sellke, M. J. Bayarri, and J. O. Berger. Calibration of p values for testing precise null hypotheses. The American Statistician, 55(1):62–71, 2001.
- R. J. Simes. An improved Bonferroni procedure for multiple tests of significance. *Biometrika*, 73(3): 751–754, 1986.
- J. D. Storey. The positive false discovery rate: a bayesian interpretation and the q-value. *The annals of statistics*, 31(6):2013–2035, 2003.
- A. Stuart, K. J. Ord, and S. Arnold. Kendall's Advanced Theory of Statistics Volume 2A: Classical Inference and the Linear Model. Hodder Education, London, 1998.
- M. S. Taqqu. A representation for self-similar processes. Stochastic Processes and their Applications, 7(1):55–64, 1978.
- R. Tibshirani. Regression shrinkage and selection via the lasso. Journal of the Royal Statistical Society Series B: Statistical Methodology, 58(1):267–288, 1996.
- J. W. Tukey. The problem of multiple comparisons. Multiple comparisons, 1953.
- A. W. Van der Vaart. Asymptotic statistics, volume 3. Cambridge university press, 2000.
- H. Van der Vaart and E. H. Yen. Weak sufficient conditions for fatou's lemma and lebesgue's dominated convergence theorem. *Mathematics Magazine*, 41(3):109–117, 1968.
- J. P. Vandenbroucke. The history of confounding. Sozial-und Präventivmedizin, 47:216–224, 2002.
- W. N. Venables and B. D. Ripley. Modern applied statistics with S. Springer, 2002.
- V. Vovk and R. Wang. E-values: Calibration, combination and applications. *The Annals of Statistics*, 49(3):1736–1754, 2021.
- E.-J. Wagenmakers and A. Ly. History and nature of the jeffreys–lindley paradox. Archive for History of Exact Sciences, 77(1):25–72, 2023.
- J. Wakefield. Bayes factors for genome-wide association studies: comparison with p-values. *Genetic Epidemiology*, 33(1):79–86, 2009.
- A. Wald. Tests of statistical hypotheses concerning several parameters when the number of observations is large. Transactions of the American Mathematical Society, 54(3):426–482, 1943.
- A. Wald. Statistical Decision Functions. Wiley, New York, 1950.
- S. S. Wilks. The large-sample distribution of the likelihood ratio for testing composite hypotheses. Annals of Mathematical Statistics, 9(1):60–62, 1938.
- D. J. Wilson. The harmonic mean p-value for combining dependent tests. Proceedings of the National Academy of Sciences, 116(4):1195–1200, 2019a.
- D. J. Wilson. Trade-offs in model averaging using multilevel tests. *Proceedings of the National Academy* of Sciences, 116(47):23384–23385, 2019b.
- A. Zellner. On assessing prior distributions and bayesian regression analysis with g-prior distributions. Bayesian inference and decision techniques, 1986.
- Q. Zhou and Y. Guan. On the null distribution of bayes factors in linear regression. Journal of the American Statistical Association, 113(523):1362–1371, 2018.

- Y. Zhou and Y. Xiao. Tail asymptotics for the extremes of bivariate Gaussian random fields. *Bernoulli*, 23(3):1566–1598, 2017.
- H. Zou and T. Hastie. Regularization and variable selection via the elastic net. Journal of the Royal Statistical Society Series B: Statistical Methodology, 67(2):301–320, 2005.
- V. Zuber, J. M. Colijn, C. Klaver, and S. Burgess. Selecting likely causal risk factors from high-throughput experiments using multivariable mendelian randomization. *Nature communications*, 11 (1):29, 2020.

Supplementary Information

1. Regularity conditions

The following regularity conditions, as listed previously by Davidson and Lever [1970] and Johnson and colleagues (Johnson [2008], Hu and Johnson [2009]), are assumed. A more detailed explanation of each condition is provided in Davidson and Lever [1970].

For almost all $y \in \mathcal{Y}$, $x \in \mathcal{X}$ and all $\theta \in \Theta$ and $j, k, l = 1, ..., \nu + \zeta$.

(a)
$$\frac{\partial \log p(y; \boldsymbol{x}, \theta)}{\partial \theta_j}$$
, $\frac{\partial^2 \log p(y; \boldsymbol{x}, \theta)}{\partial \theta_j \partial \theta_k}$ and $\frac{\partial^3 \log p(y; \boldsymbol{x}, \theta)}{\partial \theta_j \partial \theta_k \partial \theta_l}$ exist for

(b) There exists arbitrary functions, $F_j(y)$ and $F_{jk}(y)$, that are integrable over \mathcal{Y} , such that $\left|\frac{\partial \log p(y; \boldsymbol{x}, \theta)}{\partial \theta_j}\right| < F_j(y)$ and $\left|\frac{\partial^2 \log p(y; \boldsymbol{x}, \theta)}{\partial \theta_j \partial \theta_k}\right| < F_{jk}(y)$.

(c) The matrix $\mathcal{I}(\theta)$ with elements

$$\mathcal{I}(\theta)_{jk} = \mathbb{E}_{\theta} \left[\left(\frac{\partial \log p(y; \boldsymbol{x}, \theta)}{\partial \theta_j} \Big|_{\theta} \right) \left(\frac{\partial \log p(y; \boldsymbol{x}, \theta)}{\partial \theta_k} \Big|_{\theta} \right) \right]$$
(1)

is positive definite with a finite determinant.

(d)

$$\left| \frac{\partial^3 \log p(y; \boldsymbol{x}, \theta)}{\partial \theta_j \partial \theta_k \partial \theta_l} \right|_{\theta} \right| < H_{jkl}(y|\boldsymbol{x})$$
(2)

where there exists an M > 0 such that $\mathbb{E}_{\theta}[H_{jkl}(y; \boldsymbol{x})] < M < \infty$, and $\kappa, L > 0$, such that

$$E_{\theta}\left[|H_{jkl}(y;\boldsymbol{x}) - \mathbb{E}_{\theta}[H_{jkl}(y;\boldsymbol{x})]|^{1+\kappa}\right] < L < \infty$$
(3)

(e) There exists $\psi, T > 0$ such that, whenever $\parallel \theta^{**} - \theta^* \parallel \equiv \sum_{j=1}^{\nu+\zeta} \mid \theta_j^{**} - \theta_j^* \mid < \psi, \theta^{**}, \theta^* \in \Theta$

$$\mathbb{E}_{\theta} \left[\left\{ \frac{\partial^3 \log p(y; \boldsymbol{x}, \theta)}{\partial \theta_j \partial \theta_k \partial \theta_l} \Big|_{\theta} \right\}^2 \right] < T < \infty$$
(4)

(f) There exists $\eta, K > 0$, such that

$$\mathbb{E}_{\theta}\left[\left|\frac{\partial \log p(y; \boldsymbol{x}, \theta)}{\partial \theta_{j}}\right|_{\theta}\right|^{2+\eta}\right] < K < \infty$$
(5)

2. Background theory

2.1. Closed testing procedures control the familywise error rate in the strong sense. Marcus, Peritz, and Gabriel [1976] introduced the closed testing procedure (CTP) to control the frequentist familywise error rate (FWER) in the strong sense. Suppose random variable \boldsymbol{y} has a probability mass or density function $p(\boldsymbol{y}; \boldsymbol{x}, \theta)$ that depends on auxiliary data \boldsymbol{x} and parameters $\theta \in \Theta$. The aim is to test a set of hypotheses defined by $W = \{\omega_s\}$, where $\omega_s \subset \Theta$ and \boldsymbol{s} is an index. W must be closed under intersection, meaning $\omega_s, \omega_{s'} \in W$ implies $\omega_s \cap \omega_{s'} \in W$. A 'local' test controls the false positive rate (FPR) at level α_s by rejecting $\theta \in \omega_s$ when $\psi_s(\boldsymbol{y}) = 1$ (rather than 0) such that:

$$FPR_{\boldsymbol{s}} := \sup_{\boldsymbol{\theta} \in \boldsymbol{\omega}_{\boldsymbol{s}}} \Pr\left(\psi_{\boldsymbol{s}}(\boldsymbol{y}) = 1; \boldsymbol{x}, \boldsymbol{\theta}\right) = \alpha_{\boldsymbol{s}}.$$
(6)

A CTP rejects $\theta \in \omega_s$ when $\phi_s(y) = 1$, which indicates rejection of all intersection hypotheses in W:

$$\phi_{\boldsymbol{s}}(\boldsymbol{y}) = \min_{\omega_{\boldsymbol{r}} \subseteq \omega_{\boldsymbol{s}}} \psi_{\boldsymbol{r}}(\boldsymbol{y}). \tag{7}$$

By construction, rejection of the null hypothesis ω_s by the CTP implies rejection of any and all intersection hypotheses too:

$$\phi_{\boldsymbol{s}}(\boldsymbol{y}) = 1 \implies \phi_{\boldsymbol{r}}(\boldsymbol{y}) = 1 \qquad \forall \ \omega_{\boldsymbol{r}} \subseteq \omega_{\boldsymbol{s}}.$$
(8)

Sometimes this is an inherent property of the local test, in which case we have a procedure

$$\phi_{\boldsymbol{s}}(\boldsymbol{y}) = \psi_{\boldsymbol{s}}(\boldsymbol{y}) \tag{9}$$

known as a shortcut CTP. Shortcut procedures are more computationally efficient.

The purpose of the CTP is to control the FWER at a rate determined by the local tests. The FPR of test ϕ is related to the FPR of local test ψ by

$$\sup_{\theta \in \omega_{s}} \Pr\left(\phi_{s}(\boldsymbol{y}) = 1; \theta\right) = \sup_{\theta \in \omega_{s}} \Pr\left(\bigcap_{\omega_{r} \subseteq \omega_{s}} \left\{\psi_{r}(\boldsymbol{y}) = 1\right\}; \theta\right)$$
$$\leq \sup_{\theta \in \omega_{s}} \Pr\left(\psi_{s}(\boldsymbol{y}) = 1; \theta\right) = \alpha_{s}.$$
(10)

Let $\omega_{\tilde{s}}$ denote the intersection of all true null hypotheses in W. The CTP controls the FWER in the strong sense (i.e. $\tilde{s} \ge 0$) at level

FWER :=
$$\sup_{\theta \in \omega_{\tilde{s}}} \Pr\left(\bigcup_{\omega_{\tilde{s}} \subseteq \omega_{s}} \phi_{s}(\boldsymbol{y}) = 1; \boldsymbol{x}, \theta\right)$$

=
$$\sup_{\theta \in \omega_{\tilde{s}}} \Pr\left(\phi_{\tilde{s}}(\boldsymbol{y}) = 1; \boldsymbol{x}, \theta\right)$$

 $\leq \alpha_{\tilde{s}}.$ (11)

by the logic of the CTP (Equation 8) and the FPR of test ϕ (Equation 10). In other words, for a familywise error to occur, it is necessary and sufficient that $\theta \in \omega_{\tilde{s}}$ be rejected by test $\phi_{\tilde{s}}(\boldsymbol{y})$. In practice, $\omega_{\tilde{s}}$ is unknown, so we have

$$FWER \leq \max\left\{\alpha_{\boldsymbol{s}} : \omega_{\boldsymbol{s}} \in W\right\}.$$
(12)

2.2. Fixed-model: classical results. We assume that the sample size, n, is sufficiently large that the likelihood of θ constrained by model s so that $\theta \in \Theta_s$ approximates that of a Multivariate Normal distribution centered around the maximum likelihood estimate (MLE).

$$p(\boldsymbol{y};\boldsymbol{x},\boldsymbol{\theta}) \approx p(\boldsymbol{y};\boldsymbol{x},\hat{\theta}^{\boldsymbol{s}}) \exp\left\{-\frac{n}{2}\left(\theta_{\mathcal{F}_{\boldsymbol{s}}}-\hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}}\right)^{T}\mathcal{I}_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}}(\tilde{\boldsymbol{\theta}})\left(\theta_{\mathcal{F}_{\boldsymbol{s}}}-\hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}}\right)\right\}$$
$$\propto f_{\mathcal{N}_{|\boldsymbol{s}|+\zeta}}\left(\hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}};\theta_{\mathcal{F}_{\boldsymbol{s}}},n^{-1}\left[\mathcal{I}(\tilde{\boldsymbol{\theta}})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}}\right]^{-1}\right) \qquad (\boldsymbol{\theta}\in\Theta_{\boldsymbol{s}})$$
(13)

Here, $\Theta_s \subseteq \Theta$ is the parameter space of model s, constrained such that $\theta_{\mathcal{B}_1 \setminus \mathcal{B}_s} = 0$ for grand alternative model $\mathbf{1}$, $f_{\mathcal{N}_{|s|+\zeta}}$ represents the joint probability density function for a $(|s| + \zeta)$ -variate Normal distribution, and $\hat{\theta}^s$ is the MLE for model s. Furthermore, $\mathcal{I}(\theta)_{\mathcal{F}_s,\mathcal{F}_s}$ is the unit (or per-observation) Fisher information matrix, (defined in Equation 14) relating to the index set of free parameters, \mathcal{F}_s , evaluated at the true parameter value, $\tilde{\theta}$. Under the regularity conditions given in Section 1, the unit Fisher information matrix can, for discrete data y, be expressed by SI Equation 14; for continuous ythe summation is replaced by an integral.

$$\{\mathcal{I}(\theta)\}_{jk} = \frac{-1}{n} \sum_{\boldsymbol{y}} \left[\frac{\partial^2}{\partial \theta_j \, \partial \theta_k} \log p(\boldsymbol{y}; \boldsymbol{x}; \theta) \right] p(\boldsymbol{y}; \boldsymbol{x}; \theta) \qquad \{j, k = 1 \dots \nu + \zeta\}.$$
(14)

2.2.1. Motivation for the Normal approximation to the likelihood. To understand the motivation for the approximate form of the likelihood, we highlight that the theory was developed with large data sets in mind. We refer the reader to asymptotic theory (see for example Davison [2003] and Chapter 9 of Cox and Hinkley [1974]), which provides a framework for assessing properties of estimators and statistical tests in the limit as sample sizes tend to infinity. A key theorem from this work demonstrates the asymptotic normality of maximum likelihood estimators for independent data, assuming standard asymptotic theory regularity conditions (covered by those listed in Section 1). Specifically, the theorem says that the difference between the maximum likelihood estimator of a multidimensional parameter and the true value of that parameter converges in distribution to a Multivariate Normal distribution, with the zero vector as the mean, as the number of samples in the data set tends to infinity. Furthermore, the covariance matrix scales with the inverse of the unit information matrix evaluated at the true value of the parameter. In the context of our framework, this gives the following,

$$\hat{\theta}_{\mathcal{F}_{s}}^{s} \xrightarrow{d} \mathcal{N}_{|s|+\zeta} \left(\tilde{\theta}_{\mathcal{F}_{s}}, \frac{1}{n} \left[\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{s}, \mathcal{F}_{s}} \right]^{-1} \right), \qquad n \to \infty ,$$
(15)

where $N_{|\mathbf{s}|+\zeta}$ is a $|\mathbf{s}| + \zeta$ -variate Normal distribution and $\theta_{\mathcal{F}_s}$ is the unknown true value of $\theta_{\mathcal{F}_s}$. The Multivariate Normal form of the likelihood assumed in this work (Equation 21) is linked to this result and can be derived through a Taylor expansion of the log-likelihood function around the MLE in a similar manner to proof of this result (Cox and Hinkley [1974], Davison [2003]). Key steps are detailed below.

First, for $\theta \in \Theta_s$, define $l(\theta) = l(\theta; \boldsymbol{y}, \boldsymbol{x}) = \log p(\boldsymbol{y}; \boldsymbol{x}, \theta)$ to be the log-likelihood function, given \boldsymbol{y} and \boldsymbol{x} . Provided that $\| \theta - \hat{\theta}^s \| = O_p(n^{-1/2})$, a Taylor expansion of the log-likelihood function around the MLE gives

$$l(\theta) = l(\hat{\theta}^{\boldsymbol{s}}) + (\theta_{\mathcal{F}_{\boldsymbol{s}}} - \hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}})U(\hat{\theta}^{\boldsymbol{s}})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}} - \frac{1}{2}(\theta_{\mathcal{F}_{\boldsymbol{s}}} - \hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}})^{T}J(\hat{\theta}^{\boldsymbol{s}})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}}(\theta_{\mathcal{F}_{\boldsymbol{s}}} - \hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}}) + o_{p}(1).$$
(16)

Where $U(\theta)$ is a vector of first-order partial derivatives,

$$\{U(\theta)\}_{ij} = \frac{\partial l}{\partial \theta_i} \tag{17}$$

and $J(\theta)$, which is known as the observed information matrix, is a vector that incorporates the secondorder partial derivatives.

$$\{J(\theta)\}_{ij} = -\frac{\partial^2 l}{\partial \theta_i \partial \theta_j} \tag{18}$$

Crucially, by the definition of the MLE,

$$U(\hat{\theta}^s)_{\mathcal{F}_s,\mathcal{F}_s} = \mathbf{0} \tag{19}$$

Furthermore, in the context of the regularity conditions assumed here (Section 1), the unit Fisher information matrix, $I(\theta)$, is a consistent estimator of $J(\theta)/n$ [Cox and Hinkley, 1974, p. 302] and the two can be used interchangeably. In much of the theory that we discuss here on, $\mathcal{I}(\theta)$ is used, as this form is the typical form used in the theorem describing the normality of the MLE (Equation 15), which we refer to later. However, in the case of estimating confidence intervals for the model parameters (as discussed in Sections 2.2.2 and 2.3.1) the observed information is a quantity that can be calculated more easily, as it does not require expectations to be derived, and is used instead. For $\theta \in \Theta_s$, Equation 16 can therefore be simplified as follows,

$$l(\theta) = l(\hat{\theta}^{\boldsymbol{s}}) - \frac{n}{2} (\theta_{\mathcal{F}_s} - \hat{\theta}_{\mathcal{F}_s}^{\boldsymbol{s}})^T \mathcal{I}(\tilde{\theta})_{\mathcal{F}_s, \mathcal{F}_s} (\theta_{\mathcal{F}_s} - \hat{\theta}_{\mathcal{F}_s}^{\boldsymbol{s}}) + o_p(1), \qquad (\theta \in \Theta_s)$$
(20)

By exponentiating both sides of Equation 20, the expression for the Multivariate Normal form of the likelihood assumed in this theory is revealed.

$$p(\boldsymbol{y};\boldsymbol{x},\theta) = p(\boldsymbol{y};\boldsymbol{x},\hat{\theta}^{\boldsymbol{s}}) \exp\left\{-\frac{n}{2}(\theta_{\mathcal{F}_{\boldsymbol{s}}} - \hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}})^{T}\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}}(\theta_{\mathcal{F}_{\boldsymbol{s}}} - \hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}}) + o_{p}(1)\right\}, \qquad (\theta \in \Theta_{\boldsymbol{s}})$$

$$(21)$$

2.2.2. Classical point estimate and variance. We are interested in providing point estimates and confidence intervals for the effect sizes for each of the covariates included in our regression analysis, as well as for a range of nuisance parameters. In the regression problem, the classical point estimate and confidence region of the parameter $\theta_{\mathcal{F}_s}^s$ (of model s) are derived from the Normal approximation to the likelihood described above (Equation 21). Thus, the classical point estimate is precisely the MLE

$$\hat{\theta}^{\boldsymbol{s}}_{\mathcal{F}_{\boldsymbol{s}}}.\tag{22}$$

A classical variance can also be derived from the Normal approximation to the likelihood and is given by $[n \mathcal{I}(\tilde{\theta})_{\mathcal{F}_s,\mathcal{F}_s}]^{-1}$ or its consistent estimator $[J(\tilde{\theta})_{\mathcal{F}_s,\mathcal{F}_s}]^{-1}$. In the usual case, where $\tilde{\theta}$ is unknown, this quantity is also unknown. When it is required that the variance is estimated in practice (e.g. in the calculation of confidence regions), then it is typical to replace this variable with another that is a consistent estimator of it. In the large *n* scenario that we are interested in, $\hat{\theta}^s$ is a consistent estimate of $\tilde{\theta}$ [Cox and Hinkley, 1974] and therefore a more useful expression for the classical variance is given by,

$$\left[J(\hat{\theta}^s)_{\mathcal{F}_s,\mathcal{F}_s}\right]^{-1}.$$
(23)

2.2.3. Classical confidence region. Using the Normal approximation to the likelihood, it is possible to provide a $(\nu_s + \zeta)$ -dimensional $100(1 - \alpha)\%$ confidence region for the parameter $\theta_{\mathcal{F}_s}$ [Stuart et al., 1998, p. 137], given by,

$$\left\{\theta_{\mathcal{F}_{\boldsymbol{s}}}:\left(\theta_{\mathcal{F}_{\boldsymbol{s}}}-\hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}}\right)^{T}J(\hat{\theta}^{\boldsymbol{s}})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}}\left(\theta_{\mathcal{F}_{\boldsymbol{s}}}-\hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}}\right)\leq Q_{\chi^{2}_{|\boldsymbol{s}|+\zeta}}(1-\alpha)\right\},\tag{24}$$

where $Q_{\chi^2_{\nu}}(p)$ is the quantile function of the chi-squared distribution with ν degrees of freedom. For an individual parameter, the $100(1-\alpha)\%$ confidence interval simplifies to

$$\left\{\hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}\right\}_{j} \quad \pm \quad \sqrt{\left(\left[J(\hat{\theta}^{\boldsymbol{s}})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}}\right]^{-1}\right)_{jj}} Q_{z}\left(1-\frac{\alpha}{2}\right) \tag{25}$$

where $Q_z(p)$ is the quantile function of the standard Normal distribution. Note the importance of model choice for determining the standard error (the square root term): the index set \mathcal{F}_s specifies which terms are fixed (and therefore contribute no uncertainty) and which are estimated.

2.2.4. Classical hypothesis tests: p-values. To test whether there is evidence that a particular set of covariates, specified by model s, should be included in the regression model, a hypothesis test comparing the maximized likelihood ratio for the test between model s and the grand null, model 0, can be performed. When the null hypothesis is true, this test statistic can be used to generate

a p-value for the test. The maximized likelihood ratio for the test between model s and the nested grand null, model 0, is represented by

$$R_{\boldsymbol{s}} = \frac{\sup\{p(\boldsymbol{y}; \boldsymbol{x}, \theta) : \theta \in \Theta_{\boldsymbol{s}}\}}{\sup\{p(\boldsymbol{y}; \boldsymbol{x}, \theta) : \theta \in \Theta_{\boldsymbol{0}}\}}$$
(26)

$$= \frac{p(\boldsymbol{y}; \boldsymbol{x}, \hat{\theta}^{\boldsymbol{s}})}{p(\boldsymbol{y}; \boldsymbol{x}, \hat{\theta}^{\boldsymbol{0}})}.$$
(27)

This expression can be evaluated by replacing the denominator with the Normal approximation to the likelihood (MT Equation 21) evaluated at the constrained maximum likelihood $(\hat{\theta}_{\mathcal{F}_s}^0)$

$$R_{\boldsymbol{s}} = \frac{p(\boldsymbol{y}; \boldsymbol{x}, \boldsymbol{\theta}^{\boldsymbol{s}})}{p(\boldsymbol{y}; \boldsymbol{x}, \hat{\boldsymbol{\theta}}^{\boldsymbol{s}}) \exp\left\{-\frac{n}{2} \left(\hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{0}} - \hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}}\right)^{T} \mathcal{I}(\tilde{\theta})_{\mathcal{F}_{\boldsymbol{s}}, \mathcal{F}_{\boldsymbol{s}}} \left(\hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{0}} - \hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}}\right) + o_{p}(1)\right\}}$$
(28)

Therefore

$$R_{\boldsymbol{s}} = \exp\left\{\frac{n}{2}\left(\hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{0}} - \hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}}\right)^{T} \mathcal{I}(\tilde{\theta})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}}\left(\hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{0}} - \hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}}\right) + o_{p}(1)\right\}$$
(29)

This expression can be further simplified to derive one that is independent of the nuisance parameters. Although the derivation has been described previously (Cox and Hinkley [1974], Davison [2003]), the key steps are outlined here. First, a partition, compatible with the partition of the parameter vector $(\tilde{\theta})$, is defined on the per unit information matrix,

$$\mathcal{I}(\tilde{\theta})_{\mathcal{F}_s,\mathcal{F}_s} = \begin{pmatrix} \mathcal{I}(\tilde{\theta})_{\mathcal{B}_s,\mathcal{B}_s} & \mathcal{I}(\tilde{\theta})_{\mathcal{B}_s,\mathcal{F}_0} \\ \mathcal{I}(\tilde{\theta})_{\mathcal{F}_0,\mathcal{B}_s} & \mathcal{I}(\tilde{\theta})_{\mathcal{F}_0,\mathcal{F}_0} \end{pmatrix}$$
(30)

and its inverse.

$$\left[\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{s},\mathcal{F}_{s}}\right]^{-1} = \begin{pmatrix} \left(\left[\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{s},\mathcal{F}_{s}}\right]^{-1}\right)_{\mathcal{B}_{s},\mathcal{B}_{s}} & \left(\left[\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{s},\mathcal{F}_{s}}\right]^{-1}\right)_{\mathcal{B}_{s},\mathcal{F}_{0}} \\ \left(\left[\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{s},\mathcal{F}_{s}}\right]^{-1}\right)_{\mathcal{F}_{0},\mathcal{B}_{s}} & \left(\left[\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{s},\mathcal{F}_{s}}\right]^{-1}\right)_{\mathcal{F}_{0},\mathcal{F}_{0}} \end{pmatrix}$$
(31)

where the dimensions of the partition, are, in each case

$$\begin{bmatrix} |\boldsymbol{s}| \times |\boldsymbol{s}| & |\boldsymbol{s}| \times \zeta \\ \zeta \times |\boldsymbol{s}| & \zeta \times \zeta \end{bmatrix}$$
(32)

Then

$$R_{\boldsymbol{s}} = \exp\left\{\frac{n}{2} \begin{pmatrix} -\hat{\beta}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}} \\ \hat{\gamma}^{\boldsymbol{0}} - \hat{\gamma}^{\boldsymbol{s}} \end{pmatrix}^{T} \begin{pmatrix} \mathcal{I}(\tilde{\theta})_{\mathcal{B}_{\boldsymbol{s}},\mathcal{B}_{\boldsymbol{s}}} & \mathcal{I}(\tilde{\theta})_{\mathcal{B}_{\boldsymbol{s}},\mathcal{F}_{0}} \\ \mathcal{I}(\tilde{\theta})_{\mathcal{F}_{0},\mathcal{B}_{\boldsymbol{s}}} & \mathcal{I}(\tilde{\theta})_{\mathcal{F}_{0},\mathcal{F}_{0}} \end{pmatrix} \begin{pmatrix} -\hat{\beta}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}} \\ \hat{\gamma}^{\boldsymbol{0}} - \hat{\gamma}^{\boldsymbol{s}} \end{pmatrix} + o_{p}(1) \right\}$$
(33)

This can be simplified because, under the large n approximation, the MLEs of the parameters common to model **0** and model **s** are related through the following expression, derived from a Taylor expansion (see p.308 of Cox and Hinkley [1974] or p.138 of Davison [2003] for proof),

$$\hat{\boldsymbol{\gamma}}^{\mathbf{0}} - \hat{\boldsymbol{\gamma}}^{\boldsymbol{s}} = \left[\mathcal{I}(\tilde{\theta})_{\mathcal{F}_0, \mathcal{F}_0} \right]^{-1} \mathcal{I}(\tilde{\theta})_{\mathcal{F}_0, \mathcal{B}_{\boldsymbol{s}}} \hat{\beta}^{\boldsymbol{s}}_{\mathcal{B}_{\boldsymbol{s}}} + o_p(n^{-1/2})$$
(34)

giving,

$$R_{\boldsymbol{s}} = \exp\left\{\frac{n}{2}\left\{\hat{\boldsymbol{\beta}}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}}\right\}^{T}\left\{\mathcal{I}(\tilde{\boldsymbol{\theta}})_{\mathcal{B}_{\boldsymbol{s}},\mathcal{B}_{\boldsymbol{s}}} - \mathcal{I}(\tilde{\boldsymbol{\theta}})_{\mathcal{B}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{0}}}\left[\mathcal{I}(\tilde{\boldsymbol{\theta}})_{\mathcal{F}_{\boldsymbol{0}},\mathcal{F}_{\boldsymbol{0}}}\right]^{-1}\mathcal{I}(\tilde{\boldsymbol{\theta}})_{\mathcal{F}_{\boldsymbol{0}},\mathcal{B}_{\boldsymbol{s}}}\right\}\hat{\boldsymbol{\beta}}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}} + o_{p}(1)\right\}$$
(35)

Using a standard linear algebra formula (see, for example, Horn and Johnson [2012]) for the inverse of a 2×2 block matrix

$$\left(\left[\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{s},\mathcal{F}_{s}}\right]^{-1}\right)_{\mathcal{B}_{s},\mathcal{B}_{s}} = \left\{\mathcal{I}(\tilde{\theta})_{\mathcal{B}_{s},\mathcal{B}_{s}} - \mathcal{I}(\tilde{\theta})_{\mathcal{B}_{s},\mathcal{F}_{0}}\left[\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{0},\mathcal{F}_{0}}\right]^{-1} \mathcal{I}(\tilde{\theta})_{\mathcal{F}_{0},\mathcal{B}_{s}}\right\}^{-1}$$
(36)

Equation 35 can be rewritten in a form that is independent of the nuisance parameters.

$$R_{\boldsymbol{s}} = \exp\left\{\frac{n}{2}\left\{\hat{\boldsymbol{\beta}}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}}\right\}^{T}\left[\left(\left[\mathcal{I}(\tilde{\boldsymbol{\theta}})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}}\right]^{-1}\right)_{\mathcal{B}_{\boldsymbol{s}},\mathcal{B}_{\boldsymbol{s}}}\right]^{-1}\hat{\boldsymbol{\beta}}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}} + o_{p}(1)\right\}$$
(37)

By the large n approximation (15), the distribution of the MLE for model s is

$$\begin{pmatrix} \hat{\beta}_{\mathcal{B}_{s}}^{s} \\ \hat{\gamma}^{s} \end{pmatrix} \stackrel{d}{\to} N_{|s|+\zeta} \left(\begin{pmatrix} \tilde{\beta}_{\mathcal{B}_{s}} \\ \tilde{\gamma} \end{pmatrix}, \frac{1}{n} \begin{pmatrix} \left([\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{s},\mathcal{F}_{s}}]^{-1} \right)_{\mathcal{B}_{s},\mathcal{B}_{s}} & \left([\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{s},\mathcal{F}_{s}}]^{-1} \right)_{\mathcal{B}_{0},\mathcal{B}_{s}} \\ \left([\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{s},\mathcal{F}_{s}}]^{-1} \right)_{\mathcal{F}_{0},\mathcal{B}_{s}} & \left([\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{s},\mathcal{F}_{s}}]^{-1} \right)_{\mathcal{F}_{0},\mathcal{F}_{0}} \end{pmatrix} \end{pmatrix}, \qquad n \to \infty$$

$$(38)$$

The marginal distribution of the MLE of $\beta_{\mathcal{B}_s}^s$ is then multivariate Normal with mean and covariance revealed by rearrangement of Equation 38 and selection of the upper left-hand element of the multivariate covariance matrix.

$$\hat{\boldsymbol{\beta}}^{\boldsymbol{s}}_{\mathcal{B}_{\boldsymbol{s}}} \xrightarrow{d} \mathcal{N}_{|\boldsymbol{s}|} \left(\tilde{\boldsymbol{\beta}}_{\mathcal{B}_{\boldsymbol{s}}}, \frac{1}{n} \left(\left[\mathcal{I}(\tilde{\boldsymbol{\theta}})_{\mathcal{F}_{\boldsymbol{s}}, \mathcal{F}_{\boldsymbol{s}}} \right]^{-1} \right)_{\mathcal{B}_{\boldsymbol{s}}, \mathcal{B}_{\boldsymbol{s}}} \right), \qquad n \to \infty$$
(39)

In hypothesis testing, we are interested in evaluating the evidence against the null. When the null is true, $\tilde{\beta}_{\mathcal{B}_s} = 0$. Since $\tilde{\gamma}$ can always be assumed to equal **0** following reparameterization (provided **0** is not at the edge of parameter space), the null can also be specified in full as $\tilde{\theta} = \mathbf{0}$. Substituting these expressions into Equations 39 gives

$$\hat{\boldsymbol{\beta}}^{\boldsymbol{s}}_{\boldsymbol{\mathcal{B}}_{\boldsymbol{s}}} \stackrel{d}{\to} \mathcal{N}_{|\boldsymbol{s}|} \left(\boldsymbol{0}, \frac{1}{n} \left(\left[\mathcal{I}(\boldsymbol{0})_{\mathcal{F}_{\boldsymbol{s}}, \mathcal{F}_{\boldsymbol{s}}} \right]^{-1} \right)_{\mathcal{B}_{\boldsymbol{s}}, \mathcal{B}_{\boldsymbol{s}}} \right), \qquad n \to \infty$$

$$\tag{40}$$

and into Equation 37, gives

$$R_{\boldsymbol{s}} = \exp\left\{\frac{n}{2}\left\{\hat{\boldsymbol{\beta}}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}}\right\}^{T}\left[\left(\left[\mathcal{I}(\boldsymbol{0})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}}\right]^{-1}\right)_{\mathcal{B}_{\boldsymbol{s}},\mathcal{B}_{\boldsymbol{s}}}\right]^{-1}\hat{\boldsymbol{\beta}}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}} + o_{p}(1)\right\}.$$
(41)

Equations 40 and 41 lead to Wilks' Theorem (Wilks [1938]); namely, that under standard asymptotic regularity conditions (covered by those in Section 1), under the null, the limiting distribution of $2 \log R_s$ (as *n* tends to infinity) is central chi-squared with degrees of freedom equal to the difference in the number of unconstrained parameters between the alternative and the null, |s|.

$$2 \log R_{\boldsymbol{s}} \xrightarrow{a} \chi_{|\boldsymbol{s}|}^2, \qquad \theta \in \Theta_{\boldsymbol{0}}, \qquad n \to \infty.$$

$$\tag{42}$$

For a given test statistic, the *p*-value is the probability of obtaining a value of the test statistic at least as extreme as the observed result, under the assumption that the null hypothesis is correct. This leads to the standard result that for large n, the *p*-value for the nested likelihood ratio test of model s versus model $\mathbf{0}$ is given by

$$p_{\boldsymbol{s}} \rightarrow \Pr\left(\chi_{|\boldsymbol{s}|}^2 \ge 2 \log R_{\boldsymbol{s}}\right), \quad \theta \in \Theta_{\boldsymbol{0}}, \quad n \to \infty.$$
 (43)

Equivalently, the significance threshold (for R_s) for a false positive rate α is given by

$$R_{s} \ge e^{\frac{1}{2}Q_{\chi_{|s|}^{-1}}^{-1}(1-\alpha)}.$$
(44)

2.2.5. Classical hypothesis tests: power. To evaluate the power of the likelihood ratio test – the probability of rejecting the null hypothesis (model $\mathbf{0}$) when, in fact, it is false – it is necessary to consider the likelihood ratio test statistic when the alternative model (model s) is true. Following the approach of earlier works (Davidson and Lever [1970], Johnson [2008] and Hu and Johnson [2009]), a local alternatives assumption can be followed, whereby,

under model **0**:
$$\theta_{\mathcal{F}_s}^{\mathbf{0}} = \begin{pmatrix} \mathbf{0} \\ \gamma \end{pmatrix}$$
 (45)

under model
$$\boldsymbol{s}$$
: $\theta_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s},n} = \begin{pmatrix} \boldsymbol{\beta}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s},n} \\ \boldsymbol{\gamma} \end{pmatrix}$ (46)

Where

$$\beta_i^{\mathbf{s},n} = q_i^n / n^{1/2}, \text{ with } \lim_{n \to \infty} q_i^n = q_i, \ i = 1, 2..., |\mathbf{s}|.$$
 (47)

In simple terms, therefore, the local alternatives hypothesis assumes that $\beta_{\mathcal{B}_s}^{s,n}$ is a sequence of alternatives that converge to the null vector at rate $n^{1/2}$ and that interest focuses on small departures from the null. The fact that larger departures from the null become self-evident in large sample sizes motivates this scaling assumption; the focus of the statistical analysis is necessarily on the 'grey area' of small departures from the null. More importantly, the assumption is convenient because it implies the tail of the distribution of the MLE is still approximately normal near the null value, which greatly simplifies the theory.

Equation 37 gives an expression for the maximized likelihood ratio.

$$R_{\boldsymbol{s}} = \exp\left\{\frac{n}{2}\left\{\hat{\boldsymbol{\beta}}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}}\right\}^{T}\left[\left(\left[\mathcal{I}(\tilde{\boldsymbol{\theta}})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}}\right]^{-1}\right)_{\mathcal{B}_{\boldsymbol{s}},\mathcal{B}_{\boldsymbol{s}}}\right]^{-1}\hat{\boldsymbol{\beta}}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}} + o_{p}(1)\right\}$$
(48)

When the alternative is true $\tilde{\boldsymbol{\beta}}_{\mathcal{B}_s} = \boldsymbol{\beta}_{\mathcal{B}_s}^{s,n}$ and $\tilde{\theta} = \theta^{s,n}$. Therefore, Equation 48 is given by

$$R_{\boldsymbol{s}} = \exp\left\{\frac{n}{2}\left\{\hat{\boldsymbol{\beta}}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}}\right\}^{T}\left[\left(\left[\mathcal{I}(\boldsymbol{\theta}^{\boldsymbol{s},n})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}}\right]^{-1}\right)_{\mathcal{B}_{\boldsymbol{s}},\mathcal{B}_{\boldsymbol{s}}}\right]^{-1}\hat{\boldsymbol{\beta}}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}} + o_{p}(1)\right\}$$
(49)

and Equation 39 gives the marginal distribution of $\hat{\beta}^{s}_{\mathcal{B}_{s}}$.

$$\hat{\boldsymbol{\beta}}^{\boldsymbol{s}}_{\boldsymbol{\mathcal{B}}_{\boldsymbol{s}}} \stackrel{d}{\to} \mathcal{N}_{|\boldsymbol{s}|} \left(\boldsymbol{\beta}^{\boldsymbol{s},n}_{\boldsymbol{\mathcal{B}}_{\boldsymbol{s}}}, \frac{1}{n} \left(\left[\mathcal{I}(\boldsymbol{\theta}^{\boldsymbol{s},n})_{\boldsymbol{\mathcal{F}}_{\boldsymbol{s}},\boldsymbol{\mathcal{F}}_{\boldsymbol{s}}} \right]^{-1} \right)_{\boldsymbol{\mathcal{B}}_{\boldsymbol{s}},\boldsymbol{\mathcal{B}}_{\boldsymbol{s}}} \right), \qquad n \to \infty$$
(50)

It follows from Equations 49 and 50 that the distribution of $2 \log R_s$ under the sequence of local alternative models specified above converges in distribution to a non-central chi-squared distribution (see Wald [1943], Davidson and Lever [1970]) with |s| degrees of freedom

$$2\log R_{\boldsymbol{s}} \xrightarrow{d} \chi^2_{|\boldsymbol{s}|,\delta}, \qquad n \to \infty$$
(51)

where the non-centrality parameter of the chi-squared distribution is

$$\delta = n \left\{ \boldsymbol{\beta}_{\mathcal{B}_{s}}^{\boldsymbol{s},n} \right\}^{T} \left[\left(\left[\mathcal{I}(\boldsymbol{\theta}^{\boldsymbol{s},n})_{\mathcal{F}_{s},\mathcal{F}_{s}} \right]^{-1} \right)_{\mathcal{B}_{s},\mathcal{B}_{s}} \right]^{-1} \boldsymbol{\beta}_{\mathcal{B}_{s}}^{\boldsymbol{s},n} \\ = \left\{ \boldsymbol{q}_{\mathcal{B}_{s}}^{n} \right\}^{T} \left[\left(\left[\mathcal{I}(\boldsymbol{\theta}^{\boldsymbol{s},n})_{\mathcal{F}_{s},\mathcal{F}_{s}} \right]^{-1} \right)_{\mathcal{B}_{s},\mathcal{B}_{s}} \right]^{-1} \boldsymbol{q}_{\mathcal{B}_{s}}^{n}$$
(52)

Since model **0** is rejected in favour of model s when $p_s \leq \alpha$, the statistical power of the test for fixed $\tilde{\theta} \in \Theta_s$ and false positive rate α would be

$$\Pr\left(\chi_{|\boldsymbol{s}|,\delta}^2 \ge Q_{\chi_{|\boldsymbol{s}|}^2}^{-1}(1-\alpha)\right).$$
(53)

Note however, that δ is not fixed under s, otherwise it would be optimal by the Neyman-Pearson lemma (Neyman and Pearson [1933]) to use that value in the likelihood ratio test, instead of estimating it. If, instead of being fixed, the non-centrality parameter was in fact encountered with frequency approximated by

$$\delta; \mathbf{s} \sim \operatorname{Gamma}\left(\frac{|\mathbf{s}|}{2}, \frac{|\mathbf{s}|}{2\bar{\delta}_{\mathbf{s}}}\right),$$
(54)

where $\mathbb{E}(\delta; \boldsymbol{s}) = \bar{\delta}_{\boldsymbol{s}}$, then

$$2 \log R_{\boldsymbol{s}}; \, \boldsymbol{s} \stackrel{d}{\to} \left(1 + \frac{\bar{\delta}_{\boldsymbol{s}}}{|\boldsymbol{s}|} \right) \, \chi^2_{|\boldsymbol{s}|} \tag{55}$$

and the statistical power of a test with variable $\tilde{\theta}$ would be

$$\Pr\left(\chi_{|\boldsymbol{s}|}^2 \ge \left(1 + \frac{\bar{\delta}_{\boldsymbol{s}}}{|\boldsymbol{s}|}\right)^{-1} Q_{\chi_{|\boldsymbol{s}|}^2}^{-1}(1-\alpha)\right).$$
(56)

2.3. Fixed-model: Bayesian results.

2.3.1. Bayesian point estimate and variance. Following Bayes rule, the posterior distribution for the free parameters for model \boldsymbol{s} , $f(\theta_{\mathcal{F}_s} | \boldsymbol{y}, \boldsymbol{x}, \boldsymbol{s})$, is proportional to the product of the likelihood, $p(\boldsymbol{y}; \boldsymbol{x}, \theta)$ with $\theta \in \Theta_s$ (Equation 13), and the prior density, $\pi(\theta|\boldsymbol{s})$ (specified by MT Equation 16). By considering the local alternatives hypothesis, as above, we write $\theta^{\boldsymbol{s},n}$ in place of $\tilde{\theta}$ within the likelihood to give

$$f(\theta_{\mathcal{F}_{\boldsymbol{s}}} \mid \boldsymbol{y}, \boldsymbol{x}, \boldsymbol{s}) \propto p(\boldsymbol{y}; \boldsymbol{x}, \theta) \pi(\theta_{\mathcal{F}_{\boldsymbol{s}}} \mid \boldsymbol{s}), \qquad (\theta \in \boldsymbol{\Theta}_{\boldsymbol{s}})$$

$$\propto f_{\mathcal{N}_{|\boldsymbol{s}|+\zeta}} \left(\hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}} \mid \theta_{\mathcal{F}_{\boldsymbol{s}}}, n^{-1} \left[\mathcal{I}(\theta^{\boldsymbol{s}, n})_{\mathcal{F}_{\boldsymbol{s}}, \mathcal{F}_{\boldsymbol{s}}} \right]^{-1} \right) f_{\mathcal{N}_{|\boldsymbol{s}|+\zeta}} \left(\theta_{\mathcal{F}_{\boldsymbol{s}}} \mid \boldsymbol{0}_{|\boldsymbol{s}|+\zeta}, h^{-1} \left[\mathcal{I}(\boldsymbol{0})_{\mathcal{F}_{\boldsymbol{s}}, \mathcal{F}_{\boldsymbol{s}}} \right]^{-1} \right).$$

$$(57)$$

By the definition of the sequence of local alternatives (Equations 46 and 47), $\theta_{\mathcal{F}_s}^{s,n}$ is a consistent estimator of $\theta_{\mathcal{F}_s}^0 = (\mathbf{0}, \gamma)^T$. Since, through transformation, γ can be assumed to equal the zero vector, $\theta^{s,n}$ is a consistent estimator of $\mathbf{0}$. Furthermore, since \mathcal{I} is a continuous function,

$$\mathcal{I}(\theta^{s,n})_{\mathcal{F}_s,\mathcal{F}_s} \to \mathcal{I}(\mathbf{0})_{\mathcal{F}_s,\mathcal{F}_s}, \qquad n \to \infty.$$
(58)

Considering Equation 57 at the limit as *n* tends to infinity, substitution of $\mathcal{I}(\mathbf{0})_{\mathcal{F}_s,\mathcal{F}_s}$ with $\mathcal{I}(\theta^{s,n})_{\mathcal{F}_s,\mathcal{F}_s}$ allows the posterior density to be written as,

$$f(\theta_{\mathcal{F}_{\boldsymbol{s}}} \mid \boldsymbol{y}, \boldsymbol{x}, \boldsymbol{s}) \propto f_{\mathcal{N}_{|\boldsymbol{s}|+\zeta}} \left(\theta_{\mathcal{F}_{\boldsymbol{s}}} \mid \frac{n}{n+h} \hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}}, \frac{n}{n+h} \left[n \,\mathcal{I}(\theta^{\boldsymbol{s},n})_{\mathcal{F}_{\boldsymbol{s}}, \mathcal{F}_{\boldsymbol{s}}} \right]^{-1} \right) \qquad n \to \infty$$
(59)

i.e., the posterior distribution for the free parameters tends in distribution to a $(|s|+\zeta)$ -variate Normal distribution.

$$\theta_{\mathcal{F}_{\boldsymbol{s}}} \mid \boldsymbol{y}, \boldsymbol{x}, \boldsymbol{s} \stackrel{d}{\to} \mathcal{N}_{|\boldsymbol{s}|+\zeta} \left(\frac{n}{n+h} \hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}}, \frac{n}{n+h} \left[n \,\mathcal{I}(\theta^{\boldsymbol{s},n})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}} \right]^{-1} \right), \qquad n \to \infty$$

$$\tag{60}$$

For the purposes of deriving a posterior mean and variance that can be calculated in practice, the substitution of $\mathcal{I}(\theta^{s,n})_{\mathcal{F}_s,\mathcal{F}_s}$ with $J(\hat{\theta}^s)_{\mathcal{F}_s,\mathcal{F}_s}/n$, gives

$$\theta_{\mathcal{F}_{\boldsymbol{s}}} \mid \boldsymbol{y}, \boldsymbol{x}, \boldsymbol{s} \stackrel{d}{\to} \mathcal{N}_{|\boldsymbol{s}|+\zeta} \left(\frac{n}{n+h} \hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}}, \frac{n}{n+h} \left[J(\hat{\theta}^{\boldsymbol{s}})_{\mathcal{F}_{\boldsymbol{s}}, \mathcal{F}_{\boldsymbol{s}}} \right]^{-1} \right), \qquad n \to \infty$$
(61)

Thus, the posterior mean

$$\frac{n}{n+h}\hat{\theta}^{\boldsymbol{s}}_{\mathcal{F}_{\boldsymbol{s}}}\tag{62}$$

and the posterior variance

$$\frac{n}{n+h} \left[J(\hat{\theta}^{s})_{\mathcal{F}_{s},\mathcal{F}_{s}} \right]^{-1} \tag{63}$$

serve as the Bayesian point estimate and variance, respectively, and each differs from its classical counterpart (Equation 22 and 23) by a factor n/(n+h).

For the fixed parameters, the posterior variances and covariances are zero and the posterior means equal the prior means of zero: $\check{V}^{s}_{ij:i\notin\mathcal{F}_{s}} = 0, \ \check{V}^{s}_{ij:j\notin\mathcal{F}_{s}} = 0, \ \check{m}^{s}_{i:i\notin\mathcal{F}_{s}} = 0.$

2.3.2. Bayesian credibility region. From the posterior distribution derived in Equation 61, the equaltailed $100(1 - 1/(1 + \tau))\%$ credibility region is given by

$$\left\{\theta_{\mathcal{F}_{\boldsymbol{s}}}:\frac{(n+h)}{n}\left(\theta_{\mathcal{F}_{\boldsymbol{s}}}-\frac{n}{n+h}\hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}}\right)^{T}J(\hat{\theta}^{\boldsymbol{s}})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}}\left(\theta_{\mathcal{F}_{\boldsymbol{s}}}-\frac{n}{n+h}\hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}}\right)\leq Q_{\chi^{2}_{|\boldsymbol{s}|+\zeta}}\left(1-\frac{1}{1+\tau}\right)\right\}$$
(64)

and the single-parameter credibility interval is given by,

$$\frac{n}{n+h} \left\{ \hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}} \right\}_{j} \quad \pm \quad \sqrt{\frac{n}{n+h}} \left(\left[J(\hat{\theta}^{\boldsymbol{s}})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}} \right]^{-1} \right)_{jj} Q_{z} \left(1 - \frac{1}{2(1+\tau)} \right) \tag{65}$$

2.3.3. Bayesian hypothesis testing: the Bayes Factor. To derive an analytical expression for the Bayes factor, an expression for the marginal probability density function, $m(\boldsymbol{y}|\boldsymbol{x},\boldsymbol{s})$, is first derived by integrating the model likelihood (Equation 21) over the prior on θ , given model \boldsymbol{s} (MT Equation 16). As above, we assume a local alternatives hypothesis under model \boldsymbol{s} , as specified by Equations 45 to 47.

$$m(\boldsymbol{y}|\boldsymbol{x},\boldsymbol{s}) \propto \int_{\Theta_{\boldsymbol{s}}} p(\boldsymbol{y};\boldsymbol{x},\boldsymbol{\theta}) \pi(\boldsymbol{\theta}|\boldsymbol{s}) d\boldsymbol{\theta}$$

$$\propto \int_{\Theta_{\boldsymbol{s}}} f_{\mathcal{N}_{|\boldsymbol{s}|+\zeta}} \left(\hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}} \mid \boldsymbol{\theta}, n^{-1} \left[\mathcal{I}(\boldsymbol{\theta}^{\boldsymbol{s},n})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}} \right]^{-1} \right) f_{\mathcal{N}_{|\boldsymbol{s}|+\zeta}} \left(\boldsymbol{\theta} \mid \boldsymbol{0}_{|\boldsymbol{s}|+\zeta}, h^{-1} \left[\mathcal{I}(\boldsymbol{0})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}} \right]^{-1} \right) d\boldsymbol{\theta}$$
(66)

Next, though replacement of $\mathcal{I}_{\mathcal{F}_s,\mathcal{F}_s}(\theta^{s,n})$ with its consistent estimator, $\mathcal{I}_{\mathcal{F}_s,\mathcal{F}_s}(\mathbf{0})$, the marginal probability mass function can be written as a Multivariate Normal density

$$m(\boldsymbol{y}|\boldsymbol{x},\boldsymbol{s}) \propto \int_{\Theta_{\boldsymbol{s}}} f_{\mathcal{N}_{|\boldsymbol{s}|+\zeta}} \left(\theta \left| \frac{n}{n+h} \hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}}, \frac{n}{n+h} \left[n \mathcal{I}(\boldsymbol{0})_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}} \right]^{-1} \right) d\theta \\ \propto f_{\mathcal{N}_{|\boldsymbol{s}|+\zeta}} \left(\hat{\theta}_{\mathcal{F}_{\boldsymbol{s}}}^{\boldsymbol{s}} \left| \boldsymbol{0}, \boldsymbol{\Sigma} \right)$$
(67)

Where

$$\boldsymbol{\Sigma} = \left(\frac{1}{h} + \frac{1}{n}\right) \left[\mathcal{I}(\mathbf{0})_{\mathcal{F}_{s},\mathcal{F}_{s}} \right]^{-1}$$
(68)

Equation 67 can be partitioned into two Multivariate Normal distributions: one defining the marginal distribution $\hat{\beta}_{\mathcal{B}_s}^s$ and the other defining the conditional distribution of $\hat{\gamma}_{\mathcal{B}_s}^s$ given $\hat{\beta}_{\mathcal{B}_s}^s$

$$m(\boldsymbol{y}|\boldsymbol{x},\boldsymbol{s}) \approx c f_{\mathcal{N}_{|\boldsymbol{s}|+\zeta}} \left(\hat{\boldsymbol{\beta}}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}} \middle| \boldsymbol{0}, \boldsymbol{\Sigma}_{\mathcal{B}_{\boldsymbol{s}},\mathcal{B}_{\boldsymbol{s}}} \right) \times f_{\mathcal{N}_{|\boldsymbol{s}|+\zeta}} \left(\hat{\boldsymbol{\gamma}}^{\boldsymbol{s}} \middle| \boldsymbol{\Sigma}_{\mathcal{F}_{0},\mathcal{B}_{\boldsymbol{s}}} \{ \boldsymbol{\Sigma}_{\mathcal{B}_{\boldsymbol{s}},\mathcal{B}_{\boldsymbol{s}}} \}^{-1} \hat{\boldsymbol{\beta}}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}} \left[\{ \boldsymbol{\Sigma}^{-1} \}_{\mathcal{F}_{0},\mathcal{F}_{0}} \right]^{-1} \right)$$
(69)

Here c is a scale constant. An expression for $m(\boldsymbol{y}|\boldsymbol{x}, \boldsymbol{0})$ can be similarly derived by integrating the model likelihood over the prior, given model $\boldsymbol{0}$.

$$m(\boldsymbol{y}|\boldsymbol{x},\boldsymbol{0}) \approx c f_{\mathcal{N}_{|\boldsymbol{s}|+\zeta}} \left(\hat{\boldsymbol{\beta}}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}} \middle| \boldsymbol{0}, \frac{h}{n+h} \boldsymbol{\Sigma}_{\mathcal{B}_{\boldsymbol{s}},\mathcal{B}_{\boldsymbol{s}}} \right) \times f_{\mathcal{N}_{|\boldsymbol{s}|+\zeta}} \left(\hat{\boldsymbol{\gamma}}^{\boldsymbol{s}} \middle| \boldsymbol{\Sigma}_{\mathcal{F}_{0},\mathcal{B}_{\boldsymbol{s}}} \{ \boldsymbol{\Sigma}_{\mathcal{B}_{\boldsymbol{s}},\mathcal{B}_{\boldsymbol{s}}} \}^{-1} \hat{\boldsymbol{\beta}}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}}, \left[\{ \boldsymbol{\Sigma}^{-1} \}_{\mathcal{F}_{0},\mathcal{F}_{0}} \right]^{-1} \right)$$
(70)

Thus the Bayes factor for the test between model s and model 0, defined as

$$BF_{\boldsymbol{s}} = \frac{m(\boldsymbol{y}|\boldsymbol{x},\boldsymbol{s})}{m(\boldsymbol{y}|\boldsymbol{x},\boldsymbol{0})}$$
(71)

can be simplified (via Equations 69 and 70) as follows

1

L

$$BF_{s} \approx \frac{f_{\mathcal{N}}\left(\hat{\boldsymbol{\beta}}_{\mathcal{B}_{s}}^{s} \middle| \mathbf{0}, \boldsymbol{\Sigma}_{\mathcal{B}_{s}, \mathcal{B}_{s}}\right)}{f_{\mathcal{N}}\left(\hat{\boldsymbol{\beta}}_{\mathcal{B}_{s}}^{s} \middle| \mathbf{0}, \frac{h}{n+h} \boldsymbol{\Sigma}_{\mathcal{B}_{s}, \mathcal{B}_{s}}\right)}$$
$$= \frac{\left(\det\left(\frac{h}{n+h} \boldsymbol{\Sigma}_{\mathcal{B}_{s}, \mathcal{B}_{s}}\right)\right)^{1/2}}{\left(\det\left(\boldsymbol{\Sigma}_{\mathcal{B}_{s}, \mathcal{B}_{s}}\right)^{1/2}}\exp\left\{-\frac{1}{2}\left(1-\frac{n+h}{h}\right)\left\{\hat{\boldsymbol{\beta}}_{\mathcal{B}_{s}}^{s}\right\}^{T}\left\{\boldsymbol{\Sigma}_{\mathcal{B}_{s}, \mathcal{B}_{s}}\right\}^{-1}\hat{\boldsymbol{\beta}}_{\mathcal{B}_{s}}^{s}\right\}}$$
(72)

Since $\Sigma_{\mathcal{B}_{s},\mathcal{B}_{s}}$ has dimension $|s| \times |s|$, det $\left(\frac{h}{n+h}\Sigma_{\mathcal{B}_{s},\mathcal{B}_{s}}\right) = \left(\frac{h}{n+h}\right)^{|s|}$ det $(\Sigma_{\mathcal{B}_{s},\mathcal{B}_{s}})$ and therefore,

$$BF_{s} \approx \left(\frac{h}{n+h}\right)^{|s|/2} \exp\left\{\frac{n}{2h}\left\{\hat{\beta}_{\mathcal{B}_{s}}^{s}\right\}^{T}\left\{\boldsymbol{\Sigma}_{\mathcal{B}_{s},\mathcal{B}_{s}}\right\}^{-1}\hat{\beta}_{\mathcal{B}_{s}}^{s}\right\}$$
$$= \left(\frac{h}{n+h}\right)^{|s|/2} \exp\left\{\frac{n}{2h}\left(\frac{1}{h}+\frac{1}{n}\right)^{-1}\left\{\hat{\beta}_{\mathcal{B}_{s}}^{s}\right\}^{T}\left[\left(\left[\mathcal{I}(\mathbf{0})_{\mathcal{F}_{s},\mathcal{F}_{s}}\right]^{-1}\right)_{\mathcal{B}_{s},\mathcal{B}_{s}}\right]^{-1}\hat{\beta}_{\mathcal{B}_{s}}^{s}\right\}$$
(73)

Through comparison of Equation 73 with Equation 41, it is possible to write the Bayes factor as a function of the MLR, R_s , under the null hypothesis. This expression has been referred to as the likelihood ratio test-based Bayes factor (Johnson [2008], Hu and Johnson [2009]).

$$BF_{s} \approx \left(\frac{h}{n+h}\right)^{|s|/2} (R_{s})^{n/(n+h)}$$
(74)

It is noteworthy that the local alternatives assumption employed above implies that the hyperparameter, h, be proportional to the sample size $(h \propto n)$, such that $\beta_i^{s,n}$ shrinks with n, as given by definition (Equation 47).

Following the methods of Johnson and colleagues, however (Johnson [2008], Hu and Johnson [2009]), this local alternatives assumption is dropped from this point forward. Instead, h is assumed to be fixed at a constant to remove the dependence of the prior on n. Furthermore, as derived previously (see Lemma 1 of Johnson [2008]) and repeated in Section 2.3.5 for completeness, when h is a constant, the logarithm of the Bayes factor is consistent, allowing for consistency of hypothesis testing.

Notice further that because the Bayes Factor depends only on a single hyper-parameter, h, is it easy to calculate. For $n \gg h$, the main role of the hyper-parameter h is to modify the penalty on the number of free parameters in the Bayesian interpretation of the maximized likelihood ratio, R_s .

The posterior odds for the test of model s to 0 is defined to be the product of model prior odds, $\mu_{s:0} = \mu^{|s|}$ (see MT Equation 19) and the Bayes Factor (Equation 74) and can be written as follows,

$$\mathrm{PO}_{\boldsymbol{s}} \approx \mu^{|\boldsymbol{s}|} \left(\frac{h}{n+h}\right)^{|\boldsymbol{s}|/2} (R_{\boldsymbol{s}})^{n/(n+h)}$$
(75)

In the Bayesian framework, the null hypothesis, model **0**, is rejected in favor of the alternative hypothesis, model s, in the sense that the smallest $100(1 - 1/(1 + \tau))\%$ posterior model credibility set excludes model **0**, when $PO_s \ge \tau$. Model **0** is therefore rejected when the following inequality holds

$$\mu^{|\boldsymbol{s}|} \left(\frac{h}{n+h}\right)^{|\boldsymbol{s}|/2} R_{\boldsymbol{s}}^{n/(n+h)} \ge \tau \tag{76}$$

2.3.4. Classical power of the Bayesian test. Since the Bayes factor is an increasing function of the MLR, the prior implies that $\bar{\delta}_{s} = |s| n/h$, and Equation 56 can be used to calculate the classical power of the Bayesian hypothesis test.

$$\Pr\left(\chi_{|\boldsymbol{s}|}^{2} \ge \left(1 + \frac{n}{h}\right)^{-1} Q_{\chi_{|\boldsymbol{s}|}^{2}}^{-1} \left(1 - \frac{1}{1+\tau}\right)\right).$$
(77)

2.3.5. The logarithm of the Bayes factor is consistent. As demonstrated by Johnson [2008], and repeated here for completeness, the logarithm of the Bayes factor (Equation 74), as given below, is consistent.

$$\log(\mathrm{BF}_{\boldsymbol{s}}) \approx -\frac{|\boldsymbol{s}|}{2}\log\left(\frac{n}{h}+1\right) + \left(\frac{n}{n+h}\right)\log\left(R_{\boldsymbol{s}}\right)$$
(78)

This means that under model $\mathbf{0}$

$$\log(\mathrm{BF}_{\boldsymbol{s}}) \xrightarrow{p} - \infty, \qquad n \to -\infty \tag{79}$$

and under model \boldsymbol{s}

$$\log(\mathrm{BF}_{s}) \xrightarrow{p} \infty, \qquad n \to \infty$$

$$\tag{80}$$

This result follows from noticing that, when the null hypothesis is true, $2 \log R_s \xrightarrow{d} \chi^2_{|s|}$ as $n \to \infty$ (Equation 42), i.e. can be written as the sum of squares of |s| independent standard Normal distributions. The second term on the right-hand side of Equation 78 is thus bounded in probability, whereas the first tends to $-\infty$ as $n \to \infty$. Therefore, $\log(BF_s) \xrightarrow{p}{\to} -\infty$.

When the alternative hypothesis is true, however, $2 \log R_s \stackrel{d}{\to} \chi^2_{|s|,\delta}$ as $n \to \infty$ (Equation 51) i.e. represents the sum of independent and identically distributed random variables that each has an expectation that is a linear function of n (Equation 52 and MT Equation 14) and finite variance. The second term on the right-hand side of Equation 78 is therefore $O_p(n)$. As the first term is $O(-\log(n))$, the term on the right-hand side dominates the one on the left-hand side, and $\log(BF_s) \stackrel{p}{\to} \infty$.

It is noteworthy that the convergence of $\log(BF_s)$ to ∞ is $O_p(n)$ when model s is true, and to $-\infty$ is only $O_p(-\log(n))$ for fixed h. Therefore it is 'easier' to reject an under-specified versus an overspecified model.

Based upon the consistency result demonstrated here, we argue that whereas the local alternative hypothesis is useful for motivating the asymptotic theory on which the approximate methods are based, in practice h should be fixed, for example at h = 1, as in the unit information prior [Kass and Wasserman, 1995], and further that in the classical significance threshold, α , should, for example, scale with $n^{-1/2}$, to make the classical approach consistent too.

3. Proofs

3.1. Proof of Theorem 2.

Step 1. Define the type I error rate given the parameters. The type I error rate can be defined to be conditional on the true' parameter $\tilde{\theta} \in \omega_{v}$ (which specifies the 'true' model $\tilde{s} \in \mathcal{O}_{\mathcal{V}}$).

$$\Pr\left(\mathrm{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau; \ \tilde{\theta}\right) = \Pr\left(\frac{\sum_{\boldsymbol{s}\in\mathcal{A}_{\mathcal{V}}}\mathrm{PO}_{\boldsymbol{s}}}{\sum_{\boldsymbol{s}\in\mathcal{O}_{\mathcal{V}}}\mathrm{PO}_{\boldsymbol{s}}} \ge \tau; \ \tilde{\theta}\right) \quad (\text{by MT Eqn. 20}) \tag{81}$$

Let $\mathcal{T}_{\mathcal{V}} = \{ t : t \in \mathcal{A}_{\mathcal{V}}, t_j = 0 \forall j \notin \mathcal{V} \}$ define the 'basic' alternatives. Then Equation 81 can be rewritten as

$$\Pr\left(\mathrm{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau; \ \tilde{\theta}\right) = \Pr\left(\frac{\sum_{s \in \mathcal{O}_{\mathcal{V}}} \mathrm{PO}_{s} \sum_{t \in \mathcal{T}_{\mathcal{V}}} \mathrm{PO}_{s+t:s}}{\sum_{s \in \mathcal{O}_{\mathcal{V}}} \mathrm{PO}_{s}} \ge \tau; \ \tilde{\theta}\right)$$
(82)

Step 2. Simplification of the posterior odds.

Step 2a. Independence case: simplification by factorization. If we assume that the likelihood factorizes

due to independence such that

$$R_{\boldsymbol{s}} = \prod_{j=1}^{\nu} R_{\boldsymbol{e}_j}^{s_j} \qquad \forall \ \boldsymbol{s} \in \mathcal{S}$$
(83)

where $\{e_j\}_i = I(i = j)$, then

$$\mathrm{PO}_{s+t:s} = \mathrm{PO}_{s'+t:s'} \quad \forall s, s' \in \mathcal{O}_{\mathcal{V}}, t \in \mathcal{T}_{\mathcal{V}}$$

$$(84)$$

allowing Equation 82 to be rewritten as

$$\Pr\left(\mathrm{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \geq \tau; \ \tilde{\theta}\right) = \Pr\left(\frac{\sum_{\boldsymbol{s}\in\mathcal{O}_{\mathcal{V}}}\mathrm{PO}_{\boldsymbol{s}} \sum_{\boldsymbol{t}\in\mathcal{T}_{\mathcal{V}}}\mathrm{PO}_{\tilde{\boldsymbol{s}}+\boldsymbol{t}:\tilde{\boldsymbol{s}}}}{\sum_{\boldsymbol{s}\in\mathcal{O}_{\mathcal{V}}}\mathrm{PO}_{\boldsymbol{s}}} \geq \tau; \ \tilde{\theta}\right)$$
$$= \Pr\left(\sum_{\boldsymbol{t}\in\mathcal{T}_{\mathcal{V}}}\mathrm{PO}_{\tilde{\boldsymbol{s}}+\boldsymbol{t}:\tilde{\boldsymbol{s}}} \geq \tau; \ \tilde{\theta}\right). \tag{85}$$

Step 2b. General case: simplification by Bayesian consistency. If we assume identifiability, Doob's theorem ([Doob, 1949]) implies consistency of the Bayesian posterior (see also Miller [2018] and Chapter 10 of Van der Vaart [2000] for more context). We therefore assert that

$$\frac{\mathrm{PO}_{\boldsymbol{s}}}{\sum_{\boldsymbol{s}\in\mathcal{O}_{\mathcal{V}}}\mathrm{PO}_{\boldsymbol{s}}} \sim \begin{cases} 1 & \text{if } \boldsymbol{s}=\tilde{\boldsymbol{s}}\\ 0 & \text{if } \boldsymbol{s}\neq\tilde{\boldsymbol{s}} \end{cases} \quad n\to\infty$$
(86)

This implies, for $t \in \mathcal{T}_{\mathcal{V}}$ and $s \in \mathcal{O}_{\mathcal{V}} \setminus \tilde{s}$,

$$\Pr\left(\frac{\operatorname{PO}_{\boldsymbol{s}}\operatorname{PO}_{\boldsymbol{s}+\boldsymbol{t}:\boldsymbol{s}}}{\sum_{\boldsymbol{s}\in\mathcal{O}_{\mathcal{V}}}\operatorname{PO}_{\boldsymbol{s}}} \ge \tau; \ \tilde{\theta}\right) = o\left(\Pr\left(\frac{\operatorname{PO}_{\tilde{\boldsymbol{s}}}\operatorname{PO}_{\tilde{\boldsymbol{s}}+\boldsymbol{t}:\tilde{\boldsymbol{s}}}}{\sum_{\boldsymbol{s}\in\mathcal{O}_{\mathcal{V}}}\operatorname{PO}_{\boldsymbol{s}}} \ge \tau; \ \tilde{\theta}\right)\right).$$
(87)

Moreover, Wilks [1938] has shown that, assuming standard regularity conditions, as \tilde{s} is the true model, that $2 \log R_{\tilde{s}+t:\tilde{s}}$ converges in distribution to a chi-squared distribution with |t| degrees of freedom

$$2\log R_{\tilde{\boldsymbol{s}}+\boldsymbol{t}:\tilde{\boldsymbol{s}}} \stackrel{d}{\to} \chi^2_{|\boldsymbol{t}|}, \qquad \tilde{\theta} \in \Theta_{\tilde{\boldsymbol{s}}}, \qquad n \to \infty$$
(88)

or equivalently, $R_{\tilde{s}+t:\tilde{s}}$ converges in distribution to a log-gamma distribution

$$R_{\tilde{\boldsymbol{s}}+\boldsymbol{t}:\tilde{\boldsymbol{s}}} \stackrel{d}{\to} \operatorname{LG}\left(\frac{|\boldsymbol{t}|}{2},1\right), \qquad \tilde{\theta} \in \Theta_{\tilde{\boldsymbol{s}}}, \qquad n \to \infty$$

$$\tag{89}$$

By the definition of the log-gamma distribution and Karamata's theorem [Karamata, 1933] (see e.g. SI Equation 16 of Wilson [2019a]), and by Johnson [2008] (MT Eqn. 15)

$$\Pr\left(R_{\tilde{\boldsymbol{s}}+\boldsymbol{t}:\tilde{\boldsymbol{s}}} > x; \; \tilde{\theta}\right) \sim \frac{\log(x)^{|\boldsymbol{t}|/2-1}}{\Gamma\left(|\boldsymbol{t}|/2\right)} \; x^{-1}, \qquad x \to \infty$$
(90)

$$\Pr\left(\operatorname{PO}_{\tilde{\boldsymbol{s}}+\boldsymbol{t}:\tilde{\boldsymbol{s}}} \geq \tau; \; \tilde{\theta}\right) \sim \Pr\left(R_{\tilde{\boldsymbol{s}}+\boldsymbol{t}:\tilde{\boldsymbol{s}}} > \frac{\tau}{\left(\mu\sqrt{\xi_n}\right)^{|\boldsymbol{t}|}}; \; \tilde{\theta}\right), \qquad n \to \infty, \tag{91}$$

where $\xi_n = h/(n+h)$. Therefore $\text{PO}_{\tilde{s}+t:\tilde{s}}$ is a non-negative regularly varying random variable with heavy tail index converging to 1.

Remark 1.3.5 of Mikosch [1999] states, for non-negative, not necessarily independent random variables X and Y, where X is regularly varying and Pr(Y > x) = o(Pr(X > x)), that

$$\Pr(X+Y>x) \sim \Pr(X>x) \quad x \to \infty.$$
 (92)

The asymptotic forms for the posterior odds and type I error probability therefore converge on the exact forms for the independence case, giving:

$$\Pr\left(\mathrm{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau; \ \tilde{\theta}\right) \sim \Pr\left(\sum_{\boldsymbol{t}\in\mathcal{T}_{\mathcal{V}}}\mathrm{PO}_{\tilde{\boldsymbol{s}}+\boldsymbol{t}:\tilde{\boldsymbol{s}}} \ge \tau; \ \tilde{\theta}\right) \qquad n \to \infty.$$
(93)

Step 3. Removal of low-order terms. By Johnson [2008] (MT Eqn. 15) we write

$$\Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau; \ \tilde{\theta}\right) \sim \Pr\left(\sum_{\boldsymbol{t}\in\mathcal{T}_{\mathcal{V}}} \left(\mu\sqrt{\xi_n}\right)^{|\boldsymbol{t}|-1} R_{\tilde{\boldsymbol{s}}+\boldsymbol{t}:\tilde{\boldsymbol{s}}} \ge \frac{\tau}{\mu\sqrt{\xi_n}}; \ \tilde{\theta}\right) \qquad n \to \infty.$$
(94)

Whenever |t'| > |t|, we have (by Eqn. 90)

$$\frac{\Pr\left(\left(\mu\sqrt{\xi_n}\right)^{|t'|-1} R_{\tilde{s}+t':\tilde{s}} \geq \frac{\tau}{\mu\sqrt{\xi_n}}; \tilde{\theta}\right)}{\Pr\left(\left(\mu\sqrt{\xi_n}\right)^{|t|-1} R_{\tilde{s}+t:\tilde{s}} \geq \frac{\tau}{\mu\sqrt{\xi_n}}; \tilde{\theta}\right)} \sim \left(\mu\sqrt{\xi_n}\right)^{|t'|-|t|} \frac{\Gamma\left(\frac{|t|}{2}\right)}{\Gamma\left(\frac{|t'|}{2}\right)} \log\left(\frac{\tau}{\mu\sqrt{\xi_n}}\right)^{(|t'|-|t|)/2} \rightarrow 0, \qquad n \to \infty.$$
(95)

Therefore, by Remark 1.3.5 of Mikosch [1999] (Equation 92 above), we can ignore all but the one degree-of-freedom tests, giving

$$\Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau; \ \tilde{\theta}\right) \sim \Pr\left(\sum_{j \in \mathcal{V}} R_{\tilde{s}+e_j:\tilde{s}} \ge \frac{\tau}{\mu\sqrt{\xi_n}}; \ \tilde{\theta}\right) \qquad n \to \infty.$$
(96)

Here e_j is a vector of length ν such that $\{e_j\}_i = 1$ if i = j and 0 if $i \neq j$.

Step 4. Simplification of the convolution.

Step 4a. Independence case: summing non-negative regularly varying random variables. Regularly varying random variables have well-known 'fractal-like' properties: they are said to be closed under convolution. Corollary 1.3.6 of Mikosch [1999] states that if $X_1, X_2, ..., X_k$ are non-negative i.i.d. regularly varying random variables, then

$$\Pr(X_1 + X_2 + \dots + X_k > x) \sim k \Pr(X_1 > x), \quad x \to \infty.$$
 (97)

Since, by Wilks [1938], $R_{\tilde{s}+e_j:\tilde{s}}$ converges to a log-gamma distribution given $\tilde{\theta} \in \Theta_{\tilde{s}}$ (Equation 89), then

$$\Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau; \ \tilde{\theta}\right) \sim |\mathcal{V}| \Pr\left(R_{\tilde{\boldsymbol{s}}+\boldsymbol{e}_{j}:\tilde{\boldsymbol{s}}} \ge \frac{\tau}{\mu\sqrt{\xi_{n}}}; \ \tilde{\theta}\right), \qquad n \to \infty.$$
(98)

for any $j \in \mathcal{V}$.

Step 4b. General case: simplification by the Davis-Resnick condition. Davis and Resnick [1996] Lemma 2.1 states that if $X_1, ..., X_k$ are non-negative, regularly varying random variables with dependence structure such that

$$\frac{\Pr(X_i > x, X_j > x)}{\Pr(X_1 > x)} \to 0, \qquad x \to \infty, \qquad i \neq j$$
(99)

then

$$\Pr(X_1 + ... + X_k > x) \sim k \Pr(X_1 > x), \qquad n \to \infty.$$
 (100)

This result applies to the sum in Equation 96. To show this, first consider

$$R_{\tilde{\boldsymbol{s}}+\boldsymbol{t}:\tilde{\boldsymbol{s}}} = \exp\left\{\frac{n}{2} \left(\hat{\theta}_{\mathcal{F}_{\tilde{\boldsymbol{s}}+\boldsymbol{t}}}^{\tilde{\boldsymbol{s}}} - \hat{\theta}_{\mathcal{F}_{\tilde{\boldsymbol{s}}+\boldsymbol{t}}}^{\tilde{\boldsymbol{s}}+\boldsymbol{t}}\right)^{T} \mathcal{I}(\tilde{\theta})_{\mathcal{F}_{\tilde{\boldsymbol{s}}+\boldsymbol{t}},\mathcal{F}_{\tilde{\boldsymbol{s}}+\boldsymbol{t}}} \left(\hat{\theta}_{\mathcal{F}_{\tilde{\boldsymbol{s}}+\boldsymbol{t}}}^{\tilde{\boldsymbol{s}}} - \hat{\theta}_{\mathcal{F}_{\tilde{\boldsymbol{s}}+\boldsymbol{t}}}^{\tilde{\boldsymbol{s}}+\boldsymbol{t}}\right) + o_{p}(1)\right\}$$
(101)

Following Equations 29 to 39 of the background theory, it is possible to show that

$$2\log(R_{\tilde{\boldsymbol{s}}+\boldsymbol{t}:\tilde{\boldsymbol{s}}}) = U_{\tilde{\boldsymbol{s}}+\boldsymbol{t}:\tilde{\boldsymbol{s}}}^{T} U_{\tilde{\boldsymbol{s}}+\boldsymbol{t}:\tilde{\boldsymbol{s}}} + o_{p}(1)$$
(102)

where

$$U_{\tilde{s}+t;\tilde{s}} = n^{1/2} \left[\left(\left[\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{\tilde{s}+t},\mathcal{F}_{\tilde{s}+t}} \right]^{-1} \right)_{\mathcal{B}_{t},\mathcal{B}_{t}} \right]^{-1/2} \hat{\beta}_{\mathcal{B}_{t}}^{\tilde{s}+t}$$
(103)

and since \tilde{s} is the true model

$$U_{\tilde{\boldsymbol{s}}+\boldsymbol{t}:\tilde{\boldsymbol{s}}} \stackrel{d}{\to} \mathcal{N}_{|\boldsymbol{t}|}\left(\boldsymbol{0},\boldsymbol{1}\right).$$
(104)

Also, by defining the following scalar: $B_i = n^{1/2} \left[\left(\left[\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{\tilde{s}+e_i}, \mathcal{F}_{\tilde{s}+e_i}} \right]^{-1} \right)_{\mathcal{B}_{e_i}, \mathcal{B}_{e_i}} \right]^{-1/2}$, we can write

$$\begin{pmatrix} U_{\tilde{s}+e_i:\tilde{s}} \\ U_{\tilde{s}+e_j:\tilde{s}} \end{pmatrix} = \begin{pmatrix} B_i & 0 \\ 0 & B_j \end{pmatrix} \begin{pmatrix} \hat{\beta}_{\mathcal{B}_{e_i}}^{\hat{s}+e_i} \\ \hat{\beta}_{\mathcal{B}_{e_j}}^{\tilde{s}+e_j} \end{pmatrix}$$
(105)

Next, let $w = \tilde{s} + e_i + e_j$. Under the large *n* approximation, a Taylor expansion (c.f. Equation 34 of the background theory) gives,

$$\hat{\theta}_{\mathcal{F}_{\tilde{s}+e_{i}}}^{\tilde{s}+e_{i}} = \hat{\theta}_{\mathcal{F}_{\tilde{s}+e_{i}}}^{\boldsymbol{w}} + \left[\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{\tilde{s}+e_{i}},\mathcal{F}_{\tilde{s}+e_{i}}}\right]^{-1} \mathcal{I}(\tilde{\theta})_{\mathcal{F}_{w},\mathcal{B}_{e_{j}}} \hat{\beta}_{\mathcal{B}_{e_{j}}}^{\boldsymbol{w}} + o_{p}(n^{-1/2})$$
(106)

By defining the following scalar: $A_i = \left(\left[\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{\tilde{s}+e_i}, \mathcal{F}_{\tilde{s}+e_i}} \right]^{-1} \mathcal{I}(\tilde{\theta})_{\mathcal{F}_w, \mathcal{B}_{e_j}} \right)_{\mathcal{B}_{e_j}}$, element selection gives

$$\hat{\beta}_{\mathcal{B}_{e_i}}^{\hat{s}+e_i} = \hat{\beta}_{\mathcal{B}_{e_i}}^w + A_i \hat{\beta}_{\mathcal{B}_{e_j}}^w + o_p(n^{-1/2})$$
(107)

$$\implies \begin{pmatrix} \hat{\beta}_{\mathcal{B}_{e_i}}^{\hat{s}+e_i} \\ \hat{\beta}_{\mathcal{B}_{e_j}}^{\hat{s}+e_j} \end{pmatrix} \approx \begin{pmatrix} 1 & A_j \\ A_i & 1 \end{pmatrix} \begin{pmatrix} \hat{\beta}_{\mathcal{B}_{e_i}}^w \\ \hat{\beta}_{\mathcal{B}_{e_j}}^w \end{pmatrix}$$
(108)

$$\implies \begin{pmatrix} U_{\tilde{s}+e_i:\tilde{s}} \\ U_{\tilde{s}+e_j:\tilde{s}} \end{pmatrix} \approx \begin{pmatrix} B_i & 0 \\ 0 & B_j \end{pmatrix} \begin{pmatrix} 1 & A_j \\ A_i & 1 \end{pmatrix} \hat{\beta}_{\mathcal{B}_{e_i+e_j}}^{\tilde{s}+e_i+e_j} \qquad (by \text{ Eqn. 105})$$
(109)

$$= \begin{pmatrix} B_{i} & B_{i}A_{j} \\ B_{j}A_{i} & B_{j} \end{pmatrix} \left[\left(\left[\mathcal{I}(\tilde{\theta})_{\mathcal{F}_{w},\mathcal{F}_{w}} \right]^{-1} \right)_{\mathcal{B}_{\boldsymbol{e}_{i}+\boldsymbol{e}_{j}},\mathcal{B}_{\boldsymbol{e}_{i}+\boldsymbol{e}_{j}}} \right]^{1/2} U_{\tilde{\boldsymbol{s}}+\boldsymbol{e}_{i}+\boldsymbol{e}_{j}:\tilde{\boldsymbol{s}}}$$
(110)
(by Eqn. 103)

From Equation 104, $U_{\tilde{s}+e_i+e_j:\tilde{s}} \xrightarrow{d} \mathcal{N}_{|t|}(0, 1)$. Therefore, the joint distribution of the scores is a linear transformation of a standard bivariate normal distribution and, through multivariate normal affine transformation rules, is therefore also a bivariate normal distribution. Thus, for some correlation coefficient (-1 < c < 1), we can write

$$\begin{pmatrix} U_{\tilde{\mathbf{s}}+\mathbf{e}_{i}:\tilde{\mathbf{s}}} \\ U_{\tilde{\mathbf{s}}+\mathbf{e}_{j}:\tilde{\mathbf{s}}} \end{pmatrix} \stackrel{d}{\rightarrow} \mathcal{N}_{2}\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & c \\ c & 1 \end{pmatrix}\right).$$
(111)

Therefore, for $i, j \in \mathcal{V}, i \neq j$,

$$\Pr\left(R_{\tilde{\boldsymbol{s}}+\boldsymbol{e}_{i}:\tilde{\boldsymbol{s}}} > x, R_{\tilde{\boldsymbol{s}}+\boldsymbol{e}_{j}:\tilde{\boldsymbol{s}}} > x; \; \tilde{\theta}\right) = 4\Pr\left(U_{\tilde{\boldsymbol{s}}+\boldsymbol{e}_{i}:\tilde{\boldsymbol{s}}} > \sqrt{2\log(x)}, U_{\tilde{\boldsymbol{s}}+\boldsymbol{e}_{j}:\tilde{\boldsymbol{s}}} > \sqrt{2\log(x)}\right)$$
$$\sim \frac{1}{\pi} \frac{(1+c)^{2}}{\sqrt{1-c^{2}}} \log(x)^{-1} x^{-2/(1+c)}, \qquad x \to \infty, \tag{112}$$

Here, the last line is from a standard asymptotic form for the bivariate Normal distribution [Zhou and Xiao, 2017]. From Equation 90

$$\Pr\left(R_{\tilde{\boldsymbol{s}}+\boldsymbol{e}_{i}:\tilde{\boldsymbol{s}}} > x; \,\tilde{\theta}\right) \sim \frac{1}{\sqrt{\pi}} \log(x)^{-1/2} x^{-1}, \qquad x \to \infty.$$
(113)

The Davis-Resnick condition (Equation 99) is therefore satisfied because

$$\frac{\Pr\left(R_{\tilde{\boldsymbol{s}}+\boldsymbol{e}_{i}:\tilde{\boldsymbol{s}}} > \frac{\tau}{\mu\sqrt{\xi_{n}}}, R_{\tilde{\boldsymbol{s}}+\boldsymbol{e}_{j}:\tilde{\boldsymbol{s}}} > \frac{\tau}{\mu\sqrt{\xi_{n}}}; \theta\right)}{\Pr\left(R_{\tilde{\boldsymbol{s}}+\boldsymbol{e}_{k}:\tilde{\boldsymbol{s}}} > \frac{\tau}{\mu\sqrt{\xi_{n}}}; \theta\right)} \sim \frac{1}{\sqrt{\pi}} \frac{(1+c)^{2}}{\sqrt{1-c^{2}}} \log\left(\frac{\tau}{\mu\sqrt{\xi_{n}}}\right)^{-\frac{1}{2}} \left(\frac{\tau}{\mu\sqrt{\xi_{n}}}\right)^{-\frac{(1-c)}{(1+c)}} \rightarrow 0, \quad n \to \infty \quad (i, j, k \in \mathcal{V}, i \neq j) \quad (114)$$

Applying Davis and Resnick [1996] Lemma 2.1 to Equation 96 gives the same asymptotic form for the type I error probability as the independence case (Equation 98):

$$\Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau; \ \tilde{\theta}\right) \sim |\mathcal{V}| \Pr\left(R_{\tilde{s}+e_{j}:\tilde{s}} \ge \frac{\tau}{\mu\sqrt{\xi_{n}}}; \ \tilde{\theta}\right), \qquad n \to \infty.$$
(115)

for any $j \in \mathcal{V}$.

Step 5. Taking the supremum. By the definition of regular variation (Karamata [1933]; Mikosch [1999] Definition 1.1.1), a non-negative random variable X with regularly varying tail probability and tail index λ satisfies $\Pr(X > a x) \sim a^{-\lambda} \Pr(X > x)$ as $x \to \infty$ for any constant a > 0. Therefore,

$$\Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau; \ \tilde{\theta}\right) \sim \Pr\left(R_{\tilde{s}+e_{j}:\tilde{s}} \ge \frac{\tau}{|\mathcal{V}|\mu\sqrt{\xi_{n}}}; \ \tilde{\theta}\right), \qquad n \to \infty$$
(116)

By the definition of the respective distributions, if $X \stackrel{d}{=} LG(\nu/2, 1)$ then $2 \log X \stackrel{d}{=} \chi^2_{\nu}$. Finally, we have

$$\alpha_{\boldsymbol{v}} = \sup_{\boldsymbol{\theta} \in \omega_{\boldsymbol{v}}} \Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau ; \boldsymbol{\theta}\right)$$
(117)

~
$$\Pr\left(\chi_1^2 \ge 2\log\frac{\tau}{|\mathcal{V}|\,\mu\sqrt{\xi_n}}\right), \qquad n \to \infty$$
 (118)

in the sense that

$$\sup_{\theta \in \omega_{v}} \lim_{n \to \infty} \frac{\Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau ; \theta\right)}{\Pr\left(\chi_{1}^{2} \ge 2\log\frac{\tau}{|\mathcal{V}| \mu \sqrt{\xi_{n}}}\right)} = 1.$$
(119)

3.2. Proof of Theorem 3.

Proof. By the logic of a CTP (Definition 1), the FWER is controlled at or below

$$FWER \leq \max\{\alpha_{\boldsymbol{v}} : \omega_{\boldsymbol{v}} \in W\},\tag{120}$$

where the FPR for the test of null hypothesis $\omega_{\boldsymbol{v}}$ is

$$\alpha_{\boldsymbol{v}} = \sup_{\boldsymbol{\theta} \in \boldsymbol{\omega}_{\boldsymbol{v}}} \Pr\left(\psi_{\boldsymbol{v}}(\boldsymbol{y}) = 1; \boldsymbol{x}, \boldsymbol{\theta}\right), \qquad (121)$$

and $\psi_{\boldsymbol{v}}(\boldsymbol{y}) = I(\operatorname{PO}_{\omega_{\boldsymbol{v}}^{c}:\omega_{\boldsymbol{v}}} \geq \tau)$ for a Bayesian test (Definition 2).

Moreover, by Theorem 1, the Bayesian test is a shortcut CTP, so $\phi_{\boldsymbol{v}}(\boldsymbol{y}) = \psi_{\boldsymbol{v}}(\boldsymbol{y})$.

In the Doublethink model (Definition 7) we specify the null hypotheses as $\omega_{\boldsymbol{v}} = \{\theta : \beta_j = 0 \forall v_j = 0\}, \boldsymbol{v} \in 2^{\nu}$, and we use the notation $\mathrm{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} = \mathrm{PO}_{\omega_{\boldsymbol{v}}^{\mathsf{c}}:\omega_{\boldsymbol{v}}}$, where $\mathcal{V} = \{j : v_j = 0\}.$

By Theorem 2 we have

$$\alpha_{\boldsymbol{v}} = \sup_{\boldsymbol{\theta} \in \omega_{\boldsymbol{v}}} \Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau ; \boldsymbol{\theta}\right) \sim \Pr\left(\chi_{1}^{2} \ge 2\log\frac{\tau}{|\mathcal{V}|\,\mu\sqrt{\xi_{n}}}\right), \quad n \to \infty$$
(122)

meaning

$$\sup_{\theta \in \omega_{v}} \lim_{n \to \infty} \frac{\Pr\left(\operatorname{PO}_{\mathcal{A}_{v}:\mathcal{O}_{v}} \ge \tau; \theta\right)}{\Pr\left(\chi_{1}^{2} \ge 2\log\frac{\tau}{|\mathcal{V}| \mu \sqrt{\xi_{n}}}\right)} = 1.$$
(123)

The tail probability of the chi-squared distribution is monotonic decreasing, so

$$\max_{\omega_{\nu} \in W} \Pr\left(\chi_{1}^{2} \ge 2\log\frac{\tau}{|\mathcal{V}|\,\mu\sqrt{\xi_{n}}}\right) = \Pr\left(\chi_{1}^{2} \ge 2\log\frac{\tau}{\nu\,\mu\sqrt{\xi_{n}}}\right).$$
(124)

This gives

$$\alpha := \max_{\omega_{\boldsymbol{v}} \in W} \sup_{\theta \in \omega_{\boldsymbol{v}}} \Pr\left(\mathrm{PO}_{\omega_{\boldsymbol{v}}^{\mathsf{c}}:\omega_{\boldsymbol{v}}} \ge \tau; \theta\right) \sim \Pr\left(\chi_{1}^{2} \ge 2\log\frac{\tau}{\nu \mu \sqrt{\xi_{n}}}\right), \quad n \to \infty$$
(125)

meaning

$$\max_{\omega_{\boldsymbol{v}}\in W} \sup_{\theta\in\omega_{\boldsymbol{v}}} \lim_{n\to\infty} \frac{\Pr\left(\mathrm{PO}_{\omega_{\boldsymbol{v}}^{\mathsf{c}}:\omega_{\boldsymbol{v}}} \ge \tau; \theta\right)}{\Pr\left(\chi_{1}^{2} \ge 2\log\frac{\tau}{\nu\,\mu\,\sqrt{\xi_{n}}}\right)} \le 1.$$
(126)

3.3. Proof of Theorem 4. Uniform convergence in a simplified two-variable model.

Proof. Let $\theta = (\gamma, \beta_1, \beta_2)^T$, where γ is the intercept and β_1 and β_2 are the regression coefficients for the model

$$Y \stackrel{d}{=} \mathcal{N}_n\left((1, X_1, X_2) \theta, \sigma^2 I_n\right),\tag{127}$$

 σ^2 assumed known. Here, I_n is the identity matrix of dimension n. The unit Fisher information is given by:

$$\mathcal{I}(\theta) = \frac{1}{n\sigma^2} (1, X_1, X_2)^T (1, X_1, X_2) = \frac{1}{\sigma^2} \begin{pmatrix} 1 & 0 & 0\\ 0 & 1 & \rho\\ 0 & \rho & 1 \end{pmatrix}$$
(128)

And its inverse is given by:

$$\mathcal{I}(\theta)^{-1} = \sigma^2 \begin{pmatrix} 1 & 00\\ 0 & (I(\theta)_{\mathcal{B},\mathcal{B}})^{-1} \end{pmatrix} \qquad \left(\text{where } I(\theta)_{\mathcal{B},\mathcal{B}} = \begin{pmatrix} 1 & \rho\\ \rho & 1 \end{pmatrix} \right)$$
(129)

From Equation 128

$$\left[\mathcal{I}(\theta)_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}}\right]^{-1} = \begin{cases} \sigma^2 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} & \text{if } \boldsymbol{s} \in \{\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}\} \\ \mathcal{I}(\theta)^{-1} & \text{if } \boldsymbol{s} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \end{cases}$$
(130)

Following step 4b of the proof of Theorem 2, the score statistics for the models $\boldsymbol{s} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\boldsymbol{s} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ can be expressed as linear transformations of the grand alternative model, $\boldsymbol{s} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$

$$U_{\boldsymbol{s}} = n^{1/2} \left[\left(\left[\mathcal{I}(\theta)_{\mathcal{F}_{\boldsymbol{s}},\mathcal{F}_{\boldsymbol{s}}} \right]^{-1} \right)_{\mathcal{B}_{\boldsymbol{s}},\mathcal{B}_{\boldsymbol{s}}} \right]^{-1/2} \hat{\boldsymbol{\beta}}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}}, \quad (\text{Eqn. 103})$$
(131)

$$= \begin{cases} \frac{n^{1/2}}{\sigma} \hat{\boldsymbol{\beta}}_{\mathcal{B}_{\boldsymbol{s}}}^{\boldsymbol{s}} & \text{if } \boldsymbol{s} \in \{ \begin{pmatrix} 1\\ 0 \end{pmatrix}, \begin{pmatrix} 0\\ 1 \end{pmatrix} \} \\ \frac{n^{1/2}}{\sigma} \left(I(\theta)_{\mathcal{B},\mathcal{B}} \right)^{1/2} \hat{\boldsymbol{\beta}}^{\boldsymbol{s}} & \text{if } \boldsymbol{s} = \begin{pmatrix} 1\\ 1 \end{pmatrix} \end{cases}$$
(132)

Since the likelihood surface is multivariate normal

$$\hat{\boldsymbol{\beta}}_{\mathcal{B}_{s}}^{s} = \hat{\boldsymbol{\beta}}_{\mathcal{B}_{s}}^{11} - \left(\mathcal{I}(\theta)^{-1}\right)_{\mathcal{B}_{s},\mathcal{B}_{s'}} \left[\left(\mathcal{I}(\theta)^{-1}\right)_{\mathcal{B}_{s'},\mathcal{B}_{s'}} \right]^{-1} \hat{\boldsymbol{\beta}}_{\mathcal{B}_{s'}}^{11}
= \hat{\boldsymbol{\beta}}_{\mathcal{B}_{s}}^{11} + \rho \hat{\boldsymbol{\beta}}_{\mathcal{B}_{s'}}^{11} \quad \text{for } (\boldsymbol{s}, \boldsymbol{s'}) \in \left\{ \left(\begin{pmatrix} 1\\ 0 \end{pmatrix}, \begin{pmatrix} 0\\ 1 \end{pmatrix} \right), \left(\begin{pmatrix} 1\\ 1 \end{pmatrix}, \begin{pmatrix} 1\\ 1 \end{pmatrix} \right) \right\} \quad \text{(Eqn. x)}$$
(133)

Therefore,

$$\begin{pmatrix} U_{10} \\ U_{01} \end{pmatrix} = n^{1/2} I(\theta)_{\mathcal{B},\mathcal{B}} \hat{\beta}^{11} \quad \text{(by Eqns. 132 and 133)}$$

$$= I(\theta)_{\mathcal{B},\mathcal{B}} (I(\theta)_{\mathcal{B},\mathcal{B}})^{-1/2} U_{11} \quad \text{(by Eqn. 132)}$$

$$= (I(\theta)_{\mathcal{B},\mathcal{B}})^{1/2} U_{11}$$

$$= \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix}^{1/2} \begin{pmatrix} W \\ Z \end{pmatrix} \quad \text{(letting } U_{11} = \begin{pmatrix} W \\ Z \end{pmatrix})$$

$$= \begin{pmatrix} \sqrt{1-\rho^2} & \rho \\ 0 & 1 \end{pmatrix} \begin{pmatrix} W \\ Z \end{pmatrix} \quad \text{(by Cholesky decomposition)}$$

$$= \begin{pmatrix} \sqrt{1-\rho^2}W + \rho Z \\ Z \end{pmatrix}$$

$$(134)$$

Let $\tilde{\theta} = \begin{pmatrix} \tilde{\gamma} \\ \tilde{\beta} \end{pmatrix}$ be the true parameter vector. The distribution of U_{11} can be derived as follows

$$\hat{\boldsymbol{\beta}}^{11} \stackrel{d}{=} \mathcal{N}_{2} \left(\tilde{\boldsymbol{\beta}}, \frac{1}{n} [\mathcal{I}(\tilde{\boldsymbol{\theta}})^{-1}]_{\mathcal{B}, \mathcal{B}} \right) \quad (\text{c.f. Equation 39})$$

$$\implies \qquad U_{11} \stackrel{d}{=} \mathcal{N}_{2} \left(n^{\frac{1}{2}} \left([\mathcal{I}(\tilde{\boldsymbol{\theta}})^{-1}]_{\mathcal{B}, \mathcal{B}} \right)^{-\frac{1}{2}} \tilde{\boldsymbol{\beta}}, I_{2} \right) \quad (\text{Eqn. 131})$$

$$\implies \qquad \begin{pmatrix} W \\ Z \end{pmatrix} \stackrel{d}{=} \mathcal{N}_{2} \left(\frac{n^{1/2}}{\sigma} \left(\frac{1}{\rho} \frac{0}{\sqrt{1-\rho^{2}}} \right) \tilde{\boldsymbol{\beta}}, I_{2} \right)$$

$$\stackrel{d}{=} \mathcal{N}_{2} \left(\left(\frac{n^{1/2}\sigma^{-1}\tilde{\beta}_{1}}{n^{1/2}\sigma^{-1}(\rho\tilde{\beta}_{1}+\sqrt{1-\rho^{2}}\tilde{\beta}_{2})} \right), I_{2} \right) \quad (135)$$

To consider the limiting false positive rate, notice that there are three possible hypothesis tests: $\mathcal{V} = \{1, 2\}, \mathcal{V} = \{1\}$ and $\mathcal{V} = \{2\}$. Since the third of these mirrors the second theoretically, we do not consider it further. Among the remaining two hypothesis tests, the false positive rate can be calculated for the following three cases:

Case 1:
$$\tilde{s} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \ \mathcal{V} = \{1\}$$
 (136)

Case 2:
$$\tilde{s} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \ \mathcal{V} = \{1, 2\}$$
 (137)

Case 3:
$$\tilde{s} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \ \mathcal{V} = \{1\}$$
 (138)

Since $\tilde{\beta}_1 = 0$ for each case,

$$\begin{pmatrix} W \\ Z \end{pmatrix} \stackrel{d}{=} \mathcal{N}_2\left(\begin{pmatrix} 0 \\ \sqrt{n(1-\rho^2)}\tilde{\beta}_2/\sigma \end{pmatrix}, I_2 \right)$$
(Eqn. 135) (139)

Let $\check{Z} = Z - \frac{\sqrt{n(1-\rho^2)}\tilde{\beta}_2}{\sigma}$, then

$$\begin{pmatrix} W \\ \check{Z} \end{pmatrix} \stackrel{d}{=} \mathcal{N}_2(\mathbf{0}, I_2) \qquad (\text{Eqn. 135}) \tag{140}$$

The model-averaged posterior odds is given by

$$\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}(Z,W) = \frac{\sum_{s \in \mathcal{A}_{\mathcal{V}}} \operatorname{PO}_{s}(Z,W)}{\sum_{s \in \mathcal{O}_{\mathcal{V}}} \operatorname{PO}_{s}(Z,W)}$$
(MT Eqn. 20) (141)

Where

$$PO_{\boldsymbol{s}}(Z,W) = \mu^{|\boldsymbol{s}|} \left(\frac{h}{n+h}\right)^{|\boldsymbol{s}|/2} (R_{\boldsymbol{s}}(Z,W))^{n/(n+h)} \quad (Eqn. 75)$$
$$= (\mu\sqrt{\xi_n})^{|\boldsymbol{s}|} e^{U_{\boldsymbol{s}}^T U_{\boldsymbol{s}}(1-\xi_n)/2} \quad \left(\text{since } \xi_n = \frac{h}{n+h}\right) \quad (142)$$

Let $\Theta_{\boldsymbol{s}}^* = \{ \theta \in \Theta_{\boldsymbol{s}} : \ |\beta_2| \ge \beta_{\min} \}$

 $\underline{\text{Case 1 or 2}}:\; (\tilde{\boldsymbol{s}}=({}^0_0) \;\;\text{and}\; \mathcal{V}=\{1\} \;\text{or}\; \mathcal{V}=\{1,2\})$

Since $\tilde{s} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$, then $\tilde{\beta}_2 = 0$, which implies $Z = \check{Z}$. Therefore

$$\Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}(Z,W) \ge \tau; \theta\right) = \Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}(Z,W) \ge \tau; \theta\right)$$
(143)

Since the right-hand side does not depend on any unknown θ (i.e. does not depend on $\tilde{\gamma}$)

$$\sup_{\theta \in \Theta_{00}^{*}} \Pr\left(\mathrm{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau; \; \theta\right) = \Pr\left(\mathrm{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau; \; \theta\right)$$
(144)

Therefore

$$\lim_{n \to \infty} \sup_{\substack{\theta \in \Theta_{00}^* \\ \rho \in P^*}} \frac{\Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau; \theta\right)}{\Pr\left(\chi_1^2 \ge 2\log\frac{\tau}{|\mathcal{V}|\mu\sqrt{\xi_n}}\right)} = \lim_{n \to \infty} \frac{\Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau; \theta\right)}{\Pr\left(\chi_1^2 \ge 2\log\frac{\tau}{|\mathcal{V}|\mu\sqrt{\xi_n}}\right)}$$
(145)

=

$$= 1$$
 (by Ther. 2: MT Eqn. 23) (146)

Case 3:
$$(\tilde{\boldsymbol{s}} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$
 and $\mathcal{V} = \{1\}$)

Since $\tilde{\boldsymbol{s}} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$, then

$$\Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}(Z,W) \ge \tau; \; \theta\right) = \Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}\left(\check{Z} + \frac{\sqrt{n(1-\rho^2)}\tilde{\beta}_2}{\sigma}, W\right) \ge \tau; \; \theta\right)$$
(147)

i.e. the right hand side is dependent on the unknown $\tilde{\beta}_2$ – although not on $\tilde{\gamma}$. For readability let PO_{{1}} be shorthand for PO_{$\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}$} when $\mathcal{V} = \{1\}$. From Equations 141 and 142

$$\operatorname{PO}_{\{1\}}(Z,W) = \mu \sqrt{\xi_n} \, e^{(1-\xi_n)W^2/2} \, \left(\frac{e^{-(1-\xi_n)(\rho W - \sqrt{1-\rho^2 Z})^2/2} + \mu \sqrt{\xi_n}}{e^{-(1-\xi_n)Z^2/2} + \mu \sqrt{\xi_n}} \right)$$
(148)

Also let

$$\operatorname{PO}_{\{1\}}^{+}(Z,W) = \mu \sqrt{\xi_n} e^{(1-\xi_n)W^2/2} \iota_n(Z)$$
(149)

Where

$$\iota_n(Z) = 1 + \frac{1}{\mu\sqrt{\xi_n}} e^{-(1-\xi_n) \left(|\rho| \, w_c(Z) - |Z|\sqrt{1-\rho^2}\right)^2/2} \tag{150}$$

and the critical values, w_c , are given by

$$w_c(Z) = \begin{cases} \sqrt{2\log\frac{\tau}{1+\mu\sqrt{\xi_n}}} & |Z| < \frac{|\rho|}{\sqrt{1-\rho^2}}\sqrt{2\log\frac{\tau}{1+\mu\sqrt{\xi_n}}}\\ \sqrt{2\log\frac{\tau}{\mu\sqrt{\xi_n}}} & |Z| > \frac{|\rho|}{\sqrt{1-\rho^2}}\sqrt{2\log\frac{\tau}{\mu\sqrt{\xi_n}}}\\ \frac{|Z|\sqrt{1-\rho^2}}{|\rho|} & \text{otherwise} \end{cases}$$
(151)

Let $s_{\rho} = \operatorname{sgn}(\rho)$ and $s_Z = \operatorname{sgn}(Z)$. For $\log \frac{\tau}{1 + \mu \sqrt{\xi_n}} \le W^2 < \log \frac{\tau}{\mu \sqrt{\xi_n}}$, it can be shown that

$$(s_{\rho}s_{Z}W - w_{c}(Z))\left(\rho^{2}(s_{\rho}s_{Z}W + w_{c}(Z)) - 2|\rho||Z|\sqrt{1 - \rho^{2}}\right) \geq 0$$

$$\implies \left(s_{\rho}|\rho|W - s_{Z}|Z|\sqrt{1 - \rho^{2}}\right)^{2} \geq \left(|\rho|w_{c}(Z) - |Z|\sqrt{1 - \rho^{2}}\right)^{2}$$

$$\implies e^{-(1 - \xi)(\rho W - Z\sqrt{1 - \rho^{2}})^{2}/2} \leq e^{-(1 - \xi)(|\rho|w_{c}(Z) - |Z|\sqrt{1 - \rho^{2}})^{2}/2}$$

$$\implies PO_{\{1\}}(Z, W) \leq PO_{\{1\}}^{+}(Z, W)$$
(152)

For $W^2 \ge \log \frac{\tau}{\mu \sqrt{\xi_n}}$

$$\operatorname{PO}_{\{1\}}^{+}(Z,W) \ge \tau \le \operatorname{PO}_{\{1\}}(Z,W)$$
 (153)

$$\implies \Pr\left(\operatorname{PO}_{\{1\}}^{+}(Z,W) \ge \tau; \theta\right) = 1 = \Pr\left(\operatorname{PO}_{\{1\}}(Z,W) \ge \tau; \theta\right)$$
(154)

For $W^2 < \log \frac{\tau}{1 + \mu \sqrt{\xi_n}}$

$$\operatorname{PO}_{\{1\}}^{+}(Z,W) < \tau > \operatorname{PO}_{\{1\}}(Z,W)$$
 (155)

$$\implies \Pr\left(\operatorname{PO}_{\{1\}}^+(Z,W) \ge \tau; \theta\right) = 0 = \Pr\left(\operatorname{PO}_{\{1\}}(Z,W) \ge \tau; \theta\right)$$
(156)

Therefore

$$\Pr\left(\operatorname{PO}_{\{1\}}(Z,W) \ge \tau; \ \theta\right) \le \Pr\left(\operatorname{PO}_{\{1\}}^+(Z,W) \ge \tau; \ \theta\right) \qquad \forall \ (Z,W)$$
(157)

Let

$$\check{\iota}_n(\check{Z},\beta_2) := \iota_n(Z) = \iota_n\left(\check{Z} + \frac{\sqrt{n(1-\rho^2)\tilde{\beta}_2}}{\sigma}\right)$$
(158)

and let ϕ be the pdf of a standard normal random variable. Then Equation 147 can be rewritten as

$$\sup_{\substack{\theta \in \Theta_{01}^{*}\\ \rho \in P^{*}}} \Pr\left(\operatorname{PO}_{\{1\}}(Z, W) \ge \tau; \theta\right) \leq \sup_{\substack{\beta_{2}: |\beta_{2}| \ge \beta_{\min}\\ \rho \in P^{*}}} \int_{-\infty} \Pr\left(W^{2} \ge \frac{2}{1-\xi_{n}} \log\left(\tau/\mu\sqrt{\xi_{n}}\,\check{\iota}_{n}(\check{z},\beta_{2})\right)\right) \phi(\check{z}) d\check{z} \\ \leq \int_{-\infty}^{\infty} \sup_{\substack{\beta_{2}: |\beta_{2}| \ge \beta_{\min}\\ \rho \in P^{*}}} \Pr\left(W^{2} \ge \frac{2}{1-\xi_{n}} \log\left(\tau/\mu\sqrt{\xi_{n}}\,\check{\iota}_{n}(\check{z},\beta_{2})\right)\right) \phi(\check{z}) d\check{z}$$

$$(159)$$

 ∞

Notice that $\check{\iota}_n(\check{z},\beta_2)$ and therefore $\Pr\left(W^2 \ge \frac{2}{1-\xi_n} \log\left(\frac{\tau}{\mu\sqrt{\xi_n}}\check{\iota}_n(\check{z},\beta_2)\right)\right)$ is maximized w.r.t. β_2 when

$$0 = |\rho|w_{c}(\check{z}) - |\check{z}| + \sqrt{n(1-\rho^{2})}\beta_{2}/\sigma\sqrt{1-\rho^{2}}$$

$$\iff \beta_{2} = \frac{\sigma(|\rho|w_{c}(\check{z}) - \sqrt{1-\rho^{2}}|\check{z}|)}{(1-\rho^{2})\sqrt{n}}$$
(160)

$$=: \beta_2^{\text{opt}} \quad \text{(by definition)}$$

$$(161)$$

Taking account of the assumption that $|\beta_2| \ge \beta_{\min}$, the value of β_2 at the supremum, denoted $\beta_2^{\sup}(\check{z})$, is therefore given by

$$\beta_2^{\sup}(\check{z}) = \begin{cases} \beta_2^{\operatorname{opt}}(\check{z}) & \text{if } |\beta_2^{\operatorname{opt}}(\check{z})| \le \beta_{\min} \\ \operatorname{sgn}(\beta_2^{\operatorname{opt}}(\check{z}))\beta_{\min} & \text{otherwise} \end{cases}$$
(162)

allowing Equation 159 to be rewritten as

$$\sup_{\substack{\theta \in \Theta_{01}^*\\\rho \in P^*}} \Pr\left(\operatorname{PO}_{\{1\}}(Z,W) \ge \tau \; ; \; \theta\right) \leq \int_{-\infty}^{\infty} \Pr\left(\chi_1^2 \ge \frac{2}{1-\xi_n} \log\left(\tau/\mu\sqrt{\xi} \,\check{\iota}_n(\check{z},\beta_2^{\sup}(\check{z}))\right)\right) \, \phi(\check{z}) \, d\check{z} \quad (163)$$

Therefore

$$\lim_{n \to \infty} \sup_{\substack{\theta \in \Theta_{01}^{*}\\ \rho \in P^{*}}} \frac{\Pr\left(\operatorname{PO}_{\{1\}} \ge \tau \;;\; \theta\right)}{\Pr\left(\chi_{1}^{2} \ge 2\log\left(\tau/\mu\sqrt{\xi_{n}}\right)\right)} \leq \lim_{n \to \infty} \int_{-\infty}^{\infty} \frac{\Pr\left(\chi_{1}^{2} \ge \frac{2}{1-\xi_{n}}\log\left(\tau/\mu\sqrt{\xi}\,i_{n}(\tilde{z},\beta_{2}^{\sup}(\tilde{z}))\right)\right)}{\Pr\left(\chi_{1}^{2} \ge 2\log\tau/\mu\sqrt{\xi_{n}}\right)} \phi(\tilde{z}) \, d\tilde{z}$$

$$= \lim_{n \to \infty} \int_{-\infty}^{\infty} \frac{\bar{F}_{n}\left(\tau/\mu\sqrt{\xi}\,i_{n}(\tilde{z},\beta_{2}^{\sup}(\tilde{z}))\right)}{\bar{F}_{n}\left(\tau/\mu\sqrt{\xi_{n}}\right)} \phi(\tilde{z}) \, d\tilde{z} \tag{164}$$

Where $\bar{F}_n(x)$ is the tail probability of a LG $\left(\frac{1-\xi_n}{2},1\right)$ distribution (a regularly varying function at $n \to \infty$ with tail index -1). Let

$$f_n(\check{z}) = \frac{\bar{F}_n\left(\tau/\mu\sqrt{\xi_n}\,\check{\iota}_n(\check{z},\beta_2^{\sup}(\check{z}))\right)}{\bar{F}_n\left(\tau/\mu\sqrt{\xi_n}\right)}\phi(\check{z}) \tag{165}$$

and let $h_n(\check{z}) = f_{n+n_0-1}(\check{z})$ for some $n_0 \ge 0$. Therefore

$$\lim_{n \to \infty} \sup_{\substack{\theta \in \Theta_{01}^* \\ \rho \in P^*}} \frac{\Pr\left(\operatorname{PO}_{\{1\}} \ge \tau \;;\; \theta\right)}{\Pr\left(\chi_1^2 \ge 2\log\left(\frac{\tau}{\mu\sqrt{\xi_n}}\right)\right)} \le \lim_{n \to \infty} \int_{-\infty}^{\infty} f_n(\check{z}) \; d\check{z}$$
$$= \lim_{n \to \infty} \int_{-\infty}^{\infty} h_n(\check{z}) \; d\check{z} \tag{166}$$

We assert it can be shown that $\check{\iota}_n(\check{z}, \beta_2^{\sup}(\check{z}))$ and therefore $f_n(\check{z})$ is 'ultimately' monotonically decreasing with n. That is, there exists a particular n_0 (yet to be defined precisely), for which the sequence $h_n(z)$ is monotonically decreasing with n.

Since \overline{F}_n is a regularly varying function at $n \to \infty$ with tail index -1, by the theory on Potter bounds (see Theorem 11 of Kevei [2019]), there exists a $\delta > 0$ and a $A_{\delta} > 1$, such that

$$\frac{F_n(x)}{F_n(y)} \le A_{\check{\delta}} \left(\frac{x}{y}\right)^{-(1+\delta)} \qquad (\forall x < y)$$
(167)

Therefore

$$h_{n}(\check{z}) = \frac{\bar{F}_{n+n_{0}-1}\left(\tau/\mu\sqrt{\xi_{n}}\,\check{\iota}_{n+n_{0}-1}(\check{z},\beta_{2}^{\sup}(\check{z}))\right)}{\bar{F}_{n+n_{0}-1}\left(\tau/\mu\sqrt{\xi_{n}}\right)}\phi(\check{z})$$

$$\leq A_{\check{\delta}}\,\check{\iota}_{n+n_{0}-1}(\check{z},\beta_{2}^{\sup}(\check{z}))^{-(1+\check{\delta})}\,\phi(\check{z})$$

$$\leq A_{\check{\delta}}\,\check{\iota}_{n_{0}}(\check{z},\beta_{2}^{\sup}(\check{z}))^{-(1+\check{\delta})}\,\phi(\check{z})$$

$$=: g(\check{z}) \quad \text{(by definition)} \tag{168}$$

The Lebesgue dominated convergence theorem [Van der Vaart and Yen, 1968] states if $h_n : \mathbb{R} \mapsto [-\infty, \infty]$ are (Lebesgue) measurable functions such that the pointwise limit $h(x) := \lim_{n \to \infty} h_n(\check{z})$ exists and there is an integrable $g : \mathbb{R} \mapsto [0, \infty]$ with $|h_n(\check{z})| \leq g(\check{z})$ for all n and all $\check{z} \in \mathbb{R}$, then

$$\lim_{n \to \infty} \int_{-\infty}^{\infty} h_n(\check{z}) \, d\check{z} = \int_{-\infty}^{\infty} h(\check{z}) \, d\check{z}$$
(169)

Notice first that $g(\check{z}) = A_{\check{\delta}} \check{\iota}_{n_0}(\check{z}, \beta_2^{\sup}(\check{z}))^{-(1+\check{\delta})} \phi(\check{z})$ is integrable; and second that

$$h(x) := \lim_{n \to \infty} h_n(\check{z}) = \lim_{n \to \infty} f_n(\check{z}) = \phi(\check{z})$$
(170)

Therefore Equation 166 can be rewritten as

$$\lim_{n \to \infty} \sup_{\substack{\theta \in \Theta_{01}^* \\ \rho \in P^*}} \frac{\Pr\left(\operatorname{PO}_{\{1\}} \ge \tau ; \theta\right)}{\Pr\left(\chi_1^2 \ge 2\log\left(\frac{\tau}{\mu\sqrt{\xi_n}}\right)\right)} \le \int_{-\infty}^{\infty} \phi(\check{z}) \, d\check{z} = 1$$
(171)

Equations 146 and 171 cover all cases, showing that in generality

$$\lim_{n \to \infty} \sup_{\substack{\theta \in \Theta_{\tilde{s}}^* \\ \rho \in P^*}} \frac{\Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau ; \theta\right)}{\Pr\left(\chi_1^2 \ge 2\log\frac{\tau}{|\mathcal{V}|\mu\sqrt{\xi_n}}\right)} \le 1 \qquad (\forall \tilde{s})$$
(172)

Or equivalently,

$$\lim_{n \to \infty} \sup_{\substack{\theta \in \omega_v^* \\ \rho \in P^*}} \frac{\Pr\left(\operatorname{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} \ge \tau; \theta\right)}{\Pr\left(\chi_1^2 \ge 2\log\frac{\tau}{|\mathcal{V}| \mu \sqrt{\xi}}\right)} \le 1,$$
(173)

where $v_j = I(j \notin \mathcal{V})$.

4. The $\alpha \leq 0.025$ threshold in Doublethink

This section explains MT Remark 1. We consider the test statistic

$$\tilde{\text{PO}}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}} = \sum_{j\in\mathcal{V}} \text{PO}_{\tilde{\boldsymbol{s}}+\boldsymbol{e}_{j}:\tilde{\boldsymbol{s}}}, \qquad (174)$$

which is asymptotically equivalent (see proof of Theorem 2) to the posterior odds, $\text{PO}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}$, as $n \to \infty$ (Definition 10), where \tilde{s} is the 'true' model and $\{e_j\}_k = I(j = k)$. It tests the null hypothesis that $\beta_j = 0$ for all $j \in \mathcal{V}$.




FIGURE 9. Tail probabilities for the mean of $k \operatorname{LG}(1/2,1)$ random variables for $k = 1, \ldots, 10$ based on 10 million simulations (solid coloured lines) and the theory of regular variation (Equation 178; dashed coloured lines). Grey lines indicate the smallest x at which the tail probability $\Pr(\bar{X}_k \geq x)$ is greatest at k = 1.

We rewrite it as

$$\frac{\widetilde{\mathrm{PO}}_{\mathcal{A}_{\mathcal{V}}:\mathcal{O}_{\mathcal{V}}}}{|\mathcal{V}|\,\mu\,\sqrt{\xi}} = \frac{1}{|\mathcal{V}|} \sum_{j\in\mathcal{V}} R^{1-\xi}_{\tilde{\boldsymbol{s}}+\boldsymbol{e}_{j}:\tilde{\boldsymbol{s}}} \sim \frac{1}{|\mathcal{V}|} \sum_{j\in\mathcal{V}} R_{\tilde{\boldsymbol{s}}+\boldsymbol{e}_{j}:\tilde{\boldsymbol{s}}}, \tag{175}$$

as $n \to \infty$, where $\xi = h/(n+h)$.

Under independence, when \tilde{s} is true, this problem is equivalent to studying

$$\bar{X}_k = \frac{1}{k} \sum_{j=1}^k X_j,$$
(176)

where $2 \log X_j$, $j = 1 \dots k$ are independent and identically distribution chi-squared random variables with one degree of freedom. Equivalently, X_j , $j = 1 \dots k$ are independent and identically distributed log-gamma random variables with shape parameter 0.5 and scale parameter 1.

Theorem 2 and Corollary 1 are equivalent to the asymptotic approximation

$$\Pr(\bar{X}_k \ge x) \sim \Pr(\chi_1^2 \ge 2\log x), \quad x \to \infty.$$
 (177)

Whereas direct application of the theory of regular variation [Karamata, 1933] would have

$$\Pr(\bar{X}_k \ge x) \sim k \Pr(\chi_1^2 \ge 2\log k x), \qquad x \to \infty.$$
(178)

See, for example, Mikosch [1999] Corollary 1.3.6.

We used simulations to investigate whether the approximation (Equation 177) is conservative, under independence, for all k at all probabilities less than 0.0259846 (Figure 9) in the sense that

$$\Pr(\bar{X}_1 \ge x) \ge \Pr(\bar{X}_k \ge x), \quad \forall \quad k \ge 1, \ x \ge x_{\text{crit}}.$$
(179)

This is apparent in the graph because the red line (k = 1) is above all other coloured lines (k > 1) when $x \ge x_{\text{crit}}$.

We solved x_{crit} numerically based on the following empirical observation: the solution to $\Pr(X_1 \ge x) = \Pr(\bar{X}_k \ge x), x > 1$, was solved at increasingly smaller values of x as k increased. This is apparent in the graph because the red line (k = 1) crosses the pink line (k = 10) at smaller values of x than the purple line (k = 9), and so on.

Therefore it appeared to be sufficient to solve, numerically, the convolution

$$\Pr(\bar{X}_{1} \ge x_{\text{crit}}) = \Pr(\bar{X}_{2} \ge x_{\text{crit}}), \quad x_{\text{crit}} > 1$$

= $\int_{1}^{\infty} p_{\text{LG}(0.5,1)}(y) \Pr(X_{1} + y \ge 2x_{\text{crit}}) dy,$ (180)

where $p_{LG(0.5,1)}$ is the density function of a log-gamma random variable with shape parameter 0.5 and scale parameter 1. This yielded

$$x_{\text{crit}} = 11.92362,$$

 $\Pr(\bar{X}_1 \ge x_{\text{crit}}) = 0.0259846.$ (181)