

A high resolution rovibronic molecular cross-section of MgH⁺ molecular cation

Huagang Xiao^a and Tao Gao^{a*}

^a*Institute of Atomic and Molecular physics, Sichuan University, Chengdu 610065, China*

E-mail: gaotao@scu.edu.cn

ABSTRACT: The high resolution rovibronic line list of MgH⁺ molecular cation are presented in our work. The potential energy curves are calculated by the method of multireference configuration interaction plus Davidson correction (MRCI+Q) and spin-orbit coupling (SOC) effect. Spectroscopy constants are fitted and the results are in good agreement with the experiment, ensuring the accuracy of the electronic structure. On account of potential energy curves and transition dipole moments, the Franck - Condon factors and Einstein coefficients of transition are obtained. These calculations are used to obtain an accurate partition functions and line list for the molecule. Using the data obtained from the *ab initio* calculation, the absorption cross-sections under different temperatures and pressures were simulated. Our work could provide some theoretical insights into solar and cold planet spectrum.

* Corresponding author: Institute of Atomic and Molecular physics, Sichuan University, Chengdu, China

Contents

1. Introduction	1
2. Method	2
2.1 Electronic structure and transition property	2
2.2 Spectral simulation	3
3. Results	4
3.1 Electronic structure and ro-vibrational energy levels	4
3.2 Transition properties	8
3.3 Temperature-dependent partition function	11
3.4 Simulated spectra	13
4. Calculation	15
Acknowledgment.	15
Reference	15

1. Introduction

Mg is the ninth most abundant element in the universe¹ and MgH is an important astrophysical molecule²⁻⁴. MgH⁺ molecular ions may play a role in astrophysics by existing as intermediates (e.g, OH⁺⁵ and HeH⁺^{6, 7} molecular cation). As early as thirty years ago, Singh⁸ have reported that MgH⁺ is solar molecule, and its A-X transition is of great significance to the spectroscopic research of the sun and cold planets. In addition, Pearse⁹ observed the emission spectrum of MgH molecule and found a extensive system of red-degraded bands between 250 nm and 340 nm, which correctly attributed it to the ¹S - ¹S transition of the MgH⁺. Therefore, it is necessary for astrophysics to study MgH⁺ molecular ions.

The MgH⁺ molecular cation was first reported by Pearse et al. The γ -system spectrum in the Ultra- Violet spectrum of MgH molecules is the ionization of MgH⁺ ion. In 1976, Singh and Maciel¹⁰ et al discussed the possible existence of MgH⁺ molecular cation in the sun. Since then, researchers have continued to discuss this topic. However, whether MgH⁺ molecular ions exist in the solar spectrum requires the location of the line positions and dissociation energies. Therefore, it is meaningful to study the line list of MgH⁺ molecular cation using use high precision calculations. The accuracy of the molecular electronic structure is the primary guarantee for ensuring the spectrum. In 1971, Based on the experimental data, Balfou¹¹ used RKR method to conduct rotational analysis of the A¹ Σ^+ - X¹ Σ^+ and B¹ Π - X¹ Σ^+ transitions, and the partial energy level values and Franck - Condon factors of the molecule is obtained. Dalleska *et al.*¹² used Guided ion beam mass spectrometry to investigate the reaction of ground state Mg⁺ and Sr⁺ with molecular hydrogen and its isotopes, and determined the dissociation energy(1.94 ± 0.06 eV) of MgH⁺ molecular ions. In 2005, Jørgensen and Drewsen¹³ used the multiconfiguration self-consistent field (MCSCF) approach to calculate the four electronic states (X¹ Σ , A¹ Σ , B¹ Π and C¹ Σ) of the MgH⁺, and compared the rotational energy and dissociation energy with the experimental values of Balfou et al. It is noticed that the energy of the low level is in good agreement with the experimental value, but the energy of the high level has a

certain deviation from the experiment. Four years later, Aymar *et al*¹⁴ used different basis sets to calculate the potential energy curves and dipole moments of the MgH⁺ molecular ion, which showed that large basis sets could more accurate. In 2013, Khemiri *et al*¹⁵ used *ab initio* theory to extend the research of aymer et al and calculated the high excited electronic state of MgH⁺ molecular ion.

The study of MgH⁺ molecular ions is currently incomplete, such as the consideration of spin-orbit coupling effects. Our goal is to provide a high resolution line list of MgH⁺ molecular cation based on accurate electronic structures, which could increase theoretical insights for astrophysical applications.

2. Method

2.1 Electronic structure and transition property

The electronic structure and transition dipole moments are calculated using Molpro 2012 program¹⁶. The complete active space self-consistent^{17, 18} (CASSCF) and multireference configuration interaction plus Davidson corrections^{19, 20} (MRCI+Q) method are employed in the calculation. To improve accuracy, the correlation consistent polarized valence Quintuple - ζ basis set (AV5Z) is applied for Mg²¹ and H²² atoms. The diatomic molecule MgH⁺ belongs to $C_{\infty V}$ group, However, the C_{2v} subgroup is adopted limited by the Molpro code. The C_{2v} group contain four irreducible representations (a_1, b_1, b_2, a_2), which is related to the electronic state and orbital of the molecule(e.g, $\Sigma^+ = a_1, \Pi = b_1 + b_2, \Sigma^- = a_2$). In our calculation, the 1s2s2p orbital electron of Mg atom are placed into frozen space and the 3s3p orbital of and 1s orbital of H are putted into active space. Considering the calculation of excited states, electrons are excited from lower orbits to higher orbits. The 4s4p orbital of Mg atom also considered into active space. Hence, the total spaces of MgH⁺ molecular cation are expressed as ($8a_1, 3b_1, 3b_2, 0a_2$), which could ensured the accurate calculation of six $\Lambda - S$ states and eleven Ω states. Using the LEVEL program²³, the spectroscopic constants of electronic states are obtained.

Combining the potential energy curves and the transition dipole moments, the

transition properties of the molecular electronic state could be obtained. The Franck - Condon factor $f_{v'v''}$ and Einstein spontaneous emission coefficient $A_{v'v''}$ are described by the follows:

$$f_{v'v''} = \left| \int \Psi_{v'J'} \Psi_{v''J''} dr \right|^2 \quad (1)$$

$$A_{v'v''} = 3.1361891 \times 10^{-7} \frac{S(J', J'')}{2J'+1} \nu^3 \langle \Psi_{v',J'} | M(r) | \Psi_{v'',J''} \rangle \quad (2)$$

Where $v'J'$ and $v''J''$ represents upper level and lower level, $\Psi_{v'J'}$ and $\Psi_{v''J''}$ is wave functions for upper and lower levels, $M(r)$ is transition dipole moments and $S(J', J'')$ is the Hönl-London rotational intensity factor of levels.

2.2 Spectral simulation

Here, a brief description of the molecular absorption spectrum calculation is shown below. First, solving the radial Schrödinger equation of the electronic state, the vibrational level and rotational level of the molecule is obtained²⁴.

$$-\frac{\hbar^2}{2\mu} \frac{d^2 \Psi_{vJ}(r)}{dr^2} + V_J(r) \Psi_{vJ}(r) = E_{vJ} \Psi_{vJ}(r) \quad (3)$$

$$E_{vJ} = G(v) + B_v[J(J+1) - \Omega^2] - D_v[J(J+1) - \Omega^2]^2 \dots \quad (4)$$

Where E_{vJ} is the energy of ro-vibational level, $G(v)$ is energy of vibrational level, the B_v and D_v are centrifugal distortion constants.

In view of specific temperature, the temperature - dependent partition function Q (T) of the ro - vibrational energy level of the molecule is²⁵:

$$Q_{evJ} = \sum_e \sum_{v=0}^{v_{\max}} \sum_{J=\Lambda}^{J_{\max}} g_J \times \exp \left\{ -\frac{hc}{kT} [T_e + G(v) + F_v(J) - E_0] \right\} \quad (5)$$

Where T_e is vertical transition energy of electronic states, $G(v)$ is vibrational level energy, $F(v)$ is rotational level energy, E_0 is reference energy. k , h and c represent Boltzmann's constant coefficient, Planck's coefficient and the speed of light, respectively. g_J represents the degeneracy of the molecular level, which has to do with

total spin and nuclear spin of the molecule. For MgH^+ molecular cation, the nuclear spin of Mg and H atoms are 0 and 1/2, respectively. Hence, with regard to the all ro-vibrational levels, the total molecular spin is 1/2 and the spin multiplicity g_N is 2. Integrated intensity S_{ul} of each electronic vibration rotational transition spectral line:

$$S_{ul} = \frac{1}{8\pi c} (E_u - E_l)^{-2} \frac{g_u e^{-\frac{hc}{kT} E_l} \left[1 - e^{-\frac{hc}{kT} (E_u - E_l)} \right]}{Q(T)} A_{ul} \quad (6)$$

The absorption cross section σ_{ul} is closely related to the intensity S_{ul} ,

$$S_{ul} = \int_{-\infty}^{\infty} \sigma_{ul}(v) dv \quad (7)$$

In above formula, g_u represents the degeneracy of the higher level, which is related to the nuclear spin. E_u and E_l are represent the energy of higher and lower levels, respectively.

3. Results

3.1 Electronic structure and ro-vibrational energy levels

Molecular structure and properties could be described by potential energy curves (PECs). The PECs of six $\Lambda - S$ states were depicted in Fig. 1(a), which including four bond states ($X^1\Sigma^+$, $A^1\Sigma^+$, $B^1\Pi$ and $b^3\Pi$) and two repulsed states ($a^3\Sigma^+$ and $c^3\Sigma^+$). With the spin orbital couple effect (SOC), six $\Lambda - S$ states splitted into eleven Ω states and the result are shown in Fig. 1(b). Meanwhile, the Mg^+ (2P_u) splitted two energy levels ($^2P_{1/2}$ and $^2P_{3/2}$) with an energy level difference of 69.2 cm^{-1} , which is good agreement with the experiential value²⁶ (91.5 cm^{-1}).

Table 1. The dissociation relationships of Ω states

Atomic states	Ω states	Cal. energy (cm^{-1})	Expt. Energy ²⁶ (cm^{-1})
$\text{Mg}^+ (^2S_{1/2}) + \text{H} (^2S_{1/2})$	$X^1\Sigma^+_{0+}, a^3\Sigma^+_{0-}, a^3\Sigma^+_1$	0	0
$\text{Mg}^+ (^2P_{1/2}) + \text{H} (^2S_{1/2})$	$A^1\Sigma^+_{0+}, b^3\Pi_0^-, b^3\Pi_1$	34482.5	35669.3
$\text{Mg}^+ (^2P_{3/2}) + \text{H} (^2S_{1/2})$	$b^3\Pi_0^+, b^3\Pi_2, B^1\Pi_1, c^3\Sigma^+_{0-}, c^3\Sigma^+_1$	34552.1	35760.8

The spectroscopic parameters of those electronic states were fitted and were listed in Table 2. The $X^1\Sigma^+$, $A^1\Sigma^+$ and $B^1\Pi$ bond states are no splitting and it is notable that spectroscopic parameters of those states are changed little between spin-free and SOC effect. The ground state $X^1\Sigma^+$ possess a deep potential well of 16955.9 cm^{-1} , which is closer to experimental data²⁷ (16937 cm^{-1}) than other theoretical values. In addition, it found that the equilibrium length R_e , harmonic constant ω_e and rotational constant B_e of $X^1\Sigma^+$ state is better match the experimental data than other theoretical calculation. And this phenomenon also occur at the excited states $A^1\Sigma^+$ and $B^1\Pi$. The electronic state $b^3\Pi$ lying 46561.3 cm^{-1} above ground state $X^1\Sigma^+$ with a shallow potential well 4882.2 cm^{-1} (0.605 eV). The equilibrium length R_e of $b^3\Pi$ is 2.030 \AA , which is close to the data ($R_e = 2.032\text{ \AA}$) calculated by khemiri *et al* using effective core potential (ECP) + core polarization (CPP) + full configuration interaction (FCI) approach. Aymar¹⁴ *et al* also employed ECP + CPP + FCI method to calculate the electronic properties of MgH^+ molecular cation. The spectroscopic parameters of $b^3\Pi$ are far from what our calculated, however, the D_e , R_e and ω_e of $c^3\Sigma^+$ state reported by Aymar *et al* are agreement with the values of $b^3\Pi$, which may be an wrote error for them. With the SOC effect, $b^3\Pi_0^-$, $b^3\Pi_0^+$, $b^3\Pi_1$ and $b^3\Pi_2$ are splitted from $b^3\Pi$ state. Notably, the split strength of $b^3\Pi$ is inconspicuous and only 42.8 cm^{-1} energy gap generated between the $b^3\Pi_0^-$ and $b^3\Pi_2$ electronic states.

Comparing with the other theoretical calculation and experiential data, the ability of our calculation to accurately predict the electronic structure and spectroscopic constants and indicate that our present results are most sophisticated and accurate work done for the low-lying states of MgH^+ molecular cation.




Fig. 1 The potential energy curves of MgH^+ molecular cation. (a) the Λ - S states for two channels, (b) the Ω states for three channels with spin-orbital coupling effect.

Table 2. Spectroscopic parameters of the bond states for MgH^+ molecular cation from MRCI+Q+SOC calculations compared to other Theoretical and Experimental results

	D_e/cm^{-1}	T_e/cm^{-1}	$R_e/\text{\AA}$	ω_e/cm^{-1}	B_e/cm^{-1}	Reference
$X^1\Sigma^+$	16955.9	0	1.665	1677.4	6.2685	MRCI+Q
$X^1\Sigma^+_{0+}$	16955.9	0	1.665	1684.1	6.2895	MRCI+Q+SOC
	15650±500					Expt. ¹²
	16937	0	1.649	1695.3	6.411	Expt. ²⁷
	16501	0	1.635	1599		Thery ¹⁴
	14755	0	1.667	1542		Thery ¹⁵
$A^1\Sigma^+$	15837.4	35433.5	2.050	1109.6	4.1645	MRCI+Q
$A^1\Sigma^+_{0+}$	15832.4	35433.4	2.050	1115.6	4.1641	MRCI+Q+SOC
		35905	2.006	1132.7	4.330	Expt. ²⁷
	16656		2.005	1125		Thery ¹⁴
	15666	37689.1	2.001	1129		Thery ¹⁵
$B^1\Pi$	1874.1	49593.0	2.300	557.2	3.3788	MRCI+Q
$B^1\Pi_1$	1892.9	49593.2	2.300	555.7	3.3788	MRCI+Q+SOC
	1943		2.27	588	3.39	Thery ¹⁴
	1769	53009.4	2.28	549		Thery ¹⁵
$b^3\Pi$	4882.2	46561.3	2.030	951.4	4.2223	MRCI+Q

	5057		2.005	915		Thery ¹⁴ ($c^3\Sigma^+$)
	79		4.905	70		Thery ¹⁴ ($b^3\Pi$)
	4654		2.032	83		Thery ¹⁵
$b^3\Pi_0^-$	4868.5	46539.8	2.030	952.6	4.2217	MRCI+Q+SOC
$b^3\Pi_0^+$	4960.7	46539.9	2.030	921.7	4.2217	MRCI+Q+SOC
$b^3\Pi_1$	4850.4	46561.1	2.030	956.1	4.2215	MRCI+Q+SOC
$b^3\Pi_2$	4883.9	46582.6	2.030	948.3	4.2214	MRCI+Q+SOC

Fine molecular structure could bring out accurate energy level. Solving the one-dimensional Schrödinger equation, the vibrational levels and rotational levels of the ground and excited states are obtained. The radiative transition of $\Omega = 0 - 0$ is strongest, therefore, our interest mainly focus on 0 - 0 transition and the parameters of energy level for $X^1\Sigma^+_{0+}$, $A^1\Sigma^+_{0+}$ and $b^3\Pi_0^+$ states are listed in Table. 3. The ground state $X^1\Sigma^+_{0+}$ own 17 vibrational levels and the rotational level $J \leq 69$. The excited state $A^1\Sigma^+_{0+}$ also presented deep well, corresponding to a maximum vibrational quantum number ν_{max} of 19 and a maximum rotational quantum number J_{max} of 90. Notably, the gap of vibrational energy levels for different electronic state are not alike, which is due to the distinction of harmonic constant ω_e . Partial vibrational level energy are showed in Tables 3 and All rovibronic level energies are attached to the Support material. Compared with experimental data, it could found that the energy value is good agreement with experiment.

Table. 3 the vibrational quantum parameters of energy level for MgH^+ molecular cation

Electronic states	vibrational quantum number ν	E_{cal}	$E_{expt.}^{11}$
$X^1\Sigma^+_{0+}$	0	841.318	841.57
	1	2473.976	2476.13

	17	16955.954	16937
$A^1\Sigma^+_{0+}$	0	547.485	565.85

	1	1627.996	1684.81

	19	15837.419	15920
$b^3\Pi_0^+$	0	443.317	
	1	1277.759	
	
	10	4960.709	

Notes.

ν : vibrational quantum number

J : rotational quantum number

E : energy of levels/ cm^{-1}

g_N : the degenerate of levels

3.2 Transition properties

It is particularly important to understand the transition dipole moments (TDMs) when investigating the transition strength between the various electronic states. Since the $b^3\Pi \rightarrow X^1\Sigma^+$ band is forbidden, the SOC effect is considered in our calculation. The DMs of $X^1\Sigma_{0+}^+ \rightarrow X^1\Sigma_{0+}^+$, $A^1\Sigma_{0+}^+ \rightarrow X^1\Sigma_{0+}^+$ and $b^3\Pi_0^+ \rightarrow X^1\Sigma_{0+}^+$ transitions are shown in Fig. 2. The DM of $X^1\Sigma_{0+}^+ \rightarrow X^1\Sigma_{0+}^+$ transition is permanent dipole moments of the ground state $X^1\Sigma_{0+}^+$ and the max value is 1.472 a.u (3.74 Debye) when the equilibrium bond length R is 1.4 Å. The transition dipole moments (TDMs) of $A^1\Sigma_{0+}^+ \rightarrow X^1\Sigma_{0+}^+$ first increased to a maximum value of 2.1 a.u (5.34 Debye) and then slowly decreased. However, compared with the first excited state $A^1\Sigma_{0+}^+$, the TDM values of $b^3\Pi_0^+ \rightarrow X^1\Sigma_{0+}^+$ transition always near to zero in the Franck - Condon region.




Fig. 2 Transition dipole moments between $X^1\Sigma^+_{0+}$, $A^1\Sigma^+_{0+}$ and $b^3\Pi^+_{0+}$ states of MgH^+ molecular cation.

Franck - Condon factors and Einstein coefficients are importance characterization parameters for our investigation of transition properties. Relying on the potential energy curves and dipole moments, the Franck - Condon factors f_{ul} and Einstein coefficients A_{ul} of $A^1\Sigma^+_{0+} - X^1\Sigma^+_{0+}$ and $b^3\Pi^+_{0+} - X^1\Sigma^+_{0+}$ transitions are obtained using LEVEL program and the results are listed in Table. 4. It is notice that the f_{ul} not show a diagonal trend due to the difference in equilibrium bond length R between the excited states ($A^1\Sigma^+_{0+}$ and $b^3\Pi^+_{0+}$) and the ground state ($X^1\Sigma^+_{0+}$). In 1971, Balfour¹¹ evaluated the FCFs of $A^1\Sigma^+ - X^1\Sigma^+$ transition using RKR method, However, its value is slightly different from our calculation. The reason is because of the error in the equilibrium bond length R_e our calculated (our value: $R_e(X^1\Sigma^+_{0+})$ is 1.665 Å, Balfour's data: $R_e(X^1\Sigma^+_{0+})$ is 1.652 Å). It could found that the Einstein coefficients A_{ul} of $A^1\Sigma^+_{0+} - X^1\Sigma^+_{0+}$ transition is larger and the values reach $10^7 s^{-1}$, which is owing to the large TDMs. The excited state $b^3\Pi$ is lying 46561.3 cm^{-1} above ground state and the $b^3\Pi - X^1\Sigma^+$ bond is forbidden transition. Considering the spin - orbital coupling (SOC) effect, $b^3\Pi^+_{0+} - X^1\Sigma^+_{0+}$ transition is obtained in our calculation. Similar to the first excited state $A^1\Sigma^+_{0+}$, the max Franck - Condon factor is f_{02} (0.2976). The TDMs of $b^3\Pi^+_{0+} - X^1\Sigma^+_{0+}$ transition is much smaller than $A^1\Sigma^+_{0+} - X^1\Sigma^+_{0+}$ transition, which bring out minor A_{ul} of $b^3\Pi^+_{0+} - X^1\Sigma^+_{0+}$ transition.

Table. 4 The Franck - Condon factors f_{ul} and Einstein coefficients A_{ul} (s^{-1}) of two transitions for MgH^+ molecular cation.

Transition	$v' = 0$	$v' = 1$	$v' = 2$	$v' = 3$	
$A^1\Sigma^+_{0+} - X^1\Sigma^+_{0+}$	$v'' = 0$	0.0666	0.1531	0.1975	0.1876
		0.091 ^a	0.185 ^a	0.214 ^a	0.185 ^a
		1.84E+7	4.40E+7	5.88E+7	5.77E+7
	$v'' = 1$	0.2277	0.1946	0.0530	0
		0.273 ^a	0.175 ^a	0.025 ^a	0.007 ^a
		5.90E+7	5.24E+7	1.47E+7	8650.17
	$v'' = 2$	0.3268	0.0194	0.0520	0.1144
		0.337 ^a	0.002 ^a	0.093 ^a	0.114 ^a
		7.92E+7	4.86E+6	1.39E+7	3.15E+7
	$v'' = 3$	0.2493	0.0811	0.1275	0.0049
		0.216 ^a	0.149 ^a	0.097 ^a	0.002 ^a
		5.64E+7	1.94E+7	3.18E+7	1.22E+6
$b^3\Pi_{0+} - X^1\Sigma^+_{0+}$	$v'' = 0$	0.0806	0.1510	0.1685	0.1475
		58.30	221.79	363.16	382.12
	$v'' = 1$	0.2296	0.1701	0.0501	0.0021
		8.53	123.57	172.55	78.19

$v'' = 2$	0.2976	0.0213	0.0285	0.0806
	163.77	0.21	19.08	3.05
$v'' = 3$	0.2292	0.0449	0.1230	0.0364
	582.62	3.16	24.29	11.26

^a the experiential value from reference [11]

3.3 Temperature-dependent partition function

The values of partition function were calculated in 50 K - 30000 K range, whose temperature could span from low temperature, medium to high temperature to high temperature. Depending on the precise ro - vibrational level energy, the total partition function and proportion of each states are depicted in Fig. 3 and the data are listed in Support material. With the temperature increasing, the partition function Q also gradually increases. However, it could found that the molecular partition functions were almost contributed by the ground state $X^1\Sigma^+_{0+}$ below 8000 K. When the temperature continues to increase, the proportion partition functions of the excited state gradually reveal. For example, the partition function of the ground state $X^1\Sigma^+_{0+}$ and first excited state $A^1\Sigma^+_{0+}$ at temperature $T = 15000$ K calculated are 20874.3 and 1411.2, whose proportion are 89.9% and 6.1%. Moreover, it found that within a temperature of 30000 K, the sum of the partition functions of the molecule is 1, indicating that it is completely sufficient for us to calculate the electronic state of MgH^+ molecular cation.




Fig. 3 the partition function of MgH^+ molecular cation: (a) the total partition function, (b) the proportion of partition function for each electronic states.




Fig. 4 the partition function $Q(T)$ of MgH^+ molecular cation with respect to the rotational quantum number J ($v = 0$) at different temperature.

It clear found that the ground state plays a crucial role under 30000 K. Hence, the partition function of the rotational quantum number J ($v = 0$) for ground state $X^1\Sigma_0^+$ at different temperatures T is shown in Fig. 4. As the temperature increases, the

convergence of the partition with J is slower. When the temperature reaches above 8000 K, the values continue to grow, indicating that more energy levels are involved.

3.4 Simulated spectra

A ro - vibrational line lists of MgH⁺ molecular cation are computed using Duo program²⁸. As before mentioned, the Einstein coefficients of b³Π₀₊ - X¹Σ⁺₀₊ transition is smaller, hence, Our attention is mainly focused on the X¹Σ⁺₀₊ - X¹Σ⁺₀₊ transition and the A¹Σ⁺₀₊ - X¹Σ⁺₀₊ transition. Combined the .states and .trans files of the Duo program and using the EXOCROSS code²⁹, the simulations spectrum of MgH⁺ molecular cation is obtained.




Fig. 5 the absolute absorption line intensities of X¹Σ⁺₀₊ - X¹Σ⁺₀₊ transition for MgH⁺ molecular cation at T = 300 K.

A MgH⁺ stick spectrum of X¹Σ⁺₀₊ - X¹Σ⁺₀₊ transition at T = 300 K is plotted in Fig. 5, which contain the two terms ($\Delta\nu = 0$ and 1). Obviously, the $\Delta\nu = 0$ term in a pure rotational spectrum is stronger than the $\Delta\nu = 1$. In addition, It could found that the P and R branches are clearly visible in $\Delta\nu = 1$ term.

In order to make the research more systematic, the controlled variable method is adopted to investigate the molecular spectrum. First, The absorption cross sections of MgH⁺ molecular ions under different pressures at temperature T = 300 K are shown in Fig. 6. Under the same pressure(eg, 1.0 bar), the intensity of X¹Σ⁺₀₊ - X¹Σ⁺₀₊ is 10⁻³

orders of magnitude smaller than the intensity of $A^1\Sigma^+_{0+} - X^1\Sigma^+_{0+}$. The max value of cross-section of $X^1\Sigma^+_{0+} - X^1\Sigma^+_{0+}$ transition is 2.79×10^{-19} $\text{cm}^2/\text{molecule}$, which represents the transition of 0 - 0 band. However, for the $A^1\Sigma^+_{0+} - X^1\Sigma^+_{0+}$ transition, the max cross-section is 2.45×10^{-16} $\text{cm}^2/\text{molecule}$, which is generated by the 3 - 0 band. Furthermore, with the temperature remains constant and pressure increases, the peak shape of the spectrum remains unchanged, but the intensity generally decreases. And then, holding the pressure constant, the effect of temperature on the spectrum is explored and the results are presented in Fig. 7. As the temperature increases, there are more levels contribute to the absorption spectrum, which making its spectral distribution wider. Notably, with the temperature increases, the transition spectrum undergoes a shift.




Fig. 6 Absorption cross-sections of MgH^+ molecular cation under different pressures at $T = 300$ K. (a: $X^1\Sigma^+_{0+} - X^1\Sigma^+_{0+}$ transition, b: $A^1\Sigma^+_{0+} - X^1\Sigma^+_{0+}$ transition)




Fig. 7 the spectrum of MgH^+ molecular cation in various temperature at $P = 1.0$ bar. (a: $X^1\Sigma^+_{0+} -$

$X^1\Sigma^+_{0+}$ transition, b: $A^1\Sigma^+_{0+} - X^1\Sigma^+_{0+}$ transition)

4. Calculation

Here, a detailed and systematic *ab initio* spectroscopic investigation of the MgH^+ molecular cation is presented for the first time. It is crucial for spectroscopic studies of the Sun and cool stars. The potential energy curves are constructed using multireference configuration interaction and Davidson correction (MRCI+Q) method. Spin - orbital coupling effect is considered in our calculation. The spectroscopic constants we obtained are in well agreement with the experiments, and the errors are all around 1%, which could ensure the accuracy of the ro - vibrational levels. Using potential energy curves and transition dipole moments, The line list of the MgH^+ molecular cation was calculated, and the effects of different temperatures and pressures on the high-temperature spectrum were explored. The present work was conducted for applications in astrophysics.

Acknowledgment.

Thanks to Key Laboratory of High Energy Density Physics and Technology of Ministry of Education for providing the computing resources for this work.

Reference

1. K. Lodders, H. Palme and H.-P. Gail, *4.4 Abundances of the elements in the Solar System: 4, Solar System*. (2009) 712.
2. P. Weck, A. Schweitzer, P. Stancil, P. Hauschildt and K. Kirby, *The molecular line opacity of MgH in cool stellar atmospheres*, *Astrophys J.* **582** (2003) 1059.
3. N. Afram, S. V. Berdyugina, *Molecules as magnetic probes of starspots*, *Astron Astrophys.* **576** (2015) A34.
4. A. Owens, S. Dooley, L. McLaughlin, B. Tan, G. Zhang, S. N. Yurchenko and J. Tennyson, *ExoMol line lists - XLV. Rovibronic molecular line lists of calcium*

- monohydride (CaH) and magnesium monohydride (MgH), *Mon Not R Astron Soc.* **511** (2022) 5448.
5. J. N. Hodges and P. F. Bernath, *Fourier Transform Spectroscopy of the $A^3\Pi - X^3\Sigma^-$ Transition of OH^+* , *Astrophys J.* **840** (2017) 81.
 6. R. Güsten, H. Wiesemeyer, D. Neufeld, K. M. Menten, U. U. Graf, K. Jacobs, B. Klein, O. Ricken, C. Risacher and J. Stutzki, *Astrophysical detection of the helium hydride ion HeH^+* , *Nature.* **568** (2019) 357.
 7. O. Novotný, P. Wilhelm, D. Paul, A. Kalosi, S. Saurabh, A. Becker, K. Blaum, S. George, J. Göck and M. Grieser, *Quantum-state - selective electron recombination studies suggest enhanced abundance of primordial HeH^+* , *Science.* **365** (2019) 676.
 8. P. Singh, *$^{24}\text{MgH}^+$ in the solar photospheric spectrum* *Science, Astrophys Space Sci.* 102(1984) 287.
 9. R. Pearse, *The ultra-violet spectrum of magnesium hydride. —II. The many-lined γ -system*, *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character*, **125** (1929) 157.
 10. P. Singh and W. Maciel, *Possibility of $^{24}\text{MgH}^+$ in the solar atmosphere-high resolution rotation-vibration spectra*, *Sol Phys.* **49** (1976) 217.
 11. W. Balfour, *Rotational Analysis of the $A^1\Sigma^+ \rightarrow X^1\Sigma^+$ and $B^1\Pi \rightarrow X^1\Sigma^+$ Systems of $^{24}\text{MgH}^+$, $^{25}\text{MgH}^+$, and $^{26}\text{MgH}^+$* , *Can J Phys.* **50** (1972) 1082.
 12. Dalleska N F, Crellin K C, Armentrout P B, *Reactions of alkaline earth ions with hydrogen, deuterium, and hydrogen deuteride*, *J. Phys. Chem.* **13** (1993) 3123.
 13. S. Jørgensen, M. Drewsen and R. Kosloff, *Intensity and wavelength control of a single molecule reaction: Simulation of photodissociation of cold-trapped MgH^+* , *J Chem Phys.* **123** (2005) 9.
 14. M. Aymar, R. Guérout, M. Sahlaoui, O. Dulieu, *Electronic structure of the magnesium hydride molecular ion*, *J Phys B-At mol Opt.* **42** (2009) 154025.
 15. N. Khemiri, R. Dardouri, B. Oujia and F. X. Gadéa, *Ab initio investigation of electronic properties of the magnesium hydride molecular ion*, *J. Phys. Chem. A.* **117** (2013) 8915.
 16. H. J. Werner, P. J. Knowles, G. Knizia, F. R. Manby and M. Schütz, *Molpro: a general - purpose quantum chemistry program package*, *Wires Comput Mol Sci.* **2** (2012) 242.

17. P. J. Knowles and H.-J. Werner, *An efficient second-order MCSCF method for long configuration expansions*, *Chem Phys Lett.* **115** (1985) 259.
18. H. J. Werner and P. J. Knowles, *A second order multiconfiguration SCF procedure with optimum convergence*, *J Chem phys.* **82** (1985) 5053.
19. P. J. Knowles and H.-J. Werner, *An efficient method for the evaluation of coupling coefficients in configuration interaction calculations*, *Chem Phys Lett.* **145** (1988) 514.
20. S. R. Langhoff and E. R. Davidson, *Configuration interaction calculations on the nitrogen molecule*, *Int J Quantum Chem.* **8** (1974) 61.
21. B. P. Prascher, D. E. Woon, K. A. Peterson, T. H. Dunning and A. K. Wilson, *Gaussian basis sets for use in correlated molecular calculations. VII. Valence, core-valence, and scalar relativistic basis sets for Li, Be, Na, and Mg*, *Theor Chem Acc.* **128** (2011) 69.
22. R. A. Kendall, T. H. Dunning Jr and R. Harrison, *Electron affinities of the first - row atoms revisited. Systematic basis sets and wave functions*, *J Chem phys.* **96** (1992), 6796.
23. R. J. Le Roy, *LEVEL: A computer program for solving the radial Schrödinger equation for bound and quasibound levels*, *J Quant Spectrosc Ra.* **186** (2017). 167.
24. G. Herzberg, *Molecular spectra and molecular structure*, D. van Nostrand, 1945.
25. S. S. Penner and R. Landshoff, *Quantitative molecular spectroscopy and gas emissivities*, *Phys Today.* **13** (1959) 38.
26. C.E. Moore, *Atomic Energy Levels*, vol. 1, US Govt Printing Office, Washington, DC, 1971
27. K.-P. Huber, *Molecular spectra and molecular structure: IV. Constants of diatomic molecules*, Springer Science & Business Media, 2013.
28. S. N. Yurchenko, L. Lodi, J. Tennyson and A. V. Stoliarov, *Duo: a general program for calculating spectra of diatomic molecules*, *Comput Phys Commun.* **202** (2016) 262.
29. S. N. Yurchenko, A. F. Al-Refaie, J. Tennyson, *ExoCross: a general program for generating spectra from molecular line lists*, *Astron Astrophys.* **614** (2018) A131.