HOCHSCHILD AND CYCLIC HOMOLOGIES WITH BOUNDED POLES

MASAYA SATO

Abstract. We show that the classical Hochschild homology and (periodic and negative) cyclic homology groups are representable in the category of motives with modulus. We do this by constructing Hochschild homology and (periodic and negative) cyclic homologies for modulus pairs. We show a modulus version of HKR theorem, that is, there exists an isomorphism between modulus Hochschild homology and modulus Kähler differentials for affine normal crossing modulus pairs. By using the representability of modulus Hodge sheaves in the category of motives with modulus, we construct an object of the category of motives with modulus which represents modulus Hochschild homology. Similarly, We compare modulus de Rham cohomology and modulus cyclic homologies and obtain a representability of modulus cyclic homologies.

CONTENTS

1. INTRODUCTION

Let A be a commutative ring. Then there is a trace map from algebraic K -theory $K(A)$ to cyclic homology $HC(A)$, and this map induces an isomorphism

$K_n(A, I) \otimes \mathbb{Q} \cong HC_{n-1}(A, I) \otimes \mathbb{Q}$

for nilpotent ideal I [\[Goo86\]](#page-16-1). Therefore cyclic homology is important to compute algebraic K-theory. Cyclic homology is constructed from Hochschild homology $HH(A)$ as a homotopy orbits $HH(A)_{h\mathbb{T}}$. In this paper, we show Hochschild and (negative, periodic) cyclic homologies are representable in the category of motives with modulus $\text{MDM}_k^{\text{eff}}$.

Voevodsky constructed the triangulated category of motives $\mathbf{DM}_{k}^{\text{eff}}$ over a field k which unifies \mathbb{A}^1 -invariant cohomology theories. His theory has many applications, but it can not be applied to Hochschild and cyclic homologies, because they are not A 1 -invariant. In [\[KMSY21a\]](#page-16-2), [\[KMSY21b\]](#page-16-3), [\[KMSY22\]](#page-16-4), Kahn, Miyazaki, Saito, and Yamazaki constructed a triangulated category $\underline{\text{MDM}}_k^{\text{eff}}$ of motives with

2 MASAYA SATO

modulus which contains $\mathbf{DM}_{k}^{\text{eff}}$ as a full subcategory. Just as $\mathbf{DM}_{k}^{\text{eff}}$ is constructed from smooth schemes X over k, MDM^{eff} is constructed from modulus pairs $\mathcal{X} = (\overline{X}, X^{\infty})$, where \overline{X} is a separated and of finite type k-scheme, and X^{∞} is a effective Cartier divisor of \overline{X} such that $\overline{X} \setminus |X^\infty|$ is smooth over k. When k admits resolution of singularities, e.g. $char(k) = 0$ [Hir₆₄], **MDM**^{eff} can be constructed from *strict normal crossing* modulus pairs $\mathcal{X} = (\overline{X}, X^{\infty})$, that is, \overline{X} is smooth and $|X^{\infty}|$ has strict normal crossing.^{[1](#page-1-0)} An ambient morphism $f : (\overline{X}, X^{\infty}) \to (\overline{Y}, Y^{\infty})$ of modulus pairs is a morphism of schemes $f : \overline{X} \to \overline{Y}$ such that $X^{\infty} \geq f^*Y^{\infty}$. The modulus pair $\overline{\Box} := (\mathbb{P}^1, \infty)$ is called the *cube* and plays a role of unit interval instead of \mathbb{A}^1 .

In [\[KM23a\]](#page-16-6), [\[KM23b\]](#page-16-7), Kelly and Miyazaki constructed objects $\underline{M}\mathcal{O}$ and $\underline{M}\Omega^q$ in $\underline{\mathbf{M}}\mathbf{D}\mathbf{M}_{k}^{\text{eff}}$ such that they represents Zariski and Hodge cohomologies, if char $(k) = 0$. Koizumi did that if k admits resolutions of singularities $[Koi23]$. More precisely:

Theorem 1.1. For any strict normal crossing modulus pair $\mathcal{X} = (\overline{X}, X^{\infty})$, there are isomorphisms

$$
\text{Hom}_{\underline{\mathbf{M}}\mathbf{DM}_{k}^{\text{eff}}}(M(\mathcal{X}), \underline{\mathbf{M}}\mathcal{O}[n]) \cong H_{\text{Zar}}^{n}(\overline{X}, \underline{\mathbf{M}}\mathcal{O}_{\mathcal{X}})
$$

$$
\text{Hom}_{\underline{\mathbf{M}}\mathbf{DM}_{k}^{\text{eff}}}(M(\mathcal{X}), \underline{\mathbf{M}}\Omega^{q}[n]) \cong H_{\text{Zar}}^{n}(\overline{X}, \underline{\mathbf{M}}\Omega_{\mathcal{X}}^{q}),
$$

where $\underline{M}\mathcal{O}_X := \sqrt{\mathcal{I}} \otimes \mathcal{I}^{-1}$, \mathcal{I} is the ideal sheaf corresponds to the effective Cartier divisor X^{∞} , and $\underline{M}\Omega^q_{\mathcal{X}} := \underline{M}\mathcal{O}_{\mathcal{X}} \otimes \Omega^q_{\overline{\mathcal{Y}}}$ $\frac{q}{X}(\log X^{\infty}).$

If A is a smooth k-algebra, then Hochschild homology $HH_n(A/k)$ is isomorphic to the module of Kähler differentials $\Omega_{A/k}^n$, via the so-called HKR theorem, [\[HKR62\]](#page-16-9). Furthermore if k contains \mathbb{Q} ,

$$
HC_n(A/k) \cong (\Omega^n_{A/k}/d\Omega^{n-1}_{A/k}) \times H^{n-2}(\Omega^*_{A/k}) \times H^{n-4}(\Omega^*_{A/k}) \times \cdots
$$

by $[L\text{od}98, \text{Theorem 3.4.12}].$ In this paper, we define Hochschild homology $MHH(A, f)$ for affine modulus pair $(Spec A, (f))$ and show the modulus version of the HKR theorem.

Theorem 1.2 (Theorem [4.11\)](#page-11-0). Let $(Spec A, (f))$ be a strict normal crossing modulus pair such that f is a product of prime elements. Then there is an isomorphism

$$
\underline{M}\Omega^*(A,f) \cong \underline{M}HH_*(A,f).
$$

Furthermore, we define (negative, periodic) cyclic homologies for modulus pairs $MHC(A, f), MHC^{-}(A, f), MHP(A, f)$ and show isomorphims if $char(k) = 0$.

Theorem 1.3 (Theorem [5.2\)](#page-13-0). If (A, f) is strict normal crossing and $f = f_1^{r_1} \cdots f_n^{r_n}$ for prime elements $f_j \in A$, then

$$
\underline{MHC}_n(A,f) \cong (\underline{M}\Omega^n_{(A,f)}/d\underline{M}\Omega^{n-1}_{(A,f)}) \times H^{n-2}(\underline{M}\Omega^n_{(A,f)}) \times H^{n-4}(\underline{M}\Omega^n_{(A,f)}) \times \cdots
$$

\n
$$
\underline{MHC}^-(A,f) \cong \text{Ker}(\underline{M}\Omega^n_{(A,f)} \to \underline{M}\Omega^{n+1}_{(A,f)}) \times H^{n+2}(\underline{M}\Omega_{(A,f)^*}) \times H^{n+4}(\underline{M}\Omega^n_{(A,f)}) \times \cdots
$$

\n
$$
\underline{MHP}_n(A,f) \cong \prod_{p \in \mathbb{Z}} H^{2p-n}(\underline{M}\Omega^*(A,f)).
$$

Finally, we globalize this result and obtain realizations of Hochschild and (negative, periodic) cyclic homologies.

¹cf.[\[KMSY21a,](#page-16-2) Corollary 1.9.5] By the construction of $\text{MDM}_k^{\text{eff}}$, abstract blow-ups become isomorphisms.

Theorem 1.4 (Theorem [6.3\)](#page-15-0). For every strict normal crossing modulus pair $\mathcal{X} =$ $(\overline{X}, X^{\infty})$, we have the following isomorphisms

$$
\begin{aligned} &\mathrm{Hom}_{\underline{\mathbf{M}\mathbf{D}\mathbf{M}^{\mathrm{eff}}_{k}}(M(\mathcal{X}),\prod_{p\in\mathbb{Z}}\underline{\mathbf{M}}\Omega^{p}[p-n])\cong\prod_{p\in\mathbb{Z}}H^{p-n}_{\mathrm{Zar}}(\overline{X},\underline{M}\Omega^{p}_{\mathcal{X}})\cong\underline{M}HH_{n}(\mathcal{X})\\ &\mathrm{Hom}_{\underline{\mathbf{M}\mathbf{D}\mathbf{M}^{\mathrm{eff}}_{k}}(M(\mathcal{X}),\prod_{p\in\mathbb{Z}}\underline{\mathbf{M}}\Omega^{\leq p}[2p-n])\cong\prod_{p\in\mathbb{Z}}\mathbb{H}^{2p-n}_{\mathrm{Zar}}(\overline{X},\underline{M}\Omega^{\leq p}_{\mathcal{X}})\cong\underline{M}HC_{n}(\mathcal{X})\\ &\mathrm{Hom}_{\underline{\mathbf{M}\mathbf{D}\mathbf{M}^{\mathrm{eff}}_{k}}(M(\mathcal{X}),\prod_{p\in\mathbb{Z}}\underline{\mathbf{M}}\Omega^{\geq p}[2p-n])\cong\prod_{p\in\mathbb{Z}}\mathbb{H}^{2p-n}_{\mathrm{Zar}}(\overline{X},\underline{M}\Omega^{\geq p}_{\mathcal{X}})\cong\underline{M}HC_{n}^{-}(\mathcal{X})\\ &\mathrm{Hom}_{\underline{\mathbf{M}\mathbf{D}\mathbf{M}^{\mathrm{eff}}_{k}}(M(\mathcal{X}),\prod_{p\in\mathbb{Z}}\underline{\mathbf{M}}\Omega^{\bullet}[2p-n])\cong\prod_{p\in\mathbb{Z}}\mathbb{H}^{2p-n}_{\mathrm{Zar}}(\overline{X},\underline{M}\Omega^{ \bullet}_{\mathcal{X}})\cong\underline{M}HP_{n}(\mathcal{X}). \end{aligned}
$$

As a corollary, they satisfy blow-up invariance, and $\overline{\Box}$ -invariance. Furthermore, if the divisor is empty , we can recover the classical groups.

Corollarry 1.5. For any smooth, separated and of finite type k -scheme X , by taking $\mathcal{X} = (X, \emptyset)$ in Theorem [1.4,](#page-1-1) we have the following isomorphisms

$$
\begin{aligned} \operatorname{Hom}_{\operatorname{\mathbf{\underline{M}DM}}_{k}^{\operatorname{eff}}}(M(X,\varnothing),\prod_{p\in\mathbb{Z}}\operatorname{\mathbf{\underline{M} \Omega}}^{p}[p-n])&\cong \prod_{p\in\mathbb{Z}}H_{\operatorname{Zar}}^{p-n}(X,\Omega_{X}^{p})\cong HH_{n}(X)\\ \operatorname{Hom}_{\operatorname{\mathbf{\underline{M}DM}}_{k}^{\operatorname{eff}}}(M(X,\varnothing),\prod_{p\in\mathbb{Z}}\operatorname{\mathbf{\underline{M} \Omega}}^{ \leq p}[2p-n])&\cong \prod_{p\in\mathbb{Z}}\mathbb{H}_{\operatorname{Zar}}^{2p-n}(X,\Omega_{X}^{\leq p})\cong HC_{n}(X)\\ \operatorname{Hom}_{\operatorname{\mathbf{\underline{M}DM}}_{k}^{\operatorname{eff}}}(M(X,\varnothing),\prod_{p\in\mathbb{Z}}\operatorname{\mathbf{\underline{M} \Omega}}^{ \geq p}[2p-n])&\cong \prod_{p\in\mathbb{Z}}\mathbb{H}_{\operatorname{Zar}}^{2p-n}(X,\Omega_{X}^{\geq p})\cong HC_{n}^{-}(X)\\ \operatorname{Hom}_{\operatorname{\mathbf{\underline{M}DM}}_{k}^{\operatorname{eff}}}(M(X,\varnothing),\prod_{p\in\mathbb{Z}}\operatorname{\mathbf{\underline{M} \Omega}}^{ \bullet}[2p-n])&\cong \prod_{p\in\mathbb{Z}}\mathbb{H}_{\operatorname{Zar}}^{2p-n}(X,\Omega_{X}^{\bullet})\cong HP_{n}(X). \end{aligned}
$$

2. Acknowledgement

The author would like to thank Shane Kelly for proposing and supporting this research.

3. Modulus de Rham Complex

We will construct the modulus de Rham complex $\underline{M}\Omega^{\bullet}$ in the category of motives with modulus $\underline{\mathbf{M}}\mathbf{D}\mathbf{M}_{k}^{\text{eff}}$. If $\mathcal{X} = (\text{Spec}A,(f))$, then global sections of $\underline{\mathcal{M}}\mathcal{O}_{\mathcal{X}}$ and $\underline{M}\Omega^q_X$ are the following sub-A-modules of $A[f^{-1}]$ and $\Omega^q_{A[f^{-1}]}$.

Definition 3.1 ([\[KM23a,](#page-16-6) Definition 3.1], [\[KM23b,](#page-16-7) Definition 3.1]). For a k-algebra A, and a nonzero divisor $f \in A$, define

$$
\underline{M}\mathcal{O}(A,f) := \frac{\sqrt{(f)}}{f} \subseteq A[f^{-1}],
$$

 $\underline{P}\Omega_{(A,f)}^*$ to be the smallest sub-dg-algebra of $\Omega_{A[f^{-1}]}^*$ containing $A \,=\, \Omega_{A/k}^0\,$ and $d \log a := \frac{da}{a}$ for $a \in A \cap A[f^{-1}]^*$ and

$$
\underline{M}\Omega^q:=\underline{M}\mathcal{O}(A,f)\cdot \underline{P}\Omega^q_{(A,f)}\subseteq \Omega^q_{A[f^{-1}]}.
$$

We will construct the de Rham differential $d: \underline{\mathbf{M}}\Omega^q_{\mathcal{X}} \to \underline{\mathbf{M}}\Omega^{q+1}_{\mathcal{X}}$.

Lemma 3.2. If $f = f_1^{r_1} \cdots f_n^{r_n}$ is a product of prime elements, then the differential of $\Omega_{A[f^{-1}]}^{\bullet}$ induces the differential on $\underline{M}\Omega_{(A,f)}^{\bullet}$.

Proof. Since $\underline{M}\Omega^q_{(A,f)} = \underline{M}\mathcal{O}(A,f) \cdot \underline{P}\Omega^q_{(A,f)}$ $_{(A,f)}^q$ and $\underline{P}\Omega_{(A,f)}^{\bullet}$ is a complex, it is sufficient to show $d\underline{M}\mathcal{O}(A, f) \subseteq \underline{M}\Omega^1_{(A, f)}$. Since $\sqrt{f} = (f_1 \cdots f_n)A$, for $\frac{a}{f} \in \underline{M}\mathcal{O}(A, f)$, we have $a = a'g$ where $g = f_1 \cdots f_n$. Thus

$$
d\left(\frac{a}{f}\right) = \frac{da}{f} - \frac{a}{f}d\log f
$$

= $\frac{g}{f}da' + \frac{a'}{f}dg - \frac{a}{f}d\log f$
= $\frac{g}{f}da' + \frac{a}{f}d\log g - \frac{a}{f}d\log f \in \underline{M}\Omega^1_{(A,f)}$.

If $\mathcal{X} = (\overline{X}, X^{\infty})$ is a normal crossing modulus pair, then \overline{X} is smooth, in particular locally factorial, so we can define the differential $d: \underline{M}\Omega^*_{\mathcal{X}} \to \underline{M}\Omega^{*+1}_{\mathcal{X}}$ by the following lemma.

Lemma 3.3. Let SpecA be a Noetherian locally factorial affine scheme and $f \in A$ be an element. For any point $\mathfrak{p} \in \text{Spec}(A)$, there is a neighborhood $\mathfrak{p} \in \text{Spec}(A[g^{-1}])$ such that $f = f_1^{r_1} \cdots f_n^{r_n}$ for some prime elements $f_j \in A[g^{-1}]$.

Proof. Since A_p is UFD, $f = f_1^{r_1} \cdots f_n^{r_n}$ for some prime elements $f_j \in A_p$. Let $\mathfrak{q}_j =$ $(u_{j1},...,u_{jk}) \subseteq \mathfrak{p}$ be the prime ideal in A such that $\mathfrak{q}_j A_{\mathfrak{p}} = f_j A_{\mathfrak{p}}$. Thus $u_{jl} = f_j v_{jl}$ for some $v_{jl} \in A_{\mathfrak{p}}$. We can write $f_j = s_j/g$, $v_{jl} = v'_{jl}/g$, and $f_j = \sum u_{jl}(w_{jl}/g)$ for some $s_j, v'_{jl}, w_{jl} \in A$ and $g \notin \mathfrak{p}$. Therefore $f = f_1^{r_1} \cdots f_n^{r_n}$ for $f_j = s_j/g \in A[g^{-1}]$ and each f_j is a prime element because $\mathfrak{q}_j A[g^{-1}] = f_j A[g^{-1}]$ \Box

Proposition 3.4. For every integer $q \geq 0$, the differential

$$
d: \underline{M}\Omega^q \to \underline{M}\Omega^{q+1}
$$

is a morphism of sheaves with transfers.

Proof. We need to show that the back square in the diagram

$$
\begin{CD} \begin{CD} \underline{M}\Omega^q(\mathcal{Y}) @>>> \underline{M}\Omega^{q+1}(\mathcal{Y}) \\ & @>>> \Omega^q(Y^{\circ}) @>>> \Omega^{q+1}(Y^{\circ}) \\ & @>>> \Omega^q(X) @>>> \underline{M}\Omega^{q+1}(\mathcal{X}) \\ & @>>> \Omega^q(X^{\circ}) @>>> \Omega^{q+1}(X^{\circ}). \end{CD} \end{CD}
$$

commutes for every modulus correspondence $W \in \mathbf{MCor}(\mathcal{X}, \mathcal{Y})$ for every normal crossing modulus pairs $\mathcal{X} = (\overline{X}, X^{\infty})$ and $\mathcal{Y} = (\overline{Y}, Y^{\infty})$, where $X^{\circ} = \overline{X} - X^{\infty}$, $Y^{\circ} = \overline{Y} - Y^{\infty}$. The top and the bottom squares commute by the definition of the differential on $\underline{M}\Omega^*$. The left and the right squares commute by [\[KM23b,](#page-16-7) Theorem 4.3.]. Since $\underline{M}\Omega^*(\mathcal{X}) \to \Omega^*(X^{\circ})$ and $\underline{M}\Omega^*(\mathcal{Y}) \to \Omega^*(Y^{\circ})$ are injective, it is sufficient to show that the front square commutes. This case is checked in [\[KSYR16,](#page-16-11) Theorem A.4.1], and [\[LW09\]](#page-16-12). \square

Theorem 3.5. There exists a unique object $\underline{M}\Omega^{\bullet} \in \underline{M}$ DM^{eff} such that for every strict normal crossing modulus pair $\mathcal{X} = (\overline{X}, X^{\infty})$ we have

$$
\mathrm{Hom}_{\mathbf{\underline{M}\underline{D}M}^{\mathrm{eff}}_k}(M(\mathcal{X}), \mathbf{\underline{M}\Omega}^\bullet[n]) \cong \mathbb{H}_{\mathrm{Zar}}^n(\overline{X}, \mathbf{\underline{M}\Omega}^\bullet_{\mathcal{X}}).
$$

Proof. The Hodge-to-de Rham spectral sequence

$$
E_1^{p,q} = H^q_{\text{Zar}}(\overline{X}, \underline{M}\Omega_{\mathcal{X}}^p) \implies \mathbb{H}_{\text{Zar}}^{p+q}(\mathcal{X}, \underline{M}\Omega^{\bullet})
$$

and the spectral sequence of extensions

$$
E_1^{p,q}=\operatorname{Hom}\nolimits_{\operatorname{\mathbf{\underline{M}DM}}\nolimits_k^{\operatorname{eff}\nolimits}}(M(\mathcal{X}),\operatorname{\mathbf{\underline{M}\Omega}}\nolimits^p[q]) \implies \operatorname{Hom}\nolimits_{\operatorname{\mathbf{\underline{M}DM}}\nolimits_k^{\operatorname{eff}\nolimits}}(M(\mathcal{X}),\operatorname{\mathbf{\underline{M}\Omega}}\nolimits^\bullet[p+q])
$$

induce the isomorphism

$$
\mathrm{Hom}_{\underline{\mathbf{M}\mathbf{D}\mathbf{M}^{\mathrm{eff}}_k}}(M(\mathcal{X}), \underline{\mathbf{M}\mathbf{\Omega}^\bullet}[n]) \cong \mathbb{H}_{\mathrm{Zar}}^n(\overline{X}, \underline{M}\Omega^\bullet_{\mathcal{X}})
$$

by Theorem [1.1.](#page-1-2) \Box

4. Modulus Hochschild Homology

Recall that the Hochschild homology $HH_*(A)$ of an k-algebra A is the homology of the simplicial ring $HH(A)$ whose nth ring is $A^{\otimes n+1}$ and face operators d_i : $A^{\otimes n+1} \to A^{\otimes n+2}$ and degeneracy operators are

$$
d_i(a_0 \otimes \cdots a_n) = \begin{cases} a_0 \otimes \cdots \otimes a_i a_{i+1} \otimes \cdots a_n & \text{if } 0 \leq i \leq n-1 \\ a_n a_0 \otimes \cdots \otimes a_{n-1} & \text{if } i = n \end{cases}
$$

$$
s_i(a_0 \otimes \cdots a_n) = a_0 \otimes \cdots a_i \otimes 1 \otimes a_{i+1} \otimes \cdots \otimes a_n.
$$

Let *b* be its differential

$$
b(a_0\otimes\cdots\otimes a_n)=\sum_{i=0}^{n-1}(-1)^ia_0\otimes\cdots\otimes a_ia_{i+1}\cdots\otimes a_n+(-1)^na_na_0\otimes\cdots\otimes a_{n-1}.
$$

The complex $HH(A)$ has a structure of a graded commutative dg-algebra by the shuffle product ∇ . Furthermore, if A is a smooth k-algebra, then we have an HKR isomorphism of graded algebras

$$
\Omega^*_{A/k} \xrightarrow{\sim} HH_*(A).
$$

In this section, we construct the modulus Hochschild homology $\mathcal{M}HH_q$ which is a Hochschild homology for modulus pairs and show the modulus HKR theorem.

We fix a field k .

Definition 4.1. For a k-algebra A and a nonzero divisor $f \in A$, we write $\underline{PHH}(A, f)$ for the sub-simplicial ring of $HH(A[f^{-1}])$ whose nth ring is the ring of sums of elements of the form

$$
a_0m_0\otimes\cdots\otimes a_nm_n\in A[f^{-1}]^{\otimes n+1},
$$

where $a_i \in A$, $m_i \in A[f^{-1}]^*$ and $m_0 \cdots m_n \in A$. Define sub-simplicial module

$$
\underline{M}HH(A,f):=\underline{M}\mathcal{O}(A,f)\cdot \underline{P}HH(A,f)\subseteq HH(A[f^{-1}]),
$$

We call it modulus Hochschild homology. Let $\underline{PHH}_*(A, f) := H_*(\underline{PHH}(A, f)),$ $\underline{MH}_*(A, f) := H_*(\underline{MHH}(A, f))$ to be the homology groups.

Remark 4.2. By definition, $\underline{PHH}(A, f)$ is a cdga by the shuffle product ∇ .

We will compare PHH with log Hochschild homology. Before proving that, we recall some definitions about pre-log rings.

Definition 4.3. A pre-log ring (A, M, α) is a triple of a commutative ring A, a commutative monoid M, and a map of monoids $\alpha : M \to A$. A map of pre-log rings $(f, f^{\flat}) : (A, M, \alpha) \to (B, N, \beta)$ is a pair of a map of rings $f : A \to B$ and a map of monoids $f^{\flat}: M \to N$, such that

is commutative. For a surjective map of monoids $f : M \to N$, repletion of f is the pullback $M^{\text{rep}} := N \times_{N^{\text{gp}}} M^{\text{gp}}$. For any map $f : M \to N$, let $(\bigoplus_{i=1}^{n} N)^{\text{rep}}$ be the repletion of $\bigoplus_{M}^{n} N \to N$.

Proposition 4.4. Let $(R, P) \rightarrow (A, M)$ be a map of pre-log rings. If there are factorizations $M = M_1 \oplus M_2$, $P = P_1 \oplus P_2$ such that $P \to M$ is the sum of the $P_i \to M_i$, P_2 and M_2 are groups, and $R \otimes_{\mathbb{Z}[P_1]} \mathbb{Z}[M_1] \to A$ is flat, then there is an $equivalence²$ $equivalence²$ $equivalence²$

(4.1)
$$
A \otimes_{(A \otimes_R A) \otimes_{\mathbb{Z}[M \oplus_P M]} \mathbb{Z}[(M \oplus_P M)^{rep}]} A \simeq C((A, M)/(R, P)),
$$

where $C((A, M)/(R, P))$ is a complex such that $C_n((A, M)/(R, P)) = A^{\otimes_R n+1} \otimes_{\mathbb{Z}[\oplus_{P}^{n+1} M]}$ $\mathbb{Z}[(\bigoplus_{P}^{n+1} M)^{\text{rep}}]$ with Hochschild differential.

Proof. Let $B_n := A^{\otimes_R n} \otimes_{\mathbb{Z}[\oplus_P^n M]} \mathbb{Z}[(\oplus_P^n M)^{\text{rep}}]$. The Bar resolution

$$
B \to A = (\cdots \to B_3 \to B_2) \to A
$$

is a contractible augmented simplicial object. If B_{n+2} is flat over B_2 , that is, B is a B₂-flat resolution of A, and if $B_{n+2} \otimes_{B_2} A \cong B_{n+1}$, then

$$
A\otimes_{B_2}^{\mathbb{L}} A\simeq B\otimes_{B_2} A\cong C((A,M)/(R,P)).
$$

Therefore it is sufficient to show that B_{n+2} is flat over B_2 , and $B_{n+2} \otimes_{B_2} A \cong B_{n+1}$. Since $\bigoplus_{P}^{n} M = (\bigoplus_{P_1}^{n} M_1) \oplus (\bigoplus_{P_2}^{n} M_2)$, and since P_2 and M_2 are groups,

(4.2)
$$
\left(\bigoplus_{P}^{n} M\right)^{\text{rep}} = \left(\bigoplus_{P_1}^{n} M_1\right)^{\text{rep}} \oplus \left(\bigoplus_{P_2}^{n} M_2\right)^{\text{rep}} = \left(\bigoplus_{P_1}^{n} M_1\right)^{\text{rep}} \oplus \left(\bigoplus_{P_2}^{n} M_2\right).
$$

Let $R' := R \otimes_{\mathbb{Z}[P_1]} \mathbb{Z}[M_1]$. Then

(4.3)
$$
B_n \cong A^{\otimes_{R'}n} \otimes_{\mathbb{Z}[M_1]} \mathbb{Z}[(\oplus_{P_1}^n M_1)^{\text{rep}}]
$$

because

$$
B_n = A^{\otimes_R n} \otimes_{\mathbb{Z}[\oplus_{P}^n M]} \mathbb{Z}[(\oplus_{P}^n M)^{\text{rep}}]
$$

\n
$$
\cong A^{\otimes_R n} \otimes_{\mathbb{Z}[\oplus_{P_1}^n M_1]} \mathbb{Z}[(\oplus_{P_1}^n M_1)^{\text{rep}}]
$$

\n
$$
\cong A^{\otimes_{R'} n} \otimes_{R'} (R'^{\otimes_R n}) \otimes_{\mathbb{Z}[\oplus_{P_1}^n M_1]} \mathbb{Z}[(\oplus_{P_1}^n M_1)^{\text{rep}}]
$$

\n
$$
\cong A^{\otimes_{R'} n} \otimes_{\mathbb{Z}[M_1]} (\mathbb{Z}[M_1]^{\otimes_{\mathbb{Z}[P_1]^n}}) \otimes_{(\mathbb{Z}[M_1]^{\otimes_{\mathbb{Z}[P_1]^n}})} \mathbb{Z}[(\oplus_{P_1}^n M_1)^{\text{rep}}]
$$

\n
$$
\cong A^{\otimes_{R'} n} \otimes_{\mathbb{Z}[M_1]} \mathbb{Z}[(\oplus_{P_1}^n M_1)^{\text{rep}}],
$$

²This equivalence holds for any map $(R, P) \rightarrow (A, M)$ in the derived sense [\[BLPØ23,](#page-16-13) Proposition 5.6].

where the third isomorphism is induced by $(-)\otimes_R R' = (-)\otimes_R (R\otimes_{\mathbb{Z}[P_1]} \mathbb{Z}[M_1]) \cong$ $(-) \otimes_{\mathbb{Z}[P_1]} \mathbb{Z}[M_1]$ and $\mathbb{Z}[\bigoplus_{P_1}^n M_1] \cong \mathbb{Z}[M_1]^{\otimes_{\mathbb{Z}[P_1]} n}$. Since

$$
B_{n+2} \cong A^{\otimes_{R'}n} \otimes_{R'} ((A \otimes_{R'} A) \otimes_{\mathbb{Z}[M_1]} \mathbb{Z}[(M_1 \oplus_{P_1} M_1)^{\text{rep}}]) \otimes_{\mathbb{Z}[(M_1 \oplus_{P_1} M_1)^{\text{rep}}]} \mathbb{Z}[(\oplus_{P_1}^n M_1)^{\text{rep}}],
$$

for any B_2 -module N

$$
B_{n+2} \otimes_{B_2} N \cong A^{\otimes_{R'}n} \otimes_{R'} N \otimes_{\mathbb{Z}[(M_1 \oplus_{P_1} M_1)^{\text{rep}}]} \mathbb{Z}[(\oplus_{P_1}^{n+2} M_1)^{\text{rep}}].
$$

Since A is flat over R', it is sufficient to show $\mathbb{Z}[(\oplus_{P_1}^{n+2}M_1)_{\text{rep}}]$ is flat over $\mathbb{Z}[(M_1 \oplus_{P_1} M_1)]_{\text{rep}}$ (M_1) ^{rep}]. However, for any homomorphism of monoids $P \to M$, $(\bigoplus_{P}^{n+1} M)^{\text{rep}} \cong$ $M \oplus (M^{\rm gp}/P^{\rm gp})^n$ (cf. [\[BLPØ23,](#page-16-13) Lemma 5.10.]). Therefore

$$
N \otimes_{\mathbb{Z}[(M_1 \oplus_{P_1} M_1)^{\text{rep}}]} \mathbb{Z}[(\oplus_{P_1}^{n+2} M_1)^{\text{rep}}] \cong N \otimes_{\mathbb{Z}[M_1]} \mathbb{Z}[M_1 \oplus (M_1^{\text{gp}})^n].
$$

Since A is flat over R' by assumption, B_{n+2} is flat pver B_n . Furthermore, if $N = A$, that isomorphism says $B_{n+2} \otimes_{B_2} A \cong B_{n+1}$.

The left hand side of eq. (4.1) is the definition of log Hochschild homology in [BLP \emptyset 23, Definition 5.3]. We will check that if (A, f) is strict normal crossing and $f = f_1^{r_1} \cdots f_n^{r_n}$ is a product of prime elements, then we have an isomorphism

(4.4)
$$
C((A, M)/(R, P)) \xrightarrow{\sim} \underline{P}HH(A, f),
$$

where $(A, M) = (A, A \cap A[f^{-1}]^*)$, $(R, P) = (k, k^*)$, and thus we have an equivalence

(4.5)
$$
A \otimes_{(A \otimes_R A) \otimes_{\mathbb{Z}[M \oplus_P M]} \mathbb{Z}[(M \oplus_P M)^{rep}]} A \simeq \underline{P}HH(A, f).
$$

There is a canonical surjection of simplicial rings $C((A, M)/(R, P)) \to \underline{PHH}(A, f)$. To show it is injective, it is sufficient to show both inject into $C(A[f^{-1}])$, but for the target this is the definition. By assumption, $M = A^* \times f_1^{\mathbb{N}} \times \cdots \times f_n^{\mathbb{N}} \cong \mathbb{N}^n \oplus A^*$ and $P = 0 \oplus k^*$. By eq. [\(4.3\)](#page-5-2),

$$
B_{m+1} \cong A^{\otimes_{R'} m+1} \otimes_{\mathbb{Z}[M_1]} \mathbb{Z}[(M_1^m)^{rep}] \cong A^{\otimes_{R'} m+1} \otimes_{R'} R'[(M_1^m)^{rep}],
$$

where $M_1 = \mathbb{N}^n$, $M_2 = A^*$, $P_1 = 0$, $P_2 = k^*$, and $R' = k[M_1] \cong k[x_1, \ldots, x_n]$. If A is flat over $R' = k[M_1]$, then there is a injection $A^{\otimes_{R'}n+1} \otimes_{R'} R'[(M_1^n)^{rep}] \rightarrow$ $A^{\otimes_{R'}n+1}\otimes_{R'} R'[(M_1^n)^{\otimes p}] \cong A[f^{-1}]^{\otimes n+1}$. Thus to use Proposition [4.4,](#page-5-3) it remains to show the flatness of $R' \to A$.

Lemma 4.5. For any normal crossing modulus pair (A, f) such that $f = f_1^{r_1} \cdots f_n^{r_n}$ for prime elements $f_1, \ldots, f_n \in A$, there exists an étale covering $\{Spec(A_i) \to$ $Spec(A)$ such that $R' = k[x_1, \ldots, x_n] \rightarrow A \rightarrow A_i$ factors through $k[x_1, \ldots, x_n] \rightarrow$ $k[y_1, \ldots, y_m] \rightarrow A_i$ where the first map is flat, and the second map is étale. In $\label{eq:particular} {\it particular,~} R' \rightarrow A ~\, \it{is~flat.}$

Proof. First, we construct an étale covering ${\rm Spec} A_i \rightarrow {\rm Spec} A$ with étale morphisms $SpecA_i \rightarrow A^m$. By definition, there is an étale covering ${Spec(B) \rightarrow$ $Spec(A)$ with (B, f) strict normal crossing. For each maximal ideal $\mathfrak{m} \in Spec(B)$, there is a regular system of parameters $b_1, \ldots b_m \in \mathfrak{m}B_{\mathfrak{m}}$ such that $f = b_1^{s_1} \cdots b_l^{s_l}$. Then $b_1, \ldots b_m \in \mathfrak{m} - \mathfrak{m}^2$ are prime elements, and

$$
\Omega_{B_{\mathfrak{m}}/k} = \bigoplus_{i=1}^{n} B_{\mathfrak{m}} db_i.
$$

8 MASAYA SATO

By Lemma [3.3,](#page-3-0) there is a $g \notin \mathfrak{m}$ such that $b_1, \ldots, b_n \in B[g^{-1}]$ are primes, $f =$ $b_1^{s_1}\cdots b_l^{s_l}$, and

$$
\Omega_{B[g^{-1}]/k} = \bigoplus_{i=1}^{n} B[g^{-1}]db_i.
$$

Since $B' := B[g^{-1}]$ is finite type over k, there is a surjection $S := k[y_1, \ldots, y_m, z_1, \ldots, z_c] \rightarrow$ B' with $y_i \mapsto b_i$. Let I be its kernel. Since B' is smooth over k, there are short exact sequences

$$
\begin{array}{ccccccc}\n0 & \longrightarrow & I/I^2 & \longrightarrow & \Omega_{S/k} \otimes_S B' & \longrightarrow & \Omega_{B'/k} & \longrightarrow & 0 \\
& & & & & & & \\
0 & \longrightarrow & & & & & & \\
0 & \longrightarrow & & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \longrightarrow & & & & & \\
0 & \
$$

where vertical maps are isomorphimsms. So there is a basis $h_1, \ldots, h_c \in I/I^2$. ByNakayama's lemma ([\[Sta18,](#page-16-14) [00DV\(](https://stacks.math.columbia.edu/tag/00DV)3)]), there is a $h \in 1 + I$ such that $hI \subset$ $(h_1, \ldots, h_c), I[h^{-1}] = (h_1, \ldots, h_c)[h^{-1}].$ Since $h = 1$ in B' ,

$$
B' \cong S/I \cong S[h^{-1}]/I[h^{-1}] \cong k[y_1,\ldots,y_m,z_1,\ldots,z_{c+1}]/(h_1,\ldots,h_c,hz_{c+1}-1).
$$

Let $h_{c+1} := hz_{c+1} - 1$ and $J := (h_1, \ldots, h_{c+1})$. By [\[Sta18,](#page-16-14) [08JZ\(](https://stacks.math.columbia.edu/tag/08JZ)1)], J/J^2 is free with basis $h_1, \ldots h_{c+1}$. By the short exact sequence

$$
0 \to J \to S' := k[y_1, \dots, y_m, z_1, \dots, z_{c+1}] \to B' \to 0
$$

we obtain short exact sequences and vertical isomorphisms again

$$
\begin{array}{ccccccc}\n0 & \xrightarrow{\hspace{1.5cm}} & J/J^2 & \xrightarrow{\hspace{1.5cm}} & \Omega_{S'/k} \otimes_{S'} B' & \xrightarrow{\hspace{1.5cm}} & \Omega_{B'/k} & \xrightarrow{\hspace{1.5cm}} & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \\
0 & \xrightarrow{\hspace{1.5cm}} & \bigoplus_{j=1}^{c+1} B'dz_i & \xrightarrow{\hspace{1.5cm}} & (\bigoplus_{j=1}^{c+1} B'dz_j) \oplus (\bigoplus_{i=1}^{m} B'dy_i) & \xrightarrow{\hspace{1.5cm}} & \bigoplus_{i=1}^{m} B'dy_i & \xrightarrow{\hspace{1.5cm}} & 0.\n\end{array}
$$

Since left vertical map is a invertible matrix $(\partial h_i/\partial z_j)$, $B' \cong S'/J$ is a standard smooth [Stacks, 00T6]. Then $k[y_1, \ldots, y_m] \to B'$ is also standard smooth and relative dimension 0 , so it is étale.

Next, we show the above construction induces a factorization of $R' = k[x_1, \ldots, x_n] \rightarrow$ $A \to B'$. By [\[KM23a,](#page-16-6) Lemma 3.5], $(f_1 \cdots f_n)B' = \sqrt{fAB'} = \sqrt{fB'} = (b_1 \cdots b_l)B'$. Since the above argument does not change with the reordering and the multiplication by units, we may assume $f_i = \prod_{k=j_{i-1}+1}^{j_i} b_j$. Thus $R' \to B'$ factors through

$$
R' = k[x_1, \ldots, x_n] \to k[y_1, \ldots, y_m] \to B',
$$

where the first map sends x_i to $\prod_{k=j_{i-1}+1}^{j_i} y_j$. This map is a tensor product of maps of the forms $k[x] \to k[y_1, \ldots, y_d] : x \mapsto y_1 \cdots y_d$ or $k \to k[y_1, \ldots, y_d]$. These maps are flat, so the desired map is flat.

Sinceflatness is an étale local property ([\[Sta18,](#page-16-14) [03MM\]](https://stacks.math.columbia.edu/tag/03MM)), $R' \rightarrow A$ is flat. \square

We denote
$$
(A \otimes_R A)_{M \oplus_P M}^{\text{rep}} := (A \otimes_R A) \otimes_{\mathbb{Z}[M \oplus_P M]} \mathbb{Z}[(M \oplus_P M)^{\text{rep}}].
$$

Lemma 4.6. If maps of pre-log rings $(R, P) \rightarrow (A, M) \rightarrow (B, N)$ induce flat ring maps $A \to B$ and $(A \otimes_R A)_{M \oplus_P M}^{\text{rep}} \to (B \otimes_R B)_{N \oplus_P N}^{\text{rep}},$ then

$$
(A\otimes_{(A\otimes_R A)_{M\oplus_P M}^{\operatorname{rep}}A})\otimes_A B\simeq (B\otimes_A B)_{N\oplus_M N}^{\operatorname{rep}}\otimes^{\mathbb{L}}_{(B\otimes_R B)_{N\oplus_P N}^{\operatorname{rep}}B.
$$

Proof. By assumption,

$$
(A \otimes_{(A \otimes_R A)_{M \oplus_{P} M}^{rep}}^{\mathbb{L}} A) \otimes_A B \simeq (A \otimes_{(A \otimes_R A)_{M \oplus_{P} M}^{rep}}^{\mathbb{L}} B) \simeq (A \otimes_{(A \otimes_R A)_{M \oplus_{P} M}^{rep}} (B \otimes_R B)_{N \oplus_{P} N}^{rep}) \otimes_{(B \otimes_R B)_{N \oplus_{P} N}^{rep}}^{\mathbb{L}} B \simeq (B \otimes_A B)_{N \oplus_M N}^{rep} \otimes_{(B \otimes_R B)_{N \oplus_{P} N}^{rep}}^{\mathbb{L}} B,
$$

where the third equivalence is $[BLPØ23,$ Proposition 5.4.].

Proposition 4.7 (Zariski descent). If (A, f) is a normal crossing modulus pair such that $f = f_1^{r_1} \cdots f_n^{r_n}$ is a product of prime elements, then for any multiplicatively closed subset $S \subset A$, there is an equivalence

$$
\underline{P}HH(S^{-1}A, f) \simeq \underline{P}HH(A, f) \otimes_A S^{-1}A.
$$

Proof. By eq. (4.5) ,

(4.6)
$$
\underline{PHH}(A,f) \simeq A \otimes_{(A\otimes_k A)_{M\oplus_{k^*}M}}^{\mathbb{L}} A,
$$

same holds for $(S^{-1}A, f)$. To use the Lemma [4.6,](#page-7-0) we check that

(4.7)
$$
(A \otimes_k A)^{\text{rep}}_{M \oplus_{k^*} M} \to (S^{-1}A \otimes_k S^{-1}A)^{\text{rep}}_{N \oplus_{k^*} N}
$$

is flat, where $M = A \cap A[f^{-1}] *$, $N = S^{-1}A \cap S^{-1}A[f^{-1}] *$. Assume $f_1, \ldots, f_i \in N$, $f_{i+1}, \ldots, f_n \notin N$. Since $f_1, \ldots, f_i \in N$ are primes in A, the inclusion

$$
f_1^{\mathbb{Z}} \times \cdots \times f_i^{\mathbb{Z}} \to (S^{-1}A)^*
$$

has a section. Thus

$$
N = (S^{-1}A)^* \times f_{i+1}^{\mathbb{N}} \times \cdots \times f_n^{\mathbb{N}} = N_2 \times f_1^{\mathbb{Z}} \times \cdots \times f_i^{\mathbb{Z}} \times f_{i+1}^{\mathbb{N}} \times \cdots \times f_n^{\mathbb{N}}
$$

for some abelian group N_2 . Let $N_1 \cong \mathbb{Z}^i \oplus \mathbb{N}^{n-i}$ be the other summand of N. Then the map $M \to N$ is a sum of $M_1 = \prod_{j=1}^n f_j^{\mathbb{N}} \to \prod_{j=1}^i f_j^{\mathbb{Z}} \times \prod_{j=i+1}^n f_j^{\mathbb{N}} = N_1$ and $M_2 = A^* \rightarrow N_2$. We have

$$
(A \otimes_k A)^{\text{rep}}_{M \oplus_{k^*} M} \cong (A \otimes_k A) \otimes_{\mathbb{Z}[M_1 \oplus M_1]} \mathbb{Z}[(M_1 \oplus M_1)^{\text{rep}}]
$$

\n
$$
\rightarrow (S^{-1}A \otimes_k S^{-1}A) \otimes_{\mathbb{Z}[M_1 \oplus M_1]} \mathbb{Z}[(M_1 \oplus M_1)^{\text{rep}}]
$$

\n
$$
\cong (S^{-1}A \otimes_k S^{-1}A) \otimes_{\mathbb{Z}[N_1 \oplus N_1]} \mathbb{Z}[N_1 \oplus N_1] \otimes_{\mathbb{Z}[M_1 \oplus M_1]} \mathbb{Z}[(M_1 \oplus M_1)^{\text{rep}}]
$$

\n
$$
\cong (S^{-1}A \otimes_k S^{-1}A) \otimes_{\mathbb{Z}[N_1 \oplus N_1]} \mathbb{Z}[(N_1 \oplus N_1)^{\text{rep}}]
$$

\n
$$
\cong (S^{-1}A \otimes_k S^{-1}A)^{\text{rep}}_{N \oplus_{k^*} N},
$$

where eq. (4.2) implies the first and last isomorphisms. The second map is flat because $A \to S^{-1}A$ is flat. Thus to show that eq. [\(4.7\)](#page-8-0) is flat, it is sufficient to show the forth map is an isomorphism.

Since
$$
N_1 = \mathbb{Z}^i \oplus \mathbb{N}^{n-i}
$$
,
\n $(N_1 \oplus N_1)^{rep} = (\mathbb{Z}^i \oplus \mathbb{Z}^i)^{rep} \oplus (\mathbb{N}^{n-i} \oplus \mathbb{N}^{n-i})^{rep} = (\mathbb{Z}^i \oplus \mathbb{Z}^i) \oplus (\mathbb{N}^{n-i} \oplus \mathbb{N}^{n-i})^{rep}$
\nSince $M_1 = \mathbb{N}^i \oplus \mathbb{N}^{n-i}$,
\n $(N_1 \oplus N_1) \oplus_{(M_1 \oplus M_1)} (M_1 \oplus M_1)^{rep}$
\n $\cong ((\mathbb{Z}^i \oplus \mathbb{Z}^i) \oplus_{(\mathbb{N}^i \oplus \mathbb{N}^i)} (\mathbb{N}^i \oplus \mathbb{N}^i)^{rep}) \oplus ((\mathbb{N}^{n-i} \oplus \mathbb{N}^{n-i}) \oplus_{(\mathbb{N}^{n-i} \oplus \mathbb{N}^{n-i})} (\mathbb{N}^{n-i} \oplus \mathbb{N}^{n-i})^{rep})$
\n $\cong (\mathbb{Z}^i \oplus \mathbb{Z}^i) \oplus (\mathbb{N}^{n-i} \oplus \mathbb{N}^{n-i})^{rep}$

 $\cong (N_1 \oplus N_1)^{\text{rep}}.$

Since eq. (4.7) is flat, by Lemma [4.6,](#page-7-0) we have

$$
(A \otimes_{(A \otimes_k A)_{M \oplus_{k^*} M}}^{\mathbb{L}} A) \otimes_A S^{-1} A \simeq (S^{-1}A \otimes_A S^{-1}A)_{N \oplus_M N}^{\text{rep}} \otimes_{(S^{-1}A \otimes_k S^{-1}A)_{N \oplus_{k^*} N}}^{\mathbb{L}} S^{-1} A.
$$

\nHowever, $S^{-1}A \otimes_A S^{-1}A \cong S^{-1}A$, and
\n
$$
(N \oplus_M N)^{\text{rep}} = ((N_1 \oplus_{M_1} N_1) \oplus (N_2 \oplus_{M_2} N_2))^{\text{rep}}
$$
\n
$$
= ((\mathbb{Z}^i \oplus_{N^i} \mathbb{Z}^i) \oplus (\mathbb{N}^{n-i} \oplus_{N^{n-i}} \mathbb{N}^{n-i}) \oplus (N_2 \oplus_{M_2} N_2))^{\text{rep}}
$$
\n
$$
= (\mathbb{Z}^i \oplus_{N^i} \mathbb{Z}^i)^{\text{rep}} \oplus (\mathbb{N}^{n-i} \oplus_{N^{n-i}} \mathbb{N}^{n-i})^{\text{rep}} \oplus (N_2 \oplus_{M_2} N_2)^{\text{rep}}
$$
\n
$$
= \mathbb{Z}^i \oplus \mathbb{N}^{n-i} \oplus (N_2 \oplus_{M_2} N_2)
$$
\n
$$
= N \oplus_M N.
$$

\nHence $(S^{-1}A \otimes_A S^{-1}A)_{N \oplus_M N}^{\text{rep}} \cong S^{-1}A$ and
\n $(A \otimes_k A) = (S^{-1}A)_{N \oplus_M N}^{\text{rep}} \cong (S^{-1}A)_{N \oplus_M N}^{\text{rep}}$

$$
(A \otimes_{(A\otimes_k A)^{\text{rep}}_{M\oplus_{k^*}M}}^{\mathbb{L}} A) \otimes_A S^{-1}A \simeq (S^{-1}A \otimes_A S^{-1}A)^{\text{rep}}_{N\oplus_M N} \otimes_{(S^{-1}A\otimes_k S^{-1}A)^{\text{rep}}_{N\oplus_{k^*}N}}^{\mathbb{L}} S^{-1}A
$$

$$
\simeq S^{-1}A \otimes_{(S^{-1}A\otimes_k S^{-1}A)^{\text{rep}}_{N\oplus_{k^*}N}} S^{-1}A.
$$

Proposition 4.8 (Étale descent). Let (A, f) be a strict normal crossing modulus pair. Suppose that $A \rightarrow B$ is an étale morphism and f is a product of prime elements in A and B. Then

 $PHH(B, f) \simeq PHH(A, f) \otimes_A B.$

Proof. By Proposition [4.4](#page-5-3) and eq. (4.5) , $PHH(A, f)$ and $PHH(B, f)$ are equivalent to the complexes $A \otimes_{(A\otimes_k A)_{M\oplus_{k\neq M}}^{\text{rep}}}^{\mathbb{L}} A$ and $B \otimes_{(B\otimes_k B)_{N\oplus_{k\neq N}}^{\text{rep}}}^{\mathbb{L}} B$, where $M = A \cap$ $A[f^{-1}]^*$, and $N = B \cap B[f^{-1}]^*$. By Proposition [4.7,](#page-8-1) we may assume that φ : $A \rightarrow B$ is an étale homomorphism of local rings. Let m_A be a maximal ideal of A and x_1, \ldots, x_n be its regular system of parameters. By [\[Sta18,](#page-16-14) [09CC\]](https://stacks.math.columbia.edu/tag/09CC), s_1, \ldots, s_n is a regular system of parameters in Noetherian local ring (S, \mathfrak{m}) if and only if $(s_1, \ldots, s_n) = \mathfrak{m}$ and the homology of the Koszul complex $H_i(K_{\bullet}(s_1, \ldots, s_n)) = 0$ for $i \neq 0$. Since flat local homomorphisms are faithful and since $\mathfrak{m}_A B = \mathfrak{m}_B$, x_1, \ldots, x_n is also a regular system of parameters of B. Therefore $M = A \cap A[f^{-1}]^* = \bigcup_{N} M[f^{-1}]$ $A^* \times x_1^{\mathbb{N}} \times \cdots \times x_i^{\mathbb{N}}$ and $N = B \cap B[f^{-1}]^* = B^* \times x_1^{\mathbb{N}} \times \cdots \times x_i^{\mathbb{N}}$. Let $L := x_1^{\mathbb{N}} \times \cdots \times x_i^{\mathbb{N}}$. Thus

$$
(B \otimes_k B)^{\text{rep}}_{N \oplus_{k^*} N} \cong (B \otimes_k B)^{\text{rep}}_{L \oplus L}
$$

\n
$$
\cong (B \otimes_k B) \otimes_{(A \otimes_k A)} (A \otimes_k A)^{\text{rep}}_{L \oplus L}
$$

\n
$$
\cong (B \otimes_k B) \otimes_{(A \otimes_k A)} (A \otimes_k A)^{\text{rep}}_{M \oplus_{k^*} M}
$$

is flat over $(A \otimes_k A)_{M \oplus_{k^*} M}^{\text{rep}}$, where the first and third morphisms are isomorphisms by eq. [\(4.3\)](#page-5-2) for $n = 2$. By Lemma [4.6,](#page-7-0)

$$
(A\otimes^{\mathbb{L}}_{(A\otimes_R A)^{\text{rep}}_{M\oplus_P M}} A)\otimes_A B\simeq (B\otimes_A B)^{\text{rep}}_{N\oplus_M N}\otimes^{\mathbb{L}}_{(B\otimes_R B)^{\text{rep}}_{N\oplus_P N}} B.
$$

Since $N \oplus_M N \cong (B^* \oplus_{A^*} B^*) \oplus (L \oplus_L L) \cong (B^* \oplus_{A^*} B^*) \oplus L$, $(N \oplus_M N)^{\text{rep}} \cong N \oplus_M N$. Therefore $(B \otimes_A B)_{N \oplus_M N}^{\text{rep}} \cong B \otimes_A B \cong B \times C$ because $A \to B$ is étale. Hence

$$
\underline{PHH}(A, f) \otimes_A B \simeq (B \times C) \otimes_{(B \otimes_k B)^{\text{rep}}}^{\mathbb{L}} B
$$

$$
\simeq \underline{PHH}(B, f) \times (C \otimes_{(B \otimes_k B)^{\text{rep}}}^{\mathbb{L}} B).
$$

For any maximal ideal $\frak m$ of $(B\otimes_k B)^\text{rep},\ (C\otimes^{\mathbb L}_{(B\otimes_k B)^\text{rep}} B)\otimes_{(B\otimes_k B)^\text{rep}} (B\otimes_k B)^\text{rep}_\frak m \simeq$ $C_{\mathfrak{m}} \otimes_{(B\otimes_k B)^{\text{rep}}_{\mathfrak{m}}}^{\mathbb{L}} B_{\mathfrak{m}}$. Since $(B\otimes_k B)^{\text{rep}} \to B\otimes_A B \cong B \times C$ is surjective, $B_{\mathfrak{m}} = 0$ or $C_{\mathfrak{m}} = 0$. Therefore $C \otimes_{(B\otimes_k B)^{\text{rep}}}^{\mathbb{L}} B \simeq 0$ and we get an desired isomorphism. \Box

Lemma 4.9. If $M\mathcal{O}(A, f)$ is a flat A-module, then the canonical map

$$
\underline{M}\mathcal{O}(A,f) \otimes_A \underline{PHH}_q(A,f) \to \underline{M}HH_q(A,f)
$$

is an isomorphism for any $q \geq 0$. In particular if A is locally factorial, the above map is an isomorphism.

Proof. Since $M\mathcal{O}(A, f)$ is a flat A-module, $M\mathcal{O}(A, f) \otimes_A PHH_q(A, f) \cong H_q(M\mathcal{O}(A, f) \otimes_A H_q(A, f))$ $PC(A, f)$. Thus it is sufficient to show that the canonical surjection

$$
\underline{M}\mathcal{O}(A,f)\otimes \underline{PHH}(A,f)\to \underline{M}HH(A,f)
$$

of chain complexes is injective. However, there is a commutative diagram

$$
\underline{M}\mathcal{O}(A, f) \otimes_A \underline{PHH}(A, f) \longrightarrow \underline{M}HH(A, f)
$$
\n
$$
\downarrow \qquad \qquad \downarrow
$$
\n
$$
\underline{M}\mathcal{O}(A, f) \otimes_A HH(A[f^{-1}]) \longrightarrow HH(A[f^{-1}]).
$$

The left map is injective because $\mathcal{MO}(A, f)$ is flat. The bottom map is an isomorphism by $\underline{M}\mathcal{O}(A, f) \otimes_A A[f^{-1}] \cong A[f^{-1}]$. Therefore the top map is injective. \square

Proposition 4.10. For any modulus pair (A, f) , we have isomorphisms $\underline{P}\Omega^1_{(A,f)} \cong$ $\underline{PHH}_1(A, f)$ and $\underline{M}\Omega^1_{(A,f)} \cong \underline{M}HH_1(A, f)$.

Proof. Since $b(a_0 \otimes a_1) = a_0a_1 - a_1a_0 = 0$,

$$
\underline{PHH}_1(A,f)=\langle\sum c_0\otimes c_1\rangle/\langle c_0c_1\otimes c_2-c_0\otimes c_1c_2+c_2c_0\otimes c_1\rangle,
$$

where $c_i = a_i m_i$ for $a_i \in A$, $m_i \in A[f^{-1}]^*$ such that $\prod m_i \in A$. Let e : $\underline{PC}_1(A, f) := \langle \sum c_0 \otimes c_1 \rangle \rightarrow \Omega_{A[f^{-1}]}$ be the map which sends $a_0 m_0 \otimes a_1 m_1 \in$ $PC_1(A, f)$ to

$$
a_0 m_0 d(a_1 m_1) = a_0 m_0 m_1 da_1 + a_0 a_1 m_0 dm_1
$$

= $a_0 m_0 m_1 da_1 + a_0 a_1 m_0 m_1 d \log m_1$

Since $m_0m_1 \in A$ and $d \log m_1 = d \log m'_1 - d \log m''_1$, where $m_1 = \frac{m'_1}{m''_1}$ and $m'_1, m''_1 \in A$ $A \cap A[f^{-1}]^*$, e factors through $\underline{P}\Omega^1_{(A,f)}$. Since $d(c_1c_2) = c_1dc_2 + c_2dc_1$ in $\underline{P}\Omega^1_{(A,f)}$, and since

$$
b(c_0 \otimes c_1 \otimes c_2) = c_0c_1 \otimes c_2 - c_0 \otimes c_1c_2 + c_2c_0 \otimes c_1,
$$

in $\angle H H H(A, f)$, e factors through $\angle H H_1(A, f)$. Therefore e is a map from $\angle H H_1(A, f)$ to $\underline{P}\Omega^1_{(A,f)}$. Conversely, let $\psi : \underline{P}\Omega^1_{(A,f)} \to \underline{P}HH_1(A,f)$ to be the map which sends $\frac{a}{b}dc$ to $\frac{a}{b}\otimes c$. By definition, $e\psi = id$ and $\psi e = id$.

Since e and ψ commutes multiplication by an element of $A[f^{-1}]$ on the left, we can extend the isomorphism to $\underline{M}\Omega^1_{(A,f)} \cong \underline{M}HH_1(A,f)$.

By the universal property of exterior product, the morphisms $\underline{P}\Omega^0_{(A,f)} = A =$ $\underline{PHH}_0(A, f)$ and $\underline{P}\Omega^1_{(A, f)} \xrightarrow{\sim} \underline{PHH}_1(A, f)$ induce the morphism of graded-commutative algebras

(4.8)
$$
\bigwedge\nolimits_A^* \underline{P} \Omega^1_{(A,f)} \to \underline{P} HH_*(A,f).
$$

Theorem 4.11 (HKR-Theorem). Let $(Spec A, (f))$ be a strict normal crossing modulus pair such that f is a product of prime elements. Then the composition of eq. [\(4.8\)](#page-11-1) and the isomorphism constructed in [\[KM23b,](#page-16-7) Lemma 3.12]

(4.9)
$$
\underline{P}\Omega^*(A,f) \cong \bigwedge_A^* \underline{P}\Omega^1_{(A,f)} \to \underline{P}HH_*(A,f)
$$

is an isomorphism. Furthermore, there is an isomorphism

$$
\underline{M}\Omega^*(A,f) \cong \underline{M}HH_*(A,f).
$$

Proof. First, we prove that eq. [\(4.9\)](#page-11-2) is isomorphism if

$$
(A, f) = (k[x_1, \ldots, x_s, y_1, \ldots, y_t], x_1^{r_1} \cdots x_s^{r_s}).
$$

There is a commutative diagram

$$
\begin{array}{ccc}\n\underline{P}\Omega_{(A,f)}^{q} & \longrightarrow & \underline{P}HH_{q}(A,f) \\
\downarrow & & \downarrow \\
\Omega_{A[f^{-1}]}^{q} & \xrightarrow{\sim} & HH_{q}(A[f^{-1}]),\n\end{array}
$$

where the bottom map is an isomorphism by the HKR-theorem for the smooth k-algebra $A[f^{-1}]$. Here, $\underline{P}\Omega_f^q$ $_{(A,f)}^q$ is a direct summand of $\Omega_{A[f^{-1}]}^q$ and $\angle PHH_q(A, f)$ is a direct summand of $HH_q(A[f^{-1}])$. Indeed,

$$
\Omega_{A[f^{-1}]}^q = \bigoplus_{i_j \in \mathbb{Z}, \ k_l \geq 0, \ s_m + t_n = q} k \cdot x_1^{i_1} \cdots x_s^{i_s} y_1^{k_1} \cdots y_t^{k_t} dx_{s_1} \wedge \cdots \wedge dx_{s_m} \wedge dy_{t_1} \wedge \cdots \wedge dy_{t_n}
$$

$$
= \underline{P} \Omega_{(A,f)}^q \oplus D,
$$

where

$$
\underline{P}\Omega_{(A,f)}^q = \bigoplus_{i_j, k_l \ge 0, s_m + t_n = q} k \cdot x_1^{i_1} \cdots x_s^{i_s} y_1^{k_1} \cdots y_t^{k_t} \cdot d \log x_{s_1} \wedge \cdots \wedge d \log x_{s_m} \wedge dy_{t_1} \wedge \cdots \wedge dy_{t_n}
$$

$$
D^q := \bigoplus_{\exists i_{s'_m} < 0, s_m + t_n = q} k \cdot x_1^{i_1} \cdots x_s^{i_s} y_1^{k_1} \cdots y_t^{k_t} \cdot d \log x_{s_1} \wedge \cdots \wedge d \log x_{s_m} \wedge dy_{t_1} \wedge \cdots \wedge dy_{t_n}
$$

.

Also,

$$
A[f^{-1}]^{\otimes q+1} = \bigoplus_{i_j \in \mathbb{Z}^s, k_l \in (\mathbb{Z}_{\geq 0})^t} k \cdot x^{i_0} y^{k_0} \otimes \cdots \otimes x^{i_q} y^{k_q}
$$

$$
= \underline{PC}_q(A, f) \oplus E_q
$$

where $x^{i_j} = x_1^{i_{j,1}} \cdots x_s^{i_{j,s}}$, $y^{k_l} = y_1^{k_{l,1}} \cdots y_t^{k_{l,t}}$, and

$$
\underline{PC}_q(A, f) := \bigoplus_{x^{i_0} y^{k_0} \cdots x^{i_q} y^{k_q} \in A} k \cdot x^{i_0} y^{k_0} \otimes \cdots \otimes x^{i_q} y^{k_q}
$$

$$
E_q := \bigoplus_{x^{i_0} y^{k_0} \cdots x^{i_q} y^{k_q} \notin A} k \cdot x^{i_0} y^{k_0} \otimes \cdots \otimes x^{i_q} y^{k_q}.
$$

By the definition of the differential on $HH(A[f^{-1}])$, it decompose two chain complexes $HH(A[f^{-1}]) = \underline{PHH}(A, f) \oplus E_*,$ moreover $HH_q(A[f^{-1}]) = \underline{PHH}_q(A, f) \oplus$ $H_q(E)$. Since the canonical map $\Omega_{A[f^{-1}]}^q \cong HH_q(A[f^{-1}])$ sends $\underline{P}\Omega_q^q$ $\begin{array}{cc} q\\(A,f) \end{array}$ to $\underline{PHH}_{q}(A, f)$ and D^{q} to E_{q} , the canonical map $\underline{P}\Omega^{q}_{(A,f)} \to \underline{PHH}_{q}(A, f)$ is an isomorphism.

For strict normal crossing modulus pair $(Spec A, (f))$, there is an open covering $\{Spec A_k\}$ of $Spec(A)$, such that there is an étale morphism $Spec A_k \rightarrow$ $Spec(k[x_1, \ldots, x_n])$ with $(f) \times_{Spec A} Spec A_k = (x_1^{r_1} \cdots x_i^{r_i}) \times_{Spec(k[x_1, \ldots, x_n])} Spec A_k$ and x_j primes in A_k by Lemma [4.5.](#page-6-1) Therefore étale descent of $\overline{PHH_q}$ and $P\Omega^q$ $([KM23b, Proposition 3.17])$ $([KM23b, Proposition 3.17])$ $([KM23b, Proposition 3.17])$ induces the HKR isomorphism for $(Spec A, (f)).$

Since $\underline{M}\Omega_{\ell}^{q}$ $\frac{q}{(A,f)} \cong \underline{M}\mathcal{O}(A,f) \otimes_A \underline{P}\Omega^q$ $\mathcal{A}_{(A,f)}^q$ and $\mathcal{M}HH_q(A,f) \cong \mathcal{MO}(A,f) \otimes_A$ $PHH_q(A, f)$, the canonical map is an isomorphism.

5. Modulus Cyclic Homology

In this section, we will construct modulus (periodic and negative) cyclic homology. From now on, we assume characteristic of k is 0.

Before doing that, recall the cyclic homology of a flat k -algebra A . The cyclic group C_{n+1} acts on $A^{\otimes n+1}$ by

$$
t_n(a_0\otimes\cdots\otimes a_{n-1}\otimes a_n)=a_n\otimes a_1\otimes\cdots\otimes a_{n-1}.
$$

This action and simplicial structure on $HH(A)$ is compatible, that is, $HH(A)$ is a functor from the cyclic category Λ^{op} to the derived ∞ -category $\mathcal{D}(k)$ of chain complexes of k -modules (cf. [\[Lod98\]](#page-16-10)). Its geometric realization

$$
HH(A) = \varinjlim_{[n] \in \Delta^{\mathrm{op}}} A^{\otimes n+1} \in \mathrm{Fun}(B\mathbb{T}, \mathcal{D}(k))
$$

has an action by the circle $\mathbb T$. Cyclic homology and negative cylcic homology are defined by

$$
HC(A) := (HH(A))_{h\mathbb{T}}
$$

$$
HC^{-}(A) := (HH(A))^{h\mathbb{T}}
$$

the homotopy orbits and homotopy fixed points. For any $X \in \text{Fun}(B\mathbb{T}, \mathcal{D}(k)),$ there is a norm map $N: \Sigma X_{h\mathbb{T}} \to X^{h\mathbb{T}}$ (cf. [\[NS18\]](#page-16-15)), whose cofiber is called Tate cohomology and denoted by $X^{t\mathbb{T}}$. Periodic cyclic homology $HP(A)$ of A is the Tate cohomology $(HH(A))^{tT}$.

For a cyclic object $M_{\bullet} \in \text{Fun}(\Lambda^{\text{op}}, \mathcal{D}(k))$, we can define operators

$$
b := \sum_{i=0}^{n} (-1)^{i} d_{i} : M_{n} \to M_{n-1}
$$

$$
B := (1 - (-1)^{n+1} t_{n+1}) t_{n+1} s_{n} \left(\sum_{i=0}^{n} (-1)^{n} t_{n}^{i} \right) : M_{n} \to M_{n+1},
$$

and these maps satisfies $b^2 = B^2 = bB + Bb = 0$. If $M = HH(A)$, there is a commutative diagram

(5.1)
$$
A^{\otimes n} \xleftarrow{b} A^{\otimes n+1} \xrightarrow{B} A^{\otimes n+2}
$$

$$
\downarrow e \qquad \qquad \downarrow e \qquad \qquad \downarrow e
$$

$$
\Omega_A^{n-1} \xleftarrow{0} \Omega_A^n \xrightarrow{d} \Omega_A^{n+1},
$$

where the vertical morphisms $e: A^{\otimes n+1} \to \Omega^n_A$ are

(5.2)
$$
e(a_0 \otimes \cdots \otimes a_n) = \frac{1}{n!} a_0 da_1 \wedge \cdots \wedge da_n.
$$

By $[Hoy18], HP(A)$ is equivalent to the total complex of the bicomplex

(5.3) · · · ^A⊗³ ^A⊗² ^A · · · ^A⊗² ^A · · · A. b B b B b B

Similarly, if we remove all negatively (resp. positively) graded columns, then its total complex is equivalent to $HC(A)$ (resp. $HC^{-}(A)$).

If $f = f_1^{r_1} \cdots f_i^{r_i}$ is a product of prime elements, then C_{n+1} acts on the n-th group of $\underline{M}HH(A, f)$. Indeed, since $\underline{M}\mathcal{O}(A, f) = \frac{1}{f_1^{r_1-1} \cdots f_i^{r_i-1}} \cdot A$, we have

$$
(5.4)
$$
\n
$$
t\left(\frac{a_0}{f_1^{r_1-1}\cdots f_i^{r_i-1}}b_0\otimes\cdots\otimes a_nb_n\right) = a_nb_n\otimes \frac{a_0b_0}{f_1^{r_1-1}\cdots f_i^{r_i-1}}\otimes\cdots\otimes a_{n-1}b_{n-1}
$$
\n
$$
= \frac{a_n}{f_1^{r_1-1}\cdots f_i^{r_i-1}}(f_1^{r_1-1}\cdots f_i^{r_i-1}b_n)\otimes a_0\frac{b_0}{f_1^{r_1-1}\cdots f_i^{r_i-1}}\otimes\cdots\otimes a_{n-1}b_{n-1}
$$
\n
$$
\in \underline{M}HH(A, f).
$$

Thus $MHH(A, f)$ also has a circle action.

Definition 5.1. For a modulus pair (A, f) such that $f = f_1^{r_1} \cdots f_i^{r_i}$ is a product of prime elements, define

$$
\underline{MHC}(A, f) := (MHH(A, f))_{h\mathbb{T}}
$$

$$
\underline{MHC}^{-}(A, f) := (MHH(A, f))^{h\mathbb{T}}
$$

$$
\underline{MHP}(A, f) := (MHH(A, f))^{t\mathbb{T}}.
$$

We let $\underline{M}HC_n(A, f)$, $\underline{M}HC_n^-(A, f)$, and $\underline{M}HP_n(A, f)$ be their homologies, and call them modulus (negative and periodic) cyclic homologies.

Theorem 5.2. If (A, f) is strict normal crossing and $f = f_1^{r_1} \cdots f_n^{r_n}$ for prime elements $f_j \in A$, then

$$
\underline{MHC}_n(A,f) \cong (\underline{M}\Omega^n_{(A,f)}/d\underline{M}\Omega^{n-1}_{(A,f)}) \times H^{n-2}(\underline{M}\Omega^n_{(A,f)}) \times H^{n-4}(\underline{M}\Omega^n_{(A,f)}) \times \cdots
$$

\n
$$
\underline{MHC}^-(A,f) \cong \text{Ker}(\underline{M}\Omega^n_{(A,f)} \to \underline{M}\Omega^{n+1}_{(A,f)}) \times H^{n+2}(\underline{M}\Omega_{(A,f)^*}) \times H^{n+4}(\underline{M}\Omega^n_{(A,f)}) \times \cdots
$$

\n
$$
\underline{MHP}_n(A,f) \cong \prod_{p \in \mathbb{Z}} H^{2p-n}(\underline{M}\Omega^*(A,f)).
$$

Proof. We can restrict the $e: HH(A[f^{-1}]) \to \prod_{p\geq 0} \Omega_{A[f^{-1}]}^p[p]$ from eq. [\(5.2\)](#page-13-1) to subcomplexes $e: \underline{M}HH(A, f) \to \prod_{p\geq 0} \underline{M}\Omega^p_{(p)}$ $_{(A,f)}^p[p]$, because

$$
e\left(\frac{a}{f}\cdot a_0b_0\otimes a_1b_1\otimes\cdots\otimes a_nb_n\right):=\frac{a}{n!f}\cdot a_0b_0d(a_1b_1)\wedge\cdots\wedge d(a_nb_n)
$$

$$
=\frac{a}{n!f}\cdot a_0b_0\cdots b_n\sum\omega_1\wedge\cdots\wedge\omega_n\in \underline{M}\Omega^n_{(A,f)},
$$

where ω_i is da_i or $a_i d \log b_i$. The morphism e is the inverse map of the HKR isomorphism $\underline{M}\Omega_{(A,f)}^n \stackrel{\sim}{\to} \underline{M}HH_n(A,f)$ (cf. [\[Wei94,](#page-16-17) section 9.8]). Thus e is a quasi-isomorphism. Similar for periodic cyclic homology $HP(A)$ eq. [\(5.3\)](#page-13-2), by [\[Hoy18\]](#page-16-16), modulus periodic cyclic homologies are equivalent to the total complex $\text{Tot}^{\Pi}(\underline{MHH}(A, f)[\bullet], B)$, where $(\underline{MHH}(A, f)_{\bullet}, B)$ is the bicomplex associated to the chain complex of complexes

$$
(\cdots \to \underline{M}HH(A,f)[n] \xrightarrow{B} \underline{M}HH(A,f)[n-1] \to \cdots).
$$

By the commutativity of eq. (5.1) , the equivalence e commutes operators B : $\underline{MH}(A, f) \to \underline{MH}(A, f)[-1]$ and the de Rham differential $d: \prod_{p\geq 0} \underline{M}\Omega^p_{(p)}$ $_{(A,f)}^{p}[p]\rightarrow$ $(\prod_{p\geq 0} \underline{M} \Omega_p^p)$ $\binom{p}{(A,f)}[p]][-1]$, so there are equivalences

$$
\text{Tot}^{\prod}(\underline{M}HH(A,f)[\bullet],B) \xrightarrow{c} \text{Tot}^{\prod}(\prod_{p\geq 0} \underline{M}\Omega_{(A,f)}^{p}[\bullet],d) \cong \prod_{p\in \mathbb{Z}}(\underline{M}\Omega_{(A,f)}^{\bullet})^{-*}[2p].
$$

Restricting bicomplexes, we obtain equivalences

$$
\underline{MHC}(A, f) \simeq \prod_{p \in \mathbb{Z}} (\underline{M}\Omega_{(A,f)}^{\leq p})^{-*}[2p]
$$

$$
\underline{MHC}^{-}(A, f) \simeq \prod_{p \in \mathbb{Z}} (\underline{M}\Omega_{(A,f)}^{\geq p})^{-*}[2p]
$$

$$
\underline{MHP}(A, f) \simeq \prod_{p \in \mathbb{Z}} (\underline{M}\Omega_{(A,f)}^{\bullet})^{-*}[2p].
$$

If we take homologies, then we obtain the desired isomorphisms

$$
\underline{MHC}_n(A, f) \cong \prod_{p \in \mathbb{Z}} H^{2p-n}(\underline{M}\Omega_{(A,f)}^{\leq p})
$$

$$
\underline{MHC}_n^-(A, f) \cong \prod_{p \in \mathbb{Z}} H^{2p-n}(\underline{M}\Omega_{(A,f)}^{\geq p})
$$

$$
\underline{MHP}_n(A, f) \cong \prod_{p \in \mathbb{Z}} H^{2p-n}(\underline{M}\Omega_{(A,f)}^{\bullet}).
$$

 \Box

6. Realizations

We will construct objects of $\underline{M} \text{DM}_{k}^{\text{eff}}$ which represents the modulus Hochschild homology and modulus (periodic and negative) cyclic homologies.

Definition 6.1. For a modulus pair $\mathcal{X} := (\overline{X}, X^{\infty})$, let $\mathcal{M}_{\mathcal{X}} := \mathcal{O}_{\overline{X}} \cap \mathcal{O}_{X}^{*}$, where $X := \overline{X} \setminus X^{\infty}$. For a map of sheaf of rings $A \to B$, let $B^{cyc}_A(\mathcal{B})$ be the cyclic Bar construction in the category $\mathcal{Sh}(\overline{X}_{\text{Zar}}, \text{Ring})$ of sheaves of rings on $\overline{X}_{\text{Zar}}$. Let

 $B^{\text{rep}}_{\mathbb{Z}[k^*]}(\mathbb{Z}[\mathcal{M}_\mathcal{X}])$ be the subsheaf of simplicial rings $B^{\text{cyc}}_{\mathbb{Z}[k^*]}(\mathbb{Z}[\mathcal{M}_\mathcal{X}^{\text{gp}}])$ such that its n-th simplex is the monoid ring of pullback of monoids

Let $\underline{P}HH(\mathcal{X})$ be the sheaf of simplicial rings $B_k^{\text{cyc}}(\mathcal{O}_{\overline{X}}) \otimes_{B_{\mathbb{Z}[k^*]}^{\text{cyc}}(\mathbb{Z}[\mathcal{M}_\mathcal{X}])} B_{\mathbb{Z}[k^*]}^{\text{rep}}(\mathbb{Z}[\mathcal{M}_\mathcal{X}]).$ Define $\underline{M}\mathcal{H}\mathcal{H}(\mathcal{X})$ to be the simplicial $\mathcal{O}_{\overline{X}}$ -module $\underline{M}\mathcal{O}(\mathcal{X})\otimes_{\mathcal{O}_{\overline{X}}}L\mathcal{H}\mathcal{H}(\mathcal{X})$.

For a strict normal crossing modulus pair $X, MHH(X)$ has a circle action Zariski locally by eq. [\(5.4\)](#page-13-3). By Lemma [3.3,](#page-3-0) we can define a circle action on $\mathcal{MHH}(\mathcal{X})$ globally, that is, $\underline{M}\mathcal{H}\mathcal{H}(\mathcal{X}) \in \text{Fun}(\Lambda^{\text{op}}, \mathcal{D}(\overline{X}))$ is a cyclic object of $\mathcal{D}(\overline{X})$.

Definition 6.2. For strict normal crossing modulus pair \mathcal{X} , modulus Hochschild homology $\overline{MHH(X)} \in \text{Fun}(B\mathbb{T}, \mathcal{D}(k))$ of X is an image of the composition of functors^{[3](#page-15-1)}

$$
\operatorname{Fun}(\Lambda^{\mathrm{op}},\mathcal{D}(\overline{X})) \xrightarrow{|-|} \operatorname{Fun}(B\mathbb{T},\mathcal{D}(\overline{X})) \xrightarrow{R\Gamma(\overline{X},-)^{-*}} \operatorname{Fun}(B\mathbb{T},\mathcal{D}(k)).
$$

The modulus (negative and periodic) cyclic homology of $\mathcal X$ are

$$
\underline{M}HC(\mathcal{X}) := (MHH(\mathcal{X}))_{h\mathbb{T}}
$$

$$
\underline{M}HC^{-}(\mathcal{X}) := (MHH(\mathcal{X}))^{h\mathbb{T}}
$$

$$
\underline{M}HP(\mathcal{X}) := (MHH(\mathcal{X}))^{t\mathbb{T}}.
$$

Let $\underline{M}HH_n(\mathcal{X}), \underline{M}HC_n(\mathcal{X}), \underline{M}HC_n^-(\mathcal{X}),$ and $\underline{M}HP_n(\mathcal{X})$ be their homologies.

Theorem 6.3. For every strict normal crossing modulus pair $\mathcal{X} = (\overline{X}, X^{\infty})$, we have the following isomorphisms^{[4](#page-15-2)}

$$
\begin{aligned} &\mathrm{Hom}_{\underline{\mathbf{M}\mathbf{D}\mathbf{M}^{\mathrm{eff}}_{k}}(M(\mathcal{X}),\prod_{p\in\mathbb{Z}}\underline{\mathbf{M}}\Omega^{p}[p-n])\cong\prod_{p\in\mathbb{Z}}H^{p-n}_{\mathrm{Zar}}(\overline{X},\underline{M}\Omega^{p}_{\mathcal{X}})\cong\underline{M}HH_{n}(\mathcal{X})\\ &\mathrm{Hom}_{\underline{\mathbf{M}\mathbf{D}\mathbf{M}^{\mathrm{eff}}_{k}}(M(\mathcal{X}),\prod_{p\in\mathbb{Z}}\underline{\mathbf{M}}\Omega^{\leq p}[2p-n])\cong\prod_{p\in\mathbb{Z}}\mathbb{H}^{2p-n}_{\mathrm{Zar}}(\overline{X},\underline{M}\Omega^{\leq p}_{\mathcal{X}})\cong\underline{M}HC_{n}(\mathcal{X})\\ &\mathrm{Hom}_{\underline{\mathbf{M}\mathbf{D}\mathbf{M}^{\mathrm{eff}}_{k}}(M(\mathcal{X}),\prod_{p\in\mathbb{Z}}\underline{\mathbf{M}}\Omega^{\geq p}[2p-n])\cong\prod_{p\in\mathbb{Z}}\mathbb{H}^{2p-n}_{\mathrm{Zar}}(\overline{X},\underline{M}\Omega^{\geq p}_{\mathcal{X}})\cong\underline{M}HC_{n}^{-}(\mathcal{X})\\ &\mathrm{Hom}_{\underline{\mathbf{M}\mathbf{D}\mathbf{M}^{\mathrm{eff}}_{k}}(M(\mathcal{X}),\prod_{p\in\mathbb{Z}}\underline{\mathbf{M}}\Omega^{\bullet}[2p-n])\cong\prod_{p\in\mathbb{Z}}\mathbb{H}^{2p-n}_{\mathrm{Zar}}(\overline{X},\underline{M}\Omega^{ \bullet}_{\mathcal{X}})\cong\underline{M}HP_{n}(\mathcal{X}). \end{aligned}
$$

Proof. The first isomorphisms are representability of these cohomologies described in Theorem [1.1](#page-1-2) and Theorem [3.5.](#page-3-1) The map $e: \mathcal{MHH}(\mathcal{X}) \to \prod_{p\geq 0} \underline{M}\Omega^p_{\mathcal{X}}[p]$: $a_0 \otimes \cdots \otimes a_n \mapsto a_0 da_1 \wedge \cdots \wedge da_n$ is an equivalence of the chain complexes of sheaves. So we obtain the second isomorphism for $\mathcal{M}HH(\mathcal{X})$. As observed in the proof of Theorem [5.2,](#page-13-0) $MHP(X)$ is equivalent to

$$
\text{Tot}^{\prod}(\underline{M}HH(\mathcal{X})[\bullet],B) \xrightarrow{c} \text{Tot}^{\prod}((\prod_{p\geq 0}R\Gamma(\overline{X},\underline{M}\Omega_{\mathcal{X}}^{p})[p])[\bullet],R\Gamma(\overline{X},d)),
$$

³We use homological conventions for $\mathcal{D}(k)$. So the objects are chain complexes, not cochain complexes.

⁴We use cohomological conventions for $\underline{MDM}_k^{\text{eff}}$. So the objects are cochain complexes, not chain conmplexes

but right hand side is $\prod_{p\in\mathbb{Z}} R\Gamma(\overline{X}, \underline{M}\Omega^*_\mathcal{X})^{-*}[2p]$. Since cyclic (negative cyclic) homology is a total complex of non-negative (non-positive) parts of bicomplex of periodic cyclic homology, we have equivalences

$$
\underline{M}HC(\mathcal{X}) \simeq \prod_{p \in \mathbb{Z}} R\Gamma(\overline{X}, \underline{M}\Omega_{\mathcal{X}}^{\leq p})^{-*}[2p]
$$

$$
\underline{M}HC^{-}(\mathcal{X}) \simeq \prod_{p \in \mathbb{Z}} R\Gamma(\overline{X}, \underline{M}\Omega_{\mathcal{X}}^{\geq p})^{-*}[2p]
$$

$$
\underline{M}HP(\mathcal{X}) \simeq \prod_{p \in \mathbb{Z}} R\Gamma(\overline{X}, \underline{M}\Omega_{\mathcal{X}}^{\bullet})^{-*}[2p]
$$

If we take homologies, we get the desired isomorphisms. $\hfill \Box$

REFERENCES

Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro-ku, Tokyo 152-8551, Japan Email address: sato.m.bj@m.titech.ac.jp