

# PENCILS ON GENERAL COVERS OF AN ELLIPTIC CURVE

ANDREAS LEOPOLD KNUITSEN AND MARGHERITA LELLI-CHIESA

ABSTRACT. We completely describe the Brill-Noether theory of pencils on general primitive covers of elliptic curves of any degree.

## 1. INTRODUCTION

In recent years much focus has been directed toward the Brill-Noether theory of covers of  $\mathbb{P}^1$ , see, e.g., [1, 13, 8, 11, 5]. Little is however known for covers of higher genus curves, even elliptic curves. In this paper we settle the Brill-Noether theory of *pencils* of a general primitive cover of any fixed elliptic curve.

Fix any smooth irreducible elliptic curve  $E$  over  $\mathbb{C}$ . A finite cover  $\varphi : C \rightarrow E$  is called *primitive* if it does not factor through another finite cover  $\varphi' : C \rightarrow E'$  of smaller degree to an elliptic curve  $E'$ . The *Hurwitz scheme*  $\mathcal{H}_{g,k}^{\text{prim}}(E)$  parametrizes primitive simply branched covers  $\varphi : C \rightarrow E$  of degree  $k$  from some smooth irreducible curve  $C$  of genus  $g \geq 2$ . We recall that  $\mathcal{H}_{g,k}^{\text{prim}}(E)$  is smooth and irreducible of dimension  $2g - 2$  [7, 3]. We denote by  $\mathcal{M}_{g,k}(E)$  the image under the natural forgetful map of  $\mathcal{H}_{g,k}^{\text{prim}}(E)$  in the moduli space  $\mathcal{M}_g$  of genus  $g$  curves. Since any cover can be composed with an automorphism of  $E$ , the locus  $\mathcal{M}_{g,k}(E)$  is irreducible of dimension  $2g - 3$ . It turns out (cf. Proposition 4) that a general member of  $\mathcal{M}_{g,k}(E)$  admits only one  $k : 1$  cover to  $E$  up to the automorphisms of  $E$  and, if  $g > 2$ , no finite primitive maps to other elliptic curves.

As the pullback of any  $g_2^1$  on  $E$  under a degree  $k$  cover of it defines a  $g_{2k}^1$  on the covering curve, any curve in  $\mathcal{M}_{g,k}(E)$  has gonality at most  $\min\left\{2k, \left\lfloor \frac{g+3}{2} \right\rfloor\right\}$  by standard Brill-Noether theory. We will prove the following fundamental result, which answers a natural question, yet hitherto unanswered:

**Theorem 1.** *Let  $\varphi : C \rightarrow E$  define a general point in the Hurwitz scheme  $\mathcal{H}_{g,k}^{\text{prim}}(E)$  for  $g \geq 2$  and  $k \geq 2$ . Let  $G_d^1(C)^{\text{nc}}$  denote the closure of the locus of linear series in  $G_d^1(C)$  that are base point free and not composed with  $\varphi$ . Then the following hold:*

- (i)  $G_d^1(C)^{\text{nc}}$  is nonempty if and only if either  $k \geq 3$  and  $\rho(g, 1, d) \geq 0$ , or  $k = 2$  and  $d \geq g - 1$ ;
- (ii) if nonempty,  $G_d^1(C)^{\text{nc}}$  has pure dimension  $\rho(g, 1, d) := -g + 2d - 2$ .

Part (ii), and thus the consequence that  $C$  has gonality  $\min\left\{2k, \left\lfloor \frac{g+3}{2} \right\rfloor\right\}$ , was recently proved in [10, Cor. 5.3]. Up to our knowledge, the only known part of (i) was the existence of a base point free  $g_{g-1}^1$  in the case  $k = 2$  (cf. [12] and references therein, where the result is obtained for any bielliptic curve and not only for a general one).

The above theorem enables us to fully describe the Brill-Noether theory of pencils on a general curve  $C$  in  $\mathcal{M}_{g,k}(E)$ , whose  $k : 1$  map onto  $E$  we denote by  $\varphi$ . For every

$2 \leq n \leq d/k$ , we denote by  $G_{d,n}^1(C)$  the closure in  $G_d^1(C)$  of the locus of linear series  $\mathbf{g}$  with  $s := d - nk$  base points  $p_1, \dots, p_s$  such that  $\mathbf{g}(-p_1 - \dots - p_s)$  is the pullback under  $\varphi$  of a  $g_n^1$  on  $E$ . It is straightforward that  $G_{d,n}^1(C)$  is irreducible of dimension

$$(1) \quad \dim G_{d,n}^1(C) = \dim G_n^1(E) + d - nk = d - (k-2)n - 3;$$

in particular, this does not depend on  $n$  in the case  $k = 2$ . When  $\dim G_{d,n}^1(C) \geq \rho(g, 1, d)$ , that is, in the range  $2 \leq n \leq \frac{g-1-d}{k-2}$ , Theorem 1 implies that  $G_{d,n}^1(C)$  is an irreducible component of  $G_d^1(C)$ . More precisely, we obtain the following results:

**Corollary 2.** *Fix  $k \geq 3, g \geq 2$  and let  $[C] \in \mathcal{M}_{g,k}(E)$  be general. Then  $C$  has gonality*

$$\text{gon } C = \min \left\{ 2k, \left\lfloor \frac{g+3}{2} \right\rfloor \right\},$$

and for every  $d \geq \text{gon } C$  one has

$$(2) \quad \dim G_d^1(C) = \max\{\rho(g, 1, d), d - 2k + 1\}.$$

More precisely, the following hold:

- (i) if  $\rho(g, 1, d) \geq 0$ , the irreducible components of  $G_d^1(C)$  are the components of  $G_d^1(C)^{nc}$  and the loci  $G_{d,n}^1(C)$  for  $2 \leq n \leq \frac{g-1-d}{k-2}$ ;
- (ii) if  $\rho(g, 1, d) < 0$ , the irreducible components of  $G_d^1(C)$  are the loci  $G_{d,n}^1(C)$  for  $2 \leq n \leq \frac{d}{k}$ .

In particular, when  $k \geq 3$  the Brill-Noether varieties  $G_d^1(C)$  are not necessarily equidimensional. This fact highlights a big difference with the  $k = 2$  case, that we therefore state separately:

**Corollary 3.** *Let  $[C] \in \mathcal{M}_{g,2}(E)$  be general with  $g \geq 2$ . Then  $C$  has gonality*

$$\text{gon } C = \min \left\{ 4, \left\lfloor \frac{g+3}{2} \right\rfloor \right\},$$

and for every  $d \geq \text{gon } C$  the Brill-Noether variety  $G_d^1(C)$  is equidimensional of dimension

$$(3) \quad \dim G_d^1(C) = \max\{\rho(g, 1, d), d - 3\}.$$

More precisely, the following hold:

- (i) if  $d > g - 1$ , then  $G_d^1(C)$  coincides with  $G_d^1(C)^{nc}$ ;
- (ii) if  $d = g - 1$ , the irreducible components of  $G_d^1(C)$  are the components of  $G_d^1(C)^{nc}$  and the loci  $G_{d,n}^1(C)$  for  $2 \leq n \leq \frac{d}{2}$ ;
- (iii) if  $d < g - 1$ , the irreducible components of  $G_d^1(C)$  are the loci  $G_{d,n}^1(C)$  for every  $2 \leq n \leq \frac{d}{2}$ .

When  $2k < \left\lfloor \frac{g+3}{2} \right\rfloor$ , Corollaries 2 and 3 (or [10, Cor. 5.3]) imply that a general curve  $[C] \in \mathcal{M}_{g,k}(E)$  does not satisfy Aprodu's *linear growth condition* [1]. Green's Conjecture for these curves is nevertheless satisfied by the recent result [10, Thm. 5.4] (the case  $k = 3$  had been done in in [2] the case  $k = 2$  is trivial, as the curve  $C$  is 4-gonal and thus Green's conjecture reads like the ideal of the canonical curve  $C \subset \mathbb{P}^{g-1}$  being generated by quadrics).

## 2. PROOFS OF THE RESULTS

First of all, we prove what anticipated at the beginning of the introduction:

**Proposition 4.** *Fix  $g \geq 2, k \geq 2$  and any elliptic curve  $E$ . Then the following hold:*

- (i) *a general fiber of the forgetful map  $\mathcal{H}_{g,k}^{\text{prim}}(E) \rightarrow \mathcal{M}_{g,k}(E)$  has dimension 1 (corresponding to the automorphisms of  $E$ );*
- (ii) *if  $g \geq 3$ , then a general  $[C] \in \mathcal{M}_{g,k}(E)$  does not lie in any other  $\mathcal{M}_{g,k'}(E')$  (where  $k' \geq 2$  and  $E'$  is an elliptic curve possibly coinciding with  $E$  if  $k \neq k'$ ).*

*Proof.* Let  $[\varphi : C \rightarrow E] \in \mathcal{H}_{g,k}^{\text{prim}}(E)$  be general and, by contradiction, assume the existence of another primitive simply branched map  $\varphi' : C \rightarrow E'$  of degree  $k' \geq 2$  to an elliptic curve  $E'$  (where  $\varphi$  and  $\varphi'$  are required to be distinct up to the automorphisms of  $E$  in the case  $E' = E$  and  $k' = k$ ). We stress that both  $\varphi$  and  $\varphi'$  are primitive and so they are not composed one with the other. The pair  $(\varphi, \varphi')$  then defines a map from  $C$  to the abelian surface  $E \times E'$ , that can be factored as  $C \xrightarrow{\alpha} D \xrightarrow{\psi} E \times E'$ , where  $D$  is a curve of genus  $h \leq g$  (with  $D = C$  when  $h = g$ ),  $\alpha$  is a finite map of degree  $1 \leq n < \min\{k, k'\}$  dividing both  $k$  and  $k'$ , and the map  $\psi$  is birational. In particular, the image  $\psi(D)$  is a curve of numerical class  $lE + l'E'$  with  $l = k/n \geq 2$  and  $l' = k'/n \geq 2$ . By [6, Prop. 4.16], integral genus  $h$  curves in  $E \times E'$  of this numerical class move in dimension  $h$ . Modding out by the automorphisms of  $E \times E'$ , we obtain a family of dimension  $\leq h - 2$  of curves  $D$  of genus  $h \geq 2$  admitting a birational map to a curve of class  $lE + l'E'$  in  $E \times E'$ . Letting  $E'$  vary in moduli and denoting by  $\mathcal{H}_{g,n}(D)$  the Hurwitz scheme of genus  $g$  and degree  $n$  simply branched covers of a curve  $D$  as above, one computes that curves  $C$  admitting a pair  $(\varphi, \varphi')$  as described move in dimension at most

$$1 + h - 2 + \dim \mathcal{H}_{g,n}^{\text{prim}}(D) = h - 1 + 2g - 2 - n(2h - 2),$$

which is  $< 2g - 3 = \dim \mathcal{M}_{g,k}(E)$  unless  $n = 1$  and  $h = g = 2$ .  $\square$

To prove our main results, we now consider the surface

$T := E \times \mathbb{P}^1$  and let  $R$  and  $E$  denote the classes of the obvious sections of  $T$ , so that  $K_T \equiv -2E$ ,

Let  $V_g(k, d)$  denote the Severi variety of irreducible nodal curves of genus  $g$  on  $T$  of class  $kE + dR$ . Similarly, we denote by  $V^g(k, d)$  the equigeneric locus of integral curves of genus  $g$  on  $T$  of class  $kE + dR$ , so that one has an obvious inclusion  $V_g(k, d) \subset V^g(k, d)$ . Let  $\mathcal{M}_g(k, d)$  be the coarse moduli space of genus  $g$  stable maps with image of class  $kE + dR$ , and denote by  $\mathcal{M}_g(k, d)^{\text{sm}}$  the closure of the locus of maps in  $\mathcal{M}_g(k, d)$  that are smoothable, that is, can be deformed to a map from a nonsingular curve, birational onto its image (cf. [14]). By sending a stable map  $f : C \rightarrow T$  to its image  $f(C) = \overline{C}$ , one obtains a dominant morphism from the semi-normalization of  $\mathcal{M}_g(k, d)^{\text{sm}}$  to  $\overline{V^g(k, d)}$  [9, I.6]. The following result is of independent interest (part (ii) was already proved in [10, Prop. 5.2]):

**Proposition 5.** *For every  $1 \leq g \leq d(k - 1) + 1$  the following hold:*

- (i) *the variety  $V_g(k, d)$  is nonempty, smooth of dimension  $2d + g - 1$  and dense in the equigeneric locus  $V^g(k, d)$ ;*
- (ii)  *$\mathcal{M}_g(k, d)^{\text{sm}}$  is generically reduced of dimension  $2d + g - 1$ .*

*Proof.* Since  $-K_T \cdot (kE + dR) = 2d \geq 4$ , it is well-known that  $\mathcal{M}_g(k, d)^{sm}$  and  $V^g(k, d)$  are generically reduced (as a general stable map in any irreducible component is unramified) and equidimensional of dimension  $2d + g - 1$  as soon as they are nonempty, and that  $V_g(k, d)$  is dense in  $V^g(k, d)$ , cf., e.g., [4, proof of Prop. 2.2]. For  $[\overline{C}] \in V_g(k, d)$ , we denote by  $N$  the scheme of nodes of  $\overline{C}$  and by  $\nu : C \rightarrow \overline{C} \subset T$  the composition of the normalization map with the inclusion of  $\overline{C}$  in  $T$ . Since the tangent space of  $V_g(k, d)$  at the point  $[\overline{C}]$  can be identified with  $H^0(T, \mathcal{O}_T(\overline{C}) \otimes \mathcal{J}_N)$  and this is isomorphic to the  $2d + g - 1$ -dimensional space  $H^0(C, \mathcal{N}_\nu) = H^0(C, \omega_C \otimes \nu^*(-K_T))$  [6, §3.1], we conclude that  $V_g(k, d)$  is smooth. As a consequence, to prove nonemptiness we first note that  $\mathcal{M}_1(k, d)^{sm}$  is nonempty (just pick any smooth elliptic curve admitting a degree  $k$  map onto  $E$ , together with a general  $g_d^1$  on it). It follows that also  $V_1(k, d)$  is nonempty and its smoothness implies that the nodes of the curves can be smoothed independently; as a consequence,  $V_g(k, d)$  is nonempty for all  $1 \leq g \leq p_a(kE + dR) = d(k - 1) + 1$ .  $\square$

In the sequel we will make use of the following vanishing:

**Lemma 6.** *Let  $N$  be the scheme of the  $\delta := d(k - 1) + 1 - g$  nodes of a curve  $\overline{C}$  in  $V_g(k, d)$ . Then*

$$h^0(T, \mathcal{O}_T(\overline{C} + K_T) \otimes \mathcal{J}_N) = g - 1.$$

*Proof.* Let  $\mu_N : \widetilde{T}_N \rightarrow T$  be the blow up at  $N$ , with total exceptional divisor  $\mathfrak{e}$ , and let  $C$  be the normalization of  $\overline{C}$ , also coinciding with the strict transform of  $\overline{C}$  under  $\mu_N$ . Since  $C^2 = \overline{C}^2 - \delta = 2kd - (d(k - 1) + 1 - g) = kd + d + g - 1 > 0$ , we have that  $C$  is big and nef, so that

$$h^0(\mathcal{O}_{\widetilde{T}_N}(C + K_{\widetilde{T}_N})) = \chi(\mathcal{O}_{\widetilde{T}_N}(C + K_{\widetilde{T}_N})) = \frac{1}{2}C \cdot (C + K_{\widetilde{T}_N}) = g - 1.$$

But  $C + K_{\widetilde{T}_N} \sim \mu_N^* \overline{C} - 2\mathfrak{e} + \mu_N^* K_T + \mathfrak{e} = \mu_N^*(\overline{C} + K_T) - \mathfrak{e}$ , and the result follows.  $\square$

The following result is used to specialize the nodes of curves in  $V_g(k, d)$  to the nodes of some reducible curves.

**Lemma 7.** *Fix  $g \geq 5$ ,  $k \geq 3$ ,  $d \geq (g - 1)/2$  and two integers  $l_1, l_2$  such that  $l_1 + l_2 = 2d + 1 - g$  and  $0 \leq l_i \leq d - 2$  for  $i = 1, 2$ . Let*

$$f : \widetilde{X} = \widetilde{D} \cup_{Z_1} E_1 \cup_{Z_2} E_2 \rightarrow X = D \cup E_1 \cup E_2 \subset T$$

*be a genus  $g$  stable map such that  $\widetilde{D}$  is a smooth elliptic curve of genus 1 with image  $D \equiv (k - 2)E + dR$ , the curves  $E_1, E_2$  are mapped isomorphically to elements of  $|E|$ , and  $\widetilde{D}$  and  $E_i$  intersect transversally at  $d - l_i$  points denoted by  $Z_i$  for  $i = 1, 2$ . Then  $[f] \in \mathcal{M}_g(k, d)^{sm}$  and  $D + E_1 + E_2$  lies in the closure of  $V_g(k, d)$ .*

*Proof.* By Proposition 5, it is enough to show that  $f$  is smoothable. We denote by  $f_D := f|_{\widetilde{D}} : \widetilde{D} \rightarrow D \subset T$ ,  $f_{E_i} : E_i \rightarrow E_i \subset T$  for  $i = 1, 2$  and  $f_1 := f|_{\widetilde{D} \cup_{Z_1} E_1} : \widetilde{D} \cup_{Z_1} E_1 \rightarrow D \cup E_1 \subset T$  the restrictions of  $f$ . It is enough to prove that  $[f] \in \mathcal{M}_g(k, d)^{sm}$  when  $[f_D]$  is general in any irreducible component of  $\mathcal{M}_1(k - 2, d)^{sm}$ . The map  $f_1$  defines a point of  $\mathcal{M}_{1+d-l_1}(k - 1, d)$ , whose expected dimension is  $3d - l_1$ . As soon as  $l_1 \leq d - 2$ , one has  $3d - l_1 > \dim(\mathcal{M}_1(k - 2, d)^{sm} \times \mathcal{M}_1(1, 0)) = 2d + 1$ , where the equality follows from Proposition 5; as a consequence, the map  $f_1$  is smoothable, that is, it lies in a component  $\widetilde{\mathcal{M}}$  of  $\mathcal{M}_{1+d-l_1}(k - 1, d)^{sm}$ . Take now a genus  $g$ -stable map

$h : \widetilde{Y} \cup_Z E \rightarrow Y \cup E \subset T$  such that  $h_Y := h|_{\widetilde{Y}}$  corresponds to a general point of  $\widehat{\mathcal{M}}$ , the restriction  $h_E := h|_E$  defines a point of  $\mathcal{M}_1(1, 0)$  and  $Z := Y \cap E$  consists of  $d - l_2$  nodes. Then  $h$  defines a point of  $\mathcal{M}_g(k, d)$ , which has expected dimension equal to  $2d + g - 1$ . Since  $2d + g - 1 > \dim(\widehat{\mathcal{M}} \times \mathcal{M}_1(1, 0)) = 3d - l_1 + 1$  for  $l_2 \leq d - 2$ , we conclude that the map  $h$  is smoothable and thus the same holds for  $f$ .  $\square$

**Proposition 8.** *For every  $2 \leq g \leq d(k - 1) + 1$  and  $d \geq 2k$ , the forgetful map  $\psi_{g,k,d} : \mathcal{M}_g(k, d)^{sm} \dashrightarrow \mathcal{M}_g$  dominates  $\mathcal{M}_{g,k}(E)$  when  $\rho(g, 1, d) \geq 0$ .*

*Proof.* Let  $f : C \rightarrow T$  define a general point in some component of  $\mathcal{M}_g(k, d)^{sm}$  so that the image  $\overline{C} := f(C)$  is a curve with  $\delta := d(k - 1) + 1 - g$  nodes, that we denote by  $N$ . As in the proof of Lemma 6, let  $\mu_N : \widetilde{T}_N \rightarrow T$  denote the blow up at  $N$  and by  $\mathfrak{e}$  its exceptional divisor. Since  $A := f^*(\mathcal{O}_T(E)) \in G_d^1(C)$  moves in dimension  $\geq \rho(g, 1, d)$ , the fiber  $\psi_{g,k,d}^{-1}([C])$  has dimension

$$\geq \rho(g, 1, d) + \dim \text{Aut}(T) = \rho(g, 1, d) + 4 = \dim \mathcal{M}_g(k, d)^{sm} - \dim \mathcal{M}_{g,k}(E).$$

We want to show that equality holds.

Let  $\mathcal{N}_f$  denote the normal sheaf of  $f$ , fitting into

$$0 \longrightarrow \mathcal{T}_C \longrightarrow f^* \mathcal{T}_T \longrightarrow \mathcal{N}_f \longrightarrow 0.$$

The coboundary map

$$H^0(\mathcal{N}_f) \longrightarrow H^1(\mathcal{T}_C)$$

is the differential

$$d\psi_{g,k,d} : T_{[f]} \mathcal{M}_g(k, d)^{sm} \longrightarrow T_{[C]} \mathcal{M}_g$$

of the forgetful map  $\psi_{g,k,d}$  at  $[f]$ . Its kernel is isomorphic to

$$H^0(f^* \mathcal{T}_T) \simeq H^0(f^*(\mathcal{O}_T \oplus \mathcal{O}_T(2E))) \simeq H^0(\mathcal{O}_C) \oplus H^0(\mathcal{O}_C(2A)),$$

which has dimension

$$1 + h^0(\mathcal{O}_C(2A)) = 1 + (2d + 1 - g + h^0(\omega_C - 2A)) = \rho(g, 1, d) + 4 + h^0(\omega_C - 2A).$$

To finish the proof of the proposition, it will therefore be enough to prove that when  $\rho(g, 1, d) \geq 0$  the vanishing

$$(4) \quad h^0(\omega_C - 2A) = 0$$

holds for a general  $(C, A)$  where  $C$  is the domain of a general  $[f]$  in some component of  $\mathcal{M}_g(k, d)^{sm}$  and  $A = f^*(\mathcal{O}_T(E))$ . The vanishing holds for degree reasons if  $d \geq g$ , so we may assume that  $4 \leq 2k \leq d < g$ . The case  $k = 2$  thus yields  $\delta = d + 1 - g = 0$ ,  $d = g - 1$  and  $\rho(g, 1, d) = g - 4 = d - 3 \geq 1$ . The vanishing (4) thus holds as soon as  $A$  is not a theta characteristic on  $C$  and we may assume this is not the case as  $A$  moves in a positive dimensional family. From now on we will therefore assume  $6 \leq 2k \leq d < g$ .

From the short exact sequence

$$0 \longrightarrow \mathcal{O}_{\widetilde{T}_N}(K_{\widetilde{T}_N} - 2\mu_N^* E) \xrightarrow{s_N} \mathcal{O}_{\widetilde{T}_N}(K_{\widetilde{T}_N} + C - 2\mu_N^* E) \longrightarrow \omega_C - 2A \longrightarrow 0$$

and the facts that  $K_{\widetilde{T}_N} \sim \mu_N^* K_T + \mathfrak{e}$  and  $C \sim \mu_N^* \overline{C} - 2\mathfrak{e}$ , we see that (4) is equivalent to the two conditions

$$(5) \quad h^0(T, \mathcal{O}_T(\overline{C} - 2E + K_T) \otimes \mathcal{J}_N) = 0$$

and

$$(6) \quad H^1(s_N) : H^1(T, \mathcal{O}_T(-4E)) \longrightarrow H^1(T, \mathcal{O}_T(\overline{C} - 2E + K_T) \otimes \mathcal{J}_N) \text{ is injective}$$

Let  $V'$  be a component of  $\overline{V^d(k, d)}$  containing a reducible curve  $D + E_1 + E_2$  as in Lemma 7, and let  $\widehat{V}$  be a component of  $\overline{V^g(k, d)}$  containing  $V'$ . We will prove the vanishings (5) and (6) for the scheme of nodes  $N$  of a general curve  $[\overline{C}] \in \widehat{V}$ .

To prove (5), we may specialize  $N$  to a scheme of nodes of a reducible curve  $D + E_1 + E_2$  (as in Lemma 7) containing all the nodes  $N_D$  of  $D$  plus  $l_1 \leq d - 2$  nodes of  $D \cap E_1$  and  $l_2 \leq d - 2$  nodes of  $D \cap E_2$  with  $l_1 + l_2 = 2d + 1 - g$ . By Lemma 6 applied to the curve  $[D] \in V_1(k - 2, d)$ , we obtain the vanishing  $h^0(\mathcal{O}_T(\overline{C} - 2E + K_T) \otimes \mathcal{J}_{N_D}) = h^0(\mathcal{O}_T(D + K_T) \otimes \mathcal{J}_{N_D}) = 0$ , so that the vanishing (5) also holds.

We next specialize  $N$  to a length- $\delta$  subscheme  $N'$  of the scheme of nodes  $N^+$  of a general curve  $[\overline{C}^+] \in V'$ , whose normalization is denoted by  $C^+$ . Since the genus of  $\overline{C}^+$  is  $d$ , the vanishing  $h^0(\omega_{C^+} - 2A) = 0$  holds for degree reasons, whence also

$$(7) \quad h^0(T, \mathcal{O}_T(\overline{C} - 2E + K_T) \otimes \mathcal{J}_{N^+}) = 0$$

and

$$(8) \quad H^1(s_{N^+}) : H^1(T, \mathcal{O}_T(-4E)) \longrightarrow H^1(T, \mathcal{O}_T(\overline{C} - 2E + K_T) \otimes \mathcal{J}_{N^+}) \text{ is injective.}$$

Let  $W \subset N^+$  be any subset of  $g - d$  points and set  $N' := N^+ \setminus W$ . From the short exact sequence

$$0 \longrightarrow \mathcal{J}_{N^+} \longrightarrow \mathcal{J}_{N'} \longrightarrow \mathcal{O}_W \longrightarrow 0$$

we obtain the commutative diagram with exact rows:

$$\begin{array}{ccc} H^1(\mathcal{O}_T(-4E)) & \xlongequal{\quad} & H^1(\mathcal{O}_T(-4E)) \\ \downarrow H^1(s_{N^+}) & & \downarrow H^1(s_{N'}) \\ H^0(\mathcal{O}_W) \xrightarrow{\alpha_W} H^1(\mathcal{O}_T(\overline{C} - 2E + K_T) \otimes \mathcal{J}_{N^+}) & \longrightarrow & H^1(\mathcal{O}_T(\overline{C} - 2E + K_T) \otimes \mathcal{J}_{N'}) \longrightarrow 0. \end{array}$$

We will show that one may choose  $W \subset N^+ = \{n_1, \dots, n_{d(k-2)+1}\}$  appropriately so that

$$(9) \quad \text{Im } \alpha_W \cap \text{Im } H^1(s_{N^+}) = \{0\}.$$

By (8), this will imply the injectivity of  $H^1(s_{N'})$  and thus condition (6) for the nodes  $N$  of a general curve  $[\overline{C}] \in \widehat{V} \supset V'$ .

We note that  $h^1(\mathcal{O}_T(-4E)) = 3$ , so that  $\text{Im } H^1(s_{N^+}) \simeq \mathbb{C}^3$ . By (7) we have the exact sequence

$$0 \longrightarrow H^0(\mathcal{O}_T(\overline{C} - 2E + K_T)) \longrightarrow H^0(\mathcal{O}_{N^+}) \xrightarrow{\alpha_{N^+}} H^1(\mathcal{O}_T(\overline{C} - 2E + K_T) \otimes \mathcal{J}_{N^+});$$

since  $\overline{C} - 2E + K_T \equiv (k - 2)E + dR + K_T$  has no higher cohomology for  $k \geq 3$ , the image of  $\alpha_{N^+}$  has dimension

$$\begin{aligned} \deg N^+ - \chi(\mathcal{O}_T(\overline{C} - 2E + K_T)) &= d(k - 2) + 1 - d(k - 3) = d + 1 \\ &= \rho(g, 1, d) + g - d + 3 \geq g - d + 3. \end{aligned}$$

Since  $H^0(\mathcal{O}_{N^+}) \simeq \bigoplus_{i=1}^{d(k-2)+1} H^0(\mathcal{O}_{n_i})$ , we may choose a subscheme  $W \subset N^+$  of degree  $g - d$  so that (9) is satisfied.  $\square$

We can now prove the main theorem and its corollaries.

*Proof of Theorem 1.* Let  $[\varphi : C \rightarrow E] \in \mathcal{H}_{g,k}^{\text{prim}}(E)$  be general. First of all, we note that  $\varphi$  does not factor as the composition  $C \xrightarrow{\alpha} C' \xrightarrow{\varphi'} E$  of a cover  $\alpha$  of degree  $n \geq 2$  dividing  $k$  onto a curve  $C'$  of genus  $h \geq 2$ . This follows because, denoting by  $\mathcal{H}_{g,n}(C')$  the Hurwitz scheme of genus  $g$  and degree  $n$  simply branched covers of a curve  $C'$  as above, one computes that

$$\dim \mathcal{H}_{h,k/n}^{\text{prim}}(E) + \dim \mathcal{H}_{g,n}(C') = 2h - 2 + 2g - 2 - n(2h - 2) < 2g - 2 = \dim \mathcal{H}_{g,k}^{\text{prim}}(E).$$

We now assume that  $G_d^1(C)^{nc}$  is nonempty. Then we have a morphism  $C \rightarrow T$ , birational onto its image  $[\overline{C}] \in V^g(k, d)$ . Let  $\widetilde{\mathcal{G}}_d^1$  denote the family of pairs  $\{\varphi : C \rightarrow E, \mathfrak{g}\}$  (up to isomorphism) such that  $[\varphi] \in \mathcal{H}_{g,k}^{\text{prim}}(E)$  and  $\mathfrak{g} \in G_d^1(C)^{nc}$ . Proposition 5 then yields

$$\begin{aligned} \dim \widetilde{\mathcal{G}}_d^1 &\leq \dim V^g(k, d) - \dim \text{Aut } \mathbb{P}^1 = (2d + g - 1) - 3 \\ &= 2d + g - 4 = \rho(g, 1, d) + 2g - 2. \end{aligned}$$

The forgetful map  $\widetilde{\mathcal{G}}_d^1 \rightarrow \mathcal{H}_{g,k}^{\text{prim}}(E)$  is dominant by assumption, and thus its fiber  $G_d^1(C)^{nc}$  over a general  $[\varphi]$  has dimension  $\leq \rho(g, 1, d)$  and equality follows from Brill-Noether theory.

It remains to show that  $G_d^1(C)^{nc}$  is nonempty whenever  $\rho(g, 1, d) \geq 0$ . This is clearly true if  $d < 2k$  and follows from Proposition 8 when  $d \geq 2k$ .  $\square$

*Proof of Corollaries 2 and 3.* Take a general  $[C] \in \mathcal{M}_{g,k}(E)$  so that there exists a primitive degree  $k$  cover  $\varphi : C \rightarrow E$ . Let  $\mathfrak{g}$  be a  $g_h^1$  on  $C$  computing the gonality and assume  $h < \frac{g+3}{2}$ . By Theorem 1,  $\mathfrak{g}$  should be composed with  $\varphi$  and thus  $h = 2k$ . Let us now fix  $d \geq \text{gon } C$  and take a component  $G$  of  $G_d^1(C)$ , which by standard Brill-Noether theory has dimension  $\geq \rho(g, 1, d)$ . If  $G$  is contained in the closure of  $G_d^1(C)^{nc}$ , then Theorem 1 yields  $\dim G = \rho(g, 1, d)$  and, if  $k = 2$ , then  $d \geq g - 1$ . Again Theorem 1 excludes the possibility that a general  $\mathfrak{g} \in G$  has some base points  $p_1, \dots, p_s$  and  $\mathfrak{g}(-p_1 - \dots - p_s) \in G_{d-s}^1(C)^{nc}$ . It remains to treat the case where  $G$  coincides with  $G_{d,n}^1(C)$ . When  $k = 2$  the dimension of  $G_{d,n}^1(C)$  is independent on  $n$ , cf. (1), thus implying (3). For  $k > 2$ , the maximal dimension is instead obtained for  $n = 2$ , cf. again (1), and this yields (2).

As concerns points (i),(ii) in Corollary 2 and (i),(ii),(iii) in Corollary 3, we stress that  $G_d^1(C)^{nc}$  is union of irreducible components of  $G_d^1(C)$  as soon as it is nonempty. Furthermore, since  $G_d^1(C)^{nc}$  has pure dimension  $\rho(g, 1, d)$ , the locus  $G_{d,n}^1(C)$  is not contained in  $G_d^1(C)$  as soon as its dimension computed in (1) is  $\geq \rho(g, 1, d)$ . If  $G_{d,n}^1(C)$  were contained in some other  $G_{d,m}^1(C)$  with  $n \neq m$ , by dimensional reasons one should have  $n > m$ ; but then a general linear series in  $G_{d,m}^1(C)$  would have more base points than a general element of  $G_{d,n}^1(C)$ , thus contradicting the containment. As a consequence, the locus  $G_{d,n}^1(C)$  is an irreducible component of  $G_d^1(C)$  as soon as its dimension is  $\geq \rho(g, 1, d)$ .  $\square$

## REFERENCES

- [1] M. Aprodu, *Remarks on syzygies of d-gonal curves*, Math. Res. Lett. **12** (2005), 387–400.
- [2] M. Aprodu, G. Farkas, *Green's Conjecture for general covers*, Contemp. Math. **564** (2012) 211–226.

- [3] G. T. Bujokas, *Covers of an Elliptic Curve  $E$  and Curves in  $E \times \mathbb{P}^1$* , ProQuest LLC, Ann Arbor, MI, 2015, 97 pp.
- [4] L. Caporaso, J. Harris, *Counting plane curves of any genus*, Invent. Math. **131** (1998), 345–392.
- [5] K. Cook-Powell, D. Jensen, *Components of Brill-Noether loci for curves with fixed gonality*, Michigan Math. J. **71** (2022), 19–45.
- [6] T. Dedieu, E. Sernesi, *Equigeneric and equisingular families of curves on surfaces*, Publ. Mat. **61** (2017), 175–212.
- [7] D. Gabai, W. Kazez, *The classification of maps of surfaces*, Invent. Math. **90** (1987), 219–242.
- [8] D. Jensen, D. Ranganathan, *Brill-Noether theory for curves of a fixed gonality*, Forum Math. Pi **9** (2021), Paper No. e1, 33 pp.
- [9] J. Kollár, *Rational Curves on Algebraic Varieties*, Springer-Verlag, Germany, 1996.
- [10] M. Kemeny, *Projecting syzygies of curves*, Alg. Geom. **7** (2020), 561–580.
- [11] H. K. Larson, *A refined Brill-Noether theory over Hurwitz spaces*, Invent. Math. **224** (2021), 767–790.
- [12] S. S. Park, *Existence of a base-point-free pencils of degree  $g - 1$  on bielliptic curves*, Osaka J. of Math. **40** (2003), 279–285.
- [13] N. Pflueger, *Brill-Noether varieties of  $k$ -gonal curves*, Adv. Math. **312** (2017), 46–63.
- [14] R. Vakil, *A Tool for Stable Reduction of Curves on Surfaces*, Advances in Algebraic Geometry Motivated by Physics, 145–154, Amer. Math. Soc., 2001.

A. L. KNUTSEN, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BERGEN, POSTBOKS 7800, 5020 BERGEN, NORWAY

*Email address:* andreas.knutsen@math.uib.no

M. LELLI-CHESA, DIPARTIMENTO DI MATEMATICA E FISICA, UNIVERSITÀ DEGLI STUDI ROMA TRE, LARGO SAN LEONARDO MURIALDO 1, 00146 ROMA, ITALY

*Email address:* margherita.lellichiesa@uniroma3.it