REMARKABLE STRUCTURES IN INTEGRABLE PROBABILITY, I: MAX-INDEPENDENCE STRUCTURES

YACINE BARHOUMI-ANDRÉANI

ABSTRACT. We analyse in a systematic way the occurrences of a remarkable structure in the theory of integrable probability that we call a "max-independence structure", when random variables are constructed as a maximum of a sequence of independent random variables.

The list of treated examples contains : the GUE and GOE Tracy-Widom distributions, the extreme eigenvalues/eigenangles of random hermitian/unitary matrices (and in particular the historical example of the GUE extreme eigenvalues), the Hopf-Cole solution to the KPZ equation with Dirac initial condition (continuum random polymer) and the symmetric Schur measure. In this last case, the largest part of the underlying random partition is the maximum of an i.i.d randomisation of the deterministic sequence of negative integers. In the case of the GUE Tracy-Widom distribution, the laws of the random variables use a rescaling of the prolate hyperspheroidal wave functions that were introduced by Heurtley and Slepian in the context of circular optical mirrors, and in the case of the GUE, the distributions use the sigma-form of the Painlevé IV equation.

To illustrate the utility of such a structure, we rescale the largest eigenvalue of the GUE written as a maximum of N independent random variables with the classical Poisson approximation for sums of indicators. We use for this the Okamoto-Noumi-Yamada theory of the sigma-form of the Painlevé equation applied to random matrix theory by Forrester-Witte. By doing so, we find a new expression for the cumulative distribution function of the GUE Tracy-Widom distribution which is shown to be equivalent to the classical one using manipulations à la Forrester-Witte.

CONTENTS

1. Introduction	2
2. \mathcal{TW}_2 is a maximum of independent random variables	7
3. Max-independence structure in random matrix ensembles	22
4. Max-independence structure in the symmetric Schur measure	40
Appendix A. Symmetric functions	52
Appendix B. Prolate hyperspheroidal wave functions	56
Acknowledgements	59
Dedicatory	59
References	59

Date: January 17, 2024.

²⁰¹⁰ Mathematics Subject Classification. 60L70, 60F05, 15B52, 05E05, 47B35, 33E10, 33E17, 33E30.

1. INTRODUCTION

1.1. Motivations. The classical Tracy-Widom distributions are indexed by a parameter $\beta \in \{1, 2, 4\}$. Of particular interest is the case $\beta = 2$ (*GUE* Tracy-Widom distribution) that initially appeared in the fluctuations of the largest eigenvalue of a *GUE* random matrix [122, 123] and that meanwhile exhibited a universal feature in a variety of models: longuest increasing subsequence of a random uniform permutation [6], last passage percolation [73], log-Gamma polymer partition function [25, 43, 89, 92], asymmetric simple exclusion process [125, 126, 127], value in 0 of the Hopf-Cole solution to the KPZ equation [3, 28, 38, 48, 111, 112, 113], O'Connell-Yor semidiscrete directed polymer partition function [93, 24, 71], etc. Surveys documenting this rich universally class include [75, 76, 84, 85, 26, 34] and we refer to them for further references.

So far, three descriptions of the *GUE* Tracy-Widom distribution were given:

(1) its probability cumulative function is given by a Fredholm determinant [123] under the form

$$\mathbb{P}(\mathcal{TW}_2 \leqslant s) := \det(I - \mathbf{K}_{\operatorname{Ais}})_{L^2(\mathbb{R}_+)}$$
(1)

Here, \mathcal{TW}_2 is a random variable Tracy-Widom-distributed, \mathbf{K}_{Ai_s} is an operator acting on $L^2(\mathbb{R}_+)$ according to $\mathbf{K}_{Ai_s}f = \int_{\mathbb{R}_+} K_{Ai_s}(\cdot, y)f(y)dy$ (for all $f \in L^2(\mathbb{R}_+)$) and K_{Ai_s} is the "kernel"/function given by

$$K_{\operatorname{Ai}_{s}}(x,y) := \int_{\mathbb{R}_{+}} \operatorname{Ai}_{s}(x+u) \operatorname{Ai}_{s}(y+u) du, \qquad \operatorname{Ai}_{s} := \operatorname{Ai}(\cdot + s)$$
(2)

The Airy function Ai is the solution of the ordinary differential equation y''(t) = ty(t) with initial condition $y(0) = 3^{-2/3}\Gamma(\frac{2}{3})^{-1}$ and decay at infinity satisfying $y(x) \to 0$ when $x \to +\infty$. An integral representation of this function is given by

$$\operatorname{Ai}(t) = \int_{\mathbb{R}} \cos\left(t\xi + \frac{\xi^3}{3}\right) \frac{d\xi}{2\pi}$$

(2) This last probability cumulative function is also given by

$$\mathbb{P}(\mathcal{TW}_2 \leqslant s) := \exp\left(-\int_{\mathbb{R}} (x-t)_+ q(t)^2 dt\right), \qquad x_+ := \max\{x, 0\}$$
(3)

where q is the Hastings-McLeod solution of the Painlevé II equation given by¹ [123, (1.11), (1.12)]

$$q''(t) = tq(t) + 2q(t)^3 + \alpha, \qquad \alpha = 0, \qquad q(t) \mathop{\sim}_{t \to +\infty} \operatorname{Ai}(t), \qquad q(t) \mathop{\sim}_{t \to -\infty} \sqrt{-\frac{t}{2}}$$
(4)

The Painlevé II ODE admits moreover a representation as a solution of an integrable PDE given in terms of a Lax pair/isomonodromy problem discovered by

¹The general Painlevé II equation is $q''(t) = tq(t) + 2q(t)^3 + \alpha$ for $\alpha \in \mathbb{R}$; the Hastings-McLeod equation is $q''(t) = tq(t) + 2q(t)|q(t)|^{\beta}$. The Hastings-McLeod solution corresponds to $\alpha = 0/\beta = 2$ and is the only case where solutions can be bounded on \mathbb{R} , see [68].

Flaschka and Newell [53]/[105, 32.4 (iii) p. 728]. Bloemendal and Virag proved that the solution of this PDE involving half of this Lax pair can be expressed as a probability of non-explosion for the solution of a stochastic Riccatti equation related with limits of spiked matrix models [15, (1.7), (1.8)] (see also the introduction of [35]).

(3) A last description uses the Edelman-Sutton theory of stochastic operators [50, 119] formalised by Ramirez, Rider and Virag [108] and is valid not only for the classical cases $\beta \in \{1, 2, 4\}$ but for for all $\beta > 0$. Define the Stochastic Airy Operator by

$$SAO_{\beta} := \frac{d^2}{dx^2} - x - \frac{2}{\sqrt{\beta}}\dot{B}(x)$$

where \dot{B} is a white noise (the distributional derivative of a Brownian motion), i.e. a Gaussian distribution whose covariance "function" is given by $\mathbb{E}(\dot{B}(x)\dot{B}(y)) = \delta_0(x-y)$. This operator is of the Sturm-Liouville/killed diffusion type and acts on the modified Sobolev space

$$H^{1}_{0,*} := \left\{ f \in \mathcal{C}^{1}(\mathbb{R}_{+}) / f(0) = 0, \int_{\mathbb{R}_{+}} \left(f'(x)^{2} + (1+x)f(x)^{2} \right) dx < +\infty \right\}$$

The classical theory of compact operator can be proven to apply a.s. to the random operator SAO_{β} in the space $H_{0,*}^1$ [49, 87]; its eigenvalues can be written as $1 > \lambda_1(\beta) > \lambda_2(\beta) > \cdots > 0$. Using the minimax representation of Courant-Fischer, one thus has $\lambda_1(\beta) = \sup_{f \in H_{0,*}^1, \|f\|_* = 1} \langle f, SAO_{\beta}f \rangle_*$ (with obvious notations).

Passing to the limit on tri-diagonal models of random matrices yields $\mathcal{TW}_{\beta} \stackrel{\mathcal{L}}{=} \lambda_1(\beta)$ (see [50, conj. 3.10-6.5] & [108, p. 921]) and in particular, using an integration by parts

$$\mathcal{TW}_{2} \stackrel{\mathcal{L}}{=} \sup_{f \in H^{1}_{0,*}, \|f\|_{*} = 1} \left\{ \sqrt{2} \int_{\mathbb{R}_{+}} f(x)^{2} dB(x) - \int_{\mathbb{R}_{+}} \left(f'(x)^{2} + xf(x)^{2} \right) dx \right\}$$
(5)

In view of these last results, a natural question can be posed:

Question 1.1. Can one find an algebraic operation that allows to get Tracy-Widom distributions from independent random variables ?

Question 1.1 was asked e.g. during a workshop at the American Mathematics Institute organised by T. Tao and V. Vu in the fall semester 2010 [2, prob. (15)]², in [44, pb. 7, after (16)] and in [8, Qu. 2.4].

We answer question 1.1 with the following description of \mathcal{TW}_2 (theorems 1.2 and 2.3 in the sequel) :

²The question asks for a sequence of i.i.d. random variables, but they can always be obtained out of independent random variables by inversion of the cumulative distribution function, see remark 2.8.

Theorem 1.2 (\mathcal{TW}_2 is a maximum of a sequence of independent random variables). There exist independent random variables $(Z_k(Ai))_{k \ge 0}$ such that

$$\mathcal{TW}_2 \stackrel{\mathcal{L}}{=} \max_{k \ge 0} Z_k(\mathrm{Ai}) \tag{6}$$

Two random variables $Z_k(\text{Ai})$ are defined in lemma 2.1/(13) and in theorem 2.3/(19). The description given in (19) uses the $L^2(\mathbb{R}_+)$ -normalised eigenvectors of $\mathbf{H}(\text{Ai}_s)$ or equivalently the eigenvectors of the Tracy-Widom commuting differential operator $\frac{d}{dx}x\frac{d}{dx}-x(x+s)$ acting on $L^2(\mathbb{R}_+)$ [123, IV-B p. 165] which are a particular rescaling of the prolate hyperspheroidal wave functions introduced independently by Slepian [118] and Hurtley [69, 70] (see also [60], [105, ch. 30.12 p. 704] and [40, 66]); we prove this last fact and give a quick summary of results on these functions in Annex B.

1.2. Max-independence structures.

1.2.1. Detropicalised motivation : independence for traces and total parts. Two of the most classical models in integrable probability are given by the GUE_N and the Schur measure, defined respectively in (42) and (71). The eigenvalues $(\lambda_{k,N})_{1 \leq k \leq N}$ of $M_N \sim GUE_N$ satisfies the identity in law (which is in fact a deterministic identity given by the invariance by conjugacy of the trace)

$$\sum_{k=1}^{N} \boldsymbol{\lambda}_{k,N} \stackrel{\mathcal{L}}{=} \sum_{k=1}^{N} \boldsymbol{M}_{N}[k,k]$$

The sum in the RHS is composed of independent (identically distributed) Gaussians. An identity of the same type is given for the parts of the random partition $\boldsymbol{\lambda}^{\operatorname{Schur}(\boldsymbol{p},\boldsymbol{q})} := (\boldsymbol{\lambda}_{k}^{\operatorname{Schur}(\boldsymbol{p},\boldsymbol{q})})_{k \ge 1} \sim \mathbb{P}_{\operatorname{Schur}(\boldsymbol{p},\boldsymbol{q})}$ where $\boldsymbol{p} := (p_k)_k, \boldsymbol{q} := (q_k)_{k \ge 1}$ and $p_i, q_j \in (0, 1)$:

$$\sum_{k \ge 1} \boldsymbol{\lambda}_k^{\operatorname{Schur}(\boldsymbol{p}, \boldsymbol{q})} \stackrel{\mathcal{L}}{=} \sum_{i, j \ge 1} \operatorname{Geom}(p_i q_j)$$

Here, $\operatorname{Geom}(q)$ designates a random variable geometrically distributed with the convention $\mathbb{P}(\operatorname{Geom}(q) = k) = (1 - q)q^k \mathbb{1}_{\{k \in \mathbb{N}\}}$; see (110) for a proof. In the same vein as for the *GUE*, the random variables in the RHS of the previous identity are independent.

One could call such identities in law between sums of dependent random variables and sums of independent random variables a *sum-independence structure*. In this paper, we will be concerned with a *tropical* version (see [80]) of these two identities :

$$\max_{k \ge 1} \boldsymbol{\lambda}_k \stackrel{\mathcal{L}}{=} \max_{k \ge 1} W_k$$

where the random variables $(W_k)_{k\geq 1}$ in the RHS are independent. We call such an identity a max-independence structure³.

³It also includes the case of a minimum, so we could also call it a *tropical independence structure*, but in the same vein convexity relates to concavity, we will stick to this terminology.

1.2.2. A previous instance in the literature. Max-independence structures for eigenvalues of random matrix ensembles made a first apparition in the Ginibre ensemble [64]. This ensemble, noted GinE_N is the probability space $(\mathcal{M}_N(\mathbb{C}), \mathbb{P}_N)$ where $\mathcal{M}_N(\mathbb{C})$ is the space of $N \times N$ complex matrices and \mathbb{P}_N is the Gaussian measure $\mathbb{P}_N(dM) = \pi^{-N^2} e^{-\operatorname{tr}(M^*M)} dM$. If $(\lambda_{k,N})_{1 \leq k \leq N}$ denotes the eigenvalues of $G_N \sim \operatorname{GinE}_N$, ordered by modulus size, i.e. $|\lambda_{1,N}| \geq |\lambda_{2,N}| \geq \ldots$, one has the formula

$$|\boldsymbol{\lambda}_{1,N}| \stackrel{\mathcal{L}}{=} \max_{1 \leqslant k \leqslant N} \sqrt{\boldsymbol{\gamma}_k}, \qquad \boldsymbol{\gamma}_k \sim \boldsymbol{\Gamma}(k)$$
 (7)

with independent random variables $\gamma_k \stackrel{\mathcal{L}}{=} \sum_{j=1}^k e^{(j)}$, the $e^{(j)}$ being i.i.d. exponential random variables. Using (7), Rider showed that [109, thm. 1]

$$\sqrt{4N\alpha_N} \left(\frac{|\boldsymbol{\lambda}_{1,N}|}{\sqrt{N}} - 1 - \sqrt{\frac{\alpha_N}{4N}} \right) \xrightarrow[N \to +\infty]{\mathcal{L}} \operatorname{Gb}(1), \qquad \alpha_N := \log(N/2\pi) - 2\log\log(N)$$

where Gb(1) designates the Gumbel distribution, i.e. the law of $-\log e$.

See also [11, ch. 4.3.7, ex. 4.7.2, ch. 6.4] for more details and the blog [41] for a pedagogical introduction.

The presence of such an independence structure in GinE_N asks legitimately the question of its generality in any matrix ensemble. But more generally, since the universality class of \mathcal{TW}_2 previously described is so rich, a more natural question can be asked:

Question 1.3. Is the max-independence structure already present before renormalisation in all the models previously cited ?

A positive answer to question 1.3 is provided in the following cases that will be described in more details throughout the article :

- the continuum random polymer $(\S 2.3)$,
- random matrix ensembles (§ 3),
- the symmetric Schur measure (§ 4).

In the case of the symmetric Schur measure, one has a very remarkable structure : an i.i.d. randomisation of the (deterministic) sequence of integers (see § 4.5). Each of these cases has an equality in law analogous to (6). For instance, in the case of the GUE, one has

Theorem 1.4 (Theorem 3.1 in the sequel). There exist independent random variables $(W_k)_{0 \leq k \leq N-1}$ such that, if $\lambda_{1,N}$ designates the largest eigenvalue of a random matrix $H_N \sim GUE_N$,

$$\left| \boldsymbol{\lambda}_{1,N} \stackrel{\mathcal{L}}{=} \max_{0 \leqslant k \leqslant N-1} W_k \right| \tag{8}$$

While $W_0 \sim \mathcal{N}(0, 1)$, the law of W_k for $k \ge 1$ is explicitly given in theorem 3.1 but requires to use the σ -form of the Painlevé IV equation studied by Okamoto [94, 95, 96, 97],

simplified by Noumi-Yamada [91, 90] and applied in Random Matrix Theory by Forrester-Witte [56] (see also [57, 58] and [54, ch. 8 p. 328] for other cases that we will not treat in this paper).

1.3. Rescaling of the largest eigenvalue of the *GUE*. The first historical appearance of the Tracy-Widom distribution in [122, 123] was obtained by the rescaling of the largest eigenvalue of the GUE_N when $N \to +\infty$. From this perspective, the max-independence structure given by (8)/(46) is particularly interesting as one can use the classical Poisson approximation for a maximum of independent random variables: writing

$$\left\{\max_{0 \le k \le N-1} W_k \le x\right\} = \left\{\sum_{k=0}^{N-1} \mathbb{1}_{\{W_k > x\}} = 0\right\}$$

and analysing the parametric sum of independent indicators $S_N(x) := \sum_{k=0}^{N-1} \mathbb{1}_{\{W_k > x\}}$, one can perform a Poisson approximation $S_N(x) \approx \operatorname{Po}(\mu_N(x))$ where $\mu_N(x) := \mathbb{E}(S_N(x)) = \sum_{k=0}^{N-1} \mathbb{P}(W_k > x)$. The error in such an approximation is moreover explicit, given for instance by the Chen-Stein method [42]. As a result, one can try to put in a more classical probabilistic perspective the result of Tracy and Widom, for instance in the vein of Rider's analysis of the edge fluctuations in the complex Ginibre ensemble [109]. Such an analysis is performed in § 3.4 and its main result reads:

Theorem 1.5 (Theorem 3.11 in the sequel). One has

$$\mathbb{P}\left(\frac{\boldsymbol{\lambda}_{1,N} - 2\sqrt{N}}{N^{-1/6}} \leqslant \sqrt{2}t\right) \xrightarrow[N \to +\infty]{} \mathbb{P}\left(\mathcal{T}\mathcal{W}_2 \leqslant \sqrt{2}t\right) = \exp\left(-\int_{\mathbb{R}} (x-t)_+ \mathcal{Q}(t)^2 dt\right) \quad (9)$$

with $x_{+} := x \mathbb{1}_{\{x \ge 0\}}$ and

$$Q(x) := \exp\left(-\int_{x}^{+\infty} \delta\sigma\right), \qquad \delta\sigma := \sigma(\cdot; 1) - \sigma(\cdot; 0) \tag{10}$$

where $\sigma(\cdot; a)$ is the solution of the σ -form of the Painlevé II equation given in (59).

We would like to emphasize that the very structure of the limit is already present for fixed N with a function $\delta \sigma^{(N)}$ in $\mu_N(x)$ (which is in fact itself already of this form with a Stieltjes integral), and that it is preserved in the form (9) after passing to the limit.

Comparing (3) and (9), one sees that q = Q; we give a direct proof of this fact in lemma 3.12 using the Forrester-Witte theory.

1.4. Organisation of the paper. The plan of the article is as follows :

- we prove theorem (1.2) in section 2 and extend it to the case of \mathcal{TW}_1 in § 2.4 and to the continuum random polymer in § 2.3.
- We investigate independence structure in random matrix ensembles in section 3 and prove there theorems 1.4/3.1 and 1.5/3.11. We would like to draw attention

here to the problem 3.7 which is, in the author's opinion, the most important in the article, and probably the tip of a very big iceberg.

- We analyse the case of the symmetric Schur measure in section 4, starting with the Plancherel measure.
- We give in Annex A some relevant notations on symmetric functions and the Schur measure and in Annex B a summary of results on hyperspheroidal wave functions.

2. \mathcal{TW}_2 is a maximum of independent random variables

2.1. First approach.

2.1.1. Prerequisites. For an operator $\mathbf{K}_t : L^2(\mathbb{R}_+) \to L^2(\mathbb{R}_+)$ of kernel K_t , define $\dot{K}_t := \frac{d}{dt}K_t$ and $\dot{\mathbf{K}}_t : L^2(\mathbb{R}_+) \to L^2(\mathbb{R}_+)$ its associated operator. For $\phi, \psi \in L^2(\mathbb{R}_+)$, we define $\phi_t := \phi(\cdot + t), \ \psi_t := \psi(\cdot + t)$ and

$$K_t(x,y) := \int_{\mathbb{R}_+} \phi_t(x+u)\psi_t(y+u)du = \int_t^{+\infty} \phi(x+u)\psi(y+u)du$$

Define $H(\phi)$ to be the Hankel operator of symbol ϕ acting on $L^2(\mathbb{R}_+)$, i.e. the operator of kernel $(x, y) \mapsto \phi(x + y)$. Then, one has

$$oldsymbol{K}_t = oldsymbol{H}(\psi_t)oldsymbol{H}(\phi_t)$$

Set

$$\psi^*(f) := \langle f, \psi \rangle_{L^2(\mathbb{R}_+)}$$

Then,

$$\dot{K}_t(x,y) = -\phi(x+t)\psi(y+t) = -\phi_t \otimes \psi_t(x,y) \qquad \Longleftrightarrow \qquad \dot{K}_t = -\phi_t \otimes \psi_t^* \tag{11}$$

Suppose now that K_t has all its eigenvalues in (0, 1). Writing the following determinants and traces on $L^2(\mathbb{R}_+)$, we have

$$\frac{\det(I - \mathbf{K}_t)}{\det(I - \mathbf{K}_a)} = \exp\left(\operatorname{tr}(\log(I - \mathbf{K}_t) - \log(I - \mathbf{K}_a))\right)$$
$$= \exp\left(-\int_a^t \operatorname{tr}((I - \mathbf{K}_s)^{-1} \dot{\mathbf{K}}_s) ds\right)$$
$$= \exp\left(\int_a^t \operatorname{tr}((I - \mathbf{K}_s)^{-1} \phi_s \otimes \psi_s^*) ds\right)$$
$$= \exp\left(\int_a^t \left\langle (I - \mathbf{K}_s)^{-1} \phi_s, \psi_s \right\rangle_{L^2(\mathbb{R}_+)} ds\right)$$

For $a = +\infty$, as $\mathbf{K}_a = 0$, we get

$$\det(I - \mathbf{K}_t)_{L^2(\mathbb{R}_+)} = \exp\left(-\int_t^{+\infty} \left\langle (I - \mathbf{K}_s)^{-1} \phi_s, \psi_s \right\rangle_{L^2(\mathbb{R}_+)} ds\right)$$

$$= \exp\left(-\int_{t}^{+\infty} \sum_{\ell \ge 0} \left\langle \mathbf{K}_{s}^{\ell} \phi_{s}, \psi_{s} \right\rangle_{L^{2}(\mathbb{R}_{+})} ds\right)$$

And in the case where $\phi = \psi$, one finally gets

$$\det \left(I - \boldsymbol{H}(\phi_t)^2 \right)_{L^2(\mathbb{R}_+)} = \prod_{\ell \ge 0} \exp \left(-\int_t^{+\infty} \left\| \boldsymbol{H}(\phi_s)^\ell \phi_s \right\|_{L^2(\mathbb{R}_+)}^2 ds \right)$$
(12)

Suppose that $s \mapsto \|\boldsymbol{H}(\phi_s)^{\ell}\phi_s\|_{L^2(\mathbb{R}_+)}^2 \notin L^1(\mathbb{R})$. Then, one clearly gets the existence of a random variable $Z_{\ell}(\phi)$ such that

$$\mathbb{P}(Z_{\ell}(\phi) \leqslant t) = \exp\left(-\int_{t}^{+\infty} \left\|\boldsymbol{H}(\phi_{s})^{\ell}\phi_{s}\right\|_{L^{2}(\mathbb{R}_{+})}^{2} ds\right)$$
(13)

which implies that

$$\det(I - \boldsymbol{H}(\phi_t)^2)_{L^2(\mathbb{R}_+)} = \prod_{\ell \ge 0} \mathbb{P}(Z_\ell(\phi) \le t) = \mathbb{P}\left(\max_{\ell \ge 0} Z_\ell(\phi) \le t\right)$$
(14)

2.1.2. Proof of theorem 1.2. We now consider the case $\phi = \text{Ai in (13)}$. It is clear that (2) translates at the operator level into

$$\boldsymbol{K}_{\mathrm{Ai}_s} = \boldsymbol{H}(\mathrm{Ai}_s)^2 \tag{15}$$

To prove theorem 1.2, one uses (14) and the

Lemma 2.1 (Existence of a random variable $Z_{\ell}(Ai)$). One has $s \mapsto \left\| \boldsymbol{H}(Ai_s)^{\ell} Ai_s \right\|_{L^2(\mathbb{R}_+)}^2 \notin L^1(\mathbb{R})$

Proof. For $\ell = 0$, one has for s < 0

$$\|\operatorname{Ai}_{s}\|_{L^{2}(\mathbb{R}_{+})}^{2} = \int_{\mathbb{R}_{+}} \operatorname{Ai}(s+x)^{2} dx = \int_{s}^{+\infty} \operatorname{Ai}(x)^{2} dx \ge \int_{s}^{0} \operatorname{Ai}(x)^{2} dx = |s| \int_{-1}^{0} \operatorname{Ai}(x|s|)^{2} dx = \Omega(\sqrt{|s|})$$

which shows that the function is not integrable in a neighbourhood of $-\infty$.

To show that

$$C := \int_{\mathbb{R}_+} \operatorname{Ai}(x)^2 dx < \infty$$

we have used the estimate [129, (2.44) p. 14]/[79]

Ai
$$(x) = O(x^{-1/4}e^{-\frac{2}{3}x^{3/2}}), \qquad x \to +\infty$$

and to show that

$$\int_{-1}^{0} \operatorname{Ai}(x|s|)^2 dx = \Omega\left(\frac{1}{\sqrt{|s|}}\right), \qquad s \to +\infty$$

we have used [105, (9.8.1) p. 199, (9.8.20), (9.8.21) p. 200]

$$\operatorname{Ai}(-x) := M(x)\sin\theta(x), \qquad M(x)^2 := \operatorname{Ai}(-x)^2 + \operatorname{Bi}(-x)^2, \qquad \theta(x) := \arctan\left(\frac{\operatorname{Ai}(-x)}{\operatorname{Bi}(-x)}\right)$$
$$M(x)^2 \sim \frac{1}{\pi\sqrt{x}} \sum_{k \ge 0} \frac{x^{-3k}}{k!} (-1)^k \frac{1 \cdot 3 \cdot 5 \cdots (6k-1)}{96^k}, \quad \theta(x) \sim \frac{\pi}{4} + \frac{2}{3} x^{\frac{3}{2}} \left(1 - \frac{5}{32} x^{-3} + O(x^{-6})\right)$$

which gives in particular

$$\operatorname{Ai}(-x)^2 \sim \frac{1}{\pi x^{1/2}} \sin\left(\frac{\pi}{4} + \frac{2}{3}x^{\frac{3}{2}}\right)^2, \qquad x \to +\infty$$
 (16)

For $\ell = 1$, one has

$$\|\boldsymbol{H}(\mathrm{Ai}_s)\mathrm{Ai}_s\|_{L^2(\mathbb{R}_+)}^2 = \int_{\mathbb{R}_+} \left(\int_{\mathbb{R}_+} \mathrm{Ai}_s(x+y)\mathrm{Ai}_s(y)dy\right)^2 dx =: \int_{\mathbb{R}_+} K_{\mathrm{Ai}}(s+x,s)^2 dx$$

One also has

$$K_{\rm Ai}(x,y) = \frac{{\rm Ai}(x){\rm Ai}'(y) - {\rm Ai}(y){\rm Ai}'(x)}{x-y} \mathbb{1}_{\{x \neq y\}} + ({\rm Ai}'(x)^2 - x{\rm Ai}(x)^2)\mathbb{1}_{\{x=y\}}$$
(17)

and for $x \neq y$,

$$K_{\mathrm{Ai}}(x,y) = -\mathrm{Ai}(x)\mathrm{Ai}(y)\frac{\frac{\mathrm{Ai}'(y)}{\mathrm{Ai}(y)} - \frac{\mathrm{Ai}'(x)}{\mathrm{Ai}(x)}}{y-x}$$
$$= -\mathrm{Ai}(x)\mathrm{Ai}(y)\int_{0}^{1} \left(\frac{\mathrm{Ai}'}{\mathrm{Ai}}\right)'(tx+\overline{t}y)dt, \qquad \overline{t} := 1-t$$

Moreover, using $\operatorname{Ai}''(x) = x\operatorname{Ai}(x)$, one gets

$$\left(\frac{\operatorname{Ai}'}{\operatorname{Ai}}\right)'_{x} = \frac{\operatorname{Ai}''(x)\operatorname{Ai}(x) - \operatorname{Ai}'(x)^2}{\operatorname{Ai}(x)^2} = x - \left(\frac{\operatorname{Ai}'(x)}{\operatorname{Ai}(x)}\right)^2$$

hence

$$K_{\mathrm{Ai}}(x,y) = \mathrm{Ai}(x)\mathrm{Ai}(y)\left(-\frac{x+y}{2} + \int_0^1 \left(\frac{\mathrm{Ai}'}{\mathrm{Ai}}(tx+\overline{t}y)\right)^2 dt\right)$$
$$=: \mathrm{Ai}(x)\mathrm{Ai}(y)R_{\mathrm{Ai}}(x,y)$$

with

$$R_f(x,y) := -\frac{x+y}{2} + \int_0^1 \left(\frac{f'}{f}(tx+\overline{t}y)\right)^2 dt$$

This implies for s = -|s| < 0

$$\|\boldsymbol{H}(\mathrm{Ai}_s)\mathrm{Ai}_s\|_{L^2(\mathbb{R}_+)}^2 = \int_{\mathbb{R}_+} K_{\mathrm{Ai}}(s+x,s)^2 dx$$

$$= \int_{\mathbb{R}_{+}} \operatorname{Ai}(x+s)^{2} \operatorname{Ai}(s)^{2} R_{\operatorname{Ai}}(s+x,s)^{2} dx = \operatorname{Ai}(s)^{2} \int_{s}^{+\infty} \operatorname{Ai}(X)^{2} R_{\operatorname{Ai}}(X,s)^{2} dX$$

$$\geq \frac{\operatorname{Ai}(s)^{2}}{4} \int_{s}^{+\infty} \operatorname{Ai}(X)^{2} (X+s)^{2}_{-} dX = \frac{\operatorname{Ai}(s)^{2}}{4} \int_{s}^{-s} \operatorname{Ai}(X)^{2} (X+s)^{2} dX$$

$$= \frac{\operatorname{Ai}(s)^{2}}{4} |s|^{3} \int_{-1}^{1} \operatorname{Ai}(|s|u)^{2} (u-1)^{2} du$$

$$\geq \frac{\operatorname{Ai}(s)^{2}}{4} |s|^{3} \int_{-1}^{0} \operatorname{Ai}(|s|u)^{2} (1-u)^{2} du$$

$$= \Omega(s^{2}) \qquad \text{using (16).}$$

As a result, $s \mapsto \|\boldsymbol{H}(Ai_s)Ai_s\|_{L^2(\mathbb{R}_+)}^2$ is not integrable on \mathbb{R} .

In the general case, one has for $\ell=2m$ and s<0

$$\begin{split} \left\| \boldsymbol{H}(\operatorname{Ai}_{s})^{2m}\operatorname{Ai}_{s} \right\|_{L^{2}(\mathbb{R}_{+})}^{2} &= \int_{\mathbb{R}_{+}} \left(\int_{\mathbb{R}_{+}} K_{\operatorname{Ai}_{s}}^{*m}(x,y)\operatorname{Ai}_{s}(y)dy \right)^{2}dx \\ &= \int_{\mathbb{R}_{+}} \left(\int_{\mathbb{R}_{+}\times\mathbb{R}_{+}^{m-1}} K_{\operatorname{Ai}_{s}}(x,t_{1}) \prod_{j=1}^{m-2} K_{\operatorname{Ai}_{s}}(t_{j},t_{j+1}) K_{\operatorname{Ai}_{s}}(t_{m-1},y)dt \operatorname{Ai}_{s}(y)dy \right)^{2}dx \\ &= \int_{\mathbb{R}_{+}} \operatorname{Ai}_{s}(x)^{2} \left(\int_{\mathbb{R}_{+}\times\mathbb{R}_{+}^{m-1}} \prod_{j=1}^{m-2} \operatorname{Ai}_{s}(t_{j})^{2} \\ &\times R_{\operatorname{Ai}_{s}}(x,t_{1}) \prod_{j=1}^{m-2} R_{\operatorname{Ai}_{s}}(t_{j},t_{j+1}) R_{\operatorname{Ai}_{s}}(t_{m-1},y)dt \operatorname{Ai}_{s}(y)^{2}dy \right)^{2}dx \\ &\geqslant \frac{1}{2^{m}} \int_{\mathbb{R}_{+}} \operatorname{Ai}_{s}(x)^{2} \left(\int_{\mathbb{R}_{+}\times\mathbb{R}_{+}^{m-1}} \prod_{j=1}^{m-2} \operatorname{Ai}_{s}(t_{j})^{2} \\ &\times (2|s|-x-t_{1})_{+} \prod_{j=1}^{m-2} (2|s|-t_{j}-t_{j+1})_{+} (2|s|-t_{m-1}-y)_{+}dt \operatorname{Ai}_{s}(y)^{2}dy \right)^{2}dx \\ &= \frac{1}{2^{m}} \int_{[s,+\infty)} \operatorname{Ai}(X)^{2} \left(\int_{[s,+\infty)^{m}} \prod_{j=1}^{m-2} \operatorname{Ai}(T_{j})^{2} \\ &\times (X+T_{1})_{-} \prod_{j=1}^{m-2} (T_{j}+T_{j+1})_{-} (T_{m-1}+Y)_{-}dT \operatorname{Ai}(Y)^{2}dY \right)^{2}dX \\ &\geqslant \frac{|s|^{4m+1}}{2^{m}} \int_{[-1,0)} \operatorname{Ai}(|s|u)^{2} \left(\int_{[-1,0)^{m}} \prod_{j=1}^{m-2} \operatorname{Ai}(|s|\tau_{j})^{2} \right)^{2}dx \end{split}$$

$$\times (u + \tau_1)_{-} \prod_{j=1}^{m-2} (\tau_j + \tau_{j+1})_{-} (\tau_{m-1} + v)_{-} \operatorname{Ai}(|s|v)^2 dv \, d\tau \bigg)^2 du$$
$$= \Omega (|s|^{3m+1/2}) \quad \text{using (16).}$$

And for $\ell = 2m + 1$ and s < 0,

$$\begin{split} \left\|\boldsymbol{H}(\mathrm{Ai}_{s})^{2m+1}\mathrm{Ai}_{s}\right\|_{L^{2}(\mathbb{R}+)}^{2} &= \int_{\mathbb{R}+} \left(\int_{\mathbb{R}+} K_{\mathrm{Ai}_{s}}^{**} * H_{\mathrm{Ai}_{s}}(x,y)\mathrm{Ai}_{s}(y)dy\right)^{2}dx \\ &= \int_{\mathbb{R}+} \left(\int_{\mathbb{R}+\times\mathbb{R}_{+}^{m}} K_{\mathrm{Ai}_{s}}(x,t_{1})\prod_{j=1}^{m-1} K_{\mathrm{Ai}_{s}}(t_{j},t_{j+1}) H_{\mathrm{Ai}_{s}}(t_{m},y)dt \operatorname{Ai}_{s}(y)dy\right)^{2}dx \\ &= \int_{\mathbb{R}+} \left(\int_{\mathbb{R}_{+}^{m}} K_{\mathrm{Ai}_{s}}(x,t_{1})\prod_{j=1}^{m-1} K_{\mathrm{Ai}_{s}}(t_{j},t_{j+1}) K_{\mathrm{Ai}_{s}}(t_{m},0)dt\right)^{2}dx \\ &= \mathrm{Ai}_{s}(0)^{2} \int_{\mathbb{R}_{+}} \mathrm{Ai}_{s}(x)^{2} \left(\int_{\mathbb{R}_{+}^{m}} \prod_{j=1}^{m} \mathrm{Ai}_{s}(t_{j})^{2} \\ &\times R_{\mathrm{Ai}_{s}}(x,t_{1})\prod_{j=1}^{m-1} R_{\mathrm{Ai}_{s}}(t_{j},t_{j+1}) R_{\mathrm{Ai}_{s}}(t_{m},0)dt\right)^{2}dx \\ &\geq \frac{\mathrm{Ai}_{s}(0)^{2}}{2^{m}} \int_{\mathbb{R}_{+}} \mathrm{Ai}_{s}(x)^{2} \left(\int_{\mathbb{R}_{+}^{m}} \prod_{j=1}^{m} \mathrm{Ai}_{s}(t_{j})^{2} \\ &\times (2|s|-x-t_{1})_{+} \prod_{j=1}^{m} (2|s|-t_{j}-t_{j+1})_{+} (2|s|-t_{m})_{+}dt\right)^{2}dx \\ &= \frac{\mathrm{Ai}(s)^{2}}{2^{m}} \int_{[s,+\infty)} \mathrm{Ai}(X)^{2} \left(\int_{[s,+\infty)^{m}} \prod_{j=1}^{m} \mathrm{Ai}(T_{j})^{2} \\ &\times (X+T_{1})_{-} \prod_{j=1}^{m-1} (T_{j}+T_{j+1})_{-} (|s|-T_{m})_{+}dT\right)^{2}dX \\ &\geq \frac{|s|^{4m+3}}{2^{m}} \mathrm{Ai}(-|s|)^{2} \int_{[-1,0)} \mathrm{Ai}(|s|u|^{2} \left(\int_{[-1,0)^{m}} \prod_{j=1}^{m} \mathrm{Ai}(|s|t_{j})^{2} \\ &\times (u+\tau_{1})_{-} \prod_{j=1}^{m-1} (\tau_{j}+\tau_{j+1})_{-} (1-\tau_{m})_{+}d\tau\right)^{2}du \\ &= \Omega(|s|^{3m+2}) \qquad \text{using (16).} \\ \\ \end{array}$$

This concludes the proof.

11

2.2. Second approach.

2.2.1. *Prerequisites.* Before proving theorem 1.2, we introduce a slight variation of [61, lem. 1]:

Lemma 2.2 (Fixed interval Fuchs lemma). For $I \subset \mathbb{R}$, let $\mathbf{K}_t : L^2(I) \to L^2(I)$ be selfadjoint $(\mathbf{K}_t^* = \mathbf{K}_t)$. Let (g_t, λ_t) be such that $\mathbf{K}_t g_t = \lambda_t g_t$. Last, suppose that $t \mapsto (\mathbf{K}_t, g_t, \lambda_t)$ is differentiable. Then,

$$\dot{\lambda}_t = \frac{\left\langle \dot{\boldsymbol{K}}_t g_t, g_t \right\rangle_{L^2(I)}}{\left\| g_t \right\|_{L^2(I)}^2}$$

Proof. Differentiating $\mathbf{K}_t g_t = \lambda_t g_t$ yields

$$\dot{m{K}}_t g_t + m{K}_t \dot{g}_t = \dot{\lambda}_t g_t + \lambda_t \dot{g}_t$$

Taking the scalar product with g_t and dropping the index $L^2(I)$ then yields

$$\left\langle \dot{\boldsymbol{K}}_{t}g_{t},g_{t}\right\rangle +\left\langle \boldsymbol{K}_{t}\dot{g}_{t},g_{t}
ight
angle =\dot{\lambda}_{t}\left\Vert g_{t}\right\Vert ^{2}+\lambda_{t}\left\langle \dot{g}_{t},g_{t}
ight
angle$$

But

$$\begin{split} \delta_t &:= \langle \mathbf{K}_t \dot{g}_t, g_t \rangle - \lambda_t \langle \dot{g}_t, g_t \rangle \\ &= \langle \dot{g}_t, \mathbf{K}_t^* g_t \rangle - \lambda_t \langle \dot{g}_t, g_t \rangle \\ &= \langle \dot{g}_t, (\mathbf{K}_t - \lambda_t) g_t \rangle \quad \text{since } \mathbf{K}_t^* = \mathbf{K}_t \\ &= 0 \end{split}$$

hence the result.

2.2.2. Proof of theorem 1.2.

Theorem 2.3 (Max-independence structure in \mathcal{TW}_2 , bis repetita). One has

$$\mathcal{TW}_2 \stackrel{\mathcal{L}}{=} \max_{k \ge 0} Z'_k(\text{Ai}) \tag{18}$$

where $(Z'_k(Ai))_{k\geq 1}$ are independent random variables with density

$$f_{Z'_{k}(\mathrm{Ai})}(s) := \left\langle \mathrm{Ai}_{s}, \Psi_{k}^{(s)} \right\rangle_{L^{2}(\mathbb{R}_{+})}^{2}, \qquad \left\| \Psi_{k}^{(s)} \right\|_{L^{2}(\mathbb{R}_{+})}^{2} = 1$$
(19)

where $\Psi_k^{(s)}$ is the k-th eigenvector of $\boldsymbol{H}(Ai_s)$ on $L^2(\mathbb{R}_+)$ or equivalently the eigenvector of the Tracy-Widom commuting operator on $L^2(\mathbb{R}_+)$ given by

$$\mathcal{L}_{TW,s} := DXD - X(X+s) : f(x) \mapsto \frac{d}{dx} \left(x \frac{df}{dx} \right) - x(x+s)f(x)$$
(20)

Proof. Recall the Lidskii formula for trace-class operators [116, thm. 3.12.2]

$$\det(I - \mathbf{K})_{L^2(\mathbb{R}_+)} = \prod_{k \ge 1} (1 - \lambda_k), \qquad \mathbf{K}\psi_k = \lambda_k \psi_k$$
(21)

Applying this formula for $H(Ai_s)^2$ does not immediately give the result as one does not have any information on the dependency in s of the k-th eigenvalue $\lambda_k(s)$.

Nevertheless, for a normalised eigenvector g_t ($||g_t||^2_{L^2(\mathbb{R}_+)} = 1$) associated with an eigenvalue $\lambda(t)$, one gets

$$\dot{\lambda}(t) = \left\langle \dot{\boldsymbol{K}}_{\mathrm{Ai},t} g_t, g_t \right\rangle_{L^2(\mathbb{R}_+)} = -\left\langle \mathrm{Ai}_t \otimes \mathrm{Ai}_t^* g_t, g_t \right\rangle_{L^2(\mathbb{R}_+)} = -\left\langle \mathrm{Ai}_t, g_t \right\rangle_{L^2(\mathbb{R}_+)}^2$$

as, for $\mathbf{K}_{\mathrm{Ai},t} = \mathbf{H}(\mathrm{Ai}_s)^2 \curvearrowright L^2(\mathbb{R}_+)$, one has $\dot{\mathbf{K}}_{\mathrm{Ai},t} = -\mathrm{Ai}_t \otimes \mathrm{Ai}_t^*$ using (11). Moreover, $\mathbf{K}_{\mathrm{Ai},+\infty} = 0$ and $\mathbf{K}_{\mathrm{Ai},-\infty} = I$ hence $\lambda_k(\infty) = 0$ and $\lambda_k(-\infty) = 1$ and

$$\begin{split} \lambda(t) &= -\int_{t}^{+\infty} \dot{\lambda}(s) ds = +\int_{t}^{+\infty} \langle \operatorname{Ai}_{s}, g_{s} \rangle_{L^{2}(\mathbb{R}_{+})}^{2} ds, \\ &\int_{\mathbb{R}} \langle \operatorname{Ai}_{s}, g_{s} \rangle_{L^{2}(\mathbb{R}_{+})}^{2} ds = 1 \end{split}$$

The Lidskii formula then implies

$$\det \left(I - \boldsymbol{H}(\operatorname{Ai}_{t})^{2} \right)_{L^{2}(\mathbb{R}_{+})} = \prod_{k \ge 1} \left(1 - \lambda_{k}(t) \right)$$
$$= \prod_{k \ge 1} \left(1 - \int_{t}^{+\infty} \left\langle \operatorname{Ai}_{s}, \Psi_{s}^{(k)} \right\rangle_{L^{2}(\mathbb{R}_{+})}^{2} ds \right)$$
$$= \prod_{k \ge 1} \left(\int_{-\infty}^{t} \left\langle \operatorname{Ai}_{s}, \Psi_{s}^{(k)} \right\rangle_{L^{2}(\mathbb{R}_{+})}^{2} ds \right)$$
$$= \prod_{k \ge 1} \left(\int_{-\infty}^{t} f_{Z'_{k}(\operatorname{Ai})}(s) ds \right)$$
$$= \mathbb{P} \left(\max_{k \ge 1} Z'_{k}(\operatorname{Ai}) \le t \right)$$

which gives the result.

It remains to give a short description of the eigenvectors $(\Psi_k^{(s)})_{k\geq 1}$ which are the eigenvectors of $\mathcal{L}_{TW,s}$, i.e. a particular rescaling of the *prolate hyperspheroidal wave functions*. This fact is proven in Annex B.

Remark 2.4. An interesting question would be to prove (or disprove) that $Z_k(\text{Ai}) \stackrel{\mathcal{L}}{=} Z'_k(\text{Ai})$. A priori, several possible sequences can give the same result. 2.2.3. Another expression of $Z'_k(Ai)$.

Lemma 2.5 (Another expression of the law of $Z'_k(Ai)$). One also has

$$\mathbb{P}(Z'_{k}(\mathrm{Ai}) > s) := \exp\left(-\int_{-\infty}^{s} \Psi_{k}^{(t)}(0)^{2} dt\right), \qquad \left\|\Psi_{k}^{(s)}\right\|_{L^{2}(\mathbb{R}_{+})}^{2} = 1$$
(22)

Before proving lemma 2.5, we recall the original version of [61, lem. 1] :

Lemma 2.6 (Varying interval Fuchs lemma). Let $I_t := [t, +\infty) \subset \mathbb{R}$. Let $\mathbf{K} \equiv \mathbf{K}_t : L^2(I_t) \to L^2(I_t)$ be a self-adjoint $(\mathbf{K}_t^* = \mathbf{K}_t)$ such that the kernel K of \mathbf{K} does not depend on t. Let (g_t, λ_t) be such that $\mathbf{K}_t g_t = \lambda_t g_t$. Last, suppose that $t \mapsto (g_t, \lambda_t)$ is differentiable. Then,

$$\dot{\lambda}_t = -\lambda_t \frac{g_t(t)^2}{\|g_t\|_{L^2(I_t)}^2}$$

For the reader's convenience, we give a proof :

Proof. Differentiating $\mathbf{K}_t g_t = \lambda_t g_t$ yields

$$-K(\cdot,t)g_t(t) + \mathbf{K}_t \dot{g}_t = \dot{\lambda}_t g_t + \lambda_t \dot{g}_t$$

In particular

$$K(\cdot,t)g_t(t) + \dot{\lambda}_t g_t = \mathbf{K}_t \dot{g}_t - \lambda_t \dot{g}_t = (\mathbf{K}_t - \lambda_t I)\dot{g}_t \in \operatorname{Im}(\mathbf{K}_t - \lambda_t I)$$
$$\perp \ker(\mathbf{K}_t - \lambda_t I) = \mathbb{R}g_t \quad \text{as } \mathbf{K}_t^* = \mathbf{K}_t.$$

Taking the scalar product with g_t and dropping the index $L^2(I_t)$ then yields

$$0 = g_t(t) \langle K(\cdot, t), g_t \rangle + \lambda_t \langle g_t, g_t \rangle$$

= $g_t(t) \mathbf{K}_t g_t(t) + \dot{\lambda}_t ||g_t||^2$
= $\lambda_t g_t(t)^2 + \dot{\lambda}_t ||g_t||^2$

hence the result.

Remark 2.7.

• The original approach of Fuchs [61] amounts to the identity $\lambda_t \int_s^{+\infty} g_t g_s = \lambda_s \int_t^{+\infty} g_t g_s$ (with $\int_t^{+\infty} g_t^2 = 1$) which gives the result after division by t - s and the limit $t \to s$.

• The two versions of the Fuchs lemma 2.2 and 2.6 can be set on the same page if one differentiates the kernel $K_t(x, y) = \mathbb{1}_{\{x \ge t\}} K(x, y)$ in the sense of distributions (with a slight abuse of notation) :

$$\begin{split} \left\langle \dot{\boldsymbol{K}}_{t}g_{t},g_{t}\right\rangle &=-\left\langle \delta_{0}(t-\cdot)\boldsymbol{K}g_{t},g_{t}\right\rangle \\ &=-\lambda_{t}\left\langle \delta_{0}(t-\cdot)g_{t},g_{t}\right\rangle \\ &=-\lambda_{t}g_{t}(t)^{2} \end{split}$$

• If one supposes that K_t has a parametric kernel and acts on $L^2([t, +\infty))$, one easily finds the following modification of lemmas 2.2 and 2.6 :

$$\dot{\lambda}_t = -\lambda_t \frac{g_t(t)^2}{\|g_t\|_{L^2(I_t)}^2} + \frac{\left\langle \mathbf{K}_t g_t, g_t \right\rangle_{L^2(I_t)}}{\|g_t\|_{L^2(I_t)}^2}$$
(23)

• A direct corollary of lemma 2.6 is that for all $a \in \mathbb{R}$

$$\lambda_s = \lambda_a \exp\left(-\int_a^s \frac{g_t(t)^2}{\|g_t\|_{L^2(I_t)}^2} dt\right)$$

If in addition $\mathbf{K}_{-\infty} = I$ and $\mathbf{K}_{+\infty} = 0$, one can take $a = -\infty$ to get

$$\lambda_s = \exp\left(-\int_{-\infty}^s \frac{g_t(t)^2}{\|g_t\|_{L^2(I_t)}^2} dt\right), \qquad \int_{\mathbb{R}} \frac{g_t(t)^2}{\|g_t\|_{L^2(I_t)}^2} dt = +\infty$$
(24)

in which case λ_s defines a random variable Z such that $\lambda_s = \mathbb{P}(Z > s)$.

Proof of lemma 2.5. If the operator $\mathbf{K}_{Ai} \equiv \mathbf{K}_{Ai_0}$ defined in (2) acts on $L^2([s, +\infty))$, then, by an obvious change of variable :

$$\det \left(I - \boldsymbol{K}_{\mathrm{Ai}_s} \right)_{L^2(\mathbb{R}_+)} = \det \left(I - \boldsymbol{K}_{\mathrm{Ai}} \right)_{L^2([s, +\infty))}$$

The varying interval Fuchs lemma 2.6 and the Lidskii formula (21) can then be combined to give

$$\mathbb{P}(\mathcal{TW}_2 \leq s) = \det \left(I - \mathbf{K}_{\mathrm{Ai}}\right)_{L^2([s, +\infty))}$$
$$= \prod_{k \geq 1} \left(1 - \widetilde{\lambda}_k(s)\right)$$
$$= \prod_{k \geq 1} \left(1 - \exp\left(-\int_{-\infty}^s \widetilde{\Psi}_k^{(t)}(t)^2 dt\right)\right) \quad \text{with (24).}$$

Here, $\widetilde{\Psi}_k^{(t)}$ is the k-th eigenvector of $\mathbf{K}_{Ai} \curvearrowright L^2([t, +\infty))$ or equivalently of $\widetilde{\mathcal{L}}_{TW,t} := D(X-t)D - X(X-t)$. It is clear that

$$\int_{[t,+\infty)} K(x,y)f(y)dy = \lambda f(x) \quad \Longleftrightarrow \quad \int_{\mathbb{R}_+} K(X+t,Y+t)f(Y+t)dY = \lambda f(X+t)$$

As a result, $\widetilde{\Psi}_{k}^{(t)} = \Psi_{k}^{(t)}(\cdot - t)$ and $\widetilde{\Psi}_{k}^{(t)}(t) = \Psi_{k}^{(t)}(0)$. One sees moreover that $\widetilde{\lambda}_{k}(s) = \lambda_{k}(s)$, which proves that we still have an expression of the law of $Z_{k}^{\prime}(Ai)$. This concludes the proof.

Remark 2.8. Define

$$\Phi_k(t) := \int_{-\infty}^t \Psi_k^{(s)}(0)^2 ds$$

Then, writing Φ_k^{-1} for the inverse bijection of Φ_k , one has

$$Z'_k(\operatorname{Ai}) \stackrel{\mathcal{L}}{=} \Phi_k^{-1}(e), \qquad e \sim \operatorname{Exp}(1)$$

and one has expressed $\mathcal{TW}_2 \stackrel{\mathcal{L}}{=} \max_{k \ge 1} \Phi_k^{-1}(e_k)$ with an i.i.d. sequence $(e_k)_{k \ge 1}$, answering the original question of [2, prob. (15)].

2.3. Extension to operators $H(\phi_s)\mathfrak{M}_f^2H(\phi_s)$.

2.3.1. First approach. One classical generalisation of the previous result is the case where K_s is the kernel of an operator K_s acting on $L^2(\mathbb{R}_+)$ that reads

$$K_s(x,y) = \int_{\mathbb{R}} \phi_s(x+u)\phi_s(y+u)f(u)^2 du$$
(25)

with $\phi_s f \in L^2(\mathbb{R})$ for all s (say). We also suppose that K_s has all its eigenvalues in (0, 1) for all s.

Defining the multiplication operator

$$\mathfrak{M}_f: g \mapsto fg \tag{26}$$

the underlying operator reads

$$\boldsymbol{K}_{s} = \boldsymbol{H}(\phi_{s})\mathfrak{M}_{f}^{2}\boldsymbol{H}(\phi_{s}) = \boldsymbol{L}_{s}\boldsymbol{L}_{s}^{*}, \qquad \boldsymbol{L}_{s} := \boldsymbol{H}(\phi_{s})\mathfrak{M}_{f}$$
(27)

The relevant modification of (12) is

$$\det\left(I - \boldsymbol{H}(\phi_s)\mathfrak{M}_f^2\boldsymbol{H}(\phi_s)\right)_{L^2(\mathbb{R}_+)} = \prod_{\ell \ge 0} \exp\left(-\int_t^{+\infty} \left\langle (\boldsymbol{L}_s\boldsymbol{L}_s^*)^\ell \phi_s, \phi_s f^2 \right\rangle_{L^2(\mathbb{R}_+)} ds\right)$$
(28)

Usually, one does not have $L_s L_s^* = L_s^* L_s$ hence cannot write this last scalar product as $\|L_s^\ell \phi_s\|_{L^2(\mathbb{R}_+)}^2$, but the operator is clearly positive and one can define the square root $M_s := (L_s L_s^*)^{1/2}$ in which case one gets $\|M_s^\ell \phi_s\|_{L^2(\mathbb{R}_+)}^2$. Such a square root is nevertheless not easily expressible as an explicit operator such as L_s .

One classical such case is given by the value in x = 0 of the solution to the Stochastic Heat Equation (SHE) with Dirac initial condition [3, 13, 28, 38, 48, 111, 112, 113] also called continuum random polymer. This equation reads

$$\begin{cases} \partial_t u = \frac{1}{2} \Delta u + \dot{\xi} u \\ u(t=0,\cdot) = u_0 \equiv \delta_0 \end{cases}$$
(29)

If one defines $u(x,t) = e^{h(x,t)}$, then, h is the Hopf-Cole solution of the Kardar-Parisi-Zhang (KPZ) equation [13].

A key formula gives a determinantal structure for h(0, t) with $u_0 = \delta_0$ [3, 28, 38, 48, 111, 112, 113] :

$$\mathbb{P}\left(\frac{h(0,t) + \frac{t}{24}}{\gamma_t} + \frac{\operatorname{Gb}(1)}{\gamma_t} \leqslant s\right) = \det\left(I - \boldsymbol{K}_{KPZ(s,t)}\right)_{L^2(\mathbb{R}_+)}, \qquad \gamma_t := \left(\frac{t}{2}\right)^{1/3} \tag{30}$$

16

where Gb(1) is a Gumbel-distributed random variable independent of h(0,t) (for all t) and with⁴

$$K_{KPZ(s,t)}(x,y) := \int_{\mathbb{R}} \operatorname{Ai}_{s}(x+u) \operatorname{Ai}_{s}(y+u) \frac{du}{1+e^{-\gamma_{t}u}}$$
(31)

which is exactly of the type (25), the underlying operator being thus of the type (27).

It was moreover remarked by Johansson [77, prop. 1.4] that the RHS is

$$\mathbb{P}\left(\boldsymbol{\lambda}_{1}^{(t)}\leqslant s\right)$$

where $(\boldsymbol{\lambda}_{k}^{(t)})_{k\geq 1}$ is the point process given by the pointwise randomisation and re-ordering

$$(\boldsymbol{\lambda}_{k}^{(t)})_{k \ge 1} := (\boldsymbol{\lambda}_{k} + R_{k}/\gamma_{t})_{k \ge 1}^{>}, \qquad (\boldsymbol{\lambda}_{k})_{k \ge 1} \sim \mathrm{DPP}(K_{\mathrm{Ai}})$$

with $(R_k)_{k\geq 1}$ an i.i.d. sequence with law R_1 given by the Fermi factor

$$\mathbb{P}(R_1 \leqslant x) = F(x) := \frac{1}{1 + e^{-x}} \qquad \Longleftrightarrow \qquad R_1 \stackrel{\mathcal{L}}{=} \operatorname{Gb}(1) - \operatorname{Gb}'(1) \tag{32}$$

where Gb(1) and Gb'(1) are two independent Gumbel-distributed random variables. As a result,

$$\boldsymbol{\lambda}_{1}^{(t)} \stackrel{\mathcal{L}}{=} \max_{k \ge 1} \left\{ \boldsymbol{\lambda}_{k} + R_{k} / \gamma_{t} \right\}$$
(33)

This last equality in law is equivalent, after integration on the R_k s, to the tensorial identity of Fredholm determinants obtained in [3, 28, 111, 112, 113] building on [13, 125, 126, 127].

From this, one gets the equality in law

$$\frac{h(0,t) + \frac{t}{24}}{\gamma_t} + \frac{\operatorname{Gb}(1)}{\gamma_t} \stackrel{\mathcal{L}}{=} \boldsymbol{\lambda}_1^{(t)} \stackrel{\mathcal{L}}{=} \max_{k \ge 1} \left\{ \boldsymbol{\lambda}_k + R_k / \gamma_t \right\}$$

and using a slight modification of the proof of theorem 1.2, one gets the existence of independent random variables $Z_k^{(t)}(\text{Ai})$ such that

$$\boldsymbol{\lambda}_{1}^{(t)} \stackrel{\mathcal{L}}{=} \max_{k \ge 0} \left\{ Z_{k}^{(t)}(\mathrm{Ai}) \right\}$$
(34)

with

$$\mathbb{P}\Big(Z_k^{(t)}(\mathrm{Ai}) \leqslant s\Big) = \exp\left(-\int_s^{+\infty} \langle (\boldsymbol{H}(\mathrm{Ai}_u)\mathfrak{M}_{f_t}\boldsymbol{H}(\mathrm{Ai}_u))^k \mathrm{Ai}_u, \mathrm{Ai}_u f_t \rangle_{L^2(\mathbb{R}_+)} du \right), \quad f_t(x) = F(\gamma_t x)$$

The modification of the proof of lemma 2.1 goes as follows : write

$$K_{KPZ(s,t)}(x,y) = \int_{\mathbb{R}} \operatorname{Ai}(x+u) \operatorname{Ai}(y+u) \mathbb{P}(R \leqslant \gamma_t(u-s)) du$$
$$= \mathbb{E}\left(\int_{\mathbb{R}} \operatorname{Ai}(x+u) \operatorname{Ai}(y+u) \mathbb{1}_{\{R/\gamma_t+s\leqslant u\}} du\right)$$

⁴In [3, 111, 112, 113], the factor $\frac{du}{1+e^{-\gamma_t(u-s)}}$ is replaced by $\frac{du}{1-e^{-\gamma_t(u-s)}}$ with a principal value for the integral. The form we give is from [28, 38, 48] and [23, ch. 5.4.3].

$$\begin{split} &= \mathbb{E} \Biggl(\int_{s+\frac{R}{\gamma_t}}^{+\infty} \operatorname{Ai}(x+u) \operatorname{Ai}(y+u) du \Biggr) \\ &= \mathbb{E} \Bigl(K_{\operatorname{Ai},s+\frac{R}{\gamma_t}}(x,y) \Bigr) \end{split}$$

As a result,

$$\boldsymbol{K}_{KPZ(s,t)} = \mathbb{E}\left(\boldsymbol{K}_{\mathrm{Ai}_{s+R/\gamma_t}}\right) = \mathbb{E}\left(\boldsymbol{H}(\mathrm{Ai}_{s+R/\gamma_t})^2\right)$$
(35)

and, setting $R_t := \gamma_t^{-1} R_1$ and $\mathbf{K}_s \equiv \mathbf{K}_{KPZ(s,t)}$,

$$\det(I - \mathbf{K}_s) = \prod_{\ell \ge 0} \exp\left(-\mathbb{E}\left(\int_s^{+\infty} \left\langle \mathbf{H}\left(\operatorname{Ai}_{u+R_t^{(1)}}\right)^2 \cdots \mathbf{H}\left(\operatorname{Ai}_{u+R_t^{(\ell)}}\right)^2 \operatorname{Ai}_{u+R_t}, \operatorname{Ai}_{u+R_t}\right\rangle_{L^2(\mathbb{R}_+)} du\right)\right)$$

One can thus mimic the computations of lemma 2.1 a.s. for the random times $s + R_t^{(i)}$ and $s \ll 0$, which amounts to do it for $s \ll 0$, giving the same randomised estimates of $\Omega(|s|^m)$ with s replaced by $s + \min_{\ell} R_t^{(\ell)}$, and finally taking the expectation at the end, which is allowed as $R_t \stackrel{\mathcal{L}}{=} R/\gamma_t$ and R has moments at every order. Details are left to the reader.

2.3.2. Second approach. Using the fact that

$$\det (I - \boldsymbol{A}\boldsymbol{B})_{L^2(I)} = \det (I - \boldsymbol{B}\boldsymbol{A})_{L^2(I)}$$

for all trace-class operators $\boldsymbol{A}, \boldsymbol{B} \curvearrowright L^2(I)$, one gets

$$\mathbb{P}\left(\boldsymbol{\lambda}_{1}^{KPZ(t)} \leqslant s\right) = \det\left(I - \boldsymbol{K}_{KPZ(s,t)}\right)_{L^{2}(\mathbb{R}_{+})}$$
$$= \det\left(I - \boldsymbol{H}(\operatorname{Ai}_{s})\mathfrak{M}_{f_{t}^{2}}\boldsymbol{H}(\operatorname{Ai}_{s})\right)_{L^{2}(\mathbb{R}_{+})}, \qquad f_{t}(x) := \frac{1}{\sqrt{1 + e^{-\gamma_{t}x}}}, \ \gamma_{t} := \left(\frac{t}{2}\right)^{\frac{1}{3}}$$
$$= \det\left(I - \mathfrak{M}_{f_{t}}\boldsymbol{H}(\operatorname{Ai}_{s})^{2}\mathfrak{M}_{f_{t}}\right)_{L^{2}(\mathbb{R}_{+})}$$

As

$$\frac{d}{ds}\mathfrak{M}_{f_t}\boldsymbol{H}(\mathrm{Ai}_s)^2\mathfrak{M}_{f_t} = \mathfrak{M}_{f_t}\frac{d}{ds}\boldsymbol{H}(\mathrm{Ai}_s)^2\mathfrak{M}_{f_t} = \mathfrak{M}_{f_t}(\mathrm{Ai}_s \otimes \mathrm{Ai}_s^*)\mathfrak{M}_{f_t}$$
$$= (f_t\mathrm{Ai}_s) \otimes (f_t\mathrm{Ai}_s)^*$$

the application of the Fuchs lemma 2.2 to $K_{KPZ(s,t)}$ gives

$$\mathbb{P}\left(\boldsymbol{\lambda}_{1}^{KPZ(t)} \leqslant s\right) = \prod_{k \ge 1} \int_{-\infty}^{s} \left\langle f_{t} \operatorname{Ai}_{u}, \Psi_{k}^{(u)} \right\rangle_{L^{2}(\mathbb{R}_{+})}^{2} du$$

We have thus proven the :

Theorem 2.9 (Max-independence structure in the KPZ equation). Let $(Z_k^{KPZ(t)})_{k\geq 1}$ be an independent sequence of random variables with law given by the density

$$f_{Z_k^{KPZ(t)}}(s) := \left\langle f_t \operatorname{Ai}_s, \Psi_k^{(s)} \right\rangle_{L^2(\mathbb{R}_+)}^2$$
(36)

Then,

$$\boldsymbol{\lambda}_{1}^{KPZ(t)} \stackrel{\mathcal{L}}{=} \max_{k \ge 1} Z_{k}^{KPZ(t)}$$
(37)

Remark 2.10. Translating the KPZ kernel gives

$$\det \left(I - \boldsymbol{K}_{KPZ(s,t)} \right)_{L^2(\mathbb{R}_+)} = \det \left(I - \widetilde{\boldsymbol{K}}_{KPZ(s,t)} \right)_{L^2([s,+\infty))}$$

where

$$\widetilde{K}_{KPZ(s,t)}(x,y) = \int_{\mathbb{R}_+} \operatorname{Ai}(x+u) \operatorname{Ai}(y+u) f_t(u-s)^2 du$$

which would be the kernel of $\boldsymbol{H}(Ai)\mathfrak{M}_{f_t(.-s)^2}\boldsymbol{H}(Ai)$ if it were acting on $L^2(\mathbb{R}_+)$ (but since we are acting on $L^2([s, +\infty))$), this is not a correct interpretation).

The general Fuchs lemma with a parametric kernel K_s on a varying interval $L^2([s, +\infty))$ given in (23) gives then for any $\Psi^{(s)} \in \left\{\Psi_k^{(s)}\right\}_{k>1}$

$$\begin{aligned} \dot{\lambda}(s) &= -\lambda(s)\Psi^{(s)}(0)^2 + \int_{[s,+\infty)^3} \operatorname{Ai}(x+u)\operatorname{Ai}(y+u)2(f_t f'_t)(u-s)du\widetilde{\Psi}^{(s)}(x)\widetilde{\Psi}^{(s)}(y)dxdy \\ &= -\lambda(s)\Psi^{(s)}(0)^2 + \int_{[s,+\infty)} \left\langle \operatorname{Ai}_{s+u},\Psi^{(s)} \right\rangle_{L^2(\mathbb{R}_+)}^2 2(f_t f'_t)(u-s)du \end{aligned}$$

This first order affine equation $\dot{\lambda} + a\lambda = b$ can be easily integrated to give another expression of λ , of the form $\lambda_s = \int_s^{+\infty} (e^{\int_s^v a}) b(v) dv$, but it seems less interesting than (36).

Remark 2.11. Tsai [128, (1.3)] computes the large deviation functional for $h(0, t) - \frac{t}{12}$ (with initial condition δ_0). With the max-independence structure, this becomes a problem of large deviations of a max of independent random variables, and the Poisson approximation approach described in § 1.3 applies. Nevertheless, the random variables are not so explicit and a study of the prolate hyperspheroidal functions is necessary beforehand.

2.4. Extension to \mathcal{TW}_1 . Recall that the power $\alpha > 0$ of a cumulative distribution function $F_X : x \mapsto \mathbb{P}(X \leq x)$ defines a random variable $X^{(\alpha)}$. If $\alpha \in \mathbb{N}^*$, one has $X^{(\alpha)} \stackrel{\mathcal{L}}{=} \max_{1 \leq k \leq \alpha} X_k$ where $(X_k)_k$ is a sequence of i.i.d. copies of X. In the general case, it seems that such a construction is not available.

The case of the GOE Tracy-Widom distribution can then be treated analogously to the GUE one :

Theorem 2.12 (Max-independence structure in \mathcal{TW}_1). One has with independent random variables in the RHS

$$\mathcal{TW}_{1} \stackrel{\mathcal{L}}{=} \max\left\{Q^{(1/2)}, \max_{k \ge 0} Z_{k}(\mathrm{Ai})^{(1/2)}\right\}$$
(38)

where $X^{(1/2)}$ is the random variable built out of X via $\mathbb{P}(X \leq x)^{1/2} = \mathbb{P}(X^{(1/2)} \leq x)$, $(Z_k(\operatorname{Ai}))_k$ are defined in theorem 1.2 and $\mathbb{P}(Q \leq s) = e^{-\int_s^{+\infty} q(x)dx}$ with q the Hastings-McLeod solution of the Painlevé II equation defined in (4).

Proof. Using the result of [124, (53)], one has

$$\mathbb{P}(\mathcal{TW}_1 \leqslant s) = \sqrt{\mathbb{P}(\mathcal{TW}_2 \leqslant s)} e^{-\int_s^{+\infty} q(x)dx}$$
(39)

Using $\mathbb{P}(X \leq x)^{1/2} := \mathbb{P}(X^{(1/2)} \leq x)$, one thus has

$$\mathbb{P}(\mathcal{TW}_2 \leqslant s)^{1/2} = \prod_{k \ge 1} \mathbb{P}(Z_k(\operatorname{Ai}) \leqslant s)^{1/2} =: \prod_{k \ge 1} \mathbb{P}(Z_k^{(1/2)}(\operatorname{Ai}) \leqslant s) = \mathbb{P}\left(\max_{k \ge 1} Z_k(\operatorname{Ai})^{(1/2)} \leqslant s\right)$$

and it remains to show that there exists Q such that $\mathbb{P}(Q \leq s) = e^{-\int_s^{+\infty} q(x)dx}$. It is clear that $F_q(s) := e^{-\int_s^{+\infty} q(x)dx} \to 1$ when $s \to +\infty$. Proving that $F_q(s) \to 0$ when $s \to -\infty$ is equivalent to $\int_{\mathbb{R}} q = +\infty$, and showing that F_q is increasing is equivalent to $q \geq 0$ on \mathbb{R} by differentiation. Such properties are granted for q, see e.g. [68] (the fact that $\int_{\mathbb{R}} q = +\infty$ is a consequence of the asymptotics $q(t) \sim |t/2|^{1/2}$ when $t \to -\infty$).

Remark 2.13. One might be tempted to use the Ferrari-Spohn characterisation [52, Prop. 1/(16)] that reads

$$\mathbb{P}(\mathcal{TW}_1 \leqslant s) = \det(I - \boldsymbol{H}(\operatorname{Ai}_s))_{L^2(\mathbb{R}_+)}$$
(40)

but $H(Ai_s)$ is not a positive operator, and a structure similar to the one used for the *GUE* is not obvious to obtain.

Nevertheless, (40) allows to give another decomposition of the random variable Q defined in the proof of theorem 2.12 using the Fuchs lemma 2.2 :

$$\mathbb{P}(Q \leq s) = e^{-\int_{s}^{+\infty} q(x)dx} = \frac{\mathbb{P}(\mathcal{T}\mathcal{W}_{1} \leq s)^{2}}{\mathbb{P}(\mathcal{T}\mathcal{W}_{2} \leq s)} = \frac{\det(I - \boldsymbol{H}(\mathrm{Ai}_{s}))_{L^{2}(\mathbb{R}_{+})}}{\det(I - \boldsymbol{H}(\mathrm{Ai}_{s})^{2})_{L^{2}(\mathbb{R}_{+})}}$$
$$= \frac{\det(I - \boldsymbol{H}(\mathrm{Ai}_{s}))_{L^{2}(\mathbb{R}_{+})}}{\det(I + \boldsymbol{H}(\mathrm{Ai}_{s}))_{L^{2}(\mathbb{R}_{+})}}$$
$$= \det\left(I - 2\boldsymbol{H}(\mathrm{Ai}_{s})(I + \boldsymbol{H}(\mathrm{Ai}_{s}))^{-1}\right)_{L^{2}(\mathbb{R}_{+})}$$

Define

$$\boldsymbol{L}_s := 2\boldsymbol{H}(\operatorname{Ai}_s)(I + \boldsymbol{H}(\operatorname{Ai}_s))^{-1} \curvearrowright L^2(\mathbb{R}_+)$$

If $\mu_k(s)$ denotes the k-th eigenvalue of \boldsymbol{L}_s for the normalised eigenvector $\psi_{k,s}$, then

$$\mu_{k}(s) = 2 \int_{s}^{+\infty} \left\langle \frac{d}{ds} \left(\boldsymbol{H}(\operatorname{Ai}_{t})(I + \boldsymbol{H}(\operatorname{Ai}_{t}))^{-1} \right) \psi_{k,t}, \psi_{k,t} \right\rangle_{L^{2}(\mathbb{R}_{+})} dt$$
$$= 2 \int_{s}^{+\infty} \left\langle (I + \boldsymbol{H}(\operatorname{Ai}_{t}))^{-2} \operatorname{Ai}_{t} \otimes \operatorname{Ai}_{t}^{*} \psi_{k,t}, \psi_{k,t} \right\rangle_{L^{2}(\mathbb{R}_{+})} dt$$
$$= 2 \int_{s}^{+\infty} \left\langle (I + \boldsymbol{H}(\operatorname{Ai}_{t}))^{-2} \operatorname{Ai}_{t}, \psi_{k,t} \right\rangle_{L^{2}(\mathbb{R}_{+})} \left\langle \operatorname{Ai}_{t}, \psi_{k,t} \right\rangle_{L^{2}(\mathbb{R}_{+})} dt$$
$$= 2 \int_{s}^{+\infty} \left\langle (I + \boldsymbol{H}(\operatorname{Ai}_{t}))^{-1} \operatorname{Ai}_{t}, \psi_{k,t} \right\rangle_{L^{2}(\mathbb{R}_{+})} dt$$

One has moreover $\mu_k(-\infty) = 1$ as $\mathbf{L}_{-\infty} = I$ and $\mu_k(+\infty) = 0$ as $\mathbf{L}_{+\infty} = 0$. We thus deduce the existence of a random variable Q_k such that

$$\mathbb{P}(Q_k > s) = 2 \int_s^{+\infty} \left\langle (I + \boldsymbol{H}(\operatorname{Ai}_t))^{-1} \operatorname{Ai}_t, \psi_{k,t} \right\rangle_{L^2(\mathbb{R}_+)}^2 dt$$

and, with independent random variables,

$$Q \stackrel{\mathcal{L}}{=} \max_{k \ge 1} Q_k \tag{41}$$

As a result,

$$\mathcal{TW}_1 \stackrel{\mathcal{L}}{=} \max_{k \ge 1} \left\{ \max \left\{ Q_k^{(1/2)}, Z_k(\mathrm{Ai})^{(1/2)} \right\} \right\}$$

Remark 2.14. The case of the *GSE* Tracy-Widom distribution is yet to be obtained. The description similar to (39) is [124, (54)]

$$\mathbb{P}(\mathcal{TW}_4 \leqslant s/\sqrt{2}) = \sqrt{\mathbb{P}(\mathcal{TW}_2 \leqslant s)} \cosh(U(t))^2, \qquad U(t) := \frac{1}{2} \int_t^{+\infty} q$$

but the cosh term does not seem to be expressible as a probability. Other expressions are available, such as the Ferrari-Spohn one [52, (35)] that reads

$$\mathbb{P}\left(\mathcal{TW}_{4} \leqslant s/\sqrt{2}\right) = \frac{1}{2} \left(\det\left(I - \boldsymbol{H}(\operatorname{Ai}_{s})\right)_{L^{2}(\mathbb{R}_{+})} + \det\left(I + \boldsymbol{H}(\operatorname{Ai}_{s})\right)_{L^{2}(\mathbb{R}_{+})}\right)$$

but the same problem remains.

3. Max-independence structure in random matrix ensembles

3.1. The *GUE*. The Gaussian Unitary Ensemble of size N, in short GUE_N , is the probability space $(\mathcal{H}_N(\mathbb{C}), \mathbb{P}_N)$ where \mathbb{P}_N is the Gaussian measure given by

$$\mathbb{P}_N(dM) = e^{-\operatorname{tr}(MM^*)/2} \frac{dM}{(2\pi)^{N^2/4}}$$
(42)

A disintegration of $\mathbf{M} \sim GUE_N$ according to the map $M \mapsto U^* \Lambda U$ where $U \in \mathcal{U}_N(\mathbb{C})$ and Λ is a diagonal real matrix gives the law of the eigenvalues $\boldsymbol{\lambda} = (\boldsymbol{\lambda}_{1,N}, \ldots, \boldsymbol{\lambda}_{N,N})$

$$\mathbb{P}_N(\boldsymbol{\lambda} \in d\boldsymbol{x}) = \frac{1}{N! \, \mathcal{Z}_N} \Delta(\boldsymbol{x})^2 e^{-|\boldsymbol{x}|^2/2} d\boldsymbol{x}, \quad \Delta(\boldsymbol{x}) := \prod_{1 \leq i < j \leq N} (x_i - x_j), \quad \mathcal{Z}_N := (2\pi)^{N/2} \prod_{k=0}^{N-1} k!$$

In particular, supposing that $\lambda_{1,N} \ge \cdots \ge \lambda_{N,N}$, one has

$$\mathbb{P}(\boldsymbol{\lambda}_{1,N} \leqslant s) = \frac{\mathcal{Z}_N(s)}{\mathcal{Z}_N(\infty)}, \qquad \mathcal{Z}_N(s) := \frac{1}{N!} \int_{(-\infty,s)^N} \Delta(\boldsymbol{x})^2 e^{-|\boldsymbol{x}|^2/2} d\boldsymbol{x}$$

We now recall, for the reader's convenience, some classical facts about this ensemble that one can find e.g. in [4, 11, 54, 86, 121].

The Andréieff-Heine-Szegö formula [120, p. 24] reads for $f_i, g_i \in L^2(\mathbb{R}, \mu)$

$$\frac{1}{N!} \int_{\mathbb{R}^N} \det\left(f_i(t_j)\right)_{1 \le i, j \le N} \det\left(g_i(t_j)\right)_{1 \le i, j \le N} d\mu^{\otimes N}(\boldsymbol{t}) = \det\left(\int_{\mathfrak{X}} f_i g_j d\mu\right)_{1 \le i, j \le N}$$
(43)

This is the continuous version⁵ of the Cauchy-Binet formula

$$\det\left(\sum_{\ell \geqslant 1} A_{i,\ell} B_{\ell,j}\right)_{1 \leqslant i,j \leqslant N} = \sum_{1 \leqslant \ell_1 < \ell_2 < \dots < \ell_N} \det\left(A_{i,\ell_j}\right)_{1 \leqslant i,j \leqslant N} \det\left(B_{\ell_i,j}\right)_{1 \leqslant i,j \leqslant N}$$
$$= \frac{1}{N!} \sum_{1 \leqslant \ell_1 \neq \ell_2 \neq \dots \neq \ell_N} \det\left(A_{i,\ell_j}\right)_{1 \leqslant i,j \leqslant N} \det\left(B_{\ell_i,j}\right)_{1 \leqslant i,j \leqslant N}$$
$$= \frac{1}{N!} \sum_{\ell_1,\dots,\ell_N \geqslant 0} \det\left(A_{i,\ell_j}\right)_{1 \leqslant i,j \leqslant N} \det\left(B_{\ell_i,j}\right)_{1 \leqslant i,j \leqslant N}$$
(44)

In the case of $d\mu(x) := e^{-x^2/2} \frac{dx}{\sqrt{2\pi}}$, one has

$$\begin{aligned} \mathcal{Z}_N(s) &:= \frac{1}{N!} \int_{(-\infty,s)^N} \Delta(\boldsymbol{x})^2 e^{-|\boldsymbol{x}|^2/2} \frac{d\boldsymbol{x}}{(2\pi)^{N/2}} \\ &= \int_{(-\infty,s)^N} \det\left(P_{i-1}(x_j)\right)_{1 \le i,j \le N} \det\left(Q_{i-1}(t_j)\right)_{1 \le i,j \le n} e^{-|\boldsymbol{x}|^2/2} \frac{d\boldsymbol{x}}{(2\pi)^{N/2}} \end{aligned}$$

for any family of monic polynomials $(P_k, Q_k)_{k \ge 0}$ satisfying $\deg(P_k) = \deg(Q_k) = k$ (taking linear combinations in the Vandermonde determinant). The choice of the truncated

⁵Using a discrete measure μ , one gets (44) from (43); but one can pass to the limit with a Riemann sum starting from (44) to obtain (43) which makes these two formuli equivalent.

Hermite polynomials $P_k = Q_k = H_k(\cdot|s)$, i.e. the monic orthogonal polynomials for $L^2((-\infty, s), \mu)$ gives a diagonal matrix, and the Andréieff-Heine-Szegö formula (43) then implies

$$\mathcal{Z}_N(s) = \prod_{k=0}^{N-1} \int_{-\infty}^s H_k(x|s)^2 e^{-x^2/2} \frac{dx}{\sqrt{2\pi}}$$

The formula

$$\mathbb{P}(\boldsymbol{\lambda}_{1,N} \leqslant s) = \prod_{k=0}^{N-1} \frac{\|H_k(\cdot|s)\|_{L^2((-\infty,s),\mu)}^2}{\|H_k(\cdot|\infty)\|_{L^2((-\infty,\infty),\mu)}^2}$$

is *not* of the form

$$\mathbb{P}(\boldsymbol{\lambda}_{1,N} \leqslant s) = \prod_{k=0}^{N-1} \int_{-\infty}^{s} f_{W_{k}}(x) dx = \mathbb{P}\left(\max_{0 \leqslant k \leqslant N-1} W_{k} \leqslant s\right)$$

since the polynomials depend on s. The following theorem allows to circumvent this fact à la Ginibre:

Theorem 3.1 (Max-independence structure in the GUE). Let $H_k(x|t)$ be the orthogonal polynomial for $L^2((-\infty, t), e^{-x^2/2} \frac{dx}{\sqrt{2\pi}})$ (truncated Hermite polynomial). Define the law of a random variable W_k by

$$\mathbb{P}(W_k \leqslant s) = \int_{-\infty}^s \mathcal{H}_k(x)^2 e^{-x^2/2} \frac{dx}{\sqrt{2\pi} \, k!}, \qquad \mathcal{H}_k(x) := H_k(x|x) \tag{45}$$

and let $(W_k)_k$ be a sequence of independent such random variables. Then,

$$\boldsymbol{\lambda}_{1,N} \stackrel{\mathcal{L}}{=} \max_{0 \leqslant k \leqslant N-1} W_k \tag{46}$$

Proof. A differentiation of $\int_{-\infty}^{s} H_k(x|s)^2 d\mu(x)$ with $d\mu(x) := e^{-x^2/2} \frac{dx}{\sqrt{2\pi}}$ gives

$$\frac{d}{ds} \int_{-\infty}^{s} H_k(x|s)^2 e^{-x^2/2} \frac{dx}{\sqrt{2\pi}} = H_k(s|s)^2 \frac{e^{-s^2/2}}{\sqrt{2\pi}} + \int_{-\infty}^{s} \frac{\partial}{\partial s} H_k(x|s)^2 e^{-x^2/2} \frac{dx}{\sqrt{2\pi}}$$
$$= H_k(s|s)^2 \frac{e^{-s^2/2}}{\sqrt{2\pi}} + 2 \int_{-\infty}^{s} H_k(x|s) \frac{\partial}{\partial s} H_k(x|s) e^{-x^2/2} \frac{dx}{\sqrt{2\pi}}$$

Since $H_k(X|s) = X^k + \sum_{j=0}^{k-1} a_{j,k}(s) X^j$ is a monic orthogonal polynomial, $\frac{\partial}{\partial s} H_k(X|s) = \sum_{j=0}^{k-1} \frac{\partial}{\partial s} a_{j,k}(s) X^j$ is of degree k-1, hence, by orthogonality $\langle H_k, \frac{\partial}{\partial s} H_k \rangle_{L^2((-\infty,s],\mu)} = 0$. Integrating from A to s this equality gives

$$\int_{-\infty}^{s} H_k(x|s)^2 d\mu(x) - \int_{-\infty}^{A} H_k(x|A)^2 d\mu(x) = \int_{A}^{s} H_k(x|x)^2 e^{-x^2/2} \frac{dx}{\sqrt{2\pi}}$$

Now, one can make $A \to -\infty$ on the LHS which is finite, and $\int_{-\infty}^{A} H_k(x|A)^2 d\mu(x) \to 0$ since $H_k(X|A)$ is a polynomial for all A and $H_k(X|-\infty) = 0$, giving the desired result.

Note that $s \to +\infty$ in this equality gives the renormalisation constant as $||H_k(\cdot|\infty)||^2_{L^2(\mathbb{R},\mu)} = k!$ for the Gaussian measure. This concludes the proof.

Remark 3.2.

- The independence structure (7) in the Ginibre ensemble can be directly proven starting from the Andréieff-Heine-Szegö formula. Nevertheless, the differentiation step is not necessary as the orthogonal polynomials are directly the power functions on the disk. The proof extends to any complex ensemble with a circular symmetry (see e.g. [11, ch. 4.3]).
- The argument of differentiation that we use to derive (45) was also employed by Majumdar-Nadal [88, § 3.2] in the core of their rederivation of the Tracy-Widom distribution (with a method involving norms of orthogonal polynomials originally due to Gross-Matytsin [65]), but with no probabilistic considerations.
- The differentiation argument is also the one used in the Fuchs lemma 2.2/2.6, and the norms of orthogonal polynomials are the eigenvalues of the projection operator with the usual Christoffel-Darboux kernel that reads $K_N(x, y) = \sum_{k=0}^{N} H_k(x|t)H_k(y|t)$ in $L^2((-\infty, t), e^{-x^2/2} \frac{dx}{\sqrt{2\pi}})$. Normalising each term of the sum by their L^2 -norm gives a sum of weighted projectors, the weight being the eigenvalue, i.e. $||H_k(\cdot|t)||^2_{L^2((-\infty,t],\mu)}$. Equivalently, one can use the truncated Hermite polynomials in $L^2([t, +\infty), e^{-x^2/2} \frac{dx}{\sqrt{2\pi}})$ and the eigenvalue will be the norm of these polynomials, $\widetilde{H}_k(x|t)$ (say) on this set. Of course, the formuli are equivalent : $k! - ||\widetilde{H}_k(\cdot,t)||^2_{[t,+\infty)} = ||H_k(\cdot|t)||^2_{(-\infty,t]}$.

This is the exact analogue of the Fuchs lemma that one applies (after an application of the Lidskii formula). The expression of the law uses then the function $\widetilde{H}_k(x|x)^2$ in the same way (19) uses $\Psi_k^{(s)}(0)^2 = \widetilde{\Psi}_k^{(s)}(s)^2 \equiv \widetilde{\Psi}_k(s|s)^2$.

3.2. Generalisation to any matrix ensemble on the real line and the circle. The proofs of the theorems here presented are identical to the previous one and are thus omitted.

3.2.1. The real line.

Definition 3.3 (Determinantal measure). Let μ be a probability measure on a space \mathfrak{X} and let $(\varphi, \psi) := (\varphi_k, \psi_k)_{k \ge 1}$ be a family of square integrable functions of $L^2(\mathfrak{X}, \mu)$. A determinantal measure with parameters (φ, ψ, μ) is the law of random points $(\lambda_{k,N})_{1 \le k \le N}$ in \mathfrak{X} with distribution (see e.g. [82])

$$\mathbb{P}_{N}(\boldsymbol{\lambda} \in d\boldsymbol{x}) = \frac{1}{N! \det(\int_{\mathfrak{X}} \varphi_{i} \overline{\psi_{j}} d\mu)_{1 \leqslant i, j \leqslant N}} \det(\varphi_{i}(x_{j}))_{1 \leqslant i, j \leqslant N} \overline{\det(\psi_{i}(x_{j}))_{1 \leqslant i, j \leqslant N}} d\mu^{\otimes N}(\boldsymbol{x}) \quad (47)$$

We note $\lambda \sim \text{DetMeas}(\varphi, \psi, \mu | N)$ for such a random process. Note that the equality $\int_{\mathfrak{X}^N} \mathbb{P}_N(\lambda \in d\mathbf{x}) = 1$ is the Andréieff-Heine-Szegö formula (43).

Theorem 3.1 can be easily modified in the case of a general determinantal measure to give:

Theorem 3.4 (Independence structure in matrix ensembles on the real line). Consider random points $(\lambda_{k,N})_{1 \leq k \leq N}$ in \mathbb{R} with distribution

$$\mathbb{P}_N(\boldsymbol{\lambda} \in d\boldsymbol{x}) = rac{1}{N! \, \mathcal{Z}_N} \Delta(\boldsymbol{x})^2 d\mu^{\otimes N}(\boldsymbol{x}), \qquad d\mu(x) := \dot{\mu}(x) dx$$

i.e. $\boldsymbol{\lambda} \sim \text{DetMeas}(\boldsymbol{\psi}, \boldsymbol{\psi}, \mu | N)$ with $\psi_k(x) := x^{k-1}$. Then, one has

$$\boldsymbol{\lambda}_{1,N} \stackrel{\mathcal{L}}{=} \max_{1 \leqslant k \leqslant N} Y_k \tag{48}$$

where $(Y_k)_k$ is a sequence of independent random variables with distribution given by

$$\mathbb{P}(Y_{k+1} \leqslant s) = \int_{-\infty}^{s} P_k(x|x)^2 \frac{d\mu(x)}{\int_{\mathbb{R}} P_k(y|\infty)^2 d\mu(y)}$$
(49)

where $P_k(\cdot|t)$ is the orthogonal polynomial for $L^2((-\infty,t),\mu)$.

The proof is a straightforward generalisation of the previous one and is thus omitted.

3.2.2. The circle.

Theorem 3.5 (Independence structure in matrix ensembles on the circle). Consider random points $(e^{i\Theta_{k,N}})_{1 \leq k \leq N}$ in \mathbb{U} with distribution

$$\mathbb{P}_{N}(\boldsymbol{\Theta} \in d\boldsymbol{\theta}) = \frac{1}{N! \,\mathcal{Z}_{N}} \left| \Delta(e^{i\boldsymbol{\theta}}) \right|^{2} d\mu^{\otimes N}(\boldsymbol{\theta}), \qquad d\mu(\theta) := \dot{\mu}(e^{i\theta}) d\theta, \qquad \theta_{k} \in [0, 2\pi)$$

Then, one has

$$\Theta_{1,N} \stackrel{\mathcal{L}}{=} \max_{1 \leqslant k \leqslant N} A_k \tag{50}$$

where $(A_k)_k$ is a sequence of independent random variables with distribution given by

$$\mathbb{P}(A_{k+1} \leqslant s) = \int_0^s \left| P_k(e^{i\theta}|\theta) \right|^2 \frac{d\mu(\theta)}{\int_0^{2\pi} |P_k(e^{i\alpha}|2\pi)|^2 d\mu(\alpha)}$$
(51)

where $\theta \mapsto P_k(e^{i\theta}|t)$ is the monic OPUC for $L^2([0,t),\mu)$.

Remark 3.6. The Weyl integration formula reads

$$\int_{\mathcal{U}_N} \det(f(U)) dU = \frac{1}{N!} \oint_{\mathbb{U}^N} f^{\otimes N}(\boldsymbol{u}) |\Delta(\boldsymbol{u})|^2 \frac{d^* \boldsymbol{u}}{\boldsymbol{u}}$$
(52)

where dU is the normalised Haar measure on the unitary group \mathcal{U}_N and $\det(f) \in L^1(\mathcal{U}_N, dU)$.

This formula defines the *Circular Unitary Ensemble* or CUE_N [45, 46, 54] that has been studied extensively, notably using the Andréieff-Heine-Szegö formula (43) that links it to

a Toeplitz determinant of symbol f [46, fact five]

$$\int_{\mathcal{U}_N} \det(f(U)) dU = \det(\mathbb{T}_N(f)), \qquad \mathbb{T}_N(f) := \left(\left[z^{i-j} \right] f(z) \right)_{1 \le i, j \le N}$$
(53)

The previous theorem with $\dot{\mu} = 1$ treats thus the case of the CUE_N .

In view of the last results, an interesting question emerges:

Question 3.7. Can one find an almost sure construction that proves (46) or (48)?

In fact, question 3.7 is already relevant at the level of the Ginibre ensemble, as the identity in law $\lambda_1^{\text{GinE}_{N+1}} \stackrel{\mathcal{L}}{=} \max\{\lambda_1^{\text{GinE}_N}, \gamma_{N+1}\}$ is not proven with an almost sure construction.

Remark 3.8. By "almost sure construction", we mean a deterministic equality that would imply the probabilistic result once "probabilised". Such an identity was for instance given in [36] for the characteristic polynomial of a random matrix distributed according to the CUE_N . Setting $Z_{U_N}(X) := \det(XI_N - U_N)$, one has $Z_{U_N}(1) \stackrel{\mathcal{L}}{=} \prod_{k=1}^N X_k$ where X_k are explicit independent random variables. Such a decomposition comes from the decomposition into a product of transvections of a unitary matrix which implies the probabilistic identity $U_N \stackrel{\mathcal{L}}{=} H_{u_1}H_{u_2}\ldots H_{u_N}$ where the random transvections $H_{u_i}: x \mapsto$ $x - \left(1 + \frac{1-e_1^*u_i}{1-e_1^*u_i}\right) \frac{(e_1-u_i)^*x}{|e_1-u_i|^2}(e_1 - u_i)$ are independent (the vector u_i is selected uniformly on the unit sphere and $(e_k)_k$ is the canonical basis of \mathbb{C}^N). The previous identity in law comes then from the deterministic identity (see [36] for the details)

$$\det(I_N - H_{u_1}H_{u_2} \dots H_{u_N}) = (1 - e_1^*H_{u_1}e_1)\det(I_{N-1} - H_{u_2'} \dots H_{u_N'})$$

Taking the Mellin transform of this identity gives $\mathbb{E}\left(Z_{U_N}(1)^s \overline{Z_{U_N}(1)^w}\right) = \prod_{k=1}^N \mathbb{E}\left(X_k^s \overline{X_k^w}\right)$, an identity equivalent to the celebrated Selberg integral in the particular case $\beta = 2$ (see [37] for a generalisation to the $C\beta E_N$). Such decompositions allow thus to give new proofs of classical identities with almost no computations, and a better understanding of why some "magical" identities hold.

Remark 3.9. Although there exists a natural Markovian coupling between GUE_N and GUE_{N+1} , the minor process [27], the Courant-Fisher minimax characterisation of the eigenvalues of a Hermitian matrix is not likely to give the desired result when probabilised with this coupling since it can be stationary (this is seemingly not the case of $(W_k)_k$).

Of course, several other Markovian couplings are possible, although arguably less natural than the minor one. We will come back to this problem in the future.

3.3. Note on $\lambda_{N,N}$. We quickly treat the case of the minimal eigenvalue $\lambda_{N,N}$ in any previously defined matrix ensemble. This random variable is also the minimum of N independent random variables, as, using the Andréieff-Heine-Szegö formula (43), one obtains

$$\mathbb{P}(\boldsymbol{\lambda}_{N,N} \ge s) = \frac{1}{\mathcal{Z}_N(\infty)} \prod_{k=0}^{N-1} \left\| \widetilde{H}_k(\cdot|s) \right\|_{L^2([s,+\infty),\mu)}^2$$

This is the half-line $[s, +\infty)$ which is relevant in this case, and the orthogonal polynomials for the restricted measure were denoted $\widetilde{H}_k(\cdot|s)$ in remark 3.2. Using the same manipulations as in theorem 3.1, one then finds

$$\mathbb{P}(\boldsymbol{\lambda}_{N,N} \ge s) = \frac{1}{\mathcal{Z}_N(\infty)} \prod_{k=0}^{N-1} \left\| \widetilde{H}_k(\cdot|\cdot) \right\|_{L^2([s,+\infty),\mu)}^2 = \prod_{k=0}^{N-1} \frac{\left\| \widetilde{H}_k(\cdot|\cdot) \right\|_{L^2([s,+\infty),\mu)}^2}{\left\| \widetilde{H}_k(\cdot|-\infty) \right\|_{L^2(\mathbb{R},\mu)}^2}$$

In the case of the GUE_N , one has for instance with independent $(W_k)_k$

$$\boldsymbol{\lambda}_{N,N}^{(GUE)} \stackrel{\mathcal{L}}{=} \min_{0 \leqslant k \leqslant N-1} \widetilde{W}_k$$
(54)

where

$$f_{\widetilde{W}_k}(x) = \frac{\widetilde{H}_k(x|x)^2}{\sqrt{k!}} \frac{e^{-x^2/2}}{\sqrt{2\pi}}$$

Here again, the most interesting question is the equivalent of Question 3.7, i.e.

Question 3.10. Can one find an almost sure construction that proves (54) or its generalisations to other determinantal measures ?

For instance, the construction $\boldsymbol{H}_{N+1} = \boldsymbol{H}_N \oplus (\boldsymbol{X}_{N+1}, a_{N+1})$ implies a Markovian coupling. Here, \oplus designates the operation that puts \boldsymbol{H}_N in the top $N \times N$ block of an $(N+1) \times (N+1)$ matrix that has the vector \boldsymbol{X}_{N+1} (of size N) in the upper right block, its transconjugated in the lower left block and the real a_{N+1} at the (N+1, N+1) index (or the lower right block).

With such a construction (with no probability), one looks for an equality of the type

$$\lambda_{\min}(\boldsymbol{H}_{N+1}) = \min \left\{ \lambda_{\min}(\boldsymbol{H}_N), \ \mathcal{F}_{N+1}(\boldsymbol{X}_{N+1}, a_{N+1}) \right\}$$

for a certain functional \mathcal{F}_{N+1} . The probabilisation of such an identity would then imply the result, with $\widetilde{W}_N \stackrel{\mathcal{L}}{=} \mathcal{F}_{N+1}(\boldsymbol{X}_{N+1}, a_{N+1})$. We will come back to this problem in the future.

3.4. Painlevé representation of \mathcal{H}_n and a new expression of $F_{\mathcal{TW}_2}$.

3.4.1. Painlevé representation of \mathcal{H}_n . The function \mathcal{H}_n used in (45) can be described in a more explicit way.

Recall that the monic orthogonal polynomials $(P_k)_{k\geq 0}$ for $L^2([a, b], \mu)$ with $a, b \in \mathbb{R}$ are given by the following "expectations of characteristic polynomials of random matrix ensembles" [120, (2.2.10) p. 27]

$$P_n(X) = \frac{1}{n! D_n} \int_{[a,b]^n} \prod_{k=1}^n (X - t_k) \Delta(t)^2 d\mu^{\otimes n}(t), \qquad D_n := \frac{1}{n!} \int_{[a,b]^n} \Delta(t)^2 d\mu^{\otimes n}(t)$$

In particular, for

$$\mu(dx) := \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx, \qquad a := -\infty, \qquad b := t$$

one gets

$$H_n(X|t) = \frac{1}{n! D_n(t)} \int_{(-\infty,t)^n} \prod_{k=1}^n (X - t_k) \Delta(t)^2 d\mu^{\otimes n}(t), \qquad D_n(t) := \frac{1}{n!} \int_{(-\infty,t)^n} \Delta(t)^2 d\mu^{\otimes n}(t)$$

If $\mathbf{H}_n \sim GUE_n$ with eigenvalues $\lambda_{1,n} \ge \lambda_{2,n} \ge \cdots$, and if $Z_{\mathbf{H}_n}(x) := \det(xI_n - \mathbf{H}_n)$, one thus has

$$H_n(x|t) = \frac{\mathbb{E}\left(Z_{H_n}(x)\mathbb{1}_{\{\boldsymbol{\lambda}_{1,n} \leqslant t\}}\right)}{\mathbb{P}(\boldsymbol{\lambda}_{1,n} \leqslant t)} = \mathbb{E}\left(Z_{H_n}(x) \big| \boldsymbol{\lambda}_{1,n} \leqslant t\right)$$

Using the notations of $[56]^6$, define for all $a \in \mathbb{N}$

$$\widetilde{E}_n(x;a) := \frac{1}{n!} \int_{(-\infty,x)^n} \prod_{j=1}^n (x-t_j)^a \Delta(t)^2 d\mu^{\otimes n}(t)$$

Then,

$$\mathcal{H}_n(x) := H_n(x|x) = \frac{\widetilde{E}_n(x;1)}{\widetilde{E}_n(x;0)}$$
(55)

The quantities $\widetilde{E}_n(x; a)$ were studied by Forrester-Witte in [56, (1.7)]. This article is an application of the Jimbo-Miwa σ -form approach to the Painlevé IV equation [72] and the τ -function theory of the Painlevé II to VI equations due to Okamoto [94, 95, 96, 97] later refined by Noumi-Yamada [91]; see also Forrester [54, ch. 8 p. 328] for a historical and a self-contained introduction on the topic and Noumi [90] for a detailed study of P_{II} and P_{IV} .

⁶Note that the Gaussian measure is not the probabilistic one in [56] and in [54, 123], hence some slight changes of conventions.

It is proven in [56, (1.8)/(4.14)] that

$$\widetilde{E}_n(x;a) = \widetilde{E}_n(x_0;a) \exp\left(\int_{x_0}^x U_n(t;a)dt\right)$$

where $y = U_n(\cdot; a)$ satisfies the following σ -form of the Painlevé IV equation [56, (1.9)/(4.15)]

$$(\sigma - P_{IV}) : \begin{cases} (y'')^2 - 4(ty' - y)^2 + 4y'(y' - 2a)(y' + 2n) = 0\\ y(t) = -2nt - \frac{n(a+n)}{t} + O\left(\frac{1}{t^3}\right) & \text{when } t \to -\infty. \end{cases}$$
(56)

the boundary conditions being given in [56, (1.10)/(4.18)].

Notice that

$$\widetilde{E}_n(x;a) := \frac{1}{n!} \int_{(-\infty,x)^n} \prod_{j=1}^n (x-t_j)^a \Delta(t)^2 d\mu^{\otimes n}(t) \quad \underset{x \to +\infty}{\sim} \quad x^{na} \times \widetilde{E}_n(+\infty;0)$$

implying that

$$\mathcal{H}_n(x) \underset{x \to +\infty}{\sim} x^n$$

As a result, one cannot choose $x_0 = +\infty$ except in the case a = 0 (which is the case considered by Tracy and Widom).

For general x_0 , one thus has

$$\mathcal{H}_n(x)e^{-x^2/4} = \mathcal{H}_n(x_0)e^{-x_0^2/4} \exp\left(\int_{x_0}^x \left(U_n(t;1) - U_n(t;0) - \frac{t}{2}\right)dt\right)$$
(57)

For a = 0, Tracy and Widom's result

$$\frac{\boldsymbol{\lambda}_{1,N} - 2\sqrt{N}}{N^{-1/6}} \xrightarrow[N \to +\infty]{\mathcal{L}} \mathcal{TW}_2$$
(58)

gives with the notations of [56, (1.18)/(5.2), (5.3)] (where u is replaced by σ)

$$\widetilde{E}_N\left(2\sqrt{N} + N^{-\frac{1}{6}}s; a = 0\right) \xrightarrow[N \to +\infty]{} \exp\left(-\int_s^{+\infty} r(t)dt\right), \qquad r(t) =: \sigma(t; a = 0)$$

and for $a \ge 1$ [56, (1.19)/(5.6), (5.7)]

$$C(s_0) e^{-a(2\sqrt{N}+N^{-1/6}s)^2/4} \widetilde{E}_N\left(2\sqrt{N}+N^{-\frac{1}{6}}s\,;\,a\right) \xrightarrow[N\to+\infty]{} \widetilde{E}_\infty^{\text{soft}}(s_0;\,a) \exp\left(\int_{s_0}^s \sigma(t;a)dt\right)$$

were $y := \sigma(\cdot; a)$ satisfies the following σ - form of the Painlevé II equation [56, (1.17)/(5.10)]

$$(\sigma - P_{II}) \begin{cases} (y'')^2 + 4y' \left(ty' - y \right) - 4(y')^3 - a^2 = 0 \\ y(t) = \frac{t^2}{4} + \frac{4a^2 - 1}{8t} + O\left(\frac{1}{t^3}\right) & \text{when } t \to -\infty. \end{cases}$$

$$(59)$$

the boundary conditions being given in [56, (1.20)/(4.11)]

3.4.2. Required estimates for f_{W_n} .

Caveat: We will now be concerned with the "physicist convention" of Tracy-Widom [122, 123] and Forrester-Witte [56] which is also the convention of Wu-Xu-Zhao [130]. As a result, the probability expressed in theorem 1.4 is given with $t\sqrt{2}$ and the Gaussian density is $e^{-t^2}/\sqrt{\pi}$. We will set $\widehat{W}_n := \sqrt{2}W_n$ for the rescaled random variable.

We will moreover use the function σ which is denoted by u in [56] and the function p which is denoted by u in [130]. These conventions are the ones of [105, 32.6(iii) p. 729] for the Hamiltonian description of the Painlevé functions.

The Hamiltonian description goes as follows : denote by H(q, p, t) the Hamiltonian of the Painlevé II equation given by [105, 32.6.9 p. 729]

$$H(q, p, t) := \frac{p^2}{2} - \left(q^2 + \frac{t}{2}\right)p - \left(\alpha + \frac{1}{2}\right)q$$

The Hamilton-Jacobi equations $q' = \partial_p H$, $p' = -\partial_q H$ are given by [105, 32.6.10 & 32.6.11 p. 729]

$$q' = p - q^2 - \frac{t}{2}, \qquad p' = 2qp + \alpha + \frac{1}{2}$$

The equation satisfied by q is then $P_{II}(\alpha)$ given by (4) and the equation satisfied by p is the Painlevé XXXIV equation $P_{XXXIV}(\alpha)$ (numbered in this way in the original classification of Painlevé and Gambier) given by [105, 32.6.12 p. 729]

$$pp'' = \frac{(p')^2}{2} + 2p^3 - tp^2 - \frac{1}{2}(\alpha + \frac{1}{2})^2$$
(60)

Last, the equation satisfied by the function $\sigma : t \mapsto H(q(t), p(t), t)$ is σ - $P_{II}(\alpha)$ given by (59). This function allows to "reconstruct" (p, q) with the relations [105, 32.6.14 & 32.6.15 p. 729] :

$$q = \frac{4\sigma'' + 2\alpha + 1}{8\sigma'}, \qquad p = -2\sigma' \tag{61}$$

In [130, (1.6)], Wu, Xu and Zhao study

$$\boldsymbol{H}_{n}(x|\alpha,\omega) := \frac{1}{n!} \int_{\mathbb{R}^{n}} \prod_{k=1}^{n} |x - t_{j}|^{2\alpha} \left(\mathbb{1}_{\{t_{j} \leq x\}} + \omega \mathbb{1}_{\{t_{j} > x\}} \right) \Delta(\boldsymbol{t})^{2} \mu^{\otimes n}(d\boldsymbol{t}), \qquad \mu(dt) = e^{-t^{2}} \frac{dt}{\sqrt{\pi}}$$

In particular,

$$\boldsymbol{H}_n(\boldsymbol{x}|\alpha,0) := \frac{1}{n!} \int_{\mathbb{R}^n} \prod_{k=1}^n \left| (\boldsymbol{x} - \boldsymbol{t}_j)_+ \right|^{2\alpha} \Delta(\boldsymbol{t})^2 \mu^{\otimes n}(d\boldsymbol{t}) = \widetilde{E}_n(\boldsymbol{x}; \boldsymbol{a} = 2\alpha)$$

as $x_{+} = |x_{+}|$, and

$$\mathcal{H}_n(x) = \frac{\boldsymbol{H}_n(x|1/2,0)}{\boldsymbol{H}_n(x|0,0)}$$

Wu, Xu and Zhao [130, thm. 3, (1.24)] prove moreover that (in particular for $\alpha = \frac{1}{2}/a = 1$ and $\omega = 0$)

$$\frac{d}{d\mu_n} \ln \boldsymbol{H}_n(\mu_n | \alpha, \omega) = \sqrt{2n} \left(2\alpha + \frac{\sigma(s | \alpha, \omega)}{n^{1/3}} + \alpha \frac{p(s | \alpha, \omega) + s}{n^{2/3}} + O_s\left(\frac{1}{n}\right) \right),$$
$$\mu_n := \sqrt{2n} + n^{-\frac{1}{6}} \frac{s}{\sqrt{2}}$$

where $\sigma(\cdot | \alpha = 1/2, \omega = 0)$ is the σ -form of the Painlevé II function given in (59) and $p(\cdot | \alpha = \frac{1}{2}, \omega = 0)$ is solution of the Painlevé XXXIV equation with the following particular boundary condition at ∞ [130, (1.20) & (1.21)] for $\alpha \neq 0$:

$$(P_{XXXIV}) \begin{cases} p(s)p''(s) = \frac{p'(s)^2}{2} + 4p(s)^3 + 2sp(s)^2 - 2\alpha^2 \\ p(s) = \frac{\alpha}{\sqrt{s}}(1 - \alpha s^{-3/2} + O(s^{-3})), \quad s \to \infty, \quad \arg(s) \in (-\frac{\pi}{3}, \frac{\pi}{3}] \end{cases}$$
(62)

Last, [130, (1.22), (1.17)]/[56, (5.11), -]:

$$\sigma(s|\alpha,\omega=0) = -\frac{s}{2} + \frac{16\alpha^2 - 1}{8}s^{-2} + O(s^{-7/2}), \qquad s \to -\infty$$

$$\sigma(s|\alpha,\omega=0) = -\alpha\sqrt{s}\left(1 + \alpha s^{-3/2} + O(s^{-3})\right), \qquad s \to \infty, \quad \arg(s) \in (-\frac{\pi}{3}, \frac{\pi}{3}]$$
(63)

As a result, using the physicist convention for the Gaussian, one has

$$f_{W_n}(s) = \frac{\mathcal{H}_n(s)^2 e^{-s^2}}{n! \mathcal{Z}_n \sqrt{2\pi}} = \left(\frac{\mathcal{H}_n(s|1/2,0)}{\mathcal{H}_n(s|0,0)}\right)^2 \frac{e^{-s^2}}{n! \mathcal{Z}_n \sqrt{\pi}} \\ = \exp\left(2\ln\mathcal{H}_n(c) + 2\int_c^s \frac{d}{dt}\ln\mathcal{H}_n(t)dt\right) \frac{e^{-s^2}}{n! \mathcal{Z}_n \sqrt{2\pi}} \\ = \exp\left(2\ln\mathcal{H}_n(c) + 2\int_c^s \left(\frac{d}{dt}\ln\mathcal{H}_n(t|1/2,0) - \frac{d}{dt}\ln\mathcal{H}_n(t|0,0)\right)dt\right) \frac{e^{-s^2}}{n! \mathcal{Z}_n \sqrt{\pi}} \\ = \frac{\mathcal{H}_n(c)^2 e^{-c^2}}{n! \mathcal{Z}_n \sqrt{\pi}} \exp\left(2\int_c^s \left(\frac{d}{dt}\ln\mathcal{H}_n(t|1/2,0) - \frac{d}{dt}\ln\mathcal{H}_n(t|0,0) - t\right)dt\right)$$

Set

$$t_{n} := \sqrt{2n} + n^{-\frac{1}{6}} \frac{t}{\sqrt{2}}, \qquad s_{n} := \sqrt{2n} + n^{-\frac{1}{6}} \frac{s}{\sqrt{2}}, \qquad c_{n} := \sqrt{2n} + n^{-\frac{1}{6}} \frac{c}{\sqrt{2}}$$
$$\delta\sigma := \sigma(\cdot|\frac{1}{2},\omega) - \sigma(\cdot|0,\omega) = \sigma(\cdot|a=1,\omega) - \sigma(\cdot|a=0,\omega), \qquad \omega := 0$$

Then,

$$\begin{split} f_{\widehat{W}_n}(s_n) &=: \frac{\mathcal{H}_n(c_n)^2 e^{-c_n^2}}{n! \mathcal{Z}_n \sqrt{\pi}} \exp\left(2\int_c^s \left(\sqrt{2n} + \sqrt{2n} \frac{\delta \sigma^{(n)}(t_n)}{n^{1/3}} + \sqrt{2n} \frac{1}{2} \frac{p^{(n)}(t_n|\frac{1}{2}, 0) + t_n}{n^{2/3}} - t_n\right) n^{-\frac{1}{6}} dt\right) \\ &= \frac{\mathcal{H}_n(c_n)^2 e^{-c_n^2}}{n! \mathcal{Z}_n \sqrt{\pi}} \exp\left(n^{-\frac{1}{6}} 2\sqrt{2n} \int_c^s \left(1 + \frac{\delta \sigma^{(n)}(t_n)}{n^{1/3}} + \frac{p^{(n)}(t_n|\frac{1}{2}, 0) + t_n}{2n^{2/3}} - \frac{t_n}{\sqrt{2n}}\right) dt\right) \end{split}$$

$$= \frac{\mathcal{H}_{n}(c_{n})^{2}e^{-c_{n}^{2}}}{n!\mathcal{Z}_{n}\sqrt{\pi}} \exp\left(n^{\frac{1}{3}}2\sqrt{2}\int_{c}^{s}\left(\frac{\delta\sigma^{(n)}(t_{n})}{n^{1/3}} + \frac{p^{(n)}(t_{n}|\frac{1}{2},0) + t_{n}}{2n^{2/3}} - \frac{n^{-\frac{1}{6}}t/\sqrt{2}}{n^{1/2}\sqrt{2}}\right)dt\right)$$

$$= \frac{\mathcal{H}_{n}(c_{n})^{2}e^{-c_{n}^{2}}}{n!\mathcal{Z}_{n}\sqrt{\pi}} \exp\left(n^{\frac{1}{3}}2\sqrt{2}\int_{c}^{s}\left(\frac{\delta\sigma^{(n)}(t_{n})}{n^{1/3}} + \frac{p^{(n)}(t_{n}|\frac{1}{2},0) + t_{n}}{2n^{2/3}} - \frac{t}{2n^{2/3}}\right)dt\right)$$

$$= \frac{\mathcal{H}_{n}(c_{n})^{2}e^{-c_{n}^{2}}}{n!\mathcal{Z}_{n}\sqrt{\pi}} \exp\left(2\sqrt{2}\int_{c}^{s}\left(\delta\sigma(t) + \frac{p(t|\frac{1}{2},0) + t}{2n^{1/3}} - \frac{t}{2n^{1/3}} + O_{t}\left(\frac{1}{n^{2/3}}\right)\right)dt\right)$$

$$= \frac{\mathcal{H}_{n}(c_{n})^{2}e^{-c_{n}^{2}}}{n!\mathcal{Z}_{n}\sqrt{\pi}} \exp\left(2\sqrt{2}\int_{c}^{s}\left(\delta\sigma(t) + \frac{p(t|\frac{1}{2},0)}{2n^{1/3}} + O_{t}\left(\frac{1}{n^{2/3}}\right)\right)dt\right)$$

Using the second relation of (61), one then has

$$p(t|\frac{1}{2},0) = -2\frac{d}{dt}\sigma(t|\frac{1}{2},0)$$

and

$$f_{\widehat{W}_n}(s_n) = \frac{\mathcal{H}_n(c_n)^2 e^{-c_n^2 + 2\sqrt{2}n^{-1/3}\sigma(c|1/2,0)}}{n!\mathcal{Z}_n\sqrt{\pi}} \exp\left(2\sqrt{2}\int_c^s \delta\sigma(t)dt - O\left(n^{-1/3}\sigma(s|\frac{1}{2},0)\right)\right)$$
$$= \frac{\mathcal{H}_n(c_n)^2 e^{-c_n^2 + 2\sqrt{2}n^{-1/3}\sigma(c|1/2,0)}}{n!\mathcal{Z}_n\sqrt{\pi}} \exp\left(2\sqrt{2}\int_c^s \delta\sigma(t)dt\right) \left(1 - O\left(n^{-1/3}\sigma(s|\frac{1}{2},0)\right)\right)$$

namely

$$f_{\widehat{W}_{n}}(s_{n}) - f_{\widehat{W}_{n}}(c_{n})e^{2\sqrt{2}n^{-1/3}\sigma(c|1/2,0)}\exp\left(2\sqrt{2}\int_{c}^{s}\delta\sigma(t)dt\right) = -O\left(n^{-1/3}\sigma(s|\frac{1}{2},0)e^{2\sqrt{2}\int_{c}^{s}\delta\sigma(t)dt}\right)$$
(64)

The proof of theorem 3.11 will impose to choose the constant $c \equiv c_n^*$ such that

$$n^{1/6} f_{\widehat{W}_n}(c_n) e^{2\sqrt{2}n^{-1/3}\sigma(c|1/2,0)} = 1, \qquad c_n := \sqrt{2n} + n^{-\frac{1}{6}} \frac{c}{\sqrt{2}}$$

One can show that $c_n^* \to +\infty$ and that $s \mapsto \sigma(s|\frac{1}{2}, 0)e^{-2\sqrt{2}\int_s^{+\infty}\delta\sigma(t)dt} \in L^1([s, +\infty))$ for all s using the § 3.4.5.

3.4.3. Poisson approximation for maxima of independent random variables. The problem of estimating the maximum of a sequence of independent random variables $(W_k)_{1 \le k \le n}$ can be classically restated as a problem of Poisson approximation for sums of independent random variables [51, ch. 3] writing

$$\left\{\max_{1\leqslant k\leqslant n} W_k\leqslant x\right\} = \left\{\sum_{k=1}^n \mathbb{1}_{\{W_k>x\}} = 0\right\}$$

Since the random parametric indicators $B_k := \mathbb{1}_{\{W_k > x\}}$ are Bernoulli random variables of expectation $p_k := \mathbb{P}(W_k > x)$, one can proceed to a Poisson approximation of $S_n :=$ $\sum_{k=1}^{n} B_k$ [110, § 4.1]

$$\mathbb{P}(S_n = 0) - \mathbb{P}\left(\operatorname{Po}\left(\sum_{k=1}^n p_k\right) = 0\right) = O\left(\sum_{k=1}^n p_k^2 \times \min\left\{1, \frac{1}{\sum_{k=1}^n p_k}\right\}\right)$$

with

$$\mathbb{P}\left(\operatorname{Po}\left(\sum_{k=1}^{n} p_{k}\right) = 0\right) = e^{-\sum_{k=1}^{n} p_{k}}$$

In the GUE_{n+1} case, this last Poisson approximation is performed for the previous independent random variables $(W_k)_k$ defined in (45). Set

$$s_n := 2\sqrt{n} + n^{-1/6}s$$

As the first random variables do not depend on n, the first probabilities are very small : for k = O(1)

$$p_k := \mathbb{P}(W_k > s_n) \leqslant \frac{1}{s_n^2} \mathbb{E}\left(|W_k|^2\right) = O_s\left(\frac{1}{n}\right)$$

We will see in (68) that $f_{W_{m_n}}(s_n) \sim n^{-1/6} \mathcal{Q}(s+c)$ with $\mathcal{Q} \in L^2([s,+\infty))$ for all s if $m_n = n - cn^{1/3}$. As a result, with $x_n = 2\sqrt{n} + n^{-1/6}x$

$$p_{m_n} := \mathbb{P}(W_{m_n} > s_n) \leqslant \int_s^{+\infty} \left(\frac{x_n}{s_n}\right)^2 f_{W_{m_n}}(x_n) n^{-1/6} dx = O_s\left(n^{-1/3}\right)$$

The relevant part of the maximum mass is thus concentrated on the extreme random variables $(W_{[n\alpha]})_{1-\varepsilon_n \leqslant \alpha \leqslant 1}$ with $\alpha = 1 - cn^{-2/3}$ and one can neglect the first random variables to only consider the contribution of the last ones : with $m := [n(1-\varepsilon_n)] < n$, $\varepsilon = cn^{-2/3}$, one has with obvious notations

$$\delta_n := |\mathbb{P}(S_n = 0) - \mathbb{P}(S_{m \to n} = 0)| = \mathbb{P}(S_{m \to n} = 0, S_n - S_{m \to n} \ge 1)$$
$$\leqslant \mathbb{E}(S_n - S_{m \to n}) = \mathbb{E}(S_m) = \sum_{k=1}^m p_k$$

which implies

$$\mathbb{P}(S_n = 0) = \mathbb{P}(S_{m \to n} = 0) + O\left(\sum_{k=1}^m p_k\right) = \mathbb{P}(S_{m \to n} = 0) + O_s\left(n^{-1/3}\right)$$

for the choice

$$m = n - O(n^{1/3}) \tag{65}$$

For this choice of m, we also set

$$P_n := \sum_{k=1}^n p_k, \qquad P_n^* := \sum_{k=m+1}^n p_k, \qquad m := [n(1-\varepsilon_n)] = [n-cn^{1/3}]$$

so that

$$P_n = P_n^* + O_s \left(n^{-1/3} \right)$$

Last, we set

$$P_n^{(2)} := \sum_{k=1}^n p_k^2 = P_n^{*\,2} + O_s\Big(n^{-1/3}\Big), \qquad P_n^{*,2} := \sum_{k=m+1}^n p_k^2$$

If $P_n = O(1)$, the Poisson approximation amounts thus to

$$\mathbb{P}(S_n = 0) - \mathbb{P}(\operatorname{Po}(P_n) = 0) = O(P_n^{(2)}),$$

with

$$P_n = P_n^* + O_s(n^{-1/3}), \quad P_n^{(2)} = P_n^{*,2} + O_s(n^{-1/3})$$

3.4.4. A new expression of F_{TW_2} .

Theorem 3.11 (Expression of F_{TW_2}). One has

$$\mathbb{P}\left(\frac{\boldsymbol{\lambda}_{1,N} - 2\sqrt{N}}{N^{-1/6}} \leqslant \sqrt{2} s\right) \xrightarrow[N \to +\infty]{} \mathbb{P}\left(\mathcal{T}\mathcal{W}_{2} \leqslant \sqrt{2} s\right) = \exp\left(-\mathcal{Q}^{2} \ast \mathrm{id}_{+}(s)\right)$$
(66)

with $f * g(x) := \int_{\mathbb{R}} f(t)g(x-t)dt = g * f(x)$, $id_+ : x \mapsto x_+ := x \mathbb{1}_{\{x \ge 0\}}$ and

$$Q(x) := \exp\left(-\int_{x}^{+\infty} \delta\sigma\right), \qquad \delta\sigma := \sigma(\cdot; 1) - \sigma(\cdot; 0) \tag{67}$$

where $\sigma(\cdot; a)$ is the solution of the σ -form of the Painlevé II equation given in (59).

Proof. With $\widehat{W}_k := W_k/\sqrt{2}$, one has

$$\mathbb{P}\Big(\widehat{W}_{[N\alpha]} > \sqrt{2N} + N^{-1/6} 2^{-\frac{1}{2}}s\Big) = \int_{\sqrt{2N}+N^{-1/6}2^{-\frac{1}{2}}s}^{+\infty} f_{\widehat{W}_{[N\alpha]}}(x)dx$$
$$= \int_{s}^{+\infty} N^{-1/6} f_{\widehat{W}_{[N\alpha]}}(\sqrt{2N} + N^{-1/6}2^{-\frac{1}{2}}y)dy$$

and, with a Stieltjes integral,

$$P_N^* := \sum_{k=m+1}^N p_k = \int_m^N p_{[u]} du = \int_0^1 p_{[m+(N-m)v]}(N-m) dv$$

= $\int_{N^{2/3}}^0 p_{[N\alpha_N(c)]} d(N\alpha_N(c)), \qquad \alpha_N(c) := 1 - \frac{c}{N^{2/3}}$
= $\int_0^{N^{2/3}} N^{1/3} \mathbb{P}\Big(\widehat{W}_{[N-cN^{1/3}]} \ge s_N\Big) dc$
= $\int_0^{N^{2/3}} N^{1/3} \int_s^{+\infty} N^{-1/6} f_{\widehat{W}_{[N-cN^{1/3}]}}(\sqrt{2N} + 2^{-1/2}N^{-1/6}y) dy dc$

34

$$= \int_{(\mathbb{R}_+)^2} \mathbb{1}_{\left\{c \le N^{2/3}, y \ge s\right\}} N^{1/6} f_{\widehat{W}_{\left[N-cN^{1/3}\right]}}(\sqrt{2N} + 2^{-1/2}N^{-1/6}y) dy \, dc$$

We will now show that

$$N^{+1/6} f_{\widehat{W}_{\left[N-cN^{1/3}\right]}}(\sqrt{2N} + 2^{-1/2}N^{-1/6}y) \xrightarrow[N \to +\infty]{} \mathcal{Q}(y+c)^2$$

$$\tag{68}$$

This implies with the Wu-Xu-Zhao estimate (64) that $P_N^* = P_\infty^*(s) + O(N^{-1/3}P_\infty^{*,3})$ with

$$P_{\infty}^{*}(s) = \int_{s}^{+\infty} \int_{\mathbb{R}_{+}} \mathcal{Q}(y+c)^{2} dc \, dy = \int_{s}^{+\infty} \int_{y}^{+\infty} \mathcal{Q}(u)^{2} du \, dy = \int_{\mathbb{R}} \mathcal{Q}(u)^{2} (s-u)_{+} du$$
$$= \mathcal{Q}^{2} * \mathrm{id}_{+}(s)$$

and, with $\boldsymbol{\sigma}_1 = \sigma(\cdot; 1) \sim \sigma P_{II}(a=1)$ (or $\alpha = \frac{1}{2}$) defined in (59)

$$P_{\infty}^{*,3}(s) = \int_{s}^{+\infty} \int_{\mathbb{R}_{+}} \boldsymbol{\sigma}_{1}(y+c)\mathcal{Q}(y+c)^{2}dc\,dy = (\boldsymbol{\sigma}_{1}\mathcal{Q}^{2}) * \mathrm{id}_{+}(s)$$

Moreover,

$$\begin{split} P_N^{*,2} &:= \sum_{k=m+1}^n p_k^2 \\ &= \int_0^{N^{2/3}} N^{1/3} \mathbb{P} \big(W_{[N\alpha_N(c)]} \ge s_N \big)^2 dc, \qquad \alpha_N(c) := 1 - \frac{c}{N^{2/3}} \\ &= N^{-1/3} \int_0^{N^{2/3}} \Big(N^{1/3} \mathbb{P} \big(W_{[N\alpha_N(c)]} \ge s_N \big) \Big)^2 dc \\ &= O(N^{-1/3}) \times \int_{\mathbb{R}_+} \Big(\int_s^{+\infty} \mathcal{Q}(y+c)^2 dy \Big)^2 dc \\ &= O_s \Big(N^{-1/3} \Big) \end{split}$$

The fact that $\int_{s}^{+\infty} \sigma_{1}^{k} Q^{2} < \infty$ (with $k \in \{0, 1\}$) comes from the asymptotics at infinity

given in the second line of (59) (that implies⁷ $Q(x)^2 = O(x^{-2})$) and (63). The two terms $P_N^{*,2}$ and $P_N^{*,3}$, coming respectively from (64) and from the Poisson approximation give an unoptimal speed of convergence⁸, but are sufficient for our purpose.

We now prove (68). Define

$$g_N(M,y) := N^{1/3} f_{\widehat{W}_M}(\sqrt{2N} + 2^{-1/2}N^{-1/6}y), \qquad M := \left[N - cN^{1/3}\right]$$

⁷We will see in lemma 3.12 that we have in fact $\mathcal{Q} = q \sim P_{II}(0)$, hence that $\mathcal{Q}(x)^2 \sim \operatorname{Ai}(x)^2$ for $x \to +\infty$, which is much more refined than the sole difference of equivalents at infinity.

⁸The optimal speed of convergence is $O(N^{-2/3})$. It was obtained by Johnstone and Ma [78] using a very particular recentering $\mu_N \neq 2\sqrt{N}$ [78, (15)]. We are not concerned with such a problem.

Of course, the Wu-Xu-Zhao estimate (64) already gives the convergence for the right tuning of parameters, but we will first describe a more classical way of obtaining the result without the remainder, in the vein of the Forrester-Witte study [56].

Integrating at a certain x_N that will be choosen later and setting $x_N^* := \frac{x_N - 2\sqrt{N}}{N^{-1/6}}$, one has

$$g_N([N\alpha], y) = N^{1/3} g_N([N\alpha], x_N)$$
$$\exp\left(2\sqrt{2} \int_{x_N^*}^y N^{-1/6} \left(U_{[N\alpha]}(t; 1) - U_{[N\alpha]}(t; 0) - t\right)\Big|_{t=2\sqrt{N} + N^{-1/6}T} dT\right)$$

The function $y := U_{[N\alpha]}(\cdot; a)$ satisfies (56) with $n = [N\alpha]$. We thus consider

$$\sigma_N^{(\alpha)}(\cdot;a): t \mapsto \tau_N \big(U_{[N\alpha]}(\mu_N + \tau_N t;a) - a \times (\mu_N + \tau_N t) \big),$$
$$\mu_N := \sqrt{2N}, \quad \tau_N := \frac{1}{N^{1/6}\sqrt{2}}$$

Setting

$$Y(t) := \sigma_N^{(\alpha)}(\cdot; a)$$

$$Y'(t) := \frac{d}{dt} \sigma_N^{(\alpha)}(t; a) = \tau_N^2 \frac{d}{dt} U_{[N\alpha]}(\mu_N + \tau_N t; a) - a\tau_N$$

$$Y''(t) := \left(\frac{d}{dt}\right)^2 \sigma_N^{(\alpha)}(t; a) = \tau_N^3 \left(\frac{d}{dt}\right)^2 U_{[N\alpha]}(\mu_N + \tau_N t; a)$$

so that, dropping the indices N, the equation (56) becomes after multiplication by τ^6

$$0 = (Y'')^2 - 4\tau^6 \left((\mu + \tau t) \frac{Y' + a\tau}{\tau^2} - \frac{Y + a\mu + a\tau t}{\tau} \right)^2 + 4\tau^6 \frac{Y' + a\tau}{\tau^2} \left(\frac{Y' + a\tau}{\tau^2} - 2a \right) \left(\frac{Y' + a\tau}{\tau^2} + 2[N\alpha] \right)$$
$$= (Y'')^2 - 4\tau^2 \left((\mu + \tau t) Y' - \tau Y \right)^2 + 4(Y' + a\tau)(Y' + a\tau(1 - 2\tau)) \left(Y' + a\tau + 2[N\alpha] \tau^2 \right)$$
An expansion that supposes Y and its derivatives to be of order 1 shows that

$$\mathcal{L}_{N}(Y) = (Y'')^{2} + 4(Y')^{3} - 4((Y')^{2}t - YY') - a^{2} + 4(Y')^{2}(2[N\alpha]\tau^{2} - \mu^{2}\tau^{2}) + O_{Y,t}(\tau)$$

Since $\mu^2 = 2N$, $\mathcal{L}_N(Y)$ converges iff

$$N\tau^2(1-\alpha) = N^{2/3}(1-\alpha) \xrightarrow[N \to +\infty]{} c = O(1) \qquad \Longleftrightarrow \qquad \alpha \equiv \alpha_N = 1 - \frac{c + o(1)}{N^{2/3}}$$

In such a case, $\sigma_N^{(\alpha)}(\cdot; a)$ converges locally uniformly to $\sigma^{(c)}(\cdot; a) = \sigma(\cdot + c; a)$ satisfying the following shifted σ -form of the Painlevé II equation (as setting $t \leftarrow t + c$ gives back the original equation) with the same boundary condition as (59)

$$(Y'')^{2} + 4Y'((Y')^{2} - (t - c)Y' + Y) - a^{2} = 0$$

As a result, locally uniformly

$$\delta\sigma_N^{(\alpha)} := \sigma_N^{(\alpha)}(\cdot;1) - \sigma_N^{(\alpha)}(\cdot;0) \xrightarrow[N \to +\infty]{} \delta\sigma^{(c)} =: \delta\sigma(\cdot+c) = \sigma(\cdot+c;1) - \sigma(\cdot+c;0)$$

This analysis explains the relevance of the rescaling, but does not give any speed of convergence. Such a speed is required to prove the result in a probabilistic way (since the convergence of the densities does not preclude from the usual weak-* pathology of "escape of the mass at infinity"); it is furnished by the Wu-Xu-Zhao estimate (64) obtained with a Riemann-Hilbert problem.

It remains to choose x_N . We set it as the solution c to the equation

$$N^{1/6} f_{\widehat{W}_N}(c_N) e^{2\sqrt{2}N^{-1/3}\sigma_1(c)} = 1, \qquad c_N := \sqrt{2N} + N^{-\frac{1}{6}} \frac{c}{\sqrt{2}}$$

i.e.

$$N^{1/6} \mathcal{H}_N(x)^2 e^{-x^2} e^{2\sqrt{2}N^{-1/3}\sigma_1(N^{1/6}(x-\sqrt{2N}))} = \sqrt{\pi}$$

$$\iff x^2 - 2\ln\mathcal{H}_N(x) + 2\sqrt{2}N^{-1/3}\sigma_1(N^{1/6}(x-\sqrt{2N})) = \frac{\ln(N)}{6} + O(1)$$

We are just concerned with the behaviour of x_N when $N \to +\infty$ and not its unicity. As a result, if there are several such values, we choose the largest one; nevertheless, for N and x big enough $(x \gg 2\ln(N))$, there is a unique such solution, as seen using the equivalent $\mathcal{H}_N(x) \sim x^N$ and $\sigma_1(x) \sim -\sqrt{x}$ given in (63) when $x \to +\infty$.

Taking the logarithm of the equation and replacing $\mathcal{H}_N(x)$ by x^N and $\sigma_1(s)$ by $-\sqrt{s}$ yields the approximate equation

$$x_N^2 - 2N\ln(x_N) - 2\sqrt{2}N^{-1/4}\sqrt{x_N - \sqrt{2N}} = \frac{\ln(N)}{6} + O(1)$$

Since $x_N - \sqrt{2N} \ge 0$, one can neglect the term $N^{-1/4}\sqrt{x_N - \sqrt{2N}}$ for the first order of the equation. Setting $x_N^2 = Nt_N$, the equation is equivalent to $t_N - \ln(t_N) = N \ln(N) + \frac{\ln(N)}{6} + O(1)$ whose solution is given by means of the Lambert W function [105, § 4.13 p. 111], i.e. $t_N = W(N \ln(N) + \frac{\ln(N)}{6} + O(1))$. The asymptotic of this function at infinity is $W(x) = \ln(x) - \ln \ln(x) + o(1)$ [105, (4.13.10) p. 111], hence $t_N = \ln(N) + O(\ln \ln(N))$, resulting in

$$x_N = \sqrt{N\ln(N)} + O(\sqrt{N}\ln\ln(N))$$

Of course, as long as $\alpha_N = O(1)$, this behaviour is valid for $x_{[N\alpha_N]}$. In the case where α_N is close to 0, we cut the integral up to $2N^{2/3} - \varepsilon$ and choose $\varepsilon \ll N^{-1/6}$ so that $x_{[N\alpha_N]} \gg \sqrt{2N}$; the remaining integral on $(2N^{2/3} - \varepsilon, 2N^{2/3})$ is bounded by the sup of the probability (which is 1) and $\varepsilon \times N^{1/6} = o(1)$. The final result is thus

$$x_N \gg \sqrt{N} \implies x_N^* - \sqrt{2N} \to +\infty$$

This concludes the proof.

37

3.4.5. Comparison of expressions. The expression (66), namely

$$\mathbb{P}(\mathcal{TW}_2 \leqslant s) = \exp\left(-\mathcal{Q}^2 * \mathrm{id}_+(s)\right)$$

can be compared with the expression (3) that reads, with q given in (4):

$$\mathbb{P}(\mathcal{TW}_2 \leqslant s) = \exp\left(-q^2 * \mathrm{id}_+(s)\right)$$

Taking the log, differentiating twice in s and taking the square root as both functions are positive (see e.g. [105, 32.3.6 p. 727] for the positivity of q) gives the identity

q = Q

Identities between different $\boldsymbol{\sigma}_a := \boldsymbol{\sigma}(\cdot; a)$ were already noticed, for instance [56, prop. 27] gives $\boldsymbol{\sigma}_2 = \frac{\boldsymbol{\sigma}_0'}{\boldsymbol{\sigma}_0} + \boldsymbol{\sigma}_0$ and [56, (5.28)/(2.7) & (5.27)] gives $\boldsymbol{\sigma}_0' = -\boldsymbol{q}_0^2$ (see also [59, § 2.4]). This is thus natural to try to prove that $\boldsymbol{q} = \boldsymbol{\mathcal{Q}}$ with the theory furnished in [56].

Lemma 3.12 (Direct proof that q = Q). Define $\sigma_a := \sigma(\cdot; a)$ solution of (59), $q_\alpha := q(\cdot; \alpha)$ solution of (4) Then, q = Q, or, equivalently (by logarithmic differentiation and value 0 at infinity) $\frac{q'_0}{q_0} = \sigma_1 - \sigma_0$ (69)

First proof. The link between \boldsymbol{q}_{α} and $\boldsymbol{\sigma}_{a}$ is given by [56, (5.9)/(5.26)] :

$$\alpha = a - \frac{1}{2},$$

$$-2^{-1/3}\boldsymbol{\sigma}_a(-2^{-1/3}t) = \frac{1}{2}\boldsymbol{q}'_{a-1/2}(t)^2 - \frac{1}{2}\left(\boldsymbol{q}_{a-1/2}(t)^2 + \frac{t}{2}\right)^2 - a\boldsymbol{q}_{a-1/2}(t)$$

As a result, setting $\hat{t} := -2^{-1/3}t$ and specialising $a \in \{0, 1\}$ yields

$$\begin{cases} -2^{-1/3}\boldsymbol{\sigma}_{0}(\hat{t}) = \frac{1}{2}\boldsymbol{q}_{-1/2}'(t)^{2} - \frac{1}{2}\left(\boldsymbol{q}_{-1/2}(t)^{2} + \frac{t}{2}\right)^{2} \\ -2^{-1/3}\boldsymbol{\sigma}_{1}(\hat{t}) = \frac{1}{2}\boldsymbol{q}_{1/2}'(t)^{2} - \frac{1}{2}\left(\boldsymbol{q}_{1/2}(t)^{2} + \frac{t}{2}\right)^{2} - \boldsymbol{q}_{1/2}(t) \end{cases}$$

There are moreover general identities between q_0 and $q_{\pm 1/2}$ given in⁹ [56, (3.7)] :

$$\varepsilon \in \{\pm 1\}, \qquad \begin{cases} -\boldsymbol{q}_0(\hat{t})^2 = \varepsilon 2^{-1/3} \left(\boldsymbol{q}'_{\varepsilon/2}(t) - \varepsilon \boldsymbol{q}_{\varepsilon/2}(t)^2 - \frac{\varepsilon}{2} t \right) \\ \frac{\boldsymbol{q}'_0(\hat{t})}{\boldsymbol{q}_0(\hat{t})} = \varepsilon 2^{1/3} \boldsymbol{q}_{\varepsilon/2}(t) \end{cases}$$
(70)

This implies

$$2^{-1/3}(\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_0)(\hat{t}) = \boldsymbol{q}_{1/2}(t) + R$$

⁹The second equality in (70) is called the *inverse Gambier identity* in [59, (2.48)].

with

$$\begin{split} 2R &:= \boldsymbol{q}_{-1/2}'(t)^2 - \left(\boldsymbol{q}_{-1/2}(t)^2 + \frac{t}{2}\right)^2 - \left(\boldsymbol{q}_{1/2}'(t)^2 - \left(\boldsymbol{q}_{1/2}(t)^2 + \frac{t}{2}\right)^2\right) \\ &= \left(\boldsymbol{q}_{-1/2}'(t) - \boldsymbol{q}_{-1/2}(t)^2 - \frac{t}{2}\right) \left(\boldsymbol{q}_{-1/2}'(t) + \boldsymbol{q}_{-1/2}(t)^2 + \frac{t}{2}\right) \\ &- \left(\boldsymbol{q}_{1/2}'(t) - \boldsymbol{q}_{1/2}(t)^2 - \frac{t}{2}\right) \left(\boldsymbol{q}_{1/2}'(t) + \boldsymbol{q}_{1/2}(t)^2 + \frac{t}{2}\right) \\ &= \left(\boldsymbol{q}_{-1/2}'(t) - \boldsymbol{q}_{-1/2}(t)^2 - \frac{t}{2}\right) \boldsymbol{q}_0(\hat{t})^2 2^{1/3} \\ &+ \boldsymbol{q}_0(\hat{t})^2 2^{1/3} \left(\boldsymbol{q}_{1/2}'(t) + \boldsymbol{q}_{1/2}(t)^2 + \frac{t}{2}\right) \qquad \text{using (70)} \\ &= \boldsymbol{q}_0(\hat{t})^2 2^{1/3} \left(\boldsymbol{q}_{-1/2}'(t) - \boldsymbol{q}_{-1/2}(t)^2 + \boldsymbol{q}_{1/2}'(t) + \boldsymbol{q}_{1/2}(t)^2\right) \end{split}$$

The second equality in (70) gives $-\boldsymbol{q}_{-1/2}(t) = \boldsymbol{q}_{1/2}(t)$, which implies $\boldsymbol{q}'_{-1/2}(t) + \boldsymbol{q}'_{1/2}(t) = 0$ and $-\boldsymbol{q}_{-1/2}(t)^2 + \boldsymbol{q}_{1/2}(t)^2 = 0$, i.e. R = 0. The second equality in (70) then yields

$$2^{-1/3}(\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_0)(\hat{t}) = \boldsymbol{q}_{1/2}(t) = 2^{-1/3} \frac{\boldsymbol{q}_0'(t)}{\boldsymbol{q}_0(\hat{t})}$$

which concludes the proof.

Second proof. Using [56, (5.9), (5.23), prop. 19 & (3.23)] that read

$$\begin{split} H[n] &:= H(t)|_{\alpha_1 = n} \\ \boldsymbol{\sigma}_a(t) &= -2^{1/3} H(\widetilde{t})_{\alpha_1 = a}, \qquad \widetilde{t} := -2^{+1/3} t \\ q[n] &:= \boldsymbol{q}_{n + \alpha_1 + 1/2} \\ H[n+1] - H[n] &= q[n] \qquad \text{(Toda equation)} \end{split}$$

one obtains [59, (3.11)] for a = 0, namely

$$\boldsymbol{\sigma}_{n+1}(t) - \boldsymbol{\sigma}_n(t) = -2^{1/3} \boldsymbol{q}_{n+1/2}(\widetilde{t})$$

The second equality in (70) gives then the result for n = 0.

3.4.6. Ultimate remarks.

Remark 3.13. We mention the problem of large deviations for $\lambda_{1,N}$ which is studied e.g. in [35, 88]. With theorem 3.1, this problem becomes a problem of large deviations for the maximum of independent random variables, hence a problem of large deviations for sums of parametric indicators. The Poisson approximation with remainder is still available in this regime ; an alternative is provided by the renormalisation flow approach developed in the context of maxima of independent random variables in [5, 12] and which is now directly applicable to the max-independent structure of $\lambda_{1,N}$.

Remark 3.14. The analogue of the proof of theorem 1.2 would consist in using a decomposition of the type

$$\mathbb{P}(\boldsymbol{\lambda}_{1,N} \leqslant s) = \exp\left(-\sum_{k \ge 1} \int_{s}^{+\infty} Q_{k}^{(N)}(x) dx\right) = \prod_{k \ge 1} \mathbb{P}(Z_{k}(GUE_{N}) \leqslant s)$$

i.e. an equality in law with an infinite sequence of independent random variables depending on N. Mimicking the proof of theorem 1.2 could then lead directly to the convergence in law of $\frac{Z_k(GUE_N)-2\sqrt{N}}{N^{-1/6}}$ towards $Z_k(\text{Ai})$ or some \mathcal{W}_k , asking then the following problem :

Question 3.15. Can one obtain Tracy and Widom's convergence in law (58) writing

$$\frac{\boldsymbol{\lambda}_{1,N} - 2\sqrt{N}}{N^{-1/6}} \stackrel{\mathcal{L}}{=} \max_{k \ge 1} \left\{ \frac{Z_k(GUE_N) - 2\sqrt{N}}{N^{-1/6}} \right\} \xrightarrow[N \to +\infty]{} \max_{k \ge 0} \mathcal{W}_k$$

for infinite sequences of random variables $(Z_k(GUE_N))_{k\geq 1}$ and $(\mathcal{W}_k)_{k\geq 1}$?

The conjectural random variables \mathcal{W}_k obtained at the limit might differ from the $Z_k(\text{Ai})$ defined in (13) or the $Z'_k(\text{Ai})$ defined in (19) as several decompositions are a priori possible (in addition, no commuting operator for the *GUE* kernel operator acting on $L^2([s, +\infty), \mu)$ or its translated version has been found so far). We hope to come back to this problem in the future.

4. MAX-INDEPENDENCE STRUCTURE IN THE SYMMETRIC SCHUR MEASURE

The relevant notations for this section are given in Annex A.

4.1. Motivations. The Schur measure was introduced by Okounkov [99, 101] following a study of the z-measure [100] introduced by Borodin and Olshanski [32, 33] to generalise the Plancherel measure ; this last measure saw a regain of activity in relation with the Ulam problem on the longuest increasing subsequence of a random uniform permutation by Baik, Deift and Johansson [6]. Such generalisations were designed to highlight the general mechanism explaining the apparition of a determinantal structure in the Poisson-Plancherel measure and we refer to the surveys [75, 76, 104] for a zoo of models that fit into this framework. The general explanation is as follows: the Schur measure of parameters $(\mathcal{A}, \mathcal{B})$ is a measure on partitions $\mathbb{Y}_{\infty} := \bigcup_{n \geq 0} \mathbb{Y}_n$ defined by

$$\mathbb{P}_{\mathcal{A},\mathcal{B}}(\lambda) := H[-\mathcal{A}\mathcal{B}]s_{\lambda}[\mathcal{A}]s_{\lambda}[\mathcal{B}]$$
(71)

The specialisation $s_{\lambda}[\mathcal{A}]$ is written $\varphi(s_{\lambda})$ in [104] and should be understood as an "abstract" algebra morphism of Λ where the generators $(h_k)_k$, $(e_k)_k$ or $(p_k)_k$ have been attributed a particular value, the value of the Schur function resulting then from an algebraic formula expressing it with the generators, for instance the Jacobi-Trudi formula (106). This measure is a probability measure if the numbers such defined are real, for instance if $\mathcal{A} = \overline{\mathcal{B}} \subset \mathbb{C}$, or if \mathcal{A} and \mathcal{B} are positive specialisations of the Schur functions (see Annex A). The main result of interest is the determinantal character (with explicit kernel) of the shifted point process $(\lambda_k - k + a)_{k \ge 1}$ (with $a \in \{0, 1, \frac{1}{2}\}$ depending on the conventions), λ being the random partition distributed according to $\mathbb{P}_{\mathcal{A},\mathcal{B}}$. So far, seven proofs of this determinantal nature have been given and we refer to them for this fact: fermionic Gaussian "integral" [99], Riemann-Hilbert problems [22], Jacobi-Trudi formula [107], ratio of alternants formula [75], Giambelli formula and *L*-ensembles [29], coupling on Gelfand-Tsetlin graph [55] and Macdonald operator [1]. See also [30] for a discussion on the kernel.

The success of this generalisation prefigures the numerous extensions of the framework in many different directions, for instance the dynamical version of "Schur processes" [102, 103] and, in fine, the creation of the theory of integrable probability [26, 34].

Remark 4.1. We adopt here the convention of e.g. Johansson [75] that writes this measure on partitions over \mathbb{Z} , as opposed to the Okounkov convention [99, 101] (the half-infinite wedge space/fermionic Fock space has a basis indexed by Maya diagrams written on $\mathbb{Z} + \frac{1}{2}$ by convention). Such conventions are of course equivalent, but the formuli differ by some square roots ; formuli are given here for the process $(\lambda_k - k)_{k \ge 1}$ and not $(\lambda_k - k + \frac{1}{2})_{k \ge 1}$.

4.2. The Poisson-Plancherel measure. The Plancherel measure on \mathbb{Y}_n is defined by [31, 74]

$$\mathbb{P}_n(\lambda) := \frac{d_\lambda^2}{n!} \mathbb{1}_{\{\lambda \vdash n\}}$$
(72)

where d_{λ} is the dimension of the irreducible module of \mathfrak{S}_n indexed by λ (see e.g. [83, I-6]). This measure can be understood as the conditioning of a general measure \mathbb{P}_{∞} on \mathbb{Y}_{∞} by the event $\{\boldsymbol{\lambda} \vdash n\}$ (where $\boldsymbol{\lambda} \sim \mathbb{P}_{\infty}$), and the law of $|\boldsymbol{\lambda}|$ gives then the choice of the relevant randomisation for $\boldsymbol{\lambda}_n \sim \mathbb{P}_n$ to obtain a determinantal structure. Since $d_{\lambda} = n! s_{\lambda}[\mathcal{E}]$ by (105), setting $|\boldsymbol{\lambda}| \sim \operatorname{Po}(\xi)$ gives the Schur measure of parameters $\mathcal{A} = \mathcal{B} = \sqrt{\xi} \mathcal{E}$. We thus define the Poisson-Plancherel measure on \mathbb{Y}_{∞} by

$$\mathbb{P}_{\text{PoPl}(\xi)}(\lambda) := e^{-\xi} \xi^{|\lambda|} \left(\frac{d_{\lambda}}{|\lambda|!}\right)^2 \tag{73}$$

One then has

$$\widetilde{\boldsymbol{\lambda}} := (\boldsymbol{\lambda}_k - k)_{k \ge 1} \sim \mathrm{DPP}(\mathcal{K}_{\mathrm{PoPl}(\xi)})$$

implying in particular

$$\mathbb{P}(\boldsymbol{\lambda}_{1}^{(\xi)} < s) = \det(I - \mathcal{K}_{\operatorname{PoPl}(\xi)})_{\ell^{2}(s+\mathbb{N})} = \det(I - \mathcal{K}_{\operatorname{PoPl}(\xi),s})_{\ell^{2}(\mathbb{N})}$$

where $\mathcal{K}_{\text{PoPl}(\xi),s}(x,y) = \mathcal{K}_{\text{PoPl}(\xi)}(x+s,y+s).$

The corresponding specialisation of the Schur kernel $K_{\text{PoPl}(\xi)} : \mathbb{Z}^2 \to \mathbb{R}_+$ in this case is the *discrete Bessel kernel* [31, 74] that has several forms :

• the Hankel convolutive form is [74, (3.31)]

$$\mathcal{K}_{\text{PoPl}(\xi)}(x,y) := \sum_{k \ge 1} B_k^{(\xi)}(x) B_k^{(\xi)}(y), \qquad B_k^{(\xi)}(x) := J_{x+k}(2\sqrt{\xi})$$
(74)

where J_k is the k-th Bessel function of the first kind defined by :

$$J_k(x) := [z^k] e^{x(z-z^{-1})/2}$$
(75)

• the Christoffel-Darboux form is ([101, (23)], [74, (1.10)])

$$\mathcal{K}_{\text{PoPl}(\xi)}(x,y) = \sqrt{\xi} \frac{B_1^{(\xi)}(x) B_0^{(\xi)}(y) - B_0^{(\xi)}(x) B_1^{(\xi)}(y)}{x - y} \mathbb{1}_{\{x \neq y\}} + \sqrt{\xi} \dot{J}_x(2\sqrt{\xi}) \mathbb{1}_{\{x = y\}}$$
(76)

with $\dot{J}_x(t) := \frac{\partial}{\partial \alpha} J_\alpha(t) \Big|_{\alpha=x}$ (see e.g. [105, 10.2.4])

• and the *integral form* is [101, (22)]

$$\mathcal{K}_{\text{PoPl}(\xi)}(x,y) = \oint_{\mathbb{U} \times \rho \mathbb{U}} \frac{z^{-x} \omega^{-y}}{1 - z\omega} e^{\sqrt{\xi}(z - z^{-1} + \omega - \omega^{-1})} \frac{d^* z}{z} \frac{d^* \omega}{\omega}, \qquad \rho < 1$$

$$= \oint_{\mathbb{U} \times R \mathbb{U}} \frac{z^{-x} \omega^{+y}}{1 - z\omega^{-1}} e^{\sqrt{\xi}(z - z^{-1} - (\omega - \omega^{-1}))} \frac{d^* z}{z} \frac{d^* \omega}{\omega}, \qquad R > 1$$
(77)

Using the expansion $\frac{1}{1-z\omega} = \sum_{k \ge 0} (z\omega)^k$ for $z \in \mathbb{U}$ and $\omega \in \rho \mathbb{U}$ with $\rho < 1$, one obtains (74) from (77) and (75).

The Hankel convolutive form (74) is in direct analogy with the Hankel convolutive form (2) of the Airy kernel : $\mathcal{K}_{\text{PoPl}(\xi),s}$ is the square of the (discrete) Hankel convolution operator

$$\mathcal{H}(B_s^{(\xi)}): f \in \ell^2(\mathbb{N}) \mapsto \sum_{k \ge 0} B_s^{(\xi)}(\cdot + k) f(k) \in \ell^2(\mathbb{N})$$

namely

$$\mathcal{K}_{\text{PoPl}(\xi),s} = \mathcal{H}(B_s^{(\xi)})^2 \tag{78}$$

which is analogous to (15) on $L^2(\mathbb{R}_+)$.

One could hope to extend the proof of theorem 1.2 using a discrete analogue of det $(I - \mathbf{K}_t) = \exp\left(-\int_t^{+\infty} \operatorname{tr}((I - \mathbf{K}_s)^{-1}\dot{\mathbf{K}}_s)ds\right)$. Such a formula reads

$$\det(I - \mathcal{K}_N)_{\ell^2(\mathbb{N})} = \exp\left(-\sum_{\ell \ge N} \operatorname{tr}\left(\log(I - \mathcal{K}_{\ell+1}) - \log(I - \mathcal{K}_{\ell})\right)_{\ell^2(\mathbb{N})}\right)$$
(79)

To simplify it, one needs the theory of Orthogonal Polynomials on the Unit Circle (OPUC).

42

4.3. **OPUCs.** The main reference for this § is [115] from which we will only present the properties that are required in the sequel.

A family $(\Phi_k)_{k\geq 0}$ of monic polynomials (i.e. $\Phi_n(z) = z^n + \cdots$) is orthogonal in $L^2(\mathbb{U}, \mu)$ for a probability measure μ on \mathbb{U} if

$$\langle \Phi_k, \Phi_\ell \rangle_{L^2(\mu)} := \oint_{\mathbb{U}} \Phi_k \overline{\Phi_\ell} \, d\mu = \|\Phi_k\|_{L^2(\mu)}^2 \, \delta_{k,\ell}$$

Defining the anti- $L^2(\mu)$ -unitary map *,n by $P^{*,n}(z) := z^n \overline{P(\overline{z}^{-1})}$ and dropping the *n* by convention when the polynomial is of degree *n*, one can show that [115, (2.17), (2.18)]

$$\Phi_{n+1}(z) = z\Phi_n(z) - \overline{\alpha_n}\Phi_n^*(z), \qquad \alpha_n := -\overline{\Phi_{n+1}(0)}$$

The coefficient $(\alpha_k)_{k\geq 0}$ are called the *Verblunsky coefficients* and the previous recursion is called the *Szegö recursion*. One can easily prove that [115, (2.19)]

$$\|\Phi_{n+1}\|_{L^{2}(\mu)}^{2} = \left(1 - |\alpha_{n}|^{2}\right) \|\Phi_{n}\|_{L^{2}(\mu)}^{2} \implies \|\Phi_{n}\|_{L^{2}(\mu)}^{2} = \prod_{k=0}^{n-1} \left(1 - |\alpha_{k}|^{2}\right)$$
(80)

and that for a non trivial measure μ (i.e. not supported on a finite set of points) $|\alpha_k| < 1$ [115, thm. 2.1]. As a result, one has

$$\|\Phi_{n+1}\|_{L^{2}(\mu)} \leqslant \|\Phi_{n}\|_{L^{2}(\mu)} \tag{81}$$

For a probability measure μ , one defines the Toeplitz determinant $D_n(\mu)$ as

$$D_n(\mu) := \det\left(\oint_{\mathbb{U}} z^{i-j} d\mu(z)\right)_{0 \le i, j \le n}$$

Hence, using linear combinations of lines and columns one gets

$$D_n(\mu) := \det\left(\oint_{\mathbb{U}} \Phi_i(z)\overline{\Phi_j(z)}d\mu(z)\right)_{0 \le i,j \le n} = \prod_{j=0}^n \|\Phi_j\|_{L^2(\mu)}^2 = \prod_{j=0}^{n-1} (1 - |\alpha_j|^2)^{n-j}$$

This implies the Szegö theorem [115, (8.2)]

$$F(\mu) := \prod_{j \ge 0} (1 - |\alpha_j|^2) = \lim_{n \to +\infty} D_n(\mu)^{1/n} = \lim_{n \to +\infty} \frac{D_{n+1}(\mu)}{D_n(\mu)} = \lim_{n \to +\infty} \|\Phi_n\|_{L^2(\mu)}^2$$
(82)

Note that $F(\mu)$ can be equal to 0, in which case $(\alpha_k)_k \notin \ell^2(\mathbb{N})$ (see [115, thm. 2.2] for a consequence).

4.4. Independence structure in the Poisson-Plancherel measure. The Cauchy formula for the Cauchy kernel (101) in abstract alphabets \mathcal{A}, \mathcal{B} and the (restricted) orthogonality of the Schur functions (108) gives

$$\int_{\mathcal{U}_N} H\left[\widehat{\omega}(\mathcal{A}U + \mathcal{B}U^{-1})\right] dU = \int_{\mathcal{U}_N} \sum_{\lambda,\mu} s_{\lambda}[\widehat{\omega}\mathcal{A}] s_{\lambda}(U) s_{\mu}[\widehat{\omega}\mathcal{B}] s_{\mu}(U^{-1}) dU$$
$$= \sum_{\ell(\lambda) \leqslant N} s_{\lambda}[\widehat{\omega}\mathcal{A}] s_{\lambda}[\widehat{\omega}\mathcal{B}]$$
$$= \sum_{\lambda_1 \leqslant N} s_{\lambda}[\mathcal{A}] s_{\lambda}[\mathcal{B}]$$

As a result,

$$\mathbb{P}_{\mathrm{Schur}(\mathcal{A},\mathcal{B})}(\boldsymbol{\lambda}_{1} \leqslant N) = H[-\mathcal{A}\mathcal{B}] \int_{\mathcal{U}_{N}} H\left[\widehat{\omega}(\mathcal{A}U + \mathcal{B}U^{-1})\right] dU$$
(83)

Using Weyl integration formula (52) and the Andréieff-Heine-Szegö formula (43), the random unitary matrix integral in the RHS can be transformed into the Toeplitz determinant of symbol $z \mapsto H[\widehat{\omega}(z\mathcal{A}+z^{-1}\mathcal{B})]$ (see remark 3.6). The determinantal character of λ_1 allows moreover to write the LHS as a Fredholm determinant on $\ell^2(N + \mathbb{N}^*)$. The formula thus obtained is the celebrated Borodin-Okounkov formula [30] that was also discovered by Geronimo and Case the context of the inverse scattering method [39] and that meanwhile had several other proofs and generalisations [7, 10, 16, 17, 18].

Let us only consider the Plancherel case $\mathcal{A} = \mathcal{B} = \sqrt{\xi}\mathcal{E}$. One has $H[-\mathcal{A}\mathcal{B}] = H[-\xi\mathcal{E}] =$ $e^{-\xi}$, and $H[\widehat{\omega}\mathcal{A}U] = e^{p_1\left[\sqrt{\xi}U\right]} = e^{\sqrt{\xi}\operatorname{tr}(U)}$, hence

$$\mathbb{P}_{\text{PoPl}(\xi)}(\boldsymbol{\lambda}_1 \leqslant N) = e^{-\xi} \int_{\mathcal{U}_N} e^{\sqrt{\xi} \operatorname{tr}(U+U^{-1})} dU = e^{-\xi} \det \mathbb{T}_N(\mathcal{I}_{\xi}), \qquad \mathcal{I}_{\xi}(z) := e^{\sqrt{\xi}(z+z^{-1})}$$

Define the probability measure on \mathbb{U}

$$\mu_{\xi}(d\theta) := \frac{e^{\sqrt{\xi}(z+z^{-1})}}{I_0(\sqrt{\xi})} \frac{d^*z}{z}, \qquad I_0(\sqrt{\xi}) := [z^0] e^{\sqrt{\xi}(z+z^{-1})}$$

where the normalisation $I_0(\sqrt{\xi})$ is the 0-th modified Bessel function defined by

$$I_k(s) := \left[z^k\right] e^{s(z+z^{-1})}$$

We also define :

- (Φ^(ξ)_{k≥0} as the OPUCs for L²(U, μ_ξ),
 (α_k(ξ))_{k≥0} as their Verblunsky coefficients.

Then, writing the Toeplitz determinant as a product of norms of OPUCs gives

$$\mathbb{P}_{\text{PoPl}(\xi)}(\boldsymbol{\lambda}_1 \leqslant N) = e^{-\xi} \prod_{k=0}^{N-1} \oint_{\mathbb{U}} \left| \Phi_k^{(\xi)}(z) \right|^2 e^{\sqrt{\xi}(z+z^{-1})} \frac{d^*z}{z}$$

$$= e^{-\xi} \prod_{k=0}^{N-1} I_0(\sqrt{\xi}) \left\| \Phi_k^{(\xi)} \right\|_{L^2(\mathbb{U},\mu_{\xi})}^2$$
$$= e^{-\xi} \prod_{k=0}^{N-1} I_0(\sqrt{\xi}) \prod_{j=0}^{k-1} (1 - |\alpha_j(\xi)|^2)$$

Set

$$\pi_{k}(\xi) := I_{0}(\sqrt{\xi}) \left\| \Phi_{k}^{(\xi)} \right\|_{L^{2}(\mathbb{U},\mu_{\xi})}^{2} = \oint_{\mathbb{U}} \left| \Phi_{k}^{(\xi)}(z) \right|^{2} e^{\sqrt{\xi}(z+z^{-1})} \frac{d^{*}z}{z}$$

$$q_{k}(\xi) := \frac{1}{\pi_{k}(\xi)}$$
(84)

Letting $N \to +\infty$ gives the Szegö theorem (82)

$$1 = e^{-\xi} \prod_{k \ge 0} \pi_k(\xi)$$

Note that when the measure μ is not a probability measure, one needs to take into account the normalisation factor and replace the norms by the π_k s. In this setting, the Szegö theorem/the absolute convergence of the product gives

$$\pi_k(\xi) \xrightarrow[k \to +\infty]{} 1 \qquad \Longleftrightarrow \qquad \left\| \Phi_k^{(\xi)} \right\|_{L^2(\mathbb{U},\mu_{\xi})}^2 \xrightarrow[k \to +\infty]{} I_0(\sqrt{\xi})^{-1} = \prod_{j \ge 0} (1 - |\alpha_j(\xi)|^2)$$

As a consequence of the previous computations,

$$\mathbb{P}_{\text{PoPl}(\xi)}(\boldsymbol{\lambda}_1 \leqslant N) = \prod_{k \ge N} \pi_k(\xi)^{-1} =: \prod_{k \ge N} q_k(\xi) = \prod_{\ell \ge 0} q_{\ell+N}(\xi)$$

We are now ready to state the

Theorem 4.2 $(\lambda_1^{\text{PoPl}(\xi)})$ is a maximum of a sequence of independent random variables). There exist i.i.d. non negative random variables $(Z_k^{\text{PoPl}(\xi)})_{k \ge 0}$ such that

$$\boldsymbol{\lambda}_{1}^{\text{PoPl}(\xi)} \stackrel{\mathcal{L}}{=} \max_{k \ge 0} \left\{ Z_{k}^{\text{PoPl}(\xi)} - k \right\}$$
(85)

Moreover

$$Z_1^{\text{PoPl}(\xi)} \stackrel{\mathcal{L}}{=} \inf\{j \ge 0 \,/\, \forall \, m \ge j, \ B_j(\xi) = 0\}$$
(86)

where the $(B_j(\xi))_{j\geq 0}$ are independent $\{0,1\}$ -Bernoulli random variables of expectation $|\alpha_j(\xi)|^2$.

Proof. One thus has

$$\mathbb{P}_{\text{PoPl}(\xi)}(\boldsymbol{\lambda}_1 \leqslant N) = \prod_{\ell \geqslant 0} q_{\ell+N}(\xi)$$

with

$$0 \leqslant q_{\ell}(\xi) = \frac{1}{I_0(\sqrt{\xi})} \left\| \Phi_{\ell}^{(\xi)} \right\|_{L^2(\mathbb{U},\mu_{\xi})}^{-2} \xrightarrow[\ell \to +\infty]{} 1$$

Moreover, since norms of OPUCs are decreasing by (81), $(q_{\ell}(\xi))_{\ell}$ is decreasing. This can also be seen using (80), noticing that $1 - |\alpha_k(\xi)|^2 < 1$ and writing

$$q_{\ell}(\xi) = \frac{1}{I_0(\sqrt{\xi})} \left\| \Phi_{\ell}^{(\xi)} \right\|_{L^2(\mathbb{U},\mu_{\xi})}^{-2} = \prod_{k \ge 0} (1 - |\alpha_k(\xi)|^2) \times \prod_{j=0}^{\ell-1} \frac{1}{1 - |\alpha_j(\xi)|^2}$$
$$= \prod_{k \ge \ell} (1 - |\alpha_k(\xi)|^2)$$

As a result, there exists a random variable $Z^{\operatorname{PoPl}(\xi)} \in \mathbb{N}$ such that

$$q_{\ell}(\xi) = \mathbb{P}\Big(Z^{\operatorname{PoPl}(\xi)} \leqslant \ell\Big)$$

and in particular, with an i.i.d. sequence $(Z_{\ell}^{\operatorname{PoPl}(\xi)})_{\ell}$ of such random variables

$$\begin{split} \mathbb{P}_{\text{PoPl}(\xi)}(\boldsymbol{\lambda}_{1} \leqslant N) &= \prod_{\ell \geqslant 0} \mathbb{P}\left(Z_{\ell}^{\text{PoPl}(\xi)} \leqslant \ell + N\right) = \prod_{\ell \geqslant 0} \mathbb{P}\left(Z_{\ell}^{\text{PoPl}(\xi)} - \ell \leqslant N\right) \\ &= \mathbb{P}\left(\max_{\ell \geqslant 0} \left\{Z_{\ell}^{\text{PoPl}(\xi)} - \ell\right\} \leqslant N\right) \end{split}$$

Last, (80) yields

$$\mathbb{P}\left(Z_{1}^{\text{PoPl}(\xi)} = \ell\right) = q_{\ell}(\xi) - q_{\ell-1}(\xi) = \frac{1}{I_{0}(\sqrt{\xi})} \left(\frac{1}{\left\|\Phi_{\ell}^{(\xi)}\right\|^{2}} - \frac{1}{\left\|\Phi_{\ell-1}^{(\xi)}\right\|^{2}}\right)$$
$$= \frac{1}{I_{0}(\sqrt{\xi})} \times \frac{1}{\left\|\Phi_{\ell-1}^{(\xi)}\right\|^{2}} \times \left(\frac{1}{1 - |\alpha_{\ell-1}(\xi)|^{2}} - 1\right)$$
$$= \prod_{k \ge 0} (1 - |\alpha_{k}(\xi)|^{2}) \times \prod_{j=0}^{\ell-2} \frac{1}{1 - |\alpha_{j}(\xi)|^{2}} \times \frac{|\alpha_{\ell-1}(\xi)|^{2}}{1 - |\alpha_{\ell-1}(\xi)|^{2}}$$
$$=: |\alpha_{\ell-1}(\xi)|^{2} \prod_{k \ge \ell} (1 - |\alpha_{k}(\xi)|^{2})$$
$$= \mathbb{P}(B_{\ell-1}(\xi) = 1) \prod_{k \ge \ell} \mathbb{P}(B_{k}(\xi) = 0)$$
$$= \mathbb{P}(\tau_{\partial}(\xi) = \ell)$$

where

$$\boldsymbol{\tau}_{\partial}(\xi) := \inf\{j \ge 0 \,/\, B_m(\xi) = 0 \,\,\forall \, m \ge j\}$$

46

is the entrance time to the "cemetary" ∂ for the "chain" $(B_k(\xi))_k$ (here an independent sequence).

4.5. The general symmetric Schur measure. We now use (83) in the case where $\mathcal{A} = \overline{\mathcal{B}} \text{ if } \mathcal{A}, \mathcal{B} \subset \mathbb{C}, \text{ i.e. } g_k[\mathcal{A}] = \overline{g_k[\mathcal{B}]} \text{ for } g_k = p_k, e_k \text{ or } h_k. \text{ One has in particular} \\ H[\mathcal{A}\overline{\mathcal{A}}] = \exp\left(\sum_{k \ge 1} \frac{1}{k} |p_k[\mathcal{A}]|^2\right) \text{ and } H[\widehat{\omega}(z\mathcal{A} + z^{-1}\overline{\mathcal{A}})] = |H[z\widehat{\omega}\mathcal{A}]|^2 \text{ for } z \in \mathbb{U}. \\ \text{We define the probability measure on } \mathbb{U}$

$$\mu_{\mathcal{A}}(d\theta) := \frac{|H[z\widehat{\omega}\mathcal{A}]|^2}{\mathcal{Z}(\mathcal{A})} \frac{d^*z}{z}, \qquad \mathcal{Z}(\mathcal{A}) := [z^0]|H[z\widehat{\omega}\mathcal{A}]|^2$$

and we define the OPUCs for $L^2(\mathbb{U}, \mu_{\mathcal{A}})$ as $\left(\Phi_k^{(\mathcal{A})}\right)_{k\geq 0}$ and their Verblunsky coefficients as $(\alpha_k(\mathcal{A}))_{k\geq 0}$. As a result,

$$\mathbb{P}_{\mathrm{Schur}(\mathcal{A},\overline{\mathcal{A}})}(\boldsymbol{\lambda}_{1} \leq N) = H\left[-\mathcal{A}\overline{\mathcal{A}}\right] \prod_{k=0}^{N-1} \mathcal{Z}(\mathcal{A}) \prod_{j=0}^{k-1} (1 - |\alpha_{j}(\mathcal{A})|^{2})$$

We moreover suppose that the specialisation \mathcal{A} is well-defined in the sense that $0 < \infty$ $H[\mathcal{A}\overline{\mathcal{A}}] < \infty$. The Szegö theorem is then

$$H\left[\mathcal{A}\overline{\mathcal{A}}\right] = \prod_{k \ge 0} \mathcal{Z}(\mathcal{A}) \prod_{j=0}^{k-1} (1 - |\alpha_j(\mathcal{A})|^2)$$

and the absolute convergence of the product implies that

$$\mathcal{Z}(\mathcal{A}) = \prod_{j \ge 0} (1 - |\alpha_j(\mathcal{A})|^2)^{-1}$$

Thus,

$$\mathbb{P}_{\mathrm{Schur}(\mathcal{A},\overline{\mathcal{A}})}(\boldsymbol{\lambda}_{1} \leqslant N) = \prod_{k \geq N} \mathcal{Z}(\mathcal{A})^{-1} \prod_{j=0}^{k-1} (1 - |\alpha_{j}(\mathcal{A})|^{2})^{-1}$$
$$=: \prod_{k \geq N} q_{k}(\mathcal{A}) = \prod_{\ell \geq 0} q_{\ell+N}(\mathcal{A})$$

where

$$q_k(\mathcal{A}) := \mathcal{Z}(\mathcal{A})^{-1} \prod_{j=0}^{k-1} (1 - |\alpha_j(\mathcal{A})|^2)^{-1} = \prod_{j \ge k} (1 - |\alpha_j(\mathcal{A})|^2)$$

defines a random variable $Z^{\operatorname{Schur}(\mathcal{A},\overline{\mathcal{A}})}$ such that

$$q_k(\mathcal{A}) = \mathbb{P}\left(Z^{\operatorname{Schur}(\mathcal{A},\overline{\mathcal{A}})} \leqslant k\right) \quad \Longrightarrow \quad \mathbb{P}\left(Z^{\operatorname{Schur}(\mathcal{A},\overline{\mathcal{A}})} = k\right) = |\alpha_{k-1}(\mathcal{A})|^2 \prod_{j \ge k} (1 - |\alpha_j(\mathcal{A})|^2)$$

As a result, one has the

Theorem 4.3 $(\lambda_1^{\operatorname{Schur}(\mathcal{A},\overline{\mathcal{A}})})$ is a maximum of a sequence of independent random variables). For a sequence $(Z_k^{\operatorname{Schur}(\mathcal{A},\overline{\mathcal{A}})})_{k\geq 0}$ of *i.i.d.* random variables equal in law to $Z^{\operatorname{Schur}(\mathcal{A},\overline{\mathcal{A}})}$, one has

$$\boldsymbol{\lambda}_{1}^{\mathrm{Schur}(\mathcal{A},\overline{\mathcal{A}})} \stackrel{\mathcal{L}}{=} \max_{k \ge 0} \left\{ Z_{k}^{\mathrm{Schur}(\mathcal{A},\overline{\mathcal{A}})} - k \right\}$$
(87)

Moreover

$$Z_1^{\text{Schur}(\mathcal{A},\overline{\mathcal{A}})} \stackrel{\mathcal{L}}{=} \inf\{j \ge 0 \,/\, \forall \, m \ge j, \ B_j(\mathcal{A}) = 0\}$$
(88)

where the $(B_j(\mathcal{A}))_{j\geq 0}$ are independent $\{0,1\}$ -Bernoulli random variables of expectation $|\alpha_j(\mathcal{A})|^2$.

4.6. Remark on the discrete Fuchs lemma. In the vein of § 2.2, we have the :

Lemma 4.4 (Discrete unnormalised and varying interval Fuchs lemma). • For $I \subset \mathbb{N}$, let $\mathcal{K}_{\ell} : \ell^2(I) \to \ell^2(I)$ be self-adjoint. Let $(g_{\ell}, \lambda_{\ell})$ be such that $\mathcal{K}_{\ell}g_{\ell} = \lambda_{\ell}g_{\ell}$ for all $\ell \in \mathbb{N}$. Define $\Delta u_k := u_{k+1} - u_k$ and $\nabla u_k := u_k - u_{k-1}$. Then,

$$\Delta \lambda_{\ell} = \frac{\langle \Delta \mathcal{K}_{\ell} g_{\ell+1}, g_{\ell} \rangle_{\ell^{2}(I)}}{\langle g_{\ell+1}, g_{\ell} \rangle_{\ell^{2}(I)}}$$

• For $I_{\ell} := \ell + \mathbb{N} := \{\ell, \ell + 1, \dots, \}$, let $\mathcal{K} : \ell^2(I_{\ell}) \to \ell^2(I_{\ell})$ be self-adjoint with a kernel independent of ℓ . Let $(g_{\ell}, \lambda_{\ell})$ be such that $\mathcal{K}_{\ell}g_{\ell} = \lambda_{\ell}g_{\ell}$ (with an index to denote the action on $\ell^2(I_{\ell})$). Then,

$$\lambda_{\ell+1} = \lambda_{\ell} \frac{\langle g_{\ell}, g_{\ell+1} \rangle_{\ell^2(I_{\ell+1})}}{\langle g_{\ell}, g_{\ell+1} \rangle_{\ell^2(I_{\ell})}}$$

Proof. A discrete differentiation of $\mathcal{K}_{\ell}g_{\ell} = \lambda_{\ell}g_{\ell}$ yields to

 $\Delta \mathcal{K}_{\ell} g_{\ell+1} + \mathcal{K}_{\ell} \Delta g_{\ell} = \Delta \lambda_{\ell} g_{\ell+1} + \lambda_{\ell} \Delta g_{\ell} \qquad \Longleftrightarrow \qquad (\mathcal{K}_{\ell} - \lambda_{\ell} I) \Delta g_{\ell} = -\Delta (\mathcal{K}_{\ell} - \lambda_{\ell} I) g_{\ell+1}$ Since $\mathcal{K}_{\ell}^{*} = \mathcal{K}_{\ell}$ and $(\mathcal{K}_{\ell} - \lambda_{\ell} I) \Delta g_{\ell} \in \operatorname{Im}(\mathcal{K}_{\ell} - \lambda_{\ell} I)$, one has $(\mathcal{K}_{\ell} - \lambda_{\ell} I) \Delta g_{\ell} \perp \operatorname{ker}(\mathcal{K}_{\ell} - \lambda_{\ell} I) = \mathbb{R} g_{\ell}$. In the first case, one takes the scalar product with g_{ℓ} to obtain the result. In the second case, one has

$$\begin{split} (\Delta \mathcal{K}_{\ell})g_{\ell+1} &= \sum_{r \ge \ell+1} \mathcal{K}(\cdot, r)g_{\ell+1}(r) - \sum_{r \ge \ell} \mathcal{K}(\cdot, r)g_{\ell+1}(r) \\ &= -\mathcal{K}(\cdot, \ell)g_{\ell+1}(\ell) \end{split}$$

Hence, with $\langle \cdot, \cdot \rangle_{\ell} := \langle \cdot, \cdot \rangle_{\ell^2(I_{\ell})}$

$$\langle (\Delta \mathcal{K}_{\ell}) g_{\ell+1}, g_{\ell} \rangle_{\ell} = - \langle \mathcal{K}(\cdot, \ell) g_{\ell+1}(\ell), g_{\ell} \rangle_{\ell} = - \langle \mathcal{K}(\ell, \cdot), g_{\ell} \rangle_{\ell} g_{\ell+1}(\ell)$$

$$= -\mathcal{K}_{\ell}g_{\ell}(\ell)g_{\ell+1}(\ell)$$
$$= -\lambda_{\ell}g_{\ell}(\ell)g_{\ell+1}(\ell)$$

Finally

$$\Delta \lambda_{\ell} = \frac{\langle \Delta \mathcal{K}_{\ell} g_{\ell+1}, g_{\ell} \rangle_{\ell}}{\langle g_{\ell+1}, g_{\ell} \rangle_{\ell}}$$
$$= -\lambda_{\ell} \frac{g_{\ell}(\ell) g_{\ell+1}(\ell)}{\langle g_{\ell+1}, g_{\ell} \rangle_{\ell}}$$

namely

$$\begin{aligned} \lambda_{\ell+1} &= \lambda_{\ell} \bigg(1 - \frac{g_{\ell}(\ell)g_{\ell+1}(\ell)}{\langle g_{\ell+1}, g_{\ell} \rangle_{\ell}} \bigg) \\ &= \lambda_{\ell} \frac{\langle g_{\ell+1}, g_{\ell} \rangle_{\ell+1}}{\langle g_{\ell+1}, g_{\ell} \rangle_{\ell}} \end{aligned}$$

which gives the result.

Define $\psi^* = \langle \psi, \cdot \rangle_{\ell^2(I)}$ for the relevant choice of I (depending on which case we are concerned with).

In the case of the Poisson-Plancherel measure, a commuting difference operator was introduced in [31, prop. 2.12]. It reads on $\ell^2(\mathbb{N})$

$$\mathcal{D}_{\xi} := \Delta \mathfrak{M}_{\mathrm{id}} \nabla - \mathfrak{M}_{\beta_s}, \qquad \beta_s(k) := \frac{(k+s)(k+1-2\sqrt{\xi})}{\sqrt{\xi}}$$

Moreover, for $k = \left[2\sqrt{\xi} + \xi^{1/6}x\right]$ and $s = \left[2\sqrt{\xi} + \xi^{1/6}S\right]$, it converges in the weak sense to $\mathcal{L}_{TW,S} := DXD - X(X+S)$.

By (74), one has $\Delta \mathcal{K}_{\text{PoPl}(\xi),\ell} = B_{\ell}^{(\xi)} \otimes (B_{\ell}^{(\xi)})^*$ acting on $\ell^2(\mathbb{N})$, and

$$\Delta \lambda_{\ell} = \frac{\left\langle g_{\ell+1}, B_{\ell}^{(\xi)} \right\rangle_{\ell^{2}(\mathbb{N})} \left\langle g_{\ell}, B_{\ell}^{(\xi)} \right\rangle_{\ell^{2}(\mathbb{N})}}{\left\langle g_{\ell+1}, g_{\ell} \right\rangle_{\ell^{2}(\mathbb{N})}}$$

which is not known to be non negative. The same can be said about the second expression in lemma 4.4 in case of $\mathcal{K}_{\text{PoPl}(\xi)}$ acting on $\ell^2(\ell + \mathbb{N})$.

4.7. **Discussion.** The equation (79) reads

$$\det(I - \mathcal{K}_N)_{\ell^2(\mathbb{N})} = \exp\left(-\sum_{s \ge N} \operatorname{tr}\left(\log(I - \mathcal{K}_{s+1}) - \log(I - \mathcal{K}_s)\right)_{\ell^2(\mathbb{N})}\right)$$
$$= \exp\left(-\sum_{s \ge N} \operatorname{tr}\left(\log(I - (I - \mathcal{K}_s)^{-1}(\mathcal{K}_{s+1} - \mathcal{K}_s))\right)_{\ell^2(\mathbb{N})}\right)$$

the operator $I - \mathcal{K}_s$ being invertible as $\det(I - \mathcal{K}_s)_{\ell^2(\mathbb{N})} \neq 0$.

In the case of Poisson-Plancherel, the Hankel convolutive form (74)/(78) shows that

$$\mathcal{K}_N^{(\xi)} = \sum_{k \ge N} B_k^{(\xi)} \otimes (B_k^{(\xi)})^* \qquad \Longrightarrow \qquad \mathcal{K}_{s+1}^{(\xi)} - \mathcal{K}_s^{(\xi)} = -B_s^{(\xi)} \otimes (B_s^{(\xi)})^*$$

This last operator is of rank one with non zero trace, which implies

$$\det \left(I - \mathcal{K}_N^{(\xi)}\right)_{\ell^2(\mathbb{N})} = \exp\left(-\sum_{s \ge N} \operatorname{tr}\left(\log(I + (I - \mathcal{K}_s^{(\xi)})^{-1} B_s^{(\xi)} \otimes (B_s^{(\xi)})^*)\right)_{\ell^2(\mathbb{N})}\right)$$
$$= \exp\left(-\sum_{s \ge N} \log\left(1 + \left\langle(I - \mathcal{K}_s^{(\xi)})^{-1} B_s^{(\xi)}, B_s^{(\xi)}\right\rangle_{\ell^2(\mathbb{N})}\right)\right)$$
$$= \prod_{s \ge N} \frac{1}{1 + \left\langle(I - \mathcal{K}_s^{(\xi)})^{-1} B_s^{(\xi)}, B_s^{(\xi)}\right\rangle_{\ell^2(\mathbb{N})}}$$

Since det $\left(I - \mathcal{K}_N^{(\xi)}\right)_{\ell^2(\mathbb{N})} = \prod_{s \ge N} q_s(\xi)$, one gets

$$q_s(\xi) = \frac{1}{1 + \left\langle (I - \mathcal{K}_s^{(\xi)})^{-1} B_s^{(\xi)}, B_s^{(\xi)} \right\rangle_{\ell^2(\mathbb{N})}}$$
(89)

One can also write (79) as

$$\det(I - \mathcal{K}_N)_{\ell^2(\mathbb{N})} = \exp\left(+\sum_{s \ge N} \operatorname{tr}\left(\log(I + (I - \mathcal{K}_{s+1})^{-1}(\mathcal{K}_{s+1} - \mathcal{K}_s))\right)_{\ell^2(\mathbb{N})}\right)$$
$$= \exp\left(+\sum_{s \ge N} \operatorname{tr}\left(\log(I - (I - \mathcal{K}_{s+1})^{-1}B_s^{(\xi)} \otimes (B_s^{(\xi)})^*)\right)_{\ell^2(\mathbb{N})}\right)$$
$$= \prod_{s \ge N} \left(1 - \left\langle(I - \mathcal{K}_{s+1}^{(\xi)})^{-1}B_s^{(\xi)}, B_s^{(\xi)}\right\rangle_{\ell^2(\mathbb{N})}\right)$$

implying also that

$$q_s(\xi) = 1 - \left\langle (I - \mathcal{K}_{s+1}^{(\xi)})^{-1} B_s^{(\xi)}, B_s^{(\xi)} \right\rangle_{\ell^2(\mathbb{N})}$$
(90)

Several expressions for $q_s(\xi) = \mathbb{P}(Z^{\text{PoPl}(\xi)} \leq s)$ are thus given by (84), (89) and (90). To these ones, one can add the resolvant representation of Borodin [21, (4.1), (4.3)] that reads

$$q_s(\xi) = \frac{1}{\mathcal{R}_s[s,s]}, \qquad \mathcal{R}_s := \mathcal{K}_s(I - \mathcal{K}_s)^{-1} = I - (I - \mathcal{K}_s)^{-1}$$
(91)

Using the fact that $||B_s^{(\xi)}||_{\ell^2(\mathbb{N})}^2 = 1$, we see that this representation is equivalent to (89); but [21, (4.3)] gives another expression of $\mathcal{R}_s[s, s]$ using the theory of discrete Riemann-Hilbert problems for discrete integrable operators constructed in [20]. In particular, it gives a discrete Painlevé equation analogous to the continuous one given in § 3.4.

50

Such an expression can also be obtained by applying Cramer's rule for the inverse of a matrix and a block decomposition formula as in [10, first proof & (2)]. Notice that the function \mathcal{H}_N defined in (45) is also defined by squared norms of Orthogonal Polynomials on the Real Line (OPRL) and that the $q_s(\xi)$ are related with squared norms of OPUCs. The two theories are thus analogous and one can ask about similar convergence issues.

Remark 4.5. The analogue of (13) i.e.

$$\mathbb{P}(Z_{\ell}(\phi) \leqslant t) = \exp\left(-\int_{t}^{+\infty} \left\|\boldsymbol{H}(\phi_{s})^{\ell}\phi_{s}\right\|_{L^{2}(\mathbb{R}_{+})}^{2} ds\right)$$

is obtained by writing

$$\det(I - \mathcal{K}_N)_{\ell^2(\mathbb{N})} = \exp\left(\sum_{\ell \geqslant 1} \frac{1}{\ell} \sum_{s \geqslant N} \operatorname{tr}\left(\left((I - \mathcal{K}_s)^{-1} (\mathcal{K}_{s+1} - \mathcal{K}_s)\right)^\ell\right)_{\ell^2(\mathbb{N})}\right)$$
$$= \prod_{\ell \geqslant 1} \exp\left(\frac{1}{\ell} \sum_{s \geqslant N} \operatorname{tr}\left(\left((I - \mathcal{K}_s)^{-1} (-B_s^{(\xi)} \otimes (B_s^{(\xi)})^*\right)^\ell\right)_{\ell^2(\mathbb{N})}\right)$$
$$= \prod_{\ell \geqslant 1} \exp\left(-\frac{1}{\ell} \sum_{s \geqslant N} \left\langle(I - \mathcal{K}_s)^{-1} B_s^{(\xi)}, B_s^{(\xi)}\right\rangle_{\ell^2(\mathbb{N})}^\ell\right)$$
$$=: \prod_{\ell \geqslant 1} \mathbb{P}\left(\mathbf{Z}_\ell^{(\xi)} \leqslant N\right)$$

This is thus a different edge randomisation structure ! The same can be said about the GUE_N . Several such structures are thus possible.

The relevant convergence here was originally proven by Baik-Deift-Johansson [6] and later reproven by Johansson [74], Okounkov [98] and Borodin-Okounkov-Olshanski [31]. It reads

$$\frac{\boldsymbol{\lambda}_{1}^{\text{PoPl}(\xi)} - 2\sqrt{\xi}}{\xi^{1/6}} \xrightarrow[\xi \to +\infty]{\mathcal{L}} \mathcal{TW}_{2}$$

$$\tag{92}$$

With \mathcal{TW}_2 given by (6) and in the same vein as question 3.15, one can ask the :

Question 4.6. Can one prove the Baik-Deift-Johansson convergence (92) in the same vein as in theorem 3.11, using the convergence of a certain range $k \equiv k(\xi)$ of the random variables

$$\frac{Z^{\text{PoPl}(\xi)} - k(\xi) - 2\sqrt{\xi}}{\xi^{1/6}}$$

Same question for the $Z_k^{(\xi)}$ of remark 4.5.

APPENDIX A. SYMMETRIC FUNCTIONS

A.1. Generalities.

A.1.1. Series. We denote by $\mathbb{U} := \{z \in \mathbb{C} : |z| = 1\}$ the unit circle and set

$$d^*z := \frac{dz}{2i\pi}, \qquad \frac{d^*Z}{Z} := \prod_{k=1}^n \frac{d^*z_k}{z_k} \quad \text{if} \quad Z := (z_1, \dots, z_n)$$

We will use the notation

$$[z^n]f(z) = a_n$$
 if $f(z) = \sum_n a_n z^n$

to denote the *n*-th (Fourier) coefficient of a Laurent series f. Up to a dilation, one can suppose that f has a radius of convergence at least equal to 1, in which case

$$[z^n]f(z) = \oint_{\mathbb{U}} f(z) z^{-n} \frac{d^* z}{z}$$
(93)

A.1.2. Multi-index notation. We use the multi-index notation

$$X^{\alpha} := \prod_{k \ge 1} x_k^{\alpha_k} \tag{94}$$

and subsequently the notation $[X^{\alpha}]f(X)$ for the multivariate Fourier coefficient. If $N \in \mathbb{Z}$ (in particular for N = -1) and $X = (x_1, \ldots, x_k)$, we set

$$X^N := x_1^N x_2^N \dots x_k^N, \qquad X^{-1} := x_1^{-1} x_2^{-1} \dots x_k^{-1}$$

A.1.3. Partitions. The partitions of an integer n are weakly decreasing sequences $\lambda := (\lambda_1 \ge \lambda_2 \ge \cdots \ge \lambda_m)$ such that $|\lambda| := \sum_{i=1}^m \lambda_i = n$. We will write $\lambda \vdash n$ for $|\lambda| = n$. The length m of λ will be denoted by $\ell(\lambda)$. The transpose of a partition will be denoted by λ' . We refer to [83, ch. I.1] for generalities concerning them.

A.1.4. Symmetric functions and plethysm. For notations an definitions concerning symmetric functions, the reference is Macdonald [83]. We denote by s_{λ} the Schur functions, p_k the power functions, h_k the homogeneous complete symmetric functions and e_k the elementary symmetric functions. The ring of symmetric functions is denoted by $\Lambda := \mathbb{C}[p_k]_{k \ge 1} = \mathbb{C}[h_k]_{k \ge 0} = \mathbb{C}[e_k]_{k \ge 0}$.

The use of λ -ring/plethystic notations and alphabets will be done in the same vein as [81, 67], using in particular the convention of [67, § 2] that writes plethysm with a bracket [·]: the plethysm of a symmetric function f in an abstract alphabet \mathcal{A} will be denoted $f[\mathcal{A}]$. In the λ -ring vision of e.g. [81], symmetric functions are polynomial functors in the vein of [83, Appendix A p. 149]. As a result, the term functor will be sometimes used to designate some quantities.

Remark A.1. We recommend the reading of $[67, \S 2]$ to become familiar with the notion and the use of plethysm in the manipulation of symmetric functions.

A.1.5. Plethystic sum and product. Recall that $p_k(x_1, x_2, ...) := \sum_{\ell \ge 1} x_{\ell}^k$. Given the sets of variables or "alphabets" $X := \{x_k\}_{k \ge 1}$ and $Y := \{y_k\}_{k \ge 1}$, one defines

$$p_k[X+Y] = p_k[X] + p_k[Y]$$
$$p_k[X \cdot Y] = p_k[X]p_k[Y]$$

Note the difference of convention between $p_k(X+Y) = \sum_{\ell} (x_{\ell} + y_{\ell})^k$ and $p_k[X+Y]$.

With these definitions, X + Y is the alphabet obtained by concatenating X and Y and $X \cdot Y = \{x_k y_\ell\}_{k,\ell \ge 1}$ is a "tensor product" alphabet. It is clear that + and \cdot are associative operations on the alphabets. From now on, we will forget the dot when considering a tensor alphabet and we will write XY for it; we will also forget the $\{\cdot\}$ for alphabets. For instance, we write tX for $\{t\} \cdot X$.

A.1.6. The H functor. We define the generating series of the complete homogeneous symmetric functions by

$$H[X] := \sum_{n \ge 0} h_n(X) = \prod_{k \ge 1} \frac{1}{1 - x_k}$$
(95)

It clear that

$$H[X+Y] = H[X]H[Y]$$
(96)

The link between the h_k 's and the p_k 's can be summarised by the generating functions identity [83, I-2, (2.10)]¹⁰

$$H[X] = e^{P[X]}, \qquad P[X] := \sum_{k \ge 1} \frac{p_k[X]}{k}$$
 (97)

With the usual convention $p_{\lambda} := \prod_{k \ge 1} p_{\lambda_k}$, the expansion of e^P gives [83, I (2.14)]

$$H[X] = \sum_{\lambda} \frac{1}{z_{\lambda}} p_{\lambda}[X] \tag{98}$$

Replacing X by XY yields

$$\sum_{\lambda} \frac{1}{z_{\lambda}} p_{\lambda}[X] p_{\lambda}[Y] = \sum_{n \ge 0} h_n[XY] = H[XY]$$

¹⁰We slightly change the convention of [83, I-2, (2.10)] that uses instead $P(tX) = \sum_{k \ge 1} t^k p_k[X]$.

A.1.7. *Plethystic difference*. The difference of two alphabets X - Y is defined as the formal alphabet such that [83, 1-3, ex. 23]

$$H[X - Y] = \frac{H[X]}{H[Y]}$$

This last definition is equivalent to [83, I(2.10)]

$$p_k[X - Y] = p_k[X] - p_k[Y]$$

Note that $f(-X) := f(-x_1, -x_2, ...) \neq f[-X]$. To differentiate between these two operations, we define the "minus sign" alphabet

$$\varepsilon := \{-1\}\tag{99}$$

which is such that $f[\varepsilon X] = f(-X)$.

A.1.8. Involution ω . The fundamental involution ω [83, I-2, (2.7)] is defined by $\omega(h_n) = e_n$ and $\omega(e_n) = h_n$ (i.e. $\omega^2 = I$). Equivalently, it can be defined by $\omega(s_{\lambda}) = s_{\lambda'}$ [83, I-3, (3.8)]. Since $s_{\lambda}[-X] = (-1)^{|\lambda|} s_{\lambda'}[X] = s_{\lambda'}[\varepsilon X]$ (see [67] or [83, I-3, (3.10)]), we thus have $\omega(s_{\lambda}[X]) = s_{\lambda}[-\varepsilon X]$. As a result, we define the alphabet

$$\widehat{\omega} := -\varepsilon \tag{100}$$

which is such that $\omega(s_{\lambda}[X]) = s_{\lambda}[\widehat{\omega}X].$

A.1.9. Cauchy identity. The Cauchy identity is [83, I-4, (4.3)]

$$H[XY] = \sum_{\lambda} s_{\lambda}[X]s_{\lambda}[Y]$$
(101)

Acting plethystically on X with ω gives the *dual* Cauchy identity

$$H[\widehat{\omega}XY] = \sum_{\lambda} (-1)^{|\lambda|} s_{\lambda'}[X] s_{\lambda}[Y]$$
(102)

Replacing X by tX and taking $[t^n]$ in (101) and (98) gives

$$h_n[XY] = \sum_{\lambda \vdash n} \frac{1}{z_\lambda} p_\lambda[X] p_\lambda[Y] = \sum_{\lambda \vdash n} s_\lambda[X] s_\lambda[Y]$$
(103)

A.1.10. Exponential alphabet. The alphabet \mathcal{E} is defined by setting

$$p_k[\mathcal{E}] = \mathbb{1}_{\{k=1\}} \qquad \Longleftrightarrow \qquad H[t\mathcal{E}] = e^t \tag{104}$$

Schur-Weyl duality gives [83, I-7 (7.7) p. 114]

$$s_{\lambda} = \sum_{\mu \vdash |\lambda|} \chi^{\lambda}_{\mu} \frac{p_{\mu}}{z_{\mu}}$$

For $\lambda \vdash n$, one has $p_{\lambda}[\mathcal{E}] = \prod_{k} p_{\lambda_{k}}[\mathcal{E}] = \mathbb{1}_{\{\lambda=1^{n}\}}$, hence

$$s_{\lambda}[\mathcal{E}] = \sum_{\mu \vdash n} \chi_{\mu}^{\lambda} \frac{p_{\mu}[\mathcal{E}]}{z_{\mu}} = \frac{\chi_{1^n}^{\lambda}}{z_{1^n}} = \frac{\chi^{\lambda}(\mathrm{id}_{\mathfrak{S}_n})}{n!}$$

Since $\chi^{\lambda} = \operatorname{tr}(\rho^{\lambda})$ with $\rho^{\lambda} : \mathfrak{S}_n \to \operatorname{Mat}_{d_{\lambda}}(\mathbb{C})$ (or $\operatorname{End}(V^{\lambda})$ with $V^{\lambda} \simeq \mathbb{C}^{d_{\lambda}}$), the dimension d_{λ} of the irreducible module V^{λ} is given by $\chi^{\lambda}(\operatorname{id}_{\mathfrak{S}_n}) = \operatorname{tr}(I_{d_{\lambda}})$, hence

$$d_{\lambda} = |\lambda|! \, s_{\lambda}[\mathcal{E}] \tag{105}$$

A.1.11. *The Schur functions*. The Schur functions have several equivalent definitions. One is for instance the Jacobi-Trudi identity [83, I-3 (3.4)]

$$s_{\lambda} = \det \left(h_{\lambda_{i+\ell-j}} \right)_{1 \leqslant i,j \leqslant \ell}, \quad \forall \ell \geqslant \ell(\lambda)$$
(106)

which is equivalent, with the simple manipulations of e.g. $[9, \S 3.1]$ to the Fourier coefficient form

$$s_{\lambda}[X] = \left[U^{\lambda}\right] H \left[XU - U^{\varepsilon R}\right]$$
(107)

These functions satisfy the property of "restricted" orthogonality [47]/[83, VI-9 rk. 2]

$$\int_{\mathcal{U}_n} s_{\lambda}(U) s_{\mu}(U^{-1}) dU = \delta_{\lambda,\mu} \mathbb{1}_{\{\ell(\lambda) \le n\}}$$
(108)

where dU is the normalised Haar measure of the unitary group \mathcal{U}_n , whose diagonal part is given by the CUE_n measure $|\Delta(e^{i\theta})|^2 \frac{d\theta}{n!(2\pi)^n}$; see [47]/[9, § 2.3].

The Schur functions are orthogonal for the Hall scalar product $\langle \cdot, \cdot \rangle$ that can be defined by [83, I-4]

$$\langle s_{\lambda}, s_{\mu} \rangle = \delta_{\lambda,\mu} \tag{109}$$

A.2. The Schur measure and the Schur kernel. The Schur measure is defined in (71). The fact that it is a probability measure for a positive specialisation comes from the Cauchy identity (101). The law of the total parts of the measure is given by

$$\mathbb{E}_{\mathrm{Schur}(\mathcal{A},\mathcal{B})}\left(t^{|\boldsymbol{\lambda}|}\right) = H[(t-1)\mathcal{A}\mathcal{B}]$$
(110)

One has used the scaling property $s_{\lambda}[tX] = t^{|\lambda|}s_{\lambda}[X]$. In particular, for $\mathcal{A} = (p_k)_k$ and $\mathcal{B} = (q_k)_k$, one gets

$$\mathbb{E}_{\mathrm{Schur}(\mathcal{A},\mathcal{B})}\left(t^{|\boldsymbol{\lambda}|}\right) = \prod_{i,j} H[(t-1)p_iq_j] = \prod_{i,j} \frac{1-p_iq_j}{1-tp_iq_j} = \prod_{i,j} \mathbb{E}\left(t^{\mathrm{Geom}(p_iq_j)}\right)$$

hence the equality in law $|\boldsymbol{\lambda}^{\text{Schur}(\mathcal{A},\mathcal{B})}| = \sum_{i,j} \text{Geom}(p_i q_j)$ with independent random variables in the RHS. Of course, since the only ingredients needed for such an equality in law are the scaling of the symmetric functions and the Cauchy identity, it is also valid for e.g. the Macdonald measure [23], but with random variables different from the geometric ones.

The Schur kernel operator is

$$\mathcal{K}_{\mathrm{Schur}(\mathcal{A},\mathcal{B})} := \mathcal{H}(a_{\mathcal{A},\mathcal{B}})\mathcal{H}(\widetilde{a}_{\mathcal{A},\mathcal{B}}^{-1})$$
(111)

where the fundamental function $a \equiv a_{\mathcal{A},\mathcal{B}}$ is given by

$$a_{\mathcal{A},\mathcal{B}}(z) := H_{\varepsilon} \big[z\mathcal{A} - z^{-1}\mathcal{B} \big]$$
(112)

with $H_{\mathcal{A}}[X] := H[\mathcal{A}X], \ \widetilde{f}(z) := f(z^{-1}) \text{ and } f^{-1} := 1/f \text{ here.}$

Supposing that $a(z) \equiv a_{\mathcal{A},\mathcal{B}}(z) = \sum_{k \in \mathbb{Z}} \widehat{a}_k z^k$, the Schur kernel can take the following forms :

• Hankel convolutive form [10] :

$$\mathcal{K}_{\mathrm{Schur}(\mathcal{A},\mathcal{B})}[x,y] = \sum_{k \ge 1} \widehat{a}_k(x) \widehat{a}_k(y), \qquad \widehat{a}_k(x) := \widehat{a}_{k+x}$$
(113)

• Integral form [30, (2.1), (2.2)]:

$$\mathcal{K}_{\text{Schur}(\mathcal{A},\mathcal{B})}[x,y] = [z^x w^y] \frac{a(z)/a(w)}{z/w - 1} \mathbb{1}_{\{|z/w| > 1\}}$$
(114)

• Christoffel-Darboux form : see [30, rk. 2] for the expression.

Appendix B. Prolate hyperspheroidal wave functions

The main reference for this § is [118]. We also refer to [14, 60, 114]. We nevertheless use the terminology of [69, 70], i.e. "hyperspheroidal" instead of "generalised" as there are several possible generalisations of the prolate spheroidal wave functions, listed for instance in [105, § 30.12 p. 704] or [14, (3.2)]. See also [40, 66] for interesting developments of generalised prolate spheroidal wave functions in relation with the Airy kernel operator.

56

B.1. Definitions. We define the following prolate hyperspheroidal wave function operator

$$PHO_{N,c} := \frac{d}{dx}(1-x^2)\frac{d}{dx} + c^2(1-x^2) - \frac{\rho^2}{x^2}, \qquad \rho^2 \equiv \rho_N^2 := N^2 - \frac{1}{4}$$
(115)

This operator commutes in $L^2([0,1])$ with the operator $J_{N,c}: L^2([0,1]) \to L^2([0,1])$ with kernel

$$\mathcal{J}_{N,c}(x,y) := \sqrt{cxy} J_N(cxy), \qquad J_m(x) := [z^m] e^{x(z-z^{-1})/2}$$
(116)

whose origin in optics comes, amongst others, from [69, 70, 118]

"the modes in a maser interferometer with confocal spherical mirrors of circular cross section".

The common eigenvectors of $\text{PHO}_{N,c}$ and $J_{N,c}$ are called *prolate hyperspheroidal wave* functions and are denoted by $\psi_{k}^{(N,c)}$.

The operator $\text{PHO}_{1/2,c}$ is the more classical prolate spheroidal wave function operator (up to an additive constant) that commutes with the sine kernel operator \mathbf{K}_{sinc} with kernel $(x, y) \mapsto \operatorname{sinc}(\pi(x-y))$ in $L^2([-1, 1])$ [117, (23), (26)]; see Mehta [86, ch. 6 & 18] for a use of its eigenvectors in Random Matrix Theory and [105, ch. 30 p. 697] for properties of these functions. Note that the Bessel function satisfies $J_{1/2}(x) = \sqrt{\frac{2}{\pi x}} \sin(x)$, hence $\text{PHO}_{1/2,c}$ also commutes with the operator of kernel $(x, y) \mapsto \sin(cxy)$ on $L^2([0, 1])$ which is equal, up to a symmetrisation that does not change the (real) eigenvectors and eigenvalues, to the square root of \mathbf{K}_{sinc} that has kernel e^{icxy} , see e.g. [86, (6.3.17)].

The operator $\text{PHO}_{N,0}$ is a particular specialisation of an intertwinning of the Jacobi differential operator $\mathcal{L}_{\alpha,\beta} := \frac{d}{dx}(1-x^2)\frac{d}{dx} + (\beta - \alpha - x(\alpha + \beta))\frac{d}{dx}$ whose eigenfunctions are the Jacobi polynomials. More precisely, if $P_k^{(\alpha,\beta)}$ are the Jacobi polynomials [120, ch. IV], defining the odd Zernike polynomials by [114, (16)]

$$T_k^{(N)}(x) := \tau_k^{(N)} x^{N+\frac{1}{2}} P_k^{(N,0)} (1-2x^2), \qquad \tau_k^{(N)} := \sqrt{2(2k+N+1)}$$
(117)

one has

PHO_{N,0}
$$T_k^{(N)} = -\kappa_k^{(N)} T_k^{(N)}, \qquad \kappa_k^{(N)} := (N + 2k + \frac{1}{2})(N + 2k + \frac{3}{2})$$

The tri-band relation of orthogonal polynomials implies that the (semi-infinite) matrix of the endomorphism of multiplication by X^2 in the basis of the $T_k^{(N)}$ is tri-diagonal. Since the matrix of $PHO_{N,0}$ in this basis is diagonal and $PHO_{N,c} = PHO_{N,0} + c^2(1 - X^2)$, one concludes that the matrix of $PHO_{N,c}$ is tri-diagonal in the $(T_k^{(N)})_{k\geq 0}$ basis (with explicit coefficients coming from the Jacobi polynomials). As a result, expanding the eigenvectors $(\psi_{\ell}^{(N,c)})_{\ell\geq 1}$ in this basis according to

$$\psi_{\ell}^{(N,c)} = \sum_{k \ge 0} d_{\ell,k}^{(N,c)} T_k^{(N)}, \qquad \text{PHO}_{N,c} \, \psi_{\ell}^{(N,c)} = \lambda_{\ell}^{(N,c)} \psi_{\ell}^{(N,c)}$$

one obtains the semi-infinite matrix equation [114, (30)]

$$\boldsymbol{M}^{(N,c)}d_{\ell,\cdot}^{(N,c)} = \lambda_{\ell}^{(N,c)}d_{\ell,\cdot}^{(N,c)}, \qquad \boldsymbol{M}^{(N,c)} := \operatorname{Mat}_{(T_k^{(N)})_k}(\operatorname{PHO}_{N,c})$$

In the same vein as for the prolate spheroidal wave functions, one can expand $\psi_{\ell}^{(N,c)}$ and $\lambda_{\ell}^{(N,c)}$ into powers of c; see e.g. [118].

B.2. Rescaling.

Lemma B.1 (Limiting hyperspheroidal operators). Set

$$x := 1 - \frac{X}{2}N^{-2/3}, \qquad y := 1 - \frac{Y}{2}N^{-2/3}, \qquad c := 2N - sN^{1/3}$$

Then, locally uniformly in \mathbb{R}_+

$$\frac{N^{1/6}}{\sqrt{2}}\mathcal{J}_{2N,2N-sN^{1/3}}(1-XN^{1/3},1-YN^{1/3}) \xrightarrow[N\to+\infty]{} \operatorname{Ai}(X+Y+s)$$

and, on test functions that are \mathcal{C}^2 with exponential decay at infinity (say)

$$\frac{N^{-2/3}}{4} \operatorname{PHO}_{N,N-sN^{1/3}/2} \xrightarrow[N \to +\infty]{} \mathcal{L}_{TW,s} := \frac{d}{dX} X \frac{d}{dX} - X(X+s)$$

As a result, the eigenvalues of $J_{N,N-sN^{1/3}/2}$ satisfy

$$N^{-1/6}\lambda_k^{(N,N-sN^{1/3}/2)} \xrightarrow[N \to +\infty]{} \lambda_k(s)$$

where $\lambda_k(s)$ is the k-th eigenvalue of $\boldsymbol{H}(Ai_s)$ acting on $L^2(\mathbb{R}_+)$.

Remark B.2. If one defines

$$\widetilde{\mathcal{J}}_{N,c}(X,Y) := \mathcal{J}_{N,c}(e^{-X}, e^{-Y})e^{-X-Y}, \qquad \widetilde{J}_{N,c}(X) := \sqrt{c} e^{-X/2} J_N(c e^{-X})$$

and the associated operator $\widetilde{J}_{N,c}$ acting on $L^2(\mathbb{R}_+)$ with kernel $\widetilde{\mathcal{J}}_{N,c}$, then, one has

$$\widetilde{\boldsymbol{J}}_{N,c} = \boldsymbol{H}(\widetilde{J}_{N,c}) \curvearrowright L^2(\mathbb{R}_+)$$

and the previous result is equivalent to the weak convergence of operators of the Hankel type in $L^2(\mathbb{R}_+)$ given by the convergence of their symbol. This convergence can be strengthen into a strong convergence depending on the functional space in which belongs the symbol, see e.g. [106, ch. 5.4] or [19, ch. 4.2].

This operator is thus, in a certain sense, more natural than the original operator (since one remains in the framework of Hankel operators on \mathbb{R}_+). The eigenvectors of this operator are the $\psi_{\ell}^{(N,c)}(e^{-x})e^{-x}$, which is somewhat similar to the exponential change of variable of Whittaker functions [23, § 4]/[62, 63].

Proof. Using the limit [76, (4.39)]/[105, (10.19.8) p. 232]

$$\nu^{1/3} J_{2\nu}(2\nu - X\nu^{1/3}) \xrightarrow[\nu \to +\infty]{} \operatorname{Ai}(X)$$
(118)

one easily proves the first claim.

The second claim comes from the fact that $dx = -\frac{N^{-2/3}}{2}dX$ which implies

$$\frac{N^{-2/3}}{4}\frac{d}{dx}(1-x^2)\frac{d}{dx} = \frac{d}{dX}\left(X - \frac{X^2N^{-2/3}}{4}\right)\frac{d}{dX},$$

and from the expansion at the second order of

$$\frac{\rho^2}{x} = \frac{N^2 - \frac{1}{4}}{1 - XN^{-2/3}/2} = \left(N^2 - \frac{1}{4}\right) \left(1 + \frac{XN^{-2/3}}{2} + \frac{X^2N^{-4/3}}{4} + O(X^3N^{-2})\right)$$

that gets compensated from the term

$$c_N^2(1-x^2) = \left(N - \frac{sN^{-1/3}}{2}\right) \left(XN^{-2/3} - \frac{X^2N^{-4/3}}{4}\right)$$

to produce the term X(X + s) after multiplication by $\frac{N^{-2/3}}{4}$. Full details are left to the reader.

Acknowledgements

The author thanks P. Biane, G. Borot, B. Eynard, P. Ferrari, J. Heiny, O. Hénard, K. Johansson, A. N. Kirillov, A. Kuijlaars, P. Maillard, S. N. Majumdar, J. Najnudel, M. Noumi, T. Popov, D. Rosen, G. Schehr, N. Simm, C. A. Tracy, P. Tran, R. Tribe, B. Westbury, O. Zaboronski and N. Zygouras for interesting discussions, remarks and insights concerning various stages of development of this work.

A particular thanks is given to P. J. Forrester for the reference [130].

DEDICATORY

This work is dedicated to the memory of my father Ahmed Mohsen Barhoumi (10/02/1949-24/10/2023).

References

- A. Aggarwal, Correlation functions of the Schur process through Macdonald difference operators, J. Combin. Theory Ser. A, 131:88-118 (2015). 41
- 2. American Institute for Mathematics, Open problems: AIM workshop on random matrices, aimath.org/WWN/randommatrices/randommatrices.pdf (2010). 3, 16
- 3. G. Amir, I. Corwin, J. Quastel, Probability distribution of the free energy of the continuum directed random polymer in 1 + 1 dimensions, Comm. Pure Appl. Math. 64(4):466-537 (2011). 2, 16, 17
- 4. G. Anderson, A. Guionnet, O. Zeitouni, An Introduction to Random Matrices, Cambridge University press (2009). 22
- F. Angeletti, E. Bertin, P. Abry, Renormalization flow for extreme value statistics of random variables raised to a varying power, J. Phys. A 45(11):115004 (2012). 39
- J. Baik, P. Deift, K. Johansson, On the distribution of the length of the longest increasing subsequence of random permutations, J. Amer. Math. Soc. 12:1119-1178 arXiv:9810105 (1999). 2, 40, 51
- 7. J. Baik, P. Deift, E. Rains, A Fredholm determinant identity and the convergence of moments for random Young tableaux, Commun. Math. Phys. 223(3):627-672 (2001). 44

- 8. Y. Barhoumi-Andréani, Extreme value statistics for the roots of a complex Kac polynomial, arXiv:1706.07516 (2017). 3
- 9. Y. Barhoumi-Andréani, A new approach to the characteristic polynomial of a random unitary matrix, arXiv:2011.02465 (2020). 55
- E. Basor, H. Widom, A Toeplitz determinant identity of Borodin and Okounkov, Integral Equations and Operator Theory 37(4):397-401 (2000). 44, 51, 56
- J. Ben Hough, M. Krishnapur, Y. Peres, B. Virag, Zeros of Gaussian Analytic Functions and Determinantal Point Processes, American Mathematical Society, Providence (2009). 5, 22, 24
- E. Bertin, G. Györgyi, Renormalization flow in extreme value statistics, J. Stat. Mech. Theory Exp. 2010(8):P08022 (2010). 39
- L. Bertini, G. Giacomin, Stochastic Burgers and KPZ equations from particle systems, Commun. Math. Phys. 183(3):571-607 (1997). 16, 17
- V. N. Beskrovny, M. I. Kolobov, Quantum-statistical analysis of superresolution for optical systems with circular symmetry, Phys. Rev. A 78(4):043824(2008). 56
- A. Bloemendal, B. Virág, Limits of spiked random matrices I, Probab. Theory Related Fields 156(3-4):795-825 (2013). 3
- A. Böttcher, On the determinant formulas by Borodin, Okounkov, Baik, Deift and Rains, in Toeplitz Matrices and Singular Integral Equations, Pobershau (Birkhäuser, Basel), Oper. Theory Adv. Appl. 135:91-99 (2001). 44
- 17. A. Böttcher, One more proof of the Borodin-Okounkov formula for Toeplitz determinants, Integral Equations and Operator Theory 41(1):123-125 (2001). 44
- 18. A. Böttcher, Szegö via Jacobi, Linear algebra and its applications 419(2-3):656-667 (2006). 44
- A. Böttcher, B. Silbermann, Analysis of Toeplitz operators, Springer Science & Business Media (2013).
 58
- A. Borodin, Riemann-Hilbert problems and the discrete Bessel kernel, Int. Math. Res. Not. 2000(3):467-494 (2000). 50
- A. Borodin, Discrete gap probabilities and discrete Painlevé equations, Duke Math. J. 117(3):489-542 (2003). 50
- A. Borodin, Asymptotic representation theory and Riemann-Hilbert problems, In: Asymptotic Combinatorics with Applications to Mathematical Physics (pp. 3-19). Springer, Berlin, Heidelberg arXiv:0110318v1 (2003). 41
- A. Borodin, I. Corwin, *Macdonald processes*, Probab. Theory Related Fields 158(1-2):225-400 (2014). 17, 56, 58
- 24. A. Borodin, I. Corwin, P. Ferrari, Free energy fluctuations for directed polymers in random media in 1+1 dimension, Comm. Pure Appl. Math. 67(7):1129-1214 (2014). 2
- A. Borodin, I. Corwin, D. Remenik, Log-Gamma Polymer Free Energy Fluctuations via a Fredholm Determinant Identity, Commun. Math. Phys. 324(1):215-232 arXiv:1206.4573 (2013). 2
- A. Borodin, V. Gorin, Lectures on integrable probability, Lecture notes, St. Petersburg School in Probability and Statistical Physics arXiv:1212.3351 (2012). 2, 41
- A. Borodin, V. Gorin, General β-jacobi corners process and the Gaussian free field, Comm. Pure Appl. Math. 68(10):1774-1844, 2015 (2015). 26
- A. Borodin, V. Gorin, Moments match between the KPZ equation and the Airy point process, Symmetry Integrability Geom. Methods Appl. SIGMA 12:102-109, special issue in honor of Percy Deift and Craig Tracy (2016). 2, 16, 17
- A. Borodin, A. Novikov, On a conjecture of Widom, J. Phys. A: Math. Gen. 39(28):8895-8903 (2006).
 41
- A. Borodin, A. Okounkov, A Fredholm determinant formula for Toeplitz determinants, Integral Equations Operator Theory 37(4):386-396 (2000). 41, 44, 56
- A. Borodin, A. Okounkov, G. Olshanski, Asymptotics of Plancherel measures for symmetric groups, J Amer. Math. Soc. 13(3):481-515 (2000). 41, 49, 51

- 32. A. Borodin, G. Olshanski, Z-measures on partitions, Robinson-Schensted-Knuth correspondence and β = 2 ensembles, in "Random matrix models and their applications", P. M. Bleher and A. R. Its eds., MSRI Publications 40:71-94 arXiv:9905189 (2001). 40
- A. Borodin, G. Olshanski, Z-measures on partitions and their scaling limits, European J. Combin. 26(6):795-834 (2005). 40
- A. Borodin, L. Petrov, Integrable probability: from representation theory to Macdonald processes, Probab. Surveys, 11:1-58 arXiv:1310:8007 (2014). 2, 41
- 35. G. Borot, C. Nadal, *Right tail asymptotic expansion of Tracy-Widom beta laws*, Random Matrices: Theory and Applications 1(3):1250006 (**2012**). 3, 39
- P. Bourgade, C. Hughes, A. Nikeghbali, M. Yor, The characteristic polynomial of a random unitary matrix : a probabilistic approach, Duke Math. J. 145:45-69 (2008). 26
- P. Bourgade, A. Nikeghbali, A. Rouault, Circular Jacobi ensembles and deformed Verblunski coefficients, Int. Math. Res. Not. 2009(3):4357-4394 (2009). 26
- P. Calabrese, P. Le Doussal, A. Rosso, Free-energy distribution of the directed polymer at high temperature, Europhys. Lett. 90(2):20002-p1-20002-p6 (2010). 2, 16, 17
- K. M. Case, J. S. Geronimo, Scattering theory and polynomials orthogonal on the unit circle, J. Math. Phys. 20(2):299-310 (1979). 44
- 40. W. R. Casper, F. A. Grünbaum, M. Yakimov, I. Zurrián, Algebras of commuting differential operators for integral kernels of Airy type, arXiv:2112.11639 (2021). 4, 56
- 41. D. Chaffai, Aspects of the Complex Ginibre Ensemble, "Libres pensés d'un mathématicien ordinaire" djalil.chafai.net/blog/2010/11/02/aspects-of-the-complex-ginibre-ensemble (2010). 5
- 42. L. H. Y. Chen, Poisson approximation for dependent trials, Ann. Probab. 3(3):534-545 (1975). 6
- I. Corwin, N. O'Connell, T. Seppäläinen, N. Zygouras, Tropical combinatorics and Whittaker functions, Duke Math. J. 163(3):513-563 (2014).
- 44. P. Deift, Some open problems in random matrix theory and the theory of integrable systems. II, SIGMA, Symmetry, Integrability and Geometry: Methods and Applications 13:016 (2017). 3
- P. Deift, A. Its, I. Krasovsky, Toeplitz matrices and Toeplitz determinants under the impetus of the Ising model: some history and some recent results, Commun. Math. Phys. 66(9):1360-1438 (2013). 25
- P. Diaconis, Patterns in eigenvalues : the 70th Josiah Willard Gibbs lecture, Bull. Amer. Math Soc. new ser. 40(2):155-178 (2003). 25, 26
- P. Diaconis, M. Shahshahani, On the Eigenvalues of Random Matrices, J. Appl. Prob. 31:49-62 (1994).
- V. Dotsenko, Bethe ansatz derivation of the Tracy-Widom distribution for one-dimensional directed polymers, Europhys. Lett. 90(2):20003-p1-20003-p5 (2010). 2, 16, 17
- 49. F. Eckhardt, R. Gesztesy, G. Nichols, G. Teschl, Weyl-Titchmarsh theory for Sturm-Liouville operators with distributional potentials, Opuscula Math. 33(3):467-563 arXiv:1208.4677 (2013). 3
- A. Edelman, B. D. Sutton, From random matrices to stochastic operators, J. Stat. Phys. 127(6):1121-1165 (2007). 3
- 51. A. Feidt, Stein's Method for Multivariate Extremes, Ph.D. thesis, Universität Zürich arXiv:1310.2564 (2013). 32
- P. Ferrari, H. Spohn, A determinantal formula for the GOE Tracy-Widom distribution, J. Phys. A, Math. Gen. 38(33):L557-L561 (2005). 20, 21
- H. Flaschka, A. C. Newell, Monodromy and spectrum-preserving deformations, Commun. Math. Phys. 76:65-116 (1980). 3
- P. J. Forrester, Log-gases and random matrices, LMS-34, Princeton University Press (2010). 6, 22, 25, 28
- P. J. Forrester, T. Nagao, Determinantal correlations for classical projection processes, J. Stat. Mech. Theory Exp. 2011(08):P08011 (2011). 41

- 56. P. J. Forrester, N. S. Witte, Application of the τ-Function Theory of Painlevé Equations to Random Matrices: P-IV, P-II and the GUE, Commun. Math. Phys. 219(2):357-398 (2001). 6, 28, 29, 30, 31, 36, 38, 39
- 57. P. J. Forrester, N. S. Witte, Application of the τ-function theory of Painlevé Equations to random matrices: P-V, P-III, the LUE, JUE, and CUE, Comm. Pure Appl. Math. 55(6):679-727 (2002). 6
- P. J. Forrester, N. S. Witte, Boundary conditions associated with the Painlevé III' and V evaluations of some random matrix averages, J. Phys. A: Math. Gen. 39(28):8983-8995 (2006).
- P. J. Forrester, N. S. Witte, Painlevé II in random matrix theory and related fields, Constr. Approx. 41(3):589-613 (2015). 38, 39
- 60. B. R. Frieden, Evaluation, design and extrapolation methods for optical signals, based on use of the prolate functions, Progress in optics 9(VIII):311-407 (1971). 4, 56
- W.H.J. Fuchs, On the eigenvalues of an integral equation arising in the theory of band-limited signals, J. Math. Anal. Appl. 9:317-330(1964). 12, 14
- A. Gerasimov, S. Kharchev, D. Lebedev, S. Oblezin, On a Gauss-Givental representation of quantum Toda chain wave function, Int. Math. Res. Not. IMRN 2006 arXiv:0505310 (2006). 58
- A. Gerasimov, D. Lebedev, S. Oblezin, New integral representation of Whittaker functions for classical Lie groups, Russian Math. Surveys 67(1):1-100 arXiv:0705.2886 (2007). 58
- J. Ginibre, Statistical ensembles of complex, quaternion, and real matrices, J. Math. Phys. 6(3):440-449 (1965). 5
- D. J. Gross, A. Matytsin, Instanton induced large N phase transitions in two and four dimensional QCD, Nucl.Phys. B 429(1):50-74 (1994). 24
- 66. F. A. Grünbaum, B. D. Vasquez, J. P. Zubelli, *Matrix bispectrality and noncommutative algebras:* beyond the prolate spheroidals, Enseign. Math. 69(3):335-351 (**2023**). 4, 56
- M. Haiman, Macdonald polynomials and geometry, New Perspectives in Algebraic Combinatorics, MSRI Publications, https://math.berkeley.edu/~mhaiman/ftp/nfact/msri.pdf (1999). 52, 54
- P. Hastings, J. B. McLeod, A boundary value problem associated with the second Painlevé transcendent and the Korteweg-de Vries equation, Arch. Ration. Mech. Anal. 73(1):31-51 (1980). 2, 20
- J. C. Heurtley, Hyperspheroidal functions, optical resonators with circular mirrors, in: Quasi Optics, pp. 367-365 (Proc. Symp. Quasi Optics), J. Fox, Ed. New York ; proceedings of the Polytechnic Inst. of Brooklyn Symposium on Quasi-Optics, June 8-10 (1964). 4, 56, 57
- J. C. Heurtley, A theoretical study of optical resonator modes and a new class of special functions, the hyperspheroidal functions, Ph. D. dissertation, Electrical Engineering Dept., University of Rochester, Rochester, New York (1965). 4, 56, 57
- T. Imamura, T. Sasamoto, Determinantal structures in the O'Connell-Yor directed random polymer model, J. Stat. Phys. 163(4):675-671 (2016). 2
- M. Jimbo, T. Miwa, Monodromy preserving deformation of linear ordinary differential equations with rational coefficients II, Physica D, 2:407-448 (1981). 28
- K. Johansson, Shape fluctuations and random matrices, Commun. Math. Phys. 209(2):437-476 (2000).
 2
- 74. K. Johansson, Discrete orthogonal polynomial ensembles and the Plancherel measure, Ann. Math. 153(1):259-296 (2001). 41, 42, 51
- K. Johansson, Random growth and random matrices, European Congress of Mathematics, Birkhäuser Basel (2001). 2, 40, 41
- K. Johansson, Random matrices and determinantal processes, Les Houches lecture arXiv:0510038v1 (2005). 2, 40, 58
- K. Johansson, From Gumbel to Tracy-Widom, Probab. Theory Related Fields 138(1-2):75-112 (2007).
 17
- I.M. Johnstone, Z. Ma, Fast approach to the Tracy-Widom law at the edge of GOE and GUE, Ann. Appl. Prob. 22(5):1962-1988 (2012). 35

- M. Katori, H. Tanemura, Zeros of Airy function and relaxation process, J. Stat. Phys. 136:1177-1204 (2009). 8
- A. N. Kirillov, Introduction to tropical combinatorics, Physics and Combinatorics, proc. Nagoya 2000, 2nd Internat.Workshop (A. N. Kirillov and N. Liskova, eds.), World Scientific, Singapore, pp. 82-150 (2001). 4
- A. Lascoux, Symmetric functions, Course Nankai University http://www.emis.de/journals/SLC/wpapers/s68vortrag/ (2001). 52
- R. Lyons, Determinantal probability measures, Publ. Math. Inst. Hautes Études Sci. 98(1):167-212 (2003). 24
- I.G. Macdonald, Symmetric functions and Hall polynomials, Oxford Mathematical Monographs, Second edition, The Clarendon Press Oxford University Press (1995). 41, 52, 53, 54, 55
- 84. S. N. Majumdar, Random matrices, the Ulam problem, directed polymers & growth models, and sequence matching, In:Complex Systems ixxxv, J.-P. Bouchard, M. Mézard, J. Daibard eds., Oxford: Elsevier (2007). 2
- S. N. Majumdar, G. Schehr, Top eigenvalue of a random matrix: large deviations and third order phase transition, J. Stat. Mech. P01012(14):1742-5468 (2014). 2
- M.-L. Metha, Random matrices, Pure and Applied Mathematics, Elsevier Academic Press (Amsterdam), vol. 142 (2004). 22, 57
- N. Minami, Definition and self-adjointness of the stochastic Airy operator, Markov Process. Related Fields 21(3P):695-711 (2015). 3
- C. Nadal, S. N. Majumdar, A simple derivation of the Tracy-Widom distribution of the maximal eigenvalue of a Gaussian unitary random matrix, J. Stat. Mech. Theory Exp. 2011(04):P04001 (2011). 24, 39
- V.-L. Nguyen, N. Zygouras, Variants of geometric RSK, geometric PNG, and the multipoint distribution of the log-gamma polymer, Int. Math. Res. Not. 2017(15):4732-4795 (2017).
- 90. M. Noumi, Painlevé equations through symmetry, Amer. Math. Soc. Vol. 223 (2004). 6, 28
- M. Noumi, Y. Yamada, Symmetries in the fourth Painlevé equation and Okamoto polynomials, Nagoya Math. J. 153:53-86 (1999). 6, 28
- N. O'Connell, T. Seppäläinen, N. Zygouras, Geometric RSK correspondence, Whittaker functions and symmetrized random polymers, Inventiones Math. 197:361-416 (2014). 2
- N. O'Connell, M. Yor, Brownian analogues of Burke's theorem, Stochastic Processes and their Applications, 96(2):285-304 (2001). 2
- 94. K. Okamoto, On the τ-function of the Painlevé equations, Physica D: Nonlinear Phenomena 2(3):525-535 (1981). 5, 28
- 95. K. Okamoto, Studies on the Painlevé equations, I: Sixth Painlevé Equation P-VI, Ann. Mat. Pura Appl. 146(1):337-381 (1987). 5, 28
- 96. K. Okamoto, Studies on the Painlevé equations, II: Fifth Painlevé Equation P-V, Jap. J. Math. 13(1):47-76 (1985). 5, 28
- 97. K. Okamoto, Studies on the Painlevé equations, III: Second and Fourth Painlevé Equations P-II and P-IV, Math. Ann. 275(1):221-255 (1986). 5, 28
- A. Okounkov, Random matrices and random permutations, Int. Math. Res. Not. IMRN 2000(20):1043-1095 (2000). 51
- A. Okounkov, Infinite wedge and random partitions, Selecta Math. New Ser. 7(1):57-81 arXiv:9907127 (2001). 40, 41
- 100. A. Okounkov, SL(2) and z-measures, In: Random matrix models and their applications, Math. Sci. Res. Inst. Publ. 40:407-420, Cambridge Univ. Press, Cambridge (2001). 40
- 101. A. Okounkov, Symmetric functions and random partitions, Course NATO and Cambridge arXiv:0309074 (2001). 40, 41, 42

- 102. A. Okounkov, N. Reshetikhin, Correlation functions of Schur process with applications to local geometry of a random 3-dimensional Young diagram, J. Amer. Math. Soc. 16:581-603 arXiv:math.CO/0107056 (2003). 41
- 103. A. Okounkov, N. Reshetikhin, Random skew plane partitions and the Pearcey process, Commun. Math. Phys. 269(3):571-609 arXiv:math.CO/0503508 (2007). 41
- 104. G. Olshanski, *Random permutations and related topics*, The Oxford Handbook of Random Matrix Theory, ch. 25, G. Akemann, J. Baik, P. Di Francesco, eds (**2011**). 40
- 105. F. W. Olver, D. W. Lozier, R. F. Boisvert, C. W. Clark (Eds.), NIST handbook of mathematical functions, Cambridge university press (2010). 3, 4, 9, 30, 37, 38, 42, 56, 57, 58
- 106. N. Nikolski, Toeplitz matrices and operators, Cambridge University Press, vol. 182 (2020). 58
- 107. E. M. Rains, Correlation functions for symmetrized increasing subsequences, arXiv:0006097 (2000).
 41
- 108. J. Ramirez, B. Rider, B. Virág, Beta ensembles, stochastic Airy spectrum, and a diffusion, J. Amer. Math. Soc. 24(4):919-944 (2011). 3
- 109. B. Rider, A limit law at the edge of a non-Hermitian random matrix ensemble, J. Phys. A:Math. Gen. 36(12):3401-3409 (2007). 5, 6
- 110. N. Ross, Fundamentals of Stein's method, Prob. Surveys 8(1):210-293 arXiv:1109.1880 (2011). 33
- 111. T. Sasamoto, H. Spohn, One-dimensional Kardar-Parisi-Zhang equation: an exact solution and its universality, Physical review letters, 104(23):230602 (2010). 2, 16, 17
- 112. T. Sasamoto, H. Spohn, Exact height distributions for the KPZ equation with narrow wedge initial condition, Nuclear Physics B, 834(3):523-542 (2010). 2, 16, 17
- 113. T. Sasamoto, H. Spohn, The 1 + 1-dimensional Kardar-Parisi-Zhang equation and its universality class, Journal of Statistical Mechanics: Theory and Experiment, P11013(11) (2010). 2, 16, 17
- 114. Y. Shkolnisky, Prolate spheroidal wave functions on a disc integration and approximation of twodimensional bandlimited functions, Appl. Comput. Harmon. Anal. 22(2):235-256 (2007). 56, 57
- 115. B. Simon, OPUC on one foot, Bull. Amer. Math. Soc. 42(4):431-460 (2005). 43
- 116. B. Simon, A comprehensive course in Analysis, Part 4: Operator theory, American Mathematical Soc. (2008). 13
- 117. D. Slepian, H. O. Pollak, Prolate spheroidal wave functions, Fourier analysis and uncertainty I, Bell System Tech. J. 40(1):43-64 (1961). 57
- 118. D. Slepian, Prolate spheroidal wave functions, Fourier analysis and uncertainty IV: Extensions to many dimensions; generalized prolate spheroidal functions, Bell System Tech. J. 43(6):3009-3057 (1964). 4, 56, 57, 58
- 119. B. D. Sutton, *The stochastic operator approach to random matrix theory*, Ph.D. Diss. Massachusetts Institute of Technology (2005). 3
- 120. G. Szegö, Orthogonal Polynomials, Publications of the AMS (1955; 1st ed. 1939). 22, 28, 57
- 121. T. Tao, Topics in random matrix theory, Amer. Math. Soc., vol. 132 (2012). 22
- C. Tracy, H. Widom, Level-spacing distributions and the Airy kernel, Phys. Letters B 305(1-2):115-118 (1993). 2, 6, 30
- 123. C. Tracy, H. Widom, Level-spacing distributions and the Airy kernel, Commun. Math. Phys. 159(1):151-174 (1994). 2, 4, 6, 28, 30
- 124. C. Tracy, H. Widom, On orthogonal and symplectic matrix ensembles, Commun. Math. Phys. 177(3):727-754 (1996). 20, 21
- 125. C.A. Tracy, H. Widom, Integral formulas for the Asymmetric Simple Exclusion Process, Commun. Math. Phys. 279(3):815-844 arXiv:0704.2633 (2008). Erratum: Commun. Math. Phys. 304(3):875-878 (2011). 2, 17
- 126. C.A. Tracy, H. Widom, A Fredholm determinant representation in ASEP, J. Stat. Phys. 132(2):291-300 arXiv:0804.1379 (2008). 2, 17
- 127. C.A. Tracy, H. Widom, Asymptotics in ASEP with step initial condition, Commun. Math. Phys. 290(1):129-154 arXiv:0807.1713 (2009). 2, 17

- 128. L. C. Tsai, Exact lower tail large deviations of the KPZ equation, Duke Math. J. 171(9):1879-1922 arXiv:1809.03410 (2022). 19
- 129. O. Vallée, M. Soares, Airy functions and applications to physics, Imperial College Press, London (2004). 8
- 130. X.-B. Wu, S.-X. Xu, Y.-Q. Zhao, Gaussian Unitary Ensemble with boundary spectrum singularity and σ -form of the Painlevé II equation, Stud. Appl. Math. 140(2):221-251 (**2018**). 30, 31, 59

DEPARTMENT OF ALGEBRA AND LOGIC, INSTITUTE OF MATHEMATICS AND INFORMATICS, BULGAR-IAN ACADEMY OF SCIENCES, ACAD. GEORGI BONCHEV STR., BLOCK 8, 1113 SOFIA (BULGARIA). Email address: yacine.barhoumi@math.bas.bg