Renormalisation in the presence of variance blowup

January 22, 2024

Martin Hairer^{1,2}

¹ École Polytechnique Fédérale de Lausanne, 1015 Lausanne, Switzerland

 $^2\,$ Imperial College London, London SW7 2AZ, United Kingdom

Email: martin.hairer@epfl.ch

Abstract

We show that if one drives the KPZ equation by the derivative of a space-time white noise smoothened out at scale $\varepsilon \ll 1$ and multiplied by $\varepsilon^{3/4}$ then, as $\varepsilon \to 0$, solutions converge to the Cole–Hopf solutions to the KPZ equation driven by space-time white noise.

In the same vein, we also show that if one drives an SDE by fractional Brownian motion with Hurst parameter H < 1/4, smoothened out at scale $\varepsilon \ll 1$ and multiplied by $\varepsilon^{1/4-H}$ then, as $\varepsilon \to 0$, solutions converge to an SDE driven by white noise. The mechanism giving rise to both results is the same, but the proof techniques differ substantially.

MSC2020: 60H10, 60H15, 60L20, 60L30 *Keywords:* KPZ equation, fractional Brownian motion

Contents

1	Introduction		2
	1.1	The KPZ equation	3
	1.2	SDEs driven by fractional noise	4
	1.3	Notations	6
2	Convergence to KPZ		6
	2.1	Convergence of the noise	6
	2.2	Regularity structure and renormalisation	8
	2.3	Convergence of the model	10
3	Fractional Brownian motion 14		
	3.1	Integrations indexed by posets	14
	3.2	Main estimates	18
	3.3	A tool for convergence	23
	3.4	Convergence	26

INTRODUCTION

1 Introduction

The past two decades have seen a wealth of results in the general area of singular stochastic ODEs / parabolic PDEs with the advent of rough paths [Lyo98, Gub04, FH20], regularity structures [Hai14], paracontrolled calculus [GIP15], the rigorous implementation of Wilson's RG and the Polchinski's flow equation [Kup16, Duc22], etc. A common feature of all of these works (as well as their further refinements) is that their domain of validity is cleanly delimited by two main conditions. The first (and physically most relevant) condition is one of scaling subcriticality or, in QFT language, superrenormalisability. This condition, roughly speaking, guarantees that solutions to the equation of interest locally look like Gaussian processes at small scales. When this condition fails to be satisfied, i.e. in the supercritical case, one typically expects any natural approximation to the problem to converge to a "trivial" (often Gaussian) limit as its approximation scale ε is sent to zero. See for example [Frö82, Aiz82] in the context of the Φ^4 models as well as [MU18, GRZ18, DGRZ20] in the context of the KPZ equation. At the boundary, i.e. when the model is *scaling critical*, interesting phenomena can occur but depend on much finer properties of the model. For example the Φ^4 model turns out to be trivial at its critical dimension 4 [ADC21] while the KPZ equation appears to exhibit non-trivial behaviour [BC98, CSZ23b, CSZ23a].

There is however a second somewhat less widely known condition that appears in the above mentioned works as a consequence of the following. These works all exploit the fact that the solution can locally be described by a linear combination of certain multilinear expressions in the driving noise(s). When interpreted "naïvely" some of these expressions blow up as $\varepsilon \to 0$ and need to be renormalised in order to yield finite limits. This renormalisation procedure can be thought of as a way of recentering these expressions in a nonlinear way that is compatible with their algebraic structure. The recentered expressions have vanishing expectations and it is a generic fact that their variances are better behaved and typically converge to a finite limit. In certain cases however (and this only happens when considering equations driven by a noise that is less regular than the corresponding space-time white noise) it may happen that, despite being scaling subcritical, the problem is such that one of these variances also diverges as $\varepsilon \to 0$. Maybe the most prominent example of this phenomenon is given by SDEs driven by independent fractional Brownian motions B_i with Hurst parameter $H < \frac{1}{4}$. In this case, the Lévy area of natural approximations to these driving processes diverges even though, at the analytical level, the theory of rough paths would be applicable. It follows from general principles that it is possible to construct a rough path above B [LV07, NT11, Unt13], but these constructions are non-canonical (not even modulo finitely many parameters) and have pathological properties. In particular, the resulting solution flow $\Phi_{s,t}$ is not measurable with respect to the σ -algebra generated by the increments of B over [s, t].

In this article, we study two examples exploring the situation where subcriticality holds but the variance of one of the stochastic objects associated to our problem

INTRODUCTION

diverges. These are given by the KPZ equation driven by (approximations to) the spatial derivative of space-time white noise and SDEs driven by fractional Brownian motion with $H \leq \frac{1}{4}$. We will see that in both cases the limit is given by an equation in the same class but driven by white noise. See the discussion at the end of this introduction for more details regarding what one would expect in general.

1.1 The KPZ equation

It is by now well known that the KPZ equation can be derived from a large class of interface fluctuation models in the weakly asymmetric regime. The first mathematically rigorous proof of this fact dates back to the seminal work by Bertini and Giacomin [BG97] on fluctuations of the weakly asymmetric simple exclusion process which exploited a number of very specific features of the latter, in particular that it behaves "nicely" under the Cole–Hopf transform which is used there to interpret the KPZ equation itself.

One of the simplest convergence results [Hai13, Hai14, HS17] states that if u solves the PDE

$$\partial_t u = \partial_x^2 u + \sqrt{\varepsilon} (\partial_x u)^2 + \bar{\eta} , \qquad (1.1)$$

for $\bar{\eta}$ a stationary random field with good enough mixing properties, then there exists a constant C_{ε} such that $h_{\varepsilon}(t,x) = \sqrt{\varepsilon}u(t/\varepsilon^2, x/\varepsilon) - C_{\varepsilon}t$ converges as $\varepsilon \to 0$ to the KPZ equation. As one would expect, the variance of the resulting space-time white noise driving the KPZ equation is given by the space-time integral of the variance of $\bar{\eta}$. Furthermore, the leading order of the constant C_{ε} is of the form $c\varepsilon^{-1}$.

This leads naturally (at least from a mathematical perspective) to the question of what happens when the variance of $\bar{\eta}$ has vanishing integral. For example, does (1.1) admit a non-trivial scaling limit when $\bar{\eta}$ is replaced by $\partial_x \bar{\eta}$? Under the same scaling and recentering as above, we know that solutions simply converge to the solution of the deterministic equation $\partial_t h = \partial_x^2 h + (\partial_x h)^2$ but is there a scaling under which they converge to a non-trivial stochastic process? It turns out that this is indeed the case provided that we increase the strength of the nonlinearity and consider a slightly different scaling.

One aim of this article is to show that, if we consider solutions to the equation

$$\partial_t u = \partial_x^2 u + \varepsilon^{1/4} (\partial_x u)^2 + \partial_x \bar{\eta} , \qquad (1.2)$$

and set $h_{\varepsilon}(t, x) = \varepsilon^{1/4} u(t/\varepsilon^2, x/\varepsilon) - C_{\varepsilon}t$ (note the difference in scaling from the one that keeps the stochastic heat equation invariant!), then a suitable choice of C_{ε} leads again to the convergence of h_{ε} to solutions to the KPZ equation. The constant itself also behaves slightly differently: to leading order it now scales like $C_{\varepsilon} \sim \varepsilon^{-3/2}$.

After rescaling, (1.2) can be written as

$$\partial_t h_{\varepsilon} = \partial_x^2 h_{\varepsilon} + (\partial_x h_{\varepsilon})^2 + \varepsilon^{3/4} \partial_x \eta_{\varepsilon} - C_{\varepsilon} , \qquad (1.3)$$

where η_{ε} is scaled so that it converges to a white noise. In order to simplify notations and to avoid technical complications we fix a space-time white noise η on $S^1 \times \mathbf{R}$ and we assume that there exists a smooth function $\rho \colon \mathbf{R}^2 \to \mathbf{R}$ supported in the unit ball such that $\eta_{\varepsilon} = \rho_{\varepsilon} \star \eta$ where $\rho_{\varepsilon}(z) = \varepsilon^{-3}\rho(\varepsilon^{-1}z)$ and \star denotes space-time convolution. We then consider (1.3) as a Cauchy problem in $\mathscr{C}^{\alpha}(S^1)$ for some $\alpha \in (0, 1/2)$, in particular we restrict ourselves to a compact spatial domain with periodic boundary conditions. With these notations at hand, we have the following result, the proof of which will be provided in Section 2.3.

Theorem 1.1. Let $\alpha \in (0, 1/2)$, $\beta \in (0, \alpha \wedge 1/4)$, and let $h_0 \in \mathscr{C}^{\alpha}(S^1)$. Then, there exists $\sigma > 0$ and a choice of constants C_{ε} such that the solution h_{ε} to (1.3) converges in law in $\mathscr{C}(\mathbf{R}_+, \mathscr{C}^{\beta}(S^1))$ to the solution to the Cole–Hopf solution to the KPZ equation driven by space-time white noise with variance σ^2 .

A surprising feature of this result is that the limiting KPZ equation is driven by *space-time white noise* while $\partial_x \eta$, suitably rescaled, converges to the spatial derivative of such a noise. So where does this driving noise come from? The basic observation is that, writing P for the heat kernel and \star for space-time convolution, $(\partial_x P \star \partial_x \eta)^2$ suitably rescaled and recentered converges weakly to a space-time white noise and it is this noise that drives the limiting KPZ equation. Furthermore, asymptotically as $\varepsilon \to 0$, this driving noise is actually *independent* of the original field η in the sense of stable convergence [JSo₃, HL₁₅].

1.2 SDEs driven by fractional noise

We also show that a similar phenomenon arises when considering SDEs driven by fractional Brownian motion with Hurst parameter $H \leq 1/4$. Let $\{B_i\}_{i=1}^m$ be i.i.d. copies of a fractional Brownian motion with some fixed Hurst parameter $H \leq 1/4$ and let B_i^{ε} be their convolution with a mollifier at scale ε . Let furthermore V_i be m vector fields on \mathbf{R}^d (this could be replaced by a manifold) and let x^{ε} be the solution to the controlled differential equation

$$dx^{\varepsilon} = C_{\varepsilon} \sum_{i=1}^{m} V_i(x^{\varepsilon}) \, dB_i^{\varepsilon} , \qquad (1.4)$$

where we set $C_{\varepsilon} = \varepsilon^{1/4-H}$ when H < 1/4 and $C_{\varepsilon} = |\log \varepsilon|^{-1/2}$ when H = 1/4. We then have the following convergence result.

Theorem 1.2. There exists a finite constant $\sigma > 0$ and independent standard Wiener processes W_{ij} such that x^{ε} converges in law to the diffusion process xsolving

$$dx = \frac{\sigma}{2} \sum_{i < j} [V_i, V_j](x) \circ dW_{ij} . \qquad (1.5)$$

Remark 1.3. *Convergence takes place uniformly in time until the first time solutions* to (1.5) get large, in particular we do not need to assume global well-posedness. See Theorem 3.13 below for a precise formulation taking topologies into account.

INTRODUCTION

Remark 1.4. It was shown in [Unto8] that the Lévy area of two independent fractional Brownian motions with $H < \frac{1}{4}$ converges to an independent Wiener process, which strongly hints at Theorem 1.2 but is far from sufficient to prove it. A result with a somewhat similar flavour (a Brownian motion independent of the original fractional Brownian motion showing up as a second-order process) was previously obtained when H = 1/4 in [NR09, BS10].

The proof of these two results suggests the following general mechanism. Take a class of SPDEs (or SDEs) driven by noise of regularity $\alpha < 0$. What we mean here by a "class" is that we fix a rule in the sense of [BHZ19] as well as an ambient space(-time) dimension and consider all equations that can be formulated within the corresponding regularity structure. One the one hand, there then exists some α_c such that the problem is locally subcritical (in the sense of [Hai14]) when $\alpha > \alpha_c$ and fails to be so when $\alpha < \alpha_c$. On the other hand, there exists α_v such that the third condition of [CH16, Thm 2.15] is satisfied for $\alpha > \alpha_v$ and fails for $\alpha < \alpha_v$. As already mentioned, the condition $\alpha > \alpha_v$ is crucial for guaranteeing the convergence of the BPHZ lift of a regularised driving noise. When it fails, the *variance* of some of the stochastic objects relevant in the description of the solution blows up, which cannot be controlled by the addition of counterterms in the equation (these can only counteract the divergence of *expectations*).

What our result shows is that if we are in a situation where $\alpha_c < \alpha_v$ and we choose as driving noise a mollification at scale ε of a noise that is self-similar with some exponent $\beta < \alpha_v$, then we obtain a non-trivial limit driven by (space-time) white noise within the same class of equations, provided that we multiply our driving noise with $\varepsilon^{\alpha_v - \beta}$ (or a suitable negative power of $|\log \varepsilon|$ if $\beta = \alpha_v$). Indeed in the example of the KPZ equation, one has $\alpha_c = -2$ and $\alpha_v = -7/4$. Our driving noise is a mollification of the derivative of space-time white noise which is self-similar with exponent -5/2, so that this heuristic suggests that we multiply it by $\varepsilon^{3/4}$ (since 3/4 = -7/4 + 5/2) in order to obtain a non-trivial limit. In the case of SDEs/RDEs one has $\alpha_c = -1$ and $\alpha_v = -3/4$ while our driving noise has self-similarity exponent $\beta = H - 1$ (the driving noise is the derivative of our fractional Brownian motion), which again leads to the correct prediction.

Note that there is in general no particular ordering between α_c and α_v , especially since the latter is dimension-dependent while the former is not. In the case of the Φ^4 equation, one has for example $\alpha_c = -3$ and $\alpha_v = -(d + 10)/4$ so that the phenomenon described in this article is only expected to take place in spatial dimension d = 1.

One point that the reader should keep in mind is that, depending on how one interprets an equation, one may be lead to descriptions of its solutions using different regularity structures which could potentially lead to different values for α_v . For example, when writing the Navier–Stokes nonlinearity in its usual form as $(u \cdot \nabla)u$, one is lead to $\alpha_v = -(d+8)/4$. If however we remember that, thanks to the divergence free condition, it can equivalently be written as $\operatorname{div}(u \otimes u)$, we find that one actually has $\alpha_v = -(d+10)/4$. In other words, while α_v determines the

noise regularity at which the variance of some stochastic object *may* diverge, it does not guarantee that it *will* diverge since there could be additional problem-specific cancellations.

1.3 Notations

We will always use parabolic scalings so that, given $z = (t, x) \in \mathbf{R}^2$ and $\lambda \in \mathbf{R}$, we write

$$\lambda z = (\lambda^2 t, \lambda x)$$
, $|z| = \sqrt{|t|} + |x|$.

Test functions φ are always smooth, supported in the unit ball, and such that $\sup_{z} |D^{k}\varphi(z)| \leq 1$ for all $|k| \leq 10$ (say). As usual, we denote its rescaled translates by $\varphi_{z}^{\lambda}(z_{1}) = \lambda^{-3}\varphi(\lambda^{-1}(z_{1}-z))$.

2 Convergence to KPZ

Writing $h = u - \varepsilon^{3/4} P \star \partial_x \eta_{\varepsilon}$, setting ξ_{ε} as above, and setting

$$\chi_{\varepsilon} = \varepsilon^{3/4} \partial_x P \star \partial_x \eta_{\varepsilon} = \varepsilon^{3/4} \partial_x^2 P \star \rho_{\varepsilon} \star \eta ,$$

we can rewrite this equation suggestively as

$$\partial_t h = \partial_x^2 h + (\partial_x h)^2 + 2\chi_{\varepsilon} \partial_x h + \xi_{\varepsilon} . \qquad (2.1)$$

At the intuitive level, our main result then follows from Proposition 2.2 combined with the fact that $\chi_{\varepsilon} \to 0$. This is of course far from a proof since $\varepsilon^{3/4} \partial_x \eta_{\varepsilon} \to 0$, so essentially the same argument would suggest that (1.2) converges to a deterministic limit!

The main reason why the formal argument works in the case of (2.1) but fails in the case of (1.3) is that if we apply the methodology of [Hai14, CH16, BHZ20, BCCH21] to (1.3) (using the fact that, uniformly over ε , one has $\varepsilon^{3/4} \partial_x \eta_{\varepsilon} \in \mathscr{C}^{-7/4^-}$ with parabolic scaling but no better), then we end up generating a symbol containing two instances of the noise, but with degree $-3/2^-$, thus (just about) failing to satisfy the third condition of [CH16, Thm 2.15]. In fact, an analogous condition first appeared in the context of rough paths in [CQ00] and also appears for example in [LOTT22, Ass. 2.1] and the second part of [HS23, Ass 2.31]. When applying it to (2.1) on the other hand, the "worst" symbol with more than one noise that arises corresponds to $\chi_{\varepsilon} \cdot (\partial_x P \star \xi_{\varepsilon})$ and is of degree $-5/4^- > -3/2$.

2.1 Convergence of the noise

We use graphical notations similar to those of [Hai14, HS17]. In particular, we use plain nodes • for integration variables, a coloured node • for a variable taking the fixed value 0, and \bigcirc for an integration variable at which we furthermore evaluate an instance of the white noise η . We group multiple white noises together like so \bigcirc to indicate that their product should be interpreted as a Wick product.

As usual, all of our variables are space-time variables. Regarding edges (which represent kernels evaluated at the difference between the two variables represented

by the nodes they connect), we write $\cdots \cdots \cdots$ for the mollifier ρ_{ε} , \longrightarrow for a test function $\varphi^{\lambda}(t, x)$ centred at the origin and rescaled at some scale λ (if no scale is specified one takes $\lambda = 1$, different test functions are drawn with different colors), and \longrightarrow for the kernel $(t, x) \mapsto \varepsilon^{3/4} \partial_x^2 P_t(x)$. In particular, this graphical notation yields

$$\langle \eta_{\varepsilon}, \varphi \rangle = \langle 0, \langle \chi_{\varepsilon}, \varphi \rangle = \langle 0, \langle \chi_$$

An important role will be played by the covariance K_{ε} of χ_{ε} as well as the correlation between η_{ε} and χ_{ε} , so we introduce the shorthand graphical notations

 $K_{\varepsilon} = \mathsf{weight} = \longleftarrow \bullet \mathsf{weight} \bullet \mathsf{weight} \to \mathsf{weight} \bullet \mathsf{weight} \to \mathsf{weight} \bullet \mathsf{weight} \bullet \mathsf{weight} \to \mathsf{weight} \bullet \mathsf{we$

The following is a straightforward consequence of the scaling properties of ρ_{ε} and the heat kernel.

Lemma 2.1. One has $K_{\varepsilon}(z) = \varepsilon^{-3/2} K(z/\varepsilon)$ for a smooth function K satisfying the bounds $|K(z)| \leq (1+|z|)^{-3}$. Similarly, one has $\bar{K}_{\varepsilon}(z) = \varepsilon^{-9/4} \bar{K}(z/\varepsilon)$ for a smooth function \bar{K} satisfying the bounds $|\bar{K}(z)| \leq (1+|z|)^{-3}$.

Proposition 2.2. Let $\xi_{\varepsilon} = \varepsilon^{3/2} (\partial_x P \star \partial_x \eta_{\varepsilon})^2 - C_{\varepsilon}$. Then, there exists $\sigma > 0$ such that ξ_{ε} converges stably in \mathcal{C}^{α} for any $\alpha < -3/2$ to a space-time white noise ξ with variance σ^2 independent of η .

Proof. To show stable convergence, it suffices by [HL15, Prop. 3.4 (c)] to show that the pair $(\xi_{\varepsilon}, \eta_{\varepsilon})$ converges weakly in $\mathscr{C}^{\alpha} \times \mathscr{C}^{\alpha}$ to a pair of independent white noises.

We first note that ξ_{ε} and η_{ε} are decorrelated and that one has

$$\left|\mathbf{E}\langle\varphi_{z}^{\lambda},\xi_{\varepsilon}\rangle^{2}\right| = 2\left|\int\varphi_{z}^{\lambda}(z_{1})\varphi_{z}^{\lambda}(z_{2})K_{\varepsilon}^{2}(z_{1}-z_{2})\,dz_{1}\,dz_{2}\right|\,.$$

Since φ_z^{λ} is supported in a parabolic ball of radius λ , this integral is at most of order $\lambda^3 \|\varphi^{\lambda}\|_{L^{\infty}}^2 \|K_{\varepsilon}^2\|_{L^1} \approx \lambda^{-3}$. Since a similar bound holds for η_{ε} , equivalence of moments for elements in a fixed Wiener chaos combined with [FM17, Thm 1.1] yields tightness in $\mathscr{C}^{\alpha} \times \mathscr{C}^{\alpha}$.

Since K_{ε}^2 converges to some multiple of a Dirac mass, it immediately follows similarly to above that there exists $\sigma > 0$ such that

$$\lim_{\varepsilon \to 0} \mathbf{E} \langle \varphi, \xi_{\varepsilon} \rangle \langle \xi_{\varepsilon}, \psi \rangle = \sigma^2 \langle \varphi, \psi \rangle_{L^2} ,$$

and similarly for η_{ε} , for any continuous compactly supported test functions φ and ψ .

It therefore remains to show that any collection of random variables of the form $\xi_{\varepsilon}(\varphi)$ and/or $\eta_{\varepsilon}(\psi)$ converges to a jointly Gaussian limit as $\varepsilon \to 0$. For this, we

use the Nualart–Peccati fourth moment theorem [NPo5] which implies that this is the case provided that the fourth joint cumulant of any such collection of variables converges to 0 as $\varepsilon \to 0$. It follows from the diagram formula [PT11, Thm 7.1.3] that the joint cumulant $\mathbf{E}_c(\langle \varphi_1, \xi_{\varepsilon} \rangle, \ldots, \langle \varphi_4, \xi_{\varepsilon} \rangle)$ is obtained by taking four copies of the figure associated to $\langle \xi_{\varepsilon}, \varphi \rangle$ in (2.2) and summing over all ways of pairwise contracting the white noises, subject to the following constraints:

- No two noises grouped into a Wick product can be contracted.
- The resulting diagram is connected.

Clearly, the only contraction satisfying these constraints is given by



modulo permutations of the four test functions. The fact that such a term converges to 0 as $\varepsilon \to 0$ is then a consequence of Weinstein's theorem [Hai18, Prop. 2.3], noting that Lemma 2.1 implies that, for $\kappa \in [0, 3/2)$, the kernel K_{ε} is of degree $-3/2 - \kappa$ and the corresponding norm is of order ε^{κ} .

2.2 Regularity structure and renormalisation

We now describe the regularity structure associated to (2.1) by the procedure of [BHZ19] (see also [FH20, Sec. 15.2]) and we argue that the BPHZ lift of our noise only leads to constant counterterms (as opposed to additional counterterms of the type $C_{\varepsilon}\partial_x h$ which is the only other kind of counterterm that can possibly appear as a consequence of the general result of [BCCH21]).

In the language of [BHZ19], the regularity structure $(\mathcal{T}, \mathcal{G})$ in question is generated by one integration operator \mathcal{F} of regularity 2, two noises Ξ and H of respective regularities $-3/2 - \kappa$ and $-3/4 - \kappa$ for some positive $\kappa > 0$, and the normal complete rule containing the nodes $H\mathcal{F}'$ and $(\mathcal{F}')^2$. It is straightforward to verify that this rule is subcritical and also satisfies [HS23, Ass 2.31] (at least for κ small enough and we henceforth fix such a value). For the sake of intuition it helps to represent the canonical basis vectors of \mathcal{T} by trees with possible nodes given by

$$\P \simeq \Xi \;, \quad \P \simeq H \;, \quad \flat \simeq H \mathcal{J}' \;, \quad \flat \simeq \mathcal{J}' \;, \quad \curlyvee \simeq (\mathcal{J}')^2 \;. \tag{2.3}$$

We will use the convention that \mathcal{F}' annihilates Taylor monomials, so that we allow no node of type $1 \simeq \mathbf{1}$ except for the symbol 1 itself. For example, we have

$$\mathcal{J}'(H\mathcal{J}'(\Xi))\mathcal{J}'(H) = \diamondsuit{\rho}, \qquad (2.4)$$

which contains nodes of all the above types except the penultimate one and is of degree $3 + 2(-3/4 - \kappa) - 3/2 - \kappa = -3\kappa$. The prescription of [BHZ19] also requires us to include basis vectors in \mathcal{T} corresponding to the classical Taylor monomials X^k , as well as trees with additional monomials attached at their nodes.

The latter however can be completely disregarded if we restrict ourselves to solutions of a low enough regularity γ for which (2.1) is well-posed in a space of modelled distributions that are locally of class \mathfrak{D}^{γ} ($\gamma = 3/2 + 2\kappa$ will do provided that κ is small enough).

To see what degrees can possibly appear, note that each node in (2.3) has one incoming edge (at the bottom) and between zero and two outgoing edges. Furthermore, we should think of each basis vector of \mathcal{T} as represented by a tree which has one incoming edge (at the root) and no outgoing edge. As a consequence, each symbol (except 1) must contain at least one node of type \mathbb{T} or \mathbb{P} and, for each additional such node, one node of type \bigvee . As a consequence, the possible degrees that can be realised are

$$N_{\mathbf{y}} + 2N_{\mathbf{y}} - N_{\mathbf{z}}(\frac{3}{2} + \kappa) - N_{\mathbf{0}}(\frac{3}{4} + \kappa) + N_{\mathbf{y}}(\frac{1}{4} - \kappa) \;, \quad N_{\tau} \in \mathbf{N} \;, \quad N_{\mathbf{z}} + N_{\mathbf{0}} = 1 + N_{\mathbf{y}} = 0 \;.$$

Note that, using the second relation, we find that the degree of a tree τ is given by

$$\deg \tau = N_{\gamma} + N_{\Box}(\frac{1}{2} - \kappa) + N_{\Diamond}(\frac{5}{4} - \kappa) + N_{\Diamond}(\frac{1}{4} - \kappa) - 2.$$
 (2.5)

In the case of (2.4) for example one has $N_{\square} = N_{\bigcirc} = N_{\bigcirc} = N_{\curlyvee} = 1$ and $N_{\curlyvee} = 0$.

Fix now once and for all a truncation \hat{K} of the heat kernel, i.e. $\hat{K} \colon \mathbb{R}^2 \to \mathbb{R}$ is such that $\hat{K} = P$ in a ball of radius $\frac{1}{2}$ around the origin, $\hat{K} = 0$ outside of a ball of radius 1, and $\int \hat{K}(z) dz = 0$. We furthermore assume that \hat{K} is symmetric under spatial reflections, namely that

$$\tilde{K}(t, -x) = \tilde{K}(t, x) .$$

We then write $Z^{(\varepsilon)} = (\Pi^{(\varepsilon)}, \Gamma^{(\varepsilon)})$ for the canonical lift of the noise $(\eta_{\varepsilon}, \xi_{\varepsilon})$ into the regularity structure \mathcal{T} . Recall that this is built by first inductively defining a linear map $\Pi^{(\varepsilon)}$: $\mathcal{T} \to \mathscr{C}^{\infty}$ by setting $\Pi^{(\varepsilon)} 1 = 1$, $\Pi^{(\varepsilon)} = \xi_{\varepsilon}$, $\Pi^{(\varepsilon)} \odot = \eta_{\varepsilon}$, and then $\Pi^{(\varepsilon)} \mathcal{F}'(\tau) = \partial_x \hat{K} \star \Pi^{(\varepsilon)} \tau$, $\Pi^{(\varepsilon)} \bar{\tau} \tau = \Pi^{(\varepsilon)} \bar{\tau} \cdot \Pi^{(\varepsilon)} \tau$. One then defines $Z^{(\varepsilon)}$ by a suitable "recentering" procedure as in [BHZ19, Def. 6.8].

Note that each $\mathbf{\Pi}^{(\varepsilon)} \tau$ is a stationary process and we set

$$C_{\tau}^{(\varepsilon)} = \mathbf{E} \mathbf{\Pi}^{(\varepsilon)} \tau(0)$$

see [BHZ19, Eq. 6.24]. We then have the following elementary result.

Proposition 2.3. If τ is such that either $N_H^{\tau} \stackrel{\text{def}}{=} N_{\circ} + N_{\circ}$ is odd or $N_{\mathcal{F}'}^{\tau} \stackrel{\text{def}}{=} N_{\circ} + 2N_{\vee} + N_{\circ}$ is odd, then $C_{\tau}^{(\varepsilon)} = 0$.

Proof. Since ξ_{ε} is an element of the *second* (homogeneous) Wiener chaos while η_{ε} is an element of the first chaos, it follows that, whenever N_{\circ} is odd, $\Pi^{(\varepsilon)}\tau$ is a random variable belong to a sum of odd chaoses, so that its expectation vanishes.

If N_{γ} is odd then $C_{\tau}^{(\varepsilon)}$ is given by the expectation of an integral involving an odd number of instances of the kernel $\partial_x \hat{K}$. Since the latter is odd under spatial reflections but the covariances of ξ_{ε} and η_{ε} are even, it follows from reflecting all integration variables that $C_{\tau}^{(\varepsilon)} = -C_{\tau}^{(\varepsilon)}$ as claimed.

Note now that the actual renormalisation constants $\hat{C}_{\tau}^{(\varepsilon)}$ appearing in the BPHZ lift of the model (and therefore also multiplying the counterterms associated to each tree τ of negative degree) are obtained by precomposing $C^{(\varepsilon)}$ with the "negative twisted antipode" as given in [BHZ19, Prop. 6.6]. We only need to know that these are given by polynomial expressions in the $C_{\tau}^{(\varepsilon)}$ that have the property that every monomial of the form $\prod C_{\tau_i}^{(\varepsilon)}$ appearing in the definition of $\hat{C}_{\tau}^{(\varepsilon)}$ satisfies $N_H^{\tau} = \sum_i N_H^{\tau_i}$ and $N_{\mathcal{F}'}^{\tau} = \sum_i N_{\mathcal{F}'}^{\tau_i}$ (and similarly for N_{\square}). This is an immediate consequence of the definition [BHZ19, Eq. 6.25] and the definition of the "twisted antipode" in Equations 6.8–6.9 of that article. Since any sum of integers adding to an odd number must contain at least one odd summand this implies the following.

Corollary 2.4. Under the assumption of Proposition 2.3 one also has $\hat{C}_{\tau}^{(\varepsilon)} = 0$.

Recall now that, when considering (2.1) driven by the BPHZ lift of $(\eta_{\varepsilon}, \xi_{\varepsilon})$, each tree τ with deg $\tau < 0$ leads to an additional counterterm in the right-hand side of the form $\hat{C}_{\tau}^{(\varepsilon)}F_{\tau}(\partial_x h)$. We now claim that, for every such τ , one has either $F_{\tau}(u) \propto u$ or $F_{\tau}(u) \propto 1$ and, furthermore, that one has $\hat{C}_{\tau}^{(\varepsilon)} = 0$ for those τ 's such that $F_{\tau}(u) \propto u$.

Indeed, one can see from [BCCH21] (see Equation 2.12 there for the definition of the counterterms $\Upsilon^F[\tau]$, as well as Equation 3.9 and Theorem 3.25 for a description of how this relates to the renormalised equation) that, given τ , every node of τ of type \degree or \checkmark leads to a factor $\partial_x h$ while every other node leads to a constant factor, so that $F_{\tau}(u) \propto u^{N_0+N_1}$. If τ is a tree such that $N_0 + N_{\uparrow} \ge 2$, then it follows from (2.5) combined with the fact that one has the constraint $N_{\square} + N_0 \ge 1$ that deg $\tau \ge \frac{9}{4} - \kappa > 0$, so that counterterms of the type $(\partial_x h)^2$ (and higher) are indeed ruled out. The main claim of this section is the following.

Proposition 2.5. The BPHZ counterterm of (2.1) is given by $\sum_{\tau: \deg \tau < 0} \hat{C}_{\tau}^{(\varepsilon)}$.

Proof. We only need to show that $\hat{C}_{\tau}^{(\varepsilon)} = 0$ whenever τ is such that $N_{\circ} + N_{\downarrow} = 1$ and deg $\tau < 0$. Since we want to apply the criterion from Corollary 2.4, it is enough to show that if τ is such that its conditions fail (i.e. $N_{\circ} + N_{\flat}$ and $N_{\downarrow} + N_{\flat}$ are both even) and $N_{\circ} + N_{\flat} = 1$ then deg $\tau > 0$.

Let us first consider the case $N_{\gamma} = 1$. This then implies that N_{γ} is odd (and in particular at least 1) so that N_{\circ} is also odd, which leads to a contradiction with the constraint $N_{\circ} + N_{\gamma} = 1$. If $N_{\circ} = 1$, then the same argument again leads to a contradiction, thus completing the proof.

2.3 Convergence of the model

In view of the results of the previous section and the results of [Hai14, BHZ19, BCCH21], it remains to show that, in the context of a suitable regularity structure associated to (2.1), the BPHZ lift of the noises ($\chi_{\varepsilon}, \xi_{\varepsilon}$) converges to the BPHZ lift of (0, ξ) with $\xi = \lim_{\varepsilon \to 0} \xi_{\varepsilon}$ a space-time white noise.

As in [HS23], we now introduce a family of modelled distributions that will describe the derivative of our model with respect to the noise and that will be used

to inductively bound the BPHZ lift of $(\chi_{\varepsilon}, \xi_{\varepsilon})$. The only difference is that the pair $(\chi_{\varepsilon}, \xi_{\varepsilon})$ by itself doesn't appear to satisfy a spectral gap inequality. The underlying Gaussian noise η however does of course, so this is what we will be exploiting. As a consequence, our setup is slightly different from that of [HS23]. In particular, we will work with L^p -based spaces for p slightly smaller than 2 instead of the L^2 -based spaces used in [HS23, Secs 4 & 5]. Since the analytic core of that article (Section 3.2) allows for arbitrary p, this does not cause any difficulty.

Let now γ_{τ} and $\deg_2 \tau$ be defined as in [HS23, Sec. 4], and fix some p < 2(we will have to choose it very close to 2 later on, for example $p = 2 - 10^{-3}$ will do). Given any $h \in L^2$, we then define a collection of modelled distributions $H^{z,h}_{\tau;\varepsilon} \in \mathfrak{D}_p^{\gamma_{\tau},\deg_2\tau;z}$ for the canonical basis vectors τ of \mathcal{T} inductively by setting

$$H^{z,h}_{X^{k};\varepsilon} = 0 , \quad H^{z,h}_{H;\varepsilon} = \varepsilon^{3/4} (\partial_x^2 P \star \rho_{\varepsilon} \star h) \mathbf{1} , \quad H^{z,h}_{\Xi;\varepsilon} = 0 , \qquad (2.6)$$

as well as

$$H^{z,h}_{\tau\bar{\tau};\varepsilon} = H^{z,h}_{\tau;\varepsilon} f^z_{\bar{\tau}} + f^z_{\tau} H^{z,h}_{\tau\bar{\tau};\varepsilon} , \quad H^{z,h}_{\mathcal{F}\tau;\varepsilon} = \mathscr{K}^{x,p}_{\gamma_{\tau},\deg_2\tau} H^{z,h}_{\tau;\varepsilon} .$$
(2.7)

(See [HS23, Sec. 4] for more details.) As in [HS23], the second identity of (2.7) only makes sense a priori for $\gamma_{\tau} > 0$. This fails precisely when $\tau = \Xi$ (and only then).

In that case, we *postulate* that the reconstruction of $H^{z,h}_{\Xi,\varepsilon}$ is given by

$$\mathscr{R}H^{z,h}_{\Xi;\varepsilon} = 2\varepsilon^{3/4}\chi_{\varepsilon}(\partial_x^2 P \star \rho_{\varepsilon} \star h) , \qquad (2.8)$$

and we define $H^{z,h}_{\mathcal{J}\Xi;\varepsilon}$ using [HS23, Thm 3.21]. The following analogue to [HS23, Lems 4.3 & 4.5] provides the starting point for our induction.

Lemma 2.6. For any $\alpha < 3/4$ and $p \leq 2$, we have $H_{H;\varepsilon}^{z,h} \in \mathfrak{D}_p^{\alpha,\alpha;z}$ uniformly over $h \in L^2$ and there exist $\kappa > 0$ and C > 0 (depending on α) such that

$$\sup_{h \in L^2} \| H_{H;\varepsilon}^{z,h} \|_{p,\alpha,\alpha;z} \le C \varepsilon^{\kappa} , \qquad \forall \varepsilon \in (0,1] .$$

Furthermore, the right-hand side of (2.8) is almost surely a candidate for the pointed reconstruction in the sense of [HS23, Def. 3.20] of $H^{z,h}_{\Xi;\varepsilon} = 0$ viewed as an element of $\mathfrak{D}_p^{\gamma_{\tau}, \deg_2 \tau; z}$. Writing $N_{\varepsilon}(h; z)$ for the corresponding "norm" (the constant $C(f; B_z)$ appearing in that definition), one has

$$\sup_{\varepsilon \leq 1} \mathbf{E} \Big(\sup_{\|h\|=1} \sup_{z \in K} N_{\varepsilon}(h; z) \Big)^q < \infty ,$$

for all $q \geq 1$.

Proof. Regarding the first claim, it follows immediately from the definition of local Besov spaces (in 1 + 1-dimensional parabolic space-time) [HS23, Sec. 1.3] that

maps of the type $f \mapsto \varepsilon^{\alpha} \rho_{\varepsilon} \star f$ (with $\alpha \geq 0$) are uniformly bounded over $\varepsilon \in (0, 1]$ from L^p into the local Besov space \mathscr{B}_p^{α} .

Furthermore, as a consequence of [BL22, Prop. A.5], \mathscr{B}_p^{α} embeds canonically into \mathfrak{D}_p^{α} and, provided that $\alpha < 3/p$, the space \mathfrak{D}_p^{α} (with values in the polynomial sector) coincides with $\mathfrak{D}_p^{\alpha,\alpha;z}$. This latter statement can be shown in exactly the same way as [HS23, Lem. 4.3].

The second claim follows in an analogous way once we note that $\mathbf{E}\chi^2_{\varepsilon}(z) \approx \varepsilon^{-3/2}$ so that $\varepsilon^{3/4}\chi_{\varepsilon}$ belongs to every local L^p space and admits moments of all orders.

At the technical level, the main result of this section is the following.

Theorem 2.7. Let $\hat{Z}^{(\varepsilon)}$ denote the BPHZ lift of $(\xi_{\varepsilon}, \chi_{\varepsilon})$. Then, one has the convergence in law $\hat{Z}^{(\varepsilon)} \to Z$, where Z denotes the BPHZ lift of $(\xi, 0)$ with ξ as in Proposition 2.2.

Proof. We first show that $\hat{Z}^{(\varepsilon)}$ is bounded uniformly over $\varepsilon \in (0, 1]$. The proof by induction is virtually identical to that of [HS23, Thm 2.33], the only difference being the starting point for our induction. The bounds of Lemma 2.6 provide the starting point for an inductive proof that, for every $\tau \in \mathcal{T}$ there exists $k \ge 1$ and, for every $z \in \mathbb{R}^2$, there exists a compact set \mathfrak{K} containing z such that

$$\sup_{\|h\| \le 1} \sup_{\varepsilon \le 1} \|H_{\tau;\varepsilon}^{z,h}\|_{p,\gamma_{\tau},\deg_{2}\tau;z} \lesssim (1+\|\hat{Z}^{(\varepsilon)}\|_{\mathfrak{T}_{\tau};\mathfrak{K}})^{k} .$$

$$(2.9)$$

Here, γ_{τ} and deg₂ τ are defined as in [HS23, Sec. 4] and \mathcal{T}_{τ} denotes the smallest sector of \mathcal{T} such that \mathcal{T}_{τ} + span τ is also a sector. The proof of this statement is identical to that of [HS23, Prop. 4.7]. In fact, it is easier since $\tau = \Xi$ is the only element with $\gamma_{\tau} \leq 0$, so that we never need to use [HS23, Lem. 4.6] which was one of the more delicate points in that proof.

The definitions (2.6) and (2.8) guarantee that one has the identities

$$D_h \Pi_z^{(\varepsilon)} \Xi = \Re H_{\Xi;\varepsilon}^{z,h} , \qquad D_h \Pi_z^{(\varepsilon)} H = \Re H_{H;\varepsilon}^{z,h} ,$$

where D denotes the Fréchet derivative with respect to the underlying space-time white noise η in the direction $h \in L^2$. This again provides the starting point for the inductive proof that the identity $D_h \Pi_z^{(\varepsilon)} \tau = \Re H_{\tau;\varepsilon}^{z,h}$ holds for every $\tau \in \mathcal{T}$, the proof being identical to that of [HS23, Thm 4.18]. Combining these two facts with the Gaussian spectral gap inequality then yields the uniform bound on $\hat{Z}^{(\varepsilon)}$ exactly as in [HS23, Thm 2.33].

Regarding the convergence as $\varepsilon \to 0$, we again argue in a way similar to [HS23], but with significant simplifications and one additional layer of approximation. For $\delta > 0$, we define

$$\xi_{\varepsilon,\delta} = \rho_{\delta} \star \xi_{\varepsilon} , \qquad \chi_{\varepsilon,\delta} = \rho_{\delta} \star \chi_{\varepsilon} ,$$

and we denote by $\hat{Z}^{(\varepsilon,\delta)}$ the BPHZ lift of these processes. As a consequence of Proposition 2.2, of [HS23, Thm 2.33] (using the fact that ξ satisfies a spectral gap

inequality in L^2) and of the fact that all these processes are smooth, we know that

$$\lim_{\delta \to 0} \lim_{\varepsilon \to 0} \hat{Z}^{(\varepsilon,\delta)} = Z , \qquad \lim_{\delta \to 0} \hat{Z}^{(\varepsilon,\delta)} = \hat{Z}^{(\varepsilon)} ,$$

so our aim is to exchange the limits in the first bound. This is immediate if we can get a bound on the distance between $\hat{Z}^{(\varepsilon,\delta)}$ and $\hat{Z}^{(\varepsilon)}$ that goes to 0 as $\delta \to 0$, uniformly over $\varepsilon \in (0, 1]$. The proof that this is the case is virtually identical to that of [HS23, Thm 6.9], the starting point of the argument being provided by the fact that if we set

$$H^{z,h}_{H;\varepsilon,\delta} = \varepsilon^{3/4} (\rho_{\delta} \star \partial_x^2 P \star \rho_{\varepsilon} \star h) \mathbf{1} \, ,$$

as well as

$$H^{z,h}_{\Xi;\varepsilon,\delta} = 0 , \qquad \Re H^{z,h}_{\Xi;\varepsilon,\delta} = 2\varepsilon^{3/4} \rho_{\delta} \star \left(\chi_{\varepsilon} (\partial_x^2 P \star \rho_{\varepsilon} \star h) \right) ,$$

then both $\sup_{h \in L^2} ||| H^{z,h}_{H;\varepsilon} - H^{z,h}_{H;\varepsilon,\delta} ||_{p,\alpha,\alpha;z}$ and the norm of $\Re H^{z,h}_{\Xi;\varepsilon} - \Re H^{z,h}_{\Xi;\varepsilon,\delta}$ converge to 0 as $\delta \to 0$, uniformly over $\varepsilon \in (0,1]$. One then defines $H^{z,h}_{\tau;\varepsilon,\delta}$ in the same way as $H^{z,h}_{\tau;\varepsilon}$, but with all manipulations done with respect to the model given by the BPHZ lift of $(\xi_{\varepsilon,\delta}, \chi_{\varepsilon,\delta})$. By repeatedly applying [HS23, Thms 3.11, 3.19, 3.21], it is then straightforward to obtain the bound

$$\|\!\| H^{z,h}_{\tau;\varepsilon}; H^{z,h}_{\tau;\varepsilon,\delta} \|\!|_{p,\gamma\tau,\deg_2\tau;z} \lesssim o(\delta)(1+\|\hat{Z}^{(\varepsilon)}\|_{\mathcal{T}_{\tau};\mathfrak{K}}+\|\hat{Z}^{(\varepsilon)};\hat{Z}^{(\varepsilon,\delta)}\|_{\mathcal{T}_{\tau};\mathfrak{K}})^k \;,$$

with quantifiers over h, τ, z , etc as in (2.9), and o some function with $\lim_{\delta \to 0} o(\delta) = 0$. The reason why these theorems are sufficient in our case is again that $\gamma_{\tau} > 0$ for all $\tau \neq \Xi$, so we never have coefficients of negative regularity appearing in our modelled distributions.

We now have all the ingredients in place to provide a proof of the first main theorem of this article.

Proof of Theorem 1.1. Recall the construction in [CCHS22, Sec. 1.5.1] which, given an unbounded complete separable metric space X with a distinguished "origin" $o \in X$, constructs a space X^{sol} which essentially consists of continuous functions $\mathbf{R}_+ \to \hat{X}$, where \hat{X} is obtained by adjoining to X a point at infinity, such that once the function blows up it cannot be "reborn". The space X^{sol} is itself a complete separable metric space and, if f blows up at some time $\tau > 0$, then any sequence of functions g_{ε} such that $d(f(t), g_{\varepsilon}(t)) \leq \varepsilon$ for $t \leq \tau - \varepsilon$ converges to f in X^{sol} . On the other hand, if $f \in \mathscr{C}(\mathbf{R}_+, X)$, then $d(g_{\varepsilon}, f) \to 0$ if and only if the functions g_{ε} converge to f uniformly over compact time intervals.

The main result of [Hai13, Hai14, BCCH21] is that, for every $\alpha < 1/2 - \kappa$, there exists a jointly continuous map

$$\mathcal{S}: \mathscr{C}^{\alpha} \times \mathscr{M}_0 \to (\mathscr{C}^{\alpha})^{\mathrm{sol}} ,$$

FRACTIONAL BROWNIAN MOTION

such that, writing as above $\hat{Z}^{(\varepsilon)}$ for the BPHZ lift of $(\xi_{\varepsilon}, \chi_{\varepsilon})$, $\mathcal{S}(h_0, \hat{Z}^{(\varepsilon)})$ coincides with the solution to

$$\partial_t h = \partial_x^2 h + (\partial_x h)^2 + 2\chi_\varepsilon \partial_x h + \xi_\varepsilon - C_\varepsilon , \qquad (2.10)$$

with initial condition h_0 , for a suitable constant C_{ε} . Furthermore, if Z denotes the BPHZ lift of $(\xi, 0)$, then $S(h_0, Z)$ is known by [BGHZ22, Rem. 1.8] to coincide with the Cole–Hopf solution to the KPZ equation, provided that one possibly changes the value of C_{ε} by a fixed quantity of order one.

The claim then follows from Theorem 2.7, combined with the fact that solutions to the KPZ equation are global in time. $\hfill \Box$

3 Fractional Brownian motion

We now turn to the proof of convergence of SDEs driven by mollified fractional Brownian motions to a Markov process diffusing in the direction of the Lie brackets of the vector fields defining the original equation.

Let ρ be a fixed mollifier (smooth, with support in [-1, 1], integrating to 1), write $\rho_{\varepsilon}(t) = \varepsilon^{-1}\rho(t/\varepsilon)$, and set $B_{\varepsilon} = C_{\varepsilon}B \star \rho_{\varepsilon}$ where *B* denotes a two-sided *m*-dimensional fractional Brownian motion with Hurst parameter $H \in (0, 1/4]$ and C_{ε} is as in Theorem 1.2. We furthermore write $\xi_{\varepsilon} = \dot{B}_{\varepsilon}$ and we lift it to a geometric rough path in the usual way [FH20] by setting

$$\mathbb{X}_{\varepsilon}^{(k)}(s,t) = \int_{s}^{t} \int_{s}^{r_{k}} \cdots \int_{s}^{r_{2}} \xi_{\varepsilon}(r_{1}) \otimes \ldots \otimes \xi_{\varepsilon}(r_{k}) \, dr_{1} \dots dr_{k} \, .$$
(3.1)

Our proof relies on showing that $\mathbb{X}_{\varepsilon}^{(k)}$ converges to 0 for k odd, while $\mathbb{X}_{\varepsilon}^{(2n)}$ converges to the *n*-fold iterated integral of an m^2 -dimensional Wiener process W satisfying the antisymmetry constraint $W_{ij} = -W_{ji}$.

Even though the situation is quite similar to that of the KPZ equation treated in the previous section, it turns out that the proof is significantly more involved. This is mainly a consequence of the fact that, in the language of regularity structures, the symbols $\Xi_i \mathcal{F}(\Xi_j)$ which will end up representing the (time derivative of the) Wiener processes W_{ij} in the limit transform non-trivially under the action of the structure group. This is unlike the case of the KPZ equation where the second-order process ξ_{ε} can simply be represented by a single noise symbol which transforms trivially.

Our proof relies on delicate estimates on mixed moments of the quantities $\mathbb{X}_{\varepsilon}^{(k)}(s,t)$. Since the integration variables r_i appearing in the expression (3.1) are ordered like $s \leq r_1 \leq \ldots \leq r_k \leq t$, partially ordered sets (posets) will be a convenient structure used to index our integration variables.

3.1 Integrations indexed by posets

In order to express and estimate moments of (3.1), the following formalism will prove to be useful. Given a finite poset P, denote P^{\top} its maximal elements, P^{\perp} its

minimal elements, and $P^{\circ} = P \setminus (P^{\top} \cup P^{\perp})$. Given functions $s \colon P^{\perp} \to \mathbf{R}$ and $t \colon P^{\top} \to \mathbf{R}$, we then write $[s, t]_P$ for the set of all monotone functions $r \colon P \to \mathbf{R}$ such that $r \upharpoonright P^{\perp} = s$ and $r \upharpoonright P^{\top} = t$. We say that P is "linear" if every $x \in P^{\circ}$ has a unique immediate predecessor x^{\downarrow} and a unique immediate successor x^{\uparrow} . Note that, given linear posets P_1 and P_2 , their disjoint sum $P_1 \sqcup P_2$ is again linear.

Given a finite set V, we write $\mathcal{P}(V)$ for the set of pairings of V, namely the set of partitions E of V such that every element of E contains exactly two elements of V. (Note that for $\mathcal{P}(V)$ to be non-empty V needs to have an even number of elements.) Given a map $r: V \to \mathbf{R}$ and a pairing $G \in \mathcal{P}(V)$, we then set

$$K_{\varepsilon}^{(V,G)}(r) = \prod_{\{e,f\}\in G} K_{\varepsilon}(r_f - r_e) , \qquad (3.2)$$

where we set $K_{\varepsilon}(t) = \mathbf{E}\xi_{\varepsilon}(0)\xi_{\varepsilon}(t)$ and note that this expression is well-defined since K_{ε} is even.

With these notations in place, we can write down an expression for the joint moments of the components of $\mathbb{X}_{\varepsilon}^{(k)}$. This is already hinted at by noting that the integral in (3.1) runs over $[s,t]_{P[k]}$, where $P[k] = \{0,\ldots,k+1\}$ endowed with its natural order. Indeed, fix $n \geq 1$, lengths k_1,\ldots,k_n , multiindices $I_i \in$ $\{1,\ldots,m\}^{k_i}$, intervals $[s_i,t_i]$, and fix the shorthand $\mathbb{X}_{\varepsilon}[i] = \langle \mathbb{X}_{\varepsilon}^{(k_i)}(s_i,t_i), e_{I_i} \rangle$ with e_{I_i} the corresponding basis vector of $(\mathbb{R}^m)^{\otimes k_i}$. We then set $P = P[k_1] \sqcup$ $\cdots \sqcup P[k_n]$ and identify s and t with the corresponding constant functions on P^{\perp} and P^{\top} respectively. By Wick's formula, there then exists a subset $\mathcal{P}^I \subset \mathcal{P}(P^\circ)$ of the pairings of P° such that

$$\mathbf{E}\Big(\prod_{i=1}^{n} \mathbb{X}_{\varepsilon}[i]\Big) = \sum_{G \in \mathcal{P}^{I}} \int_{[s,t]_{P}} K_{\varepsilon}^{(P^{\circ},G)}(r \upharpoonright P^{\circ}) dr .$$
(3.3)

When trying to estimate such expressions, one annoying feature is that $\int |K_{\varepsilon}(t)| dt$ diverges as $\varepsilon \to 0$. As a consequence, we cannot simply replace the integrand by its absolute value but have to exploit cancellations. For this, our main tool is a systematic way of rewriting the right-hand side of (3.3) in terms of the kernel \bar{K}_{ε} given by $\bar{K}_{\varepsilon}(t) = \int_{0}^{t} K_{\varepsilon}(r) dr$.

Given a finite set V with a distinguished subset $V^{\circ} \subset V$, write $\mathcal{G}_0(V)$ for the set of all pairs (G, E) such that G is a collection of 2-element subsets of V° (undirected edges) and $E \subset V \times V$ (directed edges) without self-edges. We write $\mathcal{G}(V) \subset \mathcal{G}_0(V)$ for those graphs such that their local structure around any vertex $v \in V^{\circ}$ is furthermore of one of the following three types:

- 1. There is exactly one $g \in G$ with $v \in g$ and no edge of E touching v.
- 2. There is exactly one edge $e \in E$ pointing out of v, i.e. of the type (v, v') for some $v' \in V$, and no edge in G containing v.
- 3. There is exactly one edge $e \in E$ pointing into v, no edge in E pointing out of v, and exactly one edge in G containing v.

Furthermore, we impose that there is at most one vertex of type 3 and that vertices in $V \setminus V^{\circ}$ only have edges in E pointing into them. If we draw undirected edges in G as dark blue lines and directed edges in E as light blue arrows, examples of vertices of these types are as follows:

type 1: • , type 2: • , type 3: • .

The following is an immediate consequence of this structure.

Lemma 3.1. Each connected component of the directed graph (V, E) is of one of the following three types:

- (a) A tree directed towards its root with the root belonging to $V \setminus V^{\circ}$ and all other vertices belonging to V° .
- (b) The union of a directed cycle C of at least two vertices belonging to V° , together with finitely many trees directed towards their roots belonging to C.
- (c) A tree in V° directed towards its root which is the unique vertex of type 3 above.

Proof. Since every vertex can have at most one outgoing edge in E we can explore the graph (V, E) by following these edges. Since V is finite, we must eventually reach a directed cycle or a vertex without outgoing edge. The latter must be either of type 2 or of type 3, thus accounting for the three types of connected components. Note that a component cannot contain more than one such limit set since their basins of attraction must intersect (by connectedness) which would force such an intersection point to have at least two outgoing edges which is not allowed.

We then set, similarly to (3.2),

$$K_{\varepsilon}^{(V,G,E)}(r) = \prod_{\{e,f\}\in G} K_{\varepsilon}(r_f - r_e) \prod_{(e,f)\in E} \bar{K}_{\varepsilon}(r_f - r_e) .$$

Note that we require edges in E to be directed since \bar{K}_{ε} is odd. Consider now the case when the vertex set V is given by a linear poset P as above. It will then be convenient to consider the **Z**-module $\mathscr{G}_0^{\mathbf{Z}}(P)$ generated by $\bigcup_{\hat{P} \subset P} \mathscr{G}_0(\hat{P})$ where the union runs over sub-posets such that $\hat{P}^{\top} = P^{\top}$ and $\hat{P}^{\perp} = P^{\perp}$. We define $\mathscr{G}^{\mathbf{Z}}(P)$ analogously, except that we also quotient out the submodule spanned by graphs $(\hat{P}, G, E) \in \mathscr{G}(P)$ such that E contains a self-edge (i.e. an edge of the type e = (v, v)).

Given s, t as above, we define the Z-linear valuation $J_{\varepsilon}(s, t) \colon \mathscr{G}^{\mathbf{Z}}(P) \to \mathbf{R}$ such that

$$J_{\varepsilon}(s,t)(\hat{P},G,E) = \int_{[s,t]_{\hat{P}}} K_{\varepsilon}^{(\hat{P},G,E)}(r) dr .$$
(3.4)

This is well-defined since $K_{\varepsilon}^{(\hat{P},G,E)}$ vanishes if E contains a self-edge as a consequence of the fact that $\bar{K}_{\varepsilon}(0) = 0$. It will also be convenient to fix a total order on

FRACTIONAL BROWNIAN MOTION

P. (In order to avoid confusion, choose it compatible with the given partial order. Our construction will depend on this total order, but the resulting bounds will not.)

We then define a linear map $\mathcal{F}: \mathcal{G}^{\mathbf{Z}}(P) \to \mathcal{G}_{0}^{\mathbf{Z}}(P)$ in the following way. If G is empty, then $\mathcal{F}(\hat{P}, G, E) = (\hat{P}, G, E)$. If not, we define v^* as the unique vertex of type 3 if such a vertex exists and as the smallest (with respect to the fixed total order) vertex of type 1 otherwise. Either way, write $v_* \in \hat{P}^\circ$ for the unique element such that $g_* = \{v^*, v_*\} \in G$ which is then necessarily of type 1. We then set $\hat{P}' = \hat{P} \setminus \{v_*\}, G' = G \setminus \{g_*\}, E^{\uparrow} = E \cup \{(v^*, v_*^{\uparrow})\}, E^{\downarrow} = E \cup \{(v^*, v_*^{\downarrow})\}, as$ well as $g^{\uparrow} = (\hat{P}', G', E^{\uparrow})$, and similarly for g^{\downarrow} . With these definitions, \mathcal{F} is given by

$$\mathcal{F}(\hat{P}, G, E) = g^{\uparrow} - g^{\downarrow} . \tag{3.5}$$

This is indeed well-defined in the sense that $E^{\uparrow} \subset \hat{P}' \times \hat{P}'$ and G' consists of disjoint undirected edges of $(\hat{P}')^{\circ}$. In fact, \mathcal{F} is simply an encoding of the integration by parts formula as follows.

Proposition 3.2. The map \mathcal{F} maps $\mathcal{G}^{\mathbb{Z}}(P)$ to $\mathcal{G}^{\mathbb{Z}}(P)$ and one has the identity

$$J_{\varepsilon}(s,t)(\mathcal{F}g) = J_{\varepsilon}(s,t)(g) , \qquad (3.6)$$

for every $g \in \mathcal{G}^{\mathbb{Z}}(P)$. Furthermore, there exists N > 0 such that $\mathcal{F}^N = \mathcal{F}^{N+1}$, which we then call \mathcal{F}^{∞} .

Proof. We first argue that one has indeed $(G', E^{\uparrow}) \in \mathcal{G}(\hat{P}')$ (the proof for E^{\downarrow} being identical). Whether v^* is of type 3 or 1, v_* is necessarily of type 1 since $v_* \neq v^*$ and, if there exists any vertex of type 3 then this is v^* . Furthermore the only edge touching v^* is g_* so that, after removing it we are indeed only left with edges in \hat{P}' .

The addition of the edge (v^*, v_*^{\uparrow}) then guarantees that v^* turns into a vertex of type 2 in the graph E^{\uparrow} . The vertex v_*^{\uparrow} on the other hand has a new incoming edge which turns it into a vertex of type 3 if it was previously of type 1 and leaves it to be of type 2 (or in $\hat{P} \setminus \hat{P}^{\circ}$) otherwise.

We now turn to the proof of (3.6). Since the statement is trivial for $g = (\hat{P}, G, E)$ with G empty, we only need to consider the case when it is non-empty. For this, we first note that one has the identity

$$K_{\varepsilon}^{(P,G,E)}(r) = K_{\varepsilon}^{(P',G',E)}(r \upharpoonright \hat{P}')K_{\varepsilon}(r_{v_*} - r_{v^*}).$$

It follows that we can write

$$J_{\varepsilon}(s,t)(\hat{P},G,E) = \int_{[s,t]_{\hat{P}'}} K_{\varepsilon}^{(\hat{P}',G',E)}(r) \int_{r_{v_{*}}^{\downarrow}}^{r_{v_{*}}^{\uparrow}} K_{\varepsilon}(r'-r_{v^{*}}) dr' dr$$

=
$$\int_{[s,t]_{\hat{P}'}} K_{\varepsilon}^{(\hat{P}',G',E)}(r) (\bar{K}_{\varepsilon}(r_{v_{*}}^{\uparrow}-r_{v^{*}}) - \bar{K}_{\varepsilon}(r_{v_{*}}^{\downarrow}-r_{v^{*}})) dr$$

In order to conclude that (3.6) holds, it remains to note that $\bar{K}_{\varepsilon}(0) = 0$, so that the corresponding terms vanish when $v^* \in \{v^{\uparrow}_*, v^{\downarrow}_*\}$. The last statement follows immediately from the fact that \mathcal{F} decreases the cardinality of G by one, unless G is empty in which case it is the identity.

The following lemma provides the a priori bounds on \bar{K}_{ε} necessary for our proofs.

Lemma 3.3. The kernel \bar{K}_{ε} satisfies the following properties:

- 1. There exists a constant C such that $|\bar{K}_{\varepsilon}(t)| \leq C|t|^{-1/2}$, uniformly over $\varepsilon \in (0, 1]$ and $|t| \leq 1$.
- 2. There exists c > 0 such that, for any fixed $\delta \in (0, 1]$, one has

$$\lim_{\varepsilon \to 0} \int_{-\delta}^{\delta} |\bar{K}_{\varepsilon}(t)|^2 dt = c , \qquad \lim_{\varepsilon \to 0} \int_{-\delta}^{\delta} |\bar{K}_{\varepsilon}(t)| dt = 0 .$$
(3.7)

3. For every $\alpha > 1/2$ and every $\delta > 0$ there exist $\varepsilon > 0$ such that $|\bar{K}_{\bar{\varepsilon}}(t)| \le \delta |t|^{-\alpha}$ for all $\bar{\varepsilon} < \varepsilon$ and all $|t| \le 1$.

Proof. This is a straightforward calculation based on the fact that \bar{K}_{ε} is proportional to the convolution of ρ_{ε} with the function $t \mapsto C_{\varepsilon} |t|^{2H-1} \operatorname{sign} t$.

Remark 3.4. The second and third properties could of course be made more quantitative, but this would then lead us to always treat H = 1/4 as a special case which is something that we want to avoid.

Remark 3.5. When H < 1/4, the constant c is mollifier-dependent and given by

$$c = \int_{\mathbf{R}} (\rho \star F_H)(t)^2 \, dt \; ,$$

where $F_H(t) = 2H|t|^{2H-1}$ sign t. When H = 1/4 however one has $c = 4H^2$, independently of ρ . This is an instance of the usual fact that prefactors of logarithmic divergencies tend to be regularisation-independent.

3.2 Main estimates

Given a linear poset P and $E \in P^{\circ} \times P^{\circ}$ such that $(E, \emptyset) \in \mathcal{G}(P)$, we say that E is *full* if every connected component of E consists of a directed cycle of length 2. We furthermore say that E is *ordered* if the partial order on P descends to a partial order on the quotient P_E of P by the finest equivalence relation such that $e \sim f$ for any $(e, f) \in E$. In other words, we want the transitive closure of the smallest relation \leq on P_E such that $a \leq b$ in P implies $[a] \leq [b]$ to be again a partial order.¹

Finally, we write $E^{\leq} \subset E$ for those edges (e, f) such that $e \leq f$. We are now in a position to state the following bound, which is the main working horse for this section.

Theorem 3.6. Let P be a linear poset with $N = |P^{\circ}|$, let $E \in P^{\circ} \times P^{\circ}$ be such that $(E, \emptyset) \in \mathcal{G}(P)$, and write $g = (P, \emptyset, E)$. Let furthermore $T \in (0, 1]$ and $[s, t]_P \subset [0, T]$. Then, one has the following.

¹It is always a preorder but may in general fail to be antisymmetric.

- 1. There exists a constant C such that $|J_{\varepsilon}(s,t)(g)| \leq CT^{N/2}$, uniformly over $\varepsilon \in (0, 1]$.
- 2. If either E isn't full or E is full but unordered, then $\lim_{\varepsilon \to 0} J_{\varepsilon}(s,t)(g) = 0$.
- 3. If E is full and ordered then, writing k for the cardinality of E^{\leq} , one has

$$\lim_{\varepsilon \to 0} J_{\varepsilon}(s,t)(g) = 2^{-k} (-c)^{N/2} |[s,t]_{P_E}| , \qquad (3.8)$$

where c is as in (3.7) and $|\cdot|$ denotes Lebesgue measure on $\mathbf{R}^{P_E^{\circ}}$.

Remark 3.7. If *E* is unordered the definition of P_E still makes sense, but it is only preordered in the sense that there are $x \neq y \in P_E$ with $x \leq y \leq x$. In particular, every $r \in [s,t]_{P_E}$ must satisfy $r_x = r_y$ so that $|[s,t]_{P_E}| = 0$ and (3.8) still holds by point 2.

Proof. To obtain the first bound, write $|J_{\varepsilon}(s,t)|$ for the valuation defined like $J_{\varepsilon}(s,t)$, but with all factors of K_{ε} and \bar{K}_{ε} replaced by $|K_{\varepsilon}|$ and $|\bar{K}_{\varepsilon}|$. If $v \in P^{\circ}$ is a vertex of degree 1, we then write $g' = (P', \emptyset, E')$ with $P' = P \setminus \{v\}$ and $E' = E \setminus \{e\}$ where e is the unique edge containing v. It is then straightforward that one again has $(E', \emptyset) \in \mathcal{G}(P')$.

If such a vertex of degree 1 exists, one then has the bound

$$|J_{\varepsilon}(s,t)(g)| \le |J_{\varepsilon}(s,t)|(g) \le CT^{1/2}|J_{\varepsilon}(s,t)|(g'), \qquad (3.9)$$

as a consequence of the first bound of Lemma 3.3, which in particular implies that any integral of the form $\int_a^b K(t-c) dt$ with $a, b, c \in [0, T]$ is bounded by $CT^{1/2}$ for some C. Iterating this bound, we can reduce ourselves to the case when E consists of finitely many disjoint circles, each of which of size at least 2, going through all elements of P° .

In fact, by freezing all other variables, we can reduce ourselves to the case of one single circle so that it remains to obtain a bound of the type $I_k \leq T^{k/2}$ for any $k \geq 2$, where

$$I_k = \int_0^T \cdots \int_0^T \bar{K}_{\varepsilon}(s_1 - s_2) \cdots \bar{K}_{\varepsilon}(s_{k-1} - s_k) \bar{K}_{\varepsilon}(s_k - s_1) \, ds_1 \cdots ds_k \; .$$

By (3.7) with $\delta = T$, this is bounded by some fixed constant times

$$\int_0^T \cdots \int_0^T \bar{K}_{\varepsilon}(s_2 - s_3) \cdots \bar{K}_{\varepsilon}(s_{k-1} - s_k) \, ds_2 \cdots ds_k \; ,$$

which is in turn bounded by $T(\int_0^{2T} \bar{K}_{\varepsilon}(t) dt)^{k-2}$ which is indeed of order $T^{k/2}$ as claimed.

We now turn to the second bound in the case when E isn't full. If there exists a vertex v of degree 1, we can use the third bound of Lemma 3.3 to conclude that, for any $\delta > 0$ and $\kappa > 0$, there exists ε such that, as in (3.9) (and with g' defined in the same way as it is there), one has

$$|J_{\bar{\varepsilon}}(s,t)|(g) \leq \delta T^{1/2-\kappa} |J_{\bar{\varepsilon}}(s,t)|(g')|,$$

for any $\bar{\varepsilon} < \varepsilon$. It then follows from the first bound that $\lim_{\varepsilon \to 0} |J_{\bar{\varepsilon}}(s,t)|(g) = 0$ as desired. If there is no vertex of degree 1 then, since E isn't full by assumption, it must contain at least one cycle of length $k \ge 3$. Proceeding as in the first bound, this yields a factor $T(\int_{0}^{2T} \bar{K}_{\varepsilon}(t) dt)^{k-2}$ which converges to 0 as $\varepsilon \to 0$ by the second identity in (3.7).

Since the full but unordered case can be obtained in the same way as the third bound, we turn to that one first. We start by observing that, when $G = \emptyset$ and E is full, then N is necessarily even and the integrand of (3.4) has constant sign $(-1)^{N/2}$. This is because each of the N/2 cycles of length 2 leads to a factor of the type $\bar{K}_{\varepsilon}(r_1 - r_2)\bar{K}_{\varepsilon}(r_2 - r_1) = -|\bar{K}_{\varepsilon}(r_1 - r_2)|^2$. As a consequence, (3.8) follows if we can show that

$$\lim_{\varepsilon \to 0} |J_{\varepsilon}(s,t)|(g) = 2^{-k} c^{N/2} |[s,t]_{P_E}| .$$
(3.10)

We also note that the definition of the (pre)order on P_E is such that, if we lift any function $r: P_E \to \mathbf{R}$ to a function ιr on P by setting $(\iota r)_u = r_{[u]}$, then ιr is monotone if and only if r is monotone. The fact that E is ordered is then equivalent to the fact that there exist injective monotone functions on P_E . Write now $(s,t)_{P_E}$ for the subset of $[s,t]_{P_E}$ consisting of monotone functions that are injective on P_E^c .

Every element e of P_E° is an equivalence class consisting of two elements of P, say e_1 and e_2 .² Write U_{δ} for the set of functions $u: P_E^{\circ} \to (-\delta, \delta)$ with the additional constraint that $u_e \ge 0$ if $e_1 \le e_2$ and $u_e \le 0$ if $e_2 \le e_1$. (If $e \notin E^{\le}$ there is no constraint on the sign of u_e .)

We now claim that, given any $r \in (s,t)_{P_E}$, there exists $\delta > 0$ such that, for every $u \in U_{\delta}$, the function $(r|u): P \to \mathbf{R}$ defined by

$$(r|u)(e_1) = r(e)$$
, $(r|u)(e_2) = r(e) + u(e)$, (3.11)

and such that (r|u) coincides with s on P^{\perp} and t on P^{\top} belongs to $[s,t]_P$. For example, it suffices to take $\delta = \inf_{x \in P_E} \inf_{y \in P_E^{\circ}} |r_x - r_y|/2$. This is because we know that $(r|0) \in [s,t]_P$ and the boundary of $[s,t]_P$ consists of those r such that there exists $u \leq v$ with $r_u = r_v$. Since $(r|0) \in [s,t]_P$ and since e_1 and e_2 are incomparable for $(e_1, e_2) \notin E^{\leq}$ and satisfy $e_1 \leq e_2$ otherwise, it follows that we can move the coordinates of the second argument by δ without leaving $[s,t]_P$, provided that we satisfy the ordering constraint on E^{\leq} .

Writing $I(\varepsilon, \delta) = \int_{-\delta}^{\delta} \bar{K}_{\varepsilon}^2(u) du$, it therefore follows that

$$|J_{\bar{\varepsilon}}(s,t)|(g) \geq \int_{[s,t]_{P_E}} \int_{U_{\delta(r)}} |K^g_{\varepsilon}(r|u)| \, du \, dr \geq 2^{-k} \int_{[s,t]_{P_E}} I(\varepsilon,\delta(r))^{N/2} \, dr \; .$$

²The order here is completely arbitrary and our argument does not depend on it.

By Lemma 3.3, $I(\varepsilon, \delta(r))$ is bounded uniformly over $\varepsilon, \delta \in (0, 1]$ and satisfies $\lim_{\varepsilon \to 0} I(\varepsilon, \delta(r)) = c$ for every $\delta > 0$, so that the desired lower bound follows from the dominated convergence theorem.

To obtain the matching upper bound we first note that, in the same way as above,

$$\lim_{\varepsilon \to 0} \int_{[s,t]_{P_E}} \int_{U_T} K^g_{\varepsilon}(r|u) \, du \, dr = 2^{-k} c^{N/2} |[s,t]_{P_E}| \,. \tag{3.12}$$

This however is not sufficient since the image of $[s, t]_{P_E} \times U_T$ under $(r, u) \mapsto (r|u)$ does not fully cover $[s, t]_P$. The reason is that while every element in $[s, t]_P$ can uniquely be written in the form (r|u), it does not necessarily follow that $r \in [s, t]_{P_E}$. However, if we take an arbitrary injective function $r: P_E \to \mathbf{R}$ which coincides with s, t on P^{\perp} and P^{\top} and is such that $r \notin [s, t]_{P_E}$, then we know that $(r|0) \notin [s, t]_{P_E}$. In particular, this shows again that there exists $\delta > 0$ (depending on r) such that, for every u such that $(r|u) \in [s, t]_P$, there exists some coordinate $f \in P_E^{\circ}$ such that $|u_f| \ge \delta$. Write now $V_{\delta}^{(f)}$ for the set of maps $u: P_E^{\circ} \to [-T, T]$ such that $|u_f| \ge \delta$. It then follows that one has

$$|J_{\bar{\varepsilon}}(s,t)|(g) \leq \int_{[s,t]_{P_E}} \int_{U_T} |K^g_{\varepsilon}(r|u)| \, du \, dr + \sum_{f \in P_E^{\circ}} \int_{[s,t]_{P_E}^{c}} \int_{V^{(f)}_{\delta(r)}} |K^g_{\varepsilon}(r|u)| \, du \, dr \,,$$

where $[s, t]_{P_E}^c$ denotes the complement of $[s, t]_{P_E}$. The first term converges to the desired bound by (3.12). Regarding the second term, each summand is bounded by

$$\int_{[s,t]_{P_E}} I(\varepsilon,T)^{N/2-1} (I(\varepsilon,T) - I(\varepsilon,\delta(r))) dr ,$$

which converges to zero by the dominated convergence theorem.

It now remains to treat the case when E si full but unordered. It is then the case that for *every* injective $r: P_E \to \mathbf{R}$ (compatible with s, t) there exists $\delta > 0$ such that, for every u such that $(r|u) \in [s, t]_P$, there exists some coordinate $f \in P_E^{\circ}$ such that $|u_f| \ge \delta$. As a consequence, one has an upper bound analogous to (3.13), but with the first term missing, which proves the claim. \Box

We conclude this section by deriving a consequence of Theorem 3.6 which covers all the cases of interest to us in a way that is much easier to verify, namely in terms of graphs of the type (P, G, \emptyset) which arise naturally in our estimates, rather than the graphs of type (P, \emptyset, E) for which the theorem applies.

Given a linear finite poset P and a pairing G of P° , we say that G is "parallel" if there exists a pairing H of G such that, for any $\{\{e_1, e_2\}, \{f_1, f_2\}\} \in H$, one has (modulo relabelling one of the two edges)

$$f_1 \in \{e_1^{\uparrow}, e_1^{\downarrow}\}, \qquad f_2 \in \{e_2^{\uparrow}, e_2^{\downarrow}\}.$$
 (3.14)

We say that such a pair of edges is "crossed" if one can furthermore relabel the elements in each of the two edges in such a way that $f_1 = e_1^{\uparrow}$ and $f_2 = e_2^{\downarrow}$. The first

two graphs in (3.21) below show one pair of uncrossed parallel edges and one pair of crossed parallel edges.

Note that if G is parallel, then the pairing H is necessarily unique as a simple consequence of the finiteness of G. (We can construct H inductively by taking a minimal vertex v not yet belonging to any of the pairs of edges in H and noting that the edge of G containing v can only possibly be paired with the unique edge containing its successor v^{\uparrow} . This exhausts G after finitely many steps.) Given a parallel pairing G, we define the poset \hat{P} as the quotient of P by the equivalence relation under which, for any pair of parallel edges as in (3.14), one has $e_1 \sim f_1$ and $e_2 \sim f_2$. This is then naturally endowed with a full set E of directed edges which is such that, for any pair of edges in H, one has $([e_1], [e_2]), ([e_2], [e_1]) \in E$. We then have the following result, where we set $J(s,t) = \lim_{\varepsilon \to 0} J_{\varepsilon}(s,t)$ (which exists by Theorem 3.6).

Proposition 3.8. Let P be a linear finite poset with $|P^{\circ}| = 2N$ and let G be a pairing of P° . If G is parallel then, defining \hat{P} and E as just discussed, one has

$$J(s,t)(P,G,\emptyset) = (-1)^{\frac{N}{2}} - \ell J(s,t)(\hat{P},\emptyset,E)$$
$$= (-1)^{\ell} 2^{-k} c^{\frac{N}{2}} |[s,t]_{\hat{P}_E}| . \qquad (3.15)$$

Here, k denotes as before the cardinality of E^{\leq} while ℓ denotes the number of crossed pairs in G. If G is not parallel, then $J(s,t)(P,G,\phi) = 0$.

Proof. Denote by $\hat{\mathscr{G}}(P) \subset \mathscr{G}(P)$ the subsets of graphs (\hat{P}, G, E) such that either E contains an edge adjacent to $\hat{P} \setminus \hat{P}^\circ$ or E contains a connected component of size at least 2 which isn't a two-cycle. Since the action of \mathscr{F} always causes E to grow, we see that $\mathscr{F}: \hat{\mathscr{G}}^{\mathbb{Z}}(P) \to \hat{\mathscr{G}}^{\mathbb{Z}}(P)$ and therefore, by Theorem 3.6 and Proposition 3.2, that $\hat{\mathscr{G}}(P) \subset \ker J(s,t)$. In particular, both \mathscr{F} and J(s,t) are well-defined on $\bar{\mathscr{G}}^{\mathbb{Z}}(P) \stackrel{\text{def}}{=} \mathscr{G}^{\mathbb{Z}}(P)/\hat{\mathscr{G}}^{\mathbb{Z}}(P)$.

Assume now that the total order on P used in the definition of \mathcal{F} is compatible with the given partial order. Consider then $g = (\hat{P}, G, E) \in \mathfrak{C}(P)$ such that E contains only directed cycles of length 2 and such that G is non-empty. (In particular g contains no vertex of type 3.) Let furthermore $e = \{v^*, v_*\} \in G$ be the edge containing the smallest vertex $v^* \in \hat{P}^\circ$ of type 1 and let $f \in G$ be the edge containing the vertex $(v^*)^\uparrow$ (and $f = \emptyset$ if no such edge exists). If $f \neq \emptyset$ and f is parallel to e (i.e. one has $f = \{(v^*)^\uparrow, \hat{v}\}$ with either $\hat{v} = v^\uparrow_*$ "uncrossed" or $\hat{v} = v^\downarrow_*$ "crossed"), then we define $g' = (\hat{P}', G', E')$ by setting

$$\hat{P}' = \hat{P} \setminus \{v_*, (v^*)^{\uparrow}\}, \quad G' = G \setminus \{e, f\}, \quad E' = E \cup \{(v^*, \hat{v}), (\hat{v}, v^*)\}.$$

We claim that in $\overline{\mathscr{G}}^{\mathbf{Z}}(P)$ one then has

$$\mathcal{F}^2 g = \begin{cases} 0 & \text{if } f = \emptyset \text{ or } f \text{ isn't parallel to } e, \\ g' & \text{if the pair } (e, f) \text{ is crossed}, \\ -g' & \text{if the pair } (e, f) \text{ is uncrossed}. \end{cases}$$
(3.16)

Since $\frac{N}{2} - \ell$ is precisely the number of uncrossed pairs in *G*, this immediately implies the first identity in (3.15) (since g' is of the same type as g, so this identity can be iterated until *G* is empty), whence the claim follows from Theorem 3.6.

In order to show (3.16), note that the effect of \mathcal{F} is to delete the node v_* and to replace the edge $e \in G$ by (v^*, \hat{v}) in E with $\hat{v} \in \{v_*^{\uparrow}, v_*^{\downarrow}\}$, yielding the two terms in (3.5) (the second case being the one that comes with a minus sign). If either of these two nodes is of type 2, then this creates a connected component of E with at least two edges, but this cannot be a cycle (since v^* was of type 1) so such terms vanish in $\overline{\mathscr{G}}^{\mathbf{Z}}(P)$. If on the other hand \hat{v} is of type 1 in g, then this turns into a node of type 3 in the corresponding term of $\mathcal{F}g$. It follows that \mathcal{F}^2g is obtained by considering the (unique) edge in G of the form $f = \{\hat{v}, \tilde{v}\}$, deleting the vertex \tilde{v} , and replacing f by $(\hat{v}, \tilde{v}^{\uparrow})$ or $(\hat{v}, \tilde{v}^{\downarrow})$. Since this edge follows the edge (v^*, \hat{v}) , the only way in which we obtain a non-zero element of $\overline{\mathfrak{G}}^{\mathbf{Z}}(P)$ is if either \tilde{v}^{\uparrow} or \tilde{v}^{\downarrow} coincide with the node v^* we started from. Since v^* was the minimal node of type 1 and since \tilde{v} is necessarily of type 1, it cannot be the case that $\tilde{v}^{\uparrow} = v^*$, so the only way of getting a non-zero element is to have f parallel to e and containing $(v^*)^{\uparrow}$. Since $(v^*)^{\uparrow}$ can only be contained in one edge of G, we conclude that we have $\mathcal{F}^2 = \pm g'$ when f is parallel to e and 0 otherwise. The correct sign is obtained by tracking the two cases.

3.3 A tool for convergence

Before we turn to the convergence of the random rough paths \mathbb{X}_{ε} , we provide an estimate for quantities of the same type $J_{\varepsilon}(s,t)(g)$ as considered in (3.4), but with the addition of a second type of edges that represent cutoff functions at a different scale $\delta > 0$. (One should think of $\delta \ll 1$, but without any relation between the sizes of ε and δ .) This section can be skipped at the first read, but its results will play a crucial role in identifying the limit of the fourth-order component of \mathbb{X}_{ε} .

More precisely, we fix once and for all a smooth increasing function $\chi: \mathbf{R} \to [0, 1]$ with $\chi(0) = 0$ and $\chi(1) = 1$, and we set $\chi_{\delta}(t) = \chi(t/\delta)$. We then show a variant of Theorem 3.6 which also allows for kernels of the type $(1 - \chi_{\delta})(t - s)$ and $\chi'_{\delta}(t - s)$. Even though $1 - \chi_{\delta}$ is not symmetric, it will be convenient to also represent it by an unoriented edge which will be unambiguous since in our setting these kernels will always join two comparable vertices in our poset and the corresponding factor will always have an argument of the form $r_u - r_v$ with $v \leq u$.

To formalise this, we now assume that the set of unoriented edges G comes endowed with a decomposition $G = G_K \sqcup G_\chi$ and similarly for E. We also make the standing assumption that edges in G_K always connect vertices that are comparable. It will then be convenient to denote edges e in G_χ as $e = \{e_{\downarrow}, e_{\uparrow}\} \in G_\chi$ with $e_{\downarrow} \leq e_{\uparrow}$. We will use the same convention for edges $e \in E_\chi$. These edges furthermore come with their own orientation and we introduce the notation $(-1)^e$ to denote whether these match or not. More precisely, given $e = (e_1, e_2) \in E_\chi$, we set $(-1)^e = 1$ if $e_1 \leq e_2$ and $(-1)^e = -1$ otherwise. With all these notations at hand, we then set similarly to (3.4)

$$J_{\varepsilon,\delta}(s,t)(P,G,E) = \int_{[s,t]_{\hat{P}}} K_{\varepsilon}^{(P,G_K,E_K)}(r) H_{\delta}^{(P,G_{\chi},E_{\chi})}(r) dr , \qquad (3.17)$$

where we wrote

$$H_{\delta}^{(P,G,E)}(r) = \prod_{e \in G} (1 - \chi_{\delta})(r_{e_{\uparrow}} - r_{e_{\downarrow}}) \prod_{e \in E} (-1)^{e_{\downarrow}} \chi_{\delta}'(r_{e_{\uparrow}} - r_{e_{\downarrow}}) .$$
(3.18)

Given a finite poset P, we now write similarly to before $\mathscr{G}(P)$ for those graphs (\hat{P}, G, E) such that \hat{P} is a subset of $P, G = G_K \sqcup G_\chi$ (and similarly for E), such that edges in G_χ and E_χ are between comparable elements, and such their local structure around any vertex $v \in \hat{P}^\circ$ is of one of the following types:

- 1. There is exactly one $g \in G_K$ with $v \in g$, at most one $g \in G_{\chi}$ with $v \in g$, and no edge of E touching v.
- 2. There is exactly one edge $e \in E$ pointing out of v, i.e. of the type (v, v') for some $v' \in V$, and no edge in G_K containing v. If $e \in E_{\chi}$, then one furthermore has at least one edge in E_K pointing into v.
- 3. There is exactly one edge in E pointing into v, no edge in E pointing out of v, and exactly one edge in G_K containing v.

We again impose that there is at most one vertex of type 3 and that vertices in $V \setminus V^{\circ}$ only have edges in E pointing into them. We furthermore impose that distinct edges in G_{χ} are not comparable, i.e. if $g_i = (g_i^{\uparrow}, g_i^{\downarrow}) \in G_{\chi}$ then $g_1^{\uparrow} \leq g_2^{\uparrow}$ implies $g_1 = g_2$.³ It is easy to check that Lemma 3.1 still holds for such graphs.

We then define $\mathscr{G}^{\mathbf{Z}}(P)$ as before as the **Z**-module generated by $\mathscr{G}(P)$, quotiented by the submodule generated by graphs containing directed self-edges. Again, the valuation $J_{\varepsilon,\delta}$ is well-defined on $\mathscr{G}^{\mathbf{Z}}(P)$ as a consequence of the fact that one has $\bar{K}_{\varepsilon}(0) = \chi'_{\delta}(0) = 0$. In this setting, we can again define an integration by parts operator $\mathcal{J}: \mathscr{G}^{\mathbf{Z}}(P) \to \mathscr{G}^{\mathbf{Z}}(P)$ in a way quite similar to \mathcal{J} , but taking into account that we may not always be able to restrict our integrations over nodes adjacent to a single edge. If G_K is empty, then we do again set $\mathcal{J}(\hat{P}, G, E) = (\hat{P}, G, E)$. If not, let $v^*, v_* \in P^\circ$ be as in the definition of \mathcal{J} on page 17, but substituting G_K for G. We then define $\hat{P}' = \hat{P} \setminus \{v_*\}, G' = G \setminus \{\{v^*, v_*\}\}, E^{\uparrow} = E_K \cup \{(v^*, v_*^{\uparrow})\}$ and $E^{\downarrow} = E_K \cup \{(v^*, v_*^{\downarrow})\}$ in the same way as before. However, while it is again the case that v_* is of type 1 but there are now two possibilities, depending on whether v_* also belongs to an edge in G_{χ} or not.

If it does not, then we proceed exactly as before, setting

$$\mathcal{J}(\hat{P}, G, E) = g^{\uparrow} - g^{\downarrow} , \qquad (3.19)$$

³The fact that $g_1^{\downarrow} \leq g_2^{\downarrow}$ implies $g_1 = g_2$ is of course equivalent.

this time with $g^{\uparrow} = (\hat{P}', G', E^{\uparrow} \sqcup E_{\chi})$. If it does on the other hand, then there exists a unique $\bar{v} \in \hat{P}^{\circ}$ such that $e \stackrel{\text{def}}{=} \{v_*, \bar{v}\} \in G_{\chi}$ and we write $G'' = G' \setminus \{e\}$. In this case, we set $G^{\uparrow} = G'' \cup \{\{v_*^{\uparrow}, \bar{v}\}\}$ and similarly for G^{\downarrow} . Note that it may happen that $v_*^{\uparrow} = \bar{v}$ (resp. $v_*^{\downarrow} = \bar{v}$) in which case we simply set $G^{\uparrow} = G''$ (resp. $G^{\downarrow} = G''$).⁴ We then define \mathcal{F} as

$$\mathcal{J}(\hat{P}, G, E) = g^{\uparrow} - g^{\downarrow} + (\hat{P}, G'', E^0) , \qquad (3.20)$$

where we set $E^0 = (E_K \cup \{(v^*, v_*)\}) \sqcup (E_{\chi} \cup \{(v_*, \bar{v})\})$. We now show that the analogue of (3.6) still holds in this extended case. The case when v_* does not belong to an edge in G_{χ} is treated exactly as in Proposition 3.2. In the other case, assuming that $v_* \geq \bar{v}$, we note that (3.17) can be written as

$$J_{\varepsilon,\delta}(s,t)(\hat{P},G,E) = \int_{[s,t]_{\hat{P}'}} F_{\varepsilon,\delta}(r) \int_{r_{v_*}^{\downarrow}}^{r_{v_*}^{\uparrow}} K_{\varepsilon}(r'-r_{v^*})(1-\chi_{\delta})(r'-r_{\bar{v}}) dr' dr ,$$

where the edge g is as above and $F_{\varepsilon,\delta}$ contains all the factors corresponding to edges in G'' and E. (In the case $v_* \leq \bar{v}$ the argument of χ_{δ} changes sign.) We then integrate by parts, yielding

$$J_{\varepsilon,\delta}(s,t)(\hat{P},G,E) = \int_{[s,t]_{\hat{P}'}} F_{\varepsilon,\delta}(r)\bar{K}_{\varepsilon}(r'-r_{v^*})(1-\chi_{\delta})(r'-r_{\bar{v}})\Big|_{r_{v^*_{*}}}^{r_{v^*_{*}}} dr$$
$$+ \int_{[s,t]_{\hat{P}'}} F_{\varepsilon,\delta}(r) \int_{r_{v^*_{*}}}^{r_{v^*_{*}}} \bar{K}_{\varepsilon}(r'-r_{v^*})\chi_{\delta}'(r'-r_{\bar{v}}) dr' dr ,$$

which can indeed be identified with the three terms appearing in (3.20). Note that the additional sign appearing in the case $v_* \leq \bar{v}$ is taken care of by the factor $(-1)^e$ appearing in the definition (3.18) of the integrand.

Since the action of \mathcal{J} always decreases the number of elements in G_K by one (unless it is empty), we see again that its successive applications stabilise, so that it suffices to bound $J_{\varepsilon,\delta}(s,t)(\mathcal{J}^{\infty}(\hat{P},G,E))$ for the graphs appearing in (3.27) and (3.28). We then have the following analogue to Theorem 3.6 where we say that E is "full" if each of its connected components consists of a cycle containing exactly two edges in E_G (but possibly additional edges in E_{χ}).

Theorem 3.9. Let P be a linear poset with $N = |P^{\circ}|$, let $E = E_K \sqcup E_{\chi}$ and G_{χ} be such that $g = (P, G_{\chi}, E) \in \mathfrak{C}(P)$. For any T > 0 there exists a function o as above such that, for any fixed $s, t \leq T$,

$$|J_{\varepsilon,\delta}(s,t)(g)| \le o(\varepsilon) + o(\delta)$$
,

if E isn't full or G_{χ} contains an element linking two distinct connected components of E.

⁴It may be easier to think of elements of $\mathcal{G}(P)$ as being equivalent if the corresponding sets G_{χ} only differ by singletons.

Proof. It suffices to show that $|J_{\varepsilon,\delta}(s,t)(g)| \le o(\varepsilon)$ if E isn't full and that $|J_{\varepsilon,\delta}(s,t)(g)| \le o(\delta) + o(\varepsilon)$ if E is full and G_{χ} links (at least) two of its connected components.

The proof of the first bound is virtually identical to that of the second statement of Theorem 3.6, noting that the factors $(1 - \chi_{\delta})$ can simply be bounded by one (so edges in G_{χ} can be ignored) and that the integral of $|\chi'_{\delta}|$ is of order 1.

To prove the second bound, we note that one has for $|J_{\varepsilon,\delta}(s,t)(g)|$ the bound (3.13), but with the integrand K_{ε}^g defined in such a way that it also includes the factors $(1 - \chi_{\delta})$. Since the second term in this expression is $o(\varepsilon)$ we only need to bound the first term. We now use the fact that the order used in the definition (3.11) of (r|u) was arbitrary. We can therefore make sure that there exist two elements e, \bar{e} in P_E° (corresponding to the two cycles linked by an edge in G_{χ}) such that the corresponding elements e_1 and \bar{e}_1 appearing in (3.11) satisfy $\{e_1, \bar{e}_2\} \in G_{\chi}$. As a consequence, $(K_{\varepsilon}^g)(r|u)$ contains a factor $(1 - \chi_{\delta})(r_e - r_{\bar{e}})$, which implies that we can restrict the outer integral in the first term of (3.13) to the set of $r \in [s, t]_{P_E}$ such that $|r_e - r_{\bar{e}}| \leq \delta$. This immediately implies that it is bounded by a quantity of order δ , as required.

3.4 Convergence

We now have the tools required to show the main result of this section, namely Theorem 3.13 below, which is the precise formulation of Theorem 1.2. In order to formulate our main result regarding the convergence of \mathbb{X}_{ε} , we write \mathscr{C}_{g}^{α} for the space of \mathbb{R}^{m} -valued geometric rough paths of regularity α . As a first ingredient, we show that the \mathbb{X}_{ε} as defined in (3.1) are tight in \mathscr{C}_{g}^{α} for suitable α .

Proposition 3.10. For any $k \ge 1$ there exists a constant C such that, for any $[s,t] \subset [0,1]$, one has the bound

$$\mathbf{E}|\mathbb{X}_{\varepsilon}^{(k)}(s,t)|^2 \le C|t-s|^{k/2},$$

uniformly over $\varepsilon \in (0, 1]$. In particular, the law of $\mathbb{X}_{\varepsilon}^{(k)}$ is tight in \mathscr{C}_{g}^{α} for any $\alpha < 1/4$.

Proof. Let P be the poset consisting of two copies of $\{0, \ldots, k+1\}$. By (3.3), for any multiindex I of length k, one has a collection \mathcal{P}^I of pairings of P° such that

$$\mathbf{E}|\langle \mathbb{X}_{\varepsilon}^{(k)}(s,t),e_{I}\rangle|^{2} = \sum_{G\in\mathcal{P}^{I}}J_{\varepsilon}(s,t)(P,G,\phi) = \sum_{G\in\mathcal{P}^{I}}J_{\varepsilon}(s,t)(\mathcal{I}^{\infty}(P,G,\phi)) \ .$$

Since all the summands appearing in $\mathcal{F}^{\infty}(P, G, \emptyset)$ are of the form (\hat{P}, \emptyset, E) with $(E, \emptyset) \in \mathfrak{C}(\hat{P})$ and $|\hat{P}^{\circ}| = k$, the claim follows from the first claim of Theorem 3.6 (note that, by stationarity of increments, one can set s = 0 and t = T without loss of generality).

As a next step, we show that the components of $\mathbb{X}_{\varepsilon}^{(2)}$ converge in law to independent Wiener processes W_{ij} subject to the antisymmetry condition $W_{ji} = -W_{ij}$.

FRACTIONAL BROWNIAN MOTION

Proposition 3.11. Let W_{ij} for i < j with $i, j \in \{1, ..., m\}$ be i.i.d. standard Wiener processes, let $W_{ji} = -W_{ij}$, and set $W_{ii} = 0$. Then, for any $\alpha \in (1/5, 1/4)$, \mathbb{X}_{ε} converges in law in \mathscr{C}_{g}^{α} to the limit \mathbb{X} given as follows. One has $\mathbb{X}^{(1)} = 0$, $\mathbb{X}^{(3)} = 0$, $\mathbb{X}^{(2)}_{ij}(s,t) = \sigma \delta W_{ij}(s,t)$, and $\mathbb{X}^{(4)}_{ijk\ell}(s,t) = \sigma^2 \int_s^t \delta W_{ij}(s,r) \circ dW_{k\ell}(r)$. Here, $\sigma = \sqrt{c}$ with c as in (3.7).

Remark 3.12. In fact, the Wiener processes W_{ij} are independent of the fractional Brownian motions B_i in the sense of stable convergence [NNP16].

Proof. Since we already have tightness by Proposition 3.10, it suffices to show convergence of finite-dimensional distributions. The fact that $\lim_{\varepsilon \to 0} \mathbf{E} |\mathbb{X}_{\varepsilon}^{(k)}(s,t)|^2 = 0$ for any s < t and k odd follows immediately from Theorem 3.6 since this quantity can be written as a finite linear combination of terms of the type $J(s,t)(P, \emptyset, E)$ with $|P^{\circ}| = k$ and E cannot be full unless k is even.

Regarding $\mathbb{X}^{(2)}$, the special case m = 2 was treated in [Unto8], but since we need the general case (which doesn't appear to follow in a straightforward way) and since our proof is much shorter we give it here. Our strategy is to apply the fourth moment theorem [NPo5] which states that, when considering random variables belonging to a Wiener chaos of finite order, convergence of covariances and fourth mixed moments is sufficient to guarantee convergence of finite-dimensional distributions to a Gaussian limit. This however is a relatively straightforward consequence of Proposition 3.8. Indeed, $\mathbb{EX}_{\varepsilon;ij}^{(2)}(s_1, t_1)\mathbb{X}_{\varepsilon;k\ell}^{(2)}(s_2, t_2)$ is given by $J_{\varepsilon}(s, t)(g)$ where, pictorially, $g = (P, G, \emptyset)$ is given by

$$g = \delta_{ik}\delta_{j\ell} \xrightarrow{j}_{k_{1}} k_{2} \xrightarrow{j}_{k_{1}} k_{k_{2}} \xrightarrow{j}_{k_{1}} k_{k_{2}} \xrightarrow{j}_{k_{1}} x_{2} \xrightarrow{j}_$$

The light gray lines denote the Hasse diagram of the poset P while the pairing G is shown in blue. Note that the pairings for the first two terms are parallel (with the second one consisting of a crossed pair), while the third one is not, so that, by the first equality of Proposition 3.8, one has J(s,t)(g) = J(s,t)(g') with

$$g' = (\delta_{i\ell}\delta_{jk} - \delta_{ik}\delta_{j\ell}) \underbrace{\begin{smallmatrix} t_1 & t_2 \\ \bullet \\ \bullet \\ \bullet \\ s_1 & s_2 \end{smallmatrix}}_{s_1 & s_2}$$

where now directed edges in E are indicated as light blue arrows. It thus follows

from the second equality of Proposition 3.8 that for $i \neq j$

$$\lim_{\varepsilon \to 0} \mathbf{E} \mathbb{X}^{(2)}_{\varepsilon;ij}(s_1, t_1) \mathbb{X}^{(2)}_{\varepsilon;k\ell}(s_2, t_2) = \begin{cases} c |[s_1, t_1] \cap [s_2, t_2]| & \text{if } (ij) = (k\ell), \\ -c |[s_1, t_1] \cap [s_2, t_2]| & \text{if } (ij) = (\ell k), \\ 0 & \text{otherwise,} \end{cases}$$

with the limit also vanishing when i = j, which is precisely the covariance structure described in the claim.

To complete the proof of Gaussianity of the limit, it remains to show that all joint cumulants of order 4 of terms of the type $\mathbb{X}_{\varepsilon;ij}^{(2)}(s,t)$ vanish as $\varepsilon \to 0$. For this, let P_0 denote the totally ordered poset with 4 elements. Then, the joint cumulants are given by $J_{\varepsilon}(s,t)(g)$ where g is a linear combination of pairings of P° with $P = P_0 \sqcup P_0 \sqcup P_0 \sqcup P_0$, which cannot be written as the disjoint union of two pairings of $(P_0 \sqcup P_0)^{\circ}$. Since it is easy to see that every parallel pairing of P° can be written as such a disjoint union, the vanishing of the cumulants as $\varepsilon \to 0$ follows from Proposition 3.8.

By Skorokhod's representation theorem, we can (and will) therefore assume from now on that $\mathbb{X}_{\varepsilon}^{(2)}$ converges in probability in \mathscr{C}^{α} to $\mathbb{X}^{(2)} = \sigma \delta W$ as described in the statement. It remains to show that, for any fixed s < t, $\mathbb{X}_{\varepsilon}^{(4)}$ converges in probability to the second order iterated integrals of the Wiener processes W_{ij} . Note first that one has

$$\lim_{\varepsilon \to 0} \mathbb{X}^{(4)}_{\varepsilon; iijk} = \lim_{\varepsilon \to 0} \mathbb{X}^{(4)}_{\varepsilon; jkii} = 0 , \qquad (3.22)$$

for all indices i, j, k. This is an immediate consequence of Proposition 3.8, noting that for every parallel pairing appearing in the graphical representation of $\mathbf{E}(\mathbb{X}_{\varepsilon;iijk}^{(4)})^2$ for which the two edges containing the vertices associated to the two factors $\eta_{\varepsilon;i}$ appearing in one of the two copies of $\mathbb{X}_{\varepsilon;iijk}^{(4)}$ are crossed there is a corresponding pairing in which they are uncrossed, with both contributions cancelling out.

As a consequence, we also deduce that one has

$$\lim_{\varepsilon \to 0} \left(\mathbb{X}_{\varepsilon;ijij}^{(4)} - \frac{1}{2} (\mathbb{X}_{\varepsilon;ij}^{(2)})^2 \right) = 0 ,$$

so that, as a consequence of the convergence of $\mathbb{X}_{\varepsilon}^{(2)}$, one has indeed $\lim_{\varepsilon \to 0} \mathbb{X}_{\varepsilon;ijij}^{(4)} = \mathbb{X}_{ijij}^{(4)} \stackrel{\text{def}}{=} \frac{\sigma^2}{2} (\delta W_{ij})^2 = \sigma^2 \int_s^t \delta W_{ij}(s,r) \circ dW_{ij}(r)$ This is an immediate consequence of the shuffle identity

$$(\mathbb{X}_{\varepsilon;ij}^{(2)})^2 = 2\mathbb{X}_{\varepsilon;ijij}^{(4)} + 4\mathbb{X}_{\varepsilon;iijj}^{(4)} ,$$

combined with (3.22).

We also note that one has the identities

$$\lim_{\varepsilon \to 0} \left(\mathbb{X}_{\varepsilon; ijk\ell}^{(4)}(s,t) + \mathbb{X}_{\varepsilon; jik\ell}^{(4)}(s,t) \right) = \lim_{\varepsilon \to 0} \left(\mathbb{X}_{\varepsilon; k\ell ij}^{(4)}(s,t) + \mathbb{X}_{\varepsilon; k\ell ji}^{(4)}(s,t) \right) = 0 , \quad (3.23)$$

which can be shown in virtually the same way as (3.22). As a consequence, it remains to show that $\mathbb{X}_{\varepsilon;ijk\ell}^{(4)}(s,t)$ has the claimed limit in the cases when i, j, k, ℓ are all distinct or when j = k and i, j, ℓ are distinct.

FRACTIONAL BROWNIAN MOTION

Note now that if we set $\mathbb{B}_{\varepsilon}^{(1)}=\mathbb{X}_{\varepsilon}^{(2)}$ and define

$$\mathbb{B}_{\varepsilon}^{(2)}(s,t) = \int_{s}^{t} \mathbb{B}_{\varepsilon}^{(1)}(s,r) \otimes \partial_{r} \mathbb{B}_{\varepsilon}^{(1)}(s,r) \, dr$$

then one has the shuffle identity

$$\mathbb{B}_{\varepsilon;ijk\ell}^{(2)} = \mathbb{X}_{\varepsilon;ijk\ell}^{(4)} + \mathbb{X}_{\varepsilon;kj\ell}^{(4)} + \mathbb{X}_{\varepsilon;kij\ell}^{(4)} .$$
(3.24)

As a consequence of (3.23) the last two terms cancel out in the limit $\varepsilon \to 0$ so that it remains to show that $\mathbb{B}_{\varepsilon;ijk\ell}^{(2)}$ converges to σ^2 times the Stratonovich integral of W_{ij} against $W_{k\ell}$. The reason for considering $\mathbb{B}_{\varepsilon}^{(2)}$ rather than simply $\mathbb{X}_{\varepsilon}^{(4)}$ is that the former is already of the form of an integral of two processes converging towards Wiener processes against each other.

Let now χ_{δ} be as in Section 3.3 and let

$$\mathbb{B}_{\varepsilon,\delta}^{(2)}(s,t) = \int_s^t \int_s^r \chi_\delta(r-u) \partial_u \mathbb{B}_{\varepsilon}^{(1)}(s,u) \otimes \partial_r \mathbb{B}_{\varepsilon}^{(1)}(s,r) \, du \, dr \;. \tag{3.25}$$

It is straightforward to verify the following properties (we fix s, t throughout).

1. For any fixed $\delta > 0$, one has $\lim_{\varepsilon \to 0} \mathbb{B}^{(2)}_{\varepsilon,\delta} = \tilde{\mathbb{B}}^{(2)}_{\delta}$, where

$$\tilde{\mathbb{B}}_{\delta}^{(2)}(s,t) = \int_{s}^{t} \int_{s}^{r} \chi_{\delta}(r-u) \, dW(u) \otimes dW(r) \; .$$

This is a consequence of the convergence $\mathbb{B}_{\varepsilon}^{(1)} \to W$ in \mathscr{C}^{α} obtained in the first part, combined with the fact that the inner integral in (3.25) yields a smooth function of r for every continuous function $\mathbb{B}_{\varepsilon}^{(1)}$, so that the map $\mathbb{B}_{\varepsilon}^{(1)} \mapsto \mathbb{B}_{\varepsilon\delta}^{(2)}$ is continuous on \mathscr{C}^{α} .

2. One has $\mathbb{B}^{(2)} \stackrel{\text{def}}{=} \lim_{\delta \to 0} \tilde{\mathbb{B}}^{(2)}_{\delta} = \mathbb{X}^{(4)}$ with $\mathbb{X}^{(4)}$ as described in the statement of the proposition.

In order to conclude, it thus remains to show that the convergence $\lim_{\delta \to 0} \mathbb{B}^{(2)}_{\varepsilon,\delta} = \mathbb{B}^{(2)}_{\varepsilon}$ has good enough uniformity properties. In fact, we will show the sufficient fact that there exist a function $o: (0, 1] \to (0, 1]$ with $\lim_{\varepsilon \to 0} o(\varepsilon) = 0$ such that

$$\mathbf{E} \|\tilde{\mathbb{B}}_{\varepsilon,\delta}^{(2)}(s,t)\|^2 \le o(\varepsilon) + o(\delta) , \qquad (3.26)$$

where $\tilde{\mathbb{B}}_{\varepsilon,\delta}^{(2)} = \mathbb{B}_{\varepsilon}^{(2)} - \mathbb{B}_{\varepsilon,\delta}^{(2)}$ is given by the same expression as (3.25), but with χ_{δ} replaced by $1 - \chi_{\delta}$.

Before we proceed, let us just show how (3.26) implies the claim. For every $\delta > 0$ one has (all processes being evaluated at a fixed pair of times (s, t)):

$$\mathbf{E} \| \mathbb{X}^{(4)} - \mathbb{B}^{(2)}_{\varepsilon} \|^2 \lesssim \mathbf{E} \| \mathbb{X}^{(4)} - \tilde{\mathbb{B}}^{(2)}_{\delta} \|^2 + \mathbf{E} \| \tilde{\mathbb{B}}^{(2)}_{\delta} - \mathbb{B}^{(2)}_{\varepsilon,\delta} \|^2 + \mathbf{E} \| \tilde{\mathbb{B}}^{(2)}_{\varepsilon,\delta} \|^2$$

FRACTIONAL BROWNIAN MOTION

Given some $\bar{\varepsilon} > 0$, we first choose δ small enough so that $\mathbf{E} \| \mathbb{X}^{(4)} - \tilde{\mathbb{B}}_{\delta}^{(2)} \|^2 \leq \bar{\varepsilon}$ (which is possible by the first property) and $o(\delta) < \bar{\varepsilon}$. We then choose ε small enough so that $o(\varepsilon) < \bar{\varepsilon}$ and $\mathbf{E} \| \tilde{\mathbb{B}}_{\delta}^{(2)} - \mathbb{B}_{\varepsilon,\delta}^{(2)} \|^2 \leq \bar{\varepsilon}$ (which is possible by the second property), so that the right of the displayed equation is smaller than $4\bar{\varepsilon}$.

In the notations of Section 3.3, it follows similarly to (3.24) that one has the identity

$$\tilde{\mathbb{B}}^{(2)}_{\varepsilon,\delta;ijk\ell} = \mathbb{X}^{(4;24)}_{\varepsilon,\delta;ijk\ell} + \mathbb{X}^{(4;34)}_{\varepsilon;ikj\ell} + \mathbb{X}^{(4;34)}_{\varepsilon;kij\ell} ,$$

where we set

$$\mathbb{X}_{\varepsilon,\delta}^{(4;uv)} = \int_s^t \int_s^{r_4} \int_s^{r_3} \int_s^{r_2} (1-\chi_\delta)(r_u-r_v)\,\xi_\varepsilon(r_1)\otimes\ldots\otimes\xi_\varepsilon(r_k)\,dr_1\ldots dr_4\;.$$

It follows that, for i, j, k, ℓ all distinct, one has the identity (with the arguments (s, t) fixed and omitted for clarity)

where edges in G_K are drawn in blue as before while edges in G_{χ} are drawn in red. One furthermore has $\mathbf{E}|\tilde{\mathbb{B}}^{(2)}_{\varepsilon,\delta;ijj\ell}|^2 \leq \mathbf{E}|\tilde{\mathbb{B}}^{(2)}_{\varepsilon,\delta;ijk\ell}|^2 + J_{\varepsilon,\delta}(g)$ with

$$g = 2 () + 2 () + 2 () + 4 () .$$
 (3.28)

We will show that all the terms appearing in (3.27) and (3.28) are bounded by $o(\varepsilon) + o(\delta)$, except for the last two terms in (3.27) that have to be combined in order to take advantage of cancellations.

It follows immediately that the terms in (3.28) are all bounded by $o(\varepsilon)$ since no cycle with two elements of E_{χ} can be generated by successive applications of \mathcal{J} . Regarding (3.27), the first term is bounded by $o(\delta) + o(\varepsilon)$ since the only "full" term that can be obtained by successive applications of \mathcal{J} is



which yields a contribution at most $o(\delta)$ by Theorem 3.9.

It remains to deal with the last two terms in (3.27). It turns out that these terms are *not* separately bounded by $o(\varepsilon) + o(\delta)$, but their sum is due to cancellations. This is because, if we choose our fixed total order on the vertices such that the smallest vertex is the second one from the bottom of the left column, then one has



It is then clear from the definition of \mathcal{J} that further applications of \mathcal{J} cannot lead to a full graph, so that these terms contribute by at most $o(\varepsilon)$ by Theorem 3.9

Theorem 3.13. Given an initial condition $x_0 \in \mathbf{R}^d$, let μ_{ε} be the law of the maximal solution to (1.4), viewed as an $(\mathbf{R}^d)^{\text{sol}}$ -valued random variable. Then, μ_{ε} converges weakly to the law μ of the maximal solution to (1.5) with $\sigma^2 = c$ where c is given by (3.7).

Proof. By continuity of the solution map $\mathbf{R}^d \times \mathscr{C}_g^\alpha \to (\mathbf{R}^d)^{\text{sol}}$, it remains to show that the solution to the rough differential equation (RDE) driven by \mathbb{X} coincides with that of the Stratonovich SDE (1.5). In view of [FH20, Thm 9.1], it suffices to show the following. Assume that we are given an $\mathbf{R}^m \otimes \mathbf{R}^m$ -valued 2α -Hölder continuous geometric rough path ($\mathbb{B}^{(1)}, \mathbb{B}^{(2)}$) additionally satisfying the antisymmetry relations

$$\mathbb{B}_{ij}^{(1)} + \mathbb{B}_{ji}^{(1)} = 0 , \quad \mathbb{B}_{ijk\ell}^{(2)} + \mathbb{B}_{jik\ell}^{(2)} = 0 , \quad \mathbb{B}_{ijk\ell}^{(2)} + \mathbb{B}_{ij\ell k}^{(2)} = 0 .$$
(3.29)

In our case, \mathbb{B} is given by the Stratonovich lift of W, which indeed satisfies the additional relations (3.29) as a consequence of the fact that $W_{ji} = -W_{ij}$.

Given such a \mathbb{B} , we claim that the fourth order \mathbb{R}^m -valued rough path \mathbb{X} given by

$$\mathbb{X}^{(1)} = 0$$
, $\mathbb{X}^{(2)} = \mathbb{B}^{(1)}$, $\mathbb{X}^{(3)} = 0$, $\mathbb{X}^{(4)} = \mathbb{B}^{(2)}$, (3.30)

then satisfies $\mathbb{X} \in \mathscr{C}_g^{\alpha}$. Furthermore, for any collection of m smooth functions $V_i \colon \mathbf{R}^d \to \mathbf{R}^d$, the solution to the RDE

$$dx(t) = V_i(x) \, d\mathbb{X}_i(t) \,, \qquad x(0) = x_0 \,, \tag{3.31}$$

coincides with the solution to the RDE

$$dx(t) = \frac{1}{2} [V_i, V_j](x) d\mathbb{B}_{ij}(t) , \qquad x(0) = x_0 .$$
(3.32)

The claim of the theorem then follows at once from the fact that the RDE solutions to (3.32) with \mathbb{B} the Stratonovich lift of W do coincide with the solutions to the Stratonovich SDE (1.5) by [FH20, Thm 9.1].

The proof that solutions to (3.31) and (3.32) coincide is folklore (see [SL91, Sec. 3] for a pre-rough path statement strongly hinting at it, but it certainly goes back

even further since these kind of statements motivate the definition of Lie brackets and form the basis of geometric control theory) but this higher order version does not seem to appear in the literature, so we provide a sketch of proof. The fact that X as defined by (3.30) satisfies Chen's relations and the required analytic bounds is easy to verify for any 2α -Hölder rough path \mathbb{B} so that, in order to show that $X \in \mathscr{C}_g^{\alpha}$, it remains to verify that the shuffle relations (see for example [FH20, Eq. 2.21]) hold. This however follows immediately from the shuffle relations for \mathbb{B} , combined with the fact that we enforced (3.29).

After unpacking the definitions one finds that solutions to (3.31) are in general given by the fixed point problem

$$\begin{aligned} x(t) &= x_0 + \lim_{|\mathcal{P}_t| \to 0} \sum_{[u,v] \in \mathcal{P}_t} \left(V_i \mathbb{X}_i^{(1)} + DV_i(V_j) \mathbb{X}_{ji}^{(2)} + D^2 V_i(V_j, V_k) \mathbb{X}_{kji}^{(3)} \right. \\ &+ DV_i(DV_j(V_k)) \mathbb{X}_{kji}^{(3)} + D^3 V_i(V_j, V_k, V_\ell) \mathbb{X}_{\ell kji}^{(4)} \\ &+ DV_i(DV_j(DV_k(V_\ell))) \mathbb{X}_{\ell kji}^{(4)} + DV_i(D^2 V_j(V_k, V_\ell)) \mathbb{X}_{\ell kji}^{(4)} \\ &+ D^2 V_i(V_j, D_k(V_\ell)) (\mathbb{X}_{j\ell ki}^{(4)} + \mathbb{X}_{\ell jki}^{(4)} + \mathbb{X}_{\ell kji}^{(4)}) \Big) , \end{aligned}$$

where \mathcal{P}_t denotes a partition of [0, t] and $|\mathcal{P}_t|$ is the length of its largest subinterval. The terms involving $\mathbb{X}_i^{(1)}$ and $\mathbb{X}_i^{(3)}$ vanish by (3.30). The term involving $\mathbb{X}^{(2)}$ can be rewritten by the first identity in (3.29) as

$$\frac{1}{2} (DV_i(x_u)V_j(x_u)) (\mathbb{B}_{ji}^{(1)}(u,v) - \mathbb{B}_{ij}^{(1)}(u,v)) = \frac{1}{2} (DV_j(x_u)V_i(x_u) - DV_i(x_u)V_j(x_u)) \mathbb{B}_{ij}^{(1)}(u,v) = \frac{1}{2} [V_i, V_j](x_u) \mathbb{B}_{ij}^{(1)}(u,v) .$$

Regarding the four terms involving $\mathbb{X}^{(4)}$, the first one vanishes since $\mathbb{X}^{(4)}_{\ell k j i}$ in antisymmetric in (k, ℓ) by (3.29) and (3.30) while the prefactor is symmetric. The third one vanishes for an analogous reason. Using again (3.29) and (3.30), a straightforward calculation shows that the sum of the remaining two terms equals

$$\frac{1}{4}D[V_i, V_j]([V_k, V_\ell]) \mathbb{B}^{(2)}_{k\ell i j}$$
,

so that one indeed recovers the fixed point problem for (3.32) as required.

Remark 3.14. The convergence of course also holds at the level of flows.

References

- [ADC21] M. AIZENMAN and H. DUMINIL-COPIN. Marginal triviality of the scaling limits of critical 4D Ising and φ_4^4 models. *Annals Math.* **194**, no. 1, (2021), 163. arXiv:1912.07973. doi:10.4007/annals.2021.194.1.3.
- [Aiz82] M. AIZENMAN. Geometric analysis of φ^4 fields and Ising models. I, II. *Comm. Math. Phys.* **86**, no. 1, (1982), 1–48.

- [BC98] L. BERTINI and N. CANCRINI. The two-dimensional stochastic heat equation: renormalizing a multiplicative noise. J. Phys. A 31, no. 2, (1998), 615–622. doi:10.1088/0305-4470/31/2/019.
- [BCCH21] Y. BRUNED, A. CHANDRA, I. CHEVYREV, and M. HAIRER. Renormalising SPDEs in regularity structures. J. Eur. Math. Soc. (JEMS) 23, no. 3, (2021), 869-947. arXiv:1711.10239. doi:10.4171/jems/1025.
- [BG97] L. BERTINI and G. GIACOMIN. Stochastic Burgers and KPZ equations from particle systems. Comm. Math. Phys. 183, no. 3, (1997), 571–607. doi:10.1007/s002200050044.
- [BGHZ22] Y. BRUNED, F. GABRIEL, M. HAIRER, and L. ZAMBOTTI. Geometric stochastic heat equations. J. Amer. Math. Soc. 35, no. 1, (2022), 1–80. arXiv:1902.02884. doi:10.1090/jams/977.
- [BHZ19] Y. BRUNED, M. HAIRER, and L. ZAMBOTTI. Algebraic renormalisation of regularity structures. *Invent. Math.* **215**, no. 3, (2019), 1039–1156. arXiv:1610.08468. doi:10.1007/s00222-018-0841-x.
- [BHZ20] Y. BRUNED, M. HAIRER, and L. ZAMBOTTI. Renormalisation of stochastic partial differential equations. *Eur. Math. Soc. Newsl.*, no. 115, (2020), 7–11. doi:10.4171/news/115/3.
- [BL22] L. BROUX and D. LEE. Besov reconstruction. *Potential Analysis* (2022), 1–38. doi:10.1007/s11118-022-10028-7.
- [BS10] K. BURDZY and J. SWANSON. A change of variable formula with Itô correction term. *Ann. Probab.* **38**, no. 5, (2010), 1817–1869. doi:10.1214/09-A0P523.
- [CCHS22] A. CHANDRA, I. CHEVYREV, M. HAIRER, and H. SHEN. Langevin dynamic for the 2D Yang-Mills measure. Publ. Math. Inst. Hautes Études Sci. 136, (2022), 1–147. doi:10.1007/s10240-022-00132-0.
- [CH16] A. CHANDRA and M. HAIRER. An analytic BPHZ theorem for regularity structures. *arXiv preprint* (2016). arXiv:1612.08138.
- [CQ00] L. COUTIN and Z. QIAN. Stochastic differential equations for fractional Brownian motions. C. R. Acad. Sci. Paris Sér. I Math. 331, no. 1, (2000), 75–80. doi:10.1016/S0764-4442(00)01594-9.
- [CSZ23a] F. CARAVENNA, R. SUN, and N. ZYGOURAS. The critical 2d stochastic heat flow. *Invent. Math.* 233, no. 1, (2023), 325–460. doi:10.1007/s00222-023-01184-7.
- [CSZ23b] F. CARAVENNA, R. SUN, and N. ZYGOURAS. The critical 2d stochastic heat flow is not a Gaussian multiplicative chaos. Ann. Probab. 51, no. 6, (2023), 2265–2300. doi:10.1214/23-aop1648.
- [DGRZ20] A. DUNLAP, Y. GU, L. RYZHIK, and O. ZEITOUNI. Fluctuations of the solutions to the KPZ equation in dimensions three and higher. *Probab. Theory Related Fields* 176, no. 3-4, (2020), 1217–1258. doi:10.1007/s00440-019-00938-w.
- [Duc22] P. DUCH. Flow equation approach to singular stochastic PDEs. *arXiv preprint* (2022). arXiv:2109.11380.
- [FH20] P. K. FRIZ and M. HAIRER. A course on rough paths. Universitext. Springer, Cham, 2020, xvi+346. With an introduction to regularity structures, Second edition. doi:10.1007/978-3-030-41556-3.

- [FM17] M. FURLAN and J.-C. MOURRAT. A tightness criterion for random fields, with application to the Ising model. *Electron. J. Probab.* 22, (2017), Paper No. 97, 29. doi:10.1214/17-EJP121.
- [Frö82] J. FRÖHLICH. On the triviality of $\lambda \varphi_d^4$ theories and the approach to the critical point in d > 4 dimensions. *Nuclear Phys. B* **200**, no. 2, (1982), 281–296. doi:10.1016/0550-3213(82)90088-8.
- [GIP15] M. GUBINELLI, P. IMKELLER, and N. PERKOWSKI. Paracontrolled distributions and singular PDEs. Forum Math. Pi 3, (2015), e6, 75. arXiv:1210.2684. doi:10.1017/fmp.2015.2.
- [GRZ18] Y. GU, L. RYZHIK, and O. ZEITOUNI. The Edwards-Wilkinson limit of the random heat equation in dimensions three and higher. *Comm. Math. Phys.* **363**, no. 2, (2018), 351–388. doi:10.1007/s00220-018-3202-0.
- [Gub04] M. GUBINELLI. Controlling rough paths. J. Funct. Anal. 216, no. 1, (2004), 86–140. doi:10.1016/j.jfa.2004.01.002.
- [Hai13] M. HAIRER. Solving the KPZ equation. Ann. of Math. (2) **178**, no. 2, (2013), 559–664. doi:10.4007/annals.2013.178.2.4.
- [Hai14] M. HAIRER. A theory of regularity structures. *Invent. math.* **198**, no. 2, (2014), 269–504. arXiv:1303.5113. doi:10.1007/s00222-014-0505-4.
- [Hai18] M. HAIRER. An analyst's take on the BPHZ theorem. In Computation and combinatorics in dynamics, stochastics and control, vol. 13 of Abel Symp., 429–476. Springer, Cham, 2018. arXiv:1704.08634. doi:10.1007/978-3-030-01593-0_16.
- [HL15] E. HÄUSLER and H. LUSCHGY. Stable convergence and stable limit theorems, vol. 74 of Probability Theory and Stochastic Modelling. Springer, Cham, 2015, x+228. doi:10.1007/978-3-319-18329-9.
- [HS17] M. HAIRER and H. SHEN. A central limit theorem for the KPZ equation. *Ann. Probab.* **45**, no. 6B, (2017), 4167–4221. doi:10.1214/16-A0P1162.
- [HS23] M. HAIRER and R. STEELE. The BPHZ theorem for regularity structures via the spectral gap inequality. *arXiv preprint* (2023). arXiv:2301.10081.
- [JS03] J. JACOD and A. N. SHIRYAEV. Limit theorems for stochastic processes, vol. 288 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, second ed., 2003, xx+661. doi:10.1007/978-3-662-05265-5.
- [Kup16] A. KUPIAINEN. Renormalization group and stochastic PDEs. Annales Henri Poincaré 17(2016). doi:10.1007/s00023-015-0408-y.
- [LOTT22] P. LINARES, F. OTTO, M. TEMPELMAYR, and P. TSATSOULIS. A diagram-free approach to the stochastic estimates in regularity structures. *arXiv preprint* (2022). arXiv:2112.10739.
- [LV07] T. LYONS and N. VICTOIR. An extension theorem to rough paths. Ann. Inst. H. Poincaré C Anal. Non Linéaire 24, no. 5, (2007), 835–847. doi:10.1016/j.anihpc.2006.07.004.
- [Ly098] T. J. LYONS. Differential equations driven by rough signals. *Rev. Mat. Iberoamericana* 14, no. 2, (1998), 215–310. doi:10.4171/RMI/240.

- [MU18] J. MAGNEN and J. UNTERBERGER. The scaling limit of the KPZ equation in space dimension 3 and higher. J. Stat. Phys. **171**, no. 4, (2018), 543-598. doi:10.1007/s10955-018-2014-0.
- [NNP16] I. NOURDIN, D. NUALART, and G. PECCATI. Quantitative stable limit theorems on the Wiener space. Ann. Probab. 44, no. 1, (2016), 1–41. doi:10.1214/14-A0P965.
- [NP05] D. NUALART and G. PECCATI. Central limit theorems for sequences of multiple stochastic integrals. Ann. Probab. 33, no. 1, (2005), 177–193. doi:10.1214/009117904000000621.
- [NR09] I. NOURDIN and A. RÉVEILLAC. Asymptotic behavior of weighted quadratic variations of fractional Brownian motion: the critical case H = 1/4. Ann. Probab. 37, no. 6, (2009), 2200–2230. doi:10.1214/09-A0P473.
- [NT11] D. NUALART and S. TINDEL. A construction of the rough path above fractional Brownian motion using Volterra's representation. Ann. Probab. 39, no. 3, (2011), 1061–1096. doi:10.1214/10-A0P578.
- [PT11] G. PECCATI and M. S. TAQQU. Wiener chaos: moments, cumulants and diagrams, vol. 1 of Bocconi & Springer Series. Springer, Milan; Bocconi University Press, Milan, 2011, xiv+274. A survey with computer implementation, Supplementary material available online. doi:10.1007/978-88-470-1679-8.
- [SL91] H. SUSSMANN and W. LIU. Limits of highly oscillatory controls and the approximation of general paths by admissible trajectories. In *Proceedings of* the 30th IEEE Conference on Decision and Control, vol. 1, 437–442. 1991. doi:10.1109/CDC.1991.261338.
- [Unto8] J. UNTERBERGER. A central limit theorem for the rescaled Lévy area of twodimensional fractional Brownian motion with Hurst index H < 1/4. arXiv preprint (2008). arXiv:0808.3458.
- [Unt13] J. UNTERBERGER. A renormalized rough path over fractional Brownian motion. *Comm. Math. Phys.* **320**, no. 3, (2013), 603–636. doi:10.1007/s00220-013-1707-0.