# Probing New Physics at Cosmic Dawn with 21-cm Cosmology

Omer Zvi Katz,<sup>1</sup> Nadav Outmezguine,<sup>2,3</sup> Diego Redigolo,<sup>4</sup> and Tomer Volansky<sup>1</sup>

<sup>1</sup>School of Physics and Astronomy, Tel-Aviv University, Tel-Aviv 69978, Israel

<sup>2</sup>Berkeley Center for Theoretical Physics, University of California, Berkeley, CA 94720, U.S.A.

<sup>3</sup>Theory Group, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, U.S.A.

<sup>4</sup>INFN, Sezione di Firenze, Via Sansone 1, 50019 Sesto Fiorentino, Italy

21-cm cosmology provides an exciting opportunity to probe new physics dynamics in the early universe. In particular, a tiny sub-component of dark matter that interacts strongly with the visible sector may cool the gas in the intergalactic medium and significantly alter the expected absorption signal at Cosmic Dawn. However, the information about new physics in this observable is obscured by astrophysical systematic uncertainties. In the absence of a microscopic framework describing the astrophysical sources, these uncertainties can be encoded in a bottom up effective theory for the 21-cm observables in terms of unconstrained astrophysical fluxes. In this paper, we take a first step towards a careful assessment of the degeneracies between new physics effects and the uncertainties in these fluxes. We show that the latter can be constrained by combining measurements of the UV luminosity function, the Planck measurement of the CMB optical depth to reionization, and an upper bound on the unresolved X-ray flux. Leveraging those constraints, we demonstrate how new physics signatures can be disentangled from astrophysical effects. Focusing on the case of millicharged dark matter, we find sharp predictions, with small uncertainties within the viable parameter space.<sup>a</sup>

## I. INTRODUCTION

Cosmological observables such as the Cosmic Microwave Background (CMB) and the galaxy power spectrum provide invaluable evidence for the existence of dark matter (DM), pointing to a pressureless fluid that feebly interacts with the Standard Model. Furthermore, analysis of the same cosmological data gives detailed information regarding possible non-trivial dynamics of a so-called dark sector in which DM could reside [1–11]. Within this wealth of information from the early universe, 21-cm cosmology is tracking the behavior of the neutral hydrogen in the inter-galactic medium (IGM) at different redshifts [12]. At  $z \leq 6$ , 21-cm intensity mapping provides a new tracer of cosmic structure [13] which can complement optical surveys of Large Scale Structure [14]. For  $z \gtrsim 6$  these measurements provide a unique opportunity to probe the (rather dark) Universe since recombination and until the Epoch of Reionization – a period that remains poorly understood and challenging to investigate through alternative observables. Nevertheless, such measurements do not unfold without their inherent difficulties (see for example Ref. [12, 15, 16] for reviews on the subject).

A measurement far from the EoR, at  $z \gtrsim 30$ , is accompanied by severe experimental challenges which may be addressed by the futuristic program of lunar interferometry [16]. Conversely, in order to understand measurements at  $6 \leq z \leq 30$ , during the so-called Cosmic Dawn, non-linear structure formation and stellar evolution must be understood and hence predictions suffer from significant systematic uncertainties, even assuming standard cosmology [12, 15]. As a consequence, utilizing 21-cm measurements to detect or constrain any non-standard dark sector dynamics requires a detailed estimation of these uncertainties. This is the goal of this letter.

The interest in the Cosmic Dawn epoch has been reinvigorated by the EDGES observational result in 2018 which suggested a strong absorption signal in the global 21-cm spectrum [17]. The signal was significantly stronger than the maximal absorption signal possible within standard cosmology, hinting towards nonstandard dynamics. The EDGES result has since been disputed by the SARAS3 collaboration [18] and many subtleties in the estimation of its significance might be hiding in the assessment of the systematic uncertainties which are plaguing the global 21-cm spectrum measurements [19–22]. More information is expected to come from high-redshift 21-cm interferometry that should be able to shed light on the global signal results [23].

Irrespective of the unsettled experimental status, it is natural to ask: 'What kind of non-standard dynamics could leave detectable imprints in 21-cm observables at Cosmic Dawn?' In particular, exploring the potential for detecting imprints of a dark sector is intriguing. To date, broadly speaking, five dark sector effects have been identified:

- **Dark cooling** models in which the dark sector acts to cool the gas in the IGM through DM elastic collisions [24–31].
- **Dark heating**, where the dark sector is heating up the gas through DM annihilations or decays [32–40].
- Modification of the Raileygh-Jeans tail via a resonant conversion of dark sector particles to CMB photons [41–44].

<sup>&</sup>lt;sup>a</sup> This paper is a contribution to the proceeding of the Nobel Symposium on Dark Matter.

- Dark sector effects on the power spectrum. Such changes may delay or enhance radiation which heats up or ionizes the gas (see e.g. [45]) and therefore affects the 21-cm signal.
- Dark sector effects on star formation. In particular, suppressed stellar production [46] or variation in their evolution may impact the 21-cm signal [47].

Each of the above categories requires detailed analyses to ensure that phenomenologically viable models exist. More importantly, new physics effects may be degenerate with the inherent astrophysical uncertainties, a prevailing challenge during the Cosmic Dawn era. Within the above categories, certain scenarios exhibit complete degeneracy. For example, fuzzy or warm DM models lead to a suppressed matter power spectrum at small scales, resulting in a reduced density of Pop-III stars. As this stellar population remains observationally elusive, it is currently impossible to distinctly separate the primary effects of such dark sectors using 21-cm cosmology. The situation, however, is often more optimistic and in this letter we aim to take an initial stride in quantifying the degeneracies, concentrating specifically on the dark cooling category outlined above.

The theoretical challenges of realizing a viable dark cooling model have been addressed in the specific example of Ref. [30, 31], where a tiny fraction of the DM energy density (below the current CMB bound [48, 49]) is millicharged. If the millicharged component interacts with the rest of the cold DM through a long range interaction, the heat capacity of the cooling bath is enhanced at low redshift, thereby enhancing the signal at Cosmic Dawn. One therefore concludes that 21-cm observables have the opportunity to probe a genuinely unexplored region of the millicharge DM parameter space which lies below the current accelerator constraints [30, 50–55], but couples too strongly to be probed in regular terrestrial direct detection experiments [30, 56].

Given this well defined theoretical setup, we focus on the challenge of identifying the underlying degeneracies between the dark cooling signal and astrophysical uncertainties. In particular, the dark cooling acts to enhance the absorption signal at Cosmic Dawn well beyond the one predicted within standard cosmology. We would therefore like to ask in which region of the parameter space a measurable enhancement could occur independently of the astrophysical uncertainties. The detailed exploration of our analysis, including improved constraints on the astrophysical parameters, is presented in Ref. [57] where the different possible stellar formation models and the independent code developed to evaluate their implications on the global 21-cm spectrum are described.

Our paper is organized as follows. In Sec. II we give a brief introduction to the 21-cm global signal, discussing both standard cosmology as well as dark cooling. In Sec. III we identify the important astrophysical functions, which constitute an effective theory for 21-cm cosmology, and review the relations between them and the simplified astrophysical models. In the process we determine the corresponding astrophysical uncertainties and discuss existing and upcoming constraints. In Sec. IV we summarize some of our results. Further details and results are presented in [57].

#### II. 21-CM WITH DARK COOLING

In this section we introduce the basics of 21-cm cosmology in Sec. II A, focusing on the sky-averaged 21-cm brightness and its behavior at Cosmic Dawn. Then, we illustrate the evolution of the IGM temperature in the presence of dark cooling in Sec. II B.

#### A. 21-cm Basics

The CMB photons measured today have propagated the universe since redshift of order 1100, traversing cold neutral hydrogen clouds before reaching us. During this time, photons with a wavelnegth of 21-cm were repeatedly absorbed and emitted through the hyperfine transitions of ground state hydrogen. The resulting deviation in the tail of the CMB spectrum as measured today is usually parametrized by the differential 21-cm brightness temperature [58–62]

$$T_{21}(z) = \frac{(T_s(z) - T_{\gamma}(z))}{1 + z} \left(1 - e^{-\tau_{21}(z)}\right)$$
(1)  
$$\simeq 27x_{\rm HI} \left[1 - \frac{T_{\gamma}}{T_s}\right] \left[\frac{1 + z}{10} \frac{0.15}{\Omega_m h^2}\right]^{1/2} \left[\frac{\Omega_b h^2}{0.023}\right] \,\mathrm{mK}\,,$$

where  $T_{\gamma}$  is the CMB temperature and  $T_s$  is the effective spin temperature, quantifying the relative occupation of the two hyperfine levels of the hydrogen ground state. The second equality is obtained by assuming that the optical depth of 21-cm photons remains small, i.e.  $\tau_{21} \ll 1$ . In the absence of new physics, this condition is satisfied until the end of reionization, where the neutral fraction of hydrogen,  $x_{HI}(z) \equiv n_{HI}(z)/n_H(z)^1$  drops significantly.

The dynamics of  $T_s$  is set by spin flipping processes: i) Resonant absorption and emission of 21-cm CMB photons. This process is controlled by the induced emission and absorption rates  $B_{10} = B_{01}/3$ . ii) Hydrogen scatterings with helium, the residual ionized fraction and other hydrogen atoms. We express the collisional de-excitation

<sup>&</sup>lt;sup>1</sup> Throughout this work we use the standard notation to distinguish between the different baryonic components. In this notation: HI, HeI stand for neutral hydrogen and helium atoms, HII and HeII represent singly ionized hydrogen and helium atoms, and finally HeIII are doubly ionized helium atoms. When writing H or He it is to be understood as as the total hydrogen and helium populations, including all levels of ionization.

rate as  $C_{10} = \sum_{i=\text{HI,e,p}} n_i k_{10}^i$ , where  $n_i$  are the number densities of the different species and the  $k_{10}^i$  factors depend on the kinetic temperature of hydrogen,  $T_k$ , and are tabulated in Ref. [61, 63–65]. iii) Absorption of Ly- $\alpha$  photons followed by a de-excitation to the opposite spin state (also known as the Wouthuysen Field (WF) effect [58, 66]). Since Ly- $\alpha$  photons are emitted by stars, the corresponding de-excitation rate, parameterized by  $P_{10}$ , will depend on detailed astrophysics.

Throughout the evolution of  $T_s$ , the dominant spinflipping rate is always much grater than the Hubble expansion rate, driving the system into a steady state given by

$$T_s^{-1} = \frac{T_{\gamma}^{-1} + \bar{x}_{\alpha} T_{\alpha}^{-1} + \bar{x}_k T_k^{-1}}{1 + \bar{x}_{\alpha} + \bar{x}_k}, \qquad (2)$$

where the collisional and Ly- $\alpha$  coupling coefficients are defined as  $\bar{x}_k = C_{10}/B_{10}$  and  $\bar{x}_\alpha = P_{10}/B_{10}$ , and the Ly- $\alpha$  temperature,  $T_\alpha$ , is an effective color temperature defined through the equilibrium relation of the WF spin flip interactions, i.e.  $P_{01}/P_{10} = 3e^{-\frac{E_{21}}{T_\alpha}}$ . For completeness we also write the WF spin flipping

For completeness we also write the WF spin flipping rate in its common parametrization [67]

$$P_{10} = 4\pi \int \sigma_{10}(E) J_{\alpha}(E) dE \equiv 6\pi \lambda_{\alpha}^2 \gamma_{\alpha} \tilde{S}_{\alpha} \bar{J}_{\alpha} , \qquad (3)$$

where  $\sigma_{10} = \frac{3}{2} \lambda_{\alpha}^2 \gamma_{\alpha} \phi_{10}(E)$  is the WF de-excitation cross section,  $\lambda_{\alpha}$  is the wavelength of Ly- $\alpha$  photons,  $\gamma_{\alpha} = 0.74 \alpha_{\rm em} E_{\alpha}^3 a_0^2 \simeq 4.3 \times 10^{-7}$  eV is the line half-width at half-maximum, and  $\phi(E)$  is the corresponding line profile.  $J_{\alpha}$  is the specific number intensity (number of photons per time, energy, area and steradian) around the Ly- $\alpha$  energy, resulting from the emission of UV photons by early luminous sources.

The spectrum around the Ly- $\alpha$  line center is modified by energy transfer between Ly- $\alpha$  photons and hydrogen atoms as a result of recoil and spin exchange in WF interactions [67, 68]. This modification, together with the details of the line profile are encapsulated in  $\tilde{S}_{\alpha} = \int \phi_{10}(E) J_{\alpha}/\bar{J}_{\alpha}$ , where  $\bar{J}_{\alpha}$  is defined as the intensity at the red edge of the line and hence unaffected by the WF scatterings. In our 21-cm code, we calculate  $\tilde{S}_{\alpha}$ and  $T_{\alpha}$  following Ref. [67]<sup>2</sup>, while the intensity  $\bar{J}_{\alpha}$  depends on astrophysics and will be discussed in Sec. III B.

At the end of the Dark Ages  $(z \sim 30)$ ,  $T_s$  is coupled to  $T_{\gamma}$ , resulting in  $T_{21} = 0$ . However, at the onset of the Cosmic Dawn, star formation results in Ly- $\alpha$  fluxes which quickly become the dominant spin flipping rate, coupling  $T_s$  to  $T_{\alpha}$ . Prior to heating from astrophysical sources, one has  $T_k < T_{\alpha} < T_{\gamma}$  and thus the 21-cm signal is found in absorption. This absorption signal of Cosmic Dawn is the focus of this study. Crucially, the amplitude and width



FIG. 1. The 2cDM dark sector [30] is comprised of two DM components. The majority of DM, assumed to be cold DM (CDM), does not engage in tree-level interactions with SM particles, while the remaining small fraction,  $f_{\rm m}$ , is assumed to posses an electric charge. In addition, the two DM components interact with one another through a new long range dark force. The evolution of the 2cDM model is as follows: i) prior to hydrogen recombination, mDM is strongly coupled to baryons, evading CMB constraints as long as  $f_m \leq 4 \times 10^{-3}$  [48, 49]. During this period the long range interactions between CDM and mDM are suppressed due to the large relative velocity. ii) As the latter dissipates, mDM begins to cool through its interactions with CDM, eventually decoupling from the baryons. Acting as a large heat bath, CDM absorbs any excess heat from the mDM, always keeping it colder than the baryons which continuously deposit heat to the mDM fluid through elastic scatterings, mediated by SM photons. iii) In some regions of the 2cDM parameter space, mDM eventually couples to CDM.

of this absorption signal in standard cosmology depends on the astrophysical fluxes controlling  $T_{\alpha}$ ,  $T_{\rm k}$  and  $x_{HI}$ in Eq. (1). The goal of this study is to disentangle this dependence from new dark sector dynamics affecting  $T_{\rm k}$ , whose time evolution we describe below.

## B. The Kinetic Temperature of Hydrogen and Dark Cooling

Including both SM and BSM contributions, the kinetic temperature of baryons evolves as

$$\frac{dT_k}{d\log a} = -2T_K + \frac{2}{3H} \left( \dot{Q}_{\text{Compton}} + \dot{Q}_X + \dot{Q}_{\text{Ly}\alpha} + \dot{Q}_{\text{CMB}} + \dot{Q}_{\text{BSM}} \right), \quad (4)$$

where *a* is the scale factor,  $H(z) = \sqrt{\Omega_m} H_0 (1+z)^{3/2}$ is the Hubble rate controlling the adiabatic cooling, and the  $\dot{Q}_i$ s represent the heat transfer rates per baryon as a result of different SM and BSM processes which we briefly discuss.

At high redshifts, hydrogen is mostly ionized, CMB photons continuously Compton scatter with the residual electron fraction, coupling  $T_{\rm k}$  to  $T_{\gamma}$  through  $\dot{Q}_{\rm Compton}$ .

<sup>&</sup>lt;sup>2</sup> Ref. [67] assumed  $T_{\alpha} \gg E_{21}$  in deriving  $\tilde{S}_{\alpha}$ . Under this assumption  $P_{01} = 3P_{10}$ , and their definition is equivalent to ours.



FIG. 2. An illustration depicting the constraining procedure of the 21-cm effective theory described by three astrophysical functions: i)  $J_{\alpha}$  (light blue) - the specific intensity at the Ly- $\alpha$  line, ii)  $J_{X}$  (green) - the specific intensity at X-ray energies, and iii)  $x_{\rm HI}$  (purple) - the neutral fraction of hydrogen. These functions may be characterized using simplified astrophysical models represented as triangles with corresponding colors. All three models rely on yet another simplified model of star formation (central light grey triangle). The parameters of each simplified model are listed beneath its title. On the left side we list the data extracted from simulations and population synthesis models, which are integrated into the construction of the simplified models. On the right side, the measured observables used to constrain each simplified model are outlined, maintaining the same color coding.

However, eventually hydrogen recombines and adiabatic cooling prevails. The era of adiabatic cooling (Dark Ages) only terminates once first stars form, emitting Xray and UV radiation. The former propagates to the IGM, heating it through photoionization, while the latter redshifts or cascades to the Ly- $\alpha$  energy where the photons exchange heat with the hydrogen atoms as a result of WF scatterings. Additionally, scatterings of 21-cm CMB photons and hydrogen atoms result in spin flipping interactions affecting the hydrogen kinetic temperature which is encoded in  $\dot{Q}_{\rm CMB}$ .

The heat transfer rates  $\dot{Q}_{\rm X}$ , and  $\dot{Q}_{{\rm Ly}\alpha}$  strongly depend on the details of the astrophysical processes, including the formation of stellar objects and their emission properties. This dependence is encapsulated in specific number intensities for X-rays,  $J_{\rm X}$ , and for Ly- $\alpha$  photons,  $\bar{J}_{\alpha}$ . The modeling these intensities is described in Sec. III, but generically we have  $\dot{Q}_{\rm X} \gg \dot{Q}_{{\rm Ly}\alpha}$ .

The expressions for the Compton and X-ray heating rates are common in the 21-cm literature and can be found in Refs. [61, 69]. On the other hand, being significantly smaller, the Ly- $\alpha$  and CMB terms are often neglected. However, these rates are enhanced at low  $T_{\rm k}$ , and may therefore play a role in situations with extreme dark cooling and suppressed X-ray flux. We therefore incorporate these rates in our 21-cm code. For calculating  $\dot{Q}_{\rm Ly\alpha}$  we follow Ref. [68], but accounting for the full Ly- $\alpha$  line profile derived in Ref. [67]. For  $\dot{Q}_{\rm CMB}$  we follow Ref. [70], which corrects the original result of Ref. [71].

Finally,  $\dot{Q}_{\rm BSM}$  in Eq. (4) accounts for the dark sector contribution to the  $T_K$  evolution. Here we focus on the effect of dark cooling through DM-SM elastic scatterings. Specifically, we take as a benchmark dark cooling model the two coupled DM (2cDM) sector, first proposed in Ref. [30] whose mechanism is reviewed in Fig. 1. In this work we will assume a benchmark 2cDM model where the amount of dark cooling is maximized: the millihcrage fraction is set at the CMB bound  $f_{\rm m} = 4 \times 10^{-3} [48, 49]$ , the cold DM mass is fixed at the BBN bound  $m_{\rm C}$  = 10 MeV [72] and the mDM-CDM cross section is fixed to the maximal value allowed by CMB constraints as discussed in Ref. [30]. This leaves only two free parameters in the dark sector, the mDM mass,  $m_{\rm m}$ , and charge, Q, whose viable parameter space is shown in Fig. 4. We leave a more comprehensive study of the 2cDM parameter space to Ref. [57].

#### **III. ASTROPHYSICAL UNCERTAINTIES**

The behavior of the brightness temperature in Eq. (1) depends on three astrophysical functions: i) the Ly- $\alpha$  flux  $\bar{J}_{\alpha}$ , ii) the X-ray flux  $J_X$ , and iii) the ionized fraction  $x_{HI}$  which depends on the UV and X-ray fluxes together. If left completely undetermined, a very strong X-ray flux, a very inefficient Ly- $\alpha$  emission, or an early reionization



FIG. 3. The evolution of the 21-cm global Cosmic Dawn signal assuming only SM physics (red), and including dark cooling (blue), as predicted over the viable astrophysics and 2cDM parameter space. Each line corresponds to a specific combination of astrophysics and dark sector parameters. Here, we assume the star formation model by Ref. [73] as discussed in Sec. III A and constrain the astrophysical fluxes by imposing current measurements of the UV luminosity function [73, 74] (Sec. IIIB), the optical depth to reionization measured by Planck [75] and the measured spectra of bright quasars [76] (Sec. IIID), and the expected constraint on the unresolved X-ray flux from Chandra [77, 78], discussed in Sec. (IIIC). The **dashed black** line shows the expected boundary of these models, once results from the future Athena experiment become available [79]. For concreteness, we fix the mDM-CDM cross section to its maximal value allowed by CMB constraints, take the millicharged component relative density to be  $f_{\rm m} = 4 \times 10^{-3}$ , and fix the mass of the CDM component at  $m_{\rm C} = 10$  MeV.

can in principle make the Cosmic Dawn absorption signal arbitrarily flat for any scenario of dark cooling, making any statement about the dark sector dynamics plagued by astrophysical uncertainties.

In what follows we show that using a combination of present and future observables and simulations one can sufficiently bracket the three astrophysical functions to make robust statements about the dark sector with only few assumptions about the spectrum of astrophysical fluxes. The full procedure we follow to achieve this goal is illustrated in Fig. 2 and is detailed in the remainder of this section.

The final result of this analysis is summarized in Fig. 3 where the envelope of all possible 21-cm brightness temperature evolutions is shown for standard cosmology (red) and with the 2cDM dark cooling scenario (blue). Crucially, current constraints on the UV luminosity function [73, 74] and the optical depth to reionization [73, 75] imply that astrophysical scenarios with very inefficient Ly- $\alpha$  emission or early reionization are excluded. We also impose existing and expected constraints on the unresolved X-ray flux from Chandra [77, 78] and Athena [79] respectively. The latter is expected to improve the current constraint from Chandra by at least two orders of magnitudes.

#### A. Models of star formation

A basic ingredient in modeling the global intensity of astrophysical sources is the rate of their formation. We focus specifically on the formation of stars, and define the star formation rate density (SFRD) as the rate at which mass is converted to stars per co-moving volume. The SFRD can be expressed as an integral of the halo mass function (HMF),  $dn/dM_h^3$ :

$$\dot{\rho}^i_{\star}(t) = \frac{d}{dt} \int_0^\infty m^i_{\star}(M_h) \frac{dn}{dM_h}(M_h, t) dM_h \,, \qquad (5)$$

where the mapping function,  $m_{\star}^{*}(M_{h})$ , relates the total mass of a given halo,  $M_{h}$ , to its stellar mass. Motivated by the extrapolation of high redshift UV luminosity functions (UVLFs), the star formation efficiency in halos exceeding a threshold virial mass of  $M_{\rm cut}$  can be modeled as a powerlaw in  $M_{\rm h}$  [73, 74]. Below  $M_{\rm cut}$  – the precise value of which depends on the baryonic composition of the halo and on stellar feedback effects – star formation is highly suppressed.

To be specific, we focus on the star formation model described in Ref. [73] which considers a single star population (Pop-II stars) described by three parameters: i) the normalization of the star formation efficiency  $F_{\star}$ , ii) the power law index  $\alpha_{\star}$ , and iii) the cut-off of star formation  $M_{\rm cut}$ . More realistic star formation models, in particular including an extra population of Pop-III stars [80] will be considered in [57].

To evaluate the SFRD in our scenario we use the  $\Lambda$ CDM HMF with the prescription of [81] available in the public python toolkit COLOSSUS [82]. The cosmological parameters other than  $\tau_{reion}$  are fixed using the  $\Lambda$ CDM fit to the Planck data [75], assuming they are unaffected by the dark sector. While this assumption is certainly valid for a fraction of mDM with  $f_m < 4 \times 10^{-3}$  it would be interesting to check if in the 2cDM model, discussed in Sec. II B, the late time recoupling of CDM to mDM can lead to a sizeable suppression of small scale structures (see Ref. [45] for a related study).

## B. The Ly- $\alpha$ intensity

Ly- $\alpha$  photons are produced by non-ionizing UV photons either redshifting from below the Ly- $\beta$  line or cascading down through higher Lymann lines.  $\bar{J}_{\alpha}$  of Eq. (3) can then be written in terms of the UV emissivity,  $\epsilon_{\rm UV}$ ,

<sup>&</sup>lt;sup>3</sup> Following the modeling of Ref. [73], instead of the time derivative in Eq. 5, we divide the stellar mass by a characteristic timescale  $H^{-1}(t)t_{\star}$ , where  $t_{\star}$  is a dimensionless parameter.

which sums all the astrophysical sources weighted by their probability to end the cascade with the production of a Ly- $\alpha$  photon [67, 83]

During the Cosmic Dawn, stars are the primary source of UV photons [83, 84], with the most massive stars making the dominant contribution at the Lymann band energies [67]. Assuming that the lifetime of the stars is short with respect to variations in the star formation rate implies that the volume-averaged comoving number emissivity of UV photons is proportional to the star formation rate [83]

$$\epsilon_{\rm UV}(E,t) = \frac{\dot{\rho}_{\star}(t)}{\mu_b} \left\langle \frac{dN_*}{dE} \right\rangle \,, \tag{6}$$

where  $\epsilon_{\rm UV}$  gives the number of UV photons from all sources per unit time, energy and comoving volume,  $\mu_b$  is the average baryon mass and  $\langle \frac{dN_*}{dE} \rangle$  is the average number of photons emitted by a single baryon per unit energy, summed over all relevant stellar populations.

Since the Ly- $\alpha$  flux depends only on the spectra,  $\langle \frac{dN_*}{dE} \rangle$ , within a very small energy band, its exact shape is irrelevant and can be set by its normalization,  $N_{\rm ion}$ . Consequently, the emissivity in Eq. (6) is entirely defined by  $N_{\rm ion}\dot{\rho}_{\star}$ . Since  $N_{\rm ion}$  is degenerate with the normalization  $F_{\star}$  of  $\dot{\rho}_{\star}$ , we fix  $N_{\rm ion} = 5000$ .

Data collected by the Hubble Space Telescope (HST) can be used to map the rest frame UV luminosity function (UVLF) at  $4 \leq z \leq 10$  [74]. These UVLFs can then be used to constrain the parameters associated with stellar formation simplified models ( $F_{\star}$ ,  $\alpha_{\star}$ ,  $M_{cut}$ ) once  $N_{\text{ion}}$ is fixed. The preliminary constraints we use in our analysis were first derived in [73, 74]. The James Webb Space Telescope is expected to improve on these constraints in the near future.

#### C. The X-ray intensity

Soft X-ray photons (with energy  $E \leq 2$  keV) continuously interact with the IGM through photoionizations<sup>4</sup>. Their intensity can be derived by solving the energy transfer equation in a homogeneously absorbing, non-emitting medium [87]. The modeling of  $J_X$  can then be related to the global co-moving number emissivity,  $\epsilon_X$ , which depends on the characterization of the X-ray sources.

Based on observations of nearby starburst galaxies [88– 92], it is believed that the dominant X-ray sources during Cosmic Dawn are high-mass X-ray binaries (HMXBs). Being short lived, the emissivity (and luminosity) of the HMXBs population traces the star formation rate, and can therefore be written as in Eq. (6). In accordance with local measurements [88–92] and high redshift simulations [93, 94], the averaged photon spectrum that contributes to the IGM heating can be modelled as a single truncated power law in energy,

$$\left\langle \frac{dN_X}{dE} \right\rangle = \frac{1}{E_0} \left[ \frac{E}{E_0} \right]^{\beta_X} \Theta \left[ E - E_{\min} \right] \Theta \left[ E_{\max} - E \right], \quad (7)$$

where  $E_{\text{max}} = 2$  keV is the maximal energy and the reference energy  $E_0$  is typically related to the ratio of the luminosity and star formation rate in the corresponding energy band [61, 73, 95] which we rescale with a normalization parameter  $F_X$ .  $E_{\min}$  is the minimal energy that an X-ray photon must posses in order to escape its host galaxy. Its precise value depends on the assumed local density and metallicity of the galaxy [94]. In our scan we take  $E_{\min} \in (0.19, 0.85)$  keV, which corresponds to the  $2\sigma$  range in which the X-rays optical depth in the interstellar medium as extracted from the hydrodynamical simulations of Ref. [94] is exactly one. Finally, since the variation of  $\beta_X$  is less significant than the one of  $E_{\min}$  [96] we fix it to be  $\beta_X = -2$ , in accordance to high-redshift simulations [73, 93, 94].

An upper limit on the X-ray flux can be derived by assuming that early X-ray sources are responsible for the totality of the unresolved cosmic X-ray background (CXB) in the [0.5, 2] keV energy band, observed today by Chandra [77, 78, 97]. Defining  $z_X^{\text{un}}$  as the maximal redshift above which none of the X-ray sources are resolved by a given telescope, the following constraints from the Chandra measurement can be derived

$$\frac{1}{(1+z_X^{\rm un})^3} \int_{0.5\,\rm keV}^{2\,\rm keV} E J_{\rm X}\left(E, z_X^{\rm un}\right) dE < 0.5 \frac{\rm keV}{\rm cm^2\,\rm sec\,\rm sr} \,. \tag{8}$$

Since star formation accelerates at lower redshifts, sources at the lowest redshift dominate the CXB so that a lower  $z_X^{\text{un}}$  results in a stronger bound on  $F_X$ . Given that Chandra reported resolved soft X-ray galaxies only up to  $z \sim 1$ , we conservatively take  $z_X^{\text{unres}} = 3$ . The upper limit on the right hand side of Eq. (8) will be further improved with the future Athena experiment [79] which is expected to give a bound at least two order of magnitude tighter. Since we also expect its resolution to improve, we show the Athena expected limit taking  $z_X^{\text{unres}} = 4$ .

Estimating the impact of the constraints on the unresolved CXB on the X-ray spectrum in Eq. (7) involves the modelling of the X-ray spectrum above  $E_{\text{max}}$ . While Ref. [78] assumed a single power law, extrapolating Eq. (7) up to 14 keV, here we assume a second power law above 3.5 keV taking the softest choice consistent with simulations of HMXBs (corresponding to  $\beta_{\rm X} \approx -3.2$ ), and matching it to the soft X-ray spectrum in Eq. (7). This procedure gives a conservative constraint on the X-ray flux assuming HMXBs are the dominant X-ray sources during the Cosmic Dawn.

<sup>&</sup>lt;sup>4</sup> X-rays with energy greater than approximately 2 keV have a mean free path longer than the Hubble distance at z = 10 [85, 86]. As a consequence, only X-rays with energies below this value play a role in heating and ionizing the IGM.



FIG. 4. The parameter space of millicharged dark matter in the charge Q vs mass  $m_m$  plane, for the 2cDM model of Ref. [30] (see summary in Fig. 1). The **grey** shaded regions are excluded by BBN measurements of  $\Delta N_{\text{eff}}$  [26, 28, 29, 98], accelerator experiments [50–55], and direct detection searches collected in [30, 56]. CMB constraints on mDM-baryon and mDM-CDM interactions are accounted for in the choice of the 2cDM parameters. **Left:** The predicted minimal brightness temperature,  $T_{21}$ , of the Cosmic Dawn absorption signal for the fiducial astrophysical model of Ref. [73] (see Sec. IV for the precise choice of parameters). The minimal temperature is indicated in shades of **Yellow** while the **dashed blue** lines specify the redshift at which the minimum is achieved. The **solid black line** on the color bar marks the minimal  $T_{21}$  allowed within the Standard Model for all viable astrophysics models (see also Fig. 3), while the **solid black contour** on the main figure marks the Q,  $m_m$  values needed to reach this temperature assuming the fiducial astrophysics. **Right:** The astrophysics-independent prediction for the expected exclusion assuming a null observation with a -100 mK sensitivity and under the assumptions of a power-law SFR model fitted to UVLF data [73, 74]. In all of the shown millicharged DM parameter space, a brightness temperature as low as -100 mK can be obtained with some choice of viable astrophysics model. Within the **dark (light) purple** region,  $T_{21} < -100$  mK is predicted for *any* viable astrophysics model that satisfies the constraints discussed in Sec. III, including the X-ray constraint of Chandra [77, 78] (Athena [79]). The **yellow** line shows the predicted  $T_{21} = -100$  mK for the fiducial astrophysics scenario assumed in the left figure.

## D. The neutral hydrogen fraction

During Cosmic Dawn, astrophysical objects emit X-ray and UV photons, reionizing the universe by  $z \sim 6$  [76]. In our 21-cm code we model the X-ray and UV ionization following Refs. [99, 100] <sup>5</sup>, correcting for the total number of UV ionizing photons according to [73]. The latter assumes that the fraction of ionizing photons which escape the ionized regions surrounding the emitting source is a power law in halo mass, as suggested by high redshift simulations [107, 108].

Planck measurements of large scale CMB anisotropies measure the optical depth to reionization,  $\tau_e$ , yielding  $\tau_e = 0.054 \pm 0.0070$  at 68% C.L. [75]. This measurement can be mapped to a constraint on the ionized fraction  $x_e$  (or equivilantly the neutral fraction of hydrogen  $x_{HI}$ ) according to the relation

$$\tau_e = n_H^0 \sigma_T \int_0^{50} dz x_e(z) \frac{(1+z)^2}{H(z)} , \qquad (9)$$

where  $n_{HI}^0$  is the number density of hydrogen atoms today,  $\sigma_T$  is the Thompson cross section, and the upper integration limit is chosen to be high enough to capture the full contribution to  $\tau_e$ . In addition, missing Ly- $\alpha$  and Ly- $\beta$  photons in bright quasars spectra can also be used as a probe for reionization, setting a 1 $\sigma$  upper limit of  $x_{HI} < 0.06 + 0.05$  at z = 5.9 [76].

Since reionization is generally led by UV photons, the Planck measurement provides an upper bound on the UV photons that escape their host galaxies.

## IV. RESULTS

We are now prepared to explore the possibilities of leveraging the 21-cm global signal as a tool to unravel insights about the 2cDM dark sector. Specifically, we focus on the DM implications for the minimal temperature at Cosmic Dawn alone, leaving the utilization of the full spectral shape, which entails greater constraining and discovery powers, for the upcoming publication [57].

Without delving into the complicated systematic uncertainties which affect the 21-cm measurements, we en-

<sup>&</sup>lt;sup>5</sup> A full inhomogeneous treatment would require the use of intricate simulations (see [101–106] for examples of full numerical simulations, and [62] for the semi-analytical simulation applied in the public 21cmFAST code). However, despite its simplicity, upon comparing our results with those obtained from the semianalytical code 21cmFAST [62], we observe a high level of accuracy in reproducing the global 21-cm signal for redshifts  $z \gtrsim 10$ as well as the electron scattering optical depth to CMB.

tertain here as an illustration the possibility of a nondetection of the Cosmic Dawn peak with sensitivity of -100 mK. This assumption enables us to gauge the anticipated sensitivity to new physics in the global 21-cm spectrum, taking into account the astrophysical uncertainties.

To gear our expectations we first adopt the fiducial astrophysical model of Ref. [73] as our benchmark model with  $F_{\star} = 0.05$ ,  $\alpha_{\star} = 0.5$ ,  $M_{\rm cut} = 5 \times 10^8 M_{\odot}$   $f_{\rm X} = 1$ ,  $E_{\rm min} = 0.5$ ,  $f_{\rm esc} = 0.1$ ,  $\alpha_{\rm esc} = -0.5$ ,  $t_{\star} = 0.5$ . Within this model we then evaluate the amplitude and redshift of the Cosmic Dawn absorption signal in the mDM mass,  $m_{\rm m}$ , and charge, Q plane. The results are shown on the left of Fig. 4, where we see that most of the allowed parameter space with mDM mass below 10 GeV predicts an absorption dip which is deeper than the minimal one in standard cosmology. Furthermore, with sensitivity of -100 mK, most of the viable parameter space for the above astrophysics scenario, could be excluded.

In order to asses the systematic uncertainties on these conclusions, we perform a wide scan on the astrophysical parameters (working within the power-law SFR framework discussed in [73, 74]) together with the dark sector parameters  $(m_{\rm m}, Q)$ , systematically imposing the observational constraints discussed in Sec. III. We also evaluate the impact of the expected constraints on the unresolved CXB from Athena [79]. The envelope of all possible brightness temperatures after imposing the existing constraints is shown in Fig. 3.<sup>6</sup>

The possible variations of the signal, associated with the astrophysical uncertainties, can now be mapped onto the 2cDM parameter space, as shown on the right of Fig. 4. While for any point in the  $m_{\rm m}$ -Q plane there exists a viable astrophysics model for which the Cosmic Dawn absorption feature is below -100 mK, in part of the parameter space, such a minimum occurs for any viable astrophysics model. The dark purple region in the plot shows this part of the parameter space, scanning over all astrophysical models which satisfy the constraints of Sec. III, and taking into account only the existing Chandra X-ray limits. Within this region, a null measurement with an experimental sensitivity of -100 mK in the 21cm brightness temperature would result in a robust constraint on millicharged dark matter, *independently* of astrophysics. The lighter blue region shows the expected improvement once the limits from Athena will become

available. For comparison, the yellow line corresponding to the the contour of min  $(T_{21}) = -100$  mK in the astrophysical benchmark of the left panel is also shown. Comparing the yellow line with the dark purple region shows the importance of varying the astrophysical parameters to account for degeneracies with dark sector physics. Furthermore, we find that even under our very conservative assumption, with future X-ray measurements one could exclude this benchmark astrophysics model within standard cosmology, independently of any new physics scenario.

We note that the expected sensitivity in Fig. 4 should be complemented with data from 21-cm interferometry which have the potential of further reducing the astrophysical degeneracies (see Ref. [31] for preliminary work in this direction). The study performed so far demonstrates that 21-cm measurements can undoubtedly play a crucial role in closing the gap in the millicharge DM parameter space, in a complementary manner to future accelerators searches [53, 109–113], low-threshold direct detection experiments above ground [56] and ion traps [114, 115].

#### V. ACKNOWLEDGMENTS

We thank Andrey Mesinger, Julian Munoz and Yuxiang Qin for illuminating discussions and feedback on their code. We also thank Hongwan Liu, Andrea Caputo, Laura Lopez-Honorez and Ely Kovets for discussions. OZK thanks the Alexander Zaks scholarship for its support. The work of DR is partially financed by the PRIN 2022 – Next Generation EU. The work of TV is supported, in part, by the Israel Science Foundation (grant No. 1862/21), by the Binational Science Foundation (grant No. 2020220), by the NSF-BSF (grant No. 2021780) and by the European Research Council (ERC) under the EU Horizon 2020 Programme (ERC-CoG-2015, Proposal No. 682676 LDMThExp).

**Declaration of generative AI and AI-assisted technologies in the writing process:** During the preparation of this work the author(s) minimally used ChatGPT in order to improve the readability of a few paragraphs. After using this service, the author(s) reviewed and edited the content as needed and take(s) full responsibility for the content of the publication.

 Mikhail M. Ivanov, Evan McDonough, J. Colin Hill, Marko Simonović, Michael W. Toomey, Stephon Alexander, and Matias Zaldarriaga, "Constraining Early Dark Energy with Large-Scale Structure," Phys. Rev. D **102**, 103502 (2020), arXiv:2006.11235 [astroph.CO].

[2] Nicholas DePorzio, Weishuang Linda Xu, Julian B. Muñoz, and Cora Dvorkin, "Finding eV-scale light relics with cosmological observables," Phys. Rev. D 103, 023504 (2021), arXiv:2006.09380 [astro-ph.CO].

<sup>&</sup>lt;sup>6</sup> In detail we take  $\log_{10} F_{\star} \in [-2.09, -0.66]$ ,  $\alpha_{\star} \in [0.29, 0.70]$ ,  $\log_{10} M_{\text{cut}} \in [7.91, 9.43]$ ,  $\log_{10} f_X \in [-2, 2.1]$ ,  $E_{\min} \in [0.19, 0.85]$ ,  $\alpha_{\text{esc}} \in [-1.74, 1.18]$ ,  $\log_{10} f_{\text{esc}} \in [-1.94, 0.35]$ , and keep  $t_{\star} = 0.5$  constant given its strong degeneracy with  $F_{\star}$  (see Ref. [73]).

- [3] Weishuang Linda Xu, Julian B. Muñoz, and Cora Dvorkin, "Cosmological constraints on light but massive relics," Phys. Rev. D 105, 095029 (2022), arXiv:2107.09664 [astro-ph.CO].
- [4] Francis-Yan Cyr-Racine, Kris Sigurdson, Jesus Zavala, Torsten Bringmann, Mark Vogelsberger, and Christoph Pfrommer, "ETHOS—an effective theory of structure formation: From dark particle physics to the matter distribution of the Universe," Phys. Rev. D 93, 123527 (2016), arXiv:1512.05344 [astro-ph.CO].
- [5] Maria Archidiacono, Emanuele Castorina, Diego Redigolo, and Ennio Salvioni, "Unveiling dark fifth forces with linear cosmology," JCAP 10, 074 (2022), arXiv:2204.08484 [astro-ph.CO].
- [6] Salvatore Bottaro, Emanuele Castorina, Marco Costa, Diego Redigolo, and Ennio Salvioni, "Unveiling dark forces with the Large Scale Structure of the Universe," (2023), arXiv:2309.11496 [astro-ph.CO].
- [7] Cora Dvorkin, Kfir Blum, and Marc Kamionkowski, "Constraining Dark Matter-Baryon Scattering with Linear Cosmology," Phys. Rev. D 89, 023519 (2014), arXiv:1311.2937 [astro-ph.CO].
- [8] Vera Gluscevic and Kimberly K. Boddy, "Constraints on Scattering of keV–TeV Dark Matter with Protons in the Early Universe," Phys. Rev. Lett. **121**, 081301 (2018), arXiv:1712.07133 [astro-ph.CO].
- [9] Guido D'Amico, Leonardo Senatore, and Pierre Zhang, "Limits on wCDM from the EFTofLSS with the PyBird code," JCAP 01, 006 (2021), arXiv:2003.07956 [astroph.CO].
- [10] Weishuang Linda Xu, Cora Dvorkin, and Andrew Chael, "Probing Sub-Gev Dark Matter-Baryon Scattering with Cosmological Observables," Phys. Rev. D 97, 103530 (2018), arXiv:1802.06788 [astro-ph.CO].
- [11] Keir K. Rogers, Renée Hložek, Alex Laguë, Mikhail M. Ivanov, Oliver H. E. Philcox, Giovanni Cabass, Kazuyuki Akitsu, and David J. E. Marsh, "Ultra-light axions and the S<sub>8</sub> tension: joint constraints from the cosmic microwave background and galaxy clustering," JCAP 06, 023 (2023), arXiv:2301.08361 [astro-ph.CO].
- [12] Jonathan R. Pritchard and Abraham Loeb, "21-cm cosmology," Rept. Prog. Phys. 75, 086901 (2012), arXiv:1109.6012 [astro-ph.CO].
- [13] Jeffrey B. Peterson, Kevin Bandura, and Ue Li Pen, "The Hubble Sphere Hydrogen Survey," in 41st Rencontres de Moriond: Workshop on Cosmology: Contents and Structures of the Universe (2006) pp. 283–289, arXiv:astro-ph/0606104.
- [14] Réza Ansari *et al.* (Cosmic Visions 21 cm), "Inflation and Early Dark Energy with a Stage II Hydrogen Intensity Mapping experiment," (2018), arXiv:1810.09572 [astro-ph.CO].
- [15] Andrei Mesinger, ed., The Cosmic 21-cm Revolution, 2514-3433 (IOP Publishing, 2019).
- [16] Adrian Liu, Laura Newburgh, Benjamin Saliwanchik, and Anže Slosar (Snowmass 2021 Cosmic Frontier 5 Topical Group), "Snowmass2021 Cosmic Frontier White Paper: 21cm Radiation as a Probe of Physics Across Cosmic Ages," (2022), arXiv:2203.07864 [astro-ph.CO].
- [17] Judd D. Bowman, Alan E. E. Rogers, Raul A. Monsalve, Thomas J. Mozdzen, and Nivedita Mahesh, "An absorption profile centred at 78 megahertz in the sky-averaged spectrum," Nature 555, 67–70 (2018), arXiv:1810.05912 [astro-ph.CO].

- [18] Saurabh Singh, Jishnu Nambissan T., Ravi Subrahmanyan, N. Udaya Shankar, B. S. Girish, A. Raghunathan, R. Somashekar, K. S. Srivani, and Mayuri Sathyanarayana Rao, "On the detection of a cosmic dawn signal in the radio background," Nature Astron. 6, 607–617 (2022), arXiv:2112.06778 [astro-ph.CO].
- [19] Richard Hills, Girish Kulkarni, P. Daniel Meerburg, and Ewald Puchwein, "Concerns About Modelling of the Edges Data," Nature 564, E32–E34 (2018), arXiv:1805.01421 [astro-ph.CO].
- [20] Richard F. Bradley, Keith Tauscher, David Rapetti, and Jack O. Burns, "A Ground Plane Artifact that Induces an Absorption Profile in Averaged Spectra from Global 21-cm Measurements - with Possible Application to EDGES," Astrophys. J. 874, 153 (2019), arXiv:1810.09015 [astro-ph.IM].
- [21] Peter H. Sims and Jonathan C. Pober, "Testing for calibration systematics in the EDGES low-band data using Bayesian model selection," Mon. Not. Roy. Astron. Soc. 492, 22–38 (2020), arXiv:1910.03165 [astro-ph.CO].
- [22] Rugved Pund and Anže Slosar, "Recovering 21cm Monopole Signals Without Smoothness," (2023), arXiv:2310.06134 [astro-ph.CO].
- [23] Zara Abdurashidova *et al.* (HERA), "Improved Constraints on the 21 cm EoR Power Spectrum and the X-Ray Heating of the IGM with HERA Phase I Observations," Astrophys. J. **945**, 124 (2023), arXiv:2210.04912 [astro-ph.CO].
- [24] Hiroyuki Tashiro, Kenji Kadota, and Joseph Silk, "Effects of dark matter-baryon scattering on redshifted 21 cm signals," Phys. Rev. D 90, 083522 (2014), arXiv:1408.2571 [astro-ph.CO].
- [25] Julian B. Muñoz, Ely D. Kovetz, and Yacine Ali-Haïmoud, "Heating of Baryons due to Scattering with Dark Matter During the Dark Ages," Phys. Rev. D 92, 083528 (2015), arXiv:1509.00029 [astro-ph.CO].
- [26] Julian B. Muñoz and Abraham Loeb, "A small amount of mini-charged dark matter could cool the baryons in the early Universe," Nature 557, 684 (2018), arXiv:1802.10094 [astro-ph.CO].
- [27] Rennan Barkana, "Possible interaction between baryons and dark-matter particles revealed by the first stars," Nature 555, 71–74 (2018), arXiv:1803.06698 [astroph.CO].
- [28] Rennan Barkana, Nadav Joseph Outmezguine, Diego Redigolo, and Tomer Volansky, "Strong constraints on light dark matter interpretation of the EDGES signal," Phys. Rev. D 98, 103005 (2018), arXiv:1803.03091 [hepph].
- [29] Asher Berlin, Dan Hooper, Gordan Krnjaic, and Samuel D. McDermott, "Severely Constraining Dark Matter Interpretations of the 21-cm Anomaly," Phys. Rev. Lett. **121**, 011102 (2018), arXiv:1803.02804 [hepph].
- [30] Hongwan Liu, Nadav Joseph Outmezguine, Diego Redigolo, and Tomer Volansky, "Reviving Millicharged Dark Matter for 21-Cm Cosmology," Phys. Rev. D 100, 123011 (2019), arXiv:1908.06986 [hep-ph].
- [31] Rennan Barkana, Anastasia Fialkov, Hongwan Liu, and Nadav Joseph Outmezguine, "Anticipating a New-Physics Signal in Upcoming 21-cm Power Spectrum Observations," (2022), arXiv:2212.08082 [hep-ph].
- [32] C. Evoli, A. Mesinger, and A. Ferrara, "Unveiling the nature of dark matter with high redshift 21 cm line

experiments," Journal of Cosmology and Astroparticle Physics **2014**, 024–024 (2014).

- [33] Laura Lopez-Honorez, Olga Mena, Ángeles Moliné, Sergio Palomares-Ruiz, and Aaron C. Vincent, "The 21 cm signal and the interplay between dark matter annihilations and astrophysical processes," Journal of Cosmology and Astroparticle Physics **2016**, 004–004 (2016).
- [34] Hongwan Liu and Tracy R. Slatyer, "Implications of a 21-cm signal for dark matter annihilation and decay," Physical Review D 98 (2018), 10.1103/physrevd.98.023501.
- [35] Guido D'Amico, Paolo Panci, and Alessandro Strumia, "Bounds on dark-matter annihilations from 21-cm data," Physical Review Letters **121** (2018), 10.1103/physrevlett.121.011103.
- [36] Kingman Cheung, Jui-Lin Kuo, Kin-Wang Ng, and Yue-Lin Sming Tsai, "The impact of edges 21-cm data on dark matter interactions," Physics Letters B 789, 137–144 (2019).
- [37] Andrea Mitridate and Alessandro Podo, "Bounds on dark matter decay from 21 cm line," Journal of Cosmology and Astroparticle Physics 2018, 069–069 (2018).
- [38] Steven J. Clark, Bhaskar Dutta, Yu Gao, Yin-Zhe Ma, and Louis E. Strigari, "21 cm limits on decaying dark matter and primordial black holes," Physical Review D 98 (2018), 10.1103/physrevd.98.043006.
- [39] Sandeep Kumar Acharya, Bryce Cyr, and Jens Chluba, "The role of soft photon injection and heating in 21 cm cosmology," Monthly Notices of the Royal Astronomical Society **523**, 1908–1918 (2023).
- [40] Gaétan Facchinetti, Laura Lopez-Honorez, Yuxiang Qin, and Andrei Mesinger, "21cm signal sensitivity to dark matter decay," Journal of Cosmology and Astroparticle Physics **2024**, 005 (2024).
- [41] Maxim Pospelov, Josef Pradler, Joshua T. Ruderman, and Alfredo Urbano, "Room for New Physics in the Rayleigh-Jeans Tail of the Cosmic Microwave Background," Phys. Rev. Lett. **121**, 031103 (2018), arXiv:1803.07048 [hep-ph].
- [42] Kyrylo Bondarenko, Josef Pradler, and Anastasia Sokolenko, "Constraining dark photons and their connection to 21 cm cosmology with CMB data," Phys. Lett. B 805, 135420 (2020), arXiv:2002.08942 [astroph.CO].
- [43] Andrea Caputo, Hongwan Liu, Siddharth Mishra-Sharma, Maxim Pospelov, Joshua T. Ruderman, and Alfredo Urbano, "Edges and Endpoints in 21-cm Observations from Resonant Photon Production," Phys. Rev. Lett. **127**, 011102 (2021), arXiv:2009.03899 [astroph.CO].
- [44] Andrea Caputo, Hongwan Liu, Siddharth Mishra-Sharma, Maxim Pospelov, and Joshua T. Ruderman, "Radio excess from stimulated dark matter decay," Phys. Rev. D 107, 123033 (2023), arXiv:2206.07713 [hep-ph].
- [45] Trey Driskell, Ethan O. Nadler, Jordan Mirocha, Andrew Benson, Kimberly K. Boddy, Timothy D. Morton, Jack Lashner, Rui An, and Vera Gluscevic, "Structure formation and the global 21-cm signal in the presence of Coulomb-like dark matter-baryon interactions," Phys. Rev. D 106, 103525 (2022), arXiv:2209.04499 [astroph.CO].

- [46] Wenzer Qin, Julian B. Munoz, Hongwan Liu, and Tracy R. Slatyer, "Birth of the first stars amidst decaying and annihilating dark matter," (2023), arXiv:2308.12992 [astro-ph.CO].
- [47] Sebastian A. R. Ellis, "Premature black hole death of Population III stars by dark matter," JCAP 05, 025 (2022), arXiv:2111.02414 [astro-ph.CO].
- [48] Roland de Putter, Olivier Doré, Jérôme Gleyzes, Daniel Green, and Joel Meyers, "Dark Matter Interactions, Helium, and the Cosmic Microwave Background," Phys. Rev. Lett. **122**, 041301 (2019), arXiv:1805.11616 [astroph.CO].
- [49] Kimberly K. Boddy, Vera Gluscevic, Vivian Poulin, Ely D. Kovetz, Marc Kamionkowski, and Rennan Barkana, "Critical Assessment of Cmb Limits on Dark Matter-Baryon Scattering: New Treatment of the Relative Bulk Velocity," Phys. Rev. D 98, 123506 (2018), arXiv:1808.00001 [astro-ph.CO].
- [50] Sacha Davidson, Steen Hannestad, and Georg Raffelt, "Updated Bounds on Millicharged Particles," JHEP 05, 003 (2000), arXiv:hep-ph/0001179.
- [51] A. Badertscher, P. Crivelli, W. Fetscher, U. Gendotti, S. Gninenko, V. Postoev, A. Rubbia, V. Samoylenko, and D. Sillou, "An Improved Limit on Invisible Decays of Positronium," Phys. Rev. D 75, 032004 (2007), arXiv:hep-ex/0609059.
- [52] A. A. Prinz *et al.*, "Search for Millicharged Particles at Slac," Phys. Rev. Lett. **81**, 1175–1178 (1998), arXiv:hep-ex/9804008.
- [53] Gabriel Magill, Ryan Plestid, Maxim Pospelov, and Yu-Dai Tsai, "Millicharged Particles in Neutrino Experiments," Phys. Rev. Lett. **122**, 071801 (2019), arXiv:1806.03310 [hep-ph].
- [54] Serguei Chatrchyan, V Khachatryan, Albert M Sirunyan, A Tumasyan, W Adam, E Aguilo, T Bergauer, M Dragicevic, J Erö, C Fabjan, *et al.*, "Search for fractionally charged particles in p p collisions at s= 7 tev," Physical Review D 87, 092008 (2013).
- [55] R. Acciarri *et al.* (ArgoNeuT), "Improved Limits on Millicharged Particles Using the ArgoNeuT Experiment at Fermilab," Phys. Rev. Lett. **124**, 131801 (2020), arXiv:1911.07996 [hep-ex].
- [56] Timon Emken, Rouven Essig, Chris Kouvaris, and Mukul Sholapurkar, "Direct Detection of Strongly Interacting Sub-Gev Dark Matter via Electron Recoils," JCAP 09, 070 (2019), arXiv:1905.06348 [hep-ph].
- [57] Omer Zvi Katz, Nadav Joseph Outmezguine, Diego Redigolo, and Tomer Volansky, "On the Constraining Power of Global 21-cm Cosmology at Cosmic Dawn," In preparation.
- [58] George B. Field, "Excitation of the Hydrogen 21-CM Line," Proceedings of the IRE 46, 240–250 (1958).
- [59] George B. Field, "The Time Relaxation of a Resonance-Line Profile." Astrophys. J. 129, 551 (1959).
- [60] Piero Madau, Avery Meiksin, and Martin J. Rees, "21 Centimeter Tomography of the Intergalactic Medium at High Redshift," Astrophys. J. 475, 429–444 (1997), arXiv:astro-ph/9608010 [astro-ph].
- [61] Jonathan R. Pritchard and Steven R. Furlanetto, "21 Cm Fluctuations from Inhomogeneous X-Ray Heating Before Reionization," Mon. Not. Roy. Astron. Soc. 376, 1680–1694 (2007), arXiv:astro-ph/0607234.
- [62] Andrei Mesinger, Steven Furlanetto, and Renyue Cen, "21Cmfast: a Fast, Semi-Numerical Simulation of the

High-Redshift 21-Cm Signal," Mon. Not. Roy. Astron. Soc. **411**, 955 (2011), arXiv:1003.3878 [astro-ph.CO].

- [63] Steven Furlanetto and Michael Furlanetto, "Spin Exchange Rates in Electron-Hydrogen Collisions," Mon. Not. Roy. Astron. Soc. 374, 547–555 (2007), arXiv:astro-ph/0608067.
- [64] B. Zygelman, "Hyperfine level-changing collisions of hydrogen atoms and tomography of the dark age universe," The Astrophysical Journal 622, 1356–1362 (2005).
- [65] A. C. Allison and A. Dalgarno, "Spin change in collisions of hydrogen atoms," The Astrophysical Journal 158, 423 (1969).
- [66] S. A. Wouthuysen, "On the excitation mechanism of the 21-cm (radio-frequency) interstellar hydrogen emission line." Astron. J. 57, 31–32 (1952).
- [67] Christopher M. Hirata, "Wouthuysen-Field Coupling Strength and Application to High-Redshift 21 Cm Radiation," Mon. Not. Roy. Astron. Soc. 367, 259–274 (2006), arXiv:astro-ph/0507102.
- [68] Xue-Lei Chen and Jordi Miralda-Escude, "The Spin -Kinetic Temperature Coupling and the Heating Rate Due to Lyman - Alpha Scattering Before Reionization: Predictions for 21Cm Emission and Absorption," Astrophys. J. 602, 1–11 (2004), arXiv:astro-ph/0303395.
- [69] Andrei Mesinger, Steven Furlanetto, and Renyue Cen, "21CMFAST: a fast, seminumerical simulation of the high-redshift 21-cm signal," Monthly Notices of the Royal Astronomical Society 411, 955–972 (2011), arXiv:1003.3878 [astro-ph.CO].
- [70] A. Meiksin, "Intergalactic heating by Lyman-alpha photons including hyperfine structure corrections," Res. Notes AAS 5, 126 (2021), arXiv:2105.14516 [astroph.CO].
- [71] Tejaswi Venumadhav, Liang Dai, Alexander Kaurov, and Matias Zaldarriaga, "Heating of the Intergalactic Medium by the Cosmic Microwave Background during Cosmic Dawn," Phys. Rev. D 98, 103513 (2018), arXiv:1804.02406 [astro-ph.CO].
- [72] Céline Boehm, Matthew J. Dolan, and Christopher McCabe, "A Lower Bound on the Mass of Cold Thermal Dark Matter from Planck," JCAP 08, 041 (2013), arXiv:1303.6270 [hep-ph].
- [73] Jaehong Park, Andrei Mesinger, Bradley Greig, and Nicolas Gillet, "Inferring the astrophysics of reionization and cosmic dawn from galaxy luminosity functions and the 21-cm signal," Mon. Not. Roy. Astron. Soc. 484, 933–949 (2019), arXiv:1809.08995 [astro-ph.GA].
- [74] R. J. Bouwens, G. D. Illingworth, P. A. Oesch, M. Trenti, I. Labbé, L. Bradley, M. Carollo, P. G. van Dokkum, V. Gonzalez, B. Holwerda, M. Franx, L. Spitler, R. Smit, and D. Magee, "Uv luminosity functions at redshifts z~ 4 to z ~ 10: 10,000 galaxies fromhstlegacy fields," The Astrophysical Journal 803, 34 (2015).
- [75] N. Aghanim *et al.* (Planck), "Planck 2018 results. VI. Cosmological parameters," Astron. Astrophys. **641**, A6 (2020), [Erratum: Astron.Astrophys. 652, C4 (2021)], arXiv:1807.06209 [astro-ph.CO].
- [76] Ian McGreer, Andrei Mesinger, and Valentina D'Odorico, "Model-independent evidence in favour of an end to reionization by  $z \approx 6$ ," Mon. Not. Roy. Astron. Soc. **447**, 499–505 (2015), arXiv:1411.5375 [astro-ph.CO].

- [77] B. D. Lehmer *et al.*, "The 4 Ms Chandra Deep Field-South Number Counts Apportioned by Source Class: Pervasive Active Galactic Nuclei and the Ascent of Normal Galaxies," Astrophys. J. **752**, 46 (2012), arXiv:1204.1977 [astro-ph.CO].
- [78] Anastasia Fialkov, Aviad Cohen, Rennan Barkana, and Joseph Silk, "Constraining the redshifted 21-cm signal with the unresolved soft X-ray background," Mon. Not. Roy. Astron. Soc. 464, 3498–3508 (2017), arXiv:1602.07322 [astro-ph.CO].
- [79] Stefano Marchesi *et al.*, "Mock catalogs for the extragalactic X-ray sky: simulating AGN surveys with Athena and with the AXIS probe," Astron. Astrophys. **642**, A184 (2020), arXiv:2008.09133 [astro-ph.IM].
- [80] Julian B. Muñoz, Yuxiang Qin, Andrei Mesinger, Steven G. Murray, Bradley Greig, and Charlotte Mason, "The impact of the first galaxies on cosmic dawn and reionization," Mon. Not. Roy. Astron. Soc. 511, 3657–3681 (2022), arXiv:2110.13919 [astro-ph.CO].
- [81] Giulia Despali, Carlo Giocoli, Raul E. Angulo, Giuseppe Tormen, Ravi K. Sheth, Giacomo Baso, and Lauro Moscardini, "The universality of the virial halo mass function and models for non-universality of other halo definitions," Monthly Notices of the Royal Astronomical Society 456, 2486–2504 (2016), arXiv:1507.05627 [astro-ph.CO].
- [82] Benedikt Diemer, "COLOSSUS: A python toolkit for cosmology, large-scale structure, and dark matter halos," The Astrophysical Journal Supplement Series 239, 35 (2018).
- [83] Rennan Barkana and Abraham Loeb, "Detecting the earliest galaxies through two new sources of 21cm fluctuations," Astrophys. J. 626, 1–11 (2005), arXiv:astroph/0410129.
- [84] J. Stuart B. Wyithe, Abraham Loeb, and Chris Carilli, "Improved constraints on the neutral intergalactic hydrogen surrounding quasars at redshifts z > 6," The Astrophysical Journal **628**, 575–582 (2005).
- [85] S. Peng Oh, "Reionization by hard photons: I. x-rays from the first star clusters," Astrophys. J. 553, 499 (2001), arXiv:astro-ph/0005262.
- [86] Matthew McQuinn, "Constraints on x-ray emissions from the reionization era," Monthly Notices of the Royal Astronomical Society 426, 1349–1360 (2012).
- [87] Jordan Mirocha, "Decoding the X-Ray Properties of Pre-Reionization Era Sources," Mon. Not. Roy. Astron. Soc. 443, 1211–1223 (2014), arXiv:1406.4120 [astroph.GA].
- [88] H. J. Grimm, Marat Gilfanov, and R. Sunyaev, "High Mass X-Ray Binaries as a Star Formation Rate Indicator in Distant Galaxies," Mon. Not. Roy. Astron. Soc. **339**, 793 (2003), arXiv:astro-ph/0205371.
- [89] Piero Ranalli, Andrea Comastri, and Giancarlo Setti, "The 2–10 kev luminosity as a star formation rate indicator," Astronomy & Astrophysics **399**, 39–50 (2003).
- [90] Marat Gilfanov, H. J. Grimm, and R. Sunyaev, "L(x)-SFR relation in star forming galaxies," Mon. Not. Roy. Astron. Soc. 347, L57 (2004), arXiv:astro-ph/0301331.
- [91] Giuseppina Fabbiano, "X-ray source populations in galaxies," Ann. Rev. Astron. Astrophys. 44, 323–366 (2006), arXiv:astro-ph/0511481.
- [92] S. Mineo, M. Gilfanov, and R. Sunyaev, "X-ray emission from star-forming galaxies - I. High-mass X-ray binaries," Mon. Not. Roy. Astron. Soc. 419, 2095 (2012),

arXiv:1105.4610 [astro-ph.HE].

- [93] Tassos Fragos, Bret D. Lehmer, Smadar Naoz, Andreas Zezas, and Antara R. Basu-Zych, "Energy Feedback from X-Ray Binaries in the Early Universe," Astrophys. J. Lett. **776**, L31 (2013), arXiv:1306.1405 [astroph.CO].
- [94] Arpan Das, Andrei Mesinger, Andrea Pallottini, Andrea Ferrara, and John H. Wise, "High Mass X-Ray Binaries and the Cosmic 21-Cm Signal: Impact of Host Galaxy Absorption," Mon. Not. Roy. Astron. Soc. 469, 1166– 1174 (2017), arXiv:1702.00409 [astro-ph.CO].
- [95] Aviad Cohen, Anastasia Fialkov, Rennan Barkana, and Matan Lotem, "Charting the Parameter Space of the Global 21-Cm Signal," Mon. Not. Roy. Astron. Soc. 472, 1915–1931 (2017), arXiv:1609.02312 [astroph.CO].
- [96] Bradley Greig and Andrei Mesinger, "Simultaneously constraining the astrophysics of reionization and the epoch of heating with 21cmmc," Monthly Notices of the Royal Astronomical Society 472, 2651–2669 (2017).
- [97] N. Cappelluti *et al.*, "The nature of the unresolved extragalactic soft CXB," Mon. Not. Roy. Astron. Soc. **427**, 651 (2012), arXiv:1208.4105 [astro-ph.CO].
- [98] Cyril Creque-Sarbinowski, Lingyuan Ji, Ely D. Kovetz, and Marc Kamionkowski, "Direct millicharged dark matter cannot explain the EDGES signal," Phys. Rev. D 100, 023528 (2019), arXiv:1903.09154 [astro-ph.CO].
- [99] Piero Madau, Francesco Haardt, and Martin J Rees, "Radiative transfer in a clumpy universe. iii. the nature of cosmological ionizing sources," The Astrophysical Journal **514**, 648 (1999).
- [100] Charlotte A Mason, Rohan P Naidu, Sandro Tacchella, and Joel Leja, "Model-independent constraints on the hydrogen-ionizing emissivity at z>6," Monthly Notices of the Royal Astronomical Society 489, 2669–2676 (2019).
- [101] Geoffrey C So, Michael L Norman, Daniel R Reynolds, and John H Wise, "Fully coupled simulation of cosmic reionization. ii. recombinations, clumping factors, and the photon budget for reionization," The Astrophysical Journal **789**, 149 (2014).
- [102] Nickolay Y. Gnedin and Alexander A. Kaurov, "COS-MIC REIONIZATION ON COMPUTERS. II. REION-IZATION HISTORY AND ITS BACK-REACTION ON EARLY GALAXIES," The Astrophysical Journal **793**, 30 (2014).
- [103] Ilian T. Iliev, Garrelt Mellema, Kyungjin Ahn, Paul R. Shapiro, Yi Mao, and Ue-Li Pen, "Simulating cosmic reionization: how large a volume is large enough?" Monthly Notices of the Royal Astronomical Society 439, 725–743 (2014).

- [104] Hy Trac, Renyue Cen, and Philip Mansfield, "SCORCH. i. THE GALAXY–HALO CONNECTION IN THE FIRST BILLION YEARS," The Astrophysical Journal 813, 54 (2015).
- [105] Kristian Finlator, Laura Keating, Benjamin D Oppenheimer, Romeel Davé, and Erik Zackrisson, "Reionization in technicolor," Monthly Notices of the Royal Astronomical Society 480, 2628–2649 (2018).
- [106] R Kannan, E Garaldi, A Smith, R Pakmor, V Springel, M Vogelsberger, and L Hernquist, "Introducing the iscpithesani/scpi project: radiationmagnetohydrodynamic simulations of the epoch of reionization," Monthly Notices of the Royal Astronomical Society 511, 4005–4030 (2021).
- [107] Jan-Pieter Paardekooper, Sadegh Khochfar, and Claudio Dalla Vecchia, "The first billion years project: the escape fraction of ionizing photons in the epoch of reionization," Monthly Notices of the Royal Astronomical Society 451, 2544–2563 (2015).
- [108] Hao Xu, John H. Wise, Michael L. Norman, Kyungjin Ahn, and Brian W. O'Shea, "Galaxy properties and uv escape fractions during the epoch of reionization: Results from the renaissance simulations," The Astrophysical Journal 833, 84 (2016).
- [109] Roni Harnik, Zhen Liu, and Ornella Palamara, "Millicharged Particles in Liquid Argon Neutrino Experiments," JHEP 07, 170 (2019), arXiv:1902.03246 [hepph].
- [110] Asher Berlin, Nikita Blinov, Gordan Krnjaic, Philip Schuster, and Natalia Toro, "Dark Matter, Millicharges, Axion and Scalar Particles, Gauge Bosons, and Other New Physics with LDMX," Phys. Rev. D 99, 075001 (2019), arXiv:1807.01730 [hep-ph].
- [111] Andrew Haas, Christopher S. Hill, Eder Izaguirre, and Itay Yavin, "Looking for milli-charged particles with a new experiment at the LHC," Phys. Lett. B 746, 117– 120 (2015), arXiv:1410.6816 [hep-ph].
- [112] Austin Ball *et al.*, "A Letter of Intent to Install a milli-charged Particle Detector at LHC P5," (2016), arXiv:1607.04669 [physics.ins-det].
- [113] Kevin J. Kelly and Yu-Dai Tsai, "Proton fixedtarget scintillation experiment to search for millicharged dark matter," Phys. Rev. D 100, 015043 (2019), arXiv:1812.03998 [hep-ph].
- [114] Dmitry Budker, Peter W. Graham, Harikrishnan Ramani, Ferdinand Schmidt-Kaler, Christian Smorra, and Stefan Ulmer, "Millicharged Dark Matter Detection with Ion Traps," PRX Quantum 3, 010330 (2022), arXiv:2108.05283 [hep-ph].
- [115] Asher Berlin, Hongwan Liu, Maxim Pospelov, and Harikrishnan Ramani, "The Terrestrial Density of Strongly-Coupled Relics," (2023), arXiv:2302.06619 [hep-ph].