A PARALLEL PRECONDITIONER FOR THE ALL-AT-ONCE LINEAR SYSTEM FROM EVOLUTIONARY PDES WITH CRANK-NICOLSON DISCRETIZATION*

YONG-LIANG ZHAO[†], XIAN-MING GU[‡], AND CORNELIS W. OOSTERLEE[§]

Abstract. The Crank-Nicolson (CN) method is a well-known time integrator for evolutionary partial differential equations (PDEs) arising in many real-world applications. Since the solution at any time depends on the solution at previous time steps, the CN method is inherently difficult to parallelize. In this paper, we consider a parallel method for the solution of evolutionary PDEs with the CN scheme. Using an all-at-once approach, we can solve for all time steps simultaneously using a parallelizable over time preconditioner within a standard iterative method. Due to the diagonalization of the proposed preconditioner, we can prove that most eigenvalues of preconditioned matrices are equal to 1 and the others lie in the set: $\{z \in \mathbb{C} : 1/(1 + \alpha) < |z| < 1/(1 - \alpha)$ and $\Re e(z) > 0\}$, where $0 < \alpha < 1$ is a free parameter. Besides, the efficient implementation of the proposed preconditioner is described. Given certain conditions, we prove that the preconditioned GMRES method exhibits a mesh-independent convergence rate. Finally, we will verify both theoretical findings and the efficacy of the proposed preconditioner via numerical experiments on financial option pricing PDEs.

Key words. Evolutionary PDEs, Crank-Nicolson scheme, All-at-once discretization, Parallel-in-time preconditioning, Krylov subspace methods

MSC codes. 65L05, 65N22, 65F10

1. Introduction. Mathematical models involving evolutionary partial differential equations (PDEs) arise in many diverse applications, such as fluid flow, image processing, physics-based animation, mechanical systems, earth sciences, and mathematical finance [1, 6]. We will methodically develop tools for understanding and assessing methods and software for these problems. In this work, we are interested in solving a class of evolutionary PDEs, as follows:

(1.1)
$$\begin{cases} \partial_t u(\boldsymbol{x},t) = \mathcal{L}u(\boldsymbol{x},t) + f(\boldsymbol{x},t), & (\boldsymbol{x},t) \in \Omega \times (0,T], \\ u(\cdot,0) = u_0, & \text{in } \Omega, \end{cases}$$

where $\boldsymbol{x} = (x_1, \ldots, x_d)$ and Ω denotes a bounded, convex domain in \mathbb{R}^d (d = 1, 2, 3) with smooth boundary. The spatial operator \mathcal{L} includes (but not limited to) the following choices

(1.2)
$$\mathcal{L} = \begin{cases} \nabla \cdot (a(\boldsymbol{x})\nabla) - b(\boldsymbol{x}) \cdot \nabla - c_0(\boldsymbol{x}), & \text{self-adjoint elliptic operator,} \\ \sum_{j=1}^d \kappa_j \frac{\partial^{\beta_j}}{\partial |x_j|^{\beta_j}}, & \text{Riesz fractional operator and } \beta_j \in (1,2), \end{cases}$$

one can refer to [10, 38] for the definitions of the above operators. For simplicity, we assume that the above PDEs enjoy the well-posedness and that unique solutions exist.

As most evolutionary PDEs (1.1) cannot be solved analytically, efficient numerical methods, such as finite differences, finite elements, and spectral methods, are employed to seek approximate solutions [32]. Numerical techniques proposed in the literature for such PDEs are the so-called time-stepping schemes which solve the underlying evolutionary PDEs with space (fractional) derivative(s) by marching in time sequentially, one level after the other. As many time steps may be required to balance the numerical errors arising from the spatial discretization, such time-stepping schemes can be time-consuming. This concern motivates the recent development of parallel-in-time (PinT) numerical solutions for evolutionary PDEs, including, e.g., the inverse Laplace transform method [30], the MGRIT method [11] and the parareal method [25]. Instead of solving PDEs in a sequential manner, this class of PinT methods, i.e., the space-time methods, solves the evolutionary PDEs at all time levels simultaneously by performing an

^{*}Received by the editors DATE;

Funding: This research is supported by the Sichuan Science and Technology Program (Nos. 2022ZYD0006 and 2023NS-FSC1326) and Natural Science Foundation of Sichuan Province (No. 2022NSFSC1815). X.-M. Gu is also supported by the China Scholarship Council (CSC) and Guanghua Talent Project of Southwestern University of Finance and Economics.

[†]School of Mathematical Sciences, Sichuan Normal University, Chengdu, Sichuan 610066, P.R. China (ylzhaofde@sina.com).

[‡]Corresponding author. School of Mathematics, Southwestern University of Finance and Economics, Chengdu, Sichuan 611130, P.R. China (guxianming@live.cn, guxm@swufe.edu.cn).

[§]Mathematical Institute, Utrecht University, Budapestlaan 6, 3584 CD Utrecht, The Netherlands (c.w.oosterlee@uu.nl).

all-at-once (AaO) discretization that results into a large-scale linear system that is typically solved by preconditioned iterative solvers (e.g., preconditioned Krylov subspace methods); refer to [14, 28, 35] for details.

How to obtain the AaO linear system and establish the preconditioner is related to the choice of the time integrator for ODEs, based on the spatially semi-discretized evolutionary PDEs. Recently, McDonald et al. proposed in [28] a block circulant (BC) preconditioner to accelerate the convergence of Krylov subspace solvers for the AaO linear system arising from p-step BDF integrators for evolutionary PDEs. Parallel experiments with the BC preconditioner are reported by Goddard and Wathen in [15]. In [23], a generalized BC preconditioner (gBC) has been proposed by Lin and Ng, who introduced a parameter $\alpha \in (0,1)$ into the top-right block of the BC preconditioner that can be fine-tuned to handle efficiently the case of very small diffusion coefficients. Both the BC and gBC preconditioners use a modified diagonalization technique that is originally proposed by Maday and Rønquist [27]. Some related PinT methods and preconditioners can be found in [19, 20]. However, most of above mentioned studies focus on the temporal discretization at the integer nodes (i.e., t_k and $k \in \mathbb{Z}$) and it is not easy to extend their theoretical results to temporal discretizations at non-integer nodes (e.g., $t_{k+\theta}$ and $\theta \notin \mathbb{Z}$). Besides, for some particular PDEs, their space-discretization matrices have zero eigenvalues, which makes the corresponding BC preconditioner singular and infeasible. Although the gBC preconditioner in [23] can overcome the above deficiency, the detailed analysis of this issue is incomplete and only available for the AaO system based on the implicit Euler method and real symmetric definite space-discretization matrices.

As the implicit Crank-Nicolson (CN) scheme [3] is a popular option for evolutionary PDEs with unconditional convergence properties, we consider this method here. However, how to make the CN scheme suitable for PinT implementation and the corresponding theoretical analyses is rarely investigated. In fact, after the spatial discretization of the model (1.1), we reformulate the CN scheme for the semi-discretized system of ODEs into an AaO system, whose coefficient matrix is a block lower triangular Toeplitz matrix. Then, we estimate the condition number of the AaO system and adapt the gBC preconditioner to it. The invertibility of the gBC preconditioner is discussed differently from the work [24]. We prove that most eigenvalues of the preconditioned matrices are clustered around one and the remaining eigenvalues are located at a special set if the space-discretized matrix is negative semi-definite, which coincides with unconditional stability and convergence of the CN method. With the help of a certain condition on parameter α , we also derive a mesh-independent convergence rate for the preconditioned GMRES [9]. It is worth noting that there are some similar PinT preconditioners and the corresponding spectral analysis available for the θ -scheme (e.g., if $\theta = \frac{1}{2}$, it reduces to the CN scheme), but they assume that the spatial matrix is symmetric positive definite and only prove that the modules of the remaining eigenvalues are located at the interval $[1 - \epsilon, 1 + \epsilon]$ with $\epsilon > 0$, being unspecific about the information of the space-discretized matrix and α ; see [13], for details. Compared to the recent results, our spectral analysis of preconditioned matrices is the sharpest. We also discuss when the BC preconditioner becomes singular (see Section 5, for an example). With the help of matrix analysis, another advantage of our gBC precondition is that in practical implementation, we only need to solve the first half of a sequence of complex-shifted linear systems involving negative (semi-)definite space-discretized matrices. Estimates of the computational costs and memory requirements of the associated gBC preconditioner are conducted too. As we will illustrate in Section 5, the gBC preconditioner can be modified to handle some financial PDEs with time-varying coefficients without losing its parallelism.

The remainder of this paper is organized as follows. In the next section, the AaO linear system derived from the CN scheme for evolutionary PDEs is presented. Subsequently, the invertibility of the pertinent AaO system is proved, and its condition number is estimated. In Section 3, the gBC preconditioner is adapted and both the spectral properties and efficient implementation of the preconditioned system are analyzed rigorously. Section 4 proves that the convergence rate of the preconditioned GM-RES is independent of mesh size under certain conditions. In Section 5, numerical experiments involving European option pricing PDEs are reported to verify our theoretical results and the effectiveness of the designed parallel preconditioner. Finally, the paper closes with conclusions in Section 6.

2. The CN scheme and its AaO system. In this section, we first derive the CN scheme for approximating Eq. (1.1) and give some related analyses.

2.1. The CN scheme. For a given positive integer M, let $\tau = \frac{T}{M}$ be the time step size, to obtain time steps $t_k = k\tau$ (k = 0, 1, ..., M). Let $A \in \mathbb{R}^{N \times N}$ be the spatial discretization matrix of \mathcal{L} , via a finite difference method.

Combining the CN time integrator, the matrix form of the CN scheme of Eq. (1.1) is derived as

(2.1)
$$\frac{\boldsymbol{u}^{k+1} - \boldsymbol{u}^k}{\tau} = A \frac{\boldsymbol{u}^{k+1} + \boldsymbol{u}^k}{2} + \boldsymbol{f}^{k+\frac{1}{2}}, \quad k = 0, 1, \dots, M-1,$$

where \boldsymbol{u}^k is a lexicographically ordered vector collecting the approximate solution of $u(\cdot, t_k)$ over all the spatial grid points. Similarly, $\boldsymbol{f}^{k+\frac{1}{2}}$ is a lexicographically ordered vector collecting $f(\cdot, t_{k+\frac{1}{2}})$ over all the spatial points. Here $t_{k+\frac{1}{2}} = (k+\frac{1}{2})\tau$. To investigate the conditions for proving the unconditional stability and convergence of the CN scheme (2.1), we need the following lemma.

LEMMA 2.1. ([18]) Let the rational function R(z) be bounded for $\Re e(z) \leq 0$ and assume that the matrix $A \in \mathbb{R}^{N \times N}$ is negative semi-definite, i.e., $\langle \boldsymbol{y}, A \boldsymbol{y} \rangle \leq 0$ for all $\boldsymbol{y} \in \mathbb{R}^{N}$. Then, we have

$$||R(A)||_2 \le \sup_{\Re e(z) \le 0} |R(z)|.$$

Here $\Re e(\cdot)$ is the real part of a complex number.

At this stage, we can theoretically prove that the CN scheme (2.1) is unconditionally stable.

THEOREM 2.2. If the spatial matrix A is negative semi-definite, then the CN scheme (2.1) is unconditionally stable.

Proof. Let u^n and \tilde{u}^n be the exact and numerical solutions of (2.1), and the error $e^n = u^n - \tilde{u}^n$. Then, from (2.1), we get the error equation:

$$\left(I-\frac{\tau}{2}A\right)e^{k+1}=\left(I+\frac{\tau}{2}A\right)e^k.$$

Since A is negative semi-definite, $I - \frac{\tau}{2}A$ is invertible. Therefore, one gets

(2.2)
$$e^{k+1} = R_1\left(\frac{\tau}{2}A\right)e^k,$$

with $R_1(z) = \frac{1+z}{1-z}$. By Lemma 2.1, it follows that $\left\| R_1\left(\frac{\tau}{2}A\right) \right\|_2 \le \sup_{z \in [0, 1]} C_2$

$$\left|R_1\left(\frac{\tau}{2}A\right)\right\|_2 \le \sup_{\Re e(z) \le 0} |R(z)| = \sup_{\Re e(z) = 0} |R(z)| = 1.$$

From (2.2), we can conclude that

$$\|\boldsymbol{e}^{k+1}\|_{2} = \left\|R_{1}\left(\frac{\tau}{2}A\right)\boldsymbol{e}^{k}\right\|_{2} \le \left\|R_{1}\left(\frac{\tau}{2}A\right)\right\|_{2} \|\boldsymbol{e}^{k}\|_{2} \le \|\boldsymbol{e}^{k}\|_{2}.$$

This completes the proof.

With the help of the same conditions of Theorem 2.2 and the spatial discretization, it is easy to combine Theorem 2.2 with Lemma 2.1 to present and prove the convergence of the CN scheme (2.1). In this section, we do not yet specify which spatial discretization is chosen to obtain the spatial matrix, thus here we omit the specific convergence result of the CN scheme (2.1) for clarity.

2.2. CN AaO system. In order to construct the parallel method based on the CN scheme, we will derive the so-called AaO linear system. Before deriving the CN AaO system, several auxiliary symbols have to be introduced: **0** is the zero vector or matrix with suitable size, I_s represents an identity matrix with size N. Denote

$$oldsymbol{u} = \left[ig(oldsymbol{u}^{ op},ig(oldsymbol{u}^2ig)^{ op},ig(oldsymbol{u}^3ig)^{ op},ig(oldsymbol{h}^3ig)^{ op},ig(oldsymbol{f}^3ig)^{ op},\dots,ig(oldsymbol{f}^{M-rac{1}{2}}ig)^{ op}
ight]^{ op},$$

then, the AaO system can be written as:

$$\mathcal{M}\boldsymbol{u} = \boldsymbol{\xi} + \tau \boldsymbol{f} := \boldsymbol{b},$$

where

$$\boldsymbol{\xi} = \left[\left[\left(I_s + \frac{1}{2} \tilde{A} \right) \boldsymbol{u}^0 \right]^\top, \boldsymbol{0}^\top, \dots, \boldsymbol{0}^\top \right]^\top \quad \text{and} \quad \mathcal{M} = B_1 \otimes I_s - B_2 \otimes \tilde{A}$$

with $\tilde{A} = \tau A$, $B_1 = \text{tridiag}(-1, 1, 0) \in \mathbb{R}^{M \times M}$ and $B_2 = \frac{1}{2} \text{tridiag}(1, 1, 0) \in \mathbb{R}^{M \times M}$.

Obviously, a general sparse direct solver can be applied to system (2.3), but the computational cost will then be $\mathcal{O}(M^3N^3)$. Considering the special structure of \mathcal{M} , if a Gaussian elimination based block forward substitution method [16] is used to solve (2.3), the complexity is of order $\mathcal{O}(MN^3 + MN^2)$. This is still expensive, especially for high-dimensional problems with a fine mesh. On the other hand, we can estimate the condition number of the coefficient matrix of Eq. (2.3), this will guide us in the design of an efficient solver for such large AaO systems.

THEOREM 2.3. Suppose that $\Re(\lambda(A)) \leq 0$. Then, the following bounds hold for the critical singular values of \mathcal{M} in (2.3):

$$\sigma_{\max}(\mathcal{M}) \le 2 + \tau ||A||_2, \quad \sigma_{\min}(\mathcal{M}) \ge \frac{1}{M},$$

so that $cond(\mathcal{M}) \leq 2M + T ||A||_2$.

Proof. Since both $\Re e(\lambda(A)) \leq 0$ and $\lambda(B_2) > 0$, we can claim that $\|\mathcal{M}\mathbf{z}\|_2 \geq \|(B_1 \otimes I)\mathbf{z}\|_2$ for any vector \mathbf{z} . In particular, using also the properties of the Kronecker product, it follows that $\sigma_{\min}(\mathcal{M}) \geq \sigma_{\min}(B_1)$. The rest of this proof is similar to the idea in [5], we omit the details.

It is natural that the condition number depends linearly on all main properties of the system: the number of time steps, time interval length, and the norm of the space-discretized matrix. In order to solve such large-scale AaO linear systems efficiently, researchers prefer to use preconditioned Krylov subspace solvers, which will be discussed in the next section.

3. A PinT preconditioner and the spectra analysis. In this section, we construct an efficient PinT preconditioner for the system (2.3), and analyze the eigenvalue distribution of the preconditioned matrix. The following assumption about the matrix A is needed for this purpose.

Assumption 3.1. Assume that matrix A is negative semi-definite and diagonalizable.

It is worth noting that this assumption can easily be fulfilled such as $\mathcal{L} = -\kappa \Delta^2 + \Delta$ ($\kappa \in \mathbb{R}^+$) and Eq. (1.2).

3.1. The PinT preconditioner. We replace B_1 and B_2 by the so-called α -circulant matrices

$$C_{1}^{(\alpha)} = \begin{bmatrix} 1 & & -\alpha \\ -1 & 1 & & \\ & \ddots & \ddots & \\ & & -1 & 1 \end{bmatrix} \in \mathbb{R}^{M \times M} \text{ and } C_{2}^{(\alpha)} = \frac{1}{2} \begin{bmatrix} 1 & & \alpha \\ 1 & 1 & & \\ & \ddots & \ddots & \\ & & 1 & 1 \end{bmatrix} \in \mathbb{R}^{M \times M},$$

respectively. Here $\alpha \in (0,1)$ is a free parameter. Then, we obtain our PinT preconditioner:

$$P_{\alpha} = C_1^{(\alpha)} \otimes I_s - C_2^{(\alpha)} \otimes \tilde{A}$$

Let $\mathbb{F} = \left[\frac{\omega^{jk}}{\sqrt{M}}\right]_{j,k=0}^{M-1}$ be the discrete Fourier matrix, where $\omega = e^{-2\pi \mathbf{i}/M}(\mathbf{i} = \sqrt{-1})$. We know that α -circulant matrices $C_{\ell}^{(\alpha)}$ ($\ell = 1, 2$) can be diagonalized as $C_{\ell}^{(\alpha)} = V_{\alpha} D_{\ell} V_{\alpha}^{-1}$, with

$$V_{\alpha} = \Lambda_{\alpha}^{-1} \mathbb{F}^* \quad \text{and} \quad D_{\ell} = \operatorname{diag}\left(\sqrt{M} \mathbb{F} \Lambda_{\alpha} C_{\ell}^{(\alpha)}(:,1)\right) = \operatorname{diag}\left(\lambda_{\ell,1}, \ldots, \lambda_{\ell,M}\right),$$

where $\Lambda_{\alpha} = \operatorname{diag}\left(1, \alpha^{\frac{1}{M}}, \ldots, \alpha^{\frac{M-1}{M}}\right)$, '*' denotes the conjugate transpose of a matrix, $C_{\ell}^{(\alpha)}(:, 1)$ is the first column of $C_{\ell}^{(\alpha)}$ and

(3.2)
$$\lambda_{1,k} = 1 - \alpha^{\frac{1}{M}} e^{\frac{2(k-1)\pi \mathbf{i}}{M}}, \quad \lambda_{2,k} = \frac{1}{2} + \frac{1}{2} \alpha^{\frac{1}{M}} e^{\frac{2(k-1)\pi \mathbf{i}}{M}} \quad (k = 1, \dots, M).$$

Moreover, for $\alpha \in (0, 1)$, the real part of $\lambda_{1,k}$ is positive, i.e., $\Re e(\lambda_{1,k}) > 0$. Similarly, $\Re e(\lambda_{2,k}) > 0$.

With these decompositions, we have

$$P_{\alpha} = (V \otimes I_s) \left(D_1 \otimes I_s - D_2 \otimes \tilde{A} \right) \left(V^{-1} \otimes I_s \right).$$

Thus, for a given vector $\boldsymbol{v}, \boldsymbol{z} = P_{\alpha}^{-1} \boldsymbol{v}$ can be computed via the following three steps:

(3.3)
$$\begin{aligned} \text{step-(a)} \quad & \boldsymbol{z}_1 = \left(V^{-1} \otimes I_s\right) \boldsymbol{v}, \\ \text{step-(b)} \quad & \left(\lambda_{1,k}I_s - \lambda_{2,k}\tilde{A}\right) \boldsymbol{z}_{2,k} = \boldsymbol{z}_{1,k}, \quad k = 1, 2, \dots, M, \\ \text{step-(c)} \quad & \boldsymbol{z} = \left(V \otimes I_s\right) \boldsymbol{z}_2, \end{aligned}$$

where $\mathbf{z}_{\ell,k} = \mathbf{z}_{\ell} ((k-1)N:kN)$ ($\ell = 1, 2$) represents the k-th block of \mathbf{z}_{ℓ} . Combining $\Re e(\lambda_{\ell,k}) > 0$ ($\ell = 1, 2$) and Assumption 3.1, we know that P_{α} is nonsingular.

In (3.3), step-(a) and step-(c) can be computed efficiently via FFTs (i.e., fast Fourier transforms). In step-(b), the complex-shifted linear systems can be solved by using some sparse direct solvers, since A is generally a sparse matrix. For example, if $\mathcal{L} = \kappa \Delta$, the fast discrete sine transform (DST) is used in step-(b).

Remark 3.2. Similar to the tricks in [17, 23], we find that the complex eigenvalues/eigenvectors of $C_{\ell}^{(\alpha)}$ ($\ell = 1, 2$) appear in conjugate pairs. Thus, we only need to solve for the first $\left\lceil \frac{M+1}{2} \right\rceil$ systems in step-(b). The remaining systems are solved by conjugating the obtained solutions in order to reduce the computational cost.

Remark 3.3. For the preconditioner P_1 (i.e., $\alpha = 1$), the invertibility is equivalent to the invertibility of the complex matrices $\lambda_{1,k}I - \lambda_{2,k}\tilde{A}$ $(k = 1, 2, \dots, M)$. However, when $\alpha = 1$, then it obviously follows that $\lambda_{1,1} = 0$ and $\lambda_{2,k} = 1$, cf. Eq. (3.2). If the space-discretization matrix A of some particular PDEs is singular, it implies that the preconditioner P_1 becomes singular and fails to be a valid preconditioner. Note that this difficulty has not been discussed and shown in the existing literatures [23, 26, 28, 36].

3.2. The spectral analysis. Let

$$Q_1 = I_s - \frac{1}{2}\tilde{A}$$
 and $Q_2 = I_s + \frac{1}{2}\tilde{A}$.

Then,

$$\mathcal{M} = \begin{bmatrix} Q_1 & & & \\ -Q_2 & Q_1 & & \\ & \ddots & \ddots & \\ & & -Q_2 & Q_1 \end{bmatrix} \quad \text{and} \quad P_\alpha = \begin{bmatrix} Q_1 & & & -\alpha Q_2 \\ -Q_2 & Q_1 & & \\ & \ddots & \ddots & \\ & & -Q_2 & Q_1 \end{bmatrix}$$

From the above expressions, we can confirm that the matrix \mathcal{M} is invertible since Q_1 is invertible. With the help of these expressions, the following result is true.

LEMMA 3.4. The preconditioned matrix $P_{\alpha}^{-1}\mathcal{M}$ has (M-1)N eigenvalues equal to 1 and N eigenvalues equal to those of the matrix inverse $(I_s - \alpha J_M)^{-1}$, where $J_M = (Q_1^{-1}Q_2)^M$.

Proof. It is difficult to calculate $P_{\alpha}^{-1}\mathcal{M}$ directly. Thus, we first calculate $\mathcal{M}^{-1}P_{\alpha}$. We rewrite \mathcal{M}

and P_{α} into the form $\mathcal{M} = (I_t \otimes Q_1) \tilde{\mathcal{M}}$ and $P_{\alpha} = (I_t \otimes Q_1) \tilde{P_{\alpha}}$, where I_t is an identity matrix of size M,

$$\tilde{\mathcal{M}} = \begin{bmatrix} I_s & & & \\ -Q_1^{-1}Q_2 & I_s & & \\ & \ddots & \ddots & \\ & & -Q_1^{-1}Q_2 & I_s \end{bmatrix}, \quad \tilde{P}_{\alpha} = \begin{bmatrix} I_s & & -\alpha Q_1^{-1}Q_2 \\ -Q_1^{-1}Q_2 & I_s & & \\ & \ddots & \ddots & \\ & & -Q_1^{-1}Q_2 & I_s \end{bmatrix}.$$

By simple calculation, we have

$$\tilde{\mathcal{M}}^{-1} = \begin{bmatrix} I_s & & & \\ J_1 & I_s & & \\ \vdots & \ddots & \ddots & \\ J_{M-1} & \cdots & J_1 & I_s \end{bmatrix},$$

where $J_k = \left(Q_1^{-1}Q_2\right)^k$ (k = 1, 2, ...). Then, we arrive at

$$\mathcal{M}^{-1}P_{\alpha} = \tilde{\mathcal{M}}^{-1}\tilde{P}_{\alpha} = \begin{bmatrix} I_s & \mathbf{0} & \cdots & \mathbf{0} & -\alpha J_1 \\ & I_s & \ddots & \ddots & & -\alpha J_2 \\ & & \ddots & \ddots & & \vdots \\ & & & I_s & -\alpha J_{M-1} \\ & & & & & I_s - \alpha J_M \end{bmatrix}$$

With this at hand, we obtain

$$P_{\alpha}^{-1}\mathcal{M} = \left(\mathcal{M}^{-1}P_{\alpha}\right)^{-1} = \left(\tilde{\mathcal{M}}^{-1}\tilde{P}_{\alpha}\right)^{-1} = \begin{bmatrix} I_{s} & \mathbf{0} & \cdots & \mathbf{0} & \alpha J_{1} \left(I_{s} - \alpha J_{M}\right)^{-1} \\ I_{s} & \ddots & \ddots & \alpha J_{2} \left(I_{s} - \alpha J_{M}\right)^{-1} \\ & \ddots & \ddots & \vdots \\ & I_{s} & \alpha J_{M-1} \left(I_{s} - \alpha J_{M}\right)^{-1} \\ & & \left(I_{s} - \alpha J_{M}\right)^{-1} \end{bmatrix}.$$

The proof is completed.

It is worth mentioning that this proof is different from [28], since the Sherman-Morrison-Woodbury formula fails in our case. The reason is that the matrix Q_2 may singular. Recall the proof of Lemma 3.4, the invertibility of $I_s - \alpha J_M$ is not proved. Fortunately, it can be obtained from the following lemma.

LEMMA 3.5. The eigenvalues of $Q_1^{-1}Q_2$ satisfy

$$-1 < \Re e \left(\lambda \left(Q_1^{-1} Q_2 \right) \right) \le 1 \quad \text{and} \quad 0 \le \left| \lambda \left(Q_1^{-1} Q_2 \right) \right| \le 1.$$

Proof. From Assumption 3.1, it is not difficult to get that

$$\lambda\left(Q_1^{-1}Q_2\right) = \frac{\lambda\left(Q_2\right)}{\lambda\left(Q_1\right)} = \frac{1 + \frac{\tau}{2}\lambda\left(A\right)}{1 - \frac{\tau}{2}\lambda\left(A\right)}.$$

We know that all eigenvalues of A can be written in the form $\lambda(A) = \eta + \xi \mathbf{i}$. By Assumption 3.1, we have $\eta, \xi \in \mathbb{R}$ and $\eta \leq 0$. Then, we have

$$\frac{1 + \frac{\tau}{2}\lambda(A)}{1 - \frac{\tau}{2}\lambda(A)} = \frac{1 - \left(\frac{\tau}{2}\eta\right)^2 - \left(\frac{\tau}{2}\xi\right)^2}{\left(1 - \frac{\tau}{2}\eta\right)^2 + \left(\frac{\tau}{2}\xi\right)^2} + \frac{\tau\xi}{\left(1 - \frac{\tau}{2}\eta\right)^2 + \left(\frac{\tau}{2}\xi\right)^2}\mathbf{i}.$$

We first estimate the real part of the above expression. Obviously,

$$\frac{1 - \left(\frac{\tau}{2}\eta\right)^2 - \left(\frac{\tau}{2}\xi\right)^2}{\left(1 - \frac{\tau}{2}\eta\right)^2 + \left(\frac{\tau}{2}\xi\right)^2} = -1 + \frac{2 - \tau\eta}{\left(1 - \frac{\tau}{2}\eta\right)^2 + \left(\frac{\tau}{2}\xi\right)^2} > -1$$

On the other hand, we obtain

$$\left(1 - \frac{\tau}{2}\eta\right)^2 + \left(\frac{\tau}{2}\xi\right)^2 - \left[1 - \left(\frac{\tau}{2}\eta\right)^2 - \left(\frac{\tau}{2}\xi\right)^2\right] = -\tau\eta + \frac{\tau^2}{2}\eta^2 + \frac{\tau^2}{2}\xi^2 \ge 0$$

Thus, we arrive at

$$-1 < \Re e\left(\lambda\left(Q_1^{-1}Q_2\right)\right) \le 1.$$

Next, we turn to show the upper bound of $|\lambda(Q_1^{-1}Q_2)| \ge 0$. By some calculations, we have

$$\left[1 - \left(\frac{\tau^2}{4}\eta^2 + \frac{\tau^2}{4}\xi^2\right)\right]^2 + \tau^2\xi^2 - \left[(1 - \tau\eta) + \left(\frac{\tau^2}{4}\eta^2 + \frac{\tau^2}{4}\xi^2\right)\right]^2$$
$$= 2\tau\eta\left(\frac{\tau^2}{4}\eta^2 + \frac{\tau^2}{4}\xi^2 - \tau\eta + 1\right) \le 0.$$

This implies that $\left|\lambda\left(Q_1^{-1}Q_2\right)\right| \leq 1$.

With the help of Lemma 3.5, we can immediately show the nonsingularity of $(I_s - \alpha J_M)$.

LEMMA 3.6. Assume $\alpha \in (0,1)$. The matrix $(I_s - \alpha J_M)$ is nonsingular and its eigenvalues satisfy

 $1 - \alpha \leq \Re e \left(\lambda \left(I_s - \alpha J_M \right) \right) \leq 1 + \alpha \quad \text{and} \quad 1 - \alpha \leq \left| \lambda \left(I_s - \alpha J_M \right) \right| \leq 1 + \alpha.$

Proof. The eigenvalues of $(I_s - \alpha J_M)$ can be expressed as

$$\lambda \left(I_s - \alpha J_M \right) = 1 - \alpha \left[\lambda \left(Q_1^{-1} Q_2 \right) \right]^M.$$

Based on Lemma 3.5, we rewrite the (complex) eigenvalues of $Q_1^{-1}Q_2$ in polar form: $\lambda (Q_1^{-1}Q_2) = r (\cos \theta + \mathbf{i} \sin \theta)$, where $0 \le r \le 1$ and $\theta \in (-\pi, \pi)$. With the help of De Moivre's formula, we have

$$0 < 1 - \alpha \le \Re e \left(\lambda \left(I_s - \alpha J_M \right) \right) = 1 - \alpha r^M \cos \left(M \theta \right) \le 1 + \alpha.$$

This also implies that the matrix $(I_s - \alpha J_M)$ is nonsingular.

Using Lemma 3.5, we obtain

$$1 - \alpha \le \left| \Re e\left(\lambda \left(I_s - \alpha J_M \right) \right) \right| \le \left| \lambda \left(I_s - \alpha J_M \right) \right| \le 1 + \alpha \left| \lambda \left(Q_1^{-1} Q_2 \right) \right|^M \le 1 + \alpha.$$

This completes the proof.

Combining Lemma 3.4 and Lemma 3.6, we obtain the spectral distribution of $P_{\alpha}^{-1}\mathcal{M}$.

THEOREM 3.7. Assume $\alpha \in (0,1)$ and that the spatial matrix A fulfills Assumption 3.1. Then, the preconditioned matrix $P_{\alpha}^{-1}\mathcal{M}$ has (M-1)N eigenvalues equal to 1 and N eigenvalues distributed in the set

$$\Omega_{\alpha} = \left\{ z \in \mathbb{C} : \frac{1}{1+\alpha} \le |z| \le \frac{1}{1-\alpha} \text{ and } \Re e(z) > 0 \right\}.$$

Proof. According to Lemma 3.4, we only need to analyze the distribution of the eigenvalues of $(I_s - \alpha J_M)^{-1}$.

From Lemma 3.6, we know

$$\frac{1}{1+\alpha} \le \left| \lambda \left(\left(I_s - \alpha J_M \right)^{-1} \right) \right| \le \frac{1}{1-\alpha}$$

and $\Re \left(\lambda \left(\left(I_s - \alpha J_M \right)^{-1} \right) \right) > 0$, which completes the proof.

Moreover, if we tighten Assumption 3.1 so that the spatial matrix A is symmetric negative semidefinite, then the above result can be simplified.

COROLLARY 3.8. If the spatial matrix A is symmetric negative semi-definite and $\alpha \in (0,1)$, then the preconditioned matrix $P_{\alpha}^{-1}\mathcal{M}$ has (M-1)N eigenvalues equal to 1 and N eigenvalues distributed in the interval $\left[\frac{1}{1+\alpha}, \frac{1}{1-\alpha}\right]$.

Proof. Since the spatial matrix A is symmetric negative semi-definite, both matrices Q_1 and Q_2 are real symmetric, then $I_s - \alpha J_M$ will be real symmetric and all its (real) eigenvalues will be located in the interval $[1 - \alpha, 1 + \alpha]$, according to Lemma 3.5 and Lemma 3.6. Therefore, each eigenvalue, $z \in \mathbb{R}$, of the matrix $(I_s - \alpha J_M)^{-1}$ satisfies $z \in \left[\frac{1}{1+\alpha}, \frac{1}{1-\alpha}\right]$. Following the proof of Theorem 3.7, the target result is obtained.

Remark 3.9. Compared with the stability condition required by the CN scheme (cf. Theorem 2.2), we only need to assume that the spatial matrix A can be diagonalized to give our eigenvalue analysis for the preconditioned matrix. Reviewing Theorem 3.7, the upper bound can be reached, if the spatial matrix A has some zero eigenvalues; refer to Section 5 for a discussion.

4. Convergence analysis of the preconditioned GMRES method. Let $\boldsymbol{u}^{(k)}$ denote the numerical solution at the k-th iteration of GMRES, which is exploited to the left-preconditioned system $\mathcal{P}_{\alpha}^{-1}\mathcal{M}\boldsymbol{u}^{(k)} = \mathcal{P}_{\alpha}^{-1}\boldsymbol{b}$ with a given initial guess $\boldsymbol{u}^{(0)}$. Then, the residual vector $\boldsymbol{r}_k := \mathcal{P}_{\alpha}^{-1} \left(\boldsymbol{b} - \mathcal{M}\boldsymbol{u}^{(k)} \right)$ satisfies the minimal residual property [29]

(4.1)
$$\|\boldsymbol{r}_{k}\|_{2} = \min_{p \in \mathbb{P}_{k}} \left\| p\left(\mathcal{P}_{\alpha}^{-1}\mathcal{M}\right) \boldsymbol{r}_{0} \right\|_{2} \leq \left(\min_{p \in \mathbb{P}_{k}} \left\| p(\mathcal{P}_{\alpha}^{-1}\mathcal{M}) \right\|_{2}\right) \|\boldsymbol{r}_{0}\|_{2},$$

where \mathbb{P}_k denotes the set of polynomials p of degree k or less with p(0) = 1. For a given initial \mathbf{r}_0 , the last "worst-case" inequality may be significantly overestimated in some situations [8, 33]. In general, it is not easy to directly estimate the norm $\|p(\mathcal{P}_{\alpha}^{-1}\mathcal{M})\|_2$ from only the spectral information of the nonnormal matrix $\mathcal{P}_{\alpha}^{-1}\mathcal{M}$. However, if $\mathcal{P}_{\alpha}^{-1}\mathcal{M}$ is diagonalizable, then the above estimate (4.1) can be improved. In fact, we have the following conclusion.

THEOREM 4.1. Assume $\alpha \in (0,1)$ and the spatial matrix A fulfilling Assumption 3.1. The preconditioned matrix $P_{\alpha}^{-1}\mathcal{M}$ is diagonalizable for any $\alpha \in (0,1)$.

Proof. We first claim that the matrix $I_s - \alpha J_M$ is diagonalizable. Due to Assumption 3.1, the eigendecomposition of \tilde{A} is given by $\tilde{A} = V_a D_a V_a^{-1}$ and then

(4.2)
$$Q_1^{-1}Q_2 = V_a \left(I_s - \frac{1}{2}D_a \right)^{-1} V_a^{-1} V_a \left(I_s + \frac{1}{2}D_a \right) V_a^{-1} := V_a D_J V_a^{-1}$$

with $D_J := \left(I_s - \frac{1}{2}D_a\right)^{-1} \left(I_s + \frac{1}{2}D_a\right)$ being a diagonal matrix whose diagonal entries are the eigenvalues of $Q_1^{-1}Q_2$. Moreover, we have the following matrix diagonalization,

$$I_s - \alpha J_M = I_s - \alpha (Q_1^{-1} Q_2)^M = V_a \left[I_s - \alpha (D_J)^M \right] V_a^{-1},$$

and its inverse $(I_s - \alpha J_M)^{-1}$ can be factorized as

$$(I_s - \alpha J_M)^{-1} - I_s = V_a \left[\left(I_s - \alpha (D_J)^M \right)^{-1} - I_s \right] V_a^{-1} := V_a \Theta V_a^{-1}.$$

A tedious but simple calculation yields the following diagonalization:

$$P_{\alpha}^{-1}\mathcal{M}-\mathcal{I} = \underbrace{\begin{bmatrix} I_{s} & \mathbf{0} & \cdots & \mathbf{0} & \alpha J_{1}V_{a} \\ & I_{s} & \ddots & \ddots & \alpha J_{2}V_{a} \\ & & \ddots & \ddots & \vdots \\ & & & I_{s} & \alpha J_{M-1}V_{a} \\ & & & & V_{a} \end{bmatrix}}_{\widetilde{\Theta}} \underbrace{\begin{bmatrix} \mathbf{0} & & & \\ & \mathbf{0} & & \\ & & \ddots & \ddots & \alpha J_{2}V_{a} \\ & & & \ddots & \ddots & \alpha J_{2}V_{a} \\ & & & \ddots & \ddots & \vdots \\ & & & & I_{s} & \alpha J_{M-1}V_{a} \\ & & & & V_{a} \end{bmatrix}}^{-1},$$

which results in the desired matrix decomposition

$$P_{\alpha}^{-1}\mathcal{M} = \mathcal{S}\left(\mathcal{I} + \widetilde{\Theta}\right)\mathcal{S}^{-1} := \mathcal{S}\Upsilon\mathcal{S}^{-1}$$

with the diagonal entries of Υ being composed of the spectrum set $\sigma(\mathcal{P}_{\alpha}^{-1}\mathcal{M})$.

Based on Theorem 4.1, we naturally derive a practical upper bound

$$\begin{split} \|p(\mathcal{P}_{\alpha}^{-1}\mathcal{M})\|_{2} &= \|\mathcal{S}p(\Upsilon)\mathcal{S}^{-1}\|_{2} \\ &\leq \|\mathcal{S}\|_{2}\|p(\Upsilon)\|_{2}\|\mathcal{S}^{-1}\|_{2} \\ &= \kappa(\mathcal{S})\left(\max_{\lambda\in\sigma(\mathcal{P}_{\alpha}^{-1}\mathcal{M})}|p(\lambda)|\right), \end{split}$$

where $\kappa(\mathcal{S}) := \|\mathcal{S}\|_2 \|\mathcal{S}^{-1}\|_2$ represents the spectral condition number of the eigenvector matrix \mathcal{S} and $\sigma(\mathcal{P}_{\alpha}^{-1}\mathcal{M})$ is explicitly given in Theorem 3.7. In particular, this bound implies that a well clustered spectrum away from zero will probably yield a fast convergence rate if $\kappa(\mathcal{S})$ is not very large. According to the arguments in [28, Theorem 3], we verified that $\mathcal{P}_{\alpha}^{-1}\mathcal{M}$ is diagonalizable. Hence, the above convergence bound is indeed applicable. Nevertheless, it is still not easy to estimate $\kappa(\mathcal{S})$, especially when we just make the spatial matrix A fulfill Assumption 3.1. This phenomenon motivates us to exploit an alternative technique to find a different convergence rate bound, as explained in the next section.

In fact, for any matrix, $Z \in \mathbb{R}^{n \times n}$, we denote by $\mathcal{H}(Z) = \frac{Z+Z^{\top}}{2}$ the symmetric part of Z. Considering the convergence bound of preconditioned GMRES [7, 31] for non-symmetric positive definite systems, the relative residual norm estimate can be given as follow:

$$\frac{\|\boldsymbol{r}_k\|_2}{\|\boldsymbol{r}_0\|_2} \leq \left[1 - \frac{\lambda_{\min}^2\left(\mathcal{H}\left(\mathcal{P}_{\alpha}^{-1}\mathcal{M}\right)\right)}{\left\|\mathcal{P}_{\alpha}^{-1}\mathcal{M}\right\|_2^2}\right]^{k/2}$$

where we emphasize that achieving such a (possibly nonsharp) convergence rate necessitates the positive definiteness of the symmetric part of the preconditioned matrix, i.e., $\mathcal{H}(\mathcal{P}_{\alpha}^{-1}\mathcal{M})$, which will be shown next.

At this stage, since

$$\mathcal{P}_{\alpha} = \mathcal{M} - \alpha \begin{bmatrix} \mathbf{0} & \cdots & \mathbf{0} & Q_2 \\ \mathbf{0} & \cdots & \mathbf{0} \\ & \ddots & \vdots \\ & & & \mathbf{0} \end{bmatrix} := \mathcal{M} - \alpha \mathcal{R},$$

we have

(4.3)
$$\|\mathcal{P}_{\alpha}^{-1}\mathcal{M}\|_{2} = \|\mathcal{I} + \alpha \mathcal{P}_{\alpha}^{-1}\mathcal{R}\|_{2} \le 1 + \alpha \|\mathcal{P}_{\alpha}^{-1}\mathcal{M}\|_{2} \|\mathcal{M}^{-1}\mathcal{R}\|_{2},$$

where estimating the norm $\|\alpha \mathcal{P}_{\alpha}^{-1} \mathcal{R}\|_2$ directly poses challenges. Assuming that $\alpha \in (0, 1)$ is small enough such that $\alpha \|\mathcal{M}^{-1} \mathcal{R}\|_2 < 1/2$, it holds that

$$\|\mathcal{P}_{\alpha}^{-1}\mathcal{M}\|_{2} \leq \frac{1}{1-\alpha\|\mathcal{M}^{-1}\mathcal{R}\|_{2}}$$

Similar to the conclusion of Ref. [26], we have $\|\mathcal{P}_{\alpha}^{-1}\mathcal{M} - \mathcal{I}\|_{2} \leq \frac{\alpha \|\mathcal{M}^{-1}\mathcal{R}\|_{2}}{1-\alpha \|\mathcal{M}^{-1}\mathcal{R}\|_{2}}$ and

(4.4)
$$\lambda_{\min}\left(\mathcal{H}\left(\mathcal{P}_{\alpha}^{-1}\mathcal{M}\right)\right) = 1 + \lambda_{\min}\left(\mathcal{H}\left(\mathcal{P}_{\alpha}^{-1}\mathcal{M}-\mathcal{I}\right)\right) \ge 1 - \left\|\mathcal{H}\left(\mathcal{P}_{\alpha}^{-1}\mathcal{M}-\mathcal{I}\right)\right\|_{2} \\ \ge 1 - \left\|\mathcal{P}_{\alpha}^{-1}\mathcal{M}-\mathcal{I}\right\|_{2} > 1 - \frac{\alpha\|\mathcal{M}^{-1}\mathcal{R}\|_{2}}{1 - \alpha\|\mathcal{M}^{-1}\mathcal{R}\|_{2}} > 0,$$

due to the well-known inequalities $|\lambda_{\min}(\mathcal{H}(A))| \leq ||\mathcal{H}(A)||_2 \leq ||A||_2$ [21, p. 160] and it also implies that the symmetric part of the preconditioned matrix $\mathcal{P}_{\alpha}^{-1}\mathcal{M}$ is positive definite. Hence, combining (4.3) and

(4.4), the following estimate for the GMRES convergence rate

(4.5)
$$\frac{\|\boldsymbol{r}_k\|_2}{\|\boldsymbol{r}_0\|_2} < \left[1 - \left(\frac{1 - \frac{\alpha \|\mathcal{M}^{-1}\mathcal{R}\|_2}{1 - \alpha \|\mathcal{M}^{-1}\mathcal{R}\|_2}}{\frac{1}{1 - \alpha \|\mathcal{M}^{-1}\mathcal{R}\|_2}}\right)^2\right]^{k/2} = \left[1 - (1 - 2\alpha \|\mathcal{M}^{-1}\mathcal{R}\|_2)^2\right]^{k/2}$$

can be derived immediately.

It is not hard to see that if we want to give a restrictive convergence rate of the preconditioned GMRES, the upper bound of the norm $\|\mathcal{M}^{-1}\mathcal{R}\|_2$ should be well estimated. Unfortunately, if the spatial matrix A just fulfills Assumption 3.1, we cannot estimate the upper bound of the norm $\|\mathcal{M}^{-1}\mathcal{R}\|_2$ well, because the useful property of matrix spectral norm being invariant under orthogonal transformations is not unavailable. Next, we improve Assumption 3.1 for the case that the spatial matrix A is symmetric negative semi-definite. Then, we can obtain $\tilde{A} = V_a D_a V_a^{\top}$ and $Q_1^{-1}Q_2 = V_a D_J V_a^{\top}$, according to Eq. (4.2) and $D_J = \text{diag}(\lambda_1, \lambda_2, \ldots, \lambda_s)$ is the diagonal matrix of eigenvalues of $Q_1^{-1}Q_2$, with $\lambda_j \in [0, 1]$, $j = 1, 2, \ldots, s$, cf. Lemma 3.5. Moreover, with

(4.6)
$$\mathcal{M}^{-1}\mathcal{R} = \begin{bmatrix} K_0^{-1} & & & \\ K_1^{-1} & K_0^{-1} & & \\ \vdots & \ddots & \ddots & \\ K_{M-1}^{-1} & \cdots & K_1^{-1} & K_0^{-1} \end{bmatrix} \begin{bmatrix} \mathbf{0} & \cdots & \mathbf{0} & Q_2 \\ & \mathbf{0} & \cdots & \mathbf{0} \\ & & \ddots & \vdots \\ & & & K_1^{-1}Q_2 \\ & & \ddots & \vdots \\ & & & & K_{M-1}^{-1}Q_2 \end{bmatrix} = \begin{bmatrix} \mathbf{0} & \cdots & \mathbf{0} & G_1 \\ & \mathbf{0} & \cdots & G_2 \\ & & \ddots & \vdots \\ & & & & & G_M \end{bmatrix},$$

where $K_{\ell}^{-1} = (Q_1^{-1}Q_2)^{\ell}Q_1^{-1}$ $(\ell = 0, 1, \dots, M - 1)$ and

(4.7)
$$G_n = (Q_1^{-1}Q_2)^{n-1}Q_1^{-1}Q_2 = V_a D_J^n V_a^\top, \quad n = 1, 2, \dots, M,$$

then we have $G_n^{\top}G_n = V_a(D_J^2)^n V_a^{\top}$. At this stage, one can obtain

(4.8)
$$\|\mathcal{M}^{-1}\mathcal{R}\|_{2}^{2} = \lambda_{\max}((\mathcal{M}^{-1}\mathcal{R})^{\top}\mathcal{M}^{-1}\mathcal{R}) = \lambda_{\max}\left(\sum_{n=1}^{M} G_{n}^{\top}G_{n}\right)$$
$$\leq \left\|\sum_{n=1}^{M} (D_{J}^{2})^{n}\right\|_{2} = \max_{1 \leq j \leq N} \sum_{n=1}^{M} (\lambda_{j}^{2})^{n} \leq M,$$

due to $\lambda_i^2 \in [0, 1]$ and thus $\|\mathcal{M}^{-1}\mathcal{R}\|_2 \leq \sqrt{M}$.

In view of (4.5) and the above analysis, we can deduce the subsequent linear convergence rate of the preconditioned GMRES.

THEOREM 4.2. For any given $\delta \in (0, 1/2)$, if we select $\alpha = \delta \sqrt{\tau/T}$ such that $\alpha \|\mathcal{M}^{-1}\mathcal{R}\|_2 \leq \delta < 1/2$, then the left-preconditioned GMRES for solving $\mathcal{M}\mathbf{u} = \mathbf{b}$ with the preconditioner \mathcal{P}_{α} enjoys the following mesh-independent linear convergence rate:

(4.9)
$$\frac{\|\boldsymbol{r}_k\|_2}{\|\boldsymbol{r}_0\|_2} \le \left[1 - \left(1 - 2\alpha\sqrt{\frac{T}{\tau}}\right)^2\right]^{k/2} \le \left[1 - (1 - 2\delta)^2\right]^{k/2} = \left[2\sqrt{\delta}(1 - \delta)\right]^k,$$

where \mathbf{r}_k is the residual vector at the k-th iteration of the preconditioned GMRES.

This theorem suggests that the preconditioned GMRES iterations become independent of the mesh size when α is adequately small. It further demonstrates that a smaller α results in a faster convergence rate, as confirmed in numerical experiments. Among our tested examples, we observe that a fixed $\alpha = 0.01$

or 0.001 already provides very fast mesh-independent convergence rate. However, the theoretical condition $\alpha = \mathcal{O}(\sqrt{\tau})$ is rather restrictive, because it implies that α goes to 0 rapidly as we refine the time stepsize τ . In practical implementations, excessively small α may seriously pollute the numerical solution since unacceptable round-off errors will occur in Step-(a) and Step-(c) of Eq. (3.3); see [9, 37] for further information regarding the round-off errors arising in the diagonalization technique.

5. Numerical experiments. In this section, several numerical examples are implemented to display the convergence rates and computational efficiency of the developed block α -circulant preconditioner. All simulations are done by using MATLAB2023a on an ASUSTeK workstation with 13th Gen Intel(R) Core(TM) i9-13900K 3.00 GHz and 128GB RAM. The CPU elapsed time (in seconds) is recorded by using the timing functions tic/toc, based on the (partially) PinT implementation of the preconditioned iterative algorithms¹. Here our primary focus lies in examining the convergence properties of the algorithms, and forthcoming work will present numerical results from PinT computation in a practical parallel environment. We conduct our analysis using left-preconditioned GMRES, but in practical implementations, the right-preconditioned GMRES is favored because its termination criterion is based on the reliable true residuals, rather than the potentially significantly different preconditioned residuals [29]. The right-preconditioned GMRES solver (with restarting number being 40) in the *i*FEM package [2] is exploited with a zero initial guess and a stopping tolerance tol = 10⁻⁹ based on the reduction in relative residual norms. By "Its" we denote the number of iterations of the right-preconditioned GMRES solver; by "DoFs" the degrees of freedom, i.e., the number of unknowns.

5.1. Heston model. In the example, we consider the following Heston stochastic volatility option pricing PDE [22]

(5.1)
$$\frac{\partial u}{\partial t} = \frac{1}{2}vs^2\frac{\partial^2 u}{\partial s^2} + \sigma sv\rho\frac{\partial^2 u}{\partial s\partial v} + \frac{1}{2}\sigma^2 v\frac{\partial^2 u}{\partial v^2} + rs\frac{\partial u}{\partial s} + \kappa(\eta - v)\frac{\partial u}{\partial v} - ru,$$

where the function u(s, v, t) represents the price of a European option when at time $T - t^*$ the corresponding asset price is equal to s and its variance is v. We consider the equation on the unbounded domain

$$0 \le t \le T, \ s > 0, \ v > 0,$$

where the option final contract time T is fixed. The parameters $\kappa > 0$, $\sigma > 0$ and $\rho \in [-1, 1]$ are given. Moreover, Eq. (5.1) is usually considered under the condition $2\kappa\eta > \sigma^2$ that is known as the Feller condition. We take Eq. (5.1) together with the initial condition

(5.2)
$$u(s, v, t) = \max\{0, s - K\},\$$

where K > 0 represents the strike price of the option. The (artificial) boundaries should be set far away from the region of interest to minimize the effect on the solution. The spatial domain is restricted to the bounded set: $[0, S_{\text{max}}] \times [0, V_{\text{max}}]$ with fixed values S_{max} and V_{max} chosen large enough (set $S_{\text{max}} = 800$ and $V_{\text{max}} = 4$ in this example). In the case of a European call option, the boundary conditions are imposed as follows

(5.3)
$$u(0, v, t) = 0$$

$$(5.4) u(s, V_{\max}, t) = s$$

(5.5)
$$\frac{\partial u}{\partial s}(S_{\max}, v, t) = 1$$

The fourth boundary condition is not really a boundary condition because it involves a time derivative:

(5.6)
$$\frac{\partial u}{\partial s}(s,0,t) = \kappa \eta \frac{\partial u}{\partial v}(s,0,t) + rs \frac{\partial u}{\partial s}(s,0,t) - ru(s,0,\tau),$$

¹In our MATLAB codes, we used the **parfor** to accelerate the computational speed.

which is the artificial boundary condition at v = 0 for the Heston PDE (5.1). We perform the spatial discretization for Eqs. (5.1)–(5.5) and then it will be solved by our aforementioned PinT methodology. Here, we define the measure of relative error as

 $\operatorname{Err} = \left| \operatorname{Price}(S_0, V_0) - \operatorname{Price}_{\operatorname{ref}} \right| / \left| \operatorname{Price}_{\operatorname{ref}} \right|,$

where the reference price "Price_{ref}" is obtained by using the COS method [12] and if the point (S_0, V_0) is not a spatial mesh point, bilinear interpolation will be used to obtain the approximation of the Price at (S_0, V_0) ; see [4, p. 68] for details. Here, we consider the following sets of Heston model parameters [22]:

- Set I: $T = 1, K = 100, \kappa = 1.5, \eta = 0.04, \sigma = 0.3, r = 0.025, \rho = -0.9, S_0 = 100, V_0 = 0.5;$
- Set II: $T = 3, K = 100, \kappa = 0.6067, \eta = 0.0707, \sigma = 0.2928, r = 0.03, \rho = -0.7571, S_0 = 100, V_0 = 0.2.$

TABLE 1 Convergence of different right-preconditioned GMRES(40) methods with $N_t = N_s = 2N_v$ and $\alpha = 10^{-3}$ (Set I).

Mesh	N_t	DoFs	P_1			P_{lpha}		
			Its	CPU	Err	Its	CPU	Err
uniform	48	$55,\!296$	29	0.82	1.348e-2	4	0.09	1.348e-2
	96	442,368	25	4.76	3.213e-3	4	0.51	3.213e-3
	192	$3,\!538,\!944$	26	51.03	8.003e-4	4	5.52	8.003e-4
	384	$28,\!311,\!552$	27	664.46	1.997e-4	4	76.34	1.997e-4
nonuniform	36	23,328	21	0.29	7.679e-3	4	0.05	7.679e-3
	72	$186,\!624$	24	1.46	1.476e-3	4	0.18	1.476e-3
	144	$1,\!492,\!992$	27	14.05	4.339e-4	4	1.47	4.339e-4
	288	$11,\!943,\!936$	29	154.03	1.259e-4	4	16.01	1.259e-4

TABLE 2 Convergence of different right-preconditioned GMRES(40) methods with $N_t = N_s = 2N_v$ and $\alpha = 10^{-3}$ (Set II)

Mesh	N_t	DoFs	P_1			P_{lpha}		
			Its	CPU	Err	Its	CPU	Err
uniform	50	62,500	27	0.83	1.281e-2	4	0.10	1.281e-2
	100	500,000	27	6.94	1.725e-3	4	0.64	1.725e-3
	200	4,000,000	28	62.60	6.168e-4	4	6.28	6.169e-4
	400	$32,\!000,\!000$	28	962.07	1.550e-4	4	96.51	1.551e-4
nonuniform	40	32,000	21	0.33	2.640e-3	4	0.06	2.640e-3
	80	256,000	24	2.26	7.929e-4	4	0.29	7.929e-4
	160	2,048,000	24	16.98	1.862e-4	4	2.16	1.862e-4
	320	$16,\!384,\!000$	25	176.43	4.665e-5	4	22.05	4.665e-5

The results of right preconditioned GMRES(40) with preconditioners P_1 and P_{α} for the AaO system from the CN temporal scheme and (non-)uniform spatial discretization are listed in Table 1–Table 2 corresponding to Sets I–II, respectively. As seen from these two tables, (i) right preconditioned GMRES(40) works for the CN-type AaO system in terms of the number of iterations and numerical errors; (ii) the GMRES(40) method with preconditioner P_{α} is more efficient than that with preconditioner P_1 , in terms of computational time and iteration numbers and the former delivers an excellent mesh-independent convergence rate; (iii) a nonuniform spatial discretization using fewer grid nodes can indeed improve the accuracy of numerical solution of Heston PDEs, but it does not affect the performance of our preconditioners P_1 and P_{α} , which are analyzed via the information of the selected temporal discretization.

To support the results of Theorem 3.7, we plot the spectrum of the preconditioned matrices corresponding to the different models (i.e., Sets I–II) in Figure 1–Figure 2. These figures show that the spectrum of $P_{\alpha}^{-1}\mathcal{M}$ is more clustered around 1 than the spectrum of $P_1^{-1}\mathcal{M}$. Meanwhile, the modules of



FIG. 1. Eigenvalues of the original and preconditioned matrices (i.e., \mathcal{M} , $P_1^{-1}\mathcal{M}$ and $P_{\alpha}^{-1}\mathcal{M}$) for the model (Set I) under the uniform spatial discretization with $N_t = N_s = 2N_v = 36$ and $\alpha = 10^{-3}$.

all the eigenvalues of $P_{\alpha}^{-1}\mathcal{M}$ lie in the interval $[1/(1+\alpha), 1/(1-\alpha)]$, which is consistent with the result of Theorem 3.7.

5.2. Stochastic Alpha Beta and Rho (SABR) model. In this example, we consider a PDE formulation corresponding to the SABR model [4, 34], which is a stochastic differential equation system used for interest rates and FX modelling:

(5.7)
$$\frac{\partial u}{\partial t} = \frac{1}{2} v^2 s^{2\beta} D^{2(1-\beta)} \frac{\partial^2 u}{\partial s^2} + \rho \sigma s^\beta D^{(1-\beta)} v^2 \frac{\partial^2 u}{\partial s \partial v} + \frac{1}{2} \sigma^2 v^2 \frac{\partial^2 u}{\partial v^2} + rs \frac{\partial u}{\partial s} - ru, \quad D(t) = e^{-rt},$$

for s > 0, v > 0 and $0 \le t < T$. Here the spatial domain is restricted to the bounded set as in Subsection 5.1. Moreover, the initial condition and the first three boundary conditions are very similar to Eqs. (5.2)–(5.5), but the last boundary condition will be slightly different from Eq. (5.6), namely

$$\frac{\partial u(s,0,t)}{\partial t} = rs\frac{\partial u(s,0,t)}{\partial s} - ru(s,0,t)$$

We perform the discretization on a uniform spatial grid for the equation which will then be solved by our introduced methodology. We employ the measure following of relative error,

$$\operatorname{Err} = |\operatorname{Price}(S_0, V_0) - \operatorname{Price}_{\operatorname{ref}}| / |\operatorname{Price}_{\operatorname{ref}}|,$$

where the reference price "Price_{ref}" is taken from Problem 1 in [34, Section 4.1] and can be generated via Script_Compare_European.m in the package² and if the point (S_0, V_0) is not a spatial mesh point, bilinear interpolation will be used to obtain the approximation of Price (S_0, V_0) , see [4, p. 68] for details. We consider the following sets of SABR model parameters [34]:

²Source code is available online at https://github.com/jkirkby3/PROJ_Option_Pricing_Matlab.



FIG. 2. Eigenvalues of the original and preconditioned matrices (i.e., \mathcal{M} , $P_1^{-1}\mathcal{M}$ and $P_{\alpha}^{-1}\mathcal{M}$) for the model (Set II) under the nonuniform spatial discretization with $N_t = N_s = 2N_v = 36$ and $\alpha = 10^{-3}$.

• Set III: $S_{\text{max}} = 24$, $V_{\text{max}} = 3$, T = 2, $\beta = 0.5$, $\sigma = 0.4$, r = 0.0, $\rho = 0.0$, $K = S_0 = 0.5$, $V_0 = 0.5$; • Set IV: $S_{\text{max}} = 4$, $V_{\text{max}} = 2$, T = 10, $\beta = 0.5$, $\sigma = 0.8$, r = 0.0, $\rho = -0.6$, $K = S_0 = 0.07$, $V_0 = 0.4$.

TABLE 3 Convergence of different right-preconditioned GMRES(40) methods with $N_t = N_s = 2N_v$ and $\alpha = 10^{-3}$.

Set	N_t	DoFs	P_1			P_{lpha}			
_			Its	CPU	Err	Its	CPU	Err	
III	48	$55,\!296$	‡	_		4	0.08	1.365e-1	
	96	$442,\!368$	‡			4	0.38	3.102e-2	
	192	$3,\!538,\!944$	‡			4	3.11	7.003e-3	
	384	$28,\!311,\!552$	‡			4	33.45	1.550e-3	
IV	48	$55,\!296$	‡			4	0.09	4.760e-2	
	96	$442,\!368$	‡			4	0.40	2.191e-2	
	192	$3,\!538,\!944$	‡			4	3.68	7.892e-3	
	384	$28,\!311,\!552$	‡			4	37.61	2.839e-3	

We apply the GMRES(40) method with preconditioners P_1 and P_{α} for the AaO linear system, corresponding to the SABR model under the above parameters in Sets III–IV. The results are listed in Table 3. First of all, the symbol "‡" means that the MATLAB codes will report an error message indicating that the preconditioner P_1 is singular, i.e., the GMRES(40) method with P_1 diverges and the information of CPU and Err is unavailable, refer to Remark 3.3 for details. However, the GMRES(40) method with P_{α} converges well in terms of iteration numbers and computational time. Moreover, the number of iterations of GMRES(40) with P_{α} is robust with respect to changes of τ and h. To observe how the preconditioner P_{α} affects the spectrum of \mathcal{M} , whose eigenvalues are plotted in Figure 3. Figure 3 shows that the spectrum of $P_{\alpha}^{-1}\mathcal{M}$ is clustered around 1 and, as stated in Theorem 3.7, the modules of the eigenvalues of $P_{\alpha}^{-1}\mathcal{M}$ lie within the specific interval. As seen from Figure 4, the spatial discretization matrix Ahas some zero eigenvalues, which make the preconditioner P_1 singular, see Table 3 and Remark 3.3. In addition, due to the zero eigenvalues of A, the upper bound for estimating the spectrum of $P_{\alpha}^{-1}\mathcal{M}$ in Theorem 3.7 can be reached and thus it is a sharp bound in this case; refer to Remark 3.9 for details. In conclusion, the proposed preconditioner P_{α} is essential and efficient to accelerate the convergence of GMRES(40) for this example.



FIG. 3. Eigenvalues of the original and preconditioned matrices (i.e., \mathcal{M} and $P_{\alpha}^{-1}\mathcal{M}$) for the model (Sets III and IV) with $N_t = N_s = 2N_v = 36$ and $\alpha = 10^{-3}$



FIG. 4. Eigenvalues of the space discrete matrix \tilde{A} for the model (Sets III and IV) with $N_t = N_s = 2N_v = 36$.

On the other hand, we would like to consider the model (5.7) with $r \neq 0$. This relates to a more general case, i.e., the spatial operator \mathcal{L} in Eq. (1.1) becomes time-dependent. More precisely, $\mathcal{L}(t) = d(t)\mathcal{L}$, where d(t) > 0 is a known function, cf. (5.7). Following the same procedure as in Section 2, the CN scheme-based

AaO system of the case is obtained by changing \mathcal{M} and $\boldsymbol{\xi}$ in Eq. (2.3) to

(5.8)
$$\mathcal{M} = B_1 \otimes I_s - \left(\tilde{D}B_2\right) \otimes \tilde{A} \text{ and } \boldsymbol{\xi} = \left[\left[\left(I_s + \frac{d(t_{\frac{1}{2}})}{2} \tilde{A} \right) \boldsymbol{u}^0 \right]^\top, \boldsymbol{0}^\top, \dots, \boldsymbol{0}^\top \right]^\top,$$

respectively. Here $\tilde{D} = \text{diag}\left(d(t_{\frac{1}{2}}), d(t_{\frac{3}{2}}), \ldots, d(t_{M-\frac{1}{2}})\right)$. Since $d(t) = e^{-rt}$ is sufficiently smooth and r is often small, we can employ $\bar{d} = \frac{1}{M} \sum_{k=0}^{M-1} d(t_{k+\frac{1}{2}})$, such that $\tilde{D}B_2 \approx \bar{d}B_2$ and then only a small modification is needed for our PinT preconditioner (3.1), where $C_2^{(\alpha)}$ will be changed with $\bar{d}C_2^{(\alpha)}$. Consequently, the PinT preconditioner can be still implemented in the pattern (3.3) and the results of the right preconditioned GMRES(40) method with preconditioners P_1 and P_{α} for the AaO system (5.8) are listed in Table 4, corresponding to Set V.

TABLE 4 Convergence of different right-preconditioned GMRES(40) methods with $N_t = N_s = 2N_v$ and $\alpha = 10^{-3}$

Set	N_t	DoFs		P_1		P_{lpha}		
			Its	CPU	Err	Its	CPU	Err
V	48	$55,\!296$	100	2.25	2.282e-1	6	0.12	2.282e-1
	96	442,368	111	19.86	8.352e-2	6	0.85	8.352e-2
	192	$3,\!538,\!944$	131	179.11	9.245e-3	6	6.24	9.244e-3
	384	$28,\!311,\!552$	†			6	62.49	7.257e-4

• Set V: $S_{\text{max}} = 23, V_{\text{max}} = 3, T = 2, \beta = 0.45, \sigma = 0.4, r = 0.03, \rho = -0.65, K = S_0 = 0.5, V_0 = 0.2.$

Although our theoretical analysis in the previous sections is unavailable with this revised PinT preconditioner, due to time-varying coefficients, Table 4 shows that i) both right preconditioned GMRES(40) methods work for the AaO system (5.8) in terms of the number of iterations and numerical errors; ii) the GMRES(40) method with preconditioner P_{α} is more efficient than that with preconditioner P_1 , in terms of computational time and iteration numbers, and the former delivers a mesh-independent convergence rate. Besides, the eigenvalue plots of the original and two preconditioned matrices in this example are shown in Figure 5. Although there are no theoretical results for the eigenvalues of preconditioned matrices in this example, Figure 5 shows that the spectrum of $P_{\alpha}^{-1}\mathcal{M}$ is more clustered than the spectrum of $P_1^{-1}\mathcal{M}$, which also implies that the proposed preconditioned system with P_{α} converges faster than with P_1 .



FIG. 5. Eigenvalues of the original and preconditioned matrices (i.e., \mathcal{M} , $P_1^{-1}\mathcal{M}$ and $P_{\alpha}^{-1}\mathcal{M}$) for the model (Set V) with $N_t = N_s = 2N_v = 36$ and $\alpha = 10^{-3}$.

6. Concluding remarks. Compared to the classical time-stepping methods, a parallel numerical treatment of evolutionary PDEs has recently become popular due to the availability of massively parallel computing clusters. Built upon an all-at-once implicit CN method for discretizing the evolutionary PDEs, a parallel gBC preconditioner is proposed and the spectra of the diagonalizable preconditioned systems are analyzed rigorously. The eigenvalues of the preconditioned matrix are located within a uniformly bounded annulus whose bandwidth is explicit and only depends on the parameter α . Our eigenvalue analysis explains why the BC preconditioner P_1 may perform poorly in some cases. Utilizing the obtained eigenvalue bounds, we establish a mesh-independent convergence rate for the preconditioned GMRES, subject to specific conditions regarding the choice of α . Our theoretical findings and the efficacy of the proposed gBC preconditioner were validated by numerical results obtained from solving European option pricing PDEs.

The promising enhancement of a gBC preconditioner over the BC preconditioner deserves more exploration, especially for those celebrated iterative solvers (e.g., when solving nonsymmetric block Toeplitz systems) based on BC preconditioners. Such preliminary numerical results also reveal the potential of the gBC preconditioner when applied to evolutionary PDEs with time-varying coefficients, where the rigorous convergence analysis is worth further investigation in the future.

Acknowledgments. We would like to thank Dr. Meng Li (Zhengzhou University) who provided the computational resources to help us finish the numerical experiments in Section 5.

Conflict of interest. The authors declare that they have no conflict of interest.

Data availability. The datasets generated and analyzed during the current study are not publicly available but are available from the corresponding author on reasonable request.

REFERENCES

- [1] U. M. ASCHER, Numerical Methodsfor Evolutionary Differential Equations, SIAM, Philadelphia, PA, 2008.
- [2] L. CHEN, *iFEM: an integrated finite element methods package in MATLAB*, technical report, Department of Mathematics, University of California at Irvine, Irvine, CA, 2009. https://github.com/lyc102/ifem.
- [3] J. CRANK AND P. NICOLSON, A practical method for numerical evaluation of solutions of partial differential equations of the heat conduction type, Proc. Camb. Phil. Soc., 43 (1947), pp. 50–67.
- [4] C. S. L. DE GRAAF, Finite Difference Methods in Derivatives Pricing under Stochastic Volatility Models, master thesis, Mathematisch Instituut, Universiteit Leiden, Leiden, Netherlands, 2012.
- [5] S. DOLGOV, Tensor product methods in numerical simulation of high-dimensional dynamical problems, PhD dissertation, Universität Leipzig, Leipzig, Gemeny, 2014. 154 pages.
- [6] D. J. DUFFY, Numerical Methods in Computational Finance: A Partial Differential Equation (PDE/FDM) Approach, John Wiley & Sons, Chichester, UK, 2022.
- [7] S. C. EISENSTAT, H. C. ELMAN, AND M. H. SCHULTZ, Variational iterative methods for nonsymmetric systems of linear equations, SIAM J. Numer. Anal., 20 (1983), pp. 345–357, https://doi.org/10.1137/0720023.
- [8] M. EMBREE, How descriptive are GMRES convergence bounds?, arXiv preprint, arXiv:2209.01231 (2022), p. 35 pages. https://arxiv.org/abs/2209.01231.
- M. EMBREE, Extending Elman's bound for GMRES, arXiv preprint, (2023), p. 10 pages. https://arxiv.org/abs/2312. 15022.
- [10] L. C. EVANS, Partial Differential Equations, vol. 19 of Graduate Series in Mathematics, American Mathematical Society, Providence, RI, 2nd ed., 2010.
- [11] R. D. FALGOUT, S. FRIEDHOFF, T. V. KOLEV, S. P. MACLACHLAN, AND J. B. SCHRODER, Parallel time integration with multigrid, SIAM J. Sci. Comput., 36 (2014), pp. C635–C661, https://doi.org/10.1137/130944230.
- [12] F. FANG AND C. W. OOSTERLEE, A novel pricing method for European options based on Fourier-cosine series expansions, SIAM J. Sci. Comput., 31 (2009), pp. 826–848, https://doi.org/10.1137/080718061.
- [13] D. GAN, G.-F. ZHANG, AND Z.-Z. LIANG, Parallel preconditioning of all-at-once systems for fractional diffusion equations with Riesz fractional derivatives, SSRN Preprint, (2022), p. 23 pages, https://doi.org/10.2139/ssrn. 4234422.
- [14] M. J. GANDER, J. LIU, S.-L. WU, X. YUE, AND T. ZHOU, ParaDiag: parallel-in-time algorithms based on the diagonalization technique, arXiv preprint, arXiv:2005.09158v4 (2021), p. 22 pages. https://arxiv.org/abs/2005. 09158v4.
- [15] A. GODDARD AND A. WATHEN, A note on parallel preconditioning for all-at-once evolutionary PDEs, Electron. Trans. Numer. Anal., 51 (2019), pp. 135–150, https://doi.org/10.1553/etna_vol51s135.
- [16] G. H. GOLUB AND C. F. VAN LOAN, Matrix Computations, Johns Hopkins University Press, Baltimore, MD, 4 ed., 2013.
- [17] X.-M. GU, Y.-L. ZHAO, X.-L. ZHAO, B. CARPENTIERI, AND Y.-Y. HUANG, A note on parallel preconditioning for the all-at-once solution of Riesz fractional diffusion equations, Numer. Math. Theor. Meth. Appl., 14 (2021), pp. 893–919.

- [18] E. HAIRER AND G. WANNER, Solving Ordinary Differential Equations II: Stiff and Differential Algebraic Problems, Springer-Verlag, Berlin, Heidelberg, 1996.
- [19] S. HON, Optimal block circulant preconditioners for block Toeplitz systems with application to evolutionary PDEs, J. Comput. Appl. Math., 407 (2022), 113965, https://doi.org/10.1016/j.cam.2021.113965.
- [20] S. HON, P. Y. FUNG, J. DONG, AND S. SERRA-CAPIZZANO, A sine transform based preconditioned MINRES method for all-at-once systems from constant and variable-coefficient evolutionary PDEs, Numer. Algor., in press (2023), https://doi.org/10.1007/s11075-023-01627-5.
- [21] R. HORN AND C. JOHNSON, Topics in Matrix Analysis, Cambridge University Press, New York, NY, 1991.
- [22] K. J. IN 'T HOUT AND S. FOULON, ADI finite difference schemes for option pricing in the Heston model with correlation, Int. J. Numer. Anal. Mod., 7 (2010), pp. 303–320.
- [23] X.-L. LIN AND M. NG, An all-at-once preconditioner for evolutionary partial differential equations, SIAM J. Sci. Comput., 43 (2021), pp. A2766–A2784, https://doi.org/10.1137/20M1316354.
- [24] X.-L. LIN AND S.-L. WU, A parallel-in-time preconditioner for Crank-Nicolson discretization of a parabolic optimal control problem, arXiv preprint, arXiv:2109.12524v5 (2023), p. 30 pages. https://arxiv.org/abs/2109.12524.
- [25] J.-L. LIONS, Y. MADAY, AND G. TURINICI, Résolution d'edp par un schéma en temps «pararéel», C. R. Acad. Sci. Paris, 332 (2001), pp. 661–668, https://doi.org/10.1016/S0764-4442(00)01793-6.
- [26] J. LIU AND S.-L. WU, A fast block α-circulant preconditoner for all-at-once systems from wave equations, SIAM J. Matrix Anal. Appl., 41 (2020), pp. 1912–1943.
- [27] Y. MADAY AND E. M. RØNQUIST, Parallelization in time through tensor-product space-time solvers, C. R. Math., 346 (2008), pp. 113–118, https://doi.org/10.1016/j.crma.2007.09.012.
- [28] E. MCDONALD, J. PESTANA, AND A. WATHEN, Preconditioning and iterative solution of all-at-once systems for evolutionary partial differential equations, SIAM J. Sci. Comput., 40 (2018), pp. A1012–A1033.
- [29] Y. SAAD, Iterative Methods for Sparse Linear Systems, SIAM, Philadelphia, PA, 2nd ed., 2003.
- [30] D. SHEEN, I. H. SLOAN, AND V. THOMÉE, A parallel method for time discretization of parabolic equations based on Laplace transformation and quadrature, IMA J. Numer. Anal., 23 (2003), pp. 269–299, https://doi.org/10.1093/ imanum/23.2.269.
- [31] N. SPILLANE, Hermitian preconditioning for a class of non-Hermitian linear systems, SIAM J. Sci. Comput., to appear (2024), p. 21 pages, https://doi.org/10.48550/arXiv.2304.03546.
- [32] E. TADMOR, A review of numerical methods for nonlinear partial differential equations, Bull. Amer. Math. Soc., 49 (2012), pp. 507–554.
- [33] D. TITLEY-PELOQUIN, J. PESTANA, AND A. J. WATHEN, GMRES convergence bounds that depend on the right-handside vector, IMA J. Numer. Anal., 34 (2014), pp. 462–479.
- [34] L. VON SYDOW, S. MILOVANOVIĆ, E. LARSSON, K. IN'T HOUT, M. WIKTORSSON, C. W. OOSTERLEE, V. SHCHERBAKOV, M. WYNS, A. LEITAO, S. JAIN, H. T., AND J. WALDÉN, BENCHOP – SLV: the BENCHmarking project in option pricing – stochastic and local volatility problems, Int. J. Comput. Math., 96 (2019), pp. 1910–1923, https://doi.org/10.1080/00207160.2018.1544368.
- [35] A. WATHEN, Some observations on preconditioning for non-self-adjoint and time-dependent problems, Comput. Math. Appl., 116 (2022), pp. 176–180, https://doi.org/10.1016/j.camwa.2021.05.037.
- [36] S.-L. WU AND T. ZHOU, Parallel implementation for the two-stage SDIRK methods via diagonalization, J. Comput. Phys., 428 (2021), p. 110076, https://doi.org/10.1016/j.jcp.2020.110076.
- [37] S.-L. WU, T. ZHOU, AND Z. ZHOU, A uniform spectral analysis for a preconditioned all-at-once system from firstorder and second-order evolutionary problems, SIAM J. Matrix Anal. Appl., 43 (2022), pp. 1331–1353, https:// doi.org/10.1137/21M145358X.
- [38] G. ZASLAVSKY, Chaos, fractional kinetics, and anomalous transport, Phys. Rep., 371 (2002), pp. 461–580, https:// doi.org/10.1016/S0370-1573(02)00331-9.