

SOME HARDY AND RELlich TYPE INEQUALITIES FOR AFFINE CONNECTIONS

PENGYAN WANG¹, HUITING CHANG^{1,*}

¹School of Mathematics and Statistics, Xinyang Normal University, Xinyang, 464000, P.R. China

Abstract. In this article we study various forms of the Hardy inequality for affine connections on a complete noncompact Riemannian manifold, including the two-weight Hardy inequality, the improved Hardy inequality, the Rellich inequality, the Hardy-Poincaré inequality and the Heisenberg-Pauli-Weyl inequality. Our results improve and include many previously known results as special cases.

Keywords. affine connection; affine Laplacian; Hardy type inequality; complete manifold.

2020 Mathematics Subject Classification. 26D10, 58J05, 53B05.

1. INTRODUCTION

The classical Hardy inequality plays significant roles in many areas such as analysis, probability and partial differential equations:

$$\int_{\mathbb{R}^n} |\nabla u(x)|^2 dx \geq \left(\frac{n-2}{2}\right)^2 \int_{\mathbb{R}^n} \frac{|u(x)|^2}{|x|^2} dx,$$

where $u \in C_0^\infty(\mathbb{R}^n \setminus \{0\})$ and $n \geq 3$. A natural generalization of the Hardy inequality is the following Rellich inequality:

$$\int_{\mathbb{R}^n} |\Delta u(x)|^2 dx \geq \left(\frac{n(n-4)}{4}\right)^2 \int_{\mathbb{R}^n} \frac{|u(x)|^2}{|x|^4} dx,$$

where $u \in C_0^\infty(\mathbb{R}^n \setminus \{0\})$ and $n \geq 5$. The constant $\left(\frac{n-2}{2}\right)^2$ and $\left(\frac{n(n-4)}{4}\right)^2$ in the above inequalities are optimal and are never achieved by nontrivial functions.

In this article, we are concerned with various forms of the Hardy inequality for affine connections on a complete noncompact Riemannian manifold.

Let us provide the rigorous definitions we are going to employ for affine connections on Riemannian manifolds (see [18]).

Let (M, g) be an n -dimensional Riemannian manifold and ∇ be the Levi-Civita connection of g . Let $V = e^u$, where u is a smooth function defined on M . We call (M, g, V) a *Riemannian triple*. For two real constants λ, μ and two vector fields X, Y on M , we define a 2-parameter family of *affine connections* by

$$D_X^{\lambda, \mu} Y := \nabla_X Y + \lambda du(X)Y + \lambda du(Y)X + \mu g(X, Y)\nabla u. \quad (1.1)$$

As Li and Xia [18] have already noted, when $\lambda = \mu = 0$, $D^{\lambda, \mu}$ is a Levi-Civita connection for g ; when $\lambda = -\mu$, $D^{\lambda, \mu}$ is a Levi-Civita connection for the conformal metric $e^{2\lambda f}g$. Except for

*Corresponding author.

E-mail addresses: wangpy@xynu.edu.cn (P. Wang), changhuiting163@163.com (H. Chang).

$\lambda = \mu = 0$ and $\lambda = -\mu$, $D^{\lambda, \mu}$ may not be a Levi-Civita connection for any Riemannian metric. The definitions of the affine Ricci curvature, the affine gradient, the affine Laplacian and the affine divergence will be given in Sect. 2 below. For more details on affine connections, we refer to [8, 9] and the references therein.

Now, let's recall some relevant work about the Hardy type inequalities for affine connections (1.1). In the case $\lambda = \mu = 0$, the affine connections (1.1) reduces to the Levi-Civita connection. The affine gradient and the affine Laplacian are in consistence with the classical ones. Carron [1] first established the following Hardy inequality on Riemannian manifold:

$$\int_M \rho^\alpha |\nabla u|^2 dv_g \geq \frac{(C + \alpha - 1)^2}{4} \int_M \rho^{\alpha-2} u^2 dv_g,$$

where $\alpha \in \mathbb{R}$, $C + \alpha - 1 > 0$, $u \in C_0^\infty(M \setminus \rho^{-1}\{0\})$. The weight function ρ is nonnegative and it satisfies $|\nabla \rho| = 1$ and $\Delta \rho \geq \frac{C}{\rho}$ in the sense of distribution. Throughout this paper, dv_g, ∇ and Δ denote the volume element, the gradient and the Laplace operators on M . Under the same geometric assumptions on the weight function ρ , Kombe and Özaydin [13] established the weighted L^p -Hardy inequality:

$$\int_M \rho^\alpha |\nabla u|^p dv_g \geq \left(\frac{C + 1 + \alpha - p}{p} \right)^p \int_M \rho^{\alpha-p} u^p dv_g,$$

where $1 \leq p < \infty$, $C + 1 + \alpha - p > 0$ and $u \in C_0^\infty(M \setminus \rho^{-1}\{0\})$. Kombe and Özaydin [14] also proved a new weighted Hardy-Poincaré inequalities. They showed that if M is a complete noncompact Riemannian manifold of dimension $n > 1$ and ρ is a nonnegative function on M such that $|\nabla \rho| = 1$ and $\Delta \rho \geq \frac{C}{\rho}$ in the sense of distribution, where $C > 0$, the following inequality holds:

$$\int_M \rho^{\alpha+p} |\nabla \rho \cdot \nabla u|^p dv_g \geq \left(\frac{C + 1 + \alpha}{p} \right)^p \int_M \rho^\alpha |u|^p dv_g. \quad (1.2)$$

Kombe and Yener [15] provided a new form of the L^p -Hardy inequality involving two weight functions, that is,

$$\int_M a(x) |\nabla u|^p dv_g \geq \int_M b(x) |u|^p dv_g + c(p) \int_M a(x) |\omega|^p \left| \nabla \left(\frac{u}{\omega} \right) \right|^p dv_g$$

for suitable functions a, b and ω . Xia [23] proved the following Hardy type inequalities on a complete noncompact Riemannian manifold:

$$\int_M \rho^\alpha |\nabla u|^p \geq \left(\frac{p - C - 1 - \alpha}{p} \right)^p \int_M \rho^{\alpha-p} |u|^p - \left(\frac{p - C - 1}{p} \right)^{p-1} \int_M \rho^{\alpha-p+1} |u|^p.$$

for any $p, \alpha \in \mathbb{R}$ with $p > 1$ and $p - \alpha - C - 1 > 0$. For more results on Hardy type inequalities on Riemannian manifolds, we refer to [3, 4, 5, 10, 11, 12, 21, 20] and the references therein.

In the case $\lambda = -\mu$, the affine connections (1.1) reduces to a Levi-Civita connection for the conformal metric $e^{2\alpha f} g$. The affine gradient and the affine Laplacian are in consistence with the classical ones for the conformal metric $e^{2\alpha f} g$ (also known as the weighted Laplacian, for example, see [6, 7, 16, 22]). Du and Mao [2] proved some Hardy and Rellich type inequalities for the weighted Laplacian. Li et al. [17] established some Hardy type inequalities with remainder terms for the weighted Laplacian. While in [19], by means of various weighted Ricci curvatures, Meng et al. established several sharp Hardy type inequalities for the weighted Laplacian on closed weighted Riemannian manifolds.

As for the affine connections (1.1), to our knowledge, the only known Hardy type inequality is due to Li and Xia [18]. They showed that, under some additional conditions on the affine Ricci curvature, the L^2 -version of the Hardy type inequality holds on compact Riemannian manifolds (Theorem 5.1 in [18]).

Based on the above fact, a natural problem is for us how to generalize those Hardy type inequalities described above to the setting of complete Riemannian manifold with affine connections (1.1) and without any constraint on the affine Ricci curvature.

The motivation of this article is to answer the above problems. The primary objectives of the present paper are twofold: Firstly, we want to establish some L^2 improved Hardy type inequalities and L^p Hardy type inequalities for affine connections on a complete Riemannian manifold. Secondly, as application we would like to show some Rellich type inequalities, Hardy-Poincaré type inequalities and Heisenberg-Pauli-Weyl type inequalities for affine connections (1.1).

The rest of this paper is organized as follows. In Section 2, we recall the notations and the integration by parts under affine connections. In Section 3, we establish L^2 two-weight Hardy type inequality and L^p Hardy type inequality for affine connections. Weighted Rellich type inequalities for affine connections are discussed in Section 4. Finally, Section 5 is dedicated to the study of some inequalities of other types for affine connections.

2. PRELIMINARIES

In this section, we recall some preliminary results in Li and Xia [18].

2.1. Torsion-free affine connections $D^{\lambda,\mu}$. For two real constants λ, μ and two vector fields X, Y on M , a 2-parameter family of affine connections $D^{\lambda,\mu}$ defined by

$$D_X^{\lambda,\mu} Y := \nabla_X Y + \lambda du(X)Y + \lambda du(Y)X + \mu g(X, Y)\nabla u, \quad (2.1)$$

where u is a smooth function defined on M . One can check that $D_X^{\lambda,\mu} Y$ is *torsion-free*.

2.2. Affine gradient, affine Laplacian, affine divergence and affine Ricci curvature. Let ϕ be a smooth function on M^n . For an n -dimensional smooth Riemannian triple $(M^n, g, V = e^\mu)$.

(1) The affine gradient on M is defined by

$$\nabla^{D^{\lambda,\mu}} \phi := V^{\mu-\lambda} \nabla \phi. \quad (2.2)$$

(2) The affine Laplacian $\Delta^{D^{\lambda,\mu}} \phi$ on M is defined by

$$\Delta^{D^{\lambda,\mu}} \phi := V^{\mu-\lambda} [\Delta \phi + (2\mu + n\lambda) \langle \nabla u, \nabla \phi \rangle]. \quad (2.3)$$

(3) Let $X = \nabla^{D^{\lambda,\mu}} \phi$. The affine divergence $\operatorname{div}^{D^{\lambda,\mu}} X$ on M is defined by

$$\operatorname{div}^{D^{\lambda,\mu}} X := D_i^{\lambda,\mu} X^i = V^{\mu-\lambda} (\Delta \phi + (2\mu + n\lambda) \langle \nabla u, \nabla \phi \rangle) = \Delta^{D^{\lambda,\mu}} \phi. \quad (2.4)$$

(4) The affine Ricci curvature on M is defined by:

$$\begin{aligned} \operatorname{Ric}^{D^{\lambda,\mu}} := & \operatorname{Ric} - [(n-1)\lambda + \mu] \nabla^2 u + [(n-1)\lambda^2 - \mu^2] du \otimes du \\ & + [\mu \Delta u + (\mu^2 + (n-1)\lambda\mu) |\nabla u|^2] g. \end{aligned} \quad (2.5)$$

(5) The integration by parts with respect to affine connections as follows:

$$\int_M V^\tau \omega \Delta^{D^{\lambda,\mu}} \phi \, dv_g = - \int_M \langle \nabla \omega, \nabla^{D^{\lambda,\mu}} \phi \rangle V^\tau \, dv_g = \int_M V^\tau \phi \Delta^{D^{\lambda,\mu}} \omega \, dv_g \quad (2.6)$$

where ω and ϕ are smooth functions on M .

Lemma 2.1. ([18]) *Let W be any smooth vector field on M . Then*

$$V^\tau D_i^{\lambda, \mu} W^i = \nabla_i (V^\tau W^i) \quad (2.7)$$

is a divergent form with respect to the Riemannian volume form dv_g , where $\tau = (n+1)\lambda + \mu$ and we adopt the Einstein convention.

3. HARDY TYPE INEQUALITIES AND THEIR PROOFS

In this section, we will prove some integral inequalities of the Hardy type for affine connections $D^{\lambda, \mu}$. Our first result is an L^2 -version of the affine Hardy inequality without remainder term given as follows.

Theorem 3.1. *Let $(M^n, g, V = e^\mu)$ be an n -dimensional Riemannian triple and $\lambda, \mu \in \mathbb{R}$. Let $D = D^{\lambda, \mu}$ be the affine connection defined as in (2.1) and $\tau = (n+1)\lambda + \mu$. Let ρ be a nonnegative function on M such that $|\nabla^D \rho| = 1$ in the sense of distributions. Then for any compactly supported smooth function $\phi \in C_0^\infty(M \setminus \rho^{-1}\{0\})$, $1 < p < +\infty$, we have*

(i) *When $\Delta^D \rho \leq \frac{C}{\rho} V^{\lambda - \mu}$ in the sense of distributions, where $C > 0$ is a constant and $1 - \alpha - C > 0$, the following inequality*

$$\int_M \rho^\alpha |\nabla^D \phi|^2 V^{\tau + \lambda - \mu} dv_g \geq \left(\frac{1 - \alpha - C}{2} \right)^2 \int_M \rho^{\alpha - 2} \phi^2 V^{\tau + \lambda - \mu} dv_g \quad (3.1)$$

holds.

(ii) *When $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda - \mu}$ in the sense of distributions, where $C > 0$ is a constant and $C + \alpha - 1 > 0$, the following inequality*

$$\int_M \rho^\alpha |\nabla^D \phi|^2 V^{\tau + \lambda - \mu} dv_g \geq \left(\frac{\alpha + C - 1}{2} \right)^2 \int_M \rho^{\alpha - 2} \phi^2 V^{\tau + \lambda - \mu} dv_g \quad (3.2)$$

holds.

Proof. (i) Let $\gamma = \frac{1 - \alpha - C}{2}$. Clearly, $\gamma > 0$. By direct computation, we obtain

$$\begin{aligned} \Delta^D (\rho^{\alpha + 2\gamma}) &= V^{\mu - \lambda} (\Delta \rho^{\alpha + 2\gamma} + (2\mu + n\lambda) \langle \nabla u, \nabla \rho^{\alpha + 2\gamma} \rangle) \\ &= V^{\mu - \lambda} [(\alpha + 2\gamma) ((\alpha + 2\gamma - 1) |\nabla \rho|^2 \rho^{\alpha + 2\gamma - 2} + \rho^{\alpha + 2\gamma - 1} \Delta \rho) \\ &\quad + (2\mu + n\lambda) (\alpha + 2\gamma) \rho^{\alpha + 2\gamma - 1} \langle \nabla u, \nabla \rho \rangle] \\ &= (\alpha + 2\gamma) (\alpha + 2\gamma - 1) \rho^{\alpha + 2\gamma - 2} |\nabla^D \rho|^2 V^{\lambda - \mu} + (\alpha + 2\gamma) \rho^{\alpha + 2\gamma - 1} \Delta^D \rho \\ &= (\alpha + 2\gamma) \left((\alpha + 2\gamma - 1) |\nabla^D \rho|^2 \rho^{\alpha + 2\gamma - 2} V^{\lambda - \mu} + \rho^{\alpha + 2\gamma - 1} \Delta^D \rho \right), \end{aligned} \quad (3.3)$$

where, we use (2.3) in the first equation and also use (2.2) and (2.3) in the last second equation. Together with $|\nabla^D \rho| = 1$ and $\Delta^D \rho \leq \frac{C}{\rho} V^{\lambda - \mu}$, we can get that for $C \neq 1$, the following inequality

holds:

$$\begin{aligned}
& \frac{1}{\alpha+2\gamma} \Delta^D (\rho^{\alpha+2\gamma}) \\
&= (\alpha+2\gamma-1) |\nabla^D \rho|^2 \rho^{\alpha+2\gamma-2} V^{\lambda-\mu} + \rho^{\alpha+2\gamma-1} \Delta^D \rho \\
&\leq (\alpha+2\gamma-1+C) \rho^{\alpha+2\gamma-2} V^{\lambda-\mu} \\
&= 0.
\end{aligned} \tag{3.4}$$

Let $\phi = \rho^\gamma \psi$, where $\psi \in C_0^\infty(M \setminus \rho^{-1}\{0\})$. Simple calculation can be obtained,

$$\begin{aligned}
& \rho^\alpha |\nabla \phi|^2 \\
&= \rho^\alpha |\gamma \rho^{\gamma-1} \psi \nabla \rho + \rho^\gamma \nabla \psi|^2 \\
&= \rho^\alpha \left(\gamma^2 \rho^{2(\gamma-1)} \psi^2 |\nabla \rho|^2 + \rho^{2\gamma} |\nabla \psi|^2 + 2\gamma \rho^{2\gamma-1} \psi \langle \nabla \rho, \nabla \psi \rangle \right) \\
&= \gamma^2 \rho^{\alpha+2(\gamma-1)} \psi^2 |\nabla \rho|^2 + \rho^{\alpha+2\gamma} |\nabla \psi|^2 + 2\gamma \rho^{\alpha+2\gamma-1} \psi \langle \nabla \rho, \nabla \psi \rangle \\
&= \gamma^2 \rho^{\alpha+2(\gamma-1)} \psi^2 |\nabla \rho|^2 + \rho^{\alpha+2\gamma} |\nabla \psi|^2 + \frac{\gamma}{\alpha+2\gamma} \langle \nabla \rho^{\alpha+2\gamma}, \nabla \psi^2 \rangle \\
&= \gamma^2 \rho^{\alpha+2(\gamma-1)} \psi^2 |\nabla^D \rho|^2 V^{2(\lambda-\mu)} + \rho^{\alpha+2\gamma} |\nabla^D \psi|^2 V^{2(\lambda-\mu)} + \frac{\gamma}{\alpha+2\gamma} \langle V^{\lambda-\mu} \nabla^D \rho^{\alpha+2\gamma}, \nabla \psi^2 \rangle \\
&\geq \gamma^2 \rho^{\alpha+2(\gamma-1)} \psi^2 |\nabla^D \rho|^2 V^{2(\lambda-\mu)} + \frac{\gamma}{\alpha+2\gamma} \langle V^{\lambda-\mu} \nabla^D \rho^{\alpha+2\gamma}, \nabla \psi^2 \rangle.
\end{aligned} \tag{3.5}$$

Multiplying both sides of the above inequality by $V^{\tau+\mu-\lambda}$ and using $|\nabla^D \rho| = 1$, we get

$$\rho^\alpha |\nabla \phi|^2 V^{\tau+\mu-\lambda} \geq \gamma^2 \rho^{\alpha+2\gamma-2} \psi^2 V^{\tau+\lambda-\mu} + \frac{\gamma}{\alpha+2\gamma} \langle V^\tau \nabla^D \rho^{\alpha+2\gamma}, \nabla \psi^2 \rangle. \tag{3.6}$$

Integrating (3.6) over M yields

$$\begin{aligned}
& \int_M \rho^\alpha |\nabla \phi|^2 V^{\tau+\mu-\lambda} dv_g \\
&\geq \gamma^2 \int_M \rho^{\alpha+2\gamma-2} \psi^2 V^{\tau+\lambda-\mu} dv_g + \frac{\gamma}{\alpha+2\gamma} \int_M \langle V^\tau \nabla^D \rho^{\alpha+2\gamma}, \nabla \psi^2 \rangle dv_g.
\end{aligned} \tag{3.7}$$

We use the integration by parts (2.6) to derive that

$$\begin{aligned}
& \int_M \langle V^\tau \nabla^D \rho^{\alpha+2\gamma}, \nabla \psi^2 \rangle dv_g \\
&= \int_M V^\tau \nabla_i^D \rho^{\alpha+2\gamma} \nabla_i \psi^2 dv_g \\
&= - \int_M \psi^2 \nabla_i (V^\tau \nabla_i^D \rho^{\alpha+2\gamma}) dv_g \\
&= - \int_M \psi^2 V^\tau D_i (\nabla_i^D \rho^{\alpha+2\gamma}) dv_g \\
&= - \int_M \psi^2 V^\tau \Delta^D \rho^{\alpha+2\gamma} dv_g,
\end{aligned} \tag{3.8}$$

where, in the last second inequality, we use (2.7). Inserting (3.8) into (3.7), we attain

$$\begin{aligned} & \int_M \rho^\alpha |\nabla \phi|^2 V^{\tau+\mu-\lambda} dv_g \\ & \geq \gamma^2 \int_M \rho^{\alpha+2\gamma-2} \psi^2 V^{\tau+\lambda-\mu} dv_g - \frac{\gamma}{\alpha+2\gamma} \int_M \psi^2 V^\tau \Delta^D \rho^{\alpha+2\gamma} dv_g. \end{aligned}$$

Applying (3.4) and $\phi = \rho^\gamma \psi$, we obtain

$$\begin{aligned} & \int_M \rho^\alpha |\nabla \phi|^2 V^{\tau+\mu-\lambda} dv_g \\ & \geq \gamma^2 \int_M \rho^{\alpha+2\gamma-2} \psi^2 V^{\tau+\lambda-\mu} dv_g \\ & = \left(\frac{1-\alpha-C}{2} \right)^2 \int_M \rho^{\alpha-2} \phi^2 V^{\tau+\lambda-\mu} dv_g. \end{aligned} \tag{3.9}$$

On the other hand, clearly, when $C = 1$, by applying a similar computation as above, we can get

$$\begin{aligned} \Delta^D (\ln \rho) & = V^{\mu-\lambda} (\Delta \ln \rho + (2\mu + n\lambda) \langle \nabla u, \nabla \ln \rho \rangle) \\ & = V^{\mu-\lambda} \left(-\frac{1}{\rho^2} |\nabla \rho|^2 + \frac{1}{\rho} \Delta \rho + (2\mu + n\lambda) \frac{1}{\rho} \langle \nabla u, \nabla \rho \rangle \right) \\ & = -\frac{1}{\rho^2} |\nabla^D \rho|^2 V^{\lambda-\mu} + \frac{1}{\rho} \Delta^D \rho \end{aligned} \tag{3.10}$$

and

$$\begin{aligned} \rho^\alpha |\nabla \phi|^2 & = \gamma^2 \rho^{\alpha+2(\gamma-1)} \psi^2 |\nabla \rho|^2 + \rho^{\alpha+2\gamma} |\nabla \psi|^2 + 2\gamma \rho^{\alpha+2\gamma-1} \psi \langle \nabla \rho, \nabla \psi \rangle \\ & = \gamma^2 \rho^{\alpha+2(\gamma-1)} \psi^2 |\nabla \rho|^2 + \rho^{\alpha+2\gamma} |\nabla \psi|^2 + 2\gamma \rho^{-1} \psi \langle \nabla \rho, \nabla \psi \rangle \\ & = \gamma^2 \rho^{\alpha+2(\gamma-1)} \psi^2 |\nabla^D \rho|^2 V^{2(\lambda-\mu)} + \rho^{\alpha+2\gamma} |\nabla^D \psi|^2 V^{2(\lambda-\mu)} + 2\gamma \rho^{-1} \psi \langle \nabla \rho, \nabla \psi \rangle \\ & \geq \gamma^2 \rho^{\alpha+2(\gamma-1)} \psi^2 |\nabla^D \rho|^2 V^{2(\lambda-\mu)} + \gamma \langle \nabla \ln \rho, \nabla \psi^2 \rangle. \end{aligned} \tag{3.11}$$

Together with $|\nabla^D \rho| = 1$ and $\Delta^D \rho \leq \frac{C}{\rho} V^{\lambda-\mu}$ in (3.10), we can get that for $C = 1$, the following inequality holds:

$$\begin{aligned} \Delta^D (\ln \rho) & \leq (C-1) \frac{1}{\rho^2} V^{\lambda-\mu} \\ & = 0. \end{aligned} \tag{3.12}$$

Multiplying both sides in (3.11) by $V^{\tau+\mu-\lambda}$ and integrating over M yields

$$\begin{aligned} \int_M \rho^\alpha |\nabla \phi|^2 V^{\tau+\mu-\lambda} dv_g & \geq \gamma^2 \int_M \rho^{\alpha+2\gamma-2} \psi^2 |\nabla^D \rho|^2 V^{\tau+\lambda-\mu} dv_g \\ & \quad + \gamma \int_M \langle \nabla \ln \rho, \nabla \psi^2 \rangle V^{\tau+\mu-\lambda} dv_g. \end{aligned} \tag{3.13}$$

Applying the integration by parts, we get

$$\begin{aligned}
& \int_M \langle \nabla \ln \rho, \nabla \psi^2 \rangle V^{\tau+\mu-\lambda} dv_g \\
&= \int_M \langle V^\tau \nabla^D \ln \rho, \nabla \psi^2 \rangle dv_g \\
&= \int_M V^\tau \nabla_i^D (\ln \rho) \nabla_i \psi^2 dv_g \\
&= - \int_M \psi^2 \nabla_i (V^\tau \nabla_i^D \ln \rho) dv_g \\
&= - \int_M \psi^2 V^\tau D_i (\nabla_i^D \ln \rho) dv_g \\
&= - \int_M V^\tau \psi^2 \Delta^D (\ln \rho) dv_g.
\end{aligned} \tag{3.14}$$

Then combining (3.13) and (3.14), we obtain

$$\begin{aligned}
& \int_M \rho^\alpha |\nabla \phi|^2 V^{\tau+\mu-\lambda} dv_g \\
&\geq \gamma^2 \int_M \rho^{\alpha+2\gamma-2} \psi^2 V^{\tau+\lambda-\mu} dv_g - \gamma \int_M V^\tau \psi^2 \Delta^D (\ln \rho) dv_g,
\end{aligned} \tag{3.15}$$

where $|\nabla^D \rho| = 1$. Substituting (3.12) into (3.15), since $\gamma = -\frac{\alpha}{2} > 0$ and $\phi = \rho^\gamma \psi$, we can get

$$\begin{aligned}
& \int_M \rho^\alpha |\nabla \phi|^2 V^{\tau+\mu-\lambda} dv_g \\
&= \int_M \rho^\alpha |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \\
&\geq \left(-\frac{\alpha}{2}\right)^2 \int_M \rho^{\alpha-2} \phi^2 V^{\tau+\lambda-\mu} dv_g \\
&= \left(\frac{\alpha}{2}\right)^2 \int_M \rho^{\alpha-2} \phi^2 V^{\tau+\lambda-\mu} dv_g.
\end{aligned} \tag{3.16}$$

By (3.16), we know that (3.1) is true.

(ii) Let $\gamma = \frac{1-\alpha-C}{2}$. Clearly, $\gamma < 0$. By direct computation, we have

$$\begin{aligned}
\Delta^D (\rho^{\alpha+2\gamma}) &= V^{\mu-\lambda} (\Delta \rho^{\alpha+2\gamma} + (2\mu + n\lambda) \langle \nabla u, \nabla \rho^{\alpha+2\gamma} \rangle) \\
&= V^{\mu-\lambda} [(\alpha + 2\gamma) ((\alpha + 2\gamma - 1) |\nabla \rho|^2 \rho^{\alpha+2\gamma-2} + \rho^{\alpha+2\gamma-1} \Delta \rho) \\
&\quad + (2\mu + n\lambda) (\alpha + 2\gamma) \rho^{\alpha+2\gamma-1} \langle \nabla u, \nabla \rho \rangle] \\
&= (\alpha + 2\gamma) (\alpha + 2\gamma - 1) \rho^{\alpha+2\gamma-2} |\nabla^D \rho|^2 V^{\lambda-\mu} + (\alpha + 2\gamma) \rho^{\alpha+2\gamma-1} \Delta^D \rho \\
&= (\alpha + 2\gamma) \left((\alpha + 2\gamma - 1) |\nabla^D \rho|^2 \rho^{\alpha+2\gamma-2} V^{\lambda-\mu} + \rho^{\alpha+2\gamma-1} \Delta^D \rho \right),
\end{aligned} \tag{3.17}$$

where, we use (2.3) in the first equation and also use (2.2) and (2.3) in the last second equation. Together with $|\nabla^D \rho| = 1$ and $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$, we can get that for $C \neq 1$, the following inequality

holds:

$$\begin{aligned}
& \frac{1}{\alpha+2\gamma} \Delta^D (\rho^{\alpha+2\gamma}) \\
&= (\alpha+2\gamma-1) |\nabla^D \rho|^2 \rho^{\alpha+2\gamma-2} V^{\lambda-\mu} + \rho^{\alpha+2\gamma-1} \Delta^D \rho \\
&\geq (\alpha+2\gamma-1+C) \rho^{\alpha+2\gamma-2} V^{\lambda-\mu} \\
&= 0.
\end{aligned} \tag{3.18}$$

Let $\phi = \rho^\gamma \psi$, where $\psi \in C_0^\infty(M \setminus \rho^{-1}\{0\})$. Simple calculation can be obtained,

$$\begin{aligned}
& \rho^\alpha |\nabla \phi|^2 \\
&= \rho^\alpha |\gamma \rho^{\gamma-1} \psi \nabla \rho + \rho^\gamma \nabla \psi|^2 \\
&= \rho^\alpha \left(\gamma^2 \rho^{2(\gamma-1)} \psi^2 |\nabla \rho|^2 + \rho^{2\gamma} |\nabla \psi|^2 + 2\gamma \rho^{2\gamma-1} \psi \langle \nabla \rho, \nabla \psi \rangle \right) \\
&= \gamma^2 \rho^{\alpha+2(\gamma-1)} \psi^2 |\nabla \rho|^2 + \rho^{\alpha+2\gamma} |\nabla \psi|^2 + 2\gamma \rho^{\alpha+2\gamma-1} \psi \langle \nabla \rho, \nabla \psi \rangle \\
&= \gamma^2 \rho^{\alpha+2(\gamma-1)} \psi^2 |\nabla \rho|^2 + \rho^{\alpha+2\gamma} |\nabla \psi|^2 + \frac{\gamma}{\alpha+2\gamma} \langle \nabla \rho^{\alpha+2\gamma}, \nabla \psi^2 \rangle \\
&= \gamma^2 \rho^{\alpha+2(\gamma-1)} \psi^2 |\nabla^D \rho|^2 V^{2(\lambda-\mu)} + \rho^{\alpha+2\gamma} |\nabla^D \psi|^2 V^{2(\lambda-\mu)} + \frac{\gamma}{\alpha+2\gamma} \langle V^{\lambda-\mu} \nabla^D \rho^{\alpha+2\gamma}, \nabla \psi^2 \rangle \\
&\geq \gamma^2 \rho^{\alpha+2(\gamma-1)} \psi^2 |\nabla^D \rho|^2 V^{2(\lambda-\mu)} + \frac{\gamma}{\alpha+2\gamma} \langle V^{\lambda-\mu} \nabla^D \rho^{\alpha+2\gamma}, \nabla \psi^2 \rangle.
\end{aligned} \tag{3.19}$$

Multiplying both sides of the above inequality by $V^{\tau+\mu-\lambda}$ and using $|\nabla^D \rho| = 1$, we get

$$\rho^\alpha |\nabla \phi|^2 V^{\tau+\mu-\lambda} \geq \gamma^2 \int_M \rho^{\alpha+2\gamma-2} \psi^2 V^{\tau+\lambda-\mu} + \frac{\gamma}{\alpha+2\gamma} \langle V^\tau \nabla^D \rho^{\alpha+2\gamma}, \nabla \psi^2 \rangle. \tag{3.20}$$

Integrating (3.20) over M yields

$$\begin{aligned}
& \int_M \rho^\alpha |\nabla \phi|^2 V^{\tau+\mu-\lambda} dv_g \\
&\geq \gamma^2 \int_M \rho^{\alpha+2\gamma-2} \psi^2 V^{\tau+\lambda-\mu} dv_g + \frac{\gamma}{\alpha+2\gamma} \int_M \langle V^\tau \nabla^D \rho^{\alpha+2\gamma}, \nabla \psi^2 \rangle dv_g.
\end{aligned} \tag{3.21}$$

We use the integration by parts (2.6) to derive that

$$\begin{aligned}
& \int_M \langle V^\tau \nabla^D \rho^{\alpha+2\gamma}, \nabla \psi^2 \rangle dv_g \\
&= \int_M V^\tau \nabla_i^D \rho^{\alpha+2\gamma} \nabla_i \psi^2 dv_g \\
&= - \int_M \psi^2 \nabla_i (V^\tau \nabla_i^D \rho^{\alpha+2\gamma}) dv_g \\
&= - \int_M \psi^2 V^\tau D_i (\nabla_i^D \rho^{\alpha+2\gamma}) dv_g \\
&= - \int_M \psi^2 V^\tau \Delta^D \rho^{\alpha+2\gamma} dv_g,
\end{aligned} \tag{3.22}$$

where, in the last second inequality, we use (2.7). Inserting (3.23) into (3.21), we attain

$$\begin{aligned} & \int_M \rho^\alpha |\nabla \phi|^2 V^{\tau+\mu-\lambda} dv_g \\ & \geq \gamma^2 \int_M \rho^{\alpha+2\gamma-2} \psi^2 V^{\tau+\lambda-\mu} dv_g - \frac{\gamma}{\alpha+2\gamma} \int_M \psi^2 \Delta^D \rho^{\alpha+2\gamma} V^\tau dv_g. \end{aligned}$$

Applying (3.18) and $\phi = \rho^\gamma \psi$, we obtain

$$\begin{aligned} & \int_M \rho^\alpha |\nabla \phi|^2 V^{\tau+\mu-\lambda} dv_g \\ & \geq \gamma^2 \int_M \rho^{\alpha+2\gamma-2} \psi^2 V^{\tau+\lambda-\mu} dv_g \\ & = \left(\frac{1-\alpha-C}{2} \right)^2 \int_M \rho^{\alpha-2} \phi^2 V^{\tau+\lambda-\mu} dv_g \\ & = \left(\frac{\alpha+C-1}{2} \right)^2 \int_M \rho^{\alpha-2} \phi^2 V^{\tau+\lambda-\mu} dv_g. \end{aligned} \tag{3.23}$$

On the other hand, clearly, when $C = 1$, by applying a similar computation as above, we can get

$$\begin{aligned} \Delta^D (\ln \rho) & = V^{\mu-\lambda} (\Delta \ln \rho + (2\mu + n\lambda) \langle \nabla u, \nabla \ln \rho \rangle) \\ & = V^{\mu-\lambda} \left(-\frac{1}{\rho^2} |\nabla \rho|^2 + \frac{1}{\rho} \Delta \rho + (2\mu + n\lambda) \frac{1}{\rho} \langle \nabla u, \nabla \rho \rangle \right) \\ & = -\frac{1}{\rho^2} |\nabla^D \rho|^2 V^{\lambda-\mu} + \frac{1}{\rho} \Delta^D \rho \end{aligned} \tag{3.24}$$

and

$$\begin{aligned} \rho^\alpha |\nabla \phi|^2 & = \gamma^2 \rho^{\alpha+2(\gamma-1)} \psi^2 |\nabla \rho|^2 + \rho^{\alpha+2\gamma} |\nabla \psi|^2 + 2\gamma \rho^{\alpha+2\gamma-1} \psi \langle \nabla \rho, \nabla \psi \rangle \\ & = \gamma^2 \rho^{\alpha+2(\gamma-1)} \psi^2 |\nabla \rho|^2 + \rho^{\alpha+2\gamma} |\nabla \psi|^2 + 2\gamma \rho^{-1} \psi \langle \nabla \rho, \nabla \psi \rangle \\ & = \gamma^2 \rho^{\alpha+2(\gamma-1)} \psi^2 |\nabla^D \rho|^2 V^{2(\lambda-\mu)} + \rho^{\alpha+2\gamma} |\nabla^D \psi|^2 V^{2(\lambda-\mu)} + 2\gamma \rho^{-1} \psi \langle \nabla \rho, \nabla \psi \rangle \\ & \geq \gamma^2 \rho^{\alpha+2(\gamma-1)} \psi^2 |\nabla^D \rho|^2 V^{2(\lambda-\mu)} + \gamma \langle \nabla \ln \rho, \nabla \psi^2 \rangle. \end{aligned} \tag{3.25}$$

Together with $|\nabla^D \rho| = 1$ and $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$ in (3.10), we can get that for $C = 1$, the following inequality holds:

$$\begin{aligned} \Delta^D (\ln \rho) & \geq (C-1) \frac{1}{\rho^2} V^{\lambda-\mu} \\ & = 0. \end{aligned} \tag{3.26}$$

Multiplying both sides in (3.25) by $V^{\tau+\mu-\lambda}$ and integrating over M yields

$$\begin{aligned} & \int_M \rho^\alpha |\nabla \phi|^2 V^{\tau+\mu-\lambda} dv_g \\ & \geq \gamma^2 \int_M \rho^{\alpha+2\gamma-2} \psi^2 |\nabla^D \rho|^2 V^{\tau+\lambda-\mu} dv_g + \gamma \int_M \langle \nabla \ln \rho, \nabla \psi^2 \rangle V^{\tau+\mu-\lambda} dv_g. \end{aligned} \tag{3.27}$$

Applying the integration by parts, we get

$$\begin{aligned}
& \int_M \langle \nabla \ln \rho, \nabla \psi^2 \rangle V^{\tau+\mu-\lambda} dv_g \\
&= \int_M \langle V^\tau \nabla^D \ln \rho, \nabla \psi^2 \rangle dv_g \\
&= \int_M V^\tau \nabla_i^D (\ln \rho) \nabla_i \psi^2 dv_g \\
&= - \int_M \psi^2 \nabla_i (V^\tau \nabla_i^D \ln \rho) dv_g \\
&= - \int_M \psi^2 V^\tau D_i (\nabla_i^D \ln \rho) dv_g \\
&= - \int_M V^\tau \psi^2 \Delta^D (\ln \rho) dv_g.
\end{aligned} \tag{3.28}$$

Then combining (3.27) and (3.28), we obtain

$$\begin{aligned}
& \int_M \rho^\alpha |\nabla \phi|^2 V^{\tau+\mu-\lambda} dv_g \\
&\geq \gamma^2 \int_M \rho^{\alpha+2\gamma-2} \psi^2 V^{\tau+\lambda-\mu} dv_g - \gamma \int_M V^\tau \psi^2 \Delta^D (\ln \rho) dv_g,
\end{aligned} \tag{3.29}$$

where $|\nabla^D \rho| = 1$. Substituting (3.26) into (3.29), since $\gamma = -\frac{\alpha}{2} < 0$ and $\phi = \rho^\gamma \psi$, we can get

$$\begin{aligned}
& \int_M \rho^\alpha |\nabla \phi|^2 V^{\tau+\mu-\lambda} dv_g \\
&= \int_M \rho^\alpha |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \\
&\geq \left| -\frac{\alpha}{2} \right|^2 \int_M \rho^{\alpha-2} \phi^2 V^{\tau+\lambda-\mu} dv_g \\
&\geq \left| \frac{\alpha}{2} \right|^2 \int_M \rho^{\alpha-2} \phi^2 V^{\tau+\lambda-\mu} dv_g.
\end{aligned} \tag{3.30}$$

By (3.30), we know that (3.2) is true. This completes the proof of Theorem 3.1. \square

Our second result is L^p -version of the Hardy type inequality without remainder term given as follows.

Theorem 3.2. *Let $(M^n, g, V = e^\mu)$ be an n -dimensional Riemannian triple and $\lambda, \mu \in \mathbb{R}$. Let $D = D^{\lambda, \mu}$ be the affine connection defined as in (2.1) and $\tau = (n+1)\lambda + \mu$. Let ρ be a nonnegative function on M such that $|\nabla^D \rho| = 1$ in the sense of distributions. Then, for any $p, q \in \mathbb{R}$, $1 < p < +\infty$, $0 \leq q \leq p$, and any compactly supported smooth function $\phi \in C_0^\infty(M \setminus \rho^{-1}\{0\})$, we have*

(i) *When $\Delta^D \rho \leq \frac{C}{\rho} V^{\lambda-\mu}$ in the sense of distributions, where C is a constant and $q - C - 1 > 0$, the following inequality*

$$\int_M \frac{|\phi|^p}{\rho^q} V^{\tau+\lambda-\mu} dv_g \leq \left(\frac{p}{q-C-1} \right)^q \left(\int_M |\nabla^D \phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{q}{p}} \left(\int_M |\phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{p-q}{p}} \tag{3.31}$$

holds.

(ii) When $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$ in the sense of distributions, where C is a constant and $C+1-q > 0$, the following inequality

$$\int_M \frac{|\phi|^p}{\rho^q} V^{\tau+\lambda-\mu} dv_g \leq \left(\frac{p}{C+1-q} \right)^q \left(\int_M |\nabla^D \phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{q}{p}} \left(\int_M |\phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{p-q}{p}} \quad (3.32)$$

holds.

Proof. (i) Let $\operatorname{div}^D X = D_i X^i$ be the divergence of X with respect to D , where X is a vector field on M . By direct computation, together with assumptions $|\nabla^D \rho| = 1$ and $\Delta^D \rho \leq \frac{C}{\rho} V^{\lambda-\mu}$, we have

$$\begin{aligned} & V^\tau D_i \left(|\phi|^p \frac{\nabla_i^D \rho}{\rho^{q-1}} \right) \\ &= \nabla_i \left(V^{\tau+\mu-\lambda} |\phi|^p \frac{\nabla_i \rho}{\rho^{q-1}} \right) \\ &= \nabla_i \left(e^{u(\tau+\mu-\lambda)} |\phi|^p \frac{\nabla_i \rho}{\rho^{q-1}} \right) \\ &= (\tau + \mu - \lambda) \langle \nabla u, \nabla \rho \rangle \frac{|\phi|^p}{\rho^{q-1}} V^{\tau+\mu-\lambda} + p \frac{|\phi|^{p-1}}{\rho^{q-1}} \langle \nabla \phi, \nabla \rho \rangle V^{\tau+\mu-\lambda} \\ &\quad + \frac{|\phi|^p}{\rho^{q-1}} V^{\tau+\mu-\lambda} \Delta \rho + (1-q) |\nabla \rho|^2 \frac{|\phi|^p}{\rho^q} V^{\tau+\mu-\lambda} \\ &= \frac{|\phi|^p}{\rho^{q-1}} V^\tau \Delta^D \rho + p \frac{|\phi|^{p-1}}{\rho^{q-1}} \langle \nabla^D \phi, \nabla^D \rho \rangle V^{\tau+\lambda-\mu} + (1-q) |\nabla^D \rho|^2 \frac{|\phi|^p}{\rho^q} V^{\tau+\lambda-\mu} \\ &\leq (1+C-q) \frac{|\phi|^p}{\rho^q} V^{\tau+\lambda-\mu} + p \frac{|\phi|^{p-1}}{\rho^{q-1}} \langle \nabla^D \phi, \nabla^D \rho \rangle V^{\tau+\lambda-\mu}. \end{aligned} \quad (3.33)$$

Integrating the above inequality (3.33) over M yields

$$\begin{aligned} & (q-C-1) \int_M \frac{|\phi|^p}{\rho^q} V^{\tau+\lambda-\mu} dv_g \\ & \leq p \int_M \frac{|\phi|^{p-1}}{\rho^{q-1}} \langle \nabla^D \phi, \nabla^D \rho \rangle V^{\tau+\lambda-\mu} dv_g \\ & \leq p \int_M \frac{|\phi|^{p-1}}{\rho^{q-1}} |\nabla^D \phi| |\nabla^D \rho| V^{\tau+\lambda-\mu} dv_g \\ & = p \int_M \frac{|\phi|^{p-1}}{\rho^{q-1}} |\nabla^D \phi| V^{\tau+\lambda-\mu} dv_g. \end{aligned} \quad (3.34)$$

It follows from Hölder's inequality that

$$\begin{aligned}
& \int_M \frac{|\phi|^{p-1}}{\rho^{q-1}} |\nabla^D \phi| V^{\tau+\lambda-\mu} dv_g \\
&= \int_M |\nabla^D \phi| V^{\frac{\tau+\lambda-\mu}{p}} \frac{|\phi|^{p-1}}{\rho^{q-1}} V^{\frac{(\tau+\lambda-\mu)(p-1)}{p}} dv_g \\
&\leq \left(\int_M \left[|\nabla^D \phi| V^{\frac{\tau+\lambda-\mu}{p}} \right]^p dv_g \right)^{\frac{1}{p}} \left(\int_M \left[\frac{|\phi|^{p-1}}{\rho^{q-1}} V^{\frac{(\tau+\lambda-\mu)(p-1)}{p}} \right]^{\frac{p}{p-1}} dv_g \right)^{\frac{p-1}{p}} \\
&= \left(\int_M |\nabla^D \phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{1}{p}} \left(\int_M \frac{|\phi|^p}{\rho^{\frac{p(q-1)}{p-1}}} V^{\tau+\lambda-\mu} dv_g \right)^{\frac{p-1}{p}}
\end{aligned} \tag{3.35}$$

and

$$\begin{aligned}
& \int_M \frac{|\phi|^p}{\rho^{\frac{p(q-1)}{p-1}}} V^{\tau+\lambda-\mu} dv_g \\
&= \int_M \rho^{-\frac{p(q-1)}{p-1}} \left(|\phi|^p V^{\tau+\lambda-\mu} \right)^{\frac{p(q-1)}{(p-1)q}} \left(|\phi|^p V^{\tau+\lambda-\mu} \right)^{\left(1-\frac{p(q-1)}{(p-1)q}\right)} dv_g \\
&\leq \left(\int_M \left[\frac{\left(|\phi|^p V^{\tau+\lambda-\mu} \right)^{\frac{p(q-1)}{(p-1)q}}}{\rho^{\frac{p(q-1)}{p-1}}} \right]^{\frac{(p-1)q}{p(q-1)}} dv_g \right)^{\frac{p(q-1)}{(p-1)q}} \left(\int_M \left[\left(|\phi|^p V^{\tau+\lambda-\mu} \right)^{\left(1-\frac{p(q-1)}{(p-1)q}\right)} \right]^{\frac{1}{\left(1-\frac{p(q-1)}{(p-1)q}\right)}} dv_g \right)^{\left(1-\frac{p(q-1)}{(p-1)q}\right)} \\
&= \left(\int_M \frac{|\phi|^p V^{\tau+\lambda-\mu}}{\rho^q} dv_g \right)^{\frac{p(q-1)}{(p-1)q}} \left(\int_M |\phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\left(1-\frac{p(q-1)}{(p-1)q}\right)}.
\end{aligned} \tag{3.36}$$

Taking (3.36) into (3.35), we have

$$\begin{aligned}
& \int_M \frac{|\phi|^{p-1}}{\rho^{q-1}} |\nabla^D \phi| V^{\tau+\lambda-\mu} dv_g \\
&\leq \left(\int_M |\nabla^D \phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{1}{p}} \left(\int_M \frac{|\phi|^p V^{\tau+\lambda-\mu}}{\rho^q} dv_g \right)^{\frac{q-1}{q}} \left(\int_M |\phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\left(1-\frac{p(q-1)}{(p-1)q}\right) \frac{p-1}{p}}.
\end{aligned} \tag{3.37}$$

Then combining (3.37) and (3.34), we obtain

$$\left(\int_M \frac{|\phi|^p}{\rho^q} V^{\tau+\lambda-\mu} dv_g \right)^{\frac{1}{q}} \leq \frac{p}{q-C-1} \left(\int_M |\nabla^D \phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{1}{p}} \left(\int_M |\phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\left(1-\frac{p(q-1)}{(p-1)q}\right) \frac{p-1}{p}},$$

which is equivalent to

$$\int_M \frac{|\phi|^p}{\rho^q} V^{\tau+\lambda-\mu} dv_g \leq \left(\frac{p}{q-C-1} \right)^q \left(\int_M |\nabla^D \phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{q}{p}} \left(\int_M |\phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{p-q}{p}}.$$

(ii) By direct computation, together with assumptions $|\nabla^D \rho| = 1$ and $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$, we have

$$\begin{aligned}
& V^\tau D_i \left(|\phi|^p \frac{\nabla_i^D \rho}{\rho^{q-1}} \right) \\
&= \nabla_i \left(V^{\tau+\mu-\lambda} |\phi|^p \frac{\nabla_i \rho}{\rho^{q-1}} \right) \\
&= \nabla_i \left(e^{u(\tau+\mu-\lambda)} |\phi|^p \frac{\nabla_i \rho}{\rho^{q-1}} \right) \\
&= (\tau + \mu - \lambda) \langle \nabla u, \nabla \rho \rangle \frac{|\phi|^p}{\rho^{q-1}} V^{\tau+\mu-\lambda} + p \frac{|\phi|^{p-1}}{\rho^{q-1}} \langle \nabla \phi, \nabla \rho \rangle V^{\tau+\mu-\lambda} \\
&\quad + \frac{|\phi|^p}{\rho^{q-1}} V^{\tau+\mu-\lambda} \Delta \rho + (1-q) |\nabla \rho|^2 \frac{|\phi|^p}{\rho^q} V^{\tau+\mu-\lambda} \\
&= \frac{|\phi|^p}{\rho^{q-1}} V^\tau \Delta^D \rho + p \frac{|\phi|^{p-1}}{\rho^{q-1}} \langle \nabla^D \phi, \nabla^D \rho \rangle V^{\tau+\lambda-\mu} + (1-q) |\nabla^D \rho|^2 \frac{|\phi|^p}{\rho^q} V^{\tau+\lambda-\mu} \\
&\geq (1+C-q) \frac{|\phi|^p}{\rho^q} V^{\tau+\lambda-\mu} + p \frac{|\phi|^{p-1}}{\rho^{q-1}} \langle \nabla^D \phi, \nabla^D \rho \rangle V^{\tau+\lambda-\mu}.
\end{aligned} \tag{3.38}$$

Integrating the above inequality (3.38) over M yields

$$\begin{aligned}
& (C+1-q) \int_M \frac{|\phi|^p}{\rho^q} V^{\tau+\lambda-\mu} dv_g \\
&\leq -p \int_M \frac{|\phi|^{p-1}}{\rho^{q-1}} \langle \nabla^D \phi, \nabla^D \rho \rangle V^{\tau+\lambda-\mu} dv_g \\
&\leq \left| -p \int_M \frac{|\phi|^{p-1}}{\rho^{q-1}} \langle \nabla^D \phi, \nabla^D \rho \rangle V^{\tau+\lambda-\mu} dv_g \right| \\
&\leq p \int_M \frac{|\phi|^{p-1}}{\rho^{q-1}} |\nabla^D \phi| |\nabla^D \rho| V^{\tau+\lambda-\mu} dv_g \\
&= p \int_M \frac{|\phi|^{p-1}}{\rho^{q-1}} |\nabla^D \phi| V^{\tau+\lambda-\mu} dv_g.
\end{aligned} \tag{3.39}$$

It follows from Hölder's inequality that

$$\begin{aligned}
& \int_M \frac{|\phi|^{p-1}}{\rho^{q-1}} |\nabla^D \phi| V^{\tau+\lambda-\mu} dv_g = \int_M |\nabla^D \phi| V^{\frac{\tau+\lambda-\mu}{p}} \frac{|\phi|^{p-1}}{\rho^{q-1}} |\nabla^D \phi| V^{\frac{(\tau+\lambda-\mu)(p-1)}{p}} dv_g \\
&\leq \left(\int_M \left[|\nabla^D \phi| V^{\frac{\tau+\lambda-\mu}{p}} \right]^p dv_g \right)^{\frac{1}{p}} \left(\int_M \left[\frac{|\phi|^{p-1}}{\rho^{q-1}} |\nabla^D \phi| V^{\frac{(\tau+\lambda-\mu)(p-1)}{p}} \right]^{\frac{p}{p-1}} dv_g \right)^{\frac{p-1}{p}} \\
&= \left(\int_M |\nabla^D \phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{1}{p}} \left(\int_M \frac{|\phi|^p}{\rho^{\frac{p(q-1)}{p-1}}} V^{\tau+\lambda-\mu} dv_g \right)^{\frac{p-1}{p}}
\end{aligned} \tag{3.40}$$

and

$$\begin{aligned}
& \int_M \frac{|\phi|^p}{\rho^{\frac{p(q-1)}{p-1}}} V^{\tau+\lambda-\mu} dv_g \\
&= \int_M \rho^{-\frac{p(q-1)}{p-1}} \left(|\phi|^p V^{\tau+\lambda-\mu} \right)^{\frac{p(q-1)}{(p-1)q}} \left(|\phi|^p V^{\tau+\lambda-\mu} \right)^{\left(1-\frac{p(q-1)}{(p-1)q}\right)} dv_g \\
&\leq \left(\int_M \left[\frac{\left(|\phi|^p V^{\tau+\lambda-\mu} \right)^{\frac{p(q-1)}{(p-1)q}}}{\rho^{\frac{p(q-1)}{p-1}}} \right]^{\frac{(p-1)q}{p(q-1)}} dv_g \right)^{\frac{p(q-1)}{(p-1)q}} \left(\int_M \left[\left(|\phi|^p V^{\tau+\lambda-\mu} \right)^{\left(1-\frac{p(q-1)}{(p-1)q}\right)} \right]^{\frac{1}{\left(1-\frac{p(q-1)}{(p-1)q}\right)}} dv_g \right)^{\left(1-\frac{p(q-1)}{(p-1)q}\right)} \\
&= \left(\int_M \frac{|\phi|^p V^{\tau+\lambda-\mu}}{\rho^q} dv_g \right)^{\frac{p(q-1)}{(p-1)q}} \left(\int_M |\phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\left(1-\frac{p(q-1)}{(p-1)q}\right)}.
\end{aligned} \tag{3.41}$$

Taking (3.41) into (3.40), we have

$$\begin{aligned}
& \int_M \frac{|\phi|^{p-1}}{\rho^{q-1}} |\nabla^D \phi|^p V^{\tau+\lambda-\mu} dv_g \\
&\leq \left(\int_M |\nabla^D \phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{1}{p}} \left(\int_M \frac{|\phi|^p V^{\tau+\lambda-\mu}}{\rho^q} dv_g \right)^{\frac{q-1}{q}} \left(\int_M |\phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\left(1-\frac{p(q-1)}{(p-1)q}\right) \frac{p-1}{p}}.
\end{aligned} \tag{3.42}$$

Then combining (3.42) and (3.39), we obtain

$$\left(\int_M \frac{|\phi|^p}{\rho^q} V^{\tau+\lambda-\mu} dv_g \right)^{\frac{1}{q}} \leq \frac{p}{C+1-q} \left(\int_M |\nabla^D \phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{1}{p}} \left(\int_M |\phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\left(1-\frac{p(q-1)}{(p-1)q}\right) \frac{p-1}{p}},$$

which is equivalent to

$$\int_M \frac{|\phi|^p}{\rho^q} V^{\tau+\lambda-\mu} dv_g \leq \left(\frac{p}{C+1-q} \right)^q \left(\int_M |\nabla^D \phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{q}{p}} \left(\int_M |\phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{p-q}{p}}.$$

This completes the proof of Theorem 3.2. \square

Under affine connections $D^{\lambda,\mu}$, we now prove an L^2 -version two-weight Hardy type inequality with nonnegative remainder terms.

Theorem 3.3. *Let $(M^n, g, V = e^\mu)$ be an n -dimensional Riemannian triple and $\lambda, \mu \in \mathbb{R}$. Let $D = D^{\lambda,\mu}$ be the affine connection defined as in (2.1) and $\tau = (n+1)\lambda + \mu$. Let $a(x)$ and $b(x)$ be nonnegative weight functions in M . Let ω be a positive function satisfying the differential inequality*

$$-V^\tau D_i (a(x) \nabla_i^D \omega) \geq b(x) \omega V^{\tau+\mu-\lambda} \tag{3.43}$$

almost everywhere in M . Then the following integral inequality

$$\int_M (a(x) |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} - b(x) \phi^2 V^{\tau+\mu-\lambda}) dv_g \geq \int_M a(x) \omega^2 \left| \nabla^D \left(\frac{\phi}{\omega} \right) \right|^2 V^{\tau+\lambda-\mu} dv_g \tag{3.44}$$

holds for any $\phi \in C_0^\infty(M)$.

Proof. Let $\phi = \omega\varphi$, where $\varphi \in C_0^\infty(M)$. Simple calculation can be obtained,

$$\begin{aligned}
& |\nabla\phi|^2 \\
&= |\omega\nabla\varphi + \varphi\nabla\omega|^2 \\
&= \omega^2|\nabla\varphi|^2 + \varphi^2|\nabla\omega|^2 + 2\omega\varphi\langle\nabla\varphi, \nabla\omega\rangle \\
&= \omega^2|\nabla\varphi|^2 + \varphi^2|\nabla\omega|^2 + \omega\langle\nabla\omega, \nabla\varphi^2\rangle.
\end{aligned} \tag{3.45}$$

Multiplying both sides of (3.45) by $a(x)V^{\tau+\mu-\lambda}$ and applying the integration by parts to the last term on the right hand side of the inequality over M gives

$$\begin{aligned}
& \int_M a(x)|\nabla\phi|^2 V^{\tau+\mu-\lambda} dv_g \\
&= \int_M a(x)\varphi^2|\nabla\omega|^2 V^{\tau+\mu-\lambda} dv_g + \int_M a(x)\omega^2|\nabla\varphi|^2 V^{\tau+\mu-\lambda} dv_g \\
&\quad + \int_M a(x)\omega\langle\nabla\omega, \nabla\varphi^2\rangle V^{\tau+\mu-\lambda} dv_g \\
&= \int_M a(x)\varphi^2|\nabla\omega|^2 V^{\tau+\mu-\lambda} dv_g + \int_M a(x)\omega^2|\nabla\varphi|^2 V^{\tau+\mu-\lambda} dv_g \\
&\quad - \int_M \varphi^2\nabla_i \left(a(x)\omega V^{\tau+\mu-\lambda} \nabla_i \omega \right) dv_g \\
&= - \int_M \varphi^2\omega\nabla_i \left(a(x)V^{\tau+\mu-\lambda} \nabla_i \omega \right) dv_g + \int_M a(x)\omega^2|\nabla\varphi|^2 V^{\tau+\mu-\lambda} dv_g \\
&= - \int_M \varphi^2\omega V^\tau D_i \left(a(x)\nabla_i^D \omega \right) dv_g + \int_M a(x)\omega^2|\nabla\varphi|^2 V^{\tau+\mu-\lambda} dv_g,
\end{aligned}$$

where, in the second and third inequalities, we use the integration by parts (2.6). Since $-V^\tau D_i \left(a(x)\nabla_i^D \omega \right) \geq b(x)\omega V^{\tau+\mu-\lambda}$ then we get

$$\begin{aligned}
& \int_M a(x)|\nabla\phi|^2 V^{\tau+\mu-\lambda} dv_g \\
&\geq \int_M b(x)\phi^2 V^{\tau+\mu-\lambda} dv_g + \int_M a(x)\omega^2|\nabla\varphi|^2 V^{\tau+\mu-\lambda} dv_g.
\end{aligned}$$

Since $\varphi = \frac{\phi}{\omega}$, we obtained the desired inequality

$$\begin{aligned}
& \int_M a(x)|\nabla\phi|^2 V^{\tau+\mu-\lambda} dv_g = \int_M a(x)|\nabla^D\phi|^2 V^{\tau+\lambda-\mu} dv_g \\
&\geq \int_M b(x)\phi^2 V^{\tau+\mu-\lambda} dv_g + \int_M a(x)\omega^2 \left| \nabla \left(\frac{\phi}{\omega} \right) \right|^2 V^{\tau+\mu-\lambda} dv_g \\
&= \int_M b(x)\phi^2 V^{\tau+\mu-\lambda} dv_g + \int_M a(x)\omega^2 \left| \nabla^D \left(\frac{\phi}{\omega} \right) \right|^2 V^{\tau+\lambda-\mu} dv_g.
\end{aligned} \tag{3.46}$$

Reorganizing yields the desired equality (3.44). The proof of Theorem 3.3 is completed. \square

We now apply Theorem 3.3 to recover previously known weighted Hardy type inequalities. For instance, by considering the weight function $a(x) = \rho^\alpha$ and $\omega(x) = \rho^{-\left(\frac{c+\alpha-1}{2}\right)}$ in (3.44), we obtain the following weighted Hardy type inequality:

Corollary 3.4. *Let $(M^n, g, V = e^\mu)$ be an n -dimensional Riemannian triple and $\lambda, \mu \in \mathbb{R}$. Let $D = D^{\lambda, \mu}$ be the affine connection defined as in (2.1) and $\tau = (n+1)\lambda + \mu$. Let ρ be a nonnegative function on M such that $|\nabla^D \rho| = 1$ and $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda - \mu}$ in the sense of distribution, where $C > 1$, $C + \alpha - 1 > 0$ and $\alpha \in \mathbb{R}$. Then the following inequality*

$$\int_M \rho^\alpha |\nabla^D \phi|^2 V^{\tau + \lambda - \mu} dv_g \geq \left(\frac{C + \alpha - 1}{2} \right)^2 \int_M \rho^{\alpha - 2} \phi^2 V^{\tau + \lambda - \mu} dv_g \quad (3.47)$$

holds for any $\phi \in C_0^\infty(M \setminus \rho^{-1}\{0\})$.

Proof. By a straightforward computation, we obtain

$$\begin{aligned} & -V^\tau D_i (a(x) \nabla_i^D \omega) \\ &= -\operatorname{div} \left(V^{\tau + \mu - \lambda} a(x) \nabla \omega \right) \\ &= -\operatorname{div} \left(V^{\tau + \mu - \lambda} \rho^\alpha \nabla \rho^{-\left(\frac{C + \alpha - 1}{2}\right)} \right) \\ &= -\operatorname{div} \left(V^{\tau + \mu - \lambda} \rho^\alpha \left(-\frac{C + \alpha - 1}{2} \rho^{-\left(\frac{C + \alpha - 1}{2}\right)} \nabla \rho \right) \right) \\ &= \frac{C + \alpha - 1}{2} \operatorname{div} \left(V^{\tau + \mu - \lambda} \rho^{\alpha - \frac{C + \alpha - 1}{2}} \nabla \rho \right) \\ &= \frac{C + \alpha - 1}{2} \left((\tau + \mu - \lambda) V^{\tau + \mu - \lambda} \rho^{\alpha - \frac{C + \alpha - 1}{2}} \langle \nabla u, \nabla \rho \rangle + V^{\tau + \mu - \lambda} \rho^{\alpha - \frac{C + \alpha - 1}{2}} \Delta \rho \right) \\ &\quad + \frac{C + \alpha - 1}{2} \left(\alpha - \frac{C + \alpha - 1}{2} \right) V^{\tau + \mu - \lambda} \rho^{\alpha - \frac{C + \alpha - 1}{2} - 1} |\nabla \rho|^2 \\ &= \frac{C + \alpha - 1}{2} \left(V^\tau \rho^{\alpha - \frac{C + \alpha - 1}{2}} \Delta^D \rho + \left(\alpha - \frac{C + \alpha - 1}{2} \right) V^{\tau + \lambda - \mu} \rho^{\alpha - \frac{C + \alpha - 1}{2} - 1} \right) \\ &\geq \frac{C + \alpha - 1}{2} \left(\alpha - \frac{C + \alpha - 1}{2} + C \right) V^{\tau + \lambda - \mu} \rho^{\alpha - \frac{C + \alpha - 1}{2} - 1} \\ &= \left(\frac{C + \alpha - 1}{2} \right)^2 V^{\tau + \lambda - \mu} \rho^{\alpha - 2} \rho^{-\left(\frac{C + \alpha - 1}{2}\right)} \\ &= \left(\frac{C + \alpha - 1}{2} \right)^2 \rho^{\alpha - 2} V^{2(\lambda - \mu)} \omega V^{\tau + \mu - \lambda}. \end{aligned} \quad (3.48)$$

Since (3.43), we get

$$b(x) = \left(\frac{C + \alpha - 1}{2} \right)^2 \rho^{\alpha - 2} V^{2(\lambda - \mu)}. \quad (3.49)$$

By (3.44), we have

$$\int_M a(x) |\nabla^D \phi|^2 V^{\tau + \lambda - \mu} dv_g \geq \int_M b(x) \phi^2 V^{\tau + \mu - \lambda} dv_g. \quad (3.50)$$

Inserting (3.49) into (3.50), we obtain

$$\int_M \rho^\alpha |\nabla^D \phi|^2 V^{\tau + \lambda - \mu} dv_g \geq \left(\frac{C + \alpha - 1}{2} \right)^2 \int_M \rho^{\alpha - 2} \phi^2 V^{\tau + \lambda - \mu} dv_g.$$

This completes the proof of Corollary 3.4. \square

Remark 3.5. Particularly, when $\alpha = \gamma = 0$, then our Corollary 3.4 with respect to the affine connection $D^{\lambda, \mu}$ reduces to the Theorem 1.4 of Carron with respect to a Levi-Civita connection of g in [1].

We now choose the weight function $a(x) = \rho^\alpha$ and $\omega(x) = (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)}$ in (3.44). Then we obtain the following weighted Hardy type inequality:

Corollary 3.6. *Under the same conditions of Corollary 3.4, let ρ be a nonnegative function on M such that $|\nabla^D \rho| = 1$ and $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$ in the sense of distribution, where $C > 1$, $C + \alpha - 1 > 0$ and $\alpha \in \mathbb{R}$. Then the following inequality*

$$\int_M \rho^\alpha |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \geq (C + \alpha - 1)(C + \alpha + 1) \int_M \rho^\alpha \frac{|\phi|^2}{(1 + \rho^2)^2} V^{\tau+\lambda-\mu} dv_g \quad (3.51)$$

holds for any $\phi \in C_0^\infty(M \setminus \rho^{-1}\{0\})$.

Proof. By a straightforward computation, we obtain

$$\begin{aligned} & -V^\tau D_i (a(x) \nabla_i^D \omega) \\ = & -\operatorname{div} \left(V^{\tau+\mu-\lambda} a(x) \nabla \omega \right) \\ = & -\operatorname{div} \left(V^{\tau+\mu-\lambda} \rho^\alpha \nabla (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)} \right) \\ = & -\operatorname{div} \left(V^{\tau+\mu-\lambda} \rho^\alpha \left(-\frac{C + \alpha - 1}{2} (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)-1} 2\rho \nabla \rho \right) \right) \\ = & (C + \alpha - 1) \operatorname{div} \left(V^{\tau+\mu-\lambda} \rho^{\alpha+1} (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)-1} \nabla \rho \right) \\ = & (C + \alpha - 1) \left(V^\tau \rho^{\alpha+1} (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)-1} \Delta^D \rho + (\alpha + 1) V^{\tau+\mu-\lambda} \rho^\alpha |\nabla \rho|^2 (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)-1} \right) \\ & - (C + \alpha - 1)(C + \alpha + 1) V^{\tau+\mu-\lambda} \rho^{\alpha+2} |\nabla \rho|^2 (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)-2} \\ = & (C + \alpha - 1) \left(V^\tau \rho^{\alpha+1} (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)-1} \Delta^D \rho + (\alpha + 1) V^{\tau+\lambda-\mu} \rho^\alpha |\nabla^D \rho|^2 (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)-1} \right) \\ & - (C + \alpha - 1)(C + \alpha + 1) V^{\tau+\lambda-\mu} \rho^{\alpha+2} |\nabla^D \rho|^2 (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)-2}. \end{aligned} \quad (3.52)$$

Since $|\nabla^D \rho| = 1$ and $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$, we get

$$\begin{aligned}
& -V^\tau D_i (a(x) \nabla_i^D \omega) \\
& \geq (C + \alpha - 1) \left(C V^{\tau+\lambda-\mu} \rho^\alpha (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)-1} + (\alpha + 1) V^{\tau+\lambda-\mu} \rho^\alpha (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)-1} \right) \\
& \quad - (C + \alpha - 1)(C + \alpha + 1) V^{\tau+\lambda-\mu} \rho^{\alpha+2} (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)-2} \\
& = (C + \alpha - 1)(C + \alpha + 1) V^{\tau+\lambda-\mu} \rho^\alpha (1 + \rho^2)^{-\left(\frac{C+\alpha+1}{2}\right)} \\
& \quad - (C + \alpha - 1)(C + \alpha + 1) V^{\tau+\lambda-\mu} \rho^{\alpha+2} (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)-2} \\
& = (C + \alpha - 1)(C + \alpha + 1) V^{\tau+\lambda-\mu} \rho^\alpha (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)-1} \\
& \quad - (C + \alpha - 1)(C + \alpha + 1) V^{\tau+\lambda-\mu} \rho^\alpha (\rho^2 + 1 - 1) (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)-2} \\
& = (C + \alpha - 1)(C + \alpha + 1) V^{\tau+\lambda-\mu} \rho^\alpha (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)-2} \\
& = (C + \alpha - 1)(C + \alpha + 1) V^{\tau+\lambda-\mu} \rho^\alpha (1 + \rho^2)^{-2} (1 + \rho^2)^{-\left(\frac{C+\alpha-1}{2}\right)} \\
& = (C + \alpha - 1)(C + \alpha + 1) V^{\tau+\lambda-\mu} \rho^\alpha (1 + \rho^2)^{-2} \omega.
\end{aligned} \tag{3.53}$$

Since (3.43), we get

$$b(x) = (C + \alpha - 1)(C + \alpha + 1) \rho^\alpha (1 + \rho^2)^{-2} V^{2(\alpha-\gamma)}. \tag{3.54}$$

Inserting (3.54) into (3.50), we obtain

$$\int_M \rho^\alpha |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \geq (C + \alpha - 1)(C + \alpha + 1) \int_M \rho^\alpha \frac{|\phi|^2}{(1 + \rho^2)^2} V^{\tau+\lambda-\mu} dv_g.$$

This completes the proof of Corollary 3.6. \square

Remark 3.7. In particular, if $\alpha = \gamma = 0$ and $p = 2$, then our Corollary 3.6 with respect to the affine connection $D^{\lambda, \mu}$ reduces to the Corollary 2.3 with respect to a Levi-Civita connection of g in [15].

The following Hardy type inequality in the Euclidean setting has been proved by Ghoussoub and Moradifam [5]:

$$\int_{\mathbb{R}^n} \frac{(c + d|x|^\alpha)^\beta}{|x|^{2m}} |\nabla \phi|^2 dx \geq \left(\frac{n - 2m - 2}{2} \right)^2 \int_{\mathbb{R}^n} \frac{(c + d|x|^\alpha)^\beta}{|x|^{2m+2}} \phi^2 dx,$$

where $\phi \in C_0^\infty(\mathbb{R}^n)$, $c, d > 0$, $\alpha\beta > 0$ and $m \leq \frac{n-2}{2}$.

We now choose the pair as

$$a(x) = \frac{(c + d\rho^\alpha)^\beta}{\rho^{2m}} \quad \text{and} \quad \omega(x) = \rho^{-\left(\frac{C-2m-1}{2}\right)}.$$

Then we have the following Hardy type inequality:

Corollary 3.8. *Under the same conditions of Corollary 3.4, let ρ be a nonnegative function on M such that $|\nabla^D \rho| = 1$ and $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$ in the sense of distribution, where $C > 1$ and $C - 2m - 1 \geq 0$. Then the following inequality*

$$\int_M \frac{(c + d\rho^\alpha)^\beta}{\rho^{2m}} |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \geq \left(\frac{C - 2m - 1}{2} \right)^2 \int_M \frac{(c + d\rho^\alpha)^\beta}{\rho^{2m+2}} \phi^2 V^{\tau+\lambda-\mu} dv_g \quad (3.55)$$

holds for any $\phi \in C_0^\infty(M \setminus \rho^{-1}\{0\})$. Here $c, d > 0, \alpha\beta > 0$.

Proof. By a straightforward computation, we obtain

$$\begin{aligned} & -V^\tau D_i (a(x) \nabla_i^D \omega) \\ &= -\operatorname{div} \left(V^{\tau+\mu-\lambda} a(x) \nabla \omega \right) \\ &= -\operatorname{div} \left(V^{\tau+\mu-\lambda} \frac{(c + d\rho^\alpha)^\beta}{\rho^{2m}} \nabla \rho^{-\frac{C-2m-1}{2}} \right) \\ &= \operatorname{div} \left(V^{\tau+\mu-\lambda} \frac{(c + d\rho^\alpha)^\beta}{\rho^{2m}} \frac{C - 2m - 1}{2} \rho^{-\left(\frac{C-2m-1}{2}\right)-1} \nabla \rho \right) \\ &= \frac{C - 2m - 1}{2} (\tau + \mu - \lambda) V^{\tau+\mu-\lambda} \frac{(c + d\rho^\alpha)^\beta}{\rho^{2m}} \rho^{-\left(\frac{C-2m-1}{2}\right)-1} \langle \nabla u, \nabla \rho \rangle \\ & \quad + \frac{C - 2m - 1}{2} \frac{\beta (c + d\rho^\alpha)^{\beta-1}}{\rho^{2m}} \alpha d \rho^{\alpha-1} \rho^{-\left(\frac{C-2m-1}{2}\right)-1} |\nabla \rho|^2 V^{\tau+\mu-\lambda} \\ & \quad - \frac{2m(C - 2m - 1)}{2} \frac{(c + d\rho^\alpha)^\beta}{\rho^{2m+1}} |\nabla \rho|^2 \rho^{-\left(\frac{C-2m-1}{2}\right)-1} V^{\tau+\mu-\lambda} \\ & \quad + \frac{C - 2m - 1}{2} \frac{(c + d\rho^\alpha)^\beta}{\rho^{2m}} \left(-\frac{C - 2m + 1}{2} \right) \rho^{-\left(\frac{C-2m-1}{2}\right)-2} |\nabla \rho|^2 V^{\tau+\mu-\lambda} \\ & \quad + \frac{C - 2m - 1}{2} \frac{(c + d\rho^\alpha)^\beta}{\rho^{2m}} \rho^{-\left(\frac{C-2m-1}{2}\right)-1} V^{\tau+\mu-\lambda} \Delta \rho \\ &= \frac{C - 2m - 1}{2} \frac{\beta (c + d\rho^\alpha)^{\beta-1}}{\rho^{2m}} \alpha d \rho^{\alpha-1} \rho^{-\left(\frac{C-2m-1}{2}\right)-1} |\nabla^D \rho|^2 V^{\tau+\lambda-\mu} \\ & \quad - \frac{2m(C - 2m - 1)}{2} \frac{(c + d\rho^\alpha)^\beta}{\rho^{2m+1}} |\nabla^D \rho|^2 \rho^{-\left(\frac{C-2m-1}{2}\right)-1} V^{\tau+\lambda-\mu} \\ & \quad + \frac{C - 2m - 1}{2} \frac{(c + d\rho^\alpha)^\beta}{\rho^{2m}} \left(-\frac{C - 2m + 1}{2} \right) \rho^{-\left(\frac{C-2m-1}{2}\right)-2} |\nabla^D \rho|^2 V^{\tau+\lambda-\mu} \\ & \quad + \frac{C - 2m - 1}{2} \frac{(c + d\rho^\alpha)^\beta}{\rho^{2m}} \rho^{-\left(\frac{C-2m-1}{2}\right)-1} V^\tau \Delta^D \rho. \end{aligned} \quad (3.56)$$

Since $|\nabla^D \rho| = 1$ and $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$, we have

$$\begin{aligned} & -V^\tau D_i (a(x) \nabla_i^D \omega) \\ & \geq \left(\frac{C - 2m - 1}{2} \right)^2 \frac{(c + d\rho^\alpha)^\beta}{\rho^{2m+2}} \rho^{-\left(\frac{C-2m-1}{2}\right)} V^{\tau+\lambda-\mu}. \end{aligned} \quad (3.57)$$

Since (3.43), we get

$$b(x) = \left(\frac{C-2m-1}{2} \right)^2 \frac{(c+d\rho^\alpha)^\beta}{\rho^{2m+2}} V^{2(\alpha-\gamma)}. \quad (3.58)$$

Inserting (3.58) into (3.50), we obtain

$$\int_M \frac{(c+d\rho^\alpha)^\beta}{\rho^{2m}} |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \geq \left(\frac{C-2m-1}{2} \right)^2 \int_M \frac{(c+d\rho^\alpha)^\beta}{\rho^{2m+2}} \phi^2 V^{\tau+\lambda-\mu} dv_g.$$

This completes the proof of Corollary 3.8. \square

Remark 3.9. In particular, if $\alpha = \gamma = 0$, then our Corollary 3.8 with respect to the affine connection $D^{\lambda,\mu}$ reduces to the Corollary 2.4 with respect to a Levi-Civita connection of g in [15].

Another application of Theorem 3.3 with the special functions $a(x) = (1+\rho^2)^\alpha$ and $\omega(x) = (1+\rho^2)^{1-\alpha}$ leads to the following weighted Hardy type inequality:

Corollary 3.10. *Under the same conditions of Corollary 3.4, let ρ be a nonnegative function on M such that $|\nabla^D \rho| = 1$ and $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$ in the sense of distribution, where $C > 1$. Then, for any $\phi \in C_0^\infty(M \setminus \rho^{-1}\{0\})$ and $\alpha > 1$, we have*

$$\int_M \frac{|\nabla^D \phi|^2}{(1+\rho^2)^{-\alpha}} V^{\tau+\lambda-\mu} dv_g \geq 2(\alpha-1)(C+1) \int_M \frac{|\phi|^2}{(1+\rho^2)^{1-\alpha}} V^{\tau+\lambda-\mu} dv_g. \quad (3.59)$$

Proof. By a straightforward computation, we obtain

$$\begin{aligned} & -V^\tau D(a(x)\nabla^D \omega) \\ &= -\operatorname{div} \left(V^{\tau+\mu-\lambda} a(x) \nabla \omega \right) \\ &= -\operatorname{div} \left(V^{\tau+\mu-\lambda} (1+\rho^2)^\alpha \nabla (1+\rho^2)^{1-\alpha} \right) \\ &= -\operatorname{div} \left(2V^{\tau+\mu-\lambda} (1+\rho^2)^\alpha (1-\alpha) (1+\rho^2)^{-\alpha} \rho \nabla \rho \right) \\ &= 2(\alpha-1) \operatorname{div} \left(V^{\tau+\mu-\lambda} \rho \nabla \rho \right) \\ &= 2(\alpha-1) \left((\tau+\mu-\lambda) V^{\tau+\mu-\lambda} \rho \langle \nabla u, \nabla \rho \rangle + V^{\tau+\mu-\lambda} |\nabla \rho|^2 + V^{\tau+\mu-\lambda} \rho \Delta \rho \right) \\ &= 2(\alpha-1) \left(\rho V^\tau \Delta^D \rho + V^{\tau+\lambda-\mu} \right) \\ &\geq 2(\alpha-1)(C+1) V^{\tau+\lambda-\mu} \\ &= 2(\alpha-1)(C+1) (1+\rho^2)^{1-\alpha} (1+\rho^2)^{\alpha-1} V^{\tau+\lambda-\mu}. \end{aligned} \quad (3.60)$$

Since (3.43), we get

$$b(x) = 2(\alpha-1)(C+1) (1+\rho^2)^{\alpha-1} V^{2(\lambda-\mu)}. \quad (3.61)$$

Inserting (3.61) into (3.50), we obtain

$$\int_M \frac{|\nabla^D \phi|^2}{(1+\rho^2)^{-\alpha}} V^{\tau+\lambda-\mu} dv_g \geq 2(\alpha-1)(C+1) \int_M \frac{|\phi|^2}{(1+\rho^2)^{1-\alpha}} V^{\tau+\lambda-\mu} dv_g.$$

This completes the proof of Corollary 3.10. \square

4. RELlich-TYPE INEQUALITIES AND THEIR PROOFS

Some Rellich type inequalities with respect to $D^{\lambda,\mu}$ on Riemannian manifolds will be derived in this section. We first prove the following weighted Rellich inequality with respect to $D^{\lambda,\mu}$ on M .

Theorem 4.1. (Weighted Rellich inequality). *Let $(M^n, g, V = e^u)$ be an n -dimensional Riemannian triple and $\lambda, \mu \in \mathbb{R}$. Let $D = D^{\lambda,\mu}$ be the affine connection defined as in (2.1) and $\tau = (n+1)\lambda + \mu$. Let ρ be a nonnegative functions on M such that $|\nabla^D \rho| = 1$ and $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$ in the sense of distribution where $C > 1$, $\alpha < 2$ and $C + \alpha - 3 > 0$. Then the following inequality holds:*

$$\int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g \geq \frac{(C + \alpha - 3)^2 (C - \alpha + 1)^2}{16} \int_M \rho^{\alpha-4} \phi^2 V^{\tau+\lambda-\mu} dv_g \quad (4.1)$$

for any $\phi \in C_0^\infty(M \setminus \rho^{-1}\{0\})$.

Proof. Since $|\nabla^D \rho| = 1$ and $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$, by direct computation, we have

$$\begin{aligned} V^\tau \Delta^D \rho^{\alpha-2} &= V^\tau V^{\mu-\lambda} (\Delta \rho^{\alpha-2} + (2\mu + n\lambda) \langle \nabla u, \nabla \rho^{\alpha-2} \rangle) \\ &= V^{\tau+\mu-\lambda} [\operatorname{div}((\alpha-2)\rho^{\alpha-3} \nabla \rho) + (2\mu + n\lambda) \langle \nabla u, (\alpha-2)\rho^{\alpha-3} \nabla \rho \rangle] \\ &= V^{\tau+\mu-\lambda} [(\alpha-2)(\alpha-3)\rho^{\alpha-4} |\nabla \rho|^2 + (\alpha-2)\rho^{\alpha-3} \Delta \rho + (2\mu + n\lambda)(\alpha-2)\rho^{\alpha-3} \langle \nabla u, \nabla \rho \rangle] \\ &= V^{\tau+\lambda-\mu} [(\alpha-2)(\alpha-3)\rho^{\alpha-4} |\nabla^D \rho|^2 + (\alpha-2)\rho^{\alpha-3} \Delta^D \rho] \\ &= (\alpha-2) V^\tau [(\alpha-3)\rho^{\alpha-4} V^{\lambda-\mu} + \rho^{\alpha-3} \Delta^D \rho] \\ &\leq (\alpha-2)(\alpha-3+C)\rho^{\alpha-4} V^{\tau+\lambda-\mu}. \end{aligned}$$

Multiplying both sides of above inequality by ϕ^2 and integrating over M , we can obtain

$$(\alpha-2)(\alpha-3+C) \int_M \rho^{\alpha-4} \phi^2 V^{\tau+\lambda-\mu} dv_g \geq \int_M \phi^2 V^\tau \Delta^D \rho^{\alpha-2} dv_g. \quad (4.2)$$

Using the integration by parts again, we can get

$$\begin{aligned} &\int_M \phi^2 V^\tau \Delta^D \rho^{\alpha-2} dv_g \\ &= - \int_M \langle \nabla \phi^2, \nabla^D \rho^{\alpha-2} \rangle V^\tau dv_g \\ &= \int_M \rho^{\alpha-2} V^\tau \Delta^D \phi^2 dv_g \\ &= \int_M V^{\mu-\lambda} \rho^{\alpha-2} (\Delta \phi^2 + (2\mu + n\lambda) \langle \nabla u, \nabla \phi^2 \rangle) V^\tau dv_g \\ &= \int_M V^{\mu-\lambda} \rho^{\alpha-2} [2(\phi \Delta \phi + |\nabla \phi|^2) + (2\mu + n\lambda) 2\phi \langle \nabla u, \nabla \phi \rangle] V^\tau dv_g \\ &= 2 \int_M \rho^{\alpha-2} (\phi \Delta^D \phi + |\nabla^D \phi|^2 V^{\lambda-\mu}) V^\tau dv_g. \end{aligned} \quad (4.3)$$

Taking (4.3) into (4.2), we obtain

$$\begin{aligned} - \int_M \rho^{\alpha-2} (\phi \Delta^D \phi) V^\tau dv_g &\geq \int_M \rho^{\alpha-2} |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \\ &\quad - \frac{(\alpha-2)(\alpha-3+C)}{2} \int_M \rho^{\alpha-4} \phi^2 V^{\tau+\lambda-\mu} dv_g. \end{aligned} \quad (4.4)$$

By (3.47) in Corollary 3.4, we have

$$\int_M \rho^{\alpha-2} |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \geq \left(\frac{C+\alpha-3}{2} \right)^2 \int_M \rho^{\alpha-4} \phi^2 V^{\tau+\lambda-\mu} dv_g \quad (4.5)$$

Inserting (4.5) into (4.4), we have

$$\begin{aligned} & - \int_M \rho^{\alpha-2} (\phi \Delta^D \phi) V^\tau dv_g \\ & \geq \frac{(C+\alpha-3)^2}{2} \int_M \rho^{\alpha-4} \phi^2 V^{\tau+\lambda-\mu} dv_g - \frac{(\alpha-2)(\alpha-3+C)}{2} \int_M \rho^{\alpha-4} \phi^2 V^{\tau+\lambda-\mu} dv_g \\ & = \frac{(C+\alpha-3)(C-\alpha+1)}{2} \int_M \rho^{\alpha-4} \phi^2 V^{\tau+\lambda-\mu} dv_g. \end{aligned} \quad (4.6)$$

Using Young's inequality in (4.6), we have

$$\begin{aligned} - \int_M \rho^{\alpha-2} (\phi \Delta^D \phi) V^\tau dv_g &\leq \frac{(C+\alpha-3)(C-\alpha+1)}{4} \int_M \rho^{\alpha-4} \phi^2 V^{\tau+\lambda-\mu} dv_g \\ &\quad + \frac{4}{(C+\alpha-3)(C-\alpha+1)} \int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g. \end{aligned} \quad (4.7)$$

Then, combining (4.6) and (4.7), we know that

$$\begin{aligned} & \frac{4}{(C+\alpha-3)(C-\alpha+1)} \int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g \\ & \geq \frac{(C+\alpha-3)(C-\alpha+1)}{4} \int_M \rho^{\alpha-4} \phi^2 V^{\tau+\lambda-\mu} dv_g. \end{aligned}$$

Thus, we have

$$\int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g \geq \frac{(C+\alpha-3)^2 (C-\alpha+1)^2}{16} \int_M \rho^{\alpha-4} \phi^2 V^{\tau+\lambda-\mu} dv_g. \quad (4.8)$$

□

Then, we would like to give two weighted Rellich type inequalities with respect to $D^{\lambda,\mu}$ on M , which connect the first and the second order derivatives, as follows.

Theorem 4.2. *Let $(M^n, g, V = e^\mu)$ be an n -dimensional Riemannian triple and $\lambda, \mu \in \mathbb{R}$. Let $D = D^{\lambda,\mu}$ be the affine connection defined as in (2.1) and $\tau = (n+1)\lambda + \mu$. Let ρ be a nonnegative function on M such that $|\nabla^D \rho| = 1$ in the sense of distributions. Then for any $\phi \in C_0^\infty(M \setminus \rho^{-1}\{0\})$, we have*

(i) *When $\Delta^D \rho \leq \frac{C}{\rho} V^{\lambda-\mu}$ in the sense of distributions, where $C < 1$ is a constant, for $2 < \alpha < 3 - C$, the following inequality*

$$\int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g \geq 2(\alpha-2)(3-\alpha-C) \int_M \rho^{\alpha-2} |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \quad (4.9)$$

holds.

(ii) When $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$ in the sense of distributions, where $C > 1$ is a constant, for $\frac{7-C}{3} < \alpha < 2$, the following inequality

$$\int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g \geq \frac{(C-\alpha+1)^2}{4} \int_M \rho^{\alpha-2} |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \quad (4.10)$$

holds.

Proof. (i) Since $\alpha - 2 > 0$, $|\nabla^D \rho| = 1$ and $\Delta^D \rho \leq \frac{C}{\rho} V^{\lambda-\mu}$, by direct computation, we have

$$\begin{aligned} \Delta^D (\rho^{\alpha-2}) V^\tau &= V^\tau V^{\mu-\lambda} (\Delta \rho^{\alpha-2} + (2\mu + n\lambda) \langle \nabla u, \nabla \rho^{\alpha-2} \rangle) \\ &= V^{\tau+\mu-\lambda} [\operatorname{div}((\alpha-2)\rho^{\alpha-3} \nabla \rho) + (2\mu + n\lambda) \langle \nabla u, (\alpha-2)\rho^{\alpha-3} \nabla \rho \rangle] \\ &= V^{\tau+\mu-\lambda} [(\alpha-2)(\alpha-3)\rho^{\alpha-4} |\nabla \rho|^2 + (\alpha-2)\rho^{\alpha-3} \Delta \rho + (2\mu + n\lambda)(\alpha-2)\rho^{\alpha-3} \langle \nabla u, \nabla \rho \rangle] \\ &= V^{\tau+\lambda-\mu} [(\alpha-2)(\alpha-3)\rho^{\alpha-4} |\nabla^D \rho|^2 + (\alpha-2)\rho^{\alpha-3} \Delta^D \rho] \\ &= (\alpha-2) V^\tau [(\alpha-3)\rho^{\alpha-4} V^{\lambda-\mu} + \rho^{\alpha-3} \Delta^D \rho] \\ &\leq (\alpha-2)(\alpha-3+C)\rho^{\alpha-4} V^{\tau+\lambda-\mu}. \end{aligned}$$

Multiplying both sides of above inequality by ϕ^2 and integrating over M , we can obtain

$$(\alpha-2)(\alpha-3+C) \int_M \rho^{\alpha-4} \phi^2 V^{\tau+\lambda-\mu} dv_g \geq \int_M \phi^2 \Delta^D (\rho^{\alpha-2}) V^\tau dv_g. \quad (4.11)$$

From similar calculations in Theorem 4.1, we can get the following result:

$$\int_M \phi^2 \Delta^D (\rho^{\alpha-2}) V^\tau dv_g = \int_M 2\rho^{\alpha-2} (2\phi \Delta^D \phi + |\nabla^D \phi|^2 V^{\lambda-\mu}) V^\tau dv_g. \quad (4.12)$$

Taking (4.12) into (4.11), we obtain

$$\begin{aligned} -2 \int_M \rho^{\alpha-2} (\phi \Delta^D \phi) V^\tau dv_g &\geq 2 \int_M \rho^{\alpha-2} |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \\ &\quad - (\alpha-2)(\alpha-3+C) \int_M \rho^{\alpha-4} \phi^2 V^{\tau+\lambda-\mu} dv_g. \end{aligned} \quad (4.13)$$

On the other hand, using Young's inequality, we have

$$\begin{aligned} -2 \int_M \rho^{\alpha-2} (\phi \Delta^D \phi) V^\tau dv_g &\leq (\alpha-2)(3-\alpha-C) \int_M \rho^{\alpha-4} \phi^2 V^{\tau+\lambda-\mu} dv_g \\ &\quad + \frac{1}{(\alpha-2)(3-\alpha-C)} \int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g. \end{aligned} \quad (4.14)$$

Then, combining (4.13) and (4.14), we know that (4.9) is true.

(ii) From similar calculations in Theorem 4.1, we can get the following result:

$$\begin{aligned} - \int_M \rho^{\alpha-2} (\phi \Delta^D \phi) V^\tau dv_g &\geq \int_M \rho^{\alpha-2} |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \\ &\quad - \frac{(\alpha-2)(\alpha-3+C)}{2} \int_M \rho^{\alpha-4} \phi^2 V^{\tau+\lambda-\mu} dv_g, \end{aligned} \quad (4.15)$$

where $\frac{7-C}{3} < \alpha < 2$. Using Young's inequality in (4.15), we have

$$\begin{aligned} - \int_M \rho^{\alpha-2} (\phi \Delta^D \phi) V^\tau dv_g &\leq \varepsilon \int_M \rho^{\alpha-4} \phi^2 V^{\tau+\lambda-\mu} dv_g \\ &\quad + \frac{1}{4\varepsilon} \int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g, \end{aligned} \quad (4.16)$$

where ε is a constant to be determined and $\varepsilon > 0$. Substituting (4.16) into (4.15), we get

$$\begin{aligned} \int_M \rho^{\alpha-2} |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g &\leq B \int_M \rho^{\alpha-4} \phi^2 V^{\tau+\lambda-\mu} dv_g \\ &\quad + \frac{1}{4\varepsilon} \int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g, \end{aligned} \quad (4.17)$$

where $B = \varepsilon + \frac{(\alpha-2)(\alpha-3+C)}{2}$.

According to the sign of B , we need to consider the following three cases:

Case 1:

We choose $B < 0$ and $\varepsilon = \frac{(C-\alpha+1)^2}{16}$, then (4.17) becomes

$$\int_M \rho^{\alpha-2} |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \leq \frac{1}{4\varepsilon} \int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g,$$

which is equivalent to

$$\begin{aligned} \int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g &\geq 4\varepsilon \int_M \rho^{\alpha-2} |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \\ &= \frac{(C-\alpha+1)^2}{4} \int_M \rho^{\alpha-2} |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g. \end{aligned}$$

Case 2:

We choose $B = 0$ and $\varepsilon = -\frac{(C+\alpha-3)(\alpha-2)}{2}$, then (4.17) becomes

$$\begin{aligned} \int_M \rho^{\alpha-2} |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g &\leq \frac{1}{4\varepsilon} \int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g \\ &= \frac{1}{2(2-\alpha)(C+\alpha-3)} \int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g. \end{aligned} \quad (4.18)$$

In (4.18), since $C > 1$ and $\frac{7-C}{3} < \alpha < 2$, we can choose C close enough to 1 and α close enough to 2 such that

$$\begin{aligned} \int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g &\geq 2(2-\alpha)(C+\alpha-3) \int_M \rho^{\alpha-2} |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \\ &\geq \frac{(C-\alpha+1)^2}{4} \int_M \rho^{\alpha-2} |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g. \end{aligned}$$

Case 3:

We choose $B > 0$, using the weighted Rellich inequality (4.1), then (4.17) becomes

$$\begin{aligned} & \int_M \rho^{\alpha-2} |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \\ & \leq \left(\varepsilon + \frac{(C+\alpha-3)(\alpha-2)}{2} \right) \left(\frac{16}{(C+\alpha-3)^2(C-\alpha+1)^2} \right) \int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g \\ & \quad + \frac{1}{4\varepsilon} \int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g \\ & = P_{C,\alpha}(\varepsilon) \int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g, \end{aligned}$$

where $P_{C,\alpha}(\varepsilon) = \frac{16\varepsilon}{(C+\alpha-3)^2(C-\alpha+1)^2} + \frac{8(\alpha-2)}{(C+\alpha-3)(C-\alpha+1)^2} + \frac{1}{4\varepsilon}$. Note that the function $P_{C,\alpha}(\varepsilon)$ attains the minimum for $\varepsilon = \frac{(C+\alpha-3)(C-\alpha+1)}{8}$, and this minimum is equal to $\frac{4}{(C-\alpha+1)^2}$. Therefore we have the following inequality:

$$\int_M \rho^\alpha (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g \geq \frac{(C-\alpha+1)^2}{4} \int_M \rho^{\alpha-2} |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g.$$

This completes the proof of Theorem 4.2. \square

5. SEVERAL INEQUALITIES OF OTHER TYPES

In the last section, by applying some results we have derived in the above two sections, we can prove some inequalities of other types with respect to $D^{\lambda,\mu}$ on M . First, we can give the following Hardy-Poincaré type inequalities with respect to $D^{\lambda,\mu}$.

Theorem 5.1. *Let $(M^n, g, V = e^\mu)$ be an n -dimensional Riemannian triple and $\lambda, \mu \in \mathbb{R}$. Let $D = D^{\lambda,\mu}$ be the affine connection defined as in (2.1) and $\tau = (n+1)\lambda + \mu$. Let ρ be a nonnegative function on M such that $|\nabla^D \rho| = 1$ in the sense of distributions. Then for any $\phi \in C_0^\infty(M \setminus \rho^{-1}\{0\})$, we have*

(i) *When $\Delta^D \rho \leq \frac{C}{\rho} V^{\lambda-\mu}$ in the sense of distributions, where $C > 0$ is a constant. Let $1 < p < +\infty$ and $C + \alpha + 1 < 0$, then the following inequality*

$$\int_M \rho^{\alpha+p} |\langle \nabla^D \rho, \nabla^D \phi \rangle|^p V^{\tau+(1-2p)(\mu-\lambda)} dv_g \geq \left(\frac{|C+\alpha+1|}{p} \right)^p \int_M \rho^\alpha |\phi|^p V^{\tau+\mu-\lambda} dv_g \quad (5.1)$$

holds.

(ii) *When $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$ in the sense of distributions, where $C > 0$ is a constant. Let $1 < p < +\infty$ and $C + \alpha + 1 > 0$, then the following inequality*

$$\int_M \rho^{\alpha+p} |\langle \nabla^D \rho, \nabla^D \phi \rangle|^p V^{\tau+(1-2p)(\mu-\lambda)} dv_g \geq \left(\frac{C+\alpha+1}{p} \right)^p \int_M \rho^\alpha |\phi|^p V^{\tau+\mu-\lambda} dv_g \quad (5.2)$$

holds.

Proof. (i) It follows from $|\nabla^D \rho| = 1$ and $\Delta^D \rho \leq \frac{C}{\rho} V^{\lambda-\mu}$ that

$$\begin{aligned}
& V^\tau \operatorname{div}^D (\rho \nabla^D \rho) \\
&= \operatorname{div} (V^\tau \rho \nabla^D \rho) \\
&= \operatorname{div} (V^{\tau+\mu-\lambda} \rho \nabla \rho) \\
&= V^{\tau+\mu-\lambda} |\nabla \rho|^2 + \rho (\tau + \mu - \lambda) V^{\tau+\mu-\lambda} \langle \nabla u, \nabla \rho \rangle + \rho V^{\tau+\mu-\lambda} \Delta \rho \\
&= V^{\tau+\lambda-\mu} |\nabla^D \rho|^2 + \rho V^\tau \Delta^D \rho \\
&\leq (1+C) V^{\tau+\lambda-\mu}.
\end{aligned} \tag{5.3}$$

Multiplying (5.3) by $\rho^\alpha |\phi|^p$ and integrating over M yields

$$\begin{aligned}
& (1+C) \int_M \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \\
&\geq \int_M V^\tau \operatorname{div}^D (\rho \nabla^D \rho) \rho^\alpha |\phi|^p dv_g \\
&= \int_M \operatorname{div} (\rho V^{\tau+\mu-\lambda} \nabla \rho) \rho^\alpha |\phi|^p dv_g \\
&= - \int_M V^{\tau+\mu-\lambda} \langle \rho \nabla \rho, \nabla (\rho^\alpha |\phi|^p) \rangle dv_g \\
&= - \int_M V^{\tau+\mu-\lambda} \langle \rho \nabla \rho, \alpha \rho^{\alpha-1} \nabla \rho |\phi|^p + p |\phi|^{p-2} \phi \rho^\alpha \nabla \phi \rangle dv_g \\
&= - \alpha \int_M V^{\tau+\mu-\lambda} \rho^\alpha |\nabla \rho|^2 |\phi|^p dv_g - p \int_M \rho^{\alpha+1} |\phi|^{p-2} \phi \langle \nabla \rho, \nabla \phi \rangle V^{\tau+\mu-\lambda} dv_g \\
&= - \alpha \int_M V^{\tau+\lambda-\mu} \rho^\alpha |\phi|^p dv_g - p \int_M \rho^{\alpha+1} |\phi|^{p-2} \phi \langle \nabla \rho, \nabla \phi \rangle V^{\tau+\mu-\lambda} dv_g.
\end{aligned} \tag{5.4}$$

Since $C + \alpha + 1 < 0$, we can infer from (5.4) that

$$(C + \alpha + 1) \int_M \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \geq -p \int_M \rho^{\alpha+1} |\phi|^{p-2} \phi \langle \nabla \rho, \nabla \phi \rangle V^{\tau+\mu-\lambda} dv_g,$$

which is equivalent to

$$-(C + \alpha + 1) \int_M \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \leq p \int_M \rho^{\alpha+1} |\phi|^{p-2} \phi \langle \nabla \rho, \nabla \phi \rangle V^{\tau+\mu-\lambda} dv_g. \tag{5.5}$$

By (5.5), we can get

$$\begin{aligned}
& |C + \alpha + 1| \int_M \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \\
&\leq p \left| \int_M \rho^{\alpha+1} |\phi|^{p-2} \phi \langle \nabla \rho, \nabla \phi \rangle V^{\tau+\mu-\lambda} dv_g \right| \\
&\leq p \int_M \rho^{\alpha+1} |\phi|^{p-2} |\phi| |\langle \nabla \rho, \nabla \phi \rangle| V^{\tau+\mu-\lambda} dv_g \\
&= p \int_M \rho^{\alpha+1} |\phi|^{p-1} |\langle \nabla \rho, \nabla \phi \rangle| V^{\tau+\mu-\lambda} dv_g.
\end{aligned} \tag{5.6}$$

Applying Hölder's inequality shows that

$$\begin{aligned}
& p \int_M \rho^{\alpha+1} |\phi|^{p-1} |\langle \nabla \rho, \nabla \phi \rangle| V^{\tau+\mu-\lambda} dv_g \\
& \leq p \left[\int_M \left(\rho^{\frac{\alpha(p-1)}{p}} |\phi|^{p-1} V^{\frac{p-1}{p}(\tau+\lambda-\mu)} \right)^{\frac{p}{p-1}} dv_g \right]^{\frac{p-1}{p}} \left[\int_M \left(\rho^{\frac{\alpha+p}{p}} |\langle \nabla \rho, \nabla \phi \rangle| V^{\frac{1}{p}\tau + \left(\frac{1}{p}-2\right)(\lambda-\mu)} \right)^p dv_g \right]^{\frac{1}{p}} \\
& = p \left(\int_M \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{p-1}{p}} \left(\int_M \rho^{\alpha+p} |\langle \nabla \rho, \nabla \phi \rangle|^p V^{\tau+(1-2p)(\lambda-\mu)} dv_g \right)^{\frac{1}{p}}.
\end{aligned} \tag{5.7}$$

By Young's inequality, we have

$$\begin{aligned}
& \left(\int_M \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{p-1}{p}} \left(\int_M \rho^{\alpha+p} |\langle \nabla \rho, \nabla \phi \rangle|^p V^{\tau+(1-2p)(\lambda-\mu)} dv_g \right)^{\frac{1}{p}} \\
& \leq \frac{p-1}{p} \varepsilon^{-\frac{p}{p-1}} \int_M \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g + \frac{\varepsilon^p}{p} \int_M \rho^{\alpha+p} |\langle \nabla \rho, \nabla \phi \rangle|^p V^{\tau+(1-2p)(\lambda-\mu)} dv_g
\end{aligned} \tag{5.8}$$

for any $\varepsilon > 0$. Inserting (5.7) and (5.8) into (5.6), we obtain

$$\begin{aligned}
|C + \alpha + 1| \int_M \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g & \leq (p-1) \varepsilon^{-\frac{p}{p-1}} \int_M \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \\
& \quad + \varepsilon^p \int_M \rho^{\alpha+p} |\langle \nabla \rho, \nabla \phi \rangle|^p V^{\tau+(1-2p)(\lambda-\mu)} dv_g.
\end{aligned} \tag{5.9}$$

Thus, for any $\varepsilon > 0$, from (5.9), we have

$$\begin{aligned}
& \int_M \rho^{\alpha+p} |\langle \nabla^D \rho, \nabla^D \phi \rangle|^p V^{\tau+(1-2p)(\mu-\lambda)} dv_g \\
& \geq \varepsilon^{-p} \left(|C + \alpha + 1| - (p-1) \varepsilon^{-\frac{p}{p-1}} \right) \int_M \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g.
\end{aligned}$$

Taking

$$\varepsilon = \left(\frac{p}{|C + \alpha + 1|} \right)^{\frac{p-1}{p}}$$

in the above inequality, we can get (5.1) directly.

(ii) It follows from $|\nabla^D \rho| = 1$ and $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$ that

$$\begin{aligned}
& V^\tau \operatorname{div}^D (\rho \nabla^D \rho) \\
& = \operatorname{div} (V^\tau \rho \nabla^D \rho) \\
& = \operatorname{div} (V^{\tau+\mu-\lambda} \rho \nabla \rho) \\
& = V^{\tau+\mu-\lambda} |\nabla \rho|^2 + \rho (\tau + \mu - \lambda) V^{\tau+\mu-\lambda} \langle \nabla u, \nabla \rho \rangle + \rho V^{\tau+\mu-\lambda} \Delta \rho \\
& = V^{\tau+\lambda-\mu} |\nabla^D \rho|^2 + \rho V^\tau \Delta^D \rho \\
& \geq (1 + C) V^{\tau+\lambda-\mu}.
\end{aligned} \tag{5.10}$$

Multiplying (5.10) by $\rho^\alpha |\phi|^p$ and integrating over M yields

$$\begin{aligned}
& (1+C) \int_M \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \\
& \leq \int_M V^\tau \operatorname{div}^D(\rho \nabla^D \rho) \rho^\alpha |\phi|^p dv_g \\
& = \int_M \nabla(\rho V^{\tau+\mu-\lambda} \nabla \rho) \rho^\alpha |\phi|^p dv_g \\
& = - \int_M V^{\tau+\mu-\lambda} \langle \rho \nabla \rho, \nabla(\rho^\alpha |\phi|^p) \rangle dv_g \tag{5.11} \\
& = - \int_M V^{\tau+\mu-\lambda} \langle \rho \nabla \rho, \alpha \rho^{\alpha-1} \nabla \rho |\phi|^p + p |\phi|^{p-2} \phi \rho^\alpha \nabla \phi \rangle dv_g \\
& = - \alpha \int_M V^{\tau+\mu-\lambda} \rho^\alpha |\nabla \rho|^2 |\phi|^p dv_g - p \int_M \rho^{\alpha+1} |\phi|^{p-2} \phi \langle \nabla \rho, \nabla \phi \rangle V^{\tau+\mu-\lambda} dv_g \\
& = - \alpha \int_M V^{\tau+\lambda-\mu} \rho^\alpha |\phi|^p dv_g - p \int_M \rho^{\alpha+1} |\phi|^{p-2} \phi \langle \nabla \rho, \nabla \phi \rangle V^{\tau+\mu-\lambda} dv_g.
\end{aligned}$$

Since $C + \alpha + 1 > 0$, we can infer from (5.11) that

$$(C + \alpha + 1) \int_M \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \leq -p \int_M \rho^{\alpha+1} |\phi|^{p-2} \phi \langle \nabla \rho, \nabla \phi \rangle V^{\tau+\mu-\lambda} dv_g,$$

which is equivalent to

$$(C + \alpha + 1) \int_M \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \leq p \int_M V^{\tau+\mu-\lambda} \rho^{\alpha+1} |\phi|^{p-2} \phi \langle \nabla \rho, \nabla \phi \rangle dv_g. \tag{5.12}$$

By (5.12), we can get

$$(C + \alpha + 1) \int_M \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \leq p \int_M \rho^{\alpha+1} |\phi|^{p-1} |\langle \nabla \rho, \nabla \phi \rangle| V^{\tau+\mu-\lambda} dv_g. \tag{5.13}$$

By Hölder's and Young's inequalities, we have

$$\begin{aligned}
& p \int_M \rho^{\alpha+1} |\phi|^{p-1} |\langle \nabla \rho, \nabla \phi \rangle| V^{\tau+\mu-\lambda} dv_g \\
& \leq p \left[\int_M \left(\rho^{\frac{\alpha(p-1)}{p}} |\phi|^{p-1} V^{\frac{p-1}{p}(\tau+\lambda-\mu)} \right)^{\frac{p}{p-1}} dv_g \right]^{\frac{p-1}{p}} \left[\int_M \left(\rho^{\frac{\alpha+p}{p}} |\langle \nabla \rho, \nabla \phi \rangle| V^{\frac{1}{p}\tau + (\frac{1}{p}-2)(\lambda-\mu)} \right)^p dv_g \right]^{\frac{1}{p}} \\
& = p \left(\int_M \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{p-1}{p}} \left(\int_M \rho^{\alpha+p} |\langle \nabla \rho, \nabla \phi \rangle|^p V^{\tau+(1-2p)(\lambda-\mu)} dv_g \right)^{\frac{1}{p}} \\
& \leq (p-1) \varepsilon^{-\frac{p}{p-1}} \int_M \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g + \varepsilon^p \int_M \rho^{\alpha+p} |\langle \nabla \rho, \nabla \phi \rangle|^p V^{\tau+(1-2p)(\lambda-\mu)} dv_g. \tag{5.14}
\end{aligned}$$

Thus, for any $\varepsilon > 0$, from (5.13) and (5.14), we have

$$\begin{aligned}
& \int_M \rho^{\alpha+p} |\langle \nabla^D \rho, \nabla^D \phi \rangle|^p V^{\tau+(1-2p)(\lambda-\mu)} dv_g \\
& \geq \varepsilon^{-p} \left((C + \alpha + 1) - (p-1) \varepsilon^{-\frac{p}{p-1}} \right) \int_M \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g.
\end{aligned}$$

Taking

$$\varepsilon = \left(\frac{p}{C + \alpha + 1} \right)^{\frac{p-1}{p}}$$

in the above inequality, we can get (5.2) directly. This completes the proof of Theorem 5.1. \square

We now prove, under an additional assumption on the weight functions $\rho(x)$ and $a(x)$, the following two-weight Hardy-Poincaré inequality with a nonnegative remainder term. Our result recovers and improves the inequality (1.2).

Theorem 5.2. *Let $(M^n, g, V = e^u)$ be an n -dimensional Riemannian triple and $\lambda, \mu \in \mathbb{R}$. Let $D = D^{\lambda, \mu}$ be the affine connection defined as in (2.1) and $\tau = (n+1)\lambda + \mu$. Let ρ be a nonnegative functions on M such that $|\nabla^D \rho| = 1$, $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$ and $\langle \nabla^D \rho, \nabla^D a \rangle \geq 0$ in the sense of distribution, where $C > 1$, $C + \alpha + 1 > 0$, $\alpha \in \mathbb{R}$ and a be a nonnegative functions on M . Then the following inequality*

$$\begin{aligned} \int_M a \rho^{\alpha+p} |\langle \nabla^D \rho, \nabla^D \phi \rangle|^p V^{\tau+\lambda-\mu} dv_g &\geq \left(\frac{C + \alpha + 1}{p} \right)^p \int_M a \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \\ &\quad + \left(\frac{C + \alpha + 1}{p} \right)^{p-1} \int_M \rho^{\alpha+1} |\phi|^p \langle \nabla^D \rho, \nabla^D a \rangle V^{\tau+\lambda-\mu} dv_g \end{aligned} \quad (5.15)$$

holds for any ϕ in $C_0^\infty(M \setminus \rho^{-1}\{0\})$, $1 < p < +\infty$.

Proof. By direct computation, together with assumptions $|\nabla^D \rho| = 1$ and $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$, we have

$$\begin{aligned} V^\tau D_i(\rho \nabla_i^D \rho) &= \nabla_i(V^{\tau+\mu-\lambda} \rho \nabla_i \rho) \\ &= (\tau + \mu - \lambda) V^{\tau+\mu-\lambda} \rho \langle \nabla \rho, \nabla u \rangle + V^{\tau+\mu-\lambda} |\nabla \rho|^2 + V^{\tau+\mu-\lambda} \rho \Delta \rho \\ &= V^\tau \rho \Delta^D \rho + V^{\tau+\lambda-\mu} |\nabla^D \rho|^2 \\ &\geq (C+1) V^{\tau+\lambda-\mu}. \end{aligned} \quad (5.16)$$

Multiplying both sides of (5.16) by $a \rho^\alpha |\phi|^p$ and integrating over M yields

$$\begin{aligned} &(C+1) \int_M a \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \\ &\leq \int_M a \rho^\alpha |\phi|^p V^\tau D_i(\rho \nabla_i^D \rho) dv_g \\ &= \int_M a \rho^\alpha |\phi|^p \nabla_i(\rho V^{\tau+\mu-\lambda} \nabla_i \rho) dv_g \\ &= - \int_M V^{\tau+\mu-\lambda} \langle \rho \nabla \rho, \nabla(a \rho^\alpha |\phi|^p) \rangle dv_g \\ &= - \int_M \rho^{\alpha+1} |\phi|^p V^{\tau+\mu-\lambda} \langle \nabla \rho, \nabla a \rangle dv_g - \alpha \int_M a \rho^\alpha |\nabla \rho|^2 |\phi|^p V^{\tau+\mu-\lambda} dv_g \\ &\quad - p \int_M a \rho^{\alpha+1} |\phi|^{p-2} \phi \langle \nabla \rho, \nabla \phi \rangle V^{\tau+\mu-\lambda} dv_g. \end{aligned}$$

After rearranging the terms, the above inequality can be written in the following form

$$\begin{aligned} A &\leq -p \int_M a \rho^{\alpha+1} |\phi|^{p-2} \phi \langle \nabla \rho, \nabla \phi \rangle V^{\tau+\mu-\lambda} dv_g \\ &\leq p \int_M a \rho^{\alpha+1} |\phi|^{p-1} |\langle \nabla \rho, \nabla \phi \rangle| V^{\tau+\mu-\lambda} dv_g. \end{aligned} \quad (5.17)$$

where

$$A = (C+1+\alpha) \int_M a \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g + \int_M \rho^{\alpha+1} |\phi|^p \langle \nabla \rho, \nabla a \rangle V^{\tau+\mu-\lambda} dv_g.$$

Applying Hölder's inequality on the right hand side of (5.17), we get

$$\begin{aligned} A &\leq p \int_M a^{\frac{p-1}{p}} \rho^{\frac{\alpha(p-1)}{p}} |\phi|^{p-1} a^{\frac{1}{p}} \rho^{\frac{\alpha}{p}+1} |\langle \nabla \rho, \nabla \phi \rangle| V^{\tau+\mu-\lambda} dv_g \\ &\leq p \left[\int_M \left(a^{\frac{p-1}{p}} \rho^{\frac{\alpha(p-1)}{p}} |\phi|^{p-1} V^{\frac{(p-1)(\tau+\lambda-\mu)}{p}} \right)^{\frac{p}{p-1}} dv_g \right]^{\frac{p-1}{p}} \left[\int_M \left(a^{\frac{1}{p}} \rho^{\frac{\alpha}{p}+1} |\langle \nabla \rho, \nabla \phi \rangle| V^{\frac{\tau+(2p-1)(\mu-\lambda)}{p}} \right)^p dv_g \right]^{\frac{1}{p}} \\ &= p \left(\int_M a \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{p-1}{p}} \left(\int_M a \rho^{\alpha+p} |\langle \nabla \rho, \nabla \phi \rangle|^p V^{\tau+(2p-1)(\mu-\lambda)} dv_g \right)^{\frac{1}{p}}. \end{aligned} \quad (5.18)$$

Using Young's inequality on the right hand side of (5.18) again, we have

$$\begin{aligned} &\left(\int_M a \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \right)^{\frac{p-1}{p}} \left(\int_M a \rho^{\alpha+p} |\langle \nabla \rho, \nabla \phi \rangle|^p V^{\tau+(2p-1)(\mu-\lambda)} dv_g \right)^{\frac{1}{p}} \\ &\leq \frac{p-1}{p} \left(\frac{1}{\varepsilon^{\frac{p}{p-1}}} \int_M a \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \right) + \frac{1}{p} \left(\varepsilon^p \int_M a \rho^{\alpha+p} |\langle \nabla \rho, \nabla \phi \rangle|^p V^{\tau+(2p-1)(\mu-\lambda)} dv_g \right) \end{aligned} \quad (5.19)$$

for any $\varepsilon > 0$. Substituting (5.19) into (5.18), we can get

$$A \leq (p-1) \varepsilon^{-\frac{p}{p-1}} \int_M a \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g + \varepsilon^p \int_M a \rho^{\alpha+p} |\langle \nabla \rho, \nabla \phi \rangle|^p V^{\tau+(2p-1)(\mu-\lambda)} dv_g. \quad (5.20)$$

Reorganizing yields

$$\begin{aligned} &\left(C+1+\alpha - (p-1) \varepsilon^{-\frac{p}{p-1}} \right) \int_M a \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g + \int_M \rho^{\alpha+1} |\phi|^p \langle \nabla \rho, \nabla a \rangle V^{\tau+\mu-\lambda} dv_g \\ &\leq \varepsilon^p \int_M a \rho^{\alpha+p} |\langle \nabla \rho, \nabla \phi \rangle|^p V^{\tau+(2p-1)(\mu-\lambda)} dv_g. \end{aligned}$$

Let $f(\varepsilon) = \varepsilon^{-p} \left[C+1+\alpha - (p-1) \varepsilon^{-\frac{p}{p-1}} \right]$, we have

$$\begin{aligned} &f(\varepsilon) \int_M a \rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g + \varepsilon^{-p} \int_M \rho^{\alpha+1} |\phi|^p \langle \nabla \rho, \nabla a \rangle V^{\tau+\mu-\lambda} dv_g \\ &\leq \int_M a \rho^{\alpha+p} |\langle \nabla \rho, \nabla \phi \rangle|^p V^{\tau+(2p-1)(\mu-\lambda)} dv_g \\ &= \int_M a \rho^{\alpha+p} |\langle \nabla^D \rho, \nabla^D \phi \rangle|^p V^{\tau+\lambda-\mu} dv_g. \end{aligned}$$

Note that the function f attains the maximum for $\varepsilon_0 = \left(\frac{p}{C+\alpha+1}\right)^{\frac{p-1}{p}}$ and this maximum value is equal to $\left(\frac{C+\alpha+1}{p}\right)^p$. Therefore we obtain the desired inequality

$$\begin{aligned} \int_M a\rho^{\alpha+p} |\langle \nabla^D \rho, \nabla^D \phi \rangle|^p V^{\tau+\lambda-\mu} dv_g &\geq \left(\frac{C+\alpha+1}{p}\right)^p \int_M a\rho^\alpha |\phi|^p V^{\tau+\lambda-\mu} dv_g \\ &+ \left(\frac{C+\alpha+1}{p}\right)^{p-1} \int_M \rho^{\alpha+1} |\phi|^p \langle \nabla^D \rho, \nabla^D a \rangle V^{\tau+\lambda-\mu} dv_g. \end{aligned} \quad (5.21)$$

The proof of Theorem 5.2 is completed. \square

The following Heisenberg-Pauli-Weyl type inequalities with respect to $D^{\lambda,\mu}$ has been proved:

Theorem 5.3. *Let $(M^n, g, V = e^u)$ be an n -dimensional Riemannian triple and $\lambda, \mu \in \mathbb{R}$. Let $D = D^{\lambda,\mu}$ be the affine connection defined as in (2.1) and $\tau = (n+1)\lambda + \mu$. Let ρ be a nonnegative function on M such that $|\nabla^D \rho| = 1$ and $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$ in the sense of distributions, where $C > 0$ is a constant. Then for any compactly supported smooth function $\phi \in C_0^\infty(M \setminus \rho^{-1}\{0\})$, we have*

$$\left(\int_M \rho^2 \phi^2 V^{\tau+\lambda-\mu} dv_g \right) \left(\int_M |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \right) \geq \frac{(C+1)^2}{4} \left(\int_M \phi^2 V^{\tau+\lambda-\mu} dv_g \right)^2. \quad (5.22)$$

Moreover, for $C > 1$, we have

$$\left(\int_M \rho^4 \phi^2 V^{\tau+\lambda-\mu} dv_g \right) \left(\int_M (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g \right) \geq \frac{(C+1)^4}{16} \left(\int_M \phi^2 V^{\tau+\lambda-\mu} dv_g \right)^2. \quad (5.23)$$

Proof. It follows from $|\nabla^D \rho| = 1$ and $\Delta^D \rho \leq \frac{C}{\rho} V^{\lambda-\mu}$ that

$$\begin{aligned} \Delta^D \rho^2 &= V^{\mu-\lambda} (\Delta \rho^2 + (2\mu + n\lambda) \langle \nabla u, \nabla \rho^2 \rangle) \\ &= V^{\mu-\lambda} (2|\nabla \rho|^2 + 2\rho \Delta \rho + (2\mu + n\lambda) 2\rho \langle \nabla u, \nabla \rho \rangle) \\ &= 2|\nabla^D \rho|^2 V^{\lambda-\mu} + 2\rho \Delta^D \rho \\ &\geq (2C+2) V^{\lambda-\mu}. \end{aligned} \quad (5.24)$$

Multiplying both sides of (5.24) by $\phi^2 V^\tau$ and integrating over M yields

$$\int_M \phi^2 V^\tau \Delta^D \rho^2 dv_g \geq (2C+2) \int_M \phi^2 V^{\tau+\lambda-\mu} dv_g. \quad (5.25)$$

On the other hand, applying the integrating by parts and Hölder's inequality, we have

$$\begin{aligned}
& \int_M \phi^2 V^\tau \Delta^D \rho^2 dv_g \\
&= - \int_M \langle \nabla \rho^2, \nabla \phi^2 \rangle V^{\tau+\mu-\lambda} dv_g \\
&= - \int_M 4\rho\phi \langle \nabla \rho, \nabla \phi \rangle V^{\tau+\mu-\lambda} dv_g \\
&\leq 4 \int_M \rho |\phi| |\nabla \rho| |\nabla \phi| V^{\tau+\mu-\lambda} dv_g \\
&\leq 4 \left(\int_M |\rho\phi \nabla \rho|^2 V^{\tau+\mu-\lambda} dv_g \right)^{\frac{1}{2}} \left(\int_M |\nabla \phi|^2 V^{\tau+\mu-\lambda} dv_g \right)^{\frac{1}{2}} \\
&= 4 \left(\int_M \rho^2 \phi^2 |\nabla \rho|^2 V^{\tau+\mu-\lambda} dv_g \right)^{\frac{1}{2}} \left(\int_M |\nabla \phi|^2 V^{\tau+\mu-\lambda} dv_g \right)^{\frac{1}{2}} \\
&= 4 \left(\int_M \rho^2 \phi^2 V^{\tau+\lambda-\mu} dv_g \right)^{\frac{1}{2}} \left(\int_M |\nabla^D \phi|^2 V^{\tau+\lambda-\mu} dv_g \right)^{\frac{1}{2}}.
\end{aligned} \tag{5.26}$$

Substituting (5.26) into (5.25), we can obtain (5.22) directly.

From (5.25), we have

$$\left| \frac{1}{4} \int_M \phi^2 V^\tau \Delta^D \rho^2 dv_g \right| \geq \frac{C+1}{2} \int_M \phi^2 V^{\tau+\lambda-\mu} dv_g. \tag{5.27}$$

Using the integration by parts again, we can get

$$\begin{aligned}
& \left| \frac{1}{4} \int_M \phi^2 V^\tau \Delta^D \rho^2 dv_g \right| \\
&= \left| -\frac{1}{4} \int_M \langle \nabla^D \rho^2, \nabla \phi^2 \rangle V^\tau dv_g \right| \\
&= \left| -\frac{1}{4} \int_M \langle \nabla \rho^2, \nabla \phi^2 \rangle V^{\tau+\mu-\lambda} dv_g \right| \\
&= \left| -\frac{1}{4} \int_M 4\rho\phi \langle \nabla \rho, \nabla \phi \rangle V^{\tau+\mu-\lambda} dv_g \right| \\
&= \int_M \rho |\phi| |\nabla \rho| |\nabla \phi| V^{\tau+\mu-\lambda} dv_g \\
&\leq \left(\int_M \rho^4 \phi^2 |\nabla \rho|^2 V^{\tau+\mu-\lambda} dv_g \right)^{\frac{1}{2}} \left(\int_M \frac{|\nabla \phi|^2}{\rho^2} V^{\tau+\mu-\lambda} dv_g \right)^{\frac{1}{2}},
\end{aligned} \tag{5.28}$$

where, we use Hölder's inequality in the last equation. On the other hand, letting $\alpha = 0$ in (4.10), we have

$$\int_M (\Delta^D \phi)^2 V^{\tau+\mu-\lambda} dv_g \geq \frac{(C+1)^2}{4} \int_M \frac{|\nabla^D \phi|^2}{\rho^2} V^{\tau+\lambda-\mu} dv_g. \tag{5.29}$$

Inserting (5.27) and (5.29) into (5.28), the inequality (5.23) can be obtained naturally. This completes the proof of Theorem 5.3. \square

We now prove a weight L^p form of the Hardy type inequality with respect to $D^{\lambda, \mu}$, where our new inequality contains more remainder terms (or the improved Hardy type inequality).

Theorem 5.4. *Let $(M^n, g, V = e^\mu)$ be an n -dimensional Riemannian triple and $\lambda, \mu \in \mathbb{R}$. Let $D = D^{\lambda, \mu}$ be the affine connection defined as in (2.1) and $\tau = (n+1)\lambda + \mu$. Let δ be a positive function, and ρ and a be nonnegative functions on M such that $|\nabla^D \rho| = 1$, $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda - \mu}$, $\langle \nabla^D a, \nabla^D \rho \rangle \geq 0$ and $-V^\tau \operatorname{div}^D (a \rho^{1-C} \nabla^D \delta) \geq 0$ in the sense of distribution, where $C > 1$ is a constant, $C + \alpha - 1 > 0$ and $\alpha \in \mathbb{R}$. Then the following inequality*

$$\begin{aligned} \int_M a \rho^\alpha |\nabla^D \phi|^2 V^{\tau + \lambda - \mu} dv_g &\geq \left(\frac{C + \alpha - 1}{2} \right)^2 \int_M a \rho^{\alpha - 2} \phi^2 V^{\tau + \lambda - \mu} dv_g \\ &\quad + \frac{C + \alpha - 1}{2} \int_M \rho^{\alpha - 1} \phi^2 \langle \nabla^D a, \nabla^D \rho \rangle V^{\tau + \lambda - \mu} dv_g \\ &\quad + \frac{1}{4} \int_M a \rho^\alpha \frac{|\nabla^D \delta|^2}{\delta^2} \phi^2 V^{\tau + \lambda - \mu} dv_g \end{aligned} \quad (5.30)$$

holds for any $\phi \in C_0^\infty(M \setminus \rho^{-1}\{0\})$.

Proof. Let $\phi = \rho^\beta \psi$, where $\psi \in C_0^\infty(M \setminus \rho^{-1}\{0\})$ and $\beta < 0$. Then we have

$$\begin{aligned} |\nabla \phi|^2 &= \left| \beta \rho^{\beta - 1} \psi \nabla \rho + \rho^\beta \nabla \psi \right|^2 \\ &= \beta^2 \rho^{2(\beta - 1)} \psi^2 |\nabla \rho|^2 + \rho^{2\beta} |\nabla \psi|^2 + 2\beta \psi \rho^{2\beta - 1} \langle \nabla \rho, \nabla \psi \rangle. \end{aligned}$$

Multiplying the above inequality by $V^{\tau + \mu - \lambda}$ and using $|\nabla^D \rho| = 1$, we obtain

$$\begin{aligned} V^{\tau + \mu - \lambda} |\nabla \phi|^2 &= \beta^2 \rho^{2(\beta - 1)} \psi^2 V^{\tau + \lambda - \mu} + \rho^{2\beta} |\nabla \psi|^2 V^{\tau + \mu - \lambda} \\ &\quad + \beta \rho^{2\beta - 1} \langle \nabla \rho, \nabla \psi^2 \rangle V^{\tau + \mu - \lambda}. \end{aligned} \quad (5.31)$$

Multiplying both sides of (5.31) by $a(x) \rho^\alpha$ and applying the integration by parts over M gives

$$\begin{aligned} &\int_M a \rho^\alpha |\nabla \phi|^2 V^{\tau + \mu - \lambda} dv_g \\ &= \int_M a \beta^2 \rho^{\alpha + 2(\beta - 1)} \psi^2 V^{\tau + \lambda - \mu} dv_g + \int_M a \rho^{\alpha + 2\beta} |\nabla \psi|^2 V^{\tau + \mu - \lambda} dv_g \\ &\quad + \int_M a \beta \rho^{\alpha + 2\beta - 1} \langle \nabla \rho, \nabla \psi^2 \rangle V^{\tau + \mu - \lambda} dv_g \\ &= \beta^2 \int_M a \rho^{\alpha + 2(\beta - 1)} \psi^2 V^{\tau + \lambda - \mu} dv_g - \beta \int_M \operatorname{div} \left(a \rho^{\alpha + 2\beta - 1} V^{\tau + \mu - \lambda} \nabla \rho \right) \psi^2 dv_g \\ &\quad + \int_M a \rho^{\alpha + 2\beta} |\nabla \psi|^2 V^{\tau + \mu - \lambda} dv_g. \end{aligned} \quad (5.32)$$

Since $|\nabla^D \rho| = 1$ and $\Delta^D \rho \geq \frac{C}{\rho} V^{\lambda-\mu}$, we obtain the following inequality

$$\begin{aligned}
& \operatorname{div} \left(a \rho^{\alpha+2\beta-1} V^{\tau+\mu-\lambda} \nabla \rho \right) \\
&= \rho^{\alpha+2\beta-1} V^{\tau+\mu-\lambda} \langle \nabla \rho, \nabla a \rangle + (\alpha+2\beta-1) a \rho^{\alpha+2\beta-2} |\nabla \rho|^2 V^{\tau+\mu-\lambda} \\
&\quad + a \rho^{\alpha+2\beta-1} V^{\tau+\mu-\lambda} \Delta \rho + a(\tau+\mu-\lambda) \rho^{\alpha+2\beta-1} V^{\tau+\mu-\lambda} \langle \nabla \rho, \nabla u \rangle \\
&= \rho^{\alpha+2\beta-1} V^{\tau+\mu-\lambda} \langle \nabla \rho, \nabla a \rangle + (\alpha+2\beta-1) a \rho^{\alpha+2\beta-2} |\nabla^D \rho|^2 V^{\tau+\lambda-\mu} \\
&\quad + a \rho^{\alpha+2\beta-1} V^{\tau} \Delta^D \rho \\
&\geq (C + \alpha + 2\beta - 1) a \rho^{\alpha+2\beta-2} V^{\tau+\lambda-\mu} + \rho^{\alpha+2\beta-1} V^{\tau+\mu-\lambda} \langle \nabla \rho, \nabla a \rangle.
\end{aligned} \tag{5.33}$$

Multiplying both sides of (5.33) by $\beta \psi^2$ and integrating over M yields

$$\begin{aligned}
-\beta \int_M \operatorname{div} \left(a \rho^{\alpha+2\beta-1} V^{\tau+\mu-\lambda} \nabla \rho \right) \psi^2 dv_g &\geq -\beta (C + \alpha + 2\beta - 1) \int_M a \rho^{\alpha+2\beta-2} \psi^2 V^{\tau+\lambda-\mu} dv_g \\
&\quad - \beta \int_M \rho^{\alpha+2\beta-1} \langle \nabla a, \nabla \rho \rangle V^{\tau+\mu-\lambda} \psi^2 dv_g.
\end{aligned} \tag{5.34}$$

Hence combining the inequalities (5.32) and (5.34), we obtain

$$\begin{aligned}
& \int_M a \rho^\alpha |\nabla \phi|^2 V^{\tau+\mu-\lambda} dv_g \\
&\geq \beta^2 \int_M a \rho^{\alpha+2\beta-2} \psi^2 V^{\tau+\lambda-\mu} dv_g - \beta (C + \alpha + 2\beta - 1) \int_M a \rho^{\alpha+2\beta-2} \psi^2 V^{\tau+\lambda-\mu} dv_g \\
&\quad - \beta \int_M \rho^{\alpha+2\beta-1} \langle \nabla a, \nabla \rho \rangle \psi^2 V^{\tau+\mu-\lambda} dv_g + \int_M a \rho^{\alpha+2\beta} |\nabla \psi|^2 V^{\tau+\mu-\lambda} dv_g \\
&= f(\beta) \int_M a \rho^{\alpha+2\beta-2} \psi^2 V^{\tau+\lambda-\mu} dv_g - \beta \int_M \rho^{\alpha+2\beta-1} \langle \nabla a, \nabla \rho \rangle \psi^2 V^{\tau+\mu-\lambda} dv_g \\
&\quad + \int_M a \rho^{\alpha+2\beta} |\nabla \psi|^2 V^{\tau+\mu-\lambda} dv_g,
\end{aligned} \tag{5.35}$$

where $f(\beta) = \beta^2 - \beta (C + \alpha + 2\beta - 1)$. Since $f(\beta)$ attains the maximum for $\beta_0 = \frac{1-\alpha-C}{2}$ and this maximum value is equal to $f(\beta_0) = \left(\frac{C+\alpha-1}{2}\right)^2$, the inequality (5.35) becomes

$$\begin{aligned}
& \int_M a \rho^\alpha |\nabla \phi|^2 V^{\tau+\mu-\lambda} dv_g \\
&\geq \left(\frac{C + \alpha - 1}{2}\right)^2 \int_M a \rho^{-C-1} \psi^2 V^{\tau+\lambda-\mu} dv_g + \int_M a \rho^{1-C} |\nabla \psi|^2 V^{\tau+\mu-\lambda} dv_g \\
&\quad + \frac{C + \alpha - 1}{2} \int_M \rho^{-C} \psi^2 \langle \nabla a, \nabla \rho \rangle V^{\tau+\mu-\lambda} dv_g.
\end{aligned} \tag{5.36}$$

We now focus on the second term on the right hand side of (5.36). Let us define a new variable $\varphi := \delta^{-\frac{1}{2}}\psi$, where $\delta \in C_0^\infty(M \setminus \rho^{-1}\{0\})$ is a positive function. It is clear that

$$\begin{aligned} |\nabla\psi|^2 &= \left| \frac{1}{2}\delta^{-\frac{1}{2}}\varphi\nabla\delta + \delta^{\frac{1}{2}}\nabla\varphi \right|^2 \\ &= \frac{1}{4}\delta^{-1}|\nabla\delta|^2\varphi^2 + \delta|\nabla\varphi|^2 + \langle\nabla\delta, \nabla\varphi\rangle\varphi \\ &\geq \frac{1}{4}\delta^{-1}|\nabla\delta|^2\varphi^2 + \frac{1}{2}\langle\nabla\delta, \nabla\varphi^2\rangle. \end{aligned}$$

Therefore

$$\begin{aligned} \int_M a\rho^{1-C}|\nabla\psi|^2V^{\tau+\mu-\lambda} dv_g &\geq \frac{1}{4}\int_M a\rho^{1-C}\varphi^2\delta^{-1}|\nabla\delta|^2V^{\tau+\mu-\lambda} dv_g \\ &\quad + \frac{1}{2}\int_M a\rho^{1-C}\langle\nabla\delta, \nabla\varphi^2\rangle V^{\tau+\mu-\lambda} dv_g. \end{aligned} \quad (5.37)$$

Here, first applying the integration by parts to the second term on the right hand side of (5.37) and then using the differential inequality $-V^\tau\operatorname{div}^D(a\rho^{1-C}\nabla^D\delta) \geq 0$, we get

$$\begin{aligned} &\int_M a\rho^{1-C}\langle\nabla\delta, \nabla\varphi^2\rangle V^{\tau+\mu-\lambda} dv_g \\ &= \int_M a\rho^{1-C}\langle\nabla^D\delta, \nabla\varphi^2\rangle V^\tau dv_g \\ &= -\int_M \varphi^2\operatorname{div}(V^\tau a\rho^{1-C}\nabla^D\delta) dv_g \\ &= -\int_M \varphi^2 D_i(a\rho^{1-C}\nabla_i^D\delta) V^\tau dv_g \\ &\geq 0. \end{aligned} \quad (5.38)$$

Then combining (5.37) and (5.38), we have

$$\int_M a\rho^{1-C}|\nabla\psi|^2V^{\tau+\mu-\lambda} dv_g \geq \frac{1}{4}\int_M a\rho^{1-C}\varphi^2\delta^{-1}|\nabla\delta|^2V^{\tau+\mu-\lambda} dv_g. \quad (5.39)$$

Taking $\varphi = \delta^{-\frac{1}{2}}\rho^{\frac{C+\alpha-1}{2}}\phi$ into (5.39), then (5.39) becomes

$$\int_M a\rho^{1-C}|\nabla\psi|^2V^{\tau+\mu-\lambda} dv_g \geq \frac{1}{4}\int_M a\rho^\alpha \frac{|\nabla\delta|^2}{\delta^2} \phi^2 V^{\tau+\mu-\lambda} dv_g. \quad (5.40)$$

Finally, by using the equality $\psi = \rho^{-\beta}\phi = \rho^{\frac{C+\alpha-1}{2}}\phi$ and taking (5.40) into (5.36), we get the desired inequality (5.30). This completes the proof of Theorem 5.4. \square

Acknowledgements

The research of authors is supported by NSFC (No.12101530), the Science and Technology Project of Henan Province (No.232102310321), and the Key Scientific Research Program in Universities of Henan Province (Nos.21A110021, 22A110021) and Nanhu Scholars Program for Young Scholars of XYNU (No.2023).

REFERENCES

- [1] G. Carron, *Inégalités de Hardy sur les variétés riemanniennes non-compactes*, J. Math. Pures Appl. (9) **76** (1997), 883-891.
- [2] F. Du and J. Mao, *Hardy and Rellich type inequalities on metric measure spaces*, J. Math. Anal. Appl. **429** (2015), 354-365.
- [3] J. Flynn, L. Nguyen, G. Lu and S. Mazumdar, *Hardy's identities and inequalities on Cartan-Hadamard manifolds*, J. Geom. Anal. **33** (2023), 34pp.
- [4] G. Grillo, *Hardy and Rellich-type inequalities for metrics defined by vector fields*, Potential Anal. **18** (2003) 187-217.
- [5] N. Ghoussoub and A. Moradifam, *Bessel pairs and optimal Hardy and Hardy-Rellich inequalities*, Math. Ann. **349** (2011), 1-57.
- [6] G. Huang and H. Li, *Gradient estimates and entropy formulae of porous medium and fast diffusion equations for the Witten Laplacian*, Pacific J. Math. **268** (2014), 47-78.
- [7] G. Huang and M. Zhu, *Some integral inequalities on weighted Riemannian manifolds with boundary*, Proc. Amer. Math. Soc. **151** (2023), 4961-4970.
- [8] G. Huang and M. Zhu, *Estimates for the first eigenvalues of the affine Laplacian*, Arch. Math. (Basel) **121** (2023), no. 1, 77-87.
- [9] G. Huang, B. Ma, and M. Zhu, *Colesanti type inequalities for affine connections*, Anal. Math. Phys. **13** (2023), 15pp.
- [10] X. Huang and D. Ye, *First order Hardy inequalities revisited*, Commun. Math. Res. **38** (2022), 535-559.
- [11] Y. Jin and S. Shen, *Some L^p -Hardy and L^p -Rellich type inequalities with remainder terms*, J. Aust. Math. Soc. **113** (2022), 79-98.
- [12] I. Kombe, *Sharp weighted Rellich and uncertainty principle inequalities on Carnot groups*, Commun. Appl. Anal. **14** (2010), no.2, 251-271.
- [13] I. Kombe and M. Özaydin, *Improved Hardy and Rellich inequalities on Riemannian manifolds*, Trans. Amer. Math. Soc. **361** (2009), 6191-6203.
- [14] I. Kombe and M. Özaydin, *Hardy-Poincaré, Rellich and uncertainty principle inequalities on Riemannian manifolds*, Trans. Amer. Math. Soc. **365** (2013), 5035-5050.
- [15] I. Kombe and A. Yener, *Weighted Hardy and Rellich type inequalities on Riemannian manifolds*, Math. Nachr. **289** (2016), 994-1004.
- [16] X.-D. Li, *Liouville theorems for symmetric diffusion operators on complete Riemannian manifolds*, J. Math. Pures Appl. **84** (2005), 1295-1361.
- [17] Y. Li, A. Abolarinwa, H. Alkhalidi and A. Ali, *Some inequalities of Hardy type related to Witten Laplace operator on smooth metric measure spaces*, Mathematics **10** (2022), 4580.
- [18] J. Li, and C. Xia, *An integral formula for affine connections*, J. Geom. Anal. **27** (2017), 2539-2556.
- [19] C. Meng, H. Wang and W. Zhao, *Hardy type inequalities on closed manifolds via Ricci curvature*, Proc. R. Soc. Edinb. Sect. A Math. **151** (2021), 993-1020.
- [20] V. Nguyen, *New sharp Hardy and Rellich type inequalities on Cartan-Hadamard manifolds and their improvements*, Proc. Roy. Soc. Edinburgh Sect. A **150** (2020), no. 6, 2952-2981.
- [21] J. Vazquez and E. Zuazua, *The Hardy inequality and the asymptotic behaviour of the heat equation with an inverse-square potential*, J. Funct. Anal. **173** (2000), no. 1, 103-153.
- [22] P. Wang and F. Zeng, *Estimates for the first eigenvalue of diffusion-type operators in weighted manifolds*, J. Pseudo-Differ. Oper. Appl. **14** (2023), 18pp.
- [23] C. Xia, *Hardy and rellich type inequalities on complete manifolds*, J. Math. Anal. Appl. **409** (2014), 84-90.