

# ATOMISTIC-TO-CONTINUUM CONVERGENCE FOR QUASI-STATIC CRACK GROWTH IN BRITTLE MATERIALS

MANUEL FRIEDRICH AND JOSCHA SEUTTER

**ABSTRACT.** We study the atomistic-to-continuum limit for a model of quasi-static crack evolution driven by time-dependent boundary conditions. We consider a two-dimensional atomic mass spring system whose interactions are modeled by classical interaction potentials, supplemented by a suitable irreversibility condition accounting for the breaking of atomic bonding. In a simultaneous limit of vanishing inter-atomic distance and discretized time step, we identify a continuum model of quasi-static crack growth in brittle fracture [32] featuring an irreversibility condition, a global stability, and an energy balance. The proof of global stability relies on a careful adaptation of the jump-transfer argument in [31] to the atomistic setting.

## 1. INTRODUCTION

Since the pioneering work of GRIFFITH [43] in the 1920's, brittle materials and their fracture behavior have been subject to intensive research in mechanical engineering. The groundbreaking idea of Griffith's theory lies in regarding crack formation and propagation as the result of a competition between elastic bulk energy and a surface energy related to the increase of the crack. In their seminal paper [32], FRANC-FORT AND MARIGO adopted this viewpoint and paved the way to a variational approach to fracture, where displacements and crack paths are found by the principle of energy minimization. The proposed evolutionary model, called a *quasi-static crack evolution*, is characterized by three principles, namely an irreversibility condition, a static equilibrium at each time, and an energy balance reflecting the nondissipativity of the process. In contrast to the classical theory, this framework does not rely on prescribed crack paths and also effectively accounts for crack initiation.

Over the last two decades, the variational theory of brittle fracture has been studied in a remarkable depth, see [10] for a broad overview. Here, we mention only some of the numerous contributions [18, 22, 23, 25, 26, 31, 34] addressing the mathematical well-posedness of the model from [32] in various settings. In all results, the existence of continuous-time crack evolutions is established as the limit of suitably time-discretized approximations resulting from an iterative minimization of so-called Griffith-type functionals. Such energies comprise elastic bulk contributions for the sound regions of the body and surface terms along the crack, and are typically defined on (generalized) functions of bounded variation [5]. Among various technical difficulties that arise in the discrete-to-continuous passage in time, the most severe one is the *stability of unilateral minimality properties* [40] which ensures global minimality of displacements along the irreversible fracture process.

Besides being the limit of time-discretized solutions, quasi-static crack evolutions have been identified as effective variational limits of sequences of problems in various settings of applicative relevance, including phase-field approximation schemes [38], finite element approximation [41, 42], homogenization of brittle

---

2020 *Mathematics Subject Classification.* 49J45, 70G75, 74A25, 74R10.

*Key words and phrases.* Brittle materials, variational fracture, free-discontinuity problems, quasistatic crack propagation, irreversibility condition, atomistic systems, discrete-to-continuum limits,  $\Gamma$ -convergence.

composite materials [40], or a cohesive-to-brittle passage in the limit of infinite specimen size [39]. The goal of this work is to advance the understanding of effective theories for quasi-static crack growth by establishing a rigorous connection between atomistic and continuum models in brittle fracture.

The passage from atomistic systems to continuum models in the limit of vanishing interatomic distance has been a thriving field of research in the last years, see [8, 11] for an overview. Such an asymptotic analysis is frequently grounded on the variational tool of  $\Gamma$ -convergence [12, 21], which ensures convergence of minimizers and hence allows for establishing a rigorous relation between microscopic and macroscopic models. Among the vast body of literature, this approach has also been used to derive and validate continuum theories in brittle and cohesive fracture<sup>1</sup>, both in one [13, 16, 46] and higher dimensions [6, 15, 35, 45]. Yet, the analysis of static problems has been predominant and the evolutionary nature of crack growth has largely been neglected. Indeed, for atomistic problems in fracture mechanics, available rigorous results seem to be limited to [14] where a one-dimensional Lennard-Jones system is investigated via a minimizing movements scheme. The subject of this paper is to perform an atomistic-to-continuum analysis beyond the 1D setting accounting for irreversibility, and to derive a continuum quasi-static crack evolution in the sense of [32].

We now describe our setting. We build upon the previous work in [7] where an atomistic model for crack growth has been introduced including an irreversibility condition as a key feature. More precisely, we consider systems of interacting particles arranged in an atomic lattice occupying the bounded reference domain of a material. The interactions are modeled by classical potentials from Molecular Mechanics [1, 44], e.g., Lennard-Jones-type potentials. Following an energy minimization principle, the evolution is driven by time-dependent boundary conditions. We implement an irreversibility constraint along the evolution which resembles the use of a maximal-opening memory variable in cohesive continuum models for crack evolution [27, 39]: Whenever the distance between two atoms does not exceed a certain threshold, interactions are considered to be ‘elastic’ and can recover completely after removal of loading. On the contrary, stretched beyond the threshold, interactions are supposed to be ‘damaged’, and the system should be affected at all future times. This is modeled by a memory variable that tracks the deformation history of each interaction in the system by taking the supremum over all past time steps.

While the modeling framework in [7] is rather general, in this work we adopt a specific simple model for performing the passage to the continuum theory. More precisely, we assume that the atoms in the reference configuration are arranged in a triangular lattice and interact via nearest-neighbor potentials. Moreover, we restrict ourselves to the case of anti-plane shear displacements. As the interatomic distance  $\varepsilon$  tends to zero, the atomistic energies can then be related by means of  $\Gamma$ -convergence to a *free-discontinuity functional* of the form

$$\int_{\Omega} \Phi(\nabla u) \, dx + \int_{S(u)} \varphi(\nu_u) \, d\mathcal{H}^1, \quad (1.1)$$

where  $\Omega \subset \mathbb{R}^2$  denotes the reference configuration,  $u$  the displacement with gradient  $\nabla u$ , and  $S(u)$  stands for the crack surface with normal  $\nu_u$ . This follows by adapting the result in [35] where a vectorial variant of the present model has been investigated. In particular, the model features an elastic part with energy density  $\Phi$  derived from the molecular interaction potentials and a surface term comparable to the one-dimensional Hausdorff measure of the crack set, i.e.,  $\mathcal{H}^1(S(u))$ . A fundamental feature of the model is the anisotropic density  $\varphi$ , which is induced by the triangular lattice and favors the formation of cracks along crystallographic lines, see [36, 37] for rigorous results on crystal cleavage and [7] for numerical evidence.

---

<sup>1</sup>Already Griffith [43, p. 165] motivated his approach by heuristic considerations on the microscopic nature of fracture: ‘It is known that, in the formation of a crack in a body composed of molecules which attract one another, work must be done against the cohesive forces of the molecules on either side of the crack. This work appears as potential surface energy.’

Although we believe that in principle our analysis could be carried out for far more general lattice models, including long-range and multi-body interactions (see e.g. [6]), we prefer to consider here a simple mass-spring model, which already gives rise to a Griffith-type energy in the continuum limit with a non-degenerate elastic density and anisotropic surface terms. This allows us to focus on the challenges due to the evolutionary nature of the fracture process, in particular issues related to irreversibility.

Starting from an incremental minimization scheme with time-step size  $\delta$  and corresponding time-discrete solutions for the atomistic model, we show in our main result (Theorem 2.2) that, in the simultaneous limit  $\varepsilon, \delta \rightarrow 0$ , the atomistic evolutions converge to an irreversible quasi-static crack evolution in the sense of [32]. More precisely, we find a pair  $t \mapsto (u(t), K(t))$ , where  $u(t)$  lies in the space  $SBV(\Omega)$ , see [5], and  $K(t)$  is a rectifiable set with  $S(u(t)) \subset K(t)$ , satisfying: (1) an irreversibility condition, (2) a global stability at all times, and (3) an energy balance law, see (2.16)–(2.17) for details. As a byproduct, we show that the energies converge at all times as  $\varepsilon \rightarrow 0$ . Moreover, for suitable interpolations of the atomistic displacements that are exhibiting jumps, we obtain strong convergence of displacement gradients and convergence of crack sets in the sense of  $\sigma$ -convergence introduced in [40], see also Section 3.4.

Brittle materials frequently develop cracks already at moderately large strains without significant plastic deformation. As in [16, 35, 36], this observation is reflected in our model since either elastic displacements are small, namely of order  $\varepsilon^{1/2}$  (the exponent  $1/2$  is related to the fact that  $\Phi$  is positively homogeneous of order 2, see (2.14) below), or springs leading to fracture in the limit are elongated by a factor  $\varepsilon^{-1/2}$ . As a consequence, the limit  $\varepsilon \rightarrow 0$  involves a linearization of the elastic energy and the fracture response becomes asymptotically brittle despite the presence of cohesive forces on the atomistic level. In other words, while the surface energy in (1.1) depends on the geometry of the crack, it is not affected by its opening. To our best knowledge, our main theorem represents the first result identifying a quasi-static crack evolution not only in the atomistic-to-continuum limit but also in the limit from nonlinear-to-linearized elastic energies.

Whereas the derivation of (1.1) via  $\Gamma$ -convergence follows along the lines of [35], and the techniques to pass to the limit in energy balances are well established [22, 40], the essential novelty of the proof consists in showing the stability of the unilateral minimality property of the atomistic evolutions. This is achieved by a suitable adaptation of the jump-transfer argument from [31] to the atomistic setting. A discrete variant of this construction has indeed already been proposed in the literature in terms of discontinuous finite elements [41, 42]. However, the main challenge in the present atomistic setting lies in the fact that ‘broken’ interactions do not contribute a fixed energy, but their contribution depends also on the elongation. (Only in the continuum limit this effect vanishes, as discussed above.) In this sense, our problem is related to the work by GIACOMINI [39], where stability of unilateral minimality is shown in a cohesive-to-brittle limit. In both problems, it is essential to transfer the jump only on the part of the crack where the memory variable exceeds a certain constant. We achieve this by splitting the broken interactions on the atomistic level into two parts  $\mathcal{C}^{\text{small}}$  and  $\mathcal{C}^{\text{large}}$ , related to small and large values of the memory variable, respectively. Then, we carefully introduce interpolations exhibiting jumps only at sites associated to  $\mathcal{C}^{\text{large}}$ . With this, we can suitably adapt the original jump-transfer argument and we identify the continuum crack set by resorting to the notion of  $\sigma^p$ -convergence introduced in [22]. Eventually, although in general the concepts of  $\sigma^p$ - and  $\sigma$ -convergence are different, we show that they coincide along the evolution, cf. also [40].

Inspired by [37], the arguments involve several types of interpolations, ranging from piecewise affine interpolations to functions exhibiting jumps inside the triangles of the lattice. The use of the interpolations is vital to our reasoning since it allows us to directly employ tools for  $SBV$  functions, such as compactness and lower semicontinuity results as well as the notion of  $\sigma^p$ -convergence. This is the main reason why we focus on a simple mass-spring model. In fact, discrete-to-continuum limits for more general atomistic energies involving long-range interactions are usually based on slicing [15] or on more abstract

techniques, such as the localization method of  $\Gamma$ -convergence and integral representation [6]. Transferring our arguments for the proof of stability of unilateral minimality to these frameworks appears to require nontrivial adaptations.

Let us mention that in our result we start from time-discrete solutions for the atomistic evolutions, for existence of time-continuous evolutions on the atomistic level is not guaranteed by [7]. In fact, in [7] time-continuous solutions have only been established for a model with an additional rate-dependent dissipation term.

The paper is organized as follows. In Section 2 we introduce the atomistic model for irreversible crack growth from [7] and state our main result. Then, in Section 3 we suitably split the atomistic energy in an elastic and a crack part, and we introduce useful interpolations for atomistic displacements. Moreover, we prove compactness and lower semicontinuity results for the elastic energy and deal with different notions of set-convergence. Section 4 is devoted to the proof of our main result, where we first derive properties of the time-discrete atomistic problems and pass to the continuum limit afterwards. To ensure that the latter can be identified as a quasi-static crack evolution, we crucially rely on the above-mentioned stability of unilateral minimality whose proof is postponed to Section 5.

## 2. MODEL AND MAIN RESULTS

In this section we describe our model and present the main result on an atomistic-to continuum convergence for quasi-static crack growth.

**Atomistic anti-plane shear model.** We consider a model for anti-plane shear displacements. This corresponds to a reference configuration of the form  $U \times \mathbb{R}$ , where  $U \subset \mathbb{R}^2$  is a Lipschitz set, and the deformations for  $(x_1, x_2) \in U$  and  $x_3 \in \mathbb{R}$  are of the special form  $(x_1, x_2, x_3) \mapsto (x_1, x_2, x_3 + w(x_1, x_2))$ , where  $w: U \rightarrow \mathbb{R}$  is a scalar-valued map. Since the behavior is identical for each cross section  $\{x_3 = c\}$  for  $c \in \mathbb{R}$ , one can effectively reduce to deformations  $y: U \rightarrow \mathbb{R}^3$  of the form  $y(x_1, x_2) = (x_1, x_2, w(x_1, x_2))$ , which corresponds to the cross section  $\{x_3 = 0\}$ . We consider an atomistic model where in the reference configuration the particles in the cross section  $\{x_3 = 0\}$  are given by the points of the scaled triangular lattice  $\varepsilon\mathcal{L}$  inside  $U$ . Here,

$$\mathcal{L} := \begin{pmatrix} 1 & 1/2 \\ 0 & \sqrt{3}/2 \end{pmatrix} \mathbb{Z}^2 = \{\lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 : \lambda_1, \lambda_2 \in \mathbb{Z}\},$$

where  $\mathbf{v}_1 := e_1$  and  $\mathbf{v}_2 := (1/2, \sqrt{3}/2)^T$  are the lattice vectors, and  $\varepsilon > 0$  is a small parameter representing the length scale of the typical interatomic distances. We collect all lattice vectors in the set  $\mathcal{V} := \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_2 - \mathbf{v}_1\}$  and denote by  $\mathcal{L}_\varepsilon(U) := \varepsilon\mathcal{L} \cap U$  the particles in the reference configuration  $U$ . Now, deformations  $y: \mathcal{L}_\varepsilon(U) \rightarrow \mathbb{R}^3$  are given by  $y(x) = (x, w(x))$  for all  $x \in \mathcal{L}_\varepsilon(U)$ , where  $w: \mathcal{L}_\varepsilon(U) \rightarrow \mathbb{R}$  is a scalar displacement field in vertical direction. We model interactions between the particles by classical potentials from Molecular Mechanics [1, 44]. Following [35], we consider an associated *phenomenological energy* featuring nearest-neighbor interactions, given by

$$\mathcal{F}_\varepsilon(y) := \frac{1}{2} \sum_{(x, x') \in \text{NN}_\varepsilon(U)} W\left(\frac{|y(x) - y(x')|}{\varepsilon}\right), \quad (2.1)$$

where  $\text{NN}_\varepsilon(U)$  denotes the *nearest neighbors*

$$\text{NN}_\varepsilon(U) := \{(x, x') \in \mathcal{L}_\varepsilon(U) \times \mathcal{L}_\varepsilon(U) : |x - x'| = \varepsilon\}$$

(the factor  $\frac{1}{2}$  accounts for double counting in the sum), and  $W$  represents a *two-body interaction potential* satisfying

- (i)  $W \geq 0$  and  $W(r) = 0$  if and only if  $r = 1$ .

- (ii)  $W$  is increasing on  $(1, \infty)$ .
- (iii)  $W$  is Lipschitz on  $[1, \infty)$  and  $C^2$  on  $(1, \infty)$  with  $\lim_{r \searrow 1} W'(r) = \mu > 0$  and  $\lim_{r \searrow 1} |W''(r)| < \infty$ .
- (iv)  $\lim_{r \rightarrow \infty} W(r) = \kappa$  with  $\kappa > 0$  constant.

Here, on  $[1, \infty)$  the potential  $W$  mimics a Lennard-Jones-type potential up to a shift by  $\kappa$ , which is for convenience only to ensure that the potential is nonnegative. We emphasize that we use a linear instead of quadratic behavior close to the global minimizer 1 as this allows to derive an effective elastic energy density in (2.14) below with *quadratic* growth. We point out that discrete energies of the form (2.1) and their relation to continuum energies of Griffith-type in the sense of  $\Gamma$ -convergence have been studied intensively in the literature in a general framework including multi-body and multi-scale interactions, see e.g. [6, 15, 16]. As our focus lies on the irreversibility of crack growth, we consider here only a simple mass-spring model.

We impose Dirichlet boundary conditions on a part  $\partial_D U \subset \partial U$  of the boundary. Both in the framework of atomistic systems and of function spaces allowing for discontinuities, this is usually implemented by imposing boundary conditions in a *neighborhood* of the boundary. More precisely, we suppose that there exists another Lipschitz set  $\Omega \supset U$  with  $\partial_D U = \partial U \cap \Omega$ . Then, for a given function  $h: \Omega \rightarrow \mathbb{R}^3$ , we call deformations  $y: \mathcal{L}_\varepsilon(\Omega) \rightarrow \mathbb{R}^3$  *admissible* with respect to  $h$  if they satisfy

$$y(x) = h(x) \quad \text{for all } x \in \mathcal{L}_\varepsilon(\Omega \setminus U), \quad (2.2)$$

where similarly as before, we set  $\mathcal{L}_\varepsilon(\Omega \setminus U) = \varepsilon\mathcal{L} \cap (\Omega \setminus U)$  and  $\mathcal{L}_\varepsilon(\Omega) = \varepsilon\mathcal{L} \cap \Omega$ . From now on, deformations will always be defined on the enlarged set  $\Omega$ , and accordingly the sum in (2.1) will run over pairs of nearest-neighbor points in  $\mathcal{L}_\varepsilon(\Omega)$ , denoted by  $\text{NN}_\varepsilon(\Omega)$ .

**Evolutionary model: Irreversibility condition and memory variable.** Based on the energy (2.1), we now introduce an evolutionary model accounting for an irreversibility constraint along the fracture process in a time interval  $[0, T]$ . In analogy to [7], we implement such an irreversibility condition by incorporating a memory variable for each interacting pair of neighboring atoms  $(x, x') \in \text{NN}_\varepsilon(\Omega)$ . More precisely, suppose that for a fixed time step  $t^k \in [0, T]$  a finite family of deformations  $(y^j)_{j < k}$  at previous time steps  $(t^j)_{j < k}$  is given. Then we consider the *memory variable* related to  $(x, x') \in \text{NN}_\varepsilon(\Omega)$  by

$$M_{x,x'}((y^j)_{j < k}) := \sup_{j < k} \frac{|y^j(x') - y^j(x)|}{\varepsilon}. \quad (2.3)$$

The ‘spring’ connecting a pair  $(x, x') \in \text{NN}_\varepsilon(\Omega)$  is considered *elastic* (or *undamaged*) if  $\varepsilon^{-1}|y^j(x) - y^j(x')|$  does not exceed a given *threshold*  $\bar{R} > 1$  for all past times  $t^j < t^k$ , i.e., if  $M_{x,x'}((y^j)_{j < k}) \leq \bar{R}$ . In this case, the pair interaction can recover completely after removal of loading and thus the energy contribution of  $(x, x')$  is simply given by the stored elastic energy  $W(\varepsilon^{-1}|y(x) - y(x')|)$ . By contrast, if  $M_{x,x'}((y^j)_{j < k}) > \bar{R}$ , the spring is supposed to be *damaged* and the complete deformation history  $(y^j)_{j < k}$  should be reflected at present time. With (2.3), this leads to the energy contribution with memory

$$W(M_{x,x'}((y^j)_{j < k})) \vee W(\varepsilon^{-1}|y(x) - y(x')|),$$

where  $a \vee b := \max\{a, b\}$  for  $a, b \in \mathbb{R}$ . We refer to [7, (2.4)] for more details and mention that here we adopted as memory variable the *maximal opening*, as implemented in continuum settings, see e.g. [27, 39] or [10, Section 5.2]. Note, that also other choices would be conceivable, such as that of a cumulative increment, see e.g. [10, 20]. This, however, is not discussed in this work. Summarizing, we can express the energy contribution of a pair  $(x, x') \in \text{NN}_\varepsilon(\Omega)$  by

$$F_{x,x'}(y, (y^j)_{j < k}) := \begin{cases} W(\varepsilon^{-1}|y(x) - y(x')|) & \text{if } M_{x,x'}((y^j)_{j < k}) \leq \bar{R} \\ W(M_{x,x'}((y^j)_{j < k})) \vee W(\varepsilon^{-1}|y(x) - y(x')|) & \text{if } M_{x,x'}((y^j)_{j < k}) > \bar{R}. \end{cases}$$

Then, the assigned energy for a given history  $(y^j)_{j < k}$  of deformations and an admissible deformation  $y: \mathcal{L}_\varepsilon(\Omega) \rightarrow \mathbb{R}^3$  satisfying (2.2) (for some time-dependent boundary condition) is given by

$$\mathcal{F}_\varepsilon(y, (y^j)_{j < k}) := \frac{1}{2} \sum_{(x, x') \in \text{NN}_\varepsilon(\Omega)} F_{x, x'}(y, (y^j)_{j < k}). \quad (2.4)$$

**Formulation for rescaled displacement fields.** Since we restricted ourselves to the case of anti-plane shear, the model can be rewritten in terms of the vertical displacement  $w: \mathcal{L}_\varepsilon(\Omega) \rightarrow \mathbb{R}$ . In particular, given  $y(x_1, x_2) = (x_1, x_2, w(x_1, x_2))$ , for each  $(x, x') \in \text{NN}_\varepsilon(\Omega)$ , using that  $|x - x'| = \varepsilon$  and defining  $\Psi(s) := W(\sqrt{1 + s^2})$  for  $s \in \mathbb{R}$ , we get

$$W\left(\frac{|y(x) - y(x')|}{\varepsilon}\right) = W\left(\sqrt{1 + \left(\frac{|w(x) - w(x')|}{\varepsilon}\right)^2}\right) = \Psi\left(\frac{|w(x) - w(x')|}{\varepsilon}\right), \quad (2.5)$$

where the assumptions on  $W$  from above lead to the following properties for  $\Psi$ :

- (i')  $\Psi \geq 0$  and  $\Psi(s) = 0$  if and only if  $s = 0$ .
- (ii')  $\Psi$  is increasing on  $(0, \infty)$ .
- (iii')  $\Psi$  is  $C^2$  and Lipschitz on  $\mathbb{R}$  with  $\Psi'(0) = 0$  and  $\Psi''(0) = \mu > 0$ .
- (iv')  $\lim_{|s| \rightarrow \infty} \Psi(s) = \kappa$ .

We now discuss the scaling of the energy in terms of different displacements  $w$ . One can check that the energy of a displacement of the form  $w_1 = t\chi_V$ , for  $t \in \mathbb{R} \setminus \{0\}$  and  $V \subset \Omega$  with  $\partial V \cap \Omega$  being smooth, scales like  $\sim \varepsilon^{-1}$  as each spring crossing  $\partial V \cap \Omega$  contributes  $\Psi(t/\varepsilon) \sim \kappa$  to the energy and the number of springs crossing  $\partial V \cap \Omega$  scales like  $\varepsilon^{-1}$ . On the other hand, the energy of an affine displacement  $w_2(x) = a_\varepsilon \cdot x$ , for  $a_\varepsilon \in \mathbb{R}^2$  possibly depending on  $\varepsilon$ , scales like  $\varepsilon^{-2}\Psi(a_\varepsilon)$  since  $\#\text{NN}_\varepsilon(\Omega) \sim \varepsilon^{-2}$ .

The most interesting regime is when the contributions of the displacement with crack  $w_1$  and the elastic displacement  $w_2$  are of the same order. Since for small values  $s$  we have  $\Psi(s) \sim s^2$  (see (iii')), this suggests  $a_\varepsilon \sim \varepsilon^{1/2}$ , i.e., elastic displacements should be of (infinitesimal) small order. To account for this, we pass to *rescaled displacements*

$$u(x) := \varepsilon^{-1/2}w(x) \quad \text{for } x \in \mathcal{L}_\varepsilon(\Omega).$$

We write the energy (2.1) in terms of  $u$  and rescale by  $\varepsilon$  to ensure that for the displacements  $w_1$  and  $w_2$  considered above the energy is of order 1. Using also (2.5), this leads to

$$\mathcal{E}_\varepsilon(u) := \frac{\varepsilon}{2} \sum_{(x, x') \in \text{NN}_\varepsilon(\Omega)} \Psi\left(\frac{\varepsilon^{1/2}|u(x) - u(x')|}{\varepsilon}\right). \quad (2.6)$$

It remains to transfer the memory variable to the setting of rescaled displacements. To this end, we observe that for each pair  $(x, x') \in \text{NN}_\varepsilon(\Omega)$ , setting  $R := (\bar{R}^2 - 1)^{1/2}$ , we have

$$\frac{|y^j(x) - y^j(x')|}{\varepsilon} > \bar{R} \iff \frac{|w^j(x) - w^j(x')|}{\varepsilon} > R \iff \frac{|u^j(x) - u^j(x')|}{\varepsilon} > \varepsilon^{-1/2}R,$$

where  $(w^j)_j$  and  $(u^j)_j$  denote the history of (rescaled) displacements related to the deformations  $(y^j)_j$ . We now define the  $u$ -dependent memory variable as

$$M_{x, x'}^\varepsilon((u^j)_{j < k}) := \varepsilon^{-1/2} \sup_{j < k} |u^j(x) - u^j(x')|, \quad (2.7)$$

where we include  $\varepsilon$  in the notation to highlight the rescaled nature of the quantity. Defining

$$E_{x, x'}^\varepsilon(u, (u^j)_{j < k}) := \begin{cases} \Psi(\varepsilon^{-1/2}|u(x) - u(x')|) & \text{if } M_{x, x'}^\varepsilon((u^j)_{j < k}) \leq R \\ \Psi(M_{x, x'}^\varepsilon((u^j)_{j < k})) \vee \Psi(\varepsilon^{-1/2}|u(x) - u(x')|) & \text{if } M_{x, x'}^\varepsilon((u^j)_{j < k}) > R, \end{cases}$$

we can formulate the energy (2.4) by referring only to the rescaled displacement  $u: \mathcal{L}_\varepsilon(\Omega) \rightarrow \mathbb{R}$ , namely

$$\mathcal{E}_\varepsilon(u, (u^j)_{j < k}) := \frac{\varepsilon}{2} \sum_{\text{NN}_\varepsilon(\Omega)} E_{x,x'}^\varepsilon(u, (u^j)_{j < k}). \quad (2.8)$$

In this setting, the boundary condition in (2.2) transfers to requiring that displacements  $u$  lie in the set

$$\mathcal{A}_\varepsilon(g) := \{u: \mathcal{L}_\varepsilon(\Omega) \rightarrow \mathbb{R} : u(x) = g(x) \quad \text{for all } x \in \mathcal{L}_\varepsilon(\Omega \setminus U)\},$$

where  $g := \varepsilon^{-1/2}h \cdot e_3$ .

**2.1. Main result: Convergence of quasi-static crack growth.** In this subsection, we state our main result on the atomistic-to-continuum convergence of crack growth. We start by introducing evolutions on the atomistic level. To this end, we fix an arbitrary sequence  $(\varepsilon_n)_n \subset (0, \infty)$  with  $\varepsilon_n \rightarrow 0$  as  $n \rightarrow \infty$ . For simplicity, in the following we write  $\mathcal{E}_n$ ,  $\mathcal{E}_n$ ,  $\mathcal{L}_n(\Omega)$ ,  $\text{NN}_n(\Omega)$ , and  $\mathcal{A}_n$  in place of  $\mathcal{E}_{\varepsilon_n}$ ,  $\mathcal{E}_{\varepsilon_n}$ ,  $\mathcal{L}_{\varepsilon_n}(\Omega)$ ,  $\text{NN}_{\varepsilon_n}(\Omega)$ , and  $\mathcal{A}_{\varepsilon_n}$ .

We introduce a time discrete evolution which is driven by time-dependent boundary conditions  $g \in W^{1,1}(0, T; W^{2,\infty}(\Omega))$ . We choose a sequence  $(\delta_n)_n \subset (0, \infty)$  with  $\delta_n \rightarrow 0$  and for each  $\delta_n$  we consider the subdivision  $0 = t_n^0 < \dots < t_n^{T/\delta_n} = T$  of the interval  $[0, T]$  with step size  $\delta_n$ . (Without restriction, we assume that  $T/\delta_n \in \mathbb{N}$ .) Correspondingly, let  $(g(t_n^k))_k$  be the sequence of boundary data at different time steps  $k \in \{0, \dots, T/\delta_n\}$ . We suppose that the *initial value*  $u_n^0 \in \mathcal{A}_n(g(0))$  is a minimum configuration in the sense that

$$u_n^0 \in \operatorname{argmin}\{\mathcal{E}_n(v) : v \in \mathcal{A}_n(g(0))\}, \quad (2.9)$$

with  $\mathcal{E}_n$  as given in (2.6). We inductively define an evolution as follows: given  $(u_n^j)_{0 \leq j \leq k-1}$ , we let

$$u_n^k \in \operatorname{argmin}\{\mathcal{E}_n(v, (u_n^j)_{j < k}) : v \in \mathcal{A}_n(g(t_n^k))\}, \quad (2.10)$$

where the minimization problem is influenced by  $(u_n^j)_{j < k}$  in terms of the memory variable given in (2.7), see (2.8). We then define the evolution  $u_n: [0, T] \times \mathcal{L}_n(\Omega) \rightarrow \mathbb{R}$ , discrete in space and piecewise constant in time, by

$$u_n(t) := u_n^k \text{ for } t \in [t_n^k, t_n^{k+1}). \quad (2.11)$$

The existence of minimizers in (2.10) follows from the Direct Method and is briefly discussed in Lemma 4.1 below. We emphasize that we consider here an evolution discretized in time which is close to the spirit of the approach by FRANCFORT AND MARIGO [32]. On the contrary, the existence of a continuous-in-time solution for the energy (2.8) has been established in [7] with an additional rate-dependent dissipation term in (2.10) which allows to pass to the time-continuous limit for fixed  $\varepsilon_n$ .

Next, we associate a ‘crack set’ to the atomistic evolution by collecting the ‘broken springs’. Roughly speaking, in each triangle we draw a line between the centers of two ‘broken springs’. To this end, we let  $\mathcal{T}_n$  be the set of all closed triangles  $\Delta$  contained in  $\mathcal{L}_n(\Omega)$  and consider  $\Delta \in \mathcal{T}_n$  with vertices  $x, x', x''$ . For  $\alpha, \beta, \gamma = 1, 2, 3$  pairwise distinct, we let  $h_\Delta^\alpha$  denote the segment in  $\Delta$  between the centers of the two sides in  $\mathbf{v}_\beta$  and  $\mathbf{v}_\gamma$  direction, see Figure 1. Given  $t \in [t_n^k, t_n^{k+1})$ , we say that  $h_\Delta$  lies in  $\mathcal{K}_\Delta^k$  if  $h_\Delta = h_\Delta^\alpha$  for some  $\alpha = 1, 2, 3$  and  $\min\{M_{x,x'}^\varepsilon((u^j)_{j < k+1}), M_{x',x''}^\varepsilon((u^j)_{j < k+1})\} > R$ , where  $x, x', x''$  are labeled such that  $x - x' = \pm \varepsilon_n \mathbf{v}_\beta$  and  $x'' - x' = \pm \varepsilon_n \mathbf{v}_\gamma$ . Then, for each  $t \in [t_n^k, t_n^{k+1})$  we define

$$K_n(t) := \bigcup_{\Delta \in \mathcal{T}_n} \bigcup_{h_\Delta \in \mathcal{K}_\Delta^k} h_\Delta. \quad (2.12)$$

To formulate the main result, we introduce the *limiting continuum energy*

$$\mathcal{E}(u, K) := \int_\Omega \Phi(\nabla u) \, dx + \int_K \varphi(\nu_K) \, d\mathcal{H}^1, \quad (2.13)$$




FIGURE 1. In the first figure, the segments  $h_\Delta^\alpha, h_\Delta^\beta, h_\Delta^\gamma$  are depicted, where  $\alpha, \beta, \gamma$  always denote pairwise distinct indices in  $\{1, 2, 3\}$ . In the second and third figure, the blue lines denote the ‘broken springs’ and the red line the corresponding ‘crack set’  $\mathcal{K}_\Delta^k$ .

for each  $u \in SBV^2(\Omega)$  and each rectifiable set  $K \subset \mathbb{R}^2$  with  $\mathcal{H}^1(K) < +\infty$ , where  $\nabla u$  denotes the approximate differential and  $S(u)$  the jump set of  $u$ , which is subject to the constraint  $S(u) \tilde{\subset} K$ . (Here and in the following,  $\tilde{\subset}$  stands for inclusions up to  $\mathcal{H}^1$ -negligible sets.) For basic notation and properties of  $SBV$  functions, we refer the reader to [5]. The energy features an *elastic term* with energy density

$$\Phi(z) := \frac{\mu}{\sqrt{3}} \sum_{\mathbf{v} \in \mathcal{V}} |z \cdot \mathbf{v}|^2 \quad \text{for } z \in \mathbb{R}^2, \quad (2.14)$$

resulting from a suitable linearization of the potential  $W$ , and an anisotropic *surface energy* with density

$$\varphi(\nu) := \frac{2\kappa}{\sqrt{3}} \sum_{\mathbf{v} \in \mathcal{V}} |\nu \cdot \mathbf{v}| \quad \text{for } \nu \in \mathbb{S}^1 := \{x \in \mathbb{R}^2 : |x| = 1\} \quad (2.15)$$

where we recall  $\mathcal{V} := \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_2 - \mathbf{v}_1\}$ . Note that the latter has already appeared in the atomistic-to-continuum result [37]. Moreover, by  $AD(g, H)$  we denote all functions  $v \in SBV^2(\Omega)$  such that  $v = g$  on  $\Omega \setminus \overline{U}$  and  $S(v) \tilde{\subset} H$ . In [31] (see [22] for the case of anisotropic densities), the existence of an *irreversible quasi-static crack evolution* with respect to the boundary displacement  $g$  has been shown. This is a mapping  $t \rightarrow (u(t), K(t))$  with  $u(t) \in AD(g(t), K(t))$  for all  $t \in [0, T]$  such that the following three conditions hold:

- (a) *Irreversibility*:  $K(t_1) \tilde{\subset} K(t_2)$  for all  $0 \leq t_1 \leq t_2 \leq T$ .
- (b) *Global stability*: For every  $t \in [0, T]$ , given any  $K(t) \tilde{\subset} H$  and  $v \in AD(g(t), H)$  it holds that

$$\mathcal{E}(u(t), K(t)) \leq \mathcal{E}(v, H). \quad (2.16)$$

- (c) *Energy balance*: The function  $t \mapsto \mathcal{E}(u(t), K(t))$  is absolutely continuous and it holds that

$$\frac{d}{dt} \mathcal{E}(u(t), K(t)) = \int_{\Omega} D\Phi(\nabla u) \cdot \nabla \partial_t g(t) \, dx \quad \text{for a.e. } t \in [0, T], \quad (2.17)$$

where by  $\partial_t$  we denote the time derivative of  $g$ .

Eventually, we need to define a notion of convergence. We define

$$\Omega_n := \bigcup_{\Delta \in \mathcal{T}_n} \Delta. \quad (2.18)$$

For a sequence of discrete displacements  $u_n : \mathcal{L}_n(\Omega) \rightarrow \mathbb{R}$ , we denote by  $\tilde{u}_n : \Omega_n \rightarrow \mathbb{R}$  the interpolation of  $u_n$  which satisfies  $\tilde{u}_n = u_n$  on  $\mathcal{L}_n(\Omega)$  and is affine on each  $\Delta \in \mathcal{T}_n$ .

**Definition 2.1.** Let  $u_n: \mathcal{L}_n(\Omega) \rightarrow \mathbb{R}$  be a sequence of atomistic displacements. We say that  $u_n$  AC-converges to  $u \in SBV^2(\Omega)$  if  $\chi_{\Omega_n} \tilde{u}_n \rightarrow u$  in  $L^1(\Omega)$ .

For the crack sets, we use the notion of  $\sigma$ -convergence recalled in Definition 3.8 below, which is a suitable notion of convergence for sets related to jump sets of  $SBV$  functions. Our main result reads as follows.

**Theorem 2.2** (Atomistic-to-continuum crack growth). *There exists a quasi-static crack growth  $t \rightarrow (u(t), K(t))$  with respect to the boundary condition  $g$  such that, up to a subsequence, we have*

$$K_n(t) \text{ } \sigma\text{-converges to } K(t) \text{ for all } t \in [0, T] \text{ as } n \rightarrow \infty, \quad (2.19)$$

and for each  $t \in [0, T]$  there exists a  $t$ -dependent subsequence  $(n_k)_k$  such that

$$u_{n_k}(t) \text{ AC-converges to } u(t) \text{ as } k \rightarrow \infty. \quad (2.20)$$

Moreover, for all  $t \in [0, T]$  we have

$$\mathcal{E}_n(u_n(t), (u_n^j)_{j < k(t)}) \rightarrow \mathcal{E}(u(t), K(t)) \quad \text{as } n \rightarrow \infty, \quad (2.21)$$

where for each  $n$  the ( $n$ -dependent) index  $k(t)$  is chosen such that  $t \in [t_n^{k(t)}, t_n^{k(t)+1})$ .

*Remark 2.3.* (i) In addition to the convergence of the energy in (2.21), it holds a separate convergence of suitably defined elastic and crack energies, see (4.29).

(ii) Besides the AC-convergence stated in (2.20), also strong convergence of derivatives for suitably defined *jump interpolations* introduced in Section 3.2 holds. We refer to Lemma 3.7 for details.

(iii) Note that the exact choice of the threshold  $R$  does not affect the limiting densities  $\Phi$  and  $\varphi$  in (2.13). However, due to the memory variable on the atomistic level, the limiting quasi-static crack evolution might be sensitive to the choice of  $R$ .

The proof of this result is given in the next three sections and is structured as follows. In Section 3, we reformulate our problem in terms of the triangles of the lattice, introduce a suitable notion of elastic and crack energy on the atomistic level, and define auxiliary interpolations. We also introduce the necessary notions of convergence for crack sets, and prove some preliminary results on compactness and lower semicontinuity. Subsequently, Section 4 contains the proof of Theorem 2.2. The most technical part consists in verifying a *global stability property*, which will be analyzed in detail in Section 5.

### 3. PRELIMINARIES

In this section we collect some preliminaries that are needed for our analysis. First, we will reformulate the energy (2.8) in terms of triangles and introduce an alternative interpolation for atomistic displacements that exhibits jumps relating to the 'atomistic crack set' defined in (2.12). We also recall compactness results for discrete displacements and lower semicontinuity results for elastic energies which are essentially adaptations of statements in [35]. Eventually, we recall two types of convergence of sets and discuss their main properties.

**3.1. Reformulation of the energy in terms of triangles.** We consider the set of all triangles  $\mathcal{T}_n$  and the piecewise affine interpolation  $\tilde{v}_n: \Omega_n \rightarrow \mathbb{R}$  of an atomistic displacement  $v_n: \mathcal{L}_n(\Omega) \rightarrow \mathbb{R}$  introduced below (2.18). We also recall the set  $\mathcal{V} = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_2 - \mathbf{v}_1\}$ , representing the three directions of the triangular lattice. The gradient  $\nabla \tilde{v}_n$  is constant on each  $\triangle \in \mathcal{T}_n$ , and for convenience we denote this value

by  $(v_n)_\Delta \in \mathbb{R}^2$ . The rescaled distance between the two atoms  $x, x' \in \mathcal{L}_n(\Omega)$  in the deformed configuration can then be expressed by

$$\varepsilon_n^{-1/2} |v_n(x) - v_n(x')| = \varepsilon_n^{1/2} \left| (v_n)_\Delta \cdot \frac{x - x'}{\varepsilon_n} \right| = \varepsilon_n^{1/2} |(v_n)_\Delta \cdot \mathbf{v}|,$$

where  $\mathbf{v} \in \mathcal{V}$  is the unit vector parallel to  $x - x'$ . In fact, we can find a pair  $(\Delta, \mathbf{v}) \in \mathcal{T}_n \times \mathcal{V}$ , which unambiguously corresponds to a nearest-neighbor pair  $\{x, x'\}$ . More precisely, there is a surjective mapping  $\Xi : \mathcal{T}_n \times \mathcal{V} \times \{-1, 1\} \rightarrow \text{NN}_n(\Omega)$ , where  $\Xi(\Delta, \mathbf{v}, s)$  is given by the unique pair  $(x, x') \in \text{NN}_n(\Omega)$  such that  $x, x'$  are vertices of  $\Delta$  and  $x - x' = s\mathbf{v}$ . Note, however, except for springs at  $\partial\Omega_n$ , each pair  $(x, x') \in \text{NN}_n(\Omega)$  is associated to two different triangles.

Next, we express the memory variable in terms of  $(\Delta, \mathbf{v}) \in \mathcal{T}_n \times \mathcal{V}$ . To this end, given the time discretization  $\{0 = t_n^0 < t_n^1 < \dots < t_n^{T/\delta_n} = T\}$  of the interval  $[0, T]$  with step size  $\delta_n$ , we suppose that for  $t_n^k \in [0, T]$  the displacements  $(u_n^j)_{j < k}$  at previous time steps  $(t_n^j)_{j < k}$  have already been found. Then, recalling (2.7), we define

$$M_{n,\Delta}^{k,\mathbf{v}} := M_{x,x'}^{\varepsilon_n}((u_n^j)_{j < k}) = \varepsilon_n^{-1/2} \sup_{j < k} |u_n^j(x) - u_n^j(x')|. \quad (3.1)$$

In the sequel, we frequently need to keep track of the memory variable including competitors in the next time step. To this end, for an arbitrary displacement  $v_n : \mathcal{L}_n(\Omega) \rightarrow \mathbb{R}$  we define

$$M_{n,\Delta}^{k,\mathbf{v}}(v_n) := M_{n,\Delta}^{k,\mathbf{v}} \vee \varepsilon_n^{1/2} |(v_n)_\Delta \cdot \mathbf{v}| \quad (3.2)$$

which includes the impact of  $v_n$  on the spring in  $\Delta$  pointing in direction  $\mathbf{v} \in \mathcal{V}$  into the memory variable.

We now use this to define the notion of a 'broken triangle'. We regard a triangle  $\Delta \in \mathcal{T}_n$  as 'broken' if at least one edge has been "broken", i.e., stretched above the threshold  $R$ . This means, for a given displacement history  $(u_n^j)_{j < k}$  and an arbitrary displacement  $v_n : \mathcal{L}_n(\Omega) \rightarrow \mathbb{R}$ , we define

$$\begin{aligned} \mathcal{C}_n^k &:= \{ \Delta \in \mathcal{T}_n : M_{n,\Delta}^{k,\mathbf{v}} > R \text{ for at least one } \mathbf{v} \in \mathcal{V} \}, \\ \mathcal{C}_n^k(v_n) &:= \{ \Delta \in \mathcal{T}_n : M_{n,\Delta}^{k,\mathbf{v}}(v_n) > R \text{ for at least one } \mathbf{v} \in \mathcal{V} \}. \end{aligned} \quad (3.3)$$

Note that, whenever a triangle is not broken, i.e.,  $\Delta \notin \mathcal{C}_n^k(v_n)$ , we have by (3.2) that

$$|(v_n)_\Delta| = \left( |(v_n)_\Delta \cdot \mathbf{v}_1|^2 + |(v_n)_\Delta \cdot \frac{1}{\sqrt{3}}(\mathbf{v}_2 + \mathbf{v}_3)|^2 \right)^{1/2} \leq 2R\varepsilon_n^{-1/2}. \quad (3.4)$$

Now, we can express the energy (2.8) at time  $t_n^k$  in terms of triangles and their corresponding edges. In fact, we can separate the 'broken' triangles  $\mathcal{C}_n^k(v_n)$  from the 'intact triangles'  $\mathcal{T}_n \setminus \mathcal{C}_n^k(v_n)$ , and obtain

$$\begin{aligned} \mathcal{E}_n(v_n; (u_n^j)_{j < k}) &= \frac{\varepsilon_n}{2} \sum_{\Delta \in \mathcal{T}_n \setminus \mathcal{C}_n^k(v_n)} \sum_{\mathbf{v} \in \mathcal{V}} \Psi(\varepsilon_n^{1/2} |(v_n)_\Delta \cdot \mathbf{v}|) + \frac{\varepsilon_n}{2} \sum_{\Delta \in \mathcal{C}_n^k(v_n)} \sum_{\substack{\mathbf{v} \in \mathcal{V} \\ M_{n,\Delta}^{k,\mathbf{v}}(v_n) \leq R}} \Psi(\varepsilon_n^{1/2} |(v_n)_\Delta \cdot \mathbf{v}|) + \\ &+ \frac{\varepsilon_n}{2} \sum_{\Delta \in \mathcal{C}_n^k(v_n)} \sum_{\substack{\mathbf{v} \in \mathcal{V} \\ M_{n,\Delta}^{k,\mathbf{v}}(v_n) > R}} \Psi(M_{n,\Delta}^{k,\mathbf{v}}(v_n)) + \mathcal{E}_n^{k,\text{bdy}}(v_n), \end{aligned} \quad (3.5)$$

where  $\mathcal{E}_n^{k,\text{bdy}}$  contains the energy contributions of nearest-neighbor pairs  $(x, x')$  that are part of only one triangle and consequently only reflected with a prefactor 1/2 in the other terms of (3.5). More precisely,

we have

$$\mathcal{E}_n^{k,\text{bdy}}(v_n) = \frac{\varepsilon_n}{2} \sum_{\substack{(x,x') \in \text{NN}_n(\Omega), \\ x,x' \in \partial\Omega_n}} E_{x,x'}^{\varepsilon_n}(v_n, (u_n^j)_{j < k}), \quad (3.6)$$

where we recall the definition in (2.18). Note that the prefactor  $\frac{1}{2}$  appears again, because in the above sums all springs that do not lie at the boundary are counted twice, being associated to two different triangles. For lower bounds, we can ignore  $\mathcal{E}_n^{\text{bdy}}$  as it is clearly nonnegative. For upper bounds, however, we will need to show explicitly that the term vanishes in the limit  $n \rightarrow \infty$ , see Theorem 4.7 for details.

From now on, we simplify the notation in the following sense, whenever no confusion arises: instead of writing  $\Delta \in \mathcal{T}'_n$  below sums for a subset  $\mathcal{T}'_n \subset \mathcal{T}_n$ , we simply write  $\mathcal{T}'_n$  indicating that the sum is taken over all triangles in  $\mathcal{T}'_n$ . In a similar fashion, in sums over  $\mathcal{V}$  involving  $M_{n,\Delta}^{k,\mathbf{v}}(v_n)$ , we do not include  $\mathbf{v} \in \mathcal{V}$  in the notation.

In (3.5), we have split the energy in terms of intact and broken triangles. We further split the broken triangles in a crack part, which will lead to jumps in the limiting description as  $n \rightarrow \infty$ , and a remainder term. First, we choose a sequence of diverging thresholds  $(R_n)_n$  satisfying

$$R_n \rightarrow +\infty \quad \text{and} \quad R_n^{3/2} \varepsilon_n^{1/4} \rightarrow 0, \quad (3.7)$$

and consider the set of triangles

$$\mathcal{C}_n^{k,\text{L}}(v_n) := \{\Delta \in \mathcal{C}_n^k(v_n) : M_{n,\Delta}^{k,\mathbf{v}}(v_n) > 2R_n \text{ for at least one } \mathbf{v} \in \mathcal{V}\}. \quad (3.8)$$

Here, the superscript L indicates that the memory variable is ‘large’. By elementary geometry this implies  $M_{n,\Delta}^{k,\mathbf{v}}(v_n) > R_n$  for at least two  $\mathbf{v} \in \mathcal{V}$ , and we split the set into

$$\mathcal{C}_n^{k,\text{L}}(v_n) = \mathcal{C}_n^{k,2}(v_n) \dot{\cup} \mathcal{C}_n^{k,3}(v_n), \quad (3.9)$$

corresponding to triangles where exactly two or three  $\mathbf{v} \in \mathcal{V}$  satisfy  $M_{n,\Delta}^{k,\mathbf{v}}(v_n) > R_n$ . Then, we can write

$$\mathcal{E}_n(v_n; (u_n^j)_{j < k}) = \mathcal{E}_n^{k,\text{ela}}(v_n) + \mathcal{E}_n^{k,\text{cra}}(v_n) + \mathcal{E}_n^{k,\text{rem}}(v_n) + \mathcal{E}_n^{k,\text{bdy}}(v_n), \quad (3.10)$$

where  $\mathcal{E}_n^{k,\text{bdy}}$  is defined in (3.6), and we set

$$\begin{aligned} \mathcal{E}_n^{k,\text{ela}}(v_n) &= \frac{\varepsilon_n}{2} \sum_{\mathcal{T}_n \setminus \mathcal{C}_n^k(v_n)} \sum_{\mathbf{v} \in \mathcal{V}} \Psi(\varepsilon_n^{1/2} |(v_n)_\Delta \cdot \mathbf{v}|), \\ \mathcal{E}_n^{k,\text{cra}}(v_n) &= \frac{\varepsilon_n}{2} \sum_{\mathcal{C}_n^{k,\text{L}}(v_n)} \sum_{M_{n,\Delta}^{k,\mathbf{v}}(v_n) > R_n} \Psi(M_{n,\Delta}^{k,\mathbf{v}}(v_n)), \\ \mathcal{E}_n^{k,\text{rem}}(v_n) &= \frac{\varepsilon_n}{2} \sum_{\mathcal{C}_n^k(v_n)} \left( \sum_{R < M_{n,\Delta}^{k,\mathbf{v}}(v_n) \leq R_n} \Psi(M_{n,\Delta}^{k,\mathbf{v}}(v_n)) + \sum_{M_{n,\Delta}^{k,\mathbf{v}}(v_n) \leq R} \Psi(\varepsilon_n^{1/2} |(v_n)_\Delta \cdot \mathbf{v}|) \right) \\ &\quad + \frac{\varepsilon_n}{2} \sum_{\mathcal{C}_n^k(v_n) \setminus \mathcal{C}_n^{k,\text{L}}(v_n)} \sum_{M_{n,\Delta}^{k,\mathbf{v}}(v_n) > R_n} \Psi(M_{n,\Delta}^{k,\mathbf{v}}(v_n)). \end{aligned} \quad (3.11)$$

We will see that the contributions of the *remainder term*  $\mathcal{E}_n^{k,\text{rem}}(v_n)$  will eventually vanish in the limit  $n \rightarrow \infty$ . Still, from a technical point of view, we need delicate arguments to control its contribution in all estimates below. Our next goal is to rewrite the *elastic energy*  $\mathcal{E}_n^{k,\text{ela}}(v_n)$  and the *crack energy*  $\mathcal{E}_n^{k,\text{cra}}(v_n)$  as an integral functional which will allow us to apply lower semicontinuity results in Sobolev and *SBV* spaces. This will be achieved by introducing a jump interpolation which is inspired by the one introduced in [35].




FIGURE 2. The first picture shows the sets  $V_\Delta^\alpha$  and  $h_\Delta^\alpha$  which are used for the definition of  $K_n^{k,i}$ . The second one depicts the subdivision of a triangle into sub-triangles.

**3.2. Jump interpolation and representation of the energy.** We now define (three versions of) a ‘crack set’ related to  $\mathcal{C}_n^{k,L}(v_n)$ . To this end, as before, given  $\Delta \in \mathcal{T}_n$  and  $\alpha, \beta, \gamma = 1, 2, 3$  pairwise distinct, we let  $h_\Delta^\alpha$  denote the segment in  $\Delta$  between the centers of the both sides in direction  $\mathbf{v}_\beta$  and  $\mathbf{v}_\gamma$ , see Figure 1. Note that  $h_\Delta^\alpha$  is parallel to  $\mathbf{v}_\alpha$ . We define the set  $V_\Delta^\alpha = h_\Delta^\beta \cup h_\Delta^\gamma$ , see Figure 2, and for each  $i = 1, 2, 3$ , we let

$$K_n^{k,i}(v_n) := \bigcup_{\Delta \in \mathcal{C}_n^{k,3}(v_n)} V_\Delta^i \cup \bigcup_{\Delta \in \mathcal{C}_n^{k,2}(v_n)} h_\Delta, \quad K_n^{k,L}(v_n) := \bigcup_{i=1}^3 K_n^{k,i}(v_n), \quad (3.12)$$



where for each  $\Delta \in \mathcal{C}_n^{k,2}(v_n)$  we choose  $h_\Delta = h_\Delta^m$ , where  $m$  is the unique index with  $M_{n,\Delta}^{k,\mathbf{v}_m}(v_n) \leq R_n$ . The reason for using three different variants lies in the derivation of a lower semicontinuity result for the length of the sets, and follows the ideas in [35]. Note carefully that on triangles  $\Delta \in \mathcal{C}_n^{k,2}(v_n)$  all variants of  $K_n^{k,i}(v_n)$  coincide.

*Remark 3.1.* Recalling (2.12), one can check that the solution  $u_n^k$  at time step  $t_n^k$ , see (2.10), fulfills

$$K_n^{k,L}(u_n^k) \subset K_n(t) \quad \text{for } t \in [t_n^k, t_n^{k+1}),$$

and the difference of the sets is due to the triangles  $\Delta \in \mathcal{C}_n^k(u_n^k)$  where the numbers  $\mu_{n,\Delta}^k = \#\{\mathbf{v} \in \mathcal{V} : M_{n,\Delta}^{k,\mathbf{v}}(u_n^k) > R\}$  and  $\nu_{n,\Delta}^k = \#\{\mathbf{v} \in \mathcal{V} : M_{n,\Delta}^{k,\mathbf{v}}(u_n^k) > R_n\}$  differ. Indeed, if  $\mu_{n,\Delta}^k = \nu_{n,\Delta}^k = 3$ , then both  $K_n(t)$  and  $K_n^{k,L}(u_n^k)$  coincide with  $\bigcup_{i=1}^3 h_\Delta^i$  on  $\Delta$ . If  $\mu_{n,\Delta}^k = \nu_{n,\Delta}^k = 2$ , then there exists one spring in direction  $\mathbf{v}_i$  such that  $M_{n,\Delta}^{k,\mathbf{v}_i}(u_n^k) \leq R$ , and both  $K_n(t)$  and  $K_n^{k,L}(u_n^k)$  coincide with  $h_\Delta^i$  on  $\Delta$ . Along the proof, we will see that the difference of  $K_n(t)$  and  $K_n^{k,L}(u_n^k)$  is asymptotically vanishing for  $n \rightarrow \infty$ .

We now introduce a jump interpolation  $\hat{v}_n^k : \Omega_n \rightarrow \mathbb{R}$ , that exhibits jumps on triangles  $\Delta \in \mathcal{C}_n^k(v_n)$  and leaves  $\tilde{v}_n$  unchanged on  $\mathcal{T}_n \setminus \mathcal{C}_n^k(v_n)$ . On  $\mathcal{T}_n \setminus \mathcal{C}_n^k(v_n)$ , we simply set  $\hat{v}_n^k = \tilde{v}_n$ . Next, we consider triangles  $\Delta \in \mathcal{C}_n^{k,L}(v_n)$ . We divide  $\Delta$  in four equilateral sub-triangles, where we denote by  $\Delta_\alpha$  the triangle that is formed by taking  $h_\Delta^\alpha$  as one side, and the middle triangle with sides  $h_\Delta^\alpha$ ,  $h_\Delta^\beta$ , and  $h_\Delta^\gamma$  is denoted by  $\Delta_{\text{mid}}$ , see Figure 2. Now, if  $\Delta \in \mathcal{C}_n^{k,3}(v_n)$ , we define  $\hat{v}_n^k$  such that  $\hat{v}_n^k = v_n$  on the three vertices of  $\Delta$  and  $\nabla \hat{v}_n^k \equiv 0$  on  $\bigcup_{i=1}^3 \Delta_i$ , see Figure 3. On  $\Delta_{\text{mid}}$ , let  $\hat{v}_n^k$  be constant, attaining an arbitrary value. For  $\Delta \in \mathcal{C}_n^{k,2}(v_n)$ , assuming  $M_{n,\Delta}^{k,\mathbf{v}_\alpha}(v_n) \leq R_n$ , we define  $\hat{v}_n^k$  in such a way that  $\hat{v}_n^k$  is continuous on  $\text{int}(\Delta) \setminus h_\Delta^\alpha$ ,  $\nabla \hat{v}_n^k \equiv 0$  in  $\Delta$ , and  $\hat{v}_n^k$  coincides with  $v_n$  on two of the vertices, see Figure 4 (we choose one of the two options.) For  $\Delta \in \mathcal{C}_n^k(v_n) \setminus \mathcal{C}_n^{k,L}(v_n)$ , we define  $\hat{v}_n^k$  such that  $\hat{v}_n^k$  is constant on  $\text{int}(\Delta)$  and coincides with  $v_n$  in one of


 FIGURE 3. Construction of  $\hat{v}_n^k$  for a triangle  $\Delta \in \mathcal{C}_n^{k,3}(v_n)$  with three broken springs.

 FIGURE 4. Construction of  $\hat{v}_n^k$  in a triangle  $\Delta \in \mathcal{C}_n^{k,2}(v_n)$  with  $M_{n,\Delta}^{k,v_\alpha}(v_n) \leq R_n$ .

the three vertices (we choose one of the three options). Note that the interpolation exhibits jumps also at the boundary of broken triangles, both on  $\mathcal{C}_n^{k,L}(v_n)$  and on  $\mathcal{C}_n^k(v_n) \setminus \mathcal{C}_n^{k,L}(v_n)$ . We cover this set by

$$K_n^{k,S}(v_n) := \bigcup_{\Delta \in \mathcal{C}_n^k(v_n)} \partial\Delta, \quad (3.13)$$

where we use the superscript S to indicate that, as we will see in Remark 3.3 below, the jump height on this set is small. We also observe that, in view of (3.4), we have

$$\|\nabla \hat{v}_n^k\|_\infty \leq 2R\varepsilon_n^{-1/2}. \quad (3.14)$$

In fact, either  $\nabla \hat{v}_n^k = 0$ , namely on broken triangles, or we have  $\nabla \hat{v}_n^k = (v_n)_\Delta$  constant on a triangle  $\Delta$ .

We can now use this interpolation to obtain a convenient representation of the elastic energy. By the construction of  $\hat{v}_n^k$ , we can take the sum over all  $\mathcal{T}_n$  in the elastic energy, see (3.11), and write this term as an integral over  $\Omega_n$ . Precisely, we have

$$\mathcal{E}_n^{k,\text{ela}}(v_n) = \frac{4}{\sqrt{3}\varepsilon_n} \int_{\Omega_n} \frac{1}{2} \sum_{\mathbf{v} \in \mathcal{V}} \Psi(\varepsilon_n^{1/2} |\nabla \hat{v}_n^k(x) \cdot \mathbf{v}|) \, dx = \int_{\Omega_n} \Psi_n^{\text{cell}}(\nabla \hat{v}_n^k(x)) \, dx, \quad (3.15)$$

where we used that the area of each triangle is  $\frac{\sqrt{3}}{4}\varepsilon_n^2$  and we defined the *cell energy density*  $\Psi_n^{\text{cell}}$  by

$$\Psi_n^{\text{cell}}(z) := \frac{2}{\sqrt{3}\varepsilon_n} \sum_{\mathbf{v} \in \mathcal{V}} \Psi(\varepsilon_n^{1/2} |z \cdot \mathbf{v}|) \quad \text{for all } z \in \mathbb{R}^2. \quad (3.16)$$

Concerning the crack energy defined in (3.11), a crucial point in our analysis is the characterization of  $\mathcal{E}_n^{k,\text{cra}}(v_n)$  in terms of the sets  $K_n^{k,i}(v_n)$  introduced in (3.12). To formulate this relation, we define

$\varphi_\alpha(\nu) := (|\mathbf{v}_\beta \cdot \nu| + |\mathbf{v}_\gamma \cdot \nu|)$  for  $\alpha, \beta, \gamma \in \{1, 2, 3\}$  pairwise distinct and for  $\nu \in \mathbb{S}^1$ . Also recall the definition of  $\kappa$  in (iv).

**Proposition 3.2.** *There exists a universal constant  $C > 0$  such that, for given displacements  $(u_n^j)_{j < k}$  and an arbitrary  $v_n: \mathcal{L}_n(\Omega) \rightarrow \mathbb{R}$ , the crack energy  $\mathcal{E}_n^{k, \text{cra}}(v_n)$  satisfies*

$$\left| \mathcal{E}_n^{k, \text{cra}}(v_n) - \frac{\kappa}{\sqrt{3}} \sum_{i=1}^3 \int_{K_n^{k,i}(v_n)} \varphi_i(\nu_K) d\mathcal{H}^1 \right| \leq C |\kappa - \Psi(R_n)| \varepsilon_n \# \mathcal{C}_n^{k,L}(v_n), \quad (3.17)$$

where  $\nu_K$  denotes a unit normal to  $K_n^{k,i}(v_n)$ .

*Remark 3.3.* (i) In view of (3.7) and the properties of  $\Psi$ , the right-hand side of (3.17) will vanish for  $n \rightarrow \infty$  as we will indeed have a bound on  $\# \mathcal{C}_n^k(v_n)$  of the order  $\varepsilon_n^{-1}$ .

(ii) One can check that the interpolation  $\hat{v}_n^k$  and the crack set are related in the sense that  $S(\hat{v}_n^k) \subset K_n^{k,L}(v_n) \cup K_n^{k,S}(v_n)$  with

$$|\hat{v}_n^k(x)| \leq C \varepsilon_n^{1/2} R_n \quad \text{for all } x \in K_n^{k,S}(v_n). \quad (3.18)$$

Indeed, the inclusion follows directly from the construction of  $\hat{v}_n^k$ . Property (3.18) can be derived by using the continuity of  $\hat{v}_n^k$  at (some of) the vertices of the triangles along with (3.14), and the fact that for every pair  $x, x' \in \Delta$  such that the segment between  $x$  and  $x'$  does not intersect  $K_n^{k,L}(v_n)$  we have  $\varepsilon_n^{-1/2} |v_n(x) - v_n(x')| \leq 2R_n$ , see (3.8).

(iii) The choice in (3.7) also shows that  $R_n \varepsilon_n^{1/2} \rightarrow 0$  and hence (3.18) implies that the jumps in  $K_n^{k,S}$  are asymptotically of vanishing height. (This is the reason why this set is indicated with label ‘small’, see below (3.13).) Therefore, these jumps, artificially introduced by our interpolation, will not affect the continuum limit.

*Proof of Proposition 3.2.* The proof consists in showing that

$$\mathcal{G}_n^{k, \text{cra}}(v_n) := \kappa \frac{\varepsilon_n}{2} (2\# \mathcal{C}_n^{k,2}(v_n) + 3\# \mathcal{C}_n^{k,3}(v_n)) = \frac{\kappa}{\sqrt{3}} \sum_{i=1}^3 \int_{K_n^{k,i}(v_n)} \varphi_i(\nu_K) d\mathcal{H}^1. \quad (3.19)$$

Recalling (3.9), (3.11), and (ii'), the statement then follows from the fact that

$$\mathcal{E}_n^{k, \text{cra}}(v_n) \geq \Psi(R_n) \frac{\varepsilon_n}{2} (2\# \mathcal{C}_n^{k,2}(v_n) + 3\# \mathcal{C}_n^{k,3}(v_n)),$$

and therefore

$$|\mathcal{G}_n^{k, \text{cra}}(v_n) - \mathcal{E}_n^{k, \text{cra}}(v_n)| \leq C |\kappa - \Psi(R_n)| \varepsilon_n \# \mathcal{C}_n^{k,L}(v_n).$$

To show (3.19), we will consider the sets  $\mathcal{C}_n^{k,2}(v_n)$  and  $\mathcal{C}_n^{k,3}(v_n)$  separately.

Firstly, consider  $\Delta \in \mathcal{C}_n^{k,3}(v_n)$ . As before, we denote by  $h_\Delta^\alpha$  the segment between the midpoints of the sides in directions  $\mathbf{v}_\beta, \mathbf{v}_\gamma \in \mathcal{V}$  for pairwise distinct  $\alpha, \beta, \gamma \in \{1, 2, 3\}$ . We express the contribution inside  $\Delta$  as an integral. Recall that  $\mathcal{H}^1(h_\Delta^\alpha) = \frac{\varepsilon_n}{2}$  for all  $\alpha = 1, 2, 3$ , and note that for the normal  $\nu_\alpha$  of  $h_\Delta^\alpha$  we have  $|\mathbf{v}_\alpha \cdot \nu_\alpha| = 0$ , and  $|\mathbf{v}_\beta \cdot \nu_\alpha| = |\mathbf{v}_\gamma \cdot \nu_\alpha| = \frac{\sqrt{3}}{2}$ . Given  $\alpha = 1, 2, 3$ , the set  $h_\Delta^\beta \cup h_\Delta^\gamma = V_\Delta^\alpha$  is exactly the part of  $K_n^{k,\alpha}(v_n)$  inside  $\Delta$ . Denoting the normal to  $K_n^{k,\alpha}(v_n)$  again by  $\nu_K$ , we can rewrite

$$\frac{3}{2} = \frac{1}{\sqrt{3}\varepsilon_n} \sum_{\alpha=1}^3 \int_{K_n^{k,\alpha}(v_n) \cap \text{int}(\Delta)} (|\mathbf{v}_\beta \cdot \nu_K| + |\mathbf{v}_\gamma \cdot \nu_K|) d\mathcal{H}^1 = \frac{1}{\sqrt{3}\varepsilon_n} \sum_{\alpha=1}^3 \int_{K_n^{k,\alpha}(v_n) \cap \text{int}(\Delta)} \varphi_\alpha(\nu_K) d\mathcal{H}^1. \quad (3.20)$$

Now consider  $\Delta \in C_n^{k,2}(v_n)$ . We choose  $\mathbf{v}_m \in \mathcal{V}$  with  $M_{n,\Delta}^{k,\mathbf{v}_m}(v_n) \leq R_n$ . Then, by construction  $K_n^{k,\alpha}(v_n) \cap \text{int}(\Delta)$  coincides with  $h_\Delta^m$  for each  $\alpha = 1, 2, 3$ . Therefore, we have  $\mathcal{H}^1(K_n^{k,\alpha}(v_n) \cap \text{int}(\Delta)) = \frac{\varepsilon_n}{2}$  for  $\alpha = 1, 2, 3$  and, since  $|\nu_K \cdot \mathbf{v}_m| = 0$  and  $|\nu_K \cdot \mathbf{v}_\alpha| = \frac{\sqrt{3}}{2}$  for  $\alpha \in \{1, 2, 3\} \setminus \{m\}$ , we can check that

$$\frac{1}{\sqrt{3}\varepsilon_n} \sum_{\alpha=1}^3 \int_{K_n^{k,\alpha}(v_n) \cap \text{int}(\Delta)} \varphi_\alpha(\nu_K) d\mathcal{H}^1 = \frac{1}{\sqrt{3}\varepsilon_n} \sum_{\alpha=1}^3 \int_{K_n^{k,\alpha}(v_n) \cap \text{int}(\Delta)} (|\mathbf{v}_\beta \cdot \nu_K| + |\mathbf{v}_\gamma \cdot \nu_K|) d\mathcal{H}^1 = 1. \quad (3.21)$$

Collecting (3.20)–(3.21) we find (3.19).  $\square$

**3.3. Compactness and lower semicontinuity of elastic energies.** In this subsection, we present a compactness result for sequences with bounded energy  $\mathcal{E}_n$  (see (3.5)) with respect to the notion of AC-convergence introduced in Definition 2.1. Afterwards, we present a lower semicontinuity result for the elastic energy. We start with elementary properties of  $\Psi_n^{\text{cell}}$  defined in (3.16) and  $\Phi$  given in (2.14).

**Lemma 3.4** (Properties of  $\Psi_n^{\text{cell}}$ ). (i) *There exists a function  $\omega : \mathbb{R}^2 \rightarrow \mathbb{R}$  with  $\frac{\omega(y)}{|y|^2} \rightarrow 0$  for  $|y| \rightarrow 0$  such that*

$$\Psi_n^{\text{cell}}(y) = \Phi(y) + \varepsilon_n^{-1} \omega(\varepsilon_n^{1/2} y) \quad \text{for all } y \in \mathbb{R}^2 \text{ and all } n \in \mathbb{N}.$$

(ii) *Let  $r > 0$ . Then, there exists a constant  $c_r > 0$  depending on  $r$  such that for all  $n \in \mathbb{N}$  we have*

$$c_r |y|^2 \leq \Psi_n^{\text{cell}}(y), \quad c_r |y| \leq |D\Psi_n^{\text{cell}}(y)| \quad \text{for all } y \in \mathbb{R}^2 \text{ with } |y| \leq r\varepsilon_n^{-1/2}.$$

(iii) *Then, there exists a constant  $C > 0$  such that for all  $n \in \mathbb{N}$  we have*

$$|\Psi_n^{\text{cell}}(y)| \leq C|y|^2, \quad |D\Psi_n^{\text{cell}}(y)| \leq C|y| \quad \text{for all } y \in \mathbb{R}^2.$$

(iv) *For each  $M > 0$  and sequences  $(y_n^1)_n, (y_n^2)_n \subset \mathbb{R}^2$  with  $|y_n^1| \leq M$ ,  $|y_n^2| \leq M$ , and  $|y_n^1 - y_n^2| \rightarrow 0$ , it holds that  $|D\Psi_n^{\text{cell}}(y_n^1) - D\Psi_n^{\text{cell}}(y_n^2)| \rightarrow 0$  as  $n \rightarrow \infty$ .*

*Proof.* We start with a preliminary computation. Recalling the definition  $\Psi(s) = W(\sqrt{1+s^2})$  for  $s \in \mathbb{R}$  and the fact that  $W$  is minimized at 1 with  $W(1) = 0$  and  $\lim_{r \searrow 1} W'(r) = \mu$ , a Taylor expansion yields

$$\Psi(s) = W(\sqrt{1+s^2}) = W(1) + \mu \frac{s^2}{2} + \mathcal{O}(s^3), \quad (3.22)$$

as  $s \rightarrow 0$ , and similarly

$$\Psi'(s) = \mu s + \mathcal{O}(s^2). \quad (3.23)$$

Thus, by (3.22) there exists  $\eta > 0$  such that

$$\Psi(s) \geq \frac{\mu}{4} s^2, \quad \Psi'(s) \geq \frac{\mu}{2} s \quad \text{for all } s \in \mathbb{R} \text{ with } |s| \leq \eta. \quad (3.24)$$

First, putting together (3.22) and (3.16), we get

$$\Psi_n^{\text{cell}}(y) := \frac{\mu}{\sqrt{3}} \sum_{\mathbf{v} \in \mathcal{V}} |y \cdot \mathbf{v}|^2 + \varepsilon_n^{-1} \sum_{\mathbf{v} \in \mathcal{V}} \mathcal{O}((\varepsilon_n^{1/2} |y \cdot \mathbf{v}|)^3), \quad (3.25)$$

and thus (i) is an immediate consequence of the definition of  $\Phi$  in (2.14).

We now come to the proof of (ii). Recalling that  $\mathbf{v}_1 = (1, 0)^T$ ,  $\mathbf{v}_2 = (\frac{1}{2}, \frac{\sqrt{3}}{2})$ , and  $\mathbf{v}_3 = (-\frac{1}{2}, \frac{\sqrt{3}}{2})$  we calculate for each  $y \in \mathbb{R}^2$

$$y_1^2 = |y \cdot \mathbf{v}_1|^2, \quad y_2^2 = \frac{1}{3} |y \cdot (\mathbf{v}_2 + \mathbf{v}_3)|^2 \leq \frac{2}{3} (|y \cdot \mathbf{v}_2|^2 + |y \cdot \mathbf{v}_3|^2),$$

which yields  $|y|^2 \leq C(|y \cdot \mathbf{v}_1|^2 + |y \cdot \mathbf{v}_2|^2 + |y \cdot \mathbf{v}_3|^2)$  for a constant  $C > 0$ .

Now let  $y \in \mathbb{R}^2$  with  $|y| \leq r\varepsilon_n^{-1/2}$ . Without restriction we assume that  $\eta \leq r$  and set  $y' = \frac{\eta}{r}y$ . We then have  $|\varepsilon_n^{1/2}y' \cdot \mathbf{v}_i| \leq \eta$  for all  $i \in \{1, 2, 3\}$ , which by (3.24) yields  $|\varepsilon_n^{1/2}y' \cdot \mathbf{v}_i|^2 \leq C\Psi(|\varepsilon_n^{1/2}y' \cdot \mathbf{v}_i|)$  for a constant  $C > 0$ . Then, the previous estimate along with the definition of  $\Psi_n^{\text{cell}}$  gives us

$$|y'|^2 \leq C\varepsilon_n^{-1}(|\varepsilon_n^{1/2}y' \cdot \mathbf{v}_1|^2 + |\varepsilon_n^{1/2}y' \cdot \mathbf{v}_2|^2 + |\varepsilon_n^{1/2}y' \cdot \mathbf{v}_3|^2) \leq C\Psi_n^{\text{cell}}(y') \leq C\Psi_n^{\text{cell}}(y),$$

where the last step follows from the fact that  $\Psi$  is increasing, see (ii'). This shows the first part of (ii) for a constant depending on  $r$ . The second part of (ii) can be shown along similar lines, using the second estimate in (3.24).

Now, we prove (iii). We compute

$$D\Psi_n^{\text{cell}}(y) = \frac{2}{\sqrt{3}}\varepsilon_n^{-1/2} \sum_{\mathbf{v} \in \mathbf{V}} \Psi'(\varepsilon_n^{1/2}|y \cdot \mathbf{v}|) \frac{y \cdot \mathbf{v}}{|y \cdot \mathbf{v}|} \mathbf{v} \quad (3.26)$$

for all  $y \in \mathbb{R}^2$ . Thus, using (3.23), there exist  $C > 0$  and  $\eta > 0$  such that  $|D\Psi_n^{\text{cell}}(y)| \leq C|y|$  for all  $y \in \mathbb{R}^2$  with  $|y| \leq \eta\varepsilon_n^{-1/2}$ . If instead  $\eta\varepsilon_n^{-1/2} < |y|$ , we can use  $\Psi' \in L^\infty([0, \infty))$  by the Lipschitz continuity of  $\Psi$  to derive that

$$|D\Psi_n^{\text{cell}}(y)| \leq C\varepsilon_n^{-1/2} = C(|y|^{-1}\varepsilon_n^{-1/2})|y| \leq C|y|.$$

The bound on  $\Psi_n^{\text{cell}}$  follows in a similar way, using (3.16), (3.25), and  $\Psi \in L^\infty([0, \infty))$ . Eventually, (iv) follows directly from (3.23) and (3.26). This concludes the proof.  $\square$

We now proceed with the compactness result.

**Proposition 3.5** (Compactness). *Let  $(v_n)_n$  be a sequence of displacements with  $v_n \in \mathcal{A}(g_n)$  for a sequence of boundary conditions  $(g_n)_n$  such that  $g_n$  AC-converges to some  $g \in W^{2,\infty}(\Omega)$ . For each  $n \in \mathbb{N}$ , let  $(u_n^j)_{j < k_n}$  be a displacement history for  $k_n \in \mathbb{N}$  depending on  $n$ . Suppose that  $\mathcal{E}_n(v_n; (u_n^j)_{j < k_n}) + \|\tilde{v}_n\|_{L^\infty(\Omega_n)} \leq M$  for some  $M > 0$ . Then, there exists  $C > 0$  only depending on  $M$  such that*

$$\#\mathcal{C}_n^{k_n}(v_n) \leq \frac{C}{\varepsilon_n} \quad \text{and} \quad \mathcal{H}^1(S(\hat{v}_n^{k_n})) \leq C. \quad (3.27)$$

Moreover, there exists a function  $v \in SBV^2(\Omega)$  with  $v = g$  on  $\Omega \setminus \bar{U}$  such that, up to a subsequence (not relabeled),  $v_n$  AC-converges to  $v$  and for the jump interpolations  $\hat{v}_n^{k_n}$  we have

$$\chi_{\Omega_n} \hat{v}_n^{k_n} \rightarrow v \text{ in } L^1(\Omega) \quad \text{and} \quad \chi_{\Omega_n} \nabla \hat{v}_n^{k_n} \rightharpoonup \nabla v \text{ in } L^2(\Omega; \mathbb{R}^2). \quad (3.28)$$

*Proof.* For convenience, we simply write  $k$  in place of  $k_n$  in the proof. In view of (3.5) and (ii'), we have

$$\frac{\varepsilon_n}{2} \#\mathcal{C}_n^k(v_n) \Psi(R) \leq \mathcal{E}_n(v_n; (u_n^j)_{j < k_n}) \leq M < +\infty.$$

Hence, there exists some  $C > 0$  such that for all  $\varepsilon_n > 0$  it holds that

$$\#\mathcal{C}_n^k(v_n) \leq \frac{C}{\varepsilon_n}.$$

Recalling that by construction  $\hat{v}_n^k$  jumps only on triangles  $\Delta \in \mathcal{C}_n^k(v_n)$  and we always have  $S(\hat{v}_n^k) \cap \bar{\Delta} \leq \frac{9}{2}\varepsilon_n$ , we conclude the proof of (3.27). The bound (3.14) along with Lemma 3.4(ii) for  $r = 2R$  and (3.15) shows

$$\|\nabla \hat{v}_n^k\|_{L^2(\Omega_n)}^2 = \int_{\Omega_n} |\nabla \hat{v}_n^k(x)|^2 dx \leq C \int_{\Omega_n} \Psi_n^{\text{cell}}(\nabla \hat{v}_n^k(x)) dx = C \mathcal{E}_n^{k,\text{ela}}(v_n) \leq CM.$$

Thus, by assumption on  $v_n$  and (3.27) we derive

$$\|\nabla \hat{v}_n^k\|_{L^2(\Omega_n)} + \mathcal{H}^1(S(\hat{v}_n^k)) + \|\hat{v}_n^k\|_{L^\infty(\Omega_n)} \leq C \quad \text{for all } n \in \mathbb{N}.$$

Moreover, as  $\Omega$  is a Lipschitz set, one can check that  $\sup_n \mathcal{H}^1(\partial\Omega_n) < +\infty$ . Thus, we can apply Ambrosio's compactness theorem for  $SBV$  functions [5, Theorem 4.8] to obtain a function  $v \in SBV^2(\Omega)$  such that  $\chi_{\Omega_n} \hat{v}_n^k \rightarrow v$  in  $L^1(\Omega)$  and  $\chi_{\Omega_n} \nabla \hat{v}_n^k \rightharpoonup \nabla v$  weakly in  $L^2(\Omega; \mathbb{R}^2)$ .

By  $\mathcal{L}^2(\bigcup_{C_n^k(v_n)} \Delta) \leq \frac{\sqrt{3}}{4} \varepsilon_n^2 \#C_n^k(v_n)$ , (3.27), the bound on  $\|\tilde{v}_n\|_{L^\infty(\Omega_n)}$ , and the fact that  $\tilde{v}_n = \hat{v}_n^k$  on  $\Omega_n \setminus \bigcup_{C_n^k(v_n)} \Delta$ , we eventually also get  $\chi_{\Omega_n} \tilde{v}_n \rightarrow v$  in  $L^1(\Omega)$ , i.e.,  $v_n$  AC-converges to  $v$ . Since  $g_n$  AC-converges to  $g$  and  $v_n \in \mathcal{A}(g_n)$ , we conclude the proof by observing  $v = g$  on  $\Omega \setminus \bar{U}$ .  $\square$

Now, we prove a lower semicontinuity result for the elastic energy.

**Lemma 3.6** (Lower semicontinuity). *Let  $(v_n)_n$  be a sequence of displacements with displacement history  $(u_n^j)_{j < k_n}$  for  $k_n \in \mathbb{N}$ , satisfying  $\mathcal{E}_n(v_n; (u_n^j)_{j < k_n}) + \|\tilde{v}_n\|_{L^\infty(\Omega_n)} \leq M$  for some  $M > 0$ . Denote by  $v \in SBV^2(\Omega)$  the AC-limit given in Proposition 3.5. Then, we have*

$$\liminf_{n \rightarrow \infty} \mathcal{E}_n^{k_n, \text{ela}}(v_n) \geq \int_{\Omega} \Phi(\nabla v) \, dx.$$

*Proof.* For convenience, we again simply write  $k$  in place of  $k_n$ . Let  $\omega$  be the function from Lemma 3.4(i) satisfying  $\frac{\omega(y)}{|y|^2} \rightarrow 0$  for  $|y| \rightarrow 0$ . Then, by (3.15), by Lemma 3.4(i), and the fact that  $\Phi$  is homogeneous of degree 2, we have

$$\mathcal{E}_n^{k, \text{ela}}(v_n) = \int_{\Omega_n} \Psi_n^{\text{cell}}(\nabla \hat{v}_n^k(x)) \, dx = \int_{\Omega_n} \left( \Phi(\nabla \hat{v}_n^k) + \frac{1}{\varepsilon_n} \omega(\varepsilon_n^{1/2} \nabla \hat{v}_n^k) \right) \, dx.$$

Consider the function  $\chi_n(x) := \chi_{[0, \varepsilon_n^{-1/4})}(|\nabla \hat{v}_n^k(x)|)$  and note by (3.28) that for  $n \rightarrow \infty$  we have  $\chi_n \rightarrow 1$  in measure on  $\Omega$ . In particular, we can estimate

$$\mathcal{E}_n^{k, \text{ela}}(v_n) \geq \int_{\Omega_n} \chi_n(x) \left( \Phi(\nabla \hat{v}_n^k(x)) + \frac{1}{\varepsilon_n} \omega(\varepsilon_n^{1/2} \nabla \hat{v}_n^k(x)) \right) \, dx.$$

Note that we can express the second part of the integrand as

$$\chi_n(x) \frac{1}{\varepsilon_n} \omega(\varepsilon_n^{1/2} \nabla \hat{v}_n^k) = \chi_n(x) |\nabla \hat{v}_n^k|^2 \frac{\omega(\varepsilon_n^{1/2} \nabla \hat{v}_n^k)}{|\varepsilon_n^{1/2} \nabla \hat{v}_n^k|^2}.$$

By definition of  $\omega$  and  $\chi_n$  we deduce that  $\chi_n(x) \frac{\omega(\varepsilon_n^{1/2} \nabla \hat{v}_n^k)}{|\varepsilon_n^{1/2} \nabla \hat{v}_n^k|^2}$  converges uniformly to 0. Since by (3.28) we have  $\|\nabla \hat{v}_n^k\|_{L^2(\Omega_n)} \leq C$ , this yields

$$\int_{\Omega_n} \chi_n(x) \frac{1}{\varepsilon_n} \omega(\varepsilon_n^{1/2} \nabla \hat{v}_n^k) \, dx \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Since  $\chi_{\Omega_n} \nabla \hat{v}_n^k \rightharpoonup \nabla v$  weakly in  $L^2(\Omega; \mathbb{R}^2)$ , we have  $\chi_n \chi_{\Omega_n} \nabla \hat{v}_n^k \rightharpoonup \nabla v$  weakly in  $L^2(\Omega; \mathbb{R}^2)$ . Because  $\Phi$  defined in (2.14) is convex with  $\Phi(0) = 0$ , we get

$$\liminf_{n \rightarrow \infty} \mathcal{E}_n^{k, \text{ela}}(v_n) = \liminf_{n \rightarrow \infty} \int_{\Omega_n} \chi_n \Phi(\nabla \hat{v}_n^k) \, dx \geq \int_{\Omega} \Phi(\nabla v) \, dx. \quad (3.29)$$

This concludes the proof.  $\square$

Now, we will show that the lower semicontinuity result along with energy convergence implies strong convergence. For this we follow a standard argument, see e.g. [24, 47, 48] or [2, 37] for results in the context of fracture. More precisely, we have the following.

**Lemma 3.7** (Strong convergence). *Assume the setting of Lemma 3.6. If we have*

$$\lim_{n \rightarrow \infty} \int_{\Omega_n} \Psi_n^{\text{cell}}(\nabla \hat{v}_n^{k_n}) \, dx = \lim_{n \rightarrow \infty} \mathcal{E}_n^{k_n, \text{ela}}(v_n) = \int_{\Omega} \Phi(\nabla v) \, dx, \quad (3.30)$$

then we deduce

$$\chi_{\Omega_n} \nabla \hat{v}_n^{k_n} \rightarrow \nabla v \text{ strongly in } L^2(\Omega; \mathbb{R}^2). \quad (3.31)$$

*Proof.* As before, we simply write  $k$  in place of  $k_n$ . We first check that  $h_n := |\chi_{\Omega_n} \nabla \hat{v}_n^k|^2$  is equi-integrable on  $\Omega$ . If not, we would have

$$\lim_{M \rightarrow \infty} \limsup_{n \rightarrow \infty} \int_{\{h_n > M\}} h_n \, dx \geq \eta$$

for some  $\eta > 0$ . By a diagonal argument we then find a sequence  $M_n \rightarrow \infty$  with  $M_n \leq \varepsilon_n^{-1}$  such that

$$\limsup_{n \rightarrow \infty} \int_{\{h_n > M_n\}} h_n \, dx \geq \eta.$$

With  $\bar{\chi}_n := \chi_{\{h_n \leq M_n\}}$ , we have  $\bar{\chi}_n \rightarrow 1$  a.e. on  $\Omega$  by (3.28). Then, repeating the lower semicontinuity proof in (3.29) and using Lemma 3.4(ii) together with (3.14) we get

$$\begin{aligned} \limsup_{n \rightarrow \infty} \int_{\Omega_n} \Psi_n^{\text{cell}}(\nabla \hat{v}_n^k) \, dx &\geq \liminf_{n \rightarrow \infty} \int_{\Omega_n} \bar{\chi}_n \Psi_n^{\text{cell}}(\nabla \hat{v}_n^k) \, dx + \limsup_{n \rightarrow \infty} \int_{\Omega_n} (1 - \bar{\chi}_n) c |\nabla \hat{v}_n^k|^2 \, dx \\ &\geq \int_{\Omega} \Phi(\nabla v) \, dx + c\eta. \end{aligned}$$

This, however, contradicts (3.30) and shows that  $(h_n)_n$  is equi-integrable. As in the previous proof, we define  $\chi_n(x) = \chi_{[0, \varepsilon_n^{-1/4}]}(|\nabla \hat{v}_n^k(x)|)$ . Then, the equi-integrability and Lemma 3.4(iii) yield

$$\lim_{n \rightarrow \infty} \int_{\Omega_n} (1 - \chi_n(x)) \Psi_n^{\text{cell}}(\nabla \hat{v}_n^k) \, dx = 0, \quad \lim_{n \rightarrow \infty} \int_{\Omega_n} (1 - \chi_n(x)) \Phi(\nabla \hat{v}_n^k) \, dx = 0.$$

Along with Lemma 3.4(i) this shows

$$\left| \int_{\Omega_n} \Psi_n^{\text{cell}}(\nabla \hat{v}_n^k) \, dx - \int_{\Omega_n} \Phi(\nabla \hat{v}_n^k) \, dx \right| \rightarrow 0$$

and thus, again by (3.30),

$$\lim_{n \rightarrow \infty} \int_{\Omega} \chi_{\Omega_n} \Phi(\nabla \hat{v}_n^k) \, dx = \int_{\Omega} \Phi(\nabla v) \, dx. \quad (3.32)$$

Now, since  $\Phi$  is strictly convex and homogeneous of degree 2, see (2.14), there exists a constant  $c > 0$  such that  $y \mapsto \Phi(y) - c|y|^2$  is convex. Hence, we can estimate by (3.32)

$$\begin{aligned} \liminf_{n \rightarrow \infty} \int_{\Omega_n} -c |\nabla \hat{v}_n^k|^2 \, dx &= \liminf_{n \rightarrow \infty} \int_{\Omega_n} \left( \Phi(\nabla \hat{v}_n^k) - c |\nabla \hat{v}_n^k|^2 - \Phi(\nabla \hat{v}_n^k) \right) \, dx \\ &\geq \int_{\Omega} (\Phi(\nabla v) - c |\nabla v|^2) \, dx - \int_{\Omega} \Phi(\nabla v) \, dx = - \int_{\Omega} c |\nabla v|^2 \, dx. \end{aligned}$$

This implies  $\limsup_{n \rightarrow \infty} \|\nabla \hat{v}_n^k\|_{L^2(\Omega_n)} \leq \|\nabla v\|_{L^2(\Omega)}$ . Since  $L^2(\Omega; \mathbb{R}^2)$  is a uniformly convex Banach space, we can apply [17, Proposition 3.32] to infer that (3.31) holds.  $\square$

**3.4. Convergence of sets.** After we have discussed compactness of displacement fields and corresponding lower semicontinuity results for elastic energies in Subsection 3.3, this section is devoted to results for crack sets. We start by recalling the notion of  $\sigma$ -convergence introduced in [40].

We denote by  $\mathcal{A}(\Omega)$  all open subsets of  $\Omega$  and define the family of sets with finite perimeter in  $\Omega$  by

$$P(\Omega) := \{v \in SBV(\Omega) : u(x) \in \{0, 1\} \text{ for a.e. } x \in \Omega\}.$$

For a sequence  $(K_n)_{n \in \mathbb{N}}$  of rectifiable sets in  $\Omega$ , we consider the functionals  $\mathcal{H}_n^- : P(\Omega) \times \mathcal{A}(\Omega) \rightarrow \mathbb{R}$  defined by

$$\mathcal{H}_n^-(u, A) = \mathcal{H}^1((S(u) \setminus K_n) \cap A).$$

Following [3, Theorem 3.2], we assume that for every  $A \in \mathcal{A}(\Omega)$ ,  $\mathcal{H}_n^-(\cdot, A)$   $\Gamma$ -converges with respect to the strong topology of  $L^1(\Omega)$  to a functional  $\mathcal{H}^-(\cdot, A)$ , which by [9, Theorem 3] is of the form

$$\mathcal{H}^-(u, A) := \int_{S(u) \cap A} h^-(x, \nu) dx$$

for some function  $h^- : \Omega \times \mathbb{S}^1 \rightarrow [0, \infty)$ .

**Definition 3.8** ( $\sigma$ -convergence). Let  $(K_n)_{n \in \mathbb{N}}$  be a sequence of rectifiable sets in  $\Omega$ . We say that  $K_n$   $\sigma$ -converges to  $K$  if the functionals  $\mathcal{H}_n^-$   $\Gamma$ -converge in the strong topology of  $L^1(\Omega)$  to the functional  $\mathcal{H}^-$ , and  $K$  is the (unique) rectifiable set  $K$  in  $\Omega$  such that

$$h^-(x, \nu_K(x)) = 0 \quad \text{for } \mathcal{H}^1\text{-a.e. } x \in K,$$

and such that for every rectifiable set  $H \subset \Omega$  we have

$$h^-(x, \nu_H(x)) = 0 \quad \text{for } \mathcal{H}^1\text{-a.e. } x \in H \Rightarrow H \tilde{\subset} K.$$

Unfortunately, we cannot employ this notion in our setting, but in view of the definitions in (3.12) and (3.13), we have to deal with two parts of the jump set

$$K_n^{k,L}(v_n) = \bigcup_{i=1,2,3} K_n^{k,i}(v_n) \quad \text{and} \quad K_n^{k,S}(v_n) = \bigcup_{\mathcal{C}_n^k(v_n)} \partial \Delta, \quad (3.33)$$

where intuitively only  $K_n^{k,L}(v_n)$  should contribute to the crack in the continuum limit whereas  $K_n^{k,S}(v_n)$  should not affect it, see the comment in Remark 3.3(iii). To solve this issue, we introduce another jump interpolation which only exhibits discontinuities on  $K_n^{k,L}(v_n)$  and invoke the notion of  $\sigma^p$ -convergence, introduced in [22], where  $p$  stands for the integrability of the absolutely continuous parts of derivatives. In the following, we say that  $v_j \rightarrow v$  weakly in  $SBV^p(\Omega)$  if  $v_j, v \in SBV^p(\Omega) \cap L^\infty(\Omega)$ ,  $v_j \rightarrow v$  a.e. in  $\Omega$ ,  $\nabla v_j \rightharpoonup \nabla v$  weakly in  $L^p(\Omega; \mathbb{R}^2)$ , and  $\|v_j\|_\infty + \mathcal{H}^1(S(v_j))$  is uniformly bounded in  $j$ .

**Definition 3.9** ( $\sigma^p$ -convergence). Let  $(K_n)_{n \in \mathbb{N}}$  be a sequence of rectifiable sets in  $\Omega$ . We say that  $K_n$   $\sigma^p$ -converges to  $K \subset \Omega$  in  $\Omega$  if  $\mathcal{H}^1(K_n)$  is uniformly bounded in  $n$  and if the following is satisfied:

- (i) If a sequence  $(v_j)_j \subset SBV^p(\Omega)$  with  $v_j \rightarrow v$  weakly in  $SBV^p(\Omega)$  fulfills  $S(v_j) \tilde{\subset} K_{n_j}$  for some sequence  $n_j \rightarrow \infty$ , then  $S(v) \tilde{\subset} K$ .
- (ii) There exists a function  $v \in SBV^p(\Omega)$  and a sequence  $(v_n)_n$  that converges weakly to  $v$  in  $SBV^p(\Omega)$  such that  $S(v_n) \tilde{\subset} K_n$  for each  $n \in \mathbb{N}$  and  $K \stackrel{\cong}{=} S(v)$ , where  $\stackrel{\cong}{=}$  stands for equality up to an  $\mathcal{H}^1$ -negligible set.

*Remark 3.10.* (a) The  $\sigma^p$ -limit of any sequence is always rectifiable since  $S(u)$  is rectifiable for each  $u \in SBV(\Omega)$ .

(b) Note that  $\sigma$ -convergence is invariant under perturbations with vanishing  $\mathcal{H}^1$ -measure, i.e. given  $(\Gamma_n)_n$  and  $(\Gamma_n^*)_n$  with  $\mathcal{H}^1(\Gamma_n^*) \rightarrow 0$ , the  $\sigma$ -limits of  $(\Gamma_n)_n$  and  $(\Gamma_n \cup \Gamma_n^*)_n$ , if existent, coincide, see [40, Remark 5.2].

The same property for  $\sigma^p$  convergence does not hold in general, see [40, Remark 5.10].

(c) If  $\Gamma_n$  and  $\Gamma'_n$   $\sigma^p$ -converge to  $\Gamma$  and  $\Gamma'$  and  $\Gamma_n \subset \Gamma'_n$  for all  $n \in \mathbb{N}$ , then it follows  $\Gamma \subset \Gamma'$ , see [22, Remark 4.2].

(d) Note that the  $\sigma$ -limit of a sequence always contains its  $\sigma^p$ -limit, see [40, Corollary 5.9]. The opposite inclusion is false. See [40, Remark 5.10] for an example, where the  $\sigma^p$ -limit is strictly contained in the  $\sigma$ -limit.

In the following, we need the compactness and lower semicontinuity properties of  $\sigma^p$ -convergence proved in [22, Theorems 4.3, 4.7, and 4.8].

**Theorem 3.11.** *Let  $p \in (1, \infty)$ . Let  $t \mapsto \Gamma_n(t)$  be a sequence of increasing set functions defined on an interval  $I \subset \mathbb{R}$  with values contained in  $\Omega$ , i.e.,  $\Gamma(s) \subset \Gamma(t) \subset \Omega$  for every  $s, t \in I$  with  $s < t$ . Assume that  $\mathcal{H}^1(\Gamma_n(t))$  is bounded uniformly with respect to  $n$  and  $t$ . Then, there exist a subsequence  $(\Gamma_{n_k})_k$  and an increasing set function  $t \mapsto \Gamma(t)$  on  $I$  such that for every  $t \in I$  we have:*

- (a)  $\Gamma_{n_k}(t) \rightarrow \Gamma(t)$  in the sense of  $\sigma^p$ -convergence.
- (b) Let  $\varphi: \mathbb{R}^2 \rightarrow [0, \infty)$  be a norm on  $\mathbb{R}^2$ . Then, we have

$$\int_{\Gamma(t)} \varphi(\nu_\Gamma) d\mathcal{H}^1(s) \leq \liminf_{k \rightarrow \infty} \int_{\Gamma_{n_k}(t)} \varphi(\nu_{\Gamma_{n_k}}) d\mathcal{H}^1,$$

where  $\nu_\Gamma$  and  $\nu_{\Gamma_{n_k}}$  denote measure-theoretic unit normals to  $\Gamma(t)$  and  $\Gamma_{n_k}(t)$ , respectively.

*Remark 3.12.* The same two properties (a) and (b) also hold for  $\sigma$ -convergence in place of  $\sigma^p$ -convergence, see [40, Propositions 5.3 and 5.7].

Recall (3.33). The essential point for us will be the following property.

**Proposition 3.13** ( $\sigma^{3/2}$ -limit of crack set). *Let  $(v_n)_n$  be a sequence of atomistic displacements and, for each  $n \in \mathbb{N}$ , let  $(u_n^j)_{j < k_n}$  be a displacement history for  $k_n \in \mathbb{N}$  depending on  $n$ . Suppose that  $\mathcal{E}_n(v_n; (u_n^j)_{j < k_n}) + \|\tilde{v}_n\|_{L^\infty(\Omega_n)} \leq M$  for some  $M > 0$ . Assume that  $K_n^{k_n, L}(v_n)$   $\sigma^{3/2}$ -converges to  $K$  and that  $v_n$  AC-converges to some  $v \in SBV^2(\Omega)$ . Then, we have  $S(v) \tilde{\subset} K$ .*

*Remark 3.14.* (i) The gist is that the limit is taken with respect to  $K_n^{k_n, L}(v_n)$ , i.e., only with respect to the part of the crack set which should be relevant for the continuum limit. Note that the result is not an immediate consequence of (i) in the definition of  $\sigma^p$ -convergence as  $S(\delta_n^{k_n})$  is *not contained* in  $K_n^{k_n, L}(v_n)$ , cf. Remark 3.3(ii).

(ii) Note that the choice  $p = 3/2$  is for definiteness only and could be replaced by any  $1 < p < 2$ . Later we see that along the evolution the  $\sigma$ -limit and the  $\sigma^p$ -limit coincide, see Step 5 in the proof of Theorem 2.2.

*Proof.* For convenience, we drop the dependency of  $k$  on  $n$  in the proof and simply write  $k$  in place of  $k_n$ . The idea is to construct an interpolation  $\bar{v}_n: \Omega_n \rightarrow \mathbb{R}$  of  $v_n: \mathcal{L}_n(\Omega_n) \rightarrow \mathbb{R}$  such that

- (1)  $S(\bar{v}_n) \subset K_n^{k, L}(v_n)$ ,
- (2)  $\chi_{\Omega_n} \bar{v}_n \rightarrow v$  weakly in  $SBV^{3/2}(\Omega)$ .

Then, from item (i) of the definition of  $\sigma^{3/2}$ -convergence we obtain  $S(v) \tilde{\subset} K$ .

For triangles  $\Delta \in \mathcal{T}_n \setminus \mathcal{C}_n^{k, L}(v_n)$  we set  $\bar{v}_n := v_n$ . Recall, that by definition (see (3.8)) we then have

$$|\nabla \bar{v}_n| = |(v_n)_\Delta| \leq C \varepsilon_n^{-1/2} R_n \quad \text{on triangles } \Delta \in \mathcal{T}_n \setminus \mathcal{C}_n^{k, L}(v_n). \quad (3.34)$$

Now we consider triangles  $\Delta \in \mathcal{C}_n^{k, L}(v_n)$  and recall the division into sub-triangles introduced in Figure 2.

First, consider  $\Delta \in \mathcal{C}_n^{k,3}(v_n)$ . On each subtriangle  $\Delta_\alpha$ ,  $\alpha = 1, 2, 3$ , we define  $\bar{v}_n$  as an affine interpolation such that  $\bar{v}_n = v_n$  on the vertex and

$$\nabla \bar{v}_n \cdot \mathbf{v} = \begin{cases} \nabla \tilde{v}_n \cdot \mathbf{v} & \text{if } |\nabla \tilde{v}_n \cdot \mathbf{v}| \leq 2R_n \varepsilon_n^{-1/2}, \\ 0 & \text{if } |\nabla \tilde{v}_n \cdot \mathbf{v}| > 2R_n \varepsilon_n^{-1/2} \end{cases}, \quad \text{for } \mathbf{v} \in \{\mathbf{v}_\beta, \mathbf{v}_\gamma\} \quad (3.35)$$

where  $\alpha, \beta, \gamma$  are pairwise distinct. In  $\Delta_{\text{mid}}$ , we define  $\bar{v}_n$  as a constant.

If  $\Delta \in \mathcal{C}_n^{k,2}(v_n)$ , assuming  $M_{n,\Delta}^{k,\mathbf{v}_\alpha}(v_n) \leq R_n$ , we define  $\bar{v}_n$  on  $\Delta_\alpha, \Delta_\beta, \Delta_\gamma$  as in (3.35), and in  $\Delta_{\text{mid}}$  we choose an affine interpolation between the affine functions attained on  $h_\Delta^\beta$  and  $h_\Delta^\gamma$ .

Note that by our construction the only discontinuities in the closure of a triangle  $\Delta$  appear on  $\bigcup_{i=1}^3 h_\Delta^i$  for  $\Delta \in \mathcal{C}_n^{k,3}(v_n)$  and on  $h_\Delta^m$  for  $\Delta \in \mathcal{C}_n^{k,2}(v_n)$ , where  $m$  is the unique index with  $M_{n,\Delta}^{k,\mathbf{v}_m}(v_n) \leq R_n$ . In fact, given  $(x, x') \in \text{NN}_n(\Omega)$  with  $x - x'$  parallel to  $\mathbf{v}_\alpha$ , denote by  $\Delta$  and  $\Delta'$  the triangles adjacent to the spring. Then, we have  $\bar{v}_n(x) = v_n(x)$ ,  $\bar{v}_n(x') = v_n(x')$ , and on the corresponding sub-triangles  $\Delta_\beta \cup \Delta_\gamma \cup \Delta'_\beta \cup \Delta'_\gamma$  it either holds  $\nabla \bar{v} \cdot \mathbf{v}_\alpha = \nabla \tilde{v}_n \cdot \mathbf{v}_\alpha$  or  $\nabla \bar{v} \cdot \mathbf{v}_\alpha = 0$ , where we also use (3.8) to cover the case where one of triangles  $\Delta$  and  $\Delta'$  lie in  $\mathcal{C}_n^{k,L}(v_n)$  and the other one does not.

Thus, in view of (3.12), we get that  $S(\bar{v}_n) \subset K_n^{k,L}(v_n)$ . Moreover, the construction leads to  $|\nabla \bar{v}_n(x)| \leq CR_n \varepsilon_n^{-1/2}$  for  $x \in \Delta$  with  $\Delta \in \mathcal{C}_n^{k,L}(v_n)$ . Putting this together with (3.34) yields

$$|\nabla \bar{v}_n| \leq CR_n \varepsilon_n^{-1/2} \quad \text{a.e. on } \Omega_n.$$

Using  $\|\bar{v}_n\|_{L^\infty(\Omega_n)} \leq C$ , by construction we also have  $\|\bar{v}_n\|_{L^\infty(\Omega_n)} \leq C$ .

Now, recall that by Proposition 3.5 we have

$$\#\mathcal{C}_n^k(v_n) \leq \frac{C}{\varepsilon_n} \quad \text{and} \quad \|\nabla \hat{v}_n^k\|_{L^2(\Omega_n)}^2 \leq C. \quad (3.36)$$

Invoking (3.7) this leads to

$$\sum_{\Delta \in \mathcal{C}_n^k(v_n)} \int_\Delta |\nabla \bar{v}_n(x)|^{3/2} dx \leq C \varepsilon_n^2 \#\mathcal{C}_n^k(v_n) R_n^{3/2} \varepsilon_n^{-3/4} \leq CR_n^{3/2} \varepsilon_n^{1/4} \rightarrow 0.$$

As outside of the triangles  $\mathcal{C}_n^k(v_n)$  we have  $\bar{v}_n = \hat{v}_n^k$ , we can estimate by (3.36) and Hölder's inequality

$$\limsup_{n \rightarrow \infty} \int_{\Omega_n} |\nabla \bar{v}_n(x)|^{3/2} dx \leq \limsup_{n \rightarrow \infty} C \left( \int_{\Omega_n} |\nabla \bar{v}_n(x)|^2 dx \right)^{3/4} \leq \limsup_{n \rightarrow \infty} C \|\nabla \hat{v}_n^k\|_{L^2(\Omega_n)}^{3/2} \leq C.$$

Eventually, (3.36) also yields

$$\mathcal{H}^1(S(\bar{v}_n)) \leq \mathcal{H}^1(K_n^{k,L}(v_n)) \leq C \#\mathcal{C}_n^{k,L}(v_n) \varepsilon_n \leq C \#\mathcal{C}_n^k(v_n) \varepsilon_n \leq C.$$

Altogether, we have  $\|\nabla \bar{v}_n\|_{L^{3/2}(\Omega_n)} + \mathcal{H}^1(S(\bar{v}_n)) + \|\bar{v}_n\|_\infty < \infty$ , and thus by Ambrosio's compactness theorem and the assumption that  $v_n$  AC-converges to  $v$  we conclude  $\chi_{\Omega_n} \bar{v}_n \rightarrow v$  weakly in  $SBV^{3/2}(\Omega)$ . This proves the assertion.  $\square$

#### 4. CONVERGENCE OF QUASI-STATIC CRACK GROWTH: PROOF OF THEOREM 2.2

This section is devoted to the proof of Theorem 2.2. In Subsection 4.1, we start by considering the time-discrete atomistic problems. Afterwards, in Subsection 4.2 we establish compactness results for displacements and crack sets. Eventually, Subsection 4.3 is subject to the proof of the main result, namely the convergence to an irreversible quasi-static crack evolution  $t \mapsto (u(t), K(t))$ .

**4.1. Time-discrete atomistic evolution.** We recall the definition of the time-discrete evolution in (2.11). In particular, we denote by  $\{0 = t_n^0 < t_n^1 < \dots < t_n^{T/\delta_n} = T\}$  a time discretization of the interval  $[0, T]$ , and let  $(u_n^k)_k$  be the corresponding displacements. Recalling the boundary values  $(g(t_n^k))_k$ , we define  $g_n^k: \mathcal{L}_n(\Omega) \rightarrow \mathbb{R}$  by  $g_n^k(x) = g(t_n^k, x)$ .

**Lemma 4.1.** *The minimization problems (2.9) and (2.10) admit solutions. The solution  $t \mapsto u_n(t)$  given in (2.11) satisfies  $\sup_{t \in [0, T]} \|\tilde{u}_n(t)\|_{L^\infty(\Omega_n)} < +\infty$ .*

*Proof.* We observe that we can restrict the minimization problem to functions  $v \in \mathcal{A}_n(g(t_n^k))$  which satisfy  $|v(x)| \leq \max_{x \in \mathcal{L}_n(\Omega)} |g_n^k(x)| \leq C$  for all  $x \in \mathcal{L}_n(\Omega)$  by the monotonicity of the function  $\Psi$  and the regularity of  $g$ . As the space of displacements from  $\mathcal{L}_n(\Omega)$  to  $\mathbb{R}$  is finite dimensional, this along with the continuity of  $\Psi$  guarantees existence by the Direct Method.  $\square$

Similarly as in (2.11) we write

$$\hat{u}_n(t) := \hat{u}_n^k \quad \text{for } t \in [t_n^k, t_n^{k+1}),$$

where the interpolation  $\hat{u}_n^k$  is defined in Subsection 3.2 (for general  $v_n$  in place of  $u_n^k$ ). Recalling (2.6) and (3.5), given an arbitrary  $v_n: \mathcal{L}_n(\Omega) \rightarrow \mathbb{R}$ , we also introduce the shorthand notation

$$\mathcal{E}_n^0(v_n) := \mathcal{E}_n(v_n), \quad \mathcal{E}_n^k(v_n) := \mathcal{E}_n(v_n; (u_n^j)_{j < k}) \quad \text{for } k \geq 1. \quad (4.1)$$

**Lemma 4.2** (Bound on derivatives). *Let  $t \mapsto u_n(t)$  be the discrete evolution defined in (2.11). We have  $\mathcal{E}_n^0(u_n^0) \leq C$ . Moreover, it holds that*

$$\int_{\Omega_n} |\nabla \hat{u}_n(t)|^2 dx \leq C \quad \text{for all } t \in [0, T] \text{ and for all } n \in \mathbb{N} \quad (4.2)$$

for some  $C > 0$  depending on  $W$  and  $g$ .

*Proof.* By (2.6) and by writing the energy as an integral using (3.15)–(3.16), we get

$$\mathcal{E}_n(g_n^0) = \frac{\varepsilon_n}{2} \sum_{(x, x') \in \text{NN}_\varepsilon(\Omega)} \Psi\left(\frac{\varepsilon_n^{1/2} |g_n^0(x) - g_n^0(x')|}{\varepsilon_n}\right) = \sum_{\Delta \in \mathcal{T}_n} \int_{\Delta} \Psi_n^{\text{cell}}((g_n^0)_\Delta) + \mathcal{E}_n^{\text{bdy}}(g_n^0), \quad (4.3)$$

where  $\mathcal{E}_n^{\text{bdy}}(g_n^0)$  accounts for springs at the boundary, similarly to (3.6). As  $\mathcal{E}_n^{\text{bdy}}(g_n^0)$  is bounded and as  $(g_n^0)_\Delta$  is uniformly bounded by  $g \in W^{1,1}([0, T]; W^{2,\infty}(\Omega))$ , Lemma 3.4(iii) yields  $\mathcal{E}_n(g_n^0) \leq C$  independently of  $n$  for some  $C > 0$  large enough, which depends on  $\Phi$  (and thus on  $W$ ) and  $g$ . In particular, by (2.9) this implies  $\mathcal{E}_n(u_n^0) \leq C$ .

In a similar fashion, we obtain a control on  $u_n^k$  for  $k \geq 1$ . As  $\varepsilon_n^{1/2} \|\nabla g\|_{L^\infty([0, T]; L^\infty(\Omega))} \leq R$  for  $n$  large enough, (2.8) implies

$$\mathcal{E}_n^k(g_n^k) = \frac{\varepsilon_n}{2} \sum_{M_{x, x'}^{\varepsilon_n}, ((u_n^j)_{j < k}) \leq R} \Psi(\varepsilon_n^{-1/2} |g_n^k(x) - g_n^k(x')|) + \frac{\varepsilon_n}{2} \sum_{M_{x, x'}^{\varepsilon_n}, ((u_n^j)_{j < k}) > R} \Psi(M_{x, x'}^{\varepsilon_n}, ((u_n^j)_{j < k})),$$

where the sums run over all  $(x, x') \in \text{NN}_n(\Omega)$ . Note that by (2.10) we have  $\mathcal{E}_n^k(u_n^k) \leq \mathcal{E}_n^k(g_n^k)$ . Using the monotonicity of the memory variable in  $k$  and the monotonicity of  $\Psi$ , this implies

$$\frac{\varepsilon_n}{2} \sum_{M_{x, x'}^{\varepsilon_n}, ((u_n^j)_{j \leq k}) \leq R} \Psi(\varepsilon_n^{-1/2} |u_n^k(x) - u_n^k(x')|) \leq \frac{\varepsilon_n}{2} \sum_{M_{x, x'}^{\varepsilon_n}, ((u_n^j)_{j \leq k}) \leq R} \Psi(\varepsilon_n^{-1/2} |g_n^k(x) - g_n^k(x')|) \leq C$$

for  $C$  depending on  $\Phi$  and  $g$ , where the last step follows by repeating the argument in (4.3). Recalling the representation of the elastic energy in (3.15)–(3.16), we thus find  $\int_{\Omega_n} \Psi_n^{\text{cell}}(\nabla \hat{u}_n^k) dx \leq C$ . By (3.14)

and the property of  $\Psi_n^{\text{cell}}$  stated in Lemma 3.4(ii) we get the bound  $\int_{\Omega_n} |\nabla \hat{u}_n^k|^2 dx \leq C$ . This shows (4.2) and concludes the proof.  $\square$

We continue with an energy estimate which will be vital for the proof of (2.17) and also delivers a priori bounds for the discrete evolutions. To this end, we discretize the boundary values  $g \in W^{1,1}([0, T]; W^{2,\infty}(\Omega))$  in space. Given  $g(\tau)$  for  $\tau \in [0, T]$ , we define the function  $g_n(\tau): \mathcal{L}_n(\Omega) \rightarrow \mathbb{R}$  by  $g_n(\tau)(x) = g(\tau, x)$  for  $x \in \mathcal{L}_n(\Omega)$ . As before, by  $\tilde{g}_n(\tau): \Omega_n \rightarrow \mathbb{R}$  we denote the interpolation which is piecewise affine on the triangles of  $\mathcal{T}_n$ . In the same way, we define the time derivative  $\partial_t \tilde{g}_n(\tau)$ .

**Lemma 4.3.** *Let  $t \mapsto u_n(t)$  be the discrete evolution defined in (2.11). Let  $k = 0, \dots, (T/\delta_n) - 1$ . Then, there exists  $(\eta_n)_n$  independent of  $k$  with  $\eta_n \rightarrow 0$  such that*

$$\mathcal{E}_n^k(u_n^k) - \mathcal{E}_n^0(u_n^0) \leq \int_0^{t_n^k} \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n(\tau)) \cdot \nabla \partial_t \tilde{g}_n(\tau) dx d\tau + (1 + \varepsilon_n \# \mathcal{C}_n^k) \eta_n, \quad (4.4)$$

where  $\mathcal{C}_n^k$  is defined in (3.3).

*Proof.* The proof follows a rather standard strategy, see e.g. [22, Lemma 6.1]. Here, we include full details and in particular describe how the estimates are adapted to our discrete setting. For each step  $l$ , where  $0 < l \leq k$ , we consider the test function  $\xi_n^l := u_n^{l-1} + g_n^l - g_n^{l-1}$  and obtain by (2.10)

$$\mathcal{E}_n^l(u_n^l) \leq \mathcal{E}_n^l(\xi_n^l) = \mathcal{E}_n^l(u_n^{l-1} + g_n^l - g_n^{l-1}). \quad (4.5)$$

We aim at estimating the right-hand side of this inequality. Our goal is to prove that there exists a bounded sequence  $(\vartheta_n)_n$  in  $L^\infty([0, T] \times \Omega)$  with  $\|\vartheta_n(\tau)\|_{L^2(\Omega)} \rightarrow 0$  uniformly in  $\tau$  such that, for each  $l$ , we have

$$\begin{aligned} \mathcal{E}_n^l(u_n^{l-1} + g_n^l - g_n^{l-1}) - \mathcal{E}_n^l(u_n^{l-1}) &\leq \int_{t_n^{l-1}}^{t_n^l} \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n^{l-1} + \vartheta_n(\tau)) \cdot \nabla \partial_t \tilde{g}_n(\tau) dx d\tau \\ &\quad + C(1 + \varepsilon_n \# \mathcal{C}_n^l) \varepsilon_n^{1/2} \int_{t_n^{l-1}}^{t_n^l} \|\nabla \partial_t \tilde{g}_n(\tau)\|_{L^\infty(\Omega_n)} d\tau. \end{aligned} \quad (4.6)$$

We defer the proof of (4.6) to Step 2 below and first show how to derive the estimate (4.4) from this.

*Step 1: Proof of (4.4).* Assume (4.6) holds. Since  $\mathcal{E}_n^{l-1}(u_n^{l-1}) = \mathcal{E}_n^l(u_n^{l-1})$  for each step  $l$ , summing up over all time steps  $0 \leq l \leq k$  and using (4.5) yields a telescopic sum on the left-hand side of (4.6) such that we get

$$\mathcal{E}_n^k(u_n^k) - \mathcal{E}_n^0(u_n^0) \leq \int_0^{t_n^k} \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n(\tau) + \vartheta_n(\tau)) \cdot \nabla \partial_t \tilde{g}_n(\tau) dx d\tau + C(1 + \varepsilon_n \# \mathcal{C}_n^k) \varepsilon_n^{1/2} \quad (4.7)$$

for a constant  $C$  depending on  $g \in W^{1,1}([0, T]; W^{2,\infty}(\Omega))$ , where we used that the sets  $(\mathcal{C}_n^l)_l$  are increasing in  $l$ . By  $|D\Psi_n^{\text{cell}}(z)| \leq C|z|$  for  $z \in \mathbb{R}^2$ , see Lemma 3.4(iii), and the fact that  $(D\Psi_n^{\text{cell}})_n$  satisfies the continuity property stated in Lemma 3.4(iv), we can invoke [40, Lemma 2.4] to find

$$\left| \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n(\tau) + \vartheta_n(\tau)) \cdot \nabla \partial_t g(\tau) dx - \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n(\tau)) \cdot \nabla \partial_t g(\tau) dx \right| \rightarrow 0$$

for a.e.  $\tau \in [0, T]$ . Here, we also employed  $\|\vartheta_n(\tau)\|_{L^2(\Omega)} \rightarrow 0$  uniformly in  $\tau$ . Again using  $|D\Psi_n^{\text{cell}}(z)| \leq C|z|$  we get

$$\|D\Psi_n^{\text{cell}}(\nabla \hat{u}_n(\tau))\|_{L^2(\Omega_n)} \leq C, \quad \|D\Psi_n^{\text{cell}}(\nabla \hat{u}_n(\tau) + \vartheta_n(\tau))\|_{L^2(\Omega_n)} \leq C \quad \text{for a.e. } \tau \in [0, T], \quad (4.8)$$

where we used (4.2) and for the second estimate also that  $\|\vartheta_n(\tau)\|_{L^2(\Omega)} \rightarrow 0$  uniformly in  $\tau$ . This along with the fact that  $\chi_{\Omega_n} \nabla \partial_t \tilde{g}_n$  converges strongly to  $\nabla \partial_t g$  in  $L^1([0, T]; L^\infty(\Omega; \mathbb{R}^2))$  yields for a.e.  $\tau \in [0, T]$

$$h_n(\tau) := \left| \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n(\tau) + \vartheta_n(\tau)) \cdot \nabla \partial_t \tilde{g}_n(\tau) \, dx - \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n(\tau)) \cdot \nabla \partial_t \tilde{g}_n(\tau) \, dx \right| \rightarrow 0.$$

By (4.8) we then also get by Hölder's inequality

$$|h_n(\tau)| \leq C \|\nabla \partial_t \tilde{g}_n(\tau)\|_{L^\infty(\Omega_n)}.$$

Using again the convergence of  $\chi_{\Omega_n} \nabla \partial_t \tilde{g}_n$  to  $\nabla \partial_t g$ , we obtain  $\int_0^T h_n(\tau) \, d\tau \rightarrow 0$  as  $n \rightarrow \infty$  by dominated convergence. This along with (4.7) and setting  $\eta_n := C\varepsilon_n^{1/2} + \int_0^T h_n(\tau) \, d\tau$  shows the assertion.

*Step 2: Proof of (4.6).* We recall the notation of  $\mathcal{C}_n^l$  and  $M_{n,\Delta}^{l,\mathbf{v}}$  for all  $\Delta \in \mathcal{T}_n$  and  $\mathbf{v} \in \mathcal{V}$ , see (3.1) and (3.3). We can write the memory variable of the test function in terms of the piecewise affine interpolation for each  $\Delta \in \mathcal{C}_n^l(\xi_n^l)$  and  $\mathbf{v} \in \mathcal{V}$ , namely

$$M_{n,\Delta}^{l,\mathbf{v}}(\xi_n^l) = \varepsilon_n^{1/2} \sup_{m < l} |\nabla \tilde{u}_n^m \cdot \mathbf{v}| \vee \varepsilon_n^{1/2} |\nabla \tilde{\xi}_n^l \cdot \mathbf{v}|. \quad (4.9)$$

We estimate the last term by

$$\begin{aligned} \varepsilon_n^{1/2} |\nabla \tilde{\xi}_n^l \cdot \mathbf{v}| &= \varepsilon_n^{1/2} |\nabla(\tilde{u}_n^{l-1} + \tilde{g}_n^l - \tilde{g}_n^{l-1}) \cdot \mathbf{v}| \leq \varepsilon_n^{1/2} |\nabla \tilde{u}_n^{l-1} \cdot \mathbf{v}| + \varepsilon_n^{1/2} \|\nabla \tilde{g}_n^l - \nabla \tilde{g}_n^{l-1}\|_{L^\infty(\Omega_n)} \\ &\leq \varepsilon_n^{1/2} |\nabla \tilde{u}_n^{l-1} \cdot \mathbf{v}| + \varepsilon_n^{1/2} \int_{t_n^{l-1}}^{t_n^l} \|\partial_t \nabla \tilde{g}_n(\tau)\|_{L^\infty(\Omega_n)} \, d\tau, \end{aligned} \quad (4.10)$$

where  $\tilde{g}_n^l$  denote the piecewise affine interpolation of the atomistic boundary data  $g_n^l$ . Hence, we obtain

$$M_{n,\Delta}^{l,\mathbf{v}}(\xi_n^l) = M_{n,\Delta}^{l,\mathbf{v}}(u_n^{l-1} + g_n^l - g_n^{l-1}) \leq M_{n,\Delta}^{l,\mathbf{v}}(u_n^{l-1}) + \varepsilon_n^{1/2} \int_{t_n^{l-1}}^{t_n^l} \|\partial_t \nabla \tilde{g}_n(\tau)\|_{L^\infty(\Omega_n)} \, d\tau. \quad (4.11)$$

Recalling (3.5), we can write

$$\begin{aligned} \mathcal{E}_n^l(\xi_n^l) - \mathcal{E}_n^l(u_n^{l-1}) &= \frac{\varepsilon_n}{2} \sum_{\mathcal{T}_n \setminus \mathcal{C}_n^l} \sum_{\mathbf{v} \in \mathcal{V}} \left( \Psi(\varepsilon_n^{1/2} |(\xi_n^l)_\Delta \cdot \mathbf{v}|) - \Psi(\varepsilon_n^{1/2} |(u_n^{l-1})_\Delta \cdot \mathbf{v}|) \right) + \mathcal{G}_n^l(\xi_n^l) \\ &\quad + \mathcal{E}_n^{l,\text{bdy}}(\xi_n^l) - \mathcal{E}_n^{l,\text{bdy}}(u_n^{l-1}), \end{aligned} \quad (4.12)$$

where the boundary energy is defined in (3.6) and where we have set

$$\mathcal{G}_n^l(\xi_n^l) := \frac{\varepsilon_n}{2} \sum_{\mathcal{C}_n^l} \left( \sum_{M_{n,\Delta}^{l,\mathbf{v}} > R} (\Psi(M_{n,\Delta}^{l,\mathbf{v}}(\xi_n^l)) - \Psi(M_{n,\Delta}^{l,\mathbf{v}})) + \sum_{M_{n,\Delta}^{l,\mathbf{v}} \leq R} (\Psi(\varepsilon_n^{1/2} |(\xi_n^l)_\Delta \cdot \mathbf{v}|) - \Psi(\varepsilon_n^{1/2} |(u_n^{l-1})_\Delta \cdot \mathbf{v}|)) \right).$$

Note that here we use that  $M_{n,\Delta}^{l,\mathbf{v}}(\xi_n^l) = \varepsilon_n^{1/2} |(\xi_n^l)_\Delta \cdot \mathbf{v}|$  whenever  $M_{n,\Delta}^{l,\mathbf{v}}(\xi_n^l) > R$  and  $M_{n,\Delta}^{l,\mathbf{v}} \leq R$ , cf. (4.9). In particular, the contributions of triangles  $\Delta \in \mathcal{C}_n^l(\xi_n^l) \setminus \mathcal{C}_n^l$  are captured in the first term of (4.12).

We start by estimating the term  $\mathcal{G}_n^l(\xi_n^l)$ . In view of (4.10)–(4.11), (3.4), and the monotonicity and Lipschitz continuity of  $\Psi$ , we deduce for each  $\Delta \in \mathcal{C}_n^l$  that

$$\begin{aligned} \Psi(\varepsilon_n^{1/2} |(\xi_n^l)_\Delta \cdot \mathbf{v}|) - \Psi(\varepsilon_n^{1/2} |(u_n^{l-1})_\Delta \cdot \mathbf{v}|) &\leq L\varepsilon_n^{1/2} \int_{t_n^{l-1}}^{t_n^l} \|\partial_t \nabla \tilde{g}_n(\tau)\|_{L^\infty(\Omega_n)} \, d\tau, \\ \Psi(M_{n,\Delta}^{l,\mathbf{v}}(\xi_n^l)) - \Psi(M_{n,\Delta}^{l,\mathbf{v}}) &\leq L\varepsilon_n^{1/2} \int_{t_n^{l-1}}^{t_n^l} \|\partial_t \nabla \tilde{g}_n(\tau)\|_{L^\infty(\Omega_n)} \, d\tau, \end{aligned}$$

where  $L$  denotes the Lipschitz constant of  $\Psi$ . Here we distinguished the directions  $\mathbf{v} \in \mathcal{V}$  with  $M_{n,\Delta}^{l,\mathbf{v}} \leq R$  and  $M_{n,\Delta}^{l,\mathbf{v}} > R$ , respectively. This induces

$$\mathcal{G}_n^l(\xi_n^l) \leq C\varepsilon_n \#\mathcal{C}_n^l \varepsilon_n^{1/2} \int_{t_n^{l-1}}^{t_n^l} \|\partial_t \nabla \tilde{g}_n(\tau)\|_{L^\infty(\Omega_n)} d\tau. \quad (4.13)$$

By a similar argument, noting that  $\#\{\Delta \in \mathcal{T}_n : \partial\Delta \cap \partial\Omega_n \neq \emptyset\} \leq C\varepsilon_n^{-1}$ , we get

$$\mathcal{E}_n^{l,\text{bdy}}(\xi_n^l) - \mathcal{E}_n^{l,\text{bdy}}(u_n^{l-1}) \leq C\varepsilon_n^{1/2} \int_{t_n^{l-1}}^{t_n^l} \|\partial_t \nabla \tilde{g}_n(\tau)\|_{L^\infty(\Omega_n)} d\tau. \quad (4.14)$$

We now address the elastic energy. To this end, recall the definition of the jump interpolation  $\hat{u}_n^{l-1}$ , which outside of triangles  $\Delta \in \mathcal{C}_n^l$  coincides with the piecewise affine interpolation  $\tilde{u}_n^{l-1}$ . We define a further jump interpolation of  $\xi_n$  by

$$\check{\xi}_n^l(x) = \begin{cases} \tilde{u}_n^{l-1} + (\tilde{g}_n^l - \tilde{g}_n^{l-1}) & \text{if } x \in \Delta \in \mathcal{T}_n \setminus \mathcal{C}_n^l \\ \hat{u}_n^{l-1}, & \text{if } x \in \Delta \in \mathcal{C}_n^l. \end{cases}$$

Note carefully that for technical reasons this definition slightly differs from the interpolation  $\hat{\xi}_n$  introduced in Subsection 3.2, because on triangles  $\Delta \in \mathcal{C}_n^l(\xi_n^l) \setminus \mathcal{C}_n^l$  we use the piecewise affine interpolation  $\tilde{u}_n^{l-1}$  instead of introducing a jump, and on  $\mathcal{C}_n^l$  we simply use the interpolation of  $\hat{u}_n^{l-1}$ . Recalling the representation of the elastic energy in (3.15), in particular using the fact that  $\nabla \check{\xi}_n^l = \nabla \hat{u}_n^{l-1} = 0$  on triangles in  $\mathcal{C}_n^l$ , we obtain

$$\frac{\varepsilon_n}{2} \sum_{\mathcal{T}_n \setminus \mathcal{C}_n^l} \sum_{\mathbf{v} \in \mathcal{V}} \left( \Psi(\varepsilon_n^{1/2} |(\xi_n^l)_\Delta \cdot \mathbf{v}|) - \Psi(\varepsilon_n^{1/2} |(u_n^{l-1})_\Delta \cdot \mathbf{v}|) \right) = \int_{\Omega_n} (\Psi_n^{\text{cell}}(\nabla \check{\xi}_n^l(x)) - \Psi_n^{\text{cell}}(\nabla \hat{u}_n^{l-1}(x))) dx.$$

By the estimates (4.13)–(4.14), in view of (4.12), we hence can write

$$\mathcal{E}_n^k(\xi_n^l) - \mathcal{E}_n^k(u_n^{l-1}) \leq \int_{\Omega_n} (\Psi_n^{\text{cell}}(\nabla \check{\xi}_n^l(x)) - \Psi_n^{\text{cell}}(\nabla \hat{u}_n^{l-1}(x))) dx + C(1 + \varepsilon_n \#\mathcal{C}_n^l) \varepsilon_n^{1/2} \int_{t_n^{l-1}}^{t_n^l} \|\nabla \partial_t \tilde{g}_n(\tau)\|_{L^\infty(\Omega_n)} d\tau. \quad (4.15)$$

Now, we want to prove the existence of a function  $\vartheta_n \in L^\infty([0, T] \times \Omega)$  such that  $\|\vartheta_n(s)\|_{L^2(\Omega)} \rightarrow 0$  uniformly for  $s \in [0, T]$ , and

$$\int_{\Omega_n} (\Psi_n^{\text{cell}}(\nabla \check{\xi}_n^l) - \Psi_n^{\text{cell}}(\nabla \hat{u}_n^{l-1})) dx = \int_{t_n^{l-1}}^{t_n^l} \int_{\Omega} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n(\tau) + \vartheta_n(\tau)) \cdot \nabla \partial_t \tilde{g}_n(\tau) dx d\tau. \quad (4.16)$$

Once this is achieved, (4.15)–(4.16) imply (4.6).

Let us now show (4.16). By the mean value theorem there exists a  $\rho_n^{l-1} \in [0, 1]$  such that

$$\begin{aligned} \int_{\Omega_n} (\Psi_n^{\text{cell}}(\nabla \check{\xi}_n^l) - \Psi_n^{\text{cell}}(\nabla \hat{u}_n^{l-1})) dx &= \int_{\Omega_n} (\Psi_n^{\text{cell}}(\nabla \hat{u}_n^{l-1} + \nabla \tilde{g}_n^l - \nabla \tilde{g}_n^{l-1}) - \Psi_n^{\text{cell}}(\nabla \hat{u}_n^{l-1})) dx \\ &= \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n^{l-1} + \rho_n^{l-1}(\nabla \tilde{g}_n^l - \nabla \tilde{g}_n^{l-1})) \cdot (\nabla \tilde{g}_n^l - \nabla \tilde{g}_n^{l-1}) dx. \end{aligned}$$

In the first equality we used the fact that, by definition,  $\check{\xi}_n^l = \hat{u}_n^{l-1} + \tilde{g}_n^l - \tilde{g}_n^{l-1}$ , whenever  $\check{\xi}_n^l$  and  $\hat{u}_n^{l-1}$  differ. We define the piecewise constant function  $\vartheta_n : [0, T] \rightarrow L^2(\Omega)$  for  $s \in [t_n^{l-1}, t_n^l]$  by

$$\vartheta_n(s) := \rho_n^{l-1}(\nabla \tilde{g}_n^l - \nabla \tilde{g}_n^{l-1}) = \rho_n^{l-1} \int_{t_n^{l-1}}^{t_n^l} \nabla \partial_t \tilde{g}_n(\tau) d\tau \quad \text{on } \Omega_n,$$

and 0 outside of  $\Omega_n$ . Clearly,  $(\vartheta_n)_n$  is bounded in  $L^\infty([0, T] \times \Omega)$  since the function  $\tau \mapsto \nabla \partial_t \tilde{g}_n(\tau)$  belongs to  $L^1([0, T]; L^\infty(\Omega_n))$ . Moreover, we can use the absolute continuity of the integral and  $|t_n^l - t_n^{l-1}| = \delta_n \rightarrow 0$  to conclude that  $\|\vartheta_n(s)\|_{L^2(\Omega)} \rightarrow 0$  uniformly in  $s$ . Due to the fact that  $u_n$  and  $\vartheta_n$  are constant in time on the interval  $[t_n^{l-1}, t_n^l)$ , we then obtain (4.16) by

$$\begin{aligned} \int_{\Omega_n} (\Psi_n^{\text{cell}}(\nabla \tilde{\xi}_n^l) - \Psi_n^{\text{cell}}(\nabla \hat{u}_n^{l-1})) \, dx &= \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n^{l-1} + \vartheta_n(\tau)) \cdot \left( \int_{t_n^{l-1}}^{t_n^l} \nabla \partial_t \tilde{g}_n(\tau) \, d\tau \right) \, dx \\ &= \int_{t_n^{l-1}}^{t_n^l} \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n^{l-1} + \vartheta_n(\tau)) \cdot \nabla \partial_t \tilde{g}_n(\tau) \, dx \, d\tau. \end{aligned}$$

This shows (4.16) and concludes the proof.  $\square$

As a direct consequence, we obtain the following bound on the energy.

**Corollary 4.4** (Energy bound). *Let  $t \mapsto u_n(t)$  be the discrete evolution defined in (2.11). Then, there exists a constant  $C > 0$  only depending on  $W$  and  $g$  such that*

$$\mathcal{E}_n^k(u_n^k) \leq C \quad \text{for all } k = 0, \dots, T/\delta_n - 1 \text{ and } n \in \mathbb{N}.$$

*Proof.* We show that there is a constant  $\hat{C}$  depending on  $W$  and  $g$ , but independent of  $k$  and  $n$ , such that

$$\int_0^{t_n^k} \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n(\tau)) \cdot \nabla \partial_t \tilde{g}_n(\tau) \, dx \, d\tau \leq \hat{C}. \quad (4.17)$$

Once this is achieved, we conclude as follows. In view of (3.5) and the fact that  $\Psi(R) > 0$ , we can find a constant  $\tilde{C}$  such that  $\varepsilon_n \# \mathcal{C}_n^k \leq \tilde{C} \mathcal{E}_n^k(u_n(t))$ . Therefore, we obtain from Lemma 4.3 and  $\mathcal{E}_n^0(u_n^0) \leq C$  (see Lemma 4.2)

$$(1 - \tilde{C}\eta_n) \mathcal{E}_n^k(u_n^k) \leq \hat{C} + C + \eta_n \leq C,$$

which implies the desired bound.

To see (4.17), we argue by contradiction. If (4.17) is violated, by Hölder's inequality we find for each  $\hat{C} > 0$  some  $n \in \mathbb{N}$  large enough (depending on  $\hat{C}$ ) such that

$$\text{ess sup}_{\tau \in (0, t_n^k)} \int_{\Omega_n} |D\Psi_n^{\text{cell}}(\nabla \hat{u}_n(\tau))| \, dx > \hat{C} \|\partial_t \nabla g\|_{L^1([0, T]; L^\infty(\Omega))}^{-1}.$$

Using Lemma 3.4(iii) we find  $c > 0$  such that

$$\text{ess sup}_{\tau \in (0, t_n^k)} \int_{\Omega_n} |\nabla \hat{u}_n(\tau)| \, dx \geq c \hat{C} \|\partial_t \nabla g\|_{L^1([0, T]; L^\infty(\Omega))}^{-1}.$$

For  $\hat{C}$  large enough, using Hölder's inequality this contradicts the bound in (4.2). This shows (4.17) and concludes the proof.  $\square$

**4.2. Compactness for crack sets and displacement.** Based on the energy bound in Corollary 4.4, we can pass to the limit in the crack sets and displacements by compactness arguments. To this end, it will be convenient to express some of the quantities considered in Section 3 in terms of the time  $t$  in place of the iteration step. As before, let  $\{0 = t_n^0 < t_n^1 < \dots < t_n^{T/\delta_n} = T\}$  be a time discretization of the interval  $[0, T]$ , and let  $(u_n^j)_{j < k}$  be a corresponding displacement history. Recalling (3.5) and (4.1), for  $t \in [t_n^k, t_n^{k+1})$  we define

$$\mathcal{E}_n(v_n; t) := \mathcal{E}_n^k(v_n) = \mathcal{E}_n(v_n; (u_n^j)_{j < k}) \quad (4.18)$$




FIGURE 5. The modification in the proof of Proposition 4.5.

for each  $v_n: \mathcal{L}_n(\Omega) \rightarrow \mathbb{R}$ . We use similar notation for the parts of the energy introduced in (3.10). Given the evolution  $t \mapsto u_n(t)$  defined in (2.11) and recalling (3.33), for  $t \in [t_n^k, t_n^{k+1})$  we define the jump sets

$$K_n^i(t) := K_n^{k,i}(u_n(t)), \quad K_n^L(t) := \bigcup_{i=1}^3 K_n^i(t), \quad K_n^S(t) := K_n^{k,S}(u_n(t)). \quad (4.19)$$

One drawback of the definition of the sets in (4.19) is the ambiguity that comes from the three different variants  $i = 1, 2, 3$ . We will now prove that this ambiguity does not affect the limit, i.e., for a suitable subsequence, the  $\sigma^{3/2}$ -limits of all three variants and their union coincide.

**Proposition 4.5.** *Let  $t \mapsto u_n(t)$  be the evolution defined in (2.11). There exists an increasing set function  $t \mapsto K(t)$  on  $[0, T]$  and a subsequence (not relabeled) such that for every  $t \in [0, T]$  we have*

$$K_n^i(t) \quad \sigma^{3/2}\text{-converges to } K(t) \text{ in } \Omega \quad \text{for all } i = 1, 2, 3.$$

$$K_n^L(t) \quad \sigma^{3/2}\text{-converges to } K(t) \text{ in } \Omega.$$

*Proof.* For notational convenience, we write  $p = 3/2$  in the proof. Note that we can deduce by Proposition 3.5 and Corollary 4.4 that  $\mathcal{H}^1(K_n^L(t))$  is uniformly bounded with respect to  $n$  and  $t$ . Hence, we can apply Theorem 3.11 and thereby get a subsequence  $(K_n^L)_n$  (not relabeled) and an increasing set function  $t \mapsto K(t)$  such that

$$K_n^L(t) \quad \sigma^p\text{-converges to } K(t) \text{ in } \Omega,$$

for all  $t \in [0, T]$ . We can repeat this argument for the variants  $K_n^i(t)$  and, since  $K_n^i(t) \subset K_n^L(t)$  for  $i = 1, 2, 3$ , the  $\sigma^p$ -limits  $K^i(t)$  exist, up to passing to a further subsequence (not relabeled), and we have  $K^i(t) \subset K(t)$  by Remark 3.10(c). It suffices to show that  $K^i(t) \supset K(t)$  for each  $t \in [0, T]$ .

Without restriction we show the argument for  $i = 1$ . We fix a time  $t \in [0, T]$ , and for given  $n$  we choose  $k$  such that  $t \in [t_n^k, t_n^{k+1})$ . By the definition of  $\sigma^p$ -convergence we find a sequence  $u_n \rightarrow u$  weakly in  $SBV^p(\Omega)$  such that  $S(u_n) \subset K_n^L(t)$  and  $K(t) \subset S(u)$ . We now construct a modification  $v_n$  with  $S(v_n) \subset K_n^L(t)$ . By construction in (3.12), see also (3.9), we have  $K_n^L(t) \subset \bigcup_{\Delta \in \mathcal{C}_n^{k,L}(u_n^k)} \Delta$ . We define  $v_n = u_n$  outside of these triangles and now consider each triangle in  $\Delta \in \mathcal{C}_n^{k,L}(u_n^k)$  separately.

If  $\Delta \in \mathcal{C}_n^{k,2}(u_n)$ , then we set  $v_n = u_n$  in  $\Delta$ . If  $\Delta \in \mathcal{C}_n^{k,3}(u_n)$ , we know that  $S(u_n)$  lies in  $h_\Delta^1 \cup h_\Delta^2 \cup h_\Delta^3$ . We now explain that the function can be modified such that it jumps only on  $h_\Delta^2 \cup h_\Delta^3$ . In fact, as  $u_n \in W^{1,p}(\Omega \setminus \overline{S(u_n)})$ , we can extend  $u_n$  from the sub-triangle with edge  $h_\Delta^1$  to the middle triangle, see Figure 5, i.e., we replace  $u_n$  by  $v_n$  inside the middle triangle such that  $v_n$  is continuous on  $h_\Delta^1$ .

This construction leads to  $S(v_n) \cap \Delta \subset K_n^1(t)$ , with  $\|v_n\|_{W^{1,p}(\Delta)} \leq C\|u_n\|_{W^{1,p}(\Delta)}$ . In turn, we obtain  $\|\nabla v_n\|_{L^p(\Omega_n)} \leq C$ .

Now,  $u_n \rightarrow u$  in  $L^1(\Omega)$  along with the fact that  $\mathcal{L}^2(\{u_n \neq v_n\}) \leq \varepsilon_n^2 \#\mathcal{C}_n^k(u_n^k) \rightarrow 0$  also shows that  $v_n \rightarrow u$  in  $L^1(\Omega)$ , i.e.,  $v_n \rightarrow u$  weakly in  $SBV^p(\Omega)$ . Using part (i) of Definition 3.9 and  $S(v_n) \overset{\sim}{\subset} K_n^1(t)$  we conclude  $K(t) \overset{\sim}{=} S(u) \overset{\sim}{\subset} K^1(t)$ .  $\square$

We now proceed with a compactness result for displacements and lower semicontinuity for the energies.

**Proposition 4.6** (Compactness and lower semicontinuity). *For a fixed  $t \in [0, T]$ , let  $(u_n(t))_{n \in \mathbb{N}}$  be the evolution defined in (2.11). Let  $K(t)$  be the  $\sigma^{3/2}$ -limit of  $K_n^L(t)$  as given in Proposition 4.5. Then, there exists a subsequence  $(n_l)_l$  depending on  $t$  and some  $u \in SBV^2(\Omega)$  such that  $u_{n_l}(t)$  AC-converges to  $u$  as  $l \rightarrow \infty$ ,  $\liminf_{n \rightarrow \infty} \mathcal{E}_n(u_n(t); t) = \liminf_{l \rightarrow \infty} \mathcal{E}_{n_l}(u_{n_l}(t); t)$ , and*

$$\liminf_{l \rightarrow \infty} \mathcal{E}_{n_l}(u_{n_l}(t); t) \geq \liminf_{l \rightarrow \infty} \mathcal{E}_{n_l}^{\text{ela}}(u_{n_l}(t); t) + \liminf_{l \rightarrow \infty} \mathcal{E}_{n_l}^{\text{cra}}(u_{n_l}(t); t) \geq \mathcal{E}(u(t), K(t)).$$

*Proof.* The compactness of the displacements follows from Proposition 3.5, by using Corollary 4.4 and the  $L^\infty$ -bound on the displacements given by Lemma 4.1. In particular, the subsequence  $(n_l)_l$  can be chosen such that  $\liminf_{n \rightarrow \infty} \mathcal{E}_n(u_n(t); t) = \liminf_{l \rightarrow \infty} \mathcal{E}_{n_l}(u_{n_l}(t); t)$  holds. Then, in view of (3.10), Lemma 3.6, and the nonnegativity of  $\mathcal{E}_n^{\text{rem}}$  and  $\mathcal{E}_n^{\text{bdy}}$ , we only need to show that

$$\liminf_{n \rightarrow \infty} \mathcal{E}_n^{\text{cra}}(u_n(t); t) \geq \int_{K(t)} \varphi(\nu_{K(t)}) d\mathcal{H}^1.$$

Observe that  $\sup_k \#\mathcal{C}_n^k(u_n^k) \leq C\varepsilon_n^{-1}$  for a constant independent of  $n \in \mathbb{N}$ , by Corollary 4.4 and (3.27). Then, using Proposition 3.2,  $R_n \rightarrow \infty$  by (3.7), and  $\lim_{s \rightarrow \infty} \Psi(s) = \kappa$ , we have

$$\liminf_{n \rightarrow \infty} \mathcal{E}_n^{\text{cra}}(u_n(t); t) = \liminf_{n \rightarrow \infty} \frac{\kappa}{\sqrt{3}} \sum_{i=1}^3 \int_{K_n^i(t)} \varphi_i(\nu_{K_n^i(t)}) d\mathcal{H}^1,$$

where  $\varphi_\alpha(\nu) := (|\mathbf{v}_\beta \cdot \nu| + |\mathbf{v}_\gamma \cdot \nu|)$  for  $\nu \in \mathbb{S}^1$ , and for  $\alpha, \beta, \gamma \in \{1, 2, 3\}$  pairwise distinct. We now employ Proposition 4.5 and Theorem 3.11, i.e., the lower semicontinuity with respect to  $\sigma^{3/2}$ -convergence, to obtain

$$\liminf_{n \rightarrow \infty} \mathcal{E}_n^{\text{cra}}(u_n(t); t) \geq \frac{\kappa}{\sqrt{3}} \sum_{i=1}^3 \int_{K(t)} \varphi_i(\nu_{K(t)}) d\mathcal{H}^1.$$

By the definition of  $\varphi$  in (2.15), the proof is concluded.  $\square$

**4.3. Proof of the main result.** In this subsection we give the proof of Theorem 2.2. As a last preparation, we need the following stability result for our time-discrete atomistic energy. Recall the splitting of the energy in (3.10). For  $t \in [0, T]$  and  $n \in \mathbb{N}$ , we let  $k(t)$  be the ( $n$ -dependent) index such that  $t \in [t_n^{k(t)}, t_n^{k(t)+1})$ .

**Theorem 4.7** (Stability). *Let  $t \mapsto u_n(t)$  be the evolution defined by (2.11) and let  $K(t)$  be the  $\sigma^{3/2}$ -limit of  $K_n^L(t)$ . For each  $\psi \in SBV^2(\Omega)$  with  $\psi = g(t)$  on  $\Omega \setminus \bar{U}$  there exists a sequence  $(\psi_n)_n$  of discrete displacements with  $\psi_n \in \mathcal{A}_n(g(t_n^{k(t)}))$  such that  $\psi_n$  AC-converges to  $\psi$  and*

$$\limsup_{n \rightarrow \infty} \left( \mathcal{E}_n^{\text{cra}}(\psi_n; t) + \mathcal{E}_n^{\text{rem}}(\psi_n; t) - (\mathcal{E}_n^{\text{cra}}(u_n(t); t) + \mathcal{E}_n^{\text{rem}}(u_n(t); t)) \right) \leq \int_{S(\psi) \setminus K(t)} \varphi(\nu_\psi) d\mathcal{H}^1, \quad (4.20)$$

where  $\nu_\psi$  denotes the normal to the jump set  $S(\psi)$  of  $\psi$ . Moreover,

$$\lim_{n \rightarrow \infty} \mathcal{E}_n^{\text{ela}}(\psi_n; t) = \int_{\Omega} \Phi(\nabla \psi) dx, \quad (4.21)$$

where  $\Phi$  is given in (2.14), and we have

$$\lim_{n \rightarrow \infty} \mathcal{E}_n^{\text{bdy}}(\psi_n; t) = 0. \quad (4.22)$$

This result is essential to pass from the minimality condition (2.10) in the atomistic setting to the global stability in (2.16). For this reason, estimates of this kind are often referred to as *stability of unilateral minimizers*, see e.g. [40]. The proof will be deferred to Section 5. The strategy follows the one of the *jump transfer lemma* introduced in FRANCFORT AND LARSEN [31], see the works [22, 23, 34, 39] for several variants. Our situation is more delicate for two reasons. First, due to the underlying atomistic nature, special care is needed in the construction of the sequence  $(\psi_n)_n$ , in particular concerning the corresponding ‘broken’ springs. Secondly, whereas in [31] only the location of the crack is relevant, which in our notation means  $K_n^L(t)$ , our atomistic energy also takes into account *how much each spring* was deformed at former times. Therefore, we need to verify that the exact value of the memory variable is indeed negligible in the limit, see also [39] for a similar problem.

For the proof, we will also need the following corollary, which formally follows from Theorem 4.7 by choosing  $u_n(t) \equiv 0$  and  $K(t) = 0$ . Recall the energy  $\mathcal{E}_n$  defined in (2.6).

**Corollary 4.8** (Recovery sequence). *For each  $\psi \in SBV^2(\Omega)$  with  $\psi = g(t)$  on  $\Omega \setminus \overline{U}$  there exists a sequence  $(\psi_n)_n$  with  $\psi_n \in \mathcal{A}_n(g(t_n^{k(t)}))$  such that  $\psi_n$  AC-converges to  $\psi$  and*

$$\limsup_{n \rightarrow \infty} \mathcal{E}_n(\psi_n) \leq \mathcal{E}(\psi, S(\psi)).$$

We are now in the position to prove the main result of this paper.

*Proof of Theorem 2.2.* We split the proof into five steps.

*Step 1: Limiting evolution.* Let  $K(t)$  be the  $\sigma^{3/2}$ -limit of  $K_n^L(t)$  for  $t \in [0, T]$  given by Proposition 4.5. Since  $t \mapsto K(t)$  is an increasing set function, i.e.,  $K(t_1) \tilde{\subset} K(t_2)$  for all  $0 \leq t_1 \leq t_2 \leq T$ , the irreversibility condition (a), is satisfied. For each fixed time  $t \in [0, T]$ , by virtue of Proposition 4.6, there exists  $u(t) \in SBV^2(\Omega)$  and a subsequence depending on  $t$  (not relabeled) such that  $u_n(t)$  AC-converges to  $u(t)$ . In particular, as  $u_n(t) \in \mathcal{A}_n(g(t_n^{k(t)}))$ , this also shows that  $u(t) = g(t)$  on  $\Omega \setminus \overline{U}$ . In view of Corollary 4.4 and Lemma 4.1 we can apply Proposition 3.13 to deduce  $S(u(t)) \tilde{\subset} K(t)$ . This shows that  $u(t) \in AD(g(t), K(t))$  for all  $t \in [0, T]$ .

Next, we check that the mapping  $t \rightarrow (u(t), K(t))$  is an irreversible quasi-static crack evolution. For this, we need to confirm (2.16) and (2.17) which is content of Step 2 and Step 3–4, respectively. In the last step, we then prove the convergence of energies and show that the crack sets also converge in the sense of  $\sigma$ -convergence, see Definition 3.8.

*Step 2: Stability* (2.16). Fix  $t \in [0, T]$ . Let  $\psi \in AD(g(t), H)$  be an admissible competitor for a set  $H$  with  $K(t) \tilde{\subset} H$ . We employ Theorem 4.7 to obtain a sequence of discrete displacements  $\psi_n \in \mathcal{A}_n(g(t_n^{k(t)}))$  such that (4.20)–(4.22) are valid. By the minimality property of the solution  $u_n(t)$ , see (2.9)–(2.10), and the shorthand notation in (4.18) we have

$$\mathcal{E}_n(u_n(t); t) \leq \mathcal{E}_n(\psi_n; t).$$

We now split the energy on both sides like in (3.10): Subtracting the crack energy and the remainder term on the left-hand side and using that  $\mathcal{E}_n^{\text{bdy}}$  is nonnegative, we obtain

$$\mathcal{E}_n^{\text{ela}}(u_n(t); t) \leq \mathcal{E}_n^{\text{ela}}(\psi_n; t) + \mathcal{E}_n^{\text{cra}}(\psi_n; t) - \mathcal{E}_n^{\text{cra}}(u_n(t); t) + \mathcal{E}_n^{\text{rem}}(\psi_n; t) - \mathcal{E}_n^{\text{rem}}(u_n(t); t) + \mathcal{E}_n^{\text{bdy}}(\psi_n; t). \quad (4.23)$$

Passing to the limit in (4.23), by employing (4.20)–(4.22), we find

$$\limsup_{n \rightarrow \infty} \mathcal{E}_n^{\text{ela}}(u_n(t); t) \leq \int_{\Omega} \Phi(\nabla \psi) \, dx + \int_{S(\psi) \setminus K(t)} \varphi(\nu_{\psi}) \, d\mathcal{H}^1.$$

Since  $u_n(t)$  AC-converges to  $u(t)$  (for a  $t$ -dependent subsequence, not relabeled), employing also Lemma 4.1 and Corollary 4.4, we can use Lemma 3.6 and derive

$$\int_{\Omega} \Phi(\nabla u(t)) \, dx \leq \limsup_{n \rightarrow \infty} \mathcal{E}_n^{\text{ela}}(u_n(t); t) \leq \int_{\Omega} \Phi(\nabla \psi) \, dx + \int_{S(\psi) \setminus K(t)} \varphi(\nu_{\psi}) \, d\mathcal{H}^1. \quad (4.24)$$

Since  $\psi \in AD(g(t), H)$ , we have  $S(\psi) \tilde{\subset} H$  and thus

$$\int_{\Omega} \Phi(\nabla u(t)) \, dx \leq \int_{\Omega} \Psi(\nabla \psi) \, dx + \int_{H \setminus K(t)} \varphi(\nu_H) \, d\mathcal{H}^1.$$

Since  $K(t) \tilde{\subset} H$ , we hence conclude

$$\int_{\Omega} \Phi(\nabla u(t)) \, dx + \int_{K(t)} \varphi(\nu_{K(t)}) \, d\mathcal{H}^1 \leq \int_{\Omega} \Phi(\nabla \psi) \, dx + \int_H \varphi(\nu_H) \, d\mathcal{H}^1.$$

Recalling the definition of the limiting energy (2.13), this shows (2.16).

In particular, for  $t = 0$ , denoting by  $(\psi_n^0)_n$  a recovery sequence for  $u(0)$  from Corollary 4.8, the minimality property in (2.9) and Proposition 4.6 yield

$$\mathcal{E}(u(0), K(0)) \leq \liminf_{n \rightarrow \infty} \mathcal{E}_n(u_n(0)) \leq \limsup_{n \rightarrow \infty} \mathcal{E}_n(\psi_n^0) \leq \mathcal{E}(u(0), S(u(0))).$$

This along with  $u(0) \in AD(g(0), K(0))$ , i.e.,  $S(u(0)) \tilde{\subset} K(0)$ , shows

$$\lim_{n \rightarrow \infty} \mathcal{E}_n(u_n(0)) = \mathcal{E}(u(0), K(0)) = \mathcal{E}(u(0), S(u(0))). \quad (4.25)$$

*Step 3: Convergence of work by applied boundary load.* Next, we check that for all  $t \in [0, T]$

$$\lim_{n \rightarrow \infty} \int_0^t \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n(\tau)) \cdot \nabla \partial_t \tilde{g}_n(\tau) \, dx \, d\tau = \int_0^t \int_{\Omega} D\Phi(\nabla u(\tau)) \cdot \nabla \partial_t g(\tau) \, dx \, d\tau. \quad (4.26)$$

Let  $\tau \in [0, t)$ . Using the stability estimate (4.24) for  $\psi = u(\tau) \in AD(g(\tau), K(\tau))$ , and employing  $S(u(\tau)) \tilde{\subset} K(\tau)$  we get

$$\lim_{n \rightarrow \infty} \mathcal{E}_n^{\text{ela}}(u_n(\tau); \tau) = \int_{\Omega} \Phi(\nabla u(\tau)) \, dx \quad \text{for all } \tau \in [0, t).$$

Then (3.15) and Lemma 3.7 imply  $\chi_{\Omega_n} \nabla \hat{u}_n(\tau) \rightarrow \nabla u(\tau)$  strongly in  $L^2(\Omega; \mathbb{R}^2)$ . Thus, as in Step 1 of the proof of Lemma 4.3, we can employ [40, Lemma 2.4] to find

$$\left| \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n(\tau)) \cdot \nabla \partial_t g(\tau) \, dx - \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla u(\tau)) \cdot \nabla \partial_t g(\tau) \, dx \right| \rightarrow 0,$$

where we also used Lemma 3.4(iii),(iv) and the fact that  $\nabla \partial_t g(\tau) \in L^2(\Omega; \mathbb{R}^2)$  for almost every  $\tau \in [0, t)$ . Since  $\nabla u(\tau) \in L^2(\Omega; \mathbb{R}^2)$  and  $D\Psi_n^{\text{cell}} \rightarrow D\Phi$  pointwise, we also get by dominated convergence

$$\left| \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n(\tau)) \cdot \nabla \partial_t g(\tau) \, dx - \int_{\Omega_n} D\Phi(\nabla u(\tau)) \cdot \nabla \partial_t g(\tau) \, dx \right| \rightarrow 0.$$

At this stage, we can repeat the argumentation in (4.8), in particular using Lemma 3.4(iii), the fact that  $\chi_{\Omega_n} \nabla \partial_t \tilde{g}_n$  converges strongly to  $\nabla \partial_t g$  in  $L^1([0, T]; L^\infty(\Omega; \mathbb{R}^2))$ , and the uniform bound on  $\nabla \hat{u}_n(\tau)$  in  $L^2(\Omega_n; \mathbb{R}^2)$ , see (4.2), to obtain (4.26).

*Step 4: Energy balance* (2.17). For  $t \in [t_n^k, t_n^{k+1})$ , by Lemma 4.3 we find  $(\eta_n)_n$  with  $\eta_n \rightarrow 0$  such that

$$\begin{aligned} \mathcal{E}_n^k(u_n(t)) - \mathcal{E}_n^0(u_n^0) &\leq \int_0^{t_n^k} \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n(\tau)) \cdot \nabla \partial_t \tilde{g}_n(\tau) \, dx \, d\tau + (1 + \varepsilon_n \# \mathcal{C}_n^k) \eta_n \\ &\leq \int_0^t \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n(\tau)) \cdot \nabla \partial_t \tilde{g}_n(\tau) \, dx \, d\tau + C\eta_n + C\delta_n, \end{aligned}$$

where in the second step we have used Proposition 3.5 and Corollary 4.4 to get  $\# \mathcal{C}_n^k \leq C\varepsilon_n^{-1}$ . Furthermore, we used the first estimate in (4.8) and  $g \in W^{1,1}([0, T]; W^{2,\infty}(\Omega))$  to obtain a uniform control on the integrand, which allows to estimate the integral from  $t_n^k$  to  $t$  in terms of  $C\delta_n$ . Combining this with Proposition 4.6 and (4.25), and using that  $\mathcal{E}_n^0 = \mathcal{E}_n$ , we get by (4.26)

$$\begin{aligned} \mathcal{E}(u(t), K(t)) &\leq \liminf_{n \rightarrow \infty} \mathcal{E}_n(u_n(t); t) \leq \limsup_{n \rightarrow \infty} \mathcal{E}_n^0(u_n^0) + \limsup_{n \rightarrow \infty} \int_0^t \int_{\Omega_n} D\Psi_n^{\text{cell}}(\nabla \hat{u}_n(\tau)) \cdot \nabla \partial_t \tilde{g}_n(\tau) \, dx \, d\tau \\ &= \mathcal{E}(u(0), K(0)) + \int_0^t \int_{\Omega} D\Phi(\nabla u(\tau)) \cdot \nabla \partial_t g(\tau) \, dx \, d\tau. \end{aligned} \quad (4.27)$$

Thus, it remains to prove the reverse inequality

$$\mathcal{E}(u(t), K(t)) \geq \mathcal{E}(u(0), K(0)) + \int_0^t \int_{\Omega} D\Phi(\nabla u(\tau)) \cdot \nabla \partial_t g(\tau) \, dx \, d\tau.$$

This can be done by repeating the arguments in [22, Lemma 7.1] which are based on the global stability result (2.16) proved in *Step 2*, as well as approximation of integrals by Riemann sums and their convergence, see [22, Lemmas 5.12 and 5.7].

*Step 5: Energy convergence and  $\sigma$ -convergence of crack sets.* Combining (4.27) with the energy balance (2.17) we find

$$\mathcal{E}(u(t), K(t)) = \lim_{n \rightarrow \infty} \mathcal{E}_n(u_n(t); t) \quad \text{for all } t \in [0, T], \quad (4.28)$$

which shows (2.21). To conclude the proof it remains to show that (2.19) holds.

Consider the crack sets  $K_n(t)$  defined in (2.12). We note that  $\mathcal{H}^1(K_n(t)) \leq C$  for all  $t \in [0, T]$  and  $n \in \mathbb{N}$  by (3.27), Lemma 4.1, and Corollary 4.4. As  $K_n(t)$  is increasing in  $t$ , Remark 3.12 implies that there is a subsequence (not relabeled) and an increasing set function  $K^\sigma(t)$  such that  $K_n(t)$   $\sigma$ -converges to  $K^\sigma(t)$  for each  $t \in [0, T]$ . To conclude the proof, it suffices to check that  $K^\sigma(t) \doteq K(t)$  for all  $t \in [0, T]$ , where  $K(t)$  is given in Proposition 4.5. We note that  $K^\sigma(t) \dot{\supset} K(t)$  by Remark 3.10(d), so it suffices to check that  $K^\sigma(t) \check{\subset} K(t)$ .

Recall the splitting of  $\mathcal{E}_n(u_n(t); t)$  in (3.10). The energy convergence in (4.28) along with the lower-semicontinuity for  $\mathcal{E}_n^{\text{cra}}$  and  $\mathcal{E}_n^{\text{ela}}$  from Lemma 3.6 and Proposition 4.6, and the fact that  $\mathcal{E}_n^{\text{rem}}$  and  $\mathcal{E}_n^{\text{bdy}}$  are nonnegative show

$$\lim_{n \rightarrow \infty} \mathcal{E}_n^{\text{rem}}(u_n(t); t) = \lim_{n \rightarrow \infty} \mathcal{E}_n^{\text{bdy}}(u_n(t); t) = 0$$

and

$$\liminf_{n \rightarrow \infty} \mathcal{E}_n^{\text{ela}}(u_n(t); t) = \int_{\Omega} \Phi(\nabla u(t)) \, dx, \quad \liminf_{n \rightarrow \infty} \mathcal{E}_n^{\text{cra}}(u_n(t); t) = \int_{K(t)} \varphi(\nu_{K(t)}) \, d\mathcal{H}^1. \quad (4.29)$$

In view of Remark 3.1, we obtain

$$\mathcal{H}^1(K_n(t) \setminus K_n^{\text{L}}(t)) \leq \frac{3}{2} \varepsilon_n \# (\mathcal{C}_n^k(u_n^k) \setminus \mathcal{C}_n^{k,\text{L}}(u_n^k)) + \varepsilon_n \# \{ \mathcal{C}_n^{k,2}(u_n^k) : M_{n,\Delta}^{k,\nu_i}(u_n^k) > R \text{ for all } i = 1, 2, 3 \},$$

where the second sum is related to the triangles with (in the notation of Remark 3.1)  $\mu_{n,\Delta}^k = 3$  and  $\nu_{n,\Delta}^k = 2$ . Recalling the definition of  $\mathcal{E}_n^{\text{rem}}$  in (3.11) and using  $\Psi(R) > 0$

$$\#(\mathcal{C}_n^k(u_n^k) \setminus \mathcal{C}_n^{k,L}(u_n^k)) + \#\{\mathcal{C}_n^{k,2}(u_n^k) : M_{n,\Delta}^{k,\mathbf{v}_i}(u_n^k) > R \text{ for all } i = 1, 2, 3\} \leq C\varepsilon_n^{-1}\mathcal{E}_n^{\text{rem}}(u_n(t); t).$$

This shows  $\mathcal{H}^1(K_n(t) \setminus K_n^L(t)) \rightarrow 0$  as  $n \rightarrow \infty$ . By Remark 3.10(b) we therefore get that the  $\sigma$ -limit of  $K_n(t)$  and  $K_n^L(t)$  coincide, i.e.,  $K_n^L(t)$   $\sigma$ -converges to  $K^\sigma(t)$  for all  $t \in [0, T]$ . Arguing as in the proofs of Propositions 4.5 and 4.6, and using Remark 3.12 in place of Theorem 3.11, we get

$$\liminf_{n \rightarrow \infty} \mathcal{E}_n^{\text{cra}}(u_n(t); t) \geq \int_{K^\sigma(t)} \varphi(\nu_{K^\sigma(t)}) d\mathcal{H}^1.$$

This along with (4.29) and the inclusion  $K(t) \tilde{\subset} K^\sigma(t)$  shows  $K^\sigma(t) \tilde{=} K(t)$  and concludes the proof.  $\square$

## 5. PROOF OF THE STABILITY RESULT

This section is entirely devoted to the proof of the stability result in Theorem 4.7 and Corollary 4.8.

**Density argument.** We start by observing that it suffices to prove the statement for functions  $\psi$  with more regularity, employing a suitable density argument. Let  $\mathcal{W}(\Omega) \subset SBV(\Omega)$  be the collection of functions  $v$  such that  $S(v)$  is closed and included in a finite union of closed and connected pieces of  $C^1$ -curves and  $v$  lies in  $W^{2,\infty}(\Omega \setminus S(v))$ . For each  $v \in SBV^2(\Omega)$  with  $v = g(t)$  on  $\Omega \setminus \bar{U}$  we can choose a sequence  $(v_n)_n \subset \mathcal{W}(\Omega)$  with  $v_n = g(t)$  on  $\Omega \setminus \bar{U}$  such that

$$v_n \rightarrow v \text{ in } L^1(\Omega), \quad \|\nabla v_n - \nabla v\|_{L^2(\Omega)} \rightarrow 0, \quad \mathcal{H}^1(S(v_n) \setminus S(v)) + \mathcal{H}^1(S(v) \setminus S(v_n)) \rightarrow 0.$$

This follows from [28, Theorem C] (see also [19, Theorem 1.1] for the control on the Sobolev norms of  $\nabla v_n$ ), where a construction similar to the one performed in [38, Proposition 2.5] ensures that the boundary values are attained along the sequence. We also refer to [33, Theorem 3.2] for an analogous statement and proof for *GSBD*-functions.

With this density result at hand, we observe that it suffices to construct a sequence as in Theorem 4.7 for a function  $\psi \in \mathcal{W}(\Omega)$  with  $\psi = g(t)$  in a neighborhood of  $\Omega \setminus \bar{U}$  in  $\Omega$ . The general case then follows by a standard diagonal argument. Further, we fix  $\theta > 0$  and observe that it suffices to construct a sequence  $(\psi_n)_n$  of discrete displacements with  $\psi_n \in \mathcal{A}_n(g(t))$  such that

$$\limsup_{n \rightarrow \infty} \|\chi_{\Omega_n} \tilde{\psi}_n - \psi\|_{L^1(\Omega)} \leq C\theta, \quad (5.1)$$

$$\limsup_{n \rightarrow \infty} \left( (\mathcal{E}_n^{\text{cra}}(\psi_n; t) + \mathcal{E}_n^{\text{rem}}(\psi_n; t)) - (\mathcal{E}_n^{\text{cra}}(u_n(t); t) + \mathcal{E}_n^{\text{rem}}(u_n(t); t)) \right) \leq \int_{S(\psi) \setminus K(t)} \varphi(\nu_\psi) d\mathcal{H}^1 + C\theta, \quad (5.2)$$

$$\limsup_{n \rightarrow \infty} \mathcal{E}_n^{\text{ela}}(\psi_n; t) \leq \int_{\Omega} \Phi(\nabla \psi) dx + C\theta, \quad (5.3)$$

$$\lim_{n \rightarrow \infty} \mathcal{E}_n^{\text{bdy}}(\psi_n; t) = 0, \quad (5.4)$$

where  $\Phi$  and  $\varphi$  are given in (2.14) and (2.15), respectively, and  $C > 0$  is a universal constant. Then, the statement follows again by a diagonal argument, sending  $\theta \rightarrow 0$ . Note that, strictly speaking, proceeding in this way, the boundary values  $g(t_n^{k(t)})$  are not satisfied. Therefore, we eventually need to replace the sequence  $(\psi_n)_n$  by  $\psi_n(x) - g(t, x) + g(t_n^{k(t)}, x)$  for  $x \in \mathcal{L}_n(\Omega)$ . Due to the regularity of  $g$  and the Lipschitz continuity of  $\Psi$ , this still leads to (5.1)–(5.4), cf. also (4.6) for a similar estimate.

Now, we fix  $\psi \in \mathcal{W}(\Omega)$  with  $\psi = g(t)$  on  $\Omega \setminus \bar{U}$ , and construct a sequence  $(\psi_n)_n$  satisfying (5.1)–(5.4) and  $\psi_n \in \mathcal{A}_n(g(t))$ .

**Besicovitch covering.** Following closely the procedure in [31], we introduce a fine cover of the jump set  $S(\psi)$  with closed squares satisfying certain additional properties. For simplicity we only treat the case that  $\mathcal{H}^1(S(\psi) \cap \partial_D U) = 0$  (no jump along the boundary), for the general case follows by minor adaptations of the construction at the boundary (see [31]) which would merely overburden notation in the sequel. As a consequence, we can assume that all these squares are contained in  $U$ . We furthermore denote by  $\nu_\psi$  a measure-theoretic normal at  $S(\psi)$ .

Because of the  $\sigma^{3/2}$ -convergence of  $K_n^L(t)$  to  $K(t)$ , we have a function  $v \in SBV^{3/2}(\Omega)$  and a sequence  $(v_n)_n \subset SBV^{3/2}(\Omega)$  such that  $v_n \rightarrow v$  in  $L^1(\Omega)$ ,  $\nabla v_n \rightharpoonup \nabla v$  weakly in  $L^{3/2}(\Omega; \mathbb{R}^2)$ ,  $S(v) \doteq K(t)$ , and  $S(v_n) \tilde{\subset} K_n^L(t)$ . We can choose a suitable subset  $G_j \subset S(v) = K(t)$  as done preceding [31, (2.2)] such that

$$\mathcal{H}^1(S(v) \setminus G_j) = \mathcal{H}^1(K(t) \setminus G_j) \leq \theta \quad (5.5)$$

and for each  $x \in G_j \cap S(\psi)$  we consider closed squares  $Q_r(x)$  with sidelength  $2r$  and two sides orthogonal to  $\nu_\psi(x)$  which are contained in  $U$  and satisfy [31, (2.3), (2.5)].

For closed squares  $Q_r(x) \subset U$  with a center  $x \in S(\psi) \setminus K(t)$ , still oriented in direction  $\nu_\psi(x)$ , we can assume that for  $\mathcal{H}^1$ -a.e.  $x \in S(\psi) \setminus K(t)$  and for  $r$  sufficiently small it holds

$$\mathcal{H}^1(K(t) \cap Q_r(x)) \leq \theta r. \quad (5.6)$$

This is possible by the fact that  $K(t)$  has  $\mathcal{H}^1$ -density 0 almost everywhere in  $S(\psi) \setminus K(t)$ . As  $S(\psi)$  is contained in a finite union of closed  $C^1$ -curves, for a.e.  $x \in S(\psi)$ , possibly passing to smaller  $r$ , the above squares can be chosen such that they also satisfy

$$2r \leq \mathcal{H}^1(S(\psi) \cap Q_r(x)) \leq 4r, \quad (5.7)$$

$$S(\psi) \cap Q_{r(1+\theta)}(x) \subset \{y : |(y-x) \cdot \nu_\psi(x)| \leq \theta r\}, \quad (5.8)$$

$$|\nu_\psi(y) - \nu_\psi(x)| \leq \theta \quad \forall y \in S(\psi) \cap Q_r(x). \quad (5.9)$$

With this, we obtain a fine cover of  $\Gamma := (G_j \cap S(\psi)) \cup (S(\psi) \setminus K(t))$  to which we can apply the Besicovitch covering theorem with respect to the Radon measure  $\mathcal{L}^2 + \mathcal{H}^1|_\Gamma$ . For  $\theta > 0$  fixed as above, we hereby find a finite and disjoint subcollection  $\mathcal{B} := (Q_{r_i}(x_i))_i$ , or shortly denoted by  $(Q_i)_i$ , such that  $(Q_i)_i$  satisfy the properties mentioned above, in particular (5.7)–(5.9) and for  $x_i \notin K(t)$  also (5.6), as well as

$$\mathcal{L}^2\left(\bigcup_{\mathcal{B}} Q_i\right) \leq \theta^2, \quad \mathcal{H}^1\left(\Gamma \setminus \bigcup_{\mathcal{B}} Q_i\right) \leq \theta. \quad (5.10)$$

Here and in the following, we use  $\bigcup_{\mathcal{B}} Q_i$  as a shorthand for  $\bigcup_{Q_{r_i}(x_i) \in \mathcal{B}} Q_{r_i}(x_i)$ . Note that this implies

$$\int_{\bigcup_{\mathcal{B}} Q_i} |\nabla v| \, dx \leq C\theta, \quad \mathcal{H}^1(S(\psi) \setminus \bigcup_{\mathcal{B}} Q_i) \leq 2\theta. \quad (5.11)$$

Indeed, the first property follows from (5.10), Hölder's inequality, and  $v \in SBV^{3/2}(\Omega)$ , whereas the second is a consequence of (5.10), together with the definition of  $\Gamma$  and (5.5). Without further notice, we will frequently use that the squares are pairwise disjoint.

By  $\mathcal{B}_{\text{good}} \subset \mathcal{B}$  we denote the collection of squares  $Q_i = Q_{r_i}(x_i)$  with  $x_i \in S(\psi) \setminus K(t)$ , and similarly we let  $\mathcal{B}_{\text{bad}} \subset \mathcal{B}$  be the collection of all squares  $Q_i = Q_{r_i}(x_i)$  with  $x_i \in S(\psi) \cap G_j$ . Clearly, we have  $\mathcal{B} = \mathcal{B}_{\text{good}} \cup \mathcal{B}_{\text{bad}}$ . We also define the sets

$$B_{\text{good}} = \bigcup_{\mathcal{B}_{\text{good}}} Q_i, \quad B_{\text{bad}} = \bigcup_{\mathcal{B}_{\text{bad}}} Q_i.$$




FIGURE 6. Illustration of some notions used in the construction. Note that  $L_i^n$ ,  $R_i^n$ , and  $H_i^{n,\pm}$  are independent of  $n$  if  $Q_i \in \mathcal{B}_{\text{good}}$ .

**Good squares.** Due to (5.6) and (5.7), there is a universal constant  $C > 0$  such that

$$\mathcal{H}^1(K(t) \cap B_{\text{good}}) \leq C\mathcal{H}^1(S(\psi))\theta. \quad (5.12)$$

Next, for technical reasons we straighten the jump set  $S(\psi)$  inside of  $B_{\text{good}}$ , i.e., we construct a function  $\phi$  which jumps only on straight segments inside  $B_{\text{good}}$ . To this end, for  $Q_i = Q_{r_i}(x_i) \in \mathcal{B}_{\text{good}}$  we define

$$H_i := \{y \in Q_i : (y-x_i) \cdot \nu_\psi(x_i) = 0\}, \quad R_i := \{y \in Q_i : |(y-x_i) \cdot \nu_\psi(x_i)| \leq \theta r_i\}, \quad L_i := \partial Q_i \cap \partial R_i, \quad (5.13)$$

see Figure 6. By an extension result we can modify  $\psi$  in each square  $Q_i \in \mathcal{B}_{\text{good}}$  and obtain a function  $\phi \in W^{2,\infty}(\Omega)$  with  $\phi = \psi$  on  $\Omega \setminus B_{\text{good}}$  and  $S(\phi) \cap Q_i \subset L_i \cup H_i$  for each  $Q_i \in \mathcal{B}_{\text{good}}$ . As  $\psi \in W^{2,\infty}(\Omega \setminus \overline{S(\psi)})$  and  $\mathcal{L}^2(B_{\text{good}}) \leq C\theta$  by (5.10), this can be done such that the estimate

$$\int_{\Omega} \Phi(\nabla \phi) \, dx \leq \int_{\Omega} \Phi(\nabla \psi) \, dx + C\theta \quad (5.14)$$

holds, for some  $C$  depending on  $\psi$ . Moreover, recalling the definition  $\varphi(\nu) = \frac{2\kappa}{\sqrt{3}} \sum_{\nu \in \mathcal{V}} |\nu \cdot \mathbf{v}|$  for  $\nu \in \mathbb{S}^1$ , using the continuity of  $\varphi$ , and (5.7)–(5.9) we find

$$\int_{S(\phi)} \varphi(\nu_\phi) \, d\mathcal{H}^1 \leq \int_{S(\psi)} \varphi(\nu_\psi) \, d\mathcal{H}^1 + C\theta. \quad (5.15)$$

We now start to construct the sequence of atomistic displacements  $\psi_n : \mathcal{L}_n(\Omega) \rightarrow \mathbb{R}$  separately in the sets  $B_{\text{bad}}$  and  $\Omega_n \setminus B_{\text{bad}}$ . First, for  $x \in \mathcal{L}_n(\Omega) \setminus B_{\text{bad}}$  we will simply set  $\psi_n(x) = \phi(x)$ . On  $B_{\text{bad}}$ , however, the construction is more delicate as, due to the presence of the crack set  $K(t)$ , we cannot simply discretize the function  $\phi$ . Instead, we will employ a *transfer of jump sets* à la FRANCFORT AND LARSEN [31] to transfer the jump set of  $\phi$  that lies in  $K(t)$  onto  $K_n^L(t)$  defined in (4.19). Afterwards, we discretize the resulting displacement.

**Bad squares: jump transfer.** Let us now come to the essential points of the *jump transfer*. For all details, however, we refer to [31]. Due to the construction of the collection  $\mathcal{B}_{\text{bad}}$ , we can repeat the reasoning in [31, Theorem 2.1]. As before, let  $(v_n)_n$  be the sequence with  $S(v_n) \subset K_n^L(t)$  and  $v_n \rightarrow v$  in  $L^1(\Omega)$ ,  $\nabla v_n \rightharpoonup \nabla v$  weakly in  $L^{3/2}(\Omega; \mathbb{R}^2)$ , where  $S(v) \doteq K(t)$ . Then, it is shown that the jump set of  $\phi$




FIGURE 7. Construction of the continuous curve  $\Gamma_i^n \subset R_i^n$  from  $S(v_n)$ . The curve is related to a level set  $v_n$  and cut at the segments  $H_i^{n,\pm}$ . Note that the figure is only a schematic illustration as the jump set  $S(v_n)$  is actually contained in  $K_n^L(t)$ , which consists of straight segments of length  $\frac{\varepsilon_n}{2}$ .

can be transferred on to the ones of  $(v_n)_n$ : This means that there exist functions  $\phi_n \in SBV^2(\Omega)$  with  $\phi_n = \phi$  on  $\Omega \setminus B_{\text{bad}}$  such that

$$\|\phi_n\|_\infty \leq C\|\phi\|_\infty, \quad \|\nabla\phi_n\|_\infty \leq C\|\nabla\phi\|_\infty \quad \text{for all } n \in \mathbb{N} \quad (5.16)$$

for a universal  $C > 0$  and such that there exists an index  $N \in \mathbb{N}$  depending on  $\theta$  with

$$\mathcal{H}^1((S(\phi_n) \setminus S(v_n)) \cap B_{\text{bad}}) \leq C\theta \quad \text{for all } n \geq N. \quad (5.17)$$

The main idea of the proof is to transfer the jump of  $\phi$  in each  $Q_i \in \mathcal{B}_{\text{bad}}$  onto a continuous curve  $\Gamma_i^n \subset Q_i$  related to the boundary of a certain level set for  $v_n$ , see Figure 7 for the construction of  $\Gamma_i^n$ . (The construction provides a function with discontinuities on the boundary of a set of finite perimeter  $P_i^n \subset Q_i$ . Without restriction, up to filling holes and removing components,  $P_i^n$  can be chosen such that both  $P_i^n$  and  $Q_i \setminus P_i^n$  are connected sets (more precisely, indecomposable, see [4, Sections 4,5]). Then, by [4, Theorem 7] the boundary  $\partial^* P_i^n$  can be represented by a Jordan BV curve and, as a consequence,  $\Gamma_i^n := \partial^* P_i^n \cap Q_i$  is a continuous curve.) Here, one uses (5.11) and the  $BV$  coarea formula to show that  $\mathcal{H}^1(\Gamma_i^n \setminus S(v_n)) \leq C\theta r_i$ , see estimate [31, (2.15)].

More precisely, for each  $n \in \mathbb{N}$  and each  $Q_i := Q_{r_i}(x_i) \in \mathcal{B}_{\text{bad}}$  there are two lines  $H_i^{n,+}$  and  $H_i^{n,-}$  with normal  $\nu_\phi(x_i)$  which lie above and below the middle line  $H_i$  containing the point  $x_i$ , also with normal  $\nu_\phi(x_i)$ , such that

$$R_i^n \supset Q_i \cap \{y : |(y - x_i) \cdot \nu_\phi(x_i)| \leq 2\theta r_i\}, \quad (5.18)$$

$$\mathcal{H}^1\left(\bigcup_{\mathcal{B}_{\text{bad}}} L_i^n\right) \leq C\theta \quad (5.19)$$

for a universal constant  $C > 0$ , where  $R_i^n$  denotes the rectangular subset that lies between  $H_i^{n,+}$  and  $H_i^{n,-}$ , and  $L_i^n := \partial R_i^n \setminus (H_i^{n,+} \cup H_i^{n,-})$  denotes its lateral boundaries, cf. [31, (2.10)–(2.11)]. We again refer to Figure 6 for an illustration. The function  $\phi_n$  is then constructed such that  $\phi_n = \phi$  outside of  $\bigcup_{\mathcal{B}_{\text{bad}}} R_i^n$  and inside  $\bigcup_{\mathcal{B}_{\text{bad}}} R_i^n$  it is defined by reflection. More precisely, since  $S(\phi) \cap Q_i \subset R_i^n$  by (5.8) and (5.18), the construction in [31] actually yields

$$S(\phi_n) \cap Q_i \tilde{\subset} \Gamma_i^n \cup L_i^n \quad \text{for all } Q_i \in \mathcal{B}_{\text{bad}}. \quad (5.20)$$

We point out that, as the construction of  $\phi_n$  in [31] relies on a reflection, this leads only to a Lipschitz-regularity of  $\phi_n$  in the two components of  $Q_i \setminus \Gamma_i^n$ , see (5.16). However, for our purposes we will need that

also  $\nabla\phi_n$  is Lipschitz continuous. Therefore, we use a variant of the construction which relies on a cut-off argument and allows us to construct functions  $\phi_n$  such that, for a constant  $C > 0$ , we have

$$\nabla\phi_n \in SBV(\Omega; \mathbb{R}^2) \quad \text{with} \quad \|\nabla(\nabla\phi_n)\|_{L^\infty(\Omega)} \leq C\theta^{-1}(\min_i r_i)^{-1}\|\nabla\phi\|_\infty + C\|\nabla^2\phi\|_\infty. \quad (5.21)$$

(Cut-off constructions for  $SBV$ -functions are by now standard, see e.g. [9, Sections 2.3, 2.4], and have also already been used in related second-order problems, see [30, Lemma 4.4] or [29, Proposition 4.2].) More precisely, supposing  $\nu_\phi(x_i) = e_2$  and  $x_i = 0$  for notational convenience, we let  $T_i^\pm := \{y \in Q_i : \pm y_2 \in (\theta r_i, 2\theta r_i)\}$  and  $b_\pm(y_1) := \int_{\theta r_i}^{2\theta r_i} \phi(y_1, \pm s) ds$  for  $y_1 \in (-r_i, r_i)$ . In the connected component  $P_i^{n,+}$  of  $Q_i \setminus \Gamma_i^n$  containing  $H_i^{n,+}$ , we set  $\phi_n(y_1, y_2) = \phi(y_1, y_2)\varphi_i^+(y_1, y_2) + (1 - \varphi_i^+(y_1, y_2))b_+(y_1)$ , where  $\varphi_i^+ \in C^\infty(Q_i)$  with  $\varphi_i^+ = 1$  on  $\{y_2 > 2\theta r_i\}$  and  $\varphi_i^+ = 0$  on  $\{y_2 < \theta r_i\}$ . In the other connected components of  $Q_i \setminus \Gamma_i^n$  we perform a similar cutoff with  $b_-(y_1)$ . Omitting exact details, we mention that, to see the bounds (5.16) and (5.21), we particularly exploit  $\|\partial_1 b_\pm\|_{L^\infty(-r_i, r_i)} \leq C\|\nabla\phi\|_\infty$  and  $\|\phi(y_1, \pm \cdot) - b_\pm(y_1)\|_{L^\infty(\theta r_i, 2\theta r_i)} \leq C\theta r_i \|\nabla\phi\|_\infty$  for all  $y_1 \in (-r_i, r_i)$ , and  $\|\nabla\varphi_i^\pm\|_\infty \leq C(\theta r_i)^{-1}$ . For later convenience, up to performing a suitable small modification of the functions  $b_\pm$ , by using (5.19) one can also achieve that, for a sufficiently small constant  $c_\theta$  depending on  $\theta$ ,

$$(i) \quad \bigcup_{B_{\text{bad}}} L_i^n \cup \Gamma_i^n \tilde{\subset} S(\phi_n), \quad (ii) \quad \mathcal{H}^1(\{x \in S(\phi_n) \cap B_{\text{bad}} : |[\phi_n](x)| \leq c_\theta\}) \leq C\theta. \quad (5.22)$$

In the following, we suppose that the additional properties (5.20)–(5.22) are satisfied. As by definition of the sequence  $(v_n)_n$  we have  $S(v_n) \tilde{\subset} K_n^L(t)$ , (5.17) yields

$$\mathcal{H}^1((S(\phi_n) \setminus K_n^L(t)) \cap B_{\text{bad}}) \leq C\theta \quad \text{for all } n \geq N. \quad (5.23)$$

Moreover,  $S(\phi_n)$  is essentially closed, i.e.,  $\mathcal{H}^1(\overline{S(\phi_n)} \setminus S(\phi_n)) = 0$ , and by (5.20) and (5.22) we have

$$S(\phi_n) \cong (S(\phi) \setminus \overline{B_{\text{bad}}}) \cup \bigcup_{Q_i \in \mathcal{B}_{\text{bad}}} (L_i^n \cup \Gamma_i^n). \quad (5.24)$$

With these preparations, we can define the sequence of atomistic displacements by

$$\psi_n(x) := \phi_n(x) \quad \text{for all } x \in \mathcal{L}_n(\Omega). \quad (5.25)$$

This means in particular that  $\psi_n(x) = \phi(x)$  for  $x \in \mathcal{L}_n(\Omega) \setminus B_{\text{bad}}$  and  $\psi_n(x) = \psi(x)$  for  $x \in \mathcal{L}_n(\Omega) \setminus (B_{\text{good}} \cup B_{\text{bad}})$  by the definition of  $\phi$  below (5.13). As  $B_{\text{good}} \cup B_{\text{bad}} \subset U$ , this gives  $\psi_n(x) = g(t, x)$  for all  $x \in \mathcal{L}_n(\Omega \setminus U)$ , i.e.,  $\psi_n \in \mathcal{A}_n(g(t))$ .

**Some auxiliary lemmas.** In the following, we choose  $k \in \mathbb{N}$  such that  $t \in [t_n^k, t_n^{k+1})$ . (Clearly,  $k$  depends on  $t$  and  $n$  but we do not include this in the notation for simplicity.) We recall the definition of the energies  $\mathcal{E}_n^{k, \text{ela}}$ ,  $\mathcal{E}_n^{k, \text{cra}}$ ,  $\mathcal{E}_n^{k, \text{rem}}$ , and  $\mathcal{E}_n^{k, \text{bdy}}$  in (3.10), see also (4.18). We also recall the definition of  $M_{n, \Delta}^{k, \mathbf{v}}(v_n)$ ,  $\mathcal{C}_n^k(v_n)$ , and  $\mathcal{C}_n^{k, L}(v_n)$  in (3.2), (3.3), and (3.8). We denote by  $\mathcal{D}_n^{\mathbf{v}}(\phi_n)$  the collection of triangles such that  $\overline{S(\phi_n)}$  intersects the side of the triangle that is oriented in direction  $\mathbf{v} \in \mathcal{V}$ . We also define

$$\mathcal{D}_n(\phi_n) := \bigcup_{j=1,2,3} \mathcal{D}_n^{\mathbf{v}^j}(\phi_n) = \{\Delta : \overline{S(\phi_n)} \cap \partial\Delta \neq \emptyset\}, \quad (5.26)$$

which is the collection of triangles that are intersected by  $\overline{S(\phi_n)}$ . Note that the sets  $\mathcal{D}_n^{\mathbf{v}^j}(\phi_n)$  are obviously not disjoint in general. We now formulate three auxiliary results that will help us to prove Theorem 4.7.

**Lemma 5.1** (Small stretching of springs if there is no jump). *For each  $\mathbf{v} \in \mathcal{V}$  it holds*

$$|(\psi_n)_\Delta \cdot \mathbf{v}| \leq C \quad \text{for all } \Delta \in \mathcal{T}_n \setminus \mathcal{D}_n^{\mathbf{v}}(\phi_n) \quad (5.27)$$

for a constant  $C > 0$  depending on  $\phi$ . Moreover, for  $\varepsilon_n$  small enough, we have

$$M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) \leq M_{n,\Delta}^{k,\mathbf{v}} \vee R \leq M_{n,\Delta}^{k,\mathbf{v}}(u_n^k) \vee R \quad \text{for all } \Delta \in \mathcal{T}_n \setminus \mathcal{D}_n^{\mathbf{v}}(\phi_n). \quad (5.28)$$

Whenever a spring does not intersect the jump set of  $\phi_n$ , the stretching is necessarily controlled, see (5.27), and thus does effectively not increase the memory variable in the sense of (5.28).

**Lemma 5.2** (Small stretching of springs despite of jump). *For  $\varepsilon_n$  small enough, it holds that*

$$\varepsilon_n \sum_{\mathbf{v} \in \mathcal{V}} \# \{ \Delta \in \mathcal{D}_n^{\mathbf{v}}(\phi_n) : M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) \leq R \} \leq C\theta \quad (5.29)$$

and

$$\varepsilon_n \# (\mathcal{D}_n(\phi_n) \setminus \mathcal{C}_n^k(\psi_n)) \leq C\theta. \quad (5.30)$$

In principle, it can happen that a spring intersects the jump set of  $S(\phi_n)$ , but still the stretching is smaller than  $R$ . This is the case if the jump height  $[\phi_n](x)$  for some  $x \in S(\phi_n)$  is small. Due to the fine control on the amount of points where this can happen (use  $\phi_n = \phi$  outside of  $B_{\text{bad}}$  and (5.22)(ii)), the energy contribution of such springs can be controlled in terms of  $\theta$ .

For the third lemma, we define the collection of triangles inside  $B_{\text{bad}}$  by  $\mathcal{T}_n^{\text{bad}} = \{ \Delta \in \mathcal{T}_n : \Delta \subset B_{\text{bad}} \}$ .

**Lemma 5.3** (Bounds in  $B_{\text{bad}}$  and complement). *It holds that*

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{\varepsilon_n}{2} \sum_{(\mathcal{C}_n^k(\psi_n) \cap \mathcal{D}_n(\phi_n)) \setminus \mathcal{T}_n^{\text{bad}}} & \left( \sum_{M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) > R} \Psi(M_{n,\Delta}^{k,\mathbf{v}}(\psi_n)) - \sum_{M_{n,\Delta}^{k,\mathbf{v}}(u_n^k) > R} \Psi(M_{n,\Delta}^{k,\mathbf{v}}(u_n^k)) \right) \\ & \leq \int_{S(\phi) \setminus K(t)} \varphi(\nu_\phi) d\mathcal{H}^1 + C\theta \end{aligned} \quad (5.31)$$

and

$$\varepsilon_n \# \left\{ (\Delta, \mathbf{v}) \in (\mathcal{C}_n^k(\psi_n) \cap \mathcal{T}_n^{\text{bad}} \cap \mathcal{D}_n(\phi_n)) \times \mathcal{V} : M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) > M_{n,\Delta}^{k,\mathbf{v}}(u_n^k) \vee R \text{ and } M_{n,\Delta}^{k,\mathbf{v}}(u_n^k) \leq R_n \right\} \leq C\theta. \quad (5.32)$$

In particular, we have

$$\varepsilon_n \# \mathcal{C}_n^k(\psi_n) + \varepsilon_n \# \mathcal{C}_n^k(u_n^k) \leq C. \quad (5.33)$$

Here, we recall the shorthand notation in (5.31), namely that the first sum runs over triangles  $\Delta$  and the other sums run over directions  $\mathbf{v}$  satisfying the corresponding inequality. The two estimates (5.31) and (5.32) deal with 'broken springs' that are intersected by  $S(\phi_n)$  in  $\Omega \setminus B_{\text{bad}}$  and  $B_{\text{bad}}$ , respectively. The proof of the first estimate (5.31) resembles the construction of a recovery sequence in the atomistic-to-continuum  $\Gamma$ -convergence result [36]. In particular, outside  $B_{\text{bad}}$  no transfer of jump enters the argument. Yet, the latter is at the core of proving (5.32) which indeed fundamentally relies on (5.17). Eventually, estimate (5.33) follows from (5.27)–(5.32).

**Proof of the stability result.** We defer the proofs of the three lemmas to the end of the section and proceed with the proof of Theorem 4.7.

*Proof of Theorem 4.7.* Let  $t \in [0, T]$  be given and, for each  $n \in \mathbb{N}$ , choose  $k$  such that  $t \in [t_n^k, t_n^{k+1})$ . Let  $(\psi_n)_n$  be the sequence defined in (5.25). By construction, we clearly have  $\psi_n(x) = g(t, x)$  for  $x \in \mathcal{L}_n(\Omega \setminus U)$  as  $\phi_n$  coincides with  $\psi$  outside of  $B_{\text{good}} \cup B_{\text{bad}}$  and  $B_{\text{good}} \cup B_{\text{bad}} \subset U$ . Recalling the discussion at the beginning of the section, it suffices to show (5.1)–(5.4) for an arbitrary but fixed  $\theta > 0$ .

*Step 1: Proof of (5.1).* Consider the interpolation  $\tilde{\psi}_n$  related to  $\psi_n$  which is affine on each triangle of  $\mathcal{T}_n$ . In view of (5.25) and the regularity of  $\psi$ , one can check that

$$\chi_{\Omega_n} \tilde{\psi}_n \rightarrow \psi \quad \text{on } L^1(\Omega \setminus (B_{\text{good}} \cup B_{\text{bad}})).$$

By (5.16), (5.25), and the first property in (5.10), we also find

$$\|\chi_{\Omega_n} \tilde{\psi}_n - \psi\|_{L^1(B_{\text{good}} \cup B_{\text{bad}})} \leq C(\|\phi\|_{L^\infty(\Omega)} + \|\psi\|_{L^\infty(\Omega)})\theta^2 \leq C\theta^2.$$

This shows (5.1).

*Step 2: Proof of (5.2).* Recall the definition of  $\mathcal{E}_n^{k,\text{cra}}(\psi_n)$  and  $\mathcal{E}_n^{k,\text{rem}}(\psi_n)$  from (3.11), namely

$$\mathcal{E}_n^{k,\text{cra}}(\psi_n) = \frac{\varepsilon_n}{2} \sum_{\mathcal{C}_n^{k,\text{L}}(\psi_n)} \sum_{M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) > R_n} \Psi(M_{n,\Delta}^{k,\mathbf{v}}(\psi_n))$$

and

$$\begin{aligned} \mathcal{E}_n^{k,\text{rem}}(\psi_n) &= \frac{\varepsilon_n}{2} \sum_{\mathcal{C}_n^k(\psi_n)} \left( \sum_{R < M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) \leq R_n} \Psi(M_{n,\Delta}^{k,\mathbf{v}}(\psi_n)) + \sum_{M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) \leq R} \Psi(\varepsilon_n^{1/2} |(\psi_n)_\Delta \cdot \mathbf{v}|) \right) \\ &\quad + \frac{\varepsilon_n}{2} \sum_{\mathcal{C}_n^k(\psi_n) \setminus \mathcal{C}_n^{k,\text{L}}(\psi_n)} \sum_{M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) > R_n} \Psi(M_{n,\Delta}^{k,\mathbf{v}}(\psi_n)). \end{aligned}$$

The same definition holds for  $u_n^k$  in place of  $\psi_n$ . We then estimate the contributions by

$$\mathcal{E}_n^{k,\text{cra}}(\psi_n) - \mathcal{E}_n^{k,\text{cra}}(u_n^k) + \mathcal{E}_n^{k,\text{rem}}(\psi_n) - \mathcal{E}_n^{k,\text{rem}}(u_n^k) \leq \mathcal{F}_n^{k,1} + \mathcal{F}_n^{k,2} + \mathcal{F}_n^{k,3} + \mathcal{F}_n^{k,4},$$

where

$$\begin{aligned} \mathcal{F}_n^{k,1} &:= \frac{\varepsilon_n}{2} \sum_{(\mathcal{C}_n^k(\psi_n) \cap \mathcal{D}_n(\phi_n)) \setminus \mathcal{T}_n^{\text{bad}}} \left( \sum_{M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) > R} \Psi(M_{n,\Delta}^{k,\mathbf{v}}(\psi_n)) - \sum_{M_{n,\Delta}^{k,\mathbf{v}}(u_n^k) > R} \Psi(M_{n,\Delta}^{k,\mathbf{v}}(u_n^k)) \right), \\ \mathcal{F}_n^{k,2} &:= \frac{\varepsilon_n}{2} \sum_{\mathcal{C}_n^k(\psi_n) \cap \mathcal{D}_n(\phi_n) \cap \mathcal{T}_n^{\text{bad}}} \left( \sum_{M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) > R} \Psi(M_{n,\Delta}^{k,\mathbf{v}}(\psi_n)) - \sum_{M_{n,\Delta}^{k,\mathbf{v}}(u_n^k) > R} \Psi(M_{n,\Delta}^{k,\mathbf{v}}(u_n^k)) \right), \\ \mathcal{F}_n^{k,3} &:= \frac{\varepsilon_n}{2} \sum_{\mathcal{C}_n^k(\psi_n) \setminus \mathcal{D}_n(\phi_n)} \left( \sum_{M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) > R} \Psi(M_{n,\Delta}^{k,\mathbf{v}}(\psi_n)) - \sum_{M_{n,\Delta}^{k,\mathbf{v}}(u_n^k) > R} \Psi(M_{n,\Delta}^{k,\mathbf{v}}(u_n^k)) \right), \\ \mathcal{F}_n^{k,4} &:= \frac{\varepsilon_n}{2} \sum_{\mathcal{C}_n^k(\psi_n)} \sum_{M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) \leq R} \Psi(\varepsilon_n^{1/2} |(\psi_n)_\Delta \cdot \mathbf{v}|). \end{aligned}$$

Here, we summed up all the contributions in 'broken' triangles  $\Delta \in \mathcal{C}_n^k(\psi_n)$ . However, we dropped the contributions of related to  $u_n^k$  in triangles  $\mathcal{C}_n^k(u_n^k) \setminus \mathcal{C}_n^k(\psi_n)$  and the contributions  $\Psi(\varepsilon_n^{1/2} |(\psi_n)_\Delta \cdot \mathbf{v}|)$  for  $(\Delta, \mathbf{v}) \in \mathcal{C}_n^k(u_n^k) \times \mathcal{V}$  with  $M_{n,\Delta}^{k,\mathbf{v}}(u_n^k) \leq R$ , which are all clearly nonnegative.

We now estimate the various terms. First, (5.31) together with (5.15) implies

$$\limsup_{n \rightarrow \infty} \mathcal{F}_n^{k,1} \leq \int_{S(\psi) \setminus K(t)} \varphi(\nu_\psi) d\mathcal{H}^1 + C\theta. \quad (5.34)$$

To estimate  $\mathcal{F}_n^{k,2}$  we use (5.32). More precisely, we denote the set on the left-hand side of (5.32) by  $\Lambda_n$  for shorthand. Then, recalling that  $\Psi$  is increasing and  $\Psi \leq \kappa$  by (ii') and (iv'), we compute

$$\mathcal{F}_n^{k,2} \leq \frac{3\varepsilon_n}{2} \#((\mathcal{C}_n^k(\psi_n) \cap \mathcal{D}_n(\phi_n) \cap \mathcal{T}_n^{\text{bad}}) \setminus \Lambda_n) (\kappa - \Psi(R_n)) + \frac{3\varepsilon_n}{2} \# \Lambda_n \kappa.$$

Then, by  $\lim_{t \rightarrow \infty} \Psi(t) = \kappa$ , (5.32), and the fact that  $\#\mathcal{C}_n^k(\psi_n) \leq C\varepsilon_n^{-1}$  (see (5.33)) we conclude

$$\limsup_{n \rightarrow \infty} \mathcal{F}_n^{k,2} \leq C\theta. \quad (5.35)$$

By (5.28) and the monotonicity of  $\Psi$  we directly obtain  $\mathcal{F}_n^{k,3} \leq 0$ . Eventually, by (5.27) and (5.29) we get

$$\begin{aligned} \limsup_{n \rightarrow \infty} \mathcal{F}_n^{k,4} &\leq \limsup_{n \rightarrow \infty} \frac{\varepsilon_n}{2} \sum_{\mathbf{v} \in \mathcal{V}} \#(\mathcal{C}_n^k(\psi_n) \setminus \mathcal{D}_n^{\mathbf{v}}(\phi_n)) \Psi(\varepsilon_n^{1/2} C) \\ &\quad + \limsup_{n \rightarrow \infty} \frac{\varepsilon_n}{2} \sum_{\mathbf{v} \in \mathcal{V}} \# \{ \Delta \in \mathcal{D}_n^{\mathbf{v}}(\phi_n) : M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) \leq R \} \Psi(R) \leq C\theta, \end{aligned}$$

where we also used  $\#\mathcal{C}_n^k(\psi_n) \leq C\varepsilon_n^{-1}$  by (5.33) and  $\lim_{t \rightarrow 0} \Psi(t) = 0$ . This along with (5.34)–(5.35) concludes the proof of (5.2).

*Step 3: Proof of (5.3).* The essential point is to prove that there exists a sequence of sets  $(\Upsilon_n)_n \subset \Omega$  such that

$$\chi_{\Omega_n} \nabla \hat{\psi}_n^k - \nabla \phi_n \rightarrow 0 \text{ pointwise a.e. in } \Omega, \quad \mathcal{L}^2(\Upsilon_n) \leq C\theta\varepsilon_n, \quad \|\nabla \hat{\psi}_n^k\|_{L^\infty(\Omega_n \setminus \Upsilon_n)} \leq C. \quad (5.36)$$

Assume for the moment that (5.36) is true and recall the definition of  $\Psi_n^{\text{cell}}$  in (3.16) with  $\Psi_n^{\text{cell}} \leq C\varepsilon_n^{-1}$  on  $\mathbb{R}^2$ . Thus, using (3.15) and (5.36) it follows

$$\begin{aligned} \limsup_{n \rightarrow \infty} \mathcal{E}_n^{k,\text{ela}}(\psi_n) &= \limsup_{n \rightarrow \infty} \int_{\Omega_n} \Psi_n^{\text{cell}}(\nabla \hat{\psi}_n^k(x)) \, dx \leq \limsup_{n \rightarrow \infty} \int_{\Omega_n \setminus \Upsilon_n} \Psi_n^{\text{cell}}(\nabla \hat{\psi}_n^k(x)) \, dx + C\mathcal{L}^2(\Upsilon_n)\varepsilon_n^{-1} \\ &\leq \limsup_{n \rightarrow \infty} \int_{\Omega_n \setminus (B_{\text{bad}} \cup \Upsilon_n)} \Psi_n^{\text{cell}}(\nabla \hat{\psi}_n^k(x)) \, dx + \limsup_{n \rightarrow \infty} \int_{B_{\text{bad}} \setminus \Upsilon_n} \Psi_n^{\text{cell}}(\nabla \hat{\psi}_n^k(x)) \, dx + C\theta. \end{aligned} \quad (5.37)$$

Note that the function  $x \mapsto \Psi_n^{\text{cell}}(\nabla \hat{\psi}_n^k(x))$  is uniformly bounded on  $\Omega_n \setminus \Upsilon_n$  because of (5.36) and the fact that  $\Psi_n^{\text{cell}}$  is bounded on  $\{|y| \leq C\}$ , see Lemma 3.4(iii). Hence, by  $\mathcal{L}^2(B_{\text{bad}}) \leq \theta^2$  (see the first item in (5.10)), the second integral in (5.37) can be controlled in terms of  $C\theta$ . For the first integral we argue similarly to the proof of Lemma 3.6. The fact that  $\phi_n = \phi$  in  $\Omega \setminus B_{\text{bad}}$ , the reverse Fatou lemma, and the first item of (5.36) lead to

$$\begin{aligned} \limsup_{n \rightarrow \infty} \int_{\Omega_n \setminus (B_{\text{bad}} \cup \Upsilon_n)} \Psi_n^{\text{cell}}(\nabla \hat{\psi}_n^k(x)) \, dx &\leq \int_{\Omega \setminus B_{\text{bad}}} \Phi(\nabla \phi) \, dx + \limsup_{n \rightarrow \infty} \int_{\Omega_n \setminus (B_{\text{bad}} \cup \Upsilon_n)} |\nabla \hat{\psi}_n^k|^2 \frac{\omega(\varepsilon_n^{1/2} |\nabla \hat{\psi}_n^k|)}{(\varepsilon_n^{1/2} |\nabla \hat{\psi}_n^k|)^2} \, dx \\ &\leq \int_{\Omega \setminus B_{\text{bad}}} \Phi(\nabla \phi) \, dx. \end{aligned}$$

Here we have used Lemma 3.4(i) and the fact  $\|\nabla \hat{\psi}_n^k\|_{L^2(\Omega_n \setminus \Upsilon_n)} \leq C$  as well as  $\varepsilon_n^{1/2} |\nabla \hat{\psi}_n^k|^2 \rightarrow 0$  uniformly, which both follow from the last item in (5.36). In view of (5.14), this concludes the proof of (5.3).

Now, we proceed with the proof of (5.36). First, since  $\nabla \hat{\psi}_n^k$  is zero on each  $\Delta \in \mathcal{C}_n^k(\psi_n)$  by construction, and  $\nabla \phi_n$  is uniformly bounded on  $\Omega$ , see (5.16), we get

$$\sum_{\Delta \in \mathcal{C}_n^k(\psi_n)} \chi_\Delta (\nabla \hat{\psi}_n^k - \nabla \phi_n) \rightarrow 0 \text{ strongly in } L^1(\Omega; \mathbb{R}^2), \quad (5.38)$$

where we used  $\mathcal{L}^2(\Delta) = \frac{\sqrt{3}}{4}\varepsilon_n^2$  and  $\#\mathcal{C}_n^k(\psi_n) \leq C\varepsilon_n^{-1}$  by (5.33). Now, we consider triangles  $\Delta \in \mathcal{D}_n(\phi_n) \setminus \mathcal{C}_n^k(\psi_n)$ . Here, we have no good control on  $\nabla \hat{\psi}_n^k = \nabla \tilde{\psi}_n$ , but we can control the volume of the set in terms of  $\theta\varepsilon_n$ . Letting  $\Upsilon_n = \bigcup_{\Delta \in \mathcal{D}_n(\phi_n) \setminus \mathcal{C}_n^k(\psi_n)} \Delta$ , by (5.30) we compute

$$\mathcal{L}^2(\Upsilon_n) \leq C\varepsilon_n^2 \#(\mathcal{D}_n(\phi_n) \setminus \mathcal{C}_n^k(\psi_n)) \leq C\theta\varepsilon_n. \quad (5.39)$$

Eventually, we consider triangles  $\Delta \in \mathcal{T}_n \setminus (\mathcal{D}_n(\phi_n) \cup \mathcal{C}_n^k(\psi_n))$ . In view of (5.24), this implies that either (a)  $\Delta \not\subset B_{\text{bad}}$  with  $\Delta \cap \bigcup_{\mathcal{B}_{\text{bad}}} L_i^n = \emptyset$  or (b)  $\Delta \subset B_{\text{bad}}$ . First, assume that  $\Delta \not\subset B_{\text{bad}}$  and  $\Delta \cap \bigcup_{\mathcal{B}_{\text{bad}}} L_i^n = \emptyset$ . Then, by construction of  $\phi_n$  below (5.21), we have  $\psi_n(x) = \phi_n(x) = \phi(x)$  for all vertices  $x$  of  $\Delta$ . Since  $\phi \in W^{2,\infty}(\Omega \setminus \overline{S(\phi)})$ ,  $\nabla\phi$  is Lipschitz on  $\Delta$  and thus  $\sup_{x,x' \in \Delta} |\nabla\phi(x) - \nabla\phi(x')| \leq \|\nabla^2\phi\|_{L^\infty(\Delta)}\varepsilon_n$ . By the mean value theorem this yields

$$\|\nabla\tilde{\psi}_n - \nabla\phi\|_{L^\infty(\Delta)} \leq C\varepsilon_n \|\nabla^2\phi\|_{L^\infty(\Omega)}.$$

In a similar fashion, if  $\Delta \subset B_{\text{bad}}$ , we have  $\psi_n(x) = \phi_n(x)$  for the vertices of  $\Delta$ , and as  $\nabla\phi_n$  is Lipschitz on  $\Delta$ , we get

$$\|\nabla\tilde{\psi}_n - \nabla\phi_n\|_{L^\infty(\Delta)} \leq C\varepsilon_n \|\nabla^2\phi_n\|_{L^\infty(\Delta)} \leq C\varepsilon_n (\theta^{-1}(\min_i r_i)^{-1} \|\nabla\phi\|_\infty + \|\nabla^2\phi\|_\infty),$$

where we also used (5.21). As  $\Delta \notin \mathcal{C}_n^k(\psi_n)$ , we have  $\hat{\psi}_n^k = \tilde{\psi}_n$  on such triangles. Therefore, for fixed  $\theta > 0$ , we obtain

$$\sum_{\Delta \in \mathcal{T}_n \setminus (\mathcal{D}_n(\phi_n) \cup \mathcal{C}_n^k(\psi_n))} \chi_\Delta (\nabla\hat{\psi}_n^k - \nabla\phi_n) \rightarrow 0 \quad \text{strongly in } L^1(\Omega; \mathbb{R}^2). \quad (5.40)$$

Combining (5.38)–(5.40) with the fact that  $\|\nabla\phi_n\|_\infty \leq C$  yields (5.36).

*Step 4: Proof of (5.4).* Recall the definition of  $\mathcal{E}_n^{\text{bdy}}$  in (3.6). Let  $\mathcal{T}_n^{\text{bdy}} = \{\Delta \in \mathcal{T}_n : \partial\Delta \cap \partial\Omega_n \neq \emptyset\}$  and  $\Omega_n^{\text{bdy}} = \bigcup_{\Delta \in \mathcal{T}_n^{\text{bdy}}} \Delta$ . First, since  $B_{\text{good}} \cup B_{\text{bad}} \subset \subset \Omega$  by construction, we get  $\Delta \cap (B_{\text{good}} \cup B_{\text{bad}}) = \emptyset$  for all  $\Delta \in \mathcal{T}_n^{\text{bdy}}$ , for  $\varepsilon_n$  small enough. By  $\mathcal{H}^1(S(\psi) \cap \Omega_n^{\text{bdy}}) \rightarrow 0$  as  $n \rightarrow \infty$  and the regularity of  $S(\psi)$ , this shows that

$$\varepsilon_n \#(\mathcal{T}_n^{\text{bdy}} \cap \mathcal{D}_n(\phi_n)) = \varepsilon_n \#(\mathcal{T}_n^{\text{bdy}} \cap \mathcal{D}_n(\psi)) \rightarrow 0, \quad (5.41)$$

where  $\mathcal{D}_n(\psi)$  is defined analogous to (5.26), i.e., as the collection of triangles that are intersected by  $\overline{S(\psi)}$ . Now, the contribution of triangles in  $\mathcal{T}_n^{\text{bdy}} \cap (\mathcal{C}_n^k(\psi_n) \setminus \mathcal{D}_n(\phi_n))$  is nonpositive by (5.28) and repeating the argument for  $\mathcal{F}_n^{k,3}$  above. Furthermore, the contribution of triangles in  $\mathcal{T}_n^{\text{bdy}} \cap \mathcal{D}_n(\phi_n)$  is negligible by (5.41) and the boundedness of  $\Psi$ . Concerning the triangles  $\Delta \notin \mathcal{C}_n^k(\psi_n)$ , we can follow the lines of Step 3, in particular (5.37), where we use that also the integral over  $\Omega_n^{\text{bdy}} \setminus \Upsilon_n$  vanishes since  $\mathcal{L}^2(\Omega_n^{\text{bdy}}) \rightarrow 0$  and we have  $\mathcal{L}^2(\Upsilon_n \cap \Omega_n^{\text{bdy}})\varepsilon_n^{-1} \rightarrow 0$  by (5.41). This shows (5.4).  $\square$

**Proofs of the auxiliary lemmas.** We proceed to prove the three auxiliary results.

*Proof of Lemma 5.1.* Fix  $\mathbf{v} \in \mathcal{V}$  and  $\Delta \in \mathcal{T}_n \setminus \mathcal{D}_n^{\mathbf{v}}(\phi_n)$ . Let  $x, x'$  be the pair of vertices with  $x - x'$  parallel to  $\mathbf{v}$ . Since  $\phi_n \in W^{1,\infty}(\Omega \setminus \overline{S(\phi_n)})$ , more precisely  $L := \sup_n \|\nabla\phi_n\|_\infty < \infty$  by (5.16), and the segment between  $x$  and  $x'$  does not intersect  $\overline{S(\phi_n)}$ , we get

$$|\phi_n(x) - \phi_n(x')| \leq \|\nabla\phi_n\|_\infty \varepsilon_n \leq L\varepsilon_n.$$

As  $\psi_n = \phi_n$  on  $\mathcal{L}_n(\Omega)$ , we get

$$|(\psi_n)_\Delta \cdot \mathbf{v}| \leq L,$$

which shows (5.27). In particular, for  $\varepsilon_n$  small enough, we have  $\varepsilon_n^{1/2}L \leq R$ , and thus by (3.1)–(3.2), we find

$$M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) \leq M_{n,\Delta}^{k,\mathbf{v}} \vee R \leq M_{n,\Delta}^{k,\mathbf{v}}(u_n^k) \vee R.$$

This concludes the proof of (5.28).  $\square$

For convenience we proceed with the proof of Lemma 5.3 and address Lemma 5.2 at the end.

*Proof of Lemma 5.3.* We first prove (5.31) in Steps 1–3. Step 4 addresses (5.32), and finally in Step 5 we show (5.33). As a preparation for the proof of (5.31), we split the energy contribution on the left hand side into three terms, namely

$$\frac{\varepsilon_n}{2} \sum_{(C_n^k(\psi_n) \cap \mathcal{D}_n(\phi_n)) \setminus \mathcal{T}_n^{\text{bad}}} \left( \sum_{M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) > R} \Psi(M_{n,\Delta}^{k,\mathbf{v}}(\psi_n)) - \sum_{M_{n,\Delta}^{k,\mathbf{v}}(u_n^k) > R} \Psi(M_{n,\Delta}^{k,\mathbf{v}}(u_n^k)) \right) = \mathcal{F}_n^{\text{good}} + \mathcal{F}_n^{\text{rest}} + \mathcal{F}_n^{\text{glue}}, \quad (5.42)$$

where in each of the three terms the summation is restricted to corresponding subsets of  $\mathcal{T}_n$ , namely

$$\mathcal{T}_n^{\text{good}} = \{\Delta : \Delta \cap B_{\text{good}} \neq \emptyset\}, \quad \mathcal{T}_n^{\text{rest}} = \{\Delta : \Delta \cap (B_{\text{good}} \cup B_{\text{bad}}) = \emptyset\}, \quad \mathcal{T}_n^{\text{glue}} = \{\Delta : \Delta \cap \partial B_{\text{bad}} \neq \emptyset\}.$$

We now treat the three terms separately.

*Step 1:  $\mathcal{F}_n^{\text{good}}$ .* First, by  $\Psi \leq \kappa$  we get

$$\mathcal{F}_n^{\text{good}} \leq \sum_{\mathbf{v} \in \mathcal{V}} \frac{\kappa \varepsilon_n}{2} \#(\mathcal{D}_n^{\mathbf{v}}(\phi_n) \cap \mathcal{T}_n^{\text{good}}).$$

Recall that  $\psi_n = \phi$  on  $\mathcal{L}_n(\Omega) \setminus B_{\text{bad}}$  and  $\Delta \cap B_{\text{bad}} = \emptyset$  for all  $\Delta \in \mathcal{T}_n^{\text{good}}$  (for  $\varepsilon_n$  small enough). In particular, we get  $\mathcal{D}_n(\phi) \cap \mathcal{T}_n^{\text{good}} = \mathcal{D}_n(\phi_n) \cap \mathcal{T}_n^{\text{good}}$ , where we use the notation introduced in (5.26) also for  $\phi$ . Thus, we have

$$\mathcal{F}_n^{\text{good}} \leq \sum_{j=1}^3 \frac{\kappa \varepsilon_n}{2} \#(\mathcal{D}_n^{\mathbf{v}_j}(\phi) \cap \mathcal{T}_n^{\text{good}}). \quad (5.43)$$

Denote by  $\nu_\phi$  again the measure-theoretic unit normal to  $S(\phi)$ . For a square  $Q_i = Q_{r_i}(x_i) \in \mathcal{B}_{\text{good}}$  we now prove that

$$\#\{\Delta \in \mathcal{D}_n^{\mathbf{v}_j}(\phi) \text{ with } \Delta \cap Q_i \neq \emptyset\} \leq 2r_i \frac{4|\nu_\phi(x_i) \cdot \mathbf{v}_j|}{\sqrt{3}\varepsilon_n} + \frac{Cr_i}{\varepsilon_n} \theta. \quad (5.44)$$

By (5.8) and the construction below (5.13) we observe that, for each  $\Delta$  with  $\Delta \cap Q_i \neq \emptyset$ , we have that  $S(\phi) \cap \Delta$  is contained in the two segments forming  $L_i$  and in the segment  $H_i$ . As  $\mathcal{H}^1(L_i) = 4\theta r_i$ , the number of triangles intersecting  $L_i$  can clearly be controlled by  $C\theta r_i/\varepsilon_n$ . Therefore, it remains to consider triangles intersecting  $H_i$ . Recall that  $H_i$  is a segment with normal vector  $\nu_\phi(x_i)$ . Let  $\mathbf{v}_j^\perp$  be a unit vector orthogonal to  $\mathbf{v}_j$ , and denote by  $\Pi_j(H_i)$  the orthogonal projection of  $H_i$  onto  $\mathbb{R}\mathbf{v}_j^\perp$ . We compute  $\mathcal{H}^1(\Pi_j(H_i)) = \mathcal{H}^1(H_i)|\nu_\phi(x_i) \cdot \mathbf{v}_j| = 2r_i|\nu_\phi(x_i) \cdot \mathbf{v}_j|$ , and then by basic geometric considerations, one can estimate the amount of springs in direction  $\mathbf{v}_j$  that are crossed by  $H_i$  as  $(\frac{\sqrt{3}\varepsilon_n}{2})^{-1}2r_i|\nu_\phi(x_i) \cdot \mathbf{v}_j| + 1$ . As each spring is contained in two triangles, this implies

$$\#\{\Delta \in \mathcal{D}_n^{\mathbf{v}_j}(\phi) \text{ with } \Delta \cap H_i \neq \emptyset\} \leq 2r_i \frac{4|\nu_\phi(x_i) \cdot \mathbf{v}_j|}{\sqrt{3}\varepsilon_n} + 2.$$

For  $\varepsilon_n$  small enough with respect to  $\theta > 0$  this implies (5.44). Then, using  $\mathcal{H}^1(Q_i \cap S(\phi)) \geq 2r_i$ , recalling the definition  $\varphi(\nu) = \frac{2\kappa}{\sqrt{3}} \sum_{\mathbf{v} \in \mathcal{V}} |\nu \cdot \mathbf{v}|$  for  $\nu \in \mathbb{S}^1$ , and putting together (5.43)–(5.44), we obtain

$$\mathcal{F}_n^{\text{good}} \leq \kappa \sum_{Q_i \in \mathcal{B}_{\text{good}}} \sum_{j=1}^3 \mathcal{H}^1(S(\phi) \cap Q_i) \left( \frac{2|\nu_\phi(x_i) \cdot \mathbf{v}_j|}{\sqrt{3}} + C\theta \right) \leq \sum_{Q_i \in \mathcal{B}_{\text{good}}} \mathcal{H}^1(S(\phi) \cap Q_i) \varphi(\nu_\phi(x_i)) + C\theta$$

for a constant  $C > 0$  also depending on  $\mathcal{H}^1(S(\phi))$ . Then, due to (5.12), we also get

$$\mathcal{F}_n^{\text{good}} \leq \sum_{Q_i \in \mathcal{B}_{\text{good}}} \mathcal{H}^1((S(\phi) \setminus K(t)) \cap Q_i) \varphi(\nu_\phi(x_i)) + C\theta.$$

Eventually, using that  $\nu_\phi(x_i)$  is the normal vector to  $H_i$  and  $\mathcal{H}^1(L_i) = 4\theta r_i$  for each  $Q_i \in \mathcal{B}_{\text{good}}$ , we find

$$\mathcal{F}_n^{\text{good}} \leq \int_{(S(\phi) \setminus K(t)) \cap \mathcal{B}_{\text{good}}} \varphi(\nu_\phi) d\mathcal{H}^1 + C\theta. \quad (5.45)$$

*Step 2:  $\mathcal{F}_n^{\text{rest}}$ .* We recall that  $\psi_n = \psi$  on  $\mathcal{L}_n(\Omega) \setminus (B_{\text{good}} \cup B_{\text{bad}})$ . Repeating the argument leading to (5.43) we find

$$\mathcal{F}_n^{\text{rest}} \leq \frac{3\kappa \varepsilon_n}{2} \#(\mathcal{D}_n(\psi) \cap \mathcal{T}_n^{\text{rest}}). \quad (5.46)$$

In view of the second item of (5.11) and the regularity of  $S(\psi)$  we have  $\#(\mathcal{D}_n(\psi) \cap \mathcal{T}_n^{\text{rest}}) \leq C\theta \varepsilon_n^{-1}$ . Therefore, we have

$$\mathcal{F}_n^{\text{rest}} \leq C\theta. \quad (5.47)$$

*Step 3:  $\mathcal{F}_n^{\text{glue}}$ .* Let  $Q_i \in \mathcal{B}_{\text{bad}}$  and consider a triangle  $\Delta \in \mathcal{T}_n^{\text{glue}} \cap \mathcal{D}_n(\phi_n) \cap \mathcal{C}_n^k(\psi_n)$  with  $\Delta \cap \partial Q_i \neq \emptyset$ . Recalling the sets  $L_i^n$  introduced in (5.19), by (5.8) and (5.24) we get  $\Delta \cap S(\overline{\phi_n}) = \emptyset$  whenever  $\Delta \cap L_i^n = \emptyset$ . It is thus left to consider  $\Delta$  with  $\Delta \cap L_i^n \neq \emptyset$ . For this, we use (5.19): Because  $\bigcup_{Q_i \in \mathcal{B}_{\text{bad}}} L_i^n$  consists of straight segments, we obtain

$$\sum_{Q_i \in \mathcal{B}_{\text{bad}}} \#\{\Delta \in \mathcal{T}_n : \Delta \cap L_i^n \neq \emptyset\} \leq \frac{C}{\varepsilon_n} \sum_{Q_i \in \mathcal{B}_{\text{bad}}} \mathcal{H}^1(L_i^n) \leq \frac{C\theta}{\varepsilon_n}.$$

Since an estimate analogous to (5.43) or (5.46) also holds for  $\mathcal{F}_n^{\text{glue}}$ , this finally leads to

$$\mathcal{F}_n^{\text{glue}} \leq C\theta. \quad (5.48)$$

Recalling the splitting (5.42) and combining (5.45), (5.47), and (5.48) we conclude the proof of (5.31).

*Step 4: Proof of (5.32).* We define  $\mathcal{D}_n^* := \{\Delta \in \mathcal{T}_n^{\text{bad}} : (S(\phi_n) \setminus K_n^L(t)) \cap \Delta \neq \emptyset\}$  and first argue that it suffices to show that

$$\#\mathcal{D}_n^* \leq C\theta \varepsilon_n^{-1}. \quad (5.49)$$

To see this, we show that  $\mathcal{T}_n^{\text{bad}} \setminus \mathcal{D}_n^*$  does not intersect with the collection from (5.32). For each  $\Delta \in \mathcal{T}_n^{\text{bad}} \setminus \mathcal{D}_n^*$  and some  $\mathbf{v} \in \mathcal{V}$  we either have  $M_{n,\Delta}^{k,\mathbf{v}}(u_n^k) > R_n$  or  $M_{n,\Delta}^{k,\mathbf{v}}(u_n^k) \leq R_n$ . In the first case, the conditions in (5.32) are obviously violated. In the latter case, the segment of  $\partial\Delta$  in direction  $\mathbf{v}$  does not intersect  $K_n^L(t)$ , cf. (3.12) and (4.19), and therefore, by  $\Delta \cap (S(\phi_n) \setminus K_n^L(t)) = \emptyset$ , we conclude  $\Delta \notin \mathcal{D}_n^*(\phi_n)$ . Then, (5.28) implies  $M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) \leq M_{n,\Delta}^{k,\mathbf{v}}(u_n^k) \vee R$ , and thus the triangle does not lie in the collection given in (5.32). This shows that (5.49) induces the desired estimate.

Let us now show (5.49). To this end, we claim that by the construction of  $\psi_n$  we can find a universal constant  $C > 0$  such that for every  $Q_i \in \mathcal{B}_{\text{bad}}$  we have

$$\#\{\Delta \in \mathcal{D}_n^* : \Delta \subset Q_i\} \leq \frac{C}{\varepsilon_n} \mathcal{H}^1((S(\phi_n) \setminus K_n^L(t)) \cap Q_i). \quad (5.50)$$

Recall the curve  $\Gamma_i^n$  satisfying (5.20). Note also that  $K_n^L(t)$  consists of segments with length  $\frac{\varepsilon_n}{2}$  and each two of such segments either touch or have distance at least  $\frac{\varepsilon_n}{2}$ . Therefore, a connected component  $\Gamma_{i,l}^n$  of  $(\Gamma_i^n \setminus K_n^L(t)) \cap Q_i$  satisfies either  $\mathcal{H}^1(\Gamma_{i,l}^n) \geq \frac{\varepsilon_n}{2}$  or that the start and endpoint of  $\Gamma_{i,l}^n$  intersect the same or two consecutive segments contained in  $K_n^L(t)$ . In the latter case, in the construction of  $\phi_n$ , the part  $\Gamma_{i,l}^n$  of the curve  $\Gamma_i^n$  could be replaced by a piecewise affine curve inside  $K_n^L(t)$  such that the estimate (5.17) still holds. Therefore, in the following, we will suppose without restriction that all connected components  $(\Gamma_{i,l}^n)_l$  of  $(\Gamma_i^n \setminus K_n^L(t)) \cap Q_i$  fulfill  $\mathcal{H}^1(\Gamma_{i,l}^n) \geq \frac{\varepsilon_n}{2}$ . We now aim to prove for each of such components that

$$\#\{\Delta : \Gamma_{i,l}^n \cap \Delta \neq \emptyset\} \leq \frac{C}{\varepsilon_n} \mathcal{H}^1(\Gamma_{i,l}^n). \quad (5.51)$$

From this, we then deduce (5.50) as

$$\#\{\Delta \in \mathcal{D}_n^* : \Delta \subset Q_i\} \leq \sum_l \#\{\Delta : \Gamma_{i,l}^n \cap \Delta \neq \emptyset\} \leq \frac{C}{\varepsilon_n} \sum_l \mathcal{H}^1(\Gamma_{i,l}^n) \leq \frac{C}{\varepsilon_n} \mathcal{H}^1((S(\phi_n) \setminus K_n^L(t)) \cap Q_i).$$

In order to prove (5.51), we look at the  $\varepsilon_n$ -neighborhood of  $\Gamma_{i,l}^n$ , i.e.,

$$N_{\varepsilon_n}(\Gamma_{i,l}^n) := \{x \in \mathbb{R}^2 : \text{dist}(x, \Gamma_{i,l}^n) \leq \varepsilon_n\}. \quad (5.52)$$

As  $\mathcal{H}^1(\Gamma_{i,l}^n) \geq \frac{\varepsilon_n}{2}$ , an elementary geometric argument shows that there exists a universal constant  $C > 0$  such that

$$\mathcal{L}^2(N_{\varepsilon_n}(\Gamma_{i,l}^n)) \leq C\varepsilon_n \mathcal{H}^1(\Gamma_{i,l}^n).$$

Since  $\mathcal{L}^2(\Delta) = \frac{\sqrt{3}}{4}\varepsilon_n^2$ , we hence obtain

$$\#\{\Delta : \Gamma_{i,l}^n \cap \Delta \neq \emptyset\} \leq \#\{\Delta : \Delta \subset N_{\varepsilon_n}(\Gamma_{i,l}^n)\} \leq \frac{\mathcal{L}^2(N_{\varepsilon_n}(\Gamma_{i,l}^n))}{\mathcal{L}^2(\Delta)} \leq \frac{C}{\varepsilon_n} \mathcal{H}^1(\Gamma_{i,l}^n).$$

This yields (5.51) and therefore (5.50). In view of (5.23) and (5.50), we eventually get (5.49) by summing over the finite number of squares in  $\mathcal{B}_{\text{bad}}$ .

*Step 5: Proof of (5.33).* By Lemma 4.1 and Corollary 4.4, together with Proposition 3.5, we obtain  $\varepsilon_n \#\mathcal{C}_n^k(u_n^k) \leq C$ . It remains to estimate  $\#\mathcal{C}_n^k(\psi_n)$ . Because of Lemma 5.1, we know that  $\mathcal{C}_n^k(\psi_n) \setminus \mathcal{D}_n^{\mathbf{v}}(\phi_n) \subset \mathcal{C}_n^k(u_n^k)$  for all  $\mathbf{v} \in \mathcal{V}$  and thus  $\#\mathcal{C}_n^k(\psi_n) \setminus \mathcal{D}_n(\phi_n) \leq \#\mathcal{C}_n^k(u_n^k) \leq C\varepsilon_n^{-1}$ . It hence remains to estimate  $\#\mathcal{C}_n^k(\psi_n) \cap \mathcal{D}_n(\phi_n)$ . Due to (5.31), we obtain

$$\varepsilon_n \#\mathcal{C}_n^k(\psi_n) \cap \mathcal{D}_n(\phi_n) \setminus \mathcal{T}_n^{\text{bad}} \leq C + \kappa\varepsilon_n \#\mathcal{C}_n^k(u_n^k) + C\theta \leq C.$$

Here, we have used that contributions of triangles with  $M_{n,\Delta}^{k,\mathbf{v}}(u_n^k) > R$  for some  $\mathbf{v} \in \mathcal{V}$  can be controlled by  $\kappa\varepsilon_n \#\mathcal{C}_n^k(u_n^k)$  and the fact that  $\int_{S(\phi) \setminus K(t)} \varphi(\nu_\phi) d\mathcal{H}^1 \leq C$ . Finally, if we denote the set on the left-hand side of (5.32) by  $\Lambda_n$ , we obtain  $(\mathcal{C}_n^k(\psi_n) \cap \mathcal{D}_n(\phi_n) \cap \mathcal{T}_n^{\text{bad}}) \setminus \Lambda_n \subset \mathcal{C}_n^k(u_n^k)$ . We hence can deduce from (5.32) that

$$\varepsilon_n \#\mathcal{C}_n^k(\psi_n) \cap \mathcal{D}_n(\phi_n) \cap \mathcal{T}_n^{\text{bad}} \leq \varepsilon_n \#\mathcal{C}_n^k(u_n^k) + C\theta \leq C.$$

Putting all bounds together we conclude the proof of (5.33).  $\square$

*Proof of Lemma 5.2.* We start with (5.29). To this end, recall the construction of  $\phi_n$ , see particularly (5.24), and note that it coincides with  $\phi$  outside of  $B_{\text{bad}}$ . Therefore, for  $c_\theta$  small enough, we find that  $\mathcal{H}^1(\{x \in S(\phi) \setminus B_{\text{bad}} : |[\phi](x)| \leq c_\theta\}) \leq \theta$ . Then, using (5.22)(ii), possibly passing to a smaller  $c_\theta$  depending only on  $\theta$ , we get

$$\mathcal{H}^1(\{x \in S(\phi_n) : |[\phi_n](x)| \leq c_\theta\}) \leq C\theta \quad \text{for all } n \in \mathbb{N}, \quad (5.53)$$

where  $C > 0$  is a universal constant.

As we already found the control  $\#\mathcal{D}_n(\psi) \cap \mathcal{T}_n^{\text{rest}} \leq C\theta\varepsilon_n^{-1}$  preceding (5.47), to show (5.29) it thus suffices to consider triangles intersecting  $B_{\text{good}} \cup B_{\text{bad}}$ . We recall that  $S(\phi_n) \cap (B_{\text{good}} \cup B_{\text{bad}})$  is contained in a finite number of curves whose number is bounded uniformly in  $n$ . Therefore, there is only a bounded number of triangles intersecting more than one of these curves. Consequently, in order to show (5.29) for  $\varepsilon_n$  small enough, it suffices to consider triangles  $\Delta$  such that  $N_{\varepsilon_n}(\Delta)$  intersects exactly one of these curves, where similarly to (5.52) we define

$$N_{\varepsilon_n}(\Delta) := \{x \in \mathbb{R}^2 : \text{dist}(x, \Delta) \leq \varepsilon_n\}.$$

Further, in view of (5.49), the triangles that intersect any component  $(\Gamma_{i,l}^n)$  of  $\Gamma_i^n \setminus K_n^{k,L}$  are negligible. Hence, in order to show (5.29), it suffices to consider triangles  $\Delta$  such that  $N_{\varepsilon_n}(\Delta)$  does not intersect

any component  $(\Gamma_{i,l}^n)_{i,l}$ . Now, except for the parts  $(\Gamma_{i,l}^n)_{i,l}$  considered in (5.51), the curves are either the straight segments  $H_i = S(\phi) \cap Q_i$  and  $L_i$  for squares  $Q_i \in \mathcal{B}_{\text{good}}$ , the segments  $L_i^n$  related to  $Q_i \in \mathcal{B}_{\text{bad}}$ , or a union of intervals contained in  $K_n^{k,L}$ , see (5.24). We denote the collection of triangles that intersect one of these curves by  $\mathcal{G}_n$  and define

$$\mathcal{G}_n^{\mathbf{v}} := \{\Delta \in \mathcal{D}_n^{\mathbf{v}}(\phi_n) \cap \mathcal{G}_n : M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) \leq R\}.$$

In view of this preliminary discussion, it suffices to check  $\varepsilon_n \#\mathcal{G}_n^{\mathbf{v}} \leq C\theta$  for all  $\mathbf{v} \in \mathcal{V}$ . Let us consider a triangle  $\Delta \in \mathcal{G}_n^{\mathbf{v}}$ . We note that  $S(\phi_n) \cap N_{\varepsilon_n}(\Delta)$  is contained in a straight segment or in a union of intervals of length  $\frac{\varepsilon_n}{2}$  oriented parallel to the lattice directions. Let  $x$  and  $x'$  be the vertices of  $\Delta$  such that  $x - x'$  is in direction  $\mathbf{v}$  and let  $x_0$  be the unique point in  $\overline{S(\phi_n)}$  on the segment connecting  $x$  and  $x'$ . By the fact that  $M_{n,\Delta}^{k,\mathbf{v}}(\psi_n) \leq R$ , (5.25), and the fundamental theorem of calculus we find

$$R \geq \frac{\varepsilon_n^{1/2} |\phi_n(x) - \phi_n(x')|}{\varepsilon_n} \geq \varepsilon_n^{-1/2} (|\phi_n(x_0)| - \|\nabla \phi_n\|_{\infty} \varepsilon_n) \geq \varepsilon_n^{-1/2} |\phi_n(x_0)| - L\varepsilon_n^{1/2},$$

where we set  $L := \sup_n \|\nabla \phi_n\|_{\infty} < \infty$ , see (5.16). This shows  $|\phi_n(x_0)| \leq \varepsilon_n^{1/2} R + L\varepsilon_n$ . Since  $S(\phi_n) \cap N_{\varepsilon_n}(\Delta)$  is contained in only one of the curves of  $S(\phi_n)$ , again using  $\|\nabla \phi_n\|_{\infty} \leq L$ , we find

$$|\phi_n(z)| \leq \varepsilon_n^{1/2} R + CL\varepsilon_n \quad \text{for all } z \in S(\phi_n) \cap N_{\varepsilon_n}(\Delta).$$

(In fact,  $x_0$  and each  $z$  can be connected by two different curves on different components of  $N_{\varepsilon_n}(\Delta) \setminus \overline{S(\phi_n)}$  of length  $\sim \varepsilon_n$ .) As  $\mathcal{H}^1(S(\phi_n) \cap N_{\varepsilon_n}(\Delta))$  is at least  $\varepsilon_n/2$ , for  $\varepsilon_n$  small enough depending on  $\theta$  we get that

$$\mathcal{H}^1(\{x \in S(\phi_n) \cap N_{\varepsilon_n}(\Delta) : |\phi_n(x)| \leq c\theta\}) \geq \frac{\varepsilon_n}{2}.$$

Summing over all  $\Delta$  in  $\mathcal{G}_n^{\mathbf{v}}$  and observing that each neighborhood  $N_{\varepsilon_n}(\Delta)$  intersects only a bounded number of neighborhoods  $N_{\varepsilon_n}(\Delta')$ ,  $\Delta' \neq \Delta$ , we get

$$\varepsilon_n \#\mathcal{G}_n^{\mathbf{v}} \leq C\mathcal{H}^1(\{x \in S(\phi_n) : |\phi_n(x)| \leq c\theta\}) \leq C\theta,$$

where in the last step we used (5.53). This concludes the proof of (5.29).

Eventually, (5.30) is a simple consequence of (5.29) and the definition of  $\mathcal{D}_n(\phi_n)$  and  $\mathcal{C}_n^k(\psi_n)$ .  $\square$

We close the section with the proof of Corollary 4.8.

*Proof of Corollary 4.8.* Inspection of the proof of Theorem 4.7 shows that it is irrelevant that  $u_n(t)$  satisfies the boundary values  $g(t_n^{k(t)})$ . In fact, it only matters that  $\psi = g(t)$  on  $\Omega \setminus \overline{U}$ . Therefore, we can consider another sequence of solutions  $u_n^*(t)$  related to boundary values  $g^*(t) := 0$  for all  $t \in [0, T]$ . We obtain  $u_n^*(t) = 0$  for all  $t \in [0, T]$  and the corresponding limiting evolution  $u^*(t) = 0$  and  $K^*(t) = \emptyset$  for all  $t \in [0, T]$ . Applying Theorem 4.7 in this situation for some  $\psi \in SBV^2(\Omega)$  with  $\psi = g(t)$  on  $\Omega \setminus \overline{U}$  yields  $(\psi_n)_n$  with  $\psi_n \in \mathcal{A}_n(g(t_n^{k(t)}))$  such that  $\psi_n$  AC-converges to  $\psi$  and

$$\limsup_{n \rightarrow \infty} (\mathcal{E}_n^{\text{ela}}(\psi_n; t) + \mathcal{E}_n^{\text{cra}}(\psi_n; t) + \mathcal{E}_n^{\text{rem}}(\psi_n; t) + \mathcal{E}_n^{\text{bdy}}(\psi_n; t)) \leq \int_{\Omega} \Phi(\nabla \psi) dx + \int_{S(\psi)} \varphi(\nu_{\psi}) d\mathcal{H}^1,$$

where we used  $\mathcal{E}_n^{\text{cra}}(u_n^*(t); t) + \mathcal{E}_n^{\text{rem}}(u_n^*(t); t) = 0$  and  $K^*(t) = \emptyset$ . Now it suffices to observe that  $\mathcal{E}_n(\psi_n; t) = \mathcal{E}_n^{\text{ela}}(\psi_n; t) + \mathcal{E}_n^{\text{cra}}(\psi_n; t) + \mathcal{E}_n^{\text{rem}}(\psi_n; t) + \mathcal{E}_n^{\text{bdy}}(\psi_n; t)$  by (3.10) and the fact that  $\mathcal{E}_n(\psi_n) = \mathcal{E}_n(\psi_n; t)$  since the memory variable is not active due to  $u_n^*(t) = 0$ .  $\square$

#### ACKNOWLEDGEMENTS

This research was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - 377472739/GRK 2423/2-2023. The authors are very grateful for this support.

## REFERENCES

- [1] N.L. ALLINGER. *Molecular structure: understanding steric and electronic effects from molecular mechanics*. John Wiley & Sons (2010).
- [2] S. ALMI, E. DAVOLI, M. FRIEDRICH. *Non-interpenetration conditions in the passage from nonlinear to linearized Griffith fracture*. J. Math. Pures Appl. **175** (2023), 1–36.
- [3] L. AMBROSIO, A. BRAIDES. *Functionals defined on partitions of sets of finite perimeter, I: Integral representation and  $\Gamma$ -convergence*. J. Math. Pures Appl. **69** (1990), 285–305.
- [4] L. AMBROSIO, V. CASELLES, S. MASNOU, J. MOREL. *Connected components of sets of finite perimeter and applications to image processing*. J. Eur. Math. Soc. (JEMS) **3** (2001), 39–92.
- [5] L. AMBROSIO, N. FUSCO, D. PALLARA. *Functions of bounded variation and free discontinuity problems*. Oxford University Press, Oxford 2000.
- [6] A. BACH, A. BRAIDES, M. CICALESE. *Discrete-to-continuum limits of multi-body systems with bulk and surface long-range interactions*. SIAM J. Math. Anal. **52** (2020), 3600–3665.
- [7] R. BADAL, M. FRIEDRICH, J. SEUTTER. *Existence of quasi-static crack evolution for atomistic systems*. Forces in Mechanics **9** (2022), 100138.
- [8] X. BLANC, C. LE BRIS, P.-L. LIONS. *Atomistic to continuum limits for computational materials science*. M2AN Math. Model. Numer. Anal. **41** (2007), 391–426.
- [9] G. BOUCHITTE, I. FONSECA, G. LEONI, L. MASCARENHAUS. *A global method for relaxation  $W^{1,p}$  and in  $SBV_p$* . Arch. Ration. Mech. Anal. **165** (2002), 187–242.
- [10] B. BOURDIN, G. A. FRANCFORT, J. J. MARIGO. *The variational approach to fracture*. J. Elasticity **91** (2008), 5–148.
- [11] A. BRAIDES. *Discrete-to-continuum variational methods for lattice systems*. Proceedings of the International Congress of Mathematicians – Seoul 2014, Kyung Moon Sa, Seoul. **Vol. IV** (2014), 997–1015.
- [12] A. Braides.  *$\Gamma$ -convergence for beginners*. Oxford Lecture Series in Mathematics and its Applications, 22. Oxford University Press, New York, 2002.
- [13] A. BRAIDES, G. DAL MASO, A. GARRONI. *Variational formulation of softening phenomena in fracture mechanics: the one-dimensional case*. Arch. Ration. Mech. Anal. **146** (1999), 23–58.
- [14] A. BRAIDES, A. DEFRANCESCHI, E. VITALI. *Variational evolution of one-dimensional Lennard-Jones systems*. Netw. Heterog. Media **9** (2014), 217–238.
- [15] A. BRAIDES, M. S. GELLI. *Limits of discrete systems with long-range interactions*. J. Convex Anal. **9** (2002), 363–399.
- [16] A. BRAIDES, A. LEW, M. ORTIZ. *Effective cohesive behavior of layers of interatomic planes*. Arch. Ration. Mech. Anal. **180** (2006), 151–182.
- [17] H. BREZIS. *Functional Analysis, Sobolev Spaces and Partial Differential Equations*. Springer New York, NY.
- [18] A. CHAMBOLLE. *A density result in two-dimensional linearized elasticity, and applications*. Arch. Ration. Mech. Anal. **167** (2003), 211–233.
- [19] V. CRISMALE. *On the approximation of SBD functions and some applications*. SIAM J. Math. Anal. **51** (2019), 5011–5048.
- [20] V. CRISMALE, G. LAZZARONI, G. ORLANDO. *Cohesive fracture with irreversibility: Quasistatic evolution for a model subject to fatigue*. Math. Models Methods Appl. Sci. (M3AS) **28** (2018), 1371–1412.
- [21] G. DAL MASO. *An introduction to  $\Gamma$ -convergence*. Progress in Nonlinear Differential Equations and their Applications, 8. Birkhäuser, Boston, 1993.
- [22] G. DAL MASO, G. FRANCFORT, R. TOADER. *Quasistatic crack growth in nonlinear elasticity*. Arch. Ration. Mech. Anal. **176** (2005), 165–225.
- [23] G. DAL MASO, G. LAZZARONI. *Quasistatic crack growth in finite elasticity with non-interpenetration*. Ann. Inst. H. Poincaré Anal. Non Linéaire **27** (2010), 257–290.
- [24] G. DAL MASO, M. NEGRI, D. PERCIVALE. *Linearized elasticity as  $\Gamma$ -limit of finite elasticity*. Set-valued Anal. **10** (2002), 165–183.
- [25] G. DAL MASO, R. TOADER. *A model for the quasi-static growth of brittle fractures based on local minimization*. Math. Models Methods Appl. Sci. (M3AS) **12** (2002), 1773–1800.
- [26] G. DAL MASO, R. TOADER. *A model for the quasistatic growth of brittle fractures: existence and approximation results*. Arch. Ration. Mech. Anal. **162** (2002), 101–135.
- [27] G. DAL MASO, C. ZANINI. *Quasistatic crack growth for a cohesive zone model with prescribed crack path*. Proc. Roy. Soc. Edin. A **137** (2007), 253–279.
- [28] G. DE PHILIPPIS, N. FUSCO, A. PRATELLI. *On the approximation of SBV functions*. Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl. **28** (2017), 369–413.

- [29] A.F. DONNARUMMA.  $\Gamma$ -convergence and stochastic homogenization of second order singular perturbation models for phase transitions. J. Nonlin. Sci. to appear. arXiv: 2406.14356.
- [30] I. FONSECA, A. HAGERTY, R. PARONI. *Second-Order Structured Deformations in the Space of Functions of Bounded Hessian*. Continuum Mech. Thermodyn. **29** (2019), 2699–2734.
- [31] G. A. FRANCFORT, C. J. LARSEN. *Existence and convergence for quasi-static evolution in brittle fracture*. Comm. Pure Appl. Math. **56** (2003), 1465–1500.
- [32] G. A. FRANCFORT, J.-J. MARIGO. *Revisiting brittle fracture as an energy minimization problem*. J. Mech. Phys. Solids **46** (1998), 1319–1342.
- [33] M. FRIEDRICH, P. STEINKE, K. STINSON. *Linearization of quasistatic fracture evolution in brittle materials*. In preparation.
- [34] M. FRIEDRICH, F. SOLOMBRINO. *Quasistatic crack growth in 2d-linearized elasticity*. Ann. Inst. H. Poincaré Anal. Non Linéaire **35** (2018), 27–64.
- [35] M. FRIEDRICH, B. SCHMIDT. *On a discrete-to-continuum convergence result for a two dimensional brittle material in the small displacement regime*. Netw. Heterog. Media **10** (2015), 321–342.
- [36] M. FRIEDRICH, B. SCHMIDT. *An analysis of crystal cleavage in the passage from atomistic models to continuum theory*. Arch. Ration. Mech. Anal. **217** (2015), 263–308.
- [37] M. FRIEDRICH, B. SCHMIDT. *An atomistic-to-continuum analysis of crystal cleavage in a two-dimensional model problem*. J. Nonlin. Sci. **24** (2014), 145–183.
- [38] A. GIACOMINI. *Ambrosio-Tortorelli approximation of quasi-static evolution of brittle fractures*. Calc. Var. PDE. **22** (2005), 129–172.
- [39] A. GIACOMINI. *Size effects on quasi-static growth of cracks*. Calc. Var. PDE. **36** (2005), 1887–1928.
- [40] A. GIACOMINI, M. PONSIGLIONE. *A  $\Gamma$ -convergence Approach to Stability of Unilateral Minimality Properties in Fracture Mechanics and Applications*. Arch. Ration. Mech. Anal. **180** (2006), 399–447.
- [41] A. GIACOMINI, M. PONSIGLIONE. *A discontinuous finite element approximation of quasistatic growth of brittle fractures*. Numer. Funct. Anal. Optim. **24** (2003), 813–850.
- [42] A. GIACOMINI, M. PONSIGLIONE. *Discontinuous finite element approximation of quasistatic crack growth in finite elasticity*. Math. Models and Methods Appl. Sci. (M3AS) **16** (2006), 77–118.
- [43] A. GRIFFITH. *The phenomena of rupture and flow in solids*. Phil. Trans. Roy. Soc. London **221-A** (1920), 163–198.
- [44] E. G. LEWARS. *Computational Chemistry*. 2nd edition, Springer, 2011.
- [45] M. RUF. *Discrete stochastic approximations of the Mumford-Shah functional*. Ann. Inst. H. Poincaré C Anal. Non Linéaire **36** (2019), 887–937.
- [46] M. SCHÄFFNER, A. SCHLÖMERKEMPER. *On Lennard-Jones systems with finite range interactions and their asymptotic analysis*. Netw. Heterog. Media **13** (2018), 95–118.
- [47] B. SCHMIDT. *On the derivation of linear elasticity from atomistic models*. Netw. Heterog. Media **4** (2009), 789–812.
- [48] B. SCHMIDT. *Linear  $\Gamma$ -limits of multiwell energies in nonlinear elasticity theory*. Continuum Mech. Thermodyn. **20** (2008), 375–396.

(Manuel Friedrich) DEPARTMENT OF MATHEMATICS, FRIEDRICH-ALEXANDER UNIVERSITÄT ERLANGEN-NÜRNBERG, CAUER-STR. 11, D-91058 ERLANGEN, GERMANY  
*Email address:* [manuel.friedrich@fau.de](mailto:manuel.friedrich@fau.de)

(Joscha Seutter) DEPARTMENT OF MATHEMATICS, FRIEDRICH-ALEXANDER UNIVERSITÄT ERLANGEN-NÜRNBERG, CAUER-STR. 11, D-91058 ERLANGEN, GERMANY  
*Email address:* [joscha.seutter@fau.de](mailto:joscha.seutter@fau.de)