

On symmetries and conservation laws of a Gardner equation involving arbitrary functions

R. de la Rosa^a, M.L. Gandarias^b, M.S. Bruzón^c

^a Universidad de Cádiz, Spain (e-mail: rafael.delarosa@uca.es).

^b Universidad de Cádiz, Spain (e-mail: marialuz.gandarias@uca.es).

^c Universidad de Cádiz, Spain (e-mail: m.bruzon@uca.es).

Abstract

In this work we study a generalized variable-coefficient Gardner equation from the point of view of Lie symmetries in partial differential equations. We find conservation laws by using the multipliers method of Anco and Bluman which does not require the use of a variational principle. We also construct conservation laws using Ibragimov theorem which is based on the concept of adjoint equation for nonlinear differential equations.

Keywords: adjoint equation to nonlinear equations; conservation laws; multipliers; partial differential equations; symmetries.

1 Introduction

Nonlinear equations with variable coefficients have become increasingly important in recent years because these describe many nonlinear phenomena more realistically than equations with constant coefficients. The Gardner equation, for instance, is used in different areas of physics, such as fluid dynamics, plasma physics, quantum field theory, and it also describes a variety of wave phenomena in plasma and solid state.

In this paper, we consider the variable-coefficient Gardner equation with nonlinear terms given by

$$u_t + A(t)uu_x + C(t)u^2u_x + B(t)u_{xxx} + Q(t)u = 0, \quad (1)$$

where $A(t) \neq 0$, $B(t) \neq 0$, $C(t) \neq 0$ and $Q(t)$ are arbitrary smooth functions of t .

In [10], for $A(t) = 1$ and $C(t) = 0$, the optimal system of one-dimensional subalgebras was obtained. In [11] for the same equation with some special forms of the functions $B(t)$ and $Q(t)$ some conservation laws were constructed. Lie symmetries of equation (1) when $Q(t) = 0$, were derived in [15]. The classification of Lie symmetries obtained in [15] was enhanced in [19] by using the general extended equivalence group. In [23], adding to equation (1) the term $E(t)u_x$, where $E(t)$ is an arbitrary smooth function of t , the authors found new exact non-travelling solutions, which include soliton solutions, combined soliton solutions, triangular periodic solutions, Jacobi elliptic function solutions and combined Jacobi elliptic function solutions of equation (1). Soliton solutions of equation (1) were obtained in [20] transforming the equation to an homogeneous equation when $Q(t) = 0$ and a forcing term $R(t)$ has been added. Finally, in [9], exact solutions were obtained by using the general mapping deformation method adding a new term $E(t)u_x$ and a forcing term $R(t)$ to equation (1), where $E(t)$ and $R(t)$ are arbitrary smooth functions of t .

Lie symmetries, in general, symmetry groups, have several applications in the context of nonlinear differential equations. It is noteworthy that they are used to obtain exact solutions and conservation laws of partial differential equations [5, 6, 7, 16, 22].

In [3] Anco and Bluman gave a general treatment of a direct conservation law method for partial differential equations expressed in a standard Cauchy-Kovaleskaya form

$$u_t = G(x, u, u_x, u_{xx}, \dots, u_{nx}).$$

Nontrivial conservation laws are characterized by a multiplier λ , which has no dependence on u_t and all derivatives of u_t , satisfying

$$\hat{E}[u](\lambda u_t - \lambda G(x, u, u_x, u_{xx}, \dots, u_{nx})) = 0.$$

Here

$$\hat{E}[u] := \frac{\partial}{\partial u} - D_t \frac{\partial}{\partial u_t} - D_x \frac{\partial}{\partial u_x} + D_x^2 \frac{\partial}{\partial u_{xx}} + \dots$$

The conserved density T^t must satisfy

$$\lambda = \hat{E}[u]T^t,$$

and the flux T^x is given by

$$T^x = -D_x^{-1}(\lambda G) - \frac{\partial T^t}{\partial u_x} G + G D_x \left(\frac{\partial T^t}{\partial u_{xx}} \right) + \dots$$

In [13], Ibragimov introduced a general theorem on conservation laws which does not require existence of a classical Lagrangian and is based on the concept of an adjoint equation for nonlinear equations. In [14], Ibragimov generalized the concept of linear self-adjointness by introducing the concept of nonlinear self-adjointness of differential equations. This concept has

been recently used for constructing conservation laws [7, 18].

The aim of this work is to obtain Lie symmetries of equation (1) and construct conservation laws using both methods, the direct method proposed by Anco and Bluman [2, 3] and Ibragimov theorem [13]. We have studied Lie symmetries of equation (1) for cases $Q \neq 0$ and $Q = 0$. In order to obtain conservation laws using Ibragimov theory we have determined the subclasses of equation (1) which are nonlinearly self-adjoint.

2 Classical Symmetries

To apply the classical method to equation (1) we consider the one-parameter Lie group of infinitesimal transformations in (x, t, u) given by

$$\begin{aligned} x^* &= x + \varepsilon \xi(x, t, u) + O(\varepsilon^2), \\ t^* &= t + \varepsilon \tau(x, t, u) + O(\varepsilon^2), \\ u^* &= u + \varepsilon \eta(x, t, u) + O(\varepsilon^2), \end{aligned}$$

where ε is the group parameter. We require that this transformation leaves invariant the set of solutions of equation (1). This yields to an overdetermined, linear system of differential equations for the infinitesimals $\xi(x, t, u)$, $\tau(x, t, u)$ and $\eta(x, t, u)$. The associated Lie algebra of infinitesimal symmetries is formed by the set of vector fields of the form

$$\mathbf{v} = \xi(x, t, u)\partial_x + \tau(x, t, u)\partial_t + \eta(x, t, u)\partial_u. \quad (2)$$

Invariance of equation (1) under a Lie group of point transformations with infinitesimal generator (2) leads to a set of 18 determining equations. By simplifying this system we obtain that $\xi = \xi(x, t)$, $\tau = \tau(t)$, and $\eta = \eta(x, t, u)$ are related by the following conditions:

$$\begin{aligned} \eta_{uu} &= 0, \\ \eta_{ux} - \xi_{xx} &= 0, \\ \eta_{uuu} &= 0, \\ \eta_{uux} &= 0, \\ -\tau B_t - \tau_t B + 3\xi_x B &= 0, \\ \tau u B Q_t - \tau u B_t Q - \eta_u u B Q + 3\xi_x u B Q + \eta B Q + \eta_x u^2 B C \\ &\quad + \eta_{xxx} B^2 + \eta_x u A B + \eta_t B = 0, \\ \tau u^3 B C_t - \tau u^3 B_t C + \xi_x u^3 B C + 2\eta u^2 B C - \tau u^2 A B_t + 3\eta_{uux} u B^2 \\ &\quad - \xi_{xxx} u B^2 + \tau u^2 A_t B + 2\xi_x u^2 A B + \eta u A B - \xi_t u B = 0. \end{aligned} \quad (3)$$

In order to find Lie symmetries of the equation, we distinguish two cases: $Q \neq 0$ and $Q = 0$.

Case 1. $Q \neq 0$.

For the sake of simplicity, in Case 1 we shall consider $C(t) = 1$, obtaining the following symmetries

$$\xi = k_1 x + \beta, \quad \tau = \tau(t), \quad \eta = \frac{\beta_t}{A} + (k_1 + \alpha) u$$

where $A = A(t)$, $B = B(t)$, $Q = Q(t)$, $\alpha = \alpha(t)$, $\beta = \beta(t)$ and $\tau = \tau(t)$ must satisfy the following conditions:

$$(3k_1 - \tau_t) B - \tau B_t = 0, \quad (4)$$

$$-\tau B_t + (4k_1 + 2\alpha) B = 0, \quad (5)$$

$$2\beta_t B + \tau A A_t B - \tau A^2 B_t + (3k_1 + \alpha) A^2 B = 0, \quad (6)$$

$$\tau B Q_t - \tau B_t Q + 3k_1 B Q + \alpha_t B = 0, \quad (7)$$

$$\beta_t (A Q - A_t) + \beta_{tt} A = 0. \quad (8)$$

Equation (4) can be written as:

$$\frac{B_t}{B} = \frac{3k_1 - \tau_t}{\tau} = k, \quad (9)$$

where k is a constant. From (8) we get:

$$Q = \frac{A_t}{A} - \frac{\beta_{tt}}{\beta_t}. \quad (10)$$

Subcase 1.1. Setting $k \neq 0$, solving (9) we obtain

$$B = b_0 e^{kt}, \quad \tau(t) = k_4 e^{-kt} + \frac{3k_1}{k}. \quad (11)$$

From (5) we obtain

$$\alpha(t) = \frac{kk_4 e^{-kt} - k_1}{2}. \quad (12)$$

Substituting (10), (11) and (12) into equations (6) and (7), we get the following system

$$-2AA_t f_1 + \left(kk_1 e^{kt} + k^2 k_4 \right) A^2 - 4k e^{kt} \beta_t = 0, \quad (13)$$

$$2 \left(\frac{A_t}{A} - \frac{\beta_{tt}}{\beta_t} \right)_t f_1 - 2k^2 k_4 \left(\frac{A_t}{A} - \frac{\beta_{tt}}{\beta_t} \right) - k^3 k_4 = 0. \quad (14)$$

where $f_1(t) = 3k_1 e^{kt} + kk_4$. This system admits two solutions

$$A(t) = \frac{\sqrt{e^{kt}}}{2d_0 + kk_1} f_1^{-\frac{d_0}{3kk_1} - \frac{1}{2}} \left(2a_1 k f_1^{\frac{d_0}{3kk_1} + \frac{1}{6}} + a_0 (2d_0 + kk_1) \right), \quad (15)$$

$$\beta(t) = \frac{a_0}{8d_0 k (kk_1 - 2d_0)} f_1^{-\frac{2d_0}{3kk_1}} \left(a_0 (4d_0^2 - k^2 k_1^2) + 8a_1 d_0 k f_1^{\frac{d_0}{3kk_1} + \frac{1}{6}} \right), \quad (16)$$

and

$$A(t) = \frac{\sqrt{e^{kt}}}{2d_0 + kk_1} f_1^{-\frac{d_0}{3kk_1} - \frac{1}{2}} \left(a_0 (2d_0 + kk_1) f_1^{\frac{d_0}{3kk_1} + \frac{1}{6}} - 2a_1 k \right), \quad (17)$$

$$\beta(t) = \frac{a_1}{2(4d_0^2 - k^2 k_1^2)} f_1^{-\frac{2d_0}{3kk_1}} \left(2a_0 (2d_0 + kk_1) f_1^{\frac{d_0}{3kk_1} + \frac{1}{6}} + \frac{a_1 k}{d_0} (kk_1 - 2d_0) \right). \quad (18)$$

Lastly, substituting solution (15)-(16) or solution (17)-(18) into (10), we obtain

$$Q(t) = \frac{2d_0 e^{kt} - k^2 k_4}{2f_1}. \quad (19)$$

In the above equations, the following appointments are introduced: $a_0 \neq 0$, $a_1 \neq 0$, $b_0 \neq 0$, $d_0 \neq 0$, $k_1 \neq 0$, k_4 are arbitrary constants, $2d_0 \pm kk_1 \neq 0$.

Subcase 1.2. Setting $k = 0$, solving (9) we obtain

$$B = b_0, \quad \tau(t) = 3k_1 t + k_3. \quad (20)$$

From (5) we obtain

$$\alpha(t) = -2k_1. \quad (21)$$

Substituting (20), (21) and (10) into equations (6) and (7), we get the following system

$$-2AA_t \tau - 2k_1 A^2 - 4\beta_t = 0, \quad (22)$$

$$2\tau \left(\frac{A_t}{A} - \frac{\beta_{tt}}{\beta_t} \right)_t + 6k_1 \left(\frac{A_t}{A} - \frac{\beta_{tt}}{\beta_t} \right) = 0. \quad (23)$$

In this case, we have

$$A(t) = a_0 \tau^{-\frac{d_0}{3k_1}} + a_1 \tau^{-\frac{1}{3}}, \quad (24)$$

$$\beta(t) = \frac{1}{2} a_0 (d_0 - k_1) \tau^{-\frac{2d_0}{3k_1} - \frac{1}{3}} \left(\frac{a_0 \tau^{\frac{4}{3}}}{3k_1 - 2d_0} - \frac{a_1 \tau^{\frac{d_0}{3k_1} + 1}}{d_0 - 2k_1} \right). \quad (25)$$

Substituting solution (24)-(25) into (10), we get that

$$Q(t) = \frac{d_0}{\tau}. \quad (26)$$

In the above equations, the following appointments are introduced: $a_0 \neq 0$, $a_1, b_0 \neq 0$, $d_0 \neq 0$, $k_1 \neq 0$, k_3 are arbitrary constants, $d_0 \neq \frac{3}{2}k_1$ and $d_0 \neq 2k_1$.

Case 2. $Q = 0$.

Now, the generators are given by:

$$\xi = k_1 x + \beta, \quad \tau = \tau(t), \quad \eta = (k_1 + k_3)u + k_2,$$

where $A = A(t)$, $B = B(t)$, $C = C(t)$, $\beta = \beta(t)$ and $\tau = \tau(t)$ must satisfy the following conditions:

$$(3k_1 - \tau_t)B - \tau B_t = 0, \quad (27)$$

$$\tau B C_t - \tau B_t C + (4k_1 + 2k_3)BC = 0, \quad (28)$$

$$\tau A_t B - \tau A B_t + 2k_2 BC + (3k_1 + k_3)AB = 0, \quad (29)$$

$$k_2 A - \beta_t = 0. \quad (30)$$

Subcase 2.1. We consider $k \neq 0$. In this case we have that $B(t)$ and $\tau(t)$ are given by (11). The remaining functions are defined as follows

$$C(t) = c_0 e^{kt} f_1^{-\frac{2k_3}{3k_1} - \frac{4}{3}}, \quad (31)$$

$$A(t) = \frac{1}{k_1 + k_3} e^{kt} f_1^{-\frac{2k_3}{3k_1} - \frac{4}{3}} \left(2c_0 k_2 + a_0 (k_1 + k_3) f_1^{\frac{k_3}{3k_1} + \frac{1}{3}} \right), \quad (32)$$

$$\beta(t) = \beta_0 - \frac{a_0 k_2 f_1^{-\frac{k_3}{3k_1}}}{k k_3} - \frac{2c_0 k_2^2 f_1^{-\frac{2k_3}{3k_1} - \frac{1}{3}}}{k (k_1 + k_3) (k_1 + 2k_3)}. \quad (33)$$

where f_1 has already been previously defined as $f_1(t) = 3k_1 e^{kt} + kk_4$.

Subcase 2.2. We consider $k = 0$. Now, $B(t)$ and $\tau(t)$ are given by (20). In this case, the remaining functions are given by

$$C(t) = c_0 \tau^{-\frac{2k_3}{3k_1} - \frac{4}{3}}, \quad (34)$$

$$A(t) = \frac{1}{k_1 + k_3} \tau^{-\frac{2k_3}{3k_1} - \frac{4}{3}} \left(2c_0 k_2 + a_0 (k_1 + k_3) \tau^{\frac{k_3}{3k_1} + \frac{1}{3}} \right), \quad (35)$$

$$\beta(t) = \beta_0 - \frac{a_0 k_2 \tau^{-\frac{k_3}{3k_1}}}{k_3} - \frac{2c_0 k_2^2 \tau^{-\frac{2k_3}{3k_1} - \frac{1}{3}}}{(k_1 + k_3)(k_1 + 2k_3)}. \quad (36)$$

In Case 2, the following designations are introduced: $a_0, b_0 \neq 0, c_0 \neq 0, \beta_0, k_1 \neq 0, k_2, k_3 \neq 0, k_4$ are arbitrary constants, $k_1 \neq -k_3$ and $k_1 \neq -2k_3$. For this case the symmetries were obtained in [15].

3 Formal Lagrangian and adjoint equation

In [13] Ibragimov introduced a new theorem on conservation laws. This theorem is valid for any system of differential equations wherein the number of equations is equal to the number of dependent variables. The new theorem does not require existence of a classical Lagrangian and it is based on the concept of adjoint equation for nonlinear equations. In order to obtain the adjoint equation we use the following definition:

Definition 1. Consider an q th-order partial differential equation

$$F(x, u, u_{(1)}, \dots, u_{(s)}) = 0 \quad (37)$$

with independent variables $x = (x^1, \dots, x^n)$ and a dependent variable u , where $u_{(1)} = \{u_i\}$, $u_{(2)} = \{u_{ij}\}, \dots$ denote the sets of partial derivatives of first, second, etc. order, $u_i = \partial u / \partial x^i$, $u_{ij} = \partial^2 u / \partial x^i \partial x^j$. The formal Lagrangian is defined as

$$\mathcal{L} = v F(x, u, u_{(1)}, \dots, u_{(s)}), \quad (38)$$

where $v = v(x)$ is a new dependent variable. The adjoint equation to (37) is

$$F^*(x, u, v, u_{(1)}, v_{(1)}, \dots, u_{(s)}, v_{(s)}) = 0, \quad (39)$$

with

$$F^*(x, u, v, u_{(1)}, v_{(1)}, \dots, u_{(s)}, v_{(s)}) = \frac{\delta(vF)}{\delta u},$$

where

$$\frac{\delta}{\delta u} = \frac{\partial}{\partial u} + \sum_{s=1}^{\infty} (-1)^s D_{i_1} \cdots D_{i_s} \frac{\partial}{\partial u_{i_1 \dots i_s}},$$

denotes the variational derivatives (the Euler-Lagrange operator). Here

$$D_i = \frac{\partial}{\partial x^i} + u_i \frac{\partial}{\partial u} + u_{ij} \frac{\partial}{\partial u_j} + \cdots$$

represents the total differentiation.

Theorem 1. *The adjoint equation to equation (1) is*

$$F^* \equiv vQ - u^2 v_x C - v_{xxx} B - u v_x A - v_t. \quad (40)$$

4 Nonlinearly self-adjoint equations

In this section we use the following definition given in [14].

Definition 2. Equation (37) is said to be **nonlinearly self-adjoint** if the equation obtained from the adjoint equation (39) by the substitution

$$v = \varphi(x, u), \quad (41)$$

such that $\varphi(x, u) \neq 0$, is identical with the original equation (37), i.e.

$$F^*|_{v=\varphi} = \lambda(x, u, \dots)F. \quad (42)$$

for some differential function $\lambda = \lambda(x, u, \dots)$. If $\varphi = u$ or $\varphi = \varphi(u)$ and $\varphi'(u) \neq 0$, equation (37) is said **self-adjoint** or **quasi self-adjoint**, respectively. If $\varphi_{x^i}(x, u) \neq 0$ or $\varphi_u(x, u) \neq 0$ equation (37) is said **weak self-adjoint** [8].

Taking into account expression (40) and using (41) and its derivatives, equation (42) can be written as

$$\begin{aligned} & u_x (\lambda (-u^2 C - uA) - \varphi_u u^2 C - 3 \varphi_{uu} u_{xx} B - 3 \varphi_{u_{xx}} B - \varphi_u uA) \\ & + u_{xxx} (-\lambda B - \varphi_u B) - \lambda uQ + u_t (-\lambda - \varphi_u) + \varphi Q - \varphi_x u^2 C - \varphi_t \\ & - 3 \varphi_{ux} u_{xx} B - \varphi_{uuu} (u_x)^3 B - 3 \varphi_{uux} (u_x)^2 B - \varphi_{xxx} B - \varphi_x uA = 0. \end{aligned} \quad (43)$$

Equation (43) should be satisfied identically in all variables u_t, u_x, u_{xx}, \dots . Equating to zero the coefficients of the derivatives of u we obtain:

Theorem 2. Equation (1) with $A(t) \neq 0, B(t) \neq 0, C(t) \neq 0$ and $Q(t)$ arbitrary functions, is nonlinearly self-adjoint and

$$\varphi = c_1 e^{\int 2Q(t)dt} u + c_2 e^{\int Q(t)dt}, \quad (44)$$

with c_1 and c_2 arbitrary constants.

5 Conservation Laws

Conservation laws appear in many of physical, chemical and mechanical processes, such laws enable us to solve problems in which certain physical properties do not change over time within an isolated physical system.

A conservation law of equation (1) is a space-time divergence such that

$$D_t T^t(x, t, u, u_x, u_t, \dots) + D_x T^x(x, t, u, u_x, u_t, \dots) = 0, \quad (45)$$

on all solutions $u(x, t)$ of the equation (1). Here, T^t represents the conserved density and T^x the associated flux [4], and D_x, D_t denote the total derivative operators with respect to x and t respectively.

In this section we construct conservation laws of each case by using both methods.

5.1 Conservation laws by using the direct method of the multipliers of Anco and Bluman

We suppose, that T^t and T^x have no dependence on u_t and all derivatives of u_t .

Each conservation law (45) has an equivalent characteristic form in which has been eliminated u_t and its differential consequences from T^t and T^x by using equation (1)

$$\widehat{T}^t = T^t|_{u_t=\Delta} = T^t - \Phi,$$

$$\widehat{T}^x = T^x|_{u_t=\Delta} = T^x - \Psi,$$

where $\Delta = -Auu_x - Cu^2u_x - Bu_{xxx} - Qu$, so that

$$\left(D_t \widehat{T}^t(x, t, u, u_x, u_{xx}, \dots) + D_x \widehat{T}^x(x, t, u, u_x, u_{xx}, \dots) \right)|_{u_t=\Delta} = 0,$$

is verified on all solutions of equation (1), and where

$$D_t|_{u_t=\Delta} = \partial_t + \Delta \partial_u + D_x(\Delta) \partial_{u_x} + \dots$$

$$D_x|_{u_t=\Delta} = \partial_x + u_x \partial_u + u_{xx} \partial_{u_x} + \dots = D_x.$$

In particular, moving off of solutions, we have the identity

$$D_t = D_t|_{u_t=\Delta} + (u_t + Au u_x + Cu^2 u_x + Bu_{xxx} + Qu) \partial_u \\ + D_x(u_t + Au u_x + Cu^2 u_x + Bu_{xxx} + Qu) \partial_{u_x} + \dots$$

These expressions yield us to the characteristic form of conservation law (45)

$$D_t \widehat{T}^t(x, t, u, u_x, u_{xx}, \dots) + D_x \left(\widehat{T}^x(x, t, u, u_x, u_{xx}, \dots) + \widehat{\Psi}(x, t, u, u_x, u_t, \dots) \right) \\ = (u_t + Au u_x + Cu^2 u_x + Bu_{xxx} + Qu) \Lambda(x, t, u, u_x, u_{xx}, \dots), \quad (46)$$

where

$$\widehat{\Psi}(x, t, u, u_x, u_t, \dots) = E_{u_x}(\widehat{T}^t) (u_t + Au u_x + Cu^2 u_x + Bu_{xxx} + Qu) \\ + E_{u_{xx}}(\widehat{T}^t) D_x (u_t + Au u_x + Cu^2 u_x + Bu_{xxx} + Qu) + \dots$$

is a trivial flux [4], and the function

$$\Lambda = E_u(\widehat{T}^t),$$

is a multiplier, where $E_u = \partial_u - D_x \partial_{u_x} + D_x^2 \partial_{u_{xx}} - \dots$, denotes the (spatial) Euler operator with respect to u .

A function $\Lambda(x, t, u, u_x, u_{xx}, \dots)$ is called multiplier if it verifies that $(u_t + Au u_x + Cu^2 u_x + Bu_{xxx} + Qu) \Lambda$ is a divergence expression for all functions $u(x, t)$, not only solutions of equation (1).

In order to obtain conservation laws, we use (46) from which is deduced that all nontrivial conserved densities in the form (45) arise from multipliers Λ of equation (1), where Λ depends only on x, t, u and x derivatives of u . Divergence condition can be characterized as follows

$$\frac{\delta}{\delta u} ((u_t + Au u_x + Cu^2 u_x + Bu_{xxx} + Qu) \Lambda) = 0, \quad (47)$$

where $\frac{\delta}{\delta u} = \partial_u - D_x \partial_{u_x} - D_t \partial_{u_t} + D_x D_t \partial_{u_{xt}} + D_x^2 \partial_{u_{xx}} + \dots$, denotes the variational derivative.

Equation (47) is linear in $u_t, u_{tx}, u_{txx}, \dots$ taking the coefficient of u_t and x derivatives of u_t we obtain a system of determining equations for Λ , which yields the following equivalent equations [1, 2, 3]

$$-D_t \Lambda - D_x ((Au + Cu^2) \Lambda) - D_x^3 (B \Lambda) + (Au_x + 2Cu u_x + Q) \Lambda = 0,$$

and

$$\Lambda_u = E_u(\Lambda), \quad \Lambda_{u_x} = -E_u^{(1)}(\Lambda), \quad \Lambda_{u_{xx}} = -E_u^{(2)}(\Lambda), \dots$$

which are verified for all solutions $u(x, t)$ of equation (1).

Given a multiplier Λ , we can obtain the conserved density using a standard method [21]

$$T^t = \int_0^1 d\lambda \, u \Lambda(x, t, \lambda u, \lambda u_x, \lambda u_{xx}, \dots).$$

We have considered multipliers up to second order, i.e., $\Lambda(x, t, u, u_x, u_{xx})$. In this section we proceed to obtain conservation laws for the values obtained for $A(t)$, $B(t)$, $C(t)$ and $Q(t)$ in Case 1, Section 2.

Subcase 1.1. In this case, equation (1) is given by

$$u_t + A(t)uu_x + u^2u_x + b_0e^{kt}u_{xxx} + \frac{2d_0e^{kt} - k^2k_4}{2f_1}u = 0, \quad (48)$$

where $A(t)$ is given by (15) or (17). For equation (48) multiplier is given by

$$\Lambda = e^{-\frac{kt}{2}} f_1^{\frac{2d_0}{3kk_1}+1} \left(c_1 + c_2 e^{-\frac{kt}{2}} u \right),$$

where $f_1(t) = 3k_1e^{kt} + kk_4$. The conserved density obtained from this multiplier is

$$T^t = e^{-\frac{kt}{2}} f_1^{\frac{2d_0}{3kk_1}+1} u \left(c_1 + \frac{c_2}{2} e^{-\frac{kt}{2}} u \right).$$

The flux obtained from this multiplier depends on function $A(t)$. If $A(t)$ is given by (15), we obtain

$$\begin{aligned} T^x = & \frac{e^{-\frac{kt}{2}}}{12} f_1^{\frac{d_0}{3kk_1}} \left(3c_2 f_1^{\frac{d_0}{3kk_1}+1} \left(u^4 e^{-\frac{kt}{2}} + 2b_0 e^{\frac{kt}{2}} (2uu_{xx} - u_x^2) \right) \right. \\ & + 4(a_0c_2 + c_1)u^3 f_1 + 6a_0c_1u^2 e^{\frac{kt}{2}} f_1^{-\frac{d_0}{3kk_1}} + 12b_0c_1u_{xx}e^{kt} f_1^{\frac{1}{2}} \\ & \left. + \frac{4a_1k}{2d_0 + kk_1} u^2 \left(2c_2 u f_1^{\frac{d_0}{3kk_1}+\frac{2}{3}} + 3c_1 e^{\frac{kt}{2}} f_1^{\frac{1}{6}} \right) \right). \end{aligned}$$

If $A(t)$ is given by (17), we get

$$\begin{aligned}
T^x = & \frac{e^{-\frac{kt}{2}}}{12} f_1^{\frac{d_0}{3kk_1}} \left(3c_2 f_1^{\frac{d_0}{3kk_1}+1} \left(u^4 e^{-\frac{kt}{2}} + 2b_0 e^{\frac{kt}{2}} (2uu_{xx} - u_x^2) \right) \right. \\
& + 4c_1 f_1^{\frac{1}{2}} (u^3 + 3b_0 u_{xx} e^{kt}) + 6a_0 c_1 u^2 e^{\frac{kt}{2}} f_1^{\frac{1}{6}} + 4a_0 c_2 u^3 f_1^{\frac{d_0}{3kk_1}+\frac{2}{3}} \\
& \left. - \frac{4a_1 k}{2d_0 + kk_1} u^2 \left(2c_2 u f_1^{\frac{1}{2}} + 3c_1 e^{\frac{kt}{2}} \right) \right).
\end{aligned}$$

Subcase 1.2. In this case, equation (1) is given by

$$u_t + \left(a_0 \tau^{-\frac{d_0}{3k_1}} + a_1 \tau^{-\frac{1}{3}} \right) uu_x + u^2 u_x + b_0 u_{xxx} + \frac{d_0}{\tau} u = 0, \quad (49)$$

where τ is given by (20). For equation (49) has been obtained the following multiplier

$$\Lambda = \tau^{\frac{d_0}{3k_1}} \left(c_1 + c_2 \tau^{\frac{d_0}{3k_1}} \right).$$

The conserved density and the flux obtained from this multiplier are:

$$\begin{aligned}
T^t = & \tau^{\frac{d_0}{3k_1}} u \left(c_1 + \frac{c_2}{2} \tau^{\frac{d_0}{3k_1}} u \right), \\
T^x = & \frac{1}{12} \tau^{\frac{d_0}{3k_1}} \left(\tau^{\frac{d_0}{3k_1}} (6b_0 c_2 (2uu_{xx} - u_x^2) + 3c_2 u^4) + 6a_0 c_1 u^2 \tau^{-\frac{d_0}{3k_1}} \right. \\
& \left. + 6a_1 c_1 u^2 \tau^{-\frac{1}{3}} + 4a_1 c_2 u^3 \tau^{\frac{d_0}{3k_1}-\frac{1}{3}} + 4c_1 (u^3 + 3b_0 u_{xx}) + 4a_0 c_2 u^3 \right).
\end{aligned}$$

5.2 Conservation laws by using a general theorem on conservation laws proved by Ibragimov

In this section we construct conservation laws for the values obtained for the functions $A(t)$, $B(t)$, $C(t)$ and $Q(t)$ in Case 2, Section 2 using the following theorem on conservation laws proved in [13].

Theorem 3. Any Lie point, Lie-Bäcklund or non-local symmetry

$$\mathbf{v} = \xi^i(x, u, u_{(1)}, \dots) \frac{\partial}{\partial x^i} + \eta(x, u, u_{(1)}, \dots) \frac{\partial}{\partial u}, \quad (50)$$

of equation (37) provides a conservation law $D_i(T^i) = 0$ for the simultaneous system (37), (39). The conserved vector is given by

$$\begin{aligned} T^i &= \xi^i \mathcal{L} + W \left[\frac{\partial \mathcal{L}}{\partial u_i} - D_j \left(\frac{\partial \mathcal{L}}{\partial u_{ij}} \right) + D_j D_k \left(\frac{\partial \mathcal{L}}{\partial u_{ijk}} \right) - \dots \right] \\ &+ D_j(W) \left[\frac{\partial \mathcal{L}}{\partial u_{ij}} - D_k \left(\frac{\partial \mathcal{L}}{\partial u_{ijk}} \right) + \dots \right] + D_j D_k(W) \left[\frac{\partial \mathcal{L}}{\partial u_{ijk}} - \dots \right] + \dots, \end{aligned} \quad (51)$$

where W and \mathcal{L} are defined as follows:

$$W = \eta - \xi^j u_j, \quad \mathcal{L} = v F(x, u, u_{(1)}, \dots, u_{(s)}). \quad (52)$$

In order to apply Theorem 3 to our equation we perform the following change of notation:

$$(x^1, x^2) = (t, x), \quad (\xi^1, \xi^2) = (\tau, \xi), \quad (T^1, T^2) = (T^t, T^x).$$

Subcase 2.1. In this case, we consider equation (1) with $Q(t) = 0$, $A(t)$, $B(t)$ and $C(t)$ given by (32), (11) and (31) respectively. Consequently, the equation admits the following generator

$$v = (k_1 x + \beta(t)) \partial_x + \left(k_4 e^{-kt} + \frac{3k_1}{k} \right) \partial_t + ((k_1 + k_3)u + k_2) \partial_u, \quad (53)$$

where $\beta(t)$ is given by (33). From Theorem 2, we have that φ is given by

$$\varphi = c_1 u + c_2. \quad (54)$$

Thus, we obtain conservation law (45) with the conserved vector

$$T^t = (k_3 u + k_2)(c_1 u + c_2) + k_1 u \left(\frac{3}{2} c_1 u + 2 c_2 \right), \quad (55)$$

$$\begin{aligned}
T^x = & \frac{e^{kt}}{12(k_3 + k_1)} (6b_0 c_1 (2u u_{xx} - u_x^2) (2k_3^2 + 5k_1 k_3 + 3k_1^2) \\
& + 12b_0 u_{xx} (c_2 (k_3^2 + 3k_1 k_3 + 2k_1^2) + c_1 (k_3 + k_1)) \\
& + f_1^{-\frac{k_3}{3k_1} - 1} (4a_0 c_1 u^3 (2k_3^2 + 5k_1 k_3 + 3k_1^2) \\
& + 6a_0 c_2 u^2 (k_3^2 + 3k_1 k_3 + 2k_1^2) + 6a_0 c_1 k_2 u^2 (k_3 + k_1)) \\
& + f_1^{-\frac{2k_3}{3k_1} - \frac{4}{3}} (3c_0 c_1 u^4 (2k_3^2 + 5k_1 k_3 + 3k_1^2) + 4c_0 c_1 k_1 u^3 (5k_3 + 7k_2) \\
& + 4c_0 c_2 u^3 (k_3^2 + 3k_1 k_3 + 2k_1^2) + 12c_0 k_2 u^2 (c_2 k_3 + c_1 k_2 + 2c_2 k_1))) .
\end{aligned}$$

Subcase 2.2. Now, we consider equation (1) with $Q(t) = 0$, $A(t)$, $B(t)$ and $C(t)$ given by (35), (20) and (34) respectively. Consequently, the equation admits the following generator

$$v = (k_1 x + \beta(t)) \partial_x + \tau \partial_t + ((k_1 + k_3)u + k_2) \partial_u, \quad (56)$$

where $\tau(t)$ and $\beta(t)$ are given by (20) and (36) respectively. As in the previous case, ϕ is given by (54). Thus, we obtain conservation law (45) with the conserved vector

$$T^t = (k_3 u + k_2) (c_1 u + c_2) + k_1 u \left(\frac{3}{2} c_1 u + 2c_2 \right),$$

$$\begin{aligned}
T^x = & \frac{1}{12(k_3 + k_1)} (6b_0 c_1 (2u u_{xx} - u_x^2) (2k_3^2 + 5k_1 k_3 + 3k_1^2) \\
& + 12b_0 u_{xx} (c_2 (k_3^2 + 3k_1 k_3 + 2k_1^2) + c_1 (k_3 + k_1)) \\
& + \tau^{-\frac{k_3}{3k_1} - 1} (4a_0 c_1 u^3 (2k_3^2 + 5k_1 k_3 + 3k_1^2) \\
& + 6a_0 c_2 u^2 (k_3^2 + 3k_1 k_3 + 2k_1^2) + 6a_0 c_1 k_2 u^2 (k_3 + k_1)) \\
& + \tau^{-\frac{2k_3}{3k_1} - \frac{4}{3}} (3c_0 c_1 u^4 (2k_3^2 + 5k_1 k_3 + 3k_1^2) + 4c_0 c_1 k_1 u^3 (5k_3 + 7k_2) \\
& + 4c_0 c_2 u^3 (k_3^2 + 3k_1 k_3 + 2k_1^2) + 12c_0 k_2 u^2 (c_2 k_3 + c_1 k_2 + 2c_2 k_1))) .
\end{aligned}$$

We remark that some of these conservation laws yield conserved integrals with physical meaning. Setting in (55) $k_1 = 1$, $c_2 = \frac{1}{2}$ and $k_2 = k_3 = c_1 = 0$ we have

$$C_1 = \int_{-\infty}^{\infty} u dx,$$

which is the conserved mass for equation (1). The conserved integral arising from (55) setting $k_1 = 1$, $c_1 = \frac{2}{3}$ and $k_2 = k_3 = c_2 = 0$ gives the energy

$$C_2 = \int_{-\infty}^{\infty} u^2 dx,$$

for equation (1).

Another powerful application of conservation laws taking into account the relationship between Lie symmetries and conservation laws is the so called double reduction method given by Sjöberg [17]. This method allow us to reduce the Gardner equation to a second order ordinary differential equation. Sjöberg introduced this method in order to get solutions of a q th partial differential equation (37) from the solutions of an ordinary differential equation of order $q-1$. This method can be applied when a symmetry \mathbf{v} is associated to a conserved vector T . In accordance with the definition given by Kara and Mahomed [12] we will establish that a symmetry \mathbf{v} is associated to T if the following equation holds

$$\mathbf{v}(T^i) + T^i D_k (\xi^k) - T^k D_k (\xi^i) = 0. \quad (57)$$

In the terms of the canonical variables r , s and w , symmetry (2) becomes a translation on s , $\mathbf{v} = \frac{\partial}{\partial s}$. Thus, the conservation law can be rewritten as

$$D_s T^s + D_r T^r = 0, \quad (58)$$

with

$$T^s = \frac{T^t D_t(s) + T^x D_x(s)}{D_t(r) D_x(s) - D_x(r) D_t(s)}, \quad (59)$$

and

$$T^r = \frac{T^t D_t(r) + T^x D_x(r)}{D_t(r) D_x(s) - D_x(r) D_t(s)}. \quad (60)$$

Due to solutions of the equation (37) written in canonical variables must be invariant with respect to \mathbf{v} and T is associated with \mathbf{v} , equation (58) becomes

$$D_r T^r = 0,$$

so that

$$T^r(r, w, w_r, w_{rr}, \dots, w_{r^{q-1}}) = k, \quad k = \text{const.} \quad (61)$$

We stress that equation (61) is an ordinary differential equation of order $q-1$, whose solutions are solutions of equation (37), by writing this solution in terms of x , t and u .

In order to show the above explained procedure, let us consider the equation

$$F \equiv u_t + uu_x + u^2 u_x + u_{xxx} = 0, \quad (62)$$

It can be easily chequed that equation (62) admits the following generator

$$\mathbf{v} = \left(k_1 x - \frac{k_1}{2} t \right) \partial_x + (3k_1 t + k_2) \partial_t + \left(-k_1 u - \frac{k_1}{2} \right) \partial_u, \quad (63)$$

where k_1 and k_2 are arbitrary constants. From Theorem (50), one can get the following conserved vector for generator (63)

$$\begin{aligned} T^t &= -\frac{c_1 k_1}{2} u^2 - \frac{c_1 k_1}{2} u - \frac{c_2 k_1}{2}, \\ T^x &= -c_1 k_1 u u_{xx} - \frac{c_1 k_1}{2} u_{xx} + \frac{c_1 k_1}{2} u_x^2 \\ &\quad - \frac{c_1 k_1}{4} u^4 - \frac{c_1 k_1}{2} u^3 - \frac{c_1 k_1}{4} u^2, \end{aligned} \quad (64)$$

where c_1 and c_2 are arbitrary constants. The conserved vector (64) is not associated to symmetry (63). It can be easily seen that generator

$$\mathbf{v} = c \partial_x + \partial_t.$$

is associated to (64). The canonical coordinates are

$$r = x - ct, \quad s = t, \quad w = u.$$

We suppose without loss of generality that $k_1 = c_1 = 1$ and $c_2 = 0$. From (60) we get

$$T^r = \frac{1}{4} ((4w+2) w_{rr} - 2w_r^2 + w^4 + 2w^3 + (1-2c) w^2 - 2cw). \quad (65)$$

Setting $T^r = \frac{k}{4}$, $k = \text{const.}$, we obtain

$$(4w+2) w_{rr} - 2w_r^2 - 2cw + (1-2c) w^2 + 2w^3 + w^4 = k. \quad (66)$$

Due to the fact that (66) is an autonomous equation the substitution $w_r = p(w)$ yields the following first order ordinary differential equation

$$p' = \frac{k + 2p^2 + 2cw + (2c - 1)w^2 - 2w^3 - w^4}{4w + 2},$$

whose solutions are solutions of (62) when written in terms of x , t and u .

6 Conclusions

In this paper we have considered a generalized variable-coefficient Gardner equation. Classical symmetries of equation (1) have been obtained involving different arbitrary functions which can be used to determine similarity and exact solutions. Symmetries obtained in this work generalise those already obtained by other authors in equations which belong to the family of equations (1), such as KdV equation and other Gardner equations with time-dependent coefficients. In particular, we have considered two different cases, which in turn would lead to different subcases.

We have determined the subclasses of equation (1) which are nonlinearly self-adjoint, as well as the multipliers, of Anco and Bluman method. We have derived conservation laws by using both methods. We have shown that some of these conservation laws yields conserved integrals with physical meaning, such as mass and energy. Finally, as an example of another application of the conserved vectors, we have applied the double reduction method to get exact solutions of the Gardner equation from solutions of a second order reduced ordinary differential equation.

Acknowledgements

The authors acknowledge the financial support from Junta de Andalucía group FQM-201. The second and third author also acknowledge the support of DGICYT project MTM2009-11875 and FEDER funds.

References

- [1] S. Anco, G. Bluman, Direct construction of conservation laws from field equations, *Phys. Rev. Lett.* **78** (1997) 2869-2873.
- [2] S. Anco, G. Bluman, Direct construction method for conservation laws of partial differential equations Part I: Examples of conservation law classifications, *Eur. J. Appl. Math.* **13** (2002) 545-566.
- [3] S. Anco, G. Bluman, Direct construction method for conservation laws of partial differential equations Part II: General treatment, *Eur. J. Appl. Math.* **13** (2002) 567-585.

- [4] S. Anco et al. Symmetries and conservation laws of the generalized Krichever-Novikov equation, Preprint (2014).
- [5] M.S. Bruzón, M.L. Gandarias, Conservation laws for a family of Benjamin-Bona-Mahony-Burgers equations, NSC 2012 4th IEEE International Conference on Nonlinear Science and Complexity, Budapest, Hungary, (2012) 155–160.
- [6] M.S. Bruzón, Weak Self-Adjointness and Conservation Laws for a Family of Benjamin-Bona-Mahony-Burgers Equations, in: Discontinuity and Complexity in Nonlinear Physical Systems, Springer, 2014, pp. 23-34.
- [7] M.S. Bruzón, M.L. Gandarias, R. de la Rosa, Conservation laws of a family Reaction-Diffusion-Convection Equations, in: Localized Excitations in Nonlinear Complex Systems, Nonlinear Systems and Complexity **7**, Springer, 2014, pp. 403-417.
- [8] M. L. Gandarias, Weak self-adjoint differential equations, *J. Phys. A: Math. Theor.* **44** (2011) 262001.
- [9] B. Hong, D. Lu, New exact solutions for the generalized variable-coefficient Gardner equation with forcing term, *Appl. Math. Comput* **219** (2012) 2732–2738.
- [10] A.G. Johnpillai, C.M. Khalique, Group analysis of KdV equation with time dependent coefficients, *Appl. Math. Comput* **216** (2010) 3761–3771.
- [11] A.G. Johnpillai, C.M. Khalique, Conservation laws of KdV equation with time dependent coefficients, *Commun Nonlinear Sci Numer Simulat* **16** (2011) 3081–3089.
- [12] A.H. Kara, F.M. Mahomed, Relationship between symmetries and conservation laws, *Int. J. Theor. Phys.* **39** (2000) 23–40.
- [13] N.H. Ibragimov, A new conservation theorem, *J. Math. Anal. Appl.* **333** (2007) 311–328.
- [14] N.H. Ibragimov, Nonlinear self-adjointness and conservation laws, *J. Phys. A: Math. Theor.* **44** (2011) 432002–432010.
- [15] M. Molati, M.P. Ramollo, Symmetry classification of the Gardner equation with time-dependent coefficients arising in stratified fluids, *Commun Nonlinear Sci Numer Simulat* **17** (2012) 1542–1548.
- [16] A.G. Rasin, J. Schiff, The Gardner method for symmetries, *J. Phys. A: Math. Theor.* **46** (2013) 155202 15pp.
- [17] A. Sjöberg, Double reduction of PDEs from the association of symmetries with conservation laws with applications, *Commun Nonlinear Sci Numer Simulat* **19** (2014) 3036–3043.
- [18] R. Traciná, M.S. Bruzón, M.L. Gandarias, M. Torrisi, Nonlinear self-adjointness, conservation laws, exact solutions of a system of dispersive evolution equations, *Appl. Math. Comput.* **184** (2007) 608–616.

- [19] O. Vaneeva, O. Kuriksha, C. Sophocleous, Enhanced group classification of Gardner equations with time-dependent coefficients, *Commun Nonlinear Sci Numer Simulat* **22** (2015) 1243–1251.
- [20] A.M. Wazwaz, A study on KdV and Gardner equations with time-dependent coefficients and forcing terms, *Appl. Math. Comput.* **217** (2010) 2277–2281.
- [21] T. Wolf, An efficiency improved program LIEPDE for determining Lie-symmetries of PDEs, *Proceedings of Modern Group Analysis: advances analytical and computational methods in mathematical physics* (1993) 377–385.
- [22] B. Xu, X.Q. Liu, Classification, reduction, group invariant solutions and conservation laws of the Gardner-KP equation, *Appl. Math. Comput.* **215** (2009) 1244–1250.
- [23] L.H. Zhang, L.H. Dong, L.M. Yan, Construction of non-travelling wave solutions for the generalized variable-coefficient Gardner equation, *Appl. Math. Comput* **203** (2008) 784–791.