ON NONNEGATIVE INVARIANT QUARTICS IN TYPE A

SEBASTIAN DEBUS, CHARU GOEL, SALMA KUHLMANN, AND CORDIAN RIENER

ABSTRACT. The equivariant nonnegativity versus sums of squares question has been solved for any infinite series of essential reflection groups but type A. As a first step to a classification, we analyse A_n -invariant quartics. We prove that the cones of invariant sums of squares and nonnegative forms are equal if and only if the number of variables is at most 3 or odd.

1. INTRODUCTION

The study of nonnegative real polynomials, i.e. polynomials whose evaluation on any point is nonnegative, is a topic of interest from many perspectives, e.g. verificiation of polynomial inequalities and polynomial optimization. From complexity theoretical view the verification is NP-hard [5]. If one can write a real polynomial as a sum of squares of real polynomials, then the polynomial is clearly nonnegative. It was shown by Hilbert [13] in his celebrated theorem from 1888 that there are basically three cases where any nonnegative polynomial is a sum of squares. We formulate Hilbert's theorem in terms of *forms*, i.e. homogeneous polynomials, since any polynomial is nonnegative if and only if its homogenization is nonnegative and a sum of squares if and only if its homogenization is a sum of squares [16]. Hilbert showed that the cones of nonnegative forms and that of sums of squares of degree 2d in *n* variables are equal if and only if $(n, 2d) \in \{(2, 2d'), (n', 2), (3, 4) \mid n', d' \in \mathbb{N}\}$. Hilbert's proof was unconstructive and it took almost 80 years until the first example of a nonnegative polynomial which is not a sum of squares was given (this is the Motzkin polynomial [18]). It was then asked by Hilbert whether any nonnegative polynomial is a sum of squares of rational functions. This is known as Hilbert's 17th problem. E. Artin proved that this is true, thereby lying the cornerstone of the field of *real algebraic geometry*.

Motivated by Hilbert's 1888 theorem, several authors investigated the equivariant setting. For a group G acting on the real polynomial ring one restricts to *invariant* forms, i.e. forms which are fixed under the action of G. Choi, Lam and Reznick investigated the question for the symmetric group S_n which was completed by Goel, Kuhlmann and Reznick [11]. The signed symmetric group B_n acting on the polynomial ring via permutation of variables and switching of signs was also considered [12]. Recently, Debus and Riener considered D_n -invariant forms where D_n is the subgroup of B_n of even number of sign changes. All these groups have in common that they are *reflection groups*.

A finite group *G* is a *real reflection group* if $G \subset GL_n(\mathbb{R}^n)$ is such that the matrix group is generated by *reflections*, i.e. isometries $\mathbb{R}^n \to \mathbb{R}^n$ with a hyperplane as the set of fixed points. We usually just say that an abstract group *G* is a real reflection group and the representation of *G* is implicitly known. A real reflection group is called *essential* if no non-trivial subspace of \mathbb{R}^n is point wise fixed. It is a classical result that any real reflection group can be decomposed into a direct product of essential reflection groups. The essential real reflection groups were fully classified by Coxeter [7, 8]. There are four infinite families A_n, B_n, D_n and $I_2(m)$ and six exceptional real reflection groups H_3, H_4, F_4, E_6, E_7 , and E_8 .

For B_n , D_n and trivially $I_2(m)$ the equivariant classification of nonnegativity versus sums of squares was completed in [9]. It is a natural question to consider the remaining infinite series of essential reflection groups A_n and to study the equivariant nonnegativity versus sums of squares question. In this paper we initiate a study of A_n -invariant quartics. Although the vector space dimension of A_n -invariant quartics is only 2, we will see that the understanding is challenging. A reason for the complexity involved here is that we do not consider nonnegativity of a polynomial globally. We consider nonnegativity on a hyperplane and do consider sums of squares modulo an ideal which is in general a very difficult problem.

The paper is structured as follows. Section 2 explains the action of the group A_n on an *n*-dimensional vector space and the induced action on the polynomial ring. Following this, we examine the sets of nonnegative and sums of squares A_n invariant quartics in Section 3. We begin in Subsection 3.1 to elaborate on the difference between global nonnegativity of quartics and nonnegativity of A_n -invariant quartics. In Subsection 3.2 we provide the extremal elements of the cone of A_n -invariant nonnegative quartics before we analyse the A_n -invariant sums of squares quartics in Subsection 3.3. Finally, we present a proof of our main theorem, Theorem 3.2 in Subsection 3.4.

2. The reflection groups of type A and A_n -invariant polynomials

The real reflection group A_n is, as a group, isomorphic to the symmetric group S_{n+1} . Recall that the reflection group S_{n+1} is acting on \mathbb{R}^{n+1} via permutation of coordinates in all possible ways. We call this action the *permutation action* of the symmetric group. There is a non-trivial fixed subspace which is spanned by the vector (1, ..., 1) under the permutation action and thus the permutation action does not define an *essential* real reflection group. The action of S_{n+1} on the invariant subspace $U_n := \{a \in \mathbb{R}^{n+1} : \sum_{i=1}^n a_i = 0\}$ via permutation of coordinates defines an essential real reflection group called A_n . We also say that it is the reflection group of *type A*.

Recall that any group G acting on \mathbb{R}^n induces an action of G on the polynomial ring $\mathbb{R}[\mathbf{x}]$ in *n* variables. The action is as follows:

$$\boldsymbol{\sigma} \cdot f(\mathbf{x}) := f(\boldsymbol{\sigma}^{-1} \cdot \mathbf{x})$$

where $\mathbf{x} = (\mathbf{x}_1, \dots, \mathbf{x}_n)$ is a basis of the dual vector space of \mathbb{R}^n and $\sigma \in G$. We refer to ([4], Section 4) for details.

It is a classical result by Chevalley, Sheppard and Todd that the *invariant ring* of real polynomials under the action of a finite matrix group in $GL_n(\mathbb{R})$ is isomorphic to a polynomial ring if and only if the group is a real reflection group [6, 21].

In order to study A_n invariant forms we consider the restriction of the permutation action of the symmetric group S_{n+1} to the *n* dimensional real vector space

$$U_n = \left\{ a \in \mathbb{R}^{n+1} : \sum_i a_i = 0 \right\}.$$

Let $e_i \in \mathbb{R}^n$ denote the unit vector with 1 at the *i*-th coordinate. A linear basis of U_n is

$$u_1 = e_1 - e_2, \ldots, u_n = e_1 - e_{n+1}.$$

The group A_n acts on U_n via permutation of the e_i 's in all possible ways. We obtain an induced action on an *n*-variate polynomial ring $\mathbb{R}[\mathbf{y}]$, where \mathbf{y} is a basis of the dual vector space of U_n and on the quotient of an (n + 1)-variate polynomial ring $\mathbb{R}[\mathbf{x}]$ modulo the ideal generated by the linear polynomial $\mathbf{x}_1 + \ldots + \mathbf{x}_{n+1}$. While A_n does act on the (n + 1)-variate quotient ring $\mathbb{R}[\mathbf{x}]/(\mathbf{x}_1 + \ldots + \mathbf{x}_{n+1})$ via permutation of the \mathbf{x}_i 's, the reflection group does not permute the \mathbf{y}_i 's. We recall that those rings are isomorphic and two equivalent representations of A_n .

For two real representations *V*, *W* of a group *G* we say a linear map $\phi : V \to W$ is *G*-equivariant if $\sigma \cdot \phi(v) = \phi(\sigma \cdot w)$ for any $v \in V, w \in W, \sigma \in G$.

Proposition 2.1. The ring homomorphism $\mathbb{R}[\mathbf{y}] \to \mathbb{R}[\mathbf{x}]/(\mathbf{x}_1 + \ldots + \mathbf{x}_{n+1})$ defined by $\mathbf{y}_i \mapsto \mathbf{x}_1 - \mathbf{x}_{i+1}$, for all $1 \le i \le n$, is a A_n -equivariant isomorphism.

Proof. Recall that A_n fixes the subspace defined by $\mathbf{x}_1 + \ldots + \mathbf{x}_{n+1} = 0$. A basis of this subspace is $\mathbf{x}_1 - \mathbf{x}_{i+1}$ for $1 \le i \le n$. The basis elements and $\mathbf{x}_1 + \ldots + \mathbf{x}_{n+1}$ form a basis of the degree 1 part of $\mathbb{R}[\mathbf{x}]$. We have

$$\mathbb{R}[\mathbf{x}] \cong \mathbb{R}[\mathbf{x}_1 - \mathbf{x}_2, \dots, \mathbf{x}_1 - \mathbf{x}_{n+1}][\mathbf{x}_1 + \dots + \mathbf{x}_{n+1}]$$

and

$$\mathbb{R}[\mathbf{x}]/(\mathbf{x}_1 + \ldots + \mathbf{x}_{n+1}) \cong \mathbb{R}[\mathbf{y}]$$

With the discussion above the induced linear isomorphism is A_n -equivariant.

Since we have a ring isomorphism we have that being a sum of squares is equivalent for the image and preimage. Moreover, nonnegativity of the preimage is equivalent to nonnegativity of the image on the subspace U_n of \mathbb{R}^{n+1} .

We denote by p_k the *power sum* polynomial of degree k in the (n + 1)-variables **x**, i.e. $p_k = \sum_{i=1}^{n+1} \mathbf{x}_i^k$. It is classically known that the power sum polynomials p_2, \ldots, p_{n+1} generate the A_n -invariant ring as \mathbb{R} -algebra modulo the ideal (p_1) .

Theorem 2.2. The invariant ring of A_n is isomorphic to a polynomial ring. The invariant ring of A_n acting via permutation of the variables **x** on the (n + 1)-variate quotient ring $\mathbb{R}[\mathbf{x}]/(p_1)$ is $\mathbb{R}[\mathbf{x}]^{A_n} \simeq \mathbb{R}[p_2, \dots, p_{n+1}]$.

3. SOS VERSUS PSD FOR A_n -INVARIANT QUARTICS

In this Section we prove our main result Theorem 3.2. We mainly restrict our notation and definitions to quartics. Since the invariant ring is generated by the power sums p_2, \ldots, p_{n+1} the vector space of A_n -invariant quartics is 2 dimensional and is spanned by the quotient classes of p_2^2 and p_4 .

Definition 3.1. We call a A_n -invariant quartic in $\mathbb{R}[\mathbf{x}]/(p_1)$ nonnegative or psd if and only if any element in its quotient class in $\mathbb{R}[\mathbf{x}]$ is nonnegative on U_n . We denote the set of psd A_n -invariant quartics by P^{A_n} . We call a A_n -invariant quartic in $\mathbb{R}[\mathbf{x}]/(p_1)$ a sum of squares or sos if and only if an element in its quotient class in $\mathbb{R}[\mathbf{x}]$ is of the form $g_1^2 + \ldots + g_m^2 + p_1 \cdot g$ for some $g_1, \ldots, g_m, g \in \mathbb{R}[\mathbf{x}]$. We denote the set of all A_n -invariant sos quartics by Σ^{A_n} .

Suppose $f_1 = ap_2^2 + bp_4 + p_1 \cdot g_1$ and $f_2 = ap_2^2 + bp_4 + p_1 \cdot g_2$ are two equivalent A_n -invariant quartics. Then nonnegativity of the quotient class $f_1 \mod (p_1)$ is well defined since $p_1 = 0$ on U_n .

The sets P^{A_n} , Σ^{A_n} are pointed closed convex cones in the vector space $\mathbb{R}[\mathbf{x}]/(p_1)$.

The main result is the following.

Theorem 3.2. For $n \ge 3$ we have $P^{A_n} = \Sigma^{A_n}$ if and only if *n* is odd.

Note, we have $P^{A_n} = \Sigma^{A_n}$ by Hilbert's classification for all $n \leq 3$. We will provide a proof of Theorem 3.2 in Subsection 3.4. Our strategy is as follows. First, we calculate the extremal rays of the two-dimensional cone P^{A_n} . Second, we give a description of Σ^{A_n} using symmetry reduction. Third, we show that when *n* is even then one of the extremal rays is not a sum of squares, while for odd *n* both extremal rays are sum of squares.

To motivate the fundamental difference between S_n -invariant and A_n -invariant nonnegative quartics we start with an overview on nonnegativity in Subsection 3.1.

3.1. Global nonnegativity versus nonnegativity on U_n . We motivate the subtle difference between globally nonnegative forms and forms nonnegative on U_n in the vector space $\langle p_2^2, p_4 \rangle_{\mathbb{R}}$. For $n \ge 3$, the vector space of symmetric (n + 1)-variate quartics is five dimensional and is spanned by the following products of power sum polynomials

$$p_1^4, p_2 p_1^2, p_3 p_1, p_2^2, p_4$$

For any $n \ge 3$, there exist (n + 1)-variate symmetric quartic psd forms that are not sums of squares [11]. For instance, there exists the following uniform example [2]

$$\mathfrak{f}_n := 4p_1^4 - 5p_2p_1^2 - \frac{139}{20}p_3p_1 + 4p_2^2 + 4p_4$$

which is always nonnegative but never a sum of squares for any number of variables ≥ 4 . Note however, that restricting to the subspace U_n gives $4(p_2^2 + p_4)$. Thus f_n is a sum of squares modulo the ideal (p_1) . The form f_n can therefore not be used as a counter example for the reflection groups of type A.

We show that any psd form in the vector space $\langle p_2^2, p_4 \rangle_{\mathbb{R}}$ is a sum of squares. The proposition follows also from the nonnegativity versus sums of squares classification in type *B* [12]. The quartics result for type *B* was first observed by Choi, Lam and Reznick.

Proposition 3.3. Let $f = ap_2^2 + bp_4$ be a nonnegative (n + 1)-ary symmetric form, where $a, b \in \mathbb{R}$. Then *f* is a sum of squares.

Proof. Since *f* is an even symmetric form, nonnegativity of *f* is equivalent to nonnegativity of $ap_1^2 + bp_2$ on the probability simplex $\Delta_n := \{x \in \mathbb{R}_{\geq 0}^{n+1} : \sum_{i=1}^{n+1} x_i = 1\}$. Then $p_1 = 1$ and $\frac{1}{n+1} \le p_2 \le 1$ [1]. In particular, we need to distinguish three cases depending on the sign of *a*. We can suppose $p_2 = 1$ and $\frac{1}{n+1} \le p_4 \le 1$.

- (1) If a = 0 we have bp_4 nonnegative implies $b \ge 0$ and thus we have a sum of squares.
- (2) If a > 0 we suppose without loss of generality that a = 1 and we have $1 + bp_2 \ge 0$ on Δ_n which implies $b \ge -1$. However, the form

$$p_2^2 - p_4 = p_4 + 2\sum_{i < j} \mathbf{x}_i^2 \mathbf{x}_j^2 - p_4 = 2\sum_{i < j} \mathbf{x}_i^2 \mathbf{x}_j^2$$

on the boundary of the psd cone is clearly sos.

(3) If a < 0 we suppose a = -1 and have $-1 + bp_2 \ge 0$ on Δ_n implies $b \ge n + 1$. The form $(n + 1)p_4 - p_2^2$ on the boundary of the psd cone is a sum of squares since

$$(n+1)p_4 - p_2^2 = np_4 - 2\sum_{i < j} \mathbf{x}_i^2 \mathbf{x}_j^2 = \sum_{i < j} (\mathbf{x}_i^2 - \mathbf{x}_j^2)^2.$$

This subtle but important difference of nonnegativity on \mathbb{R}^{n+1} and on U_n has important structural consequences regarding A_n -invariant sums of squares.

3.2. **PSD** A_n **invariant quartics.** A symmetric (n + 1)-variate polynomial which is nonnegative on the linear subspace U_n must not necessarily be globally nonnegative (see e.g. the polynomial G_n for any n and F_n for any even n in Lemma 3.5). Since we are considering homogeneous invariant polynomials we have by biduality of convex cones ([3], Lemma 4.18.) the following Lemma.

Lemma 3.4. The boundary of P^{A_n} consists of the forms $f = a \cdot p_2^2 + b \cdot p_4$ for which there exists $0 \neq z \in U_n$ such that f(z) = 0.

In analogy to the proof of Proposition 3.3 we will analyse the maximum and the minimum of p_4 on the semialgebraic set defined by $p_2 = 1$ and $p_1 = 0$. **Lemma 3.5.** The extremal (n + 1)-ary A_n -invariant psd quartics are

$$G_n := p_{2,2} - \frac{1}{\beta} p_4 \text{ and } F_n := -p_{2,2} + \frac{1}{\alpha} p_4,$$

where $\beta = \frac{1 - n + n^2}{n + n^2}$ and $\alpha = \begin{cases} \frac{1}{n + 1} & \text{if } n \text{ is odd,} \\ \frac{4 + 2n + n^2}{2n + 3n^2 + n^3} & \text{if } n \text{ is even.} \end{cases}$

It follows directly from Proposition 3.3 that all of these extremal forms, but F_n when n is odd, are not globally nonnegative and cannot be a sum of squares in the polynomial ring $\mathbb{R}[\mathbf{x}]$.

Proof of Lemma 3.5. Since the quartics are homogeneous it is sufficient to analyse the minimum and maximum value of p_4 on $\mathbb{S}^n \cap U_n$. We have $p_1 = 0$ and $p_2 = 1$. This translates to the polynomial optimization problems

$$\min_{x \in \mathbb{R}} \pm p_4$$

s.t. $p_1 = 0$
 $p_2 = 1$

By a variant of Timofte's half degree principle [19, Theorem 1.1] the extremas are attained at a point with at most 2 different coordinates. The equality constraints transfer to the two equations

$$lt + (n + 1 - l)s = 0$$

$$lt^{2} + (n + 1 - l)s^{2} = 1$$

where $0 \le l \le n+1$ is an integer and $s, t \in \mathbb{R}$ are real numbers. We observe that $l \notin \{0, n+1\}$ which implies $1 \le l \le n$. For given integers *l* and *n* the equations provide unique solutions for *s* and *t* up to sign. However, inserting the solution in p_4 is independent of the signs of the coordinates and we have

$$p_4(\underbrace{t,\ldots,t}_{l \text{ times}},\underbrace{s,\ldots,s}_{n+1-l \text{ times}}) = \frac{(n+1)^2 - 3l(n+1) + 3l^2}{(n+1)(n+1-l)l}$$

~

(1

For the claim on the extremality of F_n we are left with verifying

$$\min_{1 \le l \le n, l \in \mathbb{Z}} \frac{(n+1)^2 - 3l(n+1) + 3l^2}{(n+1)(n+1-l)l} = \begin{cases} \frac{1}{n+1} & \text{if } n \text{ is odd} \\ \frac{4+2n+n^2}{2n+3n^2+n^3} & \text{if } n \text{ is even} \end{cases}$$

which we do in Lemmas 3.7 and 3.8, and verify

$$\max_{1 \le l \le n, l \in \mathbb{Z}} \frac{(n+1)^2 - 3l(n+1) + 3l^2}{(n+1)(n+1-l)l} = \frac{1-n+n^2}{n+n^2}$$

to prove that G_n is extremal. This is Lemma 3.6.

3.2.1. Verification of the extremality of G_n .

Lemma 3.6. For all
$$n \ge 3$$
 we have $\max_{1 \le l \le n, l \in \mathbb{Z}} \frac{(n+1)^2 - 3l(n+1) + 3l^2}{(n+1)(n+1-l)l}$ is attained at $l = 1$
and $l = n$, and equals $\frac{1-n+n^2}{n+n^2}$.

.

Proof. We calculate

$$\frac{(n+1)^2 - 3(n+1) + 3}{(n+1)n} \geq \frac{(n+1)^2 - 3l(n+1) + 3l^2}{(n+1)(n+1-l)l}$$

$$\Leftrightarrow (-1+l)(n-l)(n+1)^2 \geq 0.$$

Note, for $1 \le l \le n$ the inequality is tight when $l \in \{1, n\}$ and otherwise strict.

3.2.2. Verification of the extremality of F_n .

Lemma 3.7. We have

$$\min_{1 \le l \le n} \frac{(n+1)^2 - 3l(n+1) + 3l^2}{(n+1)(n+1-l)l} = \frac{1}{n+1}$$

and

$$\min_{\leq l \leq n, l \in \mathbb{Z}} \frac{(n+1)^2 - 3l(n+1) + 3l^2}{(n+1)(n+1-l)l} > \frac{1}{n+1}$$

if and only if *n* is even.

Proof. For $1 \le l \le n$ we have

$$\frac{(n+1)^2 - 3l(n+1) + 3l^2}{(n+1)(n+1-l)l} - \frac{1}{n+1} \ge 0$$

$$\iff \frac{(n+1)^2 - 3l(n+1) + 3l^2}{(n+1)(n+1-l)l} - \frac{(n+1-l)l}{(n+1)(n+1-l)l} \ge 0$$

$$\iff (n+1)^2 - 3l(n+1) + 3l^2 - (n+1-l)l \ge 0$$

$$\iff (n+1-2l)^2 \ge 0$$

The last inequality is tight on integer values $1 \le l \le n$ if and only if n + 1 is even.

Lemma 3.8. If *n* is even, then
$$\min_{\substack{1 \le l \le n, l \in \mathbb{Z}}} \frac{(n+1)^2 - 3l(n+1) + 3l^2}{(n+1)(n+1-l)l}$$
 is attained at $l = \frac{n}{2}$ and $l = \frac{n}{2} + 1$, and equals $\frac{4 + 2n + n^2}{2n + 3n^2 + n^3}$.

Proof. Evaluating $\frac{(n+1)^2 - 3l(n+1) + 3l^2}{(n+1)(n+1-l)l}$ at $l = \frac{n}{2}$ and $l = \frac{n+2}{2}$ gives $\frac{4+2n+n^2}{2n+3n^2+n^3}$.

Moreover, the denominator of

$$\frac{(n+1)^2 - 3l(n+1) + 3l^2}{(n+1)(n+1-l)l} - \frac{4+2n+n^2}{2n+3n^2+n^3} = \frac{(n+1)(4l^2 - 4l(n+1) + n(n+2))}{l(n+1-l)n(n+2)}$$

is strictly positive for all $1 \le l \le n$. The numerator is also nonnegative since

$$4l^2 - 4l(n+1) + n(n+2) = (2l - (n+1))^2 - 1 \ge 0$$

because n + 1 is odd.

3.3. **SOS** A_n -invariant quartics. Given the action of a reflection group, representation theory and invariant theory can be applied to effectively describe the invariant sums of squares cone. We briefly sketch the symmetry reduction for sums of squares invariant by a reflection group. More details can be found in [4, 9, 10, 14]. A reflection group *G* acts on the vector space $\mathbb{R}[\mathbf{x}]_d$ of all (n + 1)-variate forms of degree *d* giving it the structure of a *G*-module. We can decompose every *G*-module into a direct sum of its irreducible sub-modules to obtain its *isotypic decomposition*. Given an isotypic decomposition one constructs a *symmetry adapted basis*, which can be used to understand the invariant sums of squares of elements in $\mathbb{R}[\mathbf{x}]_{2d}$. We outline this in the following.

6

П

П

First, we note that there is a natural projection onto the invariant part of $\mathcal{F}_{n,d}$ via the so called *Reynolds-Operator* of the group *G*:

$$\mathcal{R}_G : \mathbb{R}[\mathbf{x}]_d \to \mathbb{R}[\mathbf{x}]_d^G, \ f \mapsto \frac{1}{|G|} \sum_{\sigma \in G} \sigma f.$$

Suppose that we have

$$\mathbb{R}[\mathbf{x}]_d \simeq \bigoplus_{j=1}^{\ell} \eta_j \mathcal{V}_j$$

is the isotypic decomposition of the *G* action on $\mathbb{R}[\mathbf{x}]_d$, i.e. \mathcal{V}_j are pairwise non-isomorphic irreducible *G*-modules and each occurs with multiplicity $\eta_j \in \mathbb{N}$ in $\mathbb{R}[\mathbf{x}]_d$. A symmetry adapted basis is a list

$$\{f_{11},\ldots,f_{1\eta_1},f_{21},\ldots,f_{\ell\eta_\ell}\}$$

with the property that for every *j* there are *G*-equivariant homomorphisms ϕ_{ji} which map f_{j1} to f_{ji} for all $1 \le i \le \eta_j$, and furthermore that the orbit of each f_{ji} spans an irreducible *G*-module isomorphic to \mathcal{V}_j and the set of all orbits of all f_{ji} spans $\mathbb{R}[\mathbf{x}]_d$. Given a symmetry adapted basis we can construct matrix polynomials

$$B_j := \left(\mathcal{R}_G(f_{ji_1}f_{ji_2})_{1 \le i_1, i_2 \le \eta_j} \text{ for } 1 \le j \le \ell. \right)$$

With these notations we have the following (see [9, Theorem 2.6]):

Proposition 3.9. Let $f \in \mathbb{R}[\mathbf{x}]_{2d}^G$ be an invariant form. Then f is a sum of squares if and only if there exists positive semidefinite matrices A_1, \ldots, A_ℓ such that

$$f = \sum_{j=1}^{\ell} \operatorname{Tr}(A_j B_j),$$

where the matrix polynomials B_j are constructed from a symmetry adapted bases of $\mathbb{R}[\mathbf{x}]_d$ as defined above.

Note that calculating an isotypic decomposition of $\mathbb{R}[\mathbf{x}]_d$ and a symmetry adapted basis can in principle be done with linear algebra (see [20]). For the case of finite groups Hubert and Bazan [15] constructed an algorithm to calculate equivariants which allows for an effective determination of symmetry adapted basis for all degrees. In the case when $G \in \{A_{n-1}, S_n, B_n, D_n\}$ so-called *higher Specht polynomials* can be used and their construction is completely combinatorial [9, 17].

We denote by \mathbb{S}^{λ} the Specht module associated with a partition λ .

Lemma 3.10. For $n \ge 3$, the S_{n+1} isotypic decomposition of $\mathbb{R}[\mathbf{x}]_2$ equals

$$\mathbb{R}[\mathbf{x}]_{2} = \langle p_{2} \rangle_{\mathbb{R}} \oplus \langle p_{1}^{2} \rangle_{\mathbb{R}} \oplus \langle p_{1}(\mathbf{x}_{i} - \mathbf{x}_{j}) : i < j \rangle_{\mathbb{R}}$$
$$\oplus \langle \mathbf{x}_{i}^{2} - \mathbf{x}_{j}^{2} : i < j \rangle_{\mathbb{R}} \oplus \langle (\mathbf{x}_{i} - \mathbf{x}_{j})(\mathbf{x}_{k} - \mathbf{x}_{l}) : \#\{i, j, k, l\} = 4 \rangle$$
$$2 \cdot \mathbb{S}^{(n)} \oplus 2 \cdot \mathbb{S}^{(n-1,1)} \oplus \mathbb{S}^{(n-2,2)}$$

The proof is fully computational and we calculate a symmetry adapted basis based on higher Specht polynomials [17] (and refer to [9] for details).

We apply the Reynolds-Operator of the symmetric group S_{n+1} to pairwise products of equivariants of the isotypic decomposition which do not use p_1 , since we consider sum of squares in $\mathbb{R}[\mathbf{x}]$ modulo the ideal (p_1) .

Lemma 3.11. For $n \ge 4$, we have

$$\begin{aligned} \mathcal{R}_{S_{n+1}}(p_2^2) &= p_2^2, \\ \mathcal{R}_{S_{n+1}}((\mathbf{x}_1^2 - \mathbf{x}_2^2)^2) &= \frac{2}{n}p_4 - \frac{2}{(n+1)n}p_2^2, \text{ and} \\ \mathcal{R}_{S_{n+1}}((\mathbf{x}_1 - \mathbf{x}_2)^2(\mathbf{x}_3 - \mathbf{x}_4)^2) &= \\ &= \frac{4(p_1^4 + 3p_2^2 - 4p_3p_1 + (n+1)^2(p_2^2 - p_4) + (n+1)(-2p_2p_1^2 - 3p_2^2 + 4p_3p_1 + p_4))}{(n+1)n(n-1)(n-2)}. \end{aligned}$$

Proof. We calculate the polynomials p_2^2 , $(\mathbf{x}_1^2 - \mathbf{x}_2^2)^2$, $(\mathbf{x}_1 - \mathbf{x}_2)^2(\mathbf{x}_3 - \mathbf{x}_4)^2$ and apply the Reynolds-Operator $\mathcal{R}_{S_{n+1}}$. We then obtain the right hand side of the equation in the basis of monomial symmetric polynomials. We then use the package Symmetric Function in Sage to obtain a representation in terms of the power sum polynomials.

Lemma 3.12. If $f \in \mathbb{R}[\mathbf{x}]$ is a A_n -invariant sum of squares quartic modulo the ideal (p_1) then

$$f = a(p_4 - \frac{1}{n+1}p_2^2) + b((1-n+n^2)p_2^2 - n(1+n)p_4) + p_1 \cdot g$$

for some $a, b \ge 0$ and $g \in \mathbb{R}[x]$.

Proof. Since $f \in \mathbb{R}[\mathbf{x}]$ is A_n -invariant, we can apply the Reynolds-Operator $\mathcal{R}_{A_n} = \mathcal{R}_{S_{n+1}}$ to $g_1^2 + \ldots + g_m^2$ and consider $\mathcal{R}_{S_{n+1}}(g_1^2 + \ldots + g_m^2) \mod p_1$ which has to be of the form

$$\lambda_1 p_2^2 + \lambda_2 (p_4 - \frac{1}{n+1} p_2^2) + \lambda_3 ((1-n+n^2) p_2^2 - n(1+n) p_4)$$

for some $\lambda_1, \lambda_2, \lambda_3 \ge 0$, by Lemma 3.11 and the discussion above. We have

$$(1-n)^2 p_2^2 = n(n+1)(p_4 - \frac{1}{n+1}p_2^2) + ((1-n+n^2)p_2^2 - n(1+n)p_4)$$

which proves the claim.

3.4. Proof of Theorem 3.2. We are ready to prove Theorem 3.2

Proof of Theorem 3.2. There are three statements that we want to show. First, the polynomial $G_n \in \mathbb{R}[\mathbf{x}]$ is a sum of squares modulo (p_1) for all $n \ge 3$. Second, for $n \ge 4$ odd the polynomial $F_n \in \mathbb{R}[\mathbf{x}]$ is a sum of squares modulo (p_1) . Third, for $n \ge 3$ even the polynomial $F_n \in \mathbb{R}[\mathbf{x}]$ is not a sum of squares modulo (p_1)

(1) We have

$$G_n = p_2^2 - \frac{n+n^2}{1-n+n^2} p_4$$

= $\frac{1}{1-n+n^2} ((1-n+n^2)p_2^2 - n(1+n)p_4)$

which shows that G_n is a sum of squares modulo (p_1) .

- (2) If $n \ge 4$ is odd we have F_n is a sum of squares by Proposition 3.3. This is, because α equals the global minimum of p_4 on $p_2 = 1$ and we have seen that the corresponding polynomial is a sum of squares.
- (3) For even $n \ge 4$ we have $F_n = -p_2^2 + \frac{2n+3n^2+n^3}{4+2n+n^2}p_4$. We suppose that F_n is a sum of squares modulo (p_1) . We must have

$$-p_2^2 + \frac{2n+3n^2+n^3}{4+2n+n^2}p_4 = a(p_4 - \frac{1}{n+1}p_2^2) + b((1-n+n^2)p_2^2 + (-n-n^2)p_4)$$

for some $a, b \ge 0$. Comparing the coefficients implies

$$b = -\frac{4}{4-6n+n^2+n^4}$$

which is a contradiction.

References

- [1] J. Acevedo, G. Blekherman, S. Debus, and C. Riener. *The wonderful geometry of the vandermonde map*, arXiv:2303.09512, 2023.
- [2] J. Acevedo, G. Blekherman, S. Debus, and C. Riener. *At the limit of the symmetric psd and sos cones*, in preparation.
- [3] G. Blekherman, P. A. Parrilo, R. R. Thomas. Semidefinite optimization and convex algebraic geometry, (2012) SIAM 2012.
- [4] G. Blekherman and C. Riener. Symmetric nonnegative forms and sums of squares, Discrete and Computational Geometry, 65:764–799, 2021.
- [5] L. Blum. Complexity and real computation, Springer Science & Business Media, 1998.
- [6] C. Chevalley. Invariants of finite groups generated by reflections, Amer. J. Math., 77:778-782, 1955.
- [7] H. S. M. Coxeter. Discrete groups generated by reflections, Ann. of Math. (2), 35(3):588-621, 1934.
- [8] H. S. M. Coxeter. The Complete Enumeration of Finite Groups of the Form R2i= (RiRj)kij = 1, J. London Math. Soc., 10(1):21–25, 1935.
- [9] S. Debus, C. Riener. *Reflection groups and cones of sums of squares*, Journal of Symbolic Computation, 119:112–144, 2023.
- [10] K. Gatermann, P. A. Parrilo. Symmetry groups, semidefinite programs, and sums of squares, Journal of Pure Applied Algebra ,192, (1 - 3), 95–128, 2004.
- [11] C. Goel, S. Kuhlmann, B. Reznick. On the Choi-Lam analogue of Hilbert's 1888 theorem for Symmetric forms, Linear Algebra and its Applications, 496, 114-120, 2016.
- [12] C. Goel, S. Kuhlmann, B. Reznick. The analogue of Hilbert's 1888 theorem for Even Symmetric Forms, Journal of Pure and Applied Algebra, 221, 1438-1448, 2017.
- [13] D. Hilbert. Über die Darstellung definiter Formen als Summe von Formenquadraten, Math. Ann., 32 (1888), 342-350; Ges. Abh. 2, 154-161, Springer, Berlin, reprinted by Chelsea, New York, 1981.
- [14] A. Heaton, S. Hosten, and I. Shankar. *Symmetry adapted Gram spectrahedra*, SIAM Journal on Applied Algebra and Geometry 5(1): 140-164, 2021.
- [15] E. Hubert and E. Rodriguez Bazan. Algorithms for fundamental invariants and equivariants of finite groups, Mathematics of Computation 91(337): 2459-2488, 2022.
- [16] M. Marshall. Positive Polynomials and Sum of Squares, Vol. 146, Mathematical Surveys and Monographs, AMS, 2008.
- [17] H. Morita and H.-F. Yamada. Higher Specht polynomials for the complex reflection group G (r, p, n), Hokkaido mathematical journal 27(3): 505-515, 1998.
- [18] T. S. Motzkin. The arithmetic-geometric inequality, in Inequalities, Oved Shisha (ed.) Academic Press, 205-224, 1967.
- [19] C. Riener. On the degree and half-degree principle for symmetric polynomials, Journal of Pure and Applied Algebra, 216 (4), 850-856, 2012.
- [20] J.-P. Serre. Linear representations of finite groups, Vol. 42. New York: Springer, 1977.
- [21] G. C. Shephard and J. A. Todd. Finite unitary reflection groups, Canad. J. Math., 6:274–304, 1954.

FACULTY OF MATHEMATICS, TECHNISCHE UNIVERSITÄT CHEMNITZ, GERMANY

 ${\it Email\ address:\ sebastian.debus@mathematik.tu-chemnitz.de}$

DEPARTMENT OF BASIC SCIENCES, INDIAN INSTITUTE OF INFORMATION TECHNOLOGY NAGPUR, INDIA

Email address: charugoel@iiitn.ac.in

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF KONSTANZ, GERMANY

Email address: salma.kuhlmann@uni-konstanz.de

DEPARTMENT OF MATHEMATICS AND STATISTICS, UIT THE ARCTIC UNIVERSITY OF NORWAY, NORWAY

Email address: cordian.riener@uit.no