

# Understanding perturbative and non-perturbative contributions to jets at RHIC

Isaac Mooney <sup>1,2</sup> 

<sup>1</sup> Wright Laboratory, Yale University, New Haven, CT; isaac.mooney@yale.edu

<sup>2</sup> Brookhaven National Laboratory, Upton, NY

**Abstract:** Jets, as collections of multi-scale objects, allow for insight into perturbative (high-momentum) processes, but gaining an understanding of the non-perturbative structure within jets such as hadronization effects and the underlying event has been more difficult. In observables sensitive to these non-perturbative effects, a large discrepancy is observed between partonic calculations and experimental data at RHIC. Grooming techniques such as SoftDrop reduce the impact of these non-perturbative effects and isolate the hard radiation associated with the hard scattering for more direct comparison to perturbative calculations. By analogy, CollinearDrop can be used to isolate non-perturbative contributions from a different region of the emission phase space, or the Lund plane, although no measurement of a CollinearDrop observable has yet been published. In these proceedings, we review recent progress on experimentally accessing the perturbative and non-perturbative contributions to jets and their substructure, and discuss potential future measurements at RHIC as a road map toward precision QCD at the EIC.

**Keywords:** QCD; jet; perturbative; RHIC

## 1. Introduction

Ultrarelativistic ion collisions typically produce many particles with low transverse momentum ( $p_T$ ) with respect to the incoming beams. Rarely, however, high-momentum transfers between quarks and gluons (*partons*) in the incoming nucleons occur, leading to collimated groupings of particles with high total  $p_T$ , called jets. Jets are multi-scale objects which evolve from the initial high-momentum (*hard*) scattered parton to lower energy quark and gluon daughters through successive radiation, until reaching virtualities near the hadronization scale  $\Lambda_{\text{QCD}}$ , at which point partons recombine into hadrons. Perturbative quantum chromodynamics (pQCD) calculations are applicable for high energy processes, which correspond mostly to the parton shower component occurring early in jet evolution. By contrast, soft or non-perturbative (np)QCD processes are generally not calculable from first principles. The technique of QCD on a spacetime lattice is an exception, although modeling dynamics is computationally intensive, and it is therefore inapplicable to jet studies. Partly due to difficulty theoretically describing the entirety of QCD jets, approaches were developed to avoid npQCD radiation's contribution to jet substructure observables. At the Relativistic Heavy Ion Collider (RHIC) the jet spectrum is softer than at the Large Hadron Collider (LHC), such that non-perturbative effects are more important. In Sec. 2 of these proceedings, we focus on recent measurements at RHIC using techniques which minimize the npQCD contribution. In Sec. 3, we highlight measurements and observables which either improve our understanding of the npQCD contribution within jets, or which would benefit from our improved understanding of it. Finally, in Sec. 4, we give an outlook to future progress in the study of jets in their entirety.

## 2. pQCD

The most commonly used algorithmic approach to reduction of soft, wide-angle radiation in a jet is called SoftDrop (SD) grooming [1]. Although the explicit goal is to minimize effects unrelated to a hard scattering's final state (e.g. pileup, initial state radiation, and underlying event), it is also very effective at reducing the non-perturbative

arXiv:2402.10832v1 [hep-ph] 16 Feb 2024




**Citation:** Mooney, I. Understanding perturbative and non-perturbative contributions to jets at RHIC. *Preprints* 2023, 1, 0. <https://doi.org/>



**Copyright:** © 2023 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

contribution, including hadronization<sup>1</sup>. Recent results from proton-proton ( $p+p$ ) collisions at RHIC have demonstrated good agreement with Monte Carlo (MC) event generators such as PYTHIA [3] (Figure 1a) and confirmed the assumption that SD-groomed observables are more directly comparable to perturbative, parton-level calculations (Figure 1b) although there remain discrepancies especially at small jet radii,  $R$  (Figure 1c), since the physical scales of jet production and grooming go as  $p_T R$  and  $z_{\text{cut}} p_T R^2$ , respectively, and these are closer to  $\Lambda_{\text{QCD}}$  for smaller radius.



**Figure 1.** (a) Preliminary PHENIX measurement [4] of momentum sharing fraction of the first hard split in the clustering history,  $z_g$ . Data are compared to hadron-level MC event generator PYTHIA which demonstrates good agreement. (b) STAR measurement [5] of the invariant mass (four-momentum sum of constituents) of the jet,  $M_{(g)}$ , before (red) and after (blue) SD grooming. Data are compared to a next-to-leading-logarithm accuracy (NLL) pQCD calculation (green) which qualitatively agrees with the groomed result for large jet  $p_T$  and  $R$ . (c) STAR measurement [6] of the opening angle of the first hard split,  $R_g$ . Data are also compared to an NLL calculation, which again compares better at large jet  $p_T$  and  $R$ . Data in all cases are from  $p+p$  collisions.


Another way to reduce the non-perturbative contribution within jets is the subjet method [7]. This involves reclustering a jet into subjets of smaller radius,  $r < R$ , at which point observables similar to those mentioned above (e.g. momentum sharing fraction,  $z$ , opening angle  $R$  or  $\theta$ , etc.) can be constructed between subjets rather than between prongs. One advantage compared to similar studies of hadron momentum distributions within jets is that parton-to-hadron fragmentation functions, which are non-perturbative, are not involved in the theoretical calculations [8]. From an experimental perspective, they are also useful because they reduce background contamination in heavy-ion (A+A) collisions [9].

### 3. npQCD

Although first-principles theoretical comparisons in the non-perturbative regime are often unavailable, this phase space region still deserves study. Widely used MC event generators such as PYTHIA include phenomenological modeling of hadronization and other npQCD processes, which requires tuning a parameter set to generate agreement with a large subset of collider data. PYTHIA 8.3's [3] default tune, Monash [10], did not include RHIC data which can create discrepancies with data even if the underlying perturbative physics mechanisms (matrix elements, parton shower, etc.) are modeled accurately. Specifically, subleading partonic interactions (MPI) contribute to the regularization of the partonic cross section at low- $p_T$ , controlled in PYTHIA by a parameter called  $p_{T,0}$  which enters the hard cross section as  $\hat{\sigma} = (p_T^2 + p_{T,0}^2)^{-1}$ . The relation  $p_{T,0} = p_{T,0}^{\text{ref}} (\sqrt{s}/\sqrt{s_{\text{ref}}})^{\text{ecmPow}}$  is used to extrapolate from a reference center-of-mass energy,  $\sqrt{s}$ , to another, where  $\text{ecmPow}$  is a tunable parameter. Models tuned with LHC data may fail to describe RHIC data, over an order of magnitude lower in energy, due to this simplistic extrapolation. Adjusting  $p_{T,0}^{\text{ref}}$  and  $\text{ecmPow}$  among other MPI-related parameters as part of a new tune of PYTHIA 8.3 [11] improved agreement with RHIC data, and with LHC data in some phase space regions.

In addition to improving models such that current data can be described better, observables which increase sensitivity to soft radiation have also been proposed and

investigated. One such example is CollinearDrop (CD) [12], which can be considered roughly as the complement of SoftDrop, mentioned in Sec. 2. Instead of grooming away the soft, wide-angle radiation, it is effectively the collinear radiation which is groomed away<sup>3</sup>. This anti-SD grooming results in a phase space with an enhanced non-perturbative contribution [13]. Beneficially from an experimental point of view, it is also relatively simple to obtain a CD observable if the ungroomed and groomed observables have already been measured, as it is just the difference between the two. The STAR Collaboration has a recent preliminary measurement of the CD-groomed jet mass [14] (Figure 2).



**Figure 2.** Preliminary STAR measurement of CD jet mass in  $p+p$  collisions. Data are compared to MC event generators PYTHIA and HERWIG which exhibit qualitative agreement with the data.

Another way of accessing the soft phase space within jets is to examine subsequent branches in the splitting tree [15], beyond the first hard split as defined by SD. Since the jet is evolving with subsequent splittings toward  $\Lambda_{\text{QCD}}$ , there is an expected larger npQCD contribution in later splits. In a STAR preliminary measurement, a behavior consistent with – although not conclusively caused by – this evolution was observed [16].

Finally, in  $p+p$  collisions, another promising avenue is the use of correlators of energy flow operators in the small angle limit, or more simply *energy correlators* [17–19]. Because correlations of jet constituents separated by small (large) angular distance correspond to splittings occurring at late (early) times, the energy correlators exhibit an approximate separation between perturbative and non-perturbative regimes with no need to recluster the jets or remove npQCD contributions as in the case of the SD grooming procedure. STAR has recently made a preliminary measurement of the two-point energy correlator (EEC), which agrees well in the perturbative regime with an NLL pQCD prediction, and in the non-perturbative regime with a model assuming non-interacting hadrons [20].

In heavy-ion collisions, which produce a thermal medium called the quark-gluon plasma (QGP), often parton-medium interactions are calculated analytically in the weakly coupled limit. However, recent work (see e.g. Ref. [21]) has started to incorporate multiple soft interactions with the medium into the calculation of the momentum broadening kernel. This provides a more realistic description of splitting rates within the medium, which in turn yields an accurate number of color sources for interaction with the medium.

#### 4. Discussion

This diverse set of calculations, phenomenological modeling, and experimental results improves our understanding of the full structure of jets primarily by separating perturbative and non-perturbative effects more cleanly and enhancing the contribution of one or the other. Extensions to these works are ongoing and planned, and new avenues are being explored. For example, energy correlators may be measured at higher orders [19]; in heavy-ion collisions [22]; and separated by the constituents' charge [23]. The last point is part of an effort to examine specifically non-perturbative fragmentation into hadrons, and has motivated measurement of charge correlators between the leading hadrons within jets [24].

**Funding:** This work was funded by the Office of Nuclear Physics of the U.S. Department of Energy

under award number DE SC004168 and BNL/DOE-424803.

**Data Availability Statement:** Most data mentioned above are available at <https://www.hepdata.net/>.

**Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

## Notes

- <sup>1</sup> There is still a small npQCD contribution to e.g. the groomed jet mass in the form of power corrections in the large mass region, with a larger contribution at smaller mass [2].
- <sup>2</sup>  $z_{\text{cut}}$  is the momentum fraction threshold above which a given split in the jet shower is considered hard, and is not dropped.
- <sup>3</sup> A grooming of the softest, widest-angle radiation can also be performed to reduce e.g. the effect of pile-up.

## References

1. Larkoski, A.J.; Marzani, S.; Soyez, G.; Thaler, J. Soft Drop. *JHEP* **2014**, *05*, 146.
2. Hoang, A.H.; Mantry, S.; Pathak, A.; Stewart, I.W. Nonperturbative Corrections to Soft Drop Jet Mass. *JHEP* **2019**, *12*, 002.
3. Bierlich, C.; et al. A comprehensive guide to the physics and usage of PYTHIA 8.3. *SciPost Phys. Codeb.* **2022**, *2022*, 8.
4. Belmont, R. Recent Results from PHENIX. *EPJ Web Conf.* **2023**, *276*, 06019.
5. Abdallah, M.; et al. Invariant Jet Mass Measurements in  $pp$  Collisions at  $\sqrt{s} = 200$  GeV at RHIC. *Phys. Rev. D* **2021**, *104*, 052007.
6. Adam, J.; et al. Measurement of groomed jet substructure observables in p+p collisions at  $\sqrt{s} = 200$  GeV with STAR. *Phys. Lett. B* **2020**, *811*, 135846.
7. Dai, L.; Kim, C.; Leibovich, A.K. Fragmentation of a Jet with Small Radius. *Phys. Rev. D* **2016**, *94*, 114023.
8. Kang, Z.B.; Ringer, F.; Waalewijn, W.J. The Energy Distribution of Subjets and the Jet Shape. *JHEP* **2017**, *07*, 064.
9. Abdallah, M.S.; et al. Differential measurements of jet substructure and partonic energy loss in Au+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV. *Phys. Rev. C* **2022**, *105*, 044906.
10. Skands, P.; Carrazza, S.; Rojo, J. Tuning PYTHIA 8.1: the Monash 2013 Tune. *Eur. Phys. J. C* **2014**, *74*, 3024.
11. Aguilar, M.R.; Chang, Z.; Elayavalli, R.K.; Fatemi, R.; He, Y.; Ji, Y.; Kalinkin, D.; Kelsey, M.; Mooney, I.; Verkest, V. pythia8 underlying event tune for RHIC energies. *Phys. Rev. D* **2022**, *105*, 016011.
12. Chien, Y.T.; Stewart, I.W. Collinear Drop. *JHEP* **2020**, *06*, 064.
13. Stewart, I.W.; Yao, X. Pure quark and gluon observables in collinear drop. *JHEP* **2022**, *09*, 120.
14. Song, Y. Measurement of CollinearDrop jet mass and its correlation with SoftDrop groomed jet substructure observables in  $\sqrt{s} = 200$  GeV  $pp$  collisions by STAR **2023**. [[arXiv:nucl-ex/2307.07718](https://arxiv.org/abs/2307.07718)].
15. Frye, C.; Larkoski, A.J.; Thaler, J.; Zhou, K. Casimir Meets Poisson: Improved Quark/Gluon Discrimination with Counting Observables. *JHEP* **2017**, *09*, 083.
16. Kunnawalkam Elayavalli, R. Measurement of splittings along a jet shower in  $\sqrt{s} = 200$  GeV  $pp$  collisions at STAR. *SciPost Phys. Proc.* **2022**, *8*, 059.
17. Basham, C.L.; Brown, L.S.; Ellis, S.D.; Love, S.T. Energy Correlations in electron - Positron Annihilation: Testing QCD. *Phys. Rev. Lett.* **1978**, *41*, 1585.
18. Chen, H.; Moul, I.; Zhang, X.; Zhu, H.X. Rethinking jets with energy correlators: Tracks, resummation, and analytic continuation. *Phys. Rev. D* **2020**, *102*, 054012.
19. Komsike, P.T.; Moul, I.; Thaler, J.; Zhu, H.X. Analyzing N-Point Energy Correlators inside Jets with CMS Open Data. *Phys. Rev. Lett.* **2023**, *130*, 051901.
20. Tamis, A. Measurement of Two-Point Energy Correlators Within Jets in  $pp$  Collisions at  $\sqrt{s} = 200$  GeV at STAR. In Proceedings of the 11th International Conference on Hard and Electromagnetic Probes of High-Energy Nuclear Collisions: Hard Probes 2023, 9 2023, [[arXiv:hep-ex/2309.05761](https://arxiv.org/abs/2309.05761)].
21. Moore, G.D.; Schlichting, S.; Schlusser, N.; Soudi, I. Non-perturbative phenomena in jet modification. *EPJ Web Conf.* **2022**, *258*, 05001.
22. Andres, C.; Dominguez, F.; Holguin, J.; Marquet, C.; Moul, I. A coherent view of the quark-gluon plasma from energy correlators. *JHEP* **2023**, *09*, 088.
23. Lee, K.; Moul, I. Energy Correlators Taking Charge **2023**. [[arXiv:hep-ph/2308.00746](https://arxiv.org/abs/2308.00746)].
24. Chien, Y.T.; Deshpande, A.; Mondal, M.M.; Serman, G. Probing hadronization with flavor correlations of leading particles in jets. *Phys. Rev. D* **2022**, *105*, L051502.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.